

5. 参考文献

- Aldenberg T, Slob W (1993) Confidence limits for hazardous concentrations based on logistically distributed NOEC toxicity data. *Ecotoxicology and Environmental Safety*, 25, 48–63.
- Aldenberg T, Jaworska JS (2000) Uncertainty of the hazardous concentration and fraction affected for normal species sensitivity distributions. *Ecotoxicology and Environmental Safety*, 46, 1–18.
- ANZECC (2000) Australian and New Zealand guidelines for fresh and marine water quality. Australian and New Zealand Environment and Conservation Council.
- Baird DJ, van den Brink PJ (2007) Using biological traits to predict species sensitivity to toxic substances. *Ecotoxicology and Environmental Safety*, 67, 296–301.
- Barnthouse LW, Munns WRM, Sorensen MT (2007) Population level ecological risk assessment. CRC Press.
- Campbell PJ, Arnold DJS, Brock TCM, Grandy NJ, Heger W, Heimbach F, Maund SJ, Streloke M (1999) Guidance document on higher-tier aquatic risk assessment for pesticides (HARAP). SETAC-Europe, Brussels, Belgium.
- Campbell KR, Bartell SM, Shaw JL (2000) Characterizing aquatic ecological risks from pesticides using a diquat dibromide case study. II. Approaches using quotients and distributions. *Environmental Toxicology and Chemistry*, 19, 760–774.
- Chang KH, Sakamoto M, Ha JY, Miyabara Y, Nakano S, Doi H, Hanazato T (2011) Response of the plankton community to herbicide application (triazine herbicide, simetryn) in a eutrophicated system: short-term exposure experiment using microcosms. *Limnology*, 12, 11–16.
- Chevre N, Loepfe C, Singer H, Stamm C, Fenner K, Escher BI (2006) Including mixtures in the determination of water quality criteria for herbicides in surface water. *Environmental Science and Technology*, 40, 426–435.
- de Wolf W, Sievbel-Sauer A, Lecloux A, Koch V, Holt M, Feijtel T, Comber M, Boeije G (2005) Mode of action and aquatic exposure thresholds of no concern. *Environmental Toxicology and Chemistry*, 24, 479–485.
- de Zwart, Posthuma L (2005) Complex mixture toxicity for single species and multiple species: proposed methodologies. *Environmental Toxicology and Chemistry*, 24, 2665–2676.

- de Zwart (2005) Ecological effects of pesticide use in the Netherlands: modeled and observed effects in the field ditch. *Integrated Environmental Assessment and Management*, 1, 123–134.
- Duboudin C, Ciffroy P, Magaud H (2004) Effects of data manipulation and statistical methods on species sensitivity distributions. *Environmental Toxicology and Chemistry*, 23, 489–499.
- Dyer SD, Versteeg DJ, Belanger SE, Chaney JG, Mayer FL (2006) Interspecies correlation estimates predict protective environmental concentrations. *Environmental Science and Technology*, 40, 3102–3111.
- ECB (2003) Technical guidance document on Risk Assessment. Part II Environmental risk assessment. European Chemical Bureau.
- ECETOC (2003) Aquatic hazard assessment II, Technical Report No. 91, European Centre for Ecotoxicology and Toxicology of Chemicals.
- ECOFRAM (1999) Ecological Committee on FIFRA Risk Assessment Methods (ECOFRAM) aquatic report.
- EFSA (2008a) Conclusion on the peer review of imidacloprid. *EFSA Scientific Report*, 148, 1–120.
- EFSA (2008b) Conclusion regarding the peer review of the pesticide risk assessment of the active substance etofenprox. *EFSA Scientific Report*, 213, 1–131.
- EFSA (2013a) Conclusion on the peer review of the pesticide risk assessment of the active substance [chlorantraniliprole]. *EFSA Journal*, 2013:11, 3143.
- EFSA (2013b) Guidance on tiered risk assessment for plant protection products for aquatic organisms in edge-of-field surface waters. *EFSA Journal*, 2013:11, 3290.
- EUFRAM (2006) Probabilistic approaches for assessing environmental risks of pesticides, Final report volume 1: Framework and worked examples, European FRAMework for probabilistic risk assessment of the environmental impacts of pesticides.
- Faust M, Altenburger R, Backhaus T, Blanck H, Bødeker W, Gramatica P, Hamer V, Scholze M, Vighi M, Grimme LH (2001) Predicting the joint algal toxicity of multicomponent s-triazine mixtures at low-effect concentrations of individual toxicants. *Aquatic Toxicology*, 56, 13–32.
- Faust M, Altenburger R, Backhaus T, Blanck H, Bødeker W, Gramatica P, Hamer V, Scholze M, Vighi M, Grimme LH (2003) Joint algal toxicity of 16 dissimilarly acting chemicals is predictable by the concept of independent action. *Aquatic Toxicology*, 63, 43–63.

- Forbes VE, Calow P (2002) Species sensitivity distributions revisited: a critical appraisal. *Human and Ecological Risk Assessment*, 8, 473–492.
- Freemark KE, MacQuarrie P, Swanson SM, Peterson H (1990) Development of guidelines for testing pesticide toxicity to nontarget plants for Canada. In: Wang W, Gorsuch JW, Lower WR (Eds), *Plants for toxicity assessment*. American Society for Testing and Materials, pp14–29.
- Furuhama A, Toida T, Nishikawa N, Aoki Y, Yoshioka Y, Shiraishi H (2010) Development of an ecotoxicity QSAR model for the KAshinhou Tool for Ecotoxicity (KATE) system, March 2009 version. *SAR QSAR Environmental Research*, 21, 403–413.
- Giddings JM, Biever RC, Annunziato MF, Hosmer AJ (1996) Effects of diazinon on large outdoor pond microcosms. *Environmental Toxicology and Chemistry*, 15, 618–629.
- Giddings JM, Hall LW, Solomon KR (2000) Ecological risks of diazinon from agricultural use in the Sacramento-San Joaquin River basins, California. *Risk Analysis*, 20, 545–572.
- Giddings JM, Solomon KR, Maund SJ (2001) Probabilistic risk assessment of cotton pyrethroids: II. Aquatic mesocosm and field studies. *Environmental Toxicology and Chemistry*, 20, 660–668.
- Hansen SR, Garton RR (1982) Ability of standard toxicity tests to predict the effects of the insecticide diflubenzuron on laboratory stream communities. *Canadian Journal of Fisheries and Aquatic Sciences*, 39, 1273–1288
- Hendley P, Holmes C, Kay S, Maund SJ, Travis KZ, Zhang M (2001) Probabilistic risk assessment of cotton pyrethroids: III. A spatial analysis of the Mississippi, USA, cotton landscape. *Environmental Toxicology and Chemistry*, 20, 669–678.
- Hobbs DA, Warne MSJ, Markich SJ (2005) Evaluation of criteria used to assess the quality of aquatic toxicity data. *Integrated Environmental Assessment and Management*, 1, 174–180.
- Insecticide Resistance Action Committee Modes of Action,
<http://www.iraac-online.org/modes-of-action/>.
- Fungicide Resistance Action Committee FRAC Code List 2015: Fungicides sorted by mode of action,
<http://www.frac.info/docs/default-source/publications/frac-code-list/frac-code-list-2015-finalC2AD7AA36764.pdf?sfvrsn=4>

- Herbicide Resistance Action Committee Classification of Herbicides According to Site of Action,
<http://www.hracglobal.com/pages/classificationofherbicidesiteofaction.aspx>
- Ippolito A, Todeschini R, Vighi M (2012) Sensitivity assessment of freshwater macroinvertebrates to pesticides using biological traits. *Ecotoxicology*, 21, 336–352.
- Iwafune T, Inao K, Horio T, Iwasaki N, Yokoyama A, Nagai T (2010) Behaviour of paddy pesticides and major metabolites in the Sakura River, Ibaraki, Japan. *Journal of Pesticide Science*, 35, 114–123.
- 鏡味麻衣子 (2008) ツボカビを考慮に入れた湖沼食物網の解析. *日本生態学会誌*, 58, 71–80.
- Kamo M, Naito W (2008) A novel approach to determining a population-level threshold in ecological risk assessment: A case study of zinc. *Human and Ecological Risk Assessment*, 14, 714–727.
- 環境省 生態影響試験結果一覧.
- 桐谷圭治 (2004) ただの虫を無視しない農業：生物多様性管理. 築地書館.
- 国立環境研究所 (1995) 水環境における化学物質の長期曝露による相乗的生態系影響に関する研究. *国立環境研究所特別研究報告*, SR-19-95.
- 国立環境研究所 (2012) 「平成 23 年度農薬の生物多様性への影響調査業務」報告書.
- Kroes R, Kleiner J, Renwick A (2005) The threshold of toxicological concern concept in risk assessment. *Toxicological Sciences*, 86, 226–230.
- Liess M, von der Ohe PC (2005) Analyzing effects of pesticides on invertebrate communities in streams. *Environmental Toxicology and Chemistry*, 24, 954–965.
- Maltby L, Blake N, Brock TCM, van den Brink PJ (2005) Insecticide species sensitivity distributions: The importance of test species selection and relevance to aquatic ecosystems. *Environmental Toxicology and Chemistry*, 24, 379–388.
- Maltby L, Brock TCM, van den Brink PJ (2009) Fungicide risk assessment for aquatic ecosystems: importance of interspecific variation, toxic mode of action and exposure regime. *Environmental Science and Technology*, 43, 7556–7563.
- Maund SJ, Travis KZ, Hendley P, Giddings JM, Solomon KR (2001) Probabilistic risk assessment of cotton pyrethroids: V. Combining landscape-level exposures and ecotoxicological effects data to characterize risks. *Environmental Toxicology and Chemistry*, 20, 687–692.

- Moore DRJ, Thompson RP, Rodney SI, Fischer D, Ramanarayanan T, Hall T (2009) Refined aquatic risk assessment for aldicarb in the United States. *Integrated Environmental Assessment and Management*, 6, 102-118.
- 村岡哲朗 (2009) 除草剤の生態影響評価のためのメソコズム試験「水田から流出する農薬の水系への影響」. 第9回有機化学物質研究会および第26回農薬環境動態研究会資料, (独)農業環境技術研究所編, pp61-69.
- 永井孝志、稲生圭哉、堀尾剛 (2008) 不確実性を考慮した農薬の確率論的生態リスク評価：水稲用除草剤シメトリンのケーススタディー. *日本農薬学会誌*, 33, 393-402.
- 永井孝志、稲生圭哉、横山淳史、岩船敬、堀尾剛 (2011) 11種の水稲用除草剤の確率論的生態リスク評価. *日本リスク研究学会誌*, 20, 279-291.
- Nagai T, Yokoyama A (2012) Comparison of ecological risks of insecticides for nursery-box application using species sensitivity distribution. *Journal of Pesticide Science*, 37, 233-239.
- Nagai T, Taya K, Annoh H, Ishihara S (2013) Application of a fluorometric microplate algal toxicity assays for riverine periphytic algal species. *Ecotoxicology and Environmental Safety*, 94, 37-44.
- Nagai T, Taya K (2015) Estimation of herbicide species sensitivity distribution using single-species toxicity data and information on the mode of action. *Environmental Toxicology and Chemistry*, 34, 677-684.
- Nagai T (2016) Ecological effect assessment of 68 pesticides used in Japanese paddy field using species sensitivity distribution. *Journal of Pesticide Science*, 41, 6-14.
- Naito W, Gamo Y, Yoshida K (2006) Screening-level risk assessment of di(2-ethylhexyl) phthalate for aquatic organisms using monitoring data in Japan. *Environmental Monitoring and Assessment*, 115, 451-471.
- 内藤航、加茂将史、林彬勸 (2012) 生態リスクトレードオフ評価ガイドランス Ver0.8 (予備的公開版), (独)産業技術総合研究所.
- 中西準子、花井荘輔、蒲生昌志、吉田喜久雄 (2007) リスク評価の知恵袋シリーズ2 不確実性をどう扱うか-データの外挿と分布-. 丸善.
- 日本植物防疫協会編 (2011) 農薬ハンドブック 2011. 日本植物防疫協会.
- 農業環境技術研究所 (2008) コガタシマトビケラ 1 齢幼虫を用いた農薬の急性毒性試験法マニュアル.
- 農業環境技術研究所 (2014) 河川付着藻類を用いた農薬の毒性試験マニュアル.
- 農林水産省 (2000) 農薬の登録申請に係る試験成績について (平成12年11月24日付け12農産第8147号農林水産省農産園芸局長通知) .

- OECD (1992) Report of the OECD workshop on the extrapolation of laboratory aquatic toxicity data to the real environment, OECD Environment Monograph Series No. 59. Organization for Economic Co-operation and Development.
- OECD (1995) Guidance document for aquatic effects assessment. Organization for Economic Co-operation and Development.
- Okkerman PC, vd Plassche EJ, Slooff W, van Leeuwen CJ, Canton JH (1991) Ecotoxicological effects assessment: A comparison of several extrapolation procedures. *Ecotoxicology and Environmental Safety*, 21, 182–193.
- Pestana JLT, Alexander AC, Culp JM, Baird DJ, Cessna AJ, Soares AMVM (2009) Structural and functional responses of benthic invertebrates to imidacloprid in outdoor stream mesocosms. *Environmental Pollution*, 157, 2328–2334.
- Posthuma L, Suter GW, Traas TP (2002) Species Sensitivity Distributions in Ecotoxicology (Environmental and Ecological Risk Assessment), Lewis Publishers, CRC Press.
- Ritter AM, Shaw JL, Williams WM, Travis KZ (2000) Characterizing aquatic ecological risks from pesticides using a diquat dibromide case study. I. Probabilistic exposure estimates. *Environmental Toxicology and Chemistry*, 19, 749–759.
- RIVM (2001) Guidance document on deriving environmental risk limits. National Institute of Public Health and the Environment.
- RIVM (2008a) Environmental risk limits for dimethoate. Report 601714001. National Institute of Public Health and the Environment.
- RIVM (2008b) Environmental risk limits for organophosphorous pesticides. Report 601714004. National Institute of Public Health and the Environment.
- Sánchez-Bayo F, Goka K (2006) Ecological effects of the insecticide imidacloprid and a pollutant from antidandruff shampoo in experimental rice fields. *Environmental Toxicology and Chemistry*, 25, 1677–1687.
- SANCO (2002) Guidance Document on Aquatic Ecotoxicology. European Commission.
- Sala S, Migliorati S, Monti GS, Vighi M (2012) SSD-based rating system for the classification of pesticide risk on biodiversity. *Ecotoxicology*, 21, 1050–1062.
- Smetanová S, Bláha L, Liess M, Schäfer RB, Beketov MA (2014) Do predictions from species sensitivity distributions match with field data? *Environmental Pollution*, 189, 126–133.
- Solomon KR, Baker DB, Richards RP, Dixon KR, Klaine SJ, La Point TW, Kendall RJ, Weisskopf CP, Giddings JM, Giesy JP, Hall Jr LW, Williams WM (1996) Ecological

- risk assessment of atrazine in north American surface waters. *Environmental Toxicology and Chemistry*, 15, 31-76.
- Solomon KR, Giddings JM, Maund SJ (2001) Probabilistic risk assessment of cotton pyrethroids: I. Distributional analyses of laboratory aquatic toxicity data. *Environmental Toxicology and Chemistry*, 20, 652-659.
- 田中二良編 (1978) 水生生物と農薬—急性毒性資料編—. サイエンティスト社.
- TenBrook PL, Tjeerdema RS, Hann P, Karkoski J (2009a) Methods for deriving pesticide aquatic life criteria. *Reviews of Environmental Contamination and Toxicology*, 199, 19-109.
- TenBrook PL, Palumbo AJ, Fojut TL, Hann P, Karkoski J, Tjeerdema RS (2009b) The University of California-Davis methodology for deriving aquatic life pesticide water quality criteria. *Reviews of Environmental Contamination and Toxicology*, 209, 1-155.
- Thursby G, Sankula S, Lewis M (2011) Estimating aquatic plant community hazard concentrations for pesticide effects. US Environmental Protection Agency, EPA-HQ-2011-0898-0012.
- Tomlin C (2009) Pesticide Manual 15th edition, BCPC Publications.
- Travis KZ, Hendley P (2001) Probabilistic risk assessment of cotton pyrethroids: IV. Landscape-level exposure characterization. *Environmental Toxicology and Chemistry*, 20, 679-686.
- Tremolada P, Finizio A, Villa S, Gaggi C, Vighi M (2004) Quantitative inter-specific chemical activity relationships of pesticides in the aquatic environment. *Aquatic Toxicology*, 67, 87-103.
- USDA (2005) Imidacloprid - human health and ecological risk assessment - final report. USDA Forest Service.
- USEPA (1985) Guidelines for deriving numerical national water quality criteria for the protection of aquatic organisms and their uses. US Environmental Protection Agency.
- USEPA (1997) Reregistration Eligibility Decision, Diflubenzuron. US Environmental Protection Agency.
- USEPA (2005) Aquatic life ambient water quality criteria, diazinon, final. US Environmental Protection Agency, EPA-822-R-05-006.
- USEPA (2010) Endosulfan: 2010 Environmental fate and ecological risk assessment. US Environmental Protection Agency, EPA-HQ-OPP-2002-0262.

- Vaal M, van der Wal JT, Hoekstra J, Hermens J (1997) Variation in the sensitivity of aquatic species in relation to the classification of environmental pollutants. *Chemosphere*, 35, 1311–1327.
- van den Brink PJ, Blake N, Brock TCM, Maltby L (2006) Predictive value of species sensitivity distributions for effects of herbicides in freshwater ecosystems. *Human and Ecological Risk Assessment*, 12, 645–674.
- van Wijngaarden RPA, Brock TCM, van den Brink PJ (2005) Threshold Levels for Effects of Insecticides in Freshwater Ecosystems: A Review. *Ecotoxicology*, 14, 355–380.
- Warne MSJ, Hawker DW (1995) The number of components in a mixture determines whether synergistic and antagonistic or additive toxicity predominate: The funnel hypothesis. *Ecotoxicology and Environmental Safety*, 31, 23–28.
- Wirth EF, Pennington PL, Lawton JC, DeLorenzo ME, Bearden D, Shaddrix B, Sivertsen S, Fulton MH (2004) The effects of the contemporary-use insecticide (fipronil) in an estuarine mesocosm. *Environmental Pollution*, 131, 365–371.
- 安野正之、花里孝幸、宮下衛、高村典子 (1988) 小型隔離水界による除草剤二剤の生態影響の研究. *国立公害研究所報告*, 114, 25–38.
- 吉田喜久雄、中西準子 (2006) 環境リスク解析入門 化学物質編. 東京図書.