

## 7.3 Laplace Transforms: translations & unit step functions


a lesson for MATH F302 Differential Equations

Ed Bueler, Dept. of Mathematics and Statistics, UAF

April 5, 2019

for textbook: D. Zill, *A First Course in Differential Equations with Modeling Applications*, 11th ed.

## the Laplace transform strategy



- §7.3: “operational properties” regarding translations (shifts)
  - including the *unit step function*  $\mathcal{U}(t)$
- §7.4 (next): “operational property” re *convolution*

## recall Laplace's Transform

- the definition:

$$\mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt$$

- when applying  $\mathcal{L}$  to an ODE use:

$$\mathcal{L}\{y'(t)\} = sY(s) - y(0)$$

$$\mathcal{L}\{y''(t)\} = s^2 Y(s) - sy(0) - y'(0)$$

- doing  $\mathcal{L}^{-1}$  is mostly use of a table, e.g.:

- $\mathcal{L}^{-1}\left\{\frac{1}{s-4}\right\} = e^{4t}$

- $\mathcal{L}^{-1}\left\{\frac{k}{(s-a)^2 + k^2}\right\} = e^{at} \sin kt$



Pierre-Simon Laplace  
(1749–1827)

we have a decent table

TABLE OF LAPLACE TRANSFORMS:

$\mathcal{L}\{1\} = \frac{1}{s}$	$\mathcal{L}\{te^{at}\} = \frac{1}{(s-a)^2}$
$\mathcal{L}\{t\} = \frac{1}{s^2}$	$\mathcal{L}\{t^n e^{at}\} = \frac{n!}{(s-a)^{n+1}}$
$\mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}}$	$\mathcal{L}\{e^{at} \sin(kt)\} = \frac{k}{(s-a)^2 + k^2}$
$\mathcal{L}\{t^{-1/2}\} = \frac{\sqrt{\pi}}{s^{1/2}}$	$\mathcal{L}\{e^{at} \cos(kt)\} = \frac{s-a}{(s-a)^2 + k^2}$
$\mathcal{L}\{t^{1/2}\} = \frac{\sqrt{\pi}}{2s^{3/2}}$	$\mathcal{L}\{t \sin(kt)\} = \frac{2ks}{(s^2 + k^2)^2}$
$\mathcal{L}\{t^\alpha\} = \frac{\Gamma(\alpha+1)}{s^{\alpha+1}}$	$\mathcal{L}\{t \cos(kt)\} = \frac{s^2 - k^2}{(s^2 + k^2)^2}$
$\mathcal{L}\{e^{at}\} = \frac{1}{s-a}$	$\mathcal{L}\{e^{at} f(t)\} = F(s-a)$
$\mathcal{L}\{\sin(kt)\} = \frac{k}{s^2 + k^2}$	$\mathcal{L}\{\mathcal{U}(t-a)\} = \frac{e^{-as}}{s}$
$\mathcal{L}\{\cos(kt)\} = \frac{s}{s^2 + k^2}$	$\mathcal{L}\{f(t-a)\mathcal{U}(t-a)\} = e^{-as}F(s)$
$\mathcal{L}\{\sinh(kt)\} = \frac{k}{s^2 - k^2}$	$\mathcal{L}\{g(t)\mathcal{U}(t-a)\} = e^{-as}\mathcal{L}\{g(t+a)\}$
$\mathcal{L}\{\cosh(kt)\} = \frac{s}{s^2 - k^2}$	$\mathcal{L}\{t^n f(t)\} = (-1)^n \frac{d^n}{ds^n} F(s)$
$\mathcal{L}\{f^{(n)}(t)\} = s^n F(s) - s^{n-1}f(0) - \dots - f^{(n-1)}(0)$	
$(f * g)(t) = \int_0^t f(\tau)g(t-\tau) d\tau$	
$\mathcal{L}\{f * g\} = F(s)G(s)$	

## noticeable in the table

- compare the left and right columns in this part of the table:

$$\mathcal{L}\{t\} = \frac{1}{s^2}$$

$$\mathcal{L}\{te^{at}\} = \frac{1}{(s-a)^2}$$

$$\mathcal{L}\{t^n\} = \frac{n!}{s^{n+1}}$$

$$\mathcal{L}\{t^n e^{at}\} = \frac{n!}{(s-a)^{n+1}}$$

$$\mathcal{L}\{\sin(kt)\} = \frac{k}{s^2 + k^2}$$

$$\mathcal{L}\{e^{at} \sin(kt)\} = \frac{k}{(s-a)^2 + k^2}$$

$$\mathcal{L}\{\cos(kt)\} = \frac{s}{s^2 + k^2}$$

$$\mathcal{L}\{e^{at} \cos(kt)\} = \frac{s-a}{(s-a)^2 + k^2}$$

- multiplying by  $e^{at}$  causes:  $s \rightarrow s - a$
- this is a rule!: multiplying by an exponential in  $t$  is *translation* in  $s$ :

$$\mathcal{L}\{e^{at} f(t)\} = F(s-a)$$

why?

- why does multiplying by  $e^{at}$  cause  $s \rightarrow s - a$  ?
- recall definition:

$$\mathcal{L}\{f(t)\} = F(s) = \int_0^{\infty} e^{-st} f(t) dt$$

- so:

$$\begin{aligned}\mathcal{L}\{e^{at}f(t)\} &= \int_0^{\infty} e^{-st} e^{at} f(t) dt = \int_0^{\infty} e^{-(s-a)t} f(t) dt \\ &= F(s-a)\end{aligned}$$

## examples from §7.3

- start by just going back and forth using the new rule
- *exercise 1.*

$$\mathcal{L}\{e^{2t}\sin(3t)\} =$$

- *exercise 2.*

$$\mathcal{L}^{-1}\left\{\frac{1}{s^2 - 6s + 10}\right\} =$$

## example like §7.3 #23

- exercise 3. use  $\mathcal{L}$  to solve the ODE IVP:

$$y'' + 4y' + 4y = 0, \quad y(0) = 1, \quad y'(0) = 1$$

example like §7.3 #30

- exercise 4. use  $\mathcal{L}$  to solve the ODE IVP:

$$y'' - 2y' + 5y = t, \quad y(0) = 0, \quad y'(0) = 7$$

## unit step function



- *definition.* the *unit step function* is

$$\mathcal{U}(t) = \begin{cases} 0, & t < 0 \\ 1, & t \geq 0 \end{cases}$$

- the book defines it with a translation, and only on  $[0, \infty)$

$$\mathcal{U}(t - a) = \begin{cases} 0, & 0 \leq t < a \\ 1, & t \geq a \end{cases}$$

- why? because we want to model “switching on” at time  $t = a$




$\mathcal{U}(t - a)$  helps with switching on/off

write each function in terms of unit step function(s):

- *example A.*

$$f(t) = \begin{cases} 0, & 0 \leq t < 1 \\ t^2, & t \geq 1 \end{cases}$$

- *example B.*



## Laplace transform with $\mathcal{U}(t - a)$



- $\mathcal{U}(t)$  is also called the *Heaviside* function
- easy-to-show: if  $F(s) = \mathcal{L}\{f(t)\}$  then

$$\mathcal{L}\{f(t - a)\mathcal{U}(t - a)\} = e^{-as}F(s)$$

- *show it:*



Oliver Heaviside  
(1850–1925)



## #57 in §7.3

- *exercise 5.* write the function in terms of  $\mathcal{U}$  and then find the Laplace transform:

$$f(t) = \begin{cases} 0, & 0 \leq t < 1 \\ t^2, & t \geq 1 \end{cases}$$

## second version

- the book then says:

*We are frequently confronted with the problem of finding the Laplace transform of a product of a function  $g$  and a unit step function  $\mathcal{U}(t - a)$  where the function  $g$  lacks the precise shifted form  $f(t - a)$  in Theorem 7.3.2.*

- yup, that's our problem
- 2nd form of the same rule:

$$\mathcal{L}\{g(t)\mathcal{U}(t - a)\} = e^{-at}\mathcal{L}\{g(t + a)\}$$

- it will be in the table also, when it is printed on quizzes/exams

once again

- *exercise 5.* write the function in terms of  $\mathcal{U}$  and then find the Laplace transform:

$$f(t) = \begin{cases} 0, & 0 \leq t < 1 \\ t^2, & t \geq 1 \end{cases}$$

like #66 in §7.3

- exercise 6. use Laplace transforms to solve the ODE IVP:

$$y'' + 9y = f(t), \quad y(0) = 0, \quad y'(0) = 0$$

where  $f(t) = \begin{cases} 1, & 0 \leq t < 1 \\ 0, & t \geq 1 \end{cases}$

## summary

- assume  $\mathcal{L}\{f(t)\} = F(s)$
- *1st translation theorem.*

$$\mathcal{L}\{e^{at}f(t)\} = F(s - a)$$

- *2nd translation theorem.* if  $a > 0$  then

$$\mathcal{L}\{f(t - a)\mathcal{U}(t - a)\} = e^{-as}F(s)$$

- includes easy case:  $\mathcal{L}\{\mathcal{U}(t - a)\} = \frac{e^{-as}}{s}$
- second form

$$\mathcal{L}\{g(t)\mathcal{U}(t - a)\} = e^{-as}\mathcal{L}\{g(t + a)\}$$

- these are all in the table you will get on quizzes and exams, so:  
**goal is understanding not memorizing**

## expectations

- just watching this video is *not* enough!
  - see “found online” videos and stuff at [bueler.github.io/math302/week12.html](https://bueler.github.io/math302/week12.html)
  - *read* sections 7.3 and 7.4 in the textbook
    - you can ignore “beams” and example 10 in §7.3
    - only 7.4.2 Transforms of Integrals in §7.4
  - *do* the WebAssign exercises for section 7.3
    - I will quiz on problems like these