




<h1>8</h1>	<h2>Special Senses</h2>
PART A	
PowerPoint® Lecture Slide Presentation by Jerry L. Cook, Sam Houston University	
	<p>ESSENTIALS OF HUMAN ANATOMY & PHYSIOLOGY</p> <p>EIGHTH EDITION</p> <p>ELAINE N. MARIEB</p>
Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings	

<h3>The Senses</h3>
<ul style="list-style-type: none"> ▪ General senses of touch <ul style="list-style-type: none"> ▪ Temperature ▪ Pressure ▪ Pain ▪ Special senses <ul style="list-style-type: none"> ▪ Smell ▪ Taste ▪ Sight ▪ Hearing ▪ Equilibrium
Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

<h3>The Eye and Vision</h3>
<ul style="list-style-type: none"> ▪ 70 percent of all sensory receptors are in the eyes ▪ Each eye has over a million nerve fibers ▪ Protection for the eye <ul style="list-style-type: none"> ▪ A cushion of fat
Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

<h3>Accessory Structures of the Eye</h3>
<ul style="list-style-type: none"> ▪ Eyelids ▪ Eyelashes

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings
Figure 8.1b

<h3>Accessory Structures of the Eye</h3>
<ul style="list-style-type: none"> ▪ Conjunctiva <ul style="list-style-type: none"> ▪ Membrane that lines the eyelids ▪ Connects to the surface of the eye ▪ Secretes mucus to lubricate the eye
Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

<h3>Accessory Structures of the Eye</h3>
<ul style="list-style-type: none"> ▪ Lacrimal apparatus <ul style="list-style-type: none"> ▪ Lacrimal gland ▪ Lacrimal canals

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings
Figure 8.1a

Function of the Lacrimal Apparatus

- Properties of lacrimal fluid
 - Dilute salt solution (tears)
 - Contains antibodies and lysozyme
- Protects, moistens, and lubricates the eye
- Empties into the nasal cavity

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

The Fibrous Tunic

- Sclera
 - White connective tissue layer
 - the “white of the eye”
- Cornea
 - Allows light to pass through
 - The only human tissue that can be transplanted without fear of rejection

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Choroid Layer

- Blood-rich nutritive layer
- Pigment prevents light from scattering
- Modified interiorly into two structures
 - Ciliary body – smooth muscle
 - Iris
 - Pigmented layer that gives eye color
 - Pupil – rounded opening in the iris


Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Sensory Tunic (Retina)

- Contains receptor cells (photoreceptors)
 - Rods
 - Cones
- Signals leave the retina toward the brain through the optic nerve- Cranial nerve ?

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Neurons of the Retina



Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Figure 8.4

Neurons of the Retina and Vision

- Rods
 - found towards the edges of the retina
 - Allow dim light vision and peripheral vision
 - Perception is all in gray tones

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Neurons of the Retina and Vision

- Cones
 - Detailed color vision
 - Fovea centralis – area of the retina with only cones
- No photoreceptor cells are at the optic disk, or blind spot

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Cone Sensitivity

- There are three types of cones (red, blue, green cones)
- Different cones are sensitive to different wavelengths
- Color blindness is the result of lack of one cone type




Figure 8.6

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Lens

- Biconvex crystal-like structure
- Held in place by a suspensory ligament attached to the ciliary body




Figure 8.3a

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Internal Eye Chamber Fluids

- Aqueous humor
 - Watery fluid found in chamber between the lens and cornea
 - Similar to blood plasma
 - Provides nutrients for the lens and cornea

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Internal Eye Chamber Fluids

- Vitreous humor
 - Gel-like substance behind the lens
 - Keeps the eye from collapsing
 - Lasts a lifetime and is not replaced

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Lens Accommodation

- Light must be focused to a point on the retina for optimal vision
- The eye is set for distance vision (over 20 ft away)
- The lens must change shape to focus for closer objects





Figure 8.9

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Images Formed on the Retina




Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Figure 8.10

Visual Pathway

- Photoreceptors of the retina
- Optic nerve
- Optic nerve crosses at the optic chiasma



Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Figure 8.11

Eye Reflexes

- Autonomic nervous system
 - Bright light causes pupils to constrict through action of radial and ciliary muscles
 - Viewing close objects causes accommodation

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings


The Ear

- Houses two senses
 - Hearing
 - Equilibrium (balance)
- Receptors are mechanoreceptors
- Different organs house receptors for each sense

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Anatomy of the Ear

- The ear is divided into three areas
 - Outer (external) ear
 - Middle ear
 - Inner ear




Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Figure 8.12

The External Ear

- Involved in hearing only
- Structures of the external ear
 - Pinna (auricle)
 - External auditory canal



Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Figure 8.12

The External Auditory Canal

- Lined with skin
- Ceruminous (wax) glands
- Ends at the tympanic membrane (Ear drum)

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

The Middle Ear or Tympanic Cavity

- Air-filled cavity within the temporal bone
- Only involved in the sense of hearing

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

The Middle Ear or Tympanic Cavity

- Two tubes are associated with the inner ear
 - The opening from the auditory canal is covered by the tympanic membrane
 - The auditory tube connecting the middle ear with the throat
 - Allows for equalizing pressure during yawning or swallowing
 - This tube is otherwise collapsed

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Bones of the Tympanic Cavity

- Three bones span the cavity
 - Malleus (hammer)
 - Incus (anvil)
 - Stapes (stirrip)




Figure 8.12

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Bones of the Tympanic Cavity

- Vibrations from eardrum move the malleus
- These bones transfer sound to the inner ear




Figure 8.12

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Inner Ear or Bony Labyrinth

- Includes sense organs for hearing and balance
- Filled with perilymph




Figure 8.12

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Inner Ear or Bony Labrynth

- A maze of bony chambers within the temporal bone
 - Cochlea
 - Vestibule
 - Semicircular canals




Figure 8.12

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Organs of Hearing

- Organ of Corti
 - Located in the cochlea
 - Receptors = hair cells on the basilar membrane
 - Gel-like membrane is capable of bending hair cells
 - Cochlear nerve attached to hair cells

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Organs of Hearing




Figure 8.15

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Mechanisms of Hearing

- Vibrations from sound waves move through the membrane
- Hair cells are bent by the membrane

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Mechanisms of Hearing





Figure 8.16a-b

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Organs of Equilibrium

- Receptor cells are in two structures
 - Vestibule
 - Semicircular canals



Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Figure 8.14a-b

Organs of Equilibrium

- Equilibrium has two functional parts
 - Static equilibrium
 - Dynamic equilibrium




Figure 8.14a-b

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Static Equilibrium

- Maculae – receptors in the vestibule
 - the position of the head
- Anatomy of the maculae
 - Hair cells are embedded in the otolithic membrane
 - Otoliths (tiny stones) float in a gel around the hair cells
 - Movements cause otoliths to bend the hair cells

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Function of Maculae




Figure 8.13a-b

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Dynamic Equilibrium

- Crista ampullaris – receptors in the semicircular canals
 - Tuft of hair cells
 - Cupula (gelatinous cap) covers the hair cells




Figure 8.14c

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Dynamic Equilibrium

- Action of angular head movements
 - The cupula stimulates the hair cells
 - Impulse is sent via the vestibular nerve to the cerebellum




Figure 8.14c

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Chemical Senses – Taste and Smell

- Both senses use chemoreceptors
 - Stimulated by chemicals in solution
 - Taste has four types of receptors
 - Smell can differentiate a large range of chemicals
- Both senses complement each other and respond to many of the same stimuli

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Olfaction – The Sense of Smell

- Olfactory receptors are in the roof of the nasal cavity
 - Chemicals must be dissolved in mucus for detection
- Interpretation of smells is made in the brain

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Olfactory Epithelium




Figure 8.17

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

The Sense of Taste

- Taste buds house the receptor organs
- Locations:
 - Most are on the tongue
 - Soft palate
 - Cheeks




Figure 8.18a–b

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

The Tongue and Taste

- The tongue is covered with papillae
 - Filiform papillae – sharp with no taste buds
 - Fungiform papillae – rounded with taste buds
 - Circumvallate papillae – large papillae with taste buds
- Taste buds are found on the sides of papillae

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Structure of Taste Buds

- Gustatory cells are the receptors
 - Have gustatory hairs (long microvilli)
 - Hairs are stimulated by chemicals dissolved in saliva

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Anatomy of Taste Buds




Figure 8.18

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Taste Sensations

- Sweet receptors
 - Sugars
 - Saccharine
 - Some amino acids
- Sour receptors
 - Acids
- Bitter receptors
 - Alkaloids
- Salty receptors
 - Metal ions

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings

Developmental Aspects of the Special Senses

- DO NOT COPY THIS DOWN;
INTERESTING INFORMATION ONLY
- Formed early in embryonic development
- Eyes are outgrowths of the brain
- All special senses are functional at birth

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings