CONWAY'S NAPKIN PROBLEM

ANDERS CLAESSON AND T. KYLE PETERSEN

ABSTRACT. The napkin problem was first posed by John H. Conway, and written up as a 'toughie' in "Mathematical Puzzles: A Connoisseur's Collection," by Peter Winkler. To paraphrase Winkler's book, there is a banquet dinner to be served at a mathematics conference. At a particular table, n men are to be seated around a circular table. There are n napkins, exactly one between each of the place settings. Being doubly cursed as both men and mathematicians, they are all assumed to be ignorant of table etiquette. The men come to sit at the table one at a time and in random order. When a guest sits down, he will prefer the left napkin with probability 1/2 and the right napkin with probability 1/2. If there are napkins on both sides of the place setting, he will choose the napkin he prefers. If he finds only one napkin available, he will take that napkin (though it may not be the napkin he wants). The third possibility is that no napkin is available, and the unfortunate guest is faced with the prospect of going through dinner without any napkin!

We think of the question of how many people don't get napkins as a statistic for signed permutations, where the permutation gives the order in which people sit and the sign tells us whether they initially reach left or right. We also keep track of the number of guests who get a napkin, but not the napkin they prefer. We build a generating function for the joint distribution of these statistics, and use it to answer questions like: What is the probability that every guest receives a napkin? How many guests do we expect to be without a napkin? How many guests are happy with the napkin they receive?

1. Introduction

The problem studied in this article first appeared in the book "Mathematical Puzzles: A Connoisseur's Collection," by Peter Winkler [5], and was inspired by a true story. Rather than recounting the problem and the story ourselves, we prefer to quote directly from "Mathematical Puzzles":

The Malicious Maitre D'

At a mathematics conference banquet, 48 male mathematicians, none of them knowledgeable about table etiquette, find themselves assigned to a big circular table. On the table, between each pair of settings, is a coffee cup containing a cloth napkin. As each person is seated (by the maitre d'), he takes a napkin from his left or right; if both napkins are present, he chooses randomly (but the maitre d' doesn't get to see which one he chose).

In what order should the seats be filled to maximize the expected number of mathematicians who don't get napkins?

... This problem can be traced to a particular event. Princeton mathematician John H. Conway came to Bell Labs on March 30, 2001 to give a "General Research Colloquium." At lunchtime, [Winkler] found himself sitting between Conway and computer scientist Rob Pike (now of Google), and the napkins and coffee cups were as described in the puzzle. Conway asked how many diners would be without napkins

if they were seated in *random* order, and Pike said: "Here's an easier question—what's the *worst* order?"

The problem of the Malicious Maitre D' is not horribly difficult; if you're having trouble finding a solution, you can see Winkler's book for a nice explanation. In this paper, it is Conway's problem that we focus on. Again, from [5]:

NAPKINS IN A RANDOM SETTING

Remember the conference banquet, where a bunch of mathematicians find themselves assigned to a big circular table? Again, on the table, between each pair of settings, is a coffee cup containing a cloth napkin. As each person sits down, he takes a napkin from his left or right; if both napkins are present, he chooses randomly.

This time there is no maitre d'; the seats are occupied in random order. If the number of mathematicians is large, what fraction of them (asymptotically) will end up without a napkin?

There are two proofs of the answer, $(2-\sqrt{e})^2 \approx .12339675$, given in the book. One is combinatorial, while the other, taken from Aidan Sudbury [3], is more analytical. We give two new proofs. Our first approach yields a thorough statistical analysis, while the second is more akin to the those in Winkler's book.

From our point of view, the number of people without a napkin is a statistic for signed permutations; just not one so well studied as inversions, descents, and such. We consider the order in which guests sit down at a place as a permutation of $[n] = \{1, 2, ..., n\}$, while their preference for the right or left napkin is given by a plus or minus sign. We label the places 1, 2, 3, ..., n counter-clockwise, so that place i has place i - 1 on its left, place i + 1 on its right, and place n is to the left of place 1. With this convention, the signed permutation (2, -1, 3, 4) describes the following sequence of events at a table for four. The person sitting in place 2 sits down first and takes the napkin on his left. The person in place 1 sits next, wants to take the napkin on his right, but as person 2 already took that napkin, he must take the napkin on his left. The person at place 3 sits third and takes the napkin on his right, while poor person 4 sits last to find no napkin at his place. Each signed permutation thus corresponds to a particular set of diners who will have no napkin. In particular, for each permutation there is some number of diners without napkins, and we seek to determine what proportion of signed permutations leave a given number of diners napkinless.

We remark that this problem makes for interesting mealtime conversation, and we have heard many suggestions for variations on the theme. What if instead of a math conference banquet with all men, it is a dinner party for couples, and the couples enter one at a time, lady sitting first? What if the dinner party is a mixer for singles, where now all the ladies enter first, sitting in alternating seats? What if the guests don't mind looking farther afield than simply to their immediate left and right, say, reaching as many as two places over for a napkin? Exploring answers to these questions may lead to some interesting mathematics, but in this article we stay within the original question, where the guests are all male mathematicians,

¹In fact, Sudbury's result gives the expected proportion of diners without a napkin as $\frac{(1-pe^q)(1-qe^p)}{pq}$, where p is the probability that a diner will reach to the right, and q=1-p is the probability of reaching left. In an informal survey of 55 mathematicians, we found about 69% would prefer the napkin on the left. According to Sudbury, we would thus expect about 10.58 percent of the guests to get stuck without napkins.

²This variation is actually not very interesting mathematically, as it is not difficult to see the women will always get their choice of napkin, and that 1/4 of the men are expected to be without a napkin.

they enter in a totally random order, and they are too shy to reach beyond their immediate left or right to find a napkin.

In section 2 we define the generating functions that we use throughout the paper. Section 3 explores the question of when everyone gets a napkin, and makes connections with combinatorial objects called *ordered bipartitions*, due to Dominique Foata and Doron Zeilberger [1]. This connection makes subsequent proofs much simpler, as in section 4, where we derive powerful identities involving our generating functions that ultimately lead to exact formulas. In section 5 we answer the original question of how many guests are expected to be without a napkin, as well as provide some other statistics of interest.

2. Definitions

Let us denote the set of signed permutations of [n] by C_n . For any $\pi \in C_n$, let $o(\pi)$ be the number of napkins left on the table after every guest has been seated as described by π . Note that the number of napkins left on the table is equal to the number of guests without a napkin. Furthermore, let $m(\pi)$ be the number of people who get a napkin, but not their first choice. We say a guest is napkinless if they have no napkin, and a guest is frustrated if they get a napkin, but not the napkin they originally wanted.³ Otherwise, we say the guest is happy. For example, if $\pi = (-1, -4, 3, 5, 2)$, then the napkin between places 2 and 3 is unused (guest 4 is the unlucky one), and although person 5 gets a napkin, it was not the one he wanted. Thus, $o(\pi) = 1$, $m(\pi) = 1$, and three of the guests are happy. Define

$$C_n(x,y) := \sum_{\pi \in \mathcal{C}_n} x^{o(\pi)} y^{m(\pi)}.$$

Then the generating function we are interested in is:

(1)
$$C(x,y,z) := \sum_{n\geq 1} C_n(x,y) \frac{z^n}{2^n n!} = \sum_{i,j,n\geq 0} p(i,j,n) x^i y^j z^n,$$

where p(i, j, n) is the probability that at a table for n people, i of them are napkinless, and j of them are frustrated. Our approach to deriving these probabilities is to first "straighten" the table.

Suppose that instead of a circular table with n places and n napkins, we look at a straight table with n places and n+1 napkins, so that each place has a napkin on both its left and right. If we know all that can possibly happen in this situation, then in order to determine the circular case, we just consider that the last person to enter the room sits "between" the first and last person on the linear table. Let us make this connection more precise.

Let \mathcal{B}_n be the set of all signed permutations of [n] that result in taking both napkins from the ends of the table. Similarly, \mathcal{L}_n (resp. \mathcal{R}_n) denotes the set of all signed permutations of [n] that result in the left end napkin being taken but not the right (resp. right but not left), and \mathcal{N}_n denotes those signed permutations that result in neither end napkin being taken. We note that $\mathcal{C}_n = \mathcal{B}_n \cup \mathcal{L}_n \cup \mathcal{R}_n \cup \mathcal{N}_n$.

³When discussing this problem, French mathematician Sylvie Corteel argued that if the mathematicians were French, they would never take the "incorrect" napkin. If the napkin they wanted was not there, they would simply cross their arms and refuse to eat. But of course, that is a different problem (how does the number of napkinless guests change as the proportion of French diners changes?).

Define the following polynomials, where $o(\pi)$ is still the number of napkins left on the (now linear) table, so that $o(\pi) - 1$ is the number of people without a napkin:

$$S_n(x,y) := \sum_{\pi \in \mathcal{C}_n} x^{o(\pi)-1} y^{m(\pi)}$$

$$B_n(x,y) := \sum_{\pi \in \mathcal{B}_n} x^{o(\pi)-1} y^{m(\pi)}$$

$$L_n(x,y) := \sum_{\pi \in \mathcal{L}_n} x^{o(\pi)-1} y^{m(\pi)}$$

$$R_n(x,y) := \sum_{\pi \in \mathcal{R}_n} x^{o(\pi)-1} y^{m(\pi)}$$

$$N_n(x,y) := \sum_{\pi \in \mathcal{N}_n} x^{o(\pi)-1} y^{m(\pi)}$$

By construction, we can express $S_n(x,y)$ in terms of the other polynomials:

$$S_n(x,y) = B_n(x,y) + L_n(x,y) + R_n(x,y) + N_n(x,y).$$

Further, we can observe that by symmetry $L_n(x, y) = R_n(x, y)$, giving one of our fundamental identities:

(2)
$$S_n(x,y) = B_n(x,y) + 2L_n(x,y) + N_n(x,y).$$

Now let us recast the polynomial $C_n(x,y)$ in terms of these polynomials for the linear table. First consider the case where the last person to sit down is the person sitting at place n. In other words, we are considering signed permutations of [n] of the form $(\pi_1, \pi_2, \ldots, \pi_{n-1}, \pm n)$. Then everything that has happened before n sits can be considered the result of a signed permutation of [n-1] playing out on a linear table. With this in mind, let $\pi' = (\pi_1, \pi_2, \ldots, \pi_{n-1})$.

So if person n walks in and has both napkins available, then π' must have resulted in leaving both end napkins on a linear table, i.e., $\pi' \in \mathcal{N}_{n-1}$. Whether guest n prefers the left napkin or the right napkin, he will get his choice, leaving the other napkin on the table. If person n walks in to find only one napkin available, $\pi' \in \mathcal{L}_{n-1}$ or $\pi' \in \mathcal{R}_{n-1}$, and n will take that napkin regardless of preference, getting the one he wants half the time. Finally, guest n can walk in to find both napkins already taken. In this case, we know that $\pi' \in \mathcal{B}_{n-1}$ and that guest n will be one of the napkinless guests.

If we take the sum over all signed permutations ending with $\pm n$, we then have

$$\sum_{n=1}^{\infty} x^{o(n)} y^{m(n)} = 2x B_{n-1}(x,y) + (1+y) L_{n-1}(x,y) + (1+y) R_{n-1}(x,y) + 2N_{n-1}(x,y).$$

But this sum is the same regardless of who is the last to enter the room. Notice that if $\pi = (\pi_1, \dots, \pi_{i-1}, \pm n, \pi_{i+1}, \dots, \pi_n)$, then $\pi' = (\pi_1, \dots, \pi_{i-1}, \pi_{i+1}, \dots, \pi_n)$ is still a permutation on n-1 letters, and so the same observations follow as for the i=n case. In total, we can say that for $n \geq 2$,

$$C_{n}(x,y) = n\Big(2xB_{n-1}(x,y) + (1+y)L_{n-1}(x,y) + (1+y)R_{n-1}(x,y) + 2N_{n-1}(x,y)\Big)$$

$$(3) = n\Big(2xB_{n-1}(x,y) + 2(1+y)L_{n-1}(x,y) + 2N_{n-1}(x,y)\Big).$$

Now we define generating functions for the linear table cases. We make a list (with the convention that $N_0(x, y) = S_0(x, y) = 1$):

$$S(x, y, z) := \sum_{n \ge 0} S_n(x, y) \frac{z^n}{2^n n!}$$

$$B(x, y, z) := \sum_{n \ge 1} B_n(x, y) \frac{z^n}{2^n n!}$$

$$L(x, y, z) := \sum_{n \ge 1} L_n(x, y) \frac{z^n}{2^n n!}$$

$$R(x, y, z) := \sum_{n \ge 1} R_n(x, y) \frac{z^n}{2^n n!}$$

$$N(x, y, z) := \sum_{n \ge 0} N_n(x, y) \frac{z^n}{2^n n!}$$

We have that L(x, y, z) = R(x, y, z), so we can write

(4)
$$S(x,y,z) = B(x,y,z) + 2L(x,y,z) + N(x,y,z).$$

Now we use (3) to get a similar formula,

(5)
$$C(x,y,z) = z \Big(xB(x,y,z) + (1+y)L(x,y,z) + N(x,y,z) \Big).$$

We now introduce some alternate notation. The partition of signed permutations into the sets \mathcal{B}_n , \mathcal{L}_n , \mathcal{R}_n , and \mathcal{N}_n is important, but in some situations it is possible to consider only two cases, rather than four. Specifically, let \mathcal{K}_n be the set of all signed permutations where the leftmost napkin is taken, and let \mathcal{J}_n be the set of all signed permutations where the leftmost napkin is not taken. Then we have $\mathcal{K}_n = \mathcal{B}_n \cup \mathcal{L}_n$, $\mathcal{J}_n = \mathcal{R}_n \cup \mathcal{N}_n$, and $\mathcal{K}_n \cup \mathcal{J}_n = \mathcal{C}_n$. Similarly define

$$K_n(x,y) := \sum_{\pi \in \mathcal{K}_n} x^{o(\pi)-1} y^{m(\pi)} = B_n(x,y) + L_n(x,y)$$

$$J_n(x,y) := \sum_{\pi \in \mathcal{J}_n} x^{o(\pi)-1} y^{m(\pi)} = R_n(x,y) + N_n(x,y)$$

$$K(x,y,z) := \sum_{n \ge 1} K_n(x,y) \frac{z^n}{2^n n!} = B(x,y,z) + L(x,y,z)$$

$$J(x,y,z) := \sum_{n \ge 0} J_n(x,y) \frac{z^n}{2^n n!} = R(x,y,z) + N(x,y,z).$$

So we can write

$$S(x, y, z) = K(x, y, z) + J(x, y, z),$$

when convenient. Notice however, that C(x, y, z) is not easily expressible in terms of K(x, y, z) and J(x, y, z) only.

We finish this section with some easy observations. First of all, if we set x = y = 1, then the polynomials $S_n(1,1) = 2^n n!$ simply count all the signed permutations of [n], so the

coefficient of z^n in S(1,1,z) is 1 for all n. In other words,

$$S(1,1,z) = 1 + z + z^2 + \dots = \frac{1}{1-z}.$$

Moreover, if we set x = y = 1 in C(x, y, z), then we get

$$C(1,1,z) = zS(1,1,z) = \frac{z}{1-z}.$$

3. A Warm-up problem: when does everyone have a napkin?

A natural question to ask (perhaps easier than the overall question) is: what is the probability, p(0, 1, n) = p(0, n), that at a circular table for n people, every guest has a napkin? This probability can be expressed as the number of permutations where everyone gets a napkin, divided by the total number of signed permutations, i.e.,

$$p(0,n) = \frac{1}{2^n n!} C_n(0,1).$$

So C(0,1,z) = C(0,z) is the generating function for these probabilities. (For now, we ignore the number of people who get napkins that aren't their first choice. In other words, we set y = 1 and write F(x,z) = F(x,1,z) and $F_n(x) = F_n(x,1)$ in this section unless otherwise noted.) By equation (5), we have

$$C(0,z) = z(2L(0,z) + N(0,z)),$$

but since (on a straight table with at least one person) there is always at least one person without a napkin if neither end napkin is taken, N(0, z) = 1, and

(6)
$$C(0,z) = z(2L(0,z) + 1).$$

This makes intuitive sense because if everyone is to have a napkin, they all need to take the left napkin, or all need to take the right napkin. In terms of the polynomials, $C_n(0) = L_{n-1}(0) + R_{n-1}(0)$.

Therefore we turn our attention to L(0, z). By considering what can happen to the last person to sit down, we can prove the following.

Claim 1. The function L(0,z) satisfies the following differential equation:

(7)
$$L'(0,z) = L(0,z)^2 + \frac{3}{2}L(0,z) + \frac{1}{2}.$$

Proof. First notice that at a table for 1, that person gets the napkin they want, and half the time that choice is the left napkin. Hence the constant term. Now if we equate the rest of the coefficients, we need to prove the following recurrence for any $n \ge 1$:

(8)
$$L_{n+1}(0) = 2\sum_{i=1}^{n-1} \binom{n}{i} L_i(0) L_{n-i}(0) + 3L_n(0).$$

First notice that if everyone gets a napkin and the leftmost napkin is taken but the rightmost napkin is not, then *everyone* must take the napkin on his left.

In particular, when the last person sits, everyone else has taken the left napkin. Suppose that $\pi \in \mathcal{L}_{n+1}$ and person n+1 is the last person to sit at the table. Then we have $\pi = (\pi_1, \pi_2, \dots, \pi_n, -(n+1))$ and the permutation $\pi' = (\pi_1, \dots, \pi_n)$ is in \mathcal{L}_n . If person 1 is the last to sit at the table, then we have $\pi = (\pm (n+1), \pi_2, \dots, \pi_n, \pi_{n+1})$. Because everyone on

the right of person 1 has taken the left napkin, person 1 will be forced to reach left regardless of his preference. We also know that $\pi' = (\pi_2, \dots, \pi_{n+1})$ is a signed permutation in \mathcal{L}_n . These two cases contribute the weight of $3L_n(0)$ in (8).

Now suppose that $\pi \in \mathcal{L}_{n+1}$ and person i+1, $i=1,2,\ldots,n-1$, is the last person to sit down. Then we can write $\pi = (\pi_1,\ldots,\pi_i,\pm(n+1),\pi_{i+2},\ldots,\pi_{n+1})$. Again, everyone has to take the left napkin, and so person i+1 will be forced left regardless of his preference. The i people to his left represent a permutation in \mathcal{L}_i , while the n-i people to his right represent a permutation in \mathcal{L}_{n-i} . The people to the left of place i+1 have no effect on the people to the right of place i+1. Therefore, we can shuffle these two permutations in any of $\binom{n}{i}$ ways. This case contributes the rest of the sum in (8), and the claim is proved.

Now, rather than do too much hard work to try to solve the quadratic differential equation given by (7), we compute the $L_n(0)$ for some small values: $L_1(0) = 1, L_2(0) = 3, L_3(0) = 13, L_4(0) = 75, L_5(0) = 541, \ldots$, and plug them into Sloane's Encyclopedia [2]. Luckily, we get a hit with sequence A000670! These numbers happen to be fairly well known as the "ordered Bell numbers," or the number of ordered set partitions of [n]. At the end of this section, we give a bijection between the ordered set partitions of [n] and permutations for which everyone takes the left napkin. This bijection will give us a slightly more general result.

The exponential generating function for the ordered Bell numbers is known to be

$$f(x) = \frac{1}{2 - e^x},$$

which satisfies the differential equation

$$f'(x) = 2f(x)^2 - f(x).$$

This equation is not exactly the same as ours, but we observe that if

$$f(x) = \sum_{n \ge 0} a_n \frac{x^n}{n!},$$

then

$$f(x/2) = \sum_{n>0} a_n \frac{x^n}{2^n n!},$$

and furthermore, f(0) = 1, whereas our generating function has no constant term. So it seems reasonable to guess that L(0, z) = f(z/2) - 1. It remains to check that the function f(z/2) - 1 satisfies equation (7). Checking,

$$\frac{d}{dz} \Big[f(z/2) - 1 \Big] = \frac{1}{2} f'(z/2)
= f(z/2)^2 - \frac{1}{2} f(z/2)
= (f(z/2) - 1)^2 + \frac{3}{2} (f(z/2) - 1) + \frac{1}{2},$$

as we hoped. So we can write the formula for the L(0,z) quite nicely:

$$L(0,z) = \frac{1}{2 - e^{z/2}} - 1 = \frac{e^{z/2} - 1}{2 - e^{z/2}}.$$

Now in light of equation (6), we are able to answer the question asked in this section with the following,

Theorem 1. The generating function for the probability that everyone at the circular table gets a napkin is

$$C(0,z) = \frac{ze^{z/2}}{2 - e^{z/2}}.$$

3.1. Ordered bipartitions. We finish the section with the bijection between ordered set partitions of [n] and signed permutations that correspond to everyone at a linear table taking the napkin on the left. Let p be a set partition of [n], written

$$p = \{i_{1,1}, i_{1,2}, \dots, i_{1,l_1}\}\{i_{2,1}, i_{2,2}, \dots, i_{2,l_2}\} \cdots \{i_{k,1}, i_{k,2}, \dots, i_{k,l_k}\},\$$

where $i_{j,1} > i_{j,2} > \cdots > i_{j,l_j}$ and $l_1 + l_2 + \cdots + l_k = n$. We describe the bijection with an example. Let

$$p = \{5, 2\}\{6\}\{7, 4, 1\}\{3\}.$$

First, we give a minus sign to the least element of each block, then we remove the braces, to obtain

$$\pi = (5, -2, -6, 7, 4, -1, -3),$$

a permutation corresponding to a situation where everyone takes the napkin on the left. We can see that we indeed have produced a permutation in which everyone takes the napkin on the left by thinking of π as a diagram for the entry times and napkin preferences of everyone at the table. With π as above, we see that the person seated second from the right entered first and took the napkin on the left. The person seated third from the right wanted to take that napkin, but entered fourth, and so was forced to take the napkin on his left. Because we required $i_{j,1} > i_{j,2} > \cdots > i_{j,l_j}$, anytime someone has a preference for the napkin on the right, they find that it was taken by the person to their right.

Clearly this process is reversible. Given a permutation where everyone takes the napkin on the left, it must have at least one minus sign. In particular, it must have a minus sign on 1, since the first person to enter must take the napkin on his left. Anybody with a plus sign immediately to the left of a person with a minus sign must enter after that person. And if two or more people with plus signs are sitting (consecutively) to the left of a person with a minus sign, they must arrive in order of closeness to the minus sign; the closest to the minus sign first, followed by the second closest to the minus sign, and so on. This gives us the block structure of the partition on [n]. Reading π from left to right, we separate the blocks by putting walls immediately to the right of any number appearing with a minus sign. For example, if

$$\pi = (7, -5, -6, 4, -1, 3, -2),$$

we convert this permutation into

$$p = \{7, 5\}\{6\}\{4, 1\}\{3, 2\}.$$

Now we can build the generating function for L(0, y, z) by purely combinatorial means. Taking an approach from [1] (more on that paper in a bit), let

(9)
$$H(y,z) = \frac{z}{1!} + \frac{yz^2}{2!} + \dots + \frac{y^{n-1}z^n}{n!} + \dots = (e^{yz} - 1)/y.$$

Then H(y,z) is the exponential generating function for single blocks, $\{n, n-1, \ldots, 1\}$, since for every block, only the person corresponding to the least number gets the napkin he wants,

leaving the other n-1 people frustrated. Therefore $(1-H)^{-1}=1+H+H^2+\cdots$ is the generating function for ordered sequences of blocks. Now we have

$$L(0, y, z) = \frac{1}{(1 - H(y, z/2))} - 1 = \frac{e^{yz/2} - 1}{1 + y - e^{yz/2}}.$$

Thus we can refine the formula from Theorem 1 according to how many people get the napkin they want:

$$C(0, y, z) = \frac{yze^{yz/2}}{1 + y - e^{yz/2}}.$$

We can generalize this bijection to compute the generating function for the numbers $S_n(0,1)$, i.e., the number of permutations for which everyone on the straight table would end up with a napkin. Experimentally, we compute the first few values: 1, 2, 8, 44, 308, 2612, 25988,..., and we see that Sloane's Encyclopedia [2] contains this sequence and gives its exponential generating function. We find

$$S(0,1,z) = \frac{1}{(2 - e^{z/2})^2},$$

which is simply a special case of the generating function given by (10) below.

The paper of Foata and Zeilberger [1], introduces objects called *ordered bipartitions*, which are easiest to think of as ordered set partitions with some of the subsets are underlined. A *compatible* bipartition is an ordered bipartition where all the underlined subsets are on the right, e.g.,

$$p = \{5, 2\}\{6\}\{1, 4, 7\}\{3\},\$$

where we adopt the convention that underlined subsets have their elements written in ascending order. The bijection works as follows. For every non-underlined group in p, we perform the same operation as above, while for every underlined group we perform the "opposite" operation. Specifically, we put minus signs on all but the least element before removing the braces. This produces

$$\pi = (5, -2, -6, 1, -4, -7, 3),$$

a permutation where everyone on a linear table receives a napkin. All we really need to observe is that the underlined groups correspond to the part of the table where people all take napkins on their right, while the non-underlined groups all take napkins on the left. Thus,

(10)
$$S(0, y, z) = (1 - H(y, z/2))^{-1} (1 - H(y, z/2))^{-1} = \frac{y^2}{(1 + y - e^{yz/2})^2}.$$

We can take this more general correspondence and see now that the permutations for which everyone takes the napkin on the left correspond to the ordered bipartitions with no underlined subsets, the ordered bipartitions with all subsets underlined correspond to permutations where everyone takes the right napkin, and the compatible bipartitions with at least one underlined subset and one non-underlined subset correspond to the permutations where everyone gets a napkin and both end napkins are taken. This last observation gives

$$B(0, y, z) = \frac{(e^{yz/2} - 1)^2}{(1 + y - e^{yz/2})^2}.$$

4. Ordered bipartitions and generating functions

We can embed the set of all signed permutations in the set of ordered bipartitions according to the algorithm below. Let $\varphi : \mathcal{C}_n \to \mathcal{OBP}_n$ be this map, where \mathcal{OBP}_n is the set of ordered bipartitions of [n]. The image of the injection φ will lead us to the main theorem of this section, which gives some incredible relationships between the generating functions for the linear table. Given a signed permutation π of [n], we form its image $\varphi(\pi)$ as follows:

- (1) Find the least element, $\pi(i)$ (ignoring signs), that is not already included in some subset.
- (2a) If $\pi(i)$ is positive, then underline the set including $\pi(i)$, and set j = i + 1. While $|\pi(j)| > |\pi(j-1)|$ and $\pi(j)$ negative, add $\pi(j)$ to the set containing $\pi(i)$, j = j + 1.
- (2b) If $\pi(i)$ is negative, then set j = i 1. While $|\pi(j)| > |\pi(j+1)|$ and $\pi(j)$ positive, add $\pi(j)$ to the set containing $\pi(i)$, j = j - 1.
- (3) If every element is contained in a set, then delete all minus signs and quit. Else, go to (1).

Clearly, no two permutations can be mapped to the same bipartition. We will demonstrate the injection with an example. Start with

$$\pi = (9, 1, -3, 2, 5, 6, -4, -7, 8,)$$

The number 1 is the least element, and it is positive, so its set will be underlined. We start searching to the right of 1, looking for negative numbers with norm bigger than 1. We get

$$9\ \{1,-3\}\ 2\ 5\ 6\ -4\ -7\ 8$$

Now 2 is the least element, and it is also positive, but there are no negative numbers immediately to its right, so it forms a singleton set,

$$9\ \{1,-3\}\ \{2\}\ 5\ 6\ -4\ -7\ 8$$

Now 4 is the least element not already in a set, and it is negative. So we start searching to the left of 4, looking for positive numbers with bigger norm. We get

$$9 \ \underline{\{1,-3\}} \ \underline{\{2\}} \ 5 \ \{6,-4\} \ -7 \ 8$$

The next steps give,

$$9 \underbrace{\{1, -3\}}_{\{2\}} \underbrace{\{5\}}_{\{6, -4\}} - 7 8$$

$$9 \underbrace{\{1, -3\}}_{\{2\}} \underbrace{\{5\}}_{\{6, -4\}} \{-7\} 8$$

$$9 \underbrace{\{1, -3\}}_{\{2\}} \underbrace{\{5\}}_{\{6, -4\}} \{-7\} \underbrace{\{8\}}_{\{8\}}$$

$$\underbrace{\{9\}}_{\{1, -3\}} \underbrace{\{2\}}_{\{2\}} \underbrace{\{5\}}_{\{6, -4\}} \{-7\} \underbrace{\{8\}}_{\{8\}}$$

Then we drop the minus signs to get,

$$\{9\}\ \{1,3\}\ \{2\}\ \{5\}\ \{6,4\}\ \{7\}\ \{8\}$$

so that we have an ordered bipartition of [n] where we write underlined sets in increasing order. What's more, the number of napkinless diners is exactly the number of occurrences of an underlined set followed immediately by a non-underlined set. The frustrated diners are those who are not the least element in a block, less the people without any napkin: $m(\pi) = n - l - o(\pi) + 1$, where l is the number of blocks in the bipartition.

Let $\mathcal{U}_n = \varphi(\mathcal{C}_n)$, the image set of bipartitions corresponding to all signed permutations of [n]. By examining the algorithm describing φ , we see that the set \mathcal{U}_n consists of all ordered bipartitions that never contain the patterns

$$\underline{\cdots a}$$
 $\{b\}\cdots$

where a < b, or

$$\cdots \{b\} \{c \cdots$$

where b > c.

Now we present the main theorem for this section.

Theorem 2. We have the following formulas:

(11)
$$K(x, y, z) = H(y, z/2)S(x, y, z)$$

(12)
$$B(x, y, z) = H(y, z/2)^{2} S(x, y, z)$$

(13)
$$L(x,y,z) = H(y,z/2)(1 - H(y,z/2))S(x,y,z)$$

(14)
$$N(x, y, z) = (1 - H(y, z/2))^2 S(x, y, z)$$

(15)
$$J(x, y, z) = (1 - H(y, z/2))S(x, y, z)$$

(16)
$$C(x,y,z) = z((x-y)H(y,z/2)^2 + (y-1)H(y,z/2) + 1)S(x,y,z)$$

Proof. Let \mathcal{U}_n be the image set $\varphi(\mathcal{C}_n)$. If we add a non-underlined block of size r to the left of any bipartition in \mathcal{U}_n , then clearly we get a bipartition in \mathcal{U}_{n+r} that corresponds to a permutation in \mathcal{K}_{n+r} . Furthermore, this new block will not change the number of people without a napkin (occurrences of underlined blocks immediately to the left of non-underlined blocks), and the number of new people who get a napkin they don't want is exactly r-1. Therefore,

$$K(x, y, z) = H(y, z/2)S(x, y, z).$$

Similarly, if we add a non-underlined block of size r to the left of a a bipartition in \mathcal{U}_n , and an underlined block of size s to the right, then we get a bipartition in \mathcal{U}_{n+r+s} that corresponds to a permutation in \mathcal{B}_{n+r+s} . Thus,

$$B(x, y, z) = H(y, z/2)^2 S(x, y, z).$$

Now we use the relations L = K - B, N = S - K - L, J = L + N, and C = z(xB + (1+y)L + N) to obtain the rest.

What Theorem 2 tells us is that if we can find an explicit formula for S(x, y, z), then we will have explicit formulas for all the other generating functions. We will derive such an explicit formula shortly. First we need to introduce the notion of a cyclic bipartition. A cyclic bipartition is a bipartition for which one element is distinguished, and only the cyclic ordering of the blocks matters. As a convention, we put the block containing the distinguished element at the far right if that block is not underlined, or at the far left if it is underlined. In our notation we will enclose the distinguished element in parentheses. For example,

$$\underline{\{1,6\}}$$
 $\{8,2\}$ $\underline{\{3,5\}}$ $\{9,(7),4\}$

is a cyclic bipartition, which we also write

$$7,4$$
} $\{1,6\}$ $\{8,2\}$ $\{3,5\}$ $\{9,$

so that the distinguished element is equivalently the first element in a bipartition where a block can "wrap around." Similarly to how we encoded signed permutations playing out on a straight table with ordered bipartitions, we can encode the case of the circular table with cyclic bipartitions. Here the distinguished element will correspond to a "distinguished guest," who sits in place number 1 at the circular banquet table.

Let $\varphi^{(c)}: \mathcal{C}_n \to \mathcal{OBP}_n^{(c)}$ the map encoding the circular table case, where $\mathcal{OBP}_n^{(c)}$ is the set of cyclic bipartitions of [n]. For any signed permutation π of [n], we form its image $\varphi(\pi)$ as follows (the only difference in our algorithm is that the searches are cyclic):

- (1) Find the least element, $\pi(i)$ (ignoring signs), that is not already included in some subset.
- (2a) If $\pi(i)$ is positive, then underline the set including $\pi(i)$, and set $j = i + 1 \mod n$. While $|\pi(j)| > |\pi(j-1)|$ and $\pi(j)$ negative, add $\pi(j)$ to the set containing $\pi(i)$, $j = j + 1 \mod n$.
- (2b) If $\pi(i)$ is negative, then set $j = i 1 \mod n$. While $|\pi(j)| > |\pi(j+1)|$ and $\pi(j)$ positive, add $\pi(j)$ to the set containing $\pi(i)$, $j = j - 1 \mod n$.
- (3) If every element is contained in a set, then delete all minus signs and quit. Else, go to (1).

As an example, start with the permutation

$$\pi = (-7, 1, -3, 4, -2, 5, -6).$$

The steps of the encoding are

$$-7 \ \underline{\{1, -3\}} \ 4 \ -2 \ 5 \ -6$$
$$-7 \ \underline{\{1, -3\}} \ \{4, -2\} \ 5 \ -6$$

and this last step, which requires us to perform a cyclic search,

$$-7$$
 $\{1, -3\}$ $\{4, -2\}$ $\{5, -6,$

Now we can drop the signs to get

$$7$$
 $\{1,3\}$ $\{4,2\}$ $\{5,6,$

or

$$\underline{\{5,6,(7)\}} \ \underline{\{1,3\}} \ \{4,2\}.$$

Let $\varphi^{(c)}(\mathcal{C}_n) = \mathcal{U}_n^{(c)}$ be the set of all cyclic bipartitions corresponding to signed permutations on a circular table. We will use this set of cyclic bipartitions along with \mathcal{U}_n to obtain the following theorem regarding the derivative of S(x, y, z).

Theorem 3. We have the following formula relating the generating function for the straight table and the generating function for the circular table:

(17)
$$z\frac{d}{dz}\Big[S(x,y,z)\Big] = C(x,y,z)S(x,y,z).$$

Proof. To prove equation (17), it will suffice to equate coefficients and prove

(18)
$$nS_n(x,y) = \sum_{i=0}^n \binom{n}{i} C_i(x,y) S_{n-i}(x,y).$$

We will prove (18) bijectively.

The left-hand side of equation (18) can be thought of as counting the weights of permutations corresponding to bipartitions in \mathcal{U}_n with a distinguished element, or a straight table with a distinguished guest (not necessarily in place 1). We simply put parentheses around one of the n elements of the bipartition, as in

$$\underline{\{2,8\}}$$
 $\{1\}$ $\{7,(4),3\}$ $\underline{\{9\}}$ $\underline{\{5,6\}}$

Given any bipartition with a distinguished element, we can form a pair (c, s), where c is a cyclic bipartition of a subset $A \subset [n]$ (corresponding to one in $\mathcal{U}_i^{(c)}$), and s is an ordered bipartition of $[n] \setminus A$ (corresponding to one in \mathcal{U}_{n-i}). We simply split the table in two pieces just before or after the block containing the distinguished element. If the block containing the distinguished element is not underlined, we make the split just after that block. The above example yields the pair

$$({2,8} {1} {7,(4),3}, {9} {5,6}).$$

If instead the block with the distinguished element is underlined, as in

$${5,3} {2,1} {6,(7),8} {9} {5,6},$$

we make the split before the underlined block. Now the right half becomes the circular table, and the left half is the straight table:

$$({6, (7), 8} {9} {5, 6}, {5, 3} {2, 1}).$$

By splitting the bipartition as we do, all of the guests retain their status as happy, frustrated, or napkinless. In other words, the product of the weights of the pair of tables equals the weight of original table.

Now, given any pair (c, s), where $c \in \mathcal{U}_i^{(c)}$ and $s \in \mathcal{U}_{n-i}$, we can form a bipartition in \mathcal{U}_n with a distinguished element. First, we must ensure that c and s have no common elements. Thus, in any of $\binom{n}{i}$ ways, we choose a subset $A = \{a_1 < a_2 < \cdots < a_i\} \subset [n]$, and replace k with a_k in c. For s, we replace k with b_k , where $[n] \setminus A = \{b_1 < b_2 < \cdots < b_{n-i}\}$. Now we concatenate the bipartitions. If the distinguished element of c is in an underlined block we put the straight table on the left: sc. If the distinguished element is not in an underlined block, we put the straight table on the right: cs.

Now, thanks to Theorem 2 and Theorem 3, we have:

$$S' = ((x-y)H(y,z/2)^2 + (y-1)H(y,z/2) + 1)S^2.$$

We can solve this differential equation to get:

(19)
$$S(x,y,z) = \frac{y^3}{y(1+y-e^{yz/2})^2 - x((2-e^{yz/2})^2 + yz - 1)}.$$

Now we can also obtain exact formulas for any of the generating functions discussed here by plugging equation (19) into the formulas of Theorem 2. In particular, we have

(20)
$$C(x,y,z) = \frac{yz(ye^{yz/2}(1+y-e^{yz/2})+x(e^{yz/2}-1)^2)}{y(1+y-e^{yz/2})^2-x((2-e^{yz/2})^2+yz-1)}.$$

5. The expected number of napkinless guests and other statistics

With the generating function from (20), we can in principle extract any sort of statistics related to the number of napkinless guests or frustrated guests. We will highlight a few statistical results that are of interest to us. In particular, we find the expected number of napkinless and frustrated guests, the variance for each of these distributions, and the covariance for their joint distribution.

Suppose we know exactly what the polynomial $C_n(x, y)$ is for some n. As we saw in section 3, $C_n(0, 1)$ tells us the number of permutations that result in everyone having a napkin. But now if we want to obtain the expected number of people without a napkin, we want to compute the weighted average

$$\frac{1}{2^n n!} \sum_{\pi \in C_n} o(\pi) = \frac{0 \cdot a_0 + 1 \cdot a_1 + 2 \cdot a_2 + \dots + k \cdot a_k}{2^n n!},$$

where a_i is the number of signed permutations of [n] for which i people are without a napkin, and $k = \lfloor n/3 \rfloor$ is the maximum number of people that could possibly get stuck without a napkin. In terms of the polynomial $C_n(x, 1)$, this just means that we differentiate with respect to x, set x = 1, and then divide by $2^n n!$, since

$$C'_n(x,1) = \sum_{\pi \in \mathcal{C}_n} o(\pi) x^{o(\pi)-1}$$

$$C'_n(1,1) = \sum_{\pi \in \mathcal{C}_n} o(\pi).$$

Therefore, we want to find the generating function for the numbers $E_n(o(\pi)) = \frac{1}{2^n n!} C'_n(1, 1)$, or

(21)
$$E(z) := \sum_{n\geq 0} E_n(o(\pi))z^n$$

$$= \frac{d}{dx} \Big[C(x,1,z) \Big]_{x=1}$$

$$= \frac{z(2 - e^{z/2}) (e^{z/2}(z-2) + 2)}{(1-z)^2}.$$

Expanding this series, we find the following exact formula, (which is the truncated form of the series derived by Winkler [5]).

Theorem 4. The expected number of napkinless guests on a circular table for n people is

(22)
$$E_n(o(\pi)) = n(4 - 4\exp_n(1/2) + \exp_n(1)),$$

where $\exp_n(x) = \sum_{k=0}^n x^k/k!$ is the truncated exponential function.

Proof. Let $f(n) = n(4 - 4\exp_n(1/2) + \exp_n(1))$ be the right hand side of equation (22), and let F(z) be the ordinary generating function for the numbers f(n). Note that

$$nf(n+1) = (n+1)f(n) + \frac{1 + (1/2)^{n-1}}{(n-1)!},$$

which implies

$$z\frac{d}{dz}\left[\frac{1}{z}F(z)\right] = \frac{d}{dz}(zF(z)) + e^z - e^{z/2}$$

or equivalently,

$$(1-z)F'(z) = \left(1 + \frac{1}{z}\right)F(z) + z(e^z - e^{z/2}).$$

It is easy to check that E(z), given by formula (21), satisfies this differential equation.

Corollary 1.

$$E_n(o(\pi)) = n(2 - \sqrt{e})^2 + O\left(\frac{1}{n!}\right).$$

So, the answer to Winkler's problem of napkins in a random setting is $(2 - \sqrt{e})^2 \approx 0.12339675$. It was quite a bit of work for this answer, but of course our work pays off in being able to find the following statistics as well.

We will not derive exact formulas in the rest of this section, but simply give the asymptotic results. Now, if we want to get the variance of the distribution of napkins on the table, we need to compute the sums of the squares of the number of napkins left on the table, since

$$\operatorname{Var}_n(o(\pi)) = E_n(o(\pi)^2) - E_n(o(\pi))^2,$$

and we already have $E_n(o(\pi))$. We get

$$E_n(o(\pi)^2) = \frac{1}{2^n n!} \sum_{\pi \in C_n} o(\pi)^2$$
$$= \frac{1}{2^n n!} (C''_n(1, 1) + C'_n(1, 1)).$$

Therefore the generating function for the second moment is

$$\sum_{n>0} E_n (o(\pi)^2) z^n = \frac{d^2}{dx^2} \Big[C(x, 1, z) \Big]_{x=1} + E(z),$$

and all we need to find is $\frac{d^2}{dx^2}\Big[C(x,1,z)\Big]_{x=1}$. Computing, we find:

$$C_{x^2}(1,1,z) = \frac{2z(2-e^{z/2})\Big(e^{z/2}(1-z)^2 + (2-e^{z/2})\big((e^{z/2}-1)^2(3-z) - e^z + 2z\big)\Big)}{(1-z)^3}.$$

It is straightforward to obtain asymptotic estimates for functions with a finite number of poles. See, for example, chapter 5 of Herbert Wilf's book [4]. Using such methods, we find that the variance is asymptotically $n(3-e)(2-\sqrt{e})^2 \approx n(.0347631)$.

To get the expectation and variance for the number of frustrated guests, we follow the same procedure, except now we differentiate C(1, y, z) with respect to y. The covariance is

$$E(o(\pi)m(\pi)) - E(o(\pi))E(m(\pi)),$$

so we differentiate with respect to x, then with respect to y to get the generating function for $E(o(\pi)m(\pi))$, the only piece we don't know. The statistics are summarized in Table 1. Notice in particular that if the expected number of napkinless guests is $n(2-\sqrt{e})^2$, and the expected number of frustrated guests is $n(6\sqrt{e}-e-7)$, then the expected number of happy guests is $n(4-2\sqrt{e}) \approx n(.702557)$. Seventy percent of the guests are happy!

<i>X</i> :	$o(\pi)$	$m(\pi)$
E(X)	$n(2-\sqrt{e})^2 \approx n(.12339675)$	$n(6\sqrt{e} - e - 7) \approx n(.174046)$
Var(X)	$n(3-e)(2-\sqrt{e})^2 \approx n(.0347631)$	$n(6\sqrt{e^3} - e^2 - e - 38\sqrt{e} + 46)$ $\approx n(.13138819)$
Cov(X,Y)	$n(-(2-\sqrt{e})(\sqrt{e^3}-3e-5\sqrt{e}+12)) \approx n(029239461)$	

Table 1. Statistics for napkinless and frustrated guests.

6. Another proof for the expected number of napkinless guests

Upon reviewing a draft of this paper, Ira Gessel pointed out that the generating function for the expected number of napkinless guests satisfies the following differential equation:

(23)
$$E(z) = z \frac{d}{dz} \left[\frac{(2 - e^{z/2})^2}{1 - z} \right] = z \frac{d}{dz} \left[N(1, 1, z) \right],$$

where N(1, 1, z) is the generating function for the proportion of signed permutations for which neither napkin on a straight table is taken. We will think of such permutations in terms of their image under φ : specifically, those ordered bipartitions in \mathcal{U}_n for which the leftmost block is underlined and the rightmost block is not underlined. Gessel suggested that there may be a simple combinatorial explanation for (23), and there is.

First of all, if all we want is the expected number of napkinless guests on the circular table, then because of the symmetry of the table, we have $E_n(o(\pi)) = np$, where p is the probability that any particular guest (say guest 1) has no napkin. Upon equating coefficients, equation (23) claims that $|\mathcal{N}_n| = |\mathcal{C}_n^1|$, where \mathcal{C}_n^1 is the set of all signed permutations for which guest 1 gets no napkin on a circular table.

We can give a bijection between the ordered bipartitions in $\varphi(\mathcal{N}_n)$ and the cyclic bipartitions in $\varphi^{(c)}(\mathcal{C}_n^1)$ as follows. Given an ordered bipartition for which the leftmost block is underlined and the rightmost block is not underlined, there must be a guest in the middle of the table who is napkinless. Make the leftmost such guest distinguished, and cyclically permute the blocks of the bipartition until the block with the distinguished guest is first or last, depending on whether that block is underlined. For example,

$${8} {\{(9), 6, 1\}} {\{2\}} {\{3, 5\}} {\{7, 4\}}$$

is an ordered bipartition in $\varphi(\mathcal{N}_n)$ with the leftmost napkinless guest highlighted (guest 2 in this case). We cyclically permute the blocks to get

$${2} {3,5}{7,4} {8} {(9),6,1},$$

which is a cyclic bipartition where guest 1 gets no napkin on a circular table. The inverse of this bijection is given by simply cyclically permuting the blocks until we have an ordered

bipartition for which the distinguished guest is the leftmost guest without a napkin, the leftmost block is underlined, and the rightmost block is not. It is straightforward to check that we can always achieve this state.

7. Acknowledgements

We would like to extend a sincere thank you to Don Knuth for bringing this problem to our attention. His help and encouragement were invaluable. In particular, the ideas of reducing the problem to a straight table and keeping track of frustrated guests are his. We would also like to thank the Institut Mittag-Leffler and the organizers of the spring 2005 session in algebraic combinatorics. If not for our time there, this work may never have been done. Lastly, we thank Ira Gessel, whose observation motivated Section 6.

References

- [1] D. Foata and D. Zeilberger, *Graphical major indices*, Journal of Computational and Applied Mathematics, **68** (1996), 79–101.
- [2] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, (2005), published electronically at http://www.research.att.com/~njas/sequences/.
- [3] A. Sudbury, *Inclusion-exclusion methods for treating annihilating and deposition processes*, Journal of Applied Probability, **39** (2002), 466–478.
- [4] H. S. Wilf, generatingfunctionology, Academic Press, San Diego, CA, 1994.
- [5] P. Winkler, Mathematical Puzzles: A Connoisseur's Collection, AK Peters, Natick, MA, 2004.

DIVISION OF MATHEMATICS, DEPARTMENT OF CHEMISTRY AND BIOMEDICAL SCIENCES, UNIVERSITY OF KALMAR, SWEDEN

E-mail address: anders.claesson@hik.se

DEPARTMENT OF MATHEMATICS, BRANDEIS UNIVERSITY, WALTHAM, MA, USA, 02454

E-mail address: tkpeters@brandeis.edu