

Review Notes - Solving Quadratic Equations

$$ax^2 + bx + c = 0$$

What does solve mean?

★ FIND ALL VALUES THAT MAKE THE SENTENCE TRUE!

★ How many solutions do we expect?

Up to 2 solutions [Degree = 2]

Methods for Solving Quadratic Equations:

Solving by Factoring using the Zero Product Property

Solving by using Square Roots

Solving by Quadratic Formula

How can we factor polynomials?

Factoring refers to writing something as a product.

Factoring completely means that all of the factors are relatively prime (they have a GCF of 1).


Methods of factoring:

1. Greatest Common Factor (GCF) - Any polynomial

2. Grouping - Only for 4 or 6 term polynomials

3. Trinomial Method - Only for trinomials [Quadratic]

4. Speed Factoring - Special cases only Binomials



Method 1: Factoring Out the Greatest Common Factor (GCF)

Factoring out the GCF can be done by using the distributive property.

Ex 1: Factor $12x^3 + 3x^2$.

Step 1: Find the GCF of $12x^3$ and $3x^2$.

The GCF is $3x^2$.

Step 2: Rewrite by factoring out the GCF.

$$3x^2(4x + 1)$$

Method 2: Factoring by Grouping

Ex 1: $12xy + 20x + 9y + 15$

$$(12xy + 20x) + (9y + 15)$$

Step 1: Group terms together that have a common monomial factor.

$$4x(3y + 5) + 3(3y + 5)$$

Step 2: Factor out the GCF of each group.

$$(3y + 5)(4x + 3)$$

Step 3: Find the common polynomial factor and factor it out using the distributive property.

Ex 2: $6xy + 8x - 21y - 28$

$$(6xy + 8x) + (-21y - 28)$$

$$2x(3y + 4) + (-7)(3y + 4)$$

$$(3y + 4)(2x - 7)$$

Ex 3: $4x^2z^2 - 10x^2z - 6yz + 8yz^2 - 3x^2z - 20y$

$$(4x^2z^2 - 3x^2z - 10x^2) + (8yz^2 - 6yz - 20y)$$

$$x^2(4z^2 - 3z - 10) + 2(4z^2 - 3z - 10)$$

$$(4z^2 - 3z - 10)(x^2 + 2)$$

Method 3: Factoring Using the Trinomial Method

Step 1: Write the trinomial in descending order.

Step 2: Find two numbers whose product is the same as the product of the first and third coefficients and whose sum is equal to the middle coefficient. (Make a chart.)

Step 3: Rewrite the middle term as the sum of two terms.

Step 4: Use the distributive property and factor by grouping.

Ex 1: $2x^2 - 5x - 3$

$$(2x^2 - 6x) + (x - 3)$$

$$2x(x-3) + 1(x-3)$$

$$(x-3)(2x+1)$$

$x(-6)$	$+(-5)$
$(-6)(1)$	\checkmark

Ex 2: $20y^2 + 13yz + 2z^2$

$$(20y^2 + 8yz) + (5yz + 2z^2)$$

$$4y(5y+2z) + z(5y+2z)$$

$$(5y+2z)(4y+z)$$

Method 4: Speed Factoring - Special Cases

- I. The Difference of Squares
- II. Trinomials with a lead coefficient of 1

Special Case: The Difference of Squares

Consider the product: $(a+b)(a-b)$

$$a^2 - ab + ab - b^2$$

$$a^2 - b^2$$

★ Since $(a+b)(a-b) = a^2 - b^2$, then $a^2 - b^2 = (a+b)(a-b)$.

★ $a^2 - b^2$ is called the "difference of squares."

Ex 1: $x^2 - 121$

$$(x)^2 - (11)^2$$

$$(x - 11)(x + 11)$$

Ex 2: $25x^2 - 1$

$$(5x)^2 - (1)^2$$

$$(5x - 1)(5x + 1)$$

Ex 3: $72y^2 - 50$

$$2(36y^2 - 25)$$

$$2[(6y)^2 - (5)^2]$$

$$2(6y - 5)(6y + 5)$$

Ex 4: $m^4 - 16$

$$(m^2)^2 - (4)^2$$

$$(m^2 - 4)(m^2 + 4)$$

$$[(m)^2 - (2)^2](m^2 + 4)$$

$$(m - 2)(m + 2)(m^2 + 4)$$

Special Case: Trinomials with a lead coefficient of 1

$$x^2 + bx + c$$

Find the two numbers whose product is c and whose sum is b .

These are the two numbers in the binomials.

Ex 1: $x^2 + 2x + 1$

$$(x+1)(x+1)$$

Ex 2: $x^2 - x - 12$

$$(x-4)(x+3)$$

Ex 3: $x^2 - 10x + 16$

$$(x-8)(x-2)$$

✓

Ex 4: $x^2 + 28x + 160$

$$(x+20)(x+8)$$

✓

$$\begin{array}{r|l} x(160) & +(28) \\ \hline 20(8) & \checkmark \end{array}$$

Solving Equations by Factoring - Using the Zero Product Property

The Zero Product Property:

If $xy = 0$, then either $x = 0$ or $y = 0$.

Use the zero product property to solve the following equations.

Ex 1: $x(x-1) = 0$

$x = 0$ or $x - 1 = 0$

$x = 1$

$x = 0, 1$

Ex 2: $(x-5)(x+2) = 0$

$x - 5 = 0$ or $x + 2 = 0$

$x = 5$ or $x = -2$

$x = -2, 5$

Ex 3: $5x(x-4) = 0$

$5x = 0$ or $x - 4 = 0$

$x = 0$ or $x = 4$

$x = 0, 4$

If the polynomial is not "set equal to zero", get all of the terms on one side of the equation first. Then factor the polynomial before trying to use the zero product property to solve.

Ex 4: $x^2 - 3x = 10$

$x^2 - 3x - 10 = 0$

$(x+2)(x-5) = 0$

$x + 2 = 0$ or $x - 5 = 0$

$x = -2$ or $x = 5$

$x = -2, 5$

Ex 5: $18 - 3x = x^2$

$0 = x^2 + 3x - 18$

$0 = (x+6)(x-3)$

$x = -6, 3$

Ex 6: $w^3 - w^2 = 4w - 4$

$$\underbrace{w^3 - w^2} - \underbrace{4w + 4} = 0$$

$$w^2(w-1) - 4(w-1) = 0$$

$$(w-1)(w^2-4) = 0$$

$$(w-1)(w-2)(w+2) = 0$$

$$w = -2, 1, 2$$

Ex 7: $m^3 = 121m$

$$m^3 - 121m = 0$$

$$m(m^2 - 121) = 0$$

$$m(m-11)(m+11) = 0$$

$$m = \pm 11, 0$$

$$m = -11, 0, 11$$

New Handouts:

1. Algebra Review Notes - Solving Quadratic Equations
2. Speed Factoring Packet

Homework:

Complete the Speed Factoring Packet and the Summer Assignment if necessary.