

Chapter 1




*An Introduction to
Chemistry*

By Mark Bishop

Chemistry



The science  that deals with the structure and behavior of matter

Summary of Study Strategies



The will to succeed is important, but what's more important is the will to prepare.

Bobby Knight, basketball coach


- Read the chapter in the textbook before it is covered in the lecture.
- Attend the class meetings, take notes, and participate in class discussions.
- Reread the textbook, working the exercises, and marking important sections.

More Study Strategies




- Use the chapter objectives as a focus of study.
- Use the computer-based tools that accompany the course.
- Work some of the problems at the end of the chapter.
- Ask for help when you need it.
- Review for the exam.

Scientific Method



Chapter Map



Values from Measurements




- A **value** is a quantitative description that includes both a unit and a number.
- For *100 meters*, the *meter* is a unit by which distance is measured, and the *100* is the number of units contained in the measured distance.
- **Units** are quantities defined by standards that people agree to use to compare one event or object to another.

Type	Base Unit	Abb.	Defined in terms of
length	meter	m	the fixed numerical value of the speed of light in vacuum c to be 299,792,458 when expressed in the unit m s^{-1} , where the second is defined as below.
mass	kilogram	kg	the fixed numerical value of the Planck constant, h , to be $6.62607015 \times 10^{-34}$ when expressed in the unit J s , which is equal to $\text{kg m}^2 \text{s}^{-1}$, where the meter and the second are defined in terms of c and $\Delta\nu_{\text{Cs}}$.
time	second	s	the fixed numerical value of the cesium frequency, $\Delta\nu_{\text{Cs}}$, which is the unperturbed ground-state hyperfine transition frequency of the cesium-133 atom, to be 9,192,631,770 when expressed in the unit Hz , which is equal to s^{-1} .
temperature	kelvin	K	the fixed numerical value of the Boltzmann constant, k , to be $1.380\,649 \times 10^{-23}$ when expressed in the unit J K^{-1} , which is equal to $\text{kg m}^2 \text{s}^{-2} \text{K}^{-1}$, where the kilogram, meter and second are defined in terms of h , c and $\Delta\nu_{\text{Cs}}$.

Derived Unit

1 cubic meter = 1000 liters



$$1 \text{ L} = 10^{-3} \text{ m}^3$$

$$10^3 \text{ L} = 1 \text{ m}^3$$

Some Base Units and Their Abbreviations for the International System of Measurement

Type	Base Unit	Abbreviation
Length	meter	m
Mass	gram	g
Volume	liter	L or l
Energy	joule	J

Metric Prefixes

Prefix	Abbreviation	Number
giga	G	10^9 or 1,000,000,000
mega	M	10^6 or 1,000,000
kilo	k	10^3 or 1000
centi	c	10^{-2} or 0.01
milli	m	10^{-3} or 0.001
micro	μ	10^{-6} or 0.000001
nano	n	10^{-9} or 0.000000001
pico	p	10^{-12} or 0.000000000001

Scientific Notation

- Numbers expressed in scientific notation have the following form.

$$a \times 10^b$$

Coefficient,
a number with one nonzero digit
to the left of the decimal point

Exponent, a positive or negative integer

Exponential term

Scientific Notation (Example)

- 5.5×10^{21} carbon atoms in a 0.55 carat diamond.
 - 5.5 is the coefficient
 - 10^{21} is the exponential term
 - The 21 is the exponent.
- The coefficient usually has one nonzero digit to the left of the decimal point.

Uncertainty



- The coefficient reflects the number's uncertainty.
- It is common to assume that coefficient is plus or minus one in the last position reported unless otherwise stated.
- Using this guideline, 5.5×10^{21} carbon atoms in a 0.55 carat diamond suggests that there are from 5.4×10^{21} to 5.6×10^{21} carbon atoms in the stone.

Size (Magnitude) of Number

- The exponential term shows the size or magnitude of the number.
- Positive exponents are used for large numbers. For example, the moon orbits the sun at 2.2×10^4 or 22,000 mi/hr.


$$2.2 \times 10^4 = 2.2 \times 10 \times 10 \times 10 \times 10 = 22,000$$

Size (Magnitude) of Number

- Negative exponents are used for small numbers. For example, A red blood cell has a diameter of about 5.6×10^{-4} or 0.00056 inches.

$$5.6 \times 10^{-4} = 5.6 \times \frac{1}{10^4} = \frac{5.6}{10 \times 10 \times 10 \times 10} = 0.00056$$

From Decimal Number to Scientific Notation



- Shift the decimal point until there is one nonzero number to the left of the decimal point, counting the number of positions the decimal point moves.
- Write the resulting coefficient times an exponential term in which the exponent is positive if the decimal point was moved to the left and negative if the decimal position was moved to the right. The number in the exponent is equal to the number of positions the decimal point was shifted.

From Decimal Number to Scientific Notation (Examples)

- For example, when 22,000 is converted to scientific notation, the decimal point is shifted four positions to the left so the exponential term has an exponent of 4.

$$22,000 = 2.2 \times 10^4$$

- When 0.00056 is converted to scientific notation, the decimal point is shifted four positions to the right so the exponential term has an exponent of -4.

$$0.00056 = 5.6 \times 10^{-4}$$

Scientific Notation to Decimal Number

- Shift the decimal point in the coefficient to the right if the exponent is positive and to the left if it is negative.
- The number in the exponent tells you the number of positions to shift the decimal point.

2.2×10^4 goes to 22,000

5.6×10^{-4} goes to 0.00056

Multiplying Exponential Terms

- When multiplying exponential terms, add exponents.

$$10^3 \times 10^6 = 10^{3+6} = 10^9$$

$$10^3 \times 10^{-6} = 10^{3+(-6)} = 10^{-3}$$

$$\begin{aligned} 3.2 \times 10^{-4} \times 1.5 \times 10^9 \\ &= 3.2 \times 1.5 \times 10^{-4+9} \\ &= 4.8 \times 10^5 \end{aligned}$$

When dividing exponential terms, subtract exponents.

$$\frac{10^{12}}{10^3} = 10^{12-3} = 10^9$$

$$\frac{10^6}{10^{-3}} = 10^{6-(-3)} = 10^9$$

$$\frac{9.0 \times 10^{11}}{1.5 \times 10^{-6}} = \frac{9.0}{1.5} \times 10^{11-(-6)} = 6.0 \times 10^{17}$$

$$\frac{10^2 \cdot 10^{-3}}{10^6} = 10^{2+(-3)-6} = 10^{-7}$$

$$\frac{1.5 \times 10^4 \cdot 4.0 \times 10^5}{2.0 \times 10^{12} \cdot 10^3} = \frac{1.5 \cdot 4.0}{2.0} \times 10^{4+5-12-3} = 3.0 \times 10^{-6}$$


Raising Exponential Terms to a Power

- When raising exponential terms to a power, multiply exponents.


$$(10^4)^3 = 10^{4 \cdot 3} = 10^{12}$$

$$(3 \times 10^5)^2 = (3)^2 \times (10^5)^2 = 9 \times 10^{10}$$


Length



A mile is four times around a typical high school track.



1 meter




1 centimeter




1 millimeter

Range of Lengths



Volume



1 fluid ounce (fl oz)



1 fl oz = 29.57 mL

1 mL = 0.03381 fl oz



1 milliliter (mL)
= about 20 drops

1 gal = 3.785 L



1 gallon (gal)
or 4 quarts (qt)

1 qt = 0.9464 L




1 qt or 32 fl oz

1 L = 1.057 qt
= 0.2642 gal



1 liter (L)
or 1000 mL

Range of Volumes



Mass and Weight




- **Mass** is usually defined as a measure of the amount of matter in an object. **Mass** can be defined as the property of matter that leads to gravitational attractions between objects and therefore gives rise to weight.
- **Matter** is anything that occupies a volume and has a mass.
- The **weight** of an object, on the Earth, is a measure of the force of gravitational attraction between the object and the Earth.

Comparison of the Mass and Weight of a 65 kg Person

Between Earth and Moon

	On Earth	Between Earth and Moon	On Moon
Mass	65 kg	65 kg	65 kg
Weight	637 N	≈ 0 N	$\frac{1}{6}(637 \text{ N})$ $= 106 \text{ N}$

Mass



$$1 \text{ lb} = 453.6 \text{ g}$$

$$1 \text{ kg} = 2.205 \text{ lb}$$

$$1 \text{ Mg} = 1000 \text{ kg} = 1 \text{ t}$$

$$1 \text{ oz} = 28.35 \text{ g}$$



About 2.5 grams (g) or
about 0.088 ounce (oz)




About 1 kilogram (kg) or
about 2.2 pounds (lb)





About 1 megagram (Mg) or 1 metric ton (t)


Range of Masses



Celsius and Fahrenheit Temperature



Comparing Temperature Scales



Precision and Accuracy



- **Precision** describes how closely a series of measurements of the same object resemble each other. The closer the measurements are to each other, the more precise the measurement. The precision of a measurement is not necessarily equal to its accuracy.
- **Accuracy** is a measurement's relationship to the property's true value.

Precision and Accuracy (cont.)



This archer is precise but not accurate.



This archer is precise and accurate.




This archer is imprecise and inaccurate.

Reporting Values from Measurements




- One of the conventions that scientists use for reporting numbers from measurements is to report all of the certain digits and one estimated (and thus uncertain) digit.

Graduated Cylinder




— Comparing the position of the bottom of the meniscus and the milliliter scale yields a measurement of 8.74 mL.

Graduated Cylinder Accurate to ± 0.1




— If the graduated cylinder is only accurate to ± 0.1 mL, we report 8.7 mL.

Trailing Zeros



— We report 8.00 mL to show an uncertainty of ± 0.01 mL.

Trailing Zeros (2)



— If the graduated cylinder is only accurate to ± 0.1 mL, we report 8.0 mL.

Digital Readout



Report all digits unless
otherwise instructed.

Digital Readout (2)



In many cases, it is best to round the number in the value to fewer decimal positions than displayed. For the mass displayed above, 100.432 g would indicate ± 0.001 g.