



Programme national

# Eolien et Biodiversité

LPO – ADEME - MTES

## Bibliographie

---

Programme Eolien et Biodiversité

### *Avifaune*

Geoffroy Marx – [geoffroy.marx@lpo.fr](mailto:geoffroy.marx@lpo.fr)

25/11/2020

#### *Légende*

- Eolien terrestre
- Eolien en mer

**ADEME**



Agence de l'Environnement  
et de la Maîtrise de l'Energie



# Avifaune

---

- Adams J, Kelsey EC, Felis JJ, Pereksta DM (2016). Collision and Displacement Vulnerability among Marine Birds of the California Current System Associated with Offshore Wind Energy Infrastructure. [📄](#) (p126, en)
- Al Zohbi G, Hendrick P, Bouillard P (2015). Evaluation of the impact of wind farms on birds: The case study of Lebanon. [📄](#) (p8, en)
- Allison TD, Butryn R (2019). A Summary of Bird Fatality Data in a Nationwide Database. [📄](#) (p46, en)
- Amec Wind Limited (2005). Offshore wind turbines and birds activity at Blyth. [📄](#) (p26, en)
- Arikan K, Turan SL (2017). Estimation of bird fatalities caused by wind turbines in Turkey. [📄](#) (p7, en)
- Arnett EB (2006). A preliminary evaluation on the use of dogs to recover bat fatalities at wind energy facilities. [📄](#) (p5, en)
- Arnett EB, May RF (2016). Mitigating wind energy impacts on wildlife: approaches for multiple taxa. [📄](#) (p14, en)
- Aschwanden J, Liechti F (2016). Vogelzugintensität und Anzahl Kollisionsopfer an Windenergieanlagen am Standort Le Peuchapatte (JU). [📄](#) (p74, de)
- Aschwanden J, Wanner S, Liechti F (2015). Investigation on the effectivity of bat and bird detection at a wind turbine: Final Report Bird Detection. [📄](#) (p35, en)
- Aschwanden J, Stark H, Peter D, Steuri T, Schmid B, Liechti F (2018). Bird collisions at wind turbines in a mountainous area related to bird movement intensities measured by radar. [📄](#) (p9, en)
- Aumüller R et al. (2011). Bird migration studied with remote sensing at FINO1 and FINO3. (p18, en)
- Aumüller R et al. (2013). Weichen Zugvögel Windenergieanlagen auf See aus? Eine Methode zur Untersuchung und Analyse von Reaktionen tagsüber ziehender Vogelarten auf Offshore-Windparks. [📄](#) (p11, de)
- Aumüller R et al. (2012). Beschreibung eines Vogelschlagereignisses und seiner Ursachen an einer Forschungsplattform in der Deutschen Bucht. [📄](#) (p8, de)
- Avery M et al. (1977). Weather influences on nocturnal bird mortality at a north dakota tower. [📄](#) (p9, en)
- Avery M et al. (1976). The effects of a tall tower on nocturnal bird migration - a portable ceilometer study. [📄](#) (p11, en)
- Avian Subcommittee of the National Wind Coordinating Committee (2001). Proceedings of the National Avian - Wind Power Planning Meeting IV, Carmel, CA, May 16-17, 2000. [📄](#) (p183, en)
- Avian Subcommittee of the National Wind Coordinating Committee (2000). Proceedings of national avian – wind power planning meeting III. San Diego, California, May 1998. [📄](#) (p206, en)
- Balotari-Chiebao F, Brommer JE, Niinimäki T, Laaksonen T (2015). Proximity to wind-power plants reduces the breeding success of the white-tailed eagle. [📄](#) (p8, en)
- Balotari-Chiebao F, Villers A, Ijäs A, Ovaskainen O, Repka S, Laaksonen T (2016). Post-fledging movements of white-tailed eagles: Conservation implications for wind-energy development. [📄](#) (p10, en)
- Band B (2012). Using a collision risk model to assess bird collision risks for offshore windfarms. [📄](#) (p62, en)
- Barclay RMR, Baerwald EF, Gruver JC (2007). Variation in bat and bird fatalities at wind energy facilities: assessing the effects of rotor size and tower height. [📄](#) (p8, en)
- Barrios L et al. (2004). Behavioural and environmental correlates of soaring-bird mortality at on-shore wind turbines. [📄](#) (p10, en)
- Bellebaum J, Korner-Nievergelt F, Dürr T, Mammen U (2013). Wind turbine fatalities approach a level of concern in a raptor population. [📄](#) (p7, en)
- Bennett E (2012). Observations from the Use of Dogs to Undertake Carcass Searches at Wind Facilities in Australia. [📄](#) (p10, en)
- Berg S, Iserer F, Jurczyk M, Fronczek S, Reischke N, Jung C, Braun D, Thielen D (2018). Analysis of black stork flight behaviour under different weather and land-use conditions with special consideration of existing wind turbines in the Vogelsberg SPA. [📄](#) (p219, en)
- Beroud T, Druais J, Bay Y, Ricci JC (2017). Visual counts, bioacoustics and RADAR: three methods to study waterfowl pre-nuptial migration in Southern France. [📄](#) (p14, en)

- Beston JA, Diffendorfer JE, Loss SR (2015). Insufficient Sampling to Identify Species Affected by Turbine Collisions. [📄](#) (p5, en)
- Beston JA, Diffendorfer JE, Loss SR, Johnson DH (2016). Prioritizing Avian Species for Their Risk of Population-Level Consequences from Wind Energy Development. [📄](#) (p19, en)
- Biosis Research Pty (2006). Wind farm collision risk for birds cumulative risks for threatened and migratory species. [📄](#) (p237, en)
- Biotope (2009). Étude de la migration postnuptiale par radar - Année 2008. [📄](#) (p86, fr)
- Bird Studies Canada, Canadian Wind Energy Association, Environment Canada and Ontario Ministry of Natural Resources (2018). Base de données du Suivi des populations d'oiseaux et de chauves-souris relié à l'énergie éolienne -2018. [📄](#) (p76, fr)
- Bird Studies Canada, Canadian Wind Energy Association, Environment Canada and Ontario Ministry of Natural Resources (2017). Base de données du Suivi des populations d'oiseaux et de chauves-souris relié à l'énergie éolienne - Juillet 2017. [📄](#) (p51, fr)
- Bird Studies Canada, Canadian Wind Energy Association, Environment Canada and Ontario Ministry of Natural Resources (2016). Base de données du Suivi des populations d'oiseaux et de chauves-souris relié à l'énergie éolienne - Juillet 2016. [📄](#) (p48, fr)
- Birdlife International (2015). Review and guidance on use of "shutdown-on-demand" for wind turbines to conserve migrating soaring birds in the Rift Valley/Red Sea Flyway. [📄](#) (p49, en)
- Blache S, Loose D - CORA Drôme, LPO Isère (2008). Sensibilité des busards aux parcs éoliens - Rapport technique années 2006-2007. [📄](#) (p50, fr)
- Black JE (2000). Application of weather radar to monitoring numbers of Birds in Migration. [📄](#) (p3, en)
- Bolker Ethan D., Hatch Jeremy J., Zara Catalin (2014). Modeling how windfarm geometry affects bird mortality. [📄](#) (p18, en)
- Bose a, Dürr T, Klende RA, Henle K (2019). Assessing the spatial distribution of avian collision risks at wind turbine structures in Brandenburg, Germany. [📄](#) (p21, en)
- Bose A, Dürr T, Klenke RA, Henle K (2018). Collision sensitive niche profile of the worst affected bird-groups at wind turbine structures in the Federal State of Brandenburg, Germany. [📄](#) (p13, en)
- Boutin J-M - CNERA - ONCFS (2011). Relecture du rapport "Evaluation de l'impact du parc éolien du Rochereau (Vienne) sur l'avifaune de plaine. Rapport final 2007-2010, Thomas Williamson (2011)". [📄](#) (p7, Fr)
- Brabant R, Vanermen N, Steinen E, Degraer S (2015). Towards a Cumulative Collision Risk Assessment of Local and Migrating Birds in North Sea Offshore Wind Farms. [📄](#) (p12, en)
- Bradbury G, Trinder M, Furness B, Banks AN, Caldow RWG et al. (2014). Mapping Seabird Sensitivity to Offshore Wind Farms. [📄](#) (p17, en)
- Braunischa V, Coppesa J, Bächlea S, Suchanta R (2015). Underpinning the precautionary principle with evidence: A spatial concept for guiding wind power development in endangered species' habitats. [📄](#) (p10, en)
- Bretagnolle V, Denonfoux L, Villers A (2018). Are farming and birds irreconcilable? A 21-year study of bustard nesting ecology in intensive agroecosystems. [📄](#) (p9, en)
- Bruderer B (1996). Vogelzugforschung im Bereich der Alpen 1980-1995. [📄](#) (p12, de)
- Bruderer B, Liechti F (2004). Welcher Anteil ziehender vögel fliegt im Höhenbereich von Windturbinen. (p9, de)
- Burton NHK, Thaxter CB, Cook ASCP, Austin GE, Wright LJ, Clark NA (2010). Evaluating the potential impacts of tidal power schemes on estuarine waterbirds. [📄](#) (p3, en)
- Busch M, Garthe S (2015). Approaching population thresholds in presence of uncertainty: Assessing displacement of seabirds from offshore wind farms. [📄](#) (p12, en)
- Busch M, Kannen A, Garthe S, Jessopp M (2012). Consequences of a cumulative perspective on marine environmental impacts: Offshore wind farming and seabirds at North Sea scale in context of the EU Marine Strategy Framework Directive. [📄](#) (p12, en)
- Busch M, Trautmann S, Gerlach B (2017). Overlap between breeding season distribution and wind farm risks: a spatial approach. [📄](#) (p12, en)
- Cabrera-Cruz SA, Villegas-Patracá R (2016). Response of migrating raptors to an increasing number of wind farms. [📄](#) (p15, en)
- Cadiou B et al. (2011). Cinquième recensement national des oiseaux marins nicheurs en France métropolitaine 2009-2011. 1° synthèse: bilan intermédiaire 2009-2010. [📄](#) (p62, fr)
- Camiña Á (2013). Guidelines for wind farm planning on birds and bats: Lessons learned from operating wind farms. [📄](#) (p40, en)
- Camiña Á (2007). Energía eólica y buitres negros. [📄](#) (p10, es)
- Camiña Á (2006). Las energías renovables y la conservación de aves carroñeras: el caso del buitre leonado (gyps fulvus) en el norte la península ibérica. [📄](#) (p10, es)
- Cano-Alonso LS, Strazds M, Tamás EA (2019). VII International conference on Black stork Ciconia nigra.

Conference Plenary Discussion, Conclusions and Recommendations of the VII International Conference on Black Stork *Ciconia nigra* (Doñana National Park, Spain). [📄](#) (p6, en)

- Carretea M, Sánchez-Zapatab JA, Benítezc JR, Lobónc M, Donázara JA (2009). Large scale risk-assessment of wind-farms on population viability of a globally endangered long-lived raptor. [📄](#) (p8, en)
- Carrizo et al. (2010). Understanding geographical range changes of North American breeding birds. [📄](#) (p1, en)
- Chamberlain DE et al. (2006). The effect of avoidance rates on bird mortality predictions made by wind turbine collision risk models. [📄](#) (p5, en)
- Choi DY et al. (2020). An evaluation of bird and bat mortality at wind turbines in the Northeastern United States. [📄](#) (p22, en)
- Cleasby IR, Wakefield ED, Bearhop S, Bodey TW, Votier SC, Hamer KC (2015). Three-dimensional tracking of a wide-ranging marine predator: flight heights and vulnerability to offshore wind farms. [📄](#) (p9, en)
- Collier MP, Dirksen S, Krijgsveld KL (2012). A review of methods to monitor collisions or micro-avoidance of birds with offshore wind turbines. Part 2 : Feasibility study of systems to monitor collisions. [📄](#) (p30, en)
- Collier MP, Dirksen S, Krijgsveld KL (2011). A review of methods to monitor collisions or micro-avoidance of birds with offshore wind turbines - Strategic Ornithological Support Services Project SOSS-03A. [📄](#) (p38, en)
- Cook A, Collier M (2015). Understanding avian collision rate modelling and application at the population level. [📄](#) (p45, en)
- Cook ASCP, Burton NHK (2010). A review of the potentiel impacts of Marine Aggregate extraction on Seabirds. [📄](#) (p114, en)
- Cook ASCP, Humphreys Elizabeth M, Masden Elizabeth A, Burton Niall HK (2014). The Avoidance Rates of Collision Between Birds and Offshore Turbines. [📄](#) (p263, en)
- Cook ASCP, Ward RM, Hansen WS, Larsen L (2018). Estimating Seabird Flight Height Using LiDAR. [📄](#) (p66, en)
- Cook ASCP, Wright LJ, Burton NHK - BTO (2012). A Review of flight heights and avoidance rates of birds in relation to offshore windfarms. [📄](#) (p61, en)
- Cornelis N, Strehler-Perrin C, Balsiger L et al. (2016). Evaluation des impacts résiduels cumulés des éoliennes du Jura vaudois et des régions limitrophes sur la faune ailée. [📄](#) (p70, fr)
- Dahl EL, May R, Nygård T, Åstrøm J, Diserud OH (2015). Repowering Smøla wind-power plant. An assessment of avian conflicts. [📄](#) (p46, en)
- de Lucas M et al. (2008). Collision fatality of raptors in wind farms does not depend on raptor abundance. [📄](#) (p9, en)
- de Lucas M, Ferrer M, Bechard MF, Muñoz AR (2012). Griffon vulture mortality at wind farms in southern Spain: Distribution of fatalities and active mitigation measures. [📄](#) (p6, en)
- de Lucas M, Janss GFE, Ferrer M (2004). The effects of a wind farm on birds in a migration point: the Strait of Gibraltar. [📄](#) (p13, en)
- Delprat B (2017). Projet éolien de la Combe Rougeux (52) - Evaluation du risque de collision Milan royal. [📄](#) (p29, fr)
- Department for Business Enterprise & Regulatory Reform (2007). Aerial Surveys of waterbirds in strategic wind farm areas: 2005/2006 Final report. [📄](#) (p180, en)
- Department Of Trade et Industry (2006). Aerial Surveys of waterbirds in strategic wind farm areas: 2004/2005 Final report. [📄](#) (p176, en)
- Desholm M (2009). Avian sensitivity to mortality: prioritising migratory bird species for assessment at proposed wind farms. [📄](#) (p8, en)
- Desholm M (2005). TADS investigations of avian collision risk at Nysted offshore wind farm, autumn 2004. [📄](#) (p31, en)
- Desholm M, Kahlert J (2005). Avian collision risk at an offshore wind farm. [📄](#) (p3, en)
- Devereux CL, Denny MJH, Whittingham MJ (2008). Minimal effects of wind turbines on the distribution of wintering farmland birds. [📄](#) (p6, en)
- Devoucoux P, Besnard A, Bretagnolle V (2018). Sex-dependent habitat selection in a high-density Little Bustard *Tetrax tetrax* population in southern France, and the implications for conservation. [📄](#) (p10, en)
- Dirksen S (2017). Review of methods and techniques for field validation of collision rates and avoidance amongst birds and bats at offshore wind turbines. [📄](#) (p48, en)
- Dohm R, Jennelle CS, Garvin JC, Drake D (2019). A long-term assessment of raptor displacement at a wind farm. [📄](#) (p6, en)
- Dokter A et al. (2010). Automated bird migration quantification by weather radar networks. [📄](#) (p22, en)
- Dokter AM et al. (2010). Bird migration flight altitudes studied by a network of operational weather radars. [📄](#) (p15, en)
- Dooling R (2002). Avian hearing and the Avoidance of Wind Turbines. [📄](#) (p84, en)
- Douglas DJT, Bellamy PE, Pearce-Higgins JW (2011). Changes in the abundance and distribution of upland breeding birds at an operational wind farm. [📄](#) (p7, en)

- ● Drewitt AL et al. (2006). Assessing the impacts of wind farms on birds. [📄](#) (p14, en)
- Drewitt AL, Langston RH (2008). Collision Effects of Wind-power Generators and Other Obstacles on Birds. [📄](#) (p34, en)
- Duckworth J et al. (2020). Red-throated Diver Energetics Project: Preliminary Results from 2018/19. [📄](#) (p36, en)
- Duriez O, Pilard P, Saulnier N, Bouzin M, Boudarel P, Besnard A (2018). Estimation de l'impact démographique du Parc éolien d'Aumelas sur la population héraultaise de Faucons crécerellettes. [📄](#) (p17, fr)
- Dürr T (2020). Vogelverluste an Windenergieanlagen / bird fatalities at windturbines in Europe. [📄](#) (p8, de)
- Erickson W, Johnson G, Young D, Strickland D, Good R, Bourassa M, Bay K, Sernka K (2002). Synthesis and Comparison of baseline avian and Bat use, raptor nesting and Mortality Information from proposed and existing wind developments. [📄](#) (p129, en)
- Erickson WP, Johnson GD, Strickland DM, Young DP Jr., Sernka KJ, Good RE (2001). Avian Collisions with Wind Turbines: A Summary of Existing Studies and Comparisons to Other Sources of Avian Collision Mortality in the United States. [📄](#) (p67, en)
- Erickson WP, Johnson GD, Young DP Jr. (2005). A Summary and Comparison of Bird Mortality from Anthropogenic Causes with an Emphasis on Collisions. [📄](#) (p14, en)
- Erickson WP, Strickland MD, Johnson GD, Kern JW (2000). Examples of Statistical Methods to Assess Risk of Impacts to Birds from Wind Plants. [📄](#) (p12, en)
- Erickson WP, Wolfe MM, Bay KJ, Johnson DH, Gehring JL (2014). A Comprehensive Analysis of Small-Passerine Fatalities from Collision with Turbines at Wind Energy Facilities. [📄](#) (p18, en)
- European Commission, Science For Environment Policy, Dg Environment News Alert Service (2009). Which birds are most at risk from wind farm collisions. [📄](#) (p1, en)
- Everaert J (2014). Collision risk and micro-avoidance rates of birds with wind turbines in Flanders. [📄](#) (p11, en)
- ● Everaert J, Kuijken E (2007). Wind turbines and birds in Flanders (Belgium): preliminary summary of the mortality research results. [📄](#) (p10, en)
- Everaert J, Stienen EWM (2006). Impact of wind turbines on birds in Zeebrugge (Belgium): Significant effect on breeding tern colony due to collisions. [📄](#) (p14, en)
- Exo K-M et al. (2003). Birds and offshore wind farms: a hot topic in marine ecology. [📄](#) (p4, en)
- Falavigna TJ et al. (2020). Changes in bird species composition after a wind farm installation: A case study in South America. [📄](#) (p7, en)
- Farfán MA et al. (2017). Differential recovery of habitat use by birds after wind farm installation: A multi-year comparison. [📄](#) (p8, en)
- Farque P - LPO (2013). Interactions entre oiseaux marins et parcs éolien en mer : connaissances, contexte et solutions sur les côtes françaises. [📄](#) (p56, fr)
- Fernández-Bellon D, Wilson MW, Irwin S, O'Halloran J (2018). Effects of development of wind energy and associated changes in land use on bird densities in upland areas. [📄](#) (p32, en)
- ● Fernandez-Bellon, D (2020). Limited accessibility and bias in wildlife-wind energy knowledge: A bilingual systematic review of a globally distributed bird group. [📄](#) (p11, en)
- Fielding AH et al. (2006). Spatial association as an indicator of the potential for future interactions between wind energy developments and golden eagles *Aquila chrysaetos* in Scotland. [📄](#) (p11, en)
- Fijn R, Krijgsveld K, Poot M, Dirksen S (2015). Bird Movements at Rotor Heights Measured Continuously with Vertical Radar at a Dutch Offshore Wind Farm. [📄](#) (p15, en)
- Fijn RC, Gyimesi A (2018). Behaviour related flight speeds of Sandwich Terns and their implications for wind farm collision rate modelling and impact assessment. [📄](#) (p5, en)
- Follestad A, Flagstad Ø, Nygård T, Reitan O, Schulze J (2007). Vindkraft og fugl på Smøla 2003–2006. [📄](#) (p82, no)
- Fox AD, Desholm M, Kahlert J, Christensen TK, Krag PIB (2006). Information needs to support environmental impact assessment of the effects of European marine offshore wind farms on birds. [📄](#) (p16, en)
- Furness R, Garthe S, Trinder M, Matthiopoulos J, Wanless S, Jeglinski J (2018). Nocturnal flight activity of northern gannets *Morus bassanus* and implications for modelling collision risk at offshore wind farms. [📄](#) (p6, en)
- Furness RW et al. (2013). Potential population-level consequences of marine renewables. [📄](#) (p2, en)
- Furness RW, Wade HM, Masden EA (2013). Assessing vulnerability of marine bird populations to offshore wind farms. [📄](#) (p11, en)
- Garcia DA, Canavero G, Ardenghi F, Zambon M (2015). Analysis of wind farm effects on the surrounding environment: Assessing population trends of breeding passerines. [📄](#) (p7, en)
- Garthe S, Hüppop O (2004). Scaling possible adverse effects of marine wind farms on seabirds: developing and applying a vulnerability index. [📄](#) (p4, en)
- Garthe SG, Markones Nele, Corman A-M (2017). Possible impacts of offshore wind farms on seabirds: a pilot study in Northern Gannets in the southern North Sea. [📄](#) (p6, en)



- Garvin JC et al. (2010). Response of raptors to a windfarm. [📄](#) (p11, en)
- ● Gerringer MB, Lima SL, Devault TL (2015). Evaluation of an avian radar system in a midwestern landscape. [📄](#) (p10, en)
- Gitenet P - LPO Hérault (2013). Reproduction et mortalité du Busard cendré sur un parc éolien du sud de la France. [📄](#) (p6, fr)
- Gómez-Catasús J, Garza V, Traba J (2018). Wind farms affect the occurrence, abundance and population trends of small passerine birds: The case of the Dupont's lark. [📄](#) (p31, en)
- González MA, Ena V (2011). Cantabrian Capercaillie signs disappeared after a wind farm construction. [📄](#) (p10, en)
- Goodale M, Milman A (2020). Assessing Cumulative Exposure of Northern Gannets to Offshore Wind Farms. [📄](#) (p8, en)
- Goodale MW, Stenhouse IJ, Williams KA (2014). Reducing the Adverse Effects of Offshore Wind Development on Waterbirds in the Great Lakes: A Proposed Four-Step Approach. [📄](#) (p72, en)
- Govaere A, Devos S, Elleboode C, André Y, Delcourt V, Lagrange H (2008). Etudes des mouvements d'oiseaux par radar en France - Analyse des données existantes. [📄](#) (p56, fr)
- Graff BJ (2015). An assessment of direct mortality to avifauna from wind energy facilities in North Dakota and South Dakota. [📄](#) (p88, en)
- Griffin L, Rees E, Hughes B (2010). Whooper swan *Cygnus cygnus* migration in relation to offshore wind farms. [📄](#) (p3, en)
- Grünkorn T, von Rönn J, Blew J, Nehls G, Weitekamp S, Timmermann H, Reichenbach M, Coppack T (2016). Détermination des taux de collision des oiseaux (y compris rapaces) et principes fondamentaux pour prévoir et évaluer le risque de collision dans la conception des projets éoliens (projet de recherche PROGRESS). [📄](#) (p27, fr)
- Hanagasioglu M, Aschwanden J, Bontadina F, Marcos de la Puente N (2015). Investigation of the effectiveness of bat and bird detection of the DTBat and DTBird systems at Calandawind turbine - Final Report. [📄](#) (p142, en)
- Hanssen F, May R, Nygård T (2020). High-Resolution Modeling of Uplift Landscapes can Inform Micrositing of Wind Turbines for Soaring Raptors. [📄](#) (p14, en)
- Harvey HT & Associates (2018). AWWI Technical Report: Evaluating a Commercial-Ready Technology for Raptor Detection and Deterrence at a Wind Energy Facility in California. [📄](#) (p96, en)
- ● Hausberger M, Boigné A, Lesimple C, Belin L, Henry L (2018). Wide-eyed glare scares raptors: From laboratory evidence to applied management. [📄](#) (p15, en)
- Heinänen S, Zydalis R, Kleinschmidt B, Dorsch M, Burger C, Morkunas J, Quillfeldt P, Nehls G (2020). Satellite telemetry and digital aerial surveys show strong displacement of red-throated divers (*Gavia stellata*) from offshore wind farms. [📄](#) (p36, en)
- Hernández-Pliego J, de Lucas M, Muñoz A-R, Ferrer M (2015). Effects of wind farms on Montagu's harrier (*Circus pygargus*) in southern Spain. [📄](#) (p7, en)
- Heuck C et al. (2019). Wind turbines in high quality habitat cause disproportionate increases in collision mortality of the white-tailed eagle. [📄](#) (p8, en)
- Hill R et al. (2011). Description of bird strike event and its causes at a research platform in the German Bight, North Sea. [📄](#) (p1, en)
- ● Hodos W (2003). Minimization of Motion Smear: Reducing Avian Collisions with Wind Turbines. [📄](#) (p43, en)
- Homoya W, Moore JW, Ruhl PJ, Dunning JB Jr (2017). Do American Golden-Plovers (*Pluvialis dominica*) Avoid Wind-energy Turbines in Agricultural Fields in Indiana During Spring Migration?. [📄](#) (p10, en)
- ● Horch P, Keller V (2005). Windkraftanlagen und vögel - ein Konflikt? Eine Literaturrecherche. [📄](#) (p63, de)
- Horswill C, O'Brien SH, Robinson RA (2016). Density dependence and marine bird populations: are wind farm assessments precautionary?. [📄](#) (p9, en)
- Hotker H (2006). The impact of repowering of wind farms on birds and bats. [📄](#) (p38, en)
- ● Hotker H - Compiler (2008). Birds of prey and Wind farms: analysis of problems and possible solutions. [📄](#) (p77, en)
- Hötker H et al. (2017). Birds of Prey and Wind Farms - Analysis of Problems and Possible Solutions. [📄](#) (p332, fr)
- Hötker H, Thomsen KM, Jeromin H (2006). Impacts on biodiversity of exploitation of renewable energy sources: the example of birds and bats. [📄](#) (p65, en)
- Hunt GW (2015). Addressing the Factors that Juxtapose Raptors and Wind Turbines. [📄](#) (p6, en)
- Hüppop O, Hüppop K, Dierschke J, Hill R (2016). Bird collisions at an offshore platform in the North Sea. [📄](#) (p10, en)
- Inchausti P, Bretagnolle V (2005). Predicting short-term extinction risk for the declining Little Bustard (*Tetrax tetrax*) in intensive agricultural habitats. [📄](#) (p10, en)
- Itty C, Duriez O (2018). Le suivi par GPS, une méthode efficace pour évaluer l'impact des parcs éoliens sur des espèces à fort enjeux de conservation : l'exemple de l'aigle royal (*Aquila chrysaetos*) dans le sud du massif central. [📄](#) (p7, fr)
- Jacobsen EM et al. (2019). Avoidance Behaviour of Migrating Raptors Approaching an Offshore Wind Farm. [📄](#) (p8, en)

- Jain A et al. (2007). Annual report for Maple Ridge Wind Power Project - Postconstruction bird and bat fatality study - 2006. [📄](#) (p61, en)
- Jenkins AR et al. (2018). Combining radar and direct observation to estimate pelican collision risk at a proposed wind farm on the Cape west coast, South Africa. [📄](#) (p12, en)
- Jones et al. (2009). Altamont Pass Wind Resource Area 48-Hour Search Interval Bird Fatality Study. [📄](#) (p22, en)
- Karaivanov N, Karaivanov P (2018). Research on nesting birds on the territory of Kaliakra Wind Farm. [📄](#) (p5, en)
- Katzenberger J, Sudfeld C (2019). Rotmilan une Windkraft: Negativer Zusammenhang zwischen WKA-Dichte und Bestandstrends. [📄](#) (p4, de)
- Katzner TE et al. (2016). Golden Eagle fatalities and the continental-scale consequences of local wind-energy generation. [📄](#) (p10, en)
- Kelly TA, Fielder JK (2008). A framework for mitigation of bird and bat strike risk at wind farms using avian radar and SCADA interface. [📄](#) (p7, en)
- Kelsey EC, Felis JJ, Czapanskiy M, Pereksta DM, Adams J (2018). Collision and displacement vulnerability to offshore wind energy infrastructure among marine birds of the Pacific Outer Continental Shelf. [📄](#) (p19, en)
- Khosravifard S et al. (2020). Identifying Birds' Collision Risk with Wind Turbines Using a Multidimensional Utilization Distribution Method. [📄](#) (p9, en)
- Kikuchi DM et al. (2019). Estimating Potential Costs of Cumulative Barrier Effects on Migrating Raptors: A Case Study Using Global Positioning System Tracking in Japan. [📄](#) (p15, en)
- Kitano M, Shiraki S (2013). Estimation of bird fatalities at wind farms with complex topography and vegetation in Hokkaido, Japan. [📄](#) (p8, en)
- Kleyheeg JC et al. (2018). Predicting bird collisions with wind turbines: Comparison of the new empirical Flux Collision Model with the SOSS Band model. [📄](#) (p10, en)
- Kolar PS, Bechard MJ (2016). Wind energy, nest success, and post-fledging survival of Buteo hawks. [📄](#) (p14, en)
- Korner-Nievergelt F, Brinkmann R, Niermann I, Behr O (2013). Estimating Bat and Bird Mortality Occurring at Wind Energy Turbines from Covariates and Carcass Searches Using Mixture Models. [📄](#) (p11, en)
- Korner-Nievergelt F, Brossard C, Filliger R, Gremaud J, Lugon A, Mermoud O, Schaub M, Wechsler S (2016). Effets cumulés des éoliennes du Jura vaudois et des régions limitrophes sur l'avifaune et les chiroptères : risque de collisions et de perte d'habitat pour quelques espèces d'oiseaux et de chiroptères. [📄](#) (p181, fr)
- Krijgsveld K, Dirksen S (2013). Avoidance behaviour and flight intensities of birds in an offshore wind farm in the Netherlands. [📄](#) (p1, en)
- Krijgsveld KL et al. (2011). Effect studies offshore wind farm Egmond aan Zee -Final report on fluxes, flight altitudes and behaviour of flying birds. [📄](#) (p334, en)
- Krijgsveld KL et al. (2010). Effect studies offshore wind farm Egmond aan Zee - Progress report on fluxes and behaviour of flying birds covering 2007-2008. [📄](#) (p112, en)
- Krijgsveld KL et al. (2009). Collision risk of birds with modern large wind turbines. [📄](#) (p10, en)
- Krone O, Treu G (2018). Movement patterns of white-tailed sea eagles near wind turbines. [📄](#) (p9, en)
- Krüger T, Garthe S (2001). Flight altitudes of coastal birds in relation to wind direction and speed. [📄](#) (p14, en)
- Kunz TH et al. (2007). Assessing impacts of wind-energy development on nocturnally active birds and bats: A guidance document. [📄](#) (p38, en)
- Lacrouts O, Dumoulin A - Diades Marine (2018). New technology for 3D bird detection radar. [📄](#) (p5, fr)
- Lagrange H, Rico P (2019). Evaluation of bird detection efficiency of ProBird - Tracking missed detection and false positives. [📄](#) (p19, en)
- Lalgola SM et al. (2020). Hazards of wind turbines on avifauna - a preliminary appraisal within the Indian context. [📄](#) (p14, en)
- Lampman G - New Jersey Audubon, Old Bird Inc and North East Ecological Services (2013). Post-construction wildlife monitoring at Maple Ridge wind farm - Final report. [📄](#) (p34, en)
- Lange CJ, Ballard BM, Collins DP (2018). Impacts of wind turbines on redheads in the Laguna Madre. [📄](#) (p7, en)
- Langston RHW (2013). Birds and wind projects across the pond: A UK perspective. [📄](#) (p14, en)
- Langston RHW (2010). Birds and Wind farms: where next?. [📄](#) (p4, en)
- Langston RHW (2010). Offshore wind farms and birds: Round 3 zones, extensions to Round 1 & Round 2 sites & Scottish Territorial Waters. [📄](#) (p42, en)
- Langston RHW, Pullan JD - RSPB (2004). Effects of wind farms on birds. [📄](#) (p91, en)
- Langston RHW, Teuten E (2011). Foraging ranges of northern gannets *Morus bassanus* in relation to proposed offshore wind farms in the UK:2011. [📄](#) (p19, en)
- Lapointe J et al. (2015). Habitat du faucon pèlerin dans le sud du Québec durant la période de reproduction : conséquences pour l'implantation de parcs éoliens. [📄](#) (p9, fr)
- Larsen JK, Guillemette M (2007). Effects of wind turbines on flight behaviour of wintering common



- eiders: implications for habitat use and collision risk. [\(p9, en\)](#)
- Leducq I - LPO Franche-Comté (2010). Les flux migratoires ornithologiques en Franche-Comté. Analyse et interprétation des données de Observations de la base 2010. [\(p31, fr\)](#)
  - Leopold M (2013). Are auks the ideal birds to study offshore wind farm impact on seabirds?. [\(p1, en\)](#)
  - Leopold MF, Verdaat HJP (2018). Pilot field study: observations from a fixed platform on occurrence and behaviour of common guillemots and other seabirds in offshore wind farm Luchterduinen. [\(p28, en\)](#)
  - Lie Dahl E et al. (2012). Reduced breeding success in white-tailed eagles at Smøla windfarm, western Norway, is caused by mortality and displacement. [\(p7, en\)](#)
  - Liechti F, Guélat J, Komenda-Zehnder S (2013). Modelling the spatial concentrations of bird migration to assess conflicts with wind turbines. [\(p9, en\)](#)
  - Litsgård F, Eriksson A, Wizelius T, Säfström T (2016). Pilotinstallation av DTBird-systemet i Sverige. Möjligheter med skyddssystem för fågelfaunan vid vindkraftanläggningar – erfarenheter från Sveriges första installation av DTBird. [\(p43, su\)](#)
  - Lonsdorf E, Sanders-Reed CA, Boal C, Allison TD (2018). Modeling golden eagle-vehicle collisions to design mitigation strategies. [\(p12, en\)](#)
  - Loss SR, Will T, Marra PP (2015). Direct Mortality of Birds from Anthropogenic Causes. [\(p22, en\)](#)
  - Loss SR, Will T, Marra PP (2013). Estimates of bird collision mortality at wind facilities in the contiguous United States. [\(p9, en\)](#)
  - Lovich JE (2015). Golden eagle mortality at a wind-energy facility near palm springs, California. [\(p6, en\)](#)
  - Luceole (2012). Coexistence Milan royal et parc éolien - pour une compréhension ouverte d'un problème complexe. Actes du séminaire citoyen Luceole. [\(p48, fr\)](#)
  - Lumsden LF, Moloney PD, Smales I (2019). Developing a science-based approach to defining key species of birds and bats of concern for wind farm developments in Victoria. [\(p49, en\)](#)
  - Mabey S, Paul E (2007). Impact of wind energy and related human activities on grassland and shrub-steppe birds (literature review). [\(p183, en\)](#)
  - Mackenzie A, Perrow M (2013). Individual-based population viability modelling as a tool to quantify the effect of wind farm collisions. [\(p1, en\)](#)
  - Mackenzie M et al. (2013). Wind farm displacement studies from Nysted, Denmark. [\(p1, en\)](#)
  - Madders M, Whitfield DP (2006). Upland raptors and the assessment of wind farm impacts. [\(p14, en\)](#)
  - Maggini R et al. (2010). Modelling range shifts for Swiss breeding birds. [\(p3, en\)](#)
  - Marcelino J, et al. (2017). Tracking data of the Little Bustard Tetrax tetrax in Iberia shows high anthropogenic mortality. [\(p12, en\)](#)
  - Markid L (2012). Does offshore wind farm development result in habitat loss for seabirds?. [\(p34, en\)](#)
  - Marques AT et al. (2019). Wind turbines cause functional habitat loss for migratory soaring birds. [\(p31, en\)](#)
  - Marques J, Rodrigues L, João Silva M, Santos J, Bispo R, Bernardino J (2018). Estimating Bird and Bat Fatality at Wind Farms: From Formula-Based Methods to Models to Assess Impact Significance. [\(p54, en\)](#)
  - Martin C (2015). Effectiveness of operational mitigation in reducing bat mortality and an assessment of bat and bird fatalities at the Sheffield Wind Facility, Vermont. [\(p196, en\)](#)
  - Martin GR (2010). Bird collisions: a visual or a perceptual problem?. [\(p4, en\)](#)
  - Martin GR, Shaw JM (2010). Bird collisions with power lines: Failing to see the way ahead?. [\(p8, en\)](#)
  - Martínez-Marivela I et al. (2018). Traffic Noise Pollution Does Not Influence Habitat Selection in the Endangered Little Bustard. [\(p11, en\)](#)
  - Marx G - LPO France (2018). Le parc éolien français et ses impacts sur l'avifaune - Etude des suivis de mortalité réalisés en France de 1997 à 2015. [\(p7, fr\)](#)
  - Marx G - LPO France (2017). Le parc éolien français et ses impacts sur l'avifaune - Etude des suivis de mortalité réalisés en France de 1997 à 2015. [\(p92, fr\)](#)
  - Masden E (2015). Developing an Avian Collision Risk Model to Incorporate Variability and Uncertainty. [\(p47, en\)](#)
  - Masden E (2010). Assessing the Cumulative Impacts of Wind Farms on Birds. [\(p142, en\)](#)
  - Masden EA et al. (2009). Cumulative impact assessments and bird / windfarm interactions: developing a conceptual framework. [\(p7, en\)](#)
  - Maurer WG (2018). Dancer in the Wind Farm: Automated Eagle Monitoring and Deterrence. [\(p90, en\)](#)
  - May R (2015). A Unifying Framework for the Underlying Mechanisms of Avian Avoidance of Wind Turbines. [\(p9, en\)](#)
  - May R et al. (2015). Mitigating wind-turbine induced avian mortality: Sensory, aerodynamic and cognitive constraints and options. [\(p12, en\)](#)
  - May R, Åström J, Hamre O, Dahl E (2017). Do birds in flight respond to (ultra)violet lighting?. [\(p10, en\)](#)

- May R, Hamre O, Vang R, Nygård T (2012). Evaluation of the DTBird video-system at the Smøla wind-power plant. Detection capabilities for capturing near-turbine avian behaviour. [📄](#) (p32, en)
- ● May R, Nygård T, Falkdalen U, Åström J, Hamre Ø, Stokke BG (2020). Paint it black: Efficacy of increased wind turbine rotor blade visibility to reduce avian fatalities. [📄](#) (p9, en)
- May R, Steinheim Y, Kvaløy P, Vang R, Hanssen F (2017). Performance test and verification of an off-the-shelf automated avian radar tracking system. [📄](#) (p9, en)
- ● McCann D, Bell P (2017). Visualising the Aspect-Dependent Radar Cross Section of Seabirds over a Tidal Energy Test Site Using a Commercial Marine Radar System. [📄](#) (p8, en)
- ● McClure CJW et al. (2018). State of the world's raptors: Distributions, threats, and conservation recommendations. [📄](#) (p14, en)
- ● McClure CJW, Martinson L, Allison TD (2018). Automated monitoring for birds in flight: Proof of concept with eagles at a wind power facility. [📄](#) (p8, en)
- ● McDonald C et al. (2013). Displacement effects modelling for central place foraging auks with the Forth/Tay as a case study. [📄](#) (p1, en)
- McIsaac HP (2000). Raptor Acuity and Wind Turbine Blade Conspicuity. [📄](#) (p29, en)
- McNew LB, Hunt LM, Gregory AJ, Wisely SM, Sandercock BK (2014). Effects of Wind Energy Development on Nesting Ecology of Greater Prairie-Chickens in Fragmented Grasslands. [📄](#) (p11, en)
- ● Mendel B et al. (2019). Operational offshore wind farms and associated ship traffic cause profound changes in distribution patterns of Loons (*Gavia* spp.). [📄](#) (p10, en)
- Miao R et al. (2019). Effect of wind turbines on bird abundance: A national scale analysis based on fixed effects models. [📄](#) (p10, en)
- Minderman J et al. (2014). Estimates and correlates of bird and bat mortality at small wind turbine sites. [📄](#) (p16, en)
- Montes RM (1995). Incidencia de las plantas de aerogeneradores sobre la avifauna en la comarca del Campo de Gibraltar. [📄](#) (p13, es)
- Morales MB, García JT, Arroyo B (2005). Can landscape composition changes predict spatial and annual variation of little bustard male abundance?. [📄](#) (p8, en)
- Moriguchi S, Mukai H, Komachi R, Sekijima T (2019). Wind Farm Effects on Migratory Flight of Swans and Foraging Distribution at Their Stopover Site. [📄](#) (p9, en)
- Nadaï A, Labussière O (2010). Birds, Wind and the Making of Wind Power Landscapes in Aude, Southern France. [📄](#) (p20, en)
- Navarrete LM (2011). Behavioral effects of wind farms on wintering Sandhill Cranes (*Grus canadensis*) on the Texas High Plains. [📄](#) (p93, en)
- ● Nehls G, Burger C, Kleinschmidt B, Quillfeldt P, Heinänen S, Morkunas J, Zydels R (2018). From effects to impacts : Analysing displacement of Red-throated Divers in relation to their wintering home ranges. [📄](#) (p3, fr)
- New L, Bjerre E, Millsap B, Otto Mc, Runge Mc (2015). A Collision Risk Model to Predict Avian Fatalities at Wind Facilities: An Example Using Golden Eagles, *Aquila chrysaetos*. [📄](#) (p12, en)
- ● Niemi J et al. (2019). Automatic Bird Identification for Offshore Wind Farms. [📄](#) (p17, en)
- ● Niemi J, Tantt J (2020). Deep learning-based automatic bird identification system for offshore wind farms. [📄](#) (p14, en)
- Nygard T et al. (2010). A study of white-tailed eagle *Haliaeetus albicilla* movements and mortality at a wind farm in Norway. [📄](#) (p4, en)
- ● Opiel S et al. (2011). Comparison of five modelling techniques to predict the spatial distribution and abundance of seabirds. [📄](#) (p11, en)
- Orloff SG, Flannery AF (1992). Wind turbine effects on avian activity, habitat use, and mortality in altamont pass and solano county Wind resource areas. [📄](#) (p199, en)
- Osborn Robert G, Dieter Charles D, Higgins Kenneth F (1997). Bird Flight Characteristics Near Wind Turbines in Minnesota. [📄](#) (p10, en)
- ● Owen E (2013). Connectivity between seabird SPAs and wind farms: an overview from the FAME project. [📄](#) (p1, en)
- Paula J, Augusto M, Neves T, Bispo R, Cardoso P, Mascarenhas M (2018). Comparing Field Methods Used to Determine Bird and Bat Fatalities. [📄](#) (p15, en)
- ● Pearce AT et al. (2016). Wintering Sandhill Crane exposure to wind energy development in the central and southern Great Plains, USA. [📄](#) (p12, en)
- ● Pearce-Higgins JW, Stephen L, Douse A, Langston RHW (2012). Greater impacts of wind farms on bird populations during construction than subsequent operation: results of a multi-site and multi-species analysis. [📄](#) (p9, en)
- ● Pekour MS (2002). Migrating birds: assessment of Impact on 915-MHz radar Wind Profiler Performance at the atmospheric radiation measurement program's Southern Great Plains. [📄](#) (p6, en)
- Pelikan M (2002). Radar ornithology adds to wind farm data gathering. [📄](#) (p3, en)

- Percival SM (2001). Assessment of the effects of offshore wind farms on birds. [📄](#) (p96, en)
- Perold V et al. (2020). On a collision course? The large diversity of birds killed by wind turbines in South Africa. [📄](#) (p13, en)
- Perrow MR et al. (2010). Advances in impact assessment of offshore wind farms upon seabirds: insight from studies of terns (*Sterna* and *Sternula* spp.). [📄](#) (p4, en)
- Pescador M, Gómez Ramírez JI, Perisc SJ (2019). Effectiveness of a mitigation measure for the lesser kestrel (*Falco naumanni*) in wind farms in Spain. [📄](#) (p7, en)
- Peschko V et al. (2020). Telemetry reveals strong effects of offshore wind farms on behaviour and habitat use of common guillemots (*Uria aalge*) during the breeding season. [📄](#) (p14, en)
- Petersen IK, TK Christensen, Kahlert J, Desholm M, Fox AD (2006). Final results of bird studies at the offshore wind farms at Nysted and Horns Rev, Denmark. [📄](#) (p166, en)
- Petterson J (2005). The impact of Offshore Wind farms on birds life in southern Kalmar Sound, Sweden - a final report based on studies 1999-2003. [📄](#) (p128, en)
- Plonczkier P, Simms IC (2012). Radar monitoring of migrating pink-footed geese: behavioural responses to offshore wind farm development. [📄](#) (p9, en)
- Ponchon A, Gallien F, Le Guillou G, Grémillet D (2015). Distribution en mer et utilisation de l'habitat des mouettes tridactyles nichant sur les côtes de la Manche. [📄](#) (p47, fr)
- Przybycin P et al. (2019). B-Finder System 24 month test report for T-Series. [📄](#) (p28, en)
- Reid T, Kruger S, Whitfield DP, Amar A (2015). Using spatial analyses of bearded vulture movements in southern Africa to inform wind turbine placement. [📄](#) (p12, en)
- Riols-Loyrette Christian - LPO Aude (2015). Impact de parcs éoliens sur un couple d'Aigle royal *Aquila chrysaetos* dans les Corbières. [📄](#) (p6, fr)
- Rodrigues S, Rosa L, Mascarenhas M (2018). An Overview on Methods to Assess Bird and Bat Collision Risk in Wind Farms. [📄](#) (p24, en)
- Rothery P, Newton I, Little B (2009). Observations of seabirds at offshore wind turbines near Blyth in northeast England. [📄](#) (p15, en)
- Rydell J, Engström H, Hedenström A, Larsen JK, Pettersson J, Green M (2012). The effect of wind power on birds and bats – A synthesis. [📄](#) (p152, en)
- Salamolard M, Moreau C (1999). Habitat selection by Little Bustard *Tetrax tetrax* in a cultivated area of France. [📄](#) (p10, en)
- Sansom A, Pearce-Higgins JW, Douglas DJT (2016). Negative impact of wind energy development on a breeding shorebird assessed with a BACI study design. [📄](#) (p15, en)
- Santos M et al. (2016). A spatial explicit agent based model approach to evaluate the performance of different monitoring options for mortality estimates in the scope of onshore windfarm impact assessments. [📄](#) (p10, en)
- Santos M et al. (2016). Impacts of land use and infrastructural changes on threatened Little Bustard *Tetrax tetrax* breeding populations: quantitative assessments using a recently developed spatially explicit dynamic modelling framework. [📄](#) (p18, en)
- Scacco M, Flack A, Duriez O, Wikelski M, Safi K (2018). Static landscape features predict uplift locations for soaring birds across Europe. [📄](#) (p13, en)
- Schaub Michael (2012). Spatial distribution of wind turbines is crucial for the survival of red kite populations. [📄](#) (p8, en)
- Schaub T, Klaassen RHG, Bouten W, Schaich AE, Koks BJ (2019). Collision risk of Montagu's Harriers *Circus pygargus* with wind turbines derived from high-resolution GPS tracking. [📄](#) (p36, en)
- Schmidt M, Aschwanden J, Liechti F, Wichmann G, Nemeth E (2017). Comparison of visual bird migration counts with radar estimates. [📄](#) (p15, fr)
- Schuster E, Bruns E (2020). 10 Fragen – 10 Antworten zu Detektionssystemen. [📄](#) (p14, de)
- Schuster E, Bruns E (2020). 10 questions - 10 réponses sur les systèmes de détection. [📄](#) (p17, fr)
- Schwitter P (2020). Determinants of stopover sites of migrating Swiss redkites (*Milvus milvus*). [📄](#) (p83, en)
- Scott B (2013). Seabirds and marine renewables: are we asking the right questions about indirect effects?. [📄](#) (p4, en)
- Scottish Natural Heritage (2016). Avoidance Rates for the onshore SNH Wind Farm Collision Risk Model. [📄](#) (p4, en)
- Sheppard et al. (2015). An autonomous GPS geofence alert system to curtail avian fatalities at wind farms. [📄](#) (p8, en)
- Sheppard et al. (2014). Curtailing Avian Impacts with Wind Turbines using GSM/GPS Tracking Telemetry that Incorporates Autonomous Geofence Alerts. [📄](#) (p45, en)
- Shewring MP, Vafidis JO (2017). The effectiveness of deterrent measures to minimize disturbance impacts to breeding European nightjar at an upland wind farm site in South Wales, UK. [📄](#) (p4, en)
- Silva JP (2010). Factors affecting the abundance of the little bustard *Tetrax tetrax*: implications for conservation. [📄](#) (p187, en)
- Silva JP et al. (2010). Estimating the influence of overhead transmission power lines and landscape

- context on the density of little bustard *Tetrax tetrax* breeding populations. [📄](#) (p10, en)
- Silva JP, Pinto M, Palmeirim JM (2004). Managing landscapes for the little bustard *Tetrax tetrax*: lessons from the study of winter habitat selection. [📄](#) (p8, en)
  - Singh K, Baker E, Lackner M (2014). Curtailing Wind Turbine Operations to Reduce Avian Mortality. [📄](#) (p6, en)
  - Skov H et al. (2015). Kriegers Flak Offshore Wind Farm - Birds and Bats EIA - Technical Report. [📄](#) (p198, en)
  - Skov H, Heinänen S, Norman T, Ward RM, Méndez-Roldán S, Ellis I (2018). ORJIP Bird Collision and Avoidance Study. Final report – April 2018. [📄](#) (p248, en)
  - Skov H, Heinänen S (2014). Predicting the Weather-Dependent Collision Risk for Birds at Wind Farms. [📄](#) (p19, en)
  - Skov H, Heinänen S, Norman T, Ward R, Méndez Roldan S (2018). Bird avoidance behaviour and collision impact monitoring at offshore wind farms. [📄](#) (p4, fr)
  - Smales et al. (2005). Modelled cumulative impacts on the orange-bellied parrot of wind farms across the species' range in South Eastern Australia. [📄](#) (p42, en)
  - Smales I (2015). Fauna Collisions with Wind Turbines: Effects and Impacts, Individuals and Populations. What Are We Trying to Assess?. [📄](#) (p18, en)
  - Smales I (2006). Impacts of avian collisions with wind power turbines: an overview of the modelling of cumulative risks posed by multiple wind farms. [📄](#) (p21, en)
  - Smales I (2005). Modelled cumulative impacts on the White-bellied Sea-eagle of wind farms across the species' Australian range. [📄](#) (p40, en)
  - Smales I (2005). Modelled cumulative impacts on the swift parrot of wind farms across the species' range in South Eastern Australia. [📄](#) (p41, en)
  - Smales I, Muir S (2005). Modelled cumulative impacts on the Tasmanian Wedge-tailed Eagle of wind farms across the species' range. [📄](#) (p50, en)
  - Smales I, Venosta M (2005). Risk level to select species listed under the EPBC Act, of collision at wind farms in Gippsland, Victoria. [📄](#) (p20, en)
  - Smallwood KS (2017). Long search intervals underestimate bird and bat fatalities caused by wind turbines. [📄](#) (p7, en)
  - Smallwood KS (2013). Comparing bird and bat fatality-rate estimates among North American wind-energy projects. [📄](#) (p15, en)
  - Smallwood KS (2009). Avian and Bat Fatality Rates at Old-Generation and Repowered Wind Turbines in California. [📄](#) (p10, en)
  - Smallwood KS (2007). Estimating Wind Turbine-Caused Bird Mortality. [📄](#) (p12, en)
  - Smallwood KS, Thelander C (2008). Bird Mortality in the Altamont Pass Wind Resource Area, California. [📄](#) (p10, en)
  - Smeraldo S et al. (2020). Modelling risks posed by wind turbines and power lines to soaring birds: the black stork (*Ciconia nigra*) in Italy as a case study. [📄](#) (p18, en)
  - Soanes LM et al. (2012). Individual consistency in the foraging behaviour of Northern Gannets: implications for interaction with offshore renewable energy developments. [📄](#) (p9, en)
  - Soares Maria (2014). An avian relative fatality risk index for Iberian species on wind farms based on zero inflated count models. [📄](#) (p60, en)
  - Sovacool BK (2009). Contextualizing avian mortality: a preliminary appraisal of bird and bat fatalities from wind, fossil-fuel and nuclear electricity. [📄](#) (p9, en)
  - Speakman J, Gray H, Furness L (2009). University of Aberdeen report on effects of offshore wind farms on the energy demands on seabirds - October 2009. (p23, en)
  - Stassart PM, Paquet J-Y (2012). Séminaire Milan royal / éoliennes - conclusions et perspectives. [📄](#) (p13, fr)
  - Stokke BG, Nygård T, Falkdalen U, Pedersen HC, May R (2020). Effect of tower base painting on willow ptarmigan collision rates with wind turbines. [📄](#) (p10, en)
  - Sur M, Belthoff JR, Bjerre ER, Millsap BA, Katzner T (2018). The utility of point count surveys to predict wildlife interactions with wind energy facilities: An example focused on golden eagles. [📄](#) (p8, en)
  - Suryan R, Albertani R, Polagye B (2016). A Synchronized Sensor Array for Remote Monitoring of Avian and Bat Interactions with Offshore Renewable Energy Facilities. [📄](#) (p33, en)
  - Sutter C, Martin K, Grandgent C (2012). Condor detection and alerting system. [📄](#) (p20, en)
  - Szymański P et al. (2017). The song of Skylarks *Alauda arvensis* indicates the deterioration of an acoustic environment resulting from wind farm start-up. [📄](#) (p21, en)
  - Tack JD, Noon BR, Bowen ZH, Strybos L, Fedy BC (2017). No Substitute for Survival: Perturbation Analyses Using a Golden Eagle Population Model Reveal Limits to Managing for Take. [📄](#) (p16, en)
  - Talisman Energy (2006). Effects of the demonstrator project on birds. [📄](#) (p24, en)
  - Telleria JL (2008). Potential impacts of wind farms on migratory birds crossing Spain. [📄](#) (p6, en)
  - Tesfahunegn W et al. (2020). Impact of wind energy development on birds and bats: the case of Adama wind farm, Central Ethiopia. [📄](#) (p9, en)



- Tetra Tech EC, Inc. (2018). Avian Monitoring and Mortality - Report at the National Wind Technology Center: January 2010 - February 2011. [📄](#) (p211, en)
- Thaxter C et al. (2013). Connectivity between seabird features of protected sites and offshore wind farms: Lesser Black-backed Gulls and Great Skuas through the breeding, migration and non-breeding seasons. [📄](#) (p1, en)
- Thaxter C, Ross-Smith V, Bouten W, Clark N, Conway G, Rehfish M, Burton N (2015). Seabird-wind farm interactions during the breeding season vary within and between years: A case study of lesser black-backed gull *Larus fuscus* in the UK. [📄](#) (p12, en)
- Thaxter CB et al. (2019). Avian vulnerability to wind farm collision through the year: Insights from lesser black-backed gulls (*Larus fuscus*) tracked from multiple breeding colonies. [📄](#) (p13, en)
- Therkildsen OR, Elmeros M (2015). First year post-construction monitoring of bats and birds at wind turbine test Centre Østerild. [📄](#) (p128, en)
- Therkildsen OR, Elmeros M, Kahlert J, Desholm M (2012). Baseline investigations of bats and birds at wind turbine test centre Østerild. [📄](#) (p128, en)
- Thonnerieux Y - Natur'ailes (2005). Eoliennes et Oiseaux: quelles conséquences?. [📄](#) (p6, fr)
- Tikkanen H, Rytönen S, Karlin O-P, Ollila T, Pakanen V-M, Tuohimaa H, Orell M (2018). Modelling golden eagle habitat selection and flight activity in their home ranges for safer wind farm planning. [📄](#) (p12, en)
- Tomé R, Canário F, Leitão A, Pires N, Repas M (2017). Radar Assisted Shutdown on Demand Ensures Zero Soaring Bird Mortality at a Wind Farm Located in a Migratory Flyway. [📄](#) (p14, en)
- Traxler A et al. (2004). Vogelschlag, Meideverhalten & Habitatnutzung an bestehenden Windkraftanlagen Prellenkirchen - Obersdorf - Steinberg/Prinzendorf. [📄](#) (p106, de)
- Trinder M et al. (2013). Potential impact of offshore wind farm collisions on the Gannet population of the British Isles explored using PVA. [📄](#) (p2, en)
- Valery L (2010). Etude de la répartition spatiale des oiseaux marins au large : Analyse synoptique de l'état de l'art pour le programme de connaissances Natura 2000 en mer. [📄](#) (p8, fr)
- van Kooten T, Soudijn F, Tulp I, Chen C, Benden D, and Leopold M (2019). The consequences of seabird habitat loss from offshore wind turbines, version 2 ; Displacement and population level effects in 5 selected species. [📄](#) (p117, en)
- Vanermen N, Courtens W, Daelemans R, Lens L, Müller W, Van de walle M, Verstraete H, Stienen E (2020). Attracted to the outside: a meso-scale response pattern of lesser black-backed gulls at an offshore wind farm revealed by GPS telemetry. [📄](#) (p10, en)
- Vanermen N, Onkelinx T, Courtens W, Van de walle M, Verstraete H, Stienen EWM (2014). Seabird avoidance and attraction at an offshore wind farm in the Belgian part of the North Sea. [📄](#) (p11, en)
- Vanermen N, Onkelinx T, Verschelde P, Courtens W, Van de walle M, Verstraete H, Stienen E (2014). Assessing Seabird Displacement at Offshore Wind Farms: Power Ranges of a Monitoring and Data Handling Protocol. [📄](#) (p13, en)
- Vasilakis DP et al. (2017). A balanced solution to the cumulative threat of industrialized wind farm development on cinereous vultures (*Aegypius monachus*) in south-eastern Europe. [📄](#) (p17, en)
- Vasilakis DP, Whitfield DP, Schindler S, Poirazidis KS, Kati V (2016). Reconciling endangered species conservation with wind farm development: Cinereous vultures (*Aegypius monachus*) in south-eastern Europe. [📄](#) (p8, en)
- Veltheim I, Cook S, Palmer G, Hill R, McCarthy (2019). Breeding home range movements of pre-fledged broilga chicks, *Antigone rubicunda* (Gruidae) in Victoria, Australia – Implications for wind farm planning and conservation. [📄](#) (p37, en)
- ● Veltri CJ, Klem D Jr (2005). Comparison of fatal bird injuries from collisions with towers and windows. [📄](#) (p8, en)
- Villegas-Patracá R, Cabrera-Cruz SA, Herrera-Alsina L (2014). Soaring Migratory Birds Avoid Wind Farm in the Isthmus of Tehuantepec, Southern Mexico. [📄](#) (p7, en)
- Wade H (2015). Investigating the Potential Effects of Marine Renewable Energy Developments on Seabirds. [📄](#) (p417, en)
- Wade HM et al. (2013). Habitat use by seabird species in high-velocity current flows: investigating the potential effects of tidal-stream renewable energy developments. [📄](#) (p5, en)
- Wade HM, Masden EA, Jackson AC, Furness RW (2016). Incorporating data uncertainty when estimating potential vulnerability of Scottish seabirds to marine renewable energy developments. [📄](#) (p6, en)
- Walker D et al. (2005). Resident Golden eagle ranging behaviour before and after construction of a windfarm in Argyll. [📄](#) (p9, en)
- ● Wang Sh, Wang Si, Smith P (2015). Ecological Impacts of Wind Farms on Birds: Questions, Hypotheses, and Research Needs. [📄](#) (p9, en)
- Warwick-Evans V, Atkinson P, Walkington I, Green J (2018). Predicting the impacts of wind farms on seabirds: An individual-based model. [📄](#) (p13, en)
- Watson RT, Kolar PS, Ferrer M, Nygard T, Johnston N, Hunt WG, Smit-Robinson HA, Farmer C, Huso MM, Katzner TE (2018). Raptor Interactions With Wind Energy: Case Studies From Around the World. [📄](#) (p19, en)

- Welcker J et al. (2016). Nocturnal migrants do not incur higher collision risk at wind turbines than diurnally active species. [📄](#) (p16, en)
- Whalen C, Brown M, McGee J, Powell L, Walsh E (2019). Wind turbine noise limits propagation of greater prairiechicken boom chorus, but does it matter?. [📄](#) (p13, en)
- Whalen CE, Brown MB, McGee J, Powell LA, Walsh EJ (2018). Male Greater Prairie-Chickens adjust their vocalizations in the presence of wind turbine noise. [📄](#) (p12, en)
- Whitfield D, Urquhart B (2015). Deriving an Avoidance Rate for Swans Suitable for Onshore Wind Farm Collision Risk Modelling. [📄](#) (p15, en)
- Whitfield DP, Madders M (2006). A review of the impacts of wind farms on hen harriers *circus cyaneus* and an estimation of collision avoidance rates. [📄](#) (p32, en)
- Whitfield DP, Madders M (2006). Deriving collision avoidance rates for Red kites *Milvus milvus*. [📄](#) (p14, en)
- Wiggelinkhuizen E et al. (2006). WT Bird -Bird collision recording for offshore wind farms. [📄](#) (p5, en)
- Wiggelinkhuizen E et al. (2004). WT Bird -Bird collision recording for offshore wind farms. [📄](#) (p5, en)
- Wiggelinkhuizen EJ, den Boon JH - ECN - E-Connection Project BV (2010). Monitoring of bird collisions in wind farm under offshore-like conditions using WT-BIRD system - Final report. [📄](#) (p57, en)
- Wiggelinkhuizen EJ, Rademakers LWMM, Barhorst SAM, Den Boon HJ (2006). Bird collision monitoring system for multi-megawatt wind turbines WT-Bird - Summary of prototype development and testing. [📄](#) (p17, en)
- Wildfowl Wetland & Trust (2009). Aerial Surveys of waterbirds in the UK: 2007/08, Final report. [📄](#) (p298, en)
- Williams K, Stenhouse I, Tash J, Adams E, Goodale W (2015). Pelagic Waterbird Surveys in the Great Lakes, 2012-2014: Minimizing Conflicts with Future Offshore Wind Energy Development. [📄](#) (p37, en)
- Winkelman JE (1995). Birds and the wind park near Urk: collision victims and disturbance of ducks, geese and swans. [📄](#) (p156, en)
- Winship A, Kinlan B, White T, Leirness J, Christensen J (2018). Modeling At-Sea Density of Marine Birds to Support Atlantic Marine Renewable Energy Planning. [📄](#) (p81, en)
- WWF Greece (2011). Assessing the impact of nine established wind farms on birds of prey in Thrace Greece - Annual report 2009-2010. [📄](#) (p43, en)
- WWF Greece (2011). Assessing the impact of nine established wind farms on birds of prey in Thrace, Greece. [📄](#) (p93, en)
- Xirouchakis SM et al. (2019). Estimating the Potential Mortality of Griffon Vultures (*Gyps fulvus*) Due to Wind Energy Development on the Island of Crete (Greece). [📄](#) (p18, en)
- Yoshihashi R et al. (2017). Bird detection and species classification with time-lapse images around a wind farm: Dataset construction and evaluation. [📄](#) (p13, en)
- Yoshihashi R, Kawakami R, Iida M, Naemura T - The University of Tokyo (2015). Construction of a bird image dataset for ecological investigations. [📄](#) (p5, en)
- Yoshihashi R, Kawakami R, Iida M, Naemura T - The University of Tokyo (2015). Evaluation of Bird Detection using Time-lapse Images around a Wind Farm. [📄](#) (p8, en)
- Young Jr DP, Erickson WP, Strickland MD, Good RE, Sernka KJ (2003). Comparison of Avian Responses to UV-Light-Reflective Paint on Wind Turbines. [📄](#) (p67, en)
- Zaugg S, Saporta G, Van Loon E, Schmaljohann H, Liechti F (2010). Automatic identification of bird targets with radar via patterns produced by wing flapping. [📄](#) (p13, en)
- Zehntindjiev P, Whitfield D (2016). Bird Migration Monitoring in the Saint Nikola Wind Farm, Kaliakra Region, in Autumn 2015, and an Analysis of Potential Impact after Six Years of Operation. [📄](#) (p30, en)
- Zimmerling JR, Pomeroy AC, D'Entremont MV, Francis CM (2013). Canadian Estimate of Bird Mortality Due to Collisions and Direct Habitat Loss Associated with Wind Turbine Developments. [📄](#) (p13, en)