Current Concepts in Return to Play for Concussions

Alicia Pike, MS, ATC
Bill Heinz, MD
Chad Asplund, MD
Rob Huggins, PhD, ATC

3rd Annual Collaborative Solutions for Safety in Sport National Meeting







Hot Topics

New Mexico High School football player



- Star Running Back
- Diagnosed concussion < 1 wk before championship game
- State law prohibits RTP no sooner than 1 wk following injury
- Case brought to court







Hot Topics

- New Mexico High School football player
- Connecticut High School football player
- Proposed Bill in North Carolina
- Athlete sustained concussion during Semi-Final game
- Diagnosed following day
- Athlete saw pediatrician who confirmed diagnosis and gave clearance for RTP protocol
- Parents ultimately went to School Board











NEWS ▼ WEATHER ▼

SPORTS V

ON YOUR SIDE ▼

WATCH LIVE ▼

THE CW

ABOUT ▼

Student-athlete safety bill would let parents approve return to game

Flanagan said while more education about student-athlete safety is imperative, the bill changes a key component of the Gfeller Waller Act, which is already in place.

"It gives the parents a determination that their child is fine and can return to play with a concussion," said Flanagan.

Currently, a student athlete suspected of having a concussion cannot return to play until the student is evaluated and cleared by a healthcare provider.

House Bill 116 would allow a student-athlete to return after receiving clearance from a parent or healthcare provider, which is raising some concerns.







Concussion Legislation


- Sports concussion law in effect in all 50 states plus DC
- Varies state to state
- Common components of law:
 - Removal from play
 - Return to play (RTP) procedures
 - Who can give RTP clearance?
 - Concussion education (coaches, athletes, parents)
 - Baseline testing
 - Return to learn (RTL) procedures

















What's 'Best Practice'?

Journal of Athletic Training 2014;49(2):245-265 doi: 10.4085/1062-6050-49.1.07 © by the National Athletic Trainers' Association, Inc www.natajournals.org

position statement

National Athletic Trainers' Association Position Statement:



Consensus Statement

Consensus Statement on Concussion in Sport—the 4th International Conference on Concussion in Sport Held in Zurich, November 2012

Paul McCrory, MBBS, PhD, Willem Meeuwisse, MD, PhD, Mark Aubry, MD, Bob Cantu, MD, Jiri Dvorak, MD, Ruben J. Echemendia, PhD, Lars Engebretsen, MD, PhD, Karen Johnston, MD, PhD, Jeffrey S. Kutcher, MD, Martin Raftery, MBBS, Allen Sills, MD, and





Zurich Guidelines

On-Field Evaluation

- Immediate Removal
- Evaluation by physician or healthcare provider (AT)
- SCAT 3

Return to Play

- No athlete should RTP on same day
- Gradual return to school and social activities
- Gradual return to play
 - Stepwise progression (1 step per day)

Rehabilitation Stage	Functional Exercise at Each Stage of Rehabilitation	Objective of Each Stage
1. No activity	Symptom limited physical and cognitive rest.	Recovery
Light aerobic exercise	Walking, swimming, or stationary cycling keeping intensity < 70% maximum permitted heart rate.	Increase HR
	No resistance training.	
3. Sport- specific exercise	Skating drills in ice hockey, running drills in soccer. No head impact activities.	Add movement
Noncontact training drills	Progression to more complex training drills, eg, passing drills in football and ice hockey.	Exercise, coordination, and cognitive load
	May start progressive resistance training.	
5. Full contact practice	Following medical clearance participate in normal training activities.	Restore confidence and assess functional skills by coaching staff
Return to play	Normal game play.	







Two Models

Athletic Trainer Employed

- Immediate Recognition/Removal
- AT communication/education
- Referral to 'team physician' or appropriate healthcare provider
- AT works with school personnel and physician on RTL modifications
- Physician clearance for RTP
- AT supervises athlete through RTP

No Athletic Trainer Employed

- Coach recognition of injury
- Refer (emergency room or primary care physician)
- RTL modifications not guaranteed
- Physician clearance
- RTP progression not guaranteed







Strategies

- Create a network of healthcare providers
- Identify 2-3 physicians with concussion education/knowledge for referral purposes
- Ensure collaborative relationships between key players (Medical team, athletic team, academic team)
- * Multidisciplinary Team Approach to Medical Care











Guidance for the Clinician in Rendering Pediatric Care

CLINICAL REPORT

Returning to Learning Following a Concussion

abstract



Following a concussion, it is common for children and adolescents to experience difficulties in the school setting. Cognitive difficulties, such

Mark E. Halstead, MD, FAAP, Karen McAvoy, PsyD, Cynthia D. Devore, MD, FAAP, Rebecca Carl, MD, FAAP, Michael Lee, MD, FAAP, Kelsey Logan, MD, FAAP, Council on Sports Medicine and Fitness, and Council on School Health

Guidelines for Post-Concussion Academic Accommodations









Stage	Goals/Key Ideas	Teacher's Actions	Student's Actions
I	Complete rest	 Contacted by school nurse Explanation of injury & current plan of care 	 Out of school Strict limits for use of computer, cell phone, texting, video games No Physical/Sports Activity
II	Significant deficits in processing and concentration. Cognitive activity as tolerated	 Develop lists of 3 categories for all assignments: 1. Excused: Not made up. 2. Accountable: Responsible for content, not process 3. Responsible: Must be completed and will be graded 	 In school as tolerated Observing, not participating Communicate with teachers about progress Be patient with slow recovery No Physical/Sports Activity
III	Gradual increase of time and energy, slowly resuming full workload	 Prioritize assignments with student Continue to use list above Set timeline for completion 	 Communicate with teachers on assignment progress No Physical/Sports Activity
IV	Complete resumption of normal activities	 Monitor completion of assignments Communicate w/ parents/staff when student is caught up Communicate w/ Guidance as grades are updated 	 Resume all normal activities Progress with athletic trainer – supervision resumption of participation in athletics.

Resources

- NATA Position Statement on Concussion Management
- Concussion Statement on Concussion in Sport (Zurich)
- American Academy of Pediatrics Returning to Learning Following a Concussion
- NFHS Materials
- CDC Heads Up







Questions to Ponder

- What barriers is your state currently experiencing regarding Concussion RTP and/or RTL?
- What are some of you state's successes in this area?
 How did you get there?
- What does your state need in order to implement best practices for concussion RTP procedures (including RTL)?
- Who should be allowed to make RTP decisions (Chiropractor, PT, Physician, AT, etc.)?







Thank You!

- Alicia Pike <u>alicia.pike@uconn.edu</u>
- Bill Heinz wheinz56@gmail.com
- Chad Asplund chad.asplund@gmail.com
- Rob Huggins <u>robert.huggins@uconn.edu</u>





