

# Digital tools to support the maintenance of physical activity in people with long-term conditions: A scoping review

Digital Health  
Volume 8: 1–17  
© The Author(s) 2022  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/20552076221089778  
journals.sagepub.com/home/dhj



Paul Clarkson<sup>1,2,3</sup> , Aoife Stephenson<sup>4</sup>, Chloe Grimmett<sup>1,5</sup>, Katherine Cook<sup>6</sup>, Carol Clark<sup>7</sup>, Paul E Muckelt<sup>1,3</sup>, Philip O’Gorman<sup>4</sup>, Zoe Saynor<sup>8</sup>, Jo Adams<sup>1,2,3</sup>, Maria Stokes<sup>1,2,3,5</sup> and Suzanne McDonough<sup>1,4</sup> 

## Abstract

**Objective:** This scoping review aimed to bring together and identify digital tools that support people with one or more long-term conditions to maintain physical activity and describe their components and theoretical underpinnings.

**Methods:** Searches were conducted in Cumulative Index to Nursing and Allied Health Literature, Medline, EMBASE, IEEE Xplore, PsycINFO, Scopus, Google Scholar and clinical trial databases, for studies published between 2009 and 2019, across a range of long-term conditions. Screening and data extraction was undertaken by two independent reviewers and the Preferred Reporting Items for Scoping Reviews guidelines informed the review’s conduct and reporting.

**Results:** A total of 38 results were identified from 34 studies, with the majority randomised controlled trials or protocols, with cardiovascular disease, type 2 diabetes mellitus and obesity the most common long-term conditions. Comorbidities were reported in >50% of studies but did not clearly inform intervention development. Most digital tools were web-browser-based ± wearables/trackers, telerehabilitation tools or gaming devices/components. Mobile device applications and combination short message service/activity trackers/wearables were also identified. Most interventions were supported by a facilitator, often for goal setting/feedback and/or monitoring. Physical activity maintenance outcomes were mostly reported at 9 months or 3 months post-intervention, while theoretical underpinnings were commonly social cognitive theory, the transtheoretical model and the theory of planned behaviour.

**Conclusions:** This review mapped the literature on a wide range of digital tools and long-term conditions. It identified the increasing use of digital tools, in combination with human support, to help people with long-term conditions, to maintain physical activity, commonly for under a year post-intervention. Clear gaps were the lack of digital tools for multimorbid long-term conditions, longer-term follow-ups, understanding participant’s experiences and informs future questions around effectiveness.

## Keywords

Physical activity maintenance, behaviour change, Internet, chronic, digital health, multimorbidity

Submission date: 29 June 2021; Acceptance date: 9 March 2022

<sup>1</sup>School of Health Sciences, University of Southampton, Southampton, UK

<sup>2</sup>National Institute for Health Research Applied Research Collaboration Wessex, Southampton, UK

<sup>3</sup>Centre for Sport, Exercise and Osteoarthritis Research Versus Arthritis, Southampton, UK

<sup>4</sup>School of Physiotherapy, RCSI University of Medicine and Health Sciences, Dublin, Ireland

<sup>5</sup>National Institute for Health Research, Southampton Biomedical Research Unit, Southampton, UK

<sup>6</sup>Faculty of Health and Wellbeing, School of Health and Care Professions, University of Winchester, Winchester, UK

<sup>7</sup>Department of Rehabilitation and Sport Sciences, Faculty of Health and Social Sciences, Bournemouth University, Bournemouth, UK

<sup>8</sup>Physical Activity, Health and Rehabilitation Thematic Research Group, Faculty of Science and Health, School of Sport, Health and Exercise Science, University of Portsmouth, Portsmouth, UK

### Corresponding author:

Paul Clarkson, School of Health Sciences, University of Southampton, University Road, Southampton SO17 1BJ, UK.  
Email: P.D.Clarkson@soton.ac.uk



## Introduction

Physical activity (PA) is an important part of maintaining both physical and mental health for people with one or more long-term conditions (LTCs).<sup>1,2</sup> A LTC is a broad term for a range of physical and mental health conditions ‘that cannot at present be cured but can be controlled with medication or therapies’<sup>3</sup> and is considered to last for more than 1 year.<sup>4</sup> The World Health Organization (WHO) reports that  $\geq 1.4$  billion people worldwide are not active enough and an overall lack of progress at improving PA and reducing sedentary levels over the last 20 years.<sup>4</sup> PA data from 2019/2020 for England highlights that 66.4% of adults were physically active to some extent each week.<sup>5</sup> However, when compared with data from Sports England over the same period, 72.5%–75% of people with a disability or LTCs were inactive, defined as no activity in the last 28 days at two data points (May/November).<sup>6</sup> Similar disparities in activity level between the general population and those with LTCs have also previously been reported in the research literature.<sup>7,8</sup> Previous systematic reviews and guidelines have reported the benefits of PA for people with LTCs: to reduce some symptoms, prevent complications and maintain function.<sup>7,9–11</sup>

Digital tools as defined by WHO classifications, which includes digital and mobile technologies, such as websites, mobile device applications, telehealth and wearable devices<sup>12</sup> and this is how the term will be used in this review. Digital tools offer great potential to support increasing PA and a wide range of previous systematic reviews have reported effectiveness at increasing PA levels in the short term for people with LTCs.<sup>13–15</sup> The use of digital tools for this purpose also fits with a wider long-term agenda for digital tools to support existing services in the National Health Service (NHS) and more widely and has been found to be cost effective for some services.<sup>16–18</sup> Digital tools may also be preferable to engaging with traditional services for some, given the flexibility of accessing support at a time that suits them, reducing transport-related issues<sup>19</sup> and, since the COVID-19 pandemic, infection risk.<sup>20</sup>

Preliminary searches of the literature have identified few existing systematic reviews that focus on supporting people with LTCs to maintain PA using digital tools. Of those that do exist, their scope in terms of LTCs and multimorbidity, range of digital tools and maintenance outcomes is limited. For example, five systematic reviews, mostly with single condition cohorts (e.g. cancer survivors, obesity, chronic obstructive pulmonary disease (COPD), inflammatory arthritis and a mix of chronic conditions), found few studies reporting on the use of digital tools to support maintenance outcomes with either no or limited statistical evidence of effects.<sup>21–25</sup> One reason for these findings may be that interventions that are designed to initiate change in behaviour, such as increasing PA, do not meet people’s needs when attempting to maintain PA in the community for the long term.<sup>26,27</sup>

Maintenance of PA has been conceptualised by time and intensity of PA in different studies (regular activity or statistically significant change in behaviour over 1–12 months),<sup>21,28,29</sup> behavioural automaticity or when the behaviour becomes the ‘dominant response’ in context.<sup>30,31</sup> Time-based definitions for maintenance of PA have more recently focused on 3–6 months after the end of the intervention.<sup>21,28,29</sup> Given the limited number of studies reporting maintenance of PA and the heterogeneity between studies in previous reviews,<sup>21,24</sup> we concluded that a novel scoping review would be appropriate to explore the range and depth of available literature in this area,<sup>32,33</sup> to direct future systematic reviews and/or primary research questions.

The use of theory in the development of behaviour change interventions, as part of a wider programme theory approach to intervention development<sup>34</sup> is associated with increased effectiveness.<sup>35</sup> Consequently, it is important to understand whether the theory has been used to develop digital tools, and if so, which theories are associated with the maintenance of PA. Identifying the theoretical basis and use of behaviour change techniques (BCTs) as intervention components will help support the replication of effective strategies and provide evidence to inform future intervention development.<sup>36</sup> Key theories that have previously been associated with the maintenance of health behaviours are theories of self-regulation,<sup>28,37</sup> and self-determination theory.<sup>38</sup>

Furthermore, the increasing focus on digital health in healthcare systems, both before the COVID-19 pandemic and especially since,<sup>16,39</sup> has meant that clinicians and commissioners need to understand what evidence-based digital tools are available for implementation. This scoping review will systematically map the research undertaken and planned in this area to identify tools that may be suitable for replication and to identify any existing gaps in knowledge.

This review aimed to answer the following objectives:

1. What is the ‘extent (size), range (variety) and nature (characteristics) of the evidence’<sup>40</sup> on digital tools to support the maintenance of PA for people with one or more LTCs?
2. What theoretical underpinnings are used in digital tools to promote the maintenance of PA?

## Methods

This review was conducted in accordance with guidance from the Preferred Reporting Items for Scoping Reviews (PRISMA-ScR),<sup>40</sup> the Joanna Briggs Institute,<sup>41</sup> and existing scoping review frameworks.<sup>33,42</sup> The protocol for this review is available from Protocols.io.<sup>43</sup>

## Eligibility criteria

The eligibility criteria for LTCs, PA and digital tools are shown in Table 1. The list of included LTCs was based

**Table 1.** Eligibility criteria for study inclusion.

<b>2009–2019 for full-text studies</b>	
<b>Dates</b>	<b>2017–2019 for abstracts (to avoid duplication with full texts)</b>
<b>Long-term conditions included</b>	Asthma Cardiovascular disease including atrial fibrillation, hypertension, heart failure, peripheral arterial disease, secondary prevention <sup>a</sup> of coronary heart disease Chronic kidney disease Chronic obstructive pulmonary disease Dementia Depression Type 1 or 2 diabetes mellitus Epilepsy Mental health Myocardial infarction: secondary prevention <sup>a</sup> Obesity Osteoarthritis Osteoporosis Rheumatoid arthritis Stroke/transient ischaemic attack
<b>Long-term conditions excluded</b>	Cancer, low back pain
<b>Physical activity inclusion</b>	Adults not meeting $\geq 150$ min MVPA per week
<b>Outcome timing</b>	Must have measured a physical activity outcome at least 3 months post the end of the intervention
<b>Physical activity exclusion</b>	Studies that report a reduction in sedentary time only
<b>Digital tools included<sup>12</sup></b>	Targeted client communication such as email or other messaging interventions. Web-based intervention Untargeted client communication, such as web-based or software-based interventions, including video Client -to-client communication, such as digital peer support group Personal health tracking, such as smartwatches or other activity trackers with a visual display Telemedicine systems with visual display for user On-demand information services to clients such as digital sources of information Client financial transactions such as digital incentive management Other tools that included exergaming, gamification
<b>Digital tool excluded</b>	Pedometers/accelerometers used alone without connection to another digital tool

<sup>a</sup>Preventing progression of an established condition.

MVPA: Moderate-to-vigorous physical activity.

on The Quality and Outcomes Framework (2017/2018)<sup>44</sup> and the National Institute for Health and Care Excellence PA pathways.<sup>45</sup> Broader terms (e.g. 'chronic', 'long-term condition', 'multimorbidity') in searches and studies were included if one or more of the LTCs were reported (Table 1). A small-scale pilot identified that some studies included defined LTCs as a subset of a larger sample. An a priori decision was taken to include these studies where all the other eligibility criteria were met, and results were charted for the relevant LTCs if possible. Cancer and low back pain were excluded due to existing recent reviews.<sup>21,46,47</sup>

Studies with adults ( $\geq 18$  years) who were not currently achieving the recommended levels of PA, based on the United Kingdom (UK) PA guidelines ( $\geq 150$  min of moderate-to-vigorous activity (MVPA) per week)<sup>48,49</sup> were included. Maintenance was defined as at least 3 months after the end of the intervention. While attempts were made to include studies with no contact during the maintenance period, it was recognised that this would have been too restrictive. Instead, studies were included when there was either no contact with the intervention or where a lesser version of the intervention was employed during the maintenance period. This information was charted in accordance with guidance.<sup>33</sup>

Digital tools were defined using the classification of digital health interventions from the WHO<sup>12</sup> (Table 1). All study designs were eligible for inclusion including quantitative, qualitative and mixed methods studies, protocols and conference abstracts.

### Information sources

Preliminary searches were conducted in Cumulative Index to Nursing and Allied Health Literature (CINAHL) and Medline to establish appropriate search terms. The search strategy (Supplemental material 1 Appendix A) was developed alongside an academic librarian, members of the research team and based on previously published search terms.<sup>21,28,50</sup> The search strategy was made up of keywords (e.g. digital, physical activity, maintenance and the list of LTCs (Table 1)) as well as synonyms of these terms, which were connected using Boolean operators. This search strategy was initially set up to support a search of the Medline database, before being adapted to accommodate the syntax of other databases. Comprehensive searches were undertaken in CINAHL, Medline, OVID EMBASE, IEEE Xplore, PsycINFO, Scopus and Google Scholar (to capture grey literature). Clinical trial registries (e.g. International Prospective Register of Systematic Reviews, International Standard Randomised Controlled Trial Number database, International Clinical Trials Registry Platform, European Union clinical trials register, and Clinicaltrials.gov) were also searched to ensure that ongoing and recently completed studies were not missed.


Databases were searched from 2009 to 2019, to follow on from a previous review that searched up to 2009, finding only one digital tool.<sup>29</sup> Searches were conducted between 17 and 28 January 2020.

### Study selection

Search results were transferred into endnote (Clarivate Analytics, Boston, MA). Five percent of titles were initially screened independently by two reviewers (PC, SMcD) and then discussed to determine agreement, before the remaining titles were screened by PC. A random 5% sample of titles and abstracts were screened initially by PC and SMcD, with clarifications made to the eligibility criteria. Results were transferred into the Covidence software (Veritas Health Innovation, Melbourne, Australia) for title/abstract screening by the team. Each reference was screened by four groups of two independent reviewers (PC, CC, SMcD, PM, CG, AS, JA, ZS), with conflicts highlighted through the software and decided by a verifier. The research team (PC, PM, KC, CC, CG, AS, POG, ZS) undertook the same process for full-text review but were required to select a reason for exclusion from the predefined eligibility criteria listed in Covidence. The final process of screening was undertaken by four members of the team working in two pairs (PC, SMcD; POG, AS) to determine whether the interventions were predominantly digital, based on criteria established through consensus for a related systematic review involving AS, SMcD (online Supplemental material Appendix B). Results were screened and discussed to confirm eligibility. Literature reviews that were identified as relevant to the eligibility criteria in the search results had their included studies checked against the list of included and excluded studies from the scoping review. Studies that were found to be eligible for inclusion and had not already been identified through database searches were screened and the literature reviews were excluded (Figure 1).

### Data charting process and data items

Data from the included studies were charted into an Excel sheet developed a priori based on guidelines<sup>38</sup> and previous studies.<sup>21,51</sup> This included study characteristics such as design, location, setting, primary LTC (Table 1) and comorbidities of any kind, in addition to number, age and gender of participants. Intervention description and length (defined in Table 1), inclusion/exclusion, maintenance period ( $\geq 3$  months after the end of the intervention) and any reported access to elements of the intervention during this period were recorded. Type and tool used to measure PA (objective or self-reported), and theoretical underpinning (behaviour change theory or BCTs explicitly mentioned, as the BCT taxonomy<sup>36</sup> was not used for



**Figure 1.** PRISMA flow diagram<sup>35</sup> for review phases, including results identified, excluded and reasons for exclusion.

extraction purposes due to resource limitations) were reported (full list in Supplemental material 1 Appendix C).

The charting form was piloted using one of the included papers before data extraction began, to clarify understanding of the categories. Eight members of the team were divided

into pairs to undertake data extraction (PC/KC, SMcD/ZS, CC/PM, POG/AS) with the included studies divided between them. Each reviewer independently read and extracted data into the charting form, before meeting with the other reviewer to discuss and agree on the final extraction.



Where appropriate, reviewers contacted study authors to clarify additional detail. In accordance with scoping review guidelines, critical appraisal was not undertaken.<sup>32,40</sup>

## Synthesis

The charting forms were collated into one excel sheet by PC, before collation and summarising of the data based on the objectives by PC, POG, AS, SMcD. The charted data were reviewed, summarised and clarified with the original sources. These summaries were discussed to identify the most appropriate way of presenting the results, before being sent to the wider team for review and presented at a team meeting. Data were presented descriptively using frequencies and measures of central tendency. Characteristics of the interventions included description, hardware used, intervention components, including non-digital components, type of digital tool,<sup>12</sup> (Table 1) and length of intervention. The longest length of maintenance period and any access during to the intervention during this period were synthesised. Reports of theoretical underpinnings of the interventions were collated. Theories were only extracted if they were listed as one of the 83 theories of behaviour change from the ABC of behaviour change theories,<sup>52</sup> developed by an expert group to be relevant to the design of interventions. BCTs were collated from studies that reported use of the BCT taxonomy<sup>36</sup> or its precursor by Abraham and Michie.<sup>53</sup>

## Results

Database searches identified 8206 results (Figure 1). Title review resulted in the exclusion of 6351 results. The team reviewed 1855 titles and abstracts, which resulted in 514 potentially relevant studies for full-text review. Reasons for exclusion at this stage included a lack of maintenance period, measurement of sedentary time only and studies where a pedometer was the only digital tool. During the full-text review, 457 citations were excluded, predominantly for not meeting the maintenance definition ( $n = 164$ ), not including the defined LTCs ( $n = 101$ ), or not including a digital tool ( $n = 97$ ). PA outcomes were not included in 34 results, while 41 citations were abstracts from before 2017 and therefore excluded. The team identified six potentially relevant citations from five reviews. In total, 20 reviews were excluded at this stage. After screening the six citations from the reviews, one was moved onto the digital review stage. Fifteen further citations were excluded during the digital review stage. A further 5 citations were excluded during the data extraction stage, leaving 38 results, from 34 studies to be included in the review (Supplemental material 1).

### Study characteristics

Of the 38 included papers, 19 were either randomised controlled trials (RCTs)<sup>54–71</sup> or used a quasi-experimental

design,<sup>72</sup> 14 were protocols for RCTs,<sup>73–86</sup> three were pilot or feasibility studies,<sup>87–89</sup> one study used a correlational design<sup>90</sup> with a single group, and one was a mixed-methods process evaluation<sup>91</sup> linked to one of the protocols<sup>73</sup> and RCTs.<sup>62</sup> Studies were mostly undertaken in Europe, with The Netherlands hosting the most studies (8/34), although the largest recruited sample sizes were reported in studies from North America and Australia.<sup>56,57,61,70</sup> Sample sizes at baseline ranged from  $n = 20$ –2000 overall (including anticipated samples from protocols). More than 60% of studies were undertaken since 2016 indicating increasing interest in this area. This is further exemplified by the identified protocols, which target the recruitment of a greater number of people with LTCs for future trials (Supplemental material 1).

Cardiovascular disease was the most common LTC across the sample of studies (10/35), followed by type-2 diabetes mellitus (T2DM) (7/35) (Table 2). A greater number of females were recruited overall, although three of the RCTs reported that females made up a much smaller proportion of the overall sample (10% to 20%). Mean age of participants ranged from 33.9 to 66.3 years in the intervention groups. The most common setting for referral or recruitment to use the digital tool was secondary care (10/33), followed by primary care (8/33) (Figure 2). Seven studies were defined as being undertaken in the community setting, which included community groups, referral from community-based clinicians, as well as adverts, postal invitations and word of mouth. Other settings for the included studies are shown in Figure 2. Most interventions (29/33, 88%) were designed to be used at home, while some were designed for use in a work setting ( $n = 2$ ) or within a community/local authority programme or group ( $n = 2$ ).

Types of comorbidity were reported in 16 studies, while a further three studies reported comorbidities in their samples, but did not define the number or condition/s. Figure 3 shows the studies and primary LTC, with linked comorbid condition/s. The most commonly reported comorbidities were obesity and T2DM. There was little indication in the majority of studies that the interventions were amended in any way to support these comorbidities. Two studies<sup>82,85</sup> may have adapted the intervention to account for the comorbidities reported, although this was not stated clearly.

Grey boxes show the first author of the included study and primary LTC in brackets. Blue boxes show comorbidities. Bossen et al.<sup>58</sup> is not shown as did not report specific comorbidities.

### Characteristics of the digital tools

Full details of the interventions are shown in Supplemental material 1. Digital tools were predominantly web-browser based (13/34) or used the web alongside a wearable/activity tracker or pedometer (5/34). Telerehabilitation

**Table 2.** Study characteristics.

<b>Study location (n = 34)</b>	Europe	17 (50.0%)
	<b>North America</b>	<b>8 (23.5%)</b>
	<b>Asia</b>	<b>4 (11.8%)</b>
	<b>Australia</b>	<b>4 (11.8%)</b>
	<b>Mixed continent (Europe/Asia)</b>	<b>1 (2.9%)</b>
<b>Publication date (n = 38)</b>	<b>2009–2012</b>	<b>4 (10.5%)</b>
	<b>2013–2015</b>	<b>11 (29.0%)</b>
	<b>2016–2019</b>	<b>23 (60.5%)</b>
<b>Primary LTC (n = 35)</b>	<b>Cardiovascular disease (CVD) (including hypertension, heart failure, ischaemic heart disease, angina, coronary artery disease)</b>	<b>10 (28.6%)</b>
	<b>Type 2 diabetes mellitus (T2DM)</b>	<b>7 (20.0%)</b>
	<b>Obesity</b>	<b>6 (17.1%)</b>
	<b>Chronic obstructive pulmonary disease</b>	<b>2 (5.7%)</b>
	<b>Stroke</b>	<b>2 (5.7%)</b>
	<b>Osteoarthritis</b>	<b>2 (5.7%)</b>
	<b>Depression</b>	<b>1 (2.9%)</b>
	<b>Rheumatoid arthritis</b>	<b>1 (2.9%)</b>
	<b>No single LTC reported</b>	<b>1 (2.9%)</b>
	<b>Mixed (n = 1 COPD, rheumatism, osteoporosis, chronic heart disease, musculoskeletal; n = 2 T2DM, COPD)</b>	<b>3 (8.5%)</b>

Primary LTC included n = 35 papers due to the protocol of one RCT<sup>73</sup> reporting one LTC (COPD) and the subsequent RCT and process evaluation<sup>62,91</sup> reporting a mix of conditions (type-2 diabetes mellitus and COPD).

COPD: chronic obstructive pulmonary disease; LTC: long-term condition; RCT: randomised controlled trial.

interventions were used in a further five studies, while a gaming device or an intervention that used gaming elements was used in four studies. Mobile device applications (apps) were used in three studies. Short message service (SMS) interventions with and without an activity tracker were used in two studies, and wearable devices with a connection to a website or app were included in two studies. There was a wide range of intervention lengths from 2 weeks to 12 months.

All interventions were delivered digitally, although most (22/34) included a healthcare professional (HCP) or other facilitators as an active part of the wider intervention (Table 3). A further three studies<sup>57,59,65</sup> included an HCP or facilitator to introduce the digital tool to participants

and/or set goals and provide feedback. One study<sup>62</sup> was app-based linked to a website that allowed HCPs to set goals and monitor progress. Eight interventions did not include any active contact with an HCP or facilitator.<sup>58,61,63,72,79,85,86,88</sup> The most common intervention components, reported by the author's intervention descriptions, were the use of motivational messages delivered either digitally, over the telephone or in person (21/34) and goal setting (18/34).

### *Maintenance period and measurement of PA*

Maintenance periods ranged from three to 12 months post-intervention with 9 months (11/34) and 3 months (9/34) the




Figure 2. Setting for referral of participants to digital tool and location of use. (Figure 2 shows 33 studies rather than 34 due to the setting being unclear in Barnason et al.<sup>90</sup>)

most commonly reported. Figure 4 shows the point at which the longest maintenance outcome was recorded for each study, in relation to the length of the intervention. Most

studies reported no access to the intervention during the maintenance period (18/34) or access to a lesser version of the intervention (10/34). Six studies were unclear. PA




Figure 3. List of studies with primary long-term conditions (LTCs) and linked comorbidities.



was most often objectively measured (19/34) alone or alongside a participant-reported outcome measure (PROM) (8/34). A further seven used a PROM alone. The most commonly used devices for measuring objective PA were the Actigraph accelerometer (10/27), SenseWear Armband (4/27), FitBit step counter (3/27) and GENEActiv accelerometer (2/27). The most commonly used PROMs were the International Physical Activity Questionnaire (IPAQ) (5/15) and the Short Questionnaire to Assess physical activity (SQUASH) (3/15). (Other devices or PROMs used are shown in Supplemental material 1).

### Theoretical underpinnings of interventions

Interventions were predominantly delivered using digital tools, however, some also included non-digital components (Supplemental material 1, Table 3). Theoretical underpinnings are presented for the whole study intervention as it was not possible to isolate digital/non-digital components. Fifteen interventions reported in 18 papers clearly articulated the use of behaviour change theory in the development of the intervention. Two of these interventions also had other theories associated with them, but it was unclear whether these were used in the intervention development process. A further seven studies reported the use of theory, but it was unclear whether this was specifically related to the development of the intervention. In the remaining 12 studies, there was either no theoretical underpinning reported or limited evidence to suggest the use of theory. The most commonly cited theories were social cognitive theory (SCT) ( $n = 5$ , + 1 unclear use as an underpinning), the transtheoretical model (TTM) and the theory of planned (TPB) behaviour (both  $n = 3$ , + 1 unclear use as an underpinning). BCTs<sup>36</sup> were mentioned in four studies, but it was unclear whether they were specifically used in the development of the intervention.

### Discussion

To our knowledge, this is the first scoping review to map the range and breadth of digital tools to support people with a wide range of the most prevalent LTCs to maintain PA. Over the last 20 years, our review shows that web-based digital tools continue to predominate with more recent emergence of gamification, apps and virtual environments. Interventions continue to be aimed at supporting people with a single LTC, even though a large proportion of participants also had comorbidities. Most participants were from younger age groups. The use and description of theory in the development of the tools were limited, with a lack of transparent reporting, and most studies highlighted the need for human engagement to support their use.

A novel finding of our review compared to previous reviews is the wealth of evidence we identified. There is a

significant body of evidence ( $n = 34$  studies), demonstrating the benefit of conducting a scoping review across multiple LTCs. Previous reviews with a focus on single LTCs have reported minimal use of digital tools to support the maintenance of PA,<sup>21–23</sup> while others that have focused on digital technologies for LTCs report minimal or no use of outcomes in the maintenance period,<sup>13,24,25</sup> including for one of the excluded conditions, low back pain.<sup>47</sup> Our results demonstrate an increasing interest in the use of digital tools to support people with LTCs to maintain PA over the review period and particularly since 2016. This may reflect the increased interest and guidelines advocating digital health strategies within Europe over the same period.<sup>16,92,93</sup> Most of the identified digital tools used the Internet in some form, either as the primary delivery modality, e.g. web browser-based interventions; or in an accessory capacity, such as providing visual PA metrics through an activity monitor or app. The present review identified only three studies that developed apps, which is surprising given the exponential increase in the number of available apps from commercial app stores, although many are not designed specifically for people with LTCs.<sup>94</sup> However, some apps are reported to have a limited evidence base<sup>95</sup> and it is therefore likely that our review would not have captured them, as development work is unlikely to have been published in academic journals.

The use of theory was effectively described in fewer than half of the identified studies, and identifying it proved to be a difficult task, due to inconsistencies in reporting, and we were unable to separate the digital and non-digital theoretical components. SCT, TTM and TPB were the most commonly reported theories, which is similar to other reviews of PA maintenance interventions.<sup>21,22,25</sup> Guidance on intervention development suggests that a theoretical underpinning is best practice and is associated with greater effectiveness,<sup>37,96,97</sup> while other studies show equivocal outcomes across the age and condition spectrum.<sup>98,99</sup> Michie and colleagues developed a BCT taxonomy to support fidelity in the delivery of an intervention and to identify the effective components for behaviour change to improve future intervention development.<sup>36</sup> BCTs were only reported in four studies, although their specific use as an ‘active ingredient’<sup>36</sup> was less well described. Inconsistent description of intervention components has previously been reported,<sup>100</sup> including in eHealth interventions for people with cardiovascular disease (CVD),<sup>101</sup> and was not described in a review of web-based interventions for low back pain,<sup>47</sup> reducing the potential for replication and translation of findings.<sup>102,103</sup> Given these identified limitations, we intend to explore the effective components of interventions in a future systematic review using intervention component analysis.<sup>104</sup>

While digital tools made up the primary component of interventions, additional human support (via HCP or other

**Table 3.** Healthcare professional/facilitator involvement in the study interventions.

	Active part of intervention <sup>a</sup>	Monitoring	Referral to tool or set up goals/feedback	No active intervention contact <sup>b</sup>
Vorrink, 2016		•	•	
Thorup, 2016			•	
Lorig, 2010	•			
Jones, 2016	•			
Hurkmans, 2010	•			
Lari, 2018				•
Jaarsma, 2014	•			
Jennings, 2014				•
Hawkins, 2019	•			
Dor-Haim, 2019	•			
Devi, 2014	•			
Barnason, 2016	•			
Harrison/Patel, 2019	•			
Bouwers, 2017	•			
Bossen, 2013				•
Bonn, 2018	•			
Barry, 2011	•			
Fife-Schaw, 2014	•			
Avila, 2019	•			
Olson, 2015				•
Alonso-Dominguez, 2017/2019	•			
Verwey, 2014/2016 Van der Weegen, 2015	•			
Kloek, 2014	•			
Ingram, 2018	•			
Strom, 2013			•	
Vorderstrasse, 2017				•
Reid, 2012			•	

(continued)

Table 3. Continued.

	Active part of intervention <sup>a</sup>	Monitoring	Referral to tool or set up goals/feedback	No active intervention contact <sup>b</sup>
<b>Volders, 2019</b>				•
<b>Lubans, 2009</b>	•			
<b>Givon, 2016</b>	•			
<b>Yang, 2017</b>	•			
<b>Sharma, 2019</b>				•
<b>Cox, 2018</b>	•			
<b>Banos, 2015</b>				•

<sup>a</sup>Active direct involvement during intervention period.

<sup>b</sup>May include automated reminder messages delivered digitally.




Figure 4. Length of interventions and related longest physical activity (PA) maintenance outcomes.

facilitators) was identified in most studies which have implications for staff resources needed to scale up potential solutions. Key aspects of digital interventions in our review, i.e. motivational messages and goal setting, often supported by HCPs or other facilitators, have been reported in previous reviews to support PA for people with and without LTCs.<sup>21,24,95</sup> There is debate as to whether human support is needed. Some highlight the importance of

human support to promote adoption and follow-up of Web 2.0 tools, defined as ‘participatory internet interventions’.<sup>2, 25</sup> Others, including a review of web-based interventions for low back pain have reported mixed results in terms of additional support.<sup>47</sup> Clearly, there are pros and cons to the involvement of HCPs alongside a digital tool: it can help to reduce anxiety and increase feelings of support<sup>105</sup> and build self-efficacy,<sup>106</sup> or it may lead to a

reliance on HCP input to self-manage LTCs.<sup>107</sup> While digital tools and services are often considered to be part of the solution for reducing pressure on health services,<sup>107</sup> these findings suggest that human input may still be required to support their use, further work is needed to better understand how to optimise digital tools through HCP support. Challenges remain, both in terms of ensuring the availability and digital capability of staff to meet the need of people when scaling up interventions.<sup>108</sup>

This scoping review identified digital tools which were designed for people with the most prevalent single LTCs. Previous systematic reviews in this area have also focused on the most prevalent conditions, including obesity, COPD, inflammatory arthritis and cancer survivors.<sup>15–18</sup> Comorbidities were reported in more than half of the identified studies in the present review, but there was limited evidence that the digital tools had been developed to take account of the impact of these conditions. Comorbidities were rarely reported in studies included in previous reviews, but when they were reported, approximately half of the digital tools were designed to support these comorbidities.<sup>21–24</sup> A future systematic review could explore whether this influenced the effectiveness of digital interventions.

In the present review, the upper age range of included participants was not representative of the largest proportion of people with LTCs, who are aged 65–99, based on the UK Office for National Statistics (ONS) data<sup>109</sup> and Irish Health Survey data from 2019.<sup>110</sup> This emphasises a key limitation of the current evidence and is particularly disappointing, given the greater age-associated prevalence of LTCs such as CVD and T2DM, and the greater mortality rates related to CVD and T2DM as multimorbidities.<sup>111,112</sup> Furthermore, given projections of ageing populations in Europe over the next 30 years,<sup>113</sup> it is increasingly important to include older adults in technology-related research to ensure that the needs of these groups are met. However, as is evident from the present findings, recruiting older adults is often difficult.<sup>114</sup> Reasons for this are reported to include a lack of interest, transportation issues (when required) and advice from family or clinicians against participation.<sup>114</sup> Others have highlighted that while older adults are open to the use of technologies, barriers to involvement include a lack of clear information and user support.<sup>115</sup>

This scoping review only included studies that reported a PA maintenance outcome, with the majority reporting outcomes at either 9 months or 3 months after completion of the intervention. The longest maintenance outcome was reported at 12 months, which aligns with some of the previous reviews in this area, both for digital and non-digital interventions.<sup>23–25,29</sup> However, others have reported maintenance outcomes of between 3 and 5 years,<sup>21,22</sup> although predominantly for non-digital interventions. Similar findings have been reported recently for non-digital interventions aiming at supporting longer-term

PA.<sup>116</sup> Future digital studies should therefore focus on reporting longer-term outcomes to understand their effectiveness over these longer periods.

## Strengths and limitations

The use of a scoping review methodology has enabled the identification of a coherent body of evidence of both developed and planned interventions, and their components, which would not have been possible with a systematic review. Indeed, many of the existing systematic reviews focus on a single condition and/or a narrow interpretation of digital tools.<sup>21,22,24</sup>

A strength of this scoping review is also the wide lens used to map literature across 18 LTCs as the first step to inform a systematic review and/or design of future digital interventions for maintenance of PA in people with multimorbidities. This broad approach also extended to the inclusion of digital tools and the use of a conservative definition for maintenance,<sup>21,29</sup> enabling a wide range of literature to be identified in this area. However, non-English language studies were not included, which may have meant that studies were missed, particularly given the number of studies identified in Europe. The maintenance definition was purposefully inclusive; however, studies were identified during the screening process that was highly relevant but did not exactly meet this definition and were subsequently excluded. Furthermore, our maintenance definition will have excluded interventions designed to be used during the maintenance period.

This review also aimed to explore the experiences, barriers and facilitators for people with LTCs to using digital tools to maintain PA. Unfortunately, we only identified qualitative data in one of the RCTs<sup>89</sup> and were therefore unable to address these objectives. On reflection, it may have been prudent to develop a second search strategy that focused on identifying qualitative and process evaluations of interventions or to undertake a snowballing approach after identifying the studies included.

Given the pace of change in this area, it is likely that a variety of new digital tools have been developed since our searches were conducted. Although we aimed to overcome this by including protocols for future trials, the COVID-19 pandemic has seen the development of many new digital resources through necessity, which was not captured. An example of this is the Kidney Beam tool, developed and launched mid-pandemic to support PA virtually for people with kidney disease.<sup>117</sup> The pandemic has accelerated the spread and adoption of digital resources<sup>39</sup> and progressed the digital ambitions of the NHS.<sup>16</sup> However, the adoption of digital tools by people with LTCs has traditionally been limited<sup>39</sup> and it is currently unclear whether the pandemic will lead to longer-term usage of these tools. Consequently, it will be important to understand the impact of the pandemic on longer-term usage and the associated impact on NHS resources in future research.

Scoping reviews often support the development of focused research questions for future systematic reviews or other empirical studies. This review identified an increasing use of digital tools over the past decade when compared with a previous review,<sup>29</sup> and included a considerable number of RCTs and protocols for RCTs. It would therefore be prudent to evaluate the effectiveness of these tools, alongside newly developed tools, using subgroup analyses to account for heterogeneity. Second, identification of key components of the interventions that successfully support the maintenance of PA for people with LTCs would be advantageous for future intervention development. As previously highlighted, an intervention component analysis<sup>104</sup> approach may be most appropriate to achieve this. Finally, the continual focus on single conditions and younger age groups (both in published and planned studies) highlights the potential for a future-focused systematic review to investigate the factors influencing positive effects across these conditions. Such a review would support the development of effective interventions for people with multimorbidities.

## Conclusions

This scoping review aimed to identify and map the characteristics of existing and planned studies using digital tools for supporting people with LTCs to maintain PA. Our novel finding is the wealth of evidence across the 18 LTCs identified. Digital tools were commonly designed for people with CVD, type 2 diabetes mellitus and obesity and most often delivered via web browsers, with some interventions also combining wearable devices. PA outcomes were most often reported at 9 months or 3 months after the end of the intervention. Some studies clearly articulated the use of theories of behavioural change in the development of the interventions but greater reporting transparency is needed to maximise the synthesis of findings to establish effectiveness, future adoption and spread of digital tools.

**Acknowledgements:** The literature searches were undertaken with the support of Vicky Fenerty (VF), an academic librarian at the University of Southampton.

**Declaration of conflicting interests:** The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

**Ethics approval and consent to participate:** Not applicable

**Contributorship:** PC, MS, SMcD conceived and planned the review. PC, SMcD, CC, KC, ZS, PM and VF contributed to the development of search strategy and PC conducted the searches. PC and SMcD undertook initial title review, and titles/abstracts were screened by PC, CC, SMcD, PM, CG, AS, JA, ZS. Full text review was undertaken by PC, PM, KC, CC, CG, AS, POG, ZS. Digital screening was undertaken by PC, SMcD;

POG, AS. Charting was undertaken by PC, KC, SMcD, ZS, CC, PM, POG, AS and synthesis by PC, POG, AS, SMcD. All authors have read and approved the manuscript.

**Ethical approval:** Not applicable, because this article does not contain any studies with human or animal subjects.

**Funding:** This report is independent research funded by the National Institute for Health Research ARC Wessex. The views expressed in this publication are those of the author(s) and not necessarily those of the National Institute for Health Research or the Department of Health and Social Care.

**Guarantor:** MS.

**Availability of data and materials:** All data generated or analysed during this study are included in this published article and its supplementary information files.

**Informed consent:** Not applicable, because this article does not contain any studies with human or animal subjects.

**ORCID iDs:** Paul Clarkson  <https://orcid.org/0000-0001-5955-5711>

Suzanne McDonough  <https://orcid.org/0000-0003-3758-3302>

**Trial registration:** Not applicable, because this article does not contain any clinical trials.

**Supplemental material:** Supplemental material for this article is available online.

### Supplemental material 1:

Appendix A: Medline search strategy

Appendix B: Digital review criteria

Appendix C: Summary data tables for included studies:

- Data charting items
- Study characteristics
- Characteristics of digital tools, maintenance periods and PA outcomes
- Theoretical underpinning

## References

1. Bull FC, Al-Ansari SS, Biddle S, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med* 2020; 54: 1451–1462.
2. World Health Organization. Physical activity. <https://www.who.int/dietphysicalactivity/pa/en/>. (2019, accessed 15th November 2019).
3. Department of health. Long-term conditions compendium of information: 3rd edition. In: *Department of health*. London: The Stationary Office, 2012. p. 3. [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/216528/dh\\_134486.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/216528/dh_134486.pdf). (2012, accessed 29 April 2019).



4. World Health Organisation (WHO). Physical activity fact sheet <https://www.who.int/publications/i/item/WHO-HEP-HPR-RUN-2021.2>. (2021, accessed 30 January 2022).
5. Office for Health Improvement and Disparities. Public health profiles. <https://fingertips.phe.org.uk/profile/physical-activity/data#page/1> (2022, accessed 30 January 2022).
6. Sport England. Active lives data: may 2019/20 and November 2019/20 'No activity in the last 28 days: % of all who are inactive by disability or long-term health condition'. <https://activelives.sportengland.org/Result?queryId=2817> (2022, accessed 30 January 2022)
7. Saunders DH, Sanderson M, Hayes S, et al. Physical fitness training for stroke patients. *Cochrane Database Syst Rev* 2020; 3: CD003316.
8. Moseng T, Tveter AT, Holm I, et al. Patients with musculoskeletal conditions do less vigorous physical activity and have poorer physical fitness than population controls: a cross-sectional study. *Physiotherapy* 2014; 100: 319–324
9. Rausch Osthoff A, Niedermann K, Braun J, et al. EULAR Recommendations for physical activity in people with inflammatory arthritis and osteoarthritis. *ARD* 2018; 77: 1251–1260
10. Marley J, Tully MA, Porter-Armstrong A, et al. The effectiveness of interventions aimed at increasing physical activity in adults with persistent musculoskeletal pain: a systematic review and meta-analysis. *BMC Musculoskelet Disord* 2017; 18(1): 482
11. Carson KV, Chandratilleke MG, Picot J, et al. Physical training for asthma. *Cochrane Database Syst Rev* 2013; 30: CD001116
12. World Health Organization. Classification of digital health interventions v1.0. <https://www.who.int/reproductivehealth/publications/mhealth/classification-digital-health-interventions/en/>. (2018, accessed 12 March 2020)
13. Irani I, Niyomyart A and Hickman RL. Systematic review of technology-based interventions targeting chronically ill adults and their caregivers. *WJNR* 2020; 42: 1–19
14. Berry A, McCabe CS, Muir S, et al. Digital behaviour change interventions to facilitate physical activity in osteoarthritis: a systematic review. *Phys Ther Rev* 2018; 23(3): 197–206
15. Connelly J, Kirk A, Masthoff J, et al. The use of technology to promote physical activity in type 2 diabetes management: a systematic review. *Diabetic Med* 2013; 30: 1420–1432
16. National Health Service. The NHS long term plan. <https://www.england.nhs.uk/wp-content/uploads/2017/03/NEXT-STEPS-ON-THE-NHS-FIVE-YEAR-FORWARD-VIEW.pdf>. (2019, accessed 18 March 2021).
17. World Health Organisation (WHO). Global strategy on digital health 2020–2025. <https://www.who.int/docs/defaultsource/documents/g4dhdaa2a9f352b0445bafbc79ca799dce4d.pdf> (2021, accessed 30 January 2022)
18. Jiang X, Ming WK and You JH. The cost-effectiveness of digital health interventions on the management of cardiovascular diseases: systematic review. *J Med Internet Res* 2019; 21(6): e13166.
19. Pereira PB, Johnson C and Vorderstrasse A. Internet delivered diabetes self-management education: a review. *Diabetes Technology and Therapeutics* 2015; 17: 55–63
20. Hutchings R. The impact of COVID-19 on the use of digital technology in the NHS. Briefing, Nuffield Trust (2020, accessed 30 January 2022)
21. Grimmert C, Corbett T, Brunet J, et al. Systematic review and meta-analysis of maintenance of physical activity behaviour change in cancer survivors. *IJBNPA* 2019; 16: 1–20
22. Samdal GB, Eide GE, Barth T, et al. Effective behaviour change techniques for physical activity and healthy eating in overweight and obese adults; systematic review and meta-regression analyses. *IJBNPA* 2017; 14: 42
23. McCabe C, McCann M and Brady AM. Computer and mobile technology interventions for self-management in chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2017;5 (5): CD011425.
24. Griffiths AJ, White CM, Thain PK, et al. The effect of interactive digital interventions on physical activity in people with inflammatory arthritis: a systematic review. *Rheumatol Int* 2018; 38: 1623–1634
25. Stellefson M, Chaney B, Barry AE, et al. Web 2.0 chronic disease self-management for older adults: a systematic review. *J Med Internet Res* 2013; 15: e35
26. Walker KC, Valentiner LS and Langberg H. Motivational factors for initiating, implementing, and maintaining physical activity behavior following a rehabilitation program for patients with type 2 diabetes: a longitudinal, qualitative, interview study. *Patient Prefer and Adherence* 2018; 12: 145–152
27. Nigg CR, Borrelli B, Maddock J, et al. A theory of physical activity maintenance. *Appl. Psychol Int Rev Appl. Psychol* 2008; 57: 544–560.
28. Murray MM, Brennan SF, French DP, et al. Effectiveness of physical activity interventions in achieving behaviour change maintenance in young and middle-aged adults: a systematic review and meta-analysis. *Soc Sci Med* 2017; 192: 125–133
29. Fjeldsoe B, Neuhaus M, Winkler E, et al. Systematic review of maintenance of behaviour change following physical activity and dietary interventions. *Health Psychol* 2011; 30: 99–109
30. Kwasnicka D, Dombrowski SU, White M, et al. Theoretical explanations for maintenance of behaviour change: a systematic review of behaviour theories. *Health Psychol. Rev* 2016; 10: 277–296.
31. Rothman AJ. Toward a theory-based analysis of behavioral maintenance. *Health Psychol* 2000; 19: 64–69.
32. Tricco AC, Lillie E, Zarin W, et al. A scoping review on the conduct and reporting of scoping reviews. *BMC Med Res Methodol* 2016; 16: 15.
33. Levac D, Colquhoun H and O'Brien K. Scoping studies: advancing the methodology. *Implement Sci* 2010; 5: 69.
34. Maden M, Cunliffe A, McMahon N., et al. Use of programme theory to understand the differential effects of interventions across socio-economic groups in systematic reviews—a systematic methodology review. *Syst Rev* 2017; 6: 66.
35. Gourlan M and Bernard PCB. Efficacy of theory-based interventions to promote physical activity. A meta-analysis of randomised controlled trials. *Health Psychol Rev* 2014; 1: 74.
36. Michie S, Richardson M, Johnston M, et al. The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for



- the reporting of behavior change interventions. *Ann Behav Med* 2013; 46: 81–95.
37. Strobach T, Englert C, Jekauc D, et al. Predicting adoption and maintenance of physical activity in the context of dual-process theories. *Perform Enhanc Health* 2020; 8(1): 100162
  38. Deci EL and Ryan RM. *Intrinsic motivation and self-determination in human behaviour*. New York: Planum Publishing; 1985.
  39. Organisation for the Review of Care and Health Apps (ORCHA). Digital health trends and opportunities for 2021. [https://orchahealth.com/wp-content/uploads/2021/01/COVID\\_Report\\_Jan\\_2021\\_final-version.pdf](https://orchahealth.com/wp-content/uploads/2021/01/COVID_Report_Jan_2021_final-version.pdf). (2021, accessed 19 March 2021).
  40. Tricco AC, Lillie E, Zarin W, et al. PRISMA Extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med* 2018b; 169: 467–473
  41. Peters MDJ, Godfrey C, et al. Chapter 11: scoping reviews. In: Aromataris E and Munn Z (eds). *Joanna briggs institute reviewer's manual*. Adelaide, Australia: The Joanna Briggs Institute, 2017. <https://wiki.jbi.global/display/MANUAL/Chapter+11%3A+Scoping+reviews> (2017, accessed 18 October 2019).
  42. Arksey H and O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol* 2005; 8: 19–32
  43. Clarkson P, Adams J, Muckelt P, et al. (Protocol) maintaining physical activity through the use of digital tools for people with a long-term condition/s (LTCs): a scoping review. *Protocols.io* 2020a, DOI: 10.17504/protocols.io.bf7gjrjw
  44. NHS Digital. Quality and outcomes framework (QOF), England: 2017–2018 prevalence, achievement and exceptions report. <https://digital.nhs.uk/data-and-information/publications/statistical/quality-and-outcomes-framework-achievement-prevalence-and-exceptions-data/2017-18> (2018, accessed 23 May 2019).
  45. National Institute for Health and Care Excellence (NICE). Physical activity pathway. <https://pathways.nice.org.uk/pathways/physical-activity> (2019, accessed 23 October 2019).
  46. Roberts AL, Fisher A, Smith L, et al. Digital health behaviour change interventions targeting physical activity and diet in cancer survivors: a systematic review and meta-analysis. *J Cancer Surviv* 2017; 11: 704–719
  47. Garg S, Garg D, Turin TC, et al. Web-based interventions for chronic back pain: a systematic review. *J Med Internet Res*. 2016; 18: 139
  48. UK Department of Health and Social Care. Physical activity guidelines. <https://www.gov.uk/government/collections/physical-activity-guidelines> (2019, accessed 14 November 2019).
  49. Ottenbacher AJ, Day RS, Taylor WC, et al. Long-term physical activity outcomes of home-based lifestyle interventions among breast and prostate cancer survivors. *Support Care Cancer* 2012; 20: 2483–2489
  50. Stephenson A, McDonough SM, Murphy MH, et al. Using computer, mobile and wearable technology enhanced interventions to reduce sedentary behaviour: a systematic review and meta-analysis. *IJBNPA* 2017; 14: 105.
  51. Burke S, Wurz A, Bradshaw A, et al. Physical activity and quality of life: a meta-synthesis of qualitative research. *Cancers (Basel)*. 2017; 9: 53.
  52. Michie S, West R, Campbell R, et al. *ABC Of behaviour change theories. An essential resource for researchers, policy makers and practitioners*. Sutton, UK: Silverback, 2014.
  53. Abraham C and Michie S. A taxonomy of behavior change techniques used in interventions. *Health Psychol* 2008; 27: 379–387.
  54. Lubans DR, Morgan PJ, Collins CE, et al. Exploring the mechanisms of weight loss in the SHED-IT intervention for overweight men: a mediation analysis. *Int J Behav Nutr Phys Act* 2009; 6: 76.
  55. Hurkmans EJ, van den Berg MH, Runday KH, et al. Maintenance of physical activity after internet-based physical activity interventions in patients with rheumatoid arthritis. *Rheumatology* 2010; 49: 167–172.
  56. Lorig K, Ritter PL, Laurent DD, et al. Online diabetes self-management program: a randomized study. *Diabetes Care* 2010; 33: 1275–1281.
  57. Reid RD, Morrin LI, Beaton LJ, et al. Randomized trial of an internet-based computer-tailored expert system for physical activity in patients with heart disease. *Eur J Prev Cardiol* 2012; 19: 1357–1364.
  58. Bossen D, Veenhof C, Van Beek KE, et al. Effectiveness of a web-based physical activity intervention in patients with knee and/or hip osteoarthritis: randomized controlled trial. *J Med Internet Res* 2013; 15: e257.
  59. Ström M, Uckelstam C, Andersson G, et al. Internet-delivered therapist-guided physical activity for mild to moderate depression: a randomized controlled trial. *Peer J* 2013; 1: 178
  60. Devi R, Powell J and Singh S. A web-based program improves physical activity outcomes in a primary care angina population: randomized controlled trial. *J Med Internet Res* 2014; 16: e186.
  61. Jennings CA, Vandelanotte C, Caperchione CM, et al. Effectiveness of a web-based physical activity intervention for adults with type 2 diabetes—a randomised controlled trial. *Prev. Med* 2014; 60: 33–40.
  62. van der Weegen S, Verwey R, Spreeuwenberg M, et al. It's LiFe! Mobile and web-based monitoring and feedback tool embedded in primary care increases physical activity: a cluster randomized controlled trial. *J Med Internet Res* 2015; 17: e184.
  63. Olson EA and McAuley E. Impact of a brief intervention on self-regulation, self-efficacy and physical activity in older adults with type 2 diabetes. *J Behav Med* 2015; 38: 886–898.
  64. Vorrink SN, Kort HS, Troosters T, et al. Efficacy of an mHealth intervention to stimulate physical activity in COPD patients after pulmonary rehabilitation. *Eur Resp Journal* 2016; 48: 1019–1029.
  65. Thorup C, Hansen J, GrønkJær M, et al. Cardiac patients' walking activity determined by a step counter in cardiac telerehabilitation: data from the intervention arm of a randomized controlled trial. *J Med Internet Res*. 2016; 18: 69.
  66. Givon N, Zeilig G, Weingarden H, et al. Video games used in a group setting is feasible and effective to improve indicators of physical activity in individuals with chronic stroke: a randomized controlled trial. *Clin Rehabil* 2016; 30:383–392.
  67. Yang YP, Wang CJ, Wang JJ, et al. The effects of an activity promotion system on active living in overweight subjects

- with metabolic abnormalities. *Obesity Research & Clinical Practic* 2017; 11: 718–727
68. Alonso-Domínguez R, Patino-Alonso MC, Sánchez-Aguadero N, et al. Effect of a multifactorial intervention on the increase in physical activity in subjects with type 2 diabetes mellitus: a randomized clinical trial (EMID study). *Eur J Cardiovasc Nurs* 2019; 18: 399–409.
  69. Dor-Haim H, Katzburg S and Leibowitz D. A novel digital platform for a monitored home-based cardiac rehabilitation program. *J Vis Exp* 2019;146:e59019. doi: 10.3791/59019.
  70. Patel MS, Small DS, Harrison JD, et al. Effectiveness of behaviorally designed gamification interventions with social incentives for increasing physical activity among overweight and obese adults across the United States: the STEP UP randomized clinical trial. *JAMA Intern Med* 2019; 179: 1–9.
  71. Avila A, Claes J, Buys R, et al. Home-based exercise with telemonitoring guidance in patients with coronary artery disease: does it improve long-term physical fitness? *Eur J Prev Cardiol* 2019; 27: 367–377.
  72. Lari H, Noroozi A and Tahmasebi R. Impact of short message service (SMS) education based on a health promotion model on the physical activity of patients with type II diabetes. *MJMS* 2018; 25: 67–77
  73. Verwey R, van der Weegen S, Spreeuwenberg M, et al. A monitoring and feedback tool embedded in a counselling protocol to increase physical activity of patients with COPD or type 2 diabetes in primary care: study protocol of a three-arm cluster randomised controlled trial. *BMC Fam Pract.* 2014; 15: 93.
  74. Alonso-Domínguez R, Gómez-Marcos MA, Patino-Alonso MC, et al. Effectiveness of a multifactorial intervention based on an application for smartphones, heart-healthy walks and a nutritional workshop in patients with type 2 diabetes mellitus in primary care (EMID): study protocol for a randomised controlled trial. *BMJ Open* 2017; 7: e016191.
  75. Harrison JD, Jones JM, Small DS, et al. Social incentives to encourage physical activity and understand predictors (STEP UP): design and rationale of a randomized trial among overweight and obese adults across the United States. *Contemp Clin Trials* 2019; 80: 55–60.
  76. Barry VW, McClain AC, Shuger S, et al. Using a technology-based intervention to promote weight loss in sedentary overweight or obese adults: a randomized controlled trial study design. *Diabetes Metab Syndr Obes* 2011; 4: 67–77.
  77. Kloek CJ, Bossen D, Veenhof C, et al. Effectiveness and cost-effectiveness of a blended exercise intervention for patients with hip and/or knee osteoarthritis: study protocol of a randomized controlled trial. *BMC Musculoskelet Disord* 2014; 15: 69.
  78. Fife-Schaw C, de Lusignan S, Wainwright J, et al. Comparing exercise interventions to increase persistence with physical exercise and sporting activity among people with hypertension or high normal blood pressure: study protocol for a randomised controlled trial. *Trials* 2014; 15: 36.
  79. Banos RM, Mensorio MS, Cebolla A, et al. An internet-based self-administered intervention for promoting healthy habits and weight loss in hypertensive people who are overweight or obese: a randomized controlled trial. *BMC Cardiovasc Disord* 2015; 15: 83.
  80. Jaarsma T, Klompstra L, Ben Gal T, et al. Increasing exercise capacity and quality of life of patients with heart failure through wii gaming: the rationale, design and methodology of the HF-Wii study; a multicentre randomized controlled trial. *Eur J Heart Fail* 2015; 17: 743–748.
  81. Brouwers RWM, Kraal JJ, Traa SC, et al. Effects of cardiac telerehabilitation in patients with coronary artery disease using a personalised patient-centred web application: protocol for the SmartCare-CAD randomised controlled trial. *BMC Cardiovasc Disord* 2017; 17: 46.
  82. Ingram W, Webb D, Taylor RS, et al. Multicentred randomised controlled trial of an augmented exercise referral scheme using web-based behavioural support in individuals with metabolic, musculoskeletal and mental health conditions: protocol for the e-coachER trial. *BMJ Open.* 2018; 8: e022382.
  83. Cox NS, McDonald CF, Alison JA, et al. Telerehabilitation versus traditional centre-based pulmonary rehabilitation for people with chronic respiratory disease: protocol for a randomised controlled trial. *BMC Pulm Med* 2018; 18: 71.
  84. Bonn SE, Alexandrou C, Hjörleifsdóttir Steiner K, et al. App-technology to increase physical activity among patients with diabetes type 2 - the DiaCert-study, a randomized controlled trial. *BMC Public Health* 2018; 18: 19.
  85. Volders E, Bolman CAW, de Groot RHM, et al. The effect of active Plus, a computer-tailored physical activity intervention, on cognitive functioning of elderly people with chronic illness(es) - study protocol for a randomized controlled trial. *BMC Public Health* 2019; 19: 197
  86. Sharma A, Mentz RJ, Granger BB, et al. Utilizing mobile technologies to improve physical activity and medication adherence in patients with heart failure and diabetes mellitus: rationale and design of the TARGET-HF-DM trial. *Am Heart J* 2019; 211: 22–33.
  87. Jones TM, Dear BF, Hush JM, et al. Mymoves program: feasibility and acceptability study of a remotely delivered self-management program for increasing physical activity among adults with acquired brain injury living in the community. *Phys Ther.* 2016; 96: 1982–1993.
  88. Vorderstrasse A, Melkus GD, Feinglos M, et al. Abstract 17519: virtual environment for diabetes self-management education and support: preliminary RCT outcomes. *Circulation* 2017; 136: A17519.
  89. Hawkins J, Charles JM, Edwards M, et al. Acceptability and feasibility of implementing accelerometry-based activity monitors and a linked web portal in an exercise referral scheme: feasibility randomized controlled trial. *J Medical Internet Res* 2019; 21: e12374.
  90. Barnason SA, Young L, Kupzyk K, et al. Abstract 18101: examining self-efficacy and patient activation mechanisms in a weight management intervention for overweight and obese cardiac rehabilitation patients. *Circulation* 2016; 134: 18101.
  91. Verwey R, Van der Weegen S, Spreeuwenberg M, et al. Process evaluation of physical activity counselling with and without the use of mobile technology: a mixed methods study. *Int J Nurs Stud* 2016; 53: 3–16.
  92. European Commission. Ehealth: digital health and care. [https://ec.europa.eu/health/ehealth/home\\_en](https://ec.europa.eu/health/ehealth/home_en). (2021, accessed 30 March 2021).

93. Government of the Netherlands. Government encouraging the use of eHealth. <https://www.government.nl/topics/ehealth/government-encouraging-use-of-ehealth>. (2021, accessed 30 March 2021).
  94. 42 Matters. Store stats 2021: Google play store and apple app store. <https://42matters.com/google-play-statistics-and-trends> (2021, accessed 2 March 2021).
  95. Aromatario O, Van Hoyo A, Vuillemin A, et al. How do mobile health applications support behaviour changes? A scoping review of mobile health applications relating to physical activity and eating behaviours. *Public Health* 2019; 175: 8–18.
  96. National Institute for Health and Care Excellence. Behaviour change: individual approaches. Public health guideline (PH49). <https://www.nice.org.uk/guidance/ph49>. (2014, accessed 3 March 2021).
  97. Craig P, Dieppe P, Macintyre S, et al. Developing and evaluating complex interventions: the new medical research council guidance. *BMJ (Clinical Research ed.)* 2008; 337: a1655.
  98. Romeo AV, Edney SM, Plotnikoff RC, et al. Examining social-cognitive theory constructs as mediators of behaviour change in the active team smartphone physical activity program: a mediation analysis. *BMC Public Health* 2021; 21: 88.
  99. Stacey FG, James EL, Chapman K, et al. A systematic review and meta-analysis of social cognitive theory-based physical activity and/or nutrition behavior change interventions for cancer survivors. *J Cancer Surviv* 2015; 9: 305–338.
  100. Hoffmann TC, Glasziou PP, Boutron I, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *Br Med J* 2014; 348: g1687
  101. Duff OM, Walsh DM, Furlong BA, et al. Behavior change techniques in physical activity eHealth interventions for people with cardiovascular disease: systematic review. *J Med Internet Res* 2017; 19: e281.
  102. Michie S, Fixsen D, Grimshaw JM, et al. Specifying and reporting complex behavior change interventions: the need for a scientific method. *Implement. Sci* 2009; 40: 1–6
  103. Moller AC, Merchant G, Conroy DE., et al. Applying and advancing behavior change theories and techniques in the context of a digital health revolution: proposals for more effectively realizing untapped potential. *J Behav Med* 2017; 40: 85–98
  104. Sutcliffe K, Thomas J, Stokes G, et al. Intervention component analysis (ICA): a pragmatic approach for identifying the critical features of complex interventions. *Syst Rev* 2015; 4: 40.
  105. Morton K, Dennison L, May C, et al. Using digital interventions for self-management of chronic physical health conditions: a meta-ethnography review of published studies. *Patient Educ Couns.* 2016; 100: 616–635.
  106. Lorig KR, Sobel DS, Ritter PL, et al. Effect of a self-management program on patients with chronic disease. *ECP* 2001; 4: 256–262.
  107. The Kings Fund. Technology and innovation for long-term health conditions. <https://www.kingsfund.org.uk/sites/default/files/202007/Technology%20and%20innovation%20for%20long-term%20health%20conditions%20August%202020.pdf>. (2020, accessed 2 March 2021).
  108. Castle-Clarke S, Edwards N and Buckingham H. Falling short: why the NHS is still struggling to make the most of new innovations the nuffield trust Available from: <https://www.nuffieldtrust.org.uk/research/falling-short-why-the-nhs-is-still-struggling-to-make-the-most-of-new-innovations> (2017, accessed 15 June 2021)
  109. Office for National Statistics. People with long-term health conditions, UK: January to December 2019. <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/adhocs/11478peoplewithlongtermhealthconditionsukjanuarytodecember2019>. (2020a, accessed 19 January 2021).
  110. An Phriomh-Oifig Staidrimh Central Statistics Office. Irish Health survey – main results. 2019. <https://www.cso.ie/en/releasesandpublications/ep/p-ihsmr/irishhealthsurvey2019-mainresults/healthstatus/>. (2019, accessed 30 March 2021).
  111. Roth GA, Mensah GA, Johnson CO, et al. Global burden of cardiovascular diseases and risk factors, 1990–2019: update from the GBD.2019 study. *J Am Coll Cardiol* 2019;76(25): 2982–3021. doi:10.1016/j.jacc.2020.11.010
  112. Glovaci D., Fan W. and Wong N.D. Epidemiology of diabetes Mellitus and cardiovascular disease. *Curr Cardiol Rep* 2019; 21: 21.
  113. Eurostat. Ageing Europe – statistics on population developments. [https://ec.europa.eu/eurostat/statistics-explained/index.php/Ageing\\_Europe\\_-\\_statistics\\_on\\_population\\_developments](https://ec.europa.eu/eurostat/statistics-explained/index.php/Ageing_Europe_-_statistics_on_population_developments). (2021, accessed 1 March 2021).
  114. Forsat ND, Palmowski A, Palmowski Y, et al. Recruitment and retention of older people in clinical research: a systematic literature review. *J Am Geriatr Soc* 2020; 68: 2955–2963.
  115. Vaportzis E, Clausen MG and Gow AJ. Older adults perceptions of technology and barriers to interacting with tablet computers: a focus group study. *Front Psychol* 2017; 8: 1687.
  116. Wahlich C, Chaudhry UAR, Fortescue R, et al. Effectiveness of adult community-based physical activity interventions with objective physical activity measurements and long-term follow-up: a systematic review and meta-analysis. *BMJ Open* 2020; 10: e034541
  117. Kidney Beam. Welcome to Kidney Beam. <https://beamfeelgood.com/kidney-disease>. (2021, accessed 19 March 2021).
-