



EDEXCEL INTERNATIONAL GCSE (9–1)

FURTHER PURE MATHEMATICS




Student Book

Ali Datto

Online access to your ActiveBook

Thank you for buying this Edexcel International GCSE (9–1) Further Pure Mathematics Student Book. It comes with three years' access* to your ActiveBook – an online, digital version of your textbook. You can personalise your ActiveBook with notes, highlights and links to your wider reading. It is perfect for supporting your coursework and revision activities.

*For new purchases only. If this access code has already been revealed, it may no longer be valid. If you have bought this textbook second hand, the code may already have been used by the first owner of the book.



How to access your ActiveBook

1	Scratch off panel with a coin to reveal your unique access code. Do not use a knife or other sharp object as it may damage the code.
2	Go to www.pearsonactivelearn.com
3	If you already have an ActiveLearn Digital Services account (ActiveTeach or ActiveLearn), log in and click 'I have a new access code' in the top right of the screen. <ul style="list-style-type: none">Type in the code above and select 'Activate'.
4	If you do not have an ActiveLearn Digital Services account, click 'Register'. It is free to do this. <ul style="list-style-type: none">Type in the code above and select 'Activate'.Simply follow the instructions on screen to register.

Important information

- The access code can only be used once.
- Please activate your access code as soon as possible, as it does have a 'use by date'. If your code has expired when you enter it, please contact our ActiveLearn support site at digital.support@pearson.com
- The ActiveBook will be valid for three years upon activation.

Getting help

- To check that you will be able to access an ActiveBook, go to www.pearsonactivelearn.com and choose 'Will ActiveLearn Digital Service work on my computer?'
- If you have any questions about accessing your ActiveBook, please contact our ActiveLearn support site at www.pearsonactivelearn.com/support

EDEXCEL INTERNATIONAL GCSE (9–1)

FURTHER PURE MATHEMATICS

Student Book

Ali Datto

**Greg Attwood, Keith Pledger, David Wilkins, Alistair Macpherson,
Bronwen Moran, Joseph Petran and Geoff Staley**

Published by Pearson Education Limited, 80 Strand, London, WC2R 0RL.

www.pearsonglobalschools.com

Copies of official specifications for all Pearson qualifications may be found on the website: <https://qualifications.pearson.com>

Text © Pearson Education Limited 2017

Edited by Linnet Bruce, Keith Gallick and Susan Lyons

Designed by Cobalt id

Typeset by Tech-Set Ltd, Gateshead, UK

Cover design by Pearson Education Limited

Picture research by Debbie Gallagher

Cover photo © Getty Images: Naahushi Kavuri / EyeEm

Inside front cover: **Shutterstock.com**: Dmitry Lobanov

The rights of Ali Datto, Greg Attwood, Keith Pledger, David Wilkins, Alistair Macpherson, Bronwen Moran, Joseph Petran and Geoff Staley to be identified as authors of this work have been asserted by them in accordance with the Copyright, Designs and Patents Act 1988.

First published 2017

20 19 18 17

10 9 8 7 6 5 4 3 2 1

British Library Cataloguing in Publication Data

A catalogue record for this book is available from the British Library

ISBN 978 0 435 18854 2

Copyright notice

All rights reserved. No part of this publication may be reproduced in any form or by any means (including photocopying or storing it in any medium by electronic means and whether or not transiently or incidentally to some other use of this publication) without the written permission of the copyright owner, except in accordance with the provisions of the Copyright, Designs and Patents Act 1988 or under the terms of a licence issued by the Copyright Licensing Agency, Barnard's Inn, 86 Fetter Lane, London EC4A 1EN (www.cla.co.uk). Applications for the copyright owner's written permission should be addressed to the publisher.

Printed in Slovakia by Neografia

Acknowledgements

The publisher would like to thank the following for their kind permission to reproduce their photographs:

(Key: b-bottom; c-centre; l-left; r-right; t-top)

123RF.com: ealisa 143; **Alamy Stock Photo**: Ammit 3; **Getty Images**: Christoph Hetzmannseder 2, David Crunelle / EyeEm 160, Ditto 24, EyeEm 136, hohl 74, Julia Antonio / EyeEm 60, Krzysztof Dydynski 106, Martin Barraud 124, Ozgur Donmaz 36, sebastian-julian 96, View Pictures / UIG 182; **Shutterstock.com**: Africa Studio 75, Akhenaton Images 90, Aleksandr Markin 128, andrea crisante 107t, Anette Holmberg 37, Brian Kinney 161, cigdem 148, IM_photo 123, Jacques Durocher 49t, jordache 180, Maridav 83, Miami2you 130, migrean 107b, Mikael Damkier 160, Nando 11, Nattawadee Supchapo 137, Neil Lockhart 61, Nicku 80, oksmit 125, Ondrej Prosicky 183, Patricia Hofmeester 131, Pavel L Photo and Video 154, photofriday 77t, Rawpixel.com 9, Sergey Edentod 47, Steve Noakes 25, Svetlana Privezentseva 49b, Syda Productions 77b, Viacheslav Nikolaenko 92, wavebreakmedia 121, worradirek 10, Zhukov Oleg 5

All other images © Pearson Education

Endorsement statement

In order to ensure that this resource offers high-quality support for the associated Pearson qualification, it has been through a review process by the awarding body. This process confirms that this resource fully covers the teaching and learning content of the specification or part of a specification at which it is aimed. It also confirms that it demonstrates an appropriate balance between the development of subject skills, knowledge and understanding, in addition to preparation for assessment.

Endorsement does not cover any guidance on assessment activities or processes (e.g. practice questions or advice on how to answer assessment questions), included in the resource nor does it prescribe any particular approach to the teaching or delivery of a related course.

While the publishers have made every attempt to ensure that advice on the qualification and its assessment is accurate, the official specification and associated assessment guidance materials are the only authoritative source of information and should always be referred to for definitive guidance.

Pearson examiners have not contributed to any sections in this resource relevant to examination papers for which they have responsibility.

Examiners will not use endorsed resources as a source of material for any assessment set by Pearson. Endorsement of a resource does not mean that the resource is required to achieve this Pearson qualification, nor does it mean that it is the only suitable material available to support the qualification, and any resource lists produced by the awarding body shall include this and other appropriate resources.

PREFACE	vi
CHAPTER 1: SURDS AND LOGARITHMIC FUNCTIONS	2
CHAPTER 2: THE QUADRATIC FUNCTION	24
CHAPTER 3: INEQUALITIES AND IDENTITIES	36
CHAPTER 4: SKETCHING POLYNOMIALS	60
CHAPTER 5: SEQUENCES AND SERIES	74
CHAPTER 6: THE BINOMIAL SERIES	96
CHAPTER 7: SCALAR AND VECTOR QUANTITIES	106
CHAPTER 8: RECTANGULAR CARTESIAN COORDINATES	124
CHAPTER 9: DIFFERENTIATION	136
CHAPTER 10: INTEGRATION	160
CHAPTER 11: TRIGONOMETRY	182
GLOSSARY	212
ANSWERS	216
INDEX	238

CHAPTER 1

WRITE A NUMBER EXACTLY USING A SURD	4
RATIONALISE THE DENOMINATOR OF A SURD	5
BE FAMILIAR WITH THE FUNCTIONS a^x AND $\log_b x$ AND RECOGNISE THE SHAPES OF THEIR GRAPHS	7
BE FAMILIAR WITH EXPRESSIONS OF THE TYPE e^x AND USE THEM IN GRAPHS	9
BE ABLE TO USE GRAPHS OF FUNCTIONS TO SOLVE EQUATIONS	12
WRITING AN EXPRESSION AS A LOGARITHM	14
UNDERSTAND AND USE THE LAWS OF LOGARITHMS	15
CHANGE THE BASE OF A LOGARITHM	17
SOLVE EQUATIONS OF THE FORM $a^x = b$	18
EXAM PRACTICE QUESTIONS	21
CHAPTER SUMMARY	23

CHAPTER 2

FACTORISE QUADRATIC EXPRESSIONS WHERE THE COEFFICIENT OF x^2 IS GREATER THAN 1	26
COMPLETE THE SQUARE AND USE THIS TO SOLVE QUADRATIC EQUATIONS	27
SOLVE QUADRATIC EQUATIONS USING THE QUADRATIC FORMULA	28
UNDERSTAND AND USE THE DISCRIMINANT TO IDENTIFY WHETHER THE ROOTS ARE (i) EQUAL AND REAL, (ii) UNEQUAL AND REAL OR (iii) NOT REAL	29
UNDERSTAND THE ROOTS α AND β AND HOW TO USE THEM	31
EXAM PRACTICE QUESTIONS	34
CHAPTER SUMMARY	35

CHAPTER 3

SOLVE SIMULTANEOUS EQUATIONS, ONE LINEAR AND ONE QUADRATIC	38
SOLVE LINEAR INEQUALITIES	39
SOLVE QUADRATIC INEQUALITIES	41
GRAPH LINEAR INEQUALITIES IN TWO VARIABLES	44
DIVIDE A POLYNOMIAL BY $(x \pm p)$	50

FACTORISE A POLYNOMIAL BY USING THE FACTOR THEOREM	52
USING THE REMAINDER THEOREM, FIND THE REMAINDER WHEN A POLYNOMIAL IS DIVIDED BY $(ax - b)$	54
EXAM PRACTICE QUESTIONS	56
CHAPTER SUMMARY	58

CHAPTER 4

SKETCH CUBIC CURVES OF THE FORM $y = ax^3 + bx^2 + cx + d$ OR $y = (x + a)(x + b)(x + c)$	62
SKETCH AND INTERPRET GRAPHS OF CUBIC FUNCTIONS OF THE FORM $y = x^3$	64
SKETCH THE RECIPROCAL FUNCTION $y = \frac{k}{x}$ WHERE k IS A CONSTANT	65
SKETCH CURVES OF DIFFERENT FUNCTIONS TO SHOW POINTS OF INTERSECTION AND SOLUTIONS TO EQUATIONS	67
APPLY TRANSFORMATIONS TO CURVES	69
EXAM PRACTICE QUESTIONS	71
CHAPTER SUMMARY	73

CHAPTER 5

IDENTIFY AN ARITHMETIC SEQUENCE	76
FIND THE COMMON DIFFERENCE, FIRST TERM AND n th TERM OF AN ARITHMETIC SERIES	78
FIND THE SUM OF AN ARITHMETIC SERIES AND BE ABLE TO USE Σ NOTATION	80
IDENTIFY A GEOMETRIC SEQUENCE	84
FIND THE COMMON RATIO, FIRST TERM AND n th TERM OF A GEOMETRIC SEQUENCE	85
FIND THE SUM OF A GEOMETRIC SERIES	87
FIND THE SUM TO INFINITY OF A CONVERGENT GEOMETRIC SERIES	90
EXAM PRACTICE QUESTIONS	93
CHAPTER SUMMARY	95

CHAPTER 6

USE $\binom{n}{r}$ TO WORK OUT THE COEFFICIENTS IN THE BINOMIAL EXPANSION	98
---	----

USE THE BINOMIAL EXPANSION TO EXPAND $(1 + x)^n$	100
DETERMINE THE RANGE OF VALUES FOR WHICH x IS TRUE AND VALID FOR AN EXPANSION	101
EXAM PRACTICE QUESTIONS	104
CHAPTER SUMMARY	105

CHAPTER 7

VECTOR NOTATION AND HOW TO DRAW VECTOR DIAGRAMS	108
PERFORM SIMPLE VECTOR ARITHMETIC AND UNDERSTAND THE DEFINITION OF A UNIT VECTOR	110
USE VECTORS TO DESCRIBE THE POSITION OF A POINT IN TWO DIMENSIONS	113
USE VECTORS TO DEMONSTRATE SIMPLE PROPERTIES OF GEOMETRICAL FIGURES	114
WRITE DOWN AND USE CARTESIAN COMPONENTS OF A VECTOR IN TWO DIMENSIONS	117
EXAM PRACTICE QUESTIONS	120
CHAPTER SUMMARY	123

CHAPTER 8

WORK OUT THE GRADIENT OF A STRAIGHT LINE	125
FIND THE EQUATION OF A STRAIGHT LINE	127
UNDERSTAND THE RELATIONSHIP BETWEEN PERPENDICULAR LINES	128
FIND THE DISTANCE BETWEEN TWO POINTS ON A LINE	131
FIND THE COORDINATES OF A POINT THAT DIVIDES A LINE IN A GIVEN RATIO	132
EXAM PRACTICE QUESTIONS	133
CHAPTER SUMMARY	135

CHAPTER 9

FIND THE GRADIENT FUNCTION OF A CURVE AND DIFFERENTIATE A FUNCTION THAT HAS MULTIPLE POWERS OF x	138
DIFFERENTIATE e^{ax} , $\sin ax$ AND $\cos ax$	141
USE THE CHAIN RULE TO DIFFERENTIATE MORE COMPLICATED FUNCTIONS	143
USE THE PRODUCT RULE TO DIFFERENTIATE MORE COMPLICATED FUNCTIONS	146

USE THE QUOTIENT RULE TO DIFFERENTIATE MORE COMPLICATED FUNCTIONS	148
FIND THE EQUATION OF THE TANGENT AND NORMAL TO THE CURVE $y = f(x)$	149
FIND THE STATIONARY POINTS OF A CURVE AND CALCULATE WHETHER THEY ARE MINIMUM OR MAXIMUM STATIONARY POINTS	151
APPLY WHAT YOU HAVE LEARNT ABOUT TURNING POINTS TO SOLVE PROBLEMS	153
EXAM PRACTICE QUESTIONS	156
CHAPTER SUMMARY	159

CHAPTER 10

INTEGRATION AS THE REVERSE OF DIFFERENTIATION	162
UNDERSTAND HOW CALCULUS IS RELATED TO PROBLEMS INVOLVING KINEMATICS	164
USE INTEGRATION TO FIND AREAS	166
USE INTEGRATION TO FIND A VOLUME OF REVOLUTION	171
RELATE RATES OF CHANGE TO EACH OTHER	174
EXAM PRACTICE QUESTIONS	178
CHAPTER SUMMARY	181

CHAPTER 11

MEASURE ANGLES IN RADIANs	184
CALCULATE ARC LENGTH AND THE AREA OF A CIRCLE USING RADIANs	185
UNDERSTAND THE BASIC TRIGONOMETRICAL RATIOS AND SINE, COSINE AND TANGENT GRAPHS	190
USE SINE AND COSINE RULES	193
USE SINE AND COSINE RULES TO SOLVE PROBLEMS IN 3D	197
USE TRIGONOMETRY IDENTITIES TO SOLVE PROBLEMS	199
SOLVE TRIGONOMETRIC EQUATIONS	202
USE TRIGONOMETRIC FORMULAE TO SOLVE EQUATIONS	206
EXAM PRACTICE QUESTIONS	208
CHAPTER SUMMARY	210

GLOSSARY

ANSWERS

212
216

ABOUT THIS BOOK

This book is written for students following the Edexcel International GCSE (9–1) Further Pure Maths specification and covers both years of the course. The specification and sample assessment materials for Further Pure Maths can be found on the Pearson Qualifications website.

In each chapter, there are concise explanations and worked examples, plus numerous exercises that will help you build up confidence.

There are also exam practice questions and a chapter summary to help with exam preparation. Answers to all exercises are included at the back of the book as well as a glossary of Maths-specific terminology.

Points of Interest put the maths you are about to learn in a real-world context.

Learning Objectives show what you will learn in each chapter.

Hint boxes give you tips and reminders.

THE QUADRATIC FUNCTION

CHAPTER 2 25

2 THE QUADRATIC FUNCTION

The path followed by a bottlenose dolphin jumping out of the water is called a parabola. A parabola is a **visual realisation** of the quadratic function $y = -x^2 + k$. Using this formula, scientists can calculate the height of a dolphin's jump (on the y -axis) and the distance travelled (on the x -axis).

There is no scientific agreement about why dolphins jump. Some scientists believe it is because they are trying to conserve energy, some believe it is to help them find food, and others believe they do it just for fun.

The parabola is a beautiful and elegant shape, commonly seen in nature. It is also seen in many man-made structures such as bridges and buildings.



LEARNING OBJECTIVES

- Factorise quadratic expressions where the coefficient of x^2 is greater than 1
- Complete the square and use this to solve quadratic equations
- Solve quadratic equations using the quadratic formula
- Understand and use the discriminant to identify whether the roots are (i) equal and real, (ii) unequal and real or (iii) not real
- Understand the roots α and β and know how to use them

STARTER ACTIVITIES

- Factorise
 - $6x^2 + 9x$
 - $2b^2 + 8b$
 - $9qm^2 - 27m$
 - $9xy^2 + 36x^2y$
 - $24x - 64x^2$
- Factorise
 - $x^2 + 9x + 18$
 - $x^2 - 7x + 12$
 - $x^2 - 2x - 3$
 - $x^2 + 15x + 36$
 - $x^2 + 12x + 27$
- Factorise
 - $x^2 - 9$
 - $x^2 - 25$
 - $9x^2 - 16$
 - $25x^2 - 16$

HINT

All the parts in question 3 are examples of the difference of two squares.

Examples provide a clear, instructional framework. The blue highlighted text gives further explanation of the method.

Language is graded for speakers of English as an additional language (EAL), with advanced Maths-specific terminology highlighted and defined in the glossary at the back of the book.

Key Points boxes summarise the essentials.

30 CHAPTER 2 THE QUADRATIC FUNCTION

a $2x^2 - 3x + 5 = 0$
 $a = 2, b = -3, c = 5$
 $b^2 - 4ac$
 $(-3)^2 - 4 \times 2 \times 5 = -31$
Therefore there are no real roots.

b $3x^2 - x - 1 = 0$
 $a = 3, b = -1, c = -1$
 $b^2 - 4ac = (-1)^2 - 4 \times 3 \times (-1) = 13$
Therefore there are two unequal roots.
 $x = \frac{-(-1) \pm \sqrt{13}}{2 \times 3}$ so $x = 0.768, -0.434$

c $4x^2 - 12x + 9 = 0$
 $a = 4, b = -12, c = 9$
 $(-12)^2 - 4 \times 4 \times 9 = 0$
Therefore the roots are real and equal.
 $x = \frac{-(-12)}{2 \times 4} = \frac{12}{8} = \frac{3}{2}$

EXAMPLE 7 The equation $kx^2 - 2x - 8 = 0$ has two real roots. What can you deduce about the value of the constant k ?

Since the equation has two real roots, you know that the discriminant $b^2 - 4ac$ must be greater than zero.
You substitute $a = k, b = -2$ and $c = -8$ into the inequality $b^2 - 4ac > 0$, giving
 $(-2)^2 - 4 \times k \times -8 > 0$
 $4 + 32k > 0$
 $k > -\frac{4}{32}$
 $k > -\frac{1}{8}$

EXERCISE 4 Use the discriminant to determine whether these equations have one root, two roots or no roots.

a $x^2 - 2x + 1 = 0$	b $x^2 - 3x - 2 = 0$	c $2x^2 - 3x - 4 = 0$
d $2x^2 - 4x + 5 = 0$	e $2x^2 - 4x + 2 = 0$	f $2x^2 - 7x + 3 = 0$
g $3x^2 - 6x + 5 = 0$	h $7x^2 - 144x + 57 = 0$	i $16x^2 - 2x + 3 = 0$
j $x^2 + 22x + 121 = 0$	k $5x^2 - 4x + 81 = 0$	

THE QUADRATIC FUNCTION CHAPTER 2 31

2 The equation $px^2 - 2x - 7 = 0$ has two real roots. What can you deduce about the value of p ?

3 The equation $3x^2 + 2x + m = 0$ has equal roots. Find the value of m .

UNDERSTAND THE ROOTS α AND β AND HOW TO USE THEM

If α and β are the roots of the equation $ax^2 + bx + c = 0$ then you deduce that $(x - \alpha)(x - \beta) = 0$
You can rewrite this as $x^2 - x(\alpha + \beta) + \alpha\beta = 0$
Comparing this with $ax^2 + bx + c = 0$ you can see that
 $a + b = -\frac{b}{a}$ and $a\alpha\beta = \frac{c}{a}$

KEY POINTS For the equation $ax^2 + bx + c = 0$

- The sum of roots, $\alpha + \beta = -\frac{b}{a}$
- The product of the roots, $\alpha\beta = \frac{c}{a}$

EXAMPLE 8 The roots of the equation $3x^2 + x - 6 = 0$ are α and β .

SKILLS Find an expression for $\alpha + \beta$ and an expression for $\alpha\beta$. Hence find an expression for $\alpha^2 + \beta^2$ and an expression for $\alpha^2\beta^2$. Find a quadratic equation with roots α^2 and β^2 .

a $3x^2 + x - 6 = 0$
 $x^2 + \frac{1}{3}x - 2 = 0$
Therefore sum of the roots $\alpha + \beta = -\frac{1}{3}$

Divide the equation by 3 to obtain an equation where the coefficient of x^2 is 1.
The sum of roots $\alpha + \beta = -\frac{b}{a}$

THE PRODUCT OF THE ROOTS $\alpha\beta = \frac{c}{a}$

NOTE Sometimes you will need to manipulate the expressions to help you solve the questions.

Product of the roots $\alpha\beta = -2$

b $\alpha^2 + \beta^2 = \alpha^2 + 2\alpha\beta + \beta^2$
Therefore $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$
Substituting the results from part a, gives
 $\alpha^2 + \beta^2 = \left(-\frac{1}{3}\right)^2 - 2(-2)$
 $= \frac{37}{9}$
 $\alpha^2 + \beta^2 = \alpha\beta^2 = (-2)^2 = 4$

34 CHAPTER 2 EXAM PRACTICE

EXAM PRACTICE: CHAPTER 2

1 $f(x) = 0 = 3x^2 - 10x - 2$

a Without solving the equation $f(x)=0$, form an equation, with integer coefficients which has:
i roots $\frac{2}{3}$ and $\frac{3}{2}$ [6]
ii roots $2\alpha + \beta$ and $\alpha = 2\beta$ [4]

b Solve $f(x) = 0$ using completing the square. [4]

2 The roots of a quadratic equation are α and β where $\alpha + \beta = -\frac{9}{5}$ and $\alpha\beta = -3$. Find a quadratic equation, with integer coefficients, which has roots α and β . [4]

3 Given that $\alpha + \beta = 7$ and $\alpha^2 + \beta^2 = 25$

a Show that $\alpha\beta = 12$. [2]
b Hence, or otherwise, form a quadratic equation with the integer coefficients, which has roots α and β . [3]
c Form a quadratic equation, with integer coefficients, which has roots $\frac{\alpha}{\beta}$ and $\frac{\beta}{\alpha}$. [5]

4 The equation $x^2 + (p - 3)x + (3 - 2p) = 0$, where p is a constant, has two distinct real roots.

a Show that p satisfies $p^2 + 2p - 3 > 0$ [1]
b Find the possible values of p . [2]

5 a Show that $x^2 + 6x + 11$ can be written as $(x + a)^2 + b$. [2]
b Find the value of the discriminant. [2]

6 Factorise completely

a $5x^2 + 16x + 3$ [3]
b $3x^2 - 7x + 4$ [3]

7 Solve these equations by completing the square.

a $p^2 + 3p + 2 = 0$ [3]
b $3x^2 + 13x - 10 = 0$ [3]

8 Solve these equations by using the quadratic formula.

a $5x^2 + 3x - 1 = 0$ [2]
b $12x - 5x^2 = 7$ [3]

9 $4x - 5 - x^2 = b - (x + a)^2$ where a and b are integers.

a Find the value of a and b . [2]
b Calculate the discriminant. [3]

10 Solve $\frac{4}{2x+1} - 3 = \frac{1}{4x^2-1}$ [3]

CHAPTER SUMMARY CHAPTER 2 35

CHAPTER SUMMARY: CHAPTER 2

$x^2 - y^2 = (x - y)(x + y)$ is known as the difference of two squares.

Quadratic equations can be solved by

- factorisation
- completing the square: $x^2 + bx = \left(x + \frac{b}{2}\right)^2 - \left(\frac{b}{2}\right)^2$
- using the quadratic formula: $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

The discriminant of a quadratic expression is $b^2 - 4ac$

If α and β are the roots of the equation $ax^2 + bx + c = 0$ then

- $\alpha + \beta = -\frac{b}{a}$
- $\alpha\beta = \frac{c}{a}$

Exam Practice tests cover the whole chapter and provide quick, effective feedback on your progress.

Chapter Summaries state the most important points of each chapter.

ASSESSMENT OVERVIEW

The following tables give an overview of the assessment for the Edexcel International GCSE in Further Pure Mathematics.

We recommend that you study this information closely to help ensure that you are fully prepared for this course and know exactly what to expect in the assessment.

PAPER 1	PERCENTAGE	MARK	TIME	AVAILABILITY
Written examination paper Paper code 4PM1/01C Externally set and assessed by Edexcel	50%	100	2 hours	January and June examination series First assessment June 2019
PAPER 2	PERCENTAGE	MARK	TIME	AVAILABILITY
Written examination paper Paper code 4PM1/02 Externally set and assessed by Edexcel	50%	100	2 hours	January and June examination series First assessment June 2019

CONTENT SUMMARY

- Number
- Algebra and calculus
- Geometry and calculus

ASSESSMENT

- Each paper will consist of around 11 questions with varying mark allocations per questions, which will be stated on the paper
- Each paper will contain questions from any part of the specification content, and the solution of any questions may require knowledge of more than one section of the specification content
- A formulae sheet will be included in the written examinations
- A calculator may be used in the examinations

ASSESSMENT OBJECTIVES AND WEIGHTINGS

ASSESSMENT OBJECTIVE	DESCRIPTION	% IN INTERNATIONAL GCSE
AO1	Demonstrate a confident knowledge of the techniques of pure mathematics required in the specification	30%–40%
AO2	Apply a knowledge of mathematics to the solutions of problems for which an immediate method of solution is not available and which may involve knowledge of more than one topic in the specification	20%–30%
AO3	Write clear and accurate mathematical solutions	35%–50%

RELATIONSHIP OF ASSESSMENT OBJECTIVES TO UNITS

UNIT NUMBER	ASSESSMENT OBJECTIVE		
	AO1	AO2	AO3
Paper 1	15%–20%	10%–15%	17.5%–25%
Paper 2	15%–20%	10%–15%	17.5%–25%
Total for International GCSE	30%–40%	20%–30%	35%–50%

ASSESSMENT SUMMARY

The Edexcel International GCSE in Further Pure Mathematics requires students to demonstrate application and understanding of the following topics.

Number

- Use numerical skills in a purely mathematical way and in real-life situations.

Algebra and calculus

- Use algebra and calculus to set up and solve problems.
- Develop competence and confidence when manipulating mathematical expressions.
- Construct and use graphs in a range of situations.

Geometry and trigonometry

- Understand the properties of shapes, angles and transformations.
- Use vectors and rates of change to model situations.
- Use coordinate geometry.
- Use trigonometry.

Students will be expected to have a thorough knowledge of the content common to the Pearson Edexcel International GCSE in Mathematics (Specification A) (Higher Tier) or Pearson Edexcel International GCSE in Mathematics (Specification B).

Questions may be set which assumes knowledge of some topics covered in these specifications, however knowledge of statistics and matrices will not be required.

Students will be expected to carry out arithmetic and algebraic manipulation, such as being able to change the subject of a formula and evaluate numerically the value of any variable in a formula, given the values of the other variables.

The use and notation of set theory will be adopted where appropriate.

CALCULATORS

Students will be expected to have access to a suitable electronic calculator for all examination papers. The electronic calculator should have these functions as a minimum:

$+$, $-$, \times , \div , π , x^2 , \sqrt{x} , $\frac{1}{x}$, x^y , $\ln x$, e^x , sine, cosine and tangent and their inverses in degrees and decimals of a degree or radians.

Prohibitions

Calculators with any of the following facilities are prohibited in all examinations:

- databanks
- retrieval of text or formulae
- QWERTY keyboards
- built-in symbolic algebra manipulations
- symbolic differentiation or integration.

FORMULAE SHEET

These formulae will be provided for you during the examination.

MENSURATION

Surface area of sphere $= 4\pi r^2$

Curved surface area of cone $= \pi r \times \text{slant height}$

Volume of sphere $= \frac{4}{3}\pi r^3$

SERIES

Arithmetic series

Sum to n terms $S_n = \frac{n}{2}[2a + (n-1)d]$

Geometric series

Sum to n terms, $S_n = \frac{a(1-r^n)}{(1-r)}$

Sum to infinity, $S_\infty = \frac{a}{1-r}$ $|r| < 1$

Binomial series

$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{r!}x^r + \dots$ for $|x| < 1, n \in \mathbb{Q}$

CALCULUS

Quotient rule (differentiation)

$$\frac{d}{dx}\left(\frac{f(x)}{g(x)}\right) = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2}$$

TRIGONOMETRY

Cosine rule

In triangle ABC : $a^2 = b^2 + c^2 - 2bc \cos A$

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\sin(A+B) = \sin A \cos B + \cos A \sin B$$

$$\cos(A+B) = \cos A \cos B - \sin A \sin B$$

$$\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\sin(A-B) = \sin A \cos B - \cos A \sin B$$

$$\cos(A-B) = \cos A \cos B + \sin A \sin B$$

$$\tan(A-B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

LOGARITHMS

$$\log_a x = \frac{\log_b x}{\log_b a}$$

FORMULAE TO KNOW

The following are formulae that you are expected to know and remember during the examination. These formulae **will not** be provided for you. Note that this list is not exhaustive.

LOGARITHMIC FUNCTIONS AND INDICES

$$\log_a xy = \log_a x + \log_a y$$

$$\log_a \frac{x}{y} = \log_a x - \log_a y$$

$$\log_a x^k = k \log_a x$$

$$\log_a \frac{1}{x} = -\log_a x$$

$$\log_a a = 1$$

$$\log_a 1 = 0$$

$$\log_a b = \frac{1}{\log_b a}$$

QUADRATIC EQUATIONS

$$ax^2 + bx + c = 0 \text{ has roots given by } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

When the roots of $ax^2 + bx + c = 0$ are α and β then $\alpha + \beta = -\frac{b}{a}$ and $\alpha\beta = \frac{c}{a}$ and the equation can be written $x^2 - (\alpha + \beta)x + \alpha\beta = 0$

SERIES

Arithmetic series: n th term $= l = a + (n - 1)d$

Geometric series: n th term $= ar^{n-1}$

COORDINATE GEOMETRY

The gradient of the line joining two points (x_1, y_1) and (x_2, y_2) is $\frac{y_2 - y_1}{x_2 - x_1}$

The distance d between two points (x_1, y_1) and (x_2, y_2) is given by $d^2 = (x_1 - x_2)^2 + (y_1 - y_2)^2$

The coordinates of the point dividing the line joining (x_1, y_1) and (x_2, y_2) in the ratio $m : n$ are

$$\left(\frac{nx_1 + mx_2}{m + n}, \frac{ny_1 + my_2}{m + n} \right)$$

CALCULUS

Differentiation:	function	derivative
	x^n	nx^{n-1}
	$\sin ax$	$a \cos ax$
	$\cos ax$	$-a \sin ax$
	e^{ax}	ae^{ax}
	$f(x)g(x)$	$f'(x)g(x) + f(x)g'(x)$
	$f(g(x))$	$f'(g(x))g'(x)$

Integration:	function	derivative
	x^n	$\frac{1}{n+1}x^{n+1} + c \quad n \neq -1$
	$\sin ax$	$-\frac{1}{a}\cos ax + c$
	$\cos ax$	$\frac{1}{a}\sin ax + c$
	e^{ax}	$\frac{1}{a}e^{ax} + c$

AREA AND VOLUME

Area between a curve and the x -axis $= \int_a^b y dx, y \geq 0$

$$\left| \int_a^b y dx \right|, y < 0$$

Area between a curve and the y -axis $= \int_c^d x dy, x \geq 0$

$$\left| \int_c^d x dy \right|, x < 0$$

Area between $g(x)$ and $f(x) = \int_a^b |g(x) - f(x)| dx$

Volume of revolution $= \int_a^b \pi y^2 dx$ or $\int_c^d \pi x^2 dy$

TRIGONOMETRY

Radian measure: length of arc $= r\theta$
 area of sector $= \frac{1}{2}r^2\theta$

In a triangle ABC : $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$

$$\cos^2 \theta + \sin^2 \theta = 1$$

$$\text{area of a triangle} = \frac{1}{2}ab \sin C$$

CHAPTER 1



1 SURDS AND LOGARITHMIC FUNCTIONS

The Richter scale, which describes the energy released by an earthquake, uses the base 10 logarithm as its unit. An earthquake of **magnitude 9** is 10 times as powerful as one of magnitude 8, and 100 000 times as powerful as one of magnitude 4.

The **devastating** 2004 earthquake in the Indian Ocean had a magnitude of 9. Thankfully, such events are rare. The most common earthquakes, which occur over 100 000 times a year, are magnitude 2 to 3, so humans can hardly feel them.



LEARNING OBJECTIVES

- Write a number exactly using surds
- Rationalise the denominator of a surd
- Be familiar with the **functions** a^x and $\log_b x$ and recognise the shapes of their graphs
- Be familiar with functions including e^x and similar terms, and use them in graphs
- Use graphs of functions to solve equations
- Rewrite expressions including powers using logarithms instead
- Understand and use the laws of logarithms
- Change the base of a logarithm
- Solve equations of the form $a^x = b$

STARTER ACTIVITIES

1 ► Simplify

a $y^6 \times y^5$

d $(x^2)^4$

b $2q^3 \times 4q^4$

e $(a^4)^2 \div a^3$

c $3k^2 \times 3k^7 \times 3k^{-3}$

f $64x^4y^6 \div 4xy^2$

2 ► Simplify

a $(m^3)^{\frac{1}{2}}$

d $6b^{\frac{1}{2}} \times 3b^{-\frac{1}{2}}$

b $3p^{\frac{1}{2}} \times p^3$

e $27p^{\frac{2}{3}} \div 9p^{\frac{1}{6}}$

c $28c^{\frac{2}{3}} \div 7c^{\frac{1}{3}}$

f $5y^6 \times 3y^{-7}$

3 ► Evaluate

a $16^{\frac{1}{2}}$

e $\left(\frac{6}{7}\right)^0$

b $125^{\frac{1}{3}}$

f 81^{-4}

c 8^{-2}

g $\left(\frac{9}{16}\right)^{-\frac{3}{2}}$

d $(-2)^{-3}$

WRITE A NUMBER EXACTLY USING A SURD

A surd is a number that cannot be simplified to remove a square root (or a cube root, fourth root etc). Surds are irrational numbers.

NUMBER	DECIMAL	IS IT A SURD?
$\sqrt{1}$	1	No
$\sqrt{2}$	1.414213...	Yes
$\sqrt{4}$	2	No
$\sqrt{\frac{1}{4}}$	0.5	No
$\sqrt{\frac{2}{3}}$	0.816496...	Yes

HINT

An irrational number is a number that cannot be expressed as a fraction, for example π .

You can **manipulate** surds using these rules:

$$\sqrt{ab} = \sqrt{a} \times \sqrt{b}$$

$$\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$$

EXAMPLE 1

SKILLS

CRITICAL THINKING

Simplify

a $\sqrt{12}$

b $\frac{\sqrt{20}}{2}$

c $5\sqrt{6} - 2\sqrt{24} + \sqrt{294}$

a $\sqrt{12}$

$$= \sqrt{4 \times 3}$$

$$= \sqrt{4} \times \sqrt{3}$$

$$= 2\sqrt{3}$$

Use the rule $\sqrt{ab} = \sqrt{a} \times \sqrt{b}$

$$\sqrt{4} = 2$$

b $\frac{\sqrt{20}}{2}$

$$= \frac{\sqrt{4 \times 5}}{2}$$

$$= \frac{2 \times \sqrt{5}}{2}$$

$$= \sqrt{5}$$

$$\sqrt{20} = \sqrt{4} \times \sqrt{5}$$

$$\sqrt{4} = 2$$

c $5\sqrt{6} - 2\sqrt{24} + \sqrt{294}$

$$= 5\sqrt{6} - 2\sqrt{6}\sqrt{4} + \sqrt{6}\sqrt{49}$$

$$= \sqrt{6}(5 - 2\sqrt{4} + \sqrt{49})$$

$$= \sqrt{6}(5 - 2 \times 2 + 7)$$

$$= 8\sqrt{6}$$

$\sqrt{6}$ is a common factor

Work out the square roots $\sqrt{4}$ and $\sqrt{49}$

$$5 - 4 + 7 = 8$$

EXERCISE 1

SKILLS

CRITICAL THINKING

- 1 ► Simplify without using a calculator

a $\sqrt{18}$

b $\sqrt{50}$

c $\sqrt{125}$

d $\sqrt{128}$

e $\sqrt{132}$

f $\sqrt{8625}$

- 2 ► Simplify without using a calculator

a $\frac{\sqrt{60}}{2}$

b $\frac{\sqrt{135}}{2}$

c $\frac{\sqrt{128}}{8}$

d $\frac{\sqrt{68}}{4}$

e $\frac{\sqrt{96}}{6}$

- 3 ► Simplify without using a calculator

a $6\sqrt{3} - 2\sqrt{3}$

b $7\sqrt{3} - \sqrt{12} + \sqrt{48}$

c $\sqrt{112} + 2\sqrt{172} - \sqrt{63}$

d $6\sqrt{48} - 3\sqrt{12} + 2\sqrt{27}$

e $3\sqrt{578} - \sqrt{162} + 4\sqrt{32}$

f $2\sqrt{5} \times 3\sqrt{5}$

g $6\sqrt{7} \times 4\sqrt{7}$

h $4\sqrt{8} \times 6\sqrt{8}$

- 4 ► Simplify without using a calculator

a $6(4 - \sqrt{12})$

b $9(6 - 3\sqrt{29})$

c $4(1 + \sqrt{3}) + 3(3 + 2\sqrt{3})$

d $3(\sqrt{2} - \sqrt{7}) - 5(\sqrt{2} + \sqrt{7})$

e $(4 + \sqrt{3})(4 - \sqrt{3})$

f $(2\sqrt{7} - \sqrt{6})(\sqrt{7} - 2\sqrt{6})$

g $(\sqrt{8} + \sqrt{5})(\sqrt{8} - \sqrt{5})$

- 5 ► A garden is $\sqrt{30}$ m long and $\sqrt{8}$ m wide. The garden is covered in grass except for a small rectangular pond which is $\sqrt{2}$ m long and $\sqrt{6}$ m wide.

Express the area of the pond as a percentage of the area of the garden.



- 6 ► Find the value of $2p^2 - 3pq$ when $p = \sqrt{2} + 3$ and $q = \sqrt{2} - 2$

RATIONALISE THE DENOMINATOR OF A SURD

Rationalising the denominator of a surd means removing a root from the denominator of a fraction. You will usually need to rationalise the denominator when you are asked to *simplify* it.

The rules for rationalising the denominator of a surd are:

- For fractions in the form $\frac{1}{\sqrt{a}}$, multiply the numerator and denominator by \sqrt{a}
- For fractions in the form $\frac{1}{a + \sqrt{b}}$, multiply the numerator and denominator by $a - \sqrt{b}$
- For fractions in the form $\frac{1}{a - \sqrt{b}}$, multiply the numerator and denominator by $a + \sqrt{b}$

HINT

In the denominator, the multiplication gives the difference of two squares, with the result $a^2 - b^2$, which means the surd disappears.

EXAMPLE 2

Rationalise the denominator of

a $\frac{1}{\sqrt{3}}$

b $\frac{1}{3 + \sqrt{2}}$

c $\frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} - \sqrt{2}}$

a $\frac{1}{\sqrt{3}}$

$$= \frac{1 \times \sqrt{3}}{\sqrt{3} \times \sqrt{3}}$$

$$= \frac{\sqrt{3}}{3}$$

Multiply the top and bottom by $\sqrt{3}$

$$\sqrt{3} \times \sqrt{3} = (\sqrt{3})^2 = 3$$

b $\frac{1}{3 + \sqrt{2}}$

$$= \frac{1 \times (3 - \sqrt{2})}{(3 + \sqrt{2})(3 - \sqrt{2})}$$

$$= \frac{3 - \sqrt{2}}{9 - 3\sqrt{2} + 3\sqrt{2} - 2}$$

$$= \frac{3 - \sqrt{2}}{7}$$

Multiply the top and bottom by $3 - \sqrt{2}$

$$\sqrt{2} \times \sqrt{2} = 2$$

$$9 - 2 = 7, -3\sqrt{2} + 3\sqrt{2} = 0$$

c $\frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} - \sqrt{2}}$

$$= \frac{(\sqrt{5} + \sqrt{2})(\sqrt{5} + \sqrt{2})}{(\sqrt{5} - \sqrt{2})(\sqrt{5} + \sqrt{2})}$$

$$= \frac{5 + \sqrt{5}\sqrt{2} + \sqrt{2}\sqrt{5} + 2}{5 - 2}$$

$$= \frac{7 + 2\sqrt{10}}{3}$$

Multiply the top and bottom by $\sqrt{5} + \sqrt{2}$

$$\sqrt{5}\sqrt{2} - \sqrt{2}\sqrt{5} = 0 \text{ in the denominator}$$

$$\sqrt{5}\sqrt{2} = \sqrt{10}$$

EXERCISE 2

SKILLS

EXECUTIVE
FUNCTION

1 ► Rationalise

a $\frac{1}{\sqrt{3}}$

b $\frac{1}{\sqrt{7}}$

c $\frac{2}{\sqrt{3}}$

d $\frac{\sqrt{6}}{\sqrt{3}}$

e $\frac{12}{\sqrt{3}}$

f $\frac{3\sqrt{5}}{\sqrt{3}}$

g $\frac{9\sqrt{12}}{2\sqrt{18}}$

h $\frac{1}{2 - \sqrt{3}}$

2 ► Rationalise

a $\frac{\sqrt{6}}{\sqrt{3} + \sqrt{6}}$

b $\frac{2 + \sqrt{3}}{2 - \sqrt{3}}$

c $\frac{\sqrt{2} - \sqrt{3}}{\sqrt{2} + \sqrt{3}}$

d $\frac{4\sqrt{2} - 2\sqrt{3}}{\sqrt{2} + \sqrt{3}}$

e $\frac{\sqrt{2} + 2\sqrt{5}}{\sqrt{5} - \sqrt{2}}$

f $\frac{\sqrt{7} + \sqrt{3}}{\sqrt{7} - \sqrt{3}}$

g $\frac{\sqrt{11} + 2\sqrt{5}}{\sqrt{11} + 3\sqrt{5}}$

h $\frac{2\sqrt{5} - 3\sqrt{7}}{5\sqrt{6} + 4\sqrt{2}}$

i $\frac{2 + \sqrt{10}}{\sqrt{2} + \sqrt{5}}$

j $\frac{ab}{a\sqrt{b} - b\sqrt{a}}$

k $\frac{a - b}{a\sqrt{b} - b\sqrt{a}}$

BE FAMILIAR WITH THE FUNCTIONS a^x AND $\log_a x$ AND RECOGNISE THE SHAPES OF THEIR GRAPHS

You need to be familiar with functions in the form $y = a^x$ where $a > 0$

Look at a table of values for $y = 2^x$

x	-3	-2	-1	0	1	2	3
y	$\frac{1}{8}$	$\frac{1}{4}$	$\frac{1}{2}$	1	2	4	8


Note: $2^0 = 1$

In fact a^0 is always equal to 1 if a is positive and

$$2^{-3} = \frac{1}{2^3} = \frac{1}{8}$$

a negative index turns the number into its **reciprocal**

The graph of $y = 2^x$ looks like this:



Note: the x -axis is an **asymptote** to the curve.

Other graphs of the type $y = a^x$ have similar shapes, always passing through (0, 1).

EXAMPLE 3

SKILLS

ANALYSIS

a On the same axes, **sketch** the graphs of $y = 3^x$, $y = 2^x$ and $y = 1.5^x$

b On another set of axes, sketch the graphs of $y = \left(\frac{1}{2}\right)^x$ and $y = 2^x$


a For all three graphs, $y = 1$ when $x = 0$

When $x > 0$, $3^x > 2^x > 1.5^x$

$$a^0 = 1$$

When $x < 0$, $3^x < 2^x < 1.5^x$

Work out the relative positions of the graphs




b $\frac{1}{2} = 2^{-1}$

so, $y = \left(\frac{1}{2}\right)^x$ is the same as $y = (2^{-1})^x = 2^{-x}$




$$(a^m)^n = a^{mn}$$

Therefore the graph of $y = \left(\frac{1}{2}\right)^x$ is a reflection in the y-axis of the graph of $y = 2^x$




EXAMPLE 4

If you compare the graphs of $y = 2^x$ and $y = \log_2 x$ you see the following relationship:



EXAMPLE 5

On the same set of axes sketch the graphs $y = \log_2 x$ and $y = \log_5 x$



Note:

For both graphs $y = 0$ when $x = 1$, since $\log_a 1 = 0$ for every value of a .

$\log_2 2 = 1$ so $y = \log_2 x$ passes through $(2, 1)$

and $\log_5 5 = 1$ so $y = \log_5 x$ passes through $(5, 1)$

EXERCISE 3

SKILLS

ANALYSIS REASONING

1 ▶ On the same set of axes sketch the graphs of

a $y = 5^x$

b $y = 7^x$

c $y = \left(\frac{1}{3}\right)^x$

- 2 ► On the same set of axes sketch the graphs of
- a $y = \log_5 x$ b $y = \log_7 x$
- c Write down the coordinates of the point of intersection of these two graphs.
- 3 ► On the same set of axes sketch the graphs of
- a $y = 3^x$ b $y = \log_3 x$
- 4 ► On the same set of axes sketch the graphs of
- a $y = \log_3 x$ b $y = \log_5 x$ c $y = \log_{0.5} x$ d $y = \log_{0.25} x$

BE FAMILIAR WITH EXPRESSIONS OF THE TYPE e^x AND USE THEM IN GRAPHS

Consider this example: Zainab opens an account with \$1.00. The account pays 100% interest per year. If the interest is credited once, at the end of the year, her account will contain \$2.00. How much will it contain after a year if the interest is calculated and credited more frequently? Let us investigate this more thoroughly.



HOW OFTEN INTEREST IS CREDITED INTO THE ACCOUNT	VALUE OF ACCOUNT AFTER 1 YEAR (\$)
Yearly	$\left(1 + \frac{1}{1}\right)^1 = 2$
Semi-annually	$\left(1 + \frac{1}{2}\right)^2 = 2.25$
Quarterly	$\left(1 + \frac{1}{4}\right)^4 = 2.441406...$
Monthly	$\left(1 + \frac{1}{12}\right)^{12} = 2.61303529...$
Weekly	$\left(1 + \frac{1}{52}\right)^{52} = 2.69259695...$
Daily	$\left(1 + \frac{1}{365}\right)^{365} = 2.71456748...$
Hourly	$\left(1 + \frac{1}{8760}\right)^{8760} = 2.71812669...$
Every minute	$\left(1 + \frac{1}{525\,600}\right)^{525\,600} = 2.7182154...$
Every second	$\left(1 + \frac{1}{31\,536\,000}\right)^{31\,536\,000} = 2.71828247...$

The amount in her account gets bigger and bigger the more often the interest is compounded, but the rate of growth slows. As the number of compounds increases, the calculated value appears to be approaching a fixed value. This value gets closer and closer to a fixed value of 2.7182847254..... This number is called 'e'.

The number e is called a *natural* exponential because it arises *naturally* in mathematics and has numerous real life applications.




EXAMPLE 6

Draw the graphs of e^x and e^{-x}


SKILLS

ANALYSIS
INTERPRETATION

x	-2	-1	0	1	2	3	4
e^x	0.14	0.37	1	2.7	7.4	20	55



x	-4	-3	-2	-1	0	1	2
e^{-x}	55	20	7.4	2.7	1	0.37	0.14




EXAMPLE 7

Draw the graphs of these **exponential functions**.

- a** $y = e^{2x}$
b $y = 10e^{-x}$
c $y = 3 + 4e^{\frac{1}{2}x}$


a

x	-2	-1	0	1	2
e^{2x}	0.02	0.1	1	7.4	55




b

x	-2	-1	0	1	2
$10e^{-x}$	73	27	10	3.7	1.4



c

x	-2	-1	0	1	2
$3 + 4e^{\frac{1}{2}x}$	4.5	5.4	7	8.9	13.4



On pages 7–9 you saw the connection between $y = \log_a x$ and $y = a^x$. The function $y = \log_e x$ is particularly important in mathematics and so it has a special notation:

$$\log_e x \equiv \ln x$$

Your calculator should have a special button for evaluating $\ln x$.



EXAMPLE 8

Solve these equations.

a $e^x = 3$

b $\ln x = 4$

a When $e^x = 3$

$$x = \ln 3$$

b When $\ln x = 4$

$$x = e^4$$

As you can see, the inverse of e^x is $\ln x$ (and vice versa)

EXAMPLE 9


Sketch these graphs on the same set of axes.

a $y = \ln x$

b $y = \ln(3 - x)$

c $y = 3 + \ln(2x)$

a–c



EXERCISE 4

SKILLS

ANALYSIS
REASONING

1 ▶ Sketch these graphs.

a $y = e^x + 1$

b $y = 4e^{-2x}$

c $y = 2e^x - 3$

d $y = 6 + 10^{\frac{1}{2}x}$

e $y = 100e^{-x} + 10$

2 ▶ Sketch these graphs, stating any asymptotes and intersections with the axes.

a $y = \ln(x + 1)$

b $y = 2 \ln x$

c $y = \ln(2x)$

d $y = \ln(4 - x)$

e $y = 4 + \ln(x + 2)$


EXAMPLE 10

BE ABLE TO USE GRAPHS OF FUNCTIONS TO SOLVE EQUATIONS

- a** Complete the table of values for: $y = e^{\frac{1}{2}x} - 2$
Giving your answers to two decimal places where appropriate.
- b** Draw the graph of $y = e^{\frac{1}{2}x} - 2$ for $0 \leq x \leq 5$
- c** Use your graph to estimate, to 2 significant figures, the solution of the equation $e^{\frac{1}{2}x} = 8$
Show your method clearly.
- d** By drawing a suitable line on your graph, estimate to 2 significant figures the solution to the equation $x = 2 \ln(7 - 2x)$

a

x	-1	0	1	2	3	4	5
y	-1.39	-1	-0.35	0.72	2.48	5.39	10.18



- c** $e^{\frac{1}{2}x} = 8$
 $e^{\frac{1}{2}x} - 2 = 8 - 2$
 $e^{\frac{1}{2}x} = 6$
 So the solution is the intersection of the curve $y = e^{\frac{1}{2}x}$ and $y = 6$
 $x \approx 4.15$ (In the exam you will be allowed a range of values.)

- d** $x = 2 \ln(7 - 2x)$
 $\frac{x}{2} = \ln(7 - 2x)$
 $e^{\frac{1}{2}x} = 7 - 2x$
 $e^{\frac{1}{2}x} - 2 = 7 - 2x - 2$
 $e^{\frac{1}{2}x} = 5 - 2x$

So, the solution is the intersection of the curve and $y = e^{\frac{1}{2}x}$ and the line $y = 5 - 2x$, $x \approx 2.1$

EXAMPLE 11

- a** Complete the table below of values of $y = 2 + \ln x$, giving your values of y to decimal places.

x	0.1	0.5	1	1.5	2	3	4
y	-0.3	1.31		2.41	2.69	3.10	

HINT


Make the
LHS = $e^{\frac{1}{2}x} - 2$
i.e. the equation
of the graph.
To do this, you
need to subtract
2 from 8 and
draw the line
 $y = 6$ (as shown
in the diagram).

HINT


Make the LHS
equal to the
given equation
i.e. $e^{\frac{1}{2}x} - 2$.
Draw the line
 $y = 5 - 2x$ (as
shown in the
diagram) on
your graph and
find points of
intersection.

- b** Draw the graph of $y = 2 + \ln x$ for $0.1 \leq x \leq 4$
- c** Use your graph to estimate, to 2 significant figures, the solution of the equation $\ln x = 0.5$
- d** By drawing a suitable line on your graph estimate, to 1 significant figure, the solution of the equation $x = e^{x-2}$

a	x	0.1	0.5	1	1.5	2	3	4
	y	-0.3	1.31	2	2.41	2.69	3.10	3.39




- c** $2 + \ln x = 0.5$ Make the LHS = $2 + \ln x$ i.e. the equation of the graph.
Therefore you need to add
 $\ln x = 2.5 + 2$ from 0.5 and draw the line $y = 2.5$
 $y = 2.5$



The solution is the intersection of the curve and the line $y = 2.5$. From the graph this is approximately 1.6. In the exam you will be given a small range of answers.

- d** $x = e^{x-2}$ Using the properties of logs
 $\ln x = x - 2$
 $\ln x + 2 = x - 2 + 2$ Make the LHS equal to the given equation i.e. $\ln x + 2$.



EXERCISE 5

- 1 ▶ a Draw the graph $y = 3 + 2e^{-\frac{1}{2}x}$ for $0 \leq x \leq 6$
- b Use your graph to estimate, to 2 significant figures, the solution to the equation $e^{-\frac{1}{2}x} = 0.5$, showing your method clearly.
- c By drawing a suitable line, estimate, to 2 significant figures, the solution of the equation $x = -2 \ln\left(\frac{x-2}{2}\right)$
- 2 ▶ a Draw the graph $y = 2 + \frac{1}{3}e^x$ for $-1 \leq x \leq 3$
- b Use your graph to estimate, to 2 significant figures, the solution to the equation $e^x = 12$ showing your method clearly.
- c By drawing a suitable line, estimate, to 2 significant figures, the solution to the equation $x = \ln(6 - 6x)$
- 3 ▶ a Complete the table below of values of $y = 5 \sin 2x - 2 \cos x$, giving your values of y to 2 decimal places.

x	0	15	30	45	60	75	90
y	-2	0.57		3.59	3.33		0

- b Draw the graph of $y = 5 \sin 2x - 2 \cos x$ for $0 \leq x \leq 90^\circ$
- c Use your graph to estimate, to 2 significant figures, the solution of the equation $2(1 + \cos x) = 5 \sin 2x$ showing your method clearly.

HINT

In the exam \log_{10} will be written as **lg**. This textbook uses **lg** for \log_{10}

EXAMPLE 12

SKILLS

ADAPTABILITY

WRITING AN EXPRESSION AS A LOGARITHM

$\log_a n = x$ means that $a^x = n$, where a is called the **base of the logarithm**.

- $\log_a 1 = 0$ ($a > 0$), **because** $a^0 = 1$
- $\log_a a = 1$ ($a > 0$), **because** $a^1 = a$

Write as a logarithm $2^5 = 32$

$$2^5 = 32$$

$$\text{So } \log_2 32 = 5$$

$$\text{Here } a = 2, x = 5, n = 32$$

Here 2 is the base, 5 is the logarithm. In words, you would say '2 to the power of 5 equals 32'. You would also say 'the logarithm of 32, to base 2, is 5'.

EXAMPLE 13

Rewrite using a logarithm

a $10^3 = 1000$

b $5^4 = 625$

c $2^{10} = 1024$

a $\lg 1000 = 3$

b $\log_5 625 = 4$

c $\log_2 1024 = 10$

EXAMPLE 14

Find the value of

a $\log_3 81$

b $\log_4 0.25$

c $\log_{0.5} 4$

d $\log_a (a^5)$

a $\log_3 81 = 4$

$3^4 = 81$

b $\log_4 0.25 = -1$

$4^{-1} = \frac{1}{4} = 0.25$

c $\log_{0.5} 4 = -2$

$0.5^{-2} = \left(\frac{1}{2}\right)^{-2} = 2^2 = 4$

d $\log_a (a^5) = 5$

$a^5 = a^5$

EXERCISE 6

SKILLS

CRITICAL THINKING

1 ▶ Rewrite these exponentials as logarithms.

a $4^3 = 64$

b $5^{-2} = \frac{1}{25}$

c $8^6 = 262\,144$

d $3^x = 9$

e $8^x = 1$

f $2^x = \frac{1}{4}$

2 ▶ Write these logarithms in exponential form.

a $\log_3 81 = 4$

b $\log_3 729 = 6$

c $\log_5 625 = 4$

d $\log_6 4 = \frac{1}{2}$

e $\log_3 \left(\frac{1}{27}\right) = -3$

f $\lg 1000 \cdot 0.01 = -2$

3 ▶ Without a calculator find the value of

a $\log_2 4$

b $\log_3 27$

c $\log_3 81$

d $\log_5 625$

e $\log_5 \left(\frac{1}{125}\right)$

f $\lg \sqrt{10}$

g $\log_3 \sqrt{27}$

h $\log_3 \sqrt[5]{3}$

4 ▶ Find the value of x for which

a $\log_3 x = 4$

b $\log_6 x = 3$

c $\log_x 64 = 3$

d $\log_x 16 = \frac{4}{3}$

e $\log_x 64 = \frac{2}{3}$

5 ▶ Find using your calculator

a $\lg 20$

b $\lg 14$

c $\lg 0.25$

d $\lg 0.3$

e $\lg 54.6$

UNDERSTAND AND USE THE LAWS OF LOGARITHMS

$2^5 = 32$ and $\log_2 32 = 5$

The rules of logarithms follow the **rules of indices**.

EXPONENT (POWERS)	LOGARITHMS	LAW
$c^x \times c^y = c^{x+y}$	$\log_c xy = \log_c x + \log_c y$	Multiplication Law
$c^x \div c^y = c^{x-y}$	$\log_c \frac{x}{y} = \log_c x - \log_c y$	Division Law
$(c)^q$	$\log_c (x^q) = q \log_c x$	Power Law
$\frac{1}{c} = c^{-1}$	$\log_c \left(\frac{1}{x}\right) = -\log_c x$	
$c^1 = c$	$\log_c (c) = 1$	
$c^0 = 1$	$\log_c (1) = 0$	

EXAMPLE 15

Write as a single logarithm

SKILLS

DECISION
MAKING

$$\text{a } \log_3 6 + \log_3 7 \quad \text{b } \log_2 15 - \log_2 3 \quad \text{c } 2 \log_5 3 + 3 \log_5 2 \quad \text{d } \lg 3 - 4 \lg \left(\frac{1}{2}\right)$$

$$\begin{aligned} \text{a } \log_3 (6 \times 7) \\ = \log_3 (42) \end{aligned}$$

Use the multiplication law

$$\begin{aligned} \text{b } \log_2 (15 \div 3) \\ = \log_2 5 \end{aligned}$$

Use the division law

$$\begin{aligned} \text{c } 2 \log_5 3 &= +3 \log_5 2 \\ &= \log_5 (3^2) = +\log_5 (2^3) \\ &= \log_5 9 + \log_5 8 \\ &= \log_5 72 \end{aligned}$$

Apply the power law to both expressions

Use the multiplication law

$$\begin{aligned} \text{d } \lg 3 - 4 \lg \left(\frac{1}{2}\right) \\ = \lg 3 - \lg \left(\frac{1}{2}\right)^4 \\ = \lg \left(3 \div \frac{1}{16}\right) \\ = \lg 48 \end{aligned}$$

Use the power law

Use the division law

EXAMPLE 16

Find the value of

Write in terms of $\log_a x$, $\log_a y$ and $\log_a z$

$$\text{a } \log_a (x^2 y z^3) \quad \text{b } \log_a \left(\frac{x}{y^3}\right) \quad \text{c } \log_a \left(\frac{x\sqrt{y}}{z}\right) \quad \text{d } \log_a \left(\frac{x}{a^4}\right)$$

$$\begin{aligned} \text{a } \log_a (x^2 y z^3) \\ = \log_a (x^2) + \log_a (y) + \log_a (z^3) \\ = 2 \log_a (x) + \log_a (y) + 3 \log_a (z) \end{aligned}$$

$$\begin{aligned} \text{b } \log_a \left(\frac{x}{y^3}\right) \\ = \log_a (x) - \log_a (y^3) \\ = \log_a (x) - 3 \log_a (y) \end{aligned}$$

$$\begin{aligned} \text{c } \log_a \left(\frac{x\sqrt{y}}{z}\right) \\ = \log_a (x\sqrt{y}) - \log_a (z) \\ = \log_a (x) + \log_a (\sqrt{y}) - \log_a (z) \\ = \log_a (x) + \frac{1}{2} \log_a (y) - \log_a (z) \end{aligned}$$

Use the power law $\sqrt{y} = y^{\frac{1}{2}}$

$$\begin{aligned} \text{d } \log_a \left(\frac{x}{a^4}\right) \\ = \log_a (x) - \log_a (a^4) \end{aligned}$$

$$= \log_a(x) - 4 \log_a(a)$$

$$= \log_a(x) - 4$$

$$\log_a a = 1$$

EXERCISE 7

SKILLS

CRITICAL THINKING

1 Write as a single logarithm

a $\log_4 8 + \log_4 8$

c $\log_5 27 + \log_5 3$

e $2 \log_6 9 - 10 \log_6 81$

g $\log_8 25 + \log_8 10 - 3 \log_8 5$

i $2 \lg 20 - (\lg 5 + \lg 8)$

b $\log_9 3 + \log_9 2$

d $\log_4 24 + \log_4 15 - \log_5 3$

f $\frac{1}{2} \log_2 25 + 2 \log_2 3$

h $2 \log_{12} 3 + 4 \log_{12} 2$

2 Write in terms of $\log_a x$, $\log_a y$, $\log_a z$

a $\log_a x^4 y^3 z$

b $\log_a \frac{x^6}{y^3}$

c $\log_a ((xz)^2)$

d $\log_a \frac{1}{xyz}$

e $\log_a \sqrt{xy}$

f $\log_a \sqrt{x^4 y^2 z^3}$

g $\log_a \frac{\sqrt{x^3 y^7}}{z^3}$

CHANGE THE BASE OF A LOGARITHM

Working in base a , suppose that

$$\log_a x = m$$

Writing this as a power

$$a^m = x$$

Taking logs to a different base b

$$\log_b(a^m) = \log_b(x)$$

Using the power law

$$m \log_b a = \log_b x$$

Writing m as $\log_a x$

$$\log_b x = \log_a x \times \log_b a$$

This can be written as

$$\log_a x = \frac{\log_b x}{\log_b a}$$

Using this rule, notice in particular that $\log_a b = \frac{\log_b b}{\log_b a}$,

$$\text{but } \log_b b = 1$$

$$\text{so, } \log_a b = \frac{1}{\log_b a}$$

EXAMPLE 17

SKILLS

EXECUTIVE FUNCTION

Find, to 3 significant figures, the value of $\log_8(11)$

One method is to use the change of base rule

$$\begin{aligned} \log_8 11 &= \frac{\lg 11}{\lg 8} \\ &= 1.15 \end{aligned}$$

Another method is to solve $8^x = 11$

$$\text{Let } x = \log_8(11)$$

$$8^x = 11$$

$$\lg(8^x) = \lg 11$$

$$x \lg 8 = \lg 11$$

$$x = \frac{\lg 11}{\lg 8}$$

$$x = 1.15 \text{ (3 s.f.)}$$

Take logs to base 10 of each side

Use the power law

Divide by $\lg 8$ **EXAMPLE 18**Solve the equation $\log_5 x + 6\log_x 5 = 5$

$$\log_5 x + \frac{6}{\log_5 x} = 5$$

Use change of base rule, special case

$$\text{Let } \log_5(x) = y$$

$$y + \frac{6}{y} = 5$$

$$y^2 + 6 = 5y$$

Multiply by y

$$y^2 - 5y + 6 = 0$$

$$(y - 3)(y - 2) = 0$$

$$\text{So } y = 3 \text{ or } y = 2$$

$$\log_5 x = 3 \text{ or } \log_5 x = 2$$

$$x = 5^3 \text{ or } x = 5^2$$

Write as powers

$$x = 125 \text{ or } x = 25$$

EXERCISE 8**SKILLS****EXECUTIVE
FUNCTION****1** Find, to 3 significant figures

a $\log_8 785$

b $\log_5 15$

c $\log_6 32$

d $\log_{12} 4$

e $\log_{15} \frac{1}{7}$

2 Solve, giving your answer to 3 significant figures

a $6^x = 15$

b $9^x = 751$

c $15^x = 3$

d $3^x = 17.3$

e $3^{2x} = 25$

f $4^{3x} = 64$

g $7^{3x} = 152$

3 Solve, giving your answer to 3 significant figures

a $\log_2 x = 8 + 9\log_x 2$

b $\log_6 x + 3\log_x 6 = 4$

c $\lg x + 5\log_x 10 = -6$

d $\log_2 x + \log_4 x = 2$

HINT

If no base is given in a question, you should assume base 10.

SOLVE EQUATIONS OF THE FORM $a^x = b$ You need to be able to solve equations of the form $a^x = b$ **EXAMPLE 19**Solve the equation $3^x = 20$, giving your answer to 3 significant figures.

$$3^x = 20$$

$$\lg(3^x) = \lg 20$$

Take logs to base 10 on each side

SKILLS**PROBLEM
SOLVING
ANALYSIS**

$$x \lg 3 = \lg 20$$

Use the power law

$$x = \frac{\lg 20}{\lg 3}$$

Divide by $\lg 3$

$$x = \frac{1.3010...}{0.4771...}$$

Use your calculator for logs to base 10

$$= 2.73 \text{ (3 s.f.)}$$

Or, a simpler version

$$3^x = 20$$

$$x = \log_3 20$$

$$x = 2.73$$

EXAMPLE 20Solve the equation $7^{x+1} = 3^{x+2}$

$$(x + 1) \lg 7 = (x + 2) \lg 3$$

Use the power law

$$x \lg 7 + \lg 7 = x \lg 3 + 2 \lg 3$$

Multiply out

$$x \lg 7 - x \lg 3 = 2 \lg 3 - \lg 7$$

Collect x terms on left and numerical terms on right

$$x(\lg 7 - \lg 3) = 2 \lg 3 - \lg 7$$

Factorise

$$x = \frac{2 \lg 3 - \lg 7}{\lg 7 - \lg 3}$$

Divide by $\lg 7 - \lg 3$

$$x = 0.297 \text{ (3 s.f.)}$$

EXAMPLE 21Solve the equation $5^{2x} + 7(5^x) - 30 = 0$, giving your answer to 2 decimal places.

$$\text{Let } y = 5^x$$

$$y^2 + 7y - 30 = 0$$

Use $5^{2x} = (5^x)^2 = y^2$

$$\text{So } (y + 10)(y - 3) = 0$$

$$\text{So } y = -10 \text{ or } y = 3$$

$$\text{If } y = -10, 5^x = -10, \text{ has no solution}$$

 5^x cannot be negative

$$\text{If } y = 3, 5^x = 3$$

$$\lg(5^x) = \lg 3$$

Solve as in previous examples

$$x \lg(5) = \lg 3$$

$$x = \frac{\lg 3}{\lg 5}$$

$$x = 0.683 \text{ (3 s.f.)}$$

EXERCISE 9

1 ► Solve, giving your answer to 3 significant figures

a $4^x = 12$

b $5^x = 20$

c $15^x = 175$

d $7^x = \frac{1}{4}$

e $4^{x+1} = 30$

f $7^{2x+1} = 36$

g $4^{x+1} = 8^{x+1}$

h $2^{3y-2} = 3^{2y+5}$

i $7^{2x+6} = 11^{3x-2}$

j $3^{4-3x} = 4^{x+5}$

2 ► Solve, giving your answer to 3 significant figures

a $4^{2x} + 4^x - 12 = 0$

b $6^{2x} - 10(6^x) + 8 = 0$

c $5^{2x} - 6(5^x) - 7 = 0$

d $4^{2x+1} + 7(4^x) - 15 = 0$

e $2^{2x} + 3^{2x} = 4$

f $3^{2x+1} = 26(3^x) + 9$

EXAM PRACTICE: CHAPTER 1

- 1 Simplify $\sqrt{32} + \sqrt{18}$, giving your answer in the form $p\sqrt{2}$, where p is an integer. [2]
- 2 Simplify $\frac{\sqrt{32} + \sqrt{18}}{3 + \sqrt{2}}$ giving your answer in the form $a\sqrt{2} + b$, where a and b are integers. [3]
- 3
 - a Expand and simplify $(7 + \sqrt{5})(3 - \sqrt{5})$ [2]
 - b Express $\frac{7 + \sqrt{5}}{3 + \sqrt{5}}$ in the form $a + b\sqrt{5}$, where a and b are integers. [2]
- 4 Write $\sqrt{75} - \sqrt{27}$ in the form $k\sqrt{x}$, where k and x are integers. [2]
- 5 A rectangle A has a length of $(1 + \sqrt{5})$ cm and an area of $\sqrt{80}$ cm².
Calculate the width of A in cm, giving your answer in the form $a + b\sqrt{5}$, where a and b are integers to be found. [3]
- 6 Sketch the graph of $y = 8^x$, showing the coordinates of any points at which the graph crosses the axes. [3]
- 7 Solve the equation $8^{2x} - 4(8^x) = 3$, giving your answer to 3 significant figures. [3]
- 8
 - a Given that $y = 6x^2$, Show that $\log_6 y = 1 + 2\log_6 x$ [2]
 - b Hence, or otherwise, solve the equation $1 + 2\log_3 x = \log_3(28x - 9)$, giving your answer to 3 significant figures. [3]
- 9 Find the values of x such that: $2\log_3 x - \log_3(x - 2) - 2 = 0$ [2]
- 10 Find the values of y such that: $\frac{\log_2 32 + \log_2 16}{\log_2 y} - \log_2 y = 0$ [3]
- 11 Given that $\log_b y + 3\log_b 2 = 5$, express y in terms of b in its simplest form. [2]
- 12 Solve $5^{2x} = 12(5^x) - 35$ [4]
- 13 Find, giving your answer to 3 significant figures where appropriate, the value of x for which $5^x = 10$ [3]
- 14 Given that $\log_3(3b + 1) - \log_3(a - 2) = -1$, $a > 2$, express b in terms of a . [2]
- 15 Solve $3^{3x-2} = \sqrt[3]{9}$ [4]
- 16 Solve $25^t + 5^{t+1} = 24$, giving your answer to 3 significant figures. [3]

17 Given that $\log_2 x = p$, find, in terms of p , the simplest form of

a $\log_2(16x)$, [1]

b $\log_2\left(\frac{x^4}{2}\right)$. [1]

c Hence, or otherwise, solve

$$\log_2(16x) - \log_2\left(\frac{x^4}{2}\right) = \frac{1}{2}$$

Give your answer in its simplest surd form. [3]

18 Solve $\log_3 t + \log_3 5 = \log_3(2t + 3)$ [3]

19 **a** Draw the graph $y = 2 + \ln x$ for $0.1 \leq x \leq 4$ [2]

b Use your graph to estimate, to 2 significant figures, the solution to $\ln x = 0.5$, showing clearly your method. [2]

c By drawing a suitable line, estimate to 2 significant figures, the solution of the equation $x = e^{x-2}$ [2]

CHAPTER SUMMARY: CHAPTER 1

- You can simplify expressions by using the power (indices) laws.

$c^x \times c^y = c^{x+y}$
$c^x \div c^y = c^{x-y}$
$(c^p)^q = c^{p \times q}$
$\frac{1}{c} = c^{-1}$
$c^1 = c$
$c^0 = 1$

- You can manipulate surds using these rules:

■ $\sqrt{ab} = \sqrt{a} \times \sqrt{b}$

■ $\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$

- The rules for rationalising surds are:

■ If you have a fraction in the form $\frac{1}{\sqrt{a}}$ then multiply top and bottom by \sqrt{a}

■ If you have a fraction in the form $\frac{1}{1 + \sqrt{a}}$ then multiply top and bottom by $(1 - \sqrt{a})$

■ If you have a fraction in the form $\frac{1}{1 - \sqrt{a}}$ then multiply top and bottom by $(1 + \sqrt{a})$

- $\log_a n = x$ can be rewritten as $a^x = n$ where a is the base of the logarithm.

- The laws of logarithms are:


$\log_a xy = \log_a x + \log_a y$
$\log_a \frac{x}{y} = \log_a x - \log_a y$
$\log_a (x^q) = q \log_a x$
$\log_a \left(\frac{1}{x}\right) = -\log_a x$
$\log_a(a) = 1$
$\log_a(1) = 0$

- The change of base rule for logarithms can be written as $\log_a x = \frac{\log_b x}{\log_b a}$

- From the change of base you can derive $\log_a b = \frac{1}{\log_b a}$

- The natural logarithm is defined as: $\log_e x \equiv \ln x$

- The graph of $y = e^x$ is shown below.



- The graph of $y = \ln x$ is shown below.

