



# Effect of PGPR on enzymatic activities of rice (*Oryza sativa* L.) under salt stress

Sumita Sen<sup>1,2\*</sup> and C.N. Chandrasekhar<sup>1</sup>

<sup>1</sup>Department of Agriculture, Kumarghat Horti. Sub-Division, Unakoti - 799264, Tripura, India.

<sup>2</sup>Department of Crop Physiology, Tamil Nadu Agricultural University, Coimbatore -641003, Tamil Nadu, India.

## Article History

Received 24 March, 2015  
Received in revised form 12  
May, 2015  
Accepted 18 May, 2015

**Keywords:**  
Catalase,  
Nitrate reductase,  
Peroxidase,  
Rice,  
Salt stress.

Article Type:  
Full Length Research Article

## ABSTRACT

**Salt stress makes the agricultural lands unproductive and is an obstacle towards obtaining higher crop yield. To investigate the physiological and biochemical basis of salt stress adaptation in rice plants due to inoculation with plant growth promoting bacteria, two rice genotypes namely ADT43 and IR50 treated with Plant Growth Promoting Rhizobacteria (PGPR *Pseudomonas* strains PF1 and TDK1) were subjected to 100 mM sodium chloride (NaCl), following International Rice Research Institute (IRRI) method in Neyveli Lignite Corporation (NLC) Laboratory of Department of Crop Physiology of Tamil Nadu Agricultural University. Results of our study indicate that enzymatic activities (peroxidase, catalase and nitrate reductase) required to alleviate salt stress were much higher in the plants treated with plant growth promoting rhizobacteria strains PF1 and TDK1 compared to the plants with no external treatment. Among all the treatments ADT43 treated with TDK1 showed the best performance with high enzymatic activities. Phytoremediation through root treatment with PGPR in rice seedlings could lessen the adverse impact of salt stress at later stage of the crop growth and hence increase crop yield.**

©2015 BluePen Journals Ltd. All rights reserved

## INTRODUCTION

Many crop species do not tolerate salt stress due to the accumulation of salts especially sodium chloride (NaCl) which compete with other nutrients and cause specific toxicity (Tester and Davenport, 2003). Salt stress affects adversely growth and development of a non-halophytic crop like rice. Salinity produces oxidative stress by the enhanced occurrence of damaging toxic molecule that is reactive oxygen species (ROS) (Zhu et al., 2007). Evidence suggests that membranes are the primary sites of salinity injury to cells and organelles (Candan and Tarhan, 2003) because ROS can react with unsaturated fatty acids to cause peroxidation of essential membrane lipids in plasmalemma or intracellular organelles (Karabal

et al., 2003). Plant cells produce antioxidant enzymes such as superoxide dismutase (SOD), peroxidase (POX), and catalase (CAT), and non-enzymatic antioxidants such as ascorbate, glutathione, and  $\alpha$ -tocopherol to protect the cells against oxidative stress (Del Rio et al., 2003). Plant growth promoting rhizobacteria (PGPR) are a heterogeneous group of bacteria that can be found in the rhizosphere, at root surfaces and in association with roots. PGPR synthesize hydrolytic enzymes, such as chitinases, glucanases, proteases, and lipases, which can lyse pathogenic fungal cells (Neeraja et al., 2010). Jha and Subramanian (2013) found that inoculation of plant material with PGPR can protect paddy plants against salt stress through an increase in plant growth parameters and the regulation of ion concentration and antioxidant enzymes. The objective of our present study was to study effect of PGPR treatments on enzymatic traits under salt stress condition and to study effect of

\*Corresponding author. E-mail: sumita.sen25@gmail.com. Fax: 03824-281-434.




Figure 1. Effect of salt stress on peroxidase activity in two rice genotypes.

PGPR on improvement rice tolerance to salt stress.

## MATERIALS AND METHODS

The experiment was conducted in the NLC Laboratory, Department of Crop Physiology, Tamil Nadu Agricultural University (TNAU), Coimbatore using Completely Randomized Design (CRD) as the design of experiment. Seeds of two rice genotypes (ADT43 and IR50) were divided into four sets: first set was soaked in water, second set in  $\text{CaCl}_2$  solution at 50 mM (Afzal et al., 2008), third set in PF1 (*Pseudomonas* strain) solution at 2 g per 10 L of water and fourth set in TDK1 (*Pseudomonas* strain) solution at 2 g per 10 L of water. After 12 h the seeds were transferred in sand bed and grown for 7 days. The most uniform seedlings were transferred in Yoshida nutrient solution (Yoshida et al., 1976) following modified IRRI method. At 28 days after seed germination (7 days after 100 mM NaCl treatment), the plant samples were harvested and rinsed with distilled water and kept in refrigerator for physiological and biochemical analysis. The shoots were dried at  $105^\circ\text{C}$  for 2 h and then extracted in triple acid (Nitric acid, sulphuric acid and perchloric acid in the ratio of 9:2:1 respectively).

### Peroxidase activity

Peroxidase activity was assayed according to Perur (1962) and expressed as  $\Delta \text{OD } 403 \text{ nm min}^{-1} \text{ g}^{-1}$ .

### Catalase activity

Catalase activity was determined from the change in the

absorbance at 240 nm over time of the enzyme extract in 3 ml of  $\text{H}_2\text{O}_2$  - $\text{PO}_4$  buffer (Luck, 1974) and expressed as  $\mu\text{g of H}_2\text{O}_2 \text{ min}^{-1} \text{ g}^{-1}$ .

### Nitrate reductase activity (NRase)

Nitrate Reductase (NRase) activity in the leaves was determined by adopting the method of Nicholas et al. (1976) and the enzyme activity was expressed as  $\mu\text{g of NO}_2 \text{ h}^{-1} \text{ g}^{-1}$ .

### Statistical analysis of physiological traits


Using AGRES software, significance of the observed values was determined.  $P \leq 0.05$  was considered statistically significant.

## RESULTS AND DISCUSSION

### Peroxidase and catalase activity

The peroxidase activity followed an increasing trend except for the genotype ADT43 (no external treatment) which showed comparatively less value of peroxidase activity ( $0.0394 \text{ g}^{-1} \text{ min}^{-1}$ ).

ADT43 with  $\text{CaCl}_2$  and *Pseudomonas* strain PF1 seed treatment had highest peroxidase activity ( $0.0474 \text{ g}^{-1} \text{ min}^{-1}$ ), while the genotype ADT43 with no external seed treatment recorded the lowest peroxidase activity of  $0.0394 \text{ g}^{-1} \text{ min}^{-1}$  under salt stress (Figure 1, Table 1). Genotypes and treatments significantly varied in peroxidase activity. Interaction between genotype and treatment was also found to be significant 5% probability level.



**Figure 2.** Effect of salt stress on catalase activity in two rice genotypes.

**Table 1.** Effect of salt stress on peroxidase ( $\Delta$  OD 403 nm min<sup>-1</sup> g<sup>-1</sup>) in two rice genotypes.


Treatments	ADT43	IR50
No treatment	0.0394	0.0426
CaCl <sub>2</sub>	0.0474	0.0429
PF1	0.0474	0.0446
TDK1	0.0470	0.0454
Mean	0.0453	0.0439
	<b>SEd</b>	<b>CD (P=0.05)</b>
Genotype (G)	0.0002	0.0004
Treatment (T)	0.0003	0.0005
GXT	0.0004	0.0008

**Table 2.** Effect of salt stress on catalase ( $\mu$ g of H<sub>2</sub>O<sub>2</sub> min<sup>-1</sup> g<sup>-1</sup>) in two rice genotypes.

Treatments	ADT43	IR50
No treatment	41.36	44.21
CaCl <sub>2</sub>	41.49	44.97
PF1	43.81	45.55
TDK1	54.59	52.44
Mean	45.31	46.79
	<b>SEd</b>	<b>CD (P=0.05)</b>
Genotype (G)	0.1399	0.2849
Treatment (T)	0.1978	0.4030
GXT	0.2798	0.5699

Catalase activity exhibited similar pattern of peroxidase activity. The highest enzyme activity of 54.59  $\mu$ g of H<sub>2</sub>O<sub>2</sub> min<sup>-1</sup> g<sup>-1</sup> was registered by ADT43 (treated with *Pseudomonas* strain TDK1). Catalase activity was increased in all the genotypes with seed treatment under

salt stress compared to the plants without any seed treatment. The lowest value of catalase activity (41.36  $\mu$ g of H<sub>2</sub>O<sub>2</sub> min<sup>-1</sup> g<sup>-1</sup>) was observed in ADT43 genotype without any seed treatment (Figure 2, Table 2). Genotypic variation and treatment effect were significant



**Figure 3.** Effect of salt stress on nitrate reductase activity in two rice genotypes.

**Table 3.** Effect of salt stress on nitrate reductase ( $\mu\text{g of NO}_2 \text{ h}^{-1} \text{ g}^{-1}$ ) in two rice genotypes.

Treatments	ADT43	IR50
No treatment	43.66	44.21
CaCl <sub>2</sub>	45.52	44.97
PF1	47.15	45.55
TDK1	48.64	52.44
Mean	46.24	46.79
	SEd	CD (P=0.05)
Genotype (G)	0.1419	0.2890
Treatment (T)	0.2007	0.4088
GXT	0.2838	0.5781

at  $P < 0.05$ .

The enzymatic antioxidant system is one of the protective mechanisms including superoxide dismutase, which can be found in various cell compartments and it catalyses the disproportion of two  $\text{O}_2^-$  radicals to  $\text{H}_2\text{O}_2$  and  $\text{O}_2$  (Scandalios, 1993).  $\text{H}_2\text{O}_2$  is eliminated by various antioxidant enzymes such as catalases (Scandalios, 1993) and peroxidases (Gara et al., 2003) which convert  $\text{H}_2\text{O}_2$  to water. Peroxidase is an important enzyme involved in morphogenesis and auxin oxidation. This enzyme is sensitive to environmental fluctuation and is considered as the measure of plants' resistance to abiotic stress.

In the present study, the peroxidase activity followed an increasing trend compared to control genotypes except for the genotype ADT43 grown with no external treatment which showed comparatively less value of peroxidase activity. PGPR alleviated the oxidative damage produced under water shortage (Kohler et al., 2008) confirmed by our study. Sandhya et al. (2010) reported that *Pseudomonas* inoculated maize plants showed increased antioxidant enzymes activity on exposure to drought

stress. Inoculation of lettuce (*Lactuca stiva* L.) with PGPR *Pseudomonas mendocina* augmented an antioxidant catalase under severe drought conditions, suggesting that they can be used as inoculants to alleviate the oxidative damage elicited by drought (Kohler et al., 2008).

### Nitrate reductase activity

During stress, Nitrate reductase (NRase) plays important role in reducing stress effects. It reduces nitrates to nitrite ions. Higher amount of NaCl ( $600 \text{ mM L}^{-1}$ ) reduces nitrate reductase activity considerably in leaves than in roots (Meloni et al., 2004). In the present study, among all the treatments used to treat the seeds of two genotypes IR50 genotype treated with *Pseudomonas* strain TDK1 showed highest NRase activity ( $52.44 \mu\text{g of NO}_2 \text{ hr}^{-1} \text{ g}^{-1}$ ), while the genotype ADT43 without any external seed treatment recorded the lowest NRase enzyme activity of  $43.66 \mu\text{g of NO}_2 \text{ h}^{-1} \text{ g}^{-1}$  (Figure 3, Table 3). Genotypic variation in nitrate reductase activity was significant. Different

*Pseudomonas* strain seed treatments improved the enzyme activity significantly. Interaction between genotype and treatment was significant.

The pathway of NO<sub>3</sub> assimilation is considered as the major route of conversion of inorganic N into a biologically useful organic compound. The primary step in NO<sub>3</sub> assimilation involves reduction of NO<sub>3</sub> to NO<sub>2</sub> catalyzed by the enzyme, nitrate reductase (NRase). Nitrite is subsequently reduced to NH<sub>4</sub><sup>+</sup> by the enzyme nitrite reductase. These two enzymes reduce NO<sub>3</sub> to the end product NH<sub>4</sub><sup>+</sup> which is then incorporated into amino acids. Activity of NRase in plants gives a good estimate of the N status of plants and is very often correlated with growth and yield of crops (Srivastava et al., 1988). Garg and Singla (2005) reported decrease in NRase activity in chickpea leaves and roots at all stages of the plant under salt stress. Similarly, Debouba et al (2007) found reduction in NO<sub>3</sub><sup>-</sup> content in tomato leaves affected by salinity. In the present study, among all the treatments used to treat the seeds of two genotypes IR50 genotype treated with TDK1 *Pseudomonas* strain showed highest NRase activity, while the genotype ADT43 without any external seed treatment recorded the lowest NRase enzyme activity. Genotypic variation and treatment effect was found to be significant.

## Conclusion

Plant growth promoting rhizobacteria has long been known to promote plant growth under abiotic stress conditions. Our study confirms the relationship among the antioxidative enzymes and the rhizobacteria. Peroxidase and catalase activities increased followed by *Pseudomonas* treatments. The activities of nitrate reductase were inhibited, as a function of NaCl concentration in the growth medium. The inhibition of NRase activity was comparatively more when the plants were grown under salt stress without any seed treatment. The results show that nitrate reductase is sensitive to NaCl stress. IR50 genotype treated with *Pseudomonas* strain TDK1 showed highest NRase activity. A lot of cultivated lands get salinized every year in India either due to natural or anthropogenic activities. Treating the rice seeds with PGPR prior to sowing would ameliorate the effects of salt stress to large extent.

## ACKNOWLEDGEMENTS

The authors would like to thank NLC Laboratory members of Department of Crop Physiology, Tamil Nadu Agricultural University for their technical assistance.

## REFERENCES

- Afzal I., Rauf S., Basra S. M. A. & Murtaza G. (2008). Halopriming improves vigor, metabolism of reserves and ionic contents in wheat seedlings under salt stress. *Plant Soil Environ.* 54(9):382-388.
- Candan N. & Tarhan L. (2003). The correlation between antioxidant enzyme activities and lipid peroxidation levels in *Mentha pulegium* organs grown in Ca<sup>2+</sup>, Mg<sup>2+</sup>, Cu<sup>2+</sup>, Zn<sup>2+</sup> and Mn<sup>2+</sup> stress conditions. *Plant Sci.* 163:769-779.
- Debouba M., Maaroufi-Dghimi H., Suzuki A., Ghorbel M. H. & Gouia H. (2007). Changes in growth and activity of enzymes involved in nitrate reduction and ammonium assimilation in tomato seedlings in response to NaCl stress. *Ann. Bot.* 99:1143-1151.
- Del Rio L. A., Corpas F. J., Sandalio L. M., Palma J. M. & Barroso J. B. (2003). Plant peroxisomes, reactive oxygen metabolism and nitric oxide. *IUBMB Life.* 55:71-81.
- Gara L. D., Pinto M. C. & Tommasi F. (2003). The antioxidant systems vis-à-vis reactive oxygen species during plant-pathogen interaction. *Plant Physiol. Biochem.* 41:863-870.
- Garg N. & Singla R. (2005). Nitrate reductase activity in roots and leaves of chickpea cultivars under salt stress. *Spanish J. Agric. Res.* 3(2):248-252.
- Jha Y. & Subramanian R. B. (2013). Paddy plants inoculated with PGPR show better growth physiology and nutrient content under saline conditions. *Chilean J. Agric. Res.* 73(3):213-219.
- Karabal E., Yücel M. & Öktem H. A. (2003). Antioxidant responses of tolerant and sensitive barley cultivars to boron toxicity. *Plant Sci.* 164:925-933.
- Kohler J., Tortosa G., Cegarra J., Caravaca F. & Roldan A. (2008). Impact of dom from composted "alperujo" on soil structure, AM fungi, microbial activity and growth of *Medicago sativa*. *Waste Manage.* 28(8):1423-1431.
- Luck H. (1974). In: *Methods in enzymatic analysis II* (Ed. Bergmeyer) Academic Press, New York. 885p.
- Meloni D. A., Gulotta M. R., Martínez C. A. & Oliva M. A. (2004). The effects of salt stress on growth, nitrate reduction and proline and glycinebetaine accumulation in *Prosopis alba*. *Braz. J. Plant Physiol.* 16(1):39-46.
- Neeraja C., Anil K., Purushotham P., Suma K., Sarma P., Moerschbacher B. M. & Podile A. R. (2010). Biotechnological approaches to develop bacterial chitinases as a bioshield against fungal diseases of plants. *Critic. Rev. Biotech.* 30:231-241.
- Nicholas J. C., Harper J. E. & Hageman R. H. (1976). Nitrate reductase activity in soybean. *Plant Physiol.* 58:731-735.
- Perur K. (1962). Developing salt tolerant cultivars in rice (*Oryza sativa* L.). *J. Agric. Sci.* 54:105-108.
- Sandhya V., Ali Sk. Z., Grover M., Reddy G. & Venkateswarlu B. (2010). Effect of plant growth promoting *Pseudomonas* spp. on compatible solutes, antioxidant status and plant growth of maize under drought stress. *Plant Growth Regul.* 62:21-30.
- Scandalios J. G. (1993). Oxygen stress and superoxide dismutase. *Plant Physiol.* 101:712-726.
- Srivastava J. P., Gupta S. C., Lal P., Muralia R. N. & Kumar A. (1988). Effect of salt stress on physiological and biochemical parameters in wheat. *Ann. Arid Zone.* 27:197-204.
- Tester M. & Davenport R. (2003). Na<sup>+</sup> tolerance and Na<sup>+</sup> transport in higher plants. *Ann. Bot.* 91:503-527.
- Yoshida S. D., Forno A. & Cock J. H. (1976). Laboratory manual for physiological studies of rice. IRRI, Philippines. Pp. 36-37.
- Zhu J., Fu X., Koo Y. D. & Zhu J. K. (2007). An enhancer mutant of arabidopsis salt overly sensitive 3 mediates both ion homeostasis and the oxidative stress response. *Mole. Cell Biol.* 27:5214-5224.