



## **GCE MARKING SCHEME**

**MATHEMATICS - M1-M3 & S1-S3  
AS/Advanced**

**SUMMER 2014**

## **INTRODUCTION**

The marking schemes which follow were those used by WJEC for the Summer 2014 examination in GCE MATHEMATICS - M1-M3 & S1-S3. They were finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conferences were held shortly after the papers were taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conferences was to ensure that the marking schemes were interpreted and applied in the same way by all examiners.


It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conferences, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about these marking schemes.

	<b>Page</b>
M1	1
M2	9
M3	16
S1	24
S2	28
S3	31

**M1****Q****Solution****Mark****Notes**

1(a)



Apply N2L to crate

M1    *R* and  $25g$  opposing.

Dim. Correct

$$25g - R = 25 \times 1.2$$

A1    correct equation

Any form

$$R = \underline{215} \text{ (N)}$$

A1

1(b)     $R = 25g = \underline{245} \text{ (N)}$ 

B1

**Q****Solution****Mark****Notes**


2(a) Use of  $v = u + at$  with  $u=10$ ,  $v=24$ ,  $t=21$   
 $24 = 10 + 21a$   
 $a = \frac{2}{3} (\text{ms}^{-2})$

M1 oe  
A1  
A1 accept anything derived  
from  $\frac{2}{3}$  rounded correctly

2(b)  $s = \frac{1}{2}(u + v)t$  with  $v=0$ ,  $u=24$ ,  $t=16$   
 $s = \frac{1}{2} \times 24 \times 16$   
 $s = \underline{192 \text{ (m)}}$

M1 oe  
A1  
A1

2(c)



B1 (0, 10) to (21, 24)  
B1 (21, 24) to (21+T, 24)  
B1 (21+T, 24) to (37+T, 0)  
B1 all labels, units and shape.


2(d) Area under graph = 15000  
 $0.5(10+24)21 + 24T + 192 = 15000$   
 $24T = 14451$   
 $T = \underline{602(\text{.125})}$

M1 used  
A1 ft (b)  
B1  $0.5(10+24)21$  or  $24T$   
Ft graph  
A1 Accept 600 from correct working. Cao.

<b>Q</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
3(a)	Resolve perpendicular to plane $R = mg\cos\alpha$ $F = \mu mg\cos\alpha$ $F = 0.6 \times 7 \times 9.8 \times \frac{4}{5}$ $F = \underline{32.9(28 \text{ N})}$	M1 m1	sin/cos correct expression
		A1	Accept rounding to 32.9.
3(b)	Apply N2L to A  $T + mgs\sin\alpha - F = 7a$ $T + 41.16 - 32.928 = 7a$ $T + 8.232 = 7a$  Apply N2L to B $3g - T = 3a$  $3g + 8.232 = 10a$	M1 A1 A1 M1 A1 m1	dim correct equation Friction opposes motion 4 terms. Accept cos. ft (a) dim correct equation one variable eliminated Dep on both M's
	$a = \underline{3.7(632 \text{ ms}^{-2})}$ $T = \underline{18.1(104 \text{ N})}$	A1 A1	cao cao

Q	Solution	Mark	Notes
---	----------	------	-------

4.



B1 any 1 correct moment.

Take moments about C M1 dim correct equation. oe

$$0.4R_D = 3g \times 0.6 + 12g \times 1.5 \quad \text{A1 correct equ any form}$$

$$0.4R_D = 19.8g = 194.04 \quad \text{A1 cao}$$

$$R_D = 49.5g = \underline{485.1} \text{ (N)} \quad \text{A1 cao}$$

Resolve vertically M1 equation attempted.  
Or 2<sup>nd</sup> moment equation.

$$R_D = R_C + 15g \quad \text{A1}$$

$$R_C = 34.5g = \underline{338.1} \text{ (N)} \quad \text{A1 cao}$$

#### Alternative solution

Moment equation about A/centre/B M1

Correct equation B1

Second moment equation M1

Correct equation A1

Correct method for solving simultaneously m1 Dep on both M's


$$R_C = 34.5g = \underline{338.1} \text{ (N)} \quad \text{A1 cao}$$

$$R_D = 49.5g = \underline{485.1} \text{ (N)} \quad \text{A1 cao}$$

<b>Q</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
5(a)	Resolve perpendicular to motion $20\sin 60 + T\sin 30 = 28\sin 60$ $20 \frac{\sqrt{3}}{2} + T \times \frac{1}{2} = 28 \frac{\sqrt{3}}{2}$ $T = \underline{8\sqrt{3}}$	M1 A1 A1	equation, sin/cos convincing
5(b)	N2L in direction of motion $20\cos 60 + T\cos 30 + 28\cos 60 - 16 = 80a$ $20 \times \frac{1}{2} + 8\sqrt{3} \times \frac{\sqrt{3}}{2} + 28 \times \frac{1}{2} - 16 = 80a$ $a = \underline{0.25 \text{ (ms}^{-2}\text{)}}$	M1 A2 A1	dim correct all forces and No extra force -1 each error cao
5(c)	N2L $-16 = 80a$ $a = -0.2$  Use of $v = u + at$ , $v=4$ , $u=12$ , $a=(+/-)0.2$ $4 = 12 - 0.2t$ $t = \underline{40 \text{ (s)}}$	M1 A1 m1 A1 A1	no extra force accept +/-  ft if $a < 0$ ft if $a < 0$

Q	Solution	Mark	Notes
---	----------	------	-------

6(a)



Conservation of momentum

M1 equation required  
Only one sign error.  
Ignore common factors

$$2 \times 3 - 7 \times 5 = 3v_A + 7v_B$$

$$3v_A + 7v_B = -29$$

A1

Restitution

M1  $v_B, v_A$  opposing consistent with diagram,  $+/-7$  with the 0.6.

$$v_B - v_A = -0.6(-5 - 2)$$

$$v_B - v_A = 4.2$$

A1

$$-7v_A + 7v_B = 29.4$$

$$3v_A + 7v_B = -29$$

$$10v_A = -58.4$$

m1 one variable eliminated.  
Dep on both M's.

$$v_A = (-)5.84$$

$$v_B = (-)1.64$$

A1 cao  
A1 cao

6(b) Impulse = change of momentum

M1 used

$$I = 7v_B - 7(-5)$$

$$I = -11.48 + 35$$

$$I = 23.52 \text{ (Ns)}$$


A1 ft their  $v_A$  or  $v_B$

6(c)  $3.65 = e(5.84)$

B1 ft  $v_A$  if  $> 3.65$ .

**Q****Solution****Mark****Notes**

7.



Resolve horizontally

$$T_{AB} \sin 60 = T_{AC} \sin 45$$

$$\frac{\sqrt{3}}{2} T_{AB} = \frac{1}{\sqrt{2}} T_{AC}$$

$$T_{AB} = \sqrt{\frac{2}{3}} T_{AC}$$

M1 equation, no extra force

A1

Resolve vertically

$$T_{AB} \cos 60 + T_{AC} \cos 45 = 9g$$

$$T_{AB} + \sqrt{2} T_{AC} = 18g$$

$$\sqrt{\frac{2}{3}} T_{AC} + \sqrt{2} T_{AC} = 18g$$

M1 equation, no extra force

A1

m1

$$T_{AC} = 79.(078) \text{ (N)}$$

A1 cao allow 79

$$T_{AB} = 64.(567) \text{ (N)}$$

A1 cao allow 65

Alternative MethodThird angle  $75^\circ/105^\circ$ 

B1

$$\frac{T_{AB}}{\sin 45} = \frac{9g}{\sin 75}$$

M1 sine rule attempted

$$T_{AB} = \frac{9g \times \sin 45}{\sin 75}$$

A1 si

$$T_{AB} = 64.(567) \text{ (N)}$$

A1 cao allow 65

$$\frac{T_{AC}}{\sin 60} = \frac{9g}{\sin 75}$$

M1 sine rule attempted

$$T_{AC} = \frac{9g \times \sin 60}{\sin 75}$$

A1 si

$$T_{AC} = 79.(078) \text{ (N)}$$

A1 cao allow 79

<b>Q</b>	<b>Solution</b>		<b>Mark</b>	<b>Notes</b>
8(a)	mass	$AD$	$AB$	
	$ABCD$	72	6	3 B1
	$XYZ$	12	6	2 B1
	$E$	24	3	4
	$F$	36	9	4 B1 both $E$ and $F$ correct
	Jewel	120	$x$	$y$ B1 masses in correct proportions.
8(a)(i)	Moments about $AD$			M1 masses and moments consistent.
	$120x + 12 \times 6 = 72 \times 6 + 24 \times 3 + 36 \times 9$			A1 ft table if triangle subt.
	$120x = 756$			
	$x = \frac{63}{10} = \underline{6.3\text{(cm)}}$			A1 cao
8(a)(ii)	Moments about $AB$			M1 masses & moments consistent
	$120y + 12 \times 2 = 72 \times 3 + 24 \times 4 + 36 \times 4$			A1 ft table if triangle subt.
	$120y = 432$			
	$y = \frac{18}{5} = \underline{3.6\text{(cm)}}$			A1 cao
8(b)	$PC = 12 - x$			
	$PC = \underline{5.7\text{(cm)}}$			B1 ft their $x$ if $< 12$ .

## M2

Q	Solution	Mark	Notes
1(a)	$\text{EE} = \frac{1}{2} \times \frac{\lambda x^2}{l}, \lambda=625, x=(+/-)0.1, l=0.2$ $\text{EE} = \frac{1}{2} \times \frac{625 \times 0.1^2}{0.2}$ $\text{EE} = \underline{15.625 \text{ (J)}}$	M1  A1	
1(b)	$\text{KE} = \frac{1}{2} \times 0.8v^2 (= 0.4v^2)$  $\text{WD by resistance} = 46 \times 0.1 (= 4.6)$  Work-energy Principle $\frac{1}{2} 0.8v^2 + 46 \times 0.1 = 15.625$ $0.4v^2 = 15.625 - 4.6$ $0.4v^2 = 11.025$ $v = \sqrt{\frac{11.025}{0.4}}$ $v = \underline{5.25 \text{ (ms}^{-1}\text{)}}$	B1  B1  M1  A1  A1	3 terms, no PE.  FT their EE  cao

Q	Solution	Mark	Notes
---	----------	------	-------

2(a)	$F - R = ma$ $30t^2 - 150 = 5a$ $6t^2 - 30 = a$ $\frac{dv}{dt} = 6t^{-2} - 30$	M1  A1	used, $F$ and $R$ opposing.  Answer given
(b)	$24 = \frac{6}{t^2} - 30$  $\frac{6}{t^2} = 54$  $t = \frac{1}{3}$	M1  A1	Ft (a) if same form  cao, accept 0.3.
2(c)	Integrate w.r.t. $t$ $v = -6t^{-1} - 30t (+ C)$  $t = \frac{1}{3}, v = 18$  $18 = -18 - 10 + C$ $C = 46$ $v = -6t^{-1} - 30t + 46$	M1  A1  m1	Increase in powers  m1
	When $v = 10$  $10 = -\frac{6}{t} - 30t + 46$  $5t^2 - 6t + 1 = 0$  $(5t - 1)(t - 1) = 0$  $t = \frac{1}{5}, 1$	m1  m1  A1	recognition of quadratic Some attempt to solve.  cao

<b>Q</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
----------	-----------------	-------------	--------------

3(a)  $T = \frac{P}{v}$ ,  $P = 90 \times 1000$ ,  $v = 4.8$  M1 si

$$T = \frac{90 \times 1000}{4.8}$$

$$T = 18750$$

N2L M1 dim correct, all forces  
 $T, R$  opposing.

$$T - mgsin\alpha - R = ma$$

$$18750 - 4000 \times 9.8 \times \frac{2}{49} - R = 4000 \times 1.2$$

$$R = 18750 - 1600 - 4800$$

$$R = \underline{12350 \text{ (N)}}$$

A1

A1

A1 cao

3(b) N2L with  $a = 0$  M1 all forces.

$$T = \frac{90 \times 1000}{v}$$

$$T - 1600 - 12800 = 0$$

$$v = \underline{6.25 \text{ ms}^{-1}}$$

B1 si

A1

A1

<b>Q</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
4(a)	$\mathbf{r} = \mathbf{p} + t\mathbf{v}$ $\mathbf{r}_A = (3 - t)\mathbf{i} + (5 + 2t)\mathbf{j} + (20 + t)\mathbf{k}$ $\mathbf{r}_B = (-2 + 3t)\mathbf{i} + (x - 4t)\mathbf{j} + (15 + 2t)\mathbf{k}$	M1 A1 A1	used
4(b)	$\mathbf{r}_B - \mathbf{r}_A =$ $(-5 + 4t)\mathbf{i} + (x - 5 - 6t)\mathbf{j} + (-5 + t)\mathbf{k}$	M1 A1	ft (a) similar expressions.
	$AB^2 = x^2 + y^2 + z^2$ $AB^2 = (-5 + 4t)^2 + (x - 5 - 6t)^2 + (-5 + t)^2$	M1 A1	cao
4(c)	Differentiate $\frac{dAB^2}{dt} = 2(-5 + 4t)(4) + 2(x - 5 - 6t)(-6)$ $+ 2(-5 + t)(1)$ $-40 + 32t - 12x + 60 + 72t - 10 + 2t = 0$ $106t + 10 = 12x$ When $t = 5$ $x = \underline{45}$	M1 m1 A1	powers reduced equating to 0. cao

Q	Solution	Mark	Notes
---	----------	------	-------

5(a)  $u_H = \frac{42}{2.5} = \underline{16.8 \text{ (ms}^{-1}\text{)}}$  B1

$s = u_V t + 0.5at^2, s = 3, t = 2.5, a = (\pm)9.8$  M1

$3 = 2.5u_V - 4.9 \times 2.5^2$  A1

$u_V = \underline{13.45 \text{ (ms}^{-1}\text{)}}$  A1 cao, accept 13.4, 13.5.

5(b)  $v_V = u_V + at, u_V = 13.45, a = (\pm)9.8, t = 2.5$  M1

$v_V = 13.45 - 9.8 \times 2.5$  A1 ft from (a)

$v_V = -11.05$

magnitude of vel =  $\sqrt{u_H^2 + v_V^2}$  m1

=  $\underline{20.11 \text{ (ms}^{-1}\text{)}}$  A1 cao

$\theta = \tan^{-1} \left( \frac{11.05}{16.8} \right)$  m1

$\theta = \underline{33.33^\circ}$  (below horizontal) A1 cao

5(c)  $s = ut + 0.5at^2, s = 0, u = 13.45, a = (\pm)9.8$  M1

$0 = 13.45t - 4.9t^2$

$t = 2.7449$

Distance =  $2.7449 \times 16.8$  m1

Distance = 46.11

Required distance =  $46.11 - 42 = \underline{4.11 \text{ (m)}}$  A1 cao

Q	Solution	Mark	Notes
6(a)	$\mathbf{a} = \frac{d\mathbf{v}}{dt}$  $\mathbf{a} = 8\cos 2t \mathbf{i} - 75\sin 5t \mathbf{j}$	M1  A1	differentiation attempted. Vectors required.
	At $t = \frac{3\pi}{2}$ , $(\mathbf{a} = -8\mathbf{i} + 75\mathbf{j})$	m1	substitution of $t$ .
	Magnitude of force $= 3 \times \sqrt{8^2 + 75^2}$ $= 226.28 \text{ (N)}$	M1 A1	or $\mathbf{F} = 3(-8\mathbf{i} + 75\mathbf{j})$ cao
6(b)	$\mathbf{r} = \int 4\sin 2t \mathbf{i} + 15\cos 5t \mathbf{j} dt$ $\mathbf{r} = -2\cos 2t \mathbf{i} + 3\sin 5t \mathbf{j} (+ \mathbf{c})$ At $t = 0$ , $-2\mathbf{i} + 3\mathbf{j} = -2\mathbf{i} + \mathbf{c}$ $\mathbf{c} = 3\mathbf{j}$ $\mathbf{r} = -2\cos 2t \mathbf{i} + 3\sin 5t \mathbf{j} + 3\mathbf{j}$	M1 A1  m1 A1	integration attempted cao
6(c)	Particle crosses the $y$ -axis when $-2\cos 2t = 0$	M1	
	$2t = \frac{\pi}{2}$		
	$t = \frac{\pi}{4}$	A1	cao
	Distance from origin $= 3\sin(5 \times \frac{\pi}{4}) + 3$ $= 0.88 \text{ (m)}$	m1 A1	substitute $t$ into $\mathbf{r}$ cao

<b>Q</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
7(a)	Conservation of energy $0.5m(4u)^2 = mg(2l) + 0.5mu^2$ $16u^2 = 4gl + u^2$ $u^2 = \frac{4}{15}gl$	M1 A1	
		A1	convincing
7(b)(i)	Conservation of energy $0.5m(4u)^2 = 0.5mv^2 + mgl(1 - \cos\theta)$ $v^2 = 16u^2 - 2gl + 2gl\cos\theta$ $v^2 = \frac{34}{15}gl + 2gl\cos\theta$	M1 A1 A1	
	N2L towards centre of circle $T - mg\cos\theta = \frac{mv^2}{l}$ $T = \frac{34}{15}mg + 3mg\cos\theta$ $T = \frac{mg}{15}(34 + 45\cos\theta)$	M1 A1 m1 A1	If M1s gained, substitute for $v^2$ . any correct form
7(b)(ii)	when $T = 0$ , $\cos\theta = -\frac{34}{45}$ $\theta = 139.1^\circ$	M1 A1	putting $T = 0$ in $a\cos\theta \pm b$ $Ft \cos\theta = a$ , $a < 0$ .


### M3

<b>Q</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
1(a) N2L	$500 - 100v = 1200 \frac{dv}{dt}$ $\frac{dv}{dt} = \frac{500 - 100v}{1200} = \frac{5-v}{12}$	M1	
		A1	convincing
1(b)	$\int 12 \frac{dv}{5-v} = \int dt$ $-12\ln(5-v) = t + (C)$ <p>When <math>t = 0, v = 0, C = -12\ln 5</math></p> $t = 12\ln\left(\frac{5}{5-v}\right)$ $\frac{5}{5-v} = e^{\frac{t}{12}}$ $v = 5(1 - e^{-t/12})$	M1 A1 m1 m1 A1	sep. var. ( $5-v$ ) together. correct integration allow +/-, oe inversion ft similar exp. cao
	limiting speed = 5 (ms <sup>-1</sup> )	B1	Ft similar expression
1(c)	<p>When <math>v = 4, t = 12\ln\left(\frac{5}{5-4}\right)</math></p> $t = 12\ln 5 (= 19.31s)$	M1 A1	

<b>Q</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
2(a)	Period = $\frac{2\pi}{\omega} = 2$ $k = \omega = \pi$	M1 A1	
2(b)	$x = 0.52\cos\pi t$ When $t = \frac{1}{3}$ , $x = 0.52\cos\frac{\pi}{3}$ $x = 0.26$	B1 M1 A1	for amp=0.52 allow asin/acos, c's a cao
2(c)	$0.4 = 0.52\cos\pi t$ $\cos\pi t = \frac{0.4}{0.52}$ $t = 0.22$ $t = 1.78$	M1 A1 A1	allow sin/cos cao FT $t$ , ie 2-first $t$ .
2(d)	$v^2 = \omega^2(0.52^2 - x^2)$ $v^2 = \pi^2(0.52^2 - 0.2^2)$ $v = \pi(0.48) (= 1.508 \text{ ms}^{-1})$	M1 m1 A1	used. oe sub $x = 0.2$ cao
2(e)	$\max v = a\omega$ $= 0.52\pi (= 1.634 \text{ ms}^{-1})$	M1 A1	used cao

**Q****Solution****Mark****Notes**

3



Impulse = change in momentum

M1 used

$$J = 2ucos30 - 2v$$

A1

$$J = 3v$$

B1

Eliminating  $J$ 

m1 one variable eliminated

$$3v = 2ucos30 - 2v$$

$$5v = 2ucos30$$

A1 cao

$$v = 0.4u \cos 30$$

$$v = 2.77 \text{ (ms}^{-1}\text{)} \text{(speed of } A\text{)}$$

$$J = 1.2 u \cos 30 = 8.31 \text{ (Ns)}$$

A1 ft 3 x c's  $v$ .

$$u_B = u \sin 30 = 4 \text{ (ms}^{-1}\text{)}$$

B1

$$\text{Speed of } B = \sqrt{(2.77^2 + 4^2)}$$

m1

$$\text{Speed of } B = 4.87 \text{ (ms}^{-1}\text{)}$$

A1 cao

<b>Q</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
4(a)	Auxiliary equation $2m^2 + 6m + 5 = 0$ $m = -1.5 \pm 0.5i$ C.F. is $x = e^{-1.5t}(A\sin 0.5t + B\cos 0.5t)$	B1 B1 B1	ft complex roots
	For PI, try $x = a$ $5a = 1$ $a = 0.2$	B1	ft CF + a
	GS is $x = e^{-1.5t}(A\sin 0.5t + B\cos 0.5t) + 0.2$	B1	
4(b)	$e^{-1.5t} \rightarrow 0$ as $t \rightarrow \infty$ $x$ tends to 0.2 as $t$ tends to infinity Limiting value = 0.2	M1 A1	si ft similar expression
4(c)(i)	$x = 0.5$ and $\frac{dx}{dt} = 0$ when $t = 0$ $B + 0.2 = 0.5$ $B = 0.3$	M1 A1	used cao
	$\begin{aligned}\frac{dx}{dt} &= -1.5e^{-1.5t}(A\sin 0.5t + B\cos 0.5t) \\ &\quad + e^{-1.5t}(0.5A\cos 0.5t - 0.5B\sin 0.5t)\end{aligned}$ $0 = -1.5B + 0.5A$ $A = 3B = 0.9$	B1 A1	ft similar expressions cao
	$x = e^{-1.5t}(0.9\sin 0.5t + 0.3\cos 0.5t) + 0.2$		
4(c)(ii)	When $t = \frac{\pi}{3}$ $x = e^{-\pi/2}(0.9\sin \frac{\pi}{6} + 0.3\cos \frac{\pi}{6}) + 0.2$ $x = 0.348$		
		A1	cao

<b>Q</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
5(a)	Using $F = ma$ $1200(v+3)^{-1} = 800 a$ $2v \frac{dv}{dx} = \frac{3}{v+3}$	M1  A1	convincing
5(b)	$\int 3dx = \int 2v(v+3)dv$ $3x = \frac{2v^3}{3} + 3v^2 + (C)$	M1  A1	separate variables correct integration
	$x = 0, v = 0$ , hence $C = 0$ When $v = 3$ , $3x = 18 + 27$ $x = 15$	B1 m1 A1	
5(c)	$\frac{dv}{dt} = \frac{3}{2(v+3)}$ $\int 2(v+3)dv = \int 3dt$ $v^2 + 6v = 3t + (C)$	M1  A1	
	$t = 0, v = 0$ , hence $C = 0$ When $v = 3$ $3t = 9 + 18 = 27$ $t = 9$	B1  A1	cao
5(d)(i)	$v^2 + 6v - 3t = 0$ $v = 0.5(-6 \pm \sqrt{(6^2 - 4 \times -3t)})$ $v = -3 + \sqrt{9 + 3t}$	M1  A1  A1	recognition of quadratic And attempt to solve si
(ii)	$\frac{dx}{dt} = -3 + (9 + 3t)^{\frac{1}{2}}$ $x = -3t + \frac{2}{9}(9 + 3t)^{\frac{3}{2}} + (C)$ $x = 0, t = 0$ , (hence $C = -6$ ) $x = -3t + \frac{2}{9}(9 + 3t)^{\frac{3}{2}} + (-6)$	M1  A1  m1	
	When $t = 7$ $x = -21 - 6 + 2 \times 30^{1.5}/9 = 9.5148$ $x$ is approximately 9.5	A1	cao

<b>Q</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
----------	-----------------	-------------	--------------

5(d)(ii)  $v = -3 + \sqrt{9 + 3t}$   
When  $t=7$ ,  $v = -3 + \sqrt{9+21}$   
 $v = -3 + \sqrt{30}$   
 $v = 2.4723$

M1  
A1 si


$$x = \frac{2}{9}(-2.4723)^3 + (2.4723)^2$$

m1

x = 9.51 (m) A1 cao

**Q****Solution****Mark****Notes**

6(a)



B2      B1 if one error.  
 B0      more than one error.

6(b)    Resolve vertically

$$R = 12g + 70g = 82g$$

M1      all forces

A1

6(c)    Moments about B

$$3T\sin 75 + 12g \times 4\cos 75 + 70gx \times \cos 75 = 8S\sin 75$$

M1      dim correct equation  
 All terms

A4      -1 each incorrect term  
 Accept  $T=100$ .

Resolve horizontally

$$T + F = S$$

$$F = 0.1R = 8.2g$$

$$S = T + 8.2g$$

B1      ft  $R$ B1      ft  $F$ 

$$8(8.2g + T)\sin 75 - 3T\sin 75 - 48g\cos 75 = 70gx\cos 75$$

$$5T\sin 75 =$$

$$48g\cos 75 - 65.6g\sin 75 + 70gx\cos 75$$

$$T = 100$$

$$x = 5.53 \text{ m}$$

A1      cao

<b>Q</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
----------	-----------------	-------------	--------------

**OR**

Moments about A

M1 dim correct equation  
All terms

$$5T\sin 75 + 12g \times 4\cos 75 + 70g(8-x)\cos 75 \\ + 8F\sin 75 = 8R\cos 75$$

A5 -1 each incorrect term  
Accept  $T=100$ .

$$F = 0.1R = 80.36 \text{ N}$$

B1 Ft R

$$T = 100$$

A1 cao

$$x = 5.53 \text{ m}$$

- 6(d) Ladder modelled as a rigid rod. B1

# S1

<b>Ques</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
1(a)	EITHER $P(A \cap B) = P(A) + P(B) - P(A \cup B)$ $= 0.2$	<b>M1</b> <b>A1</b>	Award M1 for using formula
(b)	This is not equal to $P(A) \times P(B)$ therefore not independent. OR Assume A,B are independent so that $P(A \cap B) = P(A) + P(B) - P(A)P(B)$ $= 0.58$ <p>Since <math>P(A \cup B) \neq 0.58</math>, A,B are not independent.</p> $P(A   B') = \frac{P(A \cap B')}{P(B')}$ $= \frac{0.3 - 0.2}{0.6}$ $= \frac{1}{6}$	<b>A1</b>  <b>M1</b> <b>A1</b> <b>A1</b>  <b>M1</b> <b>A1</b> <b>A1</b>	Award M1 for using formula  FT their $P(A \cap B)$ if independence not assumed  Accept Venn diagram
2	$np = 0.9, npq = 0.81$ Dividing, $q = 0.9, p = 0.1$ $n = 9$	<b>B1B1</b> <b>M1A1</b> <b>A1</b>	
3(a)	$P(1 \text{ of each}) =$ $\frac{3}{9} \times \frac{3}{8} \times \frac{3}{7} \times 6 \text{ or } \binom{3}{1} \times \binom{3}{1} \times \binom{3}{1} \div \binom{9}{3}$ $= \frac{9}{28}$	  <b>M1A1</b>  <b>A1</b>	M1A0 if 6 omitted
(b)	$P(2 \text{ particular colour and 1 different}) =$ $\frac{3}{9} \times \frac{2}{8} \times \frac{6}{7} \times 3 \text{ or } \binom{3}{2} \times \binom{6}{1} \div \binom{9}{3}$ $= \frac{3}{14}$ $P(2 \text{ of any colour and 1 different}) = \frac{9}{14}$	  <b>M1A1</b>  <b>A1</b>  <b>B1</b>	M1A0 if 3 omitted  Allow 3/28  FT previous line
4(a)	Let $X$ denote the number of goals scored in the first 15 minutes so that $X$ is Po(1.5) si $P(X = 2) = \frac{e^{-1.5} \times 1.5^2}{2!}$ $= 0.251$	  <b>B1</b>  <b>M1</b> <b>A1</b>	
(b)	$P(X > 2) = 1 - e^{-1.5} \left( 1 + 1.5 + \frac{1.5^2}{2!} \right)$ $= 0.191$	  <b>M1A1</b>  <b>A1</b>	Award M0 if no working seen

<b>Ques</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
5(a) (i)	Let $X$ = number of female dogs so $X$ is $B(20,0.55)$ $P(X = 12) = \binom{20}{12} \times 0.55^{12} \times 0.45^8$ $= 0.162$	<b>B1</b> <b>M1</b> <b>A1</b>	si Accept 0.4143 – 0.2520 or 0.7480 – 0.5857
(ii)	Let $Y$ = number of male dogs so $Y$ is $B(20,0.45)$ $P(8 \leq X \leq 16) = P(4 \leq Y \leq 12)$ $= 0.9420 - 0.0049$ or $0.9951 - 0.0580$ $= 0.9371$	<b>M1</b> <b>A1</b> <b>A1A1</b> <b>A1</b>	Award M0 if no working seen
(b)	Let $U$ = number of yellow dogs so $U$ is $B(60,0.05) \approx Po(3)$ $P(U < 5) = 0.8153$	<b>M1</b> <b>m1A1</b>	
6(a)	$P(\text{head}) = \frac{3}{4} \times \frac{1}{2} + \frac{1}{4} \times 1$ $= \frac{5}{8}$	<b>M1A1</b> <b>A1</b>	M1 Use of Law of Total Prob (Accept tree diagram)
(b)(i)	$P(\text{DH} \text{head}) = \frac{1/4}{5/8}$ $= \frac{2}{5} \text{ cao}$	<b>B1B1</b> <b>B1</b>	B1 num, B1 denom FT denominator from (a)
(ii)	EITHER $P(\text{head}) = \frac{3}{5} \times \frac{1}{2} + \frac{2}{5} \times 1$ $= \frac{7}{10}$ OR $P(\text{Head}) = \frac{\frac{3}{4} \times \frac{1}{2} \times \frac{1}{2} + \frac{1}{4} \times 1}{\frac{5}{8}}$ $= \frac{7}{10}$	<b>M1A1</b> <b>A1</b> <b>B1B1</b> <b>B1</b>	M1 Use of Law of Total Prob (Accept tree diagram)  B1 num, B1 denom FT denominator from (a)

<b>Ques</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
7(a)	[0,0.4]	<b>B1</b>	Allow(0,0.4)
(b)	$E(X) = 0.1 + 0.6 + 3\theta + 0.8 + 5(0.4 - \theta)$ = 3.5 - 2\theta The range is [2.7,3.5]	<b>M1</b> <b>A1</b> <b>A1</b>	FT the range from (a)
(c)	$E(X^2) = 0.1 + 1.2 + 9\theta + 3.2 + 25(0.4 - \theta)$ $\text{Var}(X) = 0.1 + 1.2 + 9\theta + 3.2 + 25(0.4 - \theta)$ – $(3.5 - 2\theta)^2$ = 2.25 – 2\theta – 4\theta <sup>2</sup> $\text{Var}(X) = 1.5$ gives $4\theta^2 + 2\theta - 0.75 = 0$ $16\theta^2 + 8\theta - 3 = 0$ $(4\theta + 3)(4\theta - 1) = 0$ $\theta = 0.25$	<b>M1A1</b> <b>M1</b> <b>A1</b> <b>M1</b> <b>A1</b> <b>M1</b> <b>A1</b>	Must be in terms of $\theta$ Allow use of formula
8(a)	EITHER the sample space contains 64 pairs of which 8 are equal OR whatever number one of them obtains, 1 number out of 8 obtained by the other one gives equality.	<b>M1</b>	
(b)	$P(\text{equal numbers}) = \frac{1}{8}$ The possible pairs are (4,8);(5,7);(6,6);(7,5);(8,4) EITHER the sample space contains 64 pairs of which 5 give a sum of 12 OR each pair has probability 1/64. $P(\text{sum} = 12) = \frac{5}{64}$	<b>A1</b> <b>B1</b> <b>M1</b> <b>A1</b>	
(c)	EITHER reduce the sample space to (4,8);(5,7);(6,6);(7,5);(8,4) OR $P(\text{equal numbers}) = \frac{P(6,6)}{P(\text{sum}=12)} = \frac{1/64}{5/64}$ Therefore $P(\text{equal numbers}) = \frac{1}{5}$	<b>M1</b> <b>A1</b>	

<b>Ques</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
9(a)(i)	$P(0.4 \leq X \leq 0.6) = F(0.6) - F(0.4)$ $= 0.261$	<b>M1</b> <b>A1</b>	
(ii)	The median $m$ satisfies $2m^3 - m^6 = 0.5$ $2m^6 - 4m^3 + 1 = 0$ $m^3 = \frac{4 \pm \sqrt{8}}{4}$ (0.293) $m = 0.664$	<b>B1</b>  <b>M1A1</b> <b>A1</b>	Award M1 for a valid attempt to solve the equation Do not award A1 if both roots given
(b)(i)	Attempting to differentiate $F(x)$ $f(x) = 6x^2 - 6x^5$	<b>M1</b> <b>A1</b>	
(ii)	$E(X^3) = \int_0^1 x^3 (6x^2 - 6x^5) dx$ $= \left[ \frac{6x^6}{6} - \frac{6x^9}{9} \right]_0^1$ $= 1/3$	<b>M1A1</b>  <b>A1</b>  <b>A1</b>	M1 for the integral of $x^3 f(x)$ A1 for completely correct although limits may be left until 2 <sup>nd</sup> line. FT their $f(x)$ if M1 awarded in (i)

## S2

<b>Ques</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
1	$\bar{x} = \frac{405.6}{8} (= 50.7)$ SE of $\bar{X} = \frac{4}{\sqrt{8}} (= 1.4142\dots)$ 90% conf limits are $50.7 \pm 1.645 \times 1.4142\dots$ giving [48.4, 53.0] cao	<b>B1</b> <b>M1A1</b> <b>M1A1</b> <b>A1</b>	M1 correct form, A1 correct z. Award M0 if no working seen
2(a)	Upper quartile = mean + $0.6745 \times \text{SD}$ $= 86.0$	<b>M1</b> <b>A1</b>	
(b)	Let $X$ =weight of an orange, $Y$ =weight of a lemon $E(\Sigma X) = 1984$ $\text{Var}(\Sigma X) = 512$ $z = \frac{2000 - 1984}{\sqrt{512}} = 0.71$	<b>B1</b> <b>B1</b> <b>M1A1</b>	Award M0 if no working seen
(c)	$\text{Prob} = 0.7611$ cao Let $U = X - 3Y$ $E(U) = -7$ $\text{Var}(U) = 64 + 9 \times 2.25 = 84.25$ We require $P(U > 0)$ $z = \frac{0 + 7}{\sqrt{84.25}} = 0.76$ $\text{Prob} = 0.2236$	<b>A1</b> <b>M1</b> <b>A1</b> <b>M1A1</b> <b>m1A1</b> <b>A1</b>	Award m0 if no working seen
3(a)	$H_0 : \mu_M = \mu_F; H_1 : \mu_M \neq \mu_F$	<b>B1</b>	
(b)	Let $X$ = male weight, $Y$ =female weight $(\sum x = 39.2; \sum y = 46.6)$ $\bar{x} = 4.9; \bar{y} = 4.66$ SE of diff of means = $\sqrt{\frac{0.5^2}{8} + \frac{0.5^2}{10}} (0.237\dots)$ $\text{Test statistic} = \frac{4.9 - 4.66}{0.237\dots}$ $= 1.01$ $\text{Prob from tables} = 0.1562$ $p\text{-value} = 0.3124$ Insufficient evidence to conclude that there is a difference in mean weight between males and females.	<b>B1B1</b> <b>M1A1</b> <b>m1</b> <b>A1</b> <b>A1</b> <b>B1</b> <b>B1</b>	Award m0 if no working seen FT line above FT their $p$ -value

<b>Ques</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
4(a)(i)	$H_0 : p = 0.6; H_1 : p < 0.6$	<b>B1</b>	
(ii)	<p>Let <math>X</math> = Number of games won  Under <math>H_0</math>, <math>X</math> is <math>B(20,0.6)</math> si  Let <math>Y</math> = Number of games lost  Under <math>H_0</math>, <math>Y</math> is <math>B(20,0.4)</math></p> $p\text{-value} = P(X \leq 7   X \text{ is } B(20,0.6))$ $= P(Y \geq 13   Y \text{ is } B(20,0.4))$ $= 0.021$ <p>Strong evidence to reject Gwilym's claim (or to accept Huw's claim).</p>	<b>B1</b> <b>B1</b> <b>B1</b> <b>M1</b> <b>A1</b> <b>A1</b> <b>B1</b>	Award M0 if no working seen FT on p-value
(b)	<p><math>X</math> is now <math>B(80,0.6)</math> (under <math>H_0) \approx N(48,19.2)</math>)  <math>p\text{-value} = P(X \leq 37   X \text{ is } N(48,19.2))</math></p> $z = \frac{37.5 - 48}{\sqrt{19.2}}$ $= -2.40$ $p\text{-value} = 0.0082$ <p>Very strong evidence to reject Gwilym's claim (or to accept Huw's claim).</p>	<b>B1B1</b> <b>M1</b> <b>A1</b> <b>A1</b> <b>A1</b> <b>B1</b>	Award M0 if no working seen Award M1A0A1 for incorrect or no continuity correction No cc ; $z = -2.51, p = 0.00604$ $36.5 ; z = -2.62, p = 0.0044$ FT on p-value only if less than 0.01
5(a)	$E(X) = E(Y) = 1.2$ $E(U) = E(X)E(Y) = 1.44$ cao	<b>B1</b> <b>B1</b>	
(b)	<p><math>\text{Var}(X) = \text{Var}(Y) = 0.96</math>  <math>E(X^2)(= E(Y^2)) = \text{Var}(X) + [E(X)]^2 = 2.4</math></p> $\text{Var}(U) = E(X^2Y^2) - [E(XY)]^2$ $= E(X^2)E(Y^2) - [E(X)E(Y)]^2$ $= 3.69$ cao	<b>B1</b> <b>M1A1</b> <b>M1</b> <b>A1</b> <b>A1</b>	FT their values from (a)
6(a)(i)	<p>Under <math>H_0</math>, <math>X</math> is <math>Po(15)</math> si  <math>P(X \leq 10) = 0.1185; P(X \geq 20) = 0.1248</math>  Significance level = 0.2433</p>	<b>B1</b> <b>B1</b> <b>B1</b>	Award B1 for either correct
(ii)	<p><math>X</math> is now <math>Poi(10)</math>  <math>P(\text{accept } H_0) = P(11 \leq X \leq 19)</math></p> $= 0.9965 - 0.5830 \text{ or } 0.4170 - 0.0035$ $= 0.4135$ cao	<b>B1</b> <b>M1</b> <b>A1</b> <b>A1</b>	Award M0 if no working seen
(b)	<p>Under <math>H_0</math>, <math>X</math> is now <math>Po(75) \approx N(75,75)</math></p> $z = \frac{91.5 - 75}{\sqrt{75}} = 1.91$ <p>Prob from tables = 0.0281  <math>p\text{-value} = 0.056</math>  Insufficient evidence to reject <math>H_0</math></p>	<b>B1</b> <b>M1A1</b> <b>A1</b> <b>A1</b> <b>B1</b>	Award M1A0 for incorrect or no continuity correction but FT further work. FT from line above FT from line above No cc gives $z = 1.96, p = .05$ 92.5 gives $z = 2.02, p = 0.0434$

<b>Ques</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
7(a)	$\begin{aligned} P(L \leq 4) &= P(A \leq 4^2) \\ &= \frac{16-15}{20-15} \\ &= 0.2 \end{aligned}$	<b>M1</b> <b>A1</b> <b>A1</b>	
(b)	$\begin{aligned} E(L) &= E(A^{1/2}) \\ &= \int_{15}^{20} a^{1/2} \times \frac{1}{5} da \\ &= \frac{2}{15} [a^{3/2}]_{15}^{20} \\ &= 4.18 \end{aligned}$	<b>M1A1</b> <b>A1</b> <b>A1</b>	Limits can be left until next line Do not accept $\sqrt{17.5} = 4.18$
(c)	$\begin{aligned} \text{Var}(L) &= E(L^2) - [E(L)]^2 \\ &= 17.5 - 4.18^2 \\ &= 0.03 \end{aligned}$	<b>M1</b> <b>A1</b> <b>A1</b>	FT their $E(L)$

### S3

Ques	Solution	Mark	Notes
1	$\bar{x} = 52.0 \text{ si}$ $\text{Variance estimate} = \frac{162480}{59} - \frac{3120^2}{60 \times 59} = 4.068$ (Accept division by 60 which gives 4.0) 90% confidence limits are $52 \pm 1.645\sqrt{4.068/60}$ giving [51.6, 52.4]	<b>B1</b> <b>M1A1</b> <b>M1A1</b> <b>A1</b>	
(a)	$H_0: \mu = 4.5; H_1: \mu \neq 4.5$	<b>B1</b>	
(b)	$\sum x = 43.6; \sum x^2 = 190.3428$ UE of $\mu = 4.36$ $\text{UE of } \sigma^2 = \frac{190.3428}{9} - \frac{43.6^2}{90}$ $= 0.0274(22\dots)$	<b>B1B1</b> <b>B1</b> <b>M1</b> <b>A1</b>	No working need be seen  Answer only no marks
(c)	$\text{test-stat} = \frac{4.36 - 4.5}{\sqrt{0.0274222../10}}$ $= -2.67 \text{ (Accept } +2.67)$ $\text{DF} = 9 \text{ si}$ Crit value = 3.25 This result suggests that we should accept $H_0$ , ie that the mean weight is 4.5 kg because $2.67 < 3.25$	<b>M1A1</b> <b>A1</b> <b>B1</b> <b>B1</b> <b>B1</b> <b>B1</b>	FT their values from (b)  Answer only no marks  FT their <i>t</i> -statistic
(a)	$\hat{p} = \frac{654}{1500} = 0.436 \text{ si}$ $\text{ESE} = \sqrt{\frac{0.436 \times 0.564}{1500}} = 0.0128.. \text{ si}$ 95% confidence limits are $0.436 \pm 1.96 \times 0.0128..$ giving [0.41, 0.46]	<b>B1</b> <b>M1A1</b> <b>M1</b> <b>A1</b> <b>A1</b>	M1 correct form A1 correct <i>z</i>
(b)	$\hat{p} = \frac{0.4348 + 0.4852}{2} = 0.46$ Number of people = $0.46 \times 1200 = 552$ $0.4852 - 0.4348 = 2z\sqrt{\frac{0.46 \times 0.54}{1200}}$ $z = 1.75$ Prob from tables = 0.0401 or 0.9599 Confidence level = 92%	<b>B1</b> <b>B1</b> <b>M1A1</b> <b>A1</b> <b>A1</b> <b>B1</b>	FT line above

<b>Ques</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
4(a) (b)	$H_0: \mu_a = \mu_b; H_1: \mu_a \neq \mu_b$ $SE = \sqrt{\frac{0.115}{80} + \frac{0.096}{70}} (= 0.053)$ $\text{Test stat} = \frac{3.65 - 3.52}{0.053} = 2.45 \quad (\text{Accept 2.46})$ $\text{Tabular value} = 0.00714 \quad (0.00695)$ $p\text{-value} = 0.01428 \quad (0.0139)$ Strong evidence to conclude that there is a difference in mean weight.	<b>B1</b> <b>M1A1</b> <b>M1A1</b> <b>A1</b> <b>A1</b> <b>A1</b> <b>B1</b> <b>B1</b>	FT their $p$ -value Accept the conclusion that the Variety B mean is greater than the Variety A mean
(c)	Estimates of the variances of the sample means are used and not exact values. The sample means are assumed to be normally distributed (using the Central Limit Theorem).	<b>B1</b> <b>B1</b>	
5(a)	$\sum x = 42, \sum x^2 = 364, \sum y = 340.6, \sum xy = 2906.4$ $S_{xy} = 2906.4 - 42 \times 340.6 / 6 = 522.2$ $S_{xx} = 364 - 42^2 / 6 = 70$ $b = \frac{522.2}{70} = 7.46$ $a = \frac{340.6 - 7.46 \times 42}{6} = 4.55$	<b>B2</b> <b>B1</b> <b>B1</b> <b>M1</b> <b>A1</b> <b>M1</b> <b>A1</b>	Minus 1 each error  Answers only no marks
(b)(i)	Unbiased estimate = $a + 5b = 41.85$	<b>B1</b>	FT their values of and $a, b$ if answer between 33.9 and 49.9
(ii)	$\text{SE of } a + 5b = 0.5 \sqrt{\frac{1}{6} + \frac{(5-7)^2}{70}} \quad (0.2365\dots)$ 95% confidence limits for $a + 5\beta$ are $41.85 \pm 1.96 \times 0.2365\dots$ giving [41.4, 42.3]	<b>M1A1</b> <b>m1A1</b> <b>A1</b>	And FT their value of $S_{xx}$
(iii)	$\text{Test stat} = \frac{7.6 - 7.46}{\sqrt{0.5^2 / 70}} = 2.34$ Critical value = 1.96 or $p\text{-value} = 0.01928$ We conclude that $\beta = 7.6$ is not consistent with the tabular values.	<b>M1A1</b> <b>A1</b> <b>B1</b>	FT their values of $b$ and $S_{xx}$ if possible. FT their test statistic FT the line above

<b>Ques</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
6(a)(i)	$E(Y) = kE(\bar{X}) = kE(X) = \frac{k\theta}{2}$ For an unbiased estimator, $k = 2$ .	<b>M1A1</b> <b>A1</b>	
(ii)	$\text{Var}(Y) = 4\text{Var}(\bar{X})$ $= \frac{4}{n}\text{Var}(X)$ $= \frac{4}{n} \times \frac{\theta^2}{12}$ $= \frac{\theta^2}{3n}$ $\text{SE} = \frac{\theta}{\sqrt{3n}}$	<b>M1</b> <b>A1</b> <b>A1</b> <b>A1</b> <b>A1</b>	FT their $k$
(b)(i)	Using $\text{Var}(Y) = E(Y^2) - [E(Y)]^2$ $E(Y^2) = \frac{\theta^2}{3n} + \theta^2$ $\neq \theta^2$ therefore not unbiased	<b>M1</b> <b>A1</b> <b>B1</b>	FT the line above
(ii)	$E(Y^2) = \theta^2 \left( \frac{3n+1}{3n} \right)$ $E \left( \frac{3nY^2}{3n+1} \right) = \theta^2$ Therefore $\frac{3nY^2}{3n+1}$ is an unbiased estimator for $\theta^2$	<b>M1</b> <b>A1</b> <b>A1</b>	



WJEC  
245 Western Avenue  
Cardiff CF5 2YX  
Tel No 029 2026 5000  
Fax 029 2057 5994  
E-mail: [exams@wjec.co.uk](mailto:exams@wjec.co.uk)  
website: [www.wjec.co.uk](http://www.wjec.co.uk)