GENETICAL RESEARCH

EDITORIAL BOARD

C. H. WADDINGTON (Chairman)
CHARLOTTE AUERBACH D. LEWIS
H. GRÜNEBERG G. PONTECORVO
W. HAYES A. ROBERTSON

EXECUTIVE EDITOR

E. C. R. REEVE

ASSISTANT EXECUTIVE EDITOR

J. O. BISHOP

Volume 7. 1966



CAMBRIDGE
AT THE UNIVERSITY PRESS
1966

Printed in Great Britain by Spottiswoode, Ballantyne & Co. Ltd.

London and Colchester

and published by the Cambridge University Press

Bentley House, 200 Euston Road, London, N.W.1

American Branch: 32 East 57th Street, New York, N.Y. 10022

Agents in Canada: The Macmillan Company of Canada Limited

CONTENTS

No. 1 (February 1966)

| Sugino, Y. Mutants of <i>Escherichia coli</i> sensitive to methylene blue and acridines | page | 1 |
|---|------|-----|
| Lyon, M. F. and Morris, T. Mutation rates at a new set of specific loci in the mouse | | 12 |
| STADLER, D. R. Glutamic dehydrogenase in revertants of am mutants in neurospora | | 18 |
| CINADER, B., DUBISKI, S. and WARDLAW, A. C. Genetics of MuBl and of a complement defect in inbred strains of mice | | 32 |
| Bohren, B. B., Hill, W. G. and Robertson, A. Some observations on asymmetrical correlated responses to selection | | 44 |
| GRÜNEBERG, H. The case for somatic crossing-over in the mouse | | 58 |
| MORTON, J. R. The multiple electrophoretic bands of mouse haemoglobins | | 76 |
| SEARLE, A. G. Curtailed, a new dominant T-allele in the house mouse | | 86 |
| SPICKETT, S. G. and Thoday, J. M. Regular responses to selection. 3. Interaction between located polygenes | | 96 |
| Bray, D. F. Variability and restrictions against inbreeding and unequal family size in control populations of $Tribolium$ | 1 | .22 |
| Lyon, M. F. Order of loci on the X-chromosome of the mouse | 1 | 30 |
| MEYNELL, E. and Datta, N. The relation of resistance transfer factors to the F-factor (sex-factor) of <i>Escherichia coli</i> K12 | 1 | .34 |
| MEYNELL, E. and DATTA, N. The nature and incidence of conjugation factors in <i>Escherichia coli</i> | 1 | .41 |

vi Contents

No. 2 (April 1966)

| Prevosti, A. Chromosomal polymorphism in western Mediterranean populations of <i>Drosophila subobscura</i> | page | 149 |
|---|------|-----|
| BLOOM, J. L. and FALCONER, D. S. 'Grizzled', a mutant in linkage group X of the mouse | | 159 |
| McNelly-Ingle, C. A., Lamb, B. C. and Frost, L. C. The effect of temperature on recombination frequency in <i>Neurospora crassa</i> | | 169 |
| CLUTTERBUCK, A. J. and ROPER, J. A. A direct determination of nuclear distribution in heterokaryons of Aspergillus nidulans | | 185 |
| Morgan, D. H. Suppression of 'purple' in Coprinus lagopus—an anomalous genetic situation | | 195 |
| Ball, C. and Roper, J. A. Studies on the inhibition and mutation of Aspergillus nidulans by acridines | , | 207 |
| GLOVER, S. W. and COLSON, C. Stable and unstable alterations of the host-induced modification properties of <i>Escherichia coli</i> B, K and C | | 223 |
| Cowan, J. W. and Lewis, D. Somatic recombination in the dikaryon of Coprinus lagopus | | 235 |
| DOWRICK, G. J. Breeding systems in tetraploid Rubus species | | 245 |
| Anderson, W. W. Genetic divergence in M. Vetukhiv's experimental populations of <i>Drosophila pseudoobscura</i> | | 255 |
| Fisher, K. W. Mechanically caused damage to Hfr cells of <i>Escherichia</i> coli K12 | | 267 |
| Schell, J. and Glover, S. W. The effect of various physiological conditions on host-controlled restriction in <i>Escherichia coli</i> K(Pl) | | 273 |
| Schell, J. and Glover, S. W. On the localization of a factor responsible for host-controlled restriction in $\textit{Escherichia coli } K(Pl)$ | | 277 |
| Reeve, E. C. R. Characteristics of some single-step mutants to chloram- phenical resistance in <i>Escherichia coli</i> K12 and their interactions with R-factor genes | | 281 |

Contents vii

No. 3 (June 1966)

| DUKE, E. J. Further studies on the inheritance of lymph proteins in Drosophila melanogaster | page | 287 |
|---|------|-----|
| Young, S. S. Y. and Lewontin, R. C. Differences in bristle-making abilities in scute and wild-type <i>Drosophila melanogaster</i> | | 295 |
| Waddington, C. H. and Robertson, E. Selection for developmental canalization | | 303 |
| LATTER, B. D. H. The interaction between effective population size and linkage intensity under artificial selection | | 313 |
| Lamb, B. C. Polarized segregation in Ascomycetes and the differential bursting of asci | | 325 |
| Warr, J. R., McVittie, A., Randall, Sir John and Hopkins, J. M. Genetic control of flagellar structure in <i>Chlamydomonas reinhardii</i> | | 335 |
| Espardellier-Joset, F. and Fournier, R. Induction of invertant males in $E.\ Coli\ \mathrm{K}12$ | i | 353 |
| Deol, M. S. The probable mode of gene action in the circling mutants of the mouse | | 363 |
| Lee, R. M. Genetic control of the hydrolysis of aromatic esters by sheep plasma A-esterase | | 373 |
| Church, R. B. and Robertson, F. W. Biochemical analysis of genetic differences in the growth of <i>Drosophila</i> | | 383 |
| Index of subjects and authors | | 409 |

PLATES should only be included where absolutely essential. They should be supplied as unmounted glossy prints; any lettering to be inserted on them is best indicated on a separate sketch. Please do not damage prints by writing heavily on their backs or by using paper clips.

DIAGRAMS should be about twice the size of the finished block, and the thickness of lines and size of points determined accordingly. They must be drawn in Indian ink on white Bristol Board or tracing linen; graph paper ruled in pale blue (but not other colours) is also acceptable. Lettering should be lightly inserted in soft pencil only, so that the printer can put in the finished lettering.

Legends to illustrations must be given on a separate sheet of paper. Each illustration must have the name of the author and figure number pencilled on the back. Plates and diagrams should be numbered separately and their position indicated on the typescript. It will hasten refereeing if a photograph of each diagram can be supplied with the carbon copy of the paper.

Tables should each be typed on a separate sheet of paper and their approximate positions in the text indicated on the typescript. Each table should be numbered and carry an appropriate title. The table should be designed, whenever possible, to be printed in the normal orientation of the text. The data should be grouped so as to make the use of rules unnecessary. Vertical rules, in particular, are expensive to print, and will only be included at the Editor's discretion.

FOOTNOTES should be avoided where possible. They can often be incorporated into the text, in parentheses.

SYMBOLS. Italic letters should generally be adopted for both gene symbols and quantities in mathematical formulae. Bold letters add to printing costs, and should only be used where they are necessary to avoid confusion.

Spelling should follow the Concise Oxford Dictionary.

REFERENCES should follow the normal usage in the journal. In the list of references at the end of the paper, titles of periodicals should be abbreviated according to the World List of Scientific Periodicals (fourth edition).

PROOFS. Two sets of single-sided page proofs, together with the typescript, of each paper will be sent to the author. The printers' marked proof should be returned after correction to the Executive Editor. Excessive alterations, other than corrections of printers' errors, may be disallowed or charged to the author. Correction should be made using the symbols in British Standard 1219: 1958, or its shortened version B.S. 1219C: 1958.

Ofference. Additional offprints may be ordered on the form sent out with proofs, provided this is returned within seven days of receipt.

VOLUME 7, NUMBER 3, 1966

CONTENTS

| DUKE, E. J. Further studies on the inheritance of lymph proteins in Drosophila melanogaster page | 287 |
|---|-----|
| Young, S. S. Y. and Lewontin, R. C. Differences in bristle-making abilities in scute and wild-type <i>Drosophila melanogaster</i> | 295 |
| Waddington, C. H. and Robertson, E. Selection for developmental canalization | 303 |
| LATTER, B. D. H. The interaction between effective population size and linkage intensity under artificial selection | 313 |
| Lamb, B. C. Polarized segregation in Ascomycetes and the differential bursting of asci | 325 |
| WARR, J. R., McVITTIE, A., RANDALL, SIR JOHN and HOPKINS, J. M. Genetic control of flagellar structure in <i>Chlamydomonas reinhardii</i> | 335 |
| ESPARDELLIER-JOSET, F. and FOURNIER, R. Induction of invertant males in E. Coli K12 | 353 |
| Deol., M. S. The probable mode of gene action in the circling mutants of the mouse | 363 |
| Lee, R. M. Genetic control of the hydrolysis of aromatic esters by sheep plasma A-esterase | 373 |
| Church, R. B. and Robertson, F. W. Biochemical analysis of genetic differences in the growth of <i>Drosophila</i> | 383 |
| Index of subjects and authors | 409 |