

HINTS TO MULTIPLE CHOICE QUESTIONS & EVALUATION TESTS



MHT-CET TRIUMPH MATHEMATICS

MULTIPLE CHOICE QUESTIONS

BASED ON STD. XI & XII SYLLABUS OF MHT-CET

Differential equations are used to determine the age of dead organisms using carbon dating technique.



At death



100% of C-14



5,730 years



50% of C-14



11,460 years



25% of C-14



17,190 years



12.5% of C-14



Target Publications Pvt. Ltd.

MHT-CET **TRIUMPH**
MATHEMATICS
Based on Std. XI & XII Syllabus of MHT-CET

HINTS TO MULTIPLE CHOICE QUESTIONS
&
EVALUATION TESTS

CONTENT

Sr. No.	Textbook Chapter No.	Chapter Name	Page No.
Std. XI			
1	2	Trigonometric Functions	1
2	3	Trigonometric Functions of Compound Angles	13
3	4	Factorization Formulae	41
4	6	Straight Line	54
5	7	Circle and Conics	79
6	1	Sets, Relations and Functions	115
7	4	Sequence and Series	138
8	11	Probability	174
Std. XII			
9	1	Mathematical Logic	196
10	2	Matrices	203
11	3	Trigonometric Functions	219
12	4	Pair of Straight Lines	270
13	5	Vectors	292
14	6	Three Dimensional Geometry	309
15	7	Line	320
16	8	Plane	338
17	9	Linear Programming	366
18	1	Continuity	383
19	2	Differentiation	405
20	3	Applications of Derivatives	458
21	4	Integration	504
22	5	Definite Integrals	558
23	6	Applications of Definite Integral	606
24	7	Differential Equations	630
25	8	Probability Distribution	671
26	9	Binomial Distribution	683



Hints



Classical Thinking

$$1. \quad \tan \theta = \frac{9}{2}$$

$$\Rightarrow \frac{\sin \theta}{\cos \theta} = \frac{9}{2}$$

$$\Rightarrow \cos \theta = \frac{2}{9} \sin \theta = \frac{2}{9} \times \frac{3}{4} = \frac{1}{6}$$

$$2. \quad 5 \sin \theta = 3 \Rightarrow \sin \theta = \frac{3}{5}$$

$$\frac{\sec \theta + \tan \theta}{\sec \theta - \tan \theta} = \frac{1 + \sin \theta}{1 - \sin \theta}$$

$$= \frac{1 + \frac{3}{5}}{1 - \frac{3}{5}}$$

$$= 4$$

$$3. \quad \frac{\sin \theta}{1 - \cot \theta} + \frac{\cos \theta}{1 - \tan \theta} = \frac{\sin \theta \cdot \sin \theta}{\sin \theta - \cos \theta} + \frac{\cos \theta \cdot \cos \theta}{\cos \theta - \sin \theta}$$

$$= \frac{\cos^2 \theta - \sin^2 \theta}{\cos \theta - \sin \theta}$$

$$= \cos \theta + \sin \theta$$

$$4. \quad \sin \theta = -\frac{1}{\sqrt{2}} \text{ and } \tan \theta = 1$$

Since, $\sin \theta$ is -ve and $\tan \theta$ is +ve in third quadrant,

$\therefore \theta$ lies in the IIIrd quadrant.

5. Since, $\sin \theta$ is -ve and $\cos \theta$ is +ve

$\therefore \theta$ lies in IVth quadrant.

$$7. \quad \sin^2 \frac{\pi}{6} + \cos^2 \frac{\pi}{3} - \tan^2 \frac{\pi}{4} = \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^2 - (1)^2$$

$$= \frac{1}{4} + \frac{1}{4} - 1 = -\frac{1}{2}$$

$$8. \quad x \sin 45^\circ \cos^2 60^\circ = \frac{\tan^2 60^\circ \operatorname{cosec} 30^\circ}{\sec 45^\circ \cot^2 30^\circ}$$

$$\Rightarrow x \cdot \frac{1}{\sqrt{2}} \cdot \frac{1}{4} = \frac{3 \cdot 2}{\sqrt{2} \cdot 3}$$

$$\Rightarrow \frac{x}{4\sqrt{2}} = \sqrt{2}$$

$$\Rightarrow x = 8$$

$$9. \quad \sin \theta = \sqrt{3} \cos \theta$$

$$\Rightarrow \frac{\sin \theta}{\cos \theta} = \sqrt{3}$$

$$\Rightarrow \tan \theta = \sqrt{3} \Rightarrow \theta = 60^\circ$$

$$10. \quad \sin(\alpha - \beta) = \frac{1}{2} = \sin 30^\circ$$

$$\Rightarrow \alpha - \beta = 30^\circ \quad \dots (i)$$

and $\cos(\alpha + \beta) = \frac{1}{2} = \cos 60^\circ$

$$\Rightarrow \alpha + \beta = 60^\circ \quad \dots (ii)$$

On solving (i) and (ii), we get
 $\alpha = 45^\circ$ and $\beta = 15^\circ$

$$11. \quad \tan \theta = \frac{20}{21}$$

Since $1 + \tan^2 \theta = \sec^2 \theta$

$$\therefore \sec^2 \theta = 1 + \frac{400}{441} = \frac{841}{441}$$

$$\Rightarrow \sec \theta = \pm \frac{29}{21}$$

$$\Rightarrow \cos \theta = \pm \frac{21}{29}$$

$$12. \quad 1 + \tan^2 \theta = \sec^2 \theta$$

$$\Rightarrow 1 + \frac{1}{10} = \sec^2 \theta$$

$$\Rightarrow \sec^2 \theta = \frac{11}{10} \Rightarrow \sec \theta = \sqrt{\frac{11}{10}}$$

.... [$\because \theta$ lies in the fourth quadrant]

$$13. \quad \sec^2 \theta = 1 + \tan^2 \theta = 1 + \frac{1}{5} = \frac{6}{5}$$

$$\Rightarrow \cos^2 \theta = \frac{5}{6}$$

$$\Rightarrow \cos \theta = \frac{\sqrt{5}}{\sqrt{6}} \quad \dots [\because \theta \text{ lies in the 1st quadrant}]$$



$$14. \quad \sec \theta + \tan \theta = \frac{1 + \sin \theta}{\cos \theta}$$

$$= \frac{1 + \sin \theta}{-\sqrt{1 - \sin^2 \theta}}$$

...[$\because \theta$ lies in the second quadrant, $\therefore \cos \theta < 0$]

$$= \frac{-\left(1 + \frac{21}{29}\right)}{\sqrt{1 - \left(\frac{21}{29}\right)^2}} = -\frac{5}{2}$$

$$15. \quad \sec^4 x - \sec^2 x = \sec^2 x (\sec^2 x - 1)$$

$$= (1 + \tan^2 x) \tan^2 x$$

$$= \tan^2 x + \tan^4 x$$

$$16. \quad \tan^2 \theta - \sin^2 \theta = \sin^2 \theta \left(\frac{1}{\cos^2 \theta} - 1 \right)$$

$$= \sin^2 \theta (\sec^2 \theta - 1)$$

$$= \sin^2 \theta \tan^2 \theta$$

Also $\sec^2 \theta \operatorname{cosec}^2 \theta \neq \sec^2 \theta - \operatorname{cosec}^2 \theta$, and
 $\operatorname{cosec}^2 \theta + \cot^2 \theta \neq \operatorname{cosec}^2 \theta \cot^2 \theta$

$$17. \quad x = \sec \theta + \tan \theta$$

$$\therefore x + \frac{1}{x} = \sec \theta + \tan \theta + \frac{1}{\sec \theta + \tan \theta}$$

$$= \sec \theta + \tan \theta + \frac{\sec \theta - \tan \theta}{\sec^2 \theta - \tan^2 \theta}$$

$$= \sec \theta + \tan \theta + \sec \theta - \tan \theta$$

...[$\because \sec^2 \theta - \tan^2 \theta = 1$]

$$= 2 \sec \theta$$

$$18. \quad \cot x + \tan x = \frac{\cos x}{\sin x} + \frac{\sin x}{\cos x}$$

$$= \frac{\cos^2 x + \sin^2 x}{\sin x \cos x}$$

$$= \frac{1}{\sin x \cos x}$$

$$= \sec x \operatorname{cosec} x$$

$$19. \quad \frac{\sin^2 20^\circ + \cos^4 20^\circ}{\sin^4 20^\circ + \cos^2 20^\circ}$$

$$= \frac{\sin^2 20^\circ + \cos^2 20^\circ (1 - \sin^2 20^\circ)}{\sin^2 20^\circ (1 - \cos^2 20^\circ) + \cos^2 20^\circ}$$

$$= \frac{1 - \sin^2 20^\circ \cos^2 20^\circ}{1 - \sin^2 20^\circ \cos^2 20^\circ} = 1$$

$$20. \quad x = a \cos \theta + b \sin \theta \quad \dots(i)$$

$$\text{and } y = a \sin \theta - b \cos \theta \quad \dots(ii)$$

Squaring (i) and (ii) and adding, we get
 $x^2 + y^2 = a^2 \cos^2 \theta + b^2 \sin^2 \theta + 2ab \cos \theta \sin \theta$
 $+ a^2 \sin^2 \theta + b^2 \cos^2 \theta - 2ab \sin \theta \cos \theta$
 $\Rightarrow x^2 + y^2 = a^2 + b^2$

$$21. \quad x = a \cos^3 \theta$$

$$\Rightarrow \left(\frac{x}{a}\right)^{\frac{1}{3}} = \cos \theta, \text{ and} \quad \dots(i)$$

$$y = b \sin^3 \theta$$

$$\Rightarrow \left(\frac{y}{b}\right)^{\frac{1}{3}} = \sin \theta \quad \dots(ii)$$

Squaring (i) and (ii) and adding, we get

$$\left(\frac{x}{a}\right)^{\frac{2}{3}} + \left(\frac{y}{b}\right)^{\frac{2}{3}} = \cos^2 \theta + \sin^2 \theta$$

$$= 1$$

$$22. \quad \cos x + \cos^2 x = 1$$

$$\Rightarrow \cos x = \sin^2 x \quad \dots[\because 1 - \cos^2 x = \sin^2 x]$$

$$\therefore \sin^2 x + \sin^4 x = \cos x + \cos^2 x = 1$$

$$23. \quad \sin x + \sin^2 x = 1$$

$$\Rightarrow \sin x = \cos^2 x$$

$$\therefore \cos^8 x + 2 \cos^6 x + \cos^4 x$$

$$= \sin^4 x + 2 \sin^3 x + \sin^2 x$$

$$= (\sin x + \sin^2 x)^2$$

$$= (1)^2 = 1$$

$$24. \quad \sec \theta = \frac{1}{2} \text{ is not possible as } |\sec \theta| \geq 1$$

25. $\tan \theta$ can have any value
 $\sin \theta$ and $\cos \theta$ cannot be numerically greater than 1.

$\sec \theta$ should be greater than 1.

\therefore Option (D) is the correct answer.

$$26. \quad \text{Since, } -1 \leq \cos \theta \leq 1$$

$$\therefore -5 \leq 5 \cos \theta \leq 5$$

$$\Rightarrow -5 + 12 \leq 5 \cos \theta + 12 \leq 5 + 12$$

$$\Rightarrow 7 \leq 5 \cos \theta + 12 \leq 17$$



Critical Thinking

$$1. \quad \frac{p \sin \theta - q \cos \theta}{p \sin \theta + q \cos \theta} = \frac{p \frac{\sin \theta}{\cos \theta} - q}{p \frac{\sin \theta}{\cos \theta} + q}$$

$$= \frac{p \tan \theta - q}{p \tan \theta + q}$$

$$= \frac{p^2 - q^2}{p^2 + q^2} \quad \dots \left[\because \tan \theta = \frac{p}{q} (\text{given}) \right]$$



$$2. \quad \cos^2 \theta + \sec^2 \theta = (\cos \theta - \sec \theta)^2 + 2 \geq 2$$

$$3. \quad \sin x + \operatorname{cosec} x = 2$$

$$\Rightarrow \sin^2 x + 1 = 2 \sin x$$

$$\Rightarrow (\sin x - 1)^2 = 0 \Rightarrow \sin x = 1$$

$$\begin{aligned} \therefore \sin^n x + \operatorname{cosec}^n x &= \sin^n x + \frac{1}{\sin^n x} \\ &= (1)^n + \frac{1}{(1)^n} = 2 \end{aligned}$$

$$4. \quad \text{Since, } 1 \text{ radian} = 57^\circ \text{ nearly and } \sin 57^\circ > \sin 1^\circ$$

$$\therefore \sin 1 > \sin 1^\circ$$

$$5. \quad \text{Since, } 1 \text{ radian} = 57^\circ \text{ nearly}$$

$$\therefore 2 \text{ radians} = 114^\circ \text{ nearly}$$

Since, 57° lies in Ist quadrant and 114° lies in IInd quadrant.

$$\therefore \tan 1 > 0 \text{ and } \tan 2 < 0$$

$$\therefore \tan 1 > \tan 2$$

$$6. \quad \cos A = \frac{\sqrt{3}}{2}$$

$$\Rightarrow \cos A = \cos 30^\circ$$

$$\Rightarrow A = 30^\circ$$

$$\therefore \tan 3A = \tan 90^\circ = \infty$$

$$7. \quad \tan(A - B) = 1 = \tan \frac{\pi}{4}$$

$$\Rightarrow A - B = \frac{\pi}{4} \quad \dots \text{(i)}$$

$$\text{and } \sec(A + B) = \frac{2}{\sqrt{3}}$$

$$\Rightarrow A + B = \frac{11\pi}{6} \quad \dots \text{(ii)}$$

From (i) and (ii), we get

$$B = \frac{19\pi}{24}$$

$$8. \quad \sin(A + B + C) = 1$$

$$\Rightarrow A + B + C = 90^\circ \quad \dots \text{(i)}$$

$$\tan(A - B) = \frac{1}{\sqrt{3}}$$

$$\Rightarrow A - B = 30^\circ \quad \dots \text{(ii)}$$

$$\sec(A + C) = 2$$

$$\Rightarrow A + C = 60^\circ \quad \dots \text{(iii)}$$

From (i), (ii) and (iii), we get

$$B = 30^\circ, A = 60^\circ, C = 0^\circ$$

$$9. \quad \cos A = \frac{3}{5} \text{ and } \cos B = \frac{4}{5}$$

Both A and B lie in the fourth quadrant.

Hence, both $\sin A$ and $\sin B$ are negative.

$$\begin{aligned} \therefore 2 \sin A + 4 \sin B &= -2\sqrt{1 - \cos^2 A} - 4\sqrt{1 - \cos^2 B} \\ &= -2\sqrt{1 - \frac{9}{25}} - 4\sqrt{1 - \frac{16}{25}} = -4 \end{aligned}$$

$$10. \quad \sec \theta + \tan \theta = \sqrt{3} \quad \dots \text{(i)}$$

$$\Rightarrow \sec \theta - \tan \theta = \frac{1}{\sqrt{3}} \quad \dots \text{(ii)}$$

$$\dots [\because \sec^2 \theta - \tan^2 \theta = 1]$$

Subtracting (ii) from (i), we get

$$2 \tan \theta = \sqrt{3} - \frac{1}{\sqrt{3}}$$

$$\Rightarrow \tan \theta = \frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{\pi}{6}$$

$$11. \quad \sqrt{\frac{1 - \sin \theta}{1 + \sin \theta}} = \sqrt{\frac{(1 - \sin \theta)^2}{1 - \sin^2 \theta}}$$

$$= \frac{1 - \sin \theta}{|\cos \theta|}$$

$$= \frac{1 - \sin \theta}{-\cos \theta} \quad \dots \left[\because \frac{\pi}{2} < \theta < \frac{3\pi}{2} \Rightarrow \cos \theta < 0 \right]$$

$$= -\sec \theta + \tan \theta$$

$$12. \quad \sqrt{\frac{1 - \sin \theta}{1 + \sin \theta}} + \sqrt{\frac{1 + \sin \theta}{1 - \sin \theta}}$$

$$= \frac{1 - \sin \theta + 1 + \sin \theta}{\sqrt{1 - \sin^2 \theta}} = \frac{2}{\sqrt{\cos^2 \theta}}$$

$$= \frac{2}{-\cos \theta} \quad \dots [\because \theta \text{ lies in the } 2^{\text{nd}} \text{ quadrant}]$$

$$= -2 \sec \theta$$

$$13. \quad \sqrt{\frac{1 - \cos \alpha}{1 + \cos \alpha}} + \sqrt{\frac{1 + \cos \alpha}{1 - \cos \alpha}}$$

$$= \frac{1 - \cos \alpha + 1 + \cos \alpha}{\sqrt{1 - \cos^2 \alpha}}$$

$$= \frac{2}{-\sin \alpha} \quad \dots \left[\because \pi < \alpha < \frac{3\pi}{2} \right]$$

$$14. \quad 3 \tan A + 4 = 0 \Rightarrow \tan A = -\frac{4}{3}$$

$$1 + \tan^2 A = \sec^2 A$$

$$\Rightarrow \sec^2 A = 1 + \frac{16}{9}$$



$$\Rightarrow \sec A = \frac{-5}{3} \quad \dots[\because \theta \text{ lies in } 2^{\text{nd}} \text{ quadrant}]$$

$$\Rightarrow \cos A = -\frac{3}{5}$$

$$\begin{aligned} \therefore \sin A &= \sqrt{1 - \cos^2 A} = \sqrt{1 - \frac{9}{25}} \\ &= \frac{4}{5} \quad \dots[\because A \text{ lies in } 2^{\text{nd}} \text{ quadrant}] \end{aligned}$$

$$\begin{aligned} \therefore 2 \cot A - 5 \cos A + \sin A \\ &= 2\left(-\frac{3}{4}\right) - 5\left(-\frac{3}{5}\right) + \frac{4}{5} \\ &= \frac{23}{10} \end{aligned}$$

$$\begin{aligned} 15. \quad \sec \theta - \tan \theta &= \frac{1}{2} \quad \dots\text{(i)} \\ \Rightarrow \sec \theta + \tan \theta &= 2 \quad \dots\text{(ii)} \\ &\dots[\because \sec^2 \theta - \tan^2 \theta = 1] \end{aligned}$$

Adding (i) and (ii), we get

$$2 \sec \theta = \frac{5}{2} \Rightarrow \sec \theta = \frac{5}{4}$$

Subtracting (ii) from (i), we get

$$2 \tan \theta = \frac{3}{2} \Rightarrow \tan \theta = \frac{3}{4}$$

Since, both $\sec \theta$ and $\tan \theta$ are positive,
 θ lies in the first quadrant.

$$\begin{aligned} 16. \quad \cos \theta + \sin \theta &= \sqrt{2} \cos \theta \\ \text{Squaring both sides, we get} \\ \therefore \cos^2 \theta + \sin^2 \theta + 2 \sin \theta \cos \theta &= 2 \cos^2 \theta \\ \Rightarrow 2 \sin \theta \cos \theta &= 2 \cos^2 \theta - 1 \quad \dots\text{(i)} \end{aligned}$$

Now

$$\begin{aligned} (\cos \theta - \sin \theta)^2 &= \cos^2 \theta + \sin^2 \theta - 2 \sin \theta \cos \theta \\ &= 1 - (2 \cos^2 \theta - 1) \end{aligned}$$

\dots [From (i)]

$$\begin{aligned} &= 2(1 - \cos^2 \theta) \\ &= 2 \sin^2 \theta \end{aligned}$$

$$\therefore \cos \theta - \sin \theta = \sqrt{2} \sin \theta$$

$$\begin{aligned} 17. \quad (\sin x - \cos x)^2 &= \sin^2 x + \cos^2 x - 2 \sin x \cos x \\ &= 1 - \{(\sin x + \cos x)^2 - (\sin^2 x + \cos^2 x)\} \\ &= 1 - (a^2 - 1) \quad \dots[\because \sin x + \cos x = a] \\ &= 2 - a^2 \end{aligned}$$

$$\therefore |\sin x - \cos x| = \sqrt{2 - a^2}$$

$$\begin{aligned} 18. \quad 3 \sin \theta + 4 \cos \theta &= 5 \\ \text{Squaring both sides, we get} \\ 9 \sin^2 \theta + 16 \cos^2 \theta + 24 \sin \theta \cos \theta &= 25 \end{aligned}$$

$$\begin{aligned} &\Rightarrow 9(1 - \cos^2 \theta) + 16(1 - \sin^2 \theta) \\ &\quad + 24 \sin \theta \cos \theta = 25 \\ &\Rightarrow 9 \cos^2 \theta + 16 \sin^2 \theta - 24 \sin \theta \cos \theta = 0 \\ &\Rightarrow (3 \cos \theta - 4 \sin \theta)^2 = 0 \\ &\Rightarrow 3 \cos \theta - 4 \sin \theta = 0 \end{aligned}$$

$$\begin{aligned} 19. \quad 2u_6 - 3u_4 \\ &= 2(\cos^6 \theta + \sin^6 \theta) - 3(\cos^4 \theta + \sin^4 \theta) \\ &= 2(1 - 3 \sin^2 \theta \cos^2 \theta) - 3(1 - 2 \sin^2 \theta \cos^2 \theta) \\ &= -1 \end{aligned}$$

$$\begin{aligned} 20. \quad \sin x + \sin^2 x &= 1 \\ \Rightarrow \sin x &= 1 - \sin^2 x \\ \Rightarrow \sin x &= \cos^2 x \end{aligned}$$

$$\begin{aligned} \therefore \cos^{12} x + 3 \cos^{10} x + 3 \cos^8 x + \cos^6 x - 2 \\ &= \sin^6 x + 3 \sin^5 x + 3 \sin^4 x + \sin^3 x - 2 \\ &= (\sin^2 x)^3 + 3(\sin^2 x)^2 \sin x + 3(\sin^2 x)(\sin x)^2 \\ &\quad + (\sin x)^3 - 2 \\ &= (\sin^2 x + \sin x)^3 - 2 \\ &= (1)^3 - 2 \quad \dots[\because \sin x + \sin^2 x = 1] \\ &= -1 \end{aligned}$$

$$\begin{aligned} 21. \quad 10 \sin^4 \alpha + 15 \cos^4 \alpha &= 6 \\ \Rightarrow 10 \sin^4 \alpha + 15 \cos^4 \alpha &= 6(\sin^2 \alpha + \cos^2 \alpha)^2 \\ \Rightarrow 10 \tan^4 \alpha + 15 &= 6(\tan^2 \alpha + 1)^2 \\ \Rightarrow (2 \tan^2 \alpha - 3)^2 &= 0 \\ \Rightarrow \tan^2 \alpha &= \frac{3}{2} \end{aligned}$$

$$\begin{aligned} \therefore 27 \operatorname{cosec}^6 \alpha + 8 \sec^6 \alpha \\ &= 27(1 + \cot^2 \alpha)^3 + 8(1 + \tan^2 \alpha)^3 \\ &= 27\left(1 + \frac{2}{3}\right)^3 + 8\left(1 + \frac{3}{2}\right)^3 = 250 \end{aligned}$$

$$\begin{aligned} 22. \quad \sec \alpha - \tan \alpha &= \frac{1 - \sin \alpha}{\cos \alpha} \\ &= \frac{1 - \frac{2pq}{p^2 + q^2}}{\sqrt{1 - \left(\frac{2pq}{p^2 + q^2}\right)^2}} \\ &= \frac{(p - q)^2}{\sqrt{(p^2 - q^2)^2}} \\ &= \frac{(p - q)^2}{p^2 - q^2} = \frac{p - q}{p + q} \end{aligned}$$



$$23. \quad \sec \theta - \tan \theta = \frac{a+1}{a-1} \quad \dots\text{(i)}$$

$$\Rightarrow \sec \theta + \tan \theta = \frac{a-1}{a+1} \quad \dots\text{(ii)}$$

$$\dots[\because \sec^2 \theta - \tan^2 \theta = 1]$$

Adding (i) and (ii), we get

$$2 \sec \theta = \frac{a+1}{a-1} + \frac{a-1}{a+1}$$

$$\Rightarrow 2 \sec \theta = \frac{(a+1)^2 + (a-1)^2}{a^2 - 1}$$

$$\Rightarrow 2 \sec \theta = \frac{2(a^2 + 1)}{a^2 - 1}$$

$$\Rightarrow \sec \theta = \frac{a^2 + 1}{a^2 - 1}$$

$$\Rightarrow \cos \theta = \frac{a^2 - 1}{a^2 + 1}$$

$$24. \quad \tan^2 \theta = \sec^2 \theta - 1$$

$$= \left(x + \frac{1}{4x}\right)^2 - 1 = \left(x - \frac{1}{4x}\right)^2$$

$$\Rightarrow \tan \theta = \pm \left(x - \frac{1}{4x}\right)$$

$$\Rightarrow \sec \theta + \tan \theta = x + \frac{1}{4x} \pm \left(x - \frac{1}{4x}\right)$$

$$\Rightarrow \sec \theta + \tan \theta = x + \frac{1}{4x} + \left(x - \frac{1}{4x}\right)$$

or

$$\sec \theta + \tan \theta = x + \frac{1}{4x} - \left(x - \frac{1}{4x}\right)$$

$$\Rightarrow \sec \theta + \tan \theta = 2x \text{ or } \frac{1}{2x}$$

$$25. \quad \sin^6\left(\frac{\pi}{49}\right) + \cos^6\left(\frac{\pi}{49}\right) - 1 + 3\sin^2\left(\frac{\pi}{49}\right)\cos^2\left(\frac{\pi}{49}\right)$$

$$= \sin^6\left(\frac{\pi}{49}\right) + \cos^6\left(\frac{\pi}{49}\right)$$

$$+ 3\sin^2\left(\frac{\pi}{49}\right)\cos^2\left(\frac{\pi}{49}\right)\left(\sin^2\frac{\pi}{49} + \cos^2\frac{\pi}{49}\right) - 1$$

$$= \left(\sin^2\frac{\pi}{49} + \cos^2\frac{\pi}{49}\right)^3 - 1 = 1 - 1 = 0$$

$$26. \quad \frac{1 - \cos \alpha + \sin \alpha}{1 + \sin \alpha} = \frac{1 - \cos \alpha + \sin \alpha}{1 + \sin \alpha} \cdot \frac{1 + \cos \alpha + \sin \alpha}{1 + \cos \alpha + \sin \alpha}$$

$$= \frac{(1 + \sin \alpha)^2 - \cos^2 \alpha}{(1 + \sin \alpha)(1 + \cos \alpha + \sin \alpha)}$$

$$= \frac{1 + 2\sin \alpha + \sin^2 \alpha - (1 - \sin^2 \alpha)}{(1 + \sin \alpha)(1 + \cos \alpha + \sin \alpha)}$$

$$= \frac{2\sin \alpha(1 + \sin \alpha)}{(1 + \sin \alpha)(1 + \cos \alpha + \sin \alpha)}$$

$$= \frac{2\sin \alpha}{1 + \cos \alpha + \sin \alpha}$$

$$= x$$

$$27. \quad 1 - \frac{\sin^2 y}{1 + \cos y} + \frac{1 + \cos y}{\sin y} - \frac{\sin y}{1 - \cos y}$$

$$= \frac{1 + \cos y - \sin^2 y}{1 + \cos y} + \frac{(1 - \cos^2 y) - \sin^2 y}{\sin y(1 - \cos y)}$$

$$= \frac{\cos^2 y + \cos y}{1 + \cos y} + \frac{\sin^2 y - \sin^2 y}{\sin y(1 - \cos y)}$$

$$= \frac{\cos y(1 + \cos y)}{1 + \cos y} + 0 = \cos y$$

$$28. \quad \frac{2\sin \theta \tan \theta(1 - \tan \theta) + 2\sin \theta \sec^2 \theta}{(1 + \tan \theta)^2}$$

$$= \frac{2\sin \theta}{(1 + \tan \theta)^2} \{\tan \theta(1 - \tan \theta) + \sec^2 \theta\}$$

$$= \frac{2\sin \theta}{(1 + \tan \theta)^2} (\tan \theta - \tan^2 \theta + 1 + \tan^2 \theta)$$

$$= \frac{2\sin \theta}{1 + \tan \theta}$$

$$29. \quad \frac{\sin^3 A - \cos^3 A}{\sin A - \cos A} + \frac{\sin A}{\sqrt{1 + \tan^2 A}} - 2 \tan A \cot A$$

$$= (\sin^2 A + \cos^2 A + \sin A \cos A) + \frac{\sin A}{|\sec A|} - 2$$

$$= 1 + \sin A \cos A - \sin A \cos A - 2$$

$$\dots[\because A \text{ is an obtuse angle} \Rightarrow \cos A < 0]$$

$$= -1$$

$$30. \quad \frac{x}{a} \cos \theta + \frac{y}{b} \sin \theta = -1 \quad \dots\text{(i)}$$

$$\text{and } \frac{x}{a} \sin \theta - \frac{y}{b} \cos \theta = 1 \quad \dots\text{(ii)}$$



Squaring (i) and (ii) and adding, we get

$$\frac{x^2}{a^2}(\sin^2 \theta + \cos^2 \theta) + \frac{y^2}{b^2}(\sin^2 \theta + \cos^2 \theta) = 2$$

$$\Rightarrow \frac{x^2}{a^2} + \frac{y^2}{b^2} = 2$$

$$31. \quad x \sin^3 \alpha + y \cos^3 \alpha = \sin \alpha \cos \alpha \quad \dots(i)$$

$$\text{and } x \sin \alpha - y \cos \alpha = 0$$

$$\Rightarrow x \sin \alpha = y \cos \alpha \quad \dots(ii)$$

\(\therefore\) From (i) and (ii), we get

$$y \cos \alpha \sin^2 \alpha + y \cos^3 \alpha = \sin \alpha \cos \alpha$$

$$\Rightarrow y \cos \alpha (\sin^2 \alpha + \cos^2 \alpha) = \sin \alpha \cos \alpha$$

$$\Rightarrow y \cos \alpha = \sin \alpha \cos \alpha$$

$$\Rightarrow y = \sin \alpha$$

$$\therefore x = \cos \alpha$$

$$\therefore x^2 + y^2 = \cos^2 \alpha + \sin^2 \alpha = 1$$

$$32. \quad m + n = a \cos^3 \alpha + 3a \cos \alpha \sin^2 \alpha \\ + 3a \cos^2 \alpha \sin \alpha + a \sin^3 \alpha \\ = a (\cos \alpha + \sin \alpha)^3$$

$$\text{Similarly, } (m - n) = a (\cos \alpha - \sin \alpha)^3$$

$$\therefore (m + n)^{2/3} + (m - n)^{2/3} \\ = a^{2/3} \{(\cos \alpha + \sin \alpha)^2 + (\cos \alpha - \sin \alpha)^2\} \\ = a^{2/3} \{2(\cos^2 \alpha + \sin^2 \alpha)\} = 2a^{2/3}$$

$$33. \quad \sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma \\ = \frac{\tan^2 \alpha}{1 + \tan^2 \alpha} + \frac{\tan^2 \beta}{1 + \tan^2 \beta} + \frac{\tan^2 \gamma}{1 + \tan^2 \gamma} \\ = \frac{x}{1+x} + \frac{y}{1+y} + \frac{z}{1+z} \\ \dots[\text{Let } x = \tan^2 \alpha, y = \tan^2 \beta, z = \tan^2 \gamma] \\ = \frac{(x+y+z) + (xy+yz+zx+2xyz) + xy+yz+zx+xyz}{(1+x)(1+y)(1+z)} \\ = \frac{x+y+z+1+xy+yz+zx+xyz}{(1+x)(1+y)(1+z)} = 1$$

$$34. \quad p + q = \frac{2 \sin \theta}{1 + \sin \theta + \cos \theta} + \frac{\cos \theta}{1 + \sin \theta} \\ = \frac{2 \sin \theta(1 + \sin \theta) + \cos \theta(1 + \sin \theta + \cos \theta)}{(1 + \sin \theta + \cos \theta)(1 + \sin \theta)} \\ = \frac{2 \sin \theta + 2 \sin^2 \theta + \cos \theta + \cos \theta \sin \theta + \cos^2 \theta}{1 + 2 \sin \theta + \sin^2 \theta + \cos \theta + \cos \theta \sin \theta} \\ = \frac{2 \sin \theta + \sin^2 \theta + (\sin^2 \theta + \cos^2 \theta) + \cos \theta + \cos \theta \sin \theta}{1 + 2 \sin \theta + \sin^2 \theta + \cos \theta + \cos \theta \sin \theta} \\ \Rightarrow p + q = 1$$

$$35. \quad xy = (\sec \phi - \tan \phi)(\operatorname{cosec} \phi + \cot \phi) \\ = \frac{1 - \sin \phi}{\cos \phi} \cdot \frac{1 + \cos \phi}{\sin \phi}$$

$$xy + 1 = \frac{1 - \sin \phi + \cos \phi - \sin \phi \cos \phi + \sin \phi \cos \phi}{\cos \phi \sin \phi}$$

$$\Rightarrow xy + 1 = \frac{1 - \sin \phi + \cos \phi}{\cos \phi \sin \phi} \quad \dots(i)$$

$$x - y = (\sec \phi - \tan \phi) - (\operatorname{cosec} \phi + \cot \phi)$$

$$= \frac{1 - \sin \phi}{\cos \phi} - \frac{1 + \cos \phi}{\sin \phi}$$

$$= \frac{\sin \phi - \sin^2 \phi - \cos \phi - \cos^2 \phi}{\cos \phi \sin \phi}$$

$$= \frac{\sin \phi - \cos \phi - (\sin^2 \phi + \cos^2 \phi)}{\cos \phi \sin \phi}$$

$$\Rightarrow x - y = \frac{\sin \phi - \cos \phi - 1}{\cos \phi \sin \phi} \quad \dots(ii)$$

Adding (i) and (ii), we get

$$xy + 1 + (x - y) = 0$$

$$\Rightarrow x = \frac{y - 1}{y + 1}$$

$$36. \quad \sin \theta_1 + \sin \theta_2 + \sin \theta_3 = 3 \\ \Rightarrow \sin \theta_1 = \sin \theta_2 = \sin \theta_3 = 1 \\ \dots[\because -1 \leq \sin x \leq 1]$$

$$\Rightarrow \theta_1 = \theta_2 = \theta_3 = \frac{\pi}{2}$$

$$\Rightarrow \cos \theta_1 + \cos \theta_2 + \cos \theta_3 = 0$$

$$37. \quad \text{Given, } (a + b)^2 = 4ab \sin^2 \theta$$

$$\Rightarrow \sin^2 \theta = \frac{(a + b)^2}{4ab} \leq 1$$

$$\Rightarrow (a + b)^2 - 4ab \leq 0$$

$$\Rightarrow (a - b)^2 \leq 0$$

$$\Rightarrow a = b$$

$$38. \quad 12 \sin \theta - 9 \sin^2 \theta = -9 \left(\sin^2 \theta - \frac{12}{9} \sin \theta \right)$$

$$= -9 \left(\sin^2 \theta - \frac{4}{3} \sin \theta \right)$$

$$= -9 \left[\sin^2 \theta - \frac{4}{3} \sin \theta + \left(\frac{2}{3} \right)^2 - \left(\frac{2}{3} \right)^2 \right]$$

$$= -9 \left(\sin \theta - \frac{2}{3} \right)^2 + 9 \times \frac{4}{9} \leq 4$$



$$\begin{aligned}
 39. \quad y &= \sin^2 \theta + \cos^4 \theta \\
 &\Rightarrow y = \cos^4 \theta - \cos^2 \theta + 1 \\
 &\Rightarrow y = \left(\cos^2 \theta - \frac{1}{2} \right)^2 + \frac{3}{4} \\
 \text{Now, } 0 &\leq \cos^2 \theta \leq 1 \\
 &\Rightarrow -\frac{1}{2} \leq \cos^2 \theta - \frac{1}{2} \leq \frac{1}{2} \\
 &\Rightarrow 0 \leq \left(\cos^2 \theta - \frac{1}{2} \right)^2 \leq \frac{1}{4} \\
 &\Rightarrow \frac{3}{4} \leq \left(\cos^2 \theta - \frac{1}{2} \right)^2 + \frac{3}{4} \leq 1 \\
 &\Rightarrow \frac{3}{4} \leq y \leq 1
 \end{aligned}$$

**Competitive Thinking**

$$\begin{aligned}
 1. \quad \frac{5 \sin \theta - 3 \cos \theta}{5 \sin \theta + 2 \cos \theta} &= \frac{5 \tan \theta - 3}{5 \tan \theta + 2} \\
 &= \frac{4 - 3}{4 + 2} \\
 &\quad \dots [\because 5 \tan \theta = 4 \text{ (given)}] \\
 &= \frac{1}{6}
 \end{aligned}$$

$$\begin{aligned}
 2. \quad \sin^2 \theta + \operatorname{cosec}^2 \theta &= (\sin \theta + \operatorname{cosec} \theta)^2 - 2 \sin \theta \operatorname{cosec} \theta \\
 &= (2)^2 - 2 \quad \dots [\because \sin \theta + \operatorname{cosec} \theta = 2 \text{ (given)}] \\
 &= 4 - 2 = 2
 \end{aligned}$$

$$\begin{aligned}
 3. \quad \sin \theta + \operatorname{cosec} \theta &= 2 \\
 \Rightarrow \sin^2 \theta + 1 &= 2 \sin \theta \quad \dots \left[\because \operatorname{cosec} \theta = \frac{1}{\sin \theta} \right] \\
 \Rightarrow \sin^2 \theta - 2 \sin \theta + 1 &= 0 \\
 \Rightarrow (\sin \theta - 1)^2 &= 0 \\
 \Rightarrow \sin \theta &= 1
 \end{aligned}$$

$$\begin{aligned}
 \therefore \sin^{10} \theta + \operatorname{cosec}^{10} \theta &= \sin^{10} \theta + \frac{1}{\sin^{10} \theta} \\
 &= (1)^{10} + \frac{1}{(1)^{10}} \\
 &= 2
 \end{aligned}$$

$$\begin{aligned}
 4. \quad \tan A + \cot A &= 4 \\
 \text{Squaring both sides, we get} \\
 \tan^2 A + \cot^2 A + 2 \tan A \cot A &= 16 \\
 \Rightarrow \tan^2 A + \cot^2 A &= 14 \\
 \text{Again, squaring both sides, we get} \\
 \tan^4 A + \cot^4 A + 2 &= 196 \\
 \Rightarrow \tan^4 A + \cot^4 A &= 194
 \end{aligned}$$

$$\begin{aligned}
 5. \quad &\text{Since, } 200^\circ \text{ lies in III}^{\text{rd}} \text{ quadrant.} \\
 &\therefore \sin 200^\circ, \cos 200^\circ \text{ are both -ve.} \\
 &\therefore \text{their sum is -ve.} \\
 6. \quad &\text{One of the factor of the given expression is} \\
 &\cos 90^\circ = 0. \\
 &\therefore \cos 1^\circ \cdot \cos 2^\circ \cdot \cos 3^\circ \dots \cos 179^\circ = 0 \\
 7. \quad &\text{Given that } x \in \left[0, \frac{\pi}{2} \right], y \in \left[0, \frac{\pi}{2} \right] \\
 &\sin x + \cos y = 2 \\
 &\text{Maximum value of } \sin x = 1 \Rightarrow x = \frac{\pi}{2} \\
 &\text{Maximum value of } \cos y = 1 \Rightarrow y = 0 \\
 &\therefore x + y = \frac{\pi}{2} + 0 = \frac{\pi}{2}
 \end{aligned}$$

$$\begin{aligned}
 58. \quad \sin \theta - \cos \theta &= 1 \quad \dots \text{(i)} \\
 \therefore (\sin \theta - \cos \theta)^2 &= 1 \\
 \therefore 1 - 2 \sin \theta \cos \theta &= 1 \\
 \therefore \sin \theta \cdot \cos \theta &= 0 \quad \dots \text{(ii)} \\
 \text{Now, } \sin^3 \theta - \cos^3 \theta &= (\sin \theta - \cos \theta) (\sin^2 \theta + \sin \theta \cos \theta + \cos^2 \theta) \\
 &= (1) (1 + \sin \theta \cdot \cos \theta) \quad \dots [\text{From (i)}] \\
 &= 1 + 0 \quad \dots [\text{From (ii)}] \\
 &= 1
 \end{aligned}$$

$$\begin{aligned}
 8. \quad \operatorname{cosec}^2 \theta &= 1 + \cot^2 \theta \\
 &= 1 + \frac{9}{16} \quad \dots \left[\because \tan \theta = -\frac{4}{3} \right] \\
 &= \frac{25}{16}
 \end{aligned}$$

$$\therefore \sin^2 \theta = \frac{1}{\operatorname{cosec}^2 \theta} = \frac{16}{25} \Rightarrow \sin \theta = \pm \frac{4}{5}$$

Both the values are acceptable.

$$\text{Since, } \tan \theta = -\frac{4}{3}$$

i.e., θ lies in 2nd or 4th quadrant.

$$\begin{aligned}
 9. \quad \sin \theta &= \frac{24}{25} \\
 \Rightarrow \cos \theta &= \frac{-7}{25}, \tan \theta = \frac{-24}{7} \\
 &\quad \dots [\because \theta \text{ lies in the } 2^{\text{nd}} \text{ quadrant}]
 \end{aligned}$$

$$\therefore \sec \theta + \tan \theta = \frac{-25}{7} - \frac{24}{7} = -7$$

$$\begin{aligned}
 10. \quad \cos \theta &= -\sqrt{1 - \left(\frac{2t}{1+t^2} \right)^2} \\
 &\quad \dots [\because \theta \text{ lies in the } 2^{\text{nd}} \text{ quadrant}]
 \end{aligned}$$



$$= -\sqrt{\frac{(1+t^2)^2 - 4t^2}{(1+t^2)^2}} = -\sqrt{\frac{(1-t^2)^2}{(1+t^2)^2}}$$

$$= \frac{-|1-t^2|}{1+t^2}$$

$$11. \sqrt{\operatorname{cosec}^2 \alpha + 2 \cot \alpha} = \sqrt{1 + \cot^2 \alpha + 2 \cot \alpha}$$

$$= |1 + \cot \alpha|$$

$$\text{But } \frac{3\pi}{4} < \alpha < \pi$$

$$\Rightarrow \cot \alpha < -1 \Rightarrow 1 + \cot \alpha < 0$$

$$\text{Hence, } |1 + \cot \alpha| = -(1 + \cot \alpha)$$

$$12. \operatorname{cosec} \theta - \cot \theta = 2017 \quad \dots(i)$$

$$\Rightarrow \frac{1}{\operatorname{cosec} \theta - \cot \theta} = \frac{1}{2017}$$

$$\Rightarrow \frac{\operatorname{cosec} \theta + \cot \theta}{(\operatorname{cosec} \theta - \cot \theta)(\operatorname{cosec} \theta + \cot \theta)} = \frac{1}{2017}$$

$$\Rightarrow \operatorname{cosec} \theta + \cot \theta = \frac{1}{2017} \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2 \operatorname{cosec} \theta = 2017 + \frac{1}{2017}$$

$$\Rightarrow \operatorname{cosec} \theta = \text{positive} \Rightarrow \sin \theta = \text{positive}$$

Subtracting (i) from (ii), we get

$$\Rightarrow 2 \cot \theta = \frac{1}{2017} - 2017$$

$$\Rightarrow \cot \theta = \text{negative} \Rightarrow \tan \theta = \text{negative}$$

\(\therefore\) \(\theta\) lies in II quadrant.

$$13. \operatorname{cosec} \theta - \cot \theta = \frac{1}{2} \quad \dots(i)$$

$$\therefore \operatorname{cosec} \theta + \cot \theta = 2 \quad \dots(ii)$$

$$\dots[\because \operatorname{cosec}^2 \theta - \cot^2 \theta = 1]$$

Adding (i) and (ii), we get

$$2 \operatorname{cosec} \theta = \frac{5}{2} \Rightarrow \sin \theta = \frac{4}{5}$$

Subtracting (ii) from (i), we get

$$2 \cot \theta = \frac{3}{2} \Rightarrow \cot \theta = \frac{3}{4}$$

$$\Rightarrow \cos \theta = \frac{3}{4} \sin \theta$$

$$= \frac{3}{4} \times \frac{4}{5} = \frac{3}{5}$$

$$14. \operatorname{cosec} A + \cot A = \frac{11}{2}$$

$$\Rightarrow \operatorname{cosec} A - \cot A = \frac{2}{11}$$

$$\therefore 2 \cot A = \frac{117}{22} \Rightarrow \tan A = \frac{44}{117}$$

$$15. \sec \theta + \tan \theta = p \quad \dots(i)$$

$$\therefore \sec \theta - \tan \theta = \frac{1}{p} \quad \dots(ii)$$

$$\therefore 2 \tan \theta = p - \frac{1}{p} \Rightarrow \tan \theta = \frac{p^2 - 1}{2p}$$

$$16. \tan \theta + \sec \theta = e^x \quad \dots(i)$$

$$\therefore \sec \theta - \tan \theta = e^{-x} \quad \dots(ii)$$

$$\therefore 2 \sec \theta = e^x + e^{-x}$$

$$\Rightarrow \cos \theta = \frac{2}{e^x + e^{-x}}$$

$$17. \sin \theta + \cos \theta = 1$$

Squaring both sides, we get

$$\sin^2 \theta + \cos^2 \theta + 2 \sin \theta \cos \theta = 1$$

$$\Rightarrow \sin \theta \cos \theta = 0$$

$$18. (3 \cos A - 5 \sin A)^2$$

$$= 9 \cos^2 A + 25 \sin^2 A - 30 \sin A \cos A$$

$$= 9(1 - \sin^2 A) + 25(1 - \cos^2 A)$$

$$- 30 \sin A \cos A$$

$$= 34 - (9 \sin^2 A + 25 \cos^2 A + 30 \sin A \cos A)$$

$$= 34 - (3 \sin A + 5 \cos A)^2$$

$$= 34 - 25 \quad \dots[\because 3 \sin A + 5 \cos A = 5]$$

$$= 9$$

$$19. \text{Given, } \sec \theta = m \text{ and } \tan \theta = n$$

$$\text{Since, } \sec^2 \theta - \tan^2 \theta = 1$$

$$\therefore (\sec \theta - \tan \theta)(\sec \theta + \tan \theta) = 1$$

$$\Rightarrow (m - n)(m + n) = 1$$

$$\Rightarrow m - n = \frac{1}{m + n} \quad \dots(i)$$

$$\therefore \frac{1}{m} \left\{ (m+n) + \frac{1}{m+n} \right\} = \frac{1}{m} (m+n+m-n)$$

$$\dots[\text{From (i)}]$$

$$= 2$$

$$20. n(m^2 - 1) = (\sec \theta + \operatorname{cosec} \theta) \cdot 2 \sin \theta \cos \theta$$

$$\dots[\because m^2 = 1 + 2 \sin \theta \cos \theta]$$

$$= \frac{\sin \theta + \cos \theta}{\sin \theta \cdot \cos \theta} \cdot 2 \sin \theta \cos \theta = 2m$$



$$21. \quad 2y \cos \theta = x \sin \theta \quad \dots(i)$$

$$\text{and } 2x \sec \theta - y \operatorname{cosec} \theta = 3$$

$$\Rightarrow \frac{2x}{\cos \theta} - \frac{y}{\sin \theta} = 3$$

$$\Rightarrow 2x \sin \theta - y \cos \theta - 3 \sin \theta \cos \theta = 0 \quad \dots(ii)$$

Solving (i) and (ii), we get

$$y = \sin \theta \text{ and } x = 2 \cos \theta$$

$$\therefore x^2 + 4y^2 = 4 \cos^2 \theta + 4 \sin^2 \theta \\ = 4 (\cos^2 \theta + \sin^2 \theta) = 4$$

$$22. \quad m + n = 2 \tan \theta, m - n = 2 \sin \theta$$

$$\therefore m^2 - n^2 = 4 \tan \theta \cdot \sin \theta \quad \dots(i)$$

$$\text{Also, } 4\sqrt{mn} = 4\sqrt{\tan^2 \theta - \sin^2 \theta} \\ = 4 \sin \theta \cdot \tan \theta \quad \dots(ii)$$

$$\therefore \text{From (i) and (ii), we get } m^2 - n^2 = 4\sqrt{mn}$$

$$23. \quad (\sec \alpha + \tan \alpha)(\sec \beta + \tan \beta)(\sec \gamma + \tan \gamma) \\ = \tan \alpha \tan \beta \tan \gamma \quad \dots(i)$$

$$\text{Let } x = (\sec \alpha - \tan \alpha)(\sec \beta - \tan \beta) \\ (\sec \gamma - \tan \gamma) \quad \dots(ii)$$

Multiplying equations (i) and (ii), we get

$$(\sec^2 \alpha - \tan^2 \alpha)(\sec^2 \beta - \tan^2 \beta)(\sec^2 \gamma - \tan^2 \gamma) \\ = x(\tan \alpha \tan \beta \tan \gamma)$$

$$\Rightarrow x = \frac{1}{\tan \alpha \tan \beta \tan \gamma}$$

$$\therefore x = \cot \alpha \cot \beta \cot \gamma$$

$$24. \quad 2P_6 - 3P_4 + 1 \\ = 2(\cos^6 \theta + \sin^6 \theta) - 3(\cos^4 \theta + \sin^4 \theta) + 1 \\ = 2(1 - 3 \sin^2 \theta \cos^2 \theta) - 3[(\sin^2 \theta + \cos^2 \theta)^2 \\ - 2 \sin^2 \theta \cos^2 \theta] + 1 \\ = 2 - 6 \sin^2 \theta \cos^2 \theta - 3(1 - 2 \sin^2 \theta \cos^2 \theta) + 1 \\ = 0$$

$$25. \quad (\sec A + \tan A - 1)(\sec A - \tan A + 1) - 2 \tan A \\ = \sec^2 A - \tan^2 A - \sec A \tan A + \sec A \\ + \sec A \tan A + \tan A - \sec A + \tan A - 1 - 2 \tan A \\ = \sec^2 A - \tan^2 A - 1 \\ = 0 \quad \dots[\because \sec^2 A - \tan^2 A = 1]$$

$$26. \quad \cos^4 \theta - \sin^4 \theta \\ = (\cos^2 \theta + \sin^2 \theta)(\cos^2 \theta - \sin^2 \theta) = \cos^2 \theta - \sin^2 \theta \\ = \cos^2 \theta - (1 - \cos^2 \theta) = 2 \cos^2 \theta - 1$$

$$27. \quad \sin^6 \theta + \cos^6 \theta + 3 \sin^2 \theta \cos^2 \theta \\ = (\sin^2 \theta + \cos^2 \theta)^3 - 3 \sin^2 \theta \cos^2 \theta \\ + 3 \sin^2 \theta \cos^2 \theta = 1$$

$$28. \quad 6(\sin^6 \theta + \cos^6 \theta) - 9(\sin^4 \theta + \cos^4 \theta) + 4 \\ = 6[(\sin^2 \theta + \cos^2 \theta)^3 - 3 \sin^2 \theta \cos^2 \theta] \\ - 9[(\sin^2 \theta + \cos^2 \theta)^2 - 2 \sin^2 \theta \cos^2 \theta] + 4 \\ = 6(1 - 3 \sin^2 \theta \cos^2 \theta) - 9(1 - 2 \sin^2 \theta \cos^2 \theta) + 4 \\ = 6 - 9 + 4 = 1$$

$$29. \quad \sin x + \sin^2 x = 1 \\ \Rightarrow \sin x = 1 - \sin^2 x \\ \Rightarrow \sin x = \cos^2 x$$

$$\therefore \cos^{12} x + 3 \cos^{10} x + 3 \cos^8 x + \cos^6 x - 1 \\ = \sin^6 x + 3 \sin^5 x + 3 \sin^4 x + \sin^3 x - 1 \\ = (\sin^2 x)^3 + 3(\sin^2 x)^2 \sin x + 3 \sin^2 x \cdot \sin^2 x \\ + (\sin x)^3 - 1 \\ = (\sin^2 x + \sin x)^3 - 1 = 1^3 - 1 = 0$$

$$30. \quad (\cos x + \sin x)^2 + k \sin x \cos x - 1 = 0 \\ \Rightarrow \cos^2 x + \sin^2 x + 2 \cos x \sin x \\ + k \sin x \cos x - 1 = 0 \\ \Rightarrow (k + 2) \cos x \sin x = 0 \\ \Rightarrow k + 2 = 0 \Rightarrow k = -2$$

$$31. \quad \text{Since, } \sec^2 \theta \geq 1 \\ \Rightarrow \frac{4xy}{(x+y)^2} \geq 1 \Rightarrow 4xy \geq (x+y)^2 \\ \Rightarrow (x-y)^2 \leq 0 \\ \text{It is possible only when } x = y \text{ and } x \neq 0.$$

$$32. \quad \text{Since, } \theta \in \left(0, \frac{\pi}{4}\right) \\ \therefore \tan \theta < 1 \text{ and } \cot \theta > 1 \\ \text{Let } \tan \theta = 1 - \lambda_1 \text{ and } \cot \theta = 1 + \lambda_2, \text{ where } \lambda_1 \\ \text{and } \lambda_2 \text{ are very small and positive.} \\ \text{Then, } t_1 = (1 - \lambda_1)^{1-\lambda_1}, t_2 = (1 - \lambda_1)^{1+\lambda_2} \\ t_3 = (1 + \lambda_2)^{1-\lambda_1}, t_4 = (1 + \lambda_2)^{1+\lambda_2} \\ \text{Clearly, } t_4 > t_3 > t_1 > t_2$$



Evaluation Test

$$\begin{aligned}
 1. \quad & \sin A = a \cos B \text{ and } \cos A = b \sin B \\
 \therefore & a^2 \cos^2 B + b^2 \sin^2 B = \sin^2 A + \cos^2 A \\
 & \Rightarrow a^2 \cos^2 B + b^2(1 - \cos^2 B) = 1 \\
 & \Rightarrow \cos^2 B = \frac{1-b^2}{a^2-b^2} \text{ and } \sin^2 B = \frac{a^2-1}{a^2-b^2} \\
 & \Rightarrow \tan^2 B = \frac{a^2-1}{1-b^2} \quad \dots(i)
 \end{aligned}$$

Again, $\sin A = a \cos B$ and $\cos A = b \sin B$

$$\Rightarrow \tan A = \frac{a}{b} \cot B$$

$$\Rightarrow \tan^2 A = \frac{a^2}{b^2} \cot^2 B$$

$$\Rightarrow \tan^2 A = \frac{a^2}{b^2} \left(\frac{1-b^2}{a^2-1} \right) \quad \dots[\text{From (i)}]$$

$$\begin{aligned}
 \therefore & (a^2-1) \tan^2 A + (1-b^2) \tan^2 B \\
 & = \frac{a^2}{b^2}(1-b^2) + (a^2-1) = \frac{a^2-b^2}{b^2}
 \end{aligned}$$

$$\begin{aligned}
 2. \quad & \text{Given, } \tan \theta = \frac{x \sin \phi}{1-x \cos \phi} \\
 & \Rightarrow \frac{1}{x} \tan \theta - \tan \theta \cos \phi = \sin \phi \\
 & \Rightarrow \frac{1}{x} = \frac{\sin \phi + \cos \phi \tan \theta}{\tan \theta}
 \end{aligned}$$

$$\text{Also, } \tan \phi = \frac{y \sin \theta}{1-y \cos \theta}$$

$$\Rightarrow \tan \phi = \frac{\sin \theta}{\frac{1}{y} - \cos \theta}$$

$$\Rightarrow \frac{1}{y} \tan \phi - \tan \phi \cos \theta = \sin \theta$$

$$\Rightarrow \frac{1}{y} \tan \phi = \sin \theta + \tan \phi \cos \theta$$

$$\Rightarrow \frac{1}{y} = \frac{\sin \theta + \tan \phi \cos \theta}{\tan \phi}$$

$$\therefore \frac{x}{y} = \left[\frac{\tan \theta}{\sin \phi + \cos \phi \tan \theta} \right] \times \left[\frac{\sin \theta + \tan \phi \cos \theta}{\tan \phi} \right]$$

$$\begin{aligned}
 & = \frac{\tan \theta}{\tan \phi} \left[\frac{\sin \theta + \cos \theta \frac{\sin \phi}{\cos \phi}}{\sin \phi + \cos \phi \frac{\sin \theta}{\cos \theta}} \right] \\
 & = \frac{\tan \theta \cos \theta}{\tan \phi \cos \phi} = \frac{\sin \theta}{\sin \phi}
 \end{aligned}$$

$$\begin{aligned}
 3. \quad & a \sin^2 x + b \cos^2 x = c \\
 & \Rightarrow a(1 - \cos^2 x) + b \cos^2 x = c \\
 & \Rightarrow (b-a) \cos^2 x = c-a \\
 & \Rightarrow (b-a) = (c-a)(1 + \tan^2 x) \\
 & \quad \quad \quad \dots[\because \sec^2 \theta = 1 + \tan^2 \theta]
 \end{aligned}$$

and $b \sin^2 y + a \cos^2 y = d$

$$\Rightarrow (a-b) \cos^2 y = d-b$$

$$\Rightarrow (a-b) = (d-b)(1 + \tan^2 y)$$

$$\therefore \tan^2 x = \frac{b-c}{c-a} \text{ and } \tan^2 y = \frac{a-d}{d-b}$$

$$\therefore \frac{\tan^2 x}{\tan^2 y} = \frac{(b-c)(d-b)}{(c-a)(a-d)} \quad \dots(i)$$

But, $a \tan x = b \tan y$

$$\Rightarrow \frac{\tan x}{\tan y} = \frac{b}{a} \quad \dots(ii)$$

From (i) and (ii), we get

$$\frac{b^2}{a^2} = \frac{(b-c)(d-b)}{(c-a)(a-d)}$$

$$\Rightarrow \frac{a^2}{b^2} = \frac{(c-a)(a-d)}{(b-c)(d-b)}$$

$$\begin{aligned}
 4. \quad & \text{Given, } \operatorname{cosec} \theta - \sin \theta = a \\
 & \Rightarrow \frac{1-\sin^2 \theta}{\sin \theta} = a \Rightarrow \frac{\cos^2 \theta}{\sin \theta} = a \quad \dots(i)
 \end{aligned}$$

and $\sec \theta - \cos \theta = b$

$$\Rightarrow \frac{1-\cos^2 \theta}{\cos \theta} = b \Rightarrow \frac{\sin^2 \theta}{\cos \theta} = b \quad \dots(ii)$$



Squaring (i) and multiplying by (ii), we get
 $\cos^3 \theta = a^2 b$

$$\Rightarrow \cos \theta = (a^2 b)^{1/3} \quad \dots(\text{iii})$$

Squaring (ii) and multiplying by (i), we get
 $\sin^3 \theta = b^2 a$

$$\Rightarrow \sin \theta = (b^2 a)^{1/3} \quad \dots(\text{iv})$$

Squaring (iii) and (iv) and adding, we get
 $1 = (a^2 b)^{2/3} + (b^2 a)^{2/3}$

$$\Rightarrow a^{4/3} b^{2/3} + b^{4/3} a^{2/3} = 1$$

$$\Rightarrow a^{2/3} b^{2/3} (a^{2/3} + b^{2/3}) = 1$$

5. $\sin x + \sin^2 x + \sin^3 x = 1$

$$\Rightarrow \sin x + \sin^3 x = 1 - \sin^2 x$$

$$\Rightarrow \sin x + \sin^3 x = \cos^2 x$$

$$\Rightarrow \sin x (1 + \sin^2 x) = \cos^2 x$$

$$\Rightarrow \sin x (2 - \cos^2 x) = \cos^2 x$$

$$\dots[\because \sin^2 \theta = 1 - \cos^2 \theta]$$

$$\Rightarrow \sin^2 x (2 - \cos^2 x)^2 = \cos^4 x$$

$$\Rightarrow (1 - \cos^2 x) (4 - 4 \cos^2 x + \cos^4 x) = \cos^4 x$$

$$\Rightarrow 4 - 4 \cos^2 x + \cos^4 x - 4 \cos^2 x + 4 \cos^4 x - \cos^6 x = \cos^4 x$$

$$\Rightarrow \cos^6 x - 4 \cos^4 x + 8 \cos^2 x = 4$$

6. Given, $\cot \theta + \tan \theta = m$

$$\Rightarrow \frac{1}{\tan \theta} + \tan \theta = m \Rightarrow 1 + \tan^2 \theta = m \tan \theta$$

$$\Rightarrow \sec^2 \theta = m \tan \theta \quad \dots(\text{i})$$

$$\text{and } \sec \theta - \cos \theta = n \Rightarrow \sec^2 \theta - 1 = n \sec \theta$$

$$\Rightarrow \tan^2 \theta = n \sec \theta$$

$$\Rightarrow \tan^4 \theta = n^2 \sec^2 \theta$$

$$\Rightarrow \tan^4 \theta = n^2 m \tan \theta \quad \dots[\text{From (i)}]$$

$$\Rightarrow \tan^3 \theta = n^2 m$$

$$\Rightarrow \tan \theta = (n^2 m)^{1/3} \quad \dots(\text{ii})$$

Putting (ii) in (i), we get

$$\sec^2 \theta = m(n^2 m)^{1/3}$$

$$\text{Since, } \sec^2 \theta - \tan^2 \theta = 1$$

$$\therefore m(mn^2)^{1/3} - (n^2 m)^{2/3} = 1$$

$$\Rightarrow m(mn^2)^{1/3} - n(nm^2)^{1/3} = 1$$

7. $\frac{\sin^4 \theta}{a} + \frac{\cos^4 \theta}{b} = \frac{1}{a+b}$

$$\Rightarrow (a+b) \left(\frac{\sin^4 \theta}{a} + \frac{\cos^4 \theta}{b} \right) = (\sin^2 \theta + \cos^2 \theta)^2$$

$$\Rightarrow \sin^4 \theta + \cos^4 \theta + \frac{b}{a} \sin^4 \theta + \frac{a}{b} \cos^4 \theta$$

$$= \sin^4 \theta + \cos^4 \theta + 2 \sin^2 \theta \cos^2 \theta$$

$$\Rightarrow \frac{b}{a} \sin^4 \theta + \frac{a}{b} \cos^4 \theta - 2 \sin^2 \theta \cos^2 \theta = 0$$

$$\Rightarrow \left(\sqrt{\frac{b}{a}} \sin^2 \theta - \sqrt{\frac{a}{b}} \cos^2 \theta \right)^2 = 0$$

$$\Rightarrow \sqrt{\frac{b}{a}} \sin^2 \theta = \sqrt{\frac{a}{b}} \cos^2 \theta$$

$$\Rightarrow b \sin^2 \theta = a \cos^2 \theta$$

$$\Rightarrow \frac{\sin^2 \theta}{a} = \frac{\cos^2 \theta}{b} = \frac{\sin^2 \theta + \cos^2 \theta}{a+b}$$

$$\Rightarrow \frac{\sin^2 \theta}{a} = \frac{\cos^2 \theta}{b} = \frac{1}{a+b}$$

$$\Rightarrow \sin^2 \theta = \frac{a}{a+b} \text{ and } \cos^2 \theta = \frac{b}{a+b}$$

Only option (B) does not satisfy these values.

Hence, option (B) is incorrect.

8. $x \sin \theta = y \cos \theta = \frac{2z \tan \theta}{1 - \tan^2 \theta}$

$$\text{Consider, } x \sin \theta = \frac{2z \tan \theta}{1 - \tan^2 \theta}$$

$$\Rightarrow x \sin \theta = \frac{2z \frac{\sin \theta}{\cos \theta}}{\frac{\cos^2 \theta - \sin^2 \theta}{\cos^2 \theta}}$$

$$\Rightarrow x \sin \theta = \frac{2z \sin \theta \cos \theta}{\cos^2 \theta - \sin^2 \theta}$$

$$\Rightarrow x = \frac{2z \cos \theta}{\cos^2 \theta - \sin^2 \theta}$$

$$\Rightarrow z = \frac{x(\cos^2 \theta - \sin^2 \theta)}{2 \cos \theta} \quad \dots(\text{i})$$

$$\text{Similarly, by solving } y \cos \theta = \frac{2z \tan \theta}{1 - \tan^2 \theta},$$

we get

$$z = \frac{y(\cos^2 \theta - \sin^2 \theta)}{2 \sin \theta} \quad \dots(\text{ii})$$

$$\text{From (i) and (ii), we get } \tan \theta = \frac{y}{x}$$

$$\therefore x \sin \theta = \frac{2 \frac{zy}{x}}{1 - \frac{y^2}{x^2}} = \frac{2xyz}{x^2 - y^2}$$



$$\Rightarrow \sin \theta = \frac{2yz}{x^2 - y^2}$$

$$\text{Similarly, } \cos \theta = \frac{2xz}{x^2 - y^2}$$

$$\text{Since, } \sin^2 \theta + \cos^2 \theta = 1$$

$$\therefore \left(\frac{2yz}{x^2 - y^2} \right)^2 + \left(\frac{2xz}{x^2 - y^2} \right)^2 = 1$$

$$\Rightarrow \frac{4z^2(x^2 + y^2)}{(x^2 - y^2)^2} = 1$$

$$\Rightarrow 4z^2(x^2 + y^2) = (x^2 - y^2)^2$$

$$9. \quad 3 \cot A = 6 \sec B = -2\sqrt{10}$$

$$\text{Consider, } \cot A = -\frac{2}{3}\sqrt{10}$$

$$\Rightarrow \cot^2 A = \frac{40}{9}$$

$$\therefore \operatorname{cosec}^2 A = \frac{49}{9}$$

$$\Rightarrow \operatorname{cosec} A = \frac{7}{3} \quad \dots \left[\because \frac{\pi}{2} < A < \pi \right]$$

$$\text{Also, } 6 \sec B = -2\sqrt{10}$$

$$\Rightarrow \cos B = -\frac{3}{\sqrt{10}}$$

$$\Rightarrow \sin B = -\sqrt{1 - \cos^2 B} \quad \dots \left[\because \pi < B < \frac{3\pi}{2} \right]$$

$$= -\sqrt{1 - \frac{9}{10}} = -\frac{1}{\sqrt{10}}$$

$$\therefore \tan B = \frac{\sin B}{\cos B} = \frac{1}{3}$$

$$\therefore \operatorname{cosec} A - \tan B = \frac{7}{3} - \frac{1}{3} = 2$$

$$10. \quad x = \sum_{n=0}^{\infty} \cos^{2n} \phi$$

$$= 1 + \cos^2 \phi + \cos^4 \phi + \dots \infty$$

$$= \frac{1}{(1 - \cos^2 \phi)} \quad \dots [\text{Since infinite G.P.}]$$

$$= \frac{1}{\sin^2 \phi}$$

$$y = \sum_{n=0}^{\infty} \sin^{2n} \phi$$

$$= 1 + \sin^2 \phi + \sin^4 \phi + \dots \infty$$

$$= \frac{1}{(1 - \sin^2 \phi)} = \frac{1}{\cos^2 \phi}$$

$$z = \sum_{n=0}^{\infty} \cos^{2n} \phi \sin^{2n} \phi$$

$$= 1 + \cos^2 \phi \sin^2 \phi + \cos^4 \phi \sin^4 \phi + \dots \infty$$

$$= \frac{1}{(1 - \cos^2 \phi \sin^2 \phi)}$$

$$\text{Now, } xyz = \frac{1}{\sin^2 \phi \cos^2 \phi (1 - \cos^2 \phi \sin^2 \phi)} \quad \dots (i)$$

$$\therefore xy + z = \frac{1}{\sin^2 \phi \cos^2 \phi} + \frac{1}{1 - \cos^2 \phi \sin^2 \phi}$$

$$= \frac{1}{\sin^2 \phi \cos^2 \phi (1 - \cos^2 \phi \sin^2 \phi)}$$

$$= xyz \quad \dots [\text{From (i)}]$$

Trigonometric Functions of Compound Angles



Hints



Classical Thinking

- $$\begin{aligned} \cos 105^\circ &= \cos (60^\circ + 45^\circ) \\ &= \cos 60^\circ \cos 45^\circ - \sin 60^\circ \sin 45^\circ \\ &= \frac{1-\sqrt{3}}{2\sqrt{2}} \end{aligned}$$
- $$\begin{aligned} \tan 15^\circ &= \tan (45^\circ - 30^\circ) \\ &= \frac{1-\frac{1}{\sqrt{3}}}{1+\frac{1}{\sqrt{3}}} \\ &\dots \left[\because \tan(A-B) = \frac{\tan A - \tan B}{1 + \tan A \tan B} \right] \\ &= \frac{\sqrt{3}-1}{\sqrt{3}+1} \times \frac{\sqrt{3}-1}{\sqrt{3}-1} \\ &= 2 - \sqrt{3} \end{aligned}$$
- $$\begin{aligned} \cos 38^\circ \cos 8^\circ + \sin 38^\circ \sin 8^\circ \\ &= \cos (38^\circ - 8^\circ) = \cos 30^\circ \end{aligned}$$
- $$\begin{aligned} \frac{1}{4} (\sqrt{3} \cos 23^\circ - \sin 23^\circ) \\ &= \frac{1}{2} (\cos 30^\circ \cos 23^\circ - \sin 30^\circ \sin 23^\circ) \\ &\dots \left[\because \cos 30^\circ = \frac{\sqrt{3}}{2}, \sin 30^\circ = \frac{1}{2} \right] \\ &= \frac{1}{2} \cos (30^\circ + 23^\circ) = \frac{1}{2} \cos 53^\circ \end{aligned}$$
- $$\begin{aligned} \tan 5A &= \tan (3A + 2A) \\ &= \frac{\tan 3A + \tan 2A}{1 - \tan 3A \tan 2A} \\ \Rightarrow \tan 5A - \tan 5A \tan 3A \tan 2A \\ &= \tan 3A + \tan 2A \\ \Rightarrow \tan 5A - \tan 3A - \tan 2A \\ &= \tan 5A \tan 3A \tan 2A \end{aligned}$$
- $$\begin{aligned} \tan (57^\circ - 12^\circ) &= \tan 45^\circ \\ \Rightarrow \frac{\tan 57^\circ - \tan 12^\circ}{1 + \tan 57^\circ \tan 12^\circ} &= 1 \end{aligned}$$

$$\begin{aligned} \Rightarrow \tan 57^\circ - \tan 12^\circ &= 1 + \tan 57^\circ \tan 12^\circ \\ \Rightarrow \tan 57^\circ - \tan 12^\circ - \tan 57^\circ \tan 12^\circ &= 1 \\ &= \tan 45^\circ \end{aligned}$$

$$\begin{aligned} 7. \quad \frac{\cos 10^\circ + \sin 10^\circ}{\cos 10^\circ - \sin 10^\circ} &= \frac{1 + \tan 10^\circ}{1 - \tan 10^\circ} \\ &= \tan (45^\circ + 10^\circ) \\ &\dots [\because \tan 45^\circ = 1] \\ &= \tan 55^\circ \end{aligned}$$

$$\begin{aligned} 8. \quad \frac{\cos 8^\circ - \sin 8^\circ}{\cos 8^\circ + \sin 8^\circ} &= \frac{1 - \tan 8^\circ}{1 + \tan 8^\circ} \\ &= \tan (45^\circ - 8^\circ) = \tan 37^\circ \end{aligned}$$

$$\begin{aligned} 9. \quad \tan (A + B) &= \frac{\tan A + \tan B}{1 - \tan A \tan B} \\ &= \frac{\frac{a}{a+1} + \frac{1}{2a+1}}{1 - \frac{a}{a+1} \cdot \frac{1}{2a+1}} \\ &= \frac{2a^2 + a + a + 1}{2a^2 + 2a + a + 1 - a} \\ &= \frac{2a^2 + 2a + 1}{2a^2 + 2a + 1} \\ &= 1 = \tan \frac{\pi}{4} \end{aligned}$$

$$\therefore A + B = \frac{\pi}{4}$$

$$\begin{aligned} 10. \quad \cot(A - B) &= \frac{1}{\tan(A - B)} \\ &= \frac{1 + \tan A \tan B}{\tan A - \tan B} \\ &= \frac{1}{\tan A - \tan B} + \frac{\tan A \tan B}{\tan A - \tan B} \\ &= \frac{1}{\tan A - \tan B} + \frac{1}{\cot B - \cot A} \\ &= \frac{1}{x} + \frac{1}{y} \end{aligned}$$



$$11. \text{ Since, } \cos^2 A - \sin^2 B = \cos(A+B) \cdot \cos(A-B)$$

$$\therefore \cos^2 48^\circ - \sin^2 12^\circ = \cos 60^\circ \cdot \cos 36^\circ$$

$$= \frac{1}{2} \left(\frac{\sqrt{5}+1}{4} \right)$$

$$= \frac{\sqrt{5}+1}{8}$$

$$12. \frac{\cot^2 15^\circ - 1}{\cot^2 15^\circ + 1} = \frac{\frac{\cos^2 15^\circ}{\sin^2 15^\circ} - 1}{\frac{\cos^2 15^\circ}{\sin^2 15^\circ} + 1}$$

$$= \frac{\cos^2 15^\circ - \sin^2 15^\circ}{\cos^2 15^\circ + \sin^2 15^\circ} = \cos(30^\circ)$$

$$[\because \cos^2 A - \sin^2 B = \cos(A+B) \cos(A-B)]$$

$$= \frac{\sqrt{3}}{2}$$

$$13. \tan(-945^\circ) = \tan[-(945^\circ)]$$

$$= -\tan[(2 \times 360^\circ + 225^\circ)]$$

$$= -\tan(225^\circ)$$

$$= -\tan 45^\circ$$

$$\dots[\because \tan(180^\circ + \theta) = \tan \theta]$$

$$= -1$$

$$14. \sin \frac{\pi}{10} \sin \frac{3\pi}{10} = \sin 18^\circ \cdot \sin 54^\circ$$

$$= \sin 18^\circ \cdot \cos 36^\circ$$

$$\dots[\because \sin(90^\circ - \theta) = \cos \theta]$$

$$= \frac{\sqrt{5}-1}{4} \cdot \frac{\sqrt{5}+1}{4}$$

$$= \frac{1}{4}$$

$$15. \sin 15^\circ + \cos 105^\circ$$

$$= \sin 15^\circ + \cos(90^\circ + 15^\circ)$$

$$= \sin 15^\circ - \sin 15^\circ$$

$$\dots[\because \cos(90^\circ + \theta) = -\sin \theta]$$

$$= 0$$

$$16. \tan(A+B) = \frac{\frac{1}{2} + \frac{1}{3}}{1 - \frac{1}{2} \cdot \frac{1}{3}} = 1$$

$$\therefore A+B = 45^\circ$$

$$\Rightarrow 2A = 90^\circ - 2B$$

$$\Rightarrow \cos 2A = \sin 2B \dots[\because \cos(90^\circ - \theta) = \sin \theta]$$

$$17. \cos 7\theta + \cos \theta = \cos(8\theta - \theta) + \cos \theta$$

$$= \cos(\pi - \theta) + \cos \theta$$

$$\dots[\because 8\theta = \pi(\text{given})]$$

$$= -\cos \theta + \cos \theta$$

$$= 0$$

$$18. \text{ Since, } A+C = 180^\circ \text{ and } B+D = 180^\circ$$

$$\therefore \cos A + \cos B = \cos(180^\circ - C) + \cos(180^\circ - D)$$

$$= -(\cos C + \cos D)$$

$$\dots[\because \cos(180^\circ - \theta) = -\cos \theta]$$

$$19. \text{ Since, ABCD is a cyclic quadrilateral.}$$

$$\therefore A+C = 180^\circ$$

$$\Rightarrow A = 180^\circ - C$$

$$\Rightarrow \cos A = \cos(180^\circ - C) = -\cos C$$

$$\Rightarrow \cos A + \cos C = 0 \dots(\text{i})$$

$$\text{Also, } B+D = 180^\circ$$

$$\Rightarrow \cos B + \cos D = 0 \dots(\text{ii})$$

$$\text{Subtracting (ii) from (i), we get}$$

$$\cos A - \cos B + \cos C - \cos D = 0$$

$$20. \sin 10^\circ + \sin 20^\circ + \sin 30^\circ + \dots + \sin 180^\circ$$

$$+ \sin(180^\circ + 10^\circ) + \sin(180^\circ + 20^\circ) + \dots$$

$$+ \sin(180^\circ + 180^\circ)$$

$$= 0 \dots[\because \sin(180^\circ + \theta) = -\sin \theta]$$

$$21. (\cos 1^\circ + \cos 179^\circ) + (\cos 2^\circ + \cos 178^\circ)$$

$$+ \dots + (\cos 89^\circ + \cos 91^\circ)$$

$$+ (\cos 90^\circ + \cos 180^\circ)$$

$$= -1 \dots[\because \cos(180^\circ - \theta) = -\cos \theta]$$

$$22. \sec\left(\frac{7\pi}{2} - A\right) = \sec\left(2\pi + \frac{3\pi}{2} - A\right)$$

$$= \sec\left(\frac{3\pi}{2} - A\right)$$

$$= -\operatorname{cosec} A$$

$$23. \frac{\cot 54^\circ}{\tan 36^\circ} + \frac{\tan 20^\circ}{\cot 70^\circ}$$

$$= \frac{\cot(90^\circ - 36^\circ)}{\tan 36^\circ} + \frac{\tan 20^\circ}{\cot(90^\circ - 20^\circ)}$$

$$= 1 + 1 \dots[\because \cot(90^\circ - \theta) = \tan \theta]$$

$$= 2$$

$$24. \frac{\cos(90^\circ + \theta) \sec(-\theta) \tan(180^\circ - \theta)}{\sin(360^\circ + \theta) \sec(180^\circ + \theta) \cot(90^\circ - \theta)}$$

$$= \frac{(-\sin \theta)(\sec \theta)(-\tan \theta)}{(\sin \theta)(-\sec \theta) \tan \theta}$$

$$= -1$$



$$25. \quad \sin^2 25^\circ + \sin^2 65^\circ = \sin^2 25^\circ + \sin^2 (90^\circ - 25^\circ) \\ = \sin^2 25^\circ + \cos^2 25^\circ \\ \dots [\because \sin(90^\circ - \theta) = \cos \theta] \\ = 1$$

$$26. \quad \sin \frac{7\pi}{8} = \sin \left(\pi - \frac{\pi}{8} \right) = \sin \frac{\pi}{8} \\ \sin \frac{5\pi}{8} = \sin \left(\pi - \frac{3\pi}{8} \right) = \sin \frac{3\pi}{8}$$

$$\therefore \sin^2 \frac{\pi}{8} + \sin^2 \frac{3\pi}{8} + \sin^2 \frac{5\pi}{8} + \sin^2 \frac{7\pi}{8} \\ = 2 \left[\sin^2 \frac{\pi}{8} + \sin^2 \frac{3\pi}{8} \right] \\ = 2 \left[\sin^2 \frac{\pi}{8} + \cos^2 \frac{\pi}{8} \right] \\ \dots \left[\because \sin \frac{3\pi}{8} = \sin \left(\frac{\pi}{2} - \frac{\pi}{8} \right) = \cos \frac{\pi}{8} \right] \\ = 2$$

$$27. \quad \cos 2\theta = 2 \cos^2 \theta - 1 \\ = 1 - 2 \sin^2 \theta \\ = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}$$

$$28. \quad \sin 4\theta = 2 \sin 2\theta \cos 2\theta \\ = 2 \cdot 2 \sin \theta \cos \theta (1 - 2 \sin^2 \theta) \\ = 4 \sin \theta (1 - 2 \sin^2 \theta) \sqrt{1 - \sin^2 \theta}$$

$$29. \quad \tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}, \quad \cos 2\theta = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}$$

$$\therefore \tan 2\theta + \sec 2\theta = \frac{2t}{1-t^2} + \frac{1+t^2}{1-t^2} \\ \dots [\because \tan \theta = t(\text{given})] \\ = \frac{(1+t)^2}{(1-t)(1+t)} = \frac{1+t}{1-t}$$

$$30. \quad \text{Given, } \sin A + \cos A = 1 \\ \text{Squaring on both sides, we get} \\ (\sin A + \cos A)^2 = 1 \\ \Rightarrow 1 + \sin 2A = 1 \\ \Rightarrow \sin 2A = 0$$

$$31. \quad 2 + 2 \cos 4\theta = 2(1 + \cos 4\theta) \\ = 4 \cos^2 2\theta \quad \dots (i)$$

$$\therefore \sqrt{2 + \sqrt{2 + 2 \cos 4\theta}} = \sqrt{2 + 2 \cos 2\theta} \quad \dots [\text{From (i)}] \\ = \sqrt{2(1 + \cos 2\theta)} \\ = \sqrt{4 \cos^2 \theta} \\ = 2 \cos \theta$$

$$32. \quad 1 + \cos^2 2A = (\cos^2 A + \sin^2 A)^2 \\ + (\cos^2 A - \sin^2 A)^2 \\ = 2(\cos^4 A + \sin^4 A)$$

$$33. \quad 1 - 2 \sin^2 \left(\frac{\pi}{4} + \theta \right) = \cos \left(\frac{\pi}{2} + 2\theta \right) \\ \dots [\because \cos 2\theta = 1 - 2 \sin^2 \theta] \\ = -\sin 2\theta$$

$$34. \quad \sin \theta \cos \theta = \frac{1}{2} (\sin 2\theta) \\ \text{Since, } -1 \leq \sin 2\theta \leq 1 \\ \Rightarrow -\frac{1}{2} \leq \frac{1}{2} (\sin 2\theta) \leq \frac{1}{2} \\ \therefore \text{Largest value is } \frac{1}{2}.$$

$$35. \quad (\sec 2A + 1) \sec^2 A \\ = \left(\frac{1 + \tan^2 A}{1 - \tan^2 A} + 1 \right) (1 + \tan^2 A) \\ = \frac{2(1 + \tan^2 A)}{1 - \tan^2 A} = 2 \sec 2A$$

$$36. \quad \operatorname{cosec} A - 2 \cot 2A \cos A \\ = \frac{1}{\sin A} - \frac{2 \cos A \cos 2A}{\sin 2A} \\ = \frac{1}{\sin A} - \frac{2 \cos A \cos 2A}{2 \sin A \cos A} \\ = \frac{1 - \cos 2A}{\sin A} = \frac{2 \sin^2 A}{\sin A} = 2 \sin A$$

$$37. \quad \cos 20^\circ \cos 40^\circ \cos 80^\circ = \frac{\sin 2^3 20^\circ}{2^3 \sin 20^\circ} \\ = \frac{\sin 160^\circ}{8 \sin 20^\circ} \\ = \frac{\sin (180^\circ - 20^\circ)}{8 \sin 20^\circ} \\ = \frac{1}{8}$$

$$38. \quad \frac{\sin \theta + \sin 2\theta}{1 + \cos \theta + \cos 2\theta} = \frac{\sin \theta + 2 \sin \theta \cos \theta}{2 \cos^2 \theta + \cos \theta} \\ = \frac{\sin \theta (1 + 2 \cos \theta)}{\cos \theta (1 + 2 \cos \theta)} \\ = \tan \theta$$

$$39. \quad \sin^3 \theta + \cos^3 \theta \\ = (\sin \theta + \cos \theta) \left(\cos^2 \theta + \sin^2 \theta - \frac{\sin 2\theta}{2} \right)$$



$$= \sqrt{(\sin \theta + \cos \theta)^2} \left(1 - \frac{\sin 2\theta}{2}\right)$$

$$\Rightarrow \sin^3 \theta + \cos^3 \theta = \sqrt{1 + \frac{3}{4} \left(1 - \frac{3}{8}\right)}$$

$$= \frac{\sqrt{7}}{2} \times \frac{5}{8} = \frac{5\sqrt{7}}{16}$$

40. Given that, $\cos 3\theta = \alpha \cos \theta + \beta \cos^3 \theta$
 But, $\cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta$
 $\Rightarrow (\alpha, \beta) = (-3, 4)$

41. Given, $\tan A = \frac{1}{2}$
 $\Rightarrow \tan 3A = \frac{3 \tan A - \tan^3 A}{1 - 3 \tan^2 A}$
 $= \frac{3 \cdot \frac{1}{2} - \frac{1}{8}}{1 - 3 \cdot \frac{1}{4}} = \frac{12 - 1}{2}$
 $= \frac{11}{2}$

42. We have, $x + \frac{1}{x} = 2 \cos \theta$
 Now, $x^3 + \frac{1}{x^3} = \left(x + \frac{1}{x}\right)^3 - 3 \left(x + \frac{1}{x}\right)$
 $= (2 \cos \theta)^3 - 3(2 \cos \theta)$
 $= 8 \cos^3 \theta - 6 \cos \theta$
 $= 2(4 \cos^3 \theta - 3 \cos \theta) = 2 \cos 3\theta$

43. $\cos \left(\frac{\alpha}{2}\right) = -\sqrt{\frac{1 + \cos \alpha}{2}}$
 $\dots \left[\because \pi < \alpha < \frac{3\pi}{2} \Rightarrow \frac{\pi}{2} < \frac{\alpha}{2} < \frac{3\pi}{4} \right]$

Now, $\cos \alpha = -\sqrt{1 - \sin^2 \alpha}$
 $\dots \left[\because \pi < \alpha < \frac{3\pi}{2} \right]$

$$= -\sqrt{1 - \frac{9}{25}} = -\frac{4}{5}$$

$$\therefore \cos \frac{\alpha}{2} = -\sqrt{\frac{1 - \frac{4}{5}}{2}} = -\frac{1}{\sqrt{10}}$$

44. $\frac{1 - t^2}{1 + t^2} = \frac{1 - \tan^2 \frac{\theta}{2}}{1 + \tan^2 \frac{\theta}{2}} = \cos \theta$

45. Given that, $\tan \frac{A}{2} = \frac{3}{2}$

$$\therefore \frac{1 + \cos A}{1 - \cos A} = \frac{2 \cos^2 \frac{A}{2}}{2 \sin^2 \frac{A}{2}} = \cot^2 \frac{A}{2}$$

$$= \left(\frac{2}{3}\right)^2 = \frac{4}{9}$$

46. $\tan \left(\frac{A}{2}\right) = \frac{\sin \left(\frac{A}{2}\right)}{\cos \left(\frac{A}{2}\right)} = \sqrt{\frac{1 - \cos A}{1 + \cos A}}$
 $= \sqrt{\frac{1 - \cos A}{1 + \cos A}}$

47. $\tan^2 \frac{\theta}{2} = \frac{1 - \cos \theta}{1 + \cos \theta}$
 $= \frac{\tan \alpha - \tan \beta}{\tan \alpha + \tan \beta}$
 $\dots \left[\because \cos \theta = \frac{\tan \beta}{\tan \alpha} \text{ (given)} \right]$
 $= \frac{\frac{\sin \alpha}{\cos \alpha} - \frac{\sin \beta}{\cos \beta}}{\frac{\sin \alpha}{\cos \alpha} + \frac{\sin \beta}{\cos \beta}}$
 $= \frac{\sin \alpha \cos \beta - \sin \beta \cos \alpha}{\sin \alpha \cos \beta + \sin \beta \cos \alpha}$
 $= \frac{\sin(\alpha - \beta)}{\sin(\alpha + \beta)}$



Critical Thinking

- $\cos(A + B) = \alpha \cos A \cos B + \beta \sin A \sin B$
 But, $\cos(A + B) = \cos A \cos B - \sin A \sin B$
 $\Rightarrow \alpha = 1, \beta = -1$
- $\cos(A + B) = \cos A \cos B - \sin A \sin B$
 $= \sqrt{1 - \frac{16}{25}} \left(-\frac{12}{13}\right) - \frac{4}{5} \sqrt{1 - \frac{144}{169}}$



$$= \frac{3}{5} \left(-\frac{12}{13} \right) - \frac{4}{5} \left(-\frac{5}{13} \right)$$

....[∵ A lies in first quadrant and
B lies in third quadrant]

$$= -\frac{16}{65}$$

3. $\sin(A + B) = \sin A \cos B + \cos A \sin B$

$$= \frac{1}{\sqrt{10}} \sqrt{1 - \frac{1}{5}} + \frac{1}{\sqrt{5}} \sqrt{1 - \frac{1}{10}}$$

$$= \frac{1}{\sqrt{10}} \sqrt{\frac{4}{5}} + \frac{1}{\sqrt{5}} \sqrt{\frac{9}{10}}$$

$$= \frac{1}{\sqrt{50}} (2 + 3) = \frac{5}{\sqrt{50}}$$

$$= \frac{1}{\sqrt{2}}$$

$$\Rightarrow \sin(A + B) = \sin \frac{\pi}{4}$$

$$\Rightarrow A + B = \frac{\pi}{4}$$

4. $\tan(\theta + \phi) = \frac{\tan \theta + \tan \phi}{1 - \tan \theta \tan \phi} = \frac{\frac{1}{2} + \frac{1}{3}}{1 - \frac{1}{2} \cdot \frac{1}{3}} = 1$

$$\tan(2\theta + \phi) = \tan[\theta + (\theta + \phi)]$$

$$= \frac{\tan \theta + \tan(\theta + \phi)}{1 - \tan \theta \tan(\theta + \phi)}$$

$$= \frac{\frac{1}{2} + 1}{1 - \frac{1}{2} \cdot 1} = 3$$

5. $\cos \theta = \frac{8}{17}$ and $0 < \theta < \frac{\pi}{2}$

$$\Rightarrow \sin \theta = \sqrt{1 - \left(\frac{8}{17}\right)^2}$$

$$= \frac{15}{17}$$

$$\cos(30^\circ + \theta) + \cos(45^\circ - \theta) + \cos(120^\circ - \theta)$$

$$= \cos 30^\circ \cos \theta - \sin 30^\circ \sin \theta + \cos 45^\circ \cos \theta$$

$$+ \sin 45^\circ \sin \theta + \cos 120^\circ \cos \theta + \sin 120^\circ \sin \theta$$

$$= \cos \theta \left(\frac{\sqrt{3}}{2} + \frac{1}{\sqrt{2}} - \frac{1}{2} \right) - \sin \theta \left(\frac{1}{2} - \frac{1}{\sqrt{2}} - \frac{\sqrt{3}}{2} \right)$$

$$= \frac{8}{17} \left(\frac{\sqrt{3}}{2} + \frac{1}{\sqrt{2}} - \frac{1}{2} \right) + \frac{15}{17} \left(\frac{\sqrt{3}}{2} + \frac{1}{\sqrt{2}} - \frac{1}{2} \right)$$

$$= \frac{23}{17} \left(\frac{\sqrt{3}-1}{2} + \frac{1}{\sqrt{2}} \right)$$

6. $A + B = \frac{\pi}{4}$

$$\Rightarrow \tan(A + B) = \tan \frac{\pi}{4}$$

$$\Rightarrow \frac{\tan A + \tan B}{1 - \tan A \tan B} = 1$$

$$\Rightarrow \tan A + \tan B + \tan A \tan B = 1$$

$$\Rightarrow (1 + \tan A)(1 + \tan B) = 2$$

7. $A + B = 45^\circ$

$$\Rightarrow \tan(A + B) = 1$$

$$\Rightarrow \tan A + \tan B = 1 - \tan A \tan B$$

$$\Rightarrow \frac{1}{\cot A} + \frac{1}{\cot B} = 1 - \frac{1}{\cot A \cot B}$$

$$\Rightarrow \cot A + \cot B = \cot A \cot B - 1$$

$$\Rightarrow \cot A \cot B - \cot A - \cot B = 1$$

$$\Rightarrow \cot A \cot B - \cot A - \cot B + 1 = 2$$

$$\Rightarrow (\cot A - 1)(\cot B - 1) = 2$$

8. $\sin(\alpha - \beta) = \sin[(\theta - \beta) - (\theta - \alpha)]$

$$= \sin(\theta - \beta) \cos(\theta - \alpha) - \cos(\theta - \beta) \sin(\theta - \alpha)$$

$$= ba - \sqrt{1 - b^2} \sqrt{1 - a^2}$$

$$\text{and } \cos(\alpha - \beta) = \cos[(\theta - \beta) - (\theta - \alpha)]$$

$$= \cos(\theta - \beta) \cos(\theta - \alpha) + \sin(\theta - \beta) \sin(\theta - \alpha)$$

$$= a\sqrt{1 - b^2} + b\sqrt{1 - a^2}$$

$$\therefore \cos^2(\alpha - \beta) + 2ab \sin(\alpha - \beta)$$

$$= (a\sqrt{1 - b^2} + b\sqrt{1 - a^2})^2 + 2ab(ab - \sqrt{1 - a^2} \sqrt{1 - b^2})$$

$$= a^2 + b^2$$

9. Let $x - y = \alpha$, $y - z = \beta$ and $z - x = \gamma$,

$$\text{then } \alpha + \beta + \gamma = 0$$

$$\Rightarrow \alpha + \beta = -\gamma$$

$$\Rightarrow \tan(\alpha + \beta) = \tan(-\gamma)$$

$$\Rightarrow \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta} = -\tan \gamma$$

$$\Rightarrow \tan \alpha + \tan \beta + \tan \gamma = \tan \alpha \tan \beta \tan \gamma$$

10. Since, $\tan 3A = \frac{\tan A + \tan 2A}{1 - \tan A \tan 2A}$

$$\Rightarrow \tan 3A - \tan 2A - \tan A$$

$$= \tan 3A \tan 2A \tan A$$



$$11. \quad \tan(20^\circ + 40^\circ) = \frac{\tan 20^\circ + \tan 40^\circ}{1 - \tan 20^\circ \tan 40^\circ}$$

$$\Rightarrow \sqrt{3} = \frac{\tan 20^\circ + \tan 40^\circ}{1 - \tan 20^\circ \tan 40^\circ}$$

$$\Rightarrow \sqrt{3} - \sqrt{3} \tan 20^\circ \tan 40^\circ = \tan 20^\circ + \tan 40^\circ$$

$$\Rightarrow \tan 20^\circ + \tan 40^\circ + \sqrt{3} \tan 20^\circ \tan 40^\circ = \sqrt{3}$$

$$12. \quad \tan\left(\frac{6\pi}{15} - \frac{\pi}{15}\right) = \tan \frac{\pi}{3}$$

$$\Rightarrow \frac{\tan \frac{6\pi}{15} - \tan \frac{\pi}{15}}{1 + \tan \frac{6\pi}{15} \tan \frac{\pi}{15}} = \tan \frac{\pi}{3}$$

$$\Rightarrow \tan \frac{6\pi}{15} - \tan \frac{\pi}{15} = \sqrt{3} + \sqrt{3} \tan \frac{6\pi}{15} \tan \frac{\pi}{15}$$

$$\Rightarrow \tan \frac{6\pi}{15} - \tan \frac{\pi}{15} - \sqrt{3} \tan \frac{6\pi}{15} \tan \frac{\pi}{15} = \sqrt{3}$$

$$\Rightarrow \tan \frac{2\pi}{5} - \tan \frac{\pi}{15} - \sqrt{3} \tan \frac{2\pi}{5} \tan \frac{\pi}{15} = \sqrt{3}$$

$$13. \quad 2 \tan(A - B) = 2 \left(\frac{\tan A - \tan B}{1 + \tan A \tan B} \right)$$

$$= 2 \left[\frac{2 \tan B + \cot B - \tan B}{1 + (2 \tan B + \cot B) \tan B} \right]$$

$$\dots [\because \tan A = 2 \tan B + \cot B]$$

$$= 2 \left[\frac{\tan B + \cot B}{2(1 + \tan^2 B)} \right]$$

$$= \frac{\cot B(\tan^2 B + 1)}{1 + \tan^2 B}$$

$$= \cot B$$

$$14. \quad \sin \alpha + \sin \beta + \sin \gamma - \sin(\alpha + \beta + \gamma)$$

$$= \sin \alpha + \sin \beta + \sin \gamma - \sin \alpha \cos \beta \cos \gamma$$

$$\quad - \cos \alpha \sin \beta \cos \gamma - \cos \alpha \cos \beta \sin \gamma$$

$$\quad + \sin \alpha \sin \beta \sin \gamma$$

$$= \sin \alpha(1 - \cos \beta \cos \gamma) + \sin \beta(1 - \cos \alpha \cos \gamma)$$

$$+ \sin \gamma(1 - \cos \alpha \cos \beta) + \sin \alpha \sin \beta \sin \gamma > 0$$

$$\therefore \sin \alpha + \sin \beta + \sin \gamma > \sin(\alpha + \beta + \gamma)$$

$$\Rightarrow \frac{\sin(\alpha + \beta + \gamma)}{\sin \alpha + \sin \beta + \sin \gamma} < 1$$

$$15. \quad \frac{1}{\tan 3A - \tan A} - \frac{1}{\cot 3A - \cot A}$$

$$= \frac{1}{\tan 3A - \tan A} + \frac{\tan A \tan 3A}{\tan 3A - \tan A}$$

$$= \frac{1}{\tan 3A - \tan A} = \frac{1}{\tan 2A} = \cot 2A$$

$$16. \quad \cos 105^\circ + \sin 105^\circ$$

$$= \cos(90^\circ + 15^\circ) + \sin(90^\circ + 15^\circ)$$

$$= \cos 15^\circ - \sin 15^\circ$$

$$= \cos(45^\circ - 30^\circ) - \sin(45^\circ - 30^\circ)$$

$$= \frac{\sqrt{3}+1}{2\sqrt{2}} - \frac{\sqrt{3}-1}{2\sqrt{2}}$$

$$= \frac{2}{2\sqrt{2}}$$

$$= \frac{1}{\sqrt{2}}$$

$$17. \quad \tan 81^\circ - \tan 63^\circ - \tan 27^\circ + \tan 9^\circ$$

$$= \{\tan(90^\circ - 9^\circ) + \tan 9^\circ\}$$

$$\quad - \{\tan(90^\circ - 27^\circ) + \tan 27^\circ\}$$

$$= (\cot 9^\circ + \tan 9^\circ) - (\cot 27^\circ + \tan 27^\circ)$$

$$= 2 \operatorname{cosec} 18^\circ - 2 \operatorname{cosec} 54^\circ$$

$$\dots [\because \tan \theta + \cot \theta = 2 \operatorname{cosec} 2\theta]$$

$$= \frac{2}{\sin 18^\circ} - \frac{2}{\sin 54^\circ}$$

$$= \frac{2}{\sin 18^\circ} - \frac{2}{\cos 36^\circ}$$

$$= \frac{2 \times 4}{\sqrt{5}-1} - \frac{2 \times 4}{\sqrt{5}+1}$$

$$= 8 \left[\frac{\sqrt{5}+1-\sqrt{5}+1}{(\sqrt{5})^2 - 1^2} \right]$$

$$= 4$$

$$18. \quad \beta + \gamma = \alpha$$

$$\Rightarrow \gamma = \alpha - \beta$$

$$\Rightarrow \tan \gamma = \tan(\alpha - \beta)$$

$$\Rightarrow \tan \gamma = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta}$$

$$\Rightarrow \tan \gamma = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \cot \alpha}$$

$$\dots \left[\because \alpha + \beta = \frac{\pi}{2}, \therefore \beta = \frac{\pi}{2} - \alpha \right]$$

$$\Rightarrow \tan \gamma = \frac{1}{2}(\tan \alpha - \tan \beta)$$

$$\Rightarrow \tan \alpha = \tan \beta + 2 \tan \gamma$$

$$19. \quad \tan\left(\frac{\pi}{4} + \theta\right) \tan\left(\frac{3\pi}{4} + \theta\right)$$

$$= \tan\left(\frac{\pi}{4} + \theta\right) \tan\left(\frac{\pi}{2} + \left(\frac{\pi}{4} + \theta\right)\right)$$



$$= \tan\left(\frac{\pi}{4} + \theta\right) \left\{ -\cot\left(\frac{\pi}{4} + \theta\right) \right\}$$

$$\dots \left[\because \tan\left(\frac{\pi}{2} + \alpha\right) = -\cot \alpha \right]$$

$$= -1$$

20. $\tan(100^\circ + 125^\circ) = \frac{\tan 100^\circ + \tan 125^\circ}{1 - \tan 100^\circ \tan 125^\circ}$

$$\Rightarrow \tan 225^\circ = \frac{\tan 100^\circ + \tan 125^\circ}{1 - \tan 100^\circ \tan 125^\circ}$$

$$\Rightarrow 1 = \frac{\tan 100^\circ + \tan 125^\circ}{1 - \tan 100^\circ \tan 125^\circ}$$

...[$\because \tan(180^\circ + 45^\circ) = \tan 45^\circ = 1$]

$$\Rightarrow \tan 100^\circ + \tan 125^\circ + \tan 100^\circ \tan 125^\circ = 1$$

21. $\frac{\cot A}{1 + \cot A} \cdot \frac{\cot B}{1 + \cot B}$

$$= \frac{1}{(1 + \tan A)(1 + \tan B)}$$

$$= \frac{1}{\tan A + \tan B + 1 + \tan A \tan B}$$

$$= \frac{1}{1 - \tan A \tan B + 1 + \tan A \tan B}$$

... $\left[\begin{array}{l} \because \tan(A + B) = \tan 225^\circ \\ \Rightarrow \tan A + \tan B = 1 - \tan A \tan B \end{array} \right]$

$$= \frac{1}{2}$$

22. Given expression

$$= \frac{-\sin(660^\circ) \tan(1050^\circ) \sec(420^\circ)}{\cos(180^\circ + 45^\circ) \operatorname{cosec}(360^\circ - 45^\circ) \cos(360^\circ + 150^\circ)}$$

$$= \frac{-\sin(7 \times 90 + 30^\circ) \tan(3 \times 360^\circ - 30^\circ) \sec(360^\circ + 60^\circ)}{(-\cos 45^\circ)(-\operatorname{cosec} 45^\circ) \cos 150^\circ}$$

$$= \frac{\cos(30^\circ)(-\tan 30^\circ) \sec 60^\circ}{(-\cos 45^\circ)(-\operatorname{cosec} 45^\circ)(-\cos 30^\circ)}$$

$$= \frac{\cos 30^\circ \tan 30^\circ \sec 60^\circ}{\cos 45^\circ \operatorname{cosec} 45^\circ \cos 30^\circ}$$

$$= \frac{\frac{1}{\sqrt{3}} \cdot 2}{\frac{1}{\sqrt{2}} \cdot \sqrt{2}}$$

$$= \frac{2}{\sqrt{3}}$$

23. $\cos^2\left(\frac{\pi}{4} - \beta\right) - \sin^2\left(\alpha - \frac{\pi}{4}\right)$

$$= \cos\left(\frac{\pi}{4} - \beta + \alpha - \frac{\pi}{4}\right) \cos\left(\frac{\pi}{4} - \beta - \alpha + \frac{\pi}{4}\right)$$

$$= \cos(\alpha - \beta) \cos\left(\frac{\pi}{2} - (\alpha + \beta)\right)$$

$$= \cos(\alpha - \beta) \sin(\alpha + \beta)$$

24. $\cos^2 \frac{\pi}{12} + \cos^2 \frac{\pi}{4} + \cos^2 \frac{5\pi}{12}$

$$= \cos^2 15^\circ + \cos^2 45^\circ + \cos^2 75^\circ$$

$$= \cos^2 15^\circ + \cos^2 75^\circ + \cos^2 45^\circ$$

$$= \cos^2 15^\circ + \sin^2 15^\circ + \left(\frac{1}{\sqrt{2}}\right)^2$$

...[$\because \cos^2 \theta = \sin^2(90^\circ - \theta)$]

$$= 1 + \frac{1}{2}$$

$$= \frac{3}{2}$$

25. $\cos 2\alpha = \frac{1-t^2}{1+t^2} = \frac{24}{25}$ (Let $t = \tan \alpha$)

$$\sin 2\beta = \frac{2T}{1+T^2} = \frac{3}{5}$$
 (Let $T = \tan \beta$)

$$\Rightarrow \cos 2\beta = \frac{4}{5}$$

Now, $\sin 4\beta = 2 \sin 2\beta \cos 2\beta$

$$= 2 \cdot \frac{3}{5} \cdot \frac{4}{5}$$

$$= \frac{24}{25}$$

$$= \cos 2\alpha$$

26. Given that, $\cos \theta = \frac{1}{2} \left(x + \frac{1}{x}\right)$

$$\Rightarrow x + \frac{1}{x} = 2 \cos \theta$$

Now, $x^2 + \frac{1}{x^2} = \left(x + \frac{1}{x}\right)^2 - 2$

$$= (2 \cos \theta)^2 - 2$$

$$= 4 \cos^2 \theta - 2 = 2 \cos 2\theta$$

$$\therefore \frac{1}{2} \left(x^2 + \frac{1}{x^2}\right) = \frac{1}{2} \times 2 \cos 2\theta = \cos 2\theta$$



$$27. \quad \sin x + \cos x = \frac{1}{5}$$

Squaring both sides

$$1 + \sin 2x = \frac{1}{25}$$

$$\Rightarrow \sin 2x = \frac{-24}{25}$$

$$\Rightarrow \frac{2 \tan x}{1 + \tan^2 x} = \frac{-24}{25}$$

$$\Rightarrow 24 \tan^2 x + 50 \tan x + 24 = 0$$

$$\Rightarrow 12 \tan^2 x + 25 \tan x + 12 = 0$$

$$\Rightarrow (3 \tan x + 4)(4 \tan x + 3) = 0$$

$$\Rightarrow \tan x = \frac{-4}{3} \text{ or } \frac{-3}{4}$$

$$28. \quad \cos x + \sin x = \frac{1}{2}$$

$$\Rightarrow (\cos x + \sin x)^2 = \frac{1}{4}$$

$$\Rightarrow 1 + \sin 2x = \frac{1}{4}$$

$$\Rightarrow \sin 2x = -\frac{3}{4}$$

$$\Rightarrow \frac{2 \tan x}{1 + \tan^2 x} = -\frac{3}{4}$$

$$\Rightarrow 3 \tan^2 x + 8 \tan x + 3 = 0$$

$$\Rightarrow \tan x = \frac{-4 \pm \sqrt{7}}{3}$$

$$29. \quad \tan\left(\frac{\pi}{4} + \theta\right) - \tan\left(\frac{\pi}{4} - \theta\right) = \frac{1 + \tan \theta}{1 - \tan \theta} - \frac{1 - \tan \theta}{1 + \tan \theta}$$

$$= \frac{4 \tan \theta}{1 - \tan^2 \theta}$$

$$= 2 \left(\frac{2 \tan \theta}{1 - \tan^2 \theta} \right)$$

$$= 2 \tan 2\theta$$

$$30. \quad \sec 2\theta = p + \tan 2\theta$$

$$\Rightarrow \sec 2\theta - \tan 2\theta = p \quad \dots(i)$$

$$\Rightarrow \sec 2\theta + \tan 2\theta = \frac{1}{p} \quad \dots(ii)$$

$$\dots[\because \sec^2 2\theta - \tan^2 2\theta = 1]$$

Adding (i) and (ii), we get

$$2 \sec 2\theta = p + \frac{1}{p}$$

$$\Rightarrow \cos 2\theta = \frac{2p}{p^2 + 1} \Rightarrow 1 - 2 \sin^2 \theta = \frac{2p}{p^2 + 1}$$

$$\Rightarrow 2 \sin^2 \theta = 1 - \frac{2p}{p^2 + 1} \Rightarrow 2 \sin^2 \theta = \frac{(p-1)^2}{p^2 + 1}$$

$$\Rightarrow \sin^2 \theta = \frac{(p-1)^2}{2(p^2 + 1)}$$

$$31. \quad \text{Given, } \tan \theta = \frac{1}{7}, \sin \phi = \frac{1}{\sqrt{10}}$$

$$\Rightarrow \sin \theta = \frac{1}{\sqrt{50}}, \cos \theta = \frac{7}{\sqrt{50}}, \cos \phi = \frac{3}{\sqrt{10}}$$

$$\therefore \cos 2\phi = 2 \cos^2 \phi - 1 = 2 \left(\frac{9}{10} \right) - 1 = \frac{4}{5}$$

$$\sin 2\phi = 2 \sin \phi \cos \phi = 2 \left(\frac{1}{\sqrt{10}} \right) \left(\frac{3}{\sqrt{10}} \right) = \frac{3}{5}$$

$$\therefore \cos(\theta + 2\phi) = \cos \theta \cos 2\phi - \sin \theta \sin 2\phi$$

$$= \frac{7}{\sqrt{50}} \cdot \frac{4}{5} - \frac{1}{\sqrt{50}} \cdot \frac{3}{5} = \frac{28}{5\sqrt{50}} - \frac{3}{5\sqrt{50}}$$

$$= \frac{25}{5\sqrt{50}} = \frac{1}{\sqrt{2}}$$

$$\therefore \theta + 2\phi = 45^\circ$$

$$32. \quad \cos(\alpha - \beta) = \cos[\theta + \alpha - (\theta + \beta)]$$

$$= \cos(\theta + \alpha) \cos(\theta + \beta)$$

$$+ \sin(\theta + \alpha) \sin(\theta + \beta)$$

$$= \sqrt{1-a^2} \sqrt{1-b^2} + ab$$

$$\text{Now, } \cos 2(\alpha - \beta) - 4ab \cos(\alpha - \beta)$$

$$= 2 \cos^2(\alpha - \beta) - 1 - 4ab \cos(\alpha - \beta)$$

$$= 2 \left(\sqrt{1-a^2} \sqrt{1-b^2} + ab \right)^2$$

$$- 4ab \left(\sqrt{1-a^2} \sqrt{1-b^2} + ab \right) - 1$$

$$= 2 \{ (1-a^2)(1-b^2) + a^2b^2$$

$$+ 2ab \sqrt{1-a^2} \sqrt{1-b^2} \}$$

$$- 4ab \left(\sqrt{1-a^2} \sqrt{1-b^2} + ab \right) - 1$$

$$= 2(1-b^2-a^2+a^2b^2) + 2a^2b^2 - 4a^2b^2 - 1$$

$$= 2(1-a^2-b^2) - 1$$

$$= 1 - 2a^2 - 2b^2$$

$$33. \quad \tan^2 \theta = 2 \tan^2 \phi + 1$$

$$\Rightarrow 1 + \tan^2 \theta = 2(1 + \tan^2 \phi)$$

$$\Rightarrow \sec^2 \theta = 2 \sec^2 \phi$$



$$\begin{aligned} \Rightarrow \cos^2 \phi &= 2 \cos^2 \theta \\ \Rightarrow \cos^2 \phi &= 1 + \cos 2\theta \\ \Rightarrow \sin^2 \phi + \cos 2\theta &= 0 \\ &\dots[\because \sin^2 \theta + \cos^2 \theta = 1] \end{aligned}$$

34. $\tan \theta - \cot \theta = a$ and $\sin \theta + \cos \theta = b$
 $\therefore (b^2 - 1)^2 (a^2 + 4)$
 $= \{(\sin \theta + \cos \theta)^2 - 1\}^2 \{(\tan \theta - \cot \theta)^2 + 4\}$
 $= (1 + \sin 2\theta - 1)^2 (\tan^2 \theta + \cot^2 \theta - 2 + 4)$
 $= \sin^2 2\theta (\operatorname{cosec}^2 \theta + \sec^2 \theta)$
 $= 4 \sin^2 \theta \cos^2 \theta \left(\frac{1}{\sin^2 \theta} + \frac{1}{\cos^2 \theta} \right)$
 $= 4$

35. Squaring and adding the given expressions, we get
 $x^2 + y^2 = 1 + 1 + 2 \cos (2A - A)$
 $\therefore \frac{x^2 + y^2 - 2}{2} = \cos A \quad \dots(i)$

Also, $\cos A + 2 \cos^2 A - 1 = y$
 $\Rightarrow (\cos A + 1)(2 \cos A - 1) = y$
 Putting the value of $\cos A$ from (i), we get
 $(x^2 + y^2)(x^2 + y^2 - 3) = 2y$

36. $y - z = a(\cos^2 x - \sin^2 x) + 4b \sin x \cos x - c(\cos^2 x - \sin^2 x)$
 $= (a - c) \cos 2x + 2b \sin 2x$
 $= (a - c) \left(\frac{1 - \tan^2 x}{1 + \tan^2 x} \right) + 2b \left(\frac{2 \tan x}{1 + \tan^2 x} \right)$
 $= (a - c) \left\{ \frac{1 - 4b^2 / (a - c)^2}{1 + 4b^2 / (a - c)^2} \right\}$
 $\quad + 2b \left\{ \frac{2.2b / (a - c)}{1 + 4b^2 / (a - c)^2} \right\}$
 $\quad \dots \left[\because \tan x = \frac{2b}{a - c} \text{ (given)} \right]$
 $= \frac{(a - c) \{ (a - c)^2 - 4b^2 \} + 8b^2 (a - c)}{(a - c)^2 + 4b^2}$

$\therefore y - z = \frac{(a - c) \{ (a - c)^2 + 4b^2 \}}{(a - c)^2 + 4b^2} = a - c$

$\Rightarrow y \neq z \quad \dots[\because a \neq c]$

$y + z = a(\cos^2 x + \sin^2 x) + c(\sin^2 x + \cos^2 x)$
 $= a + c$

37. $8 \sin \frac{x}{8} \cos \frac{x}{2} \cos \frac{x}{4} \cos \frac{x}{8}$
 $= 4 \left(2 \sin \frac{x}{8} \cos \frac{x}{8} \right) \cos \frac{x}{2} \cos \frac{x}{4}$
 $= 4 \left(\sin \frac{x}{4} \cos \frac{x}{2} \cos \frac{x}{4} \right)$
 $\dots[\because 2 \sin A \cos A = \sin 2A]$

$= 2 \left(2 \sin \frac{x}{4} \cos \frac{x}{4} \right) \cos \frac{x}{2} = 2 \sin \frac{x}{2} \cos \frac{x}{2}$
 $= \sin x$

38. $x = \cos 10^\circ \cos 20^\circ \cos 40^\circ$
 $= \frac{1}{2 \sin 10^\circ} (2 \sin 10^\circ \cos 10^\circ \cos 20^\circ \cos 40^\circ)$
 $= \frac{1}{2.2 \sin 10^\circ} (2 \sin 20^\circ \cos 20^\circ \cos 40^\circ)$
 $= \frac{1}{2.4 \sin 10^\circ} (2 \sin 40^\circ \cos 40^\circ)$
 $= \frac{1}{8 \sin 10^\circ} (\sin 80^\circ)$
 $= \frac{1}{8 \sin 10^\circ} (\cos 10^\circ) = \frac{1}{8} \cot 10^\circ$

39. Since,
 $\cos \theta \cos 2\theta \cos 2^2 \theta \dots \cos 2^{n-1} \theta = \frac{\sin 2^n \theta}{2^n \sin \theta}$
 $= \frac{\sin(\pi - \theta)}{2^n \sin \theta}$
 $\dots \left[\because \theta = \frac{\pi}{2^n + 1} \text{ (given)} \Rightarrow 2^n \theta + \theta = \pi \right]$
 $\quad \quad \quad \Rightarrow 2^n \theta = \pi - \theta$
 $= \frac{1}{2^n}$

40. $\cos \frac{\pi}{7} \cos \frac{2\pi}{7} \cos \frac{4\pi}{7} = \left[\frac{\sin \left(2^3 \cdot \frac{\pi}{7} \right)}{2^3 \sin \left(\frac{\pi}{7} \right)} \right]$
 $= \frac{\sin \frac{8\pi}{7}}{8 \sin \frac{\pi}{7}}$
 $= -\frac{1}{8} \quad \dots \left[\because \sin \frac{8\pi}{7} = \sin \left(\pi + \frac{\pi}{7} \right) = -\sin \frac{\pi}{7} \right]$



$$\begin{aligned}
 41. \quad \cos \frac{\pi}{5} \cos \frac{2\pi}{5} \cos \frac{4\pi}{5} \cos \frac{8\pi}{5} &= \frac{\sin \frac{2^4 \pi}{5}}{2^4 \sin \frac{\pi}{5}} \\
 &= \frac{\sin \frac{16\pi}{5}}{16 \sin \frac{\pi}{5}} \\
 &= \frac{\sin \left(3\pi + \frac{\pi}{5} \right)}{16 \sin \frac{\pi}{5}} \\
 &= \frac{-\sin \frac{\pi}{5}}{16 \sin \frac{\pi}{5}} = -\frac{1}{16}
 \end{aligned}$$

$$\begin{aligned}
 42. \quad \frac{1}{\sin 10^\circ} - \frac{\sqrt{3}}{\cos 10^\circ} &= \frac{\cos 10^\circ - \sqrt{3} \sin 10^\circ}{\sin 10^\circ \cos 10^\circ} \\
 &= \frac{2 \left\{ \frac{1}{2} \cos 10^\circ - \frac{\sqrt{3}}{2} \sin 10^\circ \right\}}{\sin 10^\circ \cos 10^\circ} \\
 &= \frac{2 \sin (30^\circ - 10^\circ)}{\frac{1}{2} (\sin 20^\circ)} = 4
 \end{aligned}$$

$$\begin{aligned}
 43. \quad |\tan A| < 1 \text{ and } |A| \text{ is acute.} \\
 \therefore -\frac{\pi}{4} < A < \frac{\pi}{4} \Rightarrow \cos A > \sin A \\
 \therefore \frac{\sqrt{1 + \sin 2A} + \sqrt{1 - \sin 2A}}{\sqrt{1 + \sin 2A} - \sqrt{1 - \sin 2A}} \\
 &= \frac{\sqrt{(\cos A + \sin A)^2} + \sqrt{(\cos A - \sin A)^2}}{\sqrt{(\cos A + \sin A)^2} - \sqrt{(\cos A - \sin A)^2}} \\
 &= \frac{|\cos A + \sin A| + |\cos A - \sin A|}{|\cos A + \sin A| - |\cos A - \sin A|} \\
 &= \frac{(\cos A + \sin A) + (\cos A - \sin A)}{(\cos A + \sin A) - (\cos A - \sin A)} = \cot A
 \end{aligned}$$

$$\begin{aligned}
 44. \quad \text{Given that, } \tan x &= \frac{b}{a} \\
 \therefore \sqrt{\frac{a+b}{a-b}} + \sqrt{\frac{a-b}{a+b}} &= \sqrt{\frac{1+\frac{b}{a}}{1-\frac{b}{a}}} + \sqrt{\frac{1-\frac{b}{a}}{1+\frac{b}{a}}}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{2}{\sqrt{1-\frac{b^2}{a^2}}} = \frac{2}{\sqrt{1-\tan^2 x}} \\
 &= \frac{2}{\sqrt{1-\frac{\sin^2 x}{\cos^2 x}}} = \frac{2 \cos x}{\sqrt{\cos 2x}}
 \end{aligned}$$

$$\begin{aligned}
 45. \quad \tan (60^\circ + A) \tan (60^\circ - A) \\
 &= \frac{\sin^2 60^\circ - \sin^2 A}{\cos^2 60^\circ - \sin^2 A} \\
 &= \frac{\frac{3}{4} - \left(\frac{1 - \cos 2A}{2} \right)}{\frac{1}{4} - \left(\frac{1 - \cos 2A}{2} \right)} = \frac{3 - 2 + 2 \cos 2A}{1 - 2 + 2 \cos 2A} \\
 &= \frac{2 \cos 2A + 1}{2 \cos 2A - 1}
 \end{aligned}$$

$$\begin{aligned}
 46. \quad \sin^4 \theta + \cos^4 \theta &= (\sin^2 \theta + \cos^2 \theta)^2 \\
 &\quad - 2 \sin^2 \theta \cos^2 \theta \\
 &= 1 - \frac{1}{2} (\sin 2\theta)^2
 \end{aligned}$$

$$\text{Since, } 0 \leq \sin^2 2\theta \leq 1$$

$$\begin{aligned}
 \therefore 0 &\geq -\frac{1}{2} \sin^2 2\theta \geq -\frac{1}{2} \\
 \Rightarrow 1 + 0 &\geq 1 - \frac{1}{2} \sin^2 2\theta \geq 1 - \frac{1}{2} \\
 \Rightarrow 1 &\geq \sin^4 \theta + \cos^4 \theta \geq \frac{1}{2}
 \end{aligned}$$

$$\begin{aligned}
 47. \quad 2 \sin^2 x - \cos 2x &= 4 \sin^2 x - 1 \\
 \text{and } 0 \leq \sin^2 x \leq 1 &\Leftrightarrow 0 \leq 4 \sin^2 x \leq 4 \\
 \Rightarrow -1 &\leq 4 \sin^2 x - 1 \leq 3
 \end{aligned}$$

$$\begin{aligned}
 48. \quad 3 \sin 2\theta &= 2 \sin 3\theta \\
 \Rightarrow 6 \sin \theta \cos \theta &= 2 (3 \sin \theta - 4 \sin^3 \theta) \\
 \text{Dividing by } 2 \sin \theta \neq 0, &\text{ we get} \\
 3 \cos \theta &= 3 - 4 \sin^2 \theta \\
 \Rightarrow 3 \cos \theta &= 3 - 4 (1 - \cos^2 \theta) \\
 \Rightarrow 4 \cos^2 \theta - 3 \cos \theta - 1 &= 0 \\
 \Rightarrow \cos \theta &= 1, -\frac{1}{4}
 \end{aligned}$$

$$\text{But, } 0 < \theta < \pi$$

$$\begin{aligned}
 \therefore \cos \theta &= -\frac{1}{4} \\
 \Rightarrow \sin \theta &= \sqrt{1 - \frac{1}{16}} = \frac{\sqrt{15}}{4}
 \end{aligned}$$



$$\begin{aligned}
 49. \quad & \sin 2A = \sin 3A \\
 & \Rightarrow 2 \sin A \cos A = 3 \sin A - 4 \sin^3 A \\
 & \Rightarrow \sin A = 0 \text{ or } 2 \cos A = 3 - 4 \sin^2 A \\
 & \Rightarrow A = 0 \text{ or } 2 \cos A = 3 - 4(1 - \cos^2 A) \\
 & \Rightarrow A = 0 \text{ or } 4 \cos^2 A - 2 \cos A - 1 = 0 \\
 & \Rightarrow A = 0 \text{ or } \cos A = \frac{2 \pm \sqrt{4+16}}{2 \times 4} = \frac{1 \pm \sqrt{5}}{4} \\
 & \Rightarrow A = 0^\circ \text{ or } 36^\circ \quad \dots [\because 0 \leq A \leq 90^\circ]
 \end{aligned}$$

$$\begin{aligned}
 50. \quad & \tan \theta + \tan \left(\theta + \frac{\pi}{3} \right) + \tan \left(\theta + \frac{2\pi}{3} \right) = 3 \\
 & \Rightarrow \tan \theta + \left(\frac{\tan \theta + \sqrt{3}}{1 - \sqrt{3} \tan \theta} + \frac{\tan \theta - \sqrt{3}}{1 + \sqrt{3} \tan \theta} \right) = 3 \\
 & \Rightarrow \tan \theta + \frac{8 \tan \theta}{1 - 3 \tan^2 \theta} = 3 \\
 & \Rightarrow \frac{9 \tan \theta - 3 \tan^3 \theta}{1 - 3 \tan^2 \theta} = 3 \\
 & \Rightarrow 3 \tan 3\theta = 3 \\
 & \Rightarrow \tan 3\theta = 1
 \end{aligned}$$

$$\begin{aligned}
 51. \quad & \sin \theta = -\frac{4}{5} \\
 & \Rightarrow \cos \theta = \sqrt{1 - \frac{16}{25}} = \pm \frac{3}{5} \\
 & \Rightarrow \cos \theta = \frac{-3}{5} \quad \dots [\because \theta \text{ lies in the 3rd quadrant}]
 \end{aligned}$$

$$\begin{aligned}
 \text{Since, } \cos \frac{\theta}{2} &= \pm \sqrt{\frac{1 + \cos \theta}{2}} \\
 &= \pm \sqrt{\frac{1 - 3/5}{2}} = \pm \sqrt{\frac{1}{5}}
 \end{aligned}$$

$$\therefore \cos \frac{\theta}{2} = -\frac{1}{\sqrt{5}} \quad \dots \left[\because \frac{\theta}{2} \text{ lies in the 2nd quadrant} \right]$$

$$\begin{aligned}
 52. \quad & \text{Given that, } \sec \theta = \frac{5}{4} \\
 \text{Since, } \sec \theta &= \frac{1 + \tan^2 \left(\frac{\theta}{2} \right)}{1 - \tan^2 \left(\frac{\theta}{2} \right)} \\
 \Rightarrow \frac{5}{4} &= \frac{1 + \tan^2 \left(\frac{\theta}{2} \right)}{1 - \tan^2 \left(\frac{\theta}{2} \right)}
 \end{aligned}$$

$$\Rightarrow 5 - 5 \tan^2 \left(\frac{\theta}{2} \right) = 4 + 4 \tan^2 \left(\frac{\theta}{2} \right)$$

$$\Rightarrow 9 \tan^2 \left(\frac{\theta}{2} \right) = 1 \Rightarrow \tan \left(\frac{\theta}{2} \right) = \frac{1}{3}$$

$$53. \quad \text{For } A = 133^\circ, \frac{A}{2} = 66.5^\circ$$

$$\Rightarrow \sin \frac{A}{2} > \cos \frac{A}{2} > 0$$

$$\sqrt{1 + \sin A} = \sin \frac{A}{2} + \cos \frac{A}{2} \quad \dots (i)$$

$$\text{and } \sqrt{1 - \sin A} = \sin \frac{A}{2} - \cos \frac{A}{2} \quad \dots (ii)$$

Subtracting (ii) from (i), we get

$$2 \cos \frac{A}{2} = \sqrt{1 + \sin A} - \sqrt{1 - \sin A}$$

$$54. \quad \text{Given, } \cos \theta = \frac{3}{5} \Rightarrow \sin \theta = \frac{4}{5}$$

$$\text{and } \cos \phi = \frac{4}{5} \Rightarrow \sin \phi = \frac{3}{5}$$

$$\begin{aligned}
 \therefore \cos (\theta - \phi) &= \cos \theta \cos \phi + \sin \theta \sin \phi \\
 &= \frac{3}{5} \cdot \frac{4}{5} + \frac{4}{5} \cdot \frac{3}{5} = \frac{24}{25}
 \end{aligned}$$

$$\text{But, } 2 \cos^2 \left(\frac{\theta - \phi}{2} \right) = 1 + \cos (\theta - \phi) = 1 + \frac{24}{25}$$

$$\therefore \cos^2 \left(\frac{\theta - \phi}{2} \right) = \frac{49}{50} \quad \therefore \cos \left(\frac{\theta - \phi}{2} \right) = \frac{7}{5\sqrt{2}}$$

$$\begin{aligned}
 55. \quad & (\cos \alpha + \cos \beta)^2 + (\sin \alpha + \sin \beta)^2 \\
 &= \cos^2 \alpha + \cos^2 \beta + 2 \cos \alpha \cos \beta + \sin^2 \alpha \\
 & \quad + \sin^2 \beta + 2 \sin \alpha \sin \beta \\
 &= 2 \{ 1 + \cos (\alpha - \beta) \} = 4 \cos^2 \left(\frac{\alpha - \beta}{2} \right)
 \end{aligned}$$

$$\begin{aligned}
 56. \quad & \left(\frac{\sin 2A}{1 + \cos 2A} \right) \left(\frac{\cos A}{1 + \cos A} \right) \\
 &= \left(\frac{2 \sin A \cos A}{2 \cos^2 A} \right) \left(\frac{\cos A}{1 + \cos A} \right) \\
 &= \frac{\sin A}{1 + \cos A} \\
 &= \frac{2 \sin \frac{A}{2} \cos \frac{A}{2}}{2 \cos^2 \frac{A}{2}} \\
 &= \tan \frac{A}{2}
 \end{aligned}$$



$$57. \frac{1 + \sin A - \cos A}{1 + \sin A + \cos A} = \frac{2 \sin^2 \frac{A}{2} + 2 \sin \frac{A}{2} \cos \frac{A}{2}}{2 \cos^2 \frac{A}{2} + 2 \sin \frac{A}{2} \cos \frac{A}{2}}$$

$$= \frac{2 \sin \frac{A}{2} \left(\sin \frac{A}{2} + \cos \frac{A}{2} \right)}{2 \cos \frac{A}{2} \left(\cos \frac{A}{2} + \sin \frac{A}{2} \right)}$$

$$= \tan \frac{A}{2}$$

$$58. \frac{\tan A + \sec A - 1}{\tan A - \sec A + 1} = \frac{\sin A + 1 - \cos A}{\sin A - 1 + \cos A}$$

$$= \frac{\sin A + (1 - \cos A)}{\sin A - (1 - \cos A)}$$

$$= \frac{2 \sin \frac{A}{2} \cos \frac{A}{2} + 2 \sin^2 \frac{A}{2}}{2 \sin \frac{A}{2} \cos \frac{A}{2} - 2 \sin^2 \frac{A}{2}}$$

$$= \frac{\cos \frac{A}{2} + \sin \frac{A}{2}}{\cos \frac{A}{2} - \sin \frac{A}{2}}$$

$$= \frac{\left(\cos \frac{A}{2} + \sin \frac{A}{2} \right)^2}{\cos^2 \frac{A}{2} - \sin^2 \frac{A}{2}}$$

$$= \frac{1 + \sin A}{\cos A}$$

$$59. \frac{y+1}{1-y} = \sqrt{\frac{1+\sin \theta}{1-\sin \theta}}$$

$$\Rightarrow \frac{y+1}{1-y} = \sqrt{\frac{\left(\cos \frac{\theta}{2} + \sin \frac{\theta}{2} \right)^2}{\left(\cos \frac{\theta}{2} - \sin \frac{\theta}{2} \right)^2}}$$

$$\Rightarrow \frac{1+y}{1-y} = \frac{\cos \frac{\theta}{2} + \sin \frac{\theta}{2}}{\left| \cos \frac{\theta}{2} - \sin \frac{\theta}{2} \right|}$$

$$\Rightarrow \frac{1+y}{1-y} = \frac{\cos \frac{\theta}{2} + \sin \frac{\theta}{2}}{\cos \frac{\theta}{2} - \sin \frac{\theta}{2}}$$

$$\dots \left[\begin{array}{l} \because 0 < \theta < \frac{\pi}{2} \Rightarrow 0 < \frac{\theta}{2} < \frac{\pi}{4} \\ \Rightarrow \cos \frac{\theta}{2} > \sin \frac{\theta}{2} \end{array} \right]$$

$$\Rightarrow \frac{1+y}{1-y} = \frac{1 + \tan \frac{\theta}{2}}{1 - \tan \frac{\theta}{2}}$$

$$\Rightarrow y = \tan \frac{\theta}{2}$$

60. $\tan A$ and $\tan B$ are the roots of the equation $x^2 - ax + b = 0$.

$$\therefore \tan A + \tan B = a \text{ and } \tan A \tan B = b$$

$$\therefore \tan(A+B) = \frac{a}{1-b}$$

$$\text{Now, } \sin^2(A+B) = \frac{1}{2} \{1 - \cos 2(A+B)\}$$

$$\Rightarrow \sin^2(A+B) = \frac{1}{2} \left\{ 1 - \frac{1 - \tan^2(A+B)}{1 + \tan^2(A+B)} \right\}$$

$$\Rightarrow \sin^2(A+B) = \frac{1}{2} \left\{ 1 - \frac{1 - \frac{a^2}{(1-b)^2}}{1 + \frac{a^2}{(1-b)^2}} \right\}$$

$$\Rightarrow \sin^2(A+B) = \frac{1}{2} \left\{ \frac{2a^2}{a^2 + (1-b)^2} \right\}$$

$$= \frac{a^2}{a^2 + (1-b)^2}$$



Competitive Thinking

- $$\cos\left(\frac{\pi}{4} + x\right) + \cos\left(\frac{\pi}{4} - x\right)$$

$$= \cos \frac{\pi}{4} \cos x$$

$$- \sin \frac{\pi}{4} \sin x + \cos \frac{\pi}{4} \cos x + \sin \frac{\pi}{4} \sin x$$

$$= 2 \cos \frac{\pi}{4} \cos x$$

$$= \frac{2}{\sqrt{2}} \cos x = \sqrt{2} \cos x$$
- $$2 \sin\left(\theta + \frac{\pi}{3}\right) = \cos\left(\theta - \frac{\pi}{6}\right)$$

$$\therefore 2 \left(\sin \theta \cdot \cos \frac{\pi}{3} + \cos \theta \cdot \sin \frac{\pi}{3} \right)$$

$$= \cos \theta \cdot \cos \frac{\pi}{6} + \sin \theta \cdot \sin \frac{\pi}{6}$$



$$\therefore 2 \left(\frac{\sin \theta}{2} + \frac{\sqrt{3}}{2} \cos \theta \right) = \frac{\sqrt{3}}{2} \cos \theta + \frac{1}{2} \sin \theta$$

$$\therefore \sin \theta + \sqrt{3} \cos \theta = 0$$

$$\tan \theta = -\sqrt{3}$$

$$3. \quad \cos 15^\circ - \sin 15^\circ = \sqrt{2} \left(\frac{1}{\sqrt{2}} \cos 15^\circ - \frac{1}{\sqrt{2}} \sin 15^\circ \right)$$

$$= \sqrt{2} \cos (45^\circ + 15^\circ)$$

$$= \sqrt{2} \cos 60^\circ$$

$$= \sqrt{2} \cdot \frac{1}{2} = \frac{1}{\sqrt{2}}$$

$$4. \quad \text{We have, } \sin \theta = \frac{12}{13}$$

$$\therefore \cos \theta = \sqrt{1 - \sin^2 \theta}$$

$$= \sqrt{1 - \left(\frac{12}{13} \right)^2} = \frac{5}{13} \quad \dots \left[\because 0 < \theta < \frac{\pi}{2} \right]$$

$$\text{and } \cos \phi = \frac{-3}{5}$$

$$\therefore \sin \phi = \sqrt{1 - \frac{9}{25}} = \frac{-4}{5} \quad \dots \left[\because \pi < \phi < \frac{3\pi}{2} \right]$$

$$\therefore \sin (\theta + \phi) = \sin \theta \cdot \cos \phi + \cos \theta \cdot \sin \phi$$

$$= \left(\frac{12}{13} \right) \left(\frac{-3}{5} \right) + \left(\frac{5}{13} \right) \left(\frac{-4}{5} \right)$$

$$= \frac{-36}{65} - \frac{20}{65} = \frac{-56}{65}$$

$$5. \quad \sin \alpha = \frac{15}{17}$$

$$\Rightarrow \cos \alpha = -\frac{8}{17} \quad \dots \left[\because \frac{\pi}{2} < \alpha < \pi \right]$$

$$\tan \beta = \frac{12}{5}$$

$$\Rightarrow \sin \beta = \frac{-12}{13} \text{ and } \cos \beta = \frac{-5}{13}$$

$$\dots \left[\because \pi < \beta < \frac{3\pi}{2} \right]$$

$$\therefore \sin (\beta - \alpha) = \sin \beta \cos \alpha - \cos \beta \sin \alpha = \frac{171}{221}$$

$$6. \quad 2A = (A + B) + (A - B)$$

$$\Rightarrow \tan 2A = \frac{\tan(A + B) + \tan(A - B)}{1 - \tan(A + B)\tan(A - B)}$$

$$= \frac{p + q}{1 - pq}$$

$$7. \quad \text{Given, } \cos(\alpha + \beta) = \frac{4}{5} \Rightarrow \tan(\alpha + \beta) = \frac{3}{4}$$

$$\text{and } \sin(\alpha - \beta) = \frac{5}{13} \Rightarrow \tan(\alpha - \beta) = \frac{5}{12}$$

$$\text{Now, } \tan 2\alpha = \tan[(\alpha + \beta) + (\alpha - \beta)]$$

$$= \frac{\frac{3}{4} + \frac{5}{12}}{1 - \frac{3}{4} \cdot \frac{5}{12}} = \frac{56}{33}$$

$$8. \quad \text{Given, } \cos(A - B) = \frac{3}{5}$$

$$\therefore 5 \cos A \cos B + 5 \sin A \sin B = 3 \quad \dots \text{(i)}$$

$$\text{Also, } \tan A \tan B = 2$$

$$\therefore \sin A \sin B = 2 \cos A \cos B \quad \dots \text{(ii)}$$

From (i) and (ii), we get

$$\cos A \cos B = \frac{1}{5} \text{ and } \sin A \sin B = \frac{2}{5}$$

$$9. \quad \sin \theta = 3 \sin(\theta + 2\alpha)$$

$$\Rightarrow \sin(\theta + \alpha - \alpha) = 3 \sin(\theta + \alpha + \alpha)$$

$$\Rightarrow \sin(\theta + \alpha) \cos \alpha - \cos(\theta + \alpha) \sin \alpha$$

$$= 3 \sin(\theta + \alpha) \cos \alpha + 3 \cos(\theta + \alpha) \sin \alpha$$

$$\Rightarrow -2 \sin(\theta + \alpha) \cos \alpha = 4 \cos(\theta + \alpha) \sin \alpha$$

$$\Rightarrow \frac{-\sin(\theta + \alpha)}{\cos(\theta + \alpha)} = \frac{2 \sin \alpha}{\cos \alpha}$$

$$\Rightarrow \tan(\theta + \alpha) + 2 \tan \alpha = 0$$

$$10. \quad \tan \beta = \frac{n \sin \alpha \cos \alpha}{1 - n \sin^2 \alpha}$$

$$\Rightarrow \tan \beta = \frac{n \tan \alpha}{\sec^2 \alpha - n \tan^2 \alpha}$$

$$= \frac{n \tan \alpha}{1 + \tan^2 \alpha - n \tan^2 \alpha}$$

$$\text{Now, } \tan(\alpha - \beta) = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta}$$

$$= \frac{\tan \alpha - \frac{n \tan \alpha}{1 + \tan^2 \alpha - n \tan^2 \alpha}}{1 + \tan \alpha \left(\frac{n \tan \alpha}{1 + \tan^2 \alpha - n \tan^2 \alpha} \right)}$$

$$= \frac{\tan \alpha + \tan^3 \alpha - n \tan^3 \alpha - n \tan \alpha}{1 + \tan^2 \alpha - n \tan^2 \alpha + n \tan^2 \alpha}$$

$$= \frac{\tan \alpha (1 + \tan^2 \alpha) - n \tan \alpha (1 + \tan^2 \alpha)}{(1 + \tan^2 \alpha)}$$

$$= (1 - n)(\tan \alpha)$$



11. We have, $A - B = \frac{\pi}{4}$
 $\Rightarrow \tan(A - B) = \tan \frac{\pi}{4}$
 $\Rightarrow \frac{\tan A - \tan B}{1 + \tan A \tan B} = 1$
 $\Rightarrow \tan A - \tan B - \tan A \tan B = 1$
 $\Rightarrow \tan A - \tan B - \tan A \tan B + 1 = 2$
 $\Rightarrow (1 + \tan A)(1 - \tan B) = 2$
 $\Rightarrow y = 2$
 $\therefore (y + 1)^{y+1} = (2 + 1)^{2+1} = (3)^3 = 27$
12. Since, $\cos\left(\frac{\pi}{2}\right) = 0$
 $\therefore \cos\left[\left(\frac{\pi}{4} + \theta\right) + \left(\frac{\pi}{4} - \theta\right)\right] = 0$
 $\Rightarrow \cos\left(\frac{\pi}{4} + \theta\right)\cos\left(\frac{\pi}{4} - \theta\right)$
 $\quad - \sin\left(\frac{\pi}{4} + \theta\right)\sin\left(\frac{\pi}{4} - \theta\right) = 0$
 $\Rightarrow \cos\left(\frac{\pi}{4} + \theta\right)\cos\left(\frac{\pi}{4} - \theta\right) = \sin\left(\frac{\pi}{4} + \theta\right)\sin\left(\frac{\pi}{4} - \theta\right)$
 $\Rightarrow \cot\left(\frac{\pi}{4} + \theta\right)\cot\left(\frac{\pi}{4} - \theta\right) = 1$
13. Let $\theta = \alpha + \beta$, where $\tan \alpha = \frac{1}{2}$, $\tan \beta = \frac{1}{3}$
 $\therefore \tan \theta = \tan(\alpha + \beta) = \frac{\frac{1}{2} + \frac{1}{3}}{1 - \frac{1}{2} \cdot \frac{1}{3}} = 1 \Rightarrow \theta = \frac{\pi}{4}$
14. $\cos P = \frac{1}{7} \Rightarrow \sin P = \frac{\sqrt{48}}{7}$
 $\cos Q = \frac{13}{14} \Rightarrow \sin Q = \frac{\sqrt{27}}{14}$
 $\therefore \cos(P - Q) = \cos P \cos Q + \sin P \sin Q$
 $= \frac{1}{7} \cdot \frac{13}{14} + \frac{\sqrt{48}}{7} \cdot \frac{\sqrt{27}}{14}$
 $= \frac{13 + 36}{98} = \frac{1}{2} = \cos 60^\circ$
 $\therefore P - Q = 60^\circ$

15. We have, $\sin \alpha = \frac{1}{\sqrt{5}}$
 $\therefore \cos \alpha = \sqrt{1 - \left(\frac{1}{\sqrt{5}}\right)^2} = \frac{2}{\sqrt{5}}$
 and $\sin \beta = \frac{3}{5}$
 $\therefore \cos \beta = \sqrt{1 - \left(\frac{3}{5}\right)^2} = \frac{4}{5}$
 $\therefore \sin(\beta - \alpha) = \sin \beta \cos \alpha - \cos \beta \sin \alpha$
 $= \frac{3}{5} \times \frac{2}{\sqrt{5}} - \frac{4}{5} \times \frac{1}{\sqrt{5}} = \frac{2}{5\sqrt{5}}$
 $= 0.1789$
 Now, $\sin \frac{\pi}{4} = \frac{1}{\sqrt{2}} = 0.7071$
 Since, $0 < 0.1789 < 0.7071$
 $\therefore \sin 0 < \sin(\beta - \alpha) < \sin \frac{\pi}{4}$
 $\Rightarrow 0 < (\beta - \alpha) < \frac{\pi}{4}$
16. $\tan \theta_1 = k \cot \theta_2$
 $\Rightarrow \frac{\tan \theta_1}{\cot \theta_2} = k$
 $\Rightarrow \frac{\tan \theta_1 + \cot \theta_2}{\tan \theta_1 - \cot \theta_2} = \frac{k + 1}{k - 1}$
 $\Rightarrow \frac{\sin \theta_1 \sin \theta_2 + \cos \theta_1 \cos \theta_2}{\sin \theta_1 \sin \theta_2 - \cos \theta_1 \cos \theta_2} = \frac{k + 1}{k - 1}$
 $\Rightarrow \frac{\cos \theta_1 \cos \theta_2 + \sin \theta_1 \sin \theta_2}{\cos \theta_1 \cos \theta_2 - \sin \theta_1 \sin \theta_2} = \frac{k + 1}{1 - k}$
 $\Rightarrow \frac{\cos(\theta_1 - \theta_2)}{\cos(\theta_1 + \theta_2)} = \frac{k + 1}{1 - k}$
 $\Rightarrow \frac{\cos(\theta_1 + \theta_2)}{\cos(\theta_1 - \theta_2)} = \frac{1 - k}{1 + k}$
17. $\frac{\cos 17^\circ + \sin 17^\circ}{\cos 17^\circ - \sin 17^\circ} = \frac{1 + \tan 17^\circ}{1 - \tan 17^\circ}$
 $= \frac{\tan 45^\circ + \tan 17^\circ}{1 - \tan 45^\circ \tan 17^\circ}$
 $= \tan(45^\circ + 17^\circ)$
 $= \tan 62^\circ$



18.
$$\frac{\cos 9^\circ + \sin 9^\circ}{\cos 9^\circ - \sin 9^\circ} = \frac{1 + \tan 9^\circ}{1 - \tan 9^\circ}$$

$$= \tan (45^\circ + 9^\circ)$$

$$= \tan 54^\circ$$

19. We have, $\tan \theta = \frac{\sin \alpha - \cos \alpha}{\sin \alpha + \cos \alpha}$

$$\Rightarrow \tan \theta = \frac{\frac{1}{\sqrt{2}}(\sin \alpha - \cos \alpha)}{\frac{1}{\sqrt{2}}(\sin \alpha + \cos \alpha)}$$

$$\Rightarrow \tan \theta = \frac{\sin \alpha \cos \frac{\pi}{4} - \cos \alpha \sin \frac{\pi}{4}}{\sin \alpha \sin \frac{\pi}{4} + \cos \alpha \cos \frac{\pi}{4}}$$

$$\Rightarrow \tan \theta = \frac{\sin\left(\alpha - \frac{\pi}{4}\right)}{\cos\left(\alpha - \frac{\pi}{4}\right)}$$

$$\Rightarrow \tan \theta = \tan\left(\alpha - \frac{\pi}{4}\right)$$

$$\Rightarrow \theta = \alpha - \frac{\pi}{4} \Rightarrow \alpha = \theta + \frac{\pi}{4}$$

$\therefore \sin \alpha + \cos \alpha = \sin\left(\theta + \frac{\pi}{4}\right) + \cos\left(\theta + \frac{\pi}{4}\right)$

$$= \frac{1}{\sqrt{2}} \sin \theta + \frac{1}{\sqrt{2}} \cos \theta + \frac{1}{\sqrt{2}} \cos \theta - \frac{1}{\sqrt{2}} \sin \theta$$

$$= \frac{2}{\sqrt{2}} \cos \theta = \sqrt{2} \cos \theta$$

and $\sin \alpha - \cos \alpha = \sin\left(\theta + \frac{\pi}{4}\right) - \cos\left(\theta + \frac{\pi}{4}\right)$

$$= \frac{1}{\sqrt{2}} \sin \theta + \frac{1}{\sqrt{2}} \cos \theta - \frac{1}{\sqrt{2}} \cos \theta + \frac{1}{\sqrt{2}} \sin \theta$$

$$= \frac{2}{\sqrt{2}} \sin \theta = \sqrt{2} \sin \theta$$

20. $\cos^2 45^\circ - \sin^2 15^\circ = \cos (45 + 15)^\circ$

$$= \cos (45 - 15)^\circ$$

$$\dots [\because \cos^2 A - \sin^2 B = \cos (A+B) \cos(A - B)]$$

$$= \cos 60^\circ \cos 30^\circ$$

$$= \frac{\sqrt{3}}{4}$$

21. $\cos^2\left(\frac{\pi}{6} + \theta\right) - \sin^2\left(\frac{\pi}{6} - \theta\right)$

$$= \cos\left(\frac{\pi}{6} + \theta + \frac{\pi}{6} - \theta\right) \cos\left(\frac{\pi}{6} + \theta - \frac{\pi}{6} - \theta\right)$$

$$\dots [\because \cos^2 A - \sin^2 B = \cos (A+B) \cos(A - B)]$$

$$= \cos \frac{2\pi}{6} \cos 2\theta = \frac{1}{2} \cos 2\theta$$

22. Let $f(x) = \sin\left(x + \frac{\pi}{6}\right) + \cos\left(x + \frac{\pi}{6}\right)$. Then,

$$f(x) = \sqrt{2} \left[\cos\left(x + \frac{\pi}{6}\right) \frac{1}{\sqrt{2}} + \sin\left(x + \frac{\pi}{6}\right) \frac{1}{\sqrt{2}} \right]$$

$$= \sqrt{2} \left[\cos\left(x + \frac{\pi}{6}\right) \cos \frac{\pi}{4} + \sin\left(x + \frac{\pi}{6}\right) \sin \frac{\pi}{4} \right]$$

$$= \sqrt{2} \cos\left(x + \frac{\pi}{6} - \frac{\pi}{4}\right)$$

$$\dots [\because \cos (A - B) = \cos A \cos B + \sin A \sin B]$$

$$= \sqrt{2} \cos\left(x - \frac{\pi}{12}\right)$$

Since, $-1 \leq \cos\left(x - \frac{\pi}{12}\right) \leq 1$

$\therefore f(x)$ is maximum, if $x - \frac{\pi}{12} = 0$ i.e., if $x = \frac{\pi}{12}$

23. $\sin 15^\circ = \sin(45^\circ - 30^\circ)$

$$= \sin 45^\circ \cdot \cos 30^\circ - \cos 45^\circ \cdot \sin 30^\circ$$

$$= \left(\frac{1}{\sqrt{2}}\right) \left(\frac{\sqrt{3}}{2}\right) - \left(\frac{1}{\sqrt{2}}\right) \left(\frac{1}{2}\right) = \frac{\sqrt{3} - 1}{2\sqrt{2}}$$

24. $\sin 75^\circ = \sin(90^\circ - 15^\circ)$

$$= \cos 15^\circ$$

$$= \cos (45^\circ - 30^\circ)$$

$$= \frac{\sqrt{3} + 1}{2\sqrt{2}}$$

25. $\sin 765^\circ = \sin (720 + 45)^\circ$

$$= \sin (4\pi + 45)^\circ$$

$$= \sin (2\pi + 2\pi + 45^\circ)$$

$$= \sin 45^\circ \dots [\because \sin (2\pi + \theta) = \sin \theta]$$

$$= \frac{1}{\sqrt{2}}$$

26. $\tan \theta \sin\left(\frac{\pi}{2} + \theta\right) \cos\left(\frac{\pi}{2} - \theta\right)$

$$= \tan \theta \cos \theta \sin \theta$$

$$= \sin^2 \theta$$



27. $\cot(45^\circ + \theta) \cot(45^\circ - \theta)$
 $= \tan(90^\circ - 45^\circ - \theta) \cot(45^\circ - \theta)$
 $\dots[\because \tan(90^\circ - \theta) = \cot \theta]$
 $= \tan(45^\circ - \theta) \cot(45^\circ - \theta) = 1$

28. $\tan 75^\circ - \cot 75^\circ$
 $= \tan(90^\circ - 15^\circ) - \cot 75^\circ$
 $= \cot 15^\circ - \cot 75^\circ$
 $= (2 + \sqrt{3}) - (2 - \sqrt{3})$
 $= 2\sqrt{3}$

29. $\tan 50^\circ = \tan(70^\circ - 20^\circ) = \frac{\tan 70^\circ - \tan 20^\circ}{1 + \tan 70^\circ \tan 20^\circ}$
 $\Rightarrow \tan 50^\circ + \tan 70^\circ \tan 20^\circ \tan 50^\circ$
 $= \tan 70^\circ - \tan 20^\circ$
 $\Rightarrow \tan 50^\circ + \tan 50^\circ = \tan 70^\circ - \tan 20^\circ$
 $\dots[\because \tan 70^\circ = \cot 20^\circ]$
 $\Rightarrow 2 \tan 50^\circ + \tan 20^\circ = \tan 70^\circ$

30. $\sec 50^\circ + \tan 50^\circ$
 $= \frac{1}{\cos 50^\circ} + \tan 50^\circ$
 $= \frac{\cos 20^\circ}{\cos 20^\circ \cos 50^\circ} + \tan 50^\circ$
 $= \frac{\sin 70^\circ}{\cos 20^\circ \cos 50^\circ} + \tan 50^\circ$
 $\dots[\because \cos \theta = \sin(90^\circ - \theta)]$
 $= \frac{\sin(50^\circ + 20^\circ)}{\cos 20^\circ \cos 50^\circ} + \tan 50^\circ$
 $= \frac{\sin 50^\circ \cos 20^\circ + \cos 50^\circ \sin 20^\circ}{\cos 20^\circ \cos 50^\circ} + \tan 50^\circ$
 $= \frac{\sin 50^\circ \cos 20^\circ}{\cos 20^\circ \cos 50^\circ} + \frac{\cos 50^\circ \sin 20^\circ}{\cos 20^\circ \cos 50^\circ} + \tan 50^\circ$
 $= \tan 50^\circ + \tan 20^\circ + \tan 50^\circ$
 $= 2 \tan 50^\circ + \tan 20^\circ$

31. $2 \sec 2\alpha = \tan \beta + \cot \beta$
 $\Rightarrow \frac{2}{\cos 2\alpha} = \frac{\sin \beta}{\cos \beta} + \frac{\cos \beta}{\sin \beta}$
 $= \frac{\sin^2 \beta + \cos^2 \beta}{\cos \beta \sin \beta} = \frac{1}{\cos \beta \sin \beta}$
 $\Rightarrow \cos 2\alpha = \sin 2\beta \Rightarrow \cos 2\alpha = \cos\left(\frac{\pi}{2} - 2\beta\right)$
 $\Rightarrow 2\alpha = \frac{\pi}{2} - 2\beta \Rightarrow 2\alpha + 2\beta = \frac{\pi}{2}$
 $\Rightarrow \alpha + \beta = \frac{\pi}{4}$

32. $\sin(\pi + \theta) \sin(\pi - \theta) \operatorname{cosec}^2 \theta$
 $= -\sin \theta \sin \theta \frac{1}{\sin^2 \theta}$
 $= -1$

33. Given that, ABCD is a cyclic quadrilateral.
 So, $A + C = 180^\circ \Rightarrow A = 180^\circ - C$
 $\Rightarrow \cos A = \cos(180^\circ - C) = -\cos C$
 $\Rightarrow \cos A + \cos C = 0 \dots(i)$
 Similarly, $\cos B + \cos D = 0 \dots(ii)$
 Adding, (i) and (ii), we get
 $\cos A + \cos B + \cos C + \cos D = 0$

34. $\cos(270^\circ + \theta) \cos(90^\circ - \theta) - \sin(270^\circ - \theta) \cos \theta$
 $= \sin \theta \cdot \sin \theta + \cos \theta \cdot \cos \theta = 1$

35. $\cos A + \sin(270^\circ + A) - \sin(270^\circ - A)$
 $+ \cos(180^\circ + A)$
 $= \cos A - \cos A + \cos A - \cos A = 0$

36. $\tan A + \cot(180^\circ + A) + \cot(90^\circ + A)$
 $+ \cot(360^\circ - A)$
 $= \tan A + \cot A - \tan A - \cot A = 0$

37. $\sin 1^\circ + \sin 2^\circ + \dots + \sin 359^\circ$
 $= (\sin 1^\circ + \sin 359^\circ) + (\sin 2^\circ + \sin 358^\circ) + \dots$
 $+ (\sin 179^\circ + \sin 181^\circ) + \sin 180^\circ$
 $= (\sin 1^\circ - \sin 1^\circ) + (\sin 2^\circ - \sin 2^\circ) + \dots$
 $+ (\sin 179^\circ - \sin 179^\circ) + \sin 180^\circ$
 $= 0$

38. $\sin 600^\circ \cos 330^\circ + \cos 120^\circ \sin 150^\circ$
 $= -\sin 60^\circ \cos 30^\circ - \sin 30^\circ \cos 60^\circ$
 $= -\{\sin(60^\circ + 30^\circ)\}$
 $= -1$

39. Let $f(x) = 2 \sin 3x + 3 \cos 3x$
 $\therefore f\left(\frac{5\pi}{6}\right) = 2 \sin\left(\frac{5\pi}{2}\right) + 3 \cos\left(\frac{5\pi}{2}\right)$
 $= 2 \sin\left(2\pi + \frac{\pi}{2}\right) + 3 \cos\left(2\pi + \frac{\pi}{2}\right)$
 $= 2 \sin \frac{\pi}{2} + 3 \cos \frac{\pi}{2} = 2(1) + 3(0) = 2$

40. $\frac{1 - \tan 2^\circ \cot 62^\circ}{\tan 152^\circ - \cot 88^\circ} = \frac{1 - \tan 2^\circ \cot 62^\circ}{\tan(90^\circ + 62^\circ) - \cot(90^\circ - 2^\circ)}$
 $= \frac{1 - \tan 2^\circ \cot 62^\circ}{-\cot 62^\circ - \tan 2^\circ} = \frac{\tan 62^\circ - \tan 2^\circ}{-(1 + \tan 2^\circ \tan 62^\circ)}$
 $= -\tan(62^\circ - 2^\circ)$
 $= -\tan 60^\circ = -\sqrt{3}$



$$41. \frac{\cos 12^\circ - \sin 12^\circ}{\cos 12^\circ + \sin 12^\circ} + \frac{\sin 147^\circ}{\cos 147^\circ}$$

$$= \frac{1 - \tan 12^\circ}{1 + \tan 12^\circ} + \tan 147^\circ$$

$$= \tan(45^\circ - 12^\circ) + \tan(180^\circ - 33^\circ)$$

$$= \tan 33^\circ - \tan 33^\circ = 0$$

$$42. \frac{\tan 160^\circ - \tan 110^\circ}{1 + (\tan 160^\circ)(\tan 110^\circ)}$$

$$= \frac{\tan(180^\circ - 160^\circ) - \cot(90^\circ - 110^\circ)}{1 + [\tan(180^\circ - 160^\circ) \cot(90^\circ - 110^\circ)]}$$

$$= \frac{-\tan 20^\circ + \cot 20^\circ}{1 + (-\tan 20^\circ)(-\cot 20^\circ)}$$

$$= \frac{-\lambda + \frac{1}{\lambda}}{1 + 1}$$

$$= \frac{1 - \lambda^2}{2\lambda}$$

$$43. \sin^2 17.5^\circ + \sin^2 72.5^\circ$$

$$= \sin^2 17.5^\circ + [\sin(90^\circ - 17.5^\circ)]^2$$

$$= \sin^2 17.5^\circ + \cos^2 17.5^\circ$$

$$= 1 = \tan^2 45^\circ$$

$$44. 3 \left\{ \sin^4 \left(\frac{3\pi}{2} - \alpha \right) + \sin^4(3\pi + \alpha) \right\}$$

$$- 2 \left\{ \sin^6 \left(\frac{\pi}{2} + \alpha \right) + \sin^6(5\pi - \alpha) \right\}$$

$$= 3 \{ (-\cos \alpha)^4 + (-\sin \alpha)^4 \} - 2(\cos^6 \alpha + \sin^6 \alpha)$$

$$= 3(1 - 2 \sin^2 \alpha \cos^2 \alpha) - 2(1 - 3 \sin^2 \alpha \cos^2 \alpha)$$

$$= 3 - 6 \sin^2 \alpha \cos^2 \alpha - 2 + 6 \sin^2 \alpha \cos^2 \alpha$$

$$= 3 - 2 = 1$$

$$45. \sin^2 5^\circ + \sin^2 10^\circ + \sin^2 15^\circ + \dots + \sin^2 85^\circ + \sin^2 90^\circ$$

Since, $\sin 90^\circ = 1$ or $\sin^2 90^\circ = 1$

Similarly, $\sin 45^\circ = \frac{1}{\sqrt{2}}$ or $\sin^2 45^\circ = \frac{1}{2}$ and

the angles are in A.P. of 18 terms.

Also, $\sin^2 85^\circ = [\sin(90^\circ - 5^\circ)]^2 = \cos^2 5^\circ$

Therefore from the complementary rule, we have $\sin^2 5^\circ + \sin^2 85^\circ = \sin^2 5^\circ + \cos^2 5^\circ = 1$

$$\therefore \sin^2 5^\circ + \sin^2 10^\circ + \sin^2 15^\circ + \dots + \sin^2 85^\circ + \sin^2 90^\circ$$

$$= (1 + 1 + 1 + 1 + 1 + 1 + 1 + 1) + 1 + \frac{1}{2}$$

$$= 9 \frac{1}{2}$$

$$46. \cos 15^\circ = \sqrt{\frac{1 + \cos(2 \times 15^\circ)}{2}}$$

$$= \sqrt{\frac{1 + \cos 30^\circ}{2}}$$

$$47. \sin 15^\circ = \sin(45^\circ - 30^\circ) = \frac{\sqrt{3} - 1}{2\sqrt{2}} = \text{irrational}$$

$$\cos 15^\circ = \cos(45^\circ - 30^\circ) = \frac{\sqrt{3} + 1}{2\sqrt{2}} = \text{irrational}$$

$$\sin 15^\circ \cos 15^\circ = \frac{1}{2} (2 \sin 15^\circ \cos 15^\circ)$$

$$= \frac{1}{2} \sin 30^\circ = \frac{1}{2} \cdot \frac{1}{2}$$

$$= \frac{1}{4} = \text{rational}$$

$$\sin 15^\circ \cos 75^\circ = \sin 15^\circ \sin 15^\circ$$

$$= \sin^2 15^\circ$$

$$= \left(\frac{\sqrt{3} - 1}{2\sqrt{2}} \right)^2$$

$$= \frac{4 - 2\sqrt{3}}{8} = \text{irrational}$$

$$48. \frac{1}{8} (3 - 4 \cos 2\theta + \cos 4\theta)$$

$$= \frac{1}{8} (3 - 4 \cos 2\theta + 2 \cos^2 2\theta - 1)$$

$$= \frac{1}{8} (2 \cos^2 2\theta - 4 \cos 2\theta + 2)$$

$$= \frac{1}{4} (\cos^2 2\theta - 2 \cos 2\theta + 1)$$

$$= \frac{1}{4} (\cos 2\theta - 1)^2$$

$$= \frac{1}{4} (-2 \sin^2 \theta)^2 \quad \dots [\because \cos 2\theta = 1 - 2 \sin^2 \theta]$$

$$= \frac{1}{4} (4 \sin^4 \theta) = \sin^4 \theta$$

$$49. \sec 2x - \tan 2x = \frac{1 - \sin 2x}{\cos 2x}$$

$$= \frac{(\cos x - \sin x)^2}{(\cos^2 x - \sin^2 x)}$$

$$= \frac{\cos x - \sin x}{\cos x + \sin x}$$

$$= \frac{1 - \tan x}{1 + \tan x}$$



$$\begin{aligned} &= \frac{\tan \frac{\pi}{4} - \tan x}{1 + \tan\left(\frac{\pi}{4}\right)\tan x} \\ &= \tan\left(\frac{\pi}{4} - x\right) \end{aligned}$$

$$\begin{aligned} 50. \quad \sqrt{3} \operatorname{cosec} 20^\circ - \sec 20^\circ &= \frac{\sqrt{3}}{\sin 20^\circ} - \frac{1}{\cos 20^\circ} \\ &= \frac{\sqrt{3} \cos 20^\circ - \sin 20^\circ}{\sin 20^\circ \cos 20^\circ} \\ &= \frac{2 \left[\frac{\sqrt{3}}{2} \cos 20^\circ - \frac{1}{2} \sin 20^\circ \right]}{\frac{2}{2} \sin 20^\circ \cos 20^\circ} \\ &= \frac{4 \cos(20^\circ + 30^\circ)}{\sin 40^\circ} = \frac{4 \cos 50^\circ}{\sin 40^\circ} \\ &= \frac{4 \sin 40^\circ}{\sin 40^\circ} \\ &= 4 \end{aligned}$$

$$\begin{aligned} 51. \quad \tan(1^\circ) + \tan(89^\circ) &= \tan 1^\circ + \cot 1^\circ \quad \dots [\because \tan(90^\circ - \theta) = \cot \theta] \\ &= \frac{\tan^2 1^\circ + 1}{\tan 1^\circ} \\ &= \frac{\sec^2 1^\circ}{\tan 1^\circ} = \frac{1}{\sin 1^\circ \cos 1^\circ} = \frac{2}{\sin 2^\circ} \end{aligned}$$

$$\begin{aligned} 52. \quad \cot 2\theta + \tan \theta &= \frac{1}{\tan 2\theta} + \tan \theta \\ &= \frac{1 - \tan^2 \theta}{2 \tan \theta} + \tan \theta \\ &= \frac{1 + \tan^2 \theta}{2 \tan \theta} = \frac{1}{\sin 2\theta} \\ &= \operatorname{cosec} 2\theta \end{aligned}$$

$$\begin{aligned} \text{Now, } \cot \frac{2x}{3} + \tan \frac{x}{3} &= \operatorname{cosec} \frac{kx}{3} \\ \Rightarrow \operatorname{cosec} \frac{2x}{3} &= \operatorname{cosec} \frac{kx}{3} \\ \Rightarrow k &= 2 \end{aligned}$$

$$\begin{aligned} 53. \quad 2 \sin^2 \left[\left(\frac{\pi}{2} \right) \cos^2 x \right] &= 1 - \cos(\pi \sin 2x) \\ \Rightarrow 2 \sin^2 \left[\left(\frac{\pi}{2} \right) \cos^2 x \right] &= 2 \sin^2 \left[\frac{\pi \sin 2x}{2} \right] \\ \Rightarrow \cos^2 x = \sin 2x &\Rightarrow \cos^2 x = 2 \sin x \cos x \end{aligned}$$

$$\Rightarrow \tan x = \frac{1}{2}$$

$$\text{Now, } \cos 2x = \frac{1 - \tan^2 x}{1 + \tan^2 x} = \frac{1 - \frac{1}{4}}{1 + \frac{1}{4}} = \frac{3}{5}$$

$$\begin{aligned} 54. \quad 8 \cos 2\theta + 8 \sec 2\theta &= 65 \\ \Rightarrow 8 \cos^2 2\theta + 8 &= 65 \cos 2\theta \\ \Rightarrow 8 \cos^2 2\theta - 65 \cos 2\theta + 8 &= 0 \\ \Rightarrow (\cos 2\theta - 8)(8 \cos 2\theta - 1) &= 0 \end{aligned}$$

Since, $\cos 2\theta \in [-1, 1]$

$$\therefore \cos 2\theta = \frac{1}{8}$$

$$\text{Now, } 4 \cos 4\theta = 4(2 \cos^2 2\theta - 1)$$

$$= 4 \left[2 \left(\frac{1}{8} \right)^2 - 1 \right] = -\frac{31}{8}$$

$$55. \quad 5(\tan^2 x - \cos^2 x) = 2 \cos 2x + 9$$

Put $\cos^2 x = t$

$$\therefore 5 \left(\frac{1-t}{t} - t \right) = 2(2t-1) + 9$$

$$\Rightarrow 5(1-t-t^2) = 4t^2 - 2t + 9t$$

$$\Rightarrow 9t^2 + 12t - 5 = 0$$

$$\Rightarrow 9t^2 + 15t - 3t - 5 = 0$$

$$\Rightarrow 3t(3t+5) - 1(3t+5) = 0$$

$$\Rightarrow (3t+5)(3t-1) = 0$$

$$\Rightarrow t = \frac{1}{3} \text{ or } t = \frac{-5}{3}$$

But t cannot be negative

$$\therefore t = \frac{1}{3}$$

$$\therefore \cos^2 x = \frac{1}{3}$$

$$\Rightarrow \cos 2x = 2 \cos^2 x - 1$$

$$= \frac{2}{3} - 1 = \frac{-1}{3}$$

$$\Rightarrow \cos 4x = 2 \cos^2 2x - 1$$

$$= 2 \left(\frac{-1}{3} \right)^2 - 1$$

$$= \frac{-7}{9}$$

$$56. \quad x + \frac{1}{x} = 2 \cos \alpha$$

Squaring on both sides, we get

$$x^2 + \frac{1}{x^2} + 2 = 4 \cos^2 \alpha$$



$$\Rightarrow x^2 + \frac{1}{x^2} = 4 \cos^2 \alpha - 2$$

$$\Rightarrow x^2 + \frac{1}{x^2} = 2(2 \cos^2 \alpha - 1) \\ = 2 \cos 2\alpha$$

Similarly, $x^n + \frac{1}{x^n} = 2 \cos n\alpha$

57. $\sin x + \cos x = \frac{1}{5}$

$$\Rightarrow \sin^2 x + \cos^2 x + 2 \sin x \cos x = \frac{1}{25}$$

$$\Rightarrow \sin 2x = \frac{-24}{25} \Rightarrow \cos 2x = \frac{-7}{25}$$

$$\Rightarrow \tan 2x = \frac{24}{7}$$

58. $3 \tan A - 4 = 0$

$$\Rightarrow \tan A = \frac{4}{3}$$

$$\Rightarrow \sin A = -\frac{4}{5}, \cos A = -\frac{3}{5}$$

$$\therefore 5 \sin 2A + 3 \sin A + 4 \cos A \\ = 10 \sin A \cos A + 3 \sin A + 4 \cos A \\ = 10 \left(\frac{12}{25} \right) - \frac{12}{5} - \frac{12}{5} = 0$$

59. $a \cos 2\theta + b \sin 2\theta = a \left(\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right) + b \left(\frac{2 \tan \theta}{1 + \tan^2 \theta} \right)$

$$= a \left(\frac{1 - \frac{b^2}{a^2}}{1 + \frac{b^2}{a^2}} \right) + b \left(\frac{\frac{2b}{a}}{1 + \frac{b^2}{a^2}} \right)$$

$$\dots \left[\because \tan \theta = \frac{b}{a} \text{ (given)} \right]$$

$$= a \left(\frac{a^2 - b^2}{a^2 + b^2} \right) + b \left(\frac{2ba}{a^2 + b^2} \right)$$

$$= \frac{1}{(a^2 + b^2)} (a^3 - ab^2 + 2ab^2) = \frac{a(a^2 + b^2)}{a^2 + b^2}$$

$$= a$$

60. $a \cos 2\theta + b \sin 2\theta = c$

$$\Rightarrow a \left(\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right) + b \left(\frac{2 \tan \theta}{1 + \tan^2 \theta} \right) = c$$

$$\Rightarrow a - a \tan^2 \theta + 2b \tan \theta = c + c \tan^2 \theta$$

$$\Rightarrow -(a + c) \tan^2 \theta + 2b \tan \theta + (a - c) = 0$$

$$\therefore \tan \alpha + \tan \beta = -\frac{2b}{-(c+a)} = \frac{2b}{c+a}$$

61. $6 \cos \theta + 8 \sin \theta = 9 \dots (i)$

$$\Rightarrow 8 \sin \theta = 9 - 6 \cos \theta$$

Squaring on both sides, we get

$$64 \sin^2 \theta = 81 - 108 \cos \theta + 36 \cos^2 \theta$$

$$\Rightarrow 64(1 - \cos^2 \theta) = 81 - 108 \cos \theta + 36 \cos^2 \theta$$

$$\Rightarrow 100 \cos^2 \theta - 108 \cos \theta + 17 = 0$$

$$\therefore \cos \alpha \cos \beta = \frac{17}{100}$$

From (i), $6 \cos \theta = 9 - 8 \sin \theta$

Squaring on both sides, we get

$$36 \cos^2 \theta = 81 - 144 \sin \theta + 64 \sin^2 \theta$$

$$\Rightarrow 36(1 - \sin^2 \theta) = 81 - 144 \sin \theta + 64 \sin^2 \theta$$

$$\Rightarrow 100 \sin^2 \theta - 144 \sin \theta + 45 = 0$$

$$\therefore \sin \alpha \sin \beta = \frac{45}{100}$$

Now, $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$

$$= \frac{17}{100} - \frac{45}{100} = \frac{-28}{100} = \frac{-14}{50}$$

$$\therefore \sin(\alpha + \beta) = \sqrt{1 - \cos^2(\alpha + \beta)} \\ = \sqrt{1 - \left(\frac{-14}{50} \right)^2} = \frac{1}{50} \sqrt{2500 - 196} \\ = \frac{48}{50} = \frac{24}{25}$$

62. $25 \cos^2 \alpha + 5 \cos \alpha - 12 = 0$

$$\Rightarrow \cos \alpha = \frac{-5 \pm \sqrt{25 + 1200}}{50} = \frac{-5 \pm 35}{50}$$

$$\Rightarrow \cos \alpha = \frac{-4}{5} \dots \left[\because \frac{\pi}{2} < \alpha < \pi \Rightarrow \cos \alpha < 0 \right]$$

$$\Rightarrow \sin \alpha = \sqrt{1 - \left(\frac{-4}{5} \right)^2} = \frac{3}{5}$$

$$\therefore \sin 2\alpha = 2 \sin \alpha \cos \alpha = \frac{-24}{25}$$

63. $2 \cos^2 \theta - 2 \sin^2 \theta = 1$

$$\Rightarrow 2 \cos 2\theta = 1 \Rightarrow \cos 2\theta = \frac{1}{2} = \cos 60^\circ$$

$$\Rightarrow 2\theta = 60^\circ \Rightarrow \theta = 30^\circ$$

64. $2 \sin A \cos^3 A - 2 \sin^3 A \cos A$

$$= 2 \sin A \cos A (\cos^2 A - \sin^2 A)$$

$$= 2 \sin A \cos A \cos 2A$$

$$= \sin 2A \cos 2A$$

$$= \frac{1}{2} \sin 4A$$



$$\begin{aligned}
 65. \quad & \sin^4 \frac{\pi}{8} + \sin^4 \frac{3\pi}{8} + \sin^4 \frac{5\pi}{8} + \sin^4 \frac{7\pi}{8} \\
 &= \frac{1}{4} \left[\left(2 \sin^2 \frac{\pi}{8} \right)^2 + \left(2 \sin^2 \frac{3\pi}{8} \right)^2 \right] \\
 &\quad + \frac{1}{4} \left[\left(2 \sin^2 \frac{\pi}{8} \right)^2 + \left(2 \sin^2 \frac{3\pi}{8} \right)^2 \right] \\
 &= \frac{1}{4} \left[\left(1 - \cos \frac{\pi}{4} \right)^2 + \left(1 - \cos \frac{3\pi}{4} \right)^2 \right] \\
 &\quad + \frac{1}{4} \left[\left(1 - \cos \frac{\pi}{4} \right)^2 + \left(1 - \cos \frac{3\pi}{4} \right)^2 \right] \\
 &= \frac{1}{4} \left[\left(1 - \frac{1}{\sqrt{2}} \right)^2 + \left(1 + \frac{1}{\sqrt{2}} \right)^2 \right] \\
 &\quad + \frac{1}{4} \left[\left(1 - \frac{1}{\sqrt{2}} \right)^2 + \left(1 + \frac{1}{\sqrt{2}} \right)^2 \right] \\
 &= \frac{1}{4} (3) + \frac{1}{4} (3) = \frac{3}{2}
 \end{aligned}$$

$$\begin{aligned}
 66. \quad & 3(\sin x - \cos x)^4 + 6(\sin x + \cos x)^2 \\
 &\quad + 4(\sin^6 x + \cos^6 x) \\
 &= 3(\sin^2 x + \cos^2 x - 2 \sin x \cos x)^2 + 6(\sin^2 x \\
 &\quad + \cos^2 x + 2 \sin x \cos x) + 4(\sin^6 x + \cos^6 x) \\
 &= 3(1 - \sin 2x)^2 + 6(1 + \sin 2x) \\
 &\quad + 4(1 - 3 \sin^2 x \cos^2 x) \\
 &= 3 + 3 \sin^2 2x - 6 \sin 2x + 6 + 6 \sin 2x \\
 &\quad + 4 - 3 \sin^2 2x \\
 &= 9 + 4 + 3 \sin^2 2x - 3 \sin^2 2x = 13
 \end{aligned}$$

$$\begin{aligned}
 67. \quad & \sin 2\theta + \sin 2\phi = \frac{1}{2} \quad \dots(i) \\
 & \text{and } \cos 2\theta + \cos 2\phi = \frac{3}{2} \quad \dots(ii) \\
 & \text{Squaring and adding (i) and (ii), we get} \\
 & (\sin^2 2\theta + \cos^2 2\theta) + (\sin^2 2\phi + \cos^2 2\phi) \\
 & \quad + 2(\sin 2\theta \sin 2\phi + \cos 2\theta \cos 2\phi) = \frac{1}{4} + \frac{9}{4} \\
 & \Rightarrow \cos 2\theta \cos 2\phi + \sin 2\theta \sin 2\phi = \frac{1}{4} \\
 & \Rightarrow \cos (2\theta - 2\phi) = \frac{1}{4} \\
 & \Rightarrow 2 \cos^2 (\theta - \phi) - 1 = \frac{1}{4} \\
 & \Rightarrow \cos^2 (\theta - \phi) = \frac{5}{8}
 \end{aligned}$$

$$\begin{aligned}
 68. \quad & \cos \alpha \cos 2\alpha \cos 2^2 \alpha \cos 2^3 \alpha \dots \cos 2^{n-1} \alpha \\
 &= \frac{2^n \sin \alpha \cos \alpha \cos 2\alpha \cos 2^2 \alpha \dots \cos 2^{n-1} \alpha}{2^n \sin \alpha} \\
 &= \frac{\sin \{2(2^{n-1} \alpha)\}}{2^n \sin \alpha} \\
 & \quad \text{(using } 2 \sin \theta \cos \theta = \sin 2\theta \text{ again and again)} \\
 &= \frac{\sin 2^n \alpha}{2^n \sin \alpha}
 \end{aligned}$$

$$\begin{aligned}
 69. \quad & \cos \frac{2\pi}{15} \cos \frac{4\pi}{15} \cos \frac{8\pi}{15} \cos \frac{16\pi}{15} \\
 &= \frac{\sin 2^4 \frac{2\pi}{15}}{2^4 \sin \frac{2\pi}{15}} = \frac{\sin \frac{32\pi}{15}}{16 \sin \frac{2\pi}{15}} \\
 &= \frac{\sin \left(2\pi + \frac{2\pi}{15} \right)}{16 \sin \frac{2\pi}{15}} = \frac{\sin \frac{2\pi}{15}}{16 \sin \frac{2\pi}{15}} = \frac{1}{16}
 \end{aligned}$$

$$\begin{aligned}
 70. \quad & k = \sin \frac{\pi}{18} \sin \frac{5\pi}{18} \sin \frac{7\pi}{18} \\
 &= \cos \left(\frac{\pi}{2} - \frac{\pi}{18} \right) \cos \left(\frac{\pi}{2} - \frac{5\pi}{18} \right) \cos \left(\frac{\pi}{2} - \frac{7\pi}{18} \right) \\
 &= \cos \frac{\pi}{9} \cos \frac{2\pi}{9} \cos \frac{4\pi}{9} = \frac{\sin 2^3 \frac{\pi}{9}}{2^3 \sin \frac{\pi}{9}} \\
 &= \frac{\sin \frac{8\pi}{9}}{8 \sin \frac{\pi}{9}} = \frac{\sin \left(\pi - \frac{\pi}{9} \right)}{8 \sin \frac{\pi}{9}} = \frac{1}{8}
 \end{aligned}$$

$$\begin{aligned}
 71. \quad & \sin \frac{\pi}{14} \sin \frac{3\pi}{14} \sin \frac{5\pi}{14} \sin \frac{7\pi}{14} \sin \frac{9\pi}{14} \sin \frac{11\pi}{14} \sin \frac{13\pi}{14} \\
 &= \sin \frac{\pi}{14} \sin \frac{3\pi}{14} \sin \frac{5\pi}{14} \times 1 \\
 &\quad \times \sin \left(\pi - \frac{5\pi}{14} \right) \sin \left(\pi - \frac{3\pi}{14} \right) \sin \left(\pi - \frac{\pi}{14} \right) \\
 &= \left(\sin \frac{\pi}{14} \sin \frac{3\pi}{14} \sin \frac{5\pi}{14} \right)^2 \\
 &= \left(\cos \frac{6\pi}{14} \cos \frac{4\pi}{14} \cos \frac{2\pi}{14} \right)^2 \\
 &\quad \dots \left[\because \sin \left(\frac{\pi}{2} - \theta \right) = \cos \theta \right]
 \end{aligned}$$



$$\begin{aligned}
 &= \left(\cos \frac{\pi}{7} \cos \frac{2\pi}{7} \cos \frac{3\pi}{7} \right)^2 \\
 &= \left(-\cos \frac{\pi}{7} \cos \frac{2\pi}{7} \cos \frac{4\pi}{7} \right)^2 \\
 &\quad \dots \left[\because \cos \frac{3\pi}{7} = \cos \left(\pi - \frac{4\pi}{7} \right) = -\cos \frac{4\pi}{7} \right] \\
 &= \left(-\frac{\sin \frac{2^3 \pi}{7}}{2^3 \sin \frac{\pi}{7}} \right)^2 = \frac{1}{64} \left(-\frac{\sin \frac{8\pi}{7}}{\sin \frac{\pi}{7}} \right)^2 \\
 &= \frac{1}{64} \quad \dots \left[\because \sin \frac{8\pi}{7} = \sin \left(\pi + \frac{\pi}{7} \right) = -\sin \frac{\pi}{7} \right]
 \end{aligned}$$

72. $\sin \left(\frac{31}{3} \pi \right) = \sin \left(10\pi + \frac{\pi}{3} \right)$
 $= \sin \frac{\pi}{3}$
 $= \frac{\sqrt{3}}{2}$

73. Since, $\tan \alpha + 2 \tan 2\alpha + 2^2 \tan 2^2 \alpha + \dots$
 $+ 2^n \tan 2^n \alpha + 2^{n+1} \cot 2^{n+1} \alpha = \cot \alpha \quad \forall n \in \mathbb{N}$
 Here, $\alpha = \frac{\pi}{5}$

$$\therefore \tan \frac{\pi}{5} + 2 \tan \frac{2\pi}{5} + 4 \cot \frac{4\pi}{5} = \cot \frac{\pi}{5}$$

74. $\frac{\cot x - \tan x}{\cot 2x} = \frac{\cos^2 x - \sin^2 x}{\sin x \cos x} \times \frac{\sin 2x}{\cos 2x}$
 $= \frac{2 \cos 2x}{\sin 2x} \times \frac{\sin 2x}{\cos 2x} = 2$

75. $\frac{\sec 8A - 1}{\sec 4A - 1} = \frac{1 - \cos 8A}{\cos 8A} \cdot \frac{\cos 4A}{1 - \cos 4A}$
 $= \frac{2 \sin^2 4A}{\cos 8A} \cdot \frac{\cos 4A}{2 \sin^2 2A}$
 $= \frac{2 \sin 4A \cos 4A}{\cos 8A} \cdot \frac{\sin 4A}{2 \sin^2 2A}$
 $= \tan 8A \cdot \frac{2 \sin 2A \cos 2A}{2 \sin^2 2A}$
 $= \frac{\tan 8A}{\tan 2A}$

76. $2 \tan A = 3 \tan B$
 $\Rightarrow \tan A = \frac{3}{2} \tan B = \frac{3}{2} t$ [Let $\tan B = t$]

$$\sin 2B = \frac{2t}{1+t^2}, \quad \cos 2B = \frac{1-t^2}{1+t^2}$$

$$\begin{aligned}
 \therefore \frac{\sin 2B}{5 - \cos 2B} &= \frac{\left(\frac{2t}{1+t^2} \right)}{5 - \left(\frac{1-t^2}{1+t^2} \right)} = \frac{2t}{4+6t^2} \\
 &= \frac{t}{2+3t^2} = \frac{\frac{3t}{2} - t}{1 + \frac{3t^2}{2}} \\
 &= \tan(A - B)
 \end{aligned}$$

77. $\cos 2\alpha = \frac{3 \cos 2\beta - 1}{3 - \cos 2\beta}$

By componendo – dividendo, we get

$$\frac{\cos 2\alpha + 1}{\cos 2\alpha - 1} = \frac{3 \cos 2\beta - 1 + 3 - \cos 2\beta}{3 \cos 2\beta - 1 - (3 - \cos 2\beta)}$$

$$\Rightarrow \frac{2 \cos^2 \alpha}{-2 \sin^2 \alpha} = \frac{2 + 2 \cos 2\beta}{4 \cos 2\beta - 4}$$

$$\Rightarrow \frac{-\cos^2 \alpha}{\sin^2 \alpha} = \frac{1 + \cos 2\beta}{2(\cos 2\beta - 1)} = \frac{2 \cos^2 \beta}{-4 \sin^2 \beta}$$

$$\Rightarrow \frac{\sin^2 \alpha}{\cos^2 \alpha} = \frac{2 \sin^2 \beta}{\cos^2 \beta} \Rightarrow \tan^2 \alpha = 2 \tan^2 \beta$$

$$\Rightarrow \tan \alpha = \sqrt{2} \tan \beta$$

78. Since, $\cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta$

$$\therefore \cos 3\theta = 4 \left[\frac{1}{2^3} \left(a + \frac{1}{a} \right)^3 \right] - 3 \left[\frac{1}{2} \left(a + \frac{1}{a} \right) \right]$$

$$\Rightarrow \cos 3\theta = \frac{1}{2} \left(a + \frac{1}{a} \right) \left[\left(a + \frac{1}{a} \right)^2 - 3 \right]$$

$$\Rightarrow \cos 3\theta = \frac{1}{2} \left(a^3 + \frac{1}{a^3} \right)$$

79. $\cos^3 \theta + \cos^3 (120^\circ - \theta) + \cos^3 (120^\circ + \theta)$
 $= \frac{3}{4} \cos(3\theta)$

$$\begin{aligned}
 \therefore \cos^3 10^\circ + \cos^3 110^\circ + \cos^3 130^\circ \\
 = \cos^3(10^\circ) + \cos^3(120^\circ - 10^\circ) \\
 \quad \quad \quad + \cos^3(120^\circ + 10^\circ)
 \end{aligned}$$

$$= \frac{3}{4} \cos(3 \times 10^\circ) = \frac{3}{4} \times \frac{\sqrt{3}}{2} = \frac{3\sqrt{3}}{8}$$



$$\begin{aligned}
 80. \quad \sin 6\theta &= 2 \sin 3\theta \cos 3\theta \\
 &= 2(3 \sin \theta - 4 \sin^3 \theta)(4 \cos^3 \theta - 3 \cos \theta) \\
 &= 24 \sin \theta \cos \theta (\sin^2 \theta + \cos^2 \theta) - 18 \sin \theta \cos \theta \\
 &\quad - 32 \sin^3 \theta \cos^3 \theta \\
 &= 6 \sin \theta \cos \theta - 32 \sin \theta \cos^3 \theta \sin^2 \theta \\
 &= 3 \sin 2\theta - 32 \sin \theta \cos^3 \theta (1 - \cos^2 \theta) \\
 &= 32 \cos^5 \theta \sin \theta - 32 \cos^3 \theta \sin \theta + 3 \sin 2\theta \\
 \text{Given, } \sin 6\theta &= 32 \cos^5 \theta \sin \theta - 32 \cos^3 \theta \sin \theta + 3 \sin 2\theta \\
 \therefore \text{ On comparing, we get } x &= \sin 2\theta
 \end{aligned}$$

$$\begin{aligned}
 81. \quad \tan \left(\frac{\pi}{4} + \frac{\theta}{2} \right) + \tan \left(\frac{\pi}{4} - \frac{\theta}{2} \right) \\
 &= \frac{1 + \tan \frac{\theta}{2}}{1 - \tan \frac{\theta}{2}} + \frac{1 - \tan \frac{\theta}{2}}{1 + \tan \frac{\theta}{2}} \\
 &= \frac{\left(1 + \tan \frac{\theta}{2} \right)^2 + \left(1 - \tan \frac{\theta}{2} \right)^2}{1 - \tan^2 \frac{\theta}{2}} \\
 &= 2 \left(\frac{1 + \tan^2 \frac{\theta}{2}}{1 - \tan^2 \frac{\theta}{2}} \right) \\
 &= \frac{2}{\cos \theta} \\
 &= 2 \sec \theta
 \end{aligned}$$

$$\begin{aligned}
 82. \quad \text{Given, } \tan x &= \frac{3}{4} \\
 \Rightarrow \cos x &= -\frac{4}{5} \quad \dots \left[\because \pi < x < \frac{3\pi}{2} \right] \\
 \text{Since, } 1 + \cos x &= 2 \cos^2 \frac{x}{2} \\
 \therefore 1 - \frac{4}{5} &= 2 \cos^2 \frac{x}{2} \quad \therefore \cos^2 \frac{x}{2} = \frac{1}{10} \\
 \therefore \cos \frac{x}{2} &= -\frac{1}{\sqrt{10}} \quad \dots \left[\because \pi < x < \frac{3\pi}{2} \Rightarrow \frac{\pi}{2} < \frac{x}{2} < \frac{3\pi}{4} \right]
 \end{aligned}$$

$$\begin{aligned}
 83. \quad \sin A &= \frac{4}{5} \\
 \Rightarrow \tan A &= -\frac{4}{3} \quad \dots \left[\because 90^\circ < A < 180^\circ \right] \\
 \text{Now, } \tan A &= \frac{2 \tan \frac{A}{2}}{1 - \tan^2 \frac{A}{2}} \quad (\text{Let } \tan \frac{A}{2} = P)
 \end{aligned}$$

$$\begin{aligned}
 &\Rightarrow -\frac{4}{3} = \frac{2P}{1 - P^2} \\
 &\Rightarrow 4P^2 - 6P - 4 = 0 \\
 &\Rightarrow P = \frac{-1}{2} \text{ or } P = 2
 \end{aligned}$$

$$P = \frac{-1}{2} \text{ is not possible}$$

$$\therefore P = 2 \Rightarrow \tan \frac{A}{2} = 2$$

$$84. \quad \text{Given, } \sin \frac{\theta}{2} = \sqrt{\frac{x-1}{2x}}$$

$$\therefore \cos \frac{\theta}{2} = \sqrt{1 - \sin^2 \frac{\theta}{2}} = \sqrt{\frac{x+1}{2x}}$$

$$\text{and } \tan \frac{\theta}{2} = \frac{\sqrt{x-1}}{\sqrt{x+1}}$$

$$\text{Since, } \tan \theta = \frac{2 \tan \frac{\theta}{2}}{1 - \tan^2 \frac{\theta}{2}}$$

$$\therefore \tan \theta = \sqrt{x^2 - 1}$$

$$\begin{aligned}
 85. \quad \sin x &= \frac{2 \tan \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} \\
 &= \frac{2y}{1 + y^2} \quad \dots \left[\text{Let } y = \tan \left(\frac{x}{2} \right) \right]
 \end{aligned}$$

$$\therefore \tan \frac{x}{2} = \operatorname{cosec} x - \sin x = \frac{1}{\sin x} - \sin x$$

$$\Rightarrow y = \frac{1 + y^2}{2y} - \frac{2y}{1 + y^2}$$

$$\Rightarrow 2y^2 (1 + y^2) = 1 + y^4 + 2y^2 - 4y^2$$

$$\Rightarrow 1 - y^4 - 4y^2 = 0 \Rightarrow y^4 + 4y^2 - 1 = 0$$

$$\Rightarrow y^2 = \frac{-4 \pm \sqrt{16 + 4}}{2} = \frac{-4 \pm \sqrt{20}}{2}$$

$$\Rightarrow \tan^2 \left(\frac{x}{2} \right) = -2 \pm \sqrt{5}$$

$$\text{But, } \tan^2 \left(\frac{x}{2} \right) \neq -2 - \sqrt{5}$$

$$\Rightarrow \tan^2 \left(\frac{x}{2} \right) = -2 + \sqrt{5}$$



$$86. \quad \cos \theta = \frac{\cos \alpha - \cos \beta}{1 - \cos \alpha \cos \beta}$$

By componendo – dividendo, we get

$$\frac{\cos \theta + 1}{\cos \theta - 1} = \frac{\cos \alpha - \cos \beta + 1 - \cos \alpha \cos \beta}{\cos \alpha - \cos \beta - (1 - \cos \alpha \cos \beta)}$$

$$\Rightarrow \frac{\cos \theta + 1}{\cos \theta - 1} = \frac{(1 + \cos \alpha)(1 - \cos \beta)}{-(1 - \cos \alpha)(1 + \cos \beta)}$$

$$\Rightarrow \frac{1 + \cos \theta}{1 - \cos \theta} = \frac{1 + \cos \alpha}{1 - \cos \alpha} \times \frac{1 - \cos \beta}{1 + \cos \beta}$$

$$\Rightarrow \cot^2 \frac{\theta}{2} = \cot^2 \frac{\alpha}{2} \tan^2 \frac{\beta}{2}$$

$$\Rightarrow \tan^2 \frac{\theta}{2} = \tan^2 \frac{\alpha}{2} \cot^2 \frac{\beta}{2}$$

$$\Rightarrow \tan \frac{\theta}{2} = \pm \tan \frac{\alpha}{2} \cot \frac{\beta}{2}$$

$$87. \quad \sqrt{4\cos^4 \theta + \sin^2 2\theta} + 4\cot \theta \cos^2 \left(\frac{\pi}{4} - \frac{\theta}{2} \right)$$

$$= \sqrt{4\cos^4 \theta + 4\sin^2 \theta \cos^2 \theta}$$

$$+ 4 \cot \theta \left[\frac{1 + \cos 2 \left(\frac{\pi}{4} - \frac{\theta}{2} \right)}{2} \right]$$

$$= \sqrt{4\cos^2 \theta (\cos^2 \theta + \sin^2 \theta)}$$

$$+ 2 \cot \theta \left[1 + \cos \left(\frac{\pi}{2} - \theta \right) \right]$$

$$= |2\cos \theta| + 2 \cot \theta + 2 \cos \theta$$

$$= 2 \cot \theta \quad \dots \left[\because \theta \in \left(\frac{\pi}{2}, \frac{3\pi}{2} \right) \right]$$

$$88. \quad \text{Put } \tan \left(\frac{\theta}{2} \right) = t$$

$$\therefore (m+2) \left(\frac{2t}{1+t^2} \right) + (2m-1) \left(\frac{1-t^2}{1+t^2} \right) = 2m+1$$

$$\Rightarrow (2m+4)t + (2m-1)(1-t^2) = (2m+1)(1+t^2)$$

$$\Rightarrow 4mt^2 - (2m+4)t + 2 = 0$$

$$\Rightarrow 2mt^2 - mt - 2t + 1 = 0$$

$$\Rightarrow mt(2t-1) - 1(2t-1) = 0$$

$$\Rightarrow (2t-1)(mt-1) = 0$$

$$\Rightarrow t = \frac{1}{2} \text{ or } t = \frac{1}{m}$$

If $t = \frac{1}{2}$, then

$$\tan \theta = \frac{2t}{1-t^2} = \frac{1}{1-\frac{1}{4}} = \frac{4}{3}$$

If $t = \frac{1}{m}$, then

$$\tan \theta = \frac{\frac{2}{m}}{1 - \left(\frac{1}{m^2} \right)} = \frac{2m}{m^2 - 1}$$

$$89. \quad \cos \left(\frac{\alpha - \beta}{2} \right) = 2 \cos \left(\frac{\alpha + \beta}{2} \right)$$

$$\Rightarrow \cos \frac{\alpha}{2} \cos \frac{\beta}{2} + \sin \frac{\alpha}{2} \sin \frac{\beta}{2}$$

$$= 2 \cos \frac{\alpha}{2} \cos \frac{\beta}{2} - 2 \sin \frac{\alpha}{2} \sin \frac{\beta}{2}$$

$$\Rightarrow 3 \sin \frac{\alpha}{2} \sin \frac{\beta}{2} = \cos \frac{\alpha}{2} \cos \frac{\beta}{2}$$

$$\Rightarrow \tan \frac{\alpha}{2} \tan \frac{\beta}{2} = \frac{1}{3}$$

$$90. \quad \text{Since, } \tan \frac{A}{2} = \frac{1 - \cos A}{\sin A}$$

Putting $\frac{A}{2} = \left(7\frac{1}{2} \right)^\circ$, we get

$$\tan \left(7\frac{1}{2} \right)^\circ = \frac{1 - \cos 15^\circ}{\sin 15^\circ}$$

$$= \frac{1 - \frac{\sqrt{3}+1}{2\sqrt{2}}}{\frac{\sqrt{3}-1}{2\sqrt{2}}}$$

$$= \frac{2\sqrt{2} - \sqrt{3} - 1}{\sqrt{3}-1} \times \frac{\sqrt{3}+1}{\sqrt{3}+1}$$

$$= \sqrt{6} - \sqrt{3} + \sqrt{2} - 2$$

$$91. \quad \cot \frac{A}{2} = \frac{1 + \cos A}{\sin A}$$

Putting $A = \left(7\frac{1}{2} \right)^\circ$, we get

$$\cot \left(7\frac{1}{2} \right)^\circ = \frac{1 + \cos 15^\circ}{\sin 15^\circ}$$

$$= \frac{1 + \frac{\sqrt{3}+1}{2\sqrt{2}}}{\frac{\sqrt{3}-1}{2\sqrt{2}}}$$

$$= \frac{2\sqrt{2} + \sqrt{3} + 1}{\sqrt{3}-1} \times \frac{\sqrt{3}+1}{\sqrt{3}+1}$$

$$= \sqrt{6} + \sqrt{2} + \sqrt{3} + \sqrt{4}$$



92. Since, $\sin\left(22\frac{1}{2}^\circ\right) = \frac{1}{2}\sqrt{2-\sqrt{2}} = \cos\left(67\frac{1}{2}^\circ\right)$
 and $\cos\left(22\frac{1}{2}^\circ\right) = \frac{1}{2}\sqrt{2+\sqrt{2}} = \sin\left(67\frac{1}{2}^\circ\right)$
 $\therefore \left[1 + \cos\left(22\frac{1}{2}^\circ\right)\right] \left[1 + \cos\left(67\frac{1}{2}^\circ\right)\right]$
 $\left[1 + \cos\left(112\frac{1}{2}^\circ\right)\right] \left[1 + \cos\left(157\frac{1}{2}^\circ\right)\right]$
 $= \left[1 + \cos\left(22\frac{1}{2}^\circ\right)\right] \left[1 + \cos\left(67\frac{1}{2}^\circ\right)\right]$
 $\left[1 - \sin\left(22\frac{1}{2}^\circ\right)\right] \left[1 - \sin\left(67\frac{1}{2}^\circ\right)\right]$
[$\because \cos(90^\circ + \theta) = -\sin \theta$]
 $= \left[1 + \frac{1}{2}\sqrt{2+\sqrt{2}}\right] \left[1 + \frac{1}{2}\sqrt{2-\sqrt{2}}\right]$
 $\left[1 - \frac{1}{2}\sqrt{2-\sqrt{2}}\right] \left[1 - \frac{1}{2}\sqrt{2+\sqrt{2}}\right]$
 $= \left[1 - \frac{1}{4}(2+\sqrt{2})\right] \left[1 - \frac{1}{4}(2-\sqrt{2})\right]$
 $= \frac{(4-2-\sqrt{2})(4-2+\sqrt{2})}{16}$
 $= \frac{(2-\sqrt{2})(2+\sqrt{2})}{16} = \frac{4-2}{16} = \frac{1}{8}$

93. $\tan A = \frac{1-\cos B}{\sin B} = \frac{2\sin^2\left(\frac{B}{2}\right)}{2\sin\left(\frac{B}{2}\right)\cos\left(\frac{B}{2}\right)} = \tan \frac{B}{2}$

$\Rightarrow \tan 2A = \tan B$

94. Given, $\operatorname{cosec} \theta = \frac{p+q}{p-q} \Rightarrow \frac{1}{\sin \theta} = \frac{p+q}{p-q}$

By componendo – dividendo, we get

$$\frac{1 + \sin \theta}{1 - \sin \theta} = \frac{p+q+p-q}{p+q-p+q}$$

$$\Rightarrow \left\{ \frac{\cos \frac{\theta}{2} + \sin \frac{\theta}{2}}{\cos \frac{\theta}{2} - \sin \frac{\theta}{2}} \right\}^2 = \frac{p}{q}$$

$$\Rightarrow \left\{ \frac{1 + \tan \frac{\theta}{2}}{1 - \tan \frac{\theta}{2}} \right\}^2 = \frac{p}{q} \Rightarrow \left\{ \frac{\tan \frac{\pi}{4} + \tan \frac{\theta}{2}}{1 - \tan \frac{\pi}{4} \tan \frac{\theta}{2}} \right\}^2 = \frac{p}{q}$$

$$\Rightarrow \tan^2\left(\frac{\pi}{4} + \frac{\theta}{2}\right) = \frac{p}{q} \Rightarrow \cot^2\left(\frac{\pi}{4} + \frac{\theta}{2}\right) = \frac{q}{p}$$

$$\Rightarrow \cot\left(\frac{\pi}{4} + \frac{\theta}{2}\right) = \sqrt{\frac{q}{p}}$$

95. $\frac{\sqrt{2} - \sin \alpha - \cos \alpha}{\sin \alpha - \cos \alpha}$
 $= \frac{\sqrt{2} - \sqrt{2} \left\{ \frac{1}{\sqrt{2}} \sin \alpha + \frac{1}{\sqrt{2}} \cos \alpha \right\}}{\sqrt{2} \left\{ \frac{1}{\sqrt{2}} \sin \alpha - \frac{1}{\sqrt{2}} \cos \alpha \right\}}$
 $= \frac{\sqrt{2} - \sqrt{2} \cos\left(\alpha - \frac{\pi}{4}\right)}{\sqrt{2} \sin\left(\alpha - \frac{\pi}{4}\right)}$
 $= \frac{\sqrt{2}(1 - \cos \theta)}{\sqrt{2} \sin \theta}$, where $\theta = \alpha - \frac{\pi}{4}$
 $= \frac{2\sin^2\left(\frac{\theta}{2}\right)}{2\sin\left(\frac{\theta}{2}\right)\cos\left(\frac{\theta}{2}\right)}$
 $= \tan \frac{\theta}{2} = \tan\left(\frac{\alpha}{2} - \frac{\pi}{8}\right)$ [$\because \theta = \alpha - \frac{\pi}{4}$]

96. Putting $\theta = \phi = \frac{\pi}{4}$ in the given expression,

we get

$$\cos 2\left(\frac{\pi}{2}\right) - 4 \cos\left(\frac{\pi}{2}\right) \sin\left(\frac{\pi}{4}\right) \sin\left(\frac{\pi}{4}\right) + 2\sin^2\left(\frac{\pi}{4}\right) = 0$$

Put $\theta = \phi = \frac{\pi}{4}$ in option (A), then

$$\cos 2\theta = \cos \frac{\pi}{2} = 0$$

Hence, option (A) is the correct answer.

97. Given that $\sin \theta + \sin \phi = a$ (i)
 and $\cos \theta + \cos \phi = b$ (ii)
 Squaring (i) and (ii) and adding, we get
 $2 + 2(\sin \theta \sin \phi + \cos \theta \cos \phi) = a^2 + b^2$
 $\Rightarrow 2 \cos(\theta - \phi) = a^2 + b^2 - 2$
 $\Rightarrow \cos(\theta - \phi) = \frac{a^2 + b^2 - 2}{2}$

$$\Rightarrow \frac{1 - \tan^2 \frac{\theta - \phi}{2}}{1 + \tan^2 \frac{\theta - \phi}{2}} = \frac{a^2 + b^2 - 2}{2}$$



$$\Rightarrow (a^2 + b^2) + (a^2 + b^2) \tan^2 \frac{\theta - \phi}{2} - 2 - 2 \tan^2 \frac{\theta - \phi}{2} = 2 - 2 \tan^2 \frac{\theta - \phi}{2}$$

$$\Rightarrow \frac{4 - a^2 - b^2}{a^2 + b^2} = \tan^2 \frac{\theta - \phi}{2}$$

$$\Rightarrow \tan\left(\frac{\theta - \phi}{2}\right) = \sqrt{\frac{4 - a^2 - b^2}{a^2 + b^2}}$$

Trick : Putting $\theta = \frac{\pi}{4} = \phi$, we get $\tan \frac{\theta - \phi}{2} = 0$, which is given by option (B).

98.
$$\frac{1 + \sin 2\alpha}{\cos(2\alpha - 2\pi) \tan\left(\alpha - \frac{3\pi}{4}\right)}$$

$$= \frac{1 + 2 \sin \alpha \cos \alpha}{\cos 2\alpha \left(\frac{\tan \alpha + 1}{1 - \tan \alpha}\right)}$$

$$= \frac{(1 + 2 \sin \alpha \cos \alpha)^2}{\cos 2\alpha \left(\frac{\cos \alpha + \sin \alpha}{\cos \alpha - \sin \alpha}\right)}$$

$$= \frac{(1 + 2 \sin \alpha \cos \alpha)^2}{\cos 2\alpha \left(\frac{\cos \alpha + \sin \alpha}{\cos \alpha - \sin \alpha}\right)}$$

$$= \frac{(1 + 2 \sin \alpha \cos \alpha)^2}{\cos 2\alpha \left(\frac{\cos \alpha + \sin \alpha}{\cos \alpha - \sin \alpha}\right)}$$

$$= \frac{(1 + 2 \sin \alpha \cos \alpha)^2}{\cos 2\alpha \left(\frac{\cos \alpha + \sin \alpha}{\cos \alpha - \sin \alpha}\right)}$$

99. When $\cos 4\theta = \frac{1}{3}$, then $2 \cos^2 2\theta - 1 = \frac{1}{3}$

$$\Rightarrow 2 \cos^2 2\theta = \frac{4}{3} \Rightarrow \cos^2 2\theta = \frac{2}{3}$$

$$\Rightarrow \cos 2\theta = \pm \sqrt{\frac{2}{3}} \quad \dots \left[\begin{array}{l} \because \theta \in \left(0, \frac{\pi}{4}\right) \cup \left(\frac{\pi}{4}, \frac{\pi}{2}\right), \\ \therefore 2\theta \in \left(0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right) \end{array} \right]$$

$$\therefore f\left(\frac{1}{3}\right) = f(\cos 4\theta) = \frac{2}{2 - \sec^2 \theta}$$

$$= \frac{2 \cos^2 \theta}{2 \cos^2 \theta - 1} = \frac{1 + \cos 2\theta}{\cos 2\theta}$$

$$\therefore f\left(\frac{1}{3}\right) = 1 \pm \sqrt{\frac{3}{2}} \quad \dots \left[\because \cos 2\theta = \pm \sqrt{\frac{2}{3}} \right]$$

100.
$$f_n(\theta) = \left(\tan \frac{\theta}{2}\right) (1 + \sec \theta)(1 + \sec 2\theta)$$

$$(1 + \sec 4\theta) \dots (1 + \sec 2^n \theta)$$

$$= \left(\tan \frac{\theta}{2}\right) \left(\frac{1 + \cos \theta}{\cos \theta}\right) (1 + \sec 2\theta)$$

$$(1 + \sec 4\theta) \dots (1 + \sec 2^n \theta)$$

$$= \left(\frac{\sin \frac{\theta}{2}}{\cos \frac{\theta}{2}}\right) \times \frac{2 \cos^2 \frac{\theta}{2}}{\cos \theta} (1 + \sec 2\theta)$$

$$(1 + \sec 4\theta) \dots (1 + \sec 2^n \theta)$$

$$= \frac{\sin \theta}{\cos \theta} (1 + \sec 2\theta)(1 + \sec 4\theta) \dots (1 + \sec 2^n \theta)$$

$$\dots [\because \sin 2A = 2 \sin A \cos A]$$

$$= \tan \theta \left(\frac{1 + \cos 2\theta}{\cos 2\theta}\right) (1 + \sec 4\theta) \dots (1 + \sec 2^n \theta)$$

$$= \frac{\sin \theta}{\cos \theta} \left(\frac{2 \cos^2 \theta}{\cos 2\theta}\right) (1 + \sec 4\theta) \dots (1 + \sec 2^n \theta)$$

$$= \tan 2\theta (1 + \sec 4\theta) \dots (1 + \sec 2^n \theta)$$

$$= \tan 2^n \theta$$

$$\therefore f_2\left(\frac{\pi}{16}\right) = \tan\left(2^2 \times \frac{\pi}{16}\right) = 1$$

$$f_3\left(\frac{\pi}{32}\right) = \tan\left(2^3 \times \frac{\pi}{32}\right) = 1$$

$$f_4\left(\frac{\pi}{64}\right) = \tan\left(2^4 \times \frac{\pi}{64}\right) = 1$$

$$f_5\left(\frac{\pi}{128}\right) = \tan\left(2^5 \times \frac{\pi}{128}\right) = 1$$

\therefore option (D) is incorrect.

101.
$$\sum_{n=1}^{\infty} \sin\left(\frac{n! \pi}{720}\right) = \left(\sin \frac{1! \pi}{720} + \sin \frac{2! \pi}{720} + \dots + \frac{\sin 5! \pi}{720}\right)$$

$$+ \sum_{n=6}^{\infty} \sin \frac{n! \pi}{720}$$

$$= \sin\left(\frac{\pi}{6}\right) + \sin\left(\frac{\pi}{30}\right) + \sin\left(\frac{\pi}{120}\right) + \sin\left(\frac{\pi}{360}\right)$$

$$+ \sin\left(\frac{\pi}{720}\right)$$

$$\dots \left[\because \sum_{n=6}^{\infty} \sin \frac{n! \pi}{720} = 0 \right]$$



Evaluation Test

$$1. \quad x \cos \theta = y \cos \left(\theta + \frac{2\pi}{3} \right) = z \cos \left(\theta + \frac{4\pi}{3} \right) = k \text{ (say)}$$

$$\Rightarrow \cos \theta = \frac{k}{x}, \quad \cos \left(\theta + \frac{2\pi}{3} \right) = \frac{k}{y}$$

$$\text{and } \cos \left(\theta + \frac{4\pi}{3} \right) = \frac{k}{z}$$

$$\therefore \frac{k}{x} + \frac{k}{y} + \frac{k}{z}$$

$$= \cos \theta + \cos \left(\theta + \frac{2\pi}{3} \right) + \cos \left(\theta + \frac{4\pi}{3} \right)$$

$$= \cos \theta + \cos \left(\pi - \left(\frac{\pi}{3} - \theta \right) \right) + \cos \left(\pi + \left(\frac{\pi}{3} + \theta \right) \right)$$

$$= \cos \theta - \cos \left(\frac{\pi}{3} - \theta \right) - \cos \left(\frac{\pi}{3} + \theta \right)$$

$$= \cos \theta - \left[\cos \left(\frac{\pi}{3} - \theta \right) + \cos \left(\frac{\pi}{3} + \theta \right) \right]$$

$$= \cos \theta - 2 \cos \frac{\pi}{3} \cos \theta$$

$$= \cos \theta - 2 \times \frac{1}{2} \cos \theta$$

$$\therefore \frac{k}{x} + \frac{k}{y} + \frac{k}{z} = 0$$

$$\Rightarrow \frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 0$$

$$2. \quad \tan 70^\circ - \tan 20^\circ - 2 \tan 40^\circ$$

$$= (\cot 20^\circ - \tan 20^\circ) - 2 \tan 40^\circ$$

$$\dots [\because \tan 70^\circ = \tan(90^\circ - 20^\circ) = \cot 20^\circ]$$

$$= 2 \cot 40^\circ - 2 \tan 40^\circ$$

$$\dots [\because \cot \theta - \tan \theta = 2 \cot 2\theta]$$

$$= 2(\cot 40^\circ - \tan 40^\circ)$$

$$= 2(2 \cot 80^\circ) = 4 \cot 80^\circ$$

$$= 4 \cot(90^\circ - 10^\circ) = 4 \tan 10^\circ$$

$$3. \quad \text{Given, } \sqrt{x} + \frac{1}{\sqrt{x}} = 2 \cos \theta$$

Squaring on both sides, we get

$$x + \frac{1}{x} + 2 = 4 \cos^2 \theta$$

$$\Rightarrow x + \frac{1}{x} = 4 \cos^2 \theta - 2$$

$$\Rightarrow x + \frac{1}{x} = 2(2 \cos^2 \theta - 1) = 2 \cos 2\theta$$

Again, squaring on both sides, we get

$$x^2 + \frac{1}{x^2} + 2 = 4 \cos^2 2\theta$$

$$\Rightarrow x^2 + \frac{1}{x^2} = 4 \cos^2 2\theta - 2$$

$$\Rightarrow x^2 + \frac{1}{x^2} = 2(2 \cos^2 2\theta - 1)$$

$$\Rightarrow x^2 + \frac{1}{x^2} = 2 \cos 4\theta \quad \dots (i)$$

Cubing on both sides, we get

$$\left(x^2 + \frac{1}{x^2} \right)^3 = (2 \cos 4\theta)^3$$

$$\Rightarrow x^6 + \frac{1}{x^6} + 3x^2 \times \frac{1}{x^2} \left(x^2 + \frac{1}{x^2} \right) = 8 \cos^3 4\theta$$

$$\Rightarrow x^6 + \frac{1}{x^6} + 3(2 \cos 4\theta) = 8 \cos^3 4\theta$$

....[From (i)]

$$\Rightarrow x^6 + \frac{1}{x^6} = 8 \cos^3 4\theta - 6 \cos 4\theta$$

$$= 2(4 \cos^3 4\theta - 3 \cos 4\theta)$$

$$= 2 \cos 3(4\theta)$$

$$\dots [\because \cos 3A = 4 \cos^3 A - 3 \cos A]$$

$$= 2 \cos 12\theta$$

$$4. \quad \cos^3 \theta + \cos^3(\theta + 120^\circ) + \cos^3(\theta - 120^\circ)$$

$$= \frac{\cos 3\theta + 3 \cos \theta}{4}$$

$$+ \frac{\cos(3\theta + 360^\circ) + 3 \cos(\theta + 120^\circ)}{4}$$

$$+ \frac{\cos(3\theta - 360^\circ) + 3 \cos(\theta - 120^\circ)}{4}$$

$$\dots \left[\begin{array}{l} \because \cos 3A = 4 \cos^3 A - 3 \cos A, \\ \therefore \cos^3 A = \frac{\cos 3A + 3 \cos A}{4} \end{array} \right]$$

$$= \frac{\cos 3\theta}{4} + \frac{3 \cos \theta}{4} + \frac{\cos 3\theta}{4} + \frac{3 \cos(\theta + 120^\circ)}{4}$$

$$+ \frac{\cos 3\theta}{4} + \frac{3 \cos(\theta - 120^\circ)}{4}$$

$$= \frac{3}{4} \cos 3\theta + \frac{3}{4} \{ \cos(\theta + 120^\circ)$$

$$+ \cos(\theta - 120^\circ) + \cos \theta \}$$

$$= \frac{3}{4} \cos 3\theta + \frac{3}{4} \{ 2 \cos \theta \cos 120^\circ + \cos \theta \}$$



$$\begin{aligned} &= \frac{3}{4} \cos 3\theta + \frac{3}{4} \{2\cos\theta (-\sin 30^\circ) + \cos\theta\} \\ &= \frac{3}{4} \cos 3\theta + \frac{3}{4} \left\{ 2\cos\theta \left(-\frac{1}{2}\right) + \cos\theta \right\} \\ &= \frac{3}{4} \cos 3\theta + \frac{3}{4} (-\cos\theta + \cos\theta) \\ &= \frac{3}{4} \cos 3\theta \end{aligned}$$

5. $\tan 2\alpha = \tan(\alpha + \alpha)$

$$\begin{aligned} &= \frac{\frac{1}{5} + \frac{1}{5}}{1 - \frac{1}{25}} \end{aligned}$$

$\therefore \tan 2\alpha = \frac{5}{12}$

$\tan 4\alpha = \tan(2\alpha + 2\alpha)$

$$\begin{aligned} &= \frac{\frac{5}{12} + \frac{5}{12}}{1 - \frac{25}{144}} \\ &= \frac{120}{119} \end{aligned}$$

$\therefore \tan(4\alpha - \beta) = \frac{\frac{120}{119} - \frac{1}{239}}{1 + \frac{120}{119} \times \frac{1}{239}}$

$$\begin{aligned} &= \frac{120 \times 239 - 119}{119 \times 239 + 120} \\ &= \frac{(119+1)239 - 119}{119 \times 239 + 120} \\ &= \frac{119 \times 239 + (239 - 119)}{119 \times 239 + 120} = 1 \end{aligned}$$

6. Given, $\sec(\theta + \phi)$, $\sec\theta$ and $\sec(\theta - \phi)$ are in A. P.

$\therefore 2 \sec\theta = \sec(\theta + \phi) + \sec(\theta - \phi)$

$$\Rightarrow \frac{2}{\cos\theta} = \frac{1}{\cos(\theta + \phi)} + \frac{1}{\cos(\theta - \phi)}$$

$$\Rightarrow \frac{2}{\cos\theta} = \frac{\cos(\theta - \phi) + \cos(\theta + \phi)}{\cos(\theta + \phi)\cos(\theta - \phi)}$$

$$\Rightarrow \frac{2}{\cos\theta} = \frac{2\cos\theta\cos\phi}{\cos^2\theta - \sin^2\phi}$$

... [$\because \cos(A + B)\cos(A - B) = \cos^2A - \sin^2B$]

$$\Rightarrow \cos^2\theta\cos\phi = \cos^2\theta - \sin^2\phi$$

$$\Rightarrow \cos^2\theta - \cos^2\theta\cos\phi = \sin^2\phi$$

$$\Rightarrow \cos^2\theta(1 - \cos\phi) = (1 - \cos^2\phi)$$

$$\Rightarrow \cos^2\theta = 1 + \cos\phi$$

$$\Rightarrow \cos^2\theta = 2\cos^2\frac{\phi}{2}$$

$$\Rightarrow \cos\theta = \pm\sqrt{2}\cos\frac{\phi}{2}$$

Comparing with $\cos\theta = k\cos\frac{\phi}{2}$, we get

$$k = \pm\sqrt{2}$$

7. $\sin^2\theta + 3\sin\theta\cos\theta + 5\cos^2\theta$

$$= \frac{1}{2} \{2\sin^2\theta + 3(2\sin\theta\cos\theta) + 5(2\cos^2\theta)\}$$

$$= \frac{1}{2} \{1 - \cos 2\theta + 3\sin 2\theta + 5(1 + \cos 2\theta)\}$$

$$= 3 + 2\cos 2\theta + \frac{3}{2}\sin 2\theta$$

Now,

$$-\sqrt{4 + \frac{9}{4}} \leq 2\cos 2\theta + \frac{3}{2}\sin 2\theta \leq \sqrt{4 + \frac{9}{4}}$$

$$\Rightarrow -\frac{5}{2} \leq 2\cos 2\theta + \frac{3}{2}\sin 2\theta \leq \frac{5}{2}$$

$$\Rightarrow \frac{1}{2} \leq 3 + 2\cos 2\theta + \frac{3}{2}\sin 2\theta \leq \frac{11}{2}$$

$$\Rightarrow \frac{1}{2} \leq \sin^2\theta + 3\sin\theta\cos\theta + 5\cos^2\theta \leq \frac{11}{2}$$

$$\Rightarrow \frac{2}{11} \leq \frac{1}{\sin^2\theta + 3\sin\theta\cos\theta + 5\cos^2\theta} \leq 2$$

Hence, the maximum value of the given expression is 2.

8. $\sin\alpha + \sin\beta = -\frac{21}{65}$, $\cos\alpha + \cos\beta = -\frac{27}{65}$

$\therefore (\sin\alpha + \sin\beta)^2 + (\cos\alpha + \cos\beta)^2$

$$= \left(\frac{-21}{65}\right)^2 + \left(\frac{-27}{65}\right)^2$$

$$\Rightarrow (\sin^2\alpha + \cos^2\alpha) + (\sin^2\beta + \cos^2\beta)$$

$$+ 2\sin\alpha\sin\beta + 2\cos\alpha\cos\beta = \frac{441}{(65)^2} + \frac{729}{(65)^2}$$

$$\Rightarrow 2 + 2\sin\alpha\sin\beta + 2\cos\alpha\cos\beta$$

$$= \frac{441}{(65)^2} + \frac{729}{(65)^2}$$

$$\Rightarrow 2 + 2[\cos(\alpha - \beta)] = \frac{1170}{(65)^2}$$

$$\Rightarrow 2[1 + \cos(\alpha - \beta)] = \frac{1170}{(65)^2}$$



$$\Rightarrow 2 \times 2 \cos^2\left(\frac{\alpha - \beta}{2}\right) = \frac{1170}{(65)^2}$$

$$\therefore \cos\left(\frac{\alpha - \beta}{2}\right) = -\frac{3\sqrt{130}}{130}$$

$$\dots \left[\because \frac{\pi}{2} < \frac{\alpha - \beta}{2} < \frac{3\pi}{2} \right]$$

$$= \frac{-3}{\sqrt{130}}$$

$$9. \quad \tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$$

$$\Rightarrow \tan(\alpha + \beta) = \frac{\frac{1}{1 + \frac{1}{2^x}} + \frac{1}{1 + 2^{x+1}}}{1 - \left(\frac{1}{1 + \frac{1}{2^x}}\right) \left(\frac{1}{1 + 2^{x+1}}\right)}$$

$$\Rightarrow \tan(\alpha + \beta) = \frac{2^x + 2.2^{x+x} + 2^x + 1}{1 + 2^x + 2.2^x + 2.2^{x+x} - 2^x}$$

$$\Rightarrow \tan(\alpha + \beta) = 1$$

$$\Rightarrow \tan(\alpha + \beta) = \tan \frac{\pi}{4}$$

$$\Rightarrow \alpha + \beta = \frac{\pi}{4}$$

$$\Rightarrow \alpha + \beta = \frac{\pi}{4}$$

$$10. \quad \text{We have, } \frac{3 \sin A}{\sin B} = \frac{2 \cos B}{\cos A}$$

$$\Rightarrow \frac{3 \sin A}{\cos A} = \frac{2 \cos B \sin B}{\cos^2 A}$$

$$\Rightarrow \tan A = \frac{\sin 2B}{3 \cos^2 A}$$

$$\Rightarrow \tan A = \frac{\sin 2B}{3 \cos 2B} \times \frac{\cos 2B}{\cos^2 A}$$

$$\Rightarrow \tan A = \frac{\tan 2B}{3 \cos^2 A} (2 \cos^2 B - 1)$$

$$\Rightarrow \tan A = \frac{\tan 2B}{3 \cos^2 A} (4 - 3 \cos^2 A - 1)$$

$$\dots [\because 2 \cos^2 B = 4 - 3 \cos^2 A \text{ (given)}]$$

$$\Rightarrow \tan A = \tan 2B \frac{\sin^2 A}{\cos^2 A}$$

$$\dots [\because 1 - \cos^2 A = \sin^2 A]$$

$$\Rightarrow \tan A = \tan 2B \tan^2 A$$

$$\Rightarrow \tan A \tan 2B = 1$$

$$\Rightarrow \tan A = \cot 2B$$

$$\Rightarrow \tan A = \tan \left(\frac{\pi}{2} - 2B \right)$$

$$\Rightarrow A = \frac{\pi}{2} - 2B$$

$$\Rightarrow A + 2B = \frac{\pi}{2}$$

$$11. \quad \frac{\sin^4 A}{a} + \frac{\cos^4 A}{b} = \frac{1}{a+b}$$

$$\Rightarrow \frac{(1 - \cos 2A)^2}{4a} + \frac{(1 + \cos 2A)^2}{4b} = \frac{1}{a+b}$$

$$\Rightarrow b(a+b)(1 - 2 \cos 2A + \cos^2 2A) + a(a+b)(1 + 2 \cos 2A + \cos^2 2A) = 4ab$$

$$\Rightarrow \{b(a+b) + a(a+b)\} \cos^2 2A + 2(a+b)(a-b) \cos 2A + a(a+b) + b(a+b) - 4ab = 0$$

$$\Rightarrow (a+b)^2 \cos^2 2A + 2(a+b)(a-b) \cos 2A + (a-b)^2 = 0$$

$$\Rightarrow \{(a+b) \cos 2A + (a-b)\}^2 = 0$$

$$\Rightarrow \cos 2A = \frac{b-a}{b+a} \quad \dots (i)$$

$$\therefore \frac{\sin^8 A}{a^3} + \frac{\cos^8 A}{b^3} = \frac{(1 - \cos 2A)^4}{16a^3} + \frac{(1 + \cos 2A)^4}{16b^3}$$

$$= \frac{1}{16a^3} \left(1 - \frac{b-a}{b+a}\right)^4 + \frac{1}{16b^3} \left(1 + \frac{b-a}{b+a}\right)^4$$

....[From (i)]

$$= \frac{16a^4}{16a^3(b+a)^4} + \frac{16b^4}{16b^3(b+a)^4}$$

$$= \frac{1}{(b+a)^4} (a+b) = \frac{1}{(a+b)^3}$$

$$12. \quad \sin(\pi \cos \theta) = \cos(\pi \sin \theta)$$

$$\Rightarrow \sin(\pi \cos \theta) = \sin\left(\frac{\pi}{2} + \pi \sin \theta\right)$$

$$\dots \left[\because \cos \theta = \sin\left(\frac{\pi}{2} + \theta\right) \right]$$

$$\Rightarrow \pi \cos \theta = \frac{\pi}{2} + \pi \sin \theta$$

$$\Rightarrow \cos \theta - \sin \theta = \frac{1}{2} \quad \dots (i)$$

$$\therefore \cos\left(\theta + \frac{\pi}{4}\right) = \cos \theta \cos \frac{\pi}{4} - \sin \theta \sin \frac{\pi}{4}$$

$$= \frac{1}{\sqrt{2}} (\cos \theta - \sin \theta)$$

$$= \frac{1}{2\sqrt{2}} \quad \dots [\text{From (i)}]$$

04 Factorization Formulae



Hints



Classical Thinking

$$\begin{aligned} 1. \quad \cos 5^\circ - \sin 25^\circ &= \sin(90 - 5)^\circ - \sin 25^\circ \\ &= \sin 85^\circ - \sin 25^\circ \\ &= 2 \cos 55^\circ \sin 30^\circ \\ &= \cos 55^\circ \end{aligned}$$

$$\begin{aligned} 2. \quad \cos 57^\circ + \sin 27^\circ &= \cos 57^\circ + \cos(90^\circ - 27^\circ) \\ &= \cos 57^\circ + \cos 63^\circ \\ &= 2 \cos 60^\circ \cos 3^\circ \\ &= \cos 3^\circ \end{aligned}$$

$$\begin{aligned} 3. \quad \cos 18^\circ - \sin 18^\circ &= \cos 18^\circ - \cos 72^\circ \\ &= 2 \sin 45^\circ \sin 27^\circ \\ &= \sqrt{2} \sin 27^\circ \end{aligned}$$

$$\begin{aligned} 4. \quad \cos\left(\frac{3\pi}{4} + x\right) - \cos\left(\frac{3\pi}{4} - x\right) \\ &= 2 \sin\left(\frac{\frac{3\pi}{4} + x + \frac{3\pi}{4} - x}{2}\right) \times \sin\left(\frac{\frac{3\pi}{4} - x - \frac{3\pi}{4} - x}{2}\right) \\ &= 2 \sin\left(\frac{3\pi}{4}\right) \sin(-x) \\ &= -2 \sin\left(\frac{\pi}{2} + \frac{\pi}{4}\right) \sin x \\ &= -2 \cos\left(\frac{\pi}{4}\right) \sin x \\ &= -\sqrt{2} \sin x \end{aligned}$$

$$\begin{aligned} 5. \quad (\sin 50^\circ - \sin 70^\circ) + \sin 10^\circ \\ &= 2 \cos 60^\circ \sin(-10^\circ) + \sin 10^\circ \\ &= -2 \cdot \frac{1}{2} \sin 10^\circ + \sin 10^\circ \\ &= 0 \end{aligned}$$

$$\begin{aligned} 6. \quad \cos 52^\circ + \cos 68^\circ + \cos 172^\circ \\ &= (\cos 52^\circ + \cos 172^\circ) + \cos 68^\circ \\ &= 2 \cos 112^\circ \cos 60^\circ + \cos 68^\circ \\ &= \cos 112^\circ + \cos 68^\circ \\ &= 2 \cos 90^\circ \cos 22^\circ \\ &= 0 \end{aligned}$$

$$\begin{aligned} 7. \quad \{\sin(\beta + \gamma - \alpha) + \sin(\gamma + \alpha - \beta)\} + \\ \quad \{\sin(\alpha + \beta - \gamma) - \sin(\alpha + \beta + \gamma)\} \\ &= 2 \sin \gamma \cos(\beta - \alpha) + 2 \sin(-\gamma) \cos(\alpha + \beta) \\ &= 2 \sin \gamma [\cos(\beta - \alpha) - \cos(\alpha + \beta)] \\ &= 2 \sin \gamma \cdot 2 \sin \alpha \sin \beta \\ &= 4 \sin \alpha \sin \beta \sin \gamma \end{aligned}$$

$$8. \quad \frac{\sin 3x - \sin x}{\cos 2x} = \frac{2 \cos 2x \sin x}{\cos 2x} = 2 \sin x$$

$$9. \quad \frac{\sin 5x + \sin 3x}{\cos 5x + \cos 3x} = \frac{2 \sin 4x \cos x}{2 \cos 4x \cos x} = \tan 4x$$

$$10. \quad \frac{\cos 7A + \cos 5A}{\sin 7A - \sin 5A} = \frac{2 \cos 6A \cos A}{2 \cos 6A \sin A} = \cot A$$

$$\begin{aligned} 11. \quad \frac{\sin 70^\circ + \cos 40^\circ}{\cos 70^\circ + \sin 40^\circ} &= \frac{\sin 70^\circ + \sin 50^\circ}{\cos 70^\circ + \cos 50^\circ} \\ &= \frac{2 \sin 60^\circ \cos 10^\circ}{2 \cos 60^\circ \cos 10^\circ} \\ &= \tan 60^\circ \\ &= \sqrt{3} \end{aligned}$$

$$\begin{aligned} 12. \quad \frac{\sin 3A - \cos\left(\frac{\pi}{2} - A\right)}{\cos A + \cos(\pi + 3A)} &= \frac{\sin 3A - \sin A}{\cos A - \cos 3A} \\ &= \frac{2 \cos 2A \sin A}{2 \sin 2A \sin A} \\ &= \cot 2A \end{aligned}$$

$$\begin{aligned} 13. \quad \frac{\sin 3\theta + \sin 5\theta + \sin 7\theta + \sin 9\theta}{\cos 3\theta + \cos 5\theta + \cos 7\theta + \cos 9\theta} \\ &= \frac{(\sin 3\theta + \sin 9\theta) + (\sin 5\theta + \sin 7\theta)}{(\cos 3\theta + \cos 9\theta) + (\cos 5\theta + \cos 7\theta)} \\ &= \frac{2 \sin 6\theta \cos 3\theta + 2 \sin 6\theta \cos \theta}{2 \cos 6\theta \cos 3\theta + 2 \cos 6\theta \cos \theta} \\ &= \frac{2 \sin 6\theta (\cos 3\theta + \cos \theta)}{2 \cos 6\theta (\cos 3\theta + \cos \theta)} \\ &= \tan 6\theta \end{aligned}$$



$$14. \frac{\sin(x+y)}{\sin(x-y)} = \frac{a+b}{a-b}$$

By componendo and dividendo, we get

$$\frac{\sin(x+y) + \sin(x-y)}{\sin(x+y) - \sin(x-y)} = \frac{(a+b) + (a-b)}{(a+b) - (a-b)}$$

$$\Rightarrow \frac{2 \sin x \cos y}{2 \cos x \sin y} = \frac{2a}{2b}$$

$$\Rightarrow \frac{\tan x}{\tan y} = \frac{a}{b}$$

$$15. 2 \sin 3x \cos 2x = \sin(3x+2x) + \sin(3x-2x) \\ = \sin 5x + \sin x$$

$$16. 2 \sin \frac{5\pi}{12} \cos \frac{\pi}{12} = \sin\left(\frac{5\pi}{12} + \frac{\pi}{12}\right) + \sin\left(\frac{5\pi}{12} - \frac{\pi}{12}\right) \\ = \sin \frac{\pi}{2} + \sin \frac{\pi}{3} \\ = 1 + \frac{\sqrt{3}}{2} = \frac{2+\sqrt{3}}{2}$$

$$17. \cos 75^\circ \cos 15^\circ \\ = \frac{1}{2} [2 \cos 75^\circ \cos 15^\circ] \\ = \frac{1}{2} [\cos(75^\circ + 15^\circ) + \cos(75^\circ - 15^\circ)] \\ = \frac{1}{2} [\cos 90^\circ + \cos 60^\circ] \\ = \frac{1}{4}$$

$$18. \sin(45^\circ + A) \sin(45^\circ - A) \\ = \frac{1}{2} [2 \sin(45^\circ + A) \sin(45^\circ - A)] \\ = \frac{1}{2} [\cos(45^\circ + A - 45^\circ + A) \\ - \cos(45^\circ + A + 45^\circ - A)] \\ = \frac{1}{2} [\cos 2A - \cos 90^\circ] \\ = \frac{1}{2} \cos 2A$$

$$19. 4 \sin\left(\frac{\pi}{3} + \theta\right) \sin\left(\frac{\pi}{3} - \theta\right) \\ = 2 \left[2 \sin\left(\frac{\pi}{3} + \theta\right) \sin\left(\frac{\pi}{3} - \theta\right) \right] \\ = 2 \left[\cos\left(\frac{\pi}{3} + \theta - \frac{\pi}{3} + \theta\right) - \cos\left(\frac{\pi}{3} + \theta + \frac{\pi}{3} - \theta\right) \right]$$

$$= 2 \left[\cos 2\theta - \cos\left(\frac{2\pi}{3}\right) \right]$$

$$= 2 \cos 2\theta + 1$$

$$20. \sin 18^\circ \sin 70^\circ + \sin 16^\circ \sin 36^\circ \\ = \frac{1}{2} [2 \sin 18^\circ \sin 70^\circ + 2 \sin 16^\circ \sin 36^\circ] \\ = \frac{1}{2} [\cos 52^\circ - \cos 88^\circ + \cos 20^\circ - \cos 52^\circ] \\ = \frac{1}{2} [\cos 20^\circ - \cos 88^\circ] \\ = \frac{1}{2} [2 \sin 54^\circ \sin 34^\circ] \\ = \sin 54^\circ \sin 34^\circ$$

$$21. \sin 10^\circ \sin 30^\circ \sin 50^\circ \sin 70^\circ \\ = \frac{1}{2} \cdot \frac{1}{2} (2 \sin 10^\circ \sin 50^\circ) \sin 70^\circ \\ = \frac{1}{4} (\cos 40^\circ - \cos 60^\circ) \sin 70^\circ \\ = \frac{1}{8} (2 \sin 70^\circ \cos 40^\circ - \sin 70^\circ) \\ = \frac{1}{8} (\sin 110^\circ + \sin 30^\circ - \sin 70^\circ) \\ = \frac{1}{8} (\sin 70^\circ + \frac{1}{2} - \sin 70^\circ) \\ \dots [\because \sin(180^\circ - A) = \sin A] \\ = \frac{1}{16}$$

$$22. \cos 20^\circ \cos 40^\circ \cos 60^\circ \cos 80^\circ \\ = \frac{1}{2} \cdot \frac{1}{2} (2 \cos 40^\circ \cos 20^\circ) \cos 80^\circ \\ = \frac{1}{4} (\cos 60^\circ + \cos 20^\circ) \cos 80^\circ \\ = \frac{1}{8} (\cos 80^\circ + 2 \cos 20^\circ \cos 80^\circ) \\ = \frac{1}{8} (\cos 80^\circ + \cos 100^\circ + \cos 60^\circ) \\ = \frac{1}{16} \dots [\because \cos(180^\circ - A) = -\cos A]$$

$$24. \operatorname{cosec} A (\sin B \cos C + \cos B \sin C) \\ = \operatorname{cosec} A \sin(B+C) \\ = \operatorname{cosec} A \sin(180^\circ - A) \\ = \operatorname{cosec} A \sin A \\ = 1$$



$$\begin{aligned}
25. \quad & \sin^2 \alpha + \sin^2 \beta - \sin^2 \gamma \\
& = \sin^2 \alpha + \sin(\beta - \gamma) \sin(\beta + \gamma) \\
& = \sin^2 \alpha + \sin(\pi - \alpha) \sin(\beta - \gamma) \\
& = \sin \alpha [\sin \alpha + \sin(\beta - \gamma)] \\
& = \sin \alpha [\sin(\beta + \gamma) + \sin(\beta - \gamma)] \\
& = 2 \sin \alpha \sin \beta \cos \gamma \\
26. \quad & \cos^2 A + \cos^2 B - \cos^2 C \\
& = \frac{1}{2}(1 + \cos 2A) + \frac{1}{2}(1 + \cos 2B) \\
& \quad - \frac{1}{2}(1 + \cos 2C) \\
& = \frac{1}{2} + \frac{1}{2}(\cos 2A + \cos 2B - \cos 2C) \\
& = \frac{1}{2} + \frac{1}{2}(1 - 4 \sin A \sin B \cos C) \\
& = 1 - 2 \sin A \sin B \cos C \\
27. \quad & \tan(A + B) = \tan(180^\circ - C) \\
& \Rightarrow \frac{\tan A + \tan B}{1 - \tan A \tan B} = -\tan C \\
& \Rightarrow \tan A + \tan B = -\tan C(1 - \tan A \tan B) \\
& \Rightarrow \tan A + \tan B = -\tan C + \tan A \tan B \tan C \\
& \Rightarrow \tan A + \tan B + \tan C = \tan A \tan B \tan C \\
28. \quad & \cos A = \cos B \cos C \\
& \Rightarrow \cos(\pi - B - C) = \cos B \cos C \\
& \Rightarrow -\cos(B + C) = \cos B \cos C \\
& \Rightarrow -\cos B \cos C + \sin B \sin C = \cos B \cos C \\
& \Rightarrow 2 \cos B \cos C = \sin B \sin C \\
& \Rightarrow \cot B \cot C = \frac{1}{2}
\end{aligned}$$

**Critical Thinking**

$$\begin{aligned}
1. \quad & \cos 12^\circ + \cos 84^\circ + \cos 156^\circ + \cos 132^\circ \\
& = (\cos 12^\circ + \cos 132^\circ) + (\cos 84^\circ + \cos 156^\circ) \\
& = 2 \cos 72^\circ \cos 60^\circ + 2 \cos 120^\circ \cos 36^\circ \\
& = 2 \cos 72^\circ \times \frac{1}{2} - 2 \times \frac{1}{2} \times \cos 36^\circ \\
& = \cos 72^\circ - \cos 36^\circ \\
& = \frac{\sqrt{5}-1}{4} - \frac{\sqrt{5}+1}{4} \\
& = \frac{-1}{2} \\
2. \quad & \cot 70^\circ + 4 \cos 70^\circ \\
& = \frac{\cos 70^\circ + 4 \sin 70^\circ \cos 70^\circ}{\sin 70^\circ}
\end{aligned}$$

$$\begin{aligned}
& = \frac{\cos 70^\circ + 2 \sin 140^\circ}{\sin 70^\circ} \\
& = \frac{\cos(90^\circ - 20^\circ) + 2 \sin(180^\circ - 40^\circ)}{\sin 70^\circ} \\
& = \frac{\sin 20^\circ + \sin 40^\circ + \sin 40^\circ}{\sin 70^\circ} \\
& = \frac{2 \sin 30^\circ \cos 10^\circ + \sin 40^\circ}{\sin 70^\circ} \\
& = \frac{\cos 10^\circ + \sin 40^\circ}{\sin 70^\circ} \\
& = \frac{\sin 80^\circ + \sin 40^\circ}{\sin 70^\circ} \\
& = \frac{2 \sin 60^\circ \cos 20^\circ}{\sin 70^\circ} \\
& = \sqrt{3}
\end{aligned}$$

$$\begin{aligned}
3. \quad & \cos 10x + \cos 8x + 3 \cos 4x + 3 \cos 2x \\
& = (\cos 10x + \cos 8x) + 3(\cos 4x + \cos 2x) \\
& = 2 \cos 9x \cos x + 6 \cos 3x \cos x \\
& = 2 \cos x (\cos 9x + 3 \cos 3x) \\
& = 2 \cos x [\cos(3(3x)) + 3 \cos 3x] \\
& = 2 \cos x (4 \cos^3 3x - 3 \cos 3x + 3 \cos 3x) \\
& = 8 \cos^3 3x \cos x \\
4. \quad & 1 + \cos 2x + \cos 4x + \cos 6x \\
& = (1 + \cos 6x) + (\cos 2x + \cos 4x) \\
& = 2 \cos^2 3x + 2 \cos 3x \cos x \\
& = 2 \cos 3x (\cos 3x + \cos x) \\
& = 4 \cos x \cos 2x \cos 3x \\
5. \quad & \frac{\cos 6x + 6 \cos 4x + 15 \cos 2x + 10}{\cos 5x + 5 \cos 3x + 10 \cos x} \\
& = \frac{(\cos 6x + \cos 4x) + 5(\cos 4x + \cos 2x) + 10(\cos 2x + 1)}{\cos 5x + 5 \cos 3x + 10 \cos x} \\
& = \frac{2 \cos x \cos 5x + 10 \cos x \cos 3x + 10(2 \cos^2 x - 1 + 1)}{\cos 5x + 5 \cos 3x + 10 \cos x} \\
& = \frac{2 \cos x (\cos 5x + 5 \cos 3x + 10 \cos x)}{\cos 5x + 5 \cos 3x + 10 \cos x} \\
& = 2 \cos x \\
6. \quad & \sin \theta + \sin 3\theta + \sin 2\theta = \sin \alpha \\
& \Rightarrow 2 \sin 2\theta \cos \theta + \sin 2\theta = \sin \alpha \\
& \Rightarrow \sin 2\theta (2 \cos \theta + 1) = \sin \alpha \quad \dots (i) \\
& \text{Also, } \cos \theta + \cos 3\theta + \cos 2\theta = \cos \alpha \\
& \Rightarrow 2 \cos 2\theta \cos \theta + \cos 2\theta = \cos \alpha \\
& \Rightarrow \cos 2\theta (2 \cos \theta + 1) = \cos \alpha \quad \dots (ii)
\end{aligned}$$



From (i) and (ii), we get

$$\tan 2\theta = \tan \alpha$$

$$\Rightarrow 2\theta = \alpha$$

$$\Rightarrow \theta = \frac{\alpha}{2}$$

$$\begin{aligned} 7. \quad & \cos x + \cos y + \cos \alpha = 0 \\ & \Rightarrow \cos x + \cos y = -\cos \alpha \\ & \Rightarrow 2 \cos \left(\frac{x+y}{2} \right) \cos \left(\frac{x-y}{2} \right) = -\cos \alpha \quad \dots(i) \end{aligned}$$

Also, $\sin x + \sin y + \sin \alpha = 0$

$$\Rightarrow \sin x + \sin y = -\sin \alpha$$

$$\Rightarrow 2 \sin \left(\frac{x+y}{2} \right) \cos \left(\frac{x-y}{2} \right) = -\sin \alpha \quad \dots(ii)$$

Dividing (i) by (ii), we get

$$\frac{2 \cos \left(\frac{x+y}{2} \right) \cos \left(\frac{x-y}{2} \right)}{2 \sin \left(\frac{x+y}{2} \right) \cos \left(\frac{x-y}{2} \right)} = \frac{\cos \alpha}{\sin \alpha}$$

$$\Rightarrow \cot \left(\frac{x+y}{2} \right) = \cot \alpha$$

$$\begin{aligned} 8. \quad & (\cos A + \cos B)^2 + (\sin A - \sin B)^2 \\ & = \left[2 \cos \left(\frac{A+B}{2} \right) \cos \left(\frac{A-B}{2} \right) \right]^2 \\ & \quad + \left[2 \cos \left(\frac{A+B}{2} \right) \sin \left(\frac{A-B}{2} \right) \right]^2 \\ & = 4 \cos^2 \left(\frac{A+B}{2} \right) \left[\cos^2 \left(\frac{A-B}{2} \right) + \sin^2 \left(\frac{A-B}{2} \right) \right] \\ & = 4 \cos^2 \left(\frac{A+B}{2} \right) \end{aligned}$$

$$\begin{aligned} 9. \quad & \cos^2 \alpha + \cos^2 (\alpha + 120^\circ) + \cos^2 (\alpha - 120^\circ) \\ & = \cos^2 \alpha + \{ \cos (\alpha + 120^\circ) + \cos (\alpha - 120^\circ) \}^2 \\ & \quad - 2 \cos (\alpha + 120^\circ) \cos (\alpha - 120^\circ) \\ & = \cos^2 \alpha + \{ 2 \cos \alpha \cos 120^\circ \}^2 \\ & \quad - 2 \{ \cos^2 \alpha - \sin^2 120^\circ \} \\ & = \cos^2 \alpha + \cos^2 \alpha - 2 \cos^2 \alpha + 2 \sin^2 120^\circ \\ & = 2 \sin^2 120^\circ \\ & = 2 \times \frac{3}{4} = \frac{3}{2} \end{aligned}$$

$$10. \quad \frac{\cos(A+B)}{\cos(A-B)} = \frac{\sin(C+D)}{\sin(C-D)}$$

By componendo and dividendo, we get

$$\begin{aligned} & \frac{\cos(A+B) + \cos(A-B)}{\cos(A+B) - \cos(A-B)} \\ & = \frac{\sin(C+D) + \sin(C-D)}{\sin(C+D) - \sin(C-D)} \end{aligned}$$

$$\Rightarrow \frac{2 \cos A \cos B}{-2 \sin A \sin B} = \frac{2 \sin C \cos D}{2 \cos C \sin D}$$

$$\Rightarrow -\cot A \cot B = \tan C \cot D$$

$$\Rightarrow \tan A \tan B \tan C + \tan D = 0$$

$$\begin{aligned} 11. \quad & \frac{\sin^2 A - \sin^2 B}{\sin A \cos A - \sin B \cos B} \\ & = \frac{2 \sin(A+B) \sin(A-B)}{\sin 2A - \sin 2B} \\ & \dots [\because \sin^2 A - \sin^2 B = \sin(A+B) \sin(A-B)] \\ & = \frac{2 \sin(A+B) \sin(A-B)}{2 \cos(A+B) \sin(A-B)} \\ & = \tan(A+B) \end{aligned}$$

$$\begin{aligned} 12. \quad & \text{Since, } \cos \alpha + \cos(\alpha + \beta) + \cos(\alpha + 2\beta) \\ & \quad + \dots + \cos\{\alpha + (n-1)\beta\} \\ & = \frac{\cos\left\{\alpha + (n-1)\frac{\beta}{2}\right\} \sin\left(\frac{n\beta}{2}\right)}{\sin\frac{\beta}{2}} \end{aligned}$$

$$\text{Here, } \alpha = \frac{\pi}{11} \text{ and } \beta = \frac{2\pi}{11}$$

$$\begin{aligned} \therefore & \cos \frac{\pi}{11} + \cos \frac{3\pi}{11} + \cos \frac{5\pi}{11} + \cos \frac{7\pi}{11} + \cos \frac{9\pi}{11} \\ & = \frac{\cos\left(\frac{\pi}{11} + \frac{4\pi}{11}\right) \sin\left(\frac{5\pi}{11}\right)}{\sin\left(\frac{\pi}{11}\right)} \\ & = \frac{\cos \frac{5\pi}{11} \sin \frac{5\pi}{11}}{\sin \frac{\pi}{11}} \\ & = \frac{1}{2} \frac{\sin \frac{10\pi}{11}}{\sin \frac{\pi}{11}} \quad \dots [\because 2 \sin \theta \cos \theta = \sin 2\theta] \\ & = \frac{1}{2} \end{aligned}$$

$$\begin{aligned} 13. \quad & 2 \sin \frac{5A}{2} \sin \frac{A}{2} = \cos 2A - \cos 3A \\ & = 2 \cos^2 A - 1 - 4 \cos^3 A + 3 \cos A \end{aligned}$$



$$\begin{aligned}
 &= 2 \left(\frac{9}{16} \right) - 1 - 4 \left(\frac{27}{64} \right) + 3 \left(\frac{3}{4} \right) \\
 &\quad \dots \left[\because \cos A = \frac{3}{4} \right] \\
 &= \frac{9}{8} - 1 - \frac{27}{16} + \frac{9}{4} \\
 &= \frac{11}{16}
 \end{aligned}$$

$$\begin{aligned}
 14. \quad & \sin \frac{\pi}{16} \sin \frac{3\pi}{16} \sin \frac{5\pi}{16} \sin \frac{7\pi}{16} \\
 &= \frac{1}{4} \left[2 \sin \frac{\pi}{16} \sin \frac{3\pi}{16} \cdot 2 \sin \frac{5\pi}{16} \sin \frac{7\pi}{16} \right] \\
 &= \frac{1}{4} \left[\left(\cos \frac{\pi}{8} - \cos \frac{\pi}{4} \right) \left(\cos \frac{\pi}{8} - \cos \frac{3\pi}{4} \right) \right] \\
 &= \frac{1}{4} \left[\left(\cos \frac{\pi}{8} - \frac{1}{\sqrt{2}} \right) \left(\cos \frac{\pi}{8} + \frac{1}{\sqrt{2}} \right) \right] \\
 &= \frac{1}{4} \left[\left(\cos^2 \frac{\pi}{8} - \frac{1}{2} \right) \right] \\
 &= \frac{1}{8} \left[2 \cos^2 \frac{\pi}{8} - 1 \right] = \frac{1}{8} \left[\cos \frac{\pi}{4} \right] \\
 &= \frac{1}{8} \times \frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{16}
 \end{aligned}$$

$$\begin{aligned}
 15. \quad & \sin 20^\circ \sin 40^\circ \sin 60^\circ \sin 80^\circ \\
 &= \frac{\sqrt{3}}{2} \sin 20^\circ \sin (60^\circ - 20^\circ) \sin (60^\circ + 20^\circ) \\
 &= \frac{\sqrt{3}}{2} \cdot \frac{1}{4} \sin 60^\circ \\
 &\quad \dots \left[\because \sin \theta \sin(60^\circ - \theta) \sin(60^\circ + \theta) = \frac{1}{4} \sin 3\theta \right] \\
 &= \frac{\sqrt{3}}{8} \cdot \frac{\sqrt{3}}{2} \\
 &= \frac{3}{16}
 \end{aligned}$$

$$\begin{aligned}
 16. \quad & \tan 20^\circ \tan 40^\circ \tan 60^\circ \tan 80^\circ \\
 &= \tan 20^\circ \tan (60^\circ - 20^\circ) \cdot \sqrt{3} \cdot \tan(60^\circ + 20^\circ) \\
 &= \sqrt{3} \tan 20^\circ \cdot \tan (60^\circ - 20^\circ) \cdot \tan (60^\circ + 20^\circ) \\
 &= \sqrt{3} \tan 3(20^\circ) \\
 &\quad \dots \left[\because \tan \theta \tan(60^\circ - \theta) \tan(60^\circ + \theta) = \tan 3\theta \right] \\
 &= \sqrt{3} \cdot \sqrt{3} = 3
 \end{aligned}$$

$$\begin{aligned}
 17. \quad & \frac{\tan 70^\circ - \tan 20^\circ}{\tan 50^\circ} = \frac{\frac{\sin 70^\circ}{\cos 70^\circ} - \frac{\sin 20^\circ}{\cos 20^\circ}}{\frac{\sin 50^\circ}{\cos 50^\circ}} \\
 &= \frac{\frac{\sin 70^\circ \cos 20^\circ - \cos 70^\circ \sin 20^\circ}{\cos 70^\circ \cos 20^\circ}}{\frac{\sin 50^\circ}{\cos 50^\circ}} \\
 &= \frac{\sin(70^\circ - 20^\circ) \cos 50^\circ}{\cos 70^\circ \cos 20^\circ \sin 50^\circ} \\
 &= \frac{2 \sin 50^\circ \cos 50^\circ}{2 \cos 70^\circ \cos 20^\circ \sin 50^\circ} \\
 &= \frac{2 \cos 50^\circ}{\cos 90^\circ + \cos 50^\circ} \\
 &= \frac{2 \cos 50^\circ}{0 + \cos 50^\circ} = 2
 \end{aligned}$$

$$\begin{aligned}
 18. \quad & \tan 20^\circ + 2 \tan 50^\circ - \tan 70^\circ \\
 &= \frac{\sin 20^\circ}{\cos 20^\circ} - \frac{\sin 70^\circ}{\cos 70^\circ} + 2 \tan 50^\circ \\
 &= \frac{\sin 20^\circ \cos 70^\circ - \cos 20^\circ \sin 70^\circ}{\cos 20^\circ \cos 70^\circ} + 2 \tan 50^\circ \\
 &= \frac{\sin(20^\circ - 70^\circ)}{\sin(20^\circ + 70^\circ)} + 2 \tan 50^\circ \\
 &= \frac{1}{2} [\cos(70^\circ + 20^\circ) + \cos(70^\circ - 20^\circ)] \\
 &= \frac{2 \sin(-50^\circ)}{\cos 90^\circ + \cos 50^\circ} + 2 \tan 50^\circ \\
 &= \frac{-2 \sin 50^\circ}{0 + \cos 50^\circ} + 2 \tan 50^\circ \\
 &= -2 \tan 50^\circ + 2 \tan 50^\circ \\
 &= 0
 \end{aligned}$$

$$\begin{aligned}
 19. \quad & \operatorname{cosec} 48^\circ + \operatorname{cosec} 84^\circ + \operatorname{cosec} 192^\circ + \operatorname{cosec} 384^\circ \\
 &= \frac{1}{\sin 48^\circ} + \frac{1}{\sin 84^\circ} + \frac{1}{-\sin 12^\circ} + \frac{1}{\sin 24^\circ} \\
 &= \left(\frac{1}{\sin 48^\circ} - \frac{1}{\sin 12^\circ} \right) + \left(\frac{1}{\sin 84^\circ} + \frac{1}{\sin 24^\circ} \right) \\
 &= -\frac{(\sin 48^\circ - \sin 12^\circ)}{\sin 48^\circ \sin 12^\circ} + \frac{(\sin 84^\circ + \sin 24^\circ)}{\sin 84^\circ \sin 24^\circ} \\
 &= -\frac{2 \cos 30^\circ \sin 18^\circ}{\frac{1}{2} (\cos 36^\circ - \cos 60^\circ)} \\
 &\quad + \frac{2 \sin 54^\circ \cos 30^\circ}{\frac{1}{2} (\cos 60^\circ - \cos 108^\circ)} \\
 &= \frac{4 \cos 30^\circ \sin 18^\circ}{\cos 60^\circ - \cos 36^\circ} + \frac{4 \sin 54^\circ \cos 30^\circ}{\cos 60^\circ + \sin 18^\circ}
 \end{aligned}$$



$$\begin{aligned}
 &= 4 \cos 30^\circ \left[\frac{\sin 18^\circ}{\cos 60^\circ - \cos 36^\circ} + \frac{\sin 54^\circ}{\cos 60^\circ + \sin 18^\circ} \right] \\
 &= 4 \cos 30^\circ \left[\frac{\sin 18^\circ}{\cos 60^\circ - \cos 36^\circ} + \frac{\cos 36^\circ}{\cos 60^\circ + \sin 18^\circ} \right] \\
 &= 4 \cos 30^\circ \left[\frac{\frac{\sqrt{5}-1}{4}}{\frac{1}{2} - \frac{\sqrt{5}+1}{4}} + \frac{\frac{\sqrt{5}+1}{4}}{\frac{1}{2} + \frac{\sqrt{5}-1}{4}} \right] \\
 &= 4 \cos 30^\circ (-1 + 1) = 0
 \end{aligned}$$

$$\begin{aligned}
 20. \quad &\cos \frac{2\pi}{7} + \cos \frac{4\pi}{7} + \cos \frac{6\pi}{7} + \cos \frac{7\pi}{7} \\
 &= \frac{1}{2\sin \frac{\pi}{7}} \left\{ 2\cos \frac{2\pi}{7} \sin \frac{\pi}{7} + 2\cos \frac{4\pi}{7} \sin \frac{\pi}{7} \right. \\
 &\quad \left. + 2\cos \frac{6\pi}{7} \sin \frac{\pi}{7} \right\} + \cos \pi \\
 &= \frac{1}{2\sin \frac{\pi}{7}} \left[\sin \frac{3\pi}{7} - \sin \frac{\pi}{7} + \sin \frac{5\pi}{7} - \sin \frac{3\pi}{7} \right. \\
 &\quad \left. + \sin \pi - \sin \frac{5\pi}{7} \right] + \cos \pi \\
 &= -\frac{1}{2} - 1 \\
 &= -\frac{3}{2}
 \end{aligned}$$

$$\begin{aligned}
 21. \quad &\cos^2(A - B) + \cos^2 B - 2\cos(A - B)\cos A \cos B \\
 &= \cos^2(A - B) + \cos^2 B \\
 &\quad - \cos(A - B) \{ \cos(A - B) + \cos(A + B) \} \\
 &= \cos^2 B - \cos(A - B)\cos(A + B) \\
 &= \cos^2 B - (\cos^2 A - \sin^2 B) \\
 &= 1 - \cos^2 A \\
 &\text{Hence, it depends on A.}
 \end{aligned}$$

$$\begin{aligned}
 22. \quad &\frac{A}{B} = \tan 6^\circ \tan 42^\circ \tan 66^\circ \tan 78^\circ \\
 &= \frac{(2 \sin 66^\circ \sin 6^\circ)(2 \sin 78^\circ \sin 42^\circ)}{(2 \cos 66^\circ \cos 6^\circ)(2 \cos 78^\circ \cos 42^\circ)} \\
 &= \frac{(\cos 60^\circ - \cos 72^\circ)(\cos 36^\circ - \cos 120^\circ)}{(\cos 60^\circ + \cos 72^\circ)(\cos 36^\circ + \cos 120^\circ)} \\
 &= \frac{(\cos 60^\circ - \sin 18^\circ)(\cos 36^\circ + \sin 30^\circ)}{(\cos 60^\circ + \sin 18^\circ)(\cos 36^\circ - \sin 30^\circ)} \\
 &\quad \dots [\because \cos(90^\circ + \theta) = -\sin \theta]
 \end{aligned}$$

$$\begin{aligned}
 &= \left(\frac{1 - \sqrt{5}-1}{4} \right) \left(\frac{\sqrt{5}+1}{4} + \frac{1}{2} \right) \\
 &= \left(\frac{1 + \sqrt{5}-1}{4} \right) \left(\frac{\sqrt{5}+1}{4} - \frac{1}{2} \right) \\
 &= \frac{9-5}{5-1} = 1 \\
 &\Rightarrow A = B
 \end{aligned}$$

$$\begin{aligned}
 23. \quad &\sin 2A + \sin 2B - \sin 2C \\
 &= 2 \sin A \cos A + 2 \cos(B + C) \sin(B - C) \\
 &= 2 \sin A \cos A - 2 \cos A \sin(B - C) \\
 &= 2 \cos A [\sin A - \sin(B - C)] \\
 &= 2 \cos A [\sin(B + C) - \sin(B - C)] \\
 &\quad \dots [\because \sin(B + C) = \sin A] \\
 &= 2 \cos A (2 \cos B \sin C) \\
 &= 4 \cos A \cos B \sin C \\
 &\text{Trick : Check by assuming } A = B = 45^\circ \text{ and } C = 90^\circ
 \end{aligned}$$

$$\begin{aligned}
 24. \quad &\cos 2A + \cos 2B + \cos 2C \\
 &= 2 \cos(A + B) \cos(A - B) + (2 \cos^2 C - 1) \\
 &= -1 - 2 \cos C \cos(A - B) + 2 \cos^2 C \\
 &\quad \dots [\because \cos(A + B) = -\cos C] \\
 &= -1 - 2 \cos C [\cos(A - B) + \cos(A + B)] \\
 &= -1 - 4 \cos A \cos B \cos C
 \end{aligned}$$

$$\begin{aligned}
 25. \quad &\cos 2x + \cos 2y - \cos 2z \\
 &= 2 \cos(x + y) \cos(x - y) - 2 \cos^2 z + 1 \\
 &= 2 \cos(\pi - z) \cos(x - y) - 2 \cos^2 z + 1 \\
 &= 1 - 2 \cos z \{ \cos(x - y) - \cos(x + y) \} \\
 &= 1 - 2 \cos z \cdot 2 \sin x \sin y \\
 &= 1 - 4 \sin x \sin y \cos z
 \end{aligned}$$

$$\begin{aligned}
 26. \quad &\cos 2A + \cos 2B + \cos 2C \\
 &= 2 \cos(A + B) \cos(A - B) + \cos 2C \\
 &= 2 \cos \left(\frac{3\pi}{2} - C \right) \cos(A - B) + \cos 2C \\
 &\quad \dots \left[\because A + B = \frac{3\pi}{2} - C \right] \\
 &= -2 \sin C \cos(A - B) + 1 - 2 \sin^2 C \\
 &= 1 - 2 \sin C \{ \cos(A - B) + \sin C \} \\
 &= 1 - 2 \sin C \{ \cos(A - B) - \cos(A + B) \} \\
 &= 1 - 4 \sin A \sin B \sin C \\
 &\text{Trick : Check by assuming } A = B = C = \frac{\pi}{2}
 \end{aligned}$$



$$\begin{aligned}
27. \quad & \frac{\cos A}{\sin B \sin C} + \frac{\cos B}{\sin C \sin A} + \frac{\cos C}{\sin A \sin B} \\
&= \frac{2 \sin A \cos A}{2 \sin A \sin B \sin C} + \frac{2 \sin B \cos B}{2 \sin A \sin B \sin C} \\
&\quad + \frac{2 \sin C \cos C}{2 \sin A \sin B \sin C} \\
&= \frac{\sin 2A + \sin 2B + \sin 2C}{2 \sin A \sin B \sin C} \\
&= \frac{4 \sin A \sin B \sin C}{2 \sin A \sin B \sin C} \\
&= 2
\end{aligned}$$

$$\begin{aligned}
28. \quad & \sin^2 A + \sin^2 B + \sin^2 C \\
&= 1 - \cos^2 A + 1 - \cos^2 B + \sin^2 C \\
&= 2 - \cos^2 A - (\cos^2 B - \sin^2 C) \\
&= 2 - \cos^2 A - \cos(B+C) \cos(B-C) \\
&= 2 - \cos A [\cos A - \cos(B-C)] \\
&= 2 - \cos A [-\cos(B+C) - \cos(B-C)] \\
&= 2 + \cos A \cdot 2 \cos B \cos C \\
\therefore \quad & \sin^2 A + \sin^2 B + \sin^2 C - 2 \cos A \cos B \cos C = 2
\end{aligned}$$

$$\begin{aligned}
29. \quad & \cos^2 A + \cos^2 \left(A + \frac{\pi}{3} \right) + \cos^2 \left(A - \frac{\pi}{3} \right) \\
&= \frac{1}{2} (1 + \cos 2A) + \frac{1}{2} \left\{ 1 + \cos \left(2A + \frac{2\pi}{3} \right) \right\} \\
&\quad + \frac{1}{2} \left\{ 1 + \cos \left(2A - \frac{2\pi}{3} \right) \right\} \\
&= \frac{3}{2} + \frac{1}{2} \cos 2A \\
&\quad + \frac{1}{2} \left\{ \cos \left(2A + \frac{2\pi}{3} \right) + \cos \left(2A - \frac{2\pi}{3} \right) \right\} \\
&= \frac{3}{2} + \frac{1}{2} \cos 2A + \cos 2A \cos \frac{2\pi}{3} \\
&\quad \dots [\because \cos(A+B) + \cos(A-B) = 2 \cos A \cos B] \\
&= \frac{3}{2} + \frac{1}{2} \cos 2A - \frac{1}{2} \cos 2A = \frac{3}{2}
\end{aligned}$$

$$\begin{aligned}
30. \quad & \text{We have,} \\
& \cos^2 \frac{A}{2} + \cos^2 \frac{B}{2} - \cos^2 \frac{C}{2} \\
&= \frac{1}{2} (1 + \cos A) + \frac{1}{2} (1 + \cos B) - \frac{1}{2} (1 + \cos C) \\
&= \frac{1}{2} + \frac{1}{2} (\cos A + \cos B - \cos C)
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2} + \frac{1}{2} \left[4 \cos \frac{A}{2} \cos \frac{B}{2} \sin \frac{C}{2} - 1 \right] \\
&= 2 \cos \frac{A}{2} \cos \frac{B}{2} \sin \frac{C}{2}
\end{aligned}$$

$$\begin{aligned}
31. \quad & \sin A + \sin B + \sin C \\
&= 2 \sin \frac{A+B}{2} \cos \frac{A-B}{2} + 2 \sin \frac{C}{2} \cos \frac{C}{2} \\
&= 2 \sin \left(\frac{\pi}{2} - \frac{C}{2} \right) \cos \frac{A-B}{2} \\
&\quad + 2 \cos \frac{C}{2} \sin \left(\frac{\pi}{2} - \left(\frac{A+B}{2} \right) \right) \\
&= 2 \cos \frac{C}{2} \cos \frac{A-B}{2} + 2 \cos \frac{C}{2} \cos \frac{A+B}{2} \\
&= 2 \cos \frac{C}{2} \left[\cos \frac{A-B}{2} + \cos \frac{A+B}{2} \right] \\
&= 2 \cos \frac{C}{2} \left(2 \cos \frac{A}{2} \cos \frac{B}{2} \right) \\
&= 4 \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}
\end{aligned}$$

$$\begin{aligned}
32. \quad & \frac{\sin 2A + \sin 2B + \sin 2C}{\cos A + \cos B + \cos C - 1} \\
&= \frac{4 \sin A \sin B \sin C}{1 + 4 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} - 1} \\
&= \frac{\left(2 \sin \frac{A}{2} \cos \frac{A}{2} \right) \left(2 \sin \frac{B}{2} \cos \frac{B}{2} \right) \left(2 \sin \frac{C}{2} \cos \frac{C}{2} \right)}{\sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}} \\
&\quad \dots \left[\because \sin \theta = 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2} \right] \\
&= 8 \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}
\end{aligned}$$

$$\begin{aligned}
33. \quad & \text{We have, } \alpha + \beta + \gamma = 2\pi \\
&\Rightarrow \frac{\alpha}{2} + \frac{\beta}{2} + \frac{\gamma}{2} = \pi \\
&\Rightarrow \tan \left(\frac{\alpha}{2} + \frac{\beta}{2} + \frac{\gamma}{2} \right) = \tan \pi = 0 \\
&\Rightarrow \tan \frac{\alpha}{2} + \tan \frac{\beta}{2} + \tan \frac{\gamma}{2} - \tan \frac{\alpha}{2} \tan \frac{\beta}{2} \tan \frac{\gamma}{2} = 0 \\
&\Rightarrow \tan \frac{\alpha}{2} + \tan \frac{\beta}{2} + \tan \frac{\gamma}{2} = \tan \frac{\alpha}{2} \tan \frac{\beta}{2} \tan \frac{\gamma}{2}
\end{aligned}$$



$$\begin{aligned}
 34. \quad \sum \frac{\cot A + \cot B}{\tan A + \tan B} &= \sum \frac{\frac{\cos A}{\sin A} + \frac{\cos B}{\sin B}}{\frac{\sin A}{\cos A} + \frac{\sin B}{\cos B}} \\
 &= \sum \left(\frac{\sin B \cos A + \sin A \cos B}{\sin A \sin B} \right) \left(\frac{\cos A \cos B}{\sin A \cos B + \cos A \sin B} \right) \\
 &= \sum \cot A \cot B \\
 &= \cot A \cot B + \cot B \cot C + \cot C \cot A = 1 \\
 &\left[\begin{array}{l} \because A + B + C = \pi, \\ \therefore \cot A \cot B + \cot B \cot C + \cot C \cot A = 1 \end{array} \right]
 \end{aligned}$$



Competitive Thinking

$$\begin{aligned}
 1. \quad &\sin 47^\circ + \sin 61^\circ - (\sin 11^\circ + \sin 25^\circ) \\
 &= 2 \sin 54^\circ \cos 7^\circ - 2 \sin 18^\circ \cos 7^\circ \\
 &= 2 \cos 7^\circ (\sin 54^\circ - \sin 18^\circ) \\
 &= 2 \cos 7^\circ (2 \cos 36^\circ \sin 18^\circ) \\
 &= 4 \cos 7^\circ \cdot \frac{\sqrt{5}+1}{4} \cdot \frac{\sqrt{5}-1}{4} \\
 &= \cos 7^\circ \\
 2. \quad &\cos A + \cos(240^\circ + A) + \cos(240^\circ - A) \\
 &= \cos A + 2 \cos 240^\circ \cos A \\
 &= \cos A \{1 + 2 \cos(180^\circ + 60^\circ)\} \\
 &= \cos A \left\{ 1 + 2 \left(-\frac{1}{2} \right) \right\} \\
 &= 0 \\
 3. \quad &\cos \frac{10\pi}{13} + \cos \frac{8\pi}{13} + \cos \frac{3\pi}{13} + \cos \frac{5\pi}{13} \\
 &= \left(\cos \frac{10\pi}{13} + \cos \frac{3\pi}{13} \right) + \left(\cos \frac{8\pi}{13} + \cos \frac{5\pi}{13} \right) \\
 &= 2 \cos \left(\frac{13\pi}{2 \times 13} \right) \cdot \cos \left(\frac{7\pi}{2 \times 13} \right) \\
 &\quad + 2 \cos \left(\frac{13\pi}{2 \times 13} \right) \cos \left(\frac{3\pi}{2 \times 13} \right) \\
 &= 2 \cos \frac{\pi}{2} \left(\cos \frac{7\pi}{26} + \cos \frac{3\pi}{26} \right) \\
 &= 0 \quad \dots \left[\because \cos \frac{\pi}{2} = 0 \right]
 \end{aligned}$$

$$\begin{aligned}
 4. \quad &2 \cos \frac{\pi}{13} \cos \frac{9\pi}{13} + \cos \frac{3\pi}{13} + \cos \frac{5\pi}{13} \\
 &= 2 \cos \frac{\pi}{13} \cos \frac{9\pi}{13} + 2 \cos \frac{4\pi}{13} \cos \frac{\pi}{13} \\
 &= 2 \cos \frac{\pi}{13} \left[\cos \frac{9\pi}{13} + \cos \frac{4\pi}{13} \right] \\
 &= 2 \cos \frac{\pi}{13} \left[2 \cos \frac{\pi}{2} \cos \frac{5\pi}{26} \right] \\
 &= 0 \quad \dots \left[\because \cos \frac{\pi}{2} = 0 \right]
 \end{aligned}$$

$$\begin{aligned}
 5. \quad &2 \cos x - \cos 3x - \cos 5x \\
 &= 2 \cos x (1 - \cos 4x) \\
 &= 2 \cos x \cdot 2 \sin^2 2x \\
 &= 4 \cos x \sin^2 2x \\
 &= 4 \cos x (2 \sin x \cos x)^2 \\
 &= 16 \sin^2 x \cos^3 x \\
 6. \quad &1 + \cos 10^\circ + \cos 20^\circ + \cos 30^\circ \\
 &= 2 \cos^2 5^\circ + 2 \cos 25^\circ \cos 5^\circ \\
 &= 2 \cos 5^\circ (\cos 5^\circ + \cos 25^\circ) \\
 &= 2 \cos 5^\circ (2 \cos 15^\circ \cos 10^\circ) \\
 &= 4 \cos 5^\circ \cos 10^\circ \cos 15^\circ \\
 7. \quad &1 + \cos 56^\circ + \cos 58^\circ - \cos 66^\circ \\
 &= 2 \cos^2 28^\circ + 2 \sin 62^\circ \sin 4^\circ \\
 &= 2 \cos^2 28^\circ + 2 \cos 28^\circ \cos 86^\circ \\
 &\quad \dots \left[\because \sin(90^\circ - \theta) = \cos \theta \right] \\
 &= 2 \cos 28^\circ (\cos 28^\circ + \cos 86^\circ) \\
 &= 2 \cos 28^\circ \cdot 2 \cos 57^\circ \cos 29^\circ \\
 &= 4 \cos 28^\circ \cos 29^\circ \sin 33^\circ \\
 &\quad \dots \left[\because \cos(90^\circ - \theta) = \sin \theta \right] \\
 8. \quad &\frac{\sin 85^\circ - \sin 35^\circ}{\cos 65^\circ} = \frac{2 \cos 60^\circ \sin 25^\circ}{\sin 25^\circ} \\
 &= 2 \times \frac{1}{2} \\
 &= 1 \\
 9. \quad &\frac{\sin 55^\circ - \cos 55^\circ}{\sin 10^\circ} = \frac{\sin 55^\circ - \sin 35^\circ}{\sin 10^\circ} \\
 &= \frac{2 \cos 45^\circ \sin 10^\circ}{\sin 10^\circ} = \sqrt{2} \\
 10. \quad &\tan 9^\circ - \tan 27^\circ - \tan 63^\circ + \tan 81^\circ \\
 &= \tan 9^\circ - \tan 27^\circ - \cot 27^\circ + \cot 9^\circ \\
 &\quad \dots \left[\because \tan(90^\circ - \theta) = \cot \theta \right]
 \end{aligned}$$



$$\begin{aligned}
 &= (\tan 9^\circ + \cot 9^\circ) - (\tan 27^\circ + \cot 27^\circ) \\
 &= \frac{1}{\sin 9^\circ \cos 9^\circ} - \frac{1}{\sin 27^\circ \cos 27^\circ} \\
 &= \frac{2}{\sin 18^\circ} - \frac{2}{\sin 54^\circ} \\
 &\quad \dots [\because \sin 2\theta = 2 \sin \theta \cos \theta] \\
 &= 2 \left\{ \frac{\sin 54^\circ - \sin 18^\circ}{\sin 18^\circ \sin 54^\circ} \right\} \\
 &= 2 \cdot \frac{2 \cos 36^\circ \sin 18^\circ}{\sin 18^\circ \sin 54^\circ} \\
 &= \frac{4 \cos 36^\circ}{\cos 36^\circ} \\
 &= 4
 \end{aligned}$$

$$\begin{aligned}
 11. \quad \frac{\sin A - \sin B}{\cos A + \cos B} &= \frac{2 \cdot \cos \left(\frac{A+B}{2} \right) \cdot \sin \left(\frac{A-B}{2} \right)}{2 \cdot \cos \left(\frac{A+B}{2} \right) \cdot \cos \left(\frac{A-B}{2} \right)} \\
 &= \tan \left(\frac{A-B}{2} \right)
 \end{aligned}$$

$$\begin{aligned}
 12. \quad \frac{\sin(B+A) + \cos(B-A)}{\sin(B-A) + \cos(B+A)} \\
 &= \frac{\sin(B+A) + \sin\{90^\circ - (B-A)\}}{\sin(B-A) + \sin\{90^\circ - (A+B)\}} \\
 &= \frac{2 \sin(A+45^\circ) \cos(45^\circ - B)}{2 \sin(45^\circ - A) \cos(45^\circ - B)} \\
 &= \frac{\sin(A+45^\circ)}{\sin(45^\circ - A)} = \frac{\cos A + \sin A}{\cos A - \sin A}
 \end{aligned}$$

$$\begin{aligned}
 13. \quad \sin 4A - \cos 2A &= \cos 4A - \sin 2A \\
 \Rightarrow \sin 4A + \sin 2A &= \cos 4A + \cos 2A \\
 \Rightarrow 2 \sin 3A \cos A &= 2 \cos 3A \cos A \\
 \Rightarrow \tan 3A &= 1
 \end{aligned}$$

$$\Rightarrow 3A = \frac{\pi}{4}$$

$$\Rightarrow 4A = \frac{\pi}{3}$$

$$\begin{aligned}
 \therefore \tan 4A &= \tan \frac{\pi}{3} \\
 &= \sqrt{3}
 \end{aligned}$$

$$\begin{aligned}
 14. \quad \sin x + \sin y &= \frac{1}{2} \\
 \Rightarrow 2 \sin \left(\frac{x+y}{2} \right) \cos \left(\frac{x-y}{2} \right) &= \frac{1}{2} \quad \dots (i) \\
 \cos x + \cos y &= 1
 \end{aligned}$$

$$\Rightarrow 2 \cos \left(\frac{x+y}{2} \right) \cos \left(\frac{x-y}{2} \right) = 1 \quad \dots (ii)$$

Dividing (i) by (ii), we get

$$\tan \left(\frac{x+y}{2} \right) = \frac{1}{2}$$

$$\begin{aligned}
 \text{Now, } \tan(x+y) &= \frac{2 \tan \left(\frac{x+y}{2} \right)}{1 - \tan^2 \left(\frac{x+y}{2} \right)} \\
 &= \frac{2 \left(\frac{1}{2} \right)}{1 - \frac{1}{4}} = \frac{4}{3}
 \end{aligned}$$

$$15. \quad \cos x = 3 \cos y \Rightarrow \frac{\cos x}{\cos y} = \frac{3}{1}$$

By componendo and dividendo, we get

$$\begin{aligned}
 \frac{\cos x + \cos y}{\cos x - \cos y} &= \frac{3+1}{3-1} \\
 \Rightarrow \frac{2 \cos \left(\frac{x+y}{2} \right) \cos \left(\frac{x-y}{2} \right)}{-2 \sin \left(\frac{x+y}{2} \right) \sin \left(\frac{x-y}{2} \right)} &= \frac{4}{2}
 \end{aligned}$$

$$\Rightarrow -\cot \left(\frac{x+y}{2} \right) \cot \left(\frac{x-y}{2} \right) = 2$$

$$\Rightarrow \cot \left(\frac{x+y}{2} \right) \cot \left(\frac{y-x}{2} \right) = 2$$

$$\Rightarrow 2 \tan \left(\frac{y-x}{2} \right) = \cot \left(\frac{x+y}{2} \right)$$

$$16. \quad \text{Given that, } \cos A = m \cos B$$

$$\Rightarrow \frac{m}{1} = \frac{\cos A}{\cos B}$$

By componendo and dividendo, we get

$$\begin{aligned}
 \frac{m+1}{m-1} &= \frac{\cos A + \cos B}{\cos A - \cos B} \\
 &= \frac{2 \cos \left(\frac{A+B}{2} \right) \cos \left(\frac{B-A}{2} \right)}{2 \sin \left(\frac{A+B}{2} \right) \sin \left(\frac{B-A}{2} \right)}
 \end{aligned}$$

.... [$\because \cos(B-A) = \cos(A-B)$]

$$\begin{aligned}
 &= \cot \left(\frac{A+B}{2} \right) \cot \left(\frac{B-A}{2} \right) \\
 \Rightarrow \cot \left(\frac{A+B}{2} \right) &= \frac{m+1}{m-1} \tan \left(\frac{B-A}{2} \right)
 \end{aligned}$$



$$\begin{aligned}
 17. \quad S &= \sin \theta + \sin 2\theta + \sin 3\theta + \dots + \sin n\theta \\
 \text{Since,} \\
 \sin \theta + \sin(\theta + \beta) + \sin(\theta + 2\beta) \\
 &\quad + \dots + \sin[\theta + (n-1)\beta] \\
 &= \frac{\sin \frac{n\beta}{2}}{\sin \frac{\beta}{2}} \sin \left[\theta + \left(\frac{n-1}{2} \right) \beta \right]
 \end{aligned}$$

Here, $\beta = \theta$

$$\therefore S = \frac{\sin \frac{n\theta}{2} \cdot \sin \frac{\theta(n+1)}{2}}{\sin \frac{\theta}{2}}$$

$$\begin{aligned}
 18. \quad \text{Since, } \cos \alpha + \cos 2\alpha + \cos 3\alpha + \dots + \cos n\alpha \\
 = \frac{\cos \left(\frac{n+1}{2} \right) \alpha \cdot \sin \left(\frac{n\alpha}{2} \right)}{\sin \left(\frac{\alpha}{2} \right)}
 \end{aligned}$$

Here, $n = 3$ and $\alpha = \frac{2\pi}{7}$

$$\begin{aligned}
 \therefore \cos \frac{2\pi}{7} + \cos \frac{4\pi}{7} + \cos \frac{6\pi}{7} \\
 = \frac{\cos \left(\frac{3+1}{2} \right) \left(\frac{2\pi}{7} \right) \sin \left(\frac{3 \times 2\pi}{2 \times 7} \right)}{\sin \left(\frac{2\pi}{7 \times 2} \right)} \\
 = \frac{\cos \left(\frac{4\pi}{7} \right) \cdot \sin \left(\frac{3\pi}{7} \right)}{\sin \left(\frac{\pi}{7} \right)}
 \end{aligned}$$

Since, the values of $\cos \left(\frac{4\pi}{7} \right)$, $\sin \left(\frac{3\pi}{7} \right)$ and $\sin \left(\frac{\pi}{7} \right)$ are -ve, +ve and +ve respectively.

\therefore option (C) is the correct answer.

$$19. \quad \cos A = \frac{3}{4} \Rightarrow \sin A = \frac{\sqrt{7}}{4}$$

$$\begin{aligned}
 \text{Now, } 32 \sin \frac{A}{2} \cos \frac{5A}{2} \\
 = 16 (\sin 3A - \sin 2A) \\
 = 16 (3\sin A - 4\sin^3 A - 2\sin A \cos A) \\
 = 16 \sin A (3 - 4\sin^2 A - 2\cos A) \\
 = 16 \cdot \frac{\sqrt{7}}{4} \left(3 - 4 \cdot \frac{7}{16} - 2 \cdot \frac{3}{4} \right) \\
 = -\sqrt{7}
 \end{aligned}$$

$$\begin{aligned}
 20. \quad \sin 12^\circ \sin 48^\circ \sin 54^\circ \\
 = \frac{1}{2} (\cos 36^\circ - \cos 60^\circ) \cos 36^\circ \\
 = \frac{1}{2} \left[\frac{\sqrt{5}+1}{4} - \frac{1}{2} \right] \left[\frac{\sqrt{5}+1}{4} \right] \\
 = \frac{1}{2} \left[\frac{\sqrt{5}-1}{4} \right] \left[\frac{\sqrt{5}+1}{4} \right] \\
 = \frac{5-1}{32} \\
 = \frac{1}{8}
 \end{aligned}$$

$$\begin{aligned}
 21. \quad \sin 12^\circ \sin 24^\circ \sin 48^\circ \sin 84^\circ \\
 = \frac{1}{4} (2 \sin 12^\circ \sin 48^\circ) (2 \sin 24^\circ \sin 84^\circ) \\
 = \frac{1}{2} (\cos 36^\circ - \cos 60^\circ) (\cos 60^\circ - \cos 108^\circ) \\
 = \frac{1}{4} \left(\cos 36^\circ - \frac{1}{2} \right) \left(\frac{1}{2} + \sin 18^\circ \right) \\
 = \frac{1}{4} \left\{ \frac{1}{4} (\sqrt{5}+1) - \frac{1}{2} \right\} \left\{ \frac{1}{2} + \frac{1}{4} (\sqrt{5}-1) \right\} \\
 = \frac{1}{16}
 \end{aligned}$$

Consider, $\cos 20^\circ \cos 40^\circ \cos 60^\circ \cos 80^\circ$

$$\begin{aligned}
 = \frac{1}{2} [\cos (60^\circ - 20^\circ) \cos 20^\circ \cos (60^\circ + 20^\circ)] \\
 = \frac{1}{2} \left[\frac{1}{4} \cos 3 (20^\circ) \right]
 \end{aligned}$$

$$\dots \left[\because \cos \theta \cos (60^\circ - \theta) \cos (60^\circ + \theta) = \frac{1}{4} \cos 3\theta \right]$$

$$= \frac{1}{8} \cos 60^\circ = \frac{1}{8} \times \frac{1}{2} = \frac{1}{16}$$

\therefore option (A) is the correct answer.

$$\begin{aligned}
 22. \quad \left(1 + \cos \frac{\pi}{8} \right) \left(1 + \cos \frac{7\pi}{8} \right) \left(1 + \cos \frac{3\pi}{8} \right) \left(1 + \cos \frac{5\pi}{8} \right) \\
 = \left(1 + \cos \frac{\pi}{8} \right) \left(1 - \cos \frac{\pi}{8} \right) \left(1 + \cos \frac{3\pi}{8} \right) \left(1 - \cos \frac{3\pi}{8} \right) \\
 \dots [\because \cos (\pi - \theta) = -\cos \theta] \\
 = \left(1 - \cos^2 \frac{\pi}{8} \right) \left(1 - \cos^2 \frac{3\pi}{8} \right)
 \end{aligned}$$



$$\begin{aligned}
 &= \sin^2 \frac{\pi}{8} \sin^2 \frac{3\pi}{8} \\
 &= \frac{1}{4} \left(2 \sin \frac{\pi}{8} \cdot \sin \frac{3\pi}{8} \right)^2 \\
 &= \frac{1}{4} \left(\cos \frac{\pi}{4} - \cos \frac{\pi}{2} \right)^2 \\
 &= \frac{1}{8}
 \end{aligned}$$

$$23. \quad \frac{m}{n} = \frac{\tan(\theta + 120^\circ)}{\tan(\theta - 30^\circ)}$$

$$\Rightarrow \frac{m+n}{m-n} = \frac{\tan(\theta + 120^\circ) + \tan(\theta - 30^\circ)}{\tan(\theta + 120^\circ) - \tan(\theta - 30^\circ)}$$

....[By componendo and dividendo]

$$= \frac{\sin(\theta + 120^\circ)\cos(\theta - 30^\circ) + \cos(\theta + 120^\circ)\sin(\theta - 30^\circ)}{\sin(\theta + 120^\circ)\cos(\theta - 30^\circ) - \cos(\theta + 120^\circ)\sin(\theta - 30^\circ)}$$

$$= \frac{\sin(2\theta + 90^\circ)}{\sin(150^\circ)} = \frac{\cos 2\theta}{\frac{1}{2}}$$

$$= 2 \cos 2\theta$$

$$24. \quad \cos^2 76^\circ + \cos^2 16^\circ - \cos 76^\circ \cos 16^\circ$$

$$= \frac{1}{2} [1 + \cos 152^\circ + 1 + \cos 32^\circ - \cos 92^\circ - \cos 60^\circ]$$

$$= \frac{1}{2} \left[2 - \frac{1}{2} + \cos 152^\circ + \cos 32^\circ - \cos 92^\circ \right]$$

$$= \frac{1}{2} \left[\frac{3}{2} + 2 \cos 92^\circ \cos 60^\circ - \cos 92^\circ \right]$$

$$= \frac{1}{2} \left[\frac{3}{2} + \cos 92^\circ - \cos 92^\circ \right]$$

$$= \frac{3}{4}$$

$$25. \quad \cos 2(\alpha + \beta) = 2 \cos^2(\alpha + \beta) - 1$$

$$\text{and } 2 \sin^2 \beta = 1 - \cos 2\beta$$

$$\text{Now, } 2 \sin^2 \beta + 4 \cos(\alpha + \beta) \sin \alpha \sin \beta$$

$$+ \cos 2(\alpha + \beta)$$

$$= 1 - \cos 2\beta + 4 \cos(\alpha + \beta) \sin \alpha \sin \beta$$

$$+ 2 \cos^2(\alpha + \beta) - 1$$

$$\begin{aligned}
 &= 2 \cos(\alpha + \beta)[2 \sin \alpha \sin \beta + \cos(\alpha + \beta)] - \cos 2\beta \\
 &= -\cos 2\beta + 2 \cos(\alpha + \beta) \cos(\alpha - \beta) \\
 &= -\cos 2\beta + \cos 2\alpha + \cos 2\beta \\
 &= \cos 2\alpha
 \end{aligned}$$

$$26. \quad \text{In } \triangle ABC, A + B + C = \pi$$

$$\sin 2A + \sin 2B + \sin 2C$$

$$= 2 \sin(A + B) \cos(A - B) + 2 \sin C \cos C$$

$$= 2 \sin(\pi - C) \cos(A - B) + 2 \sin C \cos\{\pi - (A + B)\}$$

$$= 2 \sin C \cos(A - B) - 2 \sin C \cos(A + B)$$

$$= 2 \sin C \{\cos(A - B) - \cos(A + B)\}$$

$$= 2 \sin C (2 \sin A \sin B)$$

$$= 4 \sin A \sin B \sin C$$

$$27. \quad \cos A = \cos B \cos C$$

$$\Rightarrow \cos[\pi - (B + C)] = \cos B \cos C$$

$$\Rightarrow -\cos(B + C) = \cos B \cos C$$

$$\Rightarrow -[\cos B \cos C - \sin B \sin C] = \cos B \cos C$$

$$\Rightarrow \sin B \sin C = 2 \cos B \cos C$$

$$\Rightarrow \tan B \tan C = 2$$

$$28. \quad \cot(A + B) = \cot(\pi - C)$$

$$\Rightarrow \frac{\cot A \cot B - 1}{\cot A + \cot B} = -\cot C$$

$$\Rightarrow \cot A \cot B - 1 = -\cot A \cot C - \cot B \cot C$$

$$\Rightarrow \cot B \cot C + \cot C \cot A + \cot A \cot B = 1$$

$$29. \quad \cot B + \cot C = \frac{\sin C \cos B + \sin B \cos C}{\sin B \sin C}$$

$$= \frac{\sin(B + C)}{\sin B \sin C}$$

$$= \frac{\sin(180^\circ - A)}{\sin B \sin C}$$

$$= \frac{\sin A}{\sin B \sin C}$$

$$\text{Similarly, } \cot C + \cot A = \frac{\sin B}{\sin C \sin A}$$

$$\text{and } \cot A + \cot B = \frac{\sin C}{\sin A \sin B}$$

$$\therefore (\cot B + \cot C)(\cot C + \cot A)(\cot A + \cot B)$$

$$= \frac{\sin A}{\sin B \sin C} \cdot \frac{\sin B}{\sin C \sin A} \cdot \frac{\sin C}{\sin A \sin B}$$

$$= \operatorname{cosec} A \operatorname{cosec} B \operatorname{cosec} C$$

$$30. \quad \text{Since, } \tan A + \tan B + \tan C = \tan A \tan B \tan C$$

$$\therefore \frac{\tan A + \tan B + \tan C}{\tan A \tan B \tan C} = 1$$



31. Given, $\angle A = \frac{\pi}{2}$
 In ΔABC , $\angle A + \angle B + \angle C = \pi$
 $\therefore \angle B + \angle C = \frac{\pi}{2}$
 $\Rightarrow B = \frac{\pi}{2} - C$
 $\Rightarrow \cos^2 B = \cos^2 \left(\frac{\pi}{2} - C \right) = \sin^2 C$
 $\therefore \cos^2 B + \cos^2 C = \sin^2 C + \cos^2 C = 1$

32. $A + B + C = 180^\circ$
 $\Rightarrow \cot \left(\frac{A}{2} + \frac{B}{2} \right) = \cot \left(90^\circ - \frac{C}{2} \right)$
 $\Rightarrow \frac{\cot \frac{A}{2} \cdot \cot \frac{B}{2} - 1}{\cot \frac{B}{2} + \cot \frac{A}{2}} = \tan \frac{C}{2} = \frac{1}{\cot \frac{C}{2}}$
 $\Rightarrow \left(\cot \frac{A}{2} \cot \frac{B}{2} - 1 \right) \cot \frac{C}{2} = \cot \frac{B}{2} + \cot \frac{A}{2}$

$$\Rightarrow \cot \frac{A}{2} \cot \frac{B}{2} \cot \frac{C}{2} = \cot \frac{C}{2} + \cot \frac{B}{2} + \cot \frac{A}{2}$$

33. $A + B + C = \pi$
 $\therefore \tan \left(\frac{A+B}{2} \right) = \tan \left(\frac{\pi}{2} - \frac{C}{2} \right)$
 $\Rightarrow \frac{\tan \frac{A}{2} + \tan \frac{B}{2}}{1 - \tan \frac{A}{2} \cdot \tan \frac{B}{2}} = \cot \frac{C}{2}$
 $\Rightarrow \frac{\frac{1}{3} + \frac{2}{3}}{1 - \frac{1}{3} \cdot \frac{2}{3}} = \cot \frac{C}{2}$
 $\Rightarrow \frac{9}{7} = \cot \frac{C}{2}$
 $\Rightarrow \tan \frac{C}{2} = \frac{7}{9}$

34. For $A = B = C = 60^\circ$, only option (C) satisfies the condition.



Evaluation Test

1. We have, $A + B + C = 180^\circ$
 $\Rightarrow \frac{A}{2} = 90^\circ - \left(\frac{B+C}{2} \right)$
 $\Rightarrow \cot \frac{A}{2} = \cot \left(90^\circ - \left(\frac{B+C}{2} \right) \right)$
 $\Rightarrow \cot \frac{A}{2} = \tan \left(\frac{B}{2} + \frac{C}{2} \right)$
 $\Rightarrow \frac{1}{\tan \frac{A}{2}} = \frac{\tan \frac{B}{2} + \tan \frac{C}{2}}{1 - \tan \frac{B}{2} \tan \frac{C}{2}}$
 $\Rightarrow 1 - \tan \frac{B}{2} \tan \frac{C}{2} = \tan \frac{A}{2} \tan \frac{B}{2} + \tan \frac{A}{2} \tan \frac{C}{2}$
 $\Rightarrow \tan \frac{A}{2} \tan \frac{B}{2} + \tan \frac{B}{2} \tan \frac{C}{2} + \tan \frac{A}{2} \tan \frac{C}{2} = 1$
 i.e., $\sum \tan \frac{A}{2} \tan \frac{B}{2} = 1$

2. $\sin^2 \alpha + \sin^2 \beta - \sin^2 \gamma$
 $= \sin^2 \alpha + \sin (\beta - \gamma) \sin (\beta + \gamma)$
 $\dots [\because \sin^2 A - \sin^2 B = \sin (A + B) \sin (A - B)]$
 $= \sin^2 \alpha + \sin (\pi - \alpha) \sin (\beta + \gamma)$
 $\dots [\because \alpha + \beta - \gamma = \pi \text{ (given)}]$

$$= \sin^2 \alpha + \sin \alpha \sin (\beta + \gamma)$$

$$= \sin \alpha \{ \sin \alpha + \sin (\beta + \gamma) \}$$

$$= \sin \alpha \{ \sin (\pi - (\beta - \gamma)) + \sin (\beta + \gamma) \}$$

$$= \sin \alpha \{ \sin (\beta - \gamma) + \sin (\beta + \gamma) \}$$

$$= \sin \alpha (2 \sin \beta \cos \gamma)$$

$$= 2 \sin \alpha \sin \beta \cos \gamma$$

3. $(\cos \theta + \cos 7\theta) + (\cos 3\theta + \cos 5\theta) = 0$
 $\Rightarrow 2 \cos 4\theta \cos 3\theta + 2 \cos 4\theta \cos \theta = 0$
 $\dots \left[\because \cos C + \cos D = 2 \cos \frac{C+D}{2} \cos \frac{C-D}{2} \right]$
 $\Rightarrow 2 \cos 4\theta (\cos 3\theta + \cos \theta) = 0$
 $\Rightarrow 4 \cos 4\theta \cos 2\theta \cos \theta = 0$
 $\Rightarrow 4 \frac{\sin 2^3 \theta}{2^3 \sin \theta} = 0$
 $\dots \left[\because \cos A \cos 2A \cos 2^2 A \cos 2^3 A \dots \cos 2^{n-1} A = \frac{\sin 2^n A}{2^n \sin A} \right]$
 $\Rightarrow \sin 8\theta = 0 \Rightarrow 8\theta = n\pi$
 $\Rightarrow \theta = \frac{n\pi}{8}$



4. Given, $\sin A + \sin B = C$
and $\cos A + \cos B = D$

$$\therefore \frac{\sin A + \sin B}{\cos A + \cos B} = \frac{C}{D}$$

$$\Rightarrow \frac{2 \sin\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)}{2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)} = \frac{C}{D}$$

$$\Rightarrow \tan\left(\frac{A+B}{2}\right) = \frac{C}{D} \quad \dots(i)$$

$$\text{Now, } \sin(A+B) = \frac{2 \tan\left(\frac{A+B}{2}\right)}{1 + \tan^2\left(\frac{A+B}{2}\right)}$$

$$\dots \left[\because \sin \theta = \frac{2 \tan \frac{\theta}{2}}{1 + \tan^2 \frac{\theta}{2}} \right]$$

$$= \frac{2 \frac{C}{D}}{1 + \frac{C^2}{D^2}} \quad \dots[\text{From (i)}]$$

$$= \frac{2CD}{C^2 + D^2}$$

5. $\frac{\sin \alpha - \sin \gamma}{\cos \gamma - \cos \alpha} = \frac{2 \cos \frac{\alpha+\gamma}{2} \sin \frac{\alpha-\gamma}{2}}{2 \sin \frac{\alpha+\gamma}{2} \sin \frac{\alpha-\gamma}{2}}$

$$= \cot \frac{\alpha+\gamma}{2}$$

But α, β, γ are in A.P. $\Rightarrow \frac{\alpha+\gamma}{2} = \beta$

$$\therefore \frac{\sin \alpha - \sin \gamma}{\cos \gamma - \cos \alpha} = \cot \beta$$

6. $\sin^3 x \sin 3x$

$$= \frac{1}{4} (3 \sin x - \sin 3x) \sin 3x$$

$$\dots \left[\begin{array}{l} \because \sin 3A = 3 \sin A - 4 \sin^3 A \\ \Rightarrow \sin^3 A = \frac{1}{4} (3 \sin A - \sin 3A) \end{array} \right]$$

$$= \frac{3}{8} (2 \sin x \sin 3x) - \frac{1}{8} (2 \sin^2 3x)$$

$$= \frac{3}{8} (\cos 2x - \cos 4x) - \frac{1}{8} (1 - \cos 6x)$$

$$= -\frac{1}{8} + \frac{3}{8} \cos 2x - \frac{3}{8} \cos 4x + \frac{1}{8} \cos 6x \quad \dots(i)$$

and $\sum_{m=0}^n c_m \cos mx$

$$= c_0 + c_1 \cos x + c_2 \cos 2x + c_3 \cos 3x$$

$$+ \dots + c_n \cos nx \quad \dots(ii)$$

But, $\sin^3 x \sin 3x = \sum_{m=0}^n c_m \cos mx$

\therefore from (i) and (ii), we get $n = 6$.

7. Given, $\sin B = \frac{1}{5} \sin(2A+B)$

$$\Rightarrow \frac{\sin(2A+B)}{\sin B} = \frac{5}{1}$$

By componendo and dividendo, we get

$$\frac{\sin(2A+B) + \sin B}{\sin(2A+B) - \sin B} = \frac{5+1}{5-1}$$

$$\Rightarrow \frac{2 \sin(A+B) \cos A}{2 \cos(A+B) \sin A} = \frac{6}{4}$$

$$\Rightarrow \frac{\tan(A+B)}{\tan A} = \frac{3}{2}$$

8. $\sin A + \sin 2A + \sin 3A = \cos A + \cos 2A + \cos 3A$

$$\therefore (\sin 3A + \sin A) + \sin 2A = (\cos 3A + \cos A) + \cos 2A$$

$$\therefore 2 \sin 2A \cos A + \sin 2A = 2 \cos 2A \cos A + \cos 2A$$

$$\therefore \sin 2A(2 \cos A + 1) = \cos 2A(2 \cos A + 1)$$

$$\therefore \tan 2A = 1$$

06 Straight Line




Hints



Classical Thinking

2. Gradient of the line which passes through (1, 0) and $(-2, \sqrt{3})$ is $m = \frac{\sqrt{3}-0}{-2-1} = -\frac{1}{\sqrt{3}}$
 $\Rightarrow \tan \theta = -\frac{1}{\sqrt{3}}$
 $\Rightarrow \theta = \tan^{-1}\left(-\frac{1}{\sqrt{3}}\right) = 150^\circ$
3. The required equation is $y + 6 = \tan 45^\circ(x - 4)$
 $\Rightarrow x - y - 10 = 0$
4. The required equation passing through (0, 0) and having gradient $m = \frac{1}{0}$, is $y = \frac{1}{0}x$
 $\Rightarrow x = 0$
5. Midpoint is (3, 4) and slope of AB = $\frac{6}{4}$
 \therefore Slope of perpendicular = $\frac{-1}{6/4} = -\frac{2}{3}$
 \therefore the required equation is $y - 4 = -\frac{2}{3}(x - 3)$
 $\Rightarrow 2x + 3y = 18$
6. $m = \frac{-1}{\frac{b'-b}{a'-a}} = \frac{a'-a}{b-b'}$
 Midpoint is $\left(\frac{a+a'}{2}, \frac{b+b'}{2}\right)$
 \therefore the required equation is $y - \left(\frac{b+b'}{2}\right) = \frac{a'-a}{b-b'} \left[x - \left(\frac{a+a'}{2}\right)\right]$
 $\Rightarrow 2(b-b')y + 2(a-a')x = b^2 - b'^2 + a^2 - a'^2$
7. Midpoint = (4, -9) and slope = $\frac{-1}{\frac{3+1}{-1-5}} = \frac{3}{2}$
 Hence, the required line is $y + 9 = \frac{3}{2}(x - 4)$
 $\Rightarrow 3x - 2y = 30$

8.



$$\text{Slope of QR} = \frac{3\sqrt{3}-0}{3-0} = \sqrt{3} \text{ i.e., } \theta = 60^\circ$$

Clearly, $\angle PQR = 120^\circ$

OQ is the angle bisector of the angle PQR, so line OQ makes 120° with the positive direction of X-axis.

Therefore, equation of the bisector of $\angle PQR$ is $y = \tan 120^\circ x \Rightarrow y = -\sqrt{3}x \Rightarrow \sqrt{3}x + y = 0$

9. $m = \frac{5-0}{-4-0} = -\frac{5}{4}$

\therefore the required equation is $5x + 4y = 0$.

10. Equation of a line passing through the given points is $\frac{y-(-6)}{-6-10} = \frac{x-(-5)}{-5-3}$
 $\Rightarrow \frac{y+6}{-16} = \frac{x+5}{-8} \Rightarrow 2x - y + 4 = 0$


11. The point of intersection is (0, 0)
 Thus, the equation of line passing through the points (0, 0) and (2, 2) is $y = x$.

12. Equation of line is $y = mx + c$
 $\Rightarrow y = (\tan 135^\circ)x - 5 \Rightarrow y = -x - 5$
 $\Rightarrow x + y + 5 = 0$

13. From the figure, $m = \tan \theta = \frac{-c}{3}$

$$\Rightarrow 3 = \frac{-c}{3}$$

$$\Rightarrow c = -9$$



Hence, the required equation is $y = 3x - 9$



14. Here, intercept on X-axis is 3 and intercept on Y-axis is -2 .

So, using double intercept form, the required equation of the line is $\frac{x}{3} - \frac{y}{2} = 1$.

15. Using double intercept form, we get

$$\frac{x}{2a \sec \theta} + \frac{y}{2a \operatorname{cosec} \theta} = 1$$

$$\Rightarrow x \cos \theta + y \sin \theta = 2a$$

16. Intersection point on X-axis is $(2x_1, 0)$ and on Y-axis is $(0, 2y_1)$. Thus, equation of line

passing through these points is $\frac{x}{x_1} + \frac{y}{y_1} = 2$.

17. Since, the given line passes through $(2, -3)$ and $(4, -5)$.

$$\therefore \frac{2}{a} - \frac{3}{b} = 1 \quad \text{and} \quad \frac{4}{a} - \frac{5}{b} = 1$$

$$\Rightarrow b = -1, a = -1$$

20. The equation of line is $\frac{x}{a} + \frac{y}{a} = 1$.

$$\Rightarrow x + y - a = 0$$

$$\therefore \text{Slope} = -\frac{\text{coefficient of } x}{\text{coefficient of } y} = -1$$

21. The required equation which passes through $(1, 2)$ and its gradient $m = 3$, is $y - 2 = 3(x - 1)$.

22. The required equation which passes through (c, d) and its gradient $-\frac{a}{b}$, is

$$y - d = -\frac{a}{b}(x - c)$$

$$\Rightarrow a(x - c) + b(y - d) = 0$$

23. The required equation passing through $(3, -4)$ and having gradient $\frac{4}{3}$ is $y + 4 = \frac{4}{3}(x - 3)$.

24. Equation of line perpendicular to $ax + by + c = 0$ is $bx - ay + \lambda = 0$ (i)
It passes through (a, b) .

$$\therefore ab - ab + \lambda = 0 \Rightarrow \lambda = 0$$

Putting $\lambda = 0$ in (i), we get $bx - ay = 0$ which is the required equation.

25. Slope of perpendicular = $\frac{-y'}{2a}$

$$\therefore \text{the required equation is } y - y' = -\frac{y'}{2a}(x - x')$$

$$\Rightarrow xy' + 2ay - 2ay' - x'y' = 0$$

26. $m_1 = \sqrt{3}, m_2 = 0$

$$\therefore \tan \theta = \left| \frac{\sqrt{3} - 0}{1 + 0} \right|$$

$$\Rightarrow \tan \theta = \sqrt{3}$$

$$\Rightarrow \theta = 60^\circ$$

27. $\theta = \tan^{-1} \left| \frac{2 - \sqrt{3} - 2 - \sqrt{3}}{1 + 4 - 3} \right| = \tan^{-1}(\sqrt{3})$

$$\Rightarrow \theta = 60^\circ$$

28. $\theta = \tan^{-1} \left| \frac{-\cot 30^\circ + \cot 60^\circ}{1 + \cot 30^\circ \cot 60^\circ} \right|$

$$= \tan^{-1} \left| \frac{\tan 60^\circ - \tan 30^\circ}{1 + \tan 30^\circ \tan 60^\circ} \right| = 30^\circ$$

29. Equation of lines are $\frac{x}{a} - \frac{y}{b} = 1$ and $\frac{x}{b} - \frac{y}{a} = 1$

$$\Rightarrow m_1 = \frac{b}{a} \quad \text{and} \quad m_2 = \frac{a}{b}$$

$$\therefore \theta = \tan^{-1} \left| \frac{\frac{b}{a} - \frac{a}{b}}{1 + \frac{b}{a} \cdot \frac{a}{b}} \right|$$

$$= \tan^{-1} \frac{b^2 - a^2}{2ab}$$

30. Let $L_1 \equiv 2x + 3y - 7 = 0$ and

$$L_2 \equiv 2x + 3y - 5 = 0$$

$$\text{Here, slope of } L_1 = \text{slope of } L_2 = -\frac{2}{3}$$

Hence, the lines are parallel.

31. Slope of given line is $\frac{1}{2}$

$$\text{Thus, } \tan 45^\circ = \pm \frac{m - \frac{1}{2}}{1 + m \cdot \frac{1}{2}} \Rightarrow m = 3 \text{ or } \frac{-1}{3}$$

Hence option (B) is correct.

33. Let $L_1 \equiv 2x + 5y - 7 = 0$ and $L_2 \equiv 2x - 5y - 9 = 0$,

$$\text{so that } m_1 = -\frac{2}{5}, m_2 = \frac{2}{5}$$

Lines are neither parallel nor perpendicular, also not coincident.

Hence, the lines are intersecting.



34.
$$\begin{vmatrix} m_1 & -1 & c_1 \\ m_2 & -1 & c_2 \\ m_3 & -1 & c_3 \end{vmatrix} = 0$$
- $$\Rightarrow \begin{vmatrix} m_1 & m_2 & m_3 \\ -1 & -1 & -1 \\ c_1 & c_2 & c_3 \end{vmatrix}$$
- $$\Rightarrow m_1(c_2 - c_3) + m_2(c_3 - c_1) + m_3(c_1 - c_2) = 0$$
35. The lines are concurrent, if
$$\begin{vmatrix} 7 & -8 & 5 \\ 3 & -4 & 5 \\ 4 & 5 & k \end{vmatrix} = 0$$
- $$\Rightarrow 7(-4k - 25) + 8(3k - 20) + 5(15 + 16) = 0$$
- $$\Rightarrow k = -45$$
36. consider
$$\begin{vmatrix} 15 & -18 & 1 \\ 12 & 10 & -3 \\ 6 & 66 & -11 \end{vmatrix}$$
- $$= 15(-110 + 198) + 18(-132 + 18) + 1(792 - 60)$$
- $$= 0$$
37. $u = a_1x + b_1y + c_1 = 0$, $v = a_2x + b_2y + c_2 = 0$
- let $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2} = c$
- $$\Rightarrow a_2 = \frac{a_1}{c}, b_2 = \frac{b_1}{c}, c_2 = \frac{c_1}{c}$$
- Given that, $u + kv = 0$
- $$\Rightarrow a_1x + b_1y + c_1 + k(a_2x + b_2y + c_2) = 0$$
- $$\Rightarrow a_1x + b_1y + c_1 + k\left(\frac{a_1}{c}\right)x + k\left(\frac{b_1}{c}\right)y + k\left(\frac{c_1}{c}\right) = 0$$
- $$\Rightarrow a_1x\left(1 + \frac{k}{c}\right) + b_1y\left(1 + \frac{k}{c}\right) + c_1\left(1 + \frac{k}{c}\right) = 0$$
- $$\Rightarrow a_1x + b_1y + c_1 = 0 = u$$
38. Required length =
$$\left| \frac{4(3) + 3(1) + 20}{5} \right| = 7$$
39. Required distance =
$$\left| \frac{-2 - 3 - 5}{\sqrt{1+1}} \right| = \frac{10}{\sqrt{2}} = 5\sqrt{2}$$
40. Required distance =
$$\left| \frac{-7}{\sqrt{12^2 + 5^2}} \right| = \frac{7}{13}$$
41. Here, equation of line is $y = x \tan \alpha + c$, $c > 0$
Length of the perpendicular drawn on line from point $(a \cos \alpha, a \sin \alpha)$ is
- $$p = \left| \frac{-a \sin \alpha + a \cos \alpha \tan \alpha + c}{\sqrt{1 + \tan^2 \alpha}} \right| = \frac{c}{\sec \alpha} = c \cos \alpha$$

42.
$$p = \left| \frac{ab}{\sqrt{a^2 + b^2}} \right|$$

$$\Rightarrow \frac{a^2 + b^2}{a^2 b^2} = \frac{1}{p^2}$$

$$\Rightarrow \frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{p^2}$$

43. Length of perpendicular is

$$\left| \frac{\frac{b}{a} - \frac{a}{b} - 1}{\sqrt{\left(\frac{1}{a}\right)^2 + \left(-\frac{1}{b}\right)^2}} \right| = \left| \frac{b^2 - a^2 - ab}{\sqrt{a^2 + b^2}} \right|$$

44. Straight line $y - y' = \frac{y'' - y'}{x'' - x'}(x - x')$

\therefore Length of perpendicular

$$= \left| \frac{x'(y'' - y') - y'(x'' - x')}{\sqrt{(x'' - x')^2 + (y'' - y')^2}} \right|$$

$$= \left| \frac{x'y'' - y'x''}{\sqrt{(x'' - x')^2 + (y'' - y')^2}} \right|$$

45. Given lines are $5x + 3y - 7 = 0$ (i)

and $15x + 9y + 14 = 0$ or

$$5x + 3y + \frac{14}{3} = 0 \quad \text{....(ii)}$$

Lines (i) and (ii) are parallel.


$$\therefore \text{Required distance} = \left| \frac{c_1 - c_2}{\sqrt{a^2 + b^2}} \right| = \left| \frac{-7 - \frac{14}{3}}{\sqrt{5^2 + 3^2}} \right|$$

$$= \left| \frac{-35}{3\sqrt{34}} \right| = \frac{35}{3\sqrt{34}}$$



Critical Thinking

- The four vertices on solving are $A(-3, 3)$, $B(1, 1)$, $C(1, -1)$ and $D(-2, -2)$.
 $m_1 =$ slope of $AC = -1$,
 $m_2 =$ slope of $BD = 1$
 $\therefore m_1 m_2 = -1$
Hence, the angle between diagonals AC and BD is 90° .
- Mid point of $(a \cos \alpha, a \sin \alpha)$ and $(a \cos \beta, a \sin \beta)$ is
 $P\left(\frac{a(\cos \alpha + \cos \beta)}{2}, \frac{a(\sin \alpha + \sin \beta)}{2}\right)$



∴ Slope of line AB is $\frac{a \sin \beta - a \sin \alpha}{a \cos \beta - a \cos \alpha} = \frac{\sin \beta - \sin \alpha}{\cos \beta - \cos \alpha} = m_1$
 and slope of OP is $\frac{\sin \alpha + \sin \beta}{\cos \alpha + \cos \beta} = m_2$
 Now, $m_1 \times m_2 = \frac{\sin^2 \beta - \sin^2 \alpha}{\cos^2 \beta - \cos^2 \alpha} = -1$
 Hence, the lines are perpendicular.

3. Slope = $\frac{8-2}{3-1} = 3$

The diagonal is $y - 2 = 3(x - 1)$
 $\Rightarrow 3x - y - 1 = 0$

4. S = midpoint of QR = $\left(\frac{6+7}{2}, \frac{-1+3}{2}\right) = \left(\frac{13}{2}, 1\right)$

∴ 'm' of PS = $\frac{2-1}{2-\frac{13}{2}} = -\frac{2}{9}$

∴ The required equation is $y + 1 = -\frac{2}{9}(x - 1)$

i.e., $2x + 9y + 7 = 0$

5. Point P(a, b) is on $3x + 2y = 13$
 So, $3a + 2b = 13$ (i)

Point Q(b, a) is on $4x - y = 5$
 So, $4b - a = 5$ (ii)

By solving (i) and (ii), we get
 $a = 3, b = 2$

Now, equation of PQ is

$$y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1)$$

$$\Rightarrow y - 2 = \frac{3-2}{2-3}(x-3) \Rightarrow y - 2 = -(x - 3)$$

$$\Rightarrow x + y = 5$$

6. Here, slope of AB = 1


$$\Rightarrow \tan \theta = m_1 = 1$$

or $\theta = 45^\circ$

∴ $\tan(45^\circ + 15^\circ) = \tan 60^\circ$


[∵ It is rotated anticlockwise so the angle will be $45^\circ + 15^\circ = 60^\circ$]

Thus, slope of new line is $\sqrt{3}$



∴ The required equation of line passing through (2, 0) and $m = \sqrt{3}$ is $y = \sqrt{3}(x - 2)$
 i.e., $y = \sqrt{3}x - 2\sqrt{3}$

7. Let ABCD be a rectangle.
 Given, A (1, 3) and C (5, 1).



Intersecting point of diagonal of a rectangle is same or at midpoint.

∴ midpoint of AC is (3, 2).
 Also, $y = 2x + c$ passes through (3, 2).
 Hence, $c = -4$


8. Since the hour, minute and second hands always pass through origin because one end of these hands is always at origin.

Now, at 4 O' clock, the hour hand makes 30° angle in fourth quadrant.

So, the equation of hour hand is

$$y = mx \Rightarrow y = -\frac{1}{\sqrt{3}}x$$

$$\Rightarrow x + \sqrt{3}y = 0$$



9. Let the co-ordinates of axes are A (a, 0) and B(0, b), but the point (-5, 4) divides the line AB in the ratio of 1 : 2.

∴ the co-ordinates of axes are $\left(-\frac{15}{2}, 0\right)$ and (0, 12).

Therefore, the equation of line passing through these coordinate axes is given by $8x - 5y + 60 = 0$

10. Let the intercept be a and 2a, then the equation of line is $\frac{x}{a} + \frac{y}{2a} = 1$, but it also passes through (1, 2), therefore $a = 2$.
 Hence, the required equation is $2x + y = 4$



11. Take two perpendicular lines as the coordinate axes. If a, b be the intercepts made by the moving line on the coordinate axes, then the equation of the line is

$$\frac{x}{a} + \frac{y}{b} = 1 \quad \dots(i)$$

$$\text{Let } \frac{1}{a} + \frac{1}{b} = \frac{1}{k}$$

$$\text{i.e., } \frac{k}{a} + \frac{k}{b} = 1 \quad \dots(ii)$$

The result (ii) shows that the straight line (i) passes through a fixed point (k, k) .

12. Given, $a + b = 14 \Rightarrow a = 14 - b$
Hence, the equation of straight line is

$$\frac{x}{14-b} + \frac{y}{b} = 1$$

Also, it passes through $(3, 4)$

$$\therefore \frac{3}{14-b} + \frac{4}{b} = 1$$

$$\Rightarrow b = 8 \text{ or } 7$$

Therefore, equations are $4x + 3y = 24$ and $x + y = 7$

13. Let the equation of the line be $\frac{x}{a} + \frac{y}{b} = 1$

The co-ordinates of the mid point of the intercept AB between the axes are $\left(\frac{a}{2}, \frac{b}{2}\right)$

$$\therefore \frac{a}{2} = 1, \frac{b}{2} = 2 \Rightarrow a = 2, b = 4$$

Hence, the equation of the line is

$$\frac{x}{2} + \frac{y}{4} = 1 \Rightarrow 2x + y = 4$$

14. A line perpendicular to the line $5x - y = 1$ is given by $x + 5y - \lambda = 0 = L$

$$\text{In intercept form } \frac{x}{\lambda} + \frac{y}{\lambda/5} = 1$$

So, area of triangle is $\frac{1}{2} \times$ (Multiplication of intercepts)

$$\Rightarrow \frac{1}{2}(\lambda) \times \left(\frac{\lambda}{5}\right) = 5$$

$$\Rightarrow \lambda = \pm 5\sqrt{2}$$

Hence, the equation of required line is $x + 5y = \pm 5\sqrt{2}$

15. Given form is $3x + 3y + 7 = 0$

$$\Rightarrow \frac{3}{\sqrt{3^2+3^2}}x + \frac{3}{\sqrt{3^2+3^2}}y + \frac{7}{\sqrt{3^2+3^2}} = 0$$

$$\Rightarrow \frac{3}{3\sqrt{2}}x + \frac{3}{3\sqrt{2}}y = \frac{-7}{3\sqrt{2}}$$

$$\therefore p = \left| \frac{-7}{3\sqrt{2}} \right| = \frac{7}{3\sqrt{2}}$$

16. Let p be the length of the perpendicular from the origin on the given line. Then its equation in normal form is

$$x \cos 30^\circ + y \sin 30^\circ = p \text{ or } \sqrt{3}x + y = 2p$$

This meets the coordinate axes at $A\left(\frac{2p}{\sqrt{3}}, 0\right)$

and $B(0, 2p)$

$$\therefore \text{Area of } \Delta OAB = \frac{1}{2} \left(\frac{2p}{\sqrt{3}}\right) 2p = \frac{2p^2}{\sqrt{3}}$$

$$\Rightarrow \frac{2p^2}{\sqrt{3}} = \frac{50}{\sqrt{3}} \Rightarrow p = \pm 5$$

Hence, the lines are $\sqrt{3}x + y \pm 10 = 0$

17. The equation of line passing through

$$A(-5, -4) \text{ is } \frac{x+5}{\cos \theta} = \frac{y+4}{\sin \theta}$$

Let $AB = r_1, AC = r_2, AD = r_3$

The co-ordinate of B is

$$(r_1 \cos \theta - 5, r_1 \sin \theta - 4)$$

which lies on $x + 3y + 2 = 0$

$$\therefore r_1 = \frac{15}{\cos \theta + 3 \sin \theta}$$

Similarly, $\frac{10}{AC} = 2 \cos \theta + \sin \theta$ and

$$\frac{6}{AD} = \cos \theta - \sin \theta$$

Putting in the given relation, we get $(2 \cos \theta + 3 \sin \theta)^2 = 0$

$$\Rightarrow \tan \theta = -\frac{2}{3}$$

$$\therefore \text{The equation of line is } y + 4 = -\frac{2}{3}(x + 5)$$

$$\Rightarrow 2x + 3y + 22 = 0$$

18. Let the required line through the point $(1, 2)$ be inclined at an angle θ to the axis of X. Then its

$$\text{equation is } \frac{x-1}{\cos \theta} = \frac{y-2}{\sin \theta} = r \quad \dots(i)$$



The co-ordinates of a point on the line (i) are $(1 + r \cos \theta, 2 + r \sin \theta)$

If this point is at a distance $\frac{\sqrt{6}}{3}$ from $(1, 2)$,

$$\text{then } r = \frac{\sqrt{6}}{3}$$

Therefore, the point is

$$\left(1 + \frac{\sqrt{6}}{3} \cos \theta, 2 + \frac{\sqrt{6}}{3} \sin \theta\right)$$

But this point lies on the line $x + y = 4$

$$\Rightarrow \frac{\sqrt{6}}{3} (\cos \theta + \sin \theta) = 1 \text{ or}$$

$$\sin \theta + \cos \theta = \frac{3}{\sqrt{6}}$$

$$\Rightarrow \frac{1}{\sqrt{2}} \sin \theta + \frac{1}{\sqrt{2}} \cos \theta = \frac{\sqrt{3}}{2}$$

....(Dividing both sides by $\sqrt{2}$)

$$\Rightarrow \sin(\theta + 45^\circ) = \sin 60^\circ \text{ or } \sin 120^\circ$$

$$\Rightarrow \theta = 15^\circ \text{ or } 75^\circ$$

19. The slope of line $x + y = 1$ is -1 .

\therefore It makes an angle of 135° with X axis.

The equation of line passing through $(1, 1)$ and making an angle of 135° is,

$$\frac{x-1}{\cos 135^\circ} = \frac{y-1}{\sin 135^\circ} = r$$

$$\Rightarrow \frac{x-1}{-1/\sqrt{2}} = \frac{y-1}{1/\sqrt{2}} = r$$

- \therefore Co-ordinates of any point on this line are

$$\left(1 - \frac{r}{\sqrt{2}}, 1 + \frac{r}{\sqrt{2}}\right)$$

If this point lies on $2x - 3y = 4$, then

$$2\left(1 - \frac{r}{\sqrt{2}}\right) - 3\left(1 + \frac{r}{\sqrt{2}}\right) = 4$$

$$\Rightarrow r = \sqrt{2}$$

$$21. \left(\frac{-2}{3a}\right) \left(\frac{-3}{4}\right) = -1 \text{ or } a = \frac{-1}{2}$$

$$22. x \cos \theta - y \sin \theta = a(\cos^4 \theta - \sin^4 \theta) = a \cos 2\theta$$

$$23. \text{Mid point} \equiv \left(\frac{1+1}{2}, \frac{3-7}{2}\right) = (1, -2)$$

Therefore, required line is

$$y + 2 = \frac{2}{3} (x - 1) \Rightarrow 2x - 3y = 8$$


24. Given, line AB makes 0 intercepts on X axis and Y axis so, $(x_1, y_1) = (0, 0)$

$$\text{Slope of perpendicular} = \frac{4}{3}$$

$$\therefore \text{Equation is } y - 0 = \frac{4}{3} (x - 0)$$

$$\Rightarrow 4x - 3y = 0$$

25. Let the equation of perpendicular bisector FN of AB is $x - y + 5 = 0$ (i)



The middle point F of AB is $\left(\frac{x_1+1}{2}, \frac{y_1-2}{2}\right)$ Which lies on line (i).

$$\therefore x_1 - y_1 = -13 \text{(ii)}$$

Also AB is perpendicular to FN. So the product of their slopes is -1 .

$$\text{i.e., } \frac{y_1+2}{x_1-1} \times 1 = -1 \text{ or } x_1 + y_1 = -1 \text{(iii)}$$

On solving (ii) and (iii), we get $B(-7, 6)$

$$\text{Similarly, } C\left(\frac{11}{5}, \frac{2}{5}\right)$$

Hence, the equation of BC is $14x + 23y - 40 = 0$

26. The equation of any line parallel to $2x + 6y + 7 = 0$ is $2x + 6y + k = 0$.

This meets the axes at $A\left(-\frac{k}{2}, 0\right)$ and

$$B\left(0, -\frac{k}{6}\right)$$

Since, $AB = 10$

$$\Rightarrow \sqrt{\frac{k^2}{4} + \frac{k^2}{36}} = 10 \Rightarrow \sqrt{\frac{10k^2}{36}} = 10$$

$$\Rightarrow 10k^2 = 3600 \Rightarrow k = \pm 6\sqrt{10}$$

Hence, there are two lines given by $2x + 6y \pm 6\sqrt{10} = 0$

$$27. \theta = \tan^{-1} \left| \frac{-\cot \alpha_1 + \cot \alpha_2}{1 + \cot \alpha_1 \cot \alpha_2} \right|$$

$$= \tan^{-1} \left| \frac{\tan \alpha_1 - \tan \alpha_2}{1 + \tan \alpha_2 \tan \alpha_1} \right| = \alpha_1 - \alpha_2$$



28. Here, equation of AB is
 $x + 4y - 4 = 0$ (i)
 and equation of BC is
 $2x + y - 22 = 0$ (ii)
 Thus angle between (i) and (ii) is given by

$$\tan^{-1} \left(\frac{-\frac{1}{4} + 2}{1 + \left(-\frac{1}{4}\right)(-2)} \right) = \tan^{-1} \frac{7}{6}$$

29. $\frac{k - (2 + \sqrt{3})}{1 + k(2 + \sqrt{3})} = \sqrt{3}$
 $\Rightarrow k - 2 - \sqrt{3} = \sqrt{3} + 2k\sqrt{3} + 3k$
 $\Rightarrow k = \frac{-2(1 + \sqrt{3})}{2(1 + \sqrt{3})} = -1$

30. Let θ be the acute angle which the line
 $y = mx + 4$ makes with the lines $y = 3x + 1$ and
 $2y = x + 3$.
 Then,

$$\tan \theta = \left| \frac{m-3}{1+3m} \right| \text{ and } \tan \theta = \left| \frac{m-\frac{1}{2}}{1+\frac{m}{2}} \right|$$

$$\Rightarrow \left| \frac{m-3}{1+3m} \right| = \left| \frac{2m-1}{m+2} \right|$$

$$\Rightarrow m^2 - m - 6 = \pm (6m^2 - m - 1)$$

$$\Rightarrow 5m^2 + 5 = 0 \text{ or } 7m^2 - 2m - 7 = 0$$

$$\Rightarrow 7m^2 - 2m - 7 = 0$$

$$\Rightarrow m = \frac{1 \pm 5\sqrt{2}}{7}$$

31. Any line through $(1, -10)$ is given by
 $y + 10 = m(x - 1)$
 Since, it makes equal angle say ' α ' with the
 given lines $7x - y + 3 = 0$ and $x + y - 3 = 0$

$$\therefore \tan \alpha = \frac{m-7}{1+7m} = \frac{m-(-1)}{1+m(-1)}$$

$$\Rightarrow m = \frac{1}{3} \text{ or } -3$$

Hence, the two possible equations of third side
 are $3x + y + 7 = 0$, $x - 3y - 31 = 0$.

32. Here,
 Slope of Ist diagonal = $m_1 = \frac{2-0}{2-0} = 1$
 $\Rightarrow \theta_1 = 45^\circ$

$$\text{Slope of II}^{\text{nd}} \text{ diagonal} = m_2 = \frac{2-0}{1-1} = \infty$$

$$\Rightarrow \theta_2 = 90^\circ$$

$$\Rightarrow \theta_2 - \theta_1 = 45^\circ = \frac{\pi}{4}$$

33. Intersection point of the line is
 $\left(\frac{ab}{a+b}, \frac{ab}{a+b} \right)$, which is satisfying all the
 equations given in options (A), (B) and (C).
 Hence, (D) is correct.


34. Putting $k = 1, 2$, we get
 $3x + 2y = 12$ (i)
 $4x + 3y = 19$ (ii)
 The given lines are not parallel.
 Hence on solving them, we get

$$x = -2, y = 9$$

Therefore, the lines pass through $(-2, 9)$

35. Slope of AC = $5/2$.
 Let m be the slope of a line inclined at an
 angle of 45° to AC,

$$\text{Then } \tan 45^\circ = \pm \frac{m - \frac{5}{2}}{1 + m \cdot \frac{5}{2}} \Rightarrow m = -\frac{7}{3}, \frac{3}{7}$$



Thus, let the slope of AB or DC be $\frac{3}{7}$ and that

of AD or BC be $-\frac{7}{3}$.

Then, equation of AB is $3x - 7y + 19 = 0$.

Also the equation of BC is $7x + 3y - 4 = 0$

On solving these equations, we get $B \left(-\frac{1}{2}, \frac{5}{2} \right)$

Now let the co-ordinates of the vertex D be
 (h, k) . Since the middle points of AC and BD
 are same

$$\therefore \frac{1}{2} \left(h - \frac{1}{2} \right) = \frac{1}{2} (3 + 1) \Rightarrow h = \frac{9}{2}$$

$$\Rightarrow \frac{1}{2} \left(k + \frac{5}{2} \right) = \frac{1}{2} (4 - 1)$$

$$\Rightarrow k = \frac{1}{2}$$

$$\text{Hence, } D = \left(\frac{9}{2}, \frac{1}{2} \right)$$



$$36. \begin{vmatrix} p-q & q-r & r-p \\ q-r & r-p & p-q \\ r-p & p-q & q-r \end{vmatrix} = \begin{vmatrix} 0 & q-r & r-p \\ 0 & r-p & p-q \\ 0 & p-q & q-r \end{vmatrix} = 0$$

....[By $C_1 \rightarrow C_1 + C_2 + C_3$]

Hence, the lines are concurrent.

37. Given lines are $3x + 4y = 5$, $5x + 4y = 4$,
 $\lambda x + 4y = 6$. These lines meet at a point if the
point of intersection of first two lines lies on
the third line.

From $3x + 4y = 5$ and $5x + 4y = 4$

$$\text{We get } x = \frac{-1}{2}, \quad y = \frac{13}{8}$$

This lies on $\lambda x + 4y = 6$, if $\lambda \left(-\frac{1}{2} \right) + 4 \left(\frac{13}{8} \right) = 6$

$$\Rightarrow \lambda = 1$$

38. If the given lines are concurrent, then

$$\begin{vmatrix} a & 1 & 1 \\ 1 & b & 1 \\ 1 & 1 & c \end{vmatrix} = 0 \Rightarrow \begin{vmatrix} a & 1-a & 1-a \\ 1 & b-1 & 0 \\ 1 & 0 & c-1 \end{vmatrix} = 0$$

[By $C_2 \rightarrow C_2 - C_1$ and $C_3 \rightarrow C_3 - C_1$]

$$\Rightarrow a(b-1)(c-1) - (b-1)(1-a) - (c-1)(1-a) = 0$$

$$\Rightarrow \frac{a}{1-a} + \frac{1}{1-b} + \frac{1}{1-c} = 0$$

....[Divide by $(1-a)(1-b)(1-c)$]

$$\Rightarrow \frac{1}{1-a} + \frac{1}{1-b} + \frac{1}{1-c} = 1$$

39. From option (B),

$$\begin{vmatrix} 1 & 2 & -10 \\ 2 & 1 & 5 \\ 5 & 4 & 0 \end{vmatrix} = 1(0 - 20) - 2(-25) - 10(3) = 0$$

Hence, option (B) is the correct answer.

40. From option (B), we get

$$\begin{vmatrix} 3 & 4 & 6 \\ 6 & 5 & 9 \\ 3 & 3 & 5 \end{vmatrix} = 3(25 - 27) - 4(3) + 6(3) = 0$$

41. The three lines are concurrent, if

$$\begin{vmatrix} 1 & 2 & -9 \\ 3 & 5 & -5 \\ a & b & -1 \end{vmatrix} = 0$$

$$\Rightarrow 35a - 22b + 1 = 0$$

which is true if the line $35x - 22y + 1 = 0$
passes through (a, b) .

42. By the given condition of $a + b + c = 0$, the
three lines reduce to

$$x - y = \frac{p}{a} \quad \text{or} \quad \frac{p}{b} \quad \text{or} \quad \frac{p}{c} \quad (p \neq 0).$$

All these lines are parallel. Hence, they do not
intersect in finite plane.

43. The point of intersection of the lines is $(1, 1)$.
and slope of the line $2y - 3x + 2 = 0$ is $\frac{3}{2}$

$$\text{Hence, the equation is } y - 1 = \frac{3}{2}(x - 1)$$

$$\Rightarrow 3x - 2y = 1$$

44. The intersection point of lines $x - 2y = 1$ and
 $x + 3y = 2$ is $\left(\frac{7}{5}, \frac{1}{5} \right)$ and the slope of required

$$\text{line} = -\frac{3}{4}$$

\therefore Equation of required line is

$$y - \frac{1}{5} = \frac{-3}{4} \left(x - \frac{7}{5} \right)$$

$$\Rightarrow \frac{3x}{4} + y = \frac{21}{20} + \frac{1}{5} \Rightarrow 3x + 4y = 5$$

$$\Rightarrow 3x + 4y - 5 = 0$$

45. The point of intersection of $5x - 6y - 1 = 0$
and $3x + 2y + 5 = 0$ is $(-1, -1)$.

Now the line perpendicular to

$3x - 5y + 11 = 0$ is $5x + 3y + k = 0$, but it
passes through $(-1, -1)$

$$\Rightarrow -5 - 3 + k = 0 \Rightarrow k = 8$$

Hence, required line is $5x + 3y + 8 = 0$.

46. Equation of line passing through point of
intersection of $x + 2y + 3 = 0$ and

$$3x + 4y + 7 = 0 \text{ is}$$

$$(x + 2y + 3) + k(3x + 4y + 7) = 0$$

$$\Rightarrow (1 + 3k)x + (2 + 4k)y + 3 + 7k = 0 \quad \dots(i)$$

$$\text{Slope of equation (i) is } m_1 = \frac{-(1 + 3k)}{2 + 4k}$$

$$\text{and slope of given line is } m_2 = \frac{-1}{-1} = 1 \quad \dots(ii)$$

Since (i) and (ii) represent perpendicular lines.

$$\therefore m_1 m_2 = -1$$

$$\therefore \frac{-(1 + 3k)}{(2 + 4k)} \times 1 = -1$$

\therefore equation of required line is

$$(x + 2y + 3) - 1(3x + 4y + 7) = 0$$

$$\Rightarrow x + y + 2 = 0$$



47. Equation of line passing through point of intersection of $u = 0$ and $v = 0$ is $u + kv = 0$

$$\therefore (x + 2y + 5) + k(3x + 4y + 1) = 0$$

It is passing through $(3, 2)$

$$\therefore (3 + 2 \times 2 + 5) + k(3 \times 3 + 4 \times 2 + 1) = 0$$

$$\therefore k = -\frac{2}{3}$$

\therefore equation of line will be

$$(x + 2y + 5) - \frac{2}{3}(3x + 4y + 1) = 0$$

$$\Rightarrow 3x + 2y - 13 = 0$$

48. Equation of line through the point of intersection of lines $2x + 3y + 1 = 0$ and

$3x - 5y - 5 = 0$ is given by

$$(2 + 3k)x + (3 - 5k)y + (1 - 5k) = 0$$

Slope of line is given by

$$\tan 45^\circ = -\frac{(2 + 3k)}{3 - 5k}$$

$$\Rightarrow k = \frac{5}{2}$$

\therefore Equation of line is $19x - 19y - 23 = 0$

49. Required line should be

$$(3x - y + 2) + \lambda(5x - 2y + 7) = 0 \quad \dots(i)$$

$$\Rightarrow (3 + 5\lambda)x - (2\lambda + 1)y + (2 + 7\lambda) = 0$$

$$\Rightarrow y = \frac{3 + 5\lambda}{2\lambda + 1}x + \frac{2 + 7\lambda}{2\lambda + 1} \quad \dots(ii)$$

As the equation (ii), has infinite slope,

$$2\lambda + 1 = 0$$

$$\Rightarrow \lambda = -1/2$$

Putting $\lambda = -1/2$ in equation (i) we have

$$(3x - y + 2) + (-1/2)(5x - 2y + 7) = 0$$

$$\Rightarrow x = 3$$

Alternate Method:

The point of intersection of $3x - y + 2 = 0$ and $5x - 2y + 7 = 0$ is $(3, 11)$

....[By solving equations simultaneously]


The required line has infinite slope (i.e. parallel to Y - axis) and passes through $(3, 11)$.

$\Rightarrow x = 3$ is required equation.

50. Equation of AD is

$$(x + y - 6) + k(x + 2y - 5) = 0$$

$$\Rightarrow (1 + k)x + (1 + 2k)y - (6 + 5k) = 0 \quad \dots(i)$$



$$\therefore \text{Slope of AD} = m_1 = \frac{-(1+k)}{(1+2k)}$$

and Slope of BC = $m_2 = -2$

$$\therefore m_1 m_2 = -1 \quad \dots[\because AD \perp BC]$$

$$\therefore k = -\frac{3}{4}$$

\therefore From (i), equation of AD is $x - 2y = 9$ (ii)

Similarly, equation of BE is $2x - y = -12$ (iii)

By solving equation (ii) and (iii), we get $x = -11, y = -10$

$$\therefore H \equiv (-11, -10)$$

51. Lengths of perpendicular from $(0,0)$ on the given lines are each equal to 2.

$$52. p_1 \cdot p_2 = \left(\frac{b\sqrt{a^2 - b^2} \cos\theta + 0 - ab}{\sqrt{b^2 \cos^2\theta + a^2 \sin^2\theta}} \right) \times \left(\frac{-b\sqrt{a^2 - b^2} \cos\theta - ab}{\sqrt{b^2 \cos^2\theta + a^2 \sin^2\theta}} \right)$$

$$= \frac{-[b^2(a^2 - b^2)\cos^2\theta - a^2b^2]}{(b^2 \cos^2\theta + a^2 \sin^2\theta)}$$

$$= \frac{b^2[a^2 - a^2 \cos^2\theta + b^2 \cos^2\theta]}{b^2 \cos^2\theta + a^2 \sin^2\theta}$$

$$= \frac{b^2[a^2 \sin^2\theta + b^2 \cos^2\theta]}{b^2 \cos^2\theta + a^2 \sin^2\theta} = b^2$$

$$53. \text{ Here, } p = \left| \frac{-k}{\sqrt{\sec^2 \alpha + \operatorname{cosec}^2 \alpha}} \right|$$

$$\text{ and } p' = \left| \frac{-k \cos 2\alpha}{\sqrt{\cos^2 \alpha + \sin^2 \alpha}} \right|$$

$$\therefore 4p^2 + p'^2 = \frac{4k^2}{\sec^2 \alpha + \operatorname{cosec}^2 \alpha} + \frac{k^2(\cos^2 \alpha - \sin^2 \alpha)^2}{1}$$



$$= 4k^2 \sin^2 \alpha \cos^2 \alpha + k^2 (\cos^4 \alpha + \sin^4 \alpha) - 2k^2 \cos^2 \alpha \sin^2 \alpha$$

$$= k^2 (\sin^2 \alpha + \cos^2 \alpha)^2$$

$$= k^2$$

54. Let the point be (h, k) , then $h + k = 4 \dots (i)$ and

$$1 = \left| \frac{4h + 3k - 10}{\sqrt{4^2 + 3^2}} \right|$$

$$\Rightarrow 4h + 3k = 15 \quad \dots (ii)$$

$$\text{and } 4h + 3k = 5 \quad \dots (iii)$$

On solving (i) and (ii), and (i) and (iii), we get the required points $(3, 1)$ and $(-7, 11)$.

55. $|AD| = \left| \frac{2 - 2 - 1}{\sqrt{1^2 + 2^2}} \right|$

$$= \frac{1}{\sqrt{5}}$$

$$\tan 60^\circ = \frac{AD}{BD}$$

$$\Rightarrow \sqrt{3} = \frac{1/\sqrt{5}}{BD}$$

$$\Rightarrow BD = \frac{1}{\sqrt{15}}$$

$\therefore BC = 2BD = 2/\sqrt{15}$

56. Equation of any line through $(0, a)$ is $y - a = m(x - 0)$ or $mx - y + a = 0 \dots (i)$

If the length of perpendicular from $(2a, 2a)$ to the line (i) is 'a', then $a = \pm \frac{m(2a) - 2a + a}{\sqrt{m^2 + 1}}$

$$\Rightarrow m = 0, \frac{4}{3}$$

Hence, the required equations of lines are $y - a = 0$, $4x - 3y + 3a = 0$

57. The equation of lines passing through $(1, 0)$ is given by $y = m(x - 1)$.

Its distance from origin is $\frac{\sqrt{3}}{2}$.

$$\Rightarrow \left| \frac{-m}{\sqrt{1+m^2}} \right| = \frac{\sqrt{3}}{2} \Rightarrow m = \pm\sqrt{3}$$

Hence, the lines are $\sqrt{3}x + y - \sqrt{3} = 0$ and $\sqrt{3}x - y - \sqrt{3} = 0$

58. Point of intersection is $(2, 3)$.
Therefore, the equation of line passing through $(2, 3)$ is $y - 3 = m(x - 2)$

$$\text{or } mx - y - (2m - 3) = 0$$

According to the condition,

$$\left| \frac{3m - 2 - (2m - 3)}{\sqrt{1 + m^2}} \right| = \frac{7}{5} \Rightarrow m = \frac{3}{4}, \frac{4}{3}$$

Hence, the equations are $3x - 4y + 6 = 0$ and $4x - 3y + 1 = 0$.

59. Slope $= -\sqrt{3}$


\therefore Line is $y = -\sqrt{3}x + c$

$$\Rightarrow \sqrt{3}x + y = c$$

$$\text{Now } \frac{c}{2} = |4|$$

$$\Rightarrow c = \pm 8$$

$$\Rightarrow x\sqrt{3} + y = \pm 8$$



60. $d = \left| \frac{8 - 3}{\sqrt{(3)^2 + (4)^2}} \right| = 1$

61. If the given lines represent the same line, then the length of the perpendiculars from the origin to the lines are equal, so that

$$\frac{c}{\sqrt{1+m^2}} = \frac{p}{\sqrt{\cos^2 \alpha + \sin^2 \alpha}}$$

$$\Rightarrow c = p\sqrt{1+m^2}$$

62. Lines $3x + 4y + 2 = 0$ and $3x + 4y + 5 = 0$ are on the same side of the origin. The distance

$$\text{between these lines is } d_1 = \left| \frac{2 - 5}{\sqrt{3^2 + 4^2}} \right| = \frac{3}{5}.$$

Lines $3x + 4y + 2 = 0$ and $3x + 4y - 5 = 0$ are on the opposite sides of the origin. The distance


$$\text{between these lines is } d_2 = \left| \frac{2 + 5}{\sqrt{3^2 + 4^2}} \right| = \frac{7}{5}.$$

Thus, $3x + 4y + 2 = 0$ divides the distance between $3x + 4y + 5 = 0$ and $3x + 4y - 5 = 0$ in the ratio $d_1 : d_2$ i.e., $3 : 7$.

63. Since, the distance between the parallel lines $x + my + n = 0$ and $x + my + n' = 0$ is same as the distance between parallel lines $mx + y + n = 0$ and $mx + y + n' = 0$. Therefore, the parallelogram is a rhombus. Since, the diagonals of a rhombus are at right angles, therefore the required angle is $\frac{\pi}{2}$.



64. Line AB will pass through (0, a) and (2a, k)




But as we are given $AB = AC$

$$\Rightarrow k = \sqrt{4a^2 + (k - a)^2} \Rightarrow k = \frac{5a}{2}$$

Hence, the required equation is $3x - 4y + 4a = 0$

65. B(0, b)



By the section formula, we get $a = -\frac{32}{3}$ and

$$b = \frac{24}{5}$$

Hence, the required equation is given by

$$\frac{x}{-(32/3)} + \frac{y}{(24/5)} = 1$$

$$\Rightarrow 9x - 20y + 96 = 0$$

66. It is given that the lines $ax + 2y + 1 = 0$, $bx + 3y + 1 = 0$ and $cx + 4y + 1 = 0$ are

$$\text{concurrent, therefore } \begin{vmatrix} a & 2 & 1 \\ b & 3 & 1 \\ c & 4 & 1 \end{vmatrix} = 0$$

$$\Rightarrow -a + 2b - c = 0 \Rightarrow 2b = a + c$$

$\Rightarrow a, b, c$ are in A. P.

67. The two lines will be identical if there exists some real number k such that

$$b^3 - c^3 = k(b - c), \quad c^3 - a^3 = k(c - a),$$

$$a^3 - b^3 = k(a - b)$$

$$\Rightarrow b - c = 0 \text{ or } b^2 + c^2 + bc = k$$

$$\Rightarrow c - a = 0 \text{ or } c^2 + a^2 + ac = k$$

$$\Rightarrow a - b = 0 \text{ or } a^2 + b^2 + ab = k$$

$$\Rightarrow b = c, \quad c = a, \quad a = b$$

$$\text{or } b^2 + c^2 + bc = c^2 + a^2 + ca$$

$$\Rightarrow b^2 - a^2 = c(a - b)$$

$$\Rightarrow b = a \text{ or } a + b + c = 0$$

$$68. \quad 2p = \frac{|0 + 0 - 1|}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2}}}$$

$$\Rightarrow \frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{4p^2} \Rightarrow \frac{1}{a^2} + \frac{1}{b^2} = \frac{2}{8p^2}$$

$$\Rightarrow a^2, 8p^2, b^2 \text{ are in H.P.}$$



Competitive Thinking

- Since, the line makes an angle of measure 30° with Y-axis. Therefore, the line will make an angle of measure 60° or -60° with X-axis.
 \therefore Slope of line = $\tan 60^\circ$ or $\tan(-60^\circ)$
 $= \sqrt{3}$ or $-\sqrt{3} = \pm \sqrt{3}$
- Here, the straight line is parallel to X-axis.
 So, the slope of such a line = 0.
- $m_1 = \frac{6+4}{-2-3} = \frac{10}{-5} = -2$ and $m_2 = \frac{-18-6}{9-(-3)} = -2$
 Hence, the lines are parallel.
- Midpoint of the line joining the points (4, -5) and (-2, 9) is $\left(\frac{4-2}{2}, \frac{-5+9}{2}\right)$ i.e., (1, 2)
 \therefore Inclination of straight line passing through point (-3, 6) and midpoint (1, 2) is
 $m = \frac{2-6}{1+3} \Rightarrow \tan \theta = -1$
 $\Rightarrow \theta = \frac{3\pi}{4}$
- The required equation of line passing through (a, b) and having gradient $m = \frac{-b}{a}$, is
 $(y - b) = \frac{-b}{a} (x - a)$
 i.e. $\frac{x}{a} + \frac{y}{b} = 2$
- The required equation of line passing through (-2, 3) and gradient $m = \frac{3}{4}$, is
 $y - 3 = \frac{3}{4} [x - (-2)]$
 i.e. $3x - 4y + 18 = 0$



7. Slope of line passing through (1, 0) and (-4, 1) = $\frac{1-0}{-4-1} = \frac{-1}{5}$

Slope of line perpendicular to the given line is $m = 5$

Equation of line passing through (-3, 5) and having slope 5 is

$$y - 5 = 5(x + 3)$$

$$\Rightarrow 5x - y + 20 = 0$$


8. Midpoint $\equiv (2, 7)$
Slope of perpendicular = -6
 \therefore the required equation is $y - 7 = -6(x - 2)$
 $\Rightarrow 6x + y - 19 = 0$

9. Midpoint of given line segment $\equiv (2, -1)$
Now, slope of the line segment = $\frac{-8}{8} = -1$
Slope of the required line segment is 1
 \therefore the required equation of line is $y + 1 = 1(x - 2)$
 $\Rightarrow x - y = 3$


10. Midpoint $\equiv (3, 2)$.
 \therefore the required equation is $y - 2 = 2(x - 3)$
 $\Rightarrow 2x - y - 4 = 0$

11. The required diagonal passes through the midpoint of AB and is perpendicular to AB. So, its equation is $y - 2 = -3(x - 2)$ or $y + 3x - 8 = 0$.

12. Co-ordinates of the vertices of the square are A(0, 0), B(0, 1), C(1, 1) and D(1, 0).



Now, the equation of AC is $y = x$ and of BD is $y - 1 = -\frac{1}{1}(x - 0) \Rightarrow x + y = 1$

13. 

From figure, diagonal BD is passing through origin, therefore its equation is given by

$$\left(y - \frac{3}{5}\right) = \frac{-(3/5)}{-(2/5)} \left(x - \frac{2}{5}\right)$$


$$\Rightarrow 3x - 2y = 0$$

14. Since, the required line will be a line passing through A and B.

$$\therefore \frac{y - 6}{6 - (-4)} = \frac{x - 1}{1 - 3}$$

$$\Rightarrow 10x - 10 = -2y + 12 \Rightarrow 5x + y - 11 = 0$$

15. Since, equation of diagonal $11x + 7y = 9$ does not pass through origin, so it cannot be the equation of the diagonal OB. Thus, on solving the equation AC with the equations OA and OC, we get $A\left(\frac{5}{3}, -\frac{4}{3}\right)$ and $C\left(\frac{-2}{3}, \frac{7}{3}\right)$



Therefore, the midpoint of AC is $\left(\frac{1}{2}, \frac{1}{2}\right)$.

Hence, the equation of OB is $y = x$ i.e. $x - y = 0$.

16. Point of intersection of $x - y + 1 = 0$ and $7x - y - 5 = 0$ is (1, 2)
Equation of diagonal passing through (-1, -2) and (1, 2) is

$$y + 2 = \frac{4}{2}(x + 1)$$

$$\Rightarrow 2x + 2 = y + 2$$


$$\Rightarrow 2x - y = 0$$

Equation of another diagonal passing through (-1, -2) and having slope $-\frac{1}{2}$ is

$$y + 2 = \frac{-1}{2}(x + 1)$$

$$\Rightarrow 2y + 4 = -x - 1$$

$$\Rightarrow x + 2y = -5$$



Point of intersection of $7x - y - 5 = 0$ and $x + 2y = -5$ is $\left(\frac{1}{3}, \frac{-8}{3}\right)$

- \therefore Answer is option (C)



$$17. \text{ Slope} = \frac{(2-1)}{1-\left(-\frac{1}{2}\right)} = \frac{1}{\left(\frac{3}{2}\right)} = \frac{2}{3}$$

So, equation of the line is $y - 2 = \frac{2}{3}(x - 1)$

$$\Rightarrow y = \frac{2}{3}x + \frac{4}{3}$$

Putting $y = 0$, to find x -intercept, $\frac{2}{3}x + \frac{4}{3} = 0$

$$\Rightarrow x = -2$$

$$\therefore x\text{-intercept} = -2$$

$$18. \text{ Let the equation be } \frac{x}{a} + \frac{y}{-a} = 1.$$

$$\Rightarrow x - y = a \quad \dots(i)$$

But, it passes through $(-3, 2)$

$$\therefore a = -3 - 2 = -5$$

Putting the value of a in (i), we get

$$x - y + 5 = 0$$

$$19. \text{ Let the equation of the line be } \frac{x}{a} + \frac{y}{b} = 1.$$

Given, $a = b$

So, equation of line is $x + y = a$

Since, this line passes through $(2, 4)$.

$$\therefore 2 + 4 = a$$

$$\Rightarrow a = 6$$

$$\therefore \text{ the required equation of line is } x + y = 6$$

i.e., $x + y - 6 = 0$

$$20. \text{ Here, } a + b = -1$$

$$\therefore \text{ required line is } \frac{x}{a} - \frac{y}{1+a} = 1 \quad \dots(i)$$

Since, line (i) passes through $(4, 3)$.

$$\therefore \frac{4}{a} - \frac{3}{1+a} = 1$$

$$\Rightarrow 4 + 4a - 3a = a + a^2$$


$$\Rightarrow a^2 = 4$$

$$\Rightarrow a = \pm 2$$

$$\therefore \text{ the required lines are } \frac{x}{2} - \frac{y}{3} = 1 \quad \text{and}$$

$$\frac{x}{-2} + \frac{y}{1} = 1.$$

$$21. \text{ Let the equation of PQ be } \frac{x}{h} + \frac{y}{k} = 1.$$



Since, the line passes through the fixed point $(2, 3)$.

$$\Rightarrow \frac{2}{h} + \frac{3}{k} = 1$$

$$\Rightarrow \text{Locus of } R(h, k) \text{ is } \frac{2}{x} + \frac{3}{y} = 1$$

$$\Rightarrow 3x + 2y = xy$$

$$22. \text{ Equation of the line has its intercepts on the X-axis and Y-axis in the ratio } 2 : 1 \text{ i.e., } 2a \text{ and } a$$

$$\therefore \frac{x}{2a} + \frac{y}{a} = 1 \Rightarrow x + 2y = 2a \quad \dots(i)$$

Line (i) also passes through midpoint of $(3, -4)$ and $(5, 2)$ i.e., $(4, -1)$

$$\therefore 4 + 2(-1) = 2a \Rightarrow a = 1$$

Hence, the equation of required line is

$$x + 2y = 2$$

$$23. \text{ Let the points of the required line on X-axis and Y-axis be } A(a, 0) \text{ and } B(0, b) \text{ respectively.}$$

Since, $\left(\frac{3}{2}, \frac{5}{2}\right)$ is midpoint of AB .

$$\therefore \frac{a+0}{2} = \frac{3}{2} \quad \text{and} \quad \frac{0+b}{2} = \frac{5}{2} \Rightarrow a = 3 \text{ and } b = 5$$

$$\therefore \text{ the equation of line is } \frac{x}{3} + \frac{y}{5} = 1$$

$$\Rightarrow 5x + 3y - 15 = 0$$


$$24. \text{ The required equation of line is}$$

$$\frac{x}{6} + \frac{y}{8} = 1 \Rightarrow 4x + 3y = 24$$

$$25. \text{ Let } P\left(\alpha = \frac{a}{2}, \beta = \frac{b}{2}\right) \text{ be the midpoint of the line joining } (a, 0) \text{ and } (0, b).$$

$$\therefore \alpha = \frac{a}{2} \Rightarrow a = 2\alpha \quad \dots(i)$$

$$\text{and } \beta = \frac{b}{2} \Rightarrow b = 2\beta \quad \dots(ii)$$



∴ Equation of a straight line cutting off intercepts a and b on X-axis and Y-axis respectively is

$$\frac{x}{a} + \frac{y}{b} = 1$$

$$\Rightarrow \frac{x}{2\alpha} + \frac{y}{2\beta} = 1 \quad \dots[\text{From (i) and (ii)}]$$

$$\Rightarrow \frac{x}{\alpha} + \frac{y}{\beta} = 2$$


26. $\frac{a+0}{2} = 4 \Rightarrow a = 8$

and $\frac{b+0}{2} = -3 \Rightarrow b = -6$


∴ the required equation of

the line is $\frac{x}{8} + \frac{y}{-6} = 1$

$$\Rightarrow \frac{3x-4y}{24} = 1 \Rightarrow 3x-4y = 24$$



27.



Point P divides AB in the ratio 2:3

$$\therefore \left(\frac{1}{2}, \frac{1}{3}\right) = \left(\frac{0+2a}{5}, \frac{3b+0}{5}\right)$$


$$\Rightarrow a = \frac{5}{4} \text{ and } b = \frac{5}{9}$$

∴ equation of line AB is

$$\frac{x}{a} + \frac{y}{b} = 1$$

i.e., $4x + 9y = 5$

28.



Point P divides AB in the ratio 3:2

$$\therefore (2, -1) = \left(\frac{0+3a}{5}, \frac{2b+0}{5}\right)$$

$$\Rightarrow a = 5 \text{ and } b = \frac{-5}{3}$$

∴ Equation of the line AB is

$$\frac{x}{a} + \frac{y}{b} = 1$$

$$\Rightarrow x - 3y - 5 = 0$$

29. Since, $px - qy = r$ intersects at X-axis and Y-axis.

$$\therefore a = \frac{r}{p} \text{ and } b = -\frac{r}{q}$$

$$\therefore a + b = \frac{r}{p} - \frac{r}{q} = r \left(\frac{q-p}{pq}\right)$$

30. Any line through the middle point $M(1, 5)$ of the intercept AB may be taken as

$$\frac{x-1}{\cos\theta} = \frac{y-5}{\sin\theta} = r \quad \dots(i)$$

Since, the points A and B are equidistant from M and on the opposite sides of it.

Therefore, if the co-ordinates of A are obtained by putting $r = d$ in (i), then the co-ordinates of B are given by putting $r = -d$.

Now, the point $A(1 + d \cos\theta, 5 + d \sin\theta)$ lies on the line $5x - y - 4 = 0$ and

point $B(1 - d \cos\theta, 5 - d \sin\theta)$ lies on the line $3x + 4y - 4 = 0$.

$$\therefore 5(1 + d \cos\theta) - (5 + d \sin\theta) - 4 = 0$$


$$\text{and } 3(1 - d \cos\theta) + 4(5 - d \sin\theta) - 4 = 0$$

Eliminating 'd', we get $\frac{\cos\theta}{35} = \frac{\sin\theta}{83}$

Hence, the required line is $\frac{x-1}{35} = \frac{y-5}{83}$ or

$$83x - 35y + 92 = 0.$$



31. Since, $m_1 m_2 = (2) \left(-\frac{1}{2}\right) = -1$
 \therefore the lines are perpendicular.
32. Here, $m_1 = -1, m_2 = -\frac{1}{k}$.
 For orthogonal lines,
 $m_1 m_2 = -1 \Rightarrow \frac{1}{k} = -1 \Rightarrow k = -1$
33. Here, $m_1 = \frac{-2}{3}$ and $m_2 = \frac{-1}{k}$
 for perpendicular lines
 $m_1 m_2 = -1$
 $\therefore \frac{-2}{3} \times \frac{-1}{k} = -1$
 $\Rightarrow k = \frac{-2}{3}$
34. $m_1 m_2 = -1$
 $\Rightarrow \left(\frac{k-3}{2-4}\right) (2) = -1 \Rightarrow 2k - 6 = 2 \Rightarrow k = 4$
35. 
 From figure,
 $\left(\frac{b/2}{a/2}\right) \left(\frac{b}{-a/2}\right) = -1$
 $\Rightarrow a^2 = 2b^2 \Rightarrow a = \pm\sqrt{2}b$
36. Since, the point $(-4, 5)$ does not lie on the diagonal $7x - y + 8 = 0$, so point will lie on the other diagonal.
 Also, diagonals are perpendicular.
 \therefore Slope of other diagonal = $\frac{-1}{7}$
 \therefore equation of the other diagonal is
 $y - 5 = -\frac{1}{7}(x + 4) \Rightarrow 7y + x = 31$
37. The equation of lines in intercept form are
 $\frac{x}{-8/a} + \frac{y}{-8/b} = 1$
 $\frac{x}{-3} + \frac{y}{2} = 1$

According to the given condition,

$$-\frac{8}{a} = -(-3) \text{ and } -\frac{8}{b} = -2$$

$$\Rightarrow a = -\frac{8}{3} \text{ and } b = 4$$

38. The equation of a line perpendicular to $x - y = 0$ is $-x - y + c = 0$... (i)
 Since, the line passes through $(3, 2)$.
 $\therefore -3 - 2 + c = 0$
 $\therefore c = 5$
 Putting $c = 5$ in (i), we get
 $x + y = 5$
39. The equation of a line perpendicular to $x + y + 1 = 0$ is $x - y + \lambda = 0$.
 Since, the line passes through the point $(1, 2)$.
 $\therefore 1 - 2 + \lambda = 0$
 $\Rightarrow \lambda = 1$
 Hence, required equation of line is
 $y - x - 1 = 0$
40. Slope of $y = 3x - 1$ is 3
 \therefore Slope of line perpendicular to the above line is
 $m = \frac{-1}{3}$
 Equation of line passing through $(1, 2)$ and having slope $(m) = \frac{-1}{3}$ is
 $(y - 2) = \frac{-1}{3}(x - 1)$
 $\Rightarrow 3y - 6 = -x + 1$
 $\Rightarrow x + 3y - 7 = 0$
41. The required equation passing through $(-1, 1)$ and having gradient $\frac{3}{2}$ is
 $y - 1 = \frac{3}{2}(x + 1) \Rightarrow 2(y - 1) = 3(x + 1)$
42. $5x - 6y - 1 = 0$... (i)
 $3x + 2y + 5 = 0$... (ii)
 On solving (i) and (ii), we get $x = -1, y = -1$
 Slope of line $3x - 5y + 11 = 0$ is $\frac{3}{5}$.
 Slope of line perpendicular to above line = $\frac{-5}{3}$
 \therefore Equation of line passing through $(-1, -1)$ and having slope $-\frac{5}{3}$ is



$$(y + 1) = -\frac{5}{3}(x + 1)$$

$$\Rightarrow 3y + 3 = -5x - 5$$

$$\Rightarrow 5x + 3y + 8 = 0$$

43. The given line is $bx - ay = ab$ (i)

It cuts X-axis at $(a, 0)$.

The equation of a line perpendicular to (i) is $ax + by = k$.

Since, the line passes through $(a, 0) \Rightarrow k = a^2$
Hence, required equation of line is $ax + by = a^2$


i.e., $\frac{x}{b} + \frac{y}{a} = \frac{a}{b}$

44. The equation of a line passing through $(2, 2)$ and perpendicular to $3x + y = 3$ is

$$y - 2 = \frac{1}{3}(x - 2) \text{ or } x - 3y + 4 = 0.$$

Putting $x = 0$ in this equation, we get $y = \frac{4}{3}$

\therefore y - intercept = $\frac{4}{3}$



In $\triangle ABC$;

$$\text{slope of BC} = \frac{0 - (-1)}{4 - 2} = \frac{1}{2}$$

$$\text{slope of AC} = \frac{0 - 3}{4 - (-2)} = \frac{-3}{6} = \frac{-1}{2}$$

Since, $AP \perp BC$ and $BQ \perp AC$,

\therefore slope of AP = -2,

slope of BQ = 2

\therefore Equation of AP is $2x + y + 1 = 0$ and equation of BQ is $2x - y - 5 = 0$

Solving the above equations, we get orthocentre, $O = (1, -3)$


Also, centroid of the triangle,

$$C = \left(\frac{-2 + 2 + 4}{3}, \frac{3 - 1 + 0}{3} \right)$$

i.e. $C = \left(\frac{4}{3}, \frac{2}{3} \right)$

\therefore Equation of line OC is $11x - y - 14 = 0$

46. Line; $x + 2y + 3 = 0$ intersects the co-ordinate axes at A $(-3, 0)$ and



Line: $x + 2y + 8 = 0$ intersects the coordinate axes at $C(-8, 0)$ and $D(0, -4)$

Since the required line divides the distance between the two lines in the ratio 1 : 2

- \therefore $(h, 0)$ divides the distance between A $(-3, 0)$ and $C(-8, 0)$ in the ratio 1 : 2

$$\therefore (h, 0) = \left(\frac{1(-8) + 2(-3)}{3}, 0 \right)$$

$$\therefore (h, 0) = \left(\frac{-14}{3}, 0 \right)$$

- \therefore the required equation of line passing through $\left(\frac{-14}{3}, 0 \right)$ and having gradient $m = \frac{-1}{2}$, is

$$(y - 0) = \frac{-1}{2} \left(x + \frac{14}{3} \right)$$

$$\therefore -3x - 6y = 14$$

Writing in normal form,

$$\frac{-3x}{\sqrt{45}} - \frac{6y}{\sqrt{45}} = \frac{14}{\sqrt{45}}$$

$$\text{i.e. } x \cos a + y \sin a = \frac{14}{\sqrt{45}}$$

$$\text{where, } \cos a = \frac{-3}{\sqrt{45}}, \sin a = \frac{-6}{\sqrt{45}}$$

$$\therefore a = \pi + \tan^{-1} \frac{\frac{-6}{\sqrt{45}}}{\frac{-3}{\sqrt{45}}}$$

$$\text{i.e. } a = \pi + \tan^{-1} 2$$



47. Here, $m_1 = -\cot \alpha$, $m_2 = \tan \beta$

$$\therefore \tan \theta = \left| \frac{-\cot \alpha - \tan \beta}{1 - \cot \alpha \tan \beta} \right|$$

$$\therefore \tan \theta = -\cot(\alpha - \beta)$$

$$\therefore \theta = \frac{\pi}{2} - \beta + \alpha$$

48. The lines are $bx + ay - ab = 0$ and $bx - ay - ab = 0$.

Hence, the required angle is


$$\theta = \tan^{-1} \left| \frac{\frac{-b}{a} - \frac{b}{a}}{1 - \frac{b^2}{a^2}} \right| = \tan^{-1} \left| \frac{2ab}{b^2 - a^2} \right|$$

$$= 2 \tan^{-1} \frac{b}{a} \quad \dots \left[\because 2 \tan^{-1} \frac{y}{x} = \tan^{-1} \left| \frac{2xy}{y^2 - x^2} \right| \right]$$

49. $\theta = 90^\circ - \tan^{-1} \left(\frac{1}{3} \right)$

$$\Rightarrow \tan \theta = \cot \left[\tan^{-1} \left(\frac{1}{3} \right) \right] = 3$$

$$\Rightarrow \theta = \tan^{-1}(3)$$



50. Given lines are $ax + by + c = 0$

and $x = \alpha t + \beta$, $y = \gamma t + \delta$

After eliminating t , we get

$$\gamma x - \alpha y + \alpha \delta - \gamma \beta = 0$$

For parallelism condition,

$$\frac{a}{\gamma} = \frac{b}{-\alpha} \Rightarrow a\alpha + b\gamma = 0$$

51. The given lines are perpendicular because

$$m_1 m_2 = (2) \left(\frac{-1}{2} \right) = -1$$

Hence, the angle between the two lines is 90° .

52. The slopes of the lines are $m_1 = \frac{-1}{2}$, $m_2 = 2$

$$\therefore m_1 m_2 = -1$$

So, the lines are perpendicular i.e., $\theta = 90^\circ$

53. Slopes of lines $\frac{x}{a} + \frac{y}{b} = 1$ and $\frac{x}{b'} + \frac{y}{a'} = 1$ are $\frac{-b}{a}$ and $\frac{-a'}{b'}$ respectively

\therefore Product of slopes is $\frac{a'b}{ab'}$

$$\text{But } \frac{1}{ab'} + \frac{1}{ba'} = 0$$

$$\Rightarrow ab' = -a'b$$

$$\Rightarrow \text{Product of slopes} = -1$$

Hence option (C)

54. The equation of a straight line passing through $(3, -2)$ is

$$y + 2 = m(x - 3) \quad \dots (i)$$

The slope of the line $\sqrt{3}x + y = 1$ is $-\sqrt{3}$

$$\text{So, } \tan 60^\circ = \pm \frac{m - (-\sqrt{3})}{1 + m(-\sqrt{3})}$$

On solving, we get

$$m = 0 \text{ or } \sqrt{3}$$

Putting the values of m in (i), the required equation of lines are $y + 2 = 0$ and

$$\sqrt{3}x - y - 2 - 3\sqrt{3} = 0$$

55. Here the lines are $x - 3 = 0$, $y - 4 = 0$ and $4x - 3y + a = 0$.

These will be concurrent, if

$$\begin{vmatrix} 1 & 0 & -3 \\ 0 & 1 & -4 \\ 4 & -3 & a \end{vmatrix} = 0 \Rightarrow a = 0$$

56. Lines are concurrent, if $\begin{vmatrix} 4 & 3 & -1 \\ 1 & -1 & 5 \\ b & 5 & -3 \end{vmatrix} = 0$

$$\Rightarrow 4(3 - 25) - 3(-3 - 5b) - 1(5 + b) = 0$$

$$\Rightarrow -88 + 9 + 15b - 5 - b = 0$$

$$\Rightarrow -84 + 14b = 0$$

$$\Rightarrow b = 6$$

57. Given lines are concurrent, if $\begin{vmatrix} 2 & 1 & -1 \\ a & 3 & -3 \\ 3 & 2 & -2 \end{vmatrix} = 0$

$$\Rightarrow - \begin{vmatrix} 2 & 1 & 1 \\ a & 3 & 3 \\ 3 & 2 & 2 \end{vmatrix} = 0$$

This is true for all values of a because C_2 and C_3 are identical.



58. Here, the given lines are

$$ax + by + c = 0$$

$$bx + cy + a = 0$$

$$cx + ay + b = 0$$

The lines will be concurrent, if
$$\begin{vmatrix} a & b & c \\ b & c & a \\ c & a & b \end{vmatrix} = 0$$

$$\Rightarrow a^3 + b^3 + c^3 - 3abc = 0$$

59. Dividing both sides of relation $3a + 2b + 4c = 0$

by 4, we get $\frac{3}{4}a + \frac{1}{2}b + c = 0$, which shows that for all values of a , b and c each member of the set of lines $ax + by + c = 0$ passes through the point $\left(\frac{3}{4}, \frac{1}{2}\right)$

60. Since, lines $x + 3y - 9 = 0$, $4x + by - 2 = 0$, and $2x - y - 4 = 0$ are concurrent

$$\begin{vmatrix} 1 & 3 & -9 \\ 4 & b & -2 \\ 2 & -1 & -4 \end{vmatrix} = 0$$

$$\therefore 1(-4b - 2) - 3(-16 + 4) - 9(-4 - 2b) = 0$$

$$\Rightarrow b = -5$$

\therefore the required line passes through $(-5, 0)$

Now, consider option (D) and $x + 3y - 9 = 0$, $4x - 5y - 2 = 0$

$$\therefore \begin{vmatrix} 1 & 3 & -9 \\ 4 & -5 & -2 \\ 1 & -4 & 5 \end{vmatrix} = 0$$

\therefore option (D) is correct

61. Required equation of line which is parallel to

$$x + 2y = 5 \text{ is } x + 2y + k = 0 \quad \dots(i)$$

Given equation of lines are

$$x + y = 2 \quad \dots(ii)$$

$$x - y = 0 \quad \dots(iii)$$

Adding (ii) and (iii), we get $2x = 2 \Rightarrow x = 1$

From (iii), we get $y = 1$

\therefore Point of intersection is $(1, 1)$.

Putting $x = 1$, $y = 1$ in (i), we get $k = -3$

\therefore the required equation of line is $x + 2y = 3$.

62. Point of intersection is $y = -\frac{21}{5}$ and $x = \frac{23}{5}$

$$\therefore 3x + 4y = \frac{3(23) + 4(-21)}{5} = \frac{69 - 84}{5} = -3$$

Hence, required line is $3x + 4y + 3 = 0$

63. The lines passing through the intersection of the lines $ax + 2by + 3b = 0$ and $bx - 2ay - 3a = 0$ is

$$ax + 2by + 3b + \lambda(bx - 2ay - 3a) = 0$$

$$\Rightarrow (a + b\lambda)x + (2b - 2a\lambda)y + 3b - 3\lambda a = 0 \dots(i)$$

Line (i) is parallel to X-axis,

$$a + b\lambda = 0 \Rightarrow \lambda = \frac{-a}{b}$$

Putting the value of λ in (i), we get

$$ax + 2by + 3b - \frac{a}{b}(bx - 2ay - 3a) = 0$$

$$\Rightarrow y\left(2b + \frac{2a^2}{b}\right) + 3b + \frac{3a^2}{b} = 0$$

$$\Rightarrow y\left(\frac{2b^2 + 2a^2}{b}\right) = -\left(\frac{3b^2 + 3a^2}{b}\right)$$

$$\Rightarrow y = \frac{-3(a^2 + b^2)}{2(b^2 + a^2)} = \frac{-3}{2}$$

So, it is $3/2$ unit below X-axis.


64. Point of intersection of the lines is $(3, -2)$

$$\text{Also, slope of perpendicular} = \frac{2}{7}$$

$$\text{Hence, the equation is } y + 2 = \frac{2}{7}(x - 3)$$

$$\Rightarrow 2x - 7y - 20 = 0$$

65. Slopes of the lines are 1 and -1



Since, the point of intersection is $(1, 1)$

Hence, the required equations are

$$y - 1 = \pm 1(x - 1)$$

66. The point of intersection of the lines

$$3x + y + 1 = 0 \text{ and } 2x - y + 3 = 0 \text{ are } \left(\frac{-4}{5}, \frac{7}{5}\right).$$

The equation of line which makes equal intercepts with the axes is $x + y = a$.

$$\therefore -\frac{4}{5} + \frac{7}{5} = a \Rightarrow a = \frac{3}{5}$$

\therefore the required equation of the line is

$$x + y - \frac{3}{5} = 0 \text{ i.e., } 5x + 5y - 3 = 0$$



67. $(a-2b)x + (a+3b)y + 3a + 4b = 0$
or $a(x+y+3) + b(-2x+3y+4) = 0$, which represents a family of straight lines through point of intersection of $x+y+3=0$ and $-2x+3y+4=0$ i.e., $(-1, -2)$.

68. Required distance = $\left| \frac{3(3) - 5(-4) - 26}{\sqrt{9+16}} \right| = \frac{3}{5}$

69. Since, $L(p, q) = \frac{ap + bq + c}{\sqrt{a^2 + b^2}}$

$$\text{and } L\left(\frac{2}{3}, \frac{1}{3}\right) + L\left(\frac{1}{3}, \frac{2}{3}\right) + L(2, 2) = 0$$

$$\therefore \frac{a\left(\frac{2}{3}\right) + b\left(\frac{1}{3}\right) + c}{\sqrt{a^2 + b^2}} + \frac{a\left(\frac{1}{3}\right) + b\left(\frac{2}{3}\right) + c}{\sqrt{a^2 + b^2}} + \frac{a(2) + b(2) + c}{\sqrt{a^2 + b^2}} = 0$$

$$\therefore \frac{3a + 3b + 3c}{\sqrt{a^2 + b^2}} = 0$$

$$\therefore a + b + c = 0 \quad \dots(i)$$

Comparing equation (i) with $ax + by + c = 0$, we get

$$a + b + c = ax + by + c$$

$$\text{i.e. } x = 1 \text{ and } y = 1$$

\therefore The line $ax + by + c = 0$ passes through the point $(1, 1)$.

70. The line is $4x - 3y - 12 = 0$.

$$\therefore \text{Required length} = \left| \frac{-12}{\sqrt{4^2 + (-3)^2}} \right| = \frac{12}{5} = 2\frac{2}{5}$$

71. Given, equation of line is

$$\frac{x \sin \alpha}{b} - \frac{y \cos \alpha}{a} - 1 = 0$$

\therefore perpendicular distance from origin

$$= \left| \frac{0 \cdot \frac{\sin \alpha}{b} - \frac{0 \cdot \cos \alpha}{a} - 1}{\sqrt{\frac{\sin^2 \alpha}{b^2} + \frac{\cos^2 \alpha}{a^2}}} \right| = \frac{|ab|}{\sqrt{a^2 \sin^2 \alpha + b^2 \cos^2 \alpha}}$$

72. Equation of the line is

$$y - 0 = \left(\frac{3-0}{-5} \right) (x-5)$$

$$\Rightarrow 3x + 5y - 15 = 0$$

$$\therefore d = \left| \frac{3(4) + 5(4) - 15}{\sqrt{3^2 + 5^2}} \right| = \frac{17}{\sqrt{34}} = \sqrt{\frac{17}{2}}$$

73. Distance of $(1, 1)$ from $3x + 4y + c = 0$ is

$$d = \left| \frac{3(1) + 4(1) + c}{\sqrt{9+16}} \right|$$

$$\Rightarrow \pm 7 = \frac{7+c}{5}$$

$$\Rightarrow c = -42, 28$$


74. Let the point be $(h, 0)$, then $a = \left| \frac{bh + 0 - ab}{\sqrt{a^2 + b^2}} \right|$

$$\Rightarrow bh = \pm a\sqrt{a^2 + b^2} + ab$$

$$\Rightarrow h = \frac{a}{b}(b \pm \sqrt{a^2 + b^2})$$

Hence, the points are $\left\{ \frac{a}{b}(b \pm \sqrt{a^2 + b^2}), 0 \right\}$.

75. From option (C),



$$BP = \sqrt{(5-3)^2 + (7+2)^2} = \sqrt{4+81} = \sqrt{85}$$

Hence, option (C) is correct.

76. $L_{12} \equiv x - 3y + 1 = 0$

$$L_{23} \equiv 2x + y - 12 = 0$$

$$L_{13} \equiv 3x - 2y - 4 = 0$$

Therefore, the required distances are

$$D_3 = \left| \frac{4 - 3 \times 4 + 1}{\sqrt{10}} \right| = \frac{7}{\sqrt{10}}$$

$$D_1 = \left| \frac{4 + 1 - 12}{\sqrt{5}} \right| = \frac{7}{\sqrt{5}}$$

$$D_2 = \left| \frac{3 \times 5 - 2 \times 2 - 4}{\sqrt{9+4}} \right| = \frac{7}{\sqrt{13}}$$

77. Let p be the length of the perpendicular from the vertex $(2, -1)$ to the base $x + y = 2$.

$$\text{Then, } p = \left| \frac{2 - 1 - 2}{\sqrt{1^2 + 1^2}} \right| = \frac{1}{\sqrt{2}}$$

If 'a' is the length of the side of triangle, then $p = a \sin 60^\circ$



$$\Rightarrow \frac{1}{\sqrt{2}} = \frac{a\sqrt{3}}{2}$$

$$\Rightarrow a = \sqrt{\frac{2}{3}}$$

$$78. \quad AD = \left| \frac{-2-2-1}{\sqrt{(2)^2 + (-1)^2}} \right| = \left| \frac{-5}{\sqrt{5}} \right| = \sqrt{5}$$

$$\text{Since, } \tan 60^\circ = \frac{AD}{BD}$$


$$\Rightarrow \sqrt{3} = \frac{\sqrt{5}}{BD}$$

$$\Rightarrow BD = \frac{\sqrt{5}}{\sqrt{3}}$$

$$\therefore BC = 2BD$$

$$= 2 \frac{\sqrt{5}}{\sqrt{3}} = \sqrt{\frac{20}{3}}$$

79.



lines $y + 2x = 1$ and $3y + 6x = 6$ are parallel to each other

$$\therefore d = \left| \frac{-1+2}{\sqrt{4+1}} \right|$$

$$d = \frac{1}{\sqrt{5}}$$

$$\text{Side of equilateral triangle} = \frac{2}{\sqrt{3}} \times d$$

$$= \frac{2}{\sqrt{3}} \times \frac{1}{\sqrt{5}} = \frac{2}{\sqrt{15}}$$

80. Point of intersection is (1, 2)

Therefore, the equation of line passing through (1, 2) is $(y - 2) = m(x - 1)$
or $mx - y + 2 - m = 0$

Since, the line is at distance of $\sqrt{5}$ from origin i.e. (0, 0),

$$\left| \frac{(0)m - (0) + 2 - m}{\sqrt{m^2 + 1}} \right| = \sqrt{5}$$

$$\Rightarrow m = \frac{-1}{2}$$

\therefore Equation of the line is $x + 2y - 5 = 0$

81. Given $b = 2a$

\therefore The equation of the line is $\frac{x}{a} + \frac{y}{b} = 1$

$$\Rightarrow \frac{x}{a} + \frac{y}{2a} = 1$$

$$\Rightarrow 2x + y = 2a$$

Distance of the line from (0, 0) is

$$d = \left| \frac{2(0) + 1(0) - 2a}{\sqrt{4+1}} \right|$$

$$\Rightarrow 1 = \left| \frac{-2a}{\sqrt{5}} \right|$$

$$\Rightarrow a = \pm \frac{\sqrt{5}}{2}$$

\therefore Equation of line is $2x + y = \pm\sqrt{5}$

82. Gradient of BC = -1 and its equation is $x + y + 4 = 0$. Therefore, the equation of line parallel to BC is $x + y + \lambda = 0$.

Also, it is $\frac{1}{2}$ unit distant from origin.

$$\text{Thus, } \frac{\lambda}{\sqrt{2}} = \frac{1}{2} \Rightarrow \lambda = \frac{\sqrt{2}}{2}$$

Hence, the required equation of line is $2x + 2y + \sqrt{2} = 0$

83. Equation of straight line parallel to $4x - 3y = 5$ is $4x - 3y = \lambda$

According to the given condition,

$$\frac{4(-1) - 3(-4) - \lambda}{\sqrt{16+9}} = \pm 1$$

$$\Rightarrow 8 - \lambda = \pm 5$$

$$\Rightarrow \lambda = 3, 13$$

\therefore the equation of one of the lines is $4x - 3y - 3 = 0$

84. Equation of AB: $4x - 3y - 17 = 0$

Equation of BC: $3x + 4y - 19 = 0$

If P(x, y) is a point on the bisector of $\angle ABC$ then,

$$\left| \frac{4x - 3y - 17}{\sqrt{(4)^2 + (-3)^2}} \right| = \left| \frac{3x + 4y - 19}{\sqrt{(3)^2 + (4)^2}} \right|$$

$\therefore 7y = x + 2$ is the required equation of the angle bisector.



85. Let point (x_1, y_1) be at distance $\sqrt{5}$ from $x - 2y + 1 = 0$

$$\therefore \sqrt{5} = \left| \frac{x_1 - 2y_1 + 1}{\sqrt{1+4}} \right|$$

$$\therefore x_1 - 2y_1 + 1 = \pm 5 \quad \dots(i)$$

Let point (x_2, y_2) be at distance $\sqrt{13}$ from $2x + 3y - 1 = 0$


$$\therefore \sqrt{13} = \left| \frac{2x_2 + 3y_2 - 1}{\sqrt{4+9}} \right|$$

$$\therefore 2x_2 + 3y_2 - 1 = \pm 13 \quad \dots(ii)$$

\therefore Equation (i) and (ii) will give us total 4 points.

\therefore answer is option (C)

86.



$$(AB) = (RQ) = \sqrt{(2-0)^2 + (0-2)^2} = \sqrt{8} = 2\sqrt{2}$$

$$(OR) = \left| \frac{1(0) + 1(0) - 2}{\sqrt{1+1}} \right| = \left| \frac{-2}{\sqrt{2}} \right| = \sqrt{2}$$

$$\therefore \text{Perpendicular distance (p)} = OR + RQ = \sqrt{2} + 2\sqrt{2} = 3\sqrt{2}$$

87. Slope of given line $ax + by + c = 0$ is $-\frac{a}{b}$.

$$\therefore -\frac{a}{b} = \pm 1 \Rightarrow a = \pm b \quad \dots(i)$$

$$\text{Distance of line } ax + by + c = 0 \text{ from } (1, -2) = \frac{|a - 2b + c|}{\sqrt{a^2 + b^2}}$$

$$\text{Distance of line } ax + by + c = 0 \text{ from } (3, 4) = \frac{|3a + 4b + c|}{\sqrt{a^2 + b^2}}$$

According to the given condition,

$$\frac{|a - 2b + c|}{\sqrt{a^2 + b^2}} = \frac{|3a + 4b + c|}{\sqrt{a^2 + b^2}}$$

$$\Rightarrow 3a + 4b + c = \pm(a - 2b + c)$$

$$\Rightarrow a + 3b = 0 \quad (\text{taking +ve}) \quad \dots(ii)$$

$$\Rightarrow 2a + b + c = 0 \quad (\text{taking -ve}) \quad \dots(iii)$$

From, (i) and (ii), we get $a = b = 0$ which is not possible so taking (i) and (iii), (taking $a = -b$) we get

$$a + c = 0 \Rightarrow c = -a$$

$$a : b : c = a : -a : -a = 1 : -1 : -1$$

$$\text{or } a = 1, b = -1, c = -1$$

From (i) and (iii) (taking $a = b$), we get

$$3a + c = 0 \Rightarrow c = -3a$$

$$a : b : c = a : a : -3a = 1 : 1 : -3$$

\therefore option (B) is the correct answer.

88. Here, the lines are $3x + 4y - 9 = 0$ and $6x + 8y - 15 = 0$ or $3x + 4y - \frac{15}{2} = 0$.

$$\therefore \text{Required distance} = \frac{\left| -9 - \left(-\frac{15}{2} \right) \right|}{\sqrt{3^2 + 4^2}} = \frac{|-3|}{10} = \frac{3}{10}$$

89. Given equation of parallel lines are $x - y + a = 0, x - y + b = 0$

$$\therefore \text{required distance} = \frac{|a - b|}{\sqrt{(1)^2 + (-1)^2}} = \frac{|a - b|}{\sqrt{2}}$$

90. Line L passes through $(13, 32)$.

$$\therefore \frac{13}{5} + \frac{32}{b} = 1$$

$$\Rightarrow b = -20$$

$$\text{So, equation of L is } \frac{x}{5} - \frac{y}{20} = 1 \Rightarrow 4x - y = 20$$

Slope of L is $m_1 = 4$.

$$\text{Slope of } \frac{x}{c} + \frac{y}{3} = 1 \text{ is } m_2 = -\frac{3}{c}$$

$$\Rightarrow -\frac{3}{c} = 4$$

$$\Rightarrow c = -\frac{3}{4}$$

$$\text{Equation of line K is } -\frac{4x}{3} + \frac{y}{3} = 1$$

$$\Rightarrow 4x - y = -3$$

$$\text{Distance between L and K is } \frac{|20 + 3|}{\sqrt{16 + 1}} = \frac{23}{\sqrt{17}}$$



91. Distance between lines $-x + y = 2$ and $x - y = 2$ is $\alpha = \left| \frac{2+2}{\sqrt{2}} \right| = 2\sqrt{2}$ (i)

Distance between lines $4x - 3y = 5$ and $6y - 8x = 1$ is

$$\beta = \left| \frac{5 - \left(\frac{-1}{2}\right)}{5} \right| = \frac{11}{10} \quad \dots(\text{ii})$$

From (i) and (ii), we get

$$\frac{\alpha}{\beta} = \frac{2\sqrt{2}}{11/10}$$

$$\Rightarrow 20\sqrt{2}\beta = 11\alpha$$

92. $(K+1)^2x + Ky - 2K^2 - 2 = 0$
 $\therefore (K^2 + 2K + 1)x + Ky - 2K^2 - 2 = 0$
 $\therefore K^2(x-2) + K(2x+y) + (x-2) = 0$
 $\therefore (K^2+1)(x-2) + K(2x+y) = 0$
 $\therefore x-2=0$ i.e. $x=2$
 and $2x+y=0$
 $\therefore 2(2)+y=0$
 $\therefore y=-4$
 \therefore The fixed point is $(2, -4)$
 \therefore The required line has slope 2 and passes through the point $(2, -4)$
 \therefore Equation of line;
 $y - (-4) = 2(x - 2)$
 $\therefore y + 4 = 2x - 4$
 $\therefore y = 2x - 8$

93. $ax + by + c = 0$ always passes through $(1, -2)$.
 $\therefore a - 2b + c = 0 \Rightarrow 2b = a + c$
 Therefore, a, b and c are in A.P.

94. Since, , m, n are in A.P.
 $\therefore 2m = +n$
 Given equation of line is $x + my = n = 0$
 Consider, option (B),
 If the point $(1, -2)$ satisfy the given equation.
 $\therefore -2m + n = 0 \Rightarrow 2m = +n$
 \Rightarrow , m, n are A.P.

95. a, b, c are in H. P., then $\frac{2}{b} = \frac{1}{a} + \frac{1}{c}$ (i)

Given, line is $\frac{x}{a} + \frac{y}{b} + \frac{1}{c} = 0$ (ii)


From (i) and (ii), we get

$$\frac{1}{a}(x-1) + \frac{1}{b}(y+2) = 0$$

Since, $a \neq 0, b \neq 0$

So, $(x-1) = 0$ and $(y+2) = 0$
 $\Rightarrow x = 1$ and $y = -2$

96.



For a triangle with side lengths a, b and c and vertices at points opposite to these sides (x_1, y_1) , (x_2, y_2) and (x_3, y_3) respectively, the incentre is given by,

$$(x_i, y_i) = \left(\frac{ax_1 + bx_2 + cx_3}{a + b + c}, \frac{ay_1 + by_2 + cy_3}{a + b + c} \right)$$

For the given triangle,

$$OA = 3 \text{ units}$$

$$OB = 4 \text{ units}$$

$$AB = \sqrt{(4-0)^2 + (0-3)^2} = 5 \text{ units}$$

\therefore Incentre

$$= \left(\frac{3(0) + 4(3) + 5(0)}{3 + 4 + 5}, \frac{3(4) + 4(0) + 5(0)}{3 + 4 + 5} \right)$$

$$= \left(\frac{12}{12}, \frac{12}{12} \right)$$

$$= (1, 1)$$

\therefore Incentre is $(1, 1)$

97. Two sides $x - 3y = 0$ and $3x + y = 0$ of the given triangle are perpendicular to each other. Therefore, its orthocentre is the point of intersection of $x - 3y = 0$ and $3x + y = 0$ i.e., $(0, 0)$.

98. The vertices of triangle are the intersection points of these given lines. The vertices of Δ are $A(0, 4)$, $B(1, 1)$, $C(4, 0)$

Now,

$$AB = \sqrt{(0-1)^2 + (4-1)^2} = \sqrt{10}$$

$$BC = \sqrt{(1-4)^2 + (1-0)^2} = \sqrt{10}$$

$$AC = \sqrt{(0-4)^2 + (4-0)^2} = 4\sqrt{2}$$

$\therefore AB = BC$

$\therefore \Delta$ is isosceles.


99. The point of intersection of the given lines are $(-1, 1)$, $(1, -1)$ and $(2/3, 2/3)$ which is the vertices of an isosceles triangle.



Evaluation Test

- Given, $f(\alpha) = x \cos \alpha + y \sin \alpha - p(\alpha)$
 $\therefore f(\beta) = x \cos \beta + y \sin \beta - p(\beta)$
 Since, both the lines are perpendicular to each other.
 $\therefore a_1 a_2 + b_1 b_2 = 0$
 $\Rightarrow \cos \alpha \cos \beta + \sin \alpha \sin \beta = 0$
 $\Rightarrow \cos(\alpha - \beta) = 0 \Rightarrow |\alpha - \beta| = \frac{\pi}{2}$


- The equations of the sides of the triangle are
 $L_1 \equiv \frac{x}{p} - \frac{y}{1+p} = -1,$
 $L_2 \equiv \frac{x}{q} - \frac{y}{1+q} = -1,$
 $L_3 \equiv y = 0$
 The coordinates of vertices are $A(-p, 0),$
 $B(-q, 0)$ and $C(pq, (1+p)(1+q)).$



The equation of the altitude through C is $x = pq$ and the equation of the altitude through B is $(1+p)y + px + pq = 0.$
 Solving these equations, we get
 $x = pq$ and $y = -pq$
 Let (h, k) be the coordinates of the orthocentre. Then,
 $h = pq$ and $k = -pq \Rightarrow k = -h$
 Hence, the locus of (h, k) is $y = -x,$ which is a straight line.

- The line $ax + by + c = 0$ meets the coordinate axes at $A\left(-\frac{c}{a}, 0\right)$ and $B\left(0, -\frac{c}{b}\right).$
 \therefore Area of $\Delta OAB = \frac{1}{2} \times OA \times OB$
 $= \frac{1}{2} \times \left|-\frac{c}{a}\right| \times \left|-\frac{c}{b}\right| = \frac{c^2}{2ab}$
 This will be constant, if a, c, b are in G.P.


- The clockwise rotation of the point $P(\cos \theta, \sin \theta)$ through an angle α takes it to the point $(\cos(\theta - \alpha), \sin(\theta - \alpha))$ and anticlockwise rotation through angle α takes P to the point $(\cos(\theta + \alpha), \sin(\theta + \alpha)).$
 \therefore options (A) and (B) are not correct.



$$\begin{aligned} \text{Slope of PQ} &= \frac{\sin(\alpha - \theta) - \sin \theta}{\cos(\alpha - \theta) - \cos \theta} \\ &= -\frac{2 \sin\left(\frac{\alpha - \theta}{2}\right) \cos \frac{\alpha}{2}}{2 \sin\left(\frac{\alpha - \theta}{2}\right) \sin \frac{\alpha}{2}} \\ &= -\cot \frac{\alpha}{2} \end{aligned}$$

This shows that PQ is perpendicular to a line with slope $\tan \frac{\alpha}{2}.$ Thus, Q can be obtained from P by taking its reflection in the line through origin with slope $\tan \frac{\alpha}{2}.$

- Let QS be the bisector of $\angle PQR.$




$$\text{Slope of QR} = \frac{3\sqrt{3} - 0}{3 - 0} = \sqrt{3}$$



- $\therefore \angle XQR = 60^\circ$
 $\Rightarrow \angle PQR = 120^\circ$
 $\Rightarrow \angle PQS = \angle SQR = 60^\circ \Rightarrow \angle XQS = 120^\circ$
 \therefore Slope of QS = $\tan 120^\circ = -\sqrt{3}$
 \therefore the equation of QS is $y = -\sqrt{3}x$ i.e., $\sqrt{3}x + y = 0$

6.



Slope of OB = $\tan\left(\frac{\pi}{4} + \alpha\right)$

- \therefore Slope of AC = $-\cot\left(\frac{\pi}{4} + \alpha\right)$
 $= -\left(\frac{\cos \alpha - \sin \alpha}{\cos \alpha + \sin \alpha}\right)$
 $= \frac{\sin \alpha - \cos \alpha}{\sin \alpha + \cos \alpha}$

\therefore the equation of AC is

$$y - a \sin \alpha = \frac{\sin \alpha - \cos \alpha}{\sin \alpha + \cos \alpha} (x - a \cos \alpha)$$

$$\Rightarrow y(\sin \alpha + \cos \alpha) + x(\cos \alpha - \sin \alpha) = a$$


7. Equation of line L passing through (1, 1) and (2, 0) is

$$y - 1 = \frac{0-1}{2-1} (x - 1)$$

$$\Rightarrow x + y = 2 \quad \dots(i)$$

\therefore Slope of L = -1

Also, slope of L' = 1 $\dots[\because L \perp L']$



Equation of line L' passing through $\left(\frac{1}{2}, 0\right)$

and having slope 1 is

$$y - 0 = 1 \left(x - \frac{1}{2}\right)$$

$$\Rightarrow 2x - 2y = 1 \quad \dots(ii)$$

Equation of Y axis, $x = 0$ $\dots(iii)$

From (i), (ii) and (iii),

the vertices of the triangle are

$$A(0, 2), B\left(0, -\frac{1}{2}\right) \text{ and } C\left(\frac{5}{4}, \frac{3}{4}\right).$$

\therefore the area of the triangle is

$$\frac{1}{2} \begin{vmatrix} 0 & 2 & 1 \\ 0 & -\frac{1}{2} & 1 \\ \frac{5}{4} & \frac{3}{4} & 1 \end{vmatrix} = \frac{25}{16} \text{ square units}$$

8. By solving $3x + 4y = 9$, $y = mx + 1$, we get

$$x = \frac{5}{3 + 4m}.$$

Now, x is an integer, if $3 + 4m = 1, -1, 5, -5$

$$\therefore m = \frac{-2}{4}, \frac{-4}{4}, \frac{2}{4}, \frac{-8}{4}.$$

Since, $m = \frac{-2}{4}, \frac{2}{4}$ do not give integral values

of m .

\therefore m has two integral values.

9. Given, the lines $ax + by + p = 0$ and

$x \cos \alpha + y \sin \alpha - p = 0$ are inclined at an

angle $\frac{\pi}{4}$.

$$\therefore \tan \frac{\pi}{4} = \left| \frac{-\frac{a}{b} + \frac{\cos \alpha}{\sin \alpha}}{1 + \frac{a \cos \alpha}{b \sin \alpha}} \right|$$

$$\Rightarrow a \cos \alpha + b \sin \alpha = -a \sin \alpha + b \cos \alpha$$

$\dots(i)$

Also, the lines $ax + by + p = 0$,

$x \cos \alpha + y \sin \alpha - p = 0$ and $x \sin \alpha - y \cos \alpha = 0$

are concurrent.



$$\therefore \begin{vmatrix} a & b & p \\ \cos \alpha & \sin \alpha & -p \\ \sin \alpha & -\cos \alpha & 0 \end{vmatrix} = 0$$

$$\Rightarrow -ap \cos \alpha - bp \sin \alpha - p = 0$$

$$\Rightarrow a \cos \alpha + b \sin \alpha = -1 \quad \dots(\text{ii})$$

From (i) and (ii), we get

$$-a \sin \alpha + b \cos \alpha = -1 \quad \dots(\text{iii})$$

Squaring (ii) and (iii) and adding, we get


$$(a \cos \alpha + b \sin \alpha)^2 + (-a \sin \alpha + b \cos \alpha)^2 = 2$$

$$\Rightarrow a^2 + b^2 = 2$$

10. Slopes of AB and BC

are -4 and $\frac{3}{4}$

respectively.



Let α be the angle between AB and BC.

$$\text{Then, } \tan \alpha = \left| \frac{-4 - \frac{3}{4}}{1 - 4\left(\frac{3}{4}\right)} \right| = \frac{19}{8} \quad \dots(\text{i})$$

Since, $AB = AC$

$$\Rightarrow \angle ABC = \angle ACB = \alpha$$

\therefore the line AC also makes an angle α with BC.

If m is the slope of the line AC, then its

$$\text{equation is } y + 7 = m(x - 2) \quad \dots(\text{ii})$$

$$\text{Now, } \tan \alpha = \pm \left[\frac{m - \frac{3}{4}}{1 + m \cdot \frac{3}{4}} \right]$$

$$\Rightarrow \frac{19}{8} = \pm \frac{4m - 3}{4 + 3m} \quad \dots[\text{From (i)}]$$

$$\Rightarrow m = -4 \text{ or } -\frac{52}{89}$$

But slope of AB is -4 , so slope of AC is $-\frac{52}{89}$.

Therefore, the equation of line AC given by (ii) is $52x + 89y + 519 = 0$.

07 Circle and Conics



Hints



Classical Thinking

2. Required equation is $(x - a)^2 + (y - a)^2 = a^2$
 $\Rightarrow x^2 + y^2 - 2ax - 2ay + a^2 = 0$

3. The equation of circle with centre (x_1, y_1) is
 $(x - x_1)^2 + (y - y_1)^2 = r^2$
 since, the circle touches both the axes,
 $x_1 = y_1 = r$

$\therefore (x - x_1)^2 + (y - x_1)^2 = x_1^2$
 $\Rightarrow x^2 + y^2 - 2x_1(x + y) + x_1^2 = 0$

4. Since, the circle touches X axis


\therefore Radius = 2.

\therefore the equation of the circle is

$$(x - 1)^2 + (y - 2)^2 = 2^2$$

$$\Rightarrow x^2 + y^2 - 2x - 4y + 1 = 0$$

5. Let O' be the centre



Now, from the figure

$$\text{Radius } (r) = \sqrt{(4)^2 + (3)^2} = 5$$

6. Extremities of diameter are $(5, 7)$ and $(1, 4)$
 Radius is half of the distance between them

\therefore Radius = $\frac{1}{2} \sqrt{(4)^2 + (3)^2}$
 $= \frac{5}{2}$

7. Using condition of point circle

$$\text{Radius} = \sqrt{g^2 + f^2 - c} = 0$$

$$\Rightarrow g^2 + f^2 = c$$

8. $(\text{Radius})^2 = g^2 + f^2 - c$
 $\Rightarrow 121 = 81 + 36 - k \Rightarrow k = -4$

9. If $c = 0$; circle passes through origin.

10. Intercept made by the circle on the X-axis
 $= 2\sqrt{g^2 - c}$
 $= 2\sqrt{9 - 9} = 0$

\therefore Intercept cut on X-axis is zero.
 Hence, circle touches X-axis.

11. Circle $x^2 + y^2 + 2gx + 2fy + c = 0$ touches X axis

\therefore radius = ordinate of centre

$$\Rightarrow \sqrt{g^2 + f^2 - c} = (-f)$$

$$\Rightarrow g^2 = c$$


12. Required conditions are $g = f = r$ and

$$\sqrt{g^2 + f^2 - c} = r$$

$$\Rightarrow g = \sqrt{c} = f = r$$

13. Both axis, as centre is $(-2, 2)$ and radius is 2.

14. Centre is $(0, -3)$ and radius = $\sqrt{0^2 + 9 - 0} = 3$



Hence, circle touches X-axis at the origin.

15. Centre $(3, 4)$ of the given circle is satisfying only $x + y = 7$

\therefore Correct answer is the option (C)

16. Here, the centre of circle $(3, -1)$ must lie on the line $x + 2by + 7 = 0$.

$\therefore 3 - 2b + 7 = 0$
 $\Rightarrow b = 5$

17. Centre of the required circle is $(-4, -5)$ and it passes through $(2, 3)$

\therefore Radius = $\sqrt{(-4 - 2)^2 + (-5 - 3)^2}$
 $= 10$

\therefore Equation of the required circle is
 $(x + 4)^2 + (y + 5)^2 = (10)^2$
 $\Rightarrow x^2 + y^2 + 8x + 10y - 59 = 0$



18. The equation of circle in third quadrant touching the coordinate axes with centre $(-a, -a)$ and radius 'a' is

$$x^2 + y^2 + 2ax + 2ay + a^2 = 0 \quad \dots(i)$$

Since, line $3x - 4y + 8 = 0$ touches the circle

- \therefore perpendicular distance from centre of the circle to the line = radius

$$\therefore \left| \frac{3(-a) - 4(-a) + 8}{\sqrt{9+16}} \right| = a$$

$$\Rightarrow a = 2$$

Substituting $a = 2$ in equation (i), we get

$$x^2 + y^2 + 4x + 4y + 4 = 0$$

This is the required equation of the circle

19. Radius of circle = $\left| \frac{2(1) - 1(-3) - 4}{\sqrt{4+1}} \right| = \frac{1}{\sqrt{5}}$

$$\therefore \text{Equation is } (x-1)^2 + (y+3)^2 = \left(\frac{1}{\sqrt{5}} \right)^2$$

$$\Rightarrow x^2 + y^2 - 2x + 6y + 10 = \frac{1}{5}$$

$$\Rightarrow 5x^2 + 5y^2 - 10x + 30y + 49 = 0$$

20. $4x^2 + 4y^2 = 9$

$$\Rightarrow x^2 + y^2 = \frac{9}{4} \Rightarrow x^2 + y^2 = \left(\frac{3}{2} \right)^2$$

$$\therefore x = \frac{3}{2} \cos \theta, y = \frac{3}{2} \sin \theta$$

21. $(x-3)^2 + (y+4)^2 = 5^2$

Comparing with $(x-h)^2 + (y-k)^2 = r^2$, we get
 $h = 3, k = -4, r = 5$

- \therefore Parametric equations are

$$x = 3 + 5 \cos \theta, y = -4 + 5 \sin \theta$$

22. Given equation can be written as

$$(x^2 + 2x + 1 - 1) + (y^2 - 4y + 4 - 4) - 4 = 0$$

$$\Rightarrow (x+1)^2 + (y-2)^2 = 3^2$$

- $\therefore h = -1, k = 2$ and $r = 3$

- \therefore Parametric form of equation are

$$x = -1 + 3 \cos \theta, y = 2 + 3 \sin \theta$$

23. $\frac{x+1}{2} = \cos \theta \quad \dots(i)$

$$\text{and } \frac{y-3}{2} = \sin \theta \quad \dots(ii)$$


Squaring (i) and (ii) and adding, we get

$$\left(\frac{x+1}{2} \right)^2 + \left(\frac{y-3}{2} \right)^2 = 1$$

$$\Rightarrow (x+1)^2 + (y-3)^2 = 4,$$

- \therefore centre is $(-1, 3)$

24. Parabola $y^2 = x$ is symmetric about X-axis.



25. $S \equiv (5,0)$

Therefore, latus rectum = $4a = 20$

26. Since, parabola $y^2 = 4ax$ passes through $(-3, 2)$

$$\therefore 4 = -12a$$

$$\Rightarrow 4a = -\frac{4}{3} = \frac{4}{3} \quad \dots[\text{Taking positive sign}]$$

27. Let the equation of parabola be $x^2 = -4ay$
 Since, parabola passing through the point

$$(-4, -2).$$

$$\therefore (-4)^2 = -4a(-2)$$

$$\Rightarrow a = 2$$

- \therefore equation becomes $x^2 = -8y$ and

$$\text{latus rectum} = 4a = 8$$

29. $y^2 = 4 \cdot \frac{1}{5}x \Rightarrow a = \frac{1}{5}$

co-ordinates of latus rectum are $(a, 2a)$ and $(a, -2a)$

$$\text{i.e., } \left(\frac{1}{5}, \frac{2}{5} \right) \text{ and } \left(\frac{1}{5}, -\frac{2}{5} \right)$$

30. $x^2 = -8y$

$$\Rightarrow a = 2$$

So, focus = $(0, -2)$

- \therefore Ends of latus rectum = $(4, -2), (-4, -2)$.

31. $y^2 = 12x$

$$\therefore a = 3$$

$$\Rightarrow \text{abscissa is } 4 - 3 = 1 \text{ and } y^2 = 12, y = \pm 2\sqrt{3}$$

Hence, points are $(1, 2\sqrt{3}), (1, -2\sqrt{3})$

32. We have, $y = 3x$

According to given condition,

$$(3x)^2 = 36x$$

$$\Rightarrow x = 4 \text{ and } y = 12$$

- \therefore Required point is $(4, 12)$

34. $y^2 + 2y + x = 0$

$$\Rightarrow y^2 + 2y + 1 = -x + 1$$

$$(y+1)^2 = -(x-1)$$

Hence, vertex is $(1, -1)$, which lies in IVth quadrant.



35. Vertex = (2, 0)
 \Rightarrow focus is (2 + 2, 0) = (4, 0)
36. $y^2 = 4y - 4x$
 $\Rightarrow y^2 - 4y + 4 = -4x + 4$
 $\Rightarrow (y - 2)^2 = -4(x - 1)$
 Comparing this equation with $Y^2 = -4aX$, we get
 $a = 1$, $X = x - 1$ and $Y = y - 2$
 Focus of the parabola is
 $X = -a$, $Y = 0$
 $\Rightarrow x - 1 = -1$, $y - 2 = 0$
 $\Rightarrow x = 0$, $y = 2$
 \therefore focus = (0, 2)
37. Equation of parabola is $x^2 - 4x - 8y + 12 = 0$
 $\Rightarrow x^2 - 4x + 4 = 8y - 8$
 $\Rightarrow (x - 2)^2 = 8(y - 1) \Rightarrow X^2 = 8Y$
 Comparing with $X^2 = 4aY$, we get $a = 2$
 \therefore Directrix is $Y = -a \Rightarrow y - 1 = -2 \Rightarrow y = -1$.
38. The parabola is $(x - 2)^2 = (3y - 6)$
 Hence, axis is $x - 2 = 0$.
39. The given equation of parabola is
 $x^2 - 4x - 8y + 12 = 0$
 $\Rightarrow x^2 - 4x = 8y - 12$
 $\Rightarrow x^2 - 4x + 4 = 8y - 12 + 4$
 $\Rightarrow (x - 2)^2 = 8(y - 1)$
 Hence, the length of latus rectum = $4a = 8$.
40. $x^2 + 5y = 0 \Rightarrow x^2 = -5y$
 On comparing with $x^2 = -4ay$, we get
 $a = \frac{5}{4}$
 End points of latus rectum of the parabola are
 $(\pm 2a, -a) = \left(\pm \frac{5}{2}, -\frac{5}{4}\right)$
41. Here, $ae = 4$ and $e = \frac{4}{5}$
 $\Rightarrow a = 5$
 Now, $b^2 = a^2(1 - e^2)$
 $\Rightarrow b^2 = 25\left(1 - \frac{16}{25}\right) = 9$
 \therefore Equation of the ellipse is $\frac{x^2}{25} + \frac{y^2}{9} = 1$.
42. Since, $e^2 = 1 - \frac{b^2}{a^2}$
 $\Rightarrow \left(\frac{2}{3}\right)^2 = 1 - \frac{b^2}{a^2}$

$$\Rightarrow b^2 = \frac{5a^2}{9} \quad \dots(i)$$

Given length of latus rectum = 5

$$\Rightarrow \frac{2b^2}{a} = 5$$

$$\Rightarrow b^2 = \frac{5a}{2} \quad \dots(ii)$$

From (i) and (ii), we get

$$\Rightarrow a^2 = \frac{81}{4}, b^2 = \frac{45}{4}$$

$$\therefore \text{Equation of ellipse is } \frac{4x^2}{81} + \frac{4y^2}{45} = 1$$

$$43. e^2 = 1 - \frac{b^2}{a^2} = 1 - \frac{28}{64}$$

$$\Rightarrow e^2 = \frac{36}{64} \Rightarrow e = \frac{3}{4}$$

$$44. \frac{x^2}{\frac{112}{16}} + \frac{y^2}{\frac{112}{7}} = 1$$

$$\therefore e = \sqrt{1 - \frac{a^2}{b^2}} = \sqrt{1 - \frac{112}{16} \cdot \frac{7}{112}} = \frac{3}{4}$$

$$45. \text{According to the condition, } \frac{2a}{e} = 3(2ae)$$

$$\Rightarrow e = \frac{1}{\sqrt{3}}$$

$$46. \text{Foci are } (\pm ae, 0)$$

\therefore According to the condition, $2ae = 2b$

$$\Rightarrow ae = b \quad \dots(i)$$

$$\text{Also, } b^2 = a^2(1 - e^2)$$

$$\Rightarrow e^2 = (1 - e^2) \quad \dots[\text{From (i)}]$$

$$\Rightarrow e = \frac{1}{\sqrt{2}}$$

$$47. \text{We have, } ae = 1 \text{ and } a = 2$$

$$\Rightarrow e = \frac{1}{2}$$

$$\text{Also, } b^2 = a^2(1 - e^2)$$

$$\Rightarrow b = 2\sqrt{1 - \frac{1}{4}} = \sqrt{3}$$

$$\therefore \text{Minor axis} = 2b = 2\sqrt{3}$$

$$49. 3x^2 + 4y^2 = 12$$

$$\Rightarrow \frac{x^2}{4} + \frac{y^2}{3} = 1$$

$$\therefore \text{Latus rectum} = \frac{2b^2}{a} = 3$$



50. Here, $a^2 = 36$, $b^2 = 49$
 Since, $b > a$
 \therefore the length of the latus rectum
 $= \frac{2a^2}{b} = 2 \times \frac{36}{7} = \frac{72}{7}$
51. We have, $e = \frac{1}{\sqrt{2}}$
 \therefore Latus rectum $= \frac{2b^2}{a} = \frac{2}{a} \times a^2(1 - e^2)$
 $= 2a \left(1 - \frac{1}{2}\right) = a$
 i.e., semi-major axis
52. We have, $2ae = 10 \Rightarrow a = \frac{10}{2 \times 5} = 8$
 Also, $b^2 = a^2(1 - e^2)$
 $\Rightarrow b = 8\sqrt{1 - \frac{25}{64}} = \sqrt{39}$
 Now, Latus rectum $= \frac{2b^2}{a} = \frac{2 \times 39}{8} = \frac{39}{4}$
53. Focal distance of any point P (x, y) on the ellipse is equal to $SP = a + ex$.
 Here, $x = a \cos \theta$
 $\therefore SP = a + ae \cos \theta$
 $= a(1 + e \cos \theta)$
54. $4x^2 + 9y^2 - 16x - 54y + 61 = 0$
 $\Rightarrow 4x^2 - 16x + 9y^2 - 54y = -61$
 $\Rightarrow 4(x^2 - 4x + 4 - 4) + 9(y^2 - 6y + 9 - 9) = -61$
 $\Rightarrow 4(x-2)^2 + 9(y-3)^2 = 36$
 $\Rightarrow \frac{(x-2)^2}{9} + \frac{(y-3)^2}{4} = 1$
 Hence, the centre is $(2, 3)$
55. Distance between foci is $4 = 2ae$
 $\Rightarrow a^2 = \frac{9}{4}$
 Also, $e^2 = 1 + \frac{b^2}{a^2}$
 $\Rightarrow \frac{16}{9} - 1 = \frac{b^2}{a^2} \Rightarrow b^2 = \frac{7}{4}$
 Centre is $(0, 4)$
 Hence, equation of hyperbola is
 $\frac{x^2}{9} - \frac{4(y-4)^2}{7} = 1$
56. Let $S(1, -1)$ be the focus and $P \equiv (x, y)$ be any point on conic.
 Now, $PS = e PM$
 $\Rightarrow \sqrt{(x-1)^2 + (y+1)^2} = \sqrt{2} \left| \frac{x-y+1}{\sqrt{1+1}} \right|$
 $\Rightarrow \sqrt{x^2 - 2x + 1 + y^2 + 2y + 1} = |x - y + 1|$
 Squaring both sides, we get
 $4x - 4y - 2xy - 1 = 0$
 $\Rightarrow 2xy - 4x + 4y + 1 = 0$
57. $2a = 8$, $2b = 6$
 Difference of focal distances of any point of the hyperbola $= 2a = 8$
58. The equation of hyperbola is $9x^2 - 16y^2 = 144$
 $\Rightarrow \frac{x^2}{16} - \frac{y^2}{9} = 1$
 $e = \frac{\sqrt{a^2 + b^2}}{a} = \frac{\sqrt{16+9}}{4} = \frac{5}{4}$
 Hence, foci are $(\pm ae, 0) \Rightarrow \left(\pm 4 \times \frac{5}{4}, 0\right)$
 i.e., $(\pm 5, 0)$.
59. Since, $e > 1$ always for hyperbola and $\frac{2}{3} < 1$.
60. $\frac{x^2}{25} - \frac{y^2}{25} = 1$
 \therefore Eccentricity $= \sqrt{2}$ as $a = b$.
61. $\frac{y^2}{k^2} - \frac{x^2}{-k} = 1$
 Also, $a^2 = b^2(e^2 - 1)$
 $\Rightarrow -k = k^2(e^2 - 1) \Rightarrow -\frac{1}{k} = e^2 - 1$
 $\Rightarrow e^2 = 1 - \frac{1}{k} \Rightarrow e = \sqrt{1 - \frac{1}{k}}$
62. Vertices $(\pm 4, 0) \equiv (\pm a, 0)$
 $\Rightarrow a = 4$
 Foci $(\pm 6, 0) \equiv (\pm ae, 0)$
 $\Rightarrow e = \frac{6}{4} = \frac{3}{2}$
63. The given equation of hyperbola is
 $16x^2 - 9y^2 = 144 \Rightarrow \frac{x^2}{9} - \frac{y^2}{16} = 1$
 \therefore L.R. $= \frac{2b^2}{a} = \frac{2 \times 16}{3} = \frac{32}{3}$



$$64. \text{ Length of latus rectum} = \frac{2b^2}{a} = \frac{2 \times 4}{\frac{2}{\sqrt{3}}} = 4\sqrt{3}$$

$$66. \text{ Equation of hyperbola is} \\ x = 8 \sec \theta, y = 8 \tan \theta$$

$$\Rightarrow \frac{x}{8} = \sec \theta, \frac{y}{8} = \tan \theta$$

$$\therefore \sec^2 \theta - \tan^2 \theta = 1$$

$$\Rightarrow \frac{x^2}{8^2} - \frac{y^2}{8^2} = 1$$

$$\text{Here, } a = 8, b = 8$$

$$\text{Here, } a = b$$

$$\therefore \text{ it is rectangular hyperbola}$$

$$\Rightarrow e = \sqrt{2}$$

$$\therefore \text{ Distance between directrices} = \frac{2a}{e} \\ = \frac{2 \times 8}{\sqrt{2}} \\ = 8\sqrt{2}$$

$$67. \text{ The given equation of hyperbola is} \\ 9x^2 - 36x - 16y^2 + 96y - 252 = 0 \\ \text{Partially differentiating with respect to } x, \text{ we get} \\ 18x - 36 = 0 \\ \Rightarrow x = 2$$

Now partially differentiating with respect to y , we get

$$-32y + 96 = 0$$

$$\Rightarrow -32y = -96 \Rightarrow y = 3$$

$$\therefore \text{ Centre} \equiv (2, 3)$$

$$68. \text{ Given equation of hyperbola is} \\ 5x^2 - 4y^2 + 20x + 8y = 4 \\ \Rightarrow 5(x^2 + 4x + 4) - 4(y^2 - 2y + 1) = 4 + 20 - 4 \\ \Rightarrow 5(x+2)^2 - 4(y-1)^2 = 20 \\ \Rightarrow \frac{(x+2)^2}{4} - \frac{(y-1)^2}{5} = 1 \\ \Rightarrow a^2 = 4, b^2 = 5 \\ e = \frac{\sqrt{a^2 + b^2}}{a} = \frac{\sqrt{4+5}}{2} = \frac{3}{2}$$

$$69. 2x = e^t + e^{-t} \text{ and } 2y = e^t - e^{-t} \\ \Rightarrow 4x^2 = e^{2t} + 2 + e^{-2t} \dots\dots\dots (i) \\ \text{and } 4y^2 = e^{2t} - 2 + e^{-2t} \dots\dots\dots (ii)$$

Subtracting (ii) from (i), we get

$$4x^2 - 4y^2 = 4$$

$$\Rightarrow x^2 - y^2 = 1$$

The equation represents hyperbola.

$$70. \text{ The equation is } (x-0)^2 + (y-0)^2 = a^2$$



Critical Thinking

$$1. \text{ Radius} = \sqrt{\cos^2 \theta + \sin^2 \theta + 8} = 3$$

$$2. \text{ The point of intersection of } 3x + y - 14 = 0 \\ \text{and } 2x + 5y - 18 = 0 \text{ is } (4, 2).$$

Centre of the circle is $(1, -2)$.

$$\therefore \text{ radius} = \sqrt{(4-1)^2 + (2+2)^2} = 5$$

$$\therefore \text{ the equation of the circle is} \\ (x-1)^2 + (y+2)^2 = 5^2$$

$$\therefore x^2 + y^2 - 2x + 4y - 20 = 0.$$

$$3. \text{ Given, } OA = 3 \text{ and} \\ OB = 4$$

$$\therefore OL = \frac{3}{2} \text{ and } CL = 2$$

By pythagoras theorem,

$$OC^2 = OL^2 + LC^2$$

$$\therefore OC^2 = \left(\frac{3}{2}\right)^2 + 2^2 \\ = \frac{25}{4}$$

$$\therefore OC = \frac{5}{2}$$

The centre of the circle is $\left(\frac{3}{2}, 2\right)$ and

$$\text{radius} = \frac{5}{2}.$$

$$\therefore \text{ the equation of the circle is}$$

$$\left(x - \frac{3}{2}\right)^2 + (y-2)^2 = \left(\frac{5}{2}\right)^2$$

$$\therefore x^2 + y^2 - 3x - 4y = 0$$

$$4. \text{ Here, } r = 10 \text{ (radius)}$$

Centre will be the point of intersection of the diameters, i.e., $(8, -2)$.

Hence, required equation is

$$(x-8)^2 + (y+2)^2 = 10^2$$

$$\Rightarrow x^2 + y^2 - 16x + 4y - 32 = 0$$

$$5. \text{ Let its centre be } (h, k), \text{ then}$$

$$h - k = 1 \dots\dots(i)$$

Also, radius $a = 3$

$$\therefore \text{ Equation of the circle is}$$

$$(x-h)^2 + (y-k)^2 = 9$$


Also, it passes through $(7, 3)$

$$\text{i.e., } (7-h)^2 + (3-k)^2 = 9 \dots\dots(ii)$$

From (i) and (ii), we get

$$h = 4, k = 3$$

$$\therefore \text{ Equation is } x^2 + y^2 - 8x - 6y + 16 = 0$$





6. Centre is $(-4, 3)$
 Radius = Distance between centres – Radius of other circle = $5 - 1 = 4$
 Hence, equation of circle is $x^2 + y^2 + 8x - 6y + 9 = 0$
7. Centre of the given circle is $(0, -1)$
 \therefore the required circle passes through $(0, -1)$.
 $\therefore r = \sqrt{(0-1)^2 + (-1+2)^2} = \sqrt{2}$
 Hence, the required equation is
 $(x-1)^2 + (y+2)^2 = (\sqrt{2})^2$
 $\Rightarrow x^2 + y^2 - 2x + 4y + 3 = 0$
8. Centre of the circle
 $x^2 + y^2 - 4x - 6y - 3 = 0$ is $C(2,3)$.
 Since, it touches the Y-axis
 $\therefore r = 2$
 Hence, required equation of the circle is
 $(x-2)^2 + (y-3)^2 = 2^2$
 $\Rightarrow x^2 + y^2 - 4x - 6y + 9 = 0$
9. Let centre of circle be (h, k) .
 Since it touches both axes, therefore $h = k = a$
 Hence, equation can be $(x-a)^2 + (y-a)^2 = a^2$
 But it also touches the line $3x + 4y = 4$
 $\therefore \frac{|3a+4a-4|}{\sqrt{9+16}} = a$
 $\Rightarrow a = 2$
 Hence, the required equation of circle is
 $(x-2)^2 + (y-2)^2 = 2^2$
 $\Rightarrow x^2 + y^2 - 4x - 4y + 4 = 0$
10. Let the centre of the required circle be (x_1, y_1) .
 Centre of given circle is $(1, 2)$ and
 $r = \sqrt{1+4+20} = 5$
 \therefore radii of both circles are same.
 \therefore Point of contact $(5, 5)$ is the mid point of the line joining the centres of both circles.
 $\therefore \frac{x_1+1}{2} = 5$ and $\frac{y_1+2}{2} = 5$
 $\Rightarrow x_1 = 9, y_1 = 8$
 Hence, the required equation is
 $(x-9)^2 + (y-8)^2 = 25$
 $\Rightarrow x^2 + y^2 - 18x - 16y + 120 = 0$
11. Equation of circle concentric to given circle is
 $x^2 + y^2 - 6x + 12y + k = 0$
 Since, area of required circle = 2 (area of given circle)
 $\Rightarrow \sqrt{9+36-k} = \sqrt{2} \sqrt{9+36-15}$
 $\Rightarrow 45 - k = 60$
 $\Rightarrow k = -15$
 Hence, the required equation of circle is
 $x^2 + y^2 - 6x + 12y - 15 = 0$
12. We have the equation of circle
 $x^2 + y^2 + 2gx + 2fy + c = 0$
 But it passes through $(0, 0)$ and $(2, 1)$
 $\therefore c = 0$
 $5 + 4g + 2f = 0 \quad \dots(i)$
 Also $\sqrt{g^2 + f^2 - c} = |g|$
 $\Rightarrow f = 0 \quad \dots[\because c = 0]$
 $\therefore g = -\frac{5}{4} \quad \dots[\text{From (i)}]$
 Hence, the equation will be $2x^2 + 2y^2 - 5x = 0$.
13. since, X-intercept = $2a$
 $\therefore 2\sqrt{g^2 - c} = 2a \quad \dots(i)$
 Also, Y-intercept = $2b$
 $\therefore 2\sqrt{f^2 - c} = 2b \quad \dots(ii)$
 On squaring (i) and (ii) and then subtracting (ii) from (i), we get
 $g^2 - f^2 = a^2 - b^2$
 Hence, the locus is
 $x^2 - y^2 = a^2 - b^2$
14. Let the equation of circle be
 $x^2 + y^2 + 2gx + 2fy + c = 0$.
 Now on passing through the given points, we get three equations
 $c = 0 \quad \dots(i)$
 $a^2 + 2ga + c = 0 \quad \dots(ii)$
 $b^2 + 2fb + c = 0 \quad \dots(iii)$
 solving equations (i), (ii) and (iii), we get
 $g = -\frac{a}{2}, f = -\frac{b}{2}$
 Hence, the centre is $\left(\frac{a}{2}, \frac{b}{2}\right)$.
15. The equation of circle through points $(0, 0)$,
 $(1, 3)$ and $(2, 4)$ is
 $x^2 + y^2 - 10x = 0$
 Point $(k, 3)$ will be on the circle, if
 $k^2 + 9 - 10k = 0$
 $\Rightarrow k^2 - 10k + 9 = 0$
 $\Rightarrow k^2 - 9k - k + 9 = 0$
 $\Rightarrow (k-1)(k-9) = 0$
 $\Rightarrow k = 1$ or $k = 9$
16. Given, $x = 2 + 3 \cos \theta$
 i.e., $x - 2 = 3 \cos \theta \quad \dots(i)$
 and $y = 3 - 3 \sin \theta$
 i.e., $y - 3 = -3 \sin \theta \quad \dots(ii)$
 Squaring and adding equation (i) and (ii), we get



$$(x-2)^2 + (y-3)^2 = 9 \cos^2 \theta + 9 \sin^2 \theta$$

$$\therefore (x-2)^2 + (y-3)^2 = 3^2$$

$$\therefore \text{centre} \equiv (2, 3), \text{radius} = 3 \text{ units}$$

$$17. \quad x = 2a \left(\frac{1-t^2}{1+t^2} \right) \quad \dots \text{(i)}$$

$$y = \frac{4at}{1+t^2} \quad \dots \text{(ii)}$$

Squaring and adding (i) and (ii), we get

$$x^2 + y^2 = 4a^2 \cdot \frac{(1-t^2)^2}{(1+t^2)^2} + \frac{16a^2t^2}{(1+t^2)^2}$$

$$= \frac{4a^2}{(1+t^2)^2} [1 - 2t^2 + t^4 + 4t^2]$$

$$= \frac{4a^2}{(1+t^2)^2} (1+t^2)^2$$

$$\therefore x^2 + y^2 = (2a)^2$$

$$\therefore \text{Radius} = 2a$$

18. The point of intersection is

$$x = a \cos \theta + b \sin \theta$$

$$y = a \sin \theta - b \cos \theta$$

$$\therefore x^2 + y^2 = a^2 + b^2$$

Hence, it is equation of a circle.

19. Since, the axis of parabola is Y axis

$$\therefore \text{Equation of parabola } x^2 = 4ay$$

Since, it passes through (6, -3)

$$\therefore 36 = -12a$$

$$\Rightarrow a = -3$$

$$\therefore \text{Equation of parabola is } x^2 = -12y$$

20. Since, the axis of parabola is vertical and its vertex is (-1, -2).

\therefore equation of parabola is

$$(x+1)^2 = 4a(y+2)$$

Also, it passes through (3, 6)

$$\therefore 16 = 4a \times 8$$

$$\Rightarrow a = \frac{1}{2} \Rightarrow (x+1)^2 = 2(y+2)$$

$$\Rightarrow x^2 + 2x - 2y - 3 = 0$$

21. $a = VS$

$$= \sqrt{(2-2)^2 + (-3+1)^2}$$

$$= 2$$

Since,


$$(x-h)^2 = -4a(y-k)$$

$$\therefore (x-2)^2 = -4 \times 2 (y+1)$$

$$\Rightarrow (x-2)^2 = -8(y+1)$$

$$\Rightarrow x^2 + 4 - 4x = -8y - 8$$

$$\Rightarrow x^2 - 4x + 8y + 12 = 0$$



22. Let any point on required parabola be P(x, y), then from definition of parabola, we get
PS = PM

$$\Rightarrow \sqrt{(x+8)^2 + (y+2)^2} = \left| \frac{2x-y-9}{\sqrt{5}} \right|$$

$$\Rightarrow 5(x^2 + 16x + 64 + y^2 + 4y + 4)$$

$$= 4x^2 + y^2 + 81 - 4xy + 18y - 36x$$

$$\Rightarrow x^2 + 4y^2 + 4xy + 116x + 2y + 259 = 0$$

23. Let (x, y) be any point on the required parabola
 \therefore by using definition of parabola, we get

$$(x-a)^2 + (y-b)^2 = \left(\frac{bx+ay-ab}{\sqrt{a^2+b^2}} \right)^2$$

$$\Rightarrow (a^2+b^2)(x^2-2ax+a^2+y^2-2yb+b^2)$$

$$= b^2x^2+a^2y^2+a^2b^2+2abxy-2a^2yb-2ab^2x$$

$$\Rightarrow a^2x^2-2a^3x+a^4+a^2b^2+b^2y^2-2yb^3+b^4$$

$$= 2abxy$$

$$\Rightarrow a^2x^2-2abxy+b^2y^2-2a^3x-2b^3y$$

$$+ a^4+a^2b^2+b^4=0$$

$$\Rightarrow (ax-by)^2-2a^3x-2b^3y+a^4+a^2b^2+b^4=0$$

24. By using definition of parabola, we get

$$SP = PM$$

$$\Rightarrow SP^2 = PM^2$$

$$\Rightarrow x^2 + y^2 = \left(\frac{x+y-4}{\sqrt{2}} \right)^2$$

$$\Rightarrow 2x^2 + 2y^2 = x^2 + y^2 + 16 + 2xy - 8y - 8x$$

$$\Rightarrow x^2 + y^2 - 2xy + 8x + 8y - 16 = 0$$

25. $(13x-1)^2 + (13y-1)^2 = k(5x-12y+1)^2$

$$\Rightarrow 169 \left[\left(x - \frac{1}{13} \right)^2 + \left(y - \frac{1}{13} \right)^2 \right] = k(5x-12y+1)^2$$

Taking square root, we get

$$\sqrt{\left(x - \frac{1}{13} \right)^2 + \left(y - \frac{1}{13} \right)^2} = \frac{\sqrt{k}(5x-12y+1)}{13}$$

Now, condition for eccentricity is PS = ePM

$$\Rightarrow PS = \sqrt{k}PM$$

$$\Rightarrow \sqrt{k} = 1 \Rightarrow k = 1$$

26. $9x^2 - 6x + 36y + 9 = 0$

$$\Rightarrow x^2 - \frac{2}{3}x + 4y + 1 = 0$$


$$\Rightarrow x^2 - \frac{2}{3}x + \frac{1}{9} + 4y + 1 - \frac{1}{9} = 0$$

$$\Rightarrow \left(x - \frac{1}{3} \right)^2 = -4 \left(y + \frac{2}{9} \right)$$

Hence, the vertex is $\left(\frac{1}{3}, -\frac{2}{9} \right)$



27. $y^2 + 8x - 12y + 20 = 0$
 $\Rightarrow y^2 - 12y = -8x - 20$
 $\Rightarrow y^2 - 12y + 36 = -8x - 20 + 36$
 $\Rightarrow (y - 6)^2 = -8x + 16$
 $\Rightarrow (y - 6)^2 = -8(x - 2)$
 \Rightarrow Vertex is (2, 6)
28. The given equation can be written as $(x - 4)^2 = 1[y - (c - 16)]$
 Therefore, the vertex of the parabola is (4, c - 16).
 The point lies on X-axis.
 $\Rightarrow c - 16 = 0$
 $\Rightarrow c = 16$
29. Given, equation can be written as $y^2 = \frac{4k}{4} \left(x - \frac{8}{k}\right)$
 The standard equation of parabola is $y^2 = 4ax$
 $\Rightarrow a = \frac{k}{4}$
 \therefore Equation of directrix is $X + \frac{k}{4} = 0$
 $\Rightarrow x - \frac{8}{k} + \frac{k}{4} = 0$
 But the given equation of directrix is $x - 1 = 0$.
 Since, both equation are same
 $\therefore \frac{8}{k} - \frac{k}{4} = 1$
 $\Rightarrow 32 - k^2 = 4k \Rightarrow k = -8, 4$
30. Since, $9y^2 - 16x - 12y - 57 = 0$
 $\Rightarrow y^2 - \frac{16}{9}x - \frac{4}{3}y - \frac{57}{9} = 0$
 $\Rightarrow y^2 - \frac{4}{3}y + \frac{4}{9} = \frac{16}{9}x + \frac{57}{9} + \frac{4}{9}$
 $\Rightarrow \left(y - \frac{2}{3}\right)^2 = \frac{16}{9} \left(x + \frac{61}{16}\right)$
 this equation can be written as $Y^2 = 4\left(\frac{4}{9}\right)X$
 Axis of the parabola is $Y = 0$
 $\Rightarrow y - \frac{2}{3} = 0$
 $\Rightarrow 3y = 2$

31. Distance between focus and directrix is $= \left| \frac{3 - 4 - 2}{\sqrt{2}} \right| = \frac{3}{\sqrt{2}}$
 Hence, latus rectum is $3\sqrt{2}$
 ...[Since, latus rectum is two times the distance between focus and directrix].
32. Since, a = distance between tangent at vertex and latus rectum
 $\therefore a = \left| \frac{-8 - (-12)}{\sqrt{1+1}} \right| = \frac{4}{\sqrt{2}}$
 \therefore Length of latus rectum = $4a = 4 \times \frac{4}{\sqrt{2}} = 8\sqrt{2}$
33. $y^2 + 2Ax + 2By + C = 0$
 $\Rightarrow y^2 + 2By + B^2 = -2Ax - C + B^2$
 $\Rightarrow (y + B)^2 = -2A \left(x + \frac{C}{2A} - \frac{B^2}{2A}\right)$
 \therefore focus $\equiv \left(\frac{-C + B^2}{2A} - \frac{A}{2}, -B\right)$
 Equation of latus rectum is $x = -a$
 $= \frac{-C + B^2}{2A} - \frac{A}{2} = \frac{B^2 - A^2 - C}{2A}$
34. 
 According to the figure, the length of latus rectum is $2(SM) = 2 \times \frac{u^2}{2g} (1 + \cos 2\alpha) = \frac{2u^2 \cos^2 \alpha}{g}$
35. Given $t = -2$
 $2y^2 = 7x$
 $\Rightarrow y^2 = \frac{7}{2}x$
 Comparing with $y^2 = 4ax$, we get $4a = \frac{7}{2} \Rightarrow a = \frac{7}{8}$
 \therefore the point is $P = (at^2, 2at) = \left(\frac{7}{8} \times 4, 2 \times \frac{7}{8} \times (-2)\right)$
 $\therefore P = \left(\frac{7}{2}, \frac{-7}{2}\right)$



36. Here, $\frac{y}{2} = t$ and $x - 2 = t^2$

$$\Rightarrow (x - 2) = \left(\frac{y}{2}\right)^2 \Rightarrow y^2 = 4(x - 2)$$

37. $x = 2 + t^2$ and $y = 2t + 1$

$$\Rightarrow t^2 = (x - 2) \text{ and } t^2 = \left(\frac{y-1}{2}\right)^2 = \frac{(y-1)^2}{4}$$

$$\Rightarrow \frac{(y-1)^2}{4} = (x - 2) \Rightarrow (y - 1)^2 = 4(x - 2)$$

\therefore Vertex is (2, 1)

38. $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

Since, it passes through (-3, 1) and (2, -2)

$$\therefore \frac{9}{a^2} + \frac{1}{b^2} = 1 \text{ and } \frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{4}$$

$$\Rightarrow a^2 = \frac{32}{3}, b^2 = \frac{32}{5}$$

Hence, required equation of ellipse is $3x^2 + 5y^2 = 32$

39. Let point P (x_1, y_1)

$$\therefore \sqrt{(x_1 + 2)^2 + y_1^2} = \frac{2}{3} \left(x_1 + \frac{9}{2}\right)$$

$$\Rightarrow (x_1 + 2)^2 + y_1^2 = \frac{4}{9} \left(x_1 + \frac{9}{2}\right)^2$$

$$\Rightarrow 9(x_1^2 + y_1^2 + 4x_1 + 4) = 4\left(x_1^2 + \frac{81}{4} + 9x_1\right)$$

$$\Rightarrow 5x_1^2 + 9y_1^2 = 45 \Rightarrow \frac{x_1^2}{9} + \frac{y_1^2}{5} = 1$$

\therefore Locus of (x_1, y_1) is $\frac{x^2}{9} + \frac{y^2}{5} = 1$, which is equation of an ellipse.

40. Foci = (3, -3)

$$\Rightarrow ae = 3 - 2 = 1$$

Vertex = (4, -3)

$$\Rightarrow a = 4 - 2 = 2$$

$$\Rightarrow e = \frac{1}{2}$$

$$\Rightarrow b = 2 \sqrt{\left(1 - \frac{1}{4}\right)} = \frac{2}{2} \sqrt{3} = \sqrt{3}$$

Therefore, equation of ellipse with centre (2, -3) is

$$\frac{(x-2)^2}{4} + \frac{(y+3)^2}{3} = 1$$

41. Vertices ($\pm 5, 0$) $\equiv (\pm a, 0)$

$$\Rightarrow a = 5$$

Foci ($\pm 4, 0$) $\equiv (\pm ae, 0)$

$$\Rightarrow e = \frac{4}{5}$$

$$\therefore b = \sqrt{a^2(1 - e^2)} = \sqrt{25 \times \frac{9}{25}} = 3$$

Hence, equation is $\frac{x^2}{25} + \frac{y^2}{9} = 1$

$$\text{i.e., } 9x^2 + 25y^2 = 225$$

42. Foci ($\pm 5, 0$) $\equiv (\pm ae, 0)$,

$$\Rightarrow ae = 5 \quad \dots(i)$$

Equation of directrix is $x = \frac{a}{e}$

$$\text{Given, } x = \frac{36}{5}$$

$$\Rightarrow \frac{a}{e} = \frac{36}{5} \quad \dots(ii)$$

$$\Rightarrow a = 6 \text{ and } e = \frac{5}{6} \quad \dots[\text{From (i) and (ii)}]$$

$$\therefore b = a\sqrt{1 - e^2} = 6\sqrt{1 - \frac{25}{36}} = \sqrt{11}$$

Hence, equation is $\frac{x^2}{36} + \frac{y^2}{11} = 1$

43. Vertex (0,7) and directrix $y = 12$

$$\therefore b = 7 \text{ and } \frac{b}{e} = 12$$

$$\Rightarrow e = \frac{7}{12}$$

Also, $a = b\sqrt{1 - e^2}$

$$\Rightarrow a = 7 \sqrt{\frac{95}{144}}$$

$$\Rightarrow a^2 = \frac{4655}{144}$$

Hence, equation of ellipse is

$$\frac{x^2}{4655/144} + \frac{y^2}{49} = 1 \text{ i.e., } 144x^2 + 95y^2 = 4655$$

44. Let the equation of ellipse be $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

\therefore It passes through (-3, 1)

$$\therefore \frac{9}{a^2} + \frac{1}{b^2} = 1$$

$$\Rightarrow 9 + \frac{a^2}{b^2} = a^2 \quad \dots(i)$$



Given, eccentricity is $\sqrt{\frac{2}{5}}$

$$\therefore \frac{2}{5} = 1 - \frac{b^2}{a^2}$$

$$\Rightarrow \frac{b^2}{a^2} = \frac{3}{5} \quad \dots \text{(ii)}$$

From equation (i) and (ii), we get

$$a^2 = \frac{32}{3}, b^2 = \frac{32}{5}$$

Hence, required equation of ellipse is $3x^2 + 5y^2 = 32$

45. Given, $\frac{2b^2}{a} = 10 \Rightarrow b^2 = 5a$ and $2b = 2ae$

$$\Rightarrow \frac{b}{a} = e$$

$$\text{Also, } b^2 = a^2(1 - e^2)$$

$$\Rightarrow e^2 = (1 - e^2) \quad \dots \left[\because e = \frac{b}{a} \right]$$

$$\Rightarrow e = \frac{1}{\sqrt{2}}$$

$$\Rightarrow b = \frac{a}{\sqrt{2}} \text{ or } b = 5\sqrt{2}, a = 10$$

Hence, equation of ellipse is

$$\frac{x^2}{(10)^2} + \frac{y^2}{(5\sqrt{2})^2} = 1$$

$$\text{i.e., } x^2 + 2y^2 = 100$$

46. We have, $2ae = 8, \frac{2a}{e} = 18$

$$\Rightarrow ae \times \frac{a}{e} = 4 \times 9$$

$$\Rightarrow a = \sqrt{4 \times 9} = 6 \text{ and } e = \frac{2}{3}$$

$$\text{Also, } b = a \sqrt{1 - e^2}$$

$$\Rightarrow b = 6 \sqrt{1 - \frac{4}{9}} = \frac{6}{3} \sqrt{5} = 2\sqrt{5}$$

$$\text{Hence, the required equation is } \frac{x^2}{36} + \frac{y^2}{20} = 1$$

$$\text{i.e., } 5x^2 + 9y^2 = 180$$

47. $\frac{x^2}{2-r} + \frac{y^2}{r-5} + 1 = 0$

$$\Rightarrow \frac{x^2}{r-2} + \frac{y^2}{5-r} = 1$$

Hence, $r > 2$ and $r < 5$

$$\Rightarrow 2 < r < 5$$

48. $16x^2 + 25y^2 = 400$

$$\Rightarrow \frac{x^2}{25} + \frac{y^2}{16} = 1$$

$$\Rightarrow e = \sqrt{1 - \frac{16}{25}} = \frac{3}{5}$$

Therefore, directrices are $x = \pm \frac{5}{3}$ or $3x = \pm 25$

49. Given that, $\frac{a}{e} - ae = 8$ and $e = \frac{1}{2}$

$$\Rightarrow a = \frac{8e}{(1 - e^2)}$$

$$= \frac{8 \cdot \frac{1}{2}}{2(3)} = \frac{16}{3}$$

$$\therefore b = \frac{16}{3} \sqrt{1 - \frac{1}{4}}$$

$$= \frac{16}{3} \cdot \frac{\sqrt{3}}{2} = \frac{8\sqrt{3}}{3}$$

Hence, the length of minor axis is $\frac{16\sqrt{3}}{3}$

50. $3x^2 + 4y^2 = 48$

$$\Rightarrow \frac{x^2}{16} + \frac{y^2}{12} = 1$$

$$\therefore a^2 = 16, b^2 = 12$$

$$\Rightarrow e = \sqrt{1 - \frac{b^2}{a^2}} = \sqrt{1 - \frac{12}{16}} = \frac{1}{2}$$

$$\therefore \text{Distance between the foci} = 2ae$$

$$= 2 \times 4 \times \frac{1}{2}$$

$$= 4$$

51. We have, $\frac{2b^2}{a} = 2ae$

$$\Rightarrow b^2 = a^2e$$

$$\Rightarrow e = \frac{b^2}{a^2} \quad \dots \text{(i)}$$

$$\text{Also, } e = \sqrt{1 - \frac{b^2}{a^2}}$$

$$\Rightarrow e^2 = 1 - e$$

$$\Rightarrow e^2 + e - 1 = 0$$

$$\therefore e = \frac{-1 \pm \sqrt{5}}{2}$$

$$\therefore e = \frac{\sqrt{5} - 1}{2}$$

....[From (i)]



52. $a = 6, b = 2\sqrt{5}$
 Now, $b^2 = a^2(1 - e^2)$
 $\Rightarrow \frac{20}{36} = (1 - e^2)$
 $\Rightarrow e = \sqrt{\frac{16}{36}} = \frac{2}{3}$
 Distance between directrix = $\frac{2a}{e} = \frac{2 \times 6}{\frac{2}{3}} = 18$
53. Comparing $\frac{x^2}{16} + \frac{y^2}{9} = 1$ with $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$,
 we get
 $a^2 = 16, b^2 = 9$
 $\Rightarrow a = 4, b = 3$
 Now, $b^2 = a^2(1 - e^2)$
 $\Rightarrow \frac{9}{16} = 1 - e^2$
 $\Rightarrow e^2 = 1 - \frac{9}{16} = \frac{7}{16}$
 $\therefore e = \frac{\sqrt{7}}{4}$
 $\therefore S = (ae, 0) = (\sqrt{7}, 0)$
 \therefore radius of the circle = $\sqrt{(0 - \sqrt{7})^2 + (3 - 0)^2}$
 $= \sqrt{16} = 4$
54. $x = 3(\cos t + \sin t), y = 4(\cos t - \sin t)$
 $\Rightarrow \frac{x}{3} = \cos t + \sin t, \frac{y}{4} = \cos t - \sin t$
 $\Rightarrow \frac{x^2}{9} = 1 + \sin 2t, \frac{y^2}{16} = 1 - \sin 2t$
 $\Rightarrow \frac{x^2}{9} + \frac{y^2}{16} = 2$
 $\Rightarrow \frac{x^2}{18} + \frac{y^2}{32} = 1$ which is an ellipse.
55. $9x^2 + 4y^2 - 6x + 4y + 1 = 0$
 $\Rightarrow 9\left(x^2 - \frac{2}{3}x + \frac{1}{9}\right) + 4\left(y^2 + y + \frac{1}{4}\right) = -1 + 1 + 1$
 $\Rightarrow 9\left(x - \frac{1}{3}\right)^2 + 4\left(y + \frac{1}{2}\right)^2 = 1$
 $\Rightarrow \frac{\left(x - \frac{1}{3}\right)^2}{\frac{1}{9}} + \frac{\left(y + \frac{1}{2}\right)^2}{\frac{1}{4}} = 1$

$$\text{Here, } a = \frac{1}{3}, b = \frac{1}{2}$$

$$\Rightarrow 2a = \frac{2}{3}, 2b = 1$$

\therefore Length of axes are $1, \frac{2}{3}$.

56. $3x^2 - 12x + 4y^2 - 8y = -4$
 $\Rightarrow 3(x^2 - 4x + 4) + 4(y^2 - 2y + 1) = -4 + 12 + 4$
 $\Rightarrow 3(x - 2)^2 + 4(y - 1)^2 = 12$
 $\Rightarrow \frac{(x - 2)^2}{4} + \frac{(y - 1)^2}{3} = 1$

$$\Rightarrow \frac{X^2}{4} + \frac{Y^2}{3} = 1$$

$$\therefore e = \sqrt{1 - \frac{3}{4}} = \frac{1}{2}$$

Foci are $x = \pm ae, y = 0$
 $\Rightarrow x - 2 = \pm 1, y - 1 = 0$
 $\Rightarrow x = 3$ or $1, y = 1$
 i.e., $(3, 1)$ and $(1, 1)$

57. $4(x^2 + 2x + 1) + 9(y^2 + 4y + 4) = 36$
 $\Rightarrow 4(x + 1)^2 + 9(y + 2)^2 = 36$
 $\Rightarrow \frac{(x + 1)^2}{9} + \frac{(y + 2)^2}{4} = 1$

Comparing with standard form, we get
 $a^2 = 9, b^2 = 4$

Now, condition for eccentricity is
 $b^2 = a^2(1 - e^2)$

$$\Rightarrow 4 = 9(1 - e^2)$$

$$\Rightarrow e^2 = 1 - \frac{4}{9} = \frac{5}{9}$$

$$\Rightarrow e = \frac{\sqrt{5}}{3}$$

58. Given, $2a = 6, 2b = 4$
 i.e., $a = 3, b = 2$

$$\text{Also, } e^2 = 1 - \frac{b^2}{a^2} = \frac{5}{9}$$

$$\Rightarrow e = \frac{\sqrt{5}}{3}$$

\therefore Distance between the pins = $2ae = 2\sqrt{5}$ cm
 and length of string = $2a + 2ae = 6 + 2\sqrt{5}$ cm



59. $2a = 7$ or $a = \frac{7}{2}$
 Also $(5, -2)$ satisfies $\frac{4}{49}(25) - \frac{51}{196}(4) = 1$
 and $a^2 = \frac{49}{4}$
 $\Rightarrow a = \frac{7}{2}$
 \therefore option (C) is correct answer
60. Centre $(0, 0)$, vertex $(4, 0) \Rightarrow a = 4$ and focus $(6, 0)$
 $\Rightarrow ae = 6$
 $\Rightarrow e = \frac{3}{2}$
 Also, $b^2 = a^2(e^2 - 1)$
 $= 20$
 Hence, required equation is $\frac{x^2}{16} - \frac{y^2}{20} = 1$
 i.e., $5x^2 - 4y^2 = 80$
61. Given that, $e = \frac{3}{2}$
 foci $= (\pm 2, 0) = (\pm ae, 0)$
 $\Rightarrow ae = 2$
 $\Rightarrow a = \frac{4}{3}$
 $\Rightarrow a^2 = \frac{16}{9}$
 Now, condition for eccentricity is $b^2 = a^2(e^2 - 1)$
 $\therefore b^2 = \frac{16}{9} \left(\frac{9}{4} - 1 \right) = \frac{16}{9} \left(\frac{5}{4} \right) = \frac{20}{9}$
 Now, equation of hyperbola is $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$
 $\Rightarrow \frac{9x^2}{16} - \frac{9y^2}{20} = 1 \Rightarrow \frac{x^2}{4} - \frac{y^2}{5} = \frac{4}{9}$
62. Given, $\frac{2b^2}{a} = 8$ and $\frac{3}{\sqrt{5}} = \sqrt{1 + \frac{b^2}{a^2}}$
 $\Rightarrow \frac{4}{5} = \frac{b^2}{a^2}$
 $\Rightarrow a = 5, b = 2\sqrt{5}$
 Hence, the required equation of hyperbola is $\frac{x^2}{25} - \frac{y^2}{20} = 1$
 i.e., $4x^2 - 5y^2 = 100$

63. Conjugate axis is 5 and distance between foci = 13
 $\Rightarrow 2b = 5$ and $2ae = 13$
 Also, $b^2 = a^2(e^2 - 1)$
 $\Rightarrow \frac{25}{4} = \frac{(13)^2}{4e^2} (e^2 - 1)$
 $\Rightarrow \frac{25}{4} = \frac{169}{4} - \frac{169}{4e^2}$
 $\Rightarrow e = \frac{13}{12}$
 $\Rightarrow a = 6, b = \frac{5}{2}$
 Hence, the required equation of hyperbola is $\frac{x^2}{36} - \frac{y^2}{25} = 1$
 i.e., $25x^2 - 144y^2 = 900$
64. Equation of hyperbola passes through (x_1, y_1)
 $\Rightarrow \frac{x_1^2}{a^2} - \frac{y_1^2}{b^2} = 1 \Rightarrow \frac{x_1^2}{a^2} - 1 = \frac{y_1^2}{b^2}$
 $\Rightarrow \frac{x_1^2 - a^2}{a^2} = \frac{y_1^2}{b^2} \Rightarrow \frac{b^2}{a^2} = \frac{y_1^2}{x_1^2 - a^2}$
 Now, $\frac{b^2}{a^2} = e^2 - 1$
 $\Rightarrow \frac{y_1^2}{x_1^2 - a^2} = e^2 - 1$
 $\Rightarrow e^2 = \frac{y_1^2 + (x_1^2 - a^2)}{x_1^2 - a^2}$
 $\Rightarrow e = \sqrt{\frac{x_1^2 - a^2 + y_1^2}{x_1^2 - a^2}} = \sqrt{\frac{a^2 - x_1^2 - y_1^2}{a^2 - x_1^2}}$
65. Center of the hyperbola is midpoint of foci.
 Hence, its center is $(1, 5)$ also distance between foci is $2ae = 10$
 $\Rightarrow a = 4 \quad \dots \left[\because e = \frac{5}{4} \right]$
 $\Rightarrow a^2 = 16$
 Now, $b^2 = a^2(e^2 - 1)$
 $= a^2e^2 - a^2 = 25 - 16 \Rightarrow b^2 = 9$
 Hence, equation of hyperbola is $\frac{(x-1)^2}{16} - \frac{(y-5)^2}{9} = 1$



66. Let $S(1, 1)$ be the focus and $P(x, y)$ be a point on the hyperbola
Now, $PS = ePM$

$$\Rightarrow \sqrt{(x-1)^2 + (y-1)^2} = \sqrt{3} \left| \frac{2x+y-1}{\sqrt{2^2+1^2}} \right|$$

Squaring both sides, we get

$$(x-1)^2 + (y-1)^2 = \frac{3}{5} (2x+y-1)^2$$

On simplification, the required equation is

$$7x^2 + 12xy - 2y^2 - 2x + 4y - 7 = 0$$

67. For ellipse, $e < 1$ and also $e' < 1$

$$\therefore e^2 + e'^2 < 2$$

For parabola, $e = 1$ and $e' = 1$

$$\therefore e^2 + e'^2 = 2$$

For hyperbola, $e > 1$ and $e' > 1$

$$\therefore e^2 + e'^2 > 2$$

Hence, it can be 3 in case of hyperbola.

68. $96x^2 - 16y^2 - 36x + 96y - 252 = 0$

$$\Rightarrow \frac{(x-2)^2}{16} - \frac{(y-3)^2}{9} = 1$$

$$\Rightarrow \frac{X^2}{16} - \frac{Y^2}{9} = 1$$

\therefore Vertices are $(X = \pm a, Y = 0)$

i.e., $(x-2 = \pm 4, y-3 = 0)$

\therefore The vertices of the hyperbola are $(6, 3)$ and $(-2, 3)$

69. $(x+1)^2 - y^2 - 1 + 5 = 0$

$$\Rightarrow -\frac{(x+1)^2}{4} + \frac{y^2}{4} = 1$$

Equation of directrices of $\frac{y^2}{b^2} - \frac{x^2}{a^2} = 1$ are

$$y = \pm \frac{b}{e}$$

Here, $b = 2$, $e = \sqrt{1+1} = \sqrt{2}$

$$\text{Hence, } y = \pm \frac{2}{\sqrt{2}}$$

$$\Rightarrow y = \pm \sqrt{2}$$

70. Given, equation of hyperbola is

$$9x^2 - 16y^2 + 72x - 32y - 16 = 0$$

$$\Rightarrow 9(x^2 + 8x) - 16(y^2 + 2y) - 16 = 0$$

$$\Rightarrow 9(x+4)^2 - 16(y+1)^2 = 144$$

$$\Rightarrow \frac{(x+4)^2}{16} - \frac{(y+1)^2}{9} = 1$$

\therefore Latus rectum = $\frac{2b^2}{a} = 2 \times \frac{9}{4} = \frac{9}{2}$

71. Squaring and subtracting, we get
 $a^2x^2 - b^2y^2 = a^2 - b^2$, which is the equation of hyperbola.

72. $2x = t + \frac{1}{t}$ and $2y = t - \frac{1}{t}$

$$\Rightarrow 4x^2 = t^2 + 2 + \frac{1}{t^2} \quad \dots(i)$$

$$\text{and } 4y^2 = t^2 - 2 + \frac{1}{t^2} \quad \dots(ii)$$

Subtracting (ii) from (i), we get

$$4x^2 - 4y^2 = 4 \Rightarrow x^2 - y^2 = 1$$

The equation is of hyperbola.

73. Multiplying both, we get

$$(bx)^2 - (ay)^2 = (ab)^2$$

$$\Rightarrow \frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

which is the standard equation of hyperbola.

74. $e^2 = 1 - \frac{b^2}{a^2} = \frac{4}{9}$

$$\Rightarrow e = \frac{2}{3}$$

$$\text{and } e'^2 = 1 + \frac{b^2}{a^2} = 1 + \frac{4}{9} = \frac{9}{4}$$

$$\Rightarrow e' = \frac{3}{2}$$

$$\therefore ee' = 1$$

75. Centres of the circles are $(0, 0)$, $(-3, 1)$ and $(6, -2)$ respectively. Line passing through any

two points say $(0, 0)$ and $(-3, 1)$ is $y = -\frac{1}{3}x$

point $(6, -2)$ lies on it.

Hence, points are collinear.



Competitive Thinking

1. Radius = Distance from origin = $\sqrt{\alpha^2 + \beta^2}$

$$\therefore (x - \alpha)^2 + (y - \beta)^2 = \alpha^2 + \beta^2$$

$$\Rightarrow x^2 + y^2 - 2\alpha x - 2\beta y = 0$$

2. Centre $(2, 2)$ and

$$r = \sqrt{(4-2)^2 + (5-2)^2}$$

$$= \sqrt{13}$$

Hence, required equation is

$$(x-2)^2 + (y-2)^2 = (\sqrt{13})^2$$

$$\Rightarrow x^2 + y^2 - 4x - 4y - 5 = 0$$



3. Let r be the radius of the circle.

Given, circumference = 10π

$$\therefore 2\pi r = 10\pi$$

$$\Rightarrow r = 5$$

\therefore the equation of the circle is

$$(x-2)^2 + (y+3)^2 = 5^2$$

$$\Rightarrow x^2 + y^2 - 4x + 6y - 12 = 0$$


4. The centre of the circle which touches each axis in first quadrant at a distance 5, will be (5, 5) and radius will be 5.

\therefore equation of the circle is

$$(x-5)^2 + (y-5)^2 = (5)^2$$

$$\Rightarrow x^2 + y^2 - 10x - 10y + 25 = 0$$

5. Since, circle touches the co-ordinate axes in III quadrant.



\therefore Radius = $-h = -k$

Hence, $h = k = -5$

\therefore Equation of circle is $(x+5)^2 + (y+5)^2 = 25$

6. Since, the circle touches

X-axis at (3, 0).

\therefore centre of the circle is (3, k).

Now, $CA^2 = CB^2$

$$\therefore (3-3)^2 + (k-0)^2$$

$$= (3-1)^2 + (k-4)^2$$

$$\therefore k^2 = 4 + k^2 - 8k + 16$$

$$\therefore k = \frac{5}{2}$$

\therefore the required equation of circle is

$$(x-3)^2 + \left(y - \frac{5}{2}\right)^2 = \left(\frac{5}{2}\right)^2$$

$$\Rightarrow x^2 + y^2 - 6x - 5y + 9 = 0$$


7. Since, the circle has centre at (1, 2) and line $x = y$ i.e. $x - y = 0$ as tangent,

$$\therefore \text{Radius of circle} = \frac{|(1)-(2)|}{\sqrt{(1)^2 + (-1)^2}}$$

$$= \frac{1}{\sqrt{2}}$$

$$\therefore \text{Area of circle} = \pi \times \left(\frac{1}{\sqrt{2}}\right)^2 = \frac{\pi}{2}$$

8. Centre is $\left(\frac{a}{2}, \frac{b}{2}\right)$ and radius = $\sqrt{\frac{a^2 + b^2}{4}}$



Hence, equation of circle is

$$x^2 + y^2 - ax - by = 0$$

9. diameter = $\sqrt{[4 - (-2)]^2 + [7 - (-1)]^2}$

$$= \sqrt{6^2 + 8^2}$$

$$= 10$$

$$\text{Centre} = \left(\frac{4 + (-2)}{2}, \frac{7 + (-1)}{2}\right) = (1, 3)$$

\therefore equation of the circle is

$$(x-1)^2 + (y-3)^2 = (5)^2$$

If this circle cuts X-axis, then

$$(x-1)^2 + (0-3)^2 = 25$$

$$\Rightarrow (x-1)^2 = 16$$

$$\Rightarrow (x-1) = \pm 4$$

$$\Rightarrow x = 5, -3$$

\therefore the points on X-axis are A(5, 0) and B(-3, 0)

$$\therefore AB = \sqrt{(5+3)^2 + 0^2} = 8$$

10. Since, the centre always lies on the diameter.

Solving $2x + 3y + 1 = 0$ and $3x - y - 4 = 0$,

the co-ordinates of the centre are (1, -1).

Given, circumference = 10π

$$\therefore 2\pi r = 10\pi \Rightarrow r = 5$$

\therefore the equation of the circle is

$$(x-1)^2 + (y+1)^2 = 5^2$$

$$\Rightarrow x^2 + y^2 - 2x + 2y - 23 = 0$$

11. Centre of circle = Point of intersection of diameters = (1, -1)

Now, area = 154

$$\Rightarrow \pi r^2 = 154 \Rightarrow r = 7$$

Hence, the equation of required circle is

$$(x-1)^2 + (y+1)^2 = 7^2$$

$$\Rightarrow x^2 + y^2 - 2x + 2y = 47$$

$$12. \quad 3x - y - 4 = 0 \quad \dots(i)$$

$$x + 3y + 2 = 0 \quad \dots(ii)$$

solving equation (i) and (ii), we get

$$(x, y) = (1, -1)$$

Since, $\pi r^2 = 154$

$$\therefore r = 7$$



∴ Equation of required circle is
 $(x-1)^2 + (y+1)^2 = 49$
 $\Rightarrow x^2 + y^2 - 2x + 2y - 47 = 0$

13. Since, the centre always lies on the diameter.
 Solving $2x + 3y = 3$ and $16x - y = 4$, we get
 co-ordinates of the centre = $\left(\frac{3}{10}, \frac{4}{5}\right)$.

The circle passes through (4, 6).

$$\begin{aligned} \therefore r^2 &= \left(4 - \frac{3}{10}\right)^2 + \left(6 - \frac{4}{5}\right)^2 \\ &= \left(\frac{37}{10}\right)^2 + \left(\frac{26}{5}\right)^2 = \frac{4073}{100} \end{aligned}$$

∴ the equation of the circle is
 $\left(x - \frac{3}{10}\right)^2 + \left(y - \frac{4}{5}\right)^2 = \frac{4073}{100}$
 $\Rightarrow 100x^2 + 100y^2 - 60x - 160y = 4000$
 $\Rightarrow 5(x^2 + y^2) - 3x - 8y = 200$

14. Let centre be (h, k). Then,
 $\sqrt{(h-2)^2 + (k-3)^2} = \sqrt{(h-4)^2 + (k-5)^2}$
 $\Rightarrow -4h + 4 - 6k + 9 = -8h + 16 - 10k + 25$
 $\Rightarrow 4h + 4k - 28 = 0$
 $\Rightarrow h + k - 7 = 0 \quad \dots(i)$


Since, centre lies on the given line.

∴ $k - 4h + 3 = 0 \quad \dots(ii)$

Solving (i) and (ii), we get
 (h, k) = (2, 5)

∴ centre is (2, 5) and
 radius = $\sqrt{(2-2)^2 + (5-3)^2} = 2$

∴ the required equation of the circle is
 $(x-2)^2 + (y-5)^2 = (2)^2$
 $\Rightarrow x^2 + y^2 - 4x - 10y + 25 = 0$



Let AB be the chord cut off by the circle on the line $2x - 5y + 18 = 0$.

Let CD be the perpendicular drawn from centre (3, -1) to AB.


$$\therefore CD = \frac{|2(3) - 5(-1) + 18|}{\sqrt{2^2 + (-5)^2}} = \sqrt{29}$$

and AD = 3

$$\therefore CA^2 = AD^2 + CD^2 = 3^2 + (\sqrt{29})^2 = 38$$

∴ the equation of the circle is
 $(x-3)^2 + (y+1)^2 = 38$.

16. ΔABC is equilateral.
 ∴ O(0, 0) is the centroid.



O divides AD in the ratio 2 : 1


$$\begin{aligned} \therefore \frac{AO}{OD} &= \frac{2}{1} \\ \Rightarrow \frac{AO}{AD - AO} &= \frac{2}{1} \Rightarrow \frac{AO}{9 - AO} = \frac{2}{1} \\ \Rightarrow AO &= 18 - 2AO \Rightarrow AO = 6 \text{ units} \end{aligned}$$

∴ radius = 6 units

∴ equation of circle is $x^2 + y^2 = 36$.

17. According to the figure, A(0, 0), B(a, 0), C(a, a) and D(0, a).

and centre is $\left(\frac{a}{2}, \frac{a}{2}\right)$.



∴ the equation of the circle is


$$\left(x - \frac{a}{2}\right)^2 + \left(y - \frac{a}{2}\right)^2 = \frac{a^2}{2}$$

$$\Rightarrow x^2 + y^2 - ax - ay = 0$$

18. Since, the circle touches Y-axis at (0, 2).
 ∴ centre of the circle is (h, 2).

Now, $CA^2 = CB^2$

$$\begin{aligned} \therefore (h-0)^2 + (2-2)^2 &= (h-(-1))^2 + (2-0)^2 \\ &= (h+1)^2 + (2-0)^2 \\ \Rightarrow h^2 &= h^2 + 2h + 1 + 4 \\ \Rightarrow 2h + 5 &= 0 \Rightarrow h = -\frac{5}{2} \end{aligned}$$



∴ equation of circle is

$$\left(x + \frac{5}{2}\right)^2 + (y-2)^2 = \left(-\frac{5}{2}\right)^2$$

$$\Rightarrow x^2 + \frac{25}{4} + 5x + y^2 - 4y + 4 = \frac{25}{4}$$


$$\Rightarrow x^2 + y^2 + 5x - 4y + 4 = 0$$

Point (-4, 0) satisfies this equation.

∴ option (D) is the correct answer.



19.



Since, the circle touches X-axis at (1, 0).

∴ centre of the circle is (1, k) and radius = k
 ∴ equation of the circle is $(x-1)^2 + (y-k)^2 = k^2$
 $\Rightarrow 1 + k^2 - 6k + 9 = k^2$
 $\Rightarrow k = \frac{5}{3}$

∴ diameter = $2k = \frac{10}{3}$

20. The equation of circle touching the coordinate axes with centre (a, a) and radius 'a' is
 $x^2 + y^2 - 2ax - 2ay + a^2 = 0$... (i)

Since, line $3x - 4y - 12 = 0$ touches the circle

∴ perpendicular distance from centre of the circle to the line = radius

∴ $\frac{|3(a) - 4(a) - 12|}{\sqrt{9+16}} = a$

$\Rightarrow a = 3$

Substituting, $a = 3$ in equation (i), we get

$x^2 + y^2 - 6x - 6y + 9 = 0$

This is the required equation of the circle.

21. The given equation represents a circle having line segment joining (x_1, y_1) and (x_2, y_2) as a diameter.

∴ the coordinates of its centre are $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$.

22. By diameter form, the required equation is

$(x - x_1)(x - x_2) + (y - y_1)(y - y_2) = 0$

∴ $(x + 4)(x - 12) + (y - 3)(y + 1) = 0$

∴ $x^2 + y^2 - 8x - 2y - 51 = 0$


23. Circle whose diametric end points are (1, 0) and (0, 1) will be of smallest radius.

∴ By using diameter form, equation of circle is

$(x - 1)(x - 0) + (y - 0)(y - 1) = 0$

$\Rightarrow x^2 + y^2 - x - y = 0$

24.



Here, the diagonals AC and BD of rectangle ABCD are diameters of the circle passing through the vertices A, B, C and D. Considering diagonal AC with end points A(a, b) and C(-a, -b), we get

Equation of circle in diameter form as,
 $(x - a)(x - (-a)) + (y - b)(y - (-b)) = 0$

∴ $x^2 - a^2 - y^2 - b^2 = 0$

∴ $x^2 + y^2 = a^2 + b^2$

26. The given equation represents a circle, if coeff. of $x^2 =$ coeff. of y^2 and coeff. of $xy = 0$

∴ $a = 2$ and $b = 0$

Also, it passes through origin.

∴ $c = 0$

27. Here, $g = 2, f = 3$ and $c = 13$

∴ $r = \sqrt{g^2 + f^2 - c}$

∴ $r = \sqrt{4 + 9 - 13} = 0$

∴ option (D) is the correct answer.

28. Consider option (A),

$x^2 + y^2 + 2ax + 2by + 2b^2 = 0$

Centre = $(-a, -b)$

∴ option (A) is the correct answer.

29. The given equation represents a circle, if coeff. of $x^2 =$ coeff. of y^2

After solving the given equation, we get

$\frac{K}{3} = \frac{1}{4} \Rightarrow K = \frac{3}{4}$

30. The given equation represents a circle, if coeff. of $xy = 0$.

∴ $2k - 1 = 0 \Rightarrow k = \frac{1}{2}$

radius = $\sqrt{(-1)^2 + (2)^2 - 3} = \sqrt{2}$



31. The given equation represents a circle, if coeff. of $xy = 0$.

$$\therefore h = 0$$

$$\text{and } \sqrt{(-3)^2 + (-1)^2 - k} = 2$$

$$\Rightarrow 10 - k = 4$$

$$\Rightarrow k = 6$$

32. Circle $x^2 + y^2 + 2gx + 2fy + c = 0$ is concentric with $x^2 + y^2 - 2x + 4y + 20 = 0$.

\therefore centre is $(1, -2)$ and

$$\text{radius} = \sqrt{(4-1)^2 + (-2+2)^2} = \sqrt{3^2 + 0^2} = 3$$

$$\text{Also, } r = \sqrt{g^2 + f^2 - c}$$

$$\therefore 3 = \sqrt{(-1)^2 + (2)^2 - c}$$

$$\therefore 9 = 1 + 4 - c$$

$$\therefore c = -4$$

33. Here, $g = \frac{-1}{4}$, $f = 0$ and $c = 0$

$$\therefore \text{centre } (-g, -f) = \left(\frac{1}{4}, 0\right)$$

$$\text{and } r = \sqrt{\frac{1}{16} + 0 - 0} = \frac{1}{4}$$

34. Let another end of the diameter be (x, y) .

Centre of the given circle is $(2, 3)$.


Since, centre is the midpoint of the diameter

$$\therefore 2 = \frac{3+x}{2}, 3 = \frac{4+y}{2}$$

$$\Rightarrow x = 1, y = 2$$

$$\Rightarrow (x, y) = (1, 2)$$

35. Let $A(x, y)$ be the required point.



given equation of circle is

$$x^2 + y^2 + 2x + 4y - 3 = 0$$

\therefore Centre = $(-1, -2)$

Since, C is the midpoint of AP.

$$\therefore A = (-3, -4)$$

36. Here, the centre of circle $(3, -1)$ must lie on the line $x + 2by + 7 = 0$.

$$\therefore 3 - 2b + 7 = 0$$

$$\Rightarrow b = 5$$

37. Given equation of circle is


$$x^2 + y^2 - 4x - 6y + 9 = 0$$

$$\Rightarrow x^2 - 4x + 4 + y^2 - 6y + 9 - 4 = 0$$

$$\Rightarrow (x - 2)^2 + (y - 3)^2 = 4$$

\therefore centre = $(2, 3)$, radius = 2

The diameter of this circle is a chord of circle with centre $O(1, 1)$.




$$OP = \sqrt{(3-1)^2 + (2-1)^2} = \sqrt{5}$$

$$QP = 2$$

$$\therefore r^2 = (\sqrt{5})^2 + 2^2 \Rightarrow r = 3$$

38.



Given circle $S_1, x^2 + y^2 + 4x + 6y - 12 = 0$

\therefore Its centre $C_1 = (-2, -3)$

radius, $r_1 = 5$

centre of circle S, is $C = (2, -3)$

From the figure, we have

Diameter, $x = -2$ of circle S_1 is the chord of circle S.

\therefore In $\triangle CC_1A$,

$C_1A = r = 5$ unit

$$CC_1 = \sqrt{(-2-2)^2 + (-3-3)^2} = 4 \text{ unit}$$


$$\therefore (CA)^2 = (CC_1)^2 + (C_1A)^2 = (4)^2 + (5)^2 = 16 + 25 = 41$$

$$\therefore CA = \sqrt{41} \text{ unit}$$

$$\therefore \text{Radius of circle S is } \sqrt{41} \text{ unit.}$$



39. Consider option (A)
 $x^2 + y^2 + 8x + 2y - 8 = 0$
 Point $(-1, 3)$ is common to both circle and lies on above circle also
 Since, point $(-1, 3)$ satisfies the equation of circle in option (A)
 \therefore correct answer is option (A)
40. $C_1 : x^2 + y^2 - 6x = 0$ (i)
 $C_2 : x^2 + y^2 - 6y = 0$ (ii)
 Solving (i) and (ii), we get
 $x = y$ (iii)
 Substituting (iii) in (i), we get
 $y = 3$
 $\therefore x = 3$
 Point on circle is $P(3, 3)$ and
 centre = $\left(\frac{3}{2}, \frac{3}{2}\right)$
 \therefore Radius = $\sqrt{\left(3 - \frac{3}{2}\right)^2 + \left(3 - \frac{3}{2}\right)^2}$
 $= \frac{3}{\sqrt{2}}$
 \therefore equation of the circle is
 $\left(x - \frac{3}{2}\right)^2 + \left(y - \frac{3}{2}\right)^2 = \frac{9}{2}$
 $\Rightarrow x^2 + y^2 - 3x - 3y = 0$
41. Putting $y = x$ in $x^2 + y^2 - 2x = 0$, we get
 $2x^2 - 2x = 0$
 $\Rightarrow x(x - 1) = 0 \Rightarrow x = 0$ or $x = 1$
 $\therefore y = 0$ or $y = 1$
 \therefore Points of intersection are $(0, 0)$ and $(1, 1)$.
 These are end points of a diameter of required circle.
 \therefore equation of required circle is
 $(x - 0)(x - 1) + (y - 0)(y - 1) = 0$
 $\Rightarrow x^2 + y^2 - x - y = 0$
42. The centres of two circles are $C_1(1, 0)$ and $C_2(-2, -4)$ and their radii are 1 and 2 units respectively.
 Let C be the centre of the required circle.
 Then, $CP = CQ = 1$.
 $\therefore CC_1 = 2$ and $CC_2 = 3$.
 Clearly, C divides $C_1 C_2$ in the ratio $2 : 3$.
 Therefore, coordinates of C are
 $\left(\frac{-4 + 3}{2 + 3}, \frac{-8 + 0}{2 + 3}\right) = \left(-\frac{1}{5}, -\frac{8}{5}\right)$.




Hence, equation of the required circle is

$$\left(x + \frac{1}{5}\right)^2 + \left(y + \frac{8}{5}\right)^2 = 1^2$$

$$\Rightarrow 5x^2 + 5y^2 + 2x + 16y + 8 = 0$$

43. Let, $P = (x, y)$
 \therefore according to the given condition
 $\frac{\sqrt{x^2 + y^2 - 2x + 4y - 20}}{\sqrt{x^2 + y^2 - 2x - 8y + 1}} = \frac{2}{1}$
 $\Rightarrow \frac{x^2 + y^2 - 2x + 4y - 20}{x^2 + y^2 - 2x - 8y + 1} = 4$
 $\Rightarrow x^2 + y^2 - 2x - 12y + 8 = 0$
44. Equation of the tangent at $(1, 7)$ to $x^2 = y - 6$ is
 $2x - y + 5 = 0$
 Centre of the given circle is $(-8, -6)$.



Perpendicular from the centre $(-8, -6)$ to $2x - y + 5 = 0$ is equal to the radius of the circle.

$$\Rightarrow \left| \frac{2(-8) - (-6) + 5}{\sqrt{2^2 + 1^2}} \right| = \sqrt{8^2 + 6^2 - c}$$


$$\Rightarrow \left| \frac{-5}{\sqrt{5}} \right| = \sqrt{100 - c} \Rightarrow c = 95$$

45. Let, $x^2 + y^2 + 2gx + 2fy + c = 0$ be the required circle.
 This circle passes through $(1, 0)$
 $\therefore 1 + 2g + c = 0$
 i.e. $2g + c = -1$ (i)
 Also, this circle is orthogonal to the circle $x^2 + y^2 - 2x + 4y + 1 = 0$ and $x^2 + y^2 + 6x - 2y + 1 = 0$
 $\therefore 2g + 4f + c = -1$ (ii), and
 $6g - f - c = 1$ (iii)
 Solving, (i), (ii) and (iii), we get
 $g = f = 0$ and $c = -1$
 \therefore Centre = $(-g, -f) = (0, 0)$



46. $x^2 + y^2 - 4x - 6y - 12 = 0$
 $C_1 = (2, 3), r = \sqrt{2^2 + 3^2 + 12} = 5$
 $x^2 + y^2 + 6x + 18y + 26 = 0$
 $C_2 = (-3, -9), r = \sqrt{9 + 81 - 26} = \sqrt{64} = 8$
 $(C_1 C_2) = r_1 + r_2$
 \therefore The circles touch externally at a single point
 \therefore Number of common tangents is 3.

47.



- $\therefore SP^2 = PM^2$
 $\Rightarrow (x-3)^2 + y^2 = \left| \frac{x+3}{\sqrt{1}} \right|^2$
 $\Rightarrow x^2 + 9 - 6x + y^2 = x^2 + 9 + 6x$
 $\Rightarrow y^2 = 12x$
48. $\sqrt{(x-1)^2 + (y+1)^2} = \left| \frac{x_1 + y_1 + 3}{\sqrt{1+1}} \right|$
 $x^2 + y^2 - 10x - 2y - 2xy - 5 = 0$
49. Equation of parabola having vertex (p, q) and focus $(p, b+q)$ is given by;
 $(x-p)^2 = 4b(y-q)$
 Given, vertex $A = (1, 1)$ and focus $S = (1, -1)$
 $\therefore p = 1, q = 1, b = -2$
 \therefore Equation of parabola is;
 $(x-1)^2 = 4(-2)(y-1)$
 i.e. $x^2 - 2x + 8y - 7 = 0$
 Only $\left(3, \frac{1}{2}\right)$ satisfies the above equation of parabola.

50. $a = 4$, vertex = $(0, 0)$, focus = $(0, -4)$
51. Given, equation is $x^2 = -8ay$
 Here, $A = 2a$
 Focus of parabola $(0, -A)$ i.e., $(0, -2a)$
 Directrix $y = A$ i.e., $y = 2a$
52. $y^2 - 4y - x + 3 = 0$
 $\Rightarrow y^2 - 4y + 4 - x + 3 - 4 = 0$
 $\Rightarrow (y-2)^2 - (x+1) = 0$
 $\Rightarrow (y-2)^2 = (x+1)$

Comparing with $Y^2 = 4aX$, we have

$$a = \frac{1}{4}, Y = y - 2, X = x + 1$$

Focus of the parabola is $Y = 0, X = a$

$$\Rightarrow y - 2 = 0, x + 1 = \frac{1}{4} \Rightarrow y = 2, x = \frac{-3}{4}$$

$$\therefore \text{focus} = \left(\frac{-3}{4}, 2 \right)$$

53. $(y+1)^2 = -8(x+2)$

Comparing this equation with $Y^2 = -4aX$, we get

$$a = 2, X = x + 2 \text{ and } Y = y + 1$$

Focus of the parabola is,

$$X = -a, Y = 0$$

$$\Rightarrow x + 2 = -2, y + 1 = 0 \Rightarrow x = -4, y = -1$$

$$\therefore \text{focus} = (-4, -1)$$

54. Parabola is $y^2 = -4ax$ (left handed parabola).

\therefore its focus is $(-a, 0)$.

\therefore option (B) is false.

56. Since, $(a, -2a) \equiv (2, -8)$

\therefore another end $(a, 2a) \equiv (2, 8)$

57. Given $y^2 = 5x$

here $(x_1, y_1) = (a, 2a)$ and $(x_2, y_2) = (a, -2a)$

$$\therefore x_1 x_2 = a^2 = \left(\frac{25}{16} \right)$$

$$\therefore a = \frac{5}{4}$$


$$y_1 y_2 = -4a^2 = \frac{-25}{4}$$

$$\therefore 4x_1 x_2 + y_1 y_2 = 0$$

58. $x^2 = 12y$

$$\Rightarrow 4a = 12$$

$$\Rightarrow a = 3$$



$$\text{Area of triangle} = \frac{1}{2} (\text{base}) (\text{height})$$

$$= \frac{1}{2} \times AB \times OS = \frac{1}{2} \times 4a \times a$$

$$= \frac{1}{2} (12)(3) = 18 \text{ sq. units}$$




59. Equation of parabola is $y^2 = 12x$
 $\therefore a = 3$
 Given $y = 6$
 \therefore substituting $y = 6$ in $y^2 = 12x$, we get
 $36 = 12x$
 $\Rightarrow x = 3$
 Now, focal distance $= |x + a| = |3 + 3| = 6$
60. Given, $y = 3x$
 Substituting $y = 3x$ in $y^2 = 18x$, we get
 $(3x)^2 = 18x$
 $\Rightarrow 9x^2 = 18x$
 $\Rightarrow x = 2$ and $y = 6$
61. Vertex $= (0, 4)$, focus $= (0, 2)$
 $\Rightarrow a = 2$
 Hence, equation of parabola is
 $(x - 0)^2 = -4 \times 2(y - 4)$
 i.e., $x^2 + 8y = 32$
62. Given, vertex of parabola $(h, k) \equiv (1, 1)$ and its focus $(a + h, k) \equiv (3, 1)$ or $a + h = 3$ or $a = 2$. The y coordinates of vertex and focus are same, therefore axis of parabola is parallel to X-axis. Thus, equation of the parabola is $(y - k)^2 = 4a(x - h)$ or $(y - 1)^2 = 4 \times 2(x - 1)$ or $(y - 1)^2 = 8(x - 1)$
63. Directrix $= x + 5 = 0$
 Focus is $(-3, 0)$
 $\Rightarrow 2a = (5 - 3) = 2$
 $\Rightarrow a = 1$
 Vertex is $\left(\frac{-3 + (-5)}{2}, 0\right) = (-4, 0)$
 Therefore, equation is $(y - 0)^2 = 4(x + 4)$
 i.e., $y^2 = 4(x + 4)$
64. Let $P(x, y)$ be any point on the parabola.
 $\therefore SP^2 = PM^2$
 $\Rightarrow (x - 5)^2 + (y - 3)^2 = \left|\frac{3x - 4y + 1}{\sqrt{9 + 16}}\right|^2$
 $\Rightarrow 25(x^2 + 25 - 10x + y^2 + 9 - 6y)$
 $\quad = 9x^2 + 16y^2 + 1 - 24xy + 6x - 8y$
 $\Rightarrow 16x^2 + 9y^2 - 256x - 142y + 24xy + 849 = 0$
 $\Rightarrow (4x + 3y)^2 - 256x - 142y + 849 = 0$
65. Equation will be of the form $y^2 = 4A(x - a)$, where $A = (a' - a)$ or $y^2 = 4(a' - a)(x - a)$.

66. Equation of parabola
 $y^2 = 8x$
 $\therefore a = 2$
 $P(2t^2, 4t)$ $A(1, 0)$
 Mid point $= (x, y) = \left(\frac{2t^2 + 1}{2}, \frac{4t + 0}{2}\right)$
 $\Rightarrow \frac{2x - 1}{2} = t^2$ and $\frac{y^2}{4} = t^2$
 $\Rightarrow \frac{y^2}{4} = \frac{2x - 1}{2} \Rightarrow y^2 = 4\left(x - \frac{1}{2}\right)$
67. Eccentricity of parabola is always 1 i.e., $e = 1$.
68. Since, vertex is the midpoint of focus and directrix.
 \therefore vertex $= \left(\frac{0 + 2}{2}, \frac{0 + 0}{2}\right) = (1, 0)$
69. $y^2 - 4y - x + 3 = 0$
 $\Rightarrow y^2 - 4y + 4 - x + 3 - 4 = 0$
 $\Rightarrow (y - 2)^2 - (x + 1) = 0 \Rightarrow (y - 2)^2 = (x + 1)$
 Comparing with $(y - k)^2 = 4a(x - h)$, we get
 $h = -1, k = 2$
 \therefore vertex $= (-1, 2)$
70. $x^2 + 4x + 2y - 7 = 0$
 $\Rightarrow x^2 + 4x + 4 = -2y + 7 + 4$
 $\Rightarrow (x + 2)^2 = -2y + 11$
 $\Rightarrow (x + 2)^2 = -2\left(y - \frac{11}{2}\right)$
 Hence, vertex is $\left(-2, \frac{11}{2}\right)$
71. The given equation of parabola is
 $y = 2x^2 + x$
 $\Rightarrow x^2 + \frac{x}{2} = \frac{y}{2}$
 $\Rightarrow \left(x + \frac{1}{4}\right)^2 = \frac{y}{2} + \frac{1}{16} \Rightarrow \left(x + \frac{1}{4}\right)^2 = \frac{1}{2}\left(y + \frac{1}{8}\right)$
 Let $X^2 = \frac{1}{2}Y$ (i)
 Here $A = \frac{1}{8}$, focus of (i) is $\left(0, \frac{1}{8}\right)$
 i.e., $X = 0, Y = \frac{1}{8}$
 $\Rightarrow x + \frac{1}{4} = 0, y + \frac{1}{8} = \frac{1}{8} \Rightarrow x = -\frac{1}{4}, y = 0$
 \therefore focus of given parabola is $\left(-\frac{1}{4}, 0\right)$.



72. Given equation of conic
 $x^2 - 6x + 4y + 1 = 0$
 $\Rightarrow x^2 - 6x + 9 + 4y + 1 - 9 = 0$
 $\Rightarrow (x - 3)^2 + 4(y - 2) = 0$
 $\Rightarrow (x - 3)^2 = -4(y - 2)$
 Comparing with $X^2 = -4aY$, we get
 $a = 1, X = x - 3, Y = y - 2$
 Focus of the parabola is $X = 0, Y = -a$
 $\Rightarrow x - 3 = 0, y - 2 = -1 \Rightarrow x = 3, y = 1$
 \therefore Focus = $(3, 1)$

73.



Slope of AB = $\frac{-3}{5}$

\therefore Slope of PQ = $\frac{5}{3}$

\therefore Equation of PQ: $y = \frac{5}{3}x + c$

i.e. $5x - 3y = -3c$

At P, $y = 0 \Rightarrow x = \frac{-3c}{5}$

At Q, $x = 0 \Rightarrow y = c$

\therefore P = $\left(\frac{-3c}{5}, 0\right)$ and Q = $(0, c)$

\therefore Equation of AQ: $\frac{y}{-c} = \frac{x-5}{5}$

i.e. $c = \frac{-5y}{x-5}$... (i)

Equation of BP: $\frac{y+3}{3} = \frac{x}{\frac{3c}{5}}$

i.e. $c = \frac{5x}{y+3}$... (ii)

Since, lines AQ and BP intersect,

$\therefore \frac{-5y}{x-5} = \frac{5x}{y+3}$

$\therefore x^2 + y^2 - 5x + 3y = 0$ is the required locus of R.

74. Given equation of parabola
 $x^2 - 2x + 3y - 2 = 0$
 $\Rightarrow x^2 - 2x + 1 = -3y + 2 + 1$
 $\Rightarrow (x - 1)^2 = -3(y - 1)$
 \therefore vertex = $(h, k) = (1, 1)$,
 focus = $(h, k + b) = \left(1, \frac{1}{4}\right)$
 \therefore distance between focus and vertex
 $= \sqrt{0 + \left(1 - \frac{1}{4}\right)^2} = \sqrt{\left(\frac{3}{4}\right)^2}$
 $= \frac{3}{4}$

75. $x^2 + 4x + 2y = 0$
 $\Rightarrow x^2 + 4x + 4 = -2y + 4$
 $\Rightarrow (x + 2)^2 = -2(y - 2)$
 \therefore Equation of directrix is $y - 2 = -\frac{1}{2}$
 $\Rightarrow y = \frac{3}{2}$
 $\Rightarrow 2y = 3$

76. Given, equation of parabola is $x^2 + 8y - 2x = 7$
 $\Rightarrow x^2 - 2x + 8y - 7 = 0$
 $\Rightarrow x^2 - 2x + 1 + 8y - 7 - 1 = 0$
 $\Rightarrow (x - 1)^2 + 8y = 8$
 $\Rightarrow (x - 1)^2 = -8(y - 1)$
 $\Rightarrow (x - 1)^2 = -4 \times 2(y - 1)$
 Here, $a = 2$
 \therefore Equation of directrix is $y - 1 = 2$ i.e., $y = 3$

77. $y^2 + 6y - 2x = -5$
 $\Rightarrow y^2 + 6y + 9 = 2x - 5 + 9$
 $\Rightarrow (y + 3)^2 = 2(x + 2)$
 \therefore vertex = $(-2, -3)$
 Here, $a = \frac{1}{2}$
 \therefore Equation of directrix is $x + 2 = -\frac{1}{2}$
 $\Rightarrow 2x + 5 = 0$

78. Given, $x^2 = 4y$... (i)
 $y^2 = 4x$... (ii)
 $\Rightarrow \frac{x^4}{16} = 4x$... [From (i) and (ii)]
 $\Rightarrow x^4 = 64x$
 $\Rightarrow x = 0, 4$
 Substituting the values of x in (ii), we get
 $y = 0, 4$
 \therefore Other point is $(4, 4)$.



79. $x^2 = 8y$
 $\Rightarrow a = 4$
 By internal division formula
 $P(x, y) = \left(t, \frac{t^2}{2}\right)$

$\therefore x = t, y = \frac{t^2}{2}$
 $\Rightarrow x^2 = 2y$


80. $y^2 = -16x$
 $\Rightarrow a = -4$
 $t = \frac{1}{2}$... (given)

$\therefore x = at^2 = (-4) \times \left(\frac{1}{2}\right)^2$
 $\Rightarrow x = -1$
 $y = 2at = 2 \times (-4) \times \left(\frac{1}{2}\right)$

$\Rightarrow y = -4$

\therefore The cartesian co-ordinates are $(-1, -4)$.

81. Equation of the tangent at $P(16, 16)$ is $x - 2y + 16 = 0$
 Equation of the normal at $P(16, 16)$ is $2x + y - 48 = 0$
 Tangent and normal intersect the axis of parabola at $A(-16, 0)$ and $B(24, 0)$ respectively.
 AB is the diameter of the circle.
 Centre of the circle is $(4, 0)$.




$m_{PC} = \frac{16-0}{16-4} = \frac{16}{12} = \frac{4}{3}$

$m_{PB} = \frac{16-0}{16-24} = \frac{16}{-8} = -2$

$\tan \theta = \left| \frac{\frac{4}{3} - (-2)}{1 + \left(\frac{4}{3}\right)(-2)} \right| = 2$

82. Given, $y = \pm x$ (i)
 $y^2 = 8x$ (ii)
 Solving (i) and (ii),
 the point of intersection are $P(8, 8)$ and $Q(8, -8)$



\therefore Length of $PQ = \sqrt{(8-8)^2 + (8+8)^2}$
 $= 16$

83. Given parabolas are $y^2 = 4ax$ (i)
 $x^2 = 4ay$ (ii)
 Putting the value of y from (ii) in (i), we get

$\frac{x^4}{16a^2} = 4ax$
 $\Rightarrow x(x^3 - 64a^3) = 0$
 $\Rightarrow x = 0, 4a$

From (ii), $y = 0, 4a$

Let $A \equiv (0, 0)$, $B \equiv (4a, 4a)$

Since, the given line $2bx + 3cy + 4d = 0$ passes through A and B .

$\therefore d = 0$ and $8ab + 12ac = 0$
 $\Rightarrow 2b + 3c = 0$ [$\because a \neq 0$]

$\therefore d^2 + (2b + 3c)^2 = 0$

84. $\frac{x^2}{36} + \frac{y^2}{16} = 1$
 Here $a^2 = 36, b^2 = 16$
 $b^2 = a^2(1 - e^2)$
 $\Rightarrow 16 = 36(1 - e^2)$
 $\Rightarrow e = \frac{\sqrt{5}}{3} = \frac{2\sqrt{5}}{6}$

85. $4x^2 + y^2 - 8x + 4y - 8 = 0$
 $\Rightarrow 4(x^2 - 2x) + y^2 + 4y = 8$
 $\Rightarrow 4(x^2 - 2x + 1) + y^2 + 4y + 4 = 16$
 $\Rightarrow 4(x-1)^2 + (y+2)^2 = 16$
 $\Rightarrow \frac{(x-1)^2}{4} + \frac{(y+2)^2}{16} = 1$
 which is an ellipse with $a^2 = 4$ and $b^2 = 16$

$\therefore e = \sqrt{1 - \frac{a^2}{b^2}} = \sqrt{1 - \frac{4}{16}} = \sqrt{\frac{12}{16}}$

$\therefore e = \frac{\sqrt{3}}{2}$



$$86. \quad x^2 + 2y^2 - 2x + 3y + 2 = 0$$

$$\therefore (x^2 - 2x + 1) + 2\left(y^2 + \frac{3}{2}y + \frac{9}{16}\right) = \frac{1}{8}$$

$$\therefore (x-1)^2 + 2\left(y + \frac{3}{4}\right)^2 = \frac{1}{8}$$

$$\therefore \frac{(x-1)^2}{\frac{1}{8}} + \frac{\left(y + \frac{3}{4}\right)^2}{\frac{1}{16}} = 1$$

\therefore Comparing this with $\frac{X^2}{a^2} + \frac{Y^2}{b^2} = 1$, we get

$$a^2 = \frac{1}{8}, b^2 = \frac{1}{16}$$

$$\therefore e = \sqrt{1 - \frac{b^2}{a^2}} = \sqrt{1 - \frac{\frac{1}{16}}{\frac{1}{8}}} = 1 - \frac{8}{16}$$

$$\therefore e = \frac{1}{\sqrt{2}}$$

87. Here, given that $2b = 10$, $2a = 8$
 $\Rightarrow b = 5$, $a = 4$

Hence, the required equation is $\frac{x^2}{16} + \frac{y^2}{25} = 1$

88. Given, centre $(0,0)$, focus $(0,3)$, $b = 5$
 Focus $(0,3)$

$$\Rightarrow be = 3$$

$$\Rightarrow e = \frac{3}{5}$$

$$\text{Also, } a = b\sqrt{1 - e^2} = 5\sqrt{1 - \frac{9}{25}} = 4$$

Hence, the required equation is $\frac{x^2}{16} + \frac{y^2}{25} = 1$

89. Given, foci $= (\pm 2, 0) = (\pm ae, 0)$
 $\Rightarrow ae = 2$

$$\text{and } e = \frac{1}{2}$$

$$\therefore a = 4$$

$$\text{Now, } b^2 = a^2(1 - e^2)$$

$$\Rightarrow b^2 = 16\left(1 - \frac{1}{4}\right)$$

$$\Rightarrow b^2 = 12$$

Hence, equation of ellipse is $\frac{x^2}{16} + \frac{y^2}{12} = 1$

$$\Rightarrow 3x^2 + 4y^2 = 48$$

90. We have, $ae = \pm\sqrt{5}$

$$\Rightarrow a = \pm\sqrt{5}\left(\frac{3}{\sqrt{5}}\right) \quad \dots \left[\because e = \frac{\sqrt{5}}{3}\right]$$

$$\Rightarrow a = \pm 3$$

$$\Rightarrow a^2 = 9$$

$$\text{Now, } b^2 = a^2(1 - e^2) = 9\left(1 - \frac{5}{9}\right) = 4$$

Hence, equation of ellipse is $\frac{x^2}{9} + \frac{y^2}{4} = 1$

$$\Rightarrow 4x^2 + 9y^2 = 36$$

$$91. \quad b^2 = a^2(1 - e^2) = a^2\left(1 - \frac{2}{5}\right) = \frac{3a^2}{5}$$

Let the equation of ellipse be

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$\Rightarrow \frac{9}{a^2} + \frac{5}{3a^2} = 1$$

$$\Rightarrow a^2 = \frac{32}{3}$$

$$\therefore b^2 = \frac{32}{5}$$

\therefore the required equation of ellipse is $3x^2 + 5y^2 - 32 = 0$

92. Since point $(-3, 1)$ satisfies equations in options (C) and (D) writing them in standard form, we have,

For option (C):

$$\frac{x^2}{32} + \frac{y^2}{32} = 1, \text{ here } a^2 > b^2$$

For option (D):

$$\frac{x^2}{48} + \frac{y^2}{16} = 1, \text{ here, } a^2 < b^2$$

Since, the ellipse has its major axis along Y-axis,

$$\therefore a^2 < b^2$$

\therefore Option (D) is correct

93. Since, directrix is parallel to Y-axis, hence axes of the ellipse are parallel to X-axis.

Let the equation of the ellipse be

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, (a > b)$$

$$\text{Now, } e^2 = 1 - \frac{b^2}{a^2}$$



$$\Rightarrow \frac{b^2}{a^2} = 1 - e^2 = 1 - \frac{1}{4} \Rightarrow \frac{b^2}{a^2} = \frac{3}{4}$$

Also, one of the directrices is $x = 4$

$$\Rightarrow \frac{a}{e} = 4 \Rightarrow a = 4e = 4 \times \frac{1}{2} = 2;$$

$$b^2 = \frac{3}{4} a^2 = \frac{3}{4} \times 4 = 3$$

\therefore Required equation of ellipse is $\frac{x^2}{4} + \frac{y^2}{3} = 1$

$$\Rightarrow 3x^2 + 4y^2 = 12$$

95. Sum of focal distances of a point in an ellipse is always equal to length of major axis of that ellipse.

96. Equation of the curve is $\frac{x^2}{5^2} + \frac{y^2}{4^2} = 1$

Here, $a = 5, b = 4$

$$\therefore PF_1 + PF_2 = 2a = 2 \times 5 = 10$$

97. The equation of the ellipse is

$$16x^2 + 25y^2 = 400 \Rightarrow \frac{x^2}{25} + \frac{y^2}{16} = 1$$


Here, $a^2 = 25, b^2 = 16$

$$e = \sqrt{1 - \frac{b^2}{a^2}} = \sqrt{1 - \frac{16}{25}} \Rightarrow e = \frac{3}{5}$$

Hence, the foci are $(\pm ae, 0) = (\pm 3, 0)$

98. Since, $\angle FBF' = \frac{\pi}{2}$ (Given)

$$\therefore \angle FBC = \angle F'BC = \frac{\pi}{4}$$




Now, $CB = CF$

$$\Rightarrow b = ae \Rightarrow b^2 = a^2 e^2$$

$$\Rightarrow a^2(1 - e^2) = a^2 e^2 \Rightarrow 1 - e^2 = e^2$$

$$\Rightarrow 2e^2 = 1 \Rightarrow e = \frac{1}{\sqrt{2}}$$

99.




$S = (ae, 0), S' = (-ae, 0)$ and $b = ae$

$$\text{Now, } a^2 e^2 = a^2(1 - e^2)$$

$$\Rightarrow e = \frac{1}{\sqrt{2}}$$

100. Since, $\triangle SBS'$ is an isosceles right angled triangle,



$$\therefore \angle SBS' = \frac{\pi}{2}, \angle BSS' = \angle BS'S = \frac{\pi}{4}$$

$$\therefore e = \frac{1}{\sqrt{2}}$$

101. Distance between the foci $= 2ae = 16$ and

$$e = \frac{1}{2}$$

\therefore Length of the major axis of the ellipse

$$= 2a = \frac{2ae}{e} = \frac{16}{\frac{1}{2}} = 32$$

102. According to the given condition,

$$\sqrt{1 - \frac{25}{169}} = \sqrt{1 - \frac{b^2}{a^2}}$$

$$\Rightarrow \frac{144}{169} = 1 - \frac{b^2}{a^2}$$

$$\Rightarrow \frac{b^2}{a^2} = \frac{25}{169}$$

$$\Rightarrow \frac{b}{a} = \frac{5}{13}$$

....[$\because a > 0, b > 0$]

$$\Rightarrow \frac{a}{b} = \frac{13}{5}$$

103. $5x^2 + 9y^2 = 45$

$$\Rightarrow \frac{x^2}{9} + \frac{y^2}{5} = 1$$

$$\therefore \text{Length of latus rectum} = \frac{2b^2}{a} = \frac{2 \times 5}{3} = \frac{10}{3}$$

104. Given, ellipse is $\frac{x^2}{\left(\frac{1}{3}\right)^2} + \frac{y^2}{\left(\frac{1}{2}\right)^2} = 1$

Here, $b > a$

$$\therefore \text{Latus rectum} = \frac{2a^2}{b} = \frac{2 \times \frac{1}{9}}{\frac{1}{2}} = \frac{4}{9}$$



105. Latus rectum = $\frac{1}{3}$ (major axis)

$$\Rightarrow \frac{2b^2}{a} = \frac{2a}{3}$$

$$\Rightarrow a^2 = 3b^2$$

$$\Rightarrow a^2 = 3a^2(1 - e^2)$$

$$\Rightarrow e = \sqrt{\frac{2}{3}}$$

106. Given, $\frac{2b^2}{a} = b$

$$\Rightarrow \frac{b}{a} = \frac{1}{2}$$

$$\Rightarrow \frac{b^2}{a^2} = \frac{1}{4}$$

Hence, $e = \sqrt{1 - \frac{b^2}{a^2}} = \frac{\sqrt{3}}{2}$

107. $x^2 = 9\cos^2\theta$
 $y^2 = 16\sin^2\theta$

$$\Rightarrow \frac{x^2}{9} + \frac{y^2}{16} = 1$$

$$\therefore e = \frac{\sqrt{7}}{4}$$

Distance between foci = $2be = 2\sqrt{7}$

108. Let (x, y) be any point on ellipse.
Then, by focus-directrix property of ellipse,

$$\sqrt{(x+1)^2 + (y-1)^2} = \frac{1}{2} \left| \frac{x-y+3}{\sqrt{1+1}} \right|$$

$$\therefore 8(x^2 + 2x + 1 + y^2 - 2y + 1) = x^2 + y^2 + 9 - 2xy - 6y + 6x$$

$$\therefore 7x^2 + 2xy + 7y^2 + 10x - 10y + 7 = 0$$

109. solving $x + y - 3 = 0$

$x - y + 1 = 0$, we get

$$(x, y) = (1, 2)$$

110. Here, $a^2 = 9$, $b^2 = 25$

Since, $b > a$

$$\therefore e = \sqrt{\frac{b^2 - a^2}{b^2}} = \sqrt{\frac{25 - 9}{25}} = \frac{4}{5}$$

111. $\frac{(x-1)^2}{2} + \left(y + \frac{3}{4}\right)^2 = \frac{1}{16}$

$$\Rightarrow \frac{(x-1)^2}{2\left(\frac{1}{16}\right)} + \frac{\left(y + \frac{3}{4}\right)^2}{\left(\frac{1}{16}\right)} = 1$$

$$\Rightarrow \frac{(x-1)^2}{\left(\frac{1}{8}\right)} + \frac{\left(y + \frac{3}{4}\right)^2}{\left(\frac{1}{16}\right)} = 1$$

$$\therefore a^2 = \frac{1}{8}, b^2 = \frac{1}{16}$$

$$b^2 = a^2(1 - e^2)$$

$$\Rightarrow \frac{1}{16} = \frac{1}{8}(1 - e^2)$$

$$\Rightarrow e^2 = 1 - \frac{1}{2}$$

$$\Rightarrow e = \frac{1}{\sqrt{2}}$$

112. $4x^2 + y^2 - 8x + 4y - 8 = 0$

$$\Rightarrow 4(x^2 - 2x) + y^2 + 4y = 8$$

$$\Rightarrow 4(x^2 - 2x + 1) + y^2 + 4y + 4 = 16$$

$$\Rightarrow 4(x-1)^2 + (y+2)^2 = 16$$

$$\Rightarrow \frac{(x-1)^2}{4} + \frac{(y+2)^2}{16} = 1$$

Comparing with $\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$, we get

$$h = 1, k = -2$$

\therefore centre of the ellipse = $(1, -2)$

113. Given equation of ellipse is

$$25x^2 + 9y^2 - 150x - 90y + 225 = 0$$

$$\Rightarrow 25(x-3)^2 + 9(y-5)^2 = 225$$

$$\Rightarrow \frac{(x-3)^2}{9} + \frac{(y-5)^2}{25} = 1$$

Since, $b > a$

$$\therefore e = \sqrt{1 - \frac{a^2}{b^2}} = \sqrt{1 - \frac{9}{25}} = \sqrt{\frac{16}{25}} = \frac{4}{5}$$

114. $x^2 + 2y^2 - 2x + 3y + 2 = 0$

$$\Rightarrow (x^2 - 2x + 1) + 2\left(y^2 + \frac{3}{2}y + \frac{9}{16}\right)$$

$$= -2 + 1 + \frac{9}{8}$$

$$\Rightarrow (x-1)^2 + 2\left(y + \frac{3}{4}\right)^2 = \frac{1}{8}$$

$$\Rightarrow \frac{(x-1)^2}{\frac{1}{8}} + \frac{\left(y + \frac{3}{4}\right)^2}{\frac{1}{16}} = 1,$$

which is an ellipse with $a^2 = \frac{1}{8}$ and $b^2 = \frac{1}{16}$



Also, $b^2 = a^2(1 - e^2)$

$$\Rightarrow \frac{1}{16} = \frac{1}{8}(1 - e^2)$$

$$\Rightarrow e^2 = 1 - \frac{1}{2}$$

$$\Rightarrow e = \frac{1}{\sqrt{2}}$$

115. $25x^2 + 4y^2 + 100x - 4y + 100 = 0$

$$\Rightarrow 25(x^2 + 4x) + 4(y^2 - y) = -100$$

$$\Rightarrow 25(x^2 + 4x + 4) + 4\left(y^2 - y + \frac{1}{4}\right) = -100 + 100 + 1$$

$$\Rightarrow 25(x + 2)^2 + 4\left(y - \frac{1}{2}\right)^2 = 1$$

$$\Rightarrow \frac{(x+2)^2}{\frac{1}{25}} + \frac{\left(y - \frac{1}{2}\right)^2}{\frac{1}{4}} = 1$$

Here $a^2 = \frac{1}{25}$, $b^2 = \frac{1}{4}$, $h = -2$, $k = \frac{1}{2}$

$$\therefore e = \frac{\sqrt{b^2 - a^2}}{b} = \frac{\sqrt{\frac{1}{4} - \frac{1}{25}}}{\frac{1}{2}} = \frac{\sqrt{21}}{5}$$

$$\therefore \text{Foci} = (h, k \pm be) = \left(-2, \frac{1}{2} \pm \frac{\sqrt{21}}{10}\right) = \left(-2, \frac{5 \pm \sqrt{21}}{10}\right)$$

116. $5x^2 + y^2 + y = 8$

$$\Rightarrow 5x^2 + y^2 + y + \frac{1}{4} = 8 + \frac{1}{4}$$

$$\Rightarrow 5x^2 + \left(y + \frac{1}{2}\right)^2 = \frac{33}{4}$$

$$\Rightarrow \frac{x^2}{\left(\frac{33}{20}\right)} + \frac{\left(y + \frac{1}{2}\right)^2}{\left(\frac{33}{4}\right)} = 1$$

The equation represent an ellipse.

117. Here, $2a = 10\text{m}$ and $2ae = 8\text{m}$

$$\therefore e = \frac{4}{5}, a = 5\text{m}$$

Now, $b^2 = a^2(1 - e^2) = 9$

$$\Rightarrow b = 3$$

Thus, required area $= \pi ab = 15\pi$ sq. metre.


118. We have, $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

$$\therefore y = \frac{b}{a} \sqrt{a^2 - x^2} \quad \dots(i)$$

Now, area of $\Delta PF_1F_2 = \frac{1}{2}(F_1F_2) \times PL$

$$= \frac{1}{2}(2ae) \times y$$

$$= ae \cdot \frac{b}{a} \sqrt{a^2 - x^2} \quad \dots[\text{From (i)}]$$



$$\therefore A = eb \sqrt{a^2 - x^2}, \text{ which is maximum when } x = 0.$$

Thus, the maximum value of A is abe .

119. Vertices $= (0, \pm 15)$, foci $= (0, \pm 20)$

$$\therefore b = 15 \text{ and } be = 20 \Rightarrow e = \frac{4}{3}$$

$$\begin{aligned} a^2 &= b^2(e^2 - 1) \\ &= 15^2 \left(\frac{16}{9} - 1\right) \\ &= 175 \end{aligned}$$

$$\therefore \text{The equation of hyperbola is}$$

$$\frac{-x^2}{175} + \frac{y^2}{225} = 1$$

$$\Rightarrow \frac{y^2}{225} - \frac{x^2}{175} = 1$$

120. Given, $ae = 2$, $e = 2$

$$\therefore a = 1$$

Now, $b^2 = a^2(e^2 - 1)$

$$\Rightarrow b^2 = 1(4 - 1)$$

$$\Rightarrow b^2 = 3$$

$$\therefore \text{the equation of hyperbola is}$$

$$\frac{x^2}{1} - \frac{y^2}{3} = 1$$

$$\Rightarrow 3x^2 - y^2 = 3$$

121. Given: $ae = 8$, $e = \sqrt{2}$

$$\therefore a = 4\sqrt{2}$$

Now, $b^2 = a^2(e^2 - 1)$

$$\Rightarrow b^2 = 32(2 - 1)$$

$$\Rightarrow b^2 = 32$$

$$\therefore \text{the equation of hyperbola is } \frac{x^2}{32} - \frac{y^2}{32} = 1$$

$$\therefore x^2 - y^2 = 32$$



122. Given, $ae = 8$ and $\frac{2b^2}{a} = 24$
 $\Rightarrow b^2 = 12a$
 Now, $b^2 = a^2(e^2 - 1)$
 $\Rightarrow 12a = a^2e^2 - a^2$
 $\Rightarrow 12a = 64 - a^2$
 $\Rightarrow a^2 + 12a - 64 = 0$
 $\Rightarrow a = 4 \quad \dots[\because a > 0]$
 $\therefore b^2 = 12(4) = 48$
 \therefore the equation of hyperbola is
 $\frac{x^2}{16} - \frac{y^2}{48} = 1 \Rightarrow 3x^2 - y^2 = 48$
123. $16x^2 - 9y^2 - 64x + 18y - 90 = 0$
 $\therefore 16(x^2 - 4x + 4) - 9(y^2 - 2y + 1) = 145$
 $\therefore \frac{(x-2)^2}{\frac{145}{16}} - \frac{(y-1)^2}{\frac{145}{9}} = 1,$
 Comparing with $\frac{X^2}{a^2} - \frac{Y^2}{b^2} = 1$, we get
 $X = x - 2, \quad Y = y - 1$
 $a^2 = \frac{145}{16}, \quad b^2 = \frac{145}{9}$
 $\therefore e = \sqrt{1 + \frac{b^2}{a^2}} = \sqrt{1 + \frac{16}{9}} = \frac{\pm 5}{3}$
 Focus of the hyperbola, $\frac{X^2}{a^2} - \frac{Y^2}{b^2} = 1$ is,
 $(X = \pm ae, Y = 0)$
 i.e. $x - 2 = \pm \left(\frac{\sqrt{145}}{4} \times \frac{5}{3}\right), y - 1 = 0$
 i.e. $x - 2 = \pm \frac{5\sqrt{145}}{12}, y = 1$
 i.e. $x = \frac{24 \pm 5\sqrt{145}}{12}, y = 1$
 \therefore Focus = $\left(\frac{24 \pm 5\sqrt{145}}{12}, 1\right)$
124. $\frac{x^2}{36} - \frac{y^2}{k^2} = 1$ is a hyperbola $\Rightarrow k^2 > 0$
 Now, $\frac{y^2}{k^2} = \frac{x^2}{36} - 1 = \frac{x^2 - 36}{36}$
 $\Rightarrow k^2 = \frac{36y^2}{x^2 - 36} > 0 \Rightarrow x^2 - 36 > 0$
 $\Rightarrow x^2 > 36$
 This is true only for point (10, 4).
 \therefore (10, 4) lies on given hyperbola.

125. $\frac{x^2}{12-k} + \frac{y^2}{8-k} = 1$
 $\Rightarrow \frac{x^2}{12-k} - \frac{y^2}{k-8} = 1$
 $\therefore 12 > k$ and $k > 8$
 $\Rightarrow 8 < k < 12$
 \therefore the given equation represents a hyperbola, if $8 < k < 12$.
126. Consider,
 $x^2 - y^2 + 3x - 2y - 43 = 0$
 Comparing with
 $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$, we get
 $a = 1, h = 0, b = -1, g = \frac{3}{2}, f = -1, c = -43$
 $\therefore \Delta = abc + 2fgh - af^2 - bg^2 - ch^2$
 $\therefore \Delta = 43 + 0 - 1 + \frac{9}{4} - 0 = \frac{177}{4}$
 $\therefore \Delta \neq 0$
 Also, $ab - h^2 = -1$
 $\therefore ab - h^2 < 0$
 $\therefore x^2 - y^2 + 3x - 2y - 43 = 0$ is the equation of hyperbola.
127. The given equation can be written as
 $\frac{x^2}{\frac{32}{3}} - \frac{y^2}{8} = 1$
 $\Rightarrow \frac{x^2}{\left(\frac{4\sqrt{2}}{\sqrt{3}}\right)^2} - \frac{y^2}{(2\sqrt{2})^2} = 1$
 $\Rightarrow a = \frac{4\sqrt{2}}{\sqrt{3}}$
 \therefore Length of transverse axis of a hyperbola
 $= 2a = 2 \times \frac{4\sqrt{2}}{\sqrt{3}} = \frac{8\sqrt{2}}{\sqrt{3}}$
128. $\frac{x^2}{9} - \frac{y^2}{4} = 1$
 $\Rightarrow a^2 = 9, b^2 = 4$
 $e = \frac{\sqrt{a^2 + b^2}}{a} = \frac{\sqrt{9+4}}{3} = \frac{\sqrt{13}}{3}$
 directrix of hyperbola is $x = \frac{a}{e}$
 $\Rightarrow x = \frac{3}{\frac{\sqrt{13}}{3}} \Rightarrow x = \frac{9}{\sqrt{13}}$
 $\Rightarrow x = \frac{3}{\sqrt{13}}$



$$129. x^2 - 4y^2 = 1$$

$$\Rightarrow \frac{x^2}{(1)^2} - \frac{y^2}{\left(\frac{1}{2}\right)^2} = 1$$

$$\Rightarrow a^2 = 1, b^2 = \left(\frac{1}{2}\right)^2$$

$$\text{Also, } b^2 = a^2(e^2 - 1)$$

$$\Rightarrow \frac{1}{4} + 1 = e^2 \Rightarrow e = \frac{\sqrt{5}}{2}$$

130. Here $a = b$, so it is a rectangular hyperbola.

Hence, eccentricity $e = \sqrt{2}$

131. Let the equation of hyperbola be $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

Since, hyperbola passes through the points $(3, 0), (3\sqrt{2}, 2)$

$$\therefore \frac{9}{a^2} - 0 = 1 \text{ and } \frac{18}{a^2} - \frac{4}{b^2} = 1$$

$$\Rightarrow a^2 = 9 \text{ and } \frac{4}{b^2} = \frac{18}{a^2} - 1$$

$$\Rightarrow a^2 = 9 \text{ and } \frac{4}{b^2} = \frac{18}{9} - 1$$

$$\Rightarrow a^2 = 9 \text{ and } b^2 = 4$$

$$\text{Now, } e = \sqrt{1 + \frac{b^2}{a^2}} = \sqrt{1 + \frac{4}{9}} = \frac{\sqrt{13}}{3}$$

132. Given length of LR = 8

$$\Rightarrow \frac{2b^2}{a} = 8$$

$$\text{Also, } 2b = \frac{1}{2} 2ae$$

$$\Rightarrow 4b^2 = a^2 e^2 \Rightarrow 4a^2(e^2 - 1) = a^2 e^2$$

$$\Rightarrow 4e^2 - e^2 = 4 \Rightarrow e = \frac{2}{\sqrt{3}}$$

133. Eccentricity of hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ is

$$e = \sqrt{1 + \frac{b^2}{a^2}}$$

$$\Rightarrow e^2 = \frac{a^2 + b^2}{a^2} \quad \dots(i)$$

Eccentricity of hyperbola $\frac{y^2}{b^2} - \frac{x^2}{a^2} = 1$ is

$$e_1 = \sqrt{1 + \frac{a^2}{b^2}}$$

$$\Rightarrow e_1^2 = \frac{b^2 + a^2}{b^2} \quad \dots(ii)$$

From (i) and (ii), we get

$$\frac{1}{e_1^2} + \frac{1}{e^2} = 1$$

134. The hyperbola is $\frac{x^2}{16} - \frac{y^2}{9} = 1$

\therefore Difference of focal distance = $2a = 8$

135. Since, distance between directrices = $\frac{2a}{e}$ and eccentricity of rectangular hyperbola = $\sqrt{2}$.

\therefore Distance between directrices = $\frac{2a}{\sqrt{2}}$

$$\text{Given, } \frac{2a}{\sqrt{2}} = 10$$

$$\Rightarrow 2a = 10\sqrt{2}$$

$$\begin{aligned} \text{Now, distance between foci} &= 2ae \\ &= (10\sqrt{2})(\sqrt{2}) \\ &= 20 \end{aligned}$$

136. Given, $x = a\left(t + \frac{1}{t}\right)$

$$\Rightarrow \frac{x}{a} = t + \frac{1}{t} \quad \dots(i)$$

$$\text{and } y = b\left(t - \frac{1}{t}\right)$$

$$\Rightarrow \frac{y}{b} = t - \frac{1}{t} \quad \dots(ii)$$

Squaring and subtracting equation (ii) from (i), we get

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = t^2 + 2 + \frac{1}{t^2} - t^2 + 2 - \frac{1}{t^2}$$

$$\Rightarrow \frac{x^2}{a^2} - \frac{y^2}{b^2} = 4$$

which represents a hyperbola.

137. $x^2 - 2x - 4y^2 + 16y - 40 = 0$

$$\Rightarrow (x^2 - 2x) - 4(y^2 - 4y) - 40 = 0$$

$$\Rightarrow (x-1)^2 - 1 - 4[(y-2)^2 - 4] - 40 = 0$$

$$\Rightarrow (x-1)^2 - 4(y-2)^2 = 25$$

$$\Rightarrow \frac{(x-1)^2}{25} - \frac{(y-2)^2}{\frac{25}{4}} = 1, \text{ which is a hyperbola.}$$



138. Given equation of lines are

$$\sqrt{3}x - y - 4\sqrt{3}k = 0$$

$$\Rightarrow \sqrt{3}x - y = 4\sqrt{3}k \quad \dots(i)$$

$$\text{and } \sqrt{3}kx + ky - 4\sqrt{3} = 0$$

$$\Rightarrow k(\sqrt{3}x + y) = 4\sqrt{3}$$

$$\Rightarrow \sqrt{3}x + y = \frac{4\sqrt{3}}{k} \quad \dots(ii)$$

Multiplying equation (i) and (ii), we get

$$(\sqrt{3}x - y)(\sqrt{3}x + y) = 4\sqrt{3}k \times \frac{4\sqrt{3}}{k}$$

$$\Rightarrow 3x^2 - y^2 = 48$$

$$\Rightarrow \frac{x^2}{(48/3)} - \frac{y^2}{48} = 1, \text{ which is a hyperbola.}$$

139. Given equation of hyperbola is

$$x^2 - y^2 + 1 = 0$$

$$\Rightarrow x^2 - y^2 = -1$$

$$\Rightarrow \frac{x^2}{1} - \frac{y^2}{1} = -1$$

$$\therefore a^2 = 1, b^2 = 1$$

$$e = \sqrt{\frac{a^2 + b^2}{b^2}} = \sqrt{\frac{1+1}{1}} = \sqrt{2}$$

$$\therefore \text{foci} \equiv (0, \pm be) \equiv (0, \pm\sqrt{2}), \text{ and}$$

$$\text{centre} \equiv (0, 0)$$

$$\text{Centre of circle} \equiv (0, 0), \text{ radius of circle} = \sqrt{2}$$

$$\therefore \text{Equation of circle is } x^2 + y^2 = 2.$$

140. Hyperbola is $\frac{x^2}{144} - \frac{y^2}{81} = \frac{1}{25}$

$$a = \sqrt{\frac{144}{25}},$$

$$b = \sqrt{\frac{81}{25}}$$

$$\therefore e_1 = \sqrt{1 + \frac{81}{144}} = \sqrt{\frac{225}{144}} = \frac{15}{12} = \frac{5}{4}$$

$$\therefore \text{foci} = (ae_1, 0) = \left(\frac{12}{5} \times \frac{5}{4}, 0\right) = (3, 0)$$

$$\therefore \text{focus of ellipse} = (4e, 0)$$

Since, focus of ellipse and hyperbola is same

$$\therefore (4e, 0) = (3, 0)$$

$$\Rightarrow e = \frac{3}{4}$$

$$\text{Hence, } b^2 = 16 \left(1 - \frac{9}{16}\right) = 7$$

141. Eccentricity of ellipse $x^2 + 9y^2 = 9$ i.e.

$$\frac{x^2}{9} + \frac{y^2}{1} = 1$$

$$(e_1)^2 = 1 - \frac{1}{9} = \frac{8}{9}$$

Now, eccentricity of hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

is,

$$(e_2)^2 = 1 + \frac{b^2}{a^2} = \frac{9}{8}$$

$$\therefore \frac{b^2}{a^2} = \frac{9}{8} - 1 \quad \therefore \frac{a^2}{b^2} = \frac{8}{1}$$

142. The equation of the ellipse is $\frac{x^2}{4} + \frac{y^2}{1} = 1$

Let e be its eccentricity.

$$\text{Then, } e = \sqrt{1 - \frac{1}{4}} = \frac{\sqrt{3}}{2}$$

The foci of the ellipse are S $(\sqrt{3}, 0)$ and

S' $(-\sqrt{3}, 0)$.

$$\text{Eccentricity of the hyperbola} = \frac{1}{e} = \frac{2}{\sqrt{3}}$$

$$\therefore b^2 = a^2 \left(\frac{4}{3} - 1\right) = \frac{a^2}{3}$$

The hyperbola passes through S $(\sqrt{3}, 0)$.

$$\therefore \frac{3}{a^2} - 0 = 1 \Rightarrow a^2 = 3 \Rightarrow a = \sqrt{3}$$

$$\therefore \text{the co-ordinates of the foci of hyperbola are } (\pm 2, 0).$$

$$143. x - 3y = 1 \quad \dots(i)$$

$$\text{and } x^2 - 4y^2 = 1 \quad \dots(ii)$$

On solving (i) and (ii), we get

$$A(1, 0) \text{ and } B\left(-\frac{13}{5}, -\frac{6}{5}\right)$$

These are the points of intersection of the straight line and hyperbola.

$$\therefore \text{Length of straight line intercepted by the hyperbola}$$


$$= \sqrt{\left(-\frac{13}{5} - 1\right)^2 + \left(-\frac{6}{5}\right)^2}$$

$$= \sqrt{\left(-\frac{18}{5}\right)^2 + \left(-\frac{6}{5}\right)^2} = \sqrt{\frac{324 + 36}{25}}$$

$$= \sqrt{\frac{360}{25}} = \frac{6}{5}\sqrt{10} \text{ units}$$



144. Given equation of circle
 $x^2 + y^2 + 2x - 2y - 2 = 0$
 $\Rightarrow (x + 1)^2 + (y - 1)^2 = 4$
 \therefore centre = $(-1, 1)$, and
 radius = 2




$$\sin 45^\circ = \frac{OM}{OA}$$

$$\Rightarrow OM = \sqrt{2}$$

\therefore locus of the mid-points of the chord is
 $(x + 1)^2 + (y - 1)^2 = (\sqrt{2})^2$
 $\Rightarrow x^2 + 2x + 1 + y^2 - 2y + 1 = 2$
 $\Rightarrow x^2 + y^2 + 2x - 2y = 0$

145.



Since, $x + y + k = 0$ touches the given circle.

$$\therefore \left| \frac{1(1) + 1(2) + k}{\sqrt{1+1}} \right| = \text{radius}$$

$$\Rightarrow \frac{3+k}{\sqrt{2}} = \pm\sqrt{2} \Rightarrow k = -1 \text{ or } k = -5$$

146. The diameter of the circle is perpendicular distance between the parallel lines (tangents)

Now, $3x - 4y + 4 = 0$ (i)
 $6x - 8y - 7 = 0$


i.e., $3x - 4y - \frac{7}{2} = 0$ (ii)

Since, equation (i) and (ii) are parallel to each other.

$$\therefore \text{diameter} = \left| \frac{4 - \left(-\frac{7}{2}\right)}{\sqrt{(3)^2 + (-4)^2}} \right| = \frac{15}{2 \times 5} = \frac{3}{2}$$

$$\therefore \text{radius} = \frac{3}{4}$$

147.



Let $M = (x, y)$

Since, $AM = 2AB$

$$\therefore \frac{AB}{AM} = \frac{1}{2}$$

\therefore B is the mid point of seg AM.

$$\therefore B = \left(\frac{x}{2}, \frac{y+3}{2} \right)$$

Since, B lies on circle $x^2 + 4x + (y - 3)^2 = 0$


$$\therefore \left(\frac{x}{2} \right)^2 + 4 \left(\frac{x}{2} \right) + \left(\frac{y+3}{2} - 3 \right)^2 = 0$$

i.e. $\frac{x^2}{4} + 2x + \frac{y-6x+9}{4} = 0$

i.e. $x^2 + y^2 + 8x - 6y + 9 = 0$
 is the required locus of M.

148. For given circle, $x^2 + y^2 - 2ax - 2ay + a^2 = 0$
 Centre = (a, a)

Also, radius = $\sqrt{a^2 + a^2 - a^2} = a$



The above circle touches $x = 0$.


149. Given equation is $x^2 + y^2 - 6x + 2y = 0$.

Centre = $(3, -1)$

Since, diameter is passing through origin and $(3, -1)$.

\therefore option (A) is the correct answer.

150.



Given, $x^2 + y^2 - 8x - 8y - 4 = 0$

$$\therefore (x^2 - 8x + 16) + (y^2 - 8y + 16) = 16 + 16 + 4$$

$$\Rightarrow (x - 4)^2 + (y - 4)^2 = 36$$

Equation of circle touching X-axis

$$(x - h)^2 + (y - k)^2 = k^2$$

Since, both circle touches externally

\therefore distance between their centre = $r_1 + r_2$

$$\sqrt{(4-h)^2 + (4-k)^2} = 6 + k$$



$$\begin{aligned} \Rightarrow (4-h)^2 + (4-k)^2 &= (6+k)^2 \\ \Rightarrow (4-h)^2 &= 36 + 12k + k^2 - 16 + 8k - k^2 \\ \Rightarrow (4-h)^2 &= 20k + 20 \end{aligned}$$

This is equation of parabola

\therefore answer is option (D)

151. Equations of tangents to the circle $x^2 + y^2 - 2x - 4y - 20 = 0$ with centre A(1, 2),

At, B(1, 7) is $y = 7$

At, C(4, -2) is $3x - 4y - 20 = 0$

These tangents intersect each other at D(16, 7).

\therefore Area of quadrilateral BACD
= Area of $\triangle ABD$ + Area of $\triangle ACD$

$$= \frac{1}{2} (AB)(BD) + \frac{1}{2} (AC)(CD)$$

...[$\because \angle ABD = \angle ACD = 90^\circ$]


$$= \frac{1}{2} \times (5) \times (15) + \frac{1}{2} \times (5) \times (15)$$

= 75 sq. units

152. $AB = \sqrt{(3+3)^2 + (3-5)^2}$

$$= \sqrt{36+4} = \sqrt{40} = 2\sqrt{10}$$

Centroid divides orthocentre and circumcentre in the ratio 2 : 1.



$$AB : BC = 2 : 1$$

$$AC = \frac{3}{2} AB$$

$$= \frac{3}{2} (2\sqrt{10}) = 3\sqrt{10}$$

$$\text{radius} = \frac{1}{2} AC = \frac{1}{2} (3\sqrt{10}) = 3\sqrt{\frac{5}{2}}$$

153. Given, $x + y = 0$ (i)

$$x^2 + y^2 + 4y = 0 \quad \dots\text{(ii)}$$

Solving (i) and (ii), we get

$$x = 0, y = 0; x = 2, y = -2$$

\therefore parabola passes through (0, 0) and (2, -2).
These points are satisfied by the parabola $y^2 = 2x$.

154. Given equation of ellipse is $\frac{x^2}{144} + \frac{y^2}{25} = 1$

$$\Rightarrow a = 12, b = 5$$

$$\text{Now, } e = \sqrt{1 - \frac{b^2}{a^2}} = \sqrt{1 - \frac{25}{144}} = \frac{\sqrt{119}}{12}$$

$$\therefore \text{ focus} = (ae, 0) = (\sqrt{119}, 0)$$

$$\therefore \text{ Radius} = \sqrt{(0 - \sqrt{119})^2 + (\sqrt{2} - 0)^2} = 11$$

155. $\frac{x^2}{16} + \frac{y^2}{12} = 1$

\therefore centre of ellipse = (0, 0)

\therefore centre of circle = (0, 0)

$$y = 2x + \sqrt{76} \quad \dots\text{(i)}$$

$$2y + x = 8 \quad \dots\text{(ii)}$$

On solving (i) and (ii), we get

$$x = \frac{8 - 2\sqrt{76}}{5}, y = \frac{16 + \sqrt{76}}{5}$$

Since (x, y) lies on the circle,

$$\begin{aligned} \therefore \text{ radius} &= \sqrt{(x-0)^2 + (y-0)^2} \\ &= \sqrt{\left(\frac{8-2\sqrt{76}}{5}\right)^2 + \left(\frac{16+\sqrt{76}}{5}\right)^2} \\ &= 2\sqrt{7} \end{aligned}$$

\therefore Equation of circle is $x^2 + y^2 = (2\sqrt{7})^2$

$$\Rightarrow x^2 + y^2 = 28$$

157. Midpoint of (4, 0) and (0, 4) is (2, 2).

Distance between (2, 2) and centre (0, 0)


$$= \sqrt{2^2 + 2^2} = 2\sqrt{2}$$

158. PQ is a chord of contact.

Equation of PQ is

$$\frac{xx_1}{9} - \frac{yy_1}{36} = 1$$

$$\Rightarrow 0 - \frac{3y}{36} = 1 \Rightarrow y = -12$$



Substituting $y = -12$ in $4x^2 - y^2 = 36$, we get $x = \pm 3\sqrt{5}$

$$P(3\sqrt{5}, -12), Q(-3\sqrt{5}, -12), T(0, 3)$$

$$PQ = 6\sqrt{5}, TR = 15$$

$$\text{Area of } \triangle PTQ = \frac{1}{2} \times PQ \times TR$$

$$= \frac{1}{2} \times 6\sqrt{5} \times 15$$

$$= 45\sqrt{5} \text{ sq. units}$$



Evaluation Test


1. Since the triangle is equilateral.
 \therefore centroid of the triangle is same as the circumcentre
 and radius of the circumcircle = $\frac{2}{3}$ (median)

$$= \frac{2}{3} (3a) = 2a$$

Hence, the equation of the circumcircle whose centre is at (0, 0) and radius 2a is
 $x^2 + y^2 = (2a)^2$
 $\Rightarrow x^2 + y^2 = 4a^2$

2. Let the other end be (t, 3 - t).
 \therefore the equation of the circle in diameter form is
 $(x - 1)(x - t) + (y - 1)(y - 3 + t) = 0$
 $\Rightarrow x^2 + y^2 - (1 + t)x - (4 - t)y + 3 = 0$
 \therefore the centre (h, k) is given by
 $h = \frac{1+t}{2}, k = \frac{4-t}{2}$
 $\Rightarrow 2h + 2k = 5$
 Hence, the locus is $2x + 2y = 5$.

3. We have, $x^2 - 8x + 12 = 0$
 $\Rightarrow (x - 6)(x - 2) = 0$
 $\Rightarrow x = 2, 6$
 and $y^2 - 14y + 45 = 0$
 $\Rightarrow (y - 5)(y - 9) = 0$
 $\Rightarrow y = 5, 9$




Since, centre of circle is inscribed in square.

- \therefore BD is the diameter of circle
 \therefore centre = (h, k) = $\left(\frac{2+6}{2}, \frac{5+9}{2}\right) = (4, 7)$

4. Let $A \equiv (x_1, y_1)$ and $B \equiv (x_2, y_2)$.
 According to the given condition,
 $x_1 + x_2 = -2a, x_1x_2 = -b^2$
 $y_1 + y_2 = -2p, y_1y_2 = -q^2$

The equation of the circle with A (x_1, y_1) and B (x_2, y_2) as the end points of diameter is
 $(x - x_1)(x - x_2) + (y - y_1)(y - y_2) = 0$
 $\Rightarrow x^2 + y^2 - x(x_1 + x_2) - y(y_1 + y_2) + x_1x_2 + y_1y_2 = 0$
 $\Rightarrow x^2 + y^2 + 2ax + 2py - b^2 - q^2 = 0$

5. Let p be the altitude, then
 $p = a \sin 60^\circ = \frac{a}{2} \sqrt{3}$.



Since, the triangle is equilateral, therefore centroid, orthocentre, circumcentre and incentre all coincide.

- \therefore radius of the inscribed circle = $\frac{p}{3} = \frac{a}{2\sqrt{3}} = r$

Let x be the side of the square inscribed, then angle in a semicircle being a right angle,
 $x^2 + x^2 = (2r)^2 = 4r^2$

$$\Rightarrow 2x^2 = \frac{4a^2}{12} = \frac{a^2}{3}$$

- \therefore required area = $x^2 = \frac{a^2}{6}$

6. Given, parabola $y = x^2$ (i)
 Straight line $y = 2x - 4$ (ii)

From (i) and (ii), $x^2 - 2x + 4 = 0$

Let $f(x) = x^2 - 2x + 4$

- $\therefore f'(x) = 2x - 2$

For least distance, $f'(x) = 0$

$$\Rightarrow 2x - 2 = 0 \Rightarrow x = 1$$

- $\therefore y = (1)^2 = 1$ [From (i)]

So the point least distance from the line is (1, 1).

7. For parabola, $y^2 = 8x$

$$\Rightarrow 4a = 8 \Rightarrow a = 2$$

vertex of $y^2 = 8x$ is $O \equiv (0, 0)$

Now, end points of latus rectum are

$$L(a, 2a); L'(a, -2a) \Rightarrow L(2, 4); L'(2, -4)$$

- \therefore the circle passes through the points (0,0), (2,4) and (2,-4).

All the three points are satisfied by the option (C).

- \therefore Option (C) is the correct answer.



8. Eccentricity of $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ is $e = \sqrt{\frac{a^2 + b^2}{a^2}}$

Eccentricity of conjugate hyperbola is

$$e' = \sqrt{\frac{a^2 + b^2}{b^2}}$$


The given equation of hyperbola can be written as

$$\frac{x^2}{1} - \frac{y^2}{\frac{1}{3}} = 1$$

Here, $a^2 = 1, b^2 = \frac{1}{3}$

$$\therefore e' = \sqrt{\frac{1 + \frac{1}{3}}{\frac{1}{3}}} = \sqrt{4} = 2$$

9. Let AB be the line of intersection of the two circles



$$x^2 + y^2 = 25 \quad \dots(i)$$

$$x^2 + y^2 - 8x + 7 = 0 \quad \dots(ii)$$

Solving (i) and (ii), we get coordinates of A and B.

Subtracting (i) from (ii), we get

$$-8x + 32 = 0 \Rightarrow x = 4$$

From (i), we get $16 + y^2 = 25$

$$\Rightarrow y^2 = 9 \Rightarrow y = \pm 3$$

Thus coordinates of A and B are (4, 3) and (4, -3).

$$\therefore \text{equation of L is } \frac{y-3}{3+3} = \frac{x-4}{4-4} \Rightarrow x-4=0$$

Also coordinates of centre C of second circle is (4, 0).

Hence, CM = Length of perpendicular from

$$C \text{ to the line } L = \left| \frac{4-4}{\sqrt{1}} \right| = 0$$

10. Here $a^2 = 16, b^2 = 9$

$$\text{But } b^2 = a^2(1 - e^2)$$

$$\therefore 9 = 16(1 - e^2)$$

$$\Rightarrow e^2 = 1 - \frac{9}{16} = \frac{7}{16} \Rightarrow e = \frac{\sqrt{7}}{4}$$

Now, foci are (ae, 0), (-ae, 0)

$$\text{i.e., } (\sqrt{7}, 0), (-\sqrt{7}, 0)$$

Centre of the circle is (0, 3)

\therefore radius of the circle is

$$= \sqrt{(\sqrt{7}-0)^2 + (0-3)^2} = \sqrt{7+9} = 4$$

11. The equation of the ellipse is $\frac{x^2}{4} + \frac{y^2}{1} = 1$.

Let e be its eccentricity.

$$\text{Then, } e = \sqrt{1 - \frac{1}{4}} = \frac{\sqrt{3}}{2}$$

The foci of the ellipse are S($\sqrt{3}$, 0) and S'(- $\sqrt{3}$, 0).

$$\text{Eccentricity of the hyperbola} = \frac{1}{e} = \frac{2}{\sqrt{3}}$$

$$\therefore b^2 = a^2 \left(\frac{4}{3} - 1 \right) = \frac{a^2}{3} \quad \dots(i)$$

The hyperbola passes through S($\sqrt{3}$, 0).

$$\therefore \frac{3}{a^2} - 0 = 1 \Rightarrow a^2 = 3$$

Putting $a^2 = 3$ in (i), we get

$$b^2 = 1$$

Hence, the equation of the hyperbola is

$$\frac{x^2}{3} - \frac{y^2}{1} = 1 \text{ i.e., } x^2 - 3y^2 = 3.$$

12. Semi minor axis = b = 2

Semi major axis = a = 4

$$\text{Equation of ellipse is } \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$\Rightarrow \frac{x^2}{16} + \frac{y^2}{4} = 1$$

$$\Rightarrow x^2 + 4y^2 = 16$$

13. Given equation of circle is

$$x^2 + y^2 - 4x - 8y - 5 = 0$$

\therefore Centre = (2, 4) and radius = $\sqrt{4+16+5} = 5$

the circle is intersecting the line $3x - 4y = m$ at two distinct points.

\therefore length of perpendicular from centre on the line < radius


$$\Rightarrow \left| \frac{6-16-m}{5} \right| < 5$$

$$\Rightarrow |10+m| < 25$$

$$\Rightarrow -25 < m+10 < 25 \Rightarrow -35 < m < 15$$



14. Let $P = (1, 0)$, $Q(-1, 0)$ and $A = (x, y)$
 Now, $\frac{AP}{AQ} = \frac{BP}{BQ} = \frac{CP}{CQ} = \frac{1}{3}$
 $\Rightarrow \frac{AP}{AQ} = \frac{1}{3}$
 $\Rightarrow 3AP = AQ \Rightarrow 9AP^2 = AQ^2$
 $\Rightarrow 9(x-1)^2 + 9y^2 = (x+1)^2 + y^2$
 $\Rightarrow 9x^2 - 18x + 9 + 9y^2 = x^2 + 2x + 1 + y^2$
 $\Rightarrow 8x^2 - 20x + 8y^2 + 8 = 0$
 $\Rightarrow x^2 + y^2 - \frac{5}{2}x + 1 = 0 \quad \dots(i)$
 Since, points A, B and C lies on the circle
 \therefore Circumcentre of ABC = Centre of Circle (i)
 $= \left(\frac{5}{4}, 0\right)$




- Given equation of ellipse is
 $x^2 + 4y^2 = 4 \Rightarrow \frac{x^2}{4} + \frac{y^2}{1} = 1 \Rightarrow a = 2, b = 1$
 $\Rightarrow P(2, 1)$
 Let the required equation of ellipse is
 $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$
 Since, the ellipse passes through $(4, 0)$.
 $\therefore a = 4$
 Also, it is passes through $P(2, 1)$.
 $\therefore \frac{4}{16} + \frac{1}{b^2} = 1$
 $\Rightarrow \frac{1}{b^2} = 1 - \frac{1}{4}$
 $\Rightarrow b^2 = \frac{4}{3}$
 \therefore equation of ellipse becomes $\frac{x^2}{16} + \frac{3y^2}{4} = 1$
 $\Rightarrow x^2 + 12y^2 = 16$

- 16.
-
- The lines $x - y - 2 = 0$ and $x - y + 2 = 0$ are parallel, and tangent to the circle.
 Distance between them = diameter of the circle

- $= \frac{2 - (-2)}{\sqrt{1^2 + 1^2}} = \frac{4}{\sqrt{2}} = 2\sqrt{2}$
 Let (h, k) be the centre of the circle.
 Since, $x + y = 0$ is the diameter.
 $\therefore h + k = 0$
 $\Rightarrow h = -k \quad \dots(i)$
 Now, perpendicular drawn from (h, k) to the $x - y - 2 = 0$ is equal to radius.
 $\therefore \left| \frac{h - k - 2}{\sqrt{2}} \right| = \sqrt{2}$
 $\therefore \left| \frac{-k - k - 2}{\sqrt{2}} \right| = \sqrt{2} \quad \dots[\text{From (i)}]$
 $\Rightarrow 2k + 2 = 2 \quad k = 0$
 $\therefore h = 0 \quad \dots[\text{From (i)}]$
 \therefore required equation of circle is

- $(x - 0)^2 + (y - 0)^2 = (\sqrt{2})^2$
 $\Rightarrow x^2 + y^2 = 2$
17. Centre of the given circle = $C(-2, 5)$
 Radius of the circle $CN = CT = \sqrt{g^2 + f^2 - c}$
 $= \sqrt{2^2 + 5^2 + 7} = \sqrt{36} = 6$
 Now, $PC = \sqrt{6^2 + 8^2} = \sqrt{100} = 10$




- We join the external point, $(4, -3)$ to the centre of the circle $(-2, 5)$. Then PT is the minimum distance, from external point P to the circle and PN is the maximum distance.
 Minimum distance = $PT = PC - CT = 10 - 6 = 4$
 Maximum distance = $PN = PC + CN = 10 + 6 = 16$
 So, sum of minimum and maximum distance
 $= 16 + 4 = 20$



18. General equation of circle
 $f(x, y) = (x - h)^2 + (y - k)^2 - a^2 = 0$
 $f(0, \lambda) = h^2 + (\lambda - k)^2 - a^2 = 0$
 $\Rightarrow \lambda^2 - 2k\lambda + k^2 + h^2 - a^2 = 0$
 Equation has equal roots (1, 1)
 \Rightarrow Sum of roots $= -\frac{b}{a} \Rightarrow 2 = 2k \Rightarrow k = 1$
 Also, $f(\lambda, 0) = (\lambda - h)^2 + (0 - k)^2 - a^2 = 0$
 $\lambda^2 - 2h\lambda + h^2 + k^2 - a^2 = 0$
 Equation has roots $\lambda = \frac{1}{2}$ and 2
 Sum of roots $= -\frac{b}{a} \Rightarrow \frac{5}{2} = 2h \Rightarrow h = \frac{5}{4}$
 \therefore Centre $(h, k) = \left(\frac{5}{4}, 1\right)$

19. Here, $y = x^2 - 4x + 3$
 $\Rightarrow y + 1 = x^2 - 4x + 4 \Rightarrow y + 1 = (x - 2)^2$
 So, the vertex is $(2, -1)$.
 and for the circle, $x^2 + (y - 3)^2 = 9$
 Centre $= (0, 3)$
 Distance between vertex and centre
 $= \sqrt{2^2 + (-4)^2} = 2\sqrt{5}$


20. Given parabola is $y^2 = 2ax$
 \therefore Focus $(a/2, 0)$ and directrix is given by
 $x = -a/2$
 Since, circle touches the directrix.
 \therefore Radius of circle = distance from the point



- $(a/2, 0)$ to the line $x = \frac{-a}{2}$
 \therefore Radius $= \frac{\left|\frac{a}{2} + \frac{a}{2}\right|}{\sqrt{1}} = a$
 \therefore Equation of circle is
 $\left(x - \frac{a}{2}\right)^2 + y^2 = a^2$ (i)
 Also, $y^2 = 2ax$ (ii)
 Solving (i) and (ii), we get
 $x = \frac{a}{2}, -\frac{3a}{2}$

Putting these values in $y^2 = 2ax$ we get
 $y = \pm a$ and $x = -3a/2$ gives imaginary values
 of y .
 \therefore Required points are $(a/2, \pm a)$.

21. Let the coordinates of P and Q are
 $(t_1^2, 2t_1), (t_2^2, 2t_2)$ on parabola $y^2 = 4x$.




Slope of OP $= \frac{2t_1 - 0}{t_1^2 - 0} = \frac{2}{t_1}$
 Slope of OQ $= \frac{2t_2 - 0}{t_2^2 - 0} = \frac{2}{t_2}$
 Slope of OP \times slope of OQ $= -1$
 $\Rightarrow \frac{2}{t_1} \cdot \frac{2}{t_2} = -1 \Rightarrow t_1 t_2 = -4$ (i)

Let the coordinates of mid point of PQ are
 (h, k)

- $\therefore t_1^2 + t_2^2 = 2h$ (ii)
 $t_1 + t_2 = k$ (iii)
 Now, $(t_1 + t_2)^2 = t_1^2 + t_2^2 + 2t_1 t_2$
 $\Rightarrow k^2 = 2h + 2(-4)$ [From (i), (ii) and (iii)]
 $\Rightarrow y^2 = 2x - 8$, which is required locus.

22. According to the given condition,
 $\sqrt{(x-2)^2 + y^2} + \sqrt{(x+2)^2 + y^2} = 8$ (i)
 Squaring on both sides, we get
 $(x-2)^2 + y^2 = 64 + (x+2)^2 + y^2$
 $-2 \times 8 \sqrt{(x+2)^2 + y^2}$
 $\Rightarrow -4x = 64 + 4x - 16 \sqrt{(x+2)^2 + y^2}$
 $\Rightarrow -x - 8 = -2 \sqrt{(x+2)^2 + y^2}$
 Put $y = 3$ in (i), we get
 $-x - 8 = -2 \sqrt{(x+2)^2 + 9}$
 Squaring on both sides, we get
 $x^2 + 64 + 16x = 4(x+2)^2 + 36$
 $3x^2 = 12$
 $\Rightarrow x^2 = 4$
 $\Rightarrow x = \pm 2$

23. Equation of auxiliary circle is $x^2 + y^2 = 9$ (i)
 Equation of AB i.e., AM is $\frac{x}{3} + \frac{y}{1} = 1$ (ii)
 Solving (i) and (ii), we get $M\left(-\frac{12}{5}, \frac{9}{5}\right)$



Now, area of $\Delta AOM = \frac{1}{2} \times OA \times MN$


$$= \frac{1}{2} \times 3 \times \frac{9}{5} = \frac{27}{10} \text{ sq. unit.}$$

24. In the given figure, S is focus whose coordinates are $(ae, 0)$.

$\therefore \Delta ABS$ is an equilateral triangle.

\therefore Area of $\Delta ABS = \frac{1}{2} \times AB \times OS = \frac{\sqrt{3}}{4} (\text{side})^2$

$\Rightarrow \frac{1}{2} \times 2b \times ae = \frac{\sqrt{3}}{4} (2b)^2$



$\Rightarrow ae = \sqrt{3} b \quad \dots(i)$

Also, $b^2 = a^2 (1 - e^2)$

$\Rightarrow \left(\frac{ae}{\sqrt{3}}\right)^2 = a^2 (1 - e^2) \quad \dots[\text{From (i)}]$

$\Rightarrow e^2 = 3 - 3e^2$

$\Rightarrow e = \frac{\sqrt{3}}{2}$

25. The auxiliary circle of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

is $x^2 + y^2 = a^2$

Area of this circle = πa^2

$\therefore \pi a^2 = 2 \times \pi ab$

$\Rightarrow a = 2b$

Eccentricity of ellipse = $\sqrt{1 - \frac{b^2}{a^2}}$

$$= \sqrt{1 - \frac{b^2}{4b^2}} = \frac{\sqrt{3}}{2}$$

26. Locus of the point P, if A and B are fixed and $PA + PB = \text{constant}$, is an ellipse.

We have, $PA + PB = 4$, which is a constant.

\therefore Locus of the point P is an ellipse.

27. Locus of the point P, if A and B are fixed and

$\angle APB = \frac{\pi}{2}$, is a circle with diameter AB.

But, we have $PA^2 + PB^2 = \text{constant}$.

\therefore Locus of the point P is a circle.

28. $y = 7x - 25 \quad \dots(i)$

and $x^2 + y^2 = 25$

$\therefore x^2 + (7x - 25)^2 = 25$

$\Rightarrow x^2 + 49x^2 + 625 - 350x = 25$

$\Rightarrow 50x^2 - 350x + 600 = 0$

$\Rightarrow x^2 - 7x + 12 = 0 \Rightarrow x = 3, 4$

Substituting $x = 3, 4$ in (i), we get

$y = 21 - 25 \Rightarrow y = -4, y = 28 - 25 \Rightarrow y = 3$

Let $A \equiv (3, -4), B \equiv (4, 3)$

Using distance formula, we get

$AB = \sqrt{(3-4)^2 + (-4-3)^2}$

$= \sqrt{1+49} = \sqrt{50} = 5\sqrt{2}$



Hints



Classical Thinking

1. Adding 1 to even integers give odd integers.
6. $A \cup B = A$ if every element of B is contained in A i.e. $B \subset A$
9. There is no real number which is both rational as well as irrational.
11. $A - B = \{x: x \in A \text{ and } x \notin B\} = A \cap B'$
12. $A - (B \cup C) = (A - B) \cap (A - C)$
15. $A = \{1, 2, 3, 4, 5, \dots\}$, $B = \{2, 4, 6, 8, \dots\}$
 $\therefore A \cap B = \{2, 4, 6, 8, \dots\}$
16. $B = \{2, 4, 6, 8, \dots\}$, $C = \{1, 3, 5, 7, \dots\}$
 $B \cap C = \phi$
17. $A = \{2, 4, 6, 8, 10, 12, 14, 16, 18, 20, \dots\}$
 $B = \{4, 8, 12, 16, 20, \dots\}$
 $A \cup B = \{2, 4, 6, 8, 10, 12, 14, 16, 18, 20, \dots\}$
19. $B \cup C = \{1, 3, 4, 5, 6, 7, 8, 9\}$
 $A \cap B = \{5, 7\}$, $A \cap C = \{4, 8\}$
 $A \cap (B \cup C) = \{4, 5, 7, 8\}$
 $(A \cap B) \cup (A \cap C) = \{4, 5, 7, 8\}$
20. $B \cap C = \{ \}$
 $A \cup (B \cap C) = \{2, 4, 5, 7, 8\}$
 $(A \cup B) \cap (A \cup C) = \{2, 4, 5, 7, 8\}$
21. $A = \{2, 4, 6, 8, 10, \dots\}$, $B = \{5, 10, 15, 20, \dots\}$
 $C = \{10, 20, 30, 40, \dots\}$
and $(A \cap B) = \{10, 20, 30, \dots\}$
 $\therefore (A \cap B) \cap C = \{10, 20, 30, \dots\}$
22. $(A \cap B)' = A' \cup B'$
23. Since A and B are disjoint,
 $\therefore A \cap B = \phi$
 $\therefore n(A \cap B) = 0$
Now $n(A \cup B) = n(A) + n(B) - n(A \cap B)$
 $= n(A) + n(B) - 0$
 $= n(A) + n(B)$.
24. Since A, B, C are disjoint sets.
 $\therefore n(A \cup B \cup C) = n(A) + n(B) + n(C) = 21$
25. $n(A) = 25$, $n(B) = 20$ and $n(A \cup B) = 35$
 $n(A \cup B) = n(A) + n(B) - n(A \cap B)$
 $35 = 25 + 20 - n(A \cap B)$
 $\Rightarrow n(A \cap B) = 10$
26. $n(U) = 100$
 $A =$ Students who play cricket, $n(A) = 60$
 $B =$ Students who play volleyball, $n(B) = 50$
 $A \cap B =$ Students who play both the games,
 $n(A \cap B) = 28$
 \therefore Number of students who play atleast one game
 $= n(A \cup B) = n(A) + n(B) - n(A \cap B) = 82$
27. $T =$ Set of members who like tea, $n(T) = 11$
 $C =$ Set of members who like coffee, $n(C) = 14$
 $\therefore n(T \cup C) = 20$
 $T \cap C' =$ Set of members who like only tea and not coffee.
 $\therefore n(T \cup C') = n(T) - n(T \cap C)$
 $T \cap C =$ Set of members who like both tea and coffee
 $\therefore n(T \cap C) = n(T) + n(C) - n(T \cup C) = 5$
 $\therefore n(T \cap C) = 5$
 $\therefore n(T \cup C') = n(T) - n(T \cap C) = 11 - 5 = 6$
28. $A \times B = \{(1, 0), (1, 1), (2, 0), (2, 1)\}$
30. $n(A \times A \times B) = n(A) \cdot n(A) \cdot n(B) = 3 \times 3 \times 4 = 36$
32. $A = \{2, 3\}$, $B = \{2, 4\}$, $C = \{4, 5\}$
 $\therefore (B \cap C) = \{4\}$
 $\therefore A \times (B \cap C) = \{2, 3\} \times \{4\} = \{(2, 4), (3, 4)\}$
33. $A \cap B = \{3\}$ and $A = \{1, 2, 3\}$
35. $A - B = \{1\}$, $B - C = \{4\}$
 $\therefore (A - B) \times (B - C) = \{(1, 4)\}$
36. Since $(a, 2)$, $(b, 3)$, $(c, 2)$, $(d, 3)$, $(e, 2)$ are elements of $A \times B$
 $\therefore a, b, c, d, e \in A$ and $2, 3 \in B$
38. $A = \{a, b\}$, $B = \{1, 2, 3\}$
 $\therefore A \times B = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$
 $B \times A = \{(1, a), (1, b), (2, a), (2, b), (3, a), (3, b), (3, c)\}$
 $\therefore (A \times B) \cap (B \times A) = \phi$
39. $(Y \times A) \cap (Y \times B) = Y \times (A \cap B) = Y \times \phi = \phi$
40. $\text{Dom}(R) = \{1, 2, 3\}$
42. Since $x \not\prec x$ therefore R is not reflexive. Also $x < y$ does not imply that $y < x$ So R is not symmetric. Let $x R y$ and $y R z$. Then, x, y and $y < z \Rightarrow x < z$ i.e., $x R z$. Hence, R is transitive.



$$44. f(x) = x^2 - 3x + 2 \Rightarrow f(-1) = (-1)^2 - 3(-1) + 2$$

$$45. f(x) = x^2 - 3x + 2 \\ f(a+h) = (a+h)^2 - 3(a+h) + 2 \\ = a^2 + (2a-3)h - 3a + 2 + h^2$$

$$46. f(x) = x^2 + \frac{1}{x} \\ f\left(\frac{1}{x}\right) = \left(\frac{1}{x}\right)^2 + \frac{1}{\left(\frac{1}{x}\right)} = \frac{1}{x^2} + x$$

$$47. f(x) = x^2 - 6x + 9, 0 \leq x \leq 4 \\ f(3) = (3)^2 - 6(3) + 9 = 0$$

$$48. f(x) = x^2 - 6x + 5, 0 \leq x \leq 4 \\ f(8) \text{ does not exist (since } x = 8 \text{ does not belong to the domain of } f).$$

$$49. f(x) = ax + 6 \Rightarrow f(1) = a(1) + 6 = a + 6 \\ f(1) = 11 \Rightarrow 11 = a + 6 \Rightarrow a = 5$$

$$50. f(a+1) - f(a-1) \\ = 4(a+1) - (a+1)^2 - [4(a-1) - (a-1)^2] \\ = 4(2-a)$$

$$52. \frac{3x^2 + 7x - 1}{3} = x^2 + \frac{7}{3}x - \frac{1}{3} \text{ is a polynomial function.}$$

55. As $f(b)$ is not defined, f is not a function.

$$56. y = 2x - 3 \Rightarrow x = \frac{y+3}{2} \\ \Rightarrow f^{-1}(y) = \frac{y+3}{2} \Rightarrow f^{-1}(x) = \frac{x+3}{2}$$

$$57. g[f(x)] = 5[f(x)] - 6 = 5x^2 - 6$$

$$58. \text{ Since } f(x) = 3x - 1, g(x) = x^2 + 1 \\ \therefore f[g(x)] = 3[g(x)] - 1 = 3[x^2 + 1] - 1 = 3x^2 + 2$$

$$59. f(f(x)) = f(x^2 + 1) = (x^2 + 1)^2 + 1 = x^4 + 2x^2 + 2$$

$$60. (f \circ g)(x) = f[g(x)] = f(x^3 + 1) = (x^3 + 1)^2$$

$$61. f\left(f\left(\frac{1}{x}\right)\right) = f\left(1 - \frac{1}{1/x}\right) = f(1-x) = \frac{x}{x-1}$$

$$62. f(x) = \frac{x-1}{x+1} \\ \Rightarrow f\left(\frac{1}{f(x)}\right) = f\left(\frac{x+1}{x-1}\right) = \frac{\frac{x+1}{x-1} - 1}{\frac{x+1}{x-1} + 1} = \frac{1}{x}$$

$$63. f[g(x)] = \frac{3[g(x)] + 4}{5[g(x)] - 7} = \frac{3\left[\frac{7x+4}{5x-3}\right] + 4}{5\left[\frac{7x+4}{5x-3}\right] - 7} = x$$

$$65. -1 \leq 5x \leq \frac{-1}{5} \Rightarrow \frac{-1}{5} \leq x \leq \frac{1}{5} \\ \text{Hence, domain is } \left[\frac{-1}{5}, \frac{1}{5}\right].$$

$$66. \text{ For } \text{Dom}(f), 5x - 7 > 0 \Rightarrow x > \frac{7}{5}$$

$$\text{Hence, } D_f = \left(\frac{7}{5}, \infty\right)$$

$$67. f(x) = \frac{(x-2)(x-1)}{(x-2)(x+3)}$$

Hence, domain is $\{x : x \in \mathbb{R}, x \neq 2, x \neq -3\}$.

$$68. \text{ For } x = -3, 3, |x^2 - 9| = 0 \\ \text{Therefore, } \log|x^2 - 9| \text{ does not exist at } x = -3, 3.$$

Hence, domain of function is $\mathbb{R} - \{-3, 3\}$

$$69. \log\left\{\frac{5x-x^2}{6}\right\} \geq 0 \Rightarrow \frac{5x-x^2}{6} \geq 1 \\ \Rightarrow x^2 - 5x + 6 \leq 0 \text{ or } (x-2)(x-3) \leq 0. \\ \text{Hence, } 2 \leq x \leq 3.$$



Critical Thinking

- \subset is a relation between two sets and \emptyset is not a set.
- There is no real number x such that $x^2 + 1 = 0$.
- Q is not a null set because $Q = \{0\}$
- Number of proper subsets of $A = 2^n - 1$
 $= 2^5 - 1$...[$\because o(A) = 5$]
 $= 32 - 1 = 31$
- $A - B$ is the set of those elements of A which are not common with B .
- $A - B = A$ iff A and B have no element in common.
- $A - B$ and $B - A$ are always disjoint and hence $A - B = B - A$ only if either of these is \emptyset i.e., if $A \subset B$ and $B \subset A$ i.e., if $A = B$.
- $A - B$, $B - A$ and $A \cap B$ are pairwise disjoint and their union is $A \cup B$.
- $A \cup B = \{1, 2, 3, 4, 6\}$
 $\Rightarrow (A \cup B)' = \{5, 7, 8\}$




13. $A = \{3, 4\}, B = \{-3, 4\}, A \cap B = \{4\}$
 14. $B = \{-3, 4\}, C = \{3, 5\}, B \cup C = \{-3, 3, 4, 5\}$
 15. $C = \{1, 3, 5, 7, \dots\}, D = \{2, 3, 5, 7, 11, \dots\}$
 $C \cap D = \{3, 5, 7, 11, \dots\}$
 16. $A = \{2, 4, 6, 8, 10, \dots\}, B = \{5, 10, 15, 20, \dots\},$
 $C = \{10, 20, 30, 40, \dots\}$
 and $(B \cup C) = \{5, 10, 15, 20, \dots\}$
 $\therefore A \cap (B \cup C) = \{10, 20, 30, \dots\}$

17. $n(A) = n(X) - n(A') = 19$
 $n(B) = n(X) - n(B') = 14$
 $n(A \cap B) = n(X) - n(A \cap B)' = 5$
 $\therefore n(A \cup B) = n(A) + n(B) - n(A \cap B) = 28$
 18. For any $(a, b) \in A \times B, a \in A$ and $b \in B$.
 Now (a, b) will belong to $B \times A$ only if $a \in B$
 and $b \in A$ and that can happen only if
 $A \cap B \neq \phi$. But, in this case $A \cap B = \phi$.
 $\therefore (A \times B) \cap (B \times A) = \phi$


19. $A \cap (A \cup B)' = A \cap (A' \cap B')$
 $\dots[\because (A \cup B)' = A' \cap B']$
 $= (A \cap A') \cap B' \dots[\text{by associative law}]$
 $= \phi \cap B', \dots[\because A \cap A' = \phi]$
 $= \phi$.

20. From Venn-Euler's diagram,



- $\therefore (A - B) \cup (B - A) \cup (A \cap B) = A \cup B$

21. From Venn-Euler's Diagram,



Clearly, $\{(A - B) \cup (B - C) \cup (C - A)\}'$
 $= A \cap B \cap C$.

22. $A = \{4, 5\}, B = \{-6, -7\}, C = \{-7, 10\}$
 $(B \cap C) = \{-7\} \Rightarrow A \cap (B \cap C) = \phi$

23. $A = \{x/6x^2 + x - 15 = 0\}$
 $\therefore 6x^2 + x - 15 = 0$
 $\Rightarrow (3x + 5)(2x - 3) = 0$

- $\therefore x = -\frac{5}{3}$ or $x = \frac{3}{2}$

$$\Rightarrow A = \left\{-\frac{5}{3}, \frac{3}{2}\right\}$$

$$\text{Similarly, } B = \left\{1, \frac{3}{2}\right\} \text{ and } C = \left\{-1, \frac{3}{2}\right\}$$

$$A \cap B \cap C = \left\{\frac{3}{2}\right\}$$

24. Let $A \equiv$ set of persons who take tea and
 $B \equiv$ set of persons who take coffee
 $n(A \cup B) = 50, n(A) = 35, n(B) = 25$
 $\therefore n(A \cap B) = 10$
 Hence, $n(A - B) = n(A) - n(A \cap B)$
 $= 35 - 10 = 25$
25. $P =$ Set of children who like pizza
 $B =$ Set of children who like burger
 $n(P) = 62, n(B) = 47, n(P \cap B) = 36$
 $(P \cap B)' =$ Set of children who like pizza but
 not burger
 $\therefore n(P \cap B)' = n(P) - n(P \cap B) = 62 - 36 = 26$.
26. $A =$ Set representing no. of consumers using
 Brand A, $n(A) = 15$
 $B =$ Set representing no. of consumers using
 Brand B, $n(B) = 20$
 $A \cap B =$ Set representing no. of consumers
 using both the brands, $n(A \cap B) = 5$
 $A \cup B =$ Set representing no. of consumers
 using atleast one brand.
 $\therefore n(A \cup B) = n(A) + n(B) - n(A \cap B) = 30$
27. Minimum value of $x = 100 - (30 + 20 + 25 + 15)$
 $= 100 - 90 = 10$.
28. $U =$ Universal set of all adults
 $M =$ Set of all males, $F =$ Set of all females
 $V =$ Set of all vegetarians
 Total number of adults = 20
 Total number of males = 8
 \therefore Total number of females = $20 - 8 = 12$
 Total number of vegetarian = 9
 Total number of male vegetarian = 5
 \therefore Total number of female vegetarian = $9 - 5 = 4$
 \therefore Total number of female non-vegetarian
 $= 12 - 4 = 8$
29. $C =$ Set of students who play chess
 $T =$ Set of students who play table tennis
 $M =$ Set of students who play carrom
 $\therefore n(X) = 120, n(C) = 46, n(T) = 30, n(M) = 40$
 $n(C \cap T) = 14, n(T \cap M) = 10, n(C \cap M) = 8,$
 $n(C \cup T \cup M)' = 30$
 $\therefore n(C \cup T \cup M) = n(X) - n(C \cup T \cup M)' = 90$



- $(C \cap T \cap M)$ = Set of students who play chess, table tennis and carrom.
 $\therefore n(C \cup T \cup M)$
 $= n(C) + n(T) + n(M) - n(C \cap T) - n(T \cap M) - n(C \cap M) + n(C \cap T \cap M)$
 $\therefore 90 = 46 + 30 + 40 - 14 - 10 - 8 + n(C \cap T \cap M)$
 $\therefore n(C \cap T \cap M) = 6$
30. Since, $y = e^x$, $y = e^{-x}$ will meet, when $e^x = e^{-x}$
 $\Rightarrow e^{2x} = 1$,
 $\therefore x = 0, y = 1$
 \therefore A and B meet on (0, 1)
 $\therefore A \cap B \neq \phi$
31. Let A denote the set of Americans, who like cheese and let B denote the set of Americans, who like apples.
 Let Population of Americans be 100.
 Then $n(A) = 63$, $n(B) = 76$
 Now, $n(A \cup B) = n(A) + n(B) - n(A \cap B)$
 $= 63 + 76 - n(A \cap B)$
 $\therefore n(A \cup B) + n(A \cap B) = 139$
 $\Rightarrow n(A \cap B) = 139 - n(A \cup B)$
 But, $n(A \cup B) \leq 100$
 $\therefore -n(A \cup B) \geq -100$
 $\therefore 139 - n(A \cup B) \geq 139 - 100 = 39$
 $\therefore n(A \cap B) \geq 39$ i.e., $39 \leq n(A \cap B)$
 Again, $A \cap B \subseteq A$, $A \cap B \subseteq B$
 $\therefore n(A \cap B) \leq n(A) = 63$ and
 $n(A \cap B) \leq n(B) = 76$
 $\therefore n(A \cap B) \leq 63$
 Then, $39 \leq n(A \cap B) \leq 63 \Rightarrow 39 \leq x \leq 63$
32. Since, $8^n - 7n - 1 = (7 + 1)^n - 7n - 1$
 $= 7^n + {}^nC_1 7^{n-1} + {}^nC_2 7^{n-2} + \dots + {}^nC_{n-1} 7 + {}^nC_n - 7n - 1$
 $= {}^nC_2 7^2 + {}^nC_3 7^3 + \dots + {}^nC_n 7^n$,
 $({}^nC_0 = {}^nC_n, {}^nC_1 = {}^nC_{n-1}$ etc.)
 $= 49[{}^nC_2 + {}^nC_3(7) + \dots + {}^nC_n 7^{n-2}]$
 $\therefore 8^n - 7n - 1$ is a multiple of 49 for $n \geq 2$
 For $n = 1$, $8^n - 7n - 1 = 8 - 7 - 1 = 0$
 For $n = 2$, $8^n - 7n - 1 = 64 - 14 - 1 = 49$
 $\therefore 8^n - 7n - 1$ is a multiple of 49 for $n \in \mathbb{N}$.
 \therefore X contains elements which are multiples of 49 and clearly Y contains all multiples of 49.
 $\therefore X \subseteq Y$
33. The given set is a cartesian product containing 6 elements. Only $A \times (B \cup C)$ contains 6 elements.
34. Here $1, 2, 3 \in A$ & $3, 5 \in B$
 $\therefore A \times B = \{1, 2, 3\} \times \{3, 5\}$
 \therefore The remaining elements are : (1, 5), (2, 3), (3, 5)
35. Clearly, A is the set of all first elements in ordered pairs in $A \times B$ and B is the set of all second elements in $A \times B$.
36. $(1, 4), (2, 6), (3, 6) \in A \times B$
 $\Rightarrow \{1, 2, 3\} \subset A$ and $\{4, 6\} \subset B$
 \therefore A has 3 elements and B has 2 elements.
37. Number of relations on the set A = Number of subsets of $(A \times A) = 2^{n^2}$, [$\because n(A \times A) = n^2$].
38. $n(A \times A) = n(A)$. $n(A) = 3^2 = 9$
 So, the total number of subsets of $A \times A$ is 2^9 and a subset of $A \times A$ is a relation over the set A.
39. Since, $(-1, 0) \in A \times A$ and $(0, 1) \in A \times A$
 $\therefore (-1, 0) \in A \times A \Rightarrow -1, 0 \in A$
 and $(0, 1) \in A \times A \Rightarrow 0, 1 \in A$
 $\therefore \{-1, 0, 1\} \in A$
41. $R_2 \subseteq A \times B$, so it is a relation from A to B.
42. Number of relations from A to B = $2^{o(A) \cdot o(B)}$
43. Since, $R = \{(x, y) \mid x, y \in \mathbb{Z}, x^2 + y^2 \leq 4\}$
 $\therefore R = \{(-2, 0), (-1, 0), (-1, 1), (0, -1), (0, 1), (0, 2), (0, -2), (1, 0), (1, 1), (2, 0)\}$
 Hence, Domain of R = $\{-2, -1, 0, 1, 2\}$.
46. Since R is an equivalence relation on set A, therefore $(a, a) \in R$ for all $a \in A$. Hence, R has at least n ordered pairs.
47. Let $(a, b) \in R$
 Then, $(a, b) \in R \Rightarrow (b, a) \in R^{-1}$
 $\Rightarrow (b, a) \in R \quad \dots [\because R = R^{-1}]$
 So, R is symmetric.
48. For any $a \in \mathbb{N}$, we find that $a|a$, therefore R is reflexive but R is not symmetric, because aRb does not imply that bRa .
49. The relation is not symmetric, because $A \subseteq B$ does not imply that $B \subset A$. But it is anti-symmetric because $A \subset B$ and $B \subset A \Rightarrow A = B$
50. The given relation is not reflexive and transitive but it is symmetric, because $x^2 + y^2 = 1 \Rightarrow y^2 + x^2 = 1$.
51. R is a relation from $\{11, 12, 13\}$ to $\{8, 10, 12\}$ defined by $y = x - 3 \Rightarrow x - y = 3$
 $\therefore R = \{11, 8\}, \{13, 10\}$.
 Hence, $R^{-1} = \{(8, 11), (10, 13)\}$
52. We have, $R = \{(1, 3); (1, 5); (2, 3); (2, 5); (3, 5); (4, 5)\}$
 $R^{-1} = \{(3, 1), (5, 1), (3, 2), (5, 2); (5, 3); (5, 4)\}$
 Hence, $\text{Ro}R^{-1} = \{(3, 3); (3, 5); (5, 3); (5, 5)\}$



53. $f(x) = f(x+1)$
 $\therefore x^2 - 2x + 3 = (x+1)^2 - 2(x+1) + 3$
 $\therefore x^2 - 2x = x^2 + 2x + 1 - 2x - 2 \Rightarrow x = 1/2$
54. $f(x) = ax^2 + bx + 2$
 $\therefore f(1) = a(1)^2 + b(1) + 2 = a + b + 2$
 But $f(1) = 3 \Rightarrow 3 = a + b + 2 \Rightarrow a + b = 1 \dots(i)$
 and $f(4) = a(4)^2 + b(4) + 2 = 16a + 4b + 2$
 But $f(4) = 42 \Rightarrow 42 = 16a + 4b + 2$
 $\therefore 40 = 16a + 4b \Rightarrow 4a + b = 10 \dots(ii)$
 By solving, (i) & (ii) $a = 3$ and $b = -2$
55. $f(x) = x + \frac{1}{x} \Rightarrow f(x^3) = x^3 + \frac{1}{x^3}$
 $\therefore [f(x)]^3 = \left(x + \frac{1}{x}\right)^3 = \left(x^3 + \frac{1}{x^3}\right) + 3\left(x + \frac{1}{x}\right)$
 $\therefore [f(x)]^3 = f(x^3) + 3f(x)$
 $\therefore [f(x)]^3 = f(x^3) + 3f\left(\frac{1}{x}\right) \Rightarrow \lambda = 3$
56. $a.f(x) + b.f\left(\frac{1}{x}\right) = \frac{1}{x} - 5$
 On replacing x by $\frac{1}{x}$, $b.f(x) + a.f\left(\frac{1}{x}\right) = x - 5$
 Solving two equations,
 $f(x) = \frac{1}{a^2 - b^2} \left(\frac{a}{x} - bx\right) - \frac{5}{a+b}$
 $\therefore f(2) = \frac{3(2b-3a)}{2(a^2-b^2)}$
58. As $f(a)$ is not unique,
 $\therefore f$ is not a function.
60. $[x] = I$ (Integers only).
61. Let $f(x) = x^2 + \sin^2 x$
 Here, $f(-x) = f(x)$
 $\therefore f(x)$ is an even function.
62. If $f(x) = \sqrt{1+x+x^2} - \sqrt{1-x+x^2}$, then
 $f(-x) = \sqrt{1-x+x^2} - \sqrt{1+x+x^2}$
 $\therefore f(-x) = -f(x)$
 So, $f(x)$ is an odd function.
63. The general expression for the function satisfying $f(x+y) = f(x)f(y)$ for all $x, y \in \mathbb{R}$ is $f(x) = [f(1)]^x = a^x$ for all $x, y \in \mathbb{R}$. [$\because f(1) = a$]
64. $f^{-1}(y) = \{x \in \mathbb{R} : y = f(x)\}$
 $\Rightarrow f^{-1}(2) = \{x \in \mathbb{R} : 2 = f(x)\}$
 $= \{x \in \mathbb{R} : x^2 - 3x + 4 = 2\}$
 $= \{x \in \mathbb{R} : x^2 - 3x + 2 = 0\} = \{1, 2\}$

$$65. f(f(x)) = \frac{1}{1-f(x)} = \frac{1}{1-\frac{1}{1-x}} = \frac{x-1}{x}$$

$$\therefore f[f(f(x))] = f\left(\frac{x-1}{x}\right) = \frac{x}{x-x+1} = x$$

$$66. f(x) = \frac{x+3}{4x-5}$$

$$\therefore f(t) = \frac{t+3}{4t-5} = \frac{\left(\frac{3+5x}{4x-1}\right)+3}{4\left(\frac{3+5x}{4x-1}\right)-5} = x$$

$$67. (\text{gof})(1) = g(f(1)) = g(4) = 8,$$

 $(\text{gof})(2) = g(f(2)) = g(5) = 7$
 and $(\text{gof})(3) = g(f(3)) = g(6) = 9$

$$68. f(x) = \frac{x-1}{x+1} \Rightarrow \frac{f(x)+1}{f(x)-1} = \frac{2x}{-2}$$

$$\Rightarrow x = \frac{1+f(x)}{1-f(x)}$$

$$\therefore f(\alpha x) = \frac{\alpha x - 1}{\alpha x + 1} = \frac{(\alpha + 1)f(x) + \alpha - 1}{(\alpha - 1)f(x) + \alpha + 1}$$

$$69. \text{ Given, } (\text{gof})\left(-\frac{5}{3}\right) - (\text{fog})\left(-\frac{5}{3}\right)$$

 $= g\left\{f\left(-\frac{5}{3}\right)\right\} - f\left\{g\left(-\frac{5}{3}\right)\right\}$
 $= g(-2) - f\left(\frac{5}{3}\right) = 2 - 1 = 1$

$$70. f(-1) = f(1) = 1$$

 \therefore function is many-one function.
 $\therefore f$ is neither one-one nor onto.

$$71. f'(x) = \frac{1}{(1+x)^2} > 0 \forall x \in [0, \infty)$$

and range $\in [0, 1)$

\Rightarrow function is one-one but not onto.

$$72. f(x) = \frac{x-2}{x-3}, x \neq 3$$

$$\text{Let } y = f(x) \Rightarrow y = \frac{x-2}{x-3} \Rightarrow x = \frac{2-3y}{1-y}$$

$$\Rightarrow y \neq 1 \Rightarrow \text{Range of } f(x) \text{ is } \mathbb{R} - \{1\}$$

So, f is onto

For one-one, let $f(x_1) = f(x_2)$

$$\Rightarrow \frac{x_1-2}{x_1-3} = \frac{x_2-2}{x_2-3} \Rightarrow x_1 = x_2$$

Hence, f is one-one.



73. Let $f(x_1) = f(x_2) \Rightarrow [x_1] = [x_2] \Rightarrow x_1 = x_2$
 {For example, if $x_1 = 1.4$, $x_2 = 1.5$, then
 $[1.4] = [1.5] = 1$ }

\therefore f is not one-one.

Also, f is not onto as its range I (set of integers) is a proper subset of its co-domain R .

74. Let $x_1, x_2 \in R$, then $f(x_1) = \cos x_1$,
 & $f(x_2) = \cos x_2$, Now $f(x_1) = f(x_2)$
 $\Rightarrow \cos x_1 = \cos x_2 \Rightarrow x_1 = 2n\pi \pm x_2$

$\Rightarrow x_1 \neq x_2$,

\therefore it is not one-one.

Again the value of f -image of x lies in between -1 to 1

$\Rightarrow f[R] = \{f(x) : -1 \leq f(x) \leq 1\}$

So other numbers of co-domain (besides -1 and 1) is not f -image. $f[R] \in R$, so it is also not onto. So this mapping is neither one-one nor onto.

75. $x^2 - 6x + 7 = (x - 3)^2 - 2$

Here, minimum value is -2 and maximum ∞ .
 Hence, range of function is $[-2, \infty)$.

76. For domain, take $\frac{x}{1+x} \geq 0$

$\therefore D_f = (-\infty, -1) \cup [0, \infty)$

77. $1 + x \geq 0$

$\Rightarrow x \geq -1$; $1 - x \geq 0$

$\Rightarrow x \leq 1, x \neq 0$

Hence, domain is $[-1, 1] - \{0\}$.

78. $y = \sin^{-1} \left[\log_3 \left(\frac{x}{3} \right) \right]$

$\therefore -1 \leq \log_3 \left(\frac{x}{3} \right) \leq 1 \quad \therefore \frac{1}{3} \leq \frac{x}{3} \leq 3$

$\therefore 1 \leq x \leq 9$

$\therefore x \in [1, 9]$

79. $f(x)$ is defined, if

$x^2 - 5x + 6 \geq 0$ and $2x + 8 - x^2 \geq 0$

$\Rightarrow (x - 2)(x - 3) \geq 0$ and $(x - 4)(x + 2) \leq 0$

$\therefore x \in (-\infty, 2] \cup [3, \infty)$ and $x \in [-2, 4]$

$\therefore x \in [-2, 2] \cup [3, 4]$

80. Domain of $f(x) = R - \{3\}$,

and for Range : $x \neq 3 \Rightarrow x < 3$ or $x > 3$

Now, $x < 3 \Rightarrow x - 3 < 0 \Rightarrow |x - 3| = -(x - 3)$

$\Rightarrow f(x) = \frac{-(x-3)}{x-3} = -1$

Similarly, for $x > 3$, $f(x) = 1$

\therefore Range $(f) = \{1, -1\}$.

81. $\text{Dom}(f) = R - \left\{ -\frac{2}{3} \right\}$

For Range (f) , let $y = f(x) = \frac{1}{3x+2}$

$\therefore 3x + 2 = \frac{1}{y} \Rightarrow x = \frac{1}{3} \left(2 - \frac{1}{y} \right)$

x is real if $y \neq 0$. Hence, $R_f = R - \{0\}$

82. $f(x)$ is defined for all $x \in R - \{0\}$.

So, $\text{dom}(f) = R - \{0\}$

Let $y = \frac{1+x^2}{x^2}$

$\Rightarrow x = \pm \sqrt{\frac{1}{y-1}}$

For x to be real, $y - 1 > 0 \Rightarrow y \in (1, \infty)$

83. $f(x)$ is defined for all $x \in R$. So, $\text{dom}(f) = R$.

Let $y = f(x) \Rightarrow y = \frac{x}{1+x^2}$

$\therefore x = \frac{1 \pm \sqrt{1-4y^2}}{2y}$

For x to be real, $1 - 4y^2 \geq 0$ and $y \neq 0$

$\Rightarrow -\frac{1}{2} \leq y \leq \frac{1}{2}$ and $y \neq 0$

84. $f(x)$ is defined for $x^2 + x - 6 \neq 0$, i.e., $x \neq -3, 2$

$\therefore \text{Dom}(f) = R - \{-3, 2\}$

Let $y = \frac{x^2 - 3x + 2}{x^2 + x - 6} = \frac{x-1}{x+3}$

$\Rightarrow x = \frac{3y+1}{y-1}$

x is real for $y - 1 \neq 0$, i.e., $y \neq 1$

Hence, range $(f) = R - \{1\}$

85. Here, $f(x) = \sqrt{x^2 + x + 1}$

$\Rightarrow y^2 = x^2 + x + 1$

$\Rightarrow x^2 + x + (1 - y^2) = 0$

$\Rightarrow x = \frac{-1 \pm \sqrt{1 - 4(1 - y^2)}}{2}$

$\Rightarrow x = \frac{-1 \pm \sqrt{4y^2 - 3}}{2}$

For x real, $4y^2 - 3 \geq 0$

$\therefore y \geq \pm \frac{\sqrt{3}}{2}$

$\therefore R_f = \left[\frac{\sqrt{3}}{2}, \infty \right)$

**Competitive Thinking**

4. $x^2 = 16 \Rightarrow x = \pm 4$
and $2x = 6 \Rightarrow x = 3$
There is no value of x which satisfies both the above equations. Thus, $A = \phi$.
5. $|2x + 3| < 7 \Rightarrow -7 < 2x + 3 < 7$
 $\Rightarrow -10 < 2x < 4 \Rightarrow -5 < x < 2 \Rightarrow 0 < x + 5 < 7$
6. **Case I:** $0 \leq x < 9$
 $2(3 - \sqrt{x}) + \sqrt{x}(\sqrt{x} - 6) + 6 = 0$
 $\Rightarrow (\sqrt{x})^2 - 8\sqrt{x} + 12 = 0$
 $\Rightarrow \sqrt{x} = 6, 2$
 $\Rightarrow x = 36, 4 \Rightarrow x = 4$
Case II: $x \geq 9$
 $2(\sqrt{x} - 3) + \sqrt{x}(\sqrt{x} - 6) + 6 = 0$
 $\Rightarrow (\sqrt{x})^2 - 4\sqrt{x} = 0$
 $\Rightarrow \sqrt{x} = 0, 4$
 $\Rightarrow x = 0, 16 \Rightarrow x = 16$
 $\Rightarrow S$ contains exactly two elements.
7. The number of non-empty subsets = $2^n - 1$
 $= 2^4 - 1 \quad \dots[\because n = 4]$
 $= 15$.
8. Power set is the set of all subsets.
 $n(A) = 5 \Rightarrow n(P(A)) = 2^5 = 32$
9. $A = \{4, 8, 12, 16, 20, 24, \dots\}$
 $B = \{6, 12, 18, 24, 30, \dots\}$
 $\therefore A \subset B = \{12, 24, \dots\} = \{x : x \text{ is a multiple of } 12\}$.
10. Given set is $\{(a, b) : 2a^2 + 3b^2 = 35, a, b \in \mathbb{Z}\}$
We can see that, $2(\pm 2)^2 + 3(\pm 3)^2 = 35$
and $2(\pm 4)^2 + 3(\pm 1)^2 = 35$
 $\therefore (2, 3), (2, -3), (-2, -3), (-2, 3), (4, 1), (4, -1), (-4, -1), (-4, 1)$ are 8 elements of the set.
 $\therefore n = 8$.
11. Let, $S = \{a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9, a_{10}\}$
 $\therefore A = \{(x, y) : x, y \in S, x \neq y\}$
 $\therefore A = \{(a_1, a_1), (a_1, a_2), (a_1, a_3), \dots (a_1, a_{10}), (a_2, a_1), (a_2, a_2), (a_2, a_3), \dots (a_2, a_{10}), \dots (a_{10}, a_1), (a_{10}, a_2), (a_{10}, a_3), \dots (a_{10}, a_{10})\}$
 $\therefore x \neq y$, removing groups with same elements
i.e. $(a_1, a_1), (a_2, a_2) \dots (a_{10}, a_{10})$, we get
 $n(A) = 9 \times 10 = 90$


12. $n(S) = 10$
 \therefore Number of subsets of S which do not contain the element 6
 $=$ number of subsets containing the remaining nine elements
 $= 2^9 = 512$
13. Since $2^m - 2^n = 56 = 8 \times 7 = 2^3 \times 7$
 $\Rightarrow 2^n(2^{m-n} - 1) = 2^3 \times 7$
 $\therefore n = 3$ and $2^{m-n} = 8 = 2^3$
 $\Rightarrow m - n = 3 \Rightarrow m - 3 = 3 \Rightarrow m = 6$
 $\therefore m = 6, n = 3$
14. $O(S) = O\left(\bigcup_{i=1}^{30} A_i\right) = \frac{1}{10}(5 \times 30) = 15$
Since, element in the union S belongs to 10 of A_i 's
Also, $O(S) = O\left(\bigcup_{j=1}^n B_j\right) = \frac{3n}{9} = \frac{n}{3}$
 $\therefore \frac{n}{3} = 15 \Rightarrow n = 45$
15. $A - (A - B) = A \cap (A \cap B^c)^c = A \cap (A^c \cup B)$
 $= \phi \cup (A \cap B) = A \cap B$
16. $A = B \cap C, B = C \cap A$
 $\Rightarrow A, B$ are equivalent sets.
 $\dots[\because A$ and B are interchangeable in both equations]
18. $A \cap X = B \cap X = \phi$
 $\therefore A$ and X, B and X are disjoint sets
Also, $A \cup X = B \cup X \Rightarrow A = B$
19. If $A \subseteq B$, then $A \cup B = B$
 $\therefore n(A \cup B) = n(B) = 6$
20. $n[(A \cap B)' \cap A]$
 $= n[(A' \cup B') \cap A]$
 \dots [By DeMorgan's law]
 $= n(A' \cap A) \cup n(B' \cap A)$
 \dots [By distributive law]
 $= n(A) - n(A \cap B) = 8 - 2 = 6$
21. $A = \{x \mid x \text{ is a root of } x^2 - 1 = 0\}$
 $= \{x \mid x \text{ is a root of } (x - 1)(x + 1) = 0\}$
 $\Rightarrow x = \pm 1$
 $B = \{x \mid x \text{ is a root of } x^2 - 2x + 1 = 0\}$
 $= \{x \mid x \text{ is a root of } (x - 1)^2 = 0\}$
 $\Rightarrow x = 1$
 $\Rightarrow A \cup B = A$



$$\begin{aligned}
 22. \quad & \text{Since, } 4^n - 3n - 1 = (3 + 1)^n - 3n - 1 \\
 & = 3^n + {}^n C_1 \cdot 3^{n-1} + {}^n C_2 \cdot 3^{n-2} \\
 & \quad + \dots + {}^n C_{n-1} \cdot 3 + {}^n C_n - 3n - 1 \\
 & = {}^n C_2 3^2 + {}^n C_3 \cdot 3^3 + \dots + {}^n C_n \cdot 3^3 \\
 & \quad \dots [{}^n C_0 = {}^n C_n, {}^n C_1 = {}^n C_{n-1}, \text{ etc.}]
 \end{aligned}$$

$\therefore 4^2 - 3n - 1$ is a multiple of 9 for $n \geq 2$.
 for $n = 1$, $4^n - 3n - 1 = 4 - 3 - 1 = 0$,
 for $n = 2$, $4^n - 3n - 1 = 16 - 6 - 1 = 9$,
 $\therefore 4^n - 3n - 1$ is a multiple of 9 for all $n \in \mathbb{N}$.
 $\therefore X$ contains elements, which are multiples of 9,
 and clearly Y contains all multiples of 9.
 $\therefore X \subseteq Y$ i.e. $X \cap Y = X$

23. $N_5 \cap N_7 = N_{35}$,
 [$\because 5$ and 7 are relatively prime numbers].



$\therefore (B - A) \cap (A \cup B)' = \phi$

25. $n(A \cup B) = n(A) + n(B) - n(A \cap B)$
 $= 12 + 9 - 4 = 17$
 Now, $n((A \cup B)^c) = n(U) - n(A \cup B)$
 $= 20 - 17 = 3$


26. $n(A \cup B) = n(A) + n(B) - n(A \cap B)$
 $= 200 + 300 - 100$
 $= 400$

$\therefore n(A' \cap B') = n(U) - n(A \cup B)$
 $= 700 - 400 = 300$






27. $n(A \cup B) = n(A) + n(B) - n(A \cap B)$
 $= 3 + 6 - n(A \cap B)$
 Since, maximum number of elements in
 $A \cap B = 3$

\therefore Minimum number of elements in
 $A \cup B = 9 - 3 = 6$



28. $(A \cup B)^c \cup (A^c \cap B)$
 $= (A^c \cap B^c) \cup (A^c \cap B)$
 $= A^c \cap (B^c \cup B)$
 $= A^c \cap U = A^c$




Since, $A - B = A - (A \cap B)$
 and $B - A = B - (A \cap B)$
 Option (D) is the correct answer.





31. $A - B = A - (A \cap B)$ is correct.
 $A = (A \cap B) \cup (A - B)$ is correct.
 (3) is false.
 \therefore (1) and (2) are true.
- 
32. $n(A - B) = n(A) - n(A \cap B) = 25 - 10 = 15$
33. $n(X) = 60, n(C) = 25, n(T) = 20, n(C \cap T) = 10$
 $\therefore n(C \cup T) = n(C) + n(T) - n(C \cap T)$
 $= 25 + 20 - 10 = 35$
 $\therefore n(C \cup T)' = n(X) - n(C \cup T) = 60 - 35 = 25$
34. $n(X \cap Y) = 12$ and these are 1, 2, 4, 5, 8, 10, 20, 25, 40, 50, 100, 200
35. Let number of newspapers be x . If every student reads one newspaper, the number of students would be $x(60) = 60x$
 Since, every student reads 5 newspapers
 \therefore Numbers of students $= \frac{x \times 60}{5} = 300, x = 25$
36. E : English speakers
 H : Hindi speakers
 $n(H \cup E) = n(H) + n(E) - n(H \cap E)$
 $= 50 + 20 - 10 = 60$
37. $n(M \text{ alone})$
 $= n(M) - n(M \cap C) - n(M \cap P) + n(M \cap P \cap C)$
- 
- $= 100 - 28 - 30 + 18 = 60$
38. $n(C) = 224, n(H) = 240, n(B) = 336$
 $n(H \cap B) = 64, n(B \cap C) = 80$
 $n(H \cap C) = 40, n(C \cap H \cap B) = 24$
 $n(C^c \cap H^c \cap B^c) = n(C \cup H \cup B)^c$
 $= n(U) - n(C \cup H \cup B)$
 $= 800 - [n(C) + n(H) + n(B) - n(H \cap C)$
 $- n(H \cap B) - n(C \cap B) + n(C \cap H \cap B)]$
 $= 800 - [224 + 240 + 336 - 64 - 80 - 40 + 24]$
 $= 800 - 640 = 160$


39. Given $n(N) = 12, n(P) = 16, n(H) = 18,$
 $n(N \cup P \cup H) = 30$ and $n(N \cap P \cap H) = 0$
 From, $n(N \cup P \cup H)$
 $= n(N) + n(P) + n(H) - n(N \cap P) - n(P \cap H)$
 $- n(N \cap H) + n(N \cap P \cap H)$
 $\therefore n(N \cap P) + n(P \cap H) + n(N \cap H) = 16$
 Now, number of pupils taking two subjects
 $= n(N \cap P) + n(P \cap H) + n(N \cap H)$
 $= 16 - 0 = 16.$
40. $n(S \cup P \cup D) = 265, n(S) = 200, n(D) = 110,$
 $n(P) = 55, n(S \cap D) = 60, n(S \cap P) = 30,$
 $n(S \cap D \cap P) = 10,$
 $n(S \cup P \cup D) = n(S) + n(D) + n(P) - n(S \cap D)$
 $- n(D \cap P) - n(P \cap S) + n(S \cap D \cap P)$
 $\therefore 265 = 200 + 110 + 55 - 60 - 30$
 $- n(P \cap D) + 10$
 $\therefore n(P \cap D) = 285 - 265 = 20$
 $\therefore n(P \cap D) - n(P \cap D \cap S) = 20 - 10 = 10$
41. $n(A) = 40\% \text{ of } 10,000 = 4,000$
 $n(B) = 20\% \text{ of } 10,000 = 2,000$
 $n(C) = 10\% \text{ of } 10,000 = 1,000$
 $n(A \cap B) = 5\% \text{ of } 10,000 = 500$
 $n(B \cap C) = 3\% \text{ of } 10,000 = 300$
 $n(C \cap A) = 4\% \text{ of } 10,000 = 400$
 $n(A \cap B \cap C) = 2\% \text{ of } 10,000 = 200$
 We want to find,
 $n(A \cap B^c \cap C^c) = n[A \cap (B \cup C)^c]$
 $= n(A) - n[A \cap (B \cup C)]$
 $= n(A) - n[(A \cap B) \cup (A \cap C)]$
 $= n(A) - [n(A \cap B) + n(A \cap C)$
 $- n(A \cap B \cap C)]$
 $= 4000 - [500 + 400 - 200]$
 $= 4000 - 700 = 3300.$
42. Since, $y = e^x$ and $y = x$ do not meet for any $x \in \mathbb{R}$
 $\therefore A \cap B = \phi$
43. $|a - 5| < 1$ and $|b - 5| < 1$
 $\Rightarrow 4 < a < 6$ and $4 < b < 6$
 $4(a - 6)^2 + 9(b - 5)^2 \leq 36$
 $\Rightarrow \frac{(a - 6)^2}{9} + \frac{(b - 5)^2}{4} \leq 1$



Set A represents square EFGH and Set B represents an ellipse.

$\Rightarrow A \subset B$

44. A = Set of all values $(x, y) : x^2 + y^2 = 25 = 5^2$



$$B = \frac{x^2}{144} + \frac{y^2}{16} = 1 \text{ i.e., } \frac{x^2}{(12)^2} + \frac{y^2}{(4)^2} = 1$$

Clearly, $A \cap B$ consists of four points.

45. Let the original set contains $(2n + 1)$ elements, then subsets of this set containing more than n elements, i.e., subsets containing $(n + 1)$ elements, $(n + 2)$ elements, $(2n + 1)$ elements.

$$\begin{aligned} \therefore \text{Required number of subsets} &= {}^{2n+1}C_{n+1} + {}^{2n+1}C_{n+2} + \dots + {}^{2n+1}C_{2n} + {}^{2n+1}C_{2n+1} \\ &= {}^{2n+1}C_n + {}^{2n+1}C_{n-1} + \dots + {}^{2n+1}C_1 + {}^{2n+1}C_0 \\ &= {}^{2n+1}C_0 + {}^{2n+1}C_1 + {}^{2n+1}C_2 + \dots + {}^{2n+1}C_{n-1} + {}^{2n+1}C_n \\ &= \frac{1}{2}[(1+1)^{2n+1}] = \frac{1}{2}[2^{2n+1}] = 2^{2n} \end{aligned}$$

46. Since, $4^n - 3n - 1 = (3 + 1)^n - 3n - 1$
 $= 3^n + {}^nC_1 3^{n-1} + {}^nC_2 3^{n-2} + \dots + {}^nC_{n-1} 3 + {}^nC_n - 3n - 1$
 $= {}^nC_2 3^2 + {}^nC_3 3^3 + \dots + {}^nC_n 3^n$
 $\dots [{}^nC_0 = {}^nC_n, {}^nC_1 = {}^nC_{n-1} \text{ etc}]$
 $= 9[{}^nC_2 + {}^nC_3(3) + \dots + {}^nC_n 3^{n-1}]$
 $\therefore 4^n - 3n - 1$ is a multiple of 9 for $n \geq 2$.
 For $n = 1, 4^n - 3n - 1 = 4 - 3 - 1 = 0$,
 For $n = 2, 4^n - 3n - 1 = 16 - 6 - 1 = 9$
 $\therefore 4^n - 3n - 1$ is a multiple of 9 for all $n \in \mathbb{N}$
 $\therefore X$ contains elements, which are multiples of 9, and clearly Y contains all multiples of 9.
 $\therefore X \subseteq Y$ i.e., $X \cup Y = Y$.

47. $R = A \times B$.
48. $A - B = \{a\}, B \cap C = \{c, d\}$
 $\therefore (A - B) \times (B \cap C) = \{a\} \times \{c, d\}$
 $= \{(a, c), (a, d)\}$
49. $R \times (P^c \cup Q^c)^c = R \times [(P^c)^c \cap (Q^c)^c]$
 $= R \times (P \cap Q) = (R \times P) \cap (R \times Q)$
50. $A = \{2, 4, 6\}; B = \{2, 3, 5\}$
 $\therefore A \times B$ contains $3 \times 3 = 9$ elements.
 Hence, number of relations from A to $B = 2^9$.
51. $n((A \times B) \cap (B \times A)) = n^2 = 99^2$.
52. $n(A) = 4, n(B) = 2$
 $\therefore n(A \times B) = 4 \times 2 = 8$
 Required numbers $= {}^8C_3 + {}^8C_4 + \dots + {}^8C_8$
 $= 2^8 - ({}^8C_0 + {}^8C_1 + {}^8C_2)$
 $= 256 - 37$
 $= 219$
53. In option (D), ordered pair $(a, d) \notin A \times B$. Thus it is not a relation from A to B .
54. A relation is equivalence if it is reflexive, symmetric and transitive.
56. Total number of reflexive relations in a set with n elements $= 2^n$.
 Therefore, total number of reflexive relation set with 4 elements $= 2^4$.
57. $x^2 - 4x^2 + 3x^2 = 0$
 $\therefore xRx \Rightarrow$ Reflexive
58. Given $A = \{2, 4, 6, 8\};$
 $R = \{(2, 4)(4, 2) (4, 6) (6, 4)\}$
 $(a, b) \in R \Rightarrow (b, a) \in R$ and also $R^{-1} = R$.
 Hence, R is symmetric.
59. For any $a \in \mathbb{N}$, we find $a|a$, therefore R is reflexive.
 But, R is not symmetric, because aRb does not imply that bRa .
60. Here, $(3, 3), (6, 6), (9, 9), (12, 12)$, [Reflexive];
 $(3, 6), (6, 12), (3, 12)$, [Transitive].
 Hence, reflexive and transitive only.
61. since $(5, 5) \notin S$.
 \therefore The relation S is not reflexive
 It is symmetric and transitive.



62.

Equivalence classes	Product
(1, 11)	1
(3, 13)	3
(5, 15)	5
(7, 17)	7
(9, 19)	9
(10, 20)	0
(12, 21) (2, 12) (2, 21)	2
(4, 14) (4, 22) (14, 22)	4
(16, 23) (6, 16) (6, 23)	6
(8, 18) (8, 24) (18, 24)	8

∴ There are 10 different equivalence classes.

63.

Given $A = \{1, 2, 3, 4\}$
 $R = \{(1, 3), (4, 2), (2, 4), (2, 3), (3, 1)\}$
 $(2, 3) \in R$ but $(3, 2) \notin R$.
Hence, R is not symmetric.
 R is not reflexive as $(1, 1) \notin R$.
 R is not a function as $(2, 4) \in R$ and $(2, 3) \in R$.
 R is not transitive as $(1, 3) \in R$ and $(3, 1) \in R$ but $(1, 1) \notin R$.

64.

For any $a \in R$, we have $a \geq a$. Therefore the relation R is reflexive, but it is not symmetric, as $(2, 1) \in R$ but $(1, 2) \notin R$. The relation R is transitive also, because $(a, b) \in R$, $(b, c) \in R$ imply that $a \geq b$ and $b \geq c$ which in turn imply that $a \geq c$.

65.

$|a - a| = 0 < 1$
∴ $aRa \forall a \in R$
∴ R is reflexive.
Again, $aRb \Rightarrow |a - b| \leq 1 \Rightarrow |b - a| \leq 1 \Rightarrow bRa$

∴ R is symmetric,

Again, $1R\frac{1}{2}$ and $\frac{1}{2}R1$ but $\frac{1}{2} \neq 1$

∴ R is not anti-symmetric.

Further, $1R2$ and $2R3$ but $1 \not R 3$,
 $[\because |1 - 3| = 2 > 1]$

∴ R is not transitive.

66.

$xpy, ypz \Rightarrow 2x + y = 41$ and $2y + z = 41$ which do not imply $2x + z = 41$

67.

for option D, $x > |x|$ is not true hence not reflexive

Take $x = 2, y = -1$, clearly $x > |y|$ but $y > |x|$ does not hold, hence not symmetric

Now, Let $x > |y|$ and $y > |z| \Rightarrow x, y > 0$.

∴ Rewriting, $x > |y|$ and $y > |z| \Rightarrow x > |z|$
Hence transitive.

68. $r = \{(a, b) \mid a, b \in R \text{ and } a - b + \sqrt{3} \text{ is an irrational no.}\}$

Here, r is reflexive as $aRa = a - a + \sqrt{3} = \sqrt{3}$ which is an irrational no.

$\sqrt{3}r1 = \sqrt{3} - 1 + \sqrt{3} = 2\sqrt{3} - 1$, which is an irrational number.

But $1r\sqrt{3} = 1 - \sqrt{3} + \sqrt{3} = 1$ which is not an irrational number.

$$\therefore \sqrt{3}r1 \Rightarrow \frac{1}{\sqrt{3}}$$

∴ r is not symmetric.

Also, r is not transitive.

Since, $\sqrt{3}r1$ and $1r2\sqrt{3} \not\Rightarrow \sqrt{3}r2\sqrt{3}$

∴ Option (B) is the correct answer.

69.

On the set R ;

$$xpy \Leftrightarrow x - y = 0 \text{ or } x - y \in Q'$$

∴ $x - x = 0 \Rightarrow xpx$ (Reflexive)

$$\text{if } x - y = 0 \Rightarrow y - x = 0 \text{ or } x - y \in Q'$$

$$\Rightarrow y - x \in Q' \text{ (Symmetric)}$$

$$\text{Take } x = 1 + \sqrt{2}; y = \sqrt{2} + \sqrt{3}; z = \sqrt{2} + 2$$

$$x - y = 1 - \sqrt{3} \in Q' \text{ and } y - z = \sqrt{3} - 2 \in Q'$$

Here xpy and ypz but x is not related to z .

∴ Not transitive

70.

Since, G. C. D. of a and a is 'a'

∴ if $a \neq 2$, then G. C. D. $\neq 2$

∴ R is not reflexive.

Let aRb

∴ G. C. D. of $a, b = 2$

i.e., $(a, b) = 2$

$$\Rightarrow (b, a) = 2 \Rightarrow \text{G. C. D. of } b, a = 2$$

∴ R is symmetric.

Again, let aRb and bRc

∴ G. C. D. of $a, b = 2$ and G. C. D. of $b, c = 2$ } $\not\Rightarrow$ G. C. D. of $(a, c) = 2$

∴ R is not transitive.

71.

Here, $R = \{(x, y) \in W \times W : \text{the words } x \text{ and } y \text{ have at least one letter in common}\}$

R is reflexive as the words x and x have all letters in common.

Hence, R is reflexive.

Also, if $(x, y) \in R$ i.e., x and y have a common letter, then y and x also have a letter in common

∴ R is symmetric.



- R is not transitive as $(x, y) \in R$ and $(y, z) \in R$ need not imply $(x, z) \in R$
 For example, let $x = \text{CANE}$, $y = \text{NEST}$ and $z = \text{WITH}$
 then $(x, y) \in R$ and $(y, z) \in R$, but $(x, z) \notin R$
- \therefore R is reflexive and symmetric but not transitive.
72. For reflexive, $\theta = \phi$, so $\sec^2\theta - \tan^2\theta = 1$,
 \therefore R is reflexive.
 For symmetric, $\sec^2\theta - \tan^2\phi = 1$
 so, $(1 + \tan^2\theta) - (\sec^2\phi - 1) \Rightarrow \sec^2\phi - \tan^2\theta = 1$
- \therefore R is symmetric
 For transitive, Let $\sec^2\theta - \tan^2\phi = 1$ (i)
 and $\sec^2\phi - \tan^2\gamma = 1$
 $\therefore 1 + \tan^2\phi - \tan^2\gamma = 1$
 $\Rightarrow \sec^2\theta - \tan^2\gamma = 1$ [From (i)]
 \therefore R is transitive.
73. $f(x) = \sqrt{x} \Rightarrow \frac{f(25)}{f(16)+f(1)} = \frac{\sqrt{25}}{\sqrt{16}+\sqrt{1}} = \frac{5}{5} = 1$
74. $f(x) = 2x, \quad x > 3$
 $= x^2, \quad 1 < x \leq 3$
 $= 3x, \quad x \leq 1$
- \therefore as, $x = -1$, $f(x) = f(-1) = 3(-1) = -3$
 as, $x = 2$, $f(x) = f(2) = (2)^2 = 4$
 as, $x = 4$, $f(x) = f(4) = 2(4) = 8$
- $\therefore f(-1) + f(2) + f(4) = 9$
75. Since, $f(x) f(y) = f(xy)$
 $\therefore f(1).f(2) = f(2)$
 $\therefore f(1).4 = 4$
 $\therefore f(1) = 1$ (i)
 Also, $f(2).f\left(\frac{1}{2}\right) = f(1)$
- $\therefore 4 \times f\left(\frac{1}{2}\right) = 1$ [From (i)]
- $\therefore f\left(\frac{1}{2}\right) = \frac{1}{4}$
76. Given, $f(x) = \cos(\log x) \Rightarrow f(y) = \cos(\log y)$
 Then, $f(x).f(y) - \frac{1}{2}\left[f\left(\frac{x}{y}\right) + f(xy)\right]$
 $= \cos(\log x) \cos(\log y) -$
 $\frac{1}{2}\left[\cos\left(\log \frac{x}{y}\right) + \cos(\log xy)\right]$
 $= \cos(\log x) \cos(\log y) -$
 $\frac{1}{2}[2 \cos(\log x) \cos(\log y)] = 0$

77. $f(x+1) - f(x) = 8x + 3$
 $\Rightarrow [b(x+1)^2 + c(x+1) + d] - (bx^2 + cx + d)$
 $= 8x + 3$
 $\Rightarrow (2b)x + (b+c) = 8x + 3$
 $\Rightarrow 2b = 8, b+c = 3$
 $\Rightarrow b = 4, c = -1$
78. $f(x+y) + f(x-y)$
 $= \frac{1}{2}[a^{x+y} + a^{-x-y} + a^{x-y} + a^{-x-y}]$
 $= \frac{1}{2}[a^x(a^y + a^{-y}) + a^{-x}(a^y + a^{-y})]$
 $= \frac{1}{2}(a^x + a^{-x})(a^y + a^{-y}) = 2f(x)f(y)$
79. $f(x) = \log\left[\frac{1+x}{1-x}\right]$
 $\therefore f\left(\frac{2x}{1+x^2}\right) = \log\left[\frac{1+\frac{2x}{1+x^2}}{1-\frac{2x}{1+x^2}}\right] = \log\left[\frac{x^2+1+2x}{x^2+1-2x}\right]$
 $= \log\left[\frac{1+x}{1-x}\right]^2 = 2 \log\left[\frac{1+x}{1-x}\right] = 2f(x)$
80. $e^{f(x)} = \frac{10+x}{10-x}, x \in (-10, 10)$
 $\Rightarrow f(x) = \log\left(\frac{10+x}{10-x}\right)$
 $\Rightarrow f\left(\frac{200x}{100+x^2}\right) = \log\left[\frac{10+\frac{200x}{100+x^2}}{10-\frac{200x}{100+x^2}}\right]$
 $= \log\left[\frac{10(10+x)}{10(10-x)}\right]^2$
 $= 2 \log\left(\frac{10+x}{10-x}\right) = 2f(x)$
- $\therefore f(x) = \frac{1}{2}f\left(\frac{200x}{100+x^2}\right) \Rightarrow k = \frac{1}{2} = 0.5$
81. $f(x) = \cos[\pi^2]x + \cos[-\pi^2]x$
 $f(x) = \cos(9x) + \cos(-10x)$
 $[\because \pi = 3.14,$
 $\therefore [9.85] = 9 \text{ and } [-9.85] = -10]$
 $= \cos(9x) + \cos(10x)$
 $= 2 \cos\left(\frac{19x}{2}\right) \cos\left(\frac{x}{2}\right)$



$$\therefore f\left(\frac{\pi}{2}\right) = 2 \cos\left(\frac{19\pi}{4}\right) \cos\left(\frac{\pi}{4}\right);$$

$$\therefore f\left(\frac{\pi}{2}\right) = 2 \times \frac{-1}{\sqrt{2}} \times \frac{1}{\sqrt{2}} = -1$$

83. $f(x) = f(-x) \Rightarrow f(0+x) = f(0-x)$ is symmetrical about $x = 0$.

$\therefore f(2+x) = f(2-x)$ is symmetrical about $x = 2$.

84. $f = \{(1, 1), (2, 4), (0, -2), (-1, -5)\}$ be a linear function from Z to Z . The function satisfies the above points, if $f(x) = 3x - 2$

85. Here, $f(x) = \log \frac{1+x}{1-x}$

$$\text{and } f(-x) = \log\left(\frac{1-x}{1+x}\right) = \log\left(\frac{1+x}{1-x}\right)^{-1}$$

$$= -\log\left(\frac{1+x}{1-x}\right) = -f(x) = f(-x)$$

$\Rightarrow f(x)$ is an odd function.

86. Since, $f(x)$ is even.

$$\therefore f(-x) = f(x)$$

$$\therefore \frac{a^{-x} - 1}{(-x)^n (a^{-x} + 1)} = \frac{a^x - 1}{x^n (a^x + 1)}$$

$$\Rightarrow \frac{1 - a^x}{(-1)^n x^n (1 + a^x)} = \frac{a^x - 1}{x^n (a^x + 1)}$$

$$\Rightarrow \frac{-1}{(-1)^n} = 1 \Rightarrow -1 = (-1)^n$$

$\therefore n = -\frac{1}{3}$ can satisfy the equation.

87. $f(-x) = \sec\left[\log\left(-x + \sqrt{1 + (-x)^2}\right)\right]$

$$= \sec\left[\log\left(-x + \sqrt{1 + x^2}\right)\right]$$

$$= \sec\left[\log\left(\sqrt{1 + x^2} - x\right)\right]$$

$$= \sec\left[\log\left(\frac{1 + x^2 - x^2}{\sqrt{1 + x^2} + x}\right)\right]$$

$$= \sec\left[\log\left(\frac{1}{\sqrt{1 + x^2} + x}\right)\right]$$

$$= \sec\left[-\log\left(\sqrt{1 + x^2} + x\right)\right]$$

$$= \sec\left[\log\left(\sqrt{1 + x^2} + x\right)\right]$$

$\therefore f(x)$ is an even function.

88. $f(x) = \sin\left(\log(x + \sqrt{1 + x^2})\right)$

$$\Rightarrow f(-x) = \sin[\log(-x + \sqrt{1 + x^2})]$$

$$\Rightarrow f(-x) = \sin \log\left(\frac{(\sqrt{1 + x^2} - x)(\sqrt{1 + x^2} + x)}{(\sqrt{1 + x^2} + x)}\right)$$

$$\Rightarrow f(-x) = \sin \log\left[\frac{1}{(x + \sqrt{1 + x^2})}\right]$$

$$\Rightarrow f(-x) = \sin\left[-\log(x + \sqrt{1 + x^2})\right]$$

$$\Rightarrow f(-x) = -\sin\left[\log(x + \sqrt{1 + x^2})\right]$$

$$\Rightarrow f(-x) = -f(x)$$

$\therefore f(x)$ is odd function.

89. $f(x) + 2f\left(\frac{1}{x}\right) = 3x \quad \dots (i)$

$$\therefore f\left(\frac{1}{x}\right) + 2f(x) = \frac{3}{x} \quad \dots (ii)$$

From (i) and (ii), we get

$$3f(x) = \frac{6}{x} - 3x$$

$$\Rightarrow f(x) = \frac{2}{x} - x \Rightarrow f(-x) = -\frac{2}{x} + x$$

Since, $f(x) = f(-x)$

$$\therefore \frac{2}{x} - x = -\frac{2}{x} + x$$

$$\Rightarrow \frac{4}{x} = 2x \Rightarrow x^2 = 2 \Rightarrow x = \pm \sqrt{2}$$

\therefore option (B) is the correct answer.

90. Given expression = $\sum_{i=0}^{98} \left[\frac{2}{3} + \frac{i}{99}\right]$

$$= \sum_{i=0}^{32} \left[\frac{2}{3} + \frac{i}{99}\right] + \sum_{i=33}^{98} \left[\frac{2}{3} + \frac{i}{99}\right]$$

$$= 0 + \sum_{i=33}^{98} \left[\frac{2}{3} + \frac{i}{99}\right]$$

$$\left[\because \frac{2}{3} \leq \frac{2}{3} + \frac{i}{99} < 1\right]$$

for $i = 0, 1, 2, \dots, 32]$

$$= 66$$

$[\because$ each term in the summation is one or more but less than 2 when $i = 33, 34, 35, \dots, 98]$



$$\begin{aligned}
 91. \quad f(x) &= \frac{1}{2}(1 + \cos 2x) + \frac{1}{2} \left[1 + \cos \left(\frac{2\pi}{3} + 2x \right) \right] \\
 &\quad - \frac{2 \cos x \cos \left(\frac{\pi}{3} + x \right)}{2} \\
 &= 1 + \frac{1}{2} \left[\cos 2x + \cos \left(\frac{2\pi}{3} + 2x \right) \right. \\
 &\quad \left. - \cos \left(2x + \frac{\pi}{3} \right) - \cos \left(\frac{\pi}{3} \right) \right] \\
 &= \frac{3}{4} + \frac{1}{2} \left[\cos 2x + \cos \left(2x + \frac{2\pi}{3} \right) \right. \\
 &\quad \left. - \cos \left(2x + \frac{\pi}{3} \right) \right] \\
 &= \frac{3}{4} + \frac{1}{2} \left[\cos 2x - 2 \sin \left(2x + \frac{\pi}{2} \right) \sin \left(\frac{\pi}{6} \right) \right] \\
 &= \frac{3}{4} + \frac{1}{2} \left[\cos 2x - 2 \sin \left(\frac{\pi}{2} + 2x \right) \cdot \frac{1}{2} \right] \\
 &= \frac{3}{4} + \frac{1}{2} [\cos 2x - \cos 2x] = \frac{3}{4}
 \end{aligned}$$

$$92. \quad y = \frac{x+2}{x-1} \Rightarrow x = \frac{3}{y-1} + 1 = \frac{y+2}{y-1} = f(y).$$

93. Function given by $f(x) = ax + b$

$$f^{-1}(x) = \frac{x-b}{a}$$

$$\text{So, } g(y) = y - 3$$

$$94. \quad f(x) = |x|$$

$$f(x) = \begin{cases} x, & \text{if } x \geq 0 \\ -x, & \text{if } x < 0 \end{cases}$$

Therefore, the function $f^{-1}(x)$ does not exist.

$$95. \quad \text{Let } f(x) = y \Rightarrow x = f^{-1}(y)$$

$$\text{Now, } y = 3x - 5$$

$$\Rightarrow x = \frac{y+5}{3}$$

$$\Rightarrow f^{-1}(y) = x = \frac{y+5}{3}$$

$$\therefore f^{-1}(x) = \frac{x+5}{3}$$

Also f is one-one and onto, so f^{-1} exists and is given by $f^{-1}(x) = \frac{x+5}{3}$.

$$96. \quad \text{Let } f(x) = y \Rightarrow x = f^{-1}(y)$$

$$\text{Now, } y = 2x + 6$$

$$\Rightarrow 2x = y - 6$$

$$\Rightarrow x = \frac{y}{2} - 3$$

$$\Rightarrow f^{-1}(y) = \frac{y}{2} - 3$$

$$\Rightarrow f^{-1}(x) = \frac{x}{2} - 3$$

$$97. \quad \text{Let } f(x) = y \Rightarrow x = f^{-1}(y)$$

$$\text{Now, } y = x^3 + 5$$

$$\Rightarrow y - 5 = x^3$$

$$\Rightarrow x = (y-5)^{\frac{1}{3}}$$

$$\Rightarrow f^{-1}(y) = (y-5)^{\frac{1}{3}}$$

$$\Rightarrow f^{-1}(x) = (x-5)^{\frac{1}{3}}$$

$$98. \quad \text{Let } f(x) = y \Rightarrow x = f^{-1}(y). \text{ Now,}$$

$$y = \frac{2x-1}{x+5}, (x \neq -5)$$

$$xy + 5y = 2x - 1 \Rightarrow 5y + 1 = 2x - xy$$

$$\Rightarrow x(2-y) = 5y + 1 \Rightarrow x = \frac{5y+1}{2-y}$$

$$\Rightarrow f^{-1}(y) = \frac{5y+1}{2-y}$$

$$\therefore f^{-1}(x) = \frac{5x+1}{2-x}, x \neq 2$$

$$99. \quad \text{We have, } f(x) = \frac{5x}{4x+5}, x \in \mathbb{R} - \left\{ \frac{5}{4} \right\}$$

$$\text{Let } f(x) = y$$

$$\Rightarrow x = f^{-1}(y)$$

$$y = \frac{5x}{4x+5}$$

$$\Rightarrow 4xy + 5y = 5x$$

$$\Rightarrow 5y = 5x - 4xy = x(5 - 4y)$$

$$\Rightarrow x = \frac{5y}{5-4y}$$

$$g(y) = f^{-1}(y) = \frac{5y}{5-4y}, x \in \mathbb{R} - \left\{ \frac{5}{4} \right\}$$



100. Given, $f(x) = 2^{x(x-1)}$
 $\Rightarrow x(x-1) = \log_2 f(x)$
 $\Rightarrow x^2 - x - \log_2 f(x) = 0$
 $\Rightarrow x = \frac{1 \pm \sqrt{1 + 4 \log_2 f(x)}}{2}$
 Only $x = \frac{1 + \sqrt{1 + 4 \log_2 f(x)}}{2}$ lies in the domain

$$\therefore f^{-1}(x) = \frac{1}{2} [1 + \sqrt{1 + 4 \log_2 x}]$$

101. Let $y = f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}} + 2$
 $\therefore y - 2 = \frac{e^{2x} - 1}{e^{2x} + 1}$
 $\Rightarrow (y - 2)e^{2x} + y - 2 = e^{2x} - 1$
 $\Rightarrow e^{2x} = \frac{1 - y}{y - 3} = \frac{y - 1}{3 - y}$

$$\Rightarrow 2x = \log_e \left(\frac{y - 1}{3 - y} \right)$$

$$\Rightarrow x = \frac{1}{2} \log_e \left(\frac{y - 1}{3 - y} \right)$$

$$\Rightarrow f^{-1}(y) = \frac{1}{2} \log_e \left(\frac{y - 1}{3 - y} \right)$$

$$\Rightarrow f^{-1}(x) = \log_e \left(\frac{x - 1}{3 - x} \right)^{\frac{1}{2}}$$

102. Let $y = f(x) = \frac{10^x - 10^{-x}}{10^x + 10^{-x}}$

$$\therefore y = \frac{10^{2x} - 1}{10^{2x} + 1}$$

$$\Rightarrow 10^{2x} = \frac{1 + y}{1 - y}$$

$$\Rightarrow 2x = \log_{10} \frac{1 + y}{1 - y}$$

$$\Rightarrow x = \frac{1}{2} \log_{10} \frac{1 + y}{1 - y}$$

$$\Rightarrow f^{-1}(y) = \frac{1}{2} \log_{10} \frac{1 + y}{1 - y}$$

$$\Rightarrow f^{-1}(x) = \frac{1}{2} \log_{10} \frac{1 + x}{1 - x}$$

103. Let $y = f(x) = \frac{16^x - 16^{-x}}{16^x + 16^{-x}}$

$$\therefore y = \frac{16^{2x} - 1}{16^{2x} + 1}$$

$$\Rightarrow 16^{2x} = \frac{1 + y}{1 - y}$$

$$\Rightarrow 2x = \log_{16} \frac{1 + y}{1 - y}$$

$$\Rightarrow x = \frac{1}{2} \log_{16} \frac{1 + y}{1 - y} \Rightarrow f^{-1}(y) = \frac{1}{2} \log_{16} \frac{1 + y}{1 - y}$$

$$\Rightarrow f^{-1}(x) = \frac{1}{2} \log_{16} \frac{1 + x}{1 - x}$$

104. $f(g(-1)) = f(-3 - 4) = f(-7) = 5 - 49 = -44$

105. $f(2) = \frac{2}{2^2 + 1} = \frac{2}{5}$

$$\therefore f(f(2)) = f\left(\frac{2}{5}\right) = \frac{\frac{2}{5}}{\left(\frac{2}{5}\right)^2 + 1} = \frac{10}{29}$$

106. Here, $f(2) = \frac{2+1}{2-1} = 3$

$$\therefore f(f(2)) = f(3) = \frac{3+1}{3-1} = \frac{4}{2} = 2$$

$$\therefore f(f(f(2))) = f(2) = \frac{2+1}{2-1} = 3$$

107. $(f \circ g)(x) = f(g(x)) = f(x^2) = \sin x^2$

108. $f(x) = \sin x + \cos x$, $g(x) = x^2$

$$\therefore f \circ g(x) = \sin x^2 + \cos x^2$$

109. $f[f(\cos 2\theta)] = f\left[\frac{1 - \cos 2\theta}{1 + \cos 2\theta}\right] = f(\tan^2 \theta)$
 $= \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} = \cos 2\theta$

110. Here, $f\left(\frac{1}{2}\right) = \left(25 - \frac{1}{16}\right)^{\frac{1}{4}} = \left(\frac{399}{16}\right)^{\frac{1}{4}}$

$$\Rightarrow f\left[f\left(\frac{1}{2}\right)\right] = f\left[\left(\frac{399}{16}\right)^{\frac{1}{4}}\right]$$

$$= \left(25 - \frac{399}{16}\right)^{\frac{1}{4}} = \left(\frac{1}{16}\right)^{\frac{1}{4}} = \frac{1}{2}$$



$$\begin{aligned}
 111. \quad f(x) &= \frac{x-1}{x+1} \\
 \Rightarrow \frac{f(x)+1}{f(x)-1} &= \frac{2x}{-2} \\
 \Rightarrow x &= \frac{f(x)+1}{1-f(x)} \\
 \therefore f(2x) &= \frac{2x-1}{2x+1} = \frac{2\left[\frac{f(x)+1}{1-f(x)}\right]-1}{2\left[\frac{f(x)+1}{1-f(x)}\right]+1} = \frac{3f(x)+1}{f(x)+3}
 \end{aligned}$$

$$\begin{aligned}
 112. \quad f(x) &= \frac{\alpha x}{x+1}; \\
 f(f(x)) &= f\left(\frac{\alpha x}{x+1}\right) = \frac{\alpha\left(\frac{\alpha x}{x+1}\right)}{\frac{\alpha x}{x+1}+1} \\
 \text{But } f(f(x)) &= x \\
 \therefore \frac{\alpha^2 x}{\alpha x + x + 1} &= x \\
 \text{L.H.S, Put } \alpha &= -1, \\
 \therefore \frac{(-1)^2 x}{(-1)x + x + 1} &= \frac{x}{-x + x + 1} = x; \\
 \therefore \alpha &= -1
 \end{aligned}$$

$$\begin{aligned}
 113. \quad \text{Given, } f(x) &= ax + b, g(x) = cx + d \\
 \text{and } f(g(x)) &= g(f(x)) \\
 \Rightarrow f(cx + d) &= g(ax + b) \\
 \Rightarrow a(cx + d) + b &= c(ax + b) + d \\
 \Rightarrow ad + b &= cb + d \\
 \Rightarrow f(d) &= g(b)
 \end{aligned}$$

$$\begin{aligned}
 114. \quad fog(x) &= f[g(x)] \\
 &= f\left(x^{\frac{1}{3}}\right) = 8\left(x^{\frac{1}{3}}\right)^3 \\
 &= 8x
 \end{aligned}$$

$$\begin{aligned}
 115. \quad (fog)(x) &= f(g(x)) = f\left(\frac{x-1}{2}\right) \\
 &= 2\left(\frac{x-1}{2}\right) + 1 = x \\
 \Rightarrow (fog)(x) &= x \Rightarrow x = (fog)^{-1}(x) \\
 \text{Hence, } (fog)^{-1}\left(\frac{1}{x}\right) &= \frac{1}{x}
 \end{aligned}$$

$$\begin{aligned}
 116. \quad f(x) &= \sin^2 x + \sin^2\left(x + \frac{\pi}{3}\right) + \cos x \cos\left(x + \frac{\pi}{3}\right) \\
 &= \sin^2 x + \left[\sin\left(x + \frac{\pi}{3}\right)\right]^2 \\
 &\quad + \cos x \left[\cos x \cos \frac{\pi}{3} - \sin x \sin \frac{\pi}{3}\right] \\
 &= \sin^2 x + \left[\sin x \cos \frac{\pi}{3} + \cos x \sin \frac{\pi}{3}\right]^2 \\
 &\quad + \cos x \left[\frac{1}{2} \cos x - \frac{\sqrt{3}}{2} \sin x\right] \\
 &= \sin^2 x + \left[\frac{\sin x}{2} + \frac{\sqrt{3}}{2} \cos x\right]^2 \\
 &\quad + \frac{\cos^2 x}{2} - \frac{\sqrt{3}}{2} \sin x \cos x \\
 &= \sin^2 x + \frac{\sin^2 x}{4} + \frac{3}{4} \cos^2 x + \frac{\cos^2 x}{2} \\
 &\quad + \frac{\sqrt{3}}{2} \sin x \cos x - \frac{\sqrt{3}}{2} \sin x \cos x \\
 &= \frac{5}{4} (\sin^2 x + \cos^2 x) = \frac{5}{4} \\
 (\text{gof})(x) &= g[f(x)] = g\left(\frac{5}{4}\right) = 1
 \end{aligned}$$

$$\begin{aligned}
 117. \quad \text{As } x - [x] &\in [0, 1), \forall x \in \mathbb{R} \\
 \therefore 0 \leq x - [x] &< 1, \forall x \in \mathbb{R} \\
 \Rightarrow 1 \leq 1 + x - [x] &< 2, \forall x \in \mathbb{R} \\
 \Rightarrow 1 \leq g(x) &< 2, \forall x \in \mathbb{R} \\
 \text{Hence, } f(g(x)) &= 1 \forall x \in \mathbb{R}
 \end{aligned}$$

$$\begin{aligned}
 118. \quad (\text{gof})(e) + (\text{fog})(\pi) &= g(f(e)) + f(g(\pi)) \\
 &= g(1) + f(0) \\
 &= -1 + 0 \\
 &= -1
 \end{aligned}$$

$$\begin{aligned}
 119. \quad g(f(x)) &= g(|x|) = [|x|] \\
 \text{and } f(g(x)) &= f([x]) = \lfloor [x] \rfloor \\
 \text{When } x \geq 0, [x] &= [x] = \lfloor [x] \rfloor \\
 \Rightarrow f(g(x)) &= g(f(x)) \\
 \text{When } x < 0, [x] &\leq x < 0 \\
 \Rightarrow \lfloor [x] \rfloor &\geq |x| \\
 \Rightarrow \lfloor [x] \rfloor &\geq |x| \geq \lfloor [x] \rfloor \quad \dots [\because [t] \leq t \text{ for all } t] \\
 \Rightarrow f(g(x)) &\geq g(f(x)) \\
 \therefore g(f(x)) &\leq f(g(x)) \text{ for all } x \in \mathbb{R}
 \end{aligned}$$



$$\begin{aligned}
 120. \quad (\text{hofog})(x) &= (\text{hof})(g(x)) \\
 &= (\text{hof})\left(\sqrt{x^2+1}\right) \\
 &= h\left(f\left(\sqrt{x^2+1}\right)\right) \\
 &= h\left[\left(\sqrt{x^2+1}\right)^2-1\right] \\
 &= h(x^2+1-1) \\
 &= h(x^2) = \begin{cases} 0, & \text{if } x=0 \\ x^2, & \text{if } x \neq 0 \end{cases}
 \end{aligned}$$

121. Given,

$$\begin{aligned}
 f(x) &= \sin^2 x + \sin^2\left(x + \frac{\pi}{3}\right) + \cos\left(x + \frac{\pi}{3}\right) \cos x \\
 &= \frac{1}{2} \left\{ 1 - \cos 2x + 1 - \cos\left(2x + \frac{2\pi}{3}\right) \right. \\
 &\quad \left. + \cos\left(2x + \frac{2\pi}{3}\right) + \cos \frac{\pi}{3} \right\}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2} \left[\frac{5}{2} - \left\{ \cos 2x + \cos\left(2x + \frac{2\pi}{3}\right) \right\} + \cos\left(2x + \frac{\pi}{3}\right) \right] \\
 &= \frac{1}{2} \left[\frac{5}{2} - 2 \cos\left(2x + \frac{\pi}{3}\right) \cos \frac{\pi}{3} + \cos\left(2x + \frac{\pi}{3}\right) \right] \\
 &= \frac{5}{4}
 \end{aligned}$$

$$\begin{aligned}
 \therefore \text{gof}(x) &= g[f(x)] = g\left(\frac{5}{4}\right) \\
 &= 1 \quad \dots \left[\because g\left(\frac{5}{4}\right) = 1 \right]
 \end{aligned}$$

Hence, $\text{gof}(x)$ is a constant function.

$$\begin{aligned}
 122. \quad (\text{gof})(x) &= \sin x^2 \Rightarrow (\text{gogof})(x) = \sin(\sin x^2) \\
 &\Rightarrow (\text{fogogof})(x) = (\sin(\sin x^2))^2 = \sin^2(\sin x^2) \\
 \text{Now, } \sin^2(\sin x^2) &= \sin(\sin x^2) \\
 &\Rightarrow \sin(\sin x^2) = 0, 1 \\
 &\Rightarrow \sin x^2 = n\pi, (4n+1)\frac{\pi}{2} \quad n \in I \\
 &\Rightarrow \sin x^2 = 0 \Rightarrow x^2 = n\pi \\
 &\Rightarrow x = \pm\sqrt{n\pi} \quad n \in W
 \end{aligned}$$

$$\begin{aligned}
 123. \quad |x| &= -x, & \text{if } x < 0 \\
 &= x, & \text{if } x \geq 0
 \end{aligned}$$

Now, $(\text{fog})(x) = f[g(x)]$

$$\begin{aligned}
 &= |g(x)| + g(x) \\
 &= ||x| - x| + |x| - x
 \end{aligned}$$

When, $x < 0$

$$\begin{aligned}
 (\text{fog})(x) &= |-x - x| + (-x) - x \\
 &= -2x - 2x = -4x
 \end{aligned}$$

124. Let $f(x)$ be periodic with period T .
Then, $f(x+T) = f(x)$ for all $x \in \mathbb{R}$
 $\Rightarrow x+T - [x+T] = x - [x]$, for all $x \in \mathbb{R}$
 $\Rightarrow x+T - x = [x+T] - [x]$
 $\Rightarrow [x+T] - [x] = T$ for all $x \in \mathbb{R}$
 $\Rightarrow T = 1, 2, 3, 4, \dots$
 The smallest value of T satisfying $f(x+T) = f(x)$ for all $x \in \mathbb{R}$ is 1.
 Hence, $f(x) = x - [x]$ has period 1.

125. $g(x)$ is neither injective nor surjective
 $(\text{gof})(x) = (e^x)^2 = e^{2x}$
 This is an injective function.

126. At $x=0$, $f(x)$ is not defined.

128. $f(x) = f(y)$
 $\Rightarrow x+2 = y+2 \Rightarrow x = y$
 \therefore Function f is one-one

129. Total number of distinct functions from $A \rightarrow A = n^n = 6^6$
 Number of bijections = $n! = 6!$
 \therefore Number of functions which are not bijections = $6^6 - 6!$

130. Number of bijective function from a set of 10 elements to itself is ${}^{10}P_{10}$.
 So, required number = $10!$

131. The total number of injective functions from a set A containing 3 elements to a set B containing 4 elements is equal to the total number of arrangements of 4 by taking 3 at a time i.e., ${}^4P_3 = 24$.

132. Number of injective mapping = ${}^5P_4 = 120$

133. $|x|$ is not one-one; x^2 is not one-one; $x^2 + 1$ is not one-one. But $2x - 5$ is one-one because $f(x) = f(y) \Rightarrow 2x - 5 = 2y - 5 \Rightarrow x = y$
 Now, $f(x) = 2x - 5$ is onto.
 $\therefore f(x) = 2x - 5$ is bijective.

134. Let $x_1, x_2 \in \mathbb{R}$ such that $f(x_1) = f(x_2)$
 $\Rightarrow x_1 = \pm x_2$
 $\therefore f(x_1) = f(x_2)$ does not imply that $x_1 = x_2$
 $\therefore f$ is not one-one.

Consider an element 2 in the co-domain \mathbb{R} .
 There does not exist any x in domain \mathbb{R} such that $f(x) = 2$.


$\therefore f$ is not onto.

135. $f'(x) = 2 + \cos x > 0$. So, $f(x)$ is strictly monotonic increasing so, $f(x)$ is one-to-one and onto.



136. Let $x, y \in \mathbb{N}$ such that $f(x) = f(y)$
 Then, $f(x) = f(y) \Rightarrow x^2 + x + 1 = y^2 + y + 1$
 $\Rightarrow (x - y)(x + y + 1) = 0$
 $\Rightarrow x = y$ or $x = (-y - 1) \notin \mathbb{N}$
 $\therefore f$ is one-one.
 Again, since for each $y \in \mathbb{N}$, there exist $x \in \mathbb{N}$
 $\therefore f$ is onto.
137. We have $f(x) = (x - 1)(x - 2)(x - 3)$ and
 $f(1) = f(2) = f(3) = 0 \Rightarrow f(x)$ is not one-one.
 For each $y \in \mathbb{R}$, there exists $x \in \mathbb{R}$ such that
 $f(x) = y$. Therefore f is onto. Hence, $f : \mathbb{R} \rightarrow \mathbb{R}$
 is onto but not one-one.

138. $f : \mathbb{N} \rightarrow \mathbb{I}$
 $f(1) = 0, f(2) = -1, f(3) = 1, f(4) = -2, f(5) = 2$
 and $f(6) = -3$ so on.



In this type of function every element of set A has unique image in set B and there is no element left in set B. Hence f is one-one and onto function.

139. $f : \mathbb{N} \rightarrow \mathbb{N}$

$$f(n) = \begin{cases} \frac{n+1}{2} & \text{if } n \text{ is odd} \\ \frac{n}{2} & \text{if } n \text{ is even} \end{cases}$$
 Now for $n = 1, f(1) = \frac{1+1}{2} = 1$
 and if $n = 2, f(2) = \frac{2}{2} = 1$
 $\therefore f(1) = f(2), \text{ But } 1 \neq 2.$
 $\therefore f(x)$ is not one-one.
 $f(x) = \frac{n+1}{2}$ if n is odd
 if $y = \frac{n+1}{2}$ then $n = 2y - 1, \forall y$
 Also, $f(x) = \frac{n}{2}$ if n is even i.e., $y = \frac{n}{2}$
 or $n = 2y \forall y$
 $\therefore f(x)$ is onto.
140. $\sigma : \mathbb{N} \rightarrow \mathbb{Z}$
 $\sigma(1) = 0, \sigma(2) = 1, \sigma(3) = -1, \sigma(4) = 2,$
 $\sigma(5) = -2, \sigma(6) = 3, \sigma(7) = -3$
 $\therefore \sigma$ is one-one and onto.

141. Function $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = e^x$. Let
 $x_1, x_2 \in \mathbb{R}$ and $f(x_1) = f(x_2)$ or $e^{x_1} = e^{x_2}$ or
 $x_1 = x_2$. Therefore f is one-one.
 Let $f(x) = e^x = y$. Taking log on both sides, we
 get $x = \log y$. We know that negative real
 numbers have no pre-image or the function is
 not onto and zero is not the image of any real
 number. Therefore function f is into.

142. Here, $(f - g)(x) = f(x) - g(x)$
 $\therefore (f - g)(x) = \begin{cases} x - 0 = x, & \text{if } x \text{ is rational} \\ 0 - x = -x, & \text{if } x \text{ is irrational} \end{cases}$

Let $k = f - g$
 Let x, y be any two distinct real numbers.

Then, $x \neq y$
 $\Rightarrow -x \neq -y$
 Now, $x \neq y$
 $\Rightarrow k(x) \neq k(y) \Rightarrow (f - g)(x) \neq (f - g)(y)$
 $\Rightarrow f - g$ is one-one.

Let y be any real number
 If y is a rational number, then
 $k(y) = y$

$\Rightarrow (f - g)(y) = y$
 If y is an irrational number, then
 $k(-y) = y$
 $\Rightarrow (f - g)(-y) = y$

Thus, every $y \in \mathbb{R}$ (co-domain) has its pre-
 image in \mathbb{R} (domain)

- $\therefore f - g : \mathbb{R} \rightarrow \mathbb{R}$ is onto.
 Hence, $f - g$ is one-one and onto.

143. Let $x, y \in \mathbb{R}$ be such that
 $f(x) = f(y)$
 $\Rightarrow x^3 + 5x + 1 = y^3 + 5y + 1$
 $\Rightarrow (x^3 - y^3) + 5(x - y) = 0$
 $\Rightarrow (x - y)(x^2 + xy + y^2 + 5) = 0$
 $\Rightarrow (x - y) \left[\left(x + \frac{y}{2}\right)^2 + \frac{3y^2}{4} + 5 \right] = 0$
 $\Rightarrow x = y$ and $\left(x + \frac{y}{2}\right)^2 + \frac{3y^2}{4} + 5 \neq 0$

- $\therefore f : \mathbb{R} \rightarrow \mathbb{R}$ is one-one
 Let y be an arbitrary element in
 \mathbb{R} (co-domain).
 Then, $f(x) = y$ i.e., $x^3 + 5x + 1 = y$ has at least
 one real root, say β in \mathbb{R}
 $\therefore \beta^3 + 5\beta + 1 = y$
 $\Rightarrow f(\beta) = y$
 Thus, for each $y \in \mathbb{R}$ there exists $\beta \in \mathbb{R}$ such
 that $f(\beta) = y$
 $\therefore f : \mathbb{R} \rightarrow \mathbb{R}$ is onto
 Hence, $f : \mathbb{R} \rightarrow \mathbb{R}$ is one-one onto.



144. Since, $f(x)$ and $g(x)$ has same domain and co-domain A and B and $f(1) = (1)^2 - 1 = 0$

$$g(1) = 2 \left| 1 - \frac{1}{2} \right| - 1 = 2 \times \frac{1}{2} - 1 = 0$$

$$f(1) = 0 = g(1), f(0) = 0 = g(0)$$

$$f(-1) = 2 = g(-1), f(2) = 2 = g(2)$$

$$A = \{-1, 0, 1, 2\}, B = \{-4, -2, 0, 2\}$$

\therefore By definition, the two function are equal $f = g$

145. $-\sqrt{1+(-\sqrt{3})^2} \leq (\sin x - \sqrt{3} \cos x) \leq \sqrt{1+(-\sqrt{3})^2}$

$$\therefore -2 \leq (\sin x - \sqrt{3} \cos x) \leq 2$$

$$\therefore -2 + 1 \leq (\sin x - \sqrt{3} \cos x + 1) \leq 2 + 1$$

$$\therefore -1 \leq (\sin x - \sqrt{3} \cos x + 1) \leq 3$$

$$\text{i.e., range} = [-1, 3]$$

\therefore For f to be onto $S = [-1, 3]$.

146. Given, $f(x) = \sin x$

$\therefore f: \mathbb{R} \rightarrow \mathbb{R}$ is neither one-one nor onto as

$$R_f = [-1, 1].$$

$$f: \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \rightarrow [-1, 1]$$

is both one-one and onto.

$$f: [0, \pi] \rightarrow [-1, 1]$$

is neither one-one nor onto as

$$R_f = [0, 1].$$

$$f: \left[0, \frac{\pi}{2}\right] \rightarrow [-1, 1] \text{ is one-one but not onto as}$$

$$R_f = [0, 1].$$

147. Let $f(x) = \frac{x^2 - 1}{x^2 + 1} = \frac{x^2 + 1 - 2}{x^2 + 1} = 1 - \frac{2}{x^2 + 1}$

$$\therefore x^2 + 1 > 1;$$

$$\therefore \frac{2}{x^2 + 1} \leq 2$$

$$\text{So } 1 - \frac{2}{x^2 + 1} \geq 1 - 2;$$

$$\therefore -1 \leq f(x) < 1$$

Thus, $f(x)$ has the minimum value equal to -1 .

148. The quantity under root is positive, when

$$-1 - \sqrt{3} \leq x \leq -1 + \sqrt{3}.$$

149. The function $f(x) = \sqrt{\log(x^2 - 6x + 6)}$ is defined, when $\log(x^2 - 6x + 6) \geq 0$

$$\Rightarrow x^2 - 6x + 6 \geq 1 \Rightarrow (x - 5)(x - 1) \geq 0$$

This inequality holds, if $x \leq 1$ or $x \geq 5$.

Hence, the domain of the function will be $(-\infty, 1] \cup [5, \infty)$.

150. Here, $x + 3 > 0$ and $x^2 + 3x + 2 \neq 0$

$$\therefore x > -3 \text{ and } (x + 1)(x + 2) \neq 0, \text{ i.e., } x \neq -1, -2.$$

$$\therefore \text{Domain} = (-3, \infty) - \{-1, -2\}.$$

151. $D_f = D_g \cap D_h$

$$\text{where } g(x) = \frac{1}{\log_{10}(1-x)} \text{ and } h(x) = \sqrt{2+x}$$

$$\text{Now, } D_g = \{x \in \mathbb{R} : 1 - x > 0, \log_{10}(1-x) \neq 0\}$$

$$= \{x \in \mathbb{R} : x < 1, 1 - x \neq 1\}$$

$$= \{x \in \mathbb{R} : x < 1, x \neq 0\}$$

$$\text{and } D_h = \{x \in \mathbb{R} : x + 2 \geq 0\}$$

$$= \{x \in \mathbb{R} : x \geq -2\}$$

$$\therefore D_f = [(-\infty, 1) - \{0\}] \cap [-2, \infty)$$

$$= [-2, 1) - \{0\}$$

152. $f(x) = \log|\log x|$, $f(x)$ is defined if $|\log x| > 0$ and $x > 0$ i.e., if $x > 0$ and $x \neq 1$

$$(\because |\log x| > 0 \text{ if } x \neq 1)$$

$$\Rightarrow x \in (0, 1) \cup (1, \infty).$$

153. $\frac{1-|x|}{2-|x|} \geq 0$

$$\Rightarrow \frac{|x|-1}{|x|-2} \geq 0$$

$$\Rightarrow |x| \leq 1 \text{ as } |x| > 2$$

$$\Rightarrow x \in (-\infty, -2) \cup (2, \infty) \cup [-1, 1]$$

154. $f(x) = \sqrt{\log \frac{1}{|\sin x|}}$

$$\Rightarrow \sin x \neq 0 \Rightarrow x \neq n\pi + (-1)^n 0$$

$$\Rightarrow x \neq n\pi. \text{ Domain of } f(x) = \mathbb{R} - \{n\pi, n \in \mathbb{I}\}.$$

155. $f(x)$ is to be defined when $x^2 - 1 > 0$

$$\Rightarrow x^2 > 1, \Rightarrow x < -1 \text{ or } x > 1 \text{ and } 3 + x > 0$$


$$\therefore x > -3 \text{ and } x \neq -2$$

$$\therefore D_f = (-3, -2) \cup (-2, -1) \cup (1, \infty).$$

156. $f(x) = e^{\sqrt{5x-3-2x^2}}$

$$\Rightarrow 5x - 3 - 2x^2 \geq 0$$

$$\Rightarrow (x-1) \left(x - \frac{3}{2}\right) \leq 0$$



$$\therefore D_f = \left[1, \frac{3}{2}\right]$$



157. To define $f(x)$, $9 - x^2 > 0 \Rightarrow |x| < 3$
 $\Rightarrow -3 < x < 3$,(i)
 and $-1 \leq (x - 3) \leq 1$
 $\Rightarrow 2 \leq x \leq 4$ (ii)
 From (i) and (ii), $2 \leq x < 3$ i.e., $[2, 3)$.

158. $-1 \leq 1 + 3x + 2x^2 \leq 1$
Case I : $2x^2 + 3x + 1 \geq -1$; $2x^2 + 3x + 2 \geq 0$
 $x = \frac{-3 \pm \sqrt{9-16}}{6} = \frac{-3 \pm i\sqrt{7}}{6}$ (imaginary).

Case II : $2x^2 + 3x + 1 \leq 1$
 $\Rightarrow 2x^2 + 3x \leq 0 \Rightarrow 2x\left(x + \frac{3}{2}\right) \leq 0$
 $\Rightarrow \frac{-3}{2} \leq x \leq 0 \Rightarrow x \in \left[-\frac{3}{2}, 0\right]$

In case I, we get imaginary value hence, rejected

\therefore Domain of function = $\left[-\frac{3}{2}, 0\right]$.

159. $f(x) = \sqrt{\cos^{-1}\left(\frac{1-|x|}{2}\right)}$

$\therefore -1 \leq \frac{1-|x|}{2} \leq 1$
 $\Rightarrow -2 - 1 \leq -|x| \leq 2 - 1$
 $\Rightarrow -3 \leq |x| \leq 1$
 $\Rightarrow -1 \leq |x| \leq 3$
 $\Rightarrow x \in [-3, 3]$

160. $-1 \leq \log_2(x^2 + 5x + 8) \leq 1$
 $\Rightarrow \frac{1}{2} \leq (x^2 + 5x + 8) \leq 2$
 $\Rightarrow x^2 + 5x + \frac{15}{2} \geq 0$
 $\Rightarrow x^2 + 2\left(\frac{5}{2}\right)x + \left(\frac{5}{2}\right)^2 - \left(\frac{5}{2}\right)^2 + \frac{15}{2} \geq 0$
 $\Rightarrow \left(x + \frac{5}{2}\right)^2 + \frac{5}{4} \geq 0$ and $x^2 + 5x + 6 \leq 0$
 $\Rightarrow (x + 3)(x + 2) \leq 0 \Rightarrow x \in [-3, -2]$

161. $f(x) = \sqrt{9 - x^2}$
 $f(0) = 3, f(3) = 0$
 $\therefore 0 \leq f(x) \leq 3$
 $\therefore x \in [0, 3]$

162. $f(x) = \frac{x+2}{|x+2|}$

$$f(x) = \begin{cases} -1, & x < -2 \\ 1, & x > -2 \end{cases}$$

\therefore Range of $f(x)$ is $\{-1, 1\}$.

163. Let $\frac{x^2 + 34x - 71}{x^2 + 2x - 7} = y$
 $\Rightarrow x^2(1 - y) + 2(17 - y)x + (7y - 71) = 0$
 For real value of x , $b^2 - 4ac \geq 0$
 $\Rightarrow y^2 - 14y + 45 \geq 0 \Rightarrow y \geq 9, y \leq 5$.

164. Dom (f) = $\mathbb{R} - \{2\}$

For Range (f), let $y = f(x) = \frac{x^2 - 4}{x - 2}$

$\therefore y = \frac{(x-2)(x+2)}{(x-2)}$

$\therefore y = (x + 2)$
 Since, Dom (f) = $\mathbb{R} - \{2\}$

$\therefore x \neq 2$
 $\therefore y \neq (2 + 2)$ i.e. $y \neq 4$
 \therefore Range (f) = $\mathbb{R} - \{4\}$

165. Let $y = \frac{x^2 - x + 4}{x^2 + x + 4}$
 $\Rightarrow (y - 1)x^2 + (y + 1)x + 4y - 4 = 0$
 For real value of x , $b^2 - 4ac \geq 0$
 $\Rightarrow (y + 1)^2 - 4(y - 1)(4y - 4) \geq 0$
 $\Rightarrow -15y^2 + 34y - 15 \geq 0$
 $\Rightarrow 15y^2 - 34y + 15 \geq 0$
 $\Rightarrow \left(y - \frac{3}{5}\right)\left(y - \frac{5}{3}\right) \leq 0$
 $\Rightarrow \frac{3}{5} \leq y \leq \frac{5}{3}$

166. Since maximum and minimum values of $\cos - \sin x$ are $\sqrt{2}$ and $-\sqrt{2}$ respectively, therefore range of $f(x)$ is $[-\sqrt{2}, \sqrt{2}]$.

167. $\cos 2x + 7 = a(2 - \sin x) \Rightarrow a = \frac{\cos 2x + 7}{2 - \sin x}$

$$\Rightarrow a = \frac{1 - 2\sin^2 x + 7}{2 - \sin x} = \frac{2(4 - \sin^2 x)}{2 - \sin x}$$

$\Rightarrow a = 2(2 + \sin x)$

$\therefore a \in [2, 6]$ [$\because -1 \leq \sin x \leq 1$]



168. Let $y = \log_e(3x^2 + 4)$

$$\Rightarrow 3x^2 + 4 = e^y \Rightarrow x^2 = \frac{e^y - 4}{3}$$

Since, $x^2 \geq 0$

$$\therefore \frac{e^y - 4}{3} \geq 0 \Rightarrow e^y - 4 \geq 0 \Rightarrow y \geq \log_e 4$$

$$\Rightarrow y \geq 2 \log_e 2$$

So, range = $[2 \log_e 2, \infty)$

169. Let $y = \log_e \sqrt{4 - x^2} \Rightarrow e^y = \sqrt{4 - x^2}$

$$\Rightarrow e^{2y} = 4 - x^2 \Rightarrow x^2 = 4 - e^{2y} \Rightarrow x = \sqrt{4 - e^{2y}}$$

$$\therefore 4 - e^{2y} \geq 0$$

$$\Rightarrow e^{2y} \leq 4 \Rightarrow 2y \leq \log_e 4$$

$$\Rightarrow y \leq \frac{1}{2} \log_e 4 \Rightarrow y \leq \log_e 2$$

$$\therefore y \in (-\infty, \log_e 2]$$

170.
$$f(\theta) = \begin{vmatrix} 1 & \tan \theta & 1 \\ -\tan \theta & 1 & \tan \theta \\ -1 & -\tan \theta & 1 \end{vmatrix}$$

$$\therefore f(\theta) = 2\sec^2 \theta \geq 2$$

$$\therefore \text{range of } f \text{ is } [2, \infty).$$

171. $f(x) = \tan \sqrt{\frac{\pi^2}{9} - x^2}$

$f(x)$ is real valued function when $\frac{\pi^2}{9} - x^2 \geq 0$

$$\Rightarrow x^2 \leq \frac{\pi^2}{9} \Rightarrow x \in \left[-\frac{\pi}{3}, \frac{\pi}{3}\right] = \text{Domain of } f(x)$$

When domain is in closed interval, we use differentiation method.

$$f'(x) = \sec^2 \sqrt{\frac{\pi^2}{9} - x^2} \cdot \frac{1}{2\sqrt{\frac{\pi^2}{9} - x^2}} (-2x)$$

When $f'(x) = 0$, $x = 0$

Finding values of $f(x)$ when $x = 0, -\frac{\pi}{3}, \frac{\pi}{3}$

[End points of domain]

$$\therefore f(0) = \tan \sqrt{\frac{\pi^2}{9}} = \sqrt{3} \text{ and } f\left(-\frac{\pi}{3}\right) = f\left(\frac{\pi}{3}\right) = 0$$

$$\therefore \text{Range of function} = [0, \sqrt{3}]$$

[Taking least value and greatest value for range]



Evaluation Test

1. Given, $g(x) = x^2 + x - 2$

and $\frac{1}{2}(\text{gof})(x) = 2x^2 - 5x + 2$

$$\Rightarrow g(f(x)) = 4x^2 - 10x + 4$$

$$\Rightarrow (f(x))^2 + f(x) - 2 = 4x^2 - 10x + 4$$

$$\Rightarrow (f(x))^2 + f(x) - (4x^2 - 10x + 6) = 0$$

$$\Rightarrow f(x) = \frac{-1 \pm \sqrt{1 + 16x^2 - 40x + 24}}{2}$$

$$= \frac{-1 \pm (4x - 5)}{2} = 2x - 3, -2x + 2$$

2. $\text{fog}(x) = x^3 - \frac{1}{x^3}$

Since, $\left(x - \frac{1}{x}\right)^3 = x^3 - \frac{1}{x^3} - 3x \cdot \frac{1}{x} \left(x - \frac{1}{x}\right)$

$$\therefore x^3 - \frac{1}{x^3} = \left(x - \frac{1}{x}\right)^3 + 3\left(x - \frac{1}{x}\right)$$

Given, $f(g(x)) = x^3 - \frac{1}{x^3}$

$$\therefore f(g(x)) = \left(x - \frac{1}{x}\right)^3 + 3\left(x - \frac{1}{x}\right)$$

$$\therefore f\left(x - \frac{1}{x}\right) = \left(x - \frac{1}{x}\right)^3 + 3\left(x - \frac{1}{x}\right)$$

Put $x - \frac{1}{x} = t$

$$\therefore f(t) = t^3 + 3t \quad \therefore f(x) = x^3 + 3x$$

$$\therefore f'(x) = 3x^2 + 3$$

3. Given, $f(x + y) = f(x) + f(y) \quad \forall x, y \in \mathbb{R}$

$$\Rightarrow f(2) = f(1+1) = f(1) + f(1) = 2f(1)$$

$$f(3) = f(2+1) = f(2) + f(1)$$

$$= 2f(1) + f(1) = 3f(1)$$

Continuing in this way, we get

$$f(r) = rf(1) \in \mathbb{N}$$

$$\therefore \sum_{r=1}^n f(r) = \sum_{r=1}^n rf(1) = f(1) \sum_{r=1}^n r$$

$$= 7(1 + 2 + 3 + \dots + n)$$

$$= \frac{7n(n+1)}{2}$$



4. Given, $f(x) = x^2 - 3$
 $\therefore f(-1) = (-1)^2 - 3 = -2$
 $\Rightarrow (f \circ f)(-1) = f(-2) = (-2)^2 - 3 = 1$
 $\Rightarrow (f \circ f \circ f)(-1) = f(1) = 1^2 - 3 = -2$ (i)
 Similarly, $(f \circ f \circ f)(0) = 33$ (ii)
 and $(f \circ f \circ f)(1) = -2$ (iii)
 From (i), (ii) and (iii), we get
 $(f \circ f \circ f)(-1) + (f \circ f \circ f)(0) + (f \circ f \circ f)(1)$
 $= -2 + 33 - 2 = 29 = f(4\sqrt{2})$

5. $x = (\sqrt{3} + 1)^5$
 $= {}^5C_0(\sqrt{3})^5 + {}^5C_1(\sqrt{3})^4 + {}^5C_2(\sqrt{3})^3$
 $\quad + {}^5C_3(\sqrt{3})^2 + {}^5C_4(\sqrt{3})^1 + {}^5C_5$
 $= (\sqrt{3})^5 + 5(9) + 10(3\sqrt{3}) + 30 + 5\sqrt{3} + 1$
 $= 76 + 44\sqrt{3} = 152.20$

$\therefore [x] = [152.20] = 152$

6. $f(x + y) = f(x) + f(y)$ (i)
 Putting $x = y = 1$ in (i), we get
 $f(2) = 2f(1)$
 $\Rightarrow f(2) = 2(5)$ [$\because f(1) = 5$ (given)]
 Putting $x = 2$ and $y = 1$ in (i), we get
 $f(3) = f(2) + f(1) = 3(5)$
 Similarly, $f(4) = 4(5)$
 $f(5) = 5(5)$

...

...

...

$f(100) = 100(5) = 500$

7. For any $x \in (0, 2)$, $x \neq x + 1$

$\therefore (x, x) \notin S$.

$\therefore S$ is not a reflexive relation.

So, S is not an equivalence relation.

$T = \{(x, y) : x - y \text{ is an integer}\}$

For any $x \in \mathbb{R}$,

$x - x = 0$, which is an integer.

$\Rightarrow (x, x) \in T$

$\therefore T$ is reflexive on \mathbb{R} .

Let $(x, y) \in T$. Then,

$x - y$ is an integer

$\Rightarrow y - x$ is an integer

$\Rightarrow (y, x) \in T$

$\therefore T$ is symmetric on \mathbb{R} .

Let $(x, y) \in T$ and $(y, z) \in T$.

Then, $x - y$ is an integer and $y - z$ is an integer

$\Rightarrow x - z$ is an integer

$\Rightarrow (x, z) \in T$

$\therefore T$ is transitive on \mathbb{R} .

So, T is an equivalence relation on \mathbb{R} .

8. Here, $(0, 3) \in R$, because $0 = 0 \times 3$

But, $(3, 0) \notin R$,

because $3 \neq (\text{any rational number}) \times 0$

So, R is not a symmetric relation and hence it is not an equivalence relation.

$S = \left\{ \left(\frac{m}{n}, \frac{p}{q} \right) : m, n, p \text{ and } q \text{ are integers} \right.$

$\left. \text{such that } n, q \neq 0 \text{ and } qm = pn \right\}$

Let $m, n \in \mathbb{Z}$ such that $n \neq 0$. Then,

$mn = nm \Rightarrow \frac{m}{n} = \frac{m}{n} \Rightarrow \left(\frac{m}{n}, \frac{m}{n} \right) \in S$

So, S is reflexive.

Let $\left(\frac{m}{n}, \frac{p}{q} \right) \in S$. Then,

$qm = pn \Rightarrow np = mq$

$\Rightarrow \left(\frac{p}{q}, \frac{m}{n} \right) \in S$

So, S is symmetric.

Let $\left(\frac{m}{n}, \frac{p}{q} \right) \in S$ and $\left(\frac{p}{q}, \frac{r}{s} \right) \in S$.

Then, $qm = pn$ and $sp = rq$

$\Rightarrow (qm)(sp) = (pn)(rq)$

$\Rightarrow sm = rn \Rightarrow \left(\frac{m}{n}, \frac{r}{s} \right) \in S$

So, S is transitive.

Hence, S is an equivalence relation.

9. $\theta \in P$

$\Rightarrow \sin \theta - \cos \theta = \sqrt{2} \cos \theta$

$\Rightarrow (\sin \theta - \cos \theta)^2 = 2 \cos^2 \theta$

$\Rightarrow \sin^2 \theta + \cos^2 \theta - 2 \sin \theta \cos \theta = 2 \cos^2 \theta$

$\Rightarrow \cos^2 \theta + 2 \sin \theta \cos \theta + \sin^2 \theta = 2 \sin^2 \theta$

$\Rightarrow (\cos \theta + \sin \theta)^2 = 2 \sin^2 \theta$

$\Rightarrow \cos \theta + \sin \theta = \sqrt{2} \sin \theta$

$\Rightarrow \theta \in Q$

$\therefore P = Q$



10. $f(x)$ is defined for

$$-1 \leq \frac{8 \cdot 3^{x-2}}{1-3^{2(x-1)}} \leq 1$$

$$\Rightarrow -1 \leq \frac{(3^2-1)(3^{x-2})}{1-3^{2x-2}} \leq 1$$

$$\Rightarrow -1 \leq \frac{3^x - 3^{x-2}}{1-3^{2x-2}} \leq 1$$

$$\Rightarrow \frac{3^x - 3^{x-2}}{1-3^{2x-2}} + 1 \geq 0 \text{ and } \frac{3^x - 3^{x-2}}{1-3^{2x-2}} - 1 \leq 0$$

$$\Rightarrow \frac{1+3^x - 3^{x-2} - 3^{2x-2}}{1-3^{2x-2}} \geq 0$$

$$\text{and } \frac{3^x - 3^{x-2} - 1 + 3^{2x-2}}{1-3^{2x-2}} \leq 0$$

$$\Rightarrow \frac{(3^x+1)(3^{x-2}-1)}{(3^x \cdot 3^{x-2}-1)} \geq 0 \text{ and } \frac{(3^x-1)(3^{x-2}+1)}{(3^{2x-2}-1)} \geq 0$$

$$\Rightarrow \frac{(3^{x-2}-1)}{(3^x \cdot 3^{x-2}-1)} \geq 0 \text{ and } \frac{(3^x-1)}{(3^{2x-2}-1)} \geq 0$$

$$\Rightarrow \frac{(3^x-3^2)}{(3^{2x}-3^2)} \geq 0 \text{ and } \frac{(3^x-1)}{(3^{2x}-3^2)} \geq 0$$

$$\Rightarrow \frac{(3^x-3^2)}{(3^x-3)} \geq 0 \text{ and } \frac{(3^x-1)}{(3^x-3)} \geq 0$$

$$\Rightarrow x \in (-\infty, 1] \cup [2, \infty) \text{ and } x \in (-\infty, 0] \cup (1, \infty)$$

$$\Rightarrow x \in (-\infty, 0] \cup [2, \infty)$$

11. $f(x-y) = f(x)f(y) - f(a-x)f(a+y)$

Putting $x=0$ and $y=0$, we get

$$f(0) = \{f(0)\}^2 - \{f(a)\}^2$$

$$\Rightarrow 1 = 1 - \{f(a)\}^2 \Rightarrow f(a) = 0$$

Now, $f(2a-x) = f(a-(x-a))$

$$= f(a)f(x-a) - f(a-a)f(a+x-a)$$

$$= f(a)f(x-a) - f(0)f(x)$$

$$= f(a)f(x-a) - f(x)$$

$$\dots [\because f(0) = 1 \text{ (given)}]$$

$$= -f(x)$$

12. $n[(A \times B) \cap (B \times A)] = n[(A \cap B) \times (B \cap A)]$

$$= n(A \cap B) \times n(B \cap A) = 3 \times 3 = 9$$

13. Given, $2 \cos^2 \theta + \sin \theta \leq 2$ and $\frac{\pi}{2} \leq \theta \leq \frac{3\pi}{2}$

$$\Rightarrow 2 - 2 \sin^2 \theta + \sin \theta \leq 2$$

$$\Rightarrow 2 \sin^2 \theta - \sin \theta \geq 0$$

$$\Rightarrow \sin \theta (2 \sin \theta - 1) \geq 0$$

$$\Rightarrow \sin \theta \geq 0 \text{ and } 2 \sin \theta - 1 \geq 0 \text{ or}$$

$$\sin \theta \leq 0 \text{ and } 2 \sin \theta - 1 \leq 0$$

Case I:

$$\sin \theta \geq 0 \text{ and } 2 \sin \theta - 1 \geq 0$$

$$\Rightarrow \sin \theta \geq 0 \text{ and } \sin \theta \geq \frac{1}{2}$$

$$\Rightarrow \frac{\pi}{6} \leq \theta \leq \frac{5\pi}{6}$$

$$\therefore A \cap B = \left\{ \theta : \frac{\pi}{2} \leq \theta \leq \frac{5\pi}{6} \right\}$$

$$\dots \left[\because B = \left\{ \theta : \frac{\pi}{2} \leq \theta \leq \frac{3\pi}{2} \right\} \right]$$

Case II:

$$\sin \theta \leq 0 \text{ and } 2 \sin \theta - 1 \leq 0$$

$$\Rightarrow \sin \theta \leq 0 \text{ and } \sin \theta \leq \frac{1}{2}$$

$$\Rightarrow \pi \leq \theta \leq 2\pi$$

$$\therefore A \cap B = \left\{ \theta : \pi \leq \theta \leq \frac{3\pi}{2} \right\}$$

$$\dots \left[\because B = \left\{ \theta : \frac{\pi}{2} \leq \theta \leq \frac{3\pi}{2} \right\} \right]$$

From Case I and II, we get

$$A \cap B = \left\{ \theta : \frac{\pi}{2} \leq \theta \leq \frac{5\pi}{6} \right\} \cup \left\{ \theta : \pi \leq \theta \leq \frac{3\pi}{2} \right\}$$

14. Given, $f(x) = \sqrt{\log_{10} \left\{ \frac{\log_{10} x}{2(3-\log_{10} x)} \right\}}$

Now, $f(x)$ is defined, if

$$\log_{10} \left\{ \frac{\log_{10} x}{2(3-\log_{10} x)} \right\} \geq 0, \frac{\log_{10} x}{2(3-\log_{10} x)} > 0$$

and $x > 0$

$$\Rightarrow \frac{\log_{10} x}{2(3-\log_{10} x)} \geq 10^0 = 1, \frac{\log_{10} x}{(3-\log_{10} x)} > 0$$

and $x > 0$

$$\Rightarrow \frac{3(\log_{10} x - 2)}{2(\log_{10} x - 3)} \leq 0, \frac{\log_{10} x}{\log_{10} x - 3} < 0 \text{ and } x > 0$$

$$\Rightarrow 2 \leq \log_{10} x < 3, 0 < \log_{10} x < 3 \text{ and } x > 0$$

$$\Rightarrow 10^2 \leq x < 10^3, 10^0 < x < 10^3 \text{ and } x > 0$$

$$\Rightarrow 10^2 \leq x < 10^3$$

$$\Rightarrow x \in [10^2, 10^3)$$



Hints



Classical Thinking

- $a = 21, d = 16 - 21 = -5$
 $t_n = a + (n - 1)d$
 $\therefore t_{15} = 21 + (15 - 1)(-5) = 21 - 70 = -49$
- $a = \sqrt{3}, d = \sqrt{12} - \sqrt{3} = \sqrt{3}$
 $\therefore t_{10} = \sqrt{3} + 9\sqrt{3} = 10\sqrt{3} = \sqrt{300}$
- Given series

$$\left(3 - \frac{1}{n}\right) + \left(3 - \frac{2}{n}\right) + \left(3 - \frac{3}{n}\right) + \dots \dots \dots \text{(A.P.)}$$

Therefore, common difference

$$d = \left(3 - \frac{2}{n}\right) - \left(3 - \frac{1}{n}\right) = -\frac{1}{n} \text{ and first term}$$

$$a = \left(3 - \frac{1}{n}\right)$$

Now, p^{th} term of the series = $a + (p - 1)d$

$$= \left(3 - \frac{1}{n}\right) + (p - 1)\left(-\frac{1}{n}\right)$$

$$= 3 - \frac{1}{n} + \frac{1}{n} - \frac{p}{n} = \left(3 - \frac{p}{n}\right)$$
- $d - c = e - d$
 $\Rightarrow 2d = e + c$
 $\Rightarrow 2d - 2c = e + c - 2c$
 $\Rightarrow 2(d - c) = e - c$
- a, b, c are in A.P., dividing by bc we get

$$\frac{a}{bc}, \frac{1}{c}, \frac{1}{b} \text{ are in A.P.}$$
- $a = 3, d = 3$

Let there be n terms.

 $\therefore 3 + (n - 1)3 = 111$
 $\Rightarrow n = 37$
- $a = 72, d = -2$

Let n^{th} term be 40.

 $\therefore t_n = a + (n - 1)d$
 $\therefore 40 = 72 + (n - 1)(-2)$
 $\Rightarrow n = 17$
- $d = -1 + 2i, t_4 = t_3 + d = 6 - 2i + (-1 + 2i) = 5$

which is purely real.

- Given that, 9^{th} term = $a + (9 - 1)d = 0$
 $\Rightarrow a + 8d = 0$
 Now, ratio of 29^{th} and 19^{th} terms

$$= \frac{a + 28d}{a + 18d} = \frac{(a + 8d) + 20d}{(a + 8d) + 10d} = \frac{20d}{10d} = \frac{2}{1}$$
- Let the first term and common difference of an A.P. be A and D respectively.
 Now, p^{th} term = $A + (p - 1)D = a$
 q^{th} term = $A + (q - 1)D = b$
 and r^{th} term = $A + (r - 1)D = c$

$$\therefore a(q - r) + b(r - p) + c(p - q)$$

$$= a\left\{\frac{b - c}{D}\right\} + b\left\{\frac{c - a}{D}\right\} + c\left\{\frac{a - b}{D}\right\}$$

$$= \frac{1}{D}(ab - ac + bc - ab + ca - bc) = 0$$
- $S_n = 3(4^n - 1)$
 $\therefore S_{n-1} = 3(4^{n-1} - 1)$
 $\therefore t_n = S_n - S_{n-1} = 3(4^n - 1) - 3(4^{n-1} - 1) = 9(4^{n-1})$
- Required sum = $1 + 3 + 5 + \dots$ upto n terms

$$= \frac{n}{2} [2 \times 1 + (n - 1)2]$$

$$= n^2$$
- Given that first term $a = 10$, last term = 50
 and sum $S = 300$

$$\therefore S = \frac{n}{2}(a +) \Rightarrow 300 = \frac{n}{2}(10 + 50) \Rightarrow n = 10$$
- $S_n = \frac{n}{2} [2a + (n - 1)d]$
 $\Rightarrow 406 = \frac{n}{2} [6 + (n - 1)4]$
 $\Rightarrow 812 = n[6 + 4n - 4] \Rightarrow 812 = 2n + 4n^2$
 $\Rightarrow 406 = 2n^2 + n \Rightarrow 2n^2 + n - 406 = 0$
 $\Rightarrow n = \frac{-1 \pm \sqrt{1 + 4 \cdot 2 \cdot 406}}{2 \cdot 2} = \frac{-1 \pm \sqrt{3249}}{4}$

$$= \frac{-1 \pm 57}{4}$$

Taking (+) sign, $n = \frac{-1 + 57}{4} = 14$



15. $S_n = 3n^2 - n$
 $\Rightarrow 3n^2 - n = \frac{n}{2} [2a + (n-1)d]$
 $\Rightarrow a = 2$
16. $S_n = \frac{n}{2} [2a + (n-1)d]$
 $\therefore S_{16} = \frac{16}{2} [2(4) + (15)d]$
 $\Rightarrow 784 = 8(8 + 15d)$
 $\Rightarrow 8 + 15d = \frac{784}{8}$
 $\Rightarrow 15d = 90$
 $\Rightarrow d = 6$
17. $t_7 = 40 \Rightarrow a + 6d = 40$
 $S_{13} = \frac{13}{2} [2a + (13-1)d] = 13(a + 6d) = 520$
18. $t_4 = a + 3d = 4$ and
 $S_7 = \frac{7}{2} [2a + (7-1)d] = \frac{7}{2} [2a + 6d]$
 $= 7(a + 3d)$
 $= 7(4) = 28$
19. The terms of given sequence are in A.P. with
 $a = 1, d = 5$ and $S_n = 148$
 $\therefore \frac{n}{2} [2a + (n-1)d] = 148 \Rightarrow n = 8$
 Now, $x = n^{\text{th}}$ term $\Rightarrow x = a + (n-1)d = 36$
20. $(x+1) + (x+4) + \dots + (x+28) = 155$
 Let n be the number of terms in the A.P. on
 L.H.S. Then,
 $x + 28 = (x+1) + (n-1)3 \Rightarrow n = 10$
 $\therefore (x+1) + (x+4) + \dots + (x+28) = 155$
 $\Rightarrow \frac{10}{2} [(x+1) + (x+28)] = 155$
 $\Rightarrow x = 1$
21. $S_5 = \frac{1}{4} (S_{10} - S_5) \Rightarrow 5S_5 = S_{10}$
 $\therefore 5 \times \frac{5}{2} (2 \times 2 + 4d) = \frac{10}{2} (2 \times 2 + 9d)$
 $\Rightarrow d = -6$
22. Here, $\frac{1}{3}, A_1, A_2, \frac{1}{24}$ will be in A.P.,
 then $A_1 - \frac{1}{3} = \frac{1}{24} - A_2$
 $\Rightarrow A_1 + A_2 = \frac{3}{8}$ (i)

Now, A_1 is a arithmetic mean of $\frac{1}{3}$ and A_2 .

$$\therefore 2A_1 = \frac{1}{3} + A_2 \Rightarrow 2A_1 - A_2 = \frac{1}{3} \text{(ii)}$$

From (i) and (ii), we get $A_1 = \frac{17}{72}$ and $A_2 = \frac{5}{36}$

23. Let the two numbers be a and b and let
 A_1, A_2, \dots, A_n be the n A.M.'s between them.
 Then $a, A_1, A_2, \dots, A_n, b$ are in A.P. and let d
 be the common difference.

Now, $T_{n+2} = b = a + (n+2-1)d$

$$\Rightarrow d = \frac{b-a}{n+1}$$

Also, $A_1 + A_2 + \dots + A_n = S_{n+1} - a$

$$= \frac{1}{2} (n+1) \left[2a + (n+1-1) \frac{(b-a)}{(n+1)} \right] - a$$

$$= \frac{n}{2} [2a + (b-a)] = \frac{n}{2} (a+b) = n \left(\frac{a+b}{2} \right)$$

$$\therefore S = Na$$

24. Let the three numbers be $a+d, a, a-d$.
 therefore, $a+d+a+a-d = 33$

$$\Rightarrow a = 11$$

$$\text{and } a(a+d)(a-d) = 792$$

$$\Rightarrow 11(121 - d^2) = 792 \Rightarrow d = 7$$

The required numbers are 4, 11, 18.

Hence, the smallest number is 4.

$$25. t_n = ar^{n-1} = 1.(2)^{n-1} = 2^{n-1}$$

26. Given sequence is $\sqrt{2}, \sqrt{10}, \sqrt{50}, \dots$

Common ratio $r = \sqrt{5}$, first term $a = \sqrt{2}$,
 then 7th term

$$t_7 = \sqrt{2}(\sqrt{5})^{7-1} = \sqrt{2}(\sqrt{5})^6 = \sqrt{2}(5)^3 = 125\sqrt{2}$$

$$28. t_n = ar^{n-1} = 1 \left(\frac{1}{2} \right)^{n-1} = \left(\frac{1}{2} \right)^{n-1}$$

$$29. a = 3, r = \frac{\left(\frac{-3}{2} \right)}{3} = \frac{-1}{2}$$

$$\Rightarrow t_n = ar^{n-1} = 3 \left(\frac{-1}{2} \right)^{n-1}$$

30. Let r be common ratio of G.P.

$$\Rightarrow t_3 = r^2, t_5 = r^4$$

$$\therefore t_3 + t_5 = 90 \Rightarrow r^2 + r^4 = 90$$

$$\Rightarrow r^2 = 9$$

$$\Rightarrow r = \pm 3$$



31. Accordingly, $ar^9 = 9$ and $ar^3 = 4$
 $r^3 = \frac{3}{2}$ and $a = \frac{8}{3}$
 \therefore 7th term i.e., $ar^6 = \frac{8}{3} \left(\frac{3}{2}\right)^2 = 6$
Trick : 7th term is equidistant from 10th and 4th
 so it will be $\sqrt{9 \times 4} = 6$.
33. $t_3 = ar^{3-1} = ar^2 = 20$ and
 $t_7 = ar^{7-1} = ar^6 = 320$
 Solving, $a = 5$ and $r = 2$
34. $a, 8, b$ are in G.P. and $a \neq b$
 $\Rightarrow \frac{8}{a} = \frac{b}{8} \Rightarrow ab = 64$
 and $a, b, -8$ are in A.P.
 $\Rightarrow b - a = -8 - b$
 $\therefore b = \left(\frac{a-8}{2}\right)$
 Solving, $a = 16$ and $b = 4$
35. $a = 5, r = 3$
 $S_n = \frac{a(r^n - 1)}{r - 1} = \frac{5(3^n - 1)}{2}$
36. $a = 3$ and $r = \frac{12}{3} = 4 > 1$
 $\therefore S_n = a \left[\frac{r^n - 1}{r - 1} \right] = 3 \left[\frac{4^n - 1}{4 - 1} \right] = 4^n - 1$
38. $a = 1, r = 3$
 $S_n = \frac{a(r^n - 1)}{r - 1}$
 $3280 = \frac{3^n - 1}{2}$
 $6561 = 3^n$
 $\Rightarrow 3^8 = 3^n \Rightarrow n = 8$
39. Let n be the number of terms needed.
 For G.P. $2, 2^2, 2^3, \dots$, $a = 2, r = 2$ and $S_n = 30$
 $S_n = \frac{a(r^n - 1)}{r - 1} \Rightarrow 30 = \frac{2(2^n - 1)}{2 - 1} \Rightarrow n = 4$
40. $S_8 = 82$ (S_4)
 Let the G.P. be $a + ar + ar^2 + \dots$, then
 $\frac{a(r^8 - 1)}{(r - 1)} = 82 \left\{ \frac{a(r^4 - 1)}{r - 1} \right\}$
 $(r^4 - 1)(r^4 + 1) = 82(r^4 - 1)$
 $\Rightarrow r^4 + 1 = 82$
 $r^4 = 81$
 $\Rightarrow r = 3$

41. $S_n = \frac{a(r^n - 1)}{r - 1}, r = 2$
 $\therefore S_8 = \frac{a(2^8 - 1)}{2 - 1} \Rightarrow a(2^8 - 1) = 510 \Rightarrow a = 2$
 $\therefore t_3 = 2(2)^{3-1} = 2(2)^2 = 8$
42. $S_n = 2 + 22 + 222 + \dots$ n terms
 $= 2 [1 + 11 + 111 + \dots$ n terms]
 $= \frac{2}{9} [(10 - 1) + (100 - 1) + (1000 - 1)$
 $+ \dots$ n terms]
 $= \frac{2}{9} \left[10 \left(\frac{10^n - 1}{10 - 1} \right) - n \right] = \frac{2}{9} \left[\frac{10}{9} (10^n - 1) - n \right]$
 $= \frac{2}{81} [10(10^n - 1) - 9n]$
43. $S_n = 0.9 + 0.99 + 0.999 + \dots$ n terms
 $= 1 - 0.1 + 1 - 0.01 + 1 - 0.001$
 $+ \dots$ n terms
 $= 1 + 1 + 1 + \dots$ n terms
 $- [0.1 + 0.01 + 0.001 + \dots$ n terms]
 $= n - 0.1 \left[\frac{1 - (0.1)^n}{1 - 0.1} \right]$
 $= n - \frac{1}{9} [1 - (0.1)^n]$
 $= \frac{9n - [1 - (0.1)^n]}{9}$
44. $a = 2, S_\infty = 6$
 Now, $S_\infty = \frac{a}{1 - r}$
 $\Rightarrow 6 = \frac{2}{1 - r}$
 $\Rightarrow 1 - r = \frac{1}{3}$
 $\Rightarrow r = 1 - \frac{1}{3}$
 $\Rightarrow r = \frac{2}{3}$
45. According to condition, $\frac{3/4}{1 - r} = \frac{4}{3}$
 $\Rightarrow r = \frac{7}{16}$
46. $\frac{g_1}{p} = \frac{q}{g_2} \Rightarrow g_1 g_2 = pq$



47. Let 1, a, b, 64
 $\Rightarrow a^2 = b$ and $b^2 = 64a$
 $\Rightarrow a = 4$ and $b = 16$
48. Let the numbers be a, ar, ar²
Sum = 70 $\Rightarrow a(1 + r + r^2) = 70$
It is given that 4a, 5ar, 4ar² are in A.P.
 $\therefore 2(5ar) = 4a + 4ar^2 \Rightarrow r = 2$ or $r = \frac{1}{2}$
Substituting values of r, a = 10 and a = 40
 \therefore The numbers are 10, 20, 40 or 40, 20, 10
49. Let numbers are $\frac{a}{r}$, a, ar
According to given conditions,
 $\frac{a}{r} \cdot a \cdot ar = 216$
 $\Rightarrow a = 6$
And, sum of product pairwise = 156
 $\Rightarrow \frac{a}{r} \cdot a + \frac{a}{r} \cdot ar + a \cdot ar = 156$
 $\Rightarrow r = 3$
Hence, numbers are 2, 6, 18.
Trick : Since $2 \times 6 \times 18 = 216$ (as given) and no other option gives the value.
50. Considering corresponding A.P.
 $a + 6d = 10$ and $a + 11d = 25 \Rightarrow d = 3, a = -8$
 $\Rightarrow t_{20} = a + 19d = -8 + 57 = 49$
Hence, 20th term of the corresponding H.P. is $\frac{1}{49}$.
53. $\therefore H < G < A$
54. (A.M.) (H.M.) = (G.M.)²
 $\Rightarrow 9 \cdot 36 = (\text{G.M.})^2 \Rightarrow \text{G.M.} = 18$
56. $G^2 = AH$
 $\Rightarrow 144 = 25H$
 $\Rightarrow H = 5.76$
58. Let $S = 1 + 3x + 5x^2 + 7x^3 + \dots$
Then, $xS = 1x + 3x^2 + 5x^3 + \dots$
 $S - xS = 1 + 2x + 2x^2 + 2x^3 + \dots$ to ∞
 $\therefore S(1 - x) = 1 + 2x + 2x^2 + 2x^3 + \dots$ to ∞
 $= 1 + \frac{2x}{1-x} = \frac{1-x+2x}{1-x}$
 $\therefore S = \frac{1+x}{(1-x)^2}$
59. Here $a = 3, d = 2$ and $r = r$
Now $S_\infty = \frac{a}{1-r} + \frac{dr}{(1-r)^2}$ ($|r| < 1$)
 $\therefore S_\infty = \frac{3}{1-r} + \frac{2r}{(1-r)^2} \quad \therefore \frac{44}{9} = \frac{3-r}{(1-r)^2}$
 $\therefore 44r^2 - 79r + 17 = 0$
 $\therefore r = \frac{1}{4}$ or $\frac{17}{11}$
But, $r \neq \frac{17}{11}$
 $\therefore r = \frac{1}{4}$
60. $\sum_{r=1}^n (2r+5) = 2 \sum_{r=1}^n r + \sum_{r=1}^n 5 = \frac{2(n)(n+1)}{2} + 5n$
 $= n(n+6)$
61. $(2^2 - 1^2) + (4^2 - 3^2) + (6^2 - 5^2) + \dots$
 $= (2^2 + 4^2 + 6^2 + \dots) - (1^2 + 3^2 + 5^2 + \dots)$
 $= \sum_{r=1}^n (2r)^2 - \sum_{r=1}^n (2r-1)^2 = \sum_{r=1}^n 4r - \sum_{r=1}^n 1$
 $= 4 \left[\frac{n(n+1)}{2} \right] - n = n(2n+1)$
62. $\frac{n(n+1)(2n+1)}{6} = 1015$
 $\therefore n(n+1)(2n+1) = 6090$
 $\Rightarrow n(n+1)(2n+1) = 14 \times 15 \times 29$
 $\Rightarrow n = 14$
63. $(31)^2 + (32)^2 + (33)^2 + \dots + (60)^2$
 $= [(1)^2 + (2)^2 + (3)^2 + \dots + (60)^2]$
 $\quad - [(1)^2 + (2)^2 + (3)^2 + \dots + (30)^2]$
 $= \sum_{r=1}^{60} r^2 - \sum_{r=1}^{30} r^2 = 64355$
64. The first factors of the terms of the given series is 1, 2, 3, 4, ..., n and second factors of the terms of the given series is 2, 3, 4,(n+1)
 $\therefore n^{\text{th}}$ term of the given series
 $= n(n+1) = n^2 + n$
Hence, sum =
 $\Sigma n^2 + \Sigma n = \frac{1}{6} n(n+1)(2n+1) + \frac{n}{2} (n+1)$
 $= \frac{1}{6} n(n+1)(2n+1+3)$
 $= \frac{1}{3} n(n+1)(n+2)$



65. $1^3 + 2^3 + 3^3 + \dots + 25^3 = \sum_{r=1}^{25} r^3$
 $= \frac{(25)^2(25+1)^2}{4}$
 $= 105625$
66. $2(1)^2 + 3(2)^2 + 4(3)^2 + \dots$ upto 10 terms
 $= \sum_{r=1}^{10} (r+1)r^2 = \sum_{r=1}^{10} r^3 + \sum_{r=1}^{10} r^2 = 3410$
67. $y = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$
 $= \log_e(1+x)$
 $\therefore 1+x = e^y \Rightarrow x = e^y - 1$
68. Sum of given series $= y + \frac{y^2}{2} + \frac{y^3}{3} + \dots$,
 where $y = x^2$.
 \therefore Sum of given series $= -\log(1-y)$
 $= -\log_e(1-x^2)$
69. $\log_e 3 - \frac{\log_e 3^2}{2^2} + \frac{\log_e 3^3}{3^2} - \frac{\log_e 3^4}{4^2} + \dots$
 $= \log_e 3 \left\{ 1 - \frac{2}{2^2} + \frac{3}{3^2} - \frac{4}{4^2} + \dots \right\}$
 $= \log_e 3 \left\{ 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots \right\}$
 $= \log_e 3 \log_e(1+1) = \log_e 3 \log_e 2$



Critical Thinking

1. $\frac{1}{1+\sqrt{x}}, \frac{1}{1-x}, \frac{1}{1-\sqrt{x}}, \dots$
 i.e., $\frac{1-\sqrt{x}}{1-x}, \frac{1}{1-x}, \frac{1+\sqrt{x}}{1-x}, \dots$,
 which is an A.P. with $d = \frac{\sqrt{x}}{1-x}$
 \therefore The fourth term $= t_3 + d = \frac{1+\sqrt{x}}{1-x} + \frac{\sqrt{x}}{1-x}$
 $= \frac{1+2\sqrt{x}}{1-x}$
2. Here, $T_n = 3n - 1$, putting $n = 1, 2, 3, 4, 5$ we get first five terms, 2, 5, 8, 11, 14
 Hence, sum is $2 + 5 + 8 + 11 + 14 = 40$.
3. Given series $3.8 + 6.11 + 9.14 + 12.17 + \dots$
 First factors are 3, 6, 9, 12 whose n^{th} term is $3n$ and second factors are 8, 11, 14, 17
 $t_n = [8 + (n-1)3] = (3n+5)$
 Hence n^{th} term of given series $= 3n(3n+5)$.

4. Required number n is the number of terms in the series $105 + 112 + 119 + \dots + 994$
 $\therefore 994 = n^{\text{th}}$ term of the above A.P.
 $\Rightarrow 994 = 105 + (n-1) \times 7$
 $\Rightarrow n = \frac{994-98}{7}$
 $\Rightarrow n = 128$
5. Given sequence is in A.P.
 $\therefore a = 8 - 6i, d = -1 + 2i$
 $\therefore t_n = a + (n-1)d = (9-n) + i(2n-8)$
 For purely imaginary term, $9-n=0$
 $\Rightarrow n = 9$
6. First term $= a, d = b - a$ and last term $= c$
 If the no. of terms is n , then
 $t_n = c = a + (n-1)(b-a) \Rightarrow \frac{c-a}{b-a} = n-1$
 Solving, $n = \frac{b+c-2a}{b-a}$
7. $d = b - a$ and if the number of terms is n , then
 $2a = a + (n-1)(b-a)$
 $\Rightarrow \frac{a}{b-a} + 1 = n \Rightarrow n = \frac{b}{b-a}$
8. a, b, c are in A.P.
 $\Rightarrow b - a = c - b \Rightarrow \frac{b-a}{c-b} = 1$
9. If D is the common difference of the A.P. a, b, c, d, e , then $b = a + D, c = a + 2D, d = a + 3D, e = a + 4D$
 $\therefore a - 4b + 6c - 4d + e$
 $= a - 4(a+D) + 6(a+2D) - 4(a+3D) + a + 4D = 0$
10. Suppose that $\angle A = x^\circ$, then $\angle B = x + 10^\circ$,
 $\angle C = x + 20^\circ$ and $\angle D = x + 30^\circ$
 So, we know that $\angle A + \angle B + \angle C + \angle D = 2\pi$
 Putting these values, we get
 $(x^\circ) + (x^\circ + 10^\circ) + (x^\circ + 20^\circ) + (x^\circ + 30^\circ) = 360^\circ$
 $\Rightarrow x = 75^\circ$
 Hence, the angles of the quadrilateral are $75^\circ, 85^\circ, 95^\circ, 105^\circ$.
11. As we know $T_n = S_n - S_{n-1}$
 $= (2n^2 + 5n) - \{2(n-1)^2 + 5(n-1)\}$
 $= 2n^2 + 5n - 2n^2 + 4n - 2 - 5n + 5$
 $= 4n + 3$



12. $t_n = S_n - S_{n-1}$
 $= \left\{ nP + \frac{n(n-1)}{2}Q \right\}$
 $\quad - \left\{ (n-1)P + \frac{(n-1)(n-2)}{2}Q \right\}$
 $= P + (n-1)Q$
 \therefore Common difference $= t_n - t_{n-1}$
 $= [P + (n-1)Q] - [P + (n-2)Q] = Q$
13. $d = \frac{1}{3} - \frac{1}{2} = \frac{-1}{6}$
 $\therefore S_9 = \frac{9}{2} \left\{ 2 \times \frac{1}{2} + (9-1) \left(\frac{-1}{6} \right) \right\} = -\frac{3}{2}$
14. Required sum $= 10 + 13 + 16 + \dots + 97$
 $= \frac{n}{2} (10 + 97) \dots (i)$
 Here, $97 = 10 + (n-1)3 \Rightarrow n = 30$
 \therefore From (i), $S_n = \frac{30}{2} (10 + 97) = 1605$
15. The smallest 3 digit no. divisible by 7 is 105 and greatest is 994.
 Given sequence is in A.P. with $d = 7$
 $\therefore 994 = 105 + (n-1)7 \Rightarrow n = 128$
 $\therefore S_n = \frac{n}{2} [2a + (n-1)d]$
 $= \frac{128}{2} [2(105) + (128-1)7] = 70336$
16. According to the given condition
 $\frac{15}{2} [10 + 14 \times d] = 390 \Rightarrow d = 3$
 Hence, middle term i.e., 8th term is given by
 $5 + 7 \times 3 = 26$
17. $= a + (n-1)d$ and
 $S_n = \frac{n}{2} (a +)$
 Eliminating a, we get
 $S_n = \frac{n}{2} \{ -(n-1)d + \} = \frac{n}{2} \{ 2 - (n-1)d \}$
18. Suppose work is completed in n days
 $\frac{n}{2} [2 \times 150 + (n-1)(-4)] = n(152 - 2n)$
 Had no worker dropped from work, total no. of workers who would have worked all the n days is $150(n-8)$
 $\therefore n(152 - 2n) = 150(n-8) \Rightarrow n = 25$

19. $d = -2$, sum $= -5$
 $\therefore -5 = \frac{5}{2} \{ 2a + 4(-2) \} \Rightarrow a = 3$
 Hence, the actual sum (when $d = 2$) is
 $\frac{5}{2} \{ 2 \times 3 + (5-1) \times 2 \} = \frac{5}{2} (6+8) = 35$
21. Here $a = S_1 = 6$
 $S_7 = 105 \Rightarrow \frac{7}{2} [2 \times 6 + (7-1)d] = 105 \Rightarrow d = 3$
 $\therefore \frac{S_n}{S_{n-3}} = \frac{\frac{n}{2} \{ 2 \times 6 + (n-1)3 \}}{\frac{(n-3)}{2} \{ 2 \times 6 + (n-4)3 \}} = \frac{n+3}{n-3}$
22. $S_{2n} = 3S_n$
 $\therefore \frac{2n}{2} [2a + (2n-1)d] = \frac{3n}{2} [2a + (n-1)d]$
 $\Rightarrow 2a = (n+1)d$
 $\therefore \frac{S_{3n}}{S_n} = \frac{\frac{3n}{2} [2a + (3n-1)d]}{\frac{n}{2} [2a + (n-1)d]} = 6$
23. Let S_n and S'_n be the sum of n terms of two A.P.'s and t_{11} and t'_{11} be the respective 11th terms, then
 $\frac{S_n}{S'_n} = \frac{\frac{n}{2} [2a + (n-1)d]}{\frac{n}{2} [2a' + (n-1)d']} = \frac{7n+1}{4n+27}$
 $\Rightarrow \frac{a + \frac{(n-1)}{2}d}{a' + \frac{(n-1)}{2}d'} = \frac{7n+1}{4n+27}$
 Now put $n = 21$,
 we get $\frac{a+10d}{a'+10d'} = \frac{t_{11}}{t'_{11}} = \frac{148}{111} = \frac{4}{3}$
24. a, b, c, are in A.P. $\Rightarrow 2b = a + c$
 Also, $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}$ are in A.P. $\Rightarrow \frac{2}{b} = \frac{1}{a} + \frac{1}{c}$
 $\therefore \frac{2}{a+c} = \frac{a+c}{ac} \Rightarrow a = c$ and $b = a$
25. $(a+2b-c)(2b+c-a)(c+a-b)$
 $= (a+a+c-c)(a+c+c-a)(2b-b)$
 $= 4abc$
 $(\because a, b, c \text{ are in A.P.}, \therefore 2b = a+c).$



26. The sum of n arithmetic mean between a and b

$$= \frac{n}{2}(a + b)$$

$$27. \frac{a^{n+1} + b^{n+1}}{a^n + b^n} = \frac{a + b}{2}$$

$$\Rightarrow a^{n+1} - ab^n + b^{n+1} - ba^n = 0$$

$$\Rightarrow (a - b)(a^n - b^n) = 0$$

$$\text{If } a^n - b^n = 0. \text{ Then } \left(\frac{a}{b}\right)^n = 1 = \left(\frac{a}{b}\right)^0$$

Hence, $n = 0$

28. The resulting progression will have $n + 2$ terms with 2 as the first term and 38 as the last term.

Therefore, the sum of the progression

$$= \frac{n+2}{2}(2+38)$$

$$= 20(n+2)$$

$$\text{By hypothesis, } 20(n+2) = 200$$

$$\Rightarrow n = 8$$

29. As, $\log 2$, $\log(2^n - 1)$ and $\log(2^n + 3)$ are in A.P.

Therefore,

$$2 \log(2^n - 1) = \log 2 + \log(2^n + 3)$$

$$2^{2n} - 4 \cdot 2^n - 5 = 0$$

$$\Rightarrow (2^n - 5)(2^n + 1) = 0$$

$$\text{As } 2^n \text{ cannot be negative, hence } 2^n - 5 = 0$$

$$\Rightarrow 2^n = 5 \text{ or } n = \log_2 5$$

30. The given numbers are in A.P.

$$\therefore 2 \log_9 (3^{1-x} + 2) = \log_3 (4 \cdot 3^x - 1) + 1$$

$$\Rightarrow 2 \log_{3^2} (3^{1-x} + 2) = \log_3 (4 \cdot 3^x - 1) + \log_3 3$$

$$\Rightarrow \frac{2}{2} \log_3 (3^{1-x} + 2) = \log_3 [3(4 \cdot 3^x - 1)]$$

$$\Rightarrow 3^{1-x} + 2 = 3(4 \cdot 3^x - 1)$$

$$\Rightarrow \frac{3}{y} + 2 = 12y - 3, \text{ where } y = 3^x$$

$$\Rightarrow 12y^2 - 5y - 3 = 0$$

$$\therefore y = \frac{-1}{3} \text{ or } \frac{3}{4} \Rightarrow 3^x = \frac{-1}{3} \text{ or } 3^x = \frac{3}{4}$$

$$\therefore x = \log_3 \left(\frac{3}{4}\right) \Rightarrow x = 1 - \log_3 4$$

31. Let the three numbers be $a - d$, a , $a + d$

$$\text{We get } a - d + a + a + d = 15$$

$$\Rightarrow a = 5$$

$$\text{and } (a - d)^2 + a^2 + (a + d)^2 = 83$$

$$\Rightarrow a^2 + d^2 - 2ad + a^2 + a^2 + d^2 + 2ad = 83$$

$$\Rightarrow 2(a^2 + d^2) + a^2 = 83$$

Putting $a = 5$

$$\Rightarrow 2(25 + d^2) + 25 = 83$$

$$\Rightarrow 2d^2 = 8$$

$$\Rightarrow d = 2$$

Thus, numbers are 3, 5, 7.

Trick :

$$\text{Since } 3 + 5 + 7 = 15 \text{ and } 3^2 + 5^2 + 7^2 = 83$$

$$32. \left(1 - \frac{1}{2}\right) + \left(1 - \frac{1}{4}\right) + \left(1 - \frac{1}{8}\right) + \dots$$

$$\therefore t_n = 1 - \left(n^{\text{th}} \text{ term of G.P. } \frac{1}{2}, \frac{1}{4}, \frac{1}{8}\right)$$

$$= 1 - \frac{1}{2} \left(\frac{1}{2}\right)^{n-1}$$

$$= 1 - \frac{1}{2^n}$$

$$33. t_3 = 4 \Rightarrow ar^2 = 4$$

$$\therefore a \times ar \times ar^2 \times ar^3 \times ar^4 = (ar^2)^5 = 4^5$$

$$34. t_3 = ar^{3-1} = ar^2 = 36 \text{ and } t_6 = ar^{6-1} = ar^5 = 972$$

Solving, $a = 4$ and $r = 3$

$$\therefore t_8 = ar^7 = 4(3)^7 = 8748$$

$$35. t_n = ar^{n-1} \text{ and } r = 2$$

$$\therefore t_n = a(2)^{n-1} \Rightarrow t_9 = a(2)^8$$

$$\therefore a(2)^8 = 128 \Rightarrow a = \frac{128}{256} = \frac{1}{2}$$

$$36. ab^2 = a(ac) \text{ and } cb^2 = c(ac)$$

$$\therefore ab^2 - cb^2 = a^2c - ac^2$$

$$\Rightarrow a(b^2 + c^2) = c(a^2 + b^2)$$

$$37. a + ar = -4 \text{ and } ar^4 = 4ar^2 \Rightarrow r^2 = 4 \Rightarrow r = \pm 2$$

Substituting $r = \pm 2$, we get $a = \frac{-4}{3}$ and $a = 4$

$$\therefore \text{Required G.P. is } \frac{-4}{3}, \frac{-8}{3}, \frac{-16}{3}, \dots$$

or $4, -8, 16, -32, \dots$

$$38. \text{The common ratio of the G.P. is } x^{n+4}$$

$$\therefore 8^{\text{th}} \text{ term} = x^{52} = x^{-4} (x^{n+4})^7$$

$$\Rightarrow 7n = 28$$

$$\Rightarrow n = 4$$

$$39. \text{Let } AR^{p-1} = a,$$

$$AR^{q-1} = b,$$

$$AR^{r-1} = c$$

So

$$a^{q-r} b^{r-p} c^{p-q} = (AR^{p-1})^{q-r} (AR^{q-1})^{r-p} (AR^{r-1})^{p-q}$$

$$= A^{(q-r+r-p+p-q)} R^{(pq-pr-qr+qr-pq-r+p+pr-rq-p+q)}$$

$$= A^0 R^0 = 1$$



$$40. \quad a = \frac{5}{2}, r = \frac{1}{2} < 1$$

$$\therefore S_n = \frac{a(1-r^n)}{1-r} = 5 \left[\frac{2^n - 1}{2^n} \right]$$

$$41. \quad \text{Given series is a G.P. with } a = \sqrt{2} \text{ and } r = \sqrt{3}$$

$$\begin{aligned} \therefore S_{10} &= \frac{\sqrt{2} \left((\sqrt{3})^{10} - 1 \right)}{\sqrt{3} - 1} = \frac{\sqrt{2}(243-1)}{\sqrt{3}-1} \times \frac{\sqrt{3}+1}{\sqrt{3}+1} \\ &= 121\sqrt{6} + 121\sqrt{2} \\ &= 121(\sqrt{6} + \sqrt{2}) \end{aligned}$$

$$42. \quad \left(1 - \frac{1}{2}\right) + \left(1 - \frac{1}{4}\right) + \left(1 - \frac{1}{8}\right) + \dots$$

$$\begin{aligned} \therefore S_n &= n - \left(\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots \text{upto } n \text{ terms} \right) \\ &= n - \frac{1 \left(1 - \left(\frac{1}{2} \right)^n \right)}{1 - \frac{1}{2}} = n - \left(1 - \frac{1}{2^n} \right) \\ &= n - 1 + 2^{-n} \end{aligned}$$

$$43. \quad t_4 = 24 \text{ and } t_9 = 768$$

$$\therefore t_4 = ar^3 \Rightarrow ar^3 = 24$$

$$\text{and } t_9 = ar^8 \Rightarrow ar^8 = 768$$

$$\text{Solving, } a = 3 \text{ and } r = 2 > 1$$

$$\therefore S_{10} = \frac{a[r^{10} - 1]}{r - 1} = \frac{3[2^{10} - 1]}{2 - 1} = 3(2^{10} - 1)$$

$$44. \quad S_{10} = 244 \quad S_5$$

$$\Rightarrow (1 - r^{10}) = 244(1 - r^5)$$

$$\Rightarrow r^{10} - 244r^5 + 243 = 0$$

$$\Rightarrow r^5 = 243 \text{ or } r^5 = 1$$

$$\Rightarrow r = 3 \text{ or } r = 1$$

$$45. \quad a_1 = 3, a_n = 96$$

$$\Rightarrow a_1 r^{n-1} = 96$$

$$\Rightarrow r^{n-1} = 32$$

$$r^{n-1} = 2^5$$

$$2^{n-1} = 2^5$$

$$\Rightarrow n - 1 = 5$$

$$\Rightarrow n = 6$$

$$46. \quad a = 7 \text{ and } ar^{n-1} = 448$$

$$\text{Now, } S_n = \frac{a(r^n - 1)}{r - 1} = 889$$

$$\Rightarrow \frac{(ar^{n-1} \cdot r - a)}{r - 1} = 889 \Rightarrow \frac{448r - 7}{r - 1} = 889$$

$$\Rightarrow r = 2$$

$$47. \quad \text{Given that } \frac{a(r^n - 1)}{r - 1} = 255 \quad (\because r > 1) \quad \dots(i)$$

$$ar^{n-1} = 128 \quad \dots(ii)$$

$$\text{and common ratio } r = 2 \quad \dots(iii)$$

From (i), (ii) and (iii), we get

$$a(2)^{n-1} = 128 \quad \dots(iv)$$

$$\text{and } \frac{a(2^n - 1)}{2 - 1} = 255 \quad \dots(v)$$

Dividing (v) by (iv), we get

$$\frac{2^n - 1}{2^{n-1}} = \frac{255}{128}$$

$$\Rightarrow 2 - 2^{-n+1} = \frac{255}{128}$$

$$\Rightarrow 2(1 - 2^{-n}) = \frac{255}{128}$$

$$\Rightarrow 2^{-n} = 2^{-8}$$

$$\Rightarrow n = 8$$

Putting $n = 8$ in equation (iv), we get $a \cdot 2^7 = 128 = 2^7$ or $a = 1$

$$48. \quad \frac{S_3}{S_6 - S_3} = \frac{125}{27} \Rightarrow \frac{S_3}{S_6} = \frac{125}{152}$$

$$\therefore \frac{a(1-r^3)}{a(1-r^6)} = \frac{125}{152} \Rightarrow \frac{1}{1+r^3} = \frac{125}{152}$$

$$\Rightarrow r^3 = \frac{27}{125} \Rightarrow r = \frac{3}{5}$$

$$49. \quad r = \frac{t_2}{t_1} = \frac{b}{a}; \text{ last term} = c$$

$$\Rightarrow ar^{n-1} = c$$

$$\Rightarrow \frac{ar^n}{r} = c$$

$$\Rightarrow ar^n = cr$$

$$\therefore S_n = \frac{a(1-r^n)}{1-r} = \frac{a - ar^n}{1-r} = \frac{a - cr}{1-r} = \frac{a - c \left(\frac{b}{a} \right)}{1 - \frac{b}{a}}$$

$$50. \quad \text{We have}$$

$$1 + a + a^2 + \dots + a^x = (1+a)(1+a^2)(1+a^4)$$

$$\Rightarrow \frac{(1 - a^{x+1})}{(1-a)} = (1+a)(1+a^2) + (1+a^4)$$

$$\Rightarrow (1 - a^{x+1}) = (1-a)(1+a)(1+a^2)(1+a^4)$$

$$\Rightarrow (1 - a^{x+1}) = (1 - a^8)$$

$$\Rightarrow x + 1 = 8$$

$$\Rightarrow x = 7$$



$$\begin{aligned}
 51. \quad S_n &= 4 + 44 + 444 + \dots \text{ to } n \text{ terms} \\
 &= \frac{4}{9} [(10-1) + (10^2-1) + (10^3-1) + \dots + (10^n-1)] \\
 &= \frac{4}{9} \{(10+10^2+10^3+\dots+10^n) - (1+1+1+\dots+n \text{ times})\} \\
 &= \frac{4}{9} \left\{ \frac{10(10^n-1)}{10-1} - n \right\} = \frac{4}{9} \left\{ \frac{10}{9}(10^n-1) - n \right\}
 \end{aligned}$$

$$\begin{aligned}
 52. \quad &1 + (1+x) + (1+x+x^2) + \dots \\
 &\quad \quad \quad + (1+x+x^2+\dots+x^{n-1}) \\
 &= \frac{1-x}{1-x} + \frac{1-x^2}{1-x} + \frac{1-x^3}{1-x} + \dots + \frac{1-x^n}{1-x} \\
 &= \frac{1}{1-x} [(1+1+\dots+n \text{ times}) \\
 &\quad \quad \quad - (x+x^2+\dots+x^n)] \\
 &= \frac{1}{1-x} \left[n - \frac{x(1-x^n)}{1-x} \right] \\
 &= \frac{n}{1-x} - \frac{x(1-x^n)}{(1-x)^2}
 \end{aligned}$$

$$\begin{aligned}
 53. \quad &\text{Infinite series } 9 - 3 + 1 - \frac{1}{3} + \dots \dots \infty \text{ is a} \\
 &\text{G.P. with } a = 9, r = \frac{-1}{3}
 \end{aligned}$$

$$\therefore S_\infty = \frac{a}{1-r} = \frac{9}{1+\left(\frac{1}{3}\right)} = \frac{9 \times 3}{4} = \frac{27}{4}$$

$$\begin{aligned}
 54. \quad &\text{Let the G.P. be } a + ar + ar^2 + \dots, |r| < 1, \\
 &\text{then } ar = 2 \text{ and } \frac{a}{1-r} = 8
 \end{aligned}$$

$$\therefore \frac{ar(1-r)}{a} = \frac{2}{8} \Rightarrow r = \frac{1}{2} \text{ and } a = 4$$

$$\begin{aligned}
 55. \quad &\frac{1}{7} + \frac{2}{7^2} + \frac{1}{7^3} + \frac{2}{7^4} + \dots \text{ upto } \infty \\
 &= \left(\frac{1}{7} + \frac{1}{7^3} + \frac{1}{7^5} + \dots \right) + 2 \left(\frac{1}{7^2} + \frac{1}{7^4} + \frac{1}{7^6} + \dots \right) \\
 &= \frac{\frac{1}{7}}{1-\frac{1}{7^2}} + \frac{2\left(\frac{1}{7^2}\right)}{1-\frac{1}{7^2}} \\
 &= \frac{3}{16}
 \end{aligned}$$

$$\begin{aligned}
 56. \quad 2.\overline{345} &= 2.3 + 0.045 + 0.00045 + \dots \\
 &= \frac{23}{10} + \frac{45}{1000} + \frac{45}{100000} + \dots
 \end{aligned}$$

From 2nd term onwards, the terms are in G.P.

$$\therefore s_\infty = \frac{a}{1-r} = \frac{\frac{45}{1000}}{1-\frac{1}{100}} = \frac{1}{22}$$

$$\therefore 2.\overline{345} = \frac{23}{10} + \frac{1}{22} = \frac{129}{55}$$

Alternate Method:

$$2.\overline{345} = 2 + \frac{345-3}{990} = 2 + \frac{342}{990} = \frac{129}{55}$$

$$\begin{aligned}
 57. \quad &4^{1/3} \cdot 4^{1/9} \cdot 4^{1/27} \dots \infty \\
 \therefore S &= 4^{1/3+1/9+1/27+\dots \infty} \\
 \Rightarrow S &= 4^{\left(\frac{1/3}{1-1/3}\right)} = 4^{\frac{1/3}{2/3}} \\
 \Rightarrow S &= 4^{1/2} \\
 \Rightarrow S &= 2
 \end{aligned}$$

$$58. \quad 5 = \frac{x}{1-r} \Rightarrow 5 - 5r = x \Rightarrow r = 1 - \frac{x}{5}$$

$$\text{As } |r| < 1 \text{ i.e., } \left| 1 - \frac{x}{5} \right| < 1$$

$$\therefore -1 < 1 - \frac{x}{5} < 1$$

$$\therefore -5 < 5 - x < 5$$

$$\therefore -10 < -x < 0$$

$$\therefore 10 > x > 0$$

$$\therefore 0 < x < 10$$

$$59. \quad A = 1 + r^z + r^{2z} + r^{3z} + \dots \infty$$

$$A = 1 + [r^z + r^{2z} + r^{3z} + \dots \infty]$$

We know that sum of infinite G.P. is

$$S_\infty = \frac{a}{1-r} \quad (-1 < r < 1)$$

$$\text{Therefore, } A = 1 + \left[\frac{r^z}{1-r^z} \right]$$

$$\Rightarrow A = \frac{1-r^z+r^z}{1-r^z} \Rightarrow A = \frac{1}{1-r^z}$$

$$\Rightarrow 1-r^z = \frac{1}{A} \Rightarrow r^z = \frac{A-1}{A}$$

$$\text{Hence, } r = \left[\frac{A-1}{A} \right]^{\frac{1}{z}}$$



60. We have, $x = \sum_{n=0}^{\infty} a^n = \frac{1}{1-a} \Rightarrow a = \frac{x-1}{x}$

$$y = \sum_{n=0}^{\infty} b^n = \frac{1}{1-b} \Rightarrow b = \frac{y-1}{y}$$

$$z = \sum_{n=0}^{\infty} a^n b^n = \frac{1}{1-ab} \Rightarrow ab = \frac{z-1}{z}$$

$$\therefore \frac{x-1}{x} \cdot \frac{y-1}{y} = \frac{z-1}{z}$$

$$\Rightarrow xy + z = zx + yz$$

61. $a = 3, r = 3$

$$\text{G.M.} = (3 \cdot 3^2 \cdot 3^3 \cdots 3^n)^{1/n}$$

$$= (3^{1+2+3+\cdots+n})^{1/n} = \left(3^{\frac{n(n+1)}{2}}\right)^{1/n} = 3^{\frac{(n+1)}{2}}$$

62. As given, $G = \sqrt{xy}$

$$\therefore \frac{1}{G^2 - x^2} + \frac{1}{G^2 - y^2} = \frac{1}{xy - x^2} + \frac{1}{xy - y^2}$$

$$= \frac{1}{x-y} \left\{ -\frac{1}{x} + \frac{1}{y} \right\} = \frac{1}{xy} = \frac{1}{G^2}$$

63. a, g_1, g_2, b are in G.P. $\Rightarrow \frac{g_1}{a} = \frac{g_2}{g_1} = \frac{b}{g_2}$

$$\therefore \frac{g_1}{a} = \frac{g_2}{g_1} \text{ and } \frac{g_2}{g_1} = \frac{b}{g_2} \Rightarrow a = \frac{g_1^2}{g_2} \text{ and } b = \frac{g_2^2}{g_1}$$

$$\therefore \frac{g_1^2}{g_2} + \frac{g_2^2}{g_1} = a + b$$

64. $\text{G.M.} = b = \sqrt{ac}$

$$\frac{1}{b-a} + \frac{1}{b-c} = \frac{1}{\sqrt{ac}-a} + \frac{1}{\sqrt{ac}-c}$$

$$= \frac{1}{\sqrt{a}[\sqrt{c}-\sqrt{a}]} + \frac{1}{\sqrt{c}[\sqrt{a}-\sqrt{c}]}$$

$$= \frac{1}{\sqrt{a}[\sqrt{c}-\sqrt{a}]} - \frac{1}{\sqrt{c}[\sqrt{c}-\sqrt{a}]}$$

$$= \frac{1}{\sqrt{ac}} = \frac{1}{b}$$

65. Let the 9 terms of a G.P. be

$$\frac{a}{r^4}, \frac{a}{r^3}, \frac{a}{r^2}, \frac{a}{r}, a, ar, ar^2, ar^3, ar^4$$

Given, fifth term $a = 2$

Hence, product of 9 terms is $a^9 = (2)^9 = 512$

66. Let the terms of given G.P. be $\frac{a}{r}, a, ar$

then product = $\frac{a}{r} \times a \times ar = 1000$

$$\frac{a}{r}, a+6, ar+7 \text{ are in A.P.}$$

$$\therefore 2(a+6) = \frac{a}{r} + ar + 7 \quad \therefore 25 = \frac{10}{r} + 10r$$

$$\therefore 2r^2 - 5r + 2 = 0$$

$$\therefore (2r-1)(r-2) = 0$$

$$\therefore r = 2, \frac{1}{2}$$

Hence, the G.P. is 5, 10, 20, ... or 20, 10, 5, ...

67. Suppose that x to be added then numbers 13, 15, 19 so that new numbers $x+13, 15+x, 19+x$ will be in H.P.

$$\Rightarrow (15+x) = \frac{2(x+13)(19+x)}{x+13+x+19}$$

$$\Rightarrow x^2 + 31x + 240 = x^2 + 32x + 247 \Rightarrow x = -7$$

68. a, b, c are in H.P. $\Rightarrow \frac{1}{a}, \frac{1}{b}, \frac{1}{c}$ are in A.P.

$$\therefore \frac{1}{b} - \frac{1}{a} = \frac{1}{c} - \frac{1}{b} \Rightarrow \frac{a-b}{b-c} = \frac{a}{c}$$

69. 7th term of corresponding A.P. is $\frac{1}{8}$ and 8th

term will be $\frac{1}{7}$

$$\Rightarrow a + 6d = \frac{1}{8} \text{ and } a + 7d = \frac{1}{7}$$

Solving these, we get $d = \frac{1}{56}$ and $a = \frac{1}{56}$

Therefore, 15th term of this A.P.

$$= \frac{1}{56} + 14 \times \frac{1}{56} = \frac{15}{56}$$

Hence, the required 15th term of the H.P. is $\frac{56}{15}$.

70. Let a be the first term and d be the common difference of the corresponding A.P.

$$p^{\text{th}} \text{ term of A.P. } (T_p) = a + (p-1)d$$

$$= \frac{1}{q} \quad \dots \text{(i)}$$

$$q^{\text{th}} \text{ term of A.P. } (T_q) = a + (q-1)d$$

$$= \frac{1}{p} \quad \dots \text{(ii)}$$



$$\text{From (i) - (ii), } (p - q)d = \frac{1}{q} - \frac{1}{p} = \frac{p - q}{pq}$$

$$\Rightarrow d = \frac{1}{pq}$$

From (i),

$$a + (p - 1)\frac{1}{pq} = \frac{1}{q} \Rightarrow a = \frac{1}{pq}$$

$$\therefore T_{pq} = a + (pq - 1)d$$

$$= \frac{1}{pq} + (pq - 1)\frac{1}{pq} = 1$$

Therefore, pq^{th} term is 1.

$$71. \text{ H.M.} = \frac{2\left(\frac{a^2}{1 - a^2b^2}\right)}{\frac{a}{1 - ab} + \frac{a}{1 + ab}} = \frac{2a^2}{2a} = a$$

$$72. c = \frac{2ab}{a + b} \Rightarrow \frac{c}{a} = \frac{2b}{a + b} \text{ and } \frac{c}{b} = \frac{2a}{a + b}$$

$$\therefore \frac{c}{a} + \frac{c}{b} = \frac{2b}{a + b} + \frac{2a}{a + b} = 2$$

$$73. H = \frac{2ab}{a + b}$$

$$\Rightarrow H - a = \frac{2ab}{a + b} - a = \frac{ab - a^2}{a + b}$$

$$\text{and } H - b = \frac{2ab}{a + b} - b = \frac{ab - b^2}{a + b}$$

$$\begin{aligned} \therefore \frac{1}{H - a} + \frac{1}{H - b} &= \frac{a + b}{ab - a^2} + \frac{a + b}{ab - b^2} \\ &= \frac{(a + b)}{(b - a)} \left[\frac{(b - a)}{ab} \right] \\ &= \frac{1}{a} + \frac{1}{b} \end{aligned}$$

$$74. H = \frac{2ab}{a + b} \Rightarrow \frac{H}{a} = \frac{2b}{a + b}$$

$$\therefore \frac{H + a}{H - a} = \frac{3b + a}{b - a}$$

$$\text{Similarly, } \frac{H + b}{H - b} = \frac{3a + b}{a - b} = -\frac{3a + b}{b - a}$$

$$\therefore \frac{H + a}{H - a} + \frac{H + b}{H - b} = \frac{2b - 2a}{b - a} = 2$$

75. a, b, c are in H.P.

$$\therefore b = \frac{2ac}{a + c}$$

$$\text{Also } b, c, d \text{ are in H.P. } \Rightarrow c = \frac{2bd}{b + d}$$

$$\text{Multiplying we get, } bc = \frac{4abcd}{(a + c)(b + d)}$$

$$\therefore ab + bc + cd + ad = 4ad$$

$$\Rightarrow ab + bc + cd = 3ad$$

76. We know that $A > G > H$

Where A is arithmetic mean, G is geometric mean and H is harmonic mean, then $A > G$

$$\Rightarrow \frac{a + b}{2} > \sqrt{ab} \text{ or } (a + b) > 2\sqrt{ab}$$

77. Let the numbers be a and b , then

$$4 = \frac{2ab}{a + b} \Rightarrow a + b = \frac{ab}{2}$$

$$A = \frac{a + b}{2} \text{ and } G = \sqrt{ab}$$

$$\text{Also, } 2A + G^2 = 27$$

$$\therefore a + b + ab = 27 \Rightarrow \frac{ab}{2} + ab = 27 \Rightarrow ab = 18$$

$$\text{and hence } a + b = 9.$$

Only option A satisfies this condition.

78. As given, $2b = a + c \Rightarrow 3^{2b} = 3^{a+c}$
or $(3^b)^2 = 3^a \cdot 3^c$ i.e. $3^a, 3^b, 3^c$ are in G.P.

$$79. \frac{T_2}{T_1} = \frac{T_3}{T_2}$$

$$\Rightarrow 2^{(b-a)x} = 2^{(c-b)x}$$

$$\Rightarrow (b - a)x = (c - b)x$$

$$\Rightarrow (b - a) = (c - b) \forall x, x \neq 0$$

$$\therefore 2^{ax+1}, 2^{bx+1}, 2^{cx+1} \text{ is a G.P., } \forall x \neq 0$$

80. a, b, c are in A.P. $\Rightarrow 2b = a + c$

Now,

$$(10^{bx+10})^2 = (10^{ax+10} \cdot 10^{cx+10})$$

$$\Rightarrow 10^{2(bx+10)} = 10^{ax+cx+20}$$

$$\Rightarrow 2(bx + 10) = ax + cx + 20, \forall x$$

$$\Rightarrow 2b = a + c \text{ i.e. } a, b, c \text{ are in A.P.}$$

Hence, these are in G.P. $\forall x$

Alternate Method :

As we know if a, b, c are in A.P., then $x^{an+r}, x^{bn+r}, x^{cn+r}$ are in G.P. for every n and r .



81. $\because a, b, c$ are in G.P. $\Rightarrow b^2 = ac \dots (i)$

Let $a^x = b^y = c^z = k$

$\Rightarrow a = k^{1/x}, b = k^{1/y}, c = k^{1/z}$

Putting these values in (i),

$$k^{2/y} = k^{1/x} \cdot k^{1/z} = k^{\frac{1}{x} + \frac{1}{z}}, \quad \frac{2}{y} = \frac{1}{x} + \frac{1}{z}$$

$\therefore \frac{1}{x}, \frac{1}{y}, \frac{1}{z}$ are in A.P. or x, y, z are in H.P.

82. Here, $\frac{\log x}{\log a}, \frac{\log x}{\log b}, \frac{\log x}{\log c}$ are in H.P.

$\Rightarrow \frac{\log a}{\log x}, \frac{\log b}{\log x}, \frac{\log c}{\log x}$ are in A.P.

$\Rightarrow \log_x a, \log_x b, \log_x c$ are in A.P.

$\Rightarrow a, b, c$ are in G.P.

83. Clearly, $x = \frac{1}{1-a}, y = \frac{1}{1-b}, z = \frac{1}{1-c}$

Since a, b, c are in A.P.

$\Rightarrow 1-a, 1-b, 1-c$ are also in A.P.

$\Rightarrow \frac{1}{1-a}, \frac{1}{1-b}, \frac{1}{1-c}$ are in H.P.

$\therefore x, y, z$ are in H.P.

84. Given that $\frac{a}{b} = \frac{9}{1}$ or $a = 9b$

Here, $H = \frac{2ab}{a+b}$ and $G = \sqrt{ab}$

$\Rightarrow H : G = \frac{2ab}{a+b} : \sqrt{ab} = \frac{2.9b^2}{10b} : 3b = \frac{3}{5}$

Hence, $G : H = 5 : 3$

85. Given that $\frac{H.M.}{G.M.} = \frac{12}{13}$

$\Rightarrow \frac{2ab}{a+b} = \frac{12}{13}$ or $\frac{a+b}{2\sqrt{ab}} = \frac{13}{12}$

$\Rightarrow \frac{(a+b) + 2\sqrt{ab}}{(a+b) - 2\sqrt{ab}} = \frac{13+12}{13-12} = \frac{25}{1}$

$\Rightarrow \frac{\sqrt{a} + \sqrt{b}}{\sqrt{a} - \sqrt{b}} = \frac{5}{1}$

$\Rightarrow \frac{(\sqrt{a} + \sqrt{b}) + (\sqrt{a} - \sqrt{b})}{(\sqrt{a} + \sqrt{b}) - (\sqrt{a} - \sqrt{b})} = \frac{5+1}{5-1}$

$\Rightarrow \frac{2\sqrt{a}}{2\sqrt{b}} = \frac{6}{4}$

$\Rightarrow \left(\frac{a}{b}\right)^{1/2} = \frac{6}{4} \Rightarrow a : b = 9 : 4$

86. Let $S = 1 + 2.2 + 3.2^2 + 4.2^3 + \dots + 100.2^{99}$
 $2S = 1.2 + 2.2^2 + 3.2^3 + \dots + 100.2^{100}$
 $S - 2S = 1 + (1.2 + 1.2^2 + 1.2^3 + \dots \text{ upto 99 terms}) - 100.2^{100}$

$\therefore S = -1 - \frac{2(2^{99} - 1)}{2-1} + 100.2^{100}$
 $= -1 - 2^{100} + 2 + 100.2^{100}$
 $= 1 + 99 \times 2^{100}$

87. Given series, let

$S_n = 1 + \frac{2}{5} + \frac{3}{5^2} + \frac{4}{5^3} + \dots + \frac{n}{5^{n-1}}$

$\frac{1}{5}S_n = \frac{1}{5} + \frac{2}{5^2} + \frac{3}{5^3} + \dots + \frac{n}{5^n}$

Subtracting,

$\left(1 - \frac{1}{5}\right)S_n = 1 + \frac{1}{5} + \frac{1}{5^2} + \frac{1}{5^3}$

$+ \dots + \text{ upto } n \text{ terms} - \frac{n}{5^n}$

$\Rightarrow \frac{4}{5}S_n = \frac{1 - \frac{1}{5^n}}{\frac{4}{5}} - \frac{n}{5^n}$

$\Rightarrow S_n = \frac{25}{16} - \frac{4n+5}{16 \times 5^{n-1}}$

88. Let $S_n = 1 + 2x + 3x^2 + 4x^3 + \dots + nx^{n-1} \dots (i)$

$xS_n = x + 2x^2 + 3x^3 + \dots + nx^n \dots (ii)$

Subtracting (ii) from (i), we get

$(1-x)S_n = 1 + x + x^2 + x^3 + \dots \text{ to } n \text{ terms} - nx^n$

$= \frac{(1-x^n)}{1-x} - nx^n$

$\Rightarrow S_n = \frac{(1-x^n) - nx^n(1-x)}{(1-x)^2}$

$= \frac{1 - (n+1)x^n + nx^{n+1}}{(1-x)^2}$

89. Let $S = 2 + 4 + 7 + 11 + 16 + \dots + t_n$

$S = 2 + 4 + 7 + 11 + 16 + \dots + t_{n-1} + t_n$

Subtracting, we get

$0 = 2 + \{2 + 3 + 4 + \dots + (t_n - t_{n-1})\} - t_n$

$\Rightarrow t_n = 1 + \{1 + 2 + 3 + 4 + \dots \text{ upto } n \text{ terms}\}$

$\Rightarrow t_n = 1 + \frac{1}{2}n(n+1)$

$= \frac{2+n^2+n}{2} = \frac{n^2+n+2}{2}$



90. We have $\sqrt{2} + \sqrt{8} + \sqrt{18} + \sqrt{32} + \dots$
 $= 1\sqrt{2} + 2\sqrt{2} + 3\sqrt{2} + 4\sqrt{2} + \dots$
 $= \sqrt{2} [1 + 2 + 3 + 4 + \dots \text{ upto 24 terms}]$
 $= \sqrt{2} \times \frac{24 \times 25}{2}$
 $= 300\sqrt{2}$

91. $t_r = \frac{1+2+3+\dots+r}{r} = \frac{\frac{r(r+1)}{2}}{r} = \frac{r+1}{2}$
 $\therefore S_n = \sum_{r=1}^n t_r = \sum_{r=1}^n \frac{r+1}{2} = \frac{1}{2} \left[\sum_{r=1}^n r + \sum_{r=1}^n 1 \right]$
 $= \frac{1}{2} \left[\frac{n(n+1)}{2} + n \right]$
 $= \frac{1}{4} (n^2 + 3n) = \frac{n(n+3)}{4}$

93. $S_n = \sum_{r=1}^n (4r-3)(4r-1)$
 $= \sum_{r=1}^n (16r^2 - 16r + 3)$
 $= \frac{16n(n+1)(2n+1)}{6} - \frac{16n(n+1)}{2} + 3n$
 $= n \left(\frac{16n^2 - 7}{3} \right)$

94. $\sum n^2 = 330 + \sum n$
 $\Rightarrow \frac{n(n+1)(2n+1)}{6} = 330 + \frac{n(n+1)}{2}$
 $\Rightarrow \frac{n(n+1)}{2} \left[\frac{2n+1}{3} - 1 \right] = 330$
 $\Rightarrow \frac{n(n+1)}{2} \cdot \frac{2(n-1)}{3} = 330$
 $\Rightarrow n(n+1)(n-1) = 990 \Rightarrow n = 10$

95. $t_r = \frac{1^3 + 2^3 + \dots + r^3}{(r+1)^2} = \frac{r^2}{4}$
 $\therefore S_n = \sum_{r=1}^n t_r = \sum_{r=1}^n \frac{r^2}{4} = \frac{1}{4} \sum_{r=1}^n r^2$
 $= \frac{1}{4} \frac{n(n+1)(2n+1)}{6}$
 $= \frac{n(n+1)(2n+1)}{24}$

96. $1^3 + 3^3 + 5^3 + \dots + 21^3$
 $= (1^3 + 2^3 + 3^3 + 4^3 + \dots + 21^3)$
 $\quad - (2^3 + 4^3 + 6^3 + \dots + 20^3)$
 $= \sum_{r=1}^{21} r^3 - 8 \sum_{r=1}^{10} r^3$
 $= \frac{(21)^2 (21+1)^2}{4} - \frac{8 \times 10^2 (10+1)^2}{4}$
 $= 29161$

97. $\sum_{n=1}^{20} (n^3) - \sum_{n=1}^{10} (n^3) = \left[\frac{n(n+1)}{2} \right]_{n=20}^2 - \left[\frac{n(n+1)}{2} \right]_{n=10}^2$
 $\Rightarrow \left[\frac{20 \times 21}{2} \right]^2 - \left[\frac{10 \times 11}{2} \right]^2$
 $= 44100 - 3025$
 $= 41075$

98. Here $T_n = \frac{1^3 + 2^3 + 3^3 + \dots + n^3}{1 + 3 + 5 + \dots \text{ upto } n \text{ terms}}$
 $= \frac{\sum n^3}{\frac{n}{2} [2 + (n-1)2]}$
 $= \frac{1}{4} \frac{n^2(n+1)^2}{n^2}$
 $= \frac{1}{4} (n^2 + 2n + 1)$

99. Given fraction $\frac{\sum_{n=1}^{12} n^3}{\sum_{n=1}^{12} n^2} = \frac{\left\{ \frac{12(12+1)}{2} \right\}^2}{\frac{12(12+1)(2 \times 12+1)}{6}}$
 $= \frac{12 \times 13}{4} \times \frac{6}{25} = \frac{234}{25}$

100. $S_1 = \frac{n(n+1)}{2}, S_2 = \frac{n(n+1)(2n+1)}{6}$
 $S_3 = \left(n \left(\frac{n+1}{2} \right) \right)^2$
 For, $S_3(1 + 8S_1) = \frac{n^2(n+1)^2}{4} \left(1 + \frac{8n(n+1)}{2} \right)$
 $= \left(\frac{n(n+1)(2n+1)}{6} \right)^2 \times 9$
 $= 9S_2^2$



$$101. \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \frac{1}{5!} + \dots = 1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \dots$$

which is the expansion of e^{-1}

$$102. e^{-x} = (1-x) + \frac{x^2}{2!} \left(1 - \frac{x}{3}\right) + \frac{x^4}{4!} \left(1 - \frac{x}{5}\right) + \dots$$

$$\therefore e^{-1} = (1-1) + \frac{1}{2!} \left(1 - \frac{1}{3}\right) + \frac{1}{4!} \left(1 - \frac{1}{5}\right) + \dots$$

$$= \frac{2}{3!} + \frac{4}{5!} + \frac{6}{7!} + \dots$$

$$103. \text{ Let } t_n = \frac{1}{(n+1)!}$$

$$S_n = \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \frac{1}{5!} + \dots$$

$$= \left[1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \frac{1}{5!} + \dots\right] - \left[1 + \frac{1}{1!}\right]$$

$$= e - (1+1) = e - 2$$

$$104. \text{ Given ratio} = \frac{\frac{1}{2} \left(e + \frac{1}{e}\right) - 1}{\frac{1}{2} \left(e - \frac{1}{e}\right)} = \frac{(e-1)^2}{(e-1)(e+1)}$$

$$= \frac{e-1}{e+1}$$

$$105. \frac{1}{1 \times 2} + \frac{1}{3 \times 4} + \frac{1}{5 \times 6} + \dots \infty$$

$$= \left(1 - \frac{1}{2}\right) + \left(\frac{1}{3} - \frac{1}{4}\right) + \left(\frac{1}{5} - \frac{1}{6}\right) + \dots \text{ to } \infty$$

$$= \log 2$$

$$106. T_n = \frac{3^n - 1}{3^n} = 1 - \left(\frac{1}{3}\right)^n$$

$$S_n = n - \sum_{n=1}^n \left(\frac{1}{3}\right)^n = n - \frac{1 \left[1 - \left(\frac{1}{3}\right)^n\right]}{\left(1 - \frac{1}{3}\right)}$$

$$= n - \frac{1}{2} (1 - 3^{-n}) = n + \frac{1}{2} (3^{-n} - 1)$$

107. The series is

$$\frac{2}{1!} + \frac{(2+5)}{2!} + \frac{(2+5+8)}{3!} + \frac{(2+5+8+11)}{4!} + \dots$$

$$\text{Hence, } T_n = \frac{(2+5+8+\dots+n \text{ terms})}{n!}$$

$$= \frac{\frac{n}{2} [2.2 + (n-1)3]}{n!}$$

$$T_n = \frac{n(3n+1)}{2(n)!}$$

$$108. \text{ Let } S = i - 2 - 3i + 4 + 5i + \dots + 100i^{100}$$

$$\Rightarrow S = i + 2i^2 + 3i^3 + 4i^4 + 5i^5 + \dots + 100i^{100}$$

$$\Rightarrow iS = i^2 + 2i^3 + 3i^4 + 4i^5 + \dots + 99i^{100} + 100i^{101}$$

$$\therefore S - iS = [i + i^2 + i^3 + i^4 + \dots + i^{100}] - 100i^{101}$$

$$\Rightarrow S(1-i) = 0 - 100i^{101} = -100i$$

$$\therefore S = \frac{-100i}{1-i} = -50i(1+i) = -50(i-1)$$

$$= 50(1-i)$$

$$109. \text{ Here, } T_r = \frac{1}{r(r+1)}, r = 1, 2, \dots, n$$

$$\Rightarrow T_r = \frac{1}{r} - \frac{1}{r+1}$$

$$\therefore \text{ Required sum} = \sum_{r=1}^n T_r$$

$$= \left(1 - \frac{1}{2}\right) + \left(\frac{1}{2} - \frac{1}{3}\right) + \left(\frac{1}{3} - \frac{1}{4}\right) + \dots + \left(\frac{1}{n} - \frac{1}{n+1}\right)$$

$$= 1 - \frac{1}{n+1} = \frac{n}{n+1}$$

110. $\sin A$, $\cos A$ and $\tan A$ are in G.P.

$$\therefore \cos^2 A = \sin A \tan A = \frac{\sin^2 A}{\cos A}$$

$$\Rightarrow \cos^3 A = \sin^2 A$$

$$\Rightarrow \cos^3 A = 1 - \cos^2 A$$

$$\Rightarrow \cos^3 A + \cos^2 A = 1$$

111. $\cos^4 \theta \sec^2 \alpha$, $\frac{1}{2}$ and $\sin^4 \theta \operatorname{cosec}^2 \alpha$ are in A.P.

$$\therefore 1 = \cos^4 \theta \sec^2 \alpha + \sin^4 \theta \operatorname{cosec}^2 \alpha$$

$$\Rightarrow \cos^4 \theta \sin^2 \alpha + \sin^4 \theta \cos^2 \alpha = \sin^2 \alpha \cos^2 \alpha$$

$$\Rightarrow (1 - \sin^2 \theta) \cos^2 \theta \sin^2 \alpha + \sin^4 \theta (1 - \sin^2 \alpha)$$

$$= \sin^2 \alpha (1 - \sin^2 \alpha)$$

$$\Rightarrow \cos^2 \theta \sin^2 \alpha + \sin^4 \theta - \sin^2 \theta \sin^2 \alpha (\cos^2 \theta + \sin^2 \theta)$$

$$= \sin^2 \alpha - \sin^4 \alpha$$

$$\Rightarrow \sin^4 \theta + \sin^4 \alpha - \sin^2 \theta \sin^2 \alpha - \sin^2 \alpha (1 - \cos^2 \theta)$$

$$= 0$$

$$\Rightarrow \sin^4 \theta + \sin^4 \alpha - 2 \sin^2 \theta \sin^2 \alpha = 0$$

$$\Rightarrow (\sin^2 \theta - \sin^2 \alpha)^2 = 0$$

$$\Rightarrow \sin^2 \theta = \sin^2 \alpha \text{ and } \cos^2 \theta = \cos^2 \alpha$$

$$\therefore \cos^8 \theta \sec^6 \alpha + \sin^8 \theta \operatorname{cosec}^6 \alpha$$

$$= \cos^2 \theta + \sin^2 \theta = 1$$

$$\Rightarrow \cos^8 \theta \sec^6 \alpha, \frac{1}{2} \text{ and } \sin^8 \theta \operatorname{cosec}^6 \alpha \text{ are in}$$

A.P.



$$112. (0.05)^{\log_{\sqrt{20}}(0.1+0.01+\dots)} = \left(\frac{1}{20}\right)^{2\log_{20}\left(\frac{0.1}{1-0.1}\right)}$$

$$= 20^{-2\log_{20}(1/9)} = 20^{2\log_{20}9}$$

$$= 20^{\log_{20}9^2} = 9^2 = 81$$

113. If $\log_{ax}x$, $\log_{bx}x$, $\log_{cx}x$ are in H.P.

Then $\frac{1}{\log_{ax}x}$, $\frac{1}{\log_{bx}x}$, $\frac{1}{\log_{cx}x}$ are in A.P.

i.e., $\log_x ax$, $\log_x bx$, $\log_x cx$ are in A.P.

$$\therefore 2 \log_x bx = \log_x ax + \log_x cx$$

$$\therefore \log_x b^2x^2 = \log_x ac \cdot x^2$$

$$\therefore b^2x^2 = ac \cdot x^2$$

$$\therefore b^2 = ac$$

$\therefore a, b, c$, are in G.P.

114. A.M. \geq G.M.

$$\therefore \frac{27^{\cos x} + 81^{\sin x}}{2} \geq \sqrt{27^{\cos x} \times 81^{\sin x}}$$

$$\Rightarrow 27^{\cos x} + 81^{\sin x} \geq 2\sqrt{3^{3\cos x + 4\sin x}}$$

$$\Rightarrow 27^{\cos x} + 81^{\sin x} \geq 2 \times \sqrt{3^{-5}}$$

$$\dots [\because -5 \leq 3 \cos x + 4 \sin x \leq 5]$$

$$\Rightarrow 27^{\cos x} + 81^{\sin x} \geq \frac{2}{9\sqrt{3}}$$

Hence, the minimum value of $27^{\cos x} + 81^{\sin x}$

$$\text{is } \frac{2}{9\sqrt{3}}.$$

115. Since, $\tan \frac{2\pi}{18}$, x , $\tan \frac{7\pi}{18}$ are in A.P. and

$\tan \frac{2\pi}{18}$, y , $\tan \frac{5\pi}{18}$ are in A.P.

$$\therefore 2x = \tan \frac{2\pi}{18} + \tan \frac{7\pi}{18} \text{ and } 2y = \tan \frac{2\pi}{18} + \tan \frac{5\pi}{18}$$

$$\Rightarrow 2x = \tan 20^\circ + \tan 70^\circ \text{ and}$$

$$2y = \tan 20^\circ + \tan 50^\circ$$

$$\Rightarrow 2x = \frac{\sin(20^\circ + 70^\circ)}{\cos 20^\circ \cos 70^\circ} \text{ and } 2y = \frac{\sin(20^\circ + 50^\circ)}{\cos 20^\circ \cos 50^\circ}$$

$$\Rightarrow 2x = \frac{\sin 90^\circ}{\sin 20^\circ \cos 20^\circ} \text{ and } 2y = \frac{\sin 70^\circ}{\cos 20^\circ \cos 50^\circ}$$

$$\Rightarrow x = \frac{1}{2 \sin 20^\circ \cos 20^\circ} \text{ and } 2y = \frac{1}{\cos 50^\circ}$$

$$\Rightarrow x = \frac{1}{\sin 40^\circ} \text{ and } 2y = \frac{1}{\sin 40^\circ}$$

$$\Rightarrow x = 2y \Rightarrow \frac{x}{y} = 2$$

116. Given, $\cos(\theta - \alpha)$, $\cos \theta$ and $\cos(\theta + \alpha)$ are in H.P.

$$\Rightarrow \frac{1}{\cos(\theta - \alpha)}, \frac{1}{\cos \theta}, \frac{1}{\cos(\theta + \alpha)} \text{ will be in A.P.}$$

$$\Rightarrow \frac{2}{\cos \theta} = \frac{1}{\cos(\theta - \alpha)} + \frac{1}{\cos(\theta + \alpha)}$$

$$= \frac{\cos(\alpha + \theta) + \cos(\theta - \alpha)}{\cos^2 \theta - \sin^2 \alpha}$$

$$\Rightarrow \frac{2}{\cos \theta} = \frac{2 \cos \theta \cos \alpha}{\cos^2 \theta - \sin^2 \alpha}$$

$$\Rightarrow \cos^2 \theta - \sin^2 \alpha = \cos^2 \theta \cos \alpha$$

$$\Rightarrow \cos^2 \theta (1 - \cos \alpha) = \sin^2 \alpha$$

$$\Rightarrow \cos^2 \theta \left(2 \sin^2 \frac{\alpha}{2}\right) = 4 \sin^2 \frac{\alpha}{2} \cos^2 \frac{\alpha}{2}$$

$$\Rightarrow \cos^2 \theta \sec^2 \frac{\alpha}{2} = 2$$

$$\Rightarrow \cos \theta \sec \frac{\alpha}{2} = \pm \sqrt{2}$$

117. $\sin \theta = \sqrt{\sin \phi \cos \phi}$

$$\Rightarrow \sin \phi \cos \phi = \sin^2 \theta$$

$$\Rightarrow \sin 2\phi = 2 \sin^2 \theta$$

$$\Rightarrow 1 - 2 \sin^2 \theta = 1 - \sin 2\phi$$

$$\Rightarrow \cos 2\theta = 1 - \cos \left(\frac{\pi}{2} - 2\phi\right)$$

$$= 2 \sin^2 \left(\frac{\pi}{4} - \phi\right)$$

118. $\cos 2B = \frac{\cos(A+C)}{\cos(A-C)} = \frac{\cos A \cos C - \sin A \sin C}{\cos A \cos C + \sin A \sin C}$

$$\Rightarrow \frac{1 - \tan^2 B}{1 + \tan^2 B} = \frac{1 - \tan A \tan C}{1 + \tan A \tan C}$$

$$\Rightarrow 1 + \tan^2 B - \tan A \tan C - \tan A \tan C \tan^2 B$$

$$= 1 - \tan^2 B + \tan A \tan C - \tan A \tan C \tan^2 B$$

$$\Rightarrow 2 \tan^2 B = 2 \tan A \tan C$$

$$\Rightarrow \tan^2 B = \tan A \tan C$$

$\therefore \tan A, \tan B, \tan C$ are in G.P.



Competitive Thinking

1. The given sequence is an A.P.

$$a = 10, d = -3$$

$$t_{30} = 10 + (30 - 1)(-3)$$

$$= -77$$



$$2. \quad \text{Given series } 27 + 9 + 5\frac{2}{5} + 3\frac{6}{7} + \dots$$

$$= 27 + \frac{27}{3} + \frac{27}{5} + \frac{27}{7} + \dots + \frac{27}{2n-1} + \dots$$

$$\text{Hence, } n^{\text{th}} \text{ term of given series } t_n = \frac{27}{2n-1}$$

$$\text{So, } t_9 = \frac{27}{2 \times 9 - 1} = \frac{27}{17} = 1\frac{10}{17}$$

3. Since, $a, 9, 3a - b$ and $3a + b$ are in A.P.

$$\therefore 9 - a = (3a + b) - (3a - b)$$

$$\Rightarrow 9 - a = 2b \Rightarrow a + 2b = 9 \quad \dots(i)$$

$$\text{Also, } 9 - a = (3a - b) - 9$$

$$\Rightarrow 4a - b = 18 \quad \dots(ii)$$

Eliminating b from (i) and (ii), we get

$$4a - 18 = (9 - a)/2$$

$$\Rightarrow 8a - 36 = 9 - a \Rightarrow 9a = 45 \Rightarrow a = 5$$

So, first 2 terms of the A.P. are 5 and 9

So, $a = 5, d = 4$

$$\therefore 2011^{\text{th}} \text{ term} = a + 2010d$$

$$= 5 + 2010 \times 4$$

$$= 8045$$

4. According to the given condition,

$$100(a + 99d) = 50(a + 49d)$$

$$\Rightarrow 2a + 198d = a + 49d$$

$$\Rightarrow a + 149d = 0$$

$$\therefore T_{150} = a + 149d = 0$$

5. $a + 1, 2a + 1, 4a - 1$ are in AP.

$$\therefore 2a + 1 - (a + 1) = 4a - 1 - (2a + 1)$$

$$\Rightarrow a = 2a - 2$$

$$\Rightarrow a = 2$$

6. It is not possible to express $a + b + 4c - 4d + e$ in terms of a .

$$7. \quad \text{Required ratio is } \frac{44}{99} = \frac{4}{9}$$

8. Given series $63 + 65 + 67 + 69 + \dots \dots (i)$

and $3 + 10 + 17 + 24 + \dots \dots (ii)$

Now from (i), m^{th} term $= (2m + 61)$ and m^{th} term of (ii) series $= (7m - 4)$

According to the given condition,

$$7m - 4 = 2m + 61$$

$$\Rightarrow 5m = 65 \Rightarrow m = 13$$

9. According to the given condition,

$$p \{a + (p - 1)d\} = q \{a + (q - 1)d\}$$

$$\Rightarrow a(p - q) + (p^2 - q^2)d + (q - p)d = 0$$

$$\Rightarrow (p - q) \{a + (p + q - 1)d\} = 0$$

$$\Rightarrow a + (p + q - 1)d = 0 \quad \dots[\because p \neq q]$$

$$\Rightarrow t_{p+q} = 0$$

10. Given that, $t_p = a + (p - 1)d = q \dots (i)$

and $t_q = a + (q - 1)d = p \dots (ii)$

$$\text{From (i) and (ii), we get } d = -\frac{(p - q)}{(p - q)} = -1$$

Putting the value of d in equation (i), we get

$$a = p + q - 1$$

$$t_r = a + (r - 1)d = (p + q - 1) + (r - 1)(-1)$$

$$= p + q - r$$

$$11. \quad t_m = a + (m - 1)d = \frac{1}{n} \text{ and}$$

$$t_n = a + (n - 1)d = \frac{1}{m}$$

$$\text{On solving, } a = \frac{1}{mn} \text{ and } d = \frac{1}{mn}$$

$$\therefore t_{mn} = a + (mn - 1)d = \frac{1}{mn} + (mn - 1)\frac{1}{mn} = 1$$

12. Let the first term be a and common difference be d .

The last 3 terms are T_{23}, T_{22} and T_{21} .

According to the given condition,

$$T_{21} + T_{22} + T_{23} = 261$$

$$\Rightarrow (a + 20d) + (a + 21d) + (a + 22d) = 261$$

$$\Rightarrow 3a + 63d = 261 \quad \dots(i)$$

Also, sum of 3 middle terms $= 141$

$$\Rightarrow T_{11} + T_{12} + T_{13} = 141$$

$$\Rightarrow (a + 10d) + (a + 11d) + (a + 12d) = 141$$

$$\Rightarrow 3a + 33d = 141 \quad \dots(ii)$$

Solving (i) and (ii), we get $a = 3$

$$13. \quad 164 = (3m^2 + 5m) - \{3(m - 1)^2 + 5(m - 1)\}$$

$$= (3m^2 + 5m) - 3m^2 + 6m - 3 - 5m + 5$$

$$\Rightarrow 164 = 6m + 2 \Rightarrow m = 27$$

$$14. \quad a = 3, d = 2$$

$$S_n = \frac{n}{2} [2a + (n - 1)d]$$

$$= \frac{n}{2} [6 + (n - 1)2]$$

$$= n(n + 2)$$

15. Series $108 + 117 + \dots + 999$ is an A.P.

Here, $a = 108, d = 9, t_n = 999$

Before 108, there are 11 multiples of 9 (and 108 is 12th multiple. 999 is 111th multiple of 9).

\Rightarrow From 108 to 999 there are 100 terms.

\Rightarrow Required sum

$$= \frac{100}{2} (108 + 999) \quad \dots \left[\because S_n = \frac{n}{2} (a +) \right]$$

$$= 50 \times 1107 = 55350$$



16. The series of all natural numbers is
3, 6, 9, 12, 99

$$\text{Here } n = \frac{99}{3} = 33, a = 3, d = 3$$

$$= 99$$

$$\therefore S_{33} = \frac{33}{2} \{3 + 99\}$$

$$= \frac{33}{2} \times 102$$

$$= 33 \times 51 = 1683$$

17. Series, 2 + 5 + 8 + 11 +
a = 2, d = 3 and let number of terms be n,

$$\text{then sum of A.P.} = \frac{n}{2} \{2a + (n-1)d\}$$

$$\Rightarrow 60100 = \frac{n}{2} \{2 \times 2 + (n-1)3\}$$

$$\Rightarrow 120200 = n(3n + 1)$$

$$\Rightarrow 3n^2 + n - 120200 = 0$$

$$\Rightarrow (n-200)(3n+601) = 0$$

$$\text{Hence, } n = 200$$

18. 12, 19, ..., 96 is the series of numbers which are of two digits and leave remainder 5 when divided by 7.

$$\text{Here, } a = 12, d = 7$$

$$\text{Last term } () = 96$$

$$S_{13} = \frac{13}{2} [12 + 96]$$

$$= \frac{13}{2} \times 108$$

$$= 702$$

19. For given series,

$$a = 1$$

$$d = 2$$

$$\therefore a_n = a + (n-1)d$$

$$\therefore 2001 = 1 + (n-1)(2)$$

$$\therefore n = 1001$$

$$\therefore S_{1001} = \frac{1001}{2} [2(1) + (1001-1) \times (2)]$$

$$\therefore S_{1001} = (1001)^2$$

20. k^{th} term = $5k + 1$

$$\therefore 1^{\text{st}} \text{ term} = a = 6$$

$$2^{\text{nd}} \text{ term} = 11$$

$$3^{\text{rd}} \text{ term} = 16$$

$$\therefore d = 5$$

$$\therefore S_{100} = \frac{100}{2} [2 \times 6 + (100-1) \times 5]$$

$$\therefore S_{100} = 50 (507)$$

21. Here, $a = ₹ 200, d = ₹ 40$

Saving in first two months = ₹ 400

Remained saving = $200 + 240 + 280 + \dots$
upto n terms

$$\Rightarrow \frac{n}{2} [400 + (n-1)40] = 11040 - 400$$

$$\Rightarrow 200n + 20n^2 - 20n = 10640$$

$$\Rightarrow 20n^2 + 180n - 10640 = 0$$

$$\Rightarrow n^2 + 9n - 532 = 0$$

$$\Rightarrow (n+28)(n-19) = 0$$

$$\Rightarrow n = 19$$

$$\therefore \text{Number of months} = 19 + 2 = 21$$

22. According to the given condition,

$$4500 = 150 \times 10$$

$$+ \{148 + 146 + \dots \text{ upto } n \text{ terms}\}$$

$$= 1500 + \frac{n}{2} \{296 + (n-1)(-2)\}$$

$$\Rightarrow n^2 - 149n + 3000 = 0$$

$$\Rightarrow (n-24)(n-125) = 0$$

$$\Rightarrow n = 24$$

$$\dots [\because n \neq 125]$$

So, total time taken = $10 + 24 = 34$ min.

23. Let the number of sides of the polygon be n.

Then the sum of interior angles of the polygon

$$= (2n-4) \frac{\pi}{2} = (n-2)\pi$$

Since, the angles are in A.P. and $a = 120^\circ, d = 5$ therefore,

$$\frac{n}{2} [2 \times 120 + (n-1)5] = (n-2)180$$

$$\Rightarrow n^2 - 25n + 144 = 0$$

$$\Rightarrow (n-9)(n-16) = 0$$

$$\Rightarrow n = 9, 16$$

But $n = 16$ gives,

$$T_{16} = a + 15d$$

$$= 120^\circ + 15.5^\circ$$

$$= 195^\circ \text{ which is impossible, as interior}$$

angle cannot be greater than 180° .

Hence, $n = 9$.

24. We have $\frac{S_{n_1}}{S_{n_2}} = \frac{2n+3}{6n+5}$

$$\Rightarrow \frac{\frac{n}{2} [2a_1 + (n-1)d_1]}{\frac{n}{2} [2a_2 + (n-1)d_2]} = \frac{2n+3}{6n+5}$$



$$\Rightarrow \frac{2 \left[a_1 + \left(\frac{n-1}{2} \right) d_1 \right]}{2 \left[a_2 + \left(\frac{n-1}{2} \right) d_2 \right]} = \frac{2n+3}{6n+5}$$

$$\Rightarrow \frac{a_1 + \left(\frac{n-1}{2} \right) d_1}{a_2 + \left(\frac{n-1}{2} \right) d_2} = \frac{2n+3}{6n+5}$$

Put $n = 25$ then $\frac{a_1 + 12d_1}{a_2 + 12d_2} = \frac{2(25)+3}{6(25)+5}$

$$\Rightarrow \frac{t_{131}}{t_{132}} = \frac{53}{155}$$

25. Let the first term be a and common difference be d .

Given, $\frac{a_1 + a_2 + \dots + a_p}{a_1 + a_2 + \dots + a_q} = \frac{p^2}{q^2}$

$$\Rightarrow \frac{pa + d[1+2+\dots+(p-1)]}{qa + d[1+2+\dots+(q-1)]} = \frac{p^2}{q^2}$$

$$\Rightarrow \frac{pa + \frac{p(p-1)}{2}d}{qa + \frac{q(q-1)}{2}d} = \frac{p^2}{q^2} \Rightarrow \frac{a + \left(\frac{p-1}{2} \right) d}{a + \left(\frac{q-1}{2} \right) d} = \frac{p}{q}$$

We have to find, $\frac{a_6}{a_{21}} = \frac{a+5d}{a+20d}$

Put $\frac{p-1}{2} = 5$ and $\frac{q-1}{2} = 20$

$$\Rightarrow p = 11 \text{ and } q = 41$$

$$\therefore \frac{a+5d}{a+20d} = \frac{11}{41}$$

26. According to the given condition,

$$\frac{n}{2} \{2a + (n-1)d\} = \frac{m}{2} \{2a + (m-1)d\}$$

$$\Rightarrow 2a(m-n) + d(m^2 - m - n^2 + n) = 0$$

$$\Rightarrow (m-n)\{2a + d(m+n-1)\} = 0$$

$$\Rightarrow 2a + (m+n-1)d = 0 \quad \dots [\because m \neq n]$$

$$\therefore S_{m+n} = \frac{m+n}{2} \{2a + (m+n-1)d\}$$

$$= \frac{m+n}{2} \{0\} = 0$$

27. Given that $S_n = nA + n^2B$
Putting $n = 1, 2, 3, \dots$ we get
 $S_1 = A + B, S_2 = 2A + 4B, S_3 = 3A + 9B$

Therefore,

$$T_1 = S_1 = A + B,$$

$$T_2 = S_2 - S_1 = A + 3B,$$

$$T_3 = S_3 - S_2 = A + 5B,$$

Hence, the sequence is

$(A + B), (A + 3B), (A + 5B), \dots$

Here, $a = A + B$ and common difference $d = 2B$

28. $S_1 = a_2 + a_4 + a_6 + a_8 + \dots + a_{100}$
 $S_2 = a_1 + a_3 + a_5 + a_7 + \dots + a_{99}$
 $\therefore S_1 - S_2 = (a_2 - a_1) + (a_4 - a_3) + \dots + (a_{100} - a_{99})$
 $= d + d + \dots + d = 50d \Rightarrow d = \frac{S_1 - S_2}{50}$

29. As given $a_2 - a_1 = a_3 - a_2 = \dots = a_n - a_{n-1} = d$
Where d is the common difference of the given A.P.

Also $a_n = a_1 + (n-1)d$

Then by rationalising each term,

$$\begin{aligned} & \frac{1}{\sqrt{a_2} + \sqrt{a_1}} + \frac{1}{\sqrt{a_3} + \sqrt{a_2}} + \dots + \frac{1}{\sqrt{a_n} + \sqrt{a_{n-1}}} \\ &= \frac{\sqrt{a_2} - \sqrt{a_1}}{a_2 - a_1} + \frac{\sqrt{a_3} - \sqrt{a_2}}{a_3 - a_2} + \dots + \frac{\sqrt{a_n} - \sqrt{a_{n-1}}}{a_n - a_{n-1}} \\ &= \frac{1}{d} (\sqrt{a_2} - \sqrt{a_1} + \sqrt{a_3} - \sqrt{a_2} + \dots + \sqrt{a_n} - \sqrt{a_{n-1}}) \end{aligned}$$

$$= \frac{1}{d} (\sqrt{a_n} - \sqrt{a_1}) = \frac{1}{d} \left(\frac{a_n - a_1}{\sqrt{a_n} + \sqrt{a_1}} \right)$$

$$= \frac{1}{d} \left\{ \frac{(n-1)d}{\sqrt{a_n} + \sqrt{a_1}} \right\} = \frac{n-1}{\sqrt{a_n} + \sqrt{a_1}}$$

30. $\frac{1}{S_1 S_2} + \frac{1}{S_2 S_3} + \dots + \frac{1}{S_{100} S_{101}} = \frac{1}{6}$
 $\Rightarrow \frac{1}{d} \left[\frac{S_2 - S_1}{S_1 S_2} + \frac{S_3 - S_2}{S_2 S_3} + \dots + \frac{S_{101} - S_{100}}{S_{100} S_{101}} \right] = \frac{1}{6}$
 $\dots [\because S_2 - S_1 = S_3 - S_2 = \dots = d]$
 $\Rightarrow \frac{1}{d} \left[\frac{1}{S_1} - \frac{1}{S_2} + \frac{1}{S_2} - \frac{1}{S_3} + \dots + \frac{1}{S_{100}} - \frac{1}{S_{101}} \right] = \frac{1}{6}$



$$\Rightarrow \frac{1}{d} \left[\frac{1}{S_1} - \frac{1}{S_{101}} \right] = \frac{1}{6} \Rightarrow \frac{1}{d} \left[\frac{1}{S_1} - \frac{1}{S_1 + 100d} \right] = \frac{1}{6}$$

$$\Rightarrow \frac{1}{d} \left[\frac{100d}{S_1(S_1 + 100d)} \right] = \frac{1}{6}$$

$$\Rightarrow S_1(S_1 + 100d) = 600 \quad \dots(i)$$

Given, $S_1 + S_{101} = 50$

$$\Rightarrow S_1 + (S_1 + 100d) = 50 \Rightarrow 2S_1 + 100d = 50$$

$$\Rightarrow S_1 + 50d = 25$$

$$\Rightarrow S_1 = 25 - 50d \quad \dots(ii)$$

Putting (ii) in (i), we get

$$(25 - 50d)(25 + 50d) = 600$$

$$\Rightarrow 625 - 2500d^2 = 600$$

$$\Rightarrow d^2 = \frac{1}{100} \Rightarrow d = \pm \frac{1}{10}$$

$$\therefore |S_1 - S_{101}| = |S_1 - (S_1 + 100d)|$$

$$= |-100d| = 100|d| \quad \dots[\because |xy| = |x| \cdot |y|]$$

$$\therefore |S_1 - S_{101}| = 10 \quad \dots[\because d = \pm 1/10]$$

31. $a_1, a_2, a_3, \dots, a_{n+1}$ are in A.P. and common difference = d

$$\text{Let } S = \frac{1}{a_1 a_2} + \frac{1}{a_2 a_3} + \dots + \frac{1}{a_n a_{n+1}}$$

$$\Rightarrow S = \frac{1}{d} \left\{ \frac{d}{a_1 a_2} + \frac{d}{a_2 a_3} + \dots + \frac{d}{a_n a_{n+1}} \right\}$$

$$\Rightarrow S = \frac{1}{d} \left\{ \frac{a_2 - a_1}{a_1 a_2} + \frac{a_3 - a_2}{a_2 a_3} + \dots + \frac{a_{n+1} - a_n}{a_n a_{n+1}} \right\}$$

$$\Rightarrow S = \frac{1}{d} \left\{ \frac{1}{a_1} - \frac{1}{a_2} + \frac{1}{a_2} - \frac{1}{a_3} + \dots + \frac{1}{a_n} - \frac{1}{a_{n+1}} \right\}$$

$$\Rightarrow S = \frac{1}{d} \left\{ \frac{1}{a_1} - \frac{1}{a_{n+1}} \right\} = \frac{1}{d} \left\{ \frac{a_{n+1} - a_1}{a_1 a_{n+1}} \right\}$$

$$\Rightarrow S = \frac{1}{d} \left(\frac{nd}{a_1 a_{n+1}} \right) = \frac{n}{a_1 a_{n+1}}$$

Trick: Check for $n = 2$.

32. Since, $a_1 = 0$

$$\therefore a_2 = d, a_3 = 2d, \dots$$

$$\therefore \left(\frac{a_3}{a_2} + \frac{a_4}{a_3} + \dots + \frac{a_n}{a_{n-1}} \right) - a_2 \left(\frac{1}{a_2} + \frac{1}{a_3} + \dots + \frac{1}{a_{n-2}} \right)$$

$$= \left(\frac{2d}{d} + \frac{3d}{2d} + \dots + \frac{(n-1)d}{(n-2)d} \right)$$

$$- d \left(\frac{1}{d} + \frac{1}{2d} + \dots + \frac{1}{(n-3)d} \right)$$

$$= \left(\frac{2}{1} + \frac{3}{2} + \dots + \frac{n-1}{n-2} \right) - \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n-3} \right)$$

$$= \left[(1+1) + \left(1 + \frac{1}{2} \right) + \dots + \left(\frac{n-1}{n-2} \right) \right]$$

$$- \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n-3} \right)$$

$$= (n-3) + \frac{n-1}{n-2} = (n-3) + 1 + \frac{1}{n-2}$$

$$= (n-2) + \frac{1}{n-2}$$

33. As given

$$d = a_2 - a_1 = a_3 - a_2 = \dots = a_n - a_{n-1}$$

$$\therefore \sin d \{ \operatorname{cosec} a_1 \operatorname{cosec} a_2 + \dots + \operatorname{cosec} a_{n-1} \operatorname{cosec} a_n \}$$

$$= \frac{\sin(a_2 - a_1)}{\sin a_1 \cdot \sin a_2} + \dots + \frac{\sin(a_n - a_{n-1})}{\sin a_{n-1} \sin a_n}$$

$$= (\cot a_1 - \cot a_2) + (\cot a_2 - \cot a_3) + \dots + (\cot a_{n-1} - \cot a_n)$$

$$= \cot a_1 - \cot a_n$$

34. $\log_3 2, \log_3(2^x - 5)$ and $\log_3 \left(2^x - \frac{7}{2} \right)$ are in A.P.

$$\Rightarrow 2 \log_3(2^x - 5) = \log_3 \left[(2) \left(2^x - \frac{7}{2} \right) \right]$$

$$\Rightarrow (2^x - 5)^2 = 2^{x+1} - 7$$

$$\Rightarrow 2^{2x} - 12 \cdot 2^x + 32 = 0$$

$$\Rightarrow x = 2, 3$$

But $x = 2$ does not hold, hence $x = 3$

35. $2 \tan^{-1} y = \tan^{-1} x + \tan^{-1} z$

$$\Rightarrow \tan^{-1} \left(\frac{2y}{1-y^2} \right) = \tan^{-1} \left(\frac{x+z}{1-xz} \right)$$

$$\Rightarrow \frac{2y}{1-y^2} = \frac{x+z}{1-xz}$$

$$\text{But } 2y = x + z \quad \dots[\because x, y, z \text{ are in A.P.}]$$

$$\therefore 1 - y^2 = 1 - xz$$

$$\Rightarrow y^2 = xz$$

$$\therefore x, y, z \text{ are both in G.P. and A.P.,}$$

$$\therefore x = y = z$$

36. Since, a, b, c are in A.P., we get

$$b - c = -d, \quad \dots(i)$$

$$c - a = 2d, \quad \dots(ii)$$

$$a - b = -d \quad \dots(iii)$$

Also, since x, y, z are in G.P., we get

$$y^2 = xz \quad \dots(iv)$$

$$\text{Now, } x^{b-c} \cdot y^{c-a} \cdot z^{a-b}$$

$$= x^{-d} \cdot y^{2d} \cdot z^{-d} \quad \dots[\text{From (i), (ii), (iii)}]$$

$$= x^{-d} \cdot (xz)^d \cdot z^{-d}$$

$$= 1$$



37. If a, b, c are in A.P., then $2b = a + c$

$$\begin{aligned} \text{So, } \frac{(a-c)^2}{(b^2-ac)} &= \frac{(a-c)^2}{\left\{\left(\frac{a+c}{2}\right)^2-ac\right\}} \\ &= \frac{4(a-c)^2}{[a^2+c^2+2ac-4ac]} \\ &= \frac{4(a-c)^2}{(a-c)^2} = 4 \end{aligned}$$

Trick: Put $a = 1, b = 2, c = 3$, then the required value is $\frac{4}{1} = 4$.

38. Let $a-d, a, a+d$ be the roots of the equation $x^3 - 12x^2 + 39x - 28 = 0$

Then, $(a-d) + a + (a+d) = 12$ and $(a-d)a(a+d) = 28$

$$\Rightarrow 3a = 12 \text{ and } a(a^2 - d^2) = 28$$

$$\Rightarrow a = 4 \text{ and } a(a^2 - d^2) = 28$$

$$\Rightarrow 16 - d^2 = 7$$

$$\Rightarrow d = \pm 3$$

39. Arithmetic mean of ${}^n C_0, {}^n C_1, {}^n C_2, \dots, {}^n C_n$ i.e. $(n+1)$ terms

$$\begin{aligned} &= \frac{{}^n C_0 + {}^n C_1 + {}^n C_2 + \dots + {}^n C_n}{n+1} \\ &= \frac{2^n}{n+1} \end{aligned}$$

40. For set a to $2b$, $2b$ is the $(n+2)$ th term

$$\therefore 2b = a + (n+1)d$$

$$\Rightarrow d = \frac{2b-a}{n+1}$$

$$\therefore m^{\text{th}} \text{ mean} = a + md = a + m \left(\frac{2b-a}{n+1} \right) \quad \dots(i)$$

For set $2a$ to b , b is the $(n+2)$ th term

$$\therefore b = 2a + (n+1)d$$

$$\Rightarrow d = \frac{b-2a}{n+1}$$

$$\therefore m^{\text{th}} \text{ mean} = 2a + md = 2a + m \left(\frac{b-2a}{n+1} \right) \quad \dots(ii)$$

\therefore From (i) and (ii)

$$a + m \left(\frac{2b-a}{n+1} \right) = 2a + m \left(\frac{b-2a}{n+1} \right)$$

$$\Rightarrow \frac{a}{b} = \frac{m}{n+1-m}$$

41. The given sequence is a G.P.

$$a = 3, r = \frac{1}{3}$$

$$t_6 = 3 \left(\frac{1}{3} \right)^{6-1}$$

$$= 3 \left(\frac{1}{3} \right)^5$$

$$= \frac{1}{81}$$

$$42. r = \frac{1}{3} \sqrt{\frac{20}{3}} \cdot \frac{9}{10} = \frac{\sqrt{60}}{10} = \sqrt{\frac{6}{10}} = \sqrt{\frac{3}{5}}$$

$$\therefore t_5 = ar^4 = \left(\frac{10}{9} \right) \left(\frac{3}{5} \right)^2 = \frac{10}{9} \cdot \frac{9}{25} = \frac{2}{5}$$

43. Given that $x, 2x+2, 3x+3$ are in G.P.

Therefore,

$$(2x+2)^2 = x(3x+3)$$

$$\Rightarrow x^2 + 5x + 4 = 0$$

$$\Rightarrow (x+4)(x+1) = 0$$

$$\Rightarrow x = -1, -4$$

Now, first term: $a = x$

and second term: $ar = 2(x+1)$

$$\Rightarrow r = \frac{2(x+1)}{x}$$

$$\text{then } 4^{\text{th}} \text{ term} = ar^3 = x \left[\frac{2(x+1)}{x} \right]^3 = \frac{8}{x^2} (x+1)^3$$

Putting, $x = -4$

$$\text{We get, } t_4 = \frac{8}{16} (-3)^3 = -\frac{27}{2} = -13.5$$

44. Let the first four terms be $a, -ar, ar^2, -ar^3$, where $r > 0, a > 0$

According to the given conditions,

$$a - ar = 12 \text{ and } ar^2 - ar^3 = 48$$

By solving, we get $r = 2$ ($r > 0$)

So, $a = -12$

$$45. t_5 = ar^4 = \frac{1}{3} \quad \dots(i)$$

$$\text{and } t_9 = ar^8 = \frac{16}{243} \quad \dots(ii)$$

Solving (i) and (ii), we get $r = \frac{2}{3}$ and $a = \frac{27}{16}$

$$\text{Now } 4^{\text{th}} \text{ term} = ar^3 = \frac{3^3}{2^4} \cdot \frac{2^3}{3^3} = \frac{1}{2}$$



46. $t_n = t_{n+1} + t_{n+2}$
 $\Rightarrow ar^{n-1} = ar^n + ar^{n+1}$
 $\Rightarrow r^{n-1} = r^n(1+r)$
 $\Rightarrow r^2 + r = 1$
 $\Rightarrow r^2 + r - 1 = 0$
 $\therefore r = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
 $= \frac{-1 \pm \sqrt{1+4}}{2}$
 $= \frac{-1 \pm \sqrt{5}}{2}$
47. Let first term and common ratio of G.P. are respectively a and r, then under condition,
 $t_n = t_{n-1} + t_{n-2}$
 $\Rightarrow ar^{n-1} = ar^{n-2} + ar^{n-3}$
 $\Rightarrow ar^{n-1} = ar^{n-1}r^{-1} + ar^{n-1}r^{-2}$
 $\Rightarrow 1 = \frac{1}{r} + \frac{1}{r^2}$
 $\Rightarrow r^2 - r - 1 = 0$
 $\Rightarrow r = \frac{1 \pm \sqrt{1+4}}{2} = \frac{1 + \sqrt{5}}{2}$
 Taking only (+) sign ($\because r > 1$)
48. Let the G.P. be a, ar, ar², ar³, ar⁴, ...
 $t_2 + t_5 = ar + ar^4 = 216$
 $\frac{t_4}{t_6} = \frac{ar^3}{ar^5} = \frac{1}{4}$
 $\Rightarrow r^2 = 4 \Rightarrow r = \pm 2$
 For r = 2,
 $a(2 + 2^4) = 216$
 $\Rightarrow a(18) = 216$
 $\Rightarrow a = \frac{216}{18} = 12$
 For r = -2,
 $a(-2 + 2^4) = 216$
 $\Rightarrow a(14) = 216$
 $\Rightarrow a = \frac{216}{14} = \frac{108}{7}$
 $\therefore a = 12$
49. Let first term of G.P. = A and common ratio = r
 We know that nth term of G.P. = Arⁿ⁻¹
 Now $t_4 = a = Ar^3$, $t_7 = b = Ar^6$ and $t_{10} = c = Ar^9$
 Relation $b^2 = ac$ is true because
 $b^2 = (Ar^6)^2 = A^2r^{12}$ and $ac = (Ar^3)(Ar^9) = A^2r^{12}$

Alternate method : As we know, if p, q, r in A.P., then pth, qth, rth terms of a G.P. are always in G.P., therefore, a, b, c will be in G.P. i.e. $b^2 = ac$.

50. The given series is a G.P. with a = i, r = -i
 $\therefore S_{100} = \frac{i(1-i^{100})}{1+i}$
 $= \frac{i(1-(i^2)^{50})}{1+i}$
 $= \frac{i(1-1)}{1+i} = 0$
51. $\frac{ar^n - a}{r-1} = 364$
 $\Rightarrow \frac{ar^{n-1} \cdot r - a}{r-1} = 364 \quad \dots(i)$
 $\Rightarrow \frac{3 \times 243 - a}{2} = 364$
 $\Rightarrow a = 1$
 Now, putting this in (i), n = 6
52. $\therefore n^{\text{th}}$ term of series = $ar^{n-1} = a(3)^{n-1} = 486 \dots(i)$
 and sum of n terms of series.
 $S_n = \frac{a(3^n - 1)}{3-1} = 728 (\because r > 1) \quad \dots(ii)$
 From (i), $a\left(\frac{3^n}{3}\right) = 486$ or $a \cdot 3^n = 3 \times 486 = 1458$
 From (ii), $a \cdot 3^n - a = 728 \times 2$
 or $a \cdot 3^n - a = 1456$
 $\therefore 1458 - a = 1456$
 $\Rightarrow a = 2$
53. $t_2 = ar = 24$
 $t_5 = ar^4 = 3$
 $\frac{t_5}{t_2} = \frac{1}{8} = r^3$
 $\Rightarrow r = \frac{1}{2}$ & a = 48
 $S_6 = \frac{a(1-r^6)}{1-r}$
 $= \frac{48\left(1 - \left(\frac{1}{2}\right)^6\right)}{1 - \frac{1}{2}} = \frac{189}{2}$



$$\begin{aligned}
54. \quad & 9 + 99 + 999 + \dots 10 \text{ terms} \\
& = (10 - 1) + (100 - 1) + (1000 - 1) \\
& \quad \quad \quad + \dots 10 \text{ terms} \\
& = (10 + 100 + 1000 + \dots 10 \text{ terms}) \\
& \quad \quad \quad - (1 + 1 + 1 + \dots 10 \text{ terms}) \\
& = (10 + 10^2 + 10^3 + \dots 10 \text{ terms}) - (10) \\
& = \frac{10(10^{10} - 1)}{10 - 1} - 10 \\
& = \frac{10(10^{10} - 1)}{9} - 10 \\
& = \frac{10(10^{10} - 1) - 90}{9} \\
& = \frac{100}{9}(10^9 - 1)
\end{aligned}$$

$$\begin{aligned}
55. \quad & \text{Series } 3 + 33 + 333 + \dots + n \text{ terms} \\
& \text{Given series can be written as,} \\
& = \frac{1}{3}[9 + 99 + 999 + \dots + n \text{ terms}] \\
& = \frac{1}{3}[(10 - 1) + (10^2 - 1) + (10^3 - 1) \\
& \quad \quad \quad + \dots + n \text{ terms}] \\
& = \frac{1}{3}[10 + 10^2 + \dots + 10^n] - \frac{1}{3}[1 + 1 + 1 \\
& \quad \quad \quad + \dots + n \text{ terms}] \\
& = \frac{1}{3} \cdot \frac{10(10^n - 1)}{10 - 1} - \frac{1}{3} \cdot n \\
& = \frac{1}{3} \left[\frac{10^{n+1} - 10}{9} - n \right] \\
& = \frac{1}{3} \left[\frac{10^{n+1} - 9n - 10}{9} \right] \\
& = \frac{1}{27} [10^{n+1} - 9n - 10]
\end{aligned}$$

$$\begin{aligned}
56. \quad & \text{Let the G.P. be } a, ar, ar^2, ar^3, \dots, ar^{48}, ar^{49} \\
& \text{i.e., } a_1 = a, a_2 = ar, a_3 = ar^2, \dots, a_{49} = ar^{48} \\
& \text{and } a_{50} = ar^{49} \\
\therefore & \frac{a_1 - a_3 + a_5 - \dots + a_{49}}{a_2 - a_4 + a_6 - \dots + a_{50}} \\
& = \frac{a - ar^2 + ar^4 - \dots + ar^{48}}{ar - ar^3 + ar^5 - \dots + ar^{49}} \\
& = \frac{a(1 - (-r^2)^{25})}{1 - (-r^2)} = \frac{1}{r} = \frac{a}{ar} = \frac{a_1}{a_2}
\end{aligned}$$

$$\begin{aligned}
57. \quad & \text{Since } n^m + 1 \text{ divides } 1 + n + n^2 + \dots + n^{127} \\
& \text{Therefore, } \frac{1 + n + n^2 + \dots + n^{127}}{n^m + 1} \text{ is an integer} \\
& \Rightarrow \frac{1 - n^{128}}{1 - n} \times \frac{1}{n^m + 1} \text{ is an integer} \\
& \Rightarrow \frac{(1 - n^{64})(1 + n^{64})}{(1 - n)(n^m + 1)} \\
& \text{is an integer, when largest } m = 64.
\end{aligned}$$

$$\begin{aligned}
58. \quad & \frac{\sqrt{2} + 1}{\sqrt{2} - 1}, \frac{1}{\sqrt{2}(\sqrt{2} - 1)}, \frac{1}{2}, \dots \\
& \text{Common ratio of the series} = \frac{1}{\sqrt{2}(\sqrt{2} + 1)} \\
\therefore \text{ sum} & = \frac{a}{1 - r} = \frac{\left(\frac{\sqrt{2} + 1}{\sqrt{2} - 1}\right)}{\left(1 - \frac{1}{\sqrt{2}(\sqrt{2} + 1)}\right)} \\
& = \frac{(\sqrt{2} + 1)}{(\sqrt{2} - 1)} \cdot \frac{\sqrt{2}(\sqrt{2} + 1)}{(1 + \sqrt{2})} \\
& = \sqrt{2}(\sqrt{2} + 1)^2
\end{aligned}$$

$$\begin{aligned}
59. \quad & \text{Clearly it is a infinite G.P. whose common} \\
& \text{ratio is } 0.24. \\
\therefore S_{\infty} & = \frac{a}{1 - r} = \frac{5.05}{1 - 0.24} = 6.64474
\end{aligned}$$

$$\begin{aligned}
60. \quad & (32)(32)^{1/6}(32)^{1/36} \dots \infty = (32)^{1 + \frac{1}{6} + \frac{1}{36} + \dots \infty} \\
& = (32)^{\frac{1}{1 - (1/6)}} = (32)^{\frac{1}{5/6}} = (32)^{\frac{6}{5}} \\
& = 2^6 = 64
\end{aligned}$$

$$\begin{aligned}
61. \quad & \text{According to the given condition,} \\
& \frac{a}{1 - r} = \frac{4}{3} \\
& \Rightarrow \frac{3}{4} \left(\frac{1}{1 - r} \right) = \frac{4}{3} \\
& \Rightarrow r = 1 - \frac{9}{16} = \frac{7}{16}
\end{aligned}$$

$$\begin{aligned}
62. \quad & \text{According to the given condition,} \\
& 4 = \frac{a}{1 - r} \\
& \Rightarrow 4 \Rightarrow a = 4 - 4r \\
& \Rightarrow 4r = 4 - a \\
& \text{Only option (D) satisfies this condition.}
\end{aligned}$$

$$\begin{aligned}
63. \quad & 3 + 3\alpha + 3\alpha^2 + 3\alpha^3 + \dots \infty = \frac{45}{8} \\
& \Rightarrow 3 \left[\frac{1}{1 - \alpha} \right] = \frac{45}{8} \Rightarrow 8 = 15(1 - \alpha) \Rightarrow \alpha = \frac{7}{15}
\end{aligned}$$



$$64. \quad y = x - x^2 + x^3 - x^4 + \dots \infty$$

Then $xy = x^2 - x^3 + x^4 - \dots \infty$

Adding, $y + xy = x + 0 + 0 \dots + 0$

$$\Rightarrow x - xy = y$$

$$\Rightarrow x(1 - y) = y$$

$$\Rightarrow x = \frac{y}{1 - y}$$

Alternate method:

$$y = \frac{x}{1 - (-x)} \Rightarrow y = \frac{x}{1 + x}$$

$$\Rightarrow y + yx = x \Rightarrow x = \frac{y}{1 - y}$$

$$65. \quad \text{Common ratio (r)} = \frac{2}{x}$$

$$\text{For sum to be finite } r < 1 \Rightarrow \frac{2}{x} < 1$$

$$\Rightarrow 2 < x$$

$$\Rightarrow x > 2$$

$$66. \quad \text{We have } \frac{a}{1 - r} = x$$

$$\text{and } \frac{a^2}{1 - r^2} = \frac{a}{1 - r} \cdot \frac{a}{1 + r} = y$$

$$\Rightarrow y = x \cdot \frac{a}{1 + r} = x \cdot \frac{x(1 - r)}{1 + r}$$

$$\Rightarrow \frac{y}{x^2} = \frac{1 - r}{1 + r} \Rightarrow \frac{x^2}{y} = \frac{1 + r}{1 - r}$$

$$\Rightarrow \frac{x^2}{y} (1 - r) = 1 + r$$

$$\Rightarrow r \left[1 + \frac{x^2}{y} \right] = -1 + \frac{x^2}{y}$$

$$\Rightarrow r = \frac{x^2 - y}{x^2 + y}$$

67. Let r be the common ratio of the G.P. Then

$$S = \frac{a}{1 - r} \Rightarrow r = 1 - \frac{a}{S}$$

Now $S_n = \text{Sum of } n \text{ terms}$

$$= a \left(\frac{1 - r^n}{1 - r} \right)$$

$$= \frac{a}{1 - r} (1 - r^n)$$

$$= S \left[1 - \left(1 - \frac{a}{S} \right)^n \right]$$

68. Since the series are in G.P., therefore

$$x = \frac{1}{1 - a} \text{ and } y = \frac{1}{1 - b}$$

$$\therefore a = \frac{x - 1}{x}, b = \frac{y - 1}{y}$$

$$\therefore 1 + ab + a^2b^2 + \dots \infty$$

$$= \frac{1}{1 - ab} = \frac{1}{1 - \frac{x-1}{x} \cdot \frac{y-1}{y}} = \frac{xy}{x + y - 1}$$

$$69. \quad 1 - \cos \alpha = \frac{1}{2 - \sqrt{2}} = 1 + \frac{1}{\sqrt{2}} \quad \dots \left[\because \frac{a}{1 - r} = 2 - \sqrt{2} \right]$$

$$\Rightarrow \cos \alpha = -\frac{1}{\sqrt{2}} = \cos \frac{3\pi}{4}$$

$$\Rightarrow \alpha = \frac{3\pi}{4}$$

$$70. \quad 1 + \sin x + \sin^2 x + \dots \text{ upto } \infty = 4 + 2\sqrt{3}$$

$$\Rightarrow \frac{1}{1 - \sin x} = 4 + 2\sqrt{3}$$

$$\Rightarrow 1 - \sin x = \frac{1}{2(2 + \sqrt{3})}$$

$$\Rightarrow \sin x = \frac{4 + 2\sqrt{3} - 1}{2(2 + \sqrt{3})}$$

$$\Rightarrow \sin x = \frac{\sqrt{3}}{2} \Rightarrow x = \frac{\pi}{3}, \frac{2\pi}{3}$$

$$71. \quad 0.2\dot{3}\dot{4} = \frac{234 - 2}{990} = \frac{232}{990}$$

$$72. \quad 0.4\dot{2}\dot{3} = \frac{423 - 4}{990} = \frac{419}{990}$$

$$73. \quad 0.14189189189\dots$$

$$= 0.14 + 0.00189 + 0.00000189 + \dots$$

$$= \frac{14}{100} + 189 \left[\frac{1}{10^5} + \frac{1}{10^8} + \dots \infty \right]$$

$$= \frac{7}{50} + 189 \left[\frac{1}{10^5} \right]$$

$$\left[1 - \left(\frac{1}{10^3} \right) \right]$$

$$= \frac{7}{50} + 189 \left[\frac{1}{10^5} \times \frac{10^3}{999} \right]$$

$$= \frac{7}{50} + \frac{189}{999 \times 100} = \frac{7}{50} + \frac{7}{3700}$$

$$= \frac{7}{50} + \frac{7}{25 \times 148} = \frac{21}{148}$$

**Alternate Method:**

$$0.14\dot{1}\ddot{8}\dot{9}$$

$$= \frac{14189 - 14}{99900} = \frac{14175}{99900} = \frac{21}{148}$$

74. Let α and β be the roots of equation $x^2 - 18x + 9 = 0$

$$\therefore \text{G.M. of } \alpha \text{ and } \beta = \sqrt{\alpha\beta} = \sqrt{9} = 3 \quad [\because \alpha\beta = 9]$$

75. Let G_1, G_2, G_3, G_4, G_5 be the G.M.'s are inserted between 486 and $\frac{2}{3}$. So total terms are 7.

$$t_n = ar^{n-1}$$

$$\Rightarrow \frac{2}{3} = 486(r)^6 \Rightarrow r = \frac{1}{3}$$

$$\text{Hence, 4}^{\text{th}} \text{ G.M. will be, } t_5 = ar^4$$

$$= 486 \left(\frac{1}{3}\right)^4$$

$$= 6$$

76. Let $a - d, a, a + d$ be three numbers in A.P.

$$\therefore a + d + a + a - d = 15$$

$$\Rightarrow a = 5$$

$a - d + 1, a + 4, a + d + 19$ are in G.P.

$$\Rightarrow 6 - d, 9, 24 + d \text{ are in G.P.}$$

$$\therefore 81 = (6 - d)(24 + d)$$

$$\Rightarrow 81 = 144 + 6d - 24d - d^2$$

$$\Rightarrow d^2 + 18d - 63 = 0$$

$$\therefore d = 3, -21$$

\therefore the numbers are 2, 5, 8 and 26, 5, -16

77. x, y, z are in G.P., then $y^2 = x.z$

$$\text{Now } a^x = b^y = c^z = m$$

$$\Rightarrow x \log_e a = y \log_e b = z \log_e c = \log_e m$$

$$\Rightarrow x = \log_a m, y = \log_b m, z = \log_c m$$

$$\text{Again as } x, y, z \text{ are in G.P., so } \frac{y}{x} = \frac{z}{y}$$

$$\Rightarrow \frac{\log_b m}{\log_a m} = \frac{\log_c m}{\log_b m}$$

$$\Rightarrow \log_b a = \log_c b$$

78. Let $a^{1/x} = b^{1/y} = c^{1/z}$

$$\Rightarrow a = k^x, b = k^y, c = k^z$$

Now, a, b, c are in G.P.

$$\Rightarrow b^2 = ac$$

$$\Rightarrow k^{2y} = k^x \cdot k^z = k^{x+z}$$

$$\Rightarrow 2y = x + z$$

$$\Rightarrow x, y, z \text{ are in A.P.}$$

79. Since, a, b, c are in G.P.

$$\therefore b^2 = ac$$

$$\Rightarrow \log_e b^2 = \log_e ac$$

$$\Rightarrow \log_e a - 2 \log_e b + \log_e c = 0$$

$$\text{Given, } (\log_e a)x^2 - (2 \log_e b)x + \log_e c = 0$$

Since, 1 satisfies this equation.

Therefore, 1 is one root and other root say β .

$$\therefore 1 \cdot \beta = \frac{\log_e c}{\log_e a}$$

$$\therefore \beta = \log_a c$$

$$80. \frac{x^{n+1} + y^{n+1}}{x^n + y^n} = \sqrt{xy} \Rightarrow x^{n+1} + y^{n+1} = \sqrt{xy} (x^n + y^n)$$

$$\Rightarrow x^{n+\frac{1}{2}} \left(x^{\frac{1}{2}} - y^{\frac{1}{2}}\right) = y^{n+\frac{1}{2}} \left(x^{\frac{1}{2}} - y^{\frac{1}{2}}\right)$$

$$\Rightarrow \left(\frac{x}{y}\right)^{n+\frac{1}{2}} = 1 \Rightarrow n = -\frac{1}{2}$$

81. $a + d, a + 4d, a + 8d$, are in G.P.

$$\Rightarrow (a + 4d)^2 = (a + d)(a + 8d)$$

$$\Rightarrow 8d^2 = ad \Rightarrow \frac{a}{d} = 8$$

$$\therefore \text{common ratio} = \frac{a + 4d}{a + d}$$

$$= \frac{8 + 4}{8 + 1} = \frac{4}{3}$$

82. Series, $2, 2\frac{1}{2}, 3\frac{1}{3}, \dots$ are in H.P.

$$\Rightarrow \frac{1}{2}, \frac{2}{5}, \frac{3}{10}, \dots \text{ will be in A.P.}$$

$$\text{Now, first term } a = \frac{1}{2} \text{ and}$$

$$\text{common difference } d = -\frac{1}{10}$$

So, 5th term of the A.P.

$$= \frac{1}{2} + (5 - 1) \left(-\frac{1}{10}\right) = \frac{1}{10}$$

Hence, 5th term of the H.P. is 10.

83. Here, 5th term of the corresponding

$$\text{A.P.} = a + 4d = 45 \quad \dots(i)$$

and 11th term of the corresponding

$$\text{A.P.} = a + 10d = 69 \quad \dots(ii)$$

From (i) and (ii), we get $a = 29, d = 4$

Therefore, 16th term of the corresponding A.P.

$$= a + 15d = 29 + 15 \times 4 = 89$$

Hence, 16th term of the H.P. is $\frac{1}{89}$.



84. Since $a_1, a_2, a_3, \dots, a_n$ are in H.P.
Therefore $\frac{1}{a_1}, \frac{1}{a_2}, \frac{1}{a_3}, \dots, \frac{1}{a_n}$ will be in A.P.
Which gives, $\frac{1}{a_2} - \frac{1}{a_1} = \frac{1}{a_3} - \frac{1}{a_2} = \dots$
$$= \frac{1}{a_n} - \frac{1}{a_{n-1}} = d$$

$$\Rightarrow \frac{a_1 - a_2}{a_1 a_2} = \frac{a_2 - a_3}{a_2 a_3} = \dots = \frac{a_{n-1} - a_n}{a_{n-1} a_n} = d$$

$$\Rightarrow a_1 - a_2 = da_1 a_2, a_2 - a_3 = da_2 a_3$$

and $a_{n-1} - a_n = da_{n-1} a_n$
Adding these, we get
$$d(a_1 a_2 + a_2 a_3 + \dots + a_{n-1} a_n)$$

$$= (a_1 + a_2 + \dots + a_{n-1}) - (a_2 + a_3 + \dots + a_n)$$

$$= a_1 - a_n \quad \dots(i)$$

Also n^{th} term of this A.P. is given by
$$\frac{1}{a_n} = \frac{1}{a_1} + (n-1)d \Rightarrow d = \frac{a_1 - a_n}{a_1 a_n (n-1)}$$

Substituting this value of d in (i)
$$(a_1 - a_n) = \frac{a_1 - a_n}{a_1 a_n (n-1)} (a_1 a_2 + a_2 a_3 + \dots + a_{n-1} a_n)$$

$$(a_1 a_2 + a_2 a_3 + \dots + a_{n-1} a_n) = a_1 a_n (n-1)$$
85. We know that, $x_n = \frac{(n+1)ab}{na+b}$
$$\therefore \text{Sixth H.M. i.e. } x_6 = \frac{7 \cdot 3 \cdot \left(\frac{6}{13}\right)}{\left(6 \cdot 3 + \frac{6}{13}\right)}$$

$$= \frac{126}{240}$$

$$= \frac{63}{120}$$
86. Let roots be α, β then
$$\alpha + \beta = -\frac{b}{a} = 10$$

$$\alpha\beta = \frac{c}{a} = 11$$

$$\text{H.M.} = \frac{2\alpha\beta}{\alpha+\beta} = \frac{11 \times 2}{10} = \frac{11}{5}$$
87. a, b, c are in H.P. $\Rightarrow b = \frac{2ac}{a+c}$
By inspection, we get (A) False (B) False (C) False

88. Since, a, b, c are in H.P.
$$\therefore \frac{2}{b} = \frac{1}{a} + \frac{1}{c} \Rightarrow b = \frac{2ac}{a+c}$$

Consider option (B),
$$\frac{1}{ca} = \frac{2\left(\frac{1}{bc} \cdot \frac{1}{ab}\right)}{\frac{1}{bc} + \frac{1}{ab}} = \frac{\left(\frac{2}{ab^2c}\right)}{\frac{a+c}{abc}}$$

$$= \frac{2(abc)}{ab^2c(a+c)} = \frac{2}{b(a+c)}$$

$$\Rightarrow ca = \frac{b(a+c)}{2}$$

$$\Rightarrow b = \frac{2ac}{a+c}$$

$$\therefore \frac{1}{bc}, \frac{1}{ca}, \frac{1}{ab} \text{ are in H.P.}$$
89. As given $H = \frac{2pq}{p+q}$
$$\therefore \frac{H}{p} + \frac{H}{q} = \frac{2q}{p+q} + \frac{2p}{p+q} = \frac{2(p+q)}{p+q} = 2$$
90. Let, the distance of school from home = d
and time taken are t_1 and t_2 .
$$\therefore t_1 = \frac{d}{x} \text{ and } t_2 = \frac{d}{y}$$

Avg. velocity = $\frac{\text{Total distance}}{\text{Total time}}$
$$= \frac{2d}{\left(\frac{d}{x} + \frac{d}{y}\right)} = \frac{2xy}{x+y}$$
, which is the H.M. of x and y .
91. If x, y, z are in H.P., then $y = \frac{2xz}{x+z}$
$$\therefore \log_e(x+z) + \log_e(x-2y+z)$$

$$= \log_e\{(x+z)(x-2y+z)\}$$

$$= \log_e\left[(x+z)\left(x+z - \frac{4xz}{x+z}\right)\right]$$

$$= \log_e[(x+z)^2 - 4xz]$$

$$= \log_e(x-z)^2$$

$$= 2 \log_e(x-z)$$
92. We have $\frac{a^{n+1} + b^{n+1}}{a^n + b^n} = \frac{2ab}{a+b}$
$$\Rightarrow a^{n+2} + ab^{n+1} + ba^{n+1} + b^{n+2} = 2a^{n+1}b + 2b^{n+1}a$$

$$\Rightarrow a^{n+1}(a-b) = b^{n+1}(a-b)$$

or $\left(\frac{a}{b}\right)^{n+1} = (1) = \left(\frac{a}{b}\right)^0$
Hence, $n = -1$



93. $f(x) = x + \frac{1}{2} = \frac{2x+1}{2}$
 $f(2x) = \frac{4x+1}{2}$
 $f(4x) = \frac{8x+1}{2}$
 $f(x), f(2x), f(4x)$ are in H.P.
 $\therefore f(2x) = \frac{2f(x)f(4x)}{f(x)+f(4x)}$
 $\Rightarrow x = 0, \frac{1}{4}$
 At $x = 0$, terms are equal, so only solution is
 $x = \frac{1}{4}$
94. Given $x_1.x_2.x_3 \dots x_n = 1$
 \therefore A.M. \geq G.M.
 $\therefore \left(\frac{x_1 + x_2 + x_3 + \dots + x_n}{n} \right) \geq (x_1.x_2.x_3 \dots x_n)^{\frac{1}{n}}$
 $= (1)^{\frac{1}{n}}$
 $= 1$
 $\therefore x_1 + x_2 + x_3 + \dots + x_n \geq n$
 $\therefore x_1 + x_2 + x_3 + \dots + x_n$ can never be less than n .
95. A.M. \geq G.M.
 $\Rightarrow \frac{a_1 + a_2 + \dots + a_{n-1} + 2a_n}{n}$
 $\geq (a_1.a_2 \dots a_{n-1}.2a_n)^{\frac{1}{n}} \geq (2c)^{\frac{1}{n}}$
 \therefore Minimum value of
 $a_1 + a_2 + \dots + a_{n-1} + 2a_n = n(2c)^{\frac{1}{n}}$
96. Since, p, q, r are in G.P.
 $\therefore q^2 = pr$
 Also, $\tan^{-1} p, \tan^{-1} q, \tan^{-1} r$ are in A.P.
 $\Rightarrow \tan^{-1} p + \tan^{-1} r = 2 \tan^{-1} q$
 $\Rightarrow p + r = 2q$
 $\Rightarrow p, q, r$ are in A.P.
 Here, p, q, r are both in A.P. and G.P.,
 which is possible only, if $p = q = r$.
97. $\frac{x+y}{2} = 3$
 $\Rightarrow x+y = 6$
 $xy = 1^2 = 1$
 $x^2 + y^2 = (x+y)^2 - 2xy$
 $= 36 - 2$
 $= 34$
98. Given that A.M. = 8 and G.M. = 5, if α, β are roots of quadratic equation, then quadratic equation is
 $x^2 - x(\alpha + \beta) + \alpha\beta = 0$
 A.M. = $\frac{\alpha + \beta}{2} = 8 \Rightarrow \alpha + \beta = 16$
 and G.M. = $\sqrt{\alpha\beta} = 5 \Rightarrow \alpha\beta = 25$
 So the required quadratic equation will be
 $x^2 - 16x + 25 = 0$.
99. $\frac{x+y}{2} = 9$
 $\Rightarrow x+y = 18$
 $xy = 4^2 = 16$
 x and y are the roots.
 \therefore The equation is
 $x^2 - 18x + 16 = 0$
100. GM = $\sqrt{ab} = 2$
 $\Rightarrow ab = 4$
 HM = $\frac{2ab}{a+b} = \frac{-8}{5}$
 $\Rightarrow \frac{8}{a+b} = \frac{-8}{5}$
 $\Rightarrow a+b = -5$
 $\Rightarrow 2a+2b = -10$
 $ab = 4 \Rightarrow (2a)(2b) = 16$
 \therefore The required quadratic equation is
 $x^2 - (2a+2b)x + (2a)(2b) = 0$
 $\Rightarrow x^2 + 10x + 16 = 0$
101. Sum of the roots of $x^2 - 2ax + b^2 = 0$ is $2a$,
 Therefore, A = A.M. of the roots = a
 Product of the roots of $x^2 - 2bx + a^2 = 0$ is a^2
 Therefore, G.M. of the roots is $G = a$
 Thus, $A = G$
102. We have, $\tan n\theta = \tan m\theta$
 $\Rightarrow n\theta = N\pi + (m\theta)$
 $\Rightarrow \theta = \frac{N\pi}{n-m}$, putting $N = 1, 2, 3, \dots$, we get
 $\frac{\pi}{n-m}, \frac{2\pi}{n-m}, \frac{3\pi}{n-m}, \dots$ which are in A.P.
 Since, common difference, $d = \frac{\pi}{n-m}$.
103. Let three numbers a, b and c in G.P., then
 $b^2 = ac$
 $\Rightarrow 2 \log_e b = \log_e a + \log_e c$ or
 $\log_e b = \frac{\log_e a + \log_e c}{2}$
 Thus, their logarithms are in A.P.



104. $225 = 3^2 \times 5^2 = d(225) = 3 \times 3 = 9$
 $1125 = 3^2 \times 5^3 = d(1125) = 3 \times 4 = 12$
 $640 = 2^7 \times 5 = d(640) = 8 \times 2 = 16$
 9, 12, 16 are in G.P.

105. If $\frac{x+y}{2}$, y , $\frac{y+z}{2}$ are in H.P., then

$$y = \frac{2\left(\frac{x+y}{2} \cdot \frac{y+z}{2}\right)}{\frac{x+y}{2} + \frac{y+z}{2}}$$

$$= \frac{\frac{2}{4}(x+y)(y+z)}{\frac{1}{2}(x+2y+z)}$$

$$y = \frac{xy + xz + y^2 + yz}{x + 2y + z}$$

$$\Rightarrow xy + 2y^2 + yz = xy + xz + y^2 + yz$$

$$\Rightarrow y^2 = xz$$

Thus, x, y, z will be in G.P.

106. $(y-x), 2(y-a), (y-z)$ are in H.P.

$$\Rightarrow \frac{1}{y-x}, \frac{1}{2(y-a)}, \frac{1}{y-z} \text{ are in A.P.}$$

$$\Rightarrow \frac{1}{2(y-a)} - \frac{1}{(y-x)} = \frac{1}{(y-z)} - \frac{1}{2(y-a)}$$

$$\Rightarrow \frac{y-x-2y+2a}{y-x} = \frac{2y-2a-y+z}{y-z}$$

$$\Rightarrow \frac{-x-y+2a}{(y-x)} = \frac{y+z-2a}{(y-z)}$$

$$\Rightarrow \frac{(x-a)+(y-a)}{(x-a)-(y-a)} = \frac{(y-a)+(z-a)}{(y-a)-(z-a)}$$

$$\Rightarrow \frac{(x-a)}{(y-a)} = \frac{(y-a)}{(z-a)}$$

$$\Rightarrow (x-a), (y-a), (z-a) \text{ are in G.P.}$$

107. $x, 1, z$ are in A.P., then $2 = x + z$ (i)
 and $4 = xz$ (ii)

Divide (ii) by (i), we get

$$\frac{x \cdot z}{x+z} = \frac{4}{2} \text{ or } \frac{2xz}{x+z} = 4$$

Hence, $x, 4, z$ will be in H.P.

108. Given, a, b, c are in G.P.

$$\Rightarrow \log_x a, \log_x b, \log_x c \text{ are in A.P.}$$

$$\Rightarrow \frac{\log a}{\log x}, \frac{\log b}{\log x}, \frac{\log c}{\log x} \text{ are in A.P.}$$

$$\Rightarrow \frac{\log x}{\log a}, \frac{\log x}{\log b}, \frac{\log x}{\log c} \text{ are in H.P.}$$

i.e., $\log_a x, \log_b x, \log_c x$ are in H.P.

109. x, y, z are in G.P.

$$\text{Hence, } y^2 = xz$$

$$\therefore 2 \log y = \log x + \log z$$

$$\Rightarrow 2(\log y + 1) = (1 + \log x) + (1 + \log z)$$

$$\Rightarrow 1 + \log x, 1 + \log y, 1 + \log z \text{ are in A.P.}$$

$$\Rightarrow \frac{1}{1 + \log x}, \frac{1}{1 + \log y}, \frac{1}{1 + \log z} \text{ are in H.P.}$$

110. Since, b^2, a^2, c^2 are in A.P.

$$\therefore a^2 - b^2 = c^2 - a^2$$

$$\Rightarrow (a-b)(a+b) = (c-a)(c+a)$$

$$\Rightarrow \frac{1}{b+c} - \frac{1}{a+b} = \frac{1}{c+a} - \frac{1}{b+c}$$

$$\Rightarrow \frac{1}{a+b}, \frac{1}{b+c}, \frac{1}{c+a} \text{ are in A.P.}$$

$$\therefore (a+b), (b+c), (c+a) \text{ are in H.P.}$$

111. Given, a, b, c are in A.P.

$$\Rightarrow 2b = a + c \Rightarrow b - c = a - b$$

Also, a^2, b^2, c^2 are in H.P.

$$\frac{1}{b^2} - \frac{1}{a^2} = \frac{1}{c^2} - \frac{1}{b^2}$$

$$\Rightarrow \frac{a^2 - b^2}{a^2 b^2} = \frac{b^2 - c^2}{b^2 c^2}$$

$$\Rightarrow (a-b)[c^2(a+b) - a^2(b+c)] = 0$$

$$\dots [\because (b-c) = (a-b)]$$

$$\Rightarrow a = b \text{ or } c^2 a + c^2 b - a^2 b - a^2 c = 0$$

$$\Rightarrow c^2 a + c^2 b - a^2 b - a^2 c = 0$$

$$\Rightarrow ac(c-a) = b(a^2 - c^2)$$

$$\Rightarrow ac = -b(c+a)$$

$$\Rightarrow -ac = b \cdot 2b$$

$$\Rightarrow b^2 = -\left(\frac{a}{2}\right)c$$

$$\therefore -\frac{a}{2}, b, c \text{ are in G.P.}$$

112. $x + y + z = 15$, if $9, x, y, z, a$ are in A.P.

$$\text{Sum} = 9 + 15 + a = \frac{5}{2}(9+a)$$

$$\Rightarrow 24 + a = \frac{5}{2}(9+a)$$

$$\Rightarrow 48 + 2a = 45 + 5a$$

$$\Rightarrow 3a = 3$$

$$\Rightarrow a = 1$$

$$\text{and } \frac{1}{x} + \frac{1}{y} + \frac{1}{z} = \frac{5}{3}, \text{ if } 9, x, y, z, a \text{ are in H.P.}$$

$$\text{Sum} = \frac{1}{9} + \frac{5}{3} + \frac{1}{a} = \frac{5}{2} \left[\frac{1}{9} + \frac{1}{a} \right] \Rightarrow a = 1$$



113. Given, a, b, c are in G.P.

$$\therefore b^2 = ac$$

$$x = \frac{a+b}{2}, y = \frac{b+c}{2}$$

$$\begin{aligned} \therefore \frac{a}{x} + \frac{c}{y} &= \frac{2a}{a+b} + \frac{2c}{b+c} \\ &= \frac{2(ab+bc+2ca)}{ab+ac+b^2+bc} \\ &= \frac{2(ab+bc+2ca)}{(ab+ac+ac+bc)} \\ &= 2 \quad \dots[\because b^2 = ac] \end{aligned}$$

114. Given that a, A_1, A_2, b are in A.P.

$$\text{Therefore, } A_1 = \frac{a+A_2}{2}, A_2 = \frac{A_1+b}{2}$$

$$\Rightarrow A_1 + A_2 = \frac{1}{2}(a+b+A_1+A_2)$$

$$\Rightarrow \frac{1}{2}(A_1 + A_2) = \frac{1}{2}(a+b) \text{ or}$$

$$A_1 + A_2 = a + b$$

and a, G_1, G_2, b are in G.P.

$$\text{Therefore, } G_1^2 = aG_2, G_2^2 = bG_1$$

$$\Rightarrow G_1^2 G_2^2 = abG_1 G_2 \Rightarrow G_1 G_2 = ab$$

$$\text{Hence, } \frac{A_1 + A_2}{G_1 G_2} = \frac{a+b}{ab}$$

Trick : Let $a = 1, b = 2$,
then $A_1 + A_2 = 1 + 2 = 3$
and $G_1 \cdot G_2 = 2 \times 1 = 2$

$$\therefore \frac{A_1 + A_2}{G_1 G_2} = \frac{3}{2}, \text{ which is given by (A).}$$

115. Given numbers a and 2 .

$$\text{A.M.} = \frac{a+2}{2} \text{ and G.M.} = \sqrt{2a}$$

According to the given condition,

$$\text{A.M.} - \text{G.M.} = 1$$

$$\Rightarrow \frac{a+2}{2} - \sqrt{2a} = 1$$

$$\Rightarrow \frac{a}{2} + 1 - 1 = \sqrt{2a}$$

$$\Rightarrow a = 2\sqrt{2a} \Rightarrow a^2 = 8a$$

$$\Rightarrow a(a-8) = 0$$

$$\Rightarrow a = 0 \text{ or } 8$$

Since, $a \neq 0$

$$\therefore a = 8$$

116. Let the two numbers be x, y .

$$\therefore x - y = 48 \quad \dots(i)$$

$$\text{and } \frac{x+y}{2} - \sqrt{xy} = 18$$

$$\Rightarrow x + y - 2\sqrt{xy} = 36$$

$$\Rightarrow 48 + y + y - 2\sqrt{(48+y)y} = 36 \dots[\text{From (i)}]$$

$$\Rightarrow 12 + 2y = 2\sqrt{y(48+y)}$$

$$\Rightarrow 6 + y = \sqrt{y(48+y)}$$

$$\Rightarrow 36 + y^2 + 12y = 48y + y^2$$

$$\Rightarrow 36y = 36 \Rightarrow y = 1$$

$$\therefore x = 48 + 1 = 49$$

117. Since, H_1, H_2 are two harmonic means between a and b .

$$\therefore \frac{1}{a}, \frac{1}{H_1}, \frac{1}{H_2}, \frac{1}{b} \text{ are in A.P.}$$

We know that $2A = a + b$ and $G^2 = ab$

$$\therefore 2 \times \frac{1}{H_1} = \frac{1}{a} + \frac{1}{H_2}$$

$$\text{Similarly, } 2 \times \frac{1}{H_2} = \frac{1}{b} + \frac{1}{H_1}$$

On adding and solving we get,

$$2\left(\frac{1}{H_1} + \frac{1}{H_2}\right) - \left(\frac{1}{H_1} + \frac{1}{H_2}\right) = \frac{1}{a} + \frac{1}{b}$$

$$\frac{1}{H_1} + \frac{1}{H_2} = \frac{a+b}{ab} = \frac{2A}{G^2}$$

118. Let a and b be two numbers.

Sum of n A.M.'s = $n \times$ single A.M.

$$\Rightarrow A_1 + A_2 = 2 \times \left(\frac{a+b}{2}\right) = a + b$$

Product of n G.M.'s = (Single G.M.) ^{n}

$$\Rightarrow G_1 \cdot G_2 = (\sqrt{ab})^2 = ab$$

$$\Rightarrow \frac{1}{a}, \frac{1}{H_1}, \frac{1}{H_2}, \frac{1}{b} \text{ are in A.P.}$$

$$\Rightarrow \frac{1}{H_1} + \frac{1}{H_2} = \frac{1}{a} + \frac{1}{b} = \frac{a+b}{ab}$$

$$\Rightarrow \frac{H_1 H_2}{H_1 + H_2} = \frac{G_1 G_2}{A_1 + A_2}$$

$$\Rightarrow \frac{G_1 G_2}{H_1 H_2} \times \frac{H_1 + H_2}{A_1 + A_2} = 1$$



119. Given, $\sqrt{ab} = 10$

$$\Rightarrow ab = 100 \text{ and } \frac{2ab}{a+b} = 8$$

$$\Rightarrow a + b = 25$$

$$\therefore a = 5, b = 20$$

120. Let the positive numbers be a_1 and a_2 .

$$a_1, A, a_2, \dots \text{ are in A.P. then } A = \frac{a_1 + a_2}{2}$$

Also, a_1, G, a_2, \dots are in G.P.

$$\therefore G = \sqrt{a_1 a_2}$$

$$\frac{1}{a_1}, \frac{1}{H}, \frac{1}{a_2}, \dots \text{ are in H.P.}$$

$$\therefore \frac{2}{H} = \frac{1}{a_1} + \frac{1}{a_2} \Rightarrow H = \frac{2a_1 a_2}{a_1 + a_2} \Rightarrow H = \frac{G^2}{A}$$

121. Given A.M. = 2(G.M.) or $\frac{1}{2}(a+b) = 2\sqrt{ab}$

$$\text{or } \frac{a+b}{2\sqrt{ab}} = \frac{2}{1}$$

$$\Rightarrow \frac{a+b+2\sqrt{ab}}{a+b-2\sqrt{ab}} = \frac{2+1}{2-1} = \frac{3}{1}$$

$$\Rightarrow \frac{(\sqrt{a} + \sqrt{b})^2}{(\sqrt{a} - \sqrt{b})^2} = \frac{3}{1}$$

$$\Rightarrow \frac{\sqrt{a} + \sqrt{b}}{\sqrt{a} - \sqrt{b}} = \frac{\sqrt{3}}{1} \Rightarrow \frac{a}{b} = \left(\frac{\sqrt{3} + 1}{\sqrt{3} - 1} \right)^2$$

$$\Rightarrow \frac{a}{b} = \frac{2 + \sqrt{3}}{2 - \sqrt{3}} \text{ or } a : b = (2 + \sqrt{3}) : (2 - \sqrt{3})$$

122. We have H.M. = $\frac{2ab}{a+b}$ and G.M. = \sqrt{ab}

$$\text{So } \frac{\text{H.M.}}{\text{G.M.}} = \frac{4}{5} \Rightarrow \frac{2ab / (a+b)}{\sqrt{ab}} = \frac{4}{5}$$

$$\Rightarrow \frac{2\sqrt{ab}}{a+b} = \frac{4}{5} \Rightarrow \frac{a+b}{2\sqrt{ab}} = \frac{5}{4}$$

$$\Rightarrow \frac{a+b+2\sqrt{ab}}{a+b-2\sqrt{ab}} = \frac{5+4}{5-4}$$

$$\Rightarrow \frac{(\sqrt{a} + \sqrt{b})^2}{(\sqrt{a} - \sqrt{b})^2} = \frac{9}{1} \Rightarrow \frac{\sqrt{a} + \sqrt{b}}{\sqrt{a} - \sqrt{b}} = \frac{3}{1}$$

$$\Rightarrow \frac{(\sqrt{a} + \sqrt{b}) + (\sqrt{a} - \sqrt{b})}{(\sqrt{a} + \sqrt{b}) - (\sqrt{a} - \sqrt{b})} = \frac{3+1}{3-1}$$

$$\Rightarrow \frac{2\sqrt{a}}{2\sqrt{b}} = \frac{4}{2} \Rightarrow \left(\frac{a}{b} \right) = 2^2 = 4$$

$$\Rightarrow a : b = 4 : 1 \text{ or } b : a = 1 : 4$$

123. According to the given condition,

$$\frac{x+y}{2} = \frac{p}{\sqrt{xy}}$$

$$\Rightarrow \frac{x+y}{2(\sqrt{xy})} = \frac{p}{q} \quad \dots \text{(i)}$$

$$\Rightarrow \frac{x^2 + y^2 + 2xy}{4xy} = \frac{p^2}{q^2}$$

$$\Rightarrow \frac{x^2 + y^2 + 2xy - 4xy}{4xy} = \frac{p^2 - q^2}{q^2}$$

$$\Rightarrow \frac{(x-y)^2}{4xy} = \frac{p^2 - q^2}{q^2}$$

$$\Rightarrow \frac{x-y}{2\sqrt{xy}} = \frac{\sqrt{p^2 - q^2}}{q} \quad \dots \text{(ii)}$$

Dividing (ii) by (i), we get

$$\frac{x+y}{x-y} = \frac{p}{\sqrt{p^2 - q^2}} \Rightarrow \frac{x}{y} = \frac{p + \sqrt{p^2 - q^2}}{p - \sqrt{p^2 - q^2}}$$

$$124. \frac{1}{t_1} + \frac{1}{t_2} + \frac{1}{t_3} + \dots + \frac{1}{t_{2003}}$$

$$= 4 \left[\frac{1}{3.4} + \frac{1}{4.5} + \frac{1}{5.6} + \dots + \frac{1}{(2005)(2006)} \right]$$

$$= 4 \left[\frac{1}{3} - \frac{1}{4} + \frac{1}{4} - \frac{1}{5} + \dots + \frac{1}{2005} - \frac{1}{2006} \right]$$

$$= 4 \left[\frac{1}{3} - \frac{1}{2006} \right]$$

$$= 4 \cdot \frac{2003}{3(2006)}$$

$$= \frac{4006}{3009}$$

125. It is an arithmetico-geometric series

$$\therefore S_{\infty} = \frac{a}{1-r} + \frac{dr}{(1-r)^2} = \frac{1}{1-\frac{1}{2}} + \frac{2 \times \frac{1}{2}}{\left(1-\frac{1}{2}\right)^2}$$

$$= 2 + 4 = 6$$

$$126. S_{\infty} = \frac{a}{1-r} + \frac{dr}{(1-r)^2}$$

$$\text{Here, } a = 1, r = \frac{1}{5}, d = 3$$

$$\therefore S_{\infty} = \frac{1}{1-\frac{1}{5}} + \frac{3 \times \frac{1}{5}}{\left(1-\frac{1}{5}\right)^2} = \frac{35}{16}$$



127. Let $S = 1 + \frac{2}{3} + \frac{6}{3^2} + \frac{10}{3^3} + \frac{14}{3^4} + \dots$ to ∞
 $\Rightarrow (S - 1) = \frac{2}{3} + \frac{6}{3^2} + \frac{10}{3^3} + \frac{14}{3^4} + \dots$ to ∞ (i)
 $\Rightarrow (S - 1) \frac{1}{3} = \frac{2}{3^2} + \frac{6}{3^3} + \frac{10}{3^4} + \dots$ to ∞ (ii)
 Subtracting (ii) from (i), we get
 $\frac{2}{3}(S - 1) = \frac{2}{3} + \frac{4}{3^2} + \frac{4}{3^3} + \frac{4}{3^4} + \dots$ to ∞
 $\Rightarrow \frac{2}{3}(S - 1) = \frac{2}{3} + \frac{4}{3^2} \frac{1}{1 - \frac{1}{3}}$
 $\Rightarrow \frac{2}{3}(S - 1) = \frac{2}{3} + \frac{2}{3} \Rightarrow S = 3$

128. $\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots = \frac{\pi^4}{90}$
 $\therefore \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots + \frac{1}{2^4} \left(\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \dots \right)$
 $= \frac{\pi^4}{90}$
 $\therefore \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \frac{1}{7^4} + \dots + \frac{1}{16} \times \frac{\pi^4}{90} = \frac{\pi^4}{90}$
 $\therefore \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \frac{1}{7^4} + \dots + \infty$
 $= \frac{\pi^4}{90} - \frac{1}{16} \left(\frac{\pi^4}{90} \right)$
 $= \frac{15}{16} \left(\frac{\pi^4}{90} \right) = \frac{\pi^4}{96}$

129. The sequence can be written as $\log a$, $(2 \log a - \log b)$, $(3 \log a - 2 \log b)$, which are in A.P. having common difference as $\log a - \log b$.

130. $\sum_{k=1}^n k(k+2) = \sum_{k=1}^n k^2 + 2 \sum_{k=1}^n k$
 $= \frac{n(n+1)(2n+1)}{6} + \frac{2n(n+1)}{2}$
 $= \frac{n(n+1)}{6} [2n+1+6]$
 $= \frac{n(n+1)(2n+7)}{6}$

131. Given series $\frac{1}{1} + \frac{1+2}{2} + \frac{1+2+3}{3} + \dots$
 So, n^{th} term of series is given by

$$t_n = \frac{1+2+3+\dots+n}{n}$$

$$= \frac{\frac{1}{2}n(n+1)}{n} = \frac{n+1}{2}$$

132. Here $t_n = \frac{n(n+1)}{2}$
 $\therefore S_n = \frac{1}{2}(\sum n^2 + \sum n) = \frac{n(n+1)(n+2)}{6}$

133. $t_n = \frac{(2n+1)}{n(n+1)(2n+1)}$
 $= \frac{6}{n(n+1)}$

$$S_n = \sum (t_n)$$

$$= \sum 6 \left[\frac{1}{n} - \frac{1}{n+1} \right]$$

$$= 6 \left[1 - \frac{1}{n+1} \right]$$

$$S_n = \frac{6n}{n+1}$$

134. General term $t_n = \frac{1^2 + 2^2 + \dots + n^2}{1 + 2 + \dots + n}$
 $= \frac{n(n+1)(2n+1)}{n(n+1)}$
 $\Rightarrow t_n = \frac{6}{n(n+1)} = \frac{1}{3} \cdot (2n+1)$

$$\therefore \sum t_n = \frac{2}{3} \sum n + \frac{1}{3} n$$

$$= \frac{2}{3} \cdot \frac{n(n+1)}{2} + \frac{1}{3} n$$

$$= \frac{1}{3} n \cdot (n+1) + \frac{1}{3} n$$

$$= \frac{n(n+2)}{3}$$

135. Mean, $\bar{x} = \frac{1 \cdot (1) + 2 \cdot (2) + 3 \cdot (3) + \dots + n \cdot (n)}{1 + 2 + 3 + \dots + n}$
 $= \frac{n(n+1)(2n+1)}{6}$
 $= \frac{n(n+1)}{2}$
 $= \frac{2n+1}{3}$



$$\begin{aligned}
 136. \quad S &= 3.6 + 4.7 + \dots \text{ upto } n - 2 \text{ terms} \\
 &= (1.4 + 2.5 + 3.6 + 4.7 + \dots \text{ upto } n \text{ terms}) - 14 \\
 &= \Sigma n(n+3) - 14 \\
 &= \frac{1}{6} (2n^3 + 12n^2 + 10n) - 14 \\
 &= \left(\frac{2n^3 + 12n^2 + 10n - 84}{6} \right),
 \end{aligned}$$

where $n = 3, 4, 5, \dots$

Trick : $S_1 = 18, S_2 = 46$

Now put in options $(n - 2) = 1, 2$ i.e. $n = 3, 4$

Option (B) gives the values.

$$\begin{aligned}
 137. \quad &\left(\frac{8}{5}\right)^2 + \left(\frac{12}{5}\right)^2 + \left(\frac{16}{5}\right)^2 + \dots \\
 &= \frac{1}{5^2} [8^2 + 12^2 + 16^2 + \dots] \\
 &= \frac{1}{5^2} [(4 \times 2)^2 + (4 \times 3)^2 + (4 \times 4)^2 + \dots] \\
 &= \frac{4^2}{5^2} [2^2 + 3^2 + 4^2 + \dots + 11^2] \\
 &= \frac{4^2}{5^2} [1^2 + 2^2 + \dots + 11^2 - 1^2] \\
 &= \frac{4^2}{5^2} \left[\frac{11(12)(23)}{6} - 1 \right] = \frac{16}{5} m
 \end{aligned}$$

$$\therefore m = \frac{1}{5} \left[\frac{3036 - 6}{6} \right] = \frac{3030}{30}$$

$$\therefore m = 101$$

$$138. \quad \sum_{k=0}^{12} a_{4k+1} = 416$$

$$\Rightarrow a_1 + a_5 + a_9 + \dots + a_{49} = 416$$

$$\Rightarrow \frac{13}{2} (2a_1 + 48d) = 416$$

$$\Rightarrow a_1 + 24d = 32 \quad \dots \text{(i)}$$

$$a_9 + a_{43} = 66$$

$$\Rightarrow 2a_1 + 50d = 66$$

$$\Rightarrow a_1 + 25d = 33 \quad \dots \text{(ii)}$$

Solving (i) and (ii), we get

$$a_1 = 8 \text{ and } d = 1$$

$$a_1^2 + a_2^2 + \dots + a_{17}^2 = 140m$$

$$\Rightarrow 8^2 + 9^2 + \dots + 24^2 = 140m$$

$$\Rightarrow (1^2 + 2^2 + \dots + 24^2)$$

$$- (1^2 + 2^2 + \dots + 7^2) = 140m$$

$$\Rightarrow \frac{24 \times 25 \times 49}{6} - \frac{7 \times 8 \times 15}{6} = 140m$$

$$\Rightarrow 4900 - 140 = 140m$$

$$\Rightarrow 140m = 4760$$

$$\Rightarrow m = 34$$

$$\begin{aligned}
 139. \quad S_n &= cn^2 \\
 S_{n-1} &= c(n-1)^2 = cn^2 + c - 2cn \\
 T_n &= 2cn - c \\
 T_n^2 &= (2cn - c)^2 = 4c^2 n^2 + c^2 - 4c^2 n \\
 \therefore \text{ Required sum} &= \sum T_n^2 \\
 &= \frac{4c^2 \cdot n(n+1)(2n+1)}{6} + nc^2 - 2c^2 n(n+1) \\
 &= \frac{2c^2 n(n+1)(2n+1) + 3nc^2 - 6c^2 n(n+1)}{3} \\
 &= \frac{nc^2 (4n^2 + 6n + 2 + 3 - 6n - 6)}{3} \\
 &= \frac{nc^2 (4n^2 - 1)}{3}
 \end{aligned}$$

$$\begin{aligned}
 140. \quad \sum_{i=1}^n \sum_{j=1}^i \sum_{k=1}^j 1 &= \sum_{i=1}^n \sum_{j=1}^i j \\
 &= \sum_{i=1}^n \left(\frac{i(i+1)}{2} \right) \\
 &= \frac{1}{2} \left[\sum_{i=1}^n i^2 + \sum_{i=1}^n i \right] \\
 &= \frac{1}{2} \left[\frac{n(n+1)(2n+1)}{6} + \frac{n(n+1)}{2} \right] \\
 &= \frac{1}{4} n(n+1) \left[\frac{2n+1}{3} + 1 \right] \\
 &= \frac{n(n+1)}{4} \left[\frac{2n+1+3}{3} \right] = \frac{n(n+1)(n+2)}{6}
 \end{aligned}$$

141. We have $S = 2 + 4 + 7 + 11 + 16 + \dots + t_n$
Again, $S = 2 + 4 + 7 + 11 + \dots + t_{n-1} + t_n$
Subtracting, we get

$$0 = 2 + \{2 + 3 + 4 + 5 + \dots + (t_n - t_{n-1})\} - t_n$$

$$t_n = 2 + \frac{1}{2} (n-1) \{(4 + (n-2)1)\}$$

$$= \frac{1}{2} (n^2 + n + 2)$$

Now,

$$S = \Sigma t_n = \frac{1}{2} \Sigma (n^2 + n + 2)$$

$$= \frac{1}{2} (\Sigma n^2 + \Sigma n + 2\Sigma 1)$$

$$= \frac{1}{2} \left\{ \frac{1}{6} n(n+1)(2n+1) + \frac{1}{2} n(n+1) + 2n \right\}$$

$$= \frac{n}{12} \{(n+1)(2n+1+3) + 12\}$$

$$= \frac{n}{6} \{(n+1)(n+2) + 6\} = \frac{n}{6} (n^2 + 3n + 8)$$



142. Sum of cubes of 'n' natural number

$$= \frac{n^2(n+1)^2}{4}$$

$$= \frac{15^2(16)^2}{4} = 14,400$$

143. $t_n = n(n+1)(n+2) = n(n^2 + 3n + 2)$
 $= n^3 + 3n^2 + 2n$ $\therefore S_n = \Sigma(n^3) + \Sigma(3n^2) + \Sigma(2n)$

$$S_n = \left[\frac{n(n+1)}{2} \right]^2 + \frac{3.n(n+1)(2n+1)}{6}$$

$$+ \frac{2.n(n+1)}{2}$$

$$S_n = \frac{1}{4} n(n+1)(n+2)(n+3)$$

144. Here, t_n of the A.P. 1, 2, 3, = n
and t_n of the A.P. 3, 5, 7, = 2n + 1 $\therefore t_n$ of given series = $n(2n + 1)^2 = 4n^3 + 4n^2 + n$

Hence,

$$S = \sum_1^{20} t_n = 4 \sum_1^{20} n^3 + 4 \sum_1^{20} n^2 + \sum_1^{20} n$$

$$= 4 \cdot \frac{1}{4} 20^2 \cdot 21^2 + 4 \cdot \frac{1}{6} 20 \cdot 21 \cdot 41 + \frac{1}{2} 20 \cdot 21$$

$$= 188090$$

145. $1^3 + 3^3 + 5^3 + 7^3 + \dots = \Sigma(2n-1)^3$

$$= \Sigma(8n^3 - 3 \cdot 4n^2 + 3 \cdot 2n - 1)$$

$$= 2n^2(n+1)^2 - 2n(n+1)(2n+1) + 3n(n+1) - n$$

$$= 2n^4 + 4n^3 + 2n^2 - 2n(2n^2 + 3n + 1)$$

$$+ 3n^2 + 3n - n$$

$$= 2n^4 + 4n^3 + 2n^2 - 4n^3 - 6n^2 - 2n$$

$$+ 3n^2 + 3n - n$$

$$= 2n^4 - n^2 = n^2(2n^2 - 1)$$

146. $(n^2 - 1^2) + 2(n^2 - 2^2) + \dots$

$$= n^2(1 + 2 + 3 + \dots) - (1 \cdot 1^2 + 2 \cdot 2^2 + 3 \cdot 3^2 + \dots)$$

$$= n^2 \sum_{r=1}^n r - \sum_{r=1}^n r^3$$

$$= \frac{n^3(n+1)}{2} - \left[\frac{n(n+1)}{2} \right]^2$$

$$= \frac{n^2(n+1)}{2} \left(\frac{n-1}{2} \right) = \frac{1}{4} n^2(n^2 - 1)$$

$$147. S_n = 1(1!) + 2(2!) + 3(3!) + \dots + n(n!) = (2-1)(1!) + (3-1)(2!) + (4-1)(3!) + \dots + [(n+1)-1](n!) = (2 \cdot 1! - 1!) + (3 \cdot 2! - 2!) + (4 \cdot 3! - 3!) + \dots + [(n+1)! - (n)!] = (2! - 1!) + (3 \cdot 2! - 2!) + (4 \cdot 3! - 3!) + \dots + [(n+1)(n!) - (n)!] = (n+1)! - 1!$$

$$148. 1^2 - 2^2 + 3^2 - 4^2 + \dots + 11^2 = (1^2 - 2^2) + (3^2 - 4^2) + \dots + (9^2 - 10^2) + 11^2$$

Now, $a^2 - b^2 = (a-b)(a+b)$

$$\therefore 1^2 - 2^2 + 3^2 - 4^2 + \dots + 11^2 = (1-2)(1+2) + (3-4)(3+4) + \dots + (9-10)(9+10) + 11^2 = (-1)[1+2+3+\dots+9+10] + 11^2 = (-1) \cdot \frac{10 \times 11}{2} + 11^2 = 66$$

149. G.M. of 1, 2, 2², 2³, ..., 2ⁿ
Here, no. of terms = (n + 1)

$$\therefore \text{G.M.} = (1 \cdot 2 \cdot 2^2 \cdot 2^3 \dots 2^n)^{\frac{1}{(n+1)}}$$

$$= (2^{0+1+2+\dots+n})^{\frac{1}{(n+1)}} = \left[2^{\frac{n(n+1)}{2}} \right]^{\frac{1}{(n+1)}}$$

$$\therefore \text{G.M.} = 2^{\frac{n}{2}}$$

$$150. 1 + 3 + 7 + \dots + t_n = 2 - 1 + 2^2 - 1 + 2^3 - 1 + \dots + 2^n - 1 = (2 + 2^2 + \dots + 2^n) - n = 2^{n+1} - 2 - n$$

151. Let n^{th} term of series is t_n , then

$$S_n = 12 + 16 + 24 + 40 + \dots + t_n$$

$$\text{Again } S_n = 12 + 16 + 24 + \dots + t_n$$

On subtraction

$$0 = (12 + 4 + 8 + 16 + \dots + \text{upto } n \text{ terms}) - t_n$$

$$\Rightarrow t_n = 12 + [4 + 8 + 16 + \dots + \text{upto } (n-1) \text{ terms}]$$

$$= 12 + \frac{4(2^{n-1} - 1)}{2-1} = 2^{n+1} + 8$$

On putting $n = 1, 2, 3, \dots$

$$t_1 = 2^2 + 8, t_2 = 2^3 + 8, t_3 = 2^4 + 8 \dots \text{etc.}$$

$$S_n = t_1 + t_2 + t_3 + \dots + t_n$$

$$= (2^2 + 2^3 + 2^4 + \dots + \text{upto } n \text{ terms})$$

$$+ (8 + 8 + 8 + \dots + \text{upto } n \text{ terms})$$

$$= \frac{2^2(2^n - 1)}{2-1} + 8n = 4(2^n - 1) + 8n$$



152. $(1 + 2) + (1 + 2 + 2^2) + \dots$ upto n terms

$$\therefore T_n = 1 + 2 + 2^2 + \dots + 2^n$$

$$\therefore T_n = \frac{1(2^{n+1} - 1)}{2 - 1} = 2^{n+1} - 1$$

$$\therefore S_n = \sum T_n = \sum (2^{n+1} - 1)$$

$$\begin{aligned} \therefore S_n &= \sum 2^{n+1} - \sum 1 \\ &= 2^2 + 2^3 + 2^4 + \dots + 2^n - (n) \\ &= 2^{n+2} - 4 - n \end{aligned}$$

153. $S = \frac{1}{2} \left(\frac{1}{2} + \frac{1}{3} \right) - \frac{1}{4} \left(\frac{1}{2^2} + \frac{1}{3^2} \right) + \frac{1}{6} \left(\frac{1}{2^3} + \frac{1}{3^3} \right) - \dots$

$$\begin{aligned} &= \frac{1}{2} \left(\frac{1}{2} - \frac{1}{2} \cdot \frac{1}{2^2} + \frac{1}{3} \cdot \frac{1}{2^3} - \dots \right) \\ &\quad + \frac{1}{2} \left(\frac{1}{3} - \frac{1}{2} \cdot \frac{1}{3^2} + \frac{1}{3} \cdot \frac{1}{3^3} - \dots \right) \end{aligned}$$

$$\begin{aligned} &= \frac{1}{2} \left[\frac{1}{2} - \frac{1}{2} \left(\frac{1}{2} \right)^2 + \frac{1}{3} \left(\frac{1}{2} \right)^3 - \dots \right] \\ &\quad + \frac{1}{2} \left[\frac{1}{3} + \frac{1}{2} \left(\frac{1}{3} \right)^2 + \frac{1}{3} \left(\frac{1}{3} \right)^3 - \dots \right] \end{aligned}$$

$$= \frac{1}{2} \log \left(1 + \frac{1}{2} \right) + \frac{1}{2} \log \left(1 + \frac{1}{3} \right)$$

$$= \frac{1}{2} \left[\log \frac{3}{2} + \log \frac{4}{3} \right]$$

$$= \frac{1}{2} \log \left(\frac{3}{2} \times \frac{4}{3} \right)$$

$$= \frac{1}{2} \log 2$$

154. Let, $S = 2 + 7 + 14 + 23 + 34 + \dots + t_n + \dots$ (i)

and $S = 2 + 7 + 14 + \dots + t_{n-1} + t_n + \dots$ (ii)

From (i) and (ii), we get

$$0 = 2 + [5 + 7 + 9 + 11 \dots + t_n - t_{n-1}] - t_n$$

$$\Rightarrow t_n = 2 + \left[\frac{n-1}{2} \{ 2 \times 5 + (n-2)2 \} \right]$$

$$\Rightarrow t_n = 2 + (n-1)(n+3)$$

Now,

put $n = 99$

$$\Rightarrow t_{99} = 2 + 98 \times 102 = 9998$$

155. $A = 1^2 + 2.2^2 + 3^2 + 2.4^2 + \dots + 2.20^2$
 $= (1^2 + 2^2 + 3^2 + \dots + 20^2)$
 $\quad + (2^2 + 4^2 + \dots + 20^2)$
 $= (1^2 + 2^2 + 3^2 + \dots + 20^2)$

$$\begin{aligned} &\quad + 4(1^2 + 2^2 + \dots + 10^2) \\ &= \frac{20 \times 21 \times 41}{6} + 4 \left(\frac{10 \times 11 \times 21}{6} \right) \\ &= 2870 + 4(385) \\ &= 4410 \end{aligned}$$

$B = 1^2 + 2.2^2 + 3^2 + 2.4^2 + \dots + 2.40^2$
 $= (1^2 + 2^2 + 3^2 + \dots + 40^2)$
 $\quad + (2^2 + 4^2 + \dots + 40^2)$
 $= (1^2 + 2^2 + 3^2 + \dots + 40^2)$

$$\begin{aligned} &\quad + 4(1^2 + 2^2 + \dots + 20^2) \\ &= \frac{40 \times 41 \times 81}{6} + 4 \left(\frac{20 \times 21 \times 41}{6} \right) \\ &= 22140 + 4(2870) \\ &= 33620 \end{aligned}$$

$$\begin{aligned} B - 2A &= 33620 - 2(4410) = 24800 \\ \Rightarrow 100 \lambda &= 24800 \Rightarrow \lambda = 248 \end{aligned}$$

156. Since, $\sin \theta$, $\cos \theta$ and $\tan \theta$ are in G.P.

$$\therefore \frac{\cos \theta}{\sin \theta} = \frac{\tan \theta}{\cos \theta} = \frac{\sin \theta}{\cos^2 \theta}$$

$$\Rightarrow \cos^3 \theta = \sin^2 \theta \quad \dots (i)$$

$$\therefore \cot^6 \theta - \cot^2 \theta = \frac{\cos^6 \theta}{\sin^6 \theta} - \frac{\cos^2 \theta}{\sin^2 \theta}$$

$$= \frac{\cos^6 \theta}{\cos^9 \theta} - \frac{\cos^2 \theta}{\cos^3 \theta} \quad \dots [\text{From (i)}]$$

$$= \frac{1}{\cos^3 \theta} - \frac{1}{\cos \theta} = \frac{1 - \cos^2 \theta}{\cos^3 \theta} = \frac{\sin^2 \theta}{\cos^3 \theta}$$

$$= 1 \quad \dots [\text{From (i)}]$$

157. Let $1 - \cos \theta = x$

\therefore the given series $= 1 + 2x + 3x^2 + 4x^3 + \dots \infty$

$$= (1 - x)^{-2}$$

$$= (1 - 1 + \cos \theta)^{-2} = \sec^2 \theta$$

$$= 1 + \tan^2 \theta = 1 + \frac{3}{2} = \frac{5}{2} \quad \dots \left[\because \tan \theta = \sqrt{\frac{3}{2}} \right]$$



158. $\cos(\theta - \alpha)$, $\cos \theta$, $\cos(\theta + \alpha)$ are in HP

$\therefore \frac{1}{\cos(\theta - \alpha)}, \frac{1}{\cos \theta}, \frac{1}{\cos(\theta + \alpha)}$ are in AP

$\therefore \frac{1}{\cos(\theta - \alpha)} + \frac{1}{\cos(\theta + \alpha)} = \frac{2}{\cos \theta}$

$$\Rightarrow \frac{\cos(\theta + \alpha) + \cos(\theta - \alpha)}{\cos(\theta + \alpha)\cos(\theta - \alpha)} = \frac{2}{\cos \theta}$$

$$\Rightarrow 2\cos \theta \cos \alpha \cdot \cos \theta = 2\cos(\theta + \alpha)\cos(\theta - \alpha)$$

$$\Rightarrow \cos^2 \theta \cos \alpha = \cos^2 \theta - \sin^2 \alpha$$

$$\Rightarrow \cos^2 \theta (\cos \alpha - 1) = -(1 - \cos^2 \alpha)$$

$$\Rightarrow \cos^2 \theta - 1 + \cos \alpha$$

159. $\sum_{k=1}^n f(a+k) = f(a+1) + f(a+2) + f(a+3) + \dots + f(a+n)$

$$= f(a)f(1) + f(a)f(2) + f(a)f(3) + \dots + f(a)f(n)$$

$$\dots [\because f(x+y) = f(x)f(y)]$$

$$\therefore \sum_{k=1}^n f(a+k) = f(a)[f(1) + f(2) + f(3) + \dots + f(n)] \quad \dots(i)$$

$$\therefore f(1) = 2$$

$$\therefore f(2) = f(1+1) = f(1) \cdot f(1) = 4$$

$$\therefore f(3) = f(2+1) = f(2) \cdot f(1) = 8$$

and so on.

\therefore substituting above values in (i), we get

$$\sum_{k=1}^n f(a+k) = f(a)[2 + 4 + 8 + \dots + f(n)]$$

$$= f(a) \cdot 2(2^n - 1)$$

$$\therefore f(a) \cdot 2(2^n - 1) = 16 \cdot (2^n - 1)$$

$$\therefore f(a) = 8$$

Since, $f(3) = 8$

$$\therefore a = 3$$



Evaluation Test

1. $a_1, a_2, a_3, \dots, a_n$ are in A.P. with common difference = 5

$$\text{i.e., } a_2 - a_1 = a_3 - a_2 = a_4 - a_3 = \dots = a_n - a_{n-1} = 5$$

$$\therefore \tan^{-1}\left(\frac{5}{1+a_1a_2}\right) + \tan^{-1}\left(\frac{5}{1+a_2a_3}\right) + \dots + \tan^{-1}\left(\frac{5}{1+a_{n-1}a_n}\right)$$

$$= \tan^{-1}\left(\frac{a_2 - a_1}{1+a_1a_2}\right) + \tan^{-1}\left(\frac{a_3 - a_2}{1+a_2a_3}\right) + \dots + \tan^{-1}\left(\frac{a_n - a_{n-1}}{1+a_{n-1}a_n}\right)$$

$$= \tan^{-1}(a_2) - \tan^{-1}(a_1) + \tan^{-1}(a_3) - \tan^{-1}(a_2) + \dots + \tan^{-1}(a_n) - \tan^{-1}(a_{n-1})$$

$$= \tan^{-1}(a_n) - \tan^{-1}(a_1) = \tan^{-1}\left(\frac{a_n - a_1}{1+a_1a_n}\right)$$

$$= \tan^{-1}\left(\frac{(n-1)5}{1+a_1a_n}\right) \quad \dots [\because a_n = a_1 + (n-1)5]$$

$$= \tan^{-1}\left(\frac{5n-5}{1+a_1a_n}\right)$$

2. Since, a_1, a_2, a_3, \dots are in H.P.

$$\therefore \frac{1}{a_1}, \frac{1}{a_2}, \frac{1}{a_3}, \dots \text{ are in A.P.}$$

$$\therefore \frac{1}{25} = \frac{1}{5} + 19d$$

$$\Rightarrow d = \frac{1}{19}\left(\frac{-4}{25}\right) = -\frac{4}{19 \times 25}$$

Since, $a_n < 0$

$$\therefore \frac{1}{5} - \frac{4}{19 \times 25} \times (n-1) < 0$$

$$\Rightarrow \frac{19 \times 5}{4} < n - 1$$

$$\Rightarrow n > 24.75$$

3. p, q, r are positive and are in A.P.

$$\therefore q = \frac{p+r}{2} \quad \dots(i)$$

Since, the roots of $px^2 + qx + r = 0$ are real

$$\therefore q^2 \geq 4pr \Rightarrow \left[\frac{p+r}{2}\right]^2 \geq 4pr \quad \dots[\text{From (i)}]$$

$$\Rightarrow p^2 + r^2 - 14pr \geq 0$$



$$\Rightarrow \left(\frac{r}{p}\right)^2 - 14\left(\frac{r}{p}\right) + 1 \geq 0$$

....[: p > 0 and p ≠ 0]

$$\Rightarrow \left(\frac{r}{p} - 7\right)^2 - 48 \geq 0$$

$$\Rightarrow \left(\frac{r}{p} - 7\right)^2 - (4\sqrt{3})^2 \geq 0$$

$$\Rightarrow \left|\frac{r}{p} - 7\right| \geq 4\sqrt{3}$$

4. Let A be the first term and R be the common ratio of the G.P. Then,

$$a = AR^{p-1}$$

$$\Rightarrow \log a = \log A + (p-1) \log R \quad \dots(i)$$

$$b = AR^{q-1}$$

$$\Rightarrow \log b = \log A + (q-1) \log R \quad \dots(ii)$$

$$c = AR^{r-1}$$

$$\Rightarrow \log c = \log A + (r-1) \log R \quad \dots(iii)$$

Multiplying (i), (ii) and (iii) by (q-r), (r-p) and (p-q) respectively and adding, we get

$$(q-r) \log a + (r-p) \log b + (p-q) \log c = 0$$

Expanding along first row, we get

$$\Delta = (q-r) \log a + (r-p) \log b + (p-q) \log c$$

$$\Rightarrow \Delta = 0$$

5. Let x_1, x_2, x_3 be a, ar, ar^2 and y_1, y_2, y_3 be b, br, br^2 .

∴ A, B, C are (a,b), (ar, br), (ar^2, br^2) resp.

$$\text{Now Slope of AB} = \frac{b}{a} = \text{slope of BC.}$$

Hence, the points are collinear.

i.e., lie on a straight line

Alternate method:

$$\text{Given } x_2 = rx_1, x_3 = r^2x_1, y_2 = ry_1, y_3 = r^2y_1$$

$$\text{Area of } \Delta = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$$

$$= \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ rx_1 & ry_1 & 1 \\ r^2x_1 & r^2y_1 & 1 \end{vmatrix}$$

$$= \frac{1}{2} x_1 y_1 \begin{vmatrix} 1 & 1 & 1 \\ r & r & 1 \\ r^2 & r^2 & 1 \end{vmatrix} = 0$$

i.e., lie on a straight line.

6. We have,

Length of a side of S_n

$$= \text{Length of a diagonal of } S_{n+1}$$

⇒ Length of a side of S_n

$$= \sqrt{2} (\text{Length of a side of } S_{n+1})$$

$$\Rightarrow \frac{\text{Length of a side of } S_{n+1}}{\text{Length of a side of } S_n} = \frac{1}{\sqrt{2}} \text{ for all } n \geq 1$$

- ∴ sides of S_1, S_2, \dots, S_n form a G.P. with common ratio $\frac{1}{\sqrt{2}}$ and first term 10.

$$\therefore \text{length of the side of } S_n = 10 \left(\frac{1}{\sqrt{2}}\right)^{n-1}$$

$$= \frac{10}{2^{\frac{n-1}{2}}}$$

$$\text{Now, area of } S_n = (\text{side})^2 = \left(\frac{10}{2^{\frac{n-1}{2}}}\right)^2$$

$$= \frac{100}{2^{n-1}}$$

But, area of $S_n < 1$

$$\Rightarrow \frac{100}{2^{n-1}} < 1$$

$$\Rightarrow 2^{n-1} > 100$$

The above inequality is satisfied if $n-1 \geq 7$ i.e., $n \geq 8$

7. We have, $T_p = a + (p-1)d = \frac{1}{q} \quad \dots(i)$

$$\text{and } T_q = a + (q-1)d = \frac{1}{p} \quad \dots(ii)$$

From (i) and (ii), we get $a = \frac{1}{pq}$ and $d = \frac{1}{pq}$

$$\begin{aligned} \therefore \text{sum of } (pq)^{\text{th}} \text{ terms} &= \frac{pq}{2} \left[\frac{2}{pq} + (pq-1) \frac{1}{pq} \right] \\ &= \frac{pq}{2} \cdot \frac{2}{pq} \left[1 + \frac{1}{2}(pq-1) \right] \\ &= \frac{2+pq-1}{2} = \frac{pq+1}{2} \end{aligned}$$



$$8. \begin{vmatrix} x+1 & x+2 & x+a \\ x+2 & x+3 & x+b \\ x+3 & x+4 & x+c \end{vmatrix}$$

Applying $C_1 \rightarrow C_1 - C_2$,

$$= \begin{vmatrix} -1 & x+2 & x+a \\ -1 & x+3 & x+b \\ -1 & x+4 & x+c \end{vmatrix}$$

$$= (-1) \begin{vmatrix} 1 & x+2 & x+a \\ 1 & x+3 & x+b \\ 1 & x+4 & x+c \end{vmatrix}$$

Applying $R_2 \rightarrow R_2 - R_1$, $R_3 \rightarrow R_3 - R_1$,

$$= (-1) \begin{vmatrix} 1 & x+2 & x+a \\ 0 & 1 & b-a \\ 0 & 2 & c-a \end{vmatrix}$$

$$= (-1)(c + a - 2b) = 0$$

....[\because a, b, c are in A.P. $\Rightarrow 2b = a + c$]

$$9. \text{ Let } S = 1 + \frac{1.3}{6} + \frac{1.3.5}{6.8} + \dots \infty$$

$$\therefore \frac{S}{4} = \frac{1}{4} + \frac{1.3}{4.6} + \frac{1.3.5}{4.6.8} + \dots \infty$$

Multiplying on both sides by $\frac{1}{2}$, we get

$$\frac{S}{8} = \frac{1}{2.4} + \frac{1.3}{2.4.6} + \frac{1.3.5}{2.4.6.8} + \dots \infty$$

$$\therefore \frac{1}{2} \frac{S}{8} = \frac{1}{2} \left[\frac{1}{2.4} + \frac{1.3}{2.4.6} + \frac{1.3.5}{2.4.6.8} + \dots \infty \right]$$

$$\Rightarrow \frac{1}{2} \frac{S}{8} = \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{4} \cdot \frac{1}{2} \cdot \frac{1.3}{4.6} \cdot \frac{1}{2} \cdot \frac{1.3.5}{4.6.8} - \dots \infty$$

$$\Rightarrow \frac{1}{2} \frac{S}{8} = 1 - \frac{1}{2} + \frac{\frac{1}{2} \left(\frac{1}{2} - 1 \right)}{1.2} - \frac{\frac{1}{2} \left(\frac{1}{2} - 1 \right) \left(\frac{1}{2} - 2 \right)}{1.2.3}$$

$$+ \frac{\frac{1}{2} \left(\frac{1}{2} - 1 \right) \left(\frac{1}{2} - 2 \right) \left(\frac{1}{2} - 3 \right)}{1.2.3.4} - \dots \infty$$

$$\Rightarrow \frac{1}{2} \frac{S}{8} = (1-1)^{\frac{1}{2}} = 0$$

$$\therefore \frac{S}{8} = \frac{1}{2} \Rightarrow S = 4$$

10. Let the first installment be a and common difference of A.P. be d.

Given, 3600 = sum of 40 terms

$$= \frac{40}{2} [2a + (40 - 1)d]$$

$$\therefore 3600 = 20(2a + 39d)$$

$$\therefore 180 = 2a + 39d \quad \dots (i)$$

After 30 installments one third of the debt is unpaid

$$\therefore \frac{3600}{3} = 1200 \text{ is unpaid and } 2400 \text{ is paid.}$$

$$\text{Now, } 2400 = \frac{30}{2} [2a + (30 - 1)d]$$

$$\therefore 160 = 2a + 29d \quad \dots (ii)$$

Subtracting (ii) from (i), we get $20 = 10d$

$$\therefore d = 2$$

From (i), $180 = 2a + 39(2)$

$$\therefore 2a = 180 - 78 = 102$$

$$\therefore a = 51$$

\therefore value of the 8th installment

$$= a + (8 - 1)d = 51 + 7(2) = ₹ 65$$

11 Probability



Hints



Classical Thinking

5. Here, $P(A) = 1$
 $\therefore P(\bar{A}) = 1 - P(A) = 0$
6. Here, $n(S) = 2 \times 2 \times 2 \times 2 = 16$
 A: Event of getting all heads
 $\Rightarrow A = \{(\text{HHHH})\}$
 $\therefore n(A) = 1$
 $\Rightarrow P(A) = \frac{1}{16}$
7. Here, $n(S) = 52$
 There is one queen of club and one king of heart
 \therefore Favourable ways = $1 + 1 = 2$
 \therefore Required Probability = $\frac{2}{52} = \frac{1}{26}$
8. Required probability = $\frac{12}{52} = \frac{3}{13}$
9. Total number of outcomes = 36
 Favourable number of outcomes = 6
 i.e., $\{(1, 1), (2, 2), (3, 3), (4, 4), (5, 5), (6, 6)\}$
 \therefore Required probability = $\frac{6}{36} = \frac{1}{6}$
10. Required probability = $\frac{3}{36} = \frac{1}{12}$
11. Required probability = $\frac{5}{25} = \frac{1}{5}$
12. Odd and perfect square (< 10) are 1, 9.
 Hence, required probability = $\frac{2}{10} = \frac{1}{5}$
13. Since there are one A, two I and one O, hence
 the required probability = $\frac{1+2+1}{11} = \frac{4}{11}$
14. Two fruits out of 6 can be chosen in ${}^6C_2 = 15$ ways.
 One mango and one apple can be chosen in
 ${}^3C_1 \times {}^3C_1 = 9$ ways
 \therefore Probability = $\frac{9}{15} = \frac{3}{5}$
15. Three persons can be chosen out of 8 in
 ${}^8C_3 = 56$ ways.
 The number of girls is more than that of the boys if either 3 girls are chosen or two girls and one boy is chosen. This can be done in
 ${}^3C_3 + {}^3C_2 \times {}^5C_1$ ways
 $= 1 + 3 \times 5 = 16$ ways.
 \therefore Required probability = $\frac{16}{56} = \frac{2}{7}$
16. Number of tickets, numbered such that it is divisible by 20 are $\frac{10000}{20} = 500$
 Hence, required probability = $\frac{500}{10000} = \frac{1}{20}$.
17. In a non-leap year, we have 365 days i.e., 52 weeks and one day. So, we may have any day of seven days.
18. Total no. of ways = $3! = 6$
 Favourable ways = 1
 \Rightarrow Probability = $\frac{1}{6}$
19. Probability of keeping at least one letter in wrong envelope = $1 - \frac{1}{n!}$
 \therefore option (B) is the correct answer.
20. Sample space when six dice are thrown = 6^6
 All dice show the same face means we are getting same number on all six dice which can be any one of the six numbers 1, 2, ..., 6.
 \therefore No. of ways of selecting a number is 6C_1 .
 \therefore Required probability = $\frac{{}^6C_1}{6^6} = \frac{1}{6^5}$
21. $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
 $\therefore \frac{5}{8} = \frac{1}{4} + \frac{1}{2} - P(A \cap B)$
 $\therefore P(A \cap B) = \frac{1}{8}$
22. Since, events are mutually exclusive, therefore
 $P(A \cap B) = 0$ i.e., $P(A \cup B) = P(A) + P(B)$
 $\Rightarrow 0.7 = 0.4 + x \Rightarrow x = \frac{3}{10}$



23. $P(A \text{ or } B) = P(A \cup B)$
 $= P(A) + P(B) - P(A \cap B)$
 $= 0.25 + 0.5 - 0.15 = 0.6$
24. $P(A) = P(A \cap B) + P(A \cup B) - P(B)$
 $= \frac{1}{3} + \frac{5}{6} - \frac{2}{3} = \frac{3}{6} = \frac{1}{2}$
25. $P(A) = 0.28, P(B) = 0.55, P(A \cap B) = 0.14$
 $P(A' \cap B') = P[(A \cup B)'] = 1 - P(A \cup B)$
 $= 1 - [P(A) + P(B) - P(A \cap B)]$
 $= 1 - (0.28 + 0.55 - 0.14) = 0.31$
26. Here, $P(A \cup B) = 0.6$ and $P(A \cap B) = 0.3$
 $\therefore P(A) + P(B) = P(A \cup B) + P(A \cap B) = 0.9$
 $\therefore P(A') + P(B') = 1 - P(A) + 1 - P(B)$
 $= 2 - 0.9 = 1.1$
27. Probability of getting either first class or second class or third class = $P(A)$
 $= \frac{2}{7} + \frac{3}{5} + \frac{1}{10}$
 $= \frac{69}{70}$
 Probability of failing = $P(A') = 1 - P(A) = \frac{1}{70}$
28. There are 4 kings, 13 hearts and a king of hearts is common to the two blocks.
 \therefore Required probability = $\frac{4+13-1}{52} = \frac{16}{52}$
29. Total number of ways = {HH, HT, TH, TT}
 $\therefore P(\text{head on first toss}) = \frac{2}{4} = \frac{1}{2} = P(A)$,
 $P(\text{head on second toss}) = \frac{2}{4} = \frac{1}{2} = P(B)$
 and $P(\text{head on both toss}) = \frac{1}{4} = P(A \cap B)$
 Hence, required probability is,
 $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
 $= \frac{1}{2} + \frac{1}{2} - \frac{1}{4} = \frac{3}{4}$
31. If A and B are independent, A' and B' are also independent.
33. $P(A/B) = \frac{P(A \cap B)}{P(B)}$
 Since, A and B are mutually exclusive.
 So, $P(A \cap B) = 0$.
 Hence, $P(A/B) = \frac{0}{P(B)} = 0$

34. $P(A/B) = \frac{P(A \cap B)}{P(B)} = \frac{0.5}{0.6} = \frac{5}{6}$
35. $P(A/B) = \frac{P(A \cap B)}{P(B)} = \frac{(3/8) + (5/8) - (3/4)}{(5/8)}$
 $= \frac{2}{5}$
36. $P(B/A) = \frac{P(A \cap B)}{P(A)} = \frac{1/4}{1/2} = \frac{1}{2}$
37. $P(A/B) = \frac{P(A \cap B)}{P(B)} = \frac{P(A \cap B)}{1 - P(B')} = \frac{0.15}{1 - 0.10}$
 $= \frac{1}{6}$
38. $P\left(\frac{\bar{A}}{\bar{B}}\right) = \frac{P(\bar{A} \cap \bar{B})}{P(\bar{B})} = \frac{P(\overline{A \cup B})}{P(\bar{B})} = \frac{1 - P(A \cup B)}{P(\bar{B})}$
39. Let E_1 be the event that man will be selected and E_2 be the event that woman will be selected. Then
 $P(E_1) = \frac{1}{2}$, So $P(\bar{E}_1) = 1 - \frac{1}{2} = \frac{1}{2}$ and
 $P(E_2) = \frac{1}{3}$, So $P(\bar{E}_2) = 1 - \frac{1}{3} = \frac{2}{3}$
 Clearly, E_1 and E_2 are independent events.
 $\therefore P(\bar{E}_1 \cap \bar{E}_2) = P(\bar{E}_1) \times P(\bar{E}_2)$
 $= \frac{1}{2} \times \frac{2}{3} = \frac{1}{3}$
40. Let A be the event of selecting bag X, B be the event of selecting bag Y and E be the event of drawing a white ball, then
 $P(A) = \frac{1}{2}, P(B) = \frac{1}{2}, P(E/A) = \frac{2}{5}$
 and $P(E/B) = \frac{4}{6} = \frac{2}{3}$
 $\therefore P(E) = P(A) P(E/A) + P(B) P(E/B)$
 $= \frac{1}{2} \cdot \frac{2}{5} + \frac{1}{2} \cdot \frac{2}{3} = \frac{8}{15}$
41. Required probability = $\frac{3}{5}$
 \therefore The probability of the occurrence = $\frac{b}{a+b}$



42. Required probability = $\frac{6}{6+5} = \frac{6}{11}$
 ... $\left[\because \text{The probability of the occurrence} = \frac{a}{a+b} \right]$
43. Here, $P(A) = \frac{3}{7}$, $P(B) = \frac{7}{12}$
 $\therefore P(A') = \frac{4}{7}$ and $P(B') = \frac{5}{12}$
 \therefore P(Problem will be considered solved even if one person solves it)
 $= 1 - [P(A') \cdot P(B')] = 1 - \frac{5}{21} = \frac{16}{21}$



Critical Thinking

- Here, $n(S) = 2 \times 2 = 4$
 A: Event of getting 2 heads or 2 tails
 $\therefore A = \{(H H), (T T)\}$
 $\Rightarrow n(A) = 2$
 $\Rightarrow P(A) = \frac{2}{4} = \frac{1}{2}$
- One card can be selected from a pack in ${}^{52}C_1$ ways.
 $\therefore n(S) = {}^{52}C_1 = 52$
 A: Event of getting a red queen
 $\therefore P(A) = P(\text{diamond queen or heart queen})$
 $= \frac{{}^2C_1}{{}^{52}C_1}$
- Favourable ways
 $= \{29, 92, 38, 83, 47, 74, 56, 65\}$
 Hence, required probability = $\frac{8}{100} = \frac{2}{25}$
- Two digits, one from each set can be selected in $9 \times 9 = 81$ ways.
 Favourable outcomes are (1, 9), (2, 8), (3, 7), (4, 6), (5, 5), (6, 4), (7, 3), (8, 2) and (9, 1).
 $\therefore n(S) = 81$
 and $n(A) = 9$
 $\therefore P(A) = \frac{9}{81} = \frac{1}{9}$
- When six dice are thrown, the total number of outcomes is 6^6 . They can show different number in ${}^6P_6 = 6!$ ways
 \therefore Required probability = $\frac{6!}{6^6} = \frac{5!}{6^5} = \frac{5}{324}$
- The sum 2 can be found in one way i.e., $\{(1, 1)\}$
 The sum 8 can be found in five ways i.e., $\{(6, 2), (5, 3), (4, 4), (3, 5), (2, 6)\}$.
 Similarly, the sum twelve can be found in one way i.e., $\{(6, 6)\}$.
 Hence, required probability = $\frac{7}{36}$.
- Between 1 and 100, there are 25 prime numbers.
 $\therefore n(S) = 98$ and $n(A) = 25$
 $\therefore P(A) = \frac{25}{98}$
- Total cases = 4
 So, probability of correct answer = $\frac{1}{4}$
- In a leap year, there are 366 days in which 52 weeks and two days. The combination of 2 days may be: Sun – Mon, Mon – Tue, Tue – Wed, Wed – Thu, Thu – Fri, Fri – Sat, Sat – Sun.
 $\therefore P(53 \text{ Sun}) = \frac{2}{7}$
- When a coin is tossed, there are two outcomes and when a dice is rolled, there are six possible outcomes.
 Hence, there are 8 (2 corresponding to head and six corresponding to tail at first toss) sample points in the sample space.
 Sample space is $\{HH, HT, T1, T2, T3, T4, T5, T6\}$.
- It six does not appear on either dice then, there are only five possible outcomes associated with one dice, the number of sample points is 5×5 .
- Since, the total '13' can't be found.
- Probabilities of H_1, H_2 and H_3 winning a race must be in the ratio 4 : 2 : 1 (due to given condition) and should also add up to 1.
- Here, $n(S) = {}^6C_2 = 15$
 If both are vowels, then they are selected in 2C_2 ways = 1.
 \therefore Required probability = $\frac{1}{15}$
- Here, $n(S) = {}^{10}C_2$
 A: Event that the watches selected are defective
 $\therefore n(A) = {}^2C_2 = 1$
 $\therefore P(A) = \frac{1}{{}^{10}C_2} = \frac{1}{45}$



16. Total no. of ways in which 2 socks can be drawn out of 9 is 9C_2 . The two socks match if either they are both black or they are both blue. So, two matching socks can be drawn in ${}^5C_2 + {}^4C_2$ ways.
- \therefore Required probability = $\frac{{}^5C_2 + {}^4C_2}{{}^9C_2}$
- $$= \frac{10+6}{36} = \frac{4}{9}$$
17. Ace is not drawn in 26 cards.
It means 26 cards are drawn from 48 cards.
- \therefore Required Probability = $\frac{{}^{48}C_{26}}{{}^{52}C_{26}}$
18. $n(S) = {}^{16}C_{11}$
A: Event that the team has exactly four bowlers.
- \therefore $n(A) = {}^6C_4 \cdot {}^{10}C_7$
- $$\Rightarrow P(A) = \frac{{}^6C_4 \cdot {}^{10}C_7}{{}^{16}C_{11}} = \frac{75}{182}$$
19. We have to select exactly 2 children
- \therefore selection contain 2 children out of 4 children and remaining 2 person can be selected from 2 women and 4 men
i.e., $4C_2 \times 6C_2$ ways
- \therefore Total favourable ways = $6 \times 15 = 90$
- \therefore Required probability = $\frac{90}{210} = \frac{3}{7}$
20. A committee of 4 can be formed in ${}^{25}C_4$ ways
A: Event that the committee contains at least 3 doctors
- \therefore $n(A) = {}^4C_3 \cdot {}^{21}C_1 + {}^4C_4 = 85$
- \therefore $P(A) = \frac{85}{{}^{25}C_4} = \frac{85}{12650} = \frac{17}{2530}$
21. Since, cards are drawn with replacement.
- \therefore Total no. of ways = 52×52 .
Now, we can choose one suit out of four in 4C_1 ways and two cards in 13×13 ways.
- \therefore Required Probability = $\frac{{}^4C_1 \times 13 \times 13}{52 \times 52} = \frac{1}{4}$
22. Besides ground floor, there are 7 floors. Since a person can leave the cabin at any of the seven floors, total no. of ways in which each of the five persons can leave the cabin at any of the 7 floors = 7^5
Five persons can leave the cabin at five different floors in ${}^7C_5 \times 5!$ ways
Hence, required probability = $\frac{{}^7C_5 \times 5!}{7^5}$
23. Here, $n(S) = 2 \times 2 \times 2 = 8$
If A is the event that there is no tail, then
 $A = \{(HHH)\}$
 $\Rightarrow n(A) = 1$
 $\Rightarrow P(A) = \frac{1}{8}$
- \therefore $P(A') = 1 - P(A) = 1 - \frac{1}{8} = \frac{7}{8}$
24. Required probability
- $$= \left(1 - \frac{1}{3}\right) \left(1 - \frac{1}{4}\right) \left(1 - \frac{1}{5}\right) = \frac{2}{3} \cdot \frac{3}{4} \cdot \frac{4}{5} = \frac{2}{5}$$
25. Out of 30 numbers from 1 to 30, three numbers can be chosen in ${}^{30}C_3$ ways.
Three consecutive numbers can be chosen in one of the following ways:
 $\{(1, 2, 3), (2, 3, 4), \dots, (28, 29, 30)\} = 28$ ways
- \therefore Probability that numbers are consecutive
- $$= \frac{28}{{}^{30}C_3} = \frac{1}{145}$$
- Hence, required probability = $1 - \frac{1}{145} = \frac{144}{145}$
26. Total no. of ways = $7!$
Arrangement of boys and girls in alternate seats is B G B G B G B
Boys can occupy seat in $4!$ ways and girls in $3!$ ways.
- \therefore Required Probability = $\frac{3! \times 4!}{7!} = \frac{1}{35}$
27. Two 3s, one 6 and one 8 can be dialled in $\frac{4!}{2!} = 12$ ways of which only one is the correct way of dialling.
- \therefore Required probability = $\frac{1}{12}$
28. As $\{(1, 1, 1), (2, 2, 2), (3, 3, 3), (4, 4, 4), (5, 5, 5), (6, 6, 6)\}$ are only favourable outcomes
- $$\Rightarrow \text{Required probability} = \frac{6}{216}$$
29. Since there are 3 A's and 2 N's.
Total no. of arrangements = $\frac{10!}{3!2!}$
Hence, the number of arrangements in which ANAND occurs without any split = 6!
- \therefore Required probability = $\frac{6!3!2!}{10!} = \frac{1}{420}$



30. 15 places are occupied. This includes the owner's car also. 14 cars are parked in 24 places of which 22 places are available (excluding the neighbouring places) and so the required probability $\frac{{}^{22}C_{14}}{{}^{24}C_{14}} = \frac{15}{92}$

31. Three numbers can be chosen out of 10 numbers in ${}^{10}C_3$ ways.

The product of two numbers, out of the three chosen numbers, will be equal to the third number, if the numbers are chosen in one of the following ways:

$$\{(2, 3, 6), (2, 4, 8), (2, 5, 10)\} = 3 \text{ ways}$$

$$\text{Hence, required probability} = \frac{3}{{}^{10}C_3} = \frac{1}{40}$$

32. 4 cards can drop out of 52 in ${}^{52}C_4$ ways. They can be one from each suit in ${}^{13}C_1 \times {}^{13}C_1 \times {}^{13}C_1 \times {}^{13}C_1 = (13 \times 13 \times 13 \times 13)$ ways.

$$\begin{aligned} \therefore \text{Required probability} &= \frac{13 \times 13 \times 13 \times 13}{{}^{52}C_4} \\ &= \frac{13 \times 13 \times 13 \times 13 \times 4!}{52 \times 51 \times 50 \times 49} \\ &= \frac{2197}{20825} \end{aligned}$$

$$33. \quad 0.7 = 0.4 + x - 0.4x$$

$$\Rightarrow x = \frac{1}{2}$$

34. Since, we have

$$\begin{aligned} P(A \cup B) + P(A \cap B) &= P(A) + P(B) \\ &= P(A) + \frac{P(A)}{2} \end{aligned}$$

$$\Rightarrow \frac{7}{8} = \frac{3P(A)}{2}$$

$$\Rightarrow P(A) = \frac{7}{12}$$

35. Since, $A \cup B = S$.

$$\therefore P(A \cup B) = P(S) = 1$$

$$\therefore 1 = P(A) + 2P(A) \quad [\because P(A \cup B) = P(A) + P(B)]$$

$$\Rightarrow 3P(A) = 1$$

$$\Rightarrow P(A) = \frac{1}{3}$$

$$\therefore P(B) = \frac{2}{3}$$

36. A: Event of obtaining an even sum and
B: Event of obtaining a sum less than five.
Since A, B are not mutually exclusive,

$$\begin{aligned} \therefore P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\ &= \frac{18}{36} + \frac{6}{36} - \frac{4}{36} = \frac{5}{9} \end{aligned}$$

[\because there are 18 ways to get an even sum i.e.

$\{(1, 1), (1, 3), (1, 5), (2, 2), (2, 4), (2, 6), (3, 1), (3, 3), (3, 5), (4, 2), (4, 4), (4, 6), (5, 1), (5, 3), (5, 5), (6, 2), (6, 4), (6, 6)\}$ and there are 6 ways to get a sum < 5 i.e., $\{(1, 3), (3, 1), (2, 2), (1, 2), (2, 1), (1, 1)\}$ and 4 ways to get an even sum < 5 i.e., $\{(1, 3), (3, 1), (2, 2), (1, 1)\}$]

37. Here, $A = \{4, 5, 6\}$

$$\Rightarrow P(A) = \frac{3}{6} = \frac{1}{2}$$

and $B = \{4, 3, 2, 1\}$

$$\Rightarrow P(B) = \frac{4}{6} = \frac{2}{3}$$

$$\therefore A \cap B = \{4\}$$

$$\Rightarrow P(A \cap B) = \frac{1}{6}$$

$$\therefore P(A \cup B) = \frac{1}{2} + \frac{2}{3} - \frac{1}{6} = 1$$

38. A is independent of itself, if

$$P(A \cap A) = P(A).P(A)$$

$$\Rightarrow P(A) = P(A)^2$$

$$\Rightarrow P(A) = 0, 1$$

39. We have $P(A + B) = P(A) + P(B) - P(AB)$

$$\Rightarrow \frac{5}{6} = \frac{1}{2} + P(B) - \frac{1}{3} \Rightarrow P(B) = \frac{4}{6} = \frac{2}{3}$$

$$\text{Thus, } P(A).P(B) = \frac{1}{2} \times \frac{2}{3} = \frac{1}{3} = P(AB)$$

Hence, events A and B are independent.

$$40. \quad \text{Let } P(A) = \frac{20}{100} = \frac{1}{5}, \quad P(B) = \frac{10}{100} = \frac{1}{10}$$

Since, events are independent and we have to find $P(A \cup B) = P(A) + P(B) - P(A).P(B)$

$$= \frac{1}{5} + \frac{1}{10} - \frac{1}{5} \times \frac{1}{10}$$

$$= \frac{3}{10} - \frac{1}{50} = \frac{14}{50} \times 100 = 28\%$$



41. In a leap year, there are 366 days in which 52 weeks and two days. The combination of 2 days may be: Sun-Mon, Mon-Tue, Tue-Wed, Wed-Thu, Thu-Fri, Fri-Sat, Sat-Sun.

$$\therefore P(53 \text{ Fri}) = \frac{2}{7}; P(53 \text{ Sat}) = \frac{2}{7}$$

There is one combination in common i.e., (Fri-Sat)

$$\therefore P(53 \text{ Fri and } 53 \text{ Sat}) = \frac{1}{7}$$

$$\begin{aligned} \therefore P(53 \text{ Fri or } 53 \text{ Sat}) &= P(53 \text{ Fri}) + P(53 \text{ Sat}) \\ &\quad - P(53 \text{ Fri and } 53 \text{ Sat}) \\ &= \frac{2}{7} + \frac{2}{7} - \frac{1}{7} = \frac{3}{7} \end{aligned}$$

42. Here, $P(A) = P(B) = 2P(C)$,
and $P(A) + P(B) + P(C) = 1$

$$\Rightarrow P(C) = \frac{1}{5} \text{ and } P(A) = P(B) = \frac{2}{5}$$

$$\text{Hence, } P(A \cup B) = P(A) + P(B) = \frac{2}{5} + \frac{2}{5} = \frac{4}{5}$$

43. For both to be boys, the probability

$$= \left(\frac{1}{2}\right)\left(\frac{1}{2}\right) = \frac{1}{4}$$

44. We have to consider order for IIT

$$\therefore \text{Required probability} = \frac{10}{20} \times \frac{9}{19} \times \frac{10}{18} = \frac{5}{38}$$

45. In the word 'MULTIPLE' there are 3 vowels, out of total of 8, 1 vowel can be chosen in 3C_1 ways. In the word 'CHOICE' there are 3 vowels, out of the total of 6, 1 vowel can be chosen in 3C_1 ways.

$$\therefore \text{Required probability} = \frac{{}^3C_1}{8} \times \frac{{}^3C_1}{6} = \frac{3}{16}$$

46. A total of 7 and a total of 9 cannot occur simultaneously.

$$\begin{aligned} \therefore P(\text{total of } 7 \text{ or } 9) \\ &= P(\text{total of } 7) + P(\text{total of } 9) = \frac{6}{36} + \frac{4}{36} = \frac{5}{18} \end{aligned}$$

(A total of 7 and a total of 9 cannot occur simultaneously)

$$47. \frac{1}{2} \times \frac{2}{5} + \frac{1}{2} \times \frac{3}{10} = \frac{7}{20}$$

$$48. P(G) = \frac{25}{80}, P(R) = \frac{10}{80}, P(I) = \frac{20}{80}$$

Since events are independent,

$$\begin{aligned} \therefore P(\text{selecting rich and intelligent girls}) \\ &= P(G) \cdot P(R) \cdot P(I) = \frac{5}{512} \end{aligned}$$

$$\begin{aligned} 49. P(A' \cup B') &= P[(A \cap B)'] \\ &= 1 - P(A \cap B) = 1 - \frac{1}{4} = \frac{3}{4} \end{aligned}$$

$$\begin{aligned} 50. P(A' \cap B') &= \frac{1}{3} \\ \Rightarrow P[(A \cup B)'] &= \frac{1}{3} \end{aligned}$$

$$\begin{aligned} \therefore 1 - P(A \cup B) &= \frac{1}{3} \\ \Rightarrow P(A \cup B) &= 1 - \frac{1}{3} = \frac{2}{3} \end{aligned}$$

$$\therefore P(A) + P(B) - P(A \cap B) = \frac{2}{3}$$

$$\begin{aligned} \therefore p + 2p - \frac{1}{2} &= \frac{2}{3} \\ \Rightarrow 3p &= \frac{2}{3} + \frac{1}{2} = \frac{7}{6} \Rightarrow p = \frac{7}{18} \end{aligned}$$

51. Required Probability

$$\begin{aligned} &= P[(A \cap B') \cup (A' \cap B)] \\ &= P(A \cap B') + P(A' \cap B) \\ &= P(A) - P(A \cap B) + P(B) - P(A \cap B) \\ &= P(A) + P(B) - 2P(A \cap B) \end{aligned}$$

52. P(neither A nor B)

$$\begin{aligned} &= 1 - P(\text{either A or B}) = 1 - P(A \cup B) \\ &= 1 - [P(A) + P(B) - P(A \cap B)] \\ &= 1 - 0.25 - 0.50 + 0.14 = 0.39 \end{aligned}$$

53. M: Event that student passed in Mathematics.

E: Event that student passed in Electronics

$$\therefore n(M) = 30, n(E) = 20, n(M \cap E) = 10, n(S) = 80.$$

$$\therefore P(M) = \frac{30}{80}, P(E) = \frac{20}{80}, P(M \cap E) = \frac{10}{80}$$

$$\begin{aligned} \therefore P(M \cup E) &= P(M) + P(E) - P(M \cap E) \\ &= \frac{30}{80} + \frac{20}{80} - \frac{10}{80} = \frac{1}{2} \end{aligned}$$

\therefore P(Student has passed in none of the subject)

$$= P[(M \cup E)'] = 1 - P(M \cup E) = 1 - \frac{1}{2} = \frac{1}{2}$$

$$\begin{aligned} 54. P(\text{neither } E_1 \text{ nor } E_2 \text{ occurs}) &= P(E_1' \cap E_2') \\ &= P(E_1')P(E_2') \\ &= (1 - p_1)(1 - p_2) \end{aligned}$$



55. $P(M) = \frac{1}{4} \Rightarrow P(M') = \frac{3}{4}$
 and $P(W) = \frac{1}{3} \Rightarrow P(W') = \frac{2}{3}$
 Both events are independent so that probability that no one will be alive is
 $P(W' \cap M') = P(W') P(M') = \frac{3}{4} \times \frac{2}{3} = \frac{1}{2}$

56. Here, $P(A) = p$
 $\Rightarrow P(\bar{A}) = 1 - p$
 and $P(B) = q \Rightarrow P(\bar{B}) = 1 - q$
 Probability that one person is alive is the sum of two cases A dies B lives and A lives B dies
 $= p(1 - q) + q(1 - p) = p + q - 2pq$

57. Here, $P(A) = 0.6$; $P(B) = 0.9$
 \therefore Required probability
 $= P(A) \cdot P(\bar{B}) + P(B) \cdot P(\bar{A}) = (0.6)(0.1) + (0.9)(0.4)$
 $= 0.06 + 0.36 = 42$

58. Since, E and F are independent
 $\therefore P(E \cap F) = P(E) P(F)$
 $\Rightarrow P(E) P(F) = \frac{1}{12}$
 Now, E and F are independent
 $\therefore E'$ and F' are also independent
 $\therefore P(E' \cap F') = P(E') \cdot P(F') = \frac{1}{2}$

$\therefore [1 - P(E)] \cdot [1 - P(F)] = \frac{1}{2}$
 $\therefore 1 - P(E) - P(F) + P(E) \cdot P(F) = \frac{1}{2}$
 $\therefore 1 - P(E) - P(F) + \frac{1}{12} = \frac{1}{2}$
 $\Rightarrow P(E) + P(F) = \frac{7}{12}$
 Solving, $P(E) = \frac{1}{4}$, $P(F) = \frac{1}{3}$

59. $P\left(\frac{A}{B}\right) = \frac{P(A \cap B)}{P(B)} = \frac{P(A) \cdot P(B)}{P(B)} = P(A)$.

60. Since, $A \subseteq B \Rightarrow A \cap B = B \cap A = A$
 Hence, $P\left(\frac{B}{A}\right) = \frac{P(B \cap A)}{P(A)} = \frac{P(A)}{P(A)} = 1$

61. $P\left(\frac{\bar{B}}{\bar{A}}\right) = \frac{1 - P(A \cup B)}{P(\bar{A})} = \frac{1 - \frac{23}{60}}{1 - \frac{1}{3}} = \frac{37}{60} \times \frac{3}{2} = \frac{37}{40}$

62. A: Brown hair
 $\Rightarrow P(A) = \frac{40}{100}$
 B: Brown eyes
 $\Rightarrow P(B) = \frac{25}{100}$

$\therefore P(A \cap B) = \frac{15}{100}$

$\therefore P(B/A) = \frac{P(A \cap B)}{P(A)} = \frac{\frac{15}{100}}{\frac{40}{100}} = \frac{3}{8}$

63. $P(A \cap B) = P(A) \cdot P(B/A) = \frac{1}{4} \times \frac{1}{2} = \frac{1}{8}$

Since, $P(A \cap B) = P(B) P(A/B)$

$\therefore \frac{1}{8} = P(B) \times \frac{1}{4}$

$\Rightarrow P(B) = \frac{1}{2}$

$\therefore P(A) \cdot P(B) = \frac{1}{4} \times \frac{1}{2} = \frac{1}{8} = P(A \cap B)$

\therefore A and B are independent

64. It is based on Baye's theorem.

Probability of picked bag A, i.e., $P(A) = \frac{1}{2}$

Probability of picked bag B, i.e., $P(B) = \frac{1}{2}$

Probability of green ball picked from bag A

$= P(A) \cdot P\left(\frac{G}{A}\right) = \frac{1}{2} \times \frac{4}{7} = \frac{2}{7}$

Probability of green ball picked from bag B

$= P(B) \cdot P\left(\frac{G}{B}\right) = \frac{1}{2} \times \frac{3}{7} = \frac{3}{14}$

\therefore Total probability of green ball $= \frac{2}{7} + \frac{3}{14} = \frac{1}{2}$

\therefore Probability of fact that green ball is drawn from bag B

$= \frac{P(B)P\left(\frac{G}{B}\right)}{P(A)P\left(\frac{G}{A}\right) + P(B)P\left(\frac{G}{B}\right)} = \frac{\frac{1}{2} \times \frac{3}{7}}{\frac{1}{2} \times \frac{4}{7} + \frac{1}{2} \times \frac{3}{7}} = \frac{3}{7}$



65. Consider the following events :

A → Ball drawn is black;

E₁ → Bag I is chosen;

E₂ → Bag II is chosen and

E₃ → Bag III is chosen.

$$\text{Then } P(E_1) = P(E_2) = P(E_3) = \frac{1}{3}, P\left(\frac{A}{E_1}\right) = \frac{3}{5}$$

$$P\left(\frac{A}{E_2}\right) = \frac{1}{5}, P\left(\frac{A}{E_3}\right) = \frac{7}{10}$$

$$\therefore \text{ Required probability} = P\left(\frac{E_3}{A}\right)$$

$$\begin{aligned} &= \frac{P(E_3)P\left(\frac{A}{E_3}\right)}{P(E_1)P\left(\frac{A}{E_1}\right) + P(E_2)P\left(\frac{A}{E_2}\right) + P(E_3)P\left(\frac{A}{E_3}\right)} \\ &= \frac{7}{15} \end{aligned}$$

66. Let E denote the event that a six occurs and A is the event that the man reports that it is a '6', we have

$$P(E) = \frac{1}{6}, P(E') = \frac{5}{6}, P(A/E) = \frac{3}{4} \text{ and}$$

$$P(A/E') = \frac{1}{4}$$

∴ From Baye's theorem,

$$\begin{aligned} P(E/A) &= \frac{P(E).P\left(\frac{A}{E}\right)}{P(E).P\left(\frac{A}{E}\right) + P(E').P\left(\frac{A}{E'}\right)} \\ &= \frac{\frac{1}{6} \times \frac{3}{4}}{\frac{1}{6} \times \frac{3}{4} + \frac{5}{6} \times \frac{1}{4}} = \frac{3}{8} \end{aligned}$$

67. We define the following events :

A₁ : He knows the answer.

A₂ : He does not know the answer.

E : He gets the correct answer.

$$\text{Then } P(A_1) = \frac{9}{10}, P(A_2) = 1 - \frac{9}{10} = \frac{1}{10},$$

$$P\left(\frac{E}{A_1}\right) = 1 \text{ and } P\left(\frac{E}{A_2}\right) = \frac{1}{4}$$

∴ Required probability is

$$P\left(\frac{A_2}{E}\right) = \frac{P(A_2)P\left(\frac{E}{A_2}\right)}{P(A_1)P\left(\frac{E}{A_1}\right) + P(A_2)P\left(\frac{E}{A_2}\right)} = \frac{1}{37}$$

68. We define the following events :

A₁ : Selecting a pair of consecutive letter from the word LONDON.

A₂ : Selecting a pair of consecutive letters from the word CLIFTON.

E : Selecting a pair of letters 'ON'.

Then $P(A_1 \cap E) = \frac{2}{5}$; as there are 5 pairs of consecutive letters out of which 2 are ON.

$P(A_2 \cap E) = \frac{1}{6}$; as there are 6 pairs of consecutive letters of which one is ON.

∴ The required probability is

$$\begin{aligned} P\left(\frac{A_1}{E}\right) &= \frac{P(A_1 \cap E)}{P(A_1 \cap E) + P(A_2 \cap E)} = \frac{\frac{2}{5}}{\frac{2}{5} + \frac{1}{6}} \\ &= \frac{12}{17} \end{aligned}$$

$$69. \text{ Required probability} = \frac{5}{5+3} = \frac{5}{8}$$

$$\left[\begin{array}{l} \therefore \text{ If odds in favours of an event are } a : b, \\ \text{ then the probability of non - occurrence} \\ \text{ of that event is } \frac{b}{a+b} \end{array} \right]$$

$$70. \text{ Required probability} = \frac{4}{4+5} = \frac{4}{9}$$

71. Let p be the probability of the other event. Then the probability of the first event is $\frac{2}{3}p$.

$$\therefore \frac{p}{p + \frac{2}{3}p} = \frac{3}{3+2}$$

∴ odds in favour of the other are 3 : 2



72. Probabilities of winning the race by three horses are $\frac{1}{3}$, $\frac{1}{4}$ and $\frac{1}{5}$.

$$\text{Hence, required probability} = \frac{1}{3} + \frac{1}{4} + \frac{1}{5} = \frac{47}{60}$$

73. Required probability = $\frac{1}{2} \times \frac{4}{7} + \frac{1}{2} \times \frac{6}{8} = \frac{37}{56}$

74. Probability of the card being a spade or an ace = $\frac{16}{52} = \frac{4}{13}$. Hence, odds in favour is 4 : 9.

So, the odds against his winning is 9 : 4

75. We have ratio of the ships A, B and C for arriving safely are 2 : 5, 3 : 7 and 6 : 11 respectively.

\therefore The probability of ship A for arriving safely = $\frac{2}{2+5} = \frac{2}{7}$

Similarly, for B = $\frac{3}{3+7} = \frac{3}{10}$ and for

$$C = \frac{6}{6+11} = \frac{6}{17}$$

\therefore Probability of all the ships for arriving safely = $\frac{2}{7} \times \frac{3}{10} \times \frac{6}{17} = \frac{18}{595}$.

76. Let A and B be two given events. The odds against A are 5:2, therefore $P(A) = \frac{2}{7}$.

And the odds in favour of B are 6:5,

$$\text{therefore } P(B) = \frac{6}{11}$$

\therefore The required probability = $1 - P(\bar{A})P(\bar{B})$
 $= 1 - \left(1 - \frac{2}{7}\right)\left(1 - \frac{6}{11}\right) = \frac{52}{77}$



Competitive Thinking

1. $n(S) = 36$
 $E = \{(1, 4), (4, 1), (2, 3), (3, 2)\}$

$$\therefore P(E) = \frac{4}{36} = \frac{1}{9}$$

2. Required probability = $\frac{26}{36} = \frac{13}{18}$

3. Required probability = $\frac{4}{36} = \frac{1}{9}$

4. Total number of ways = 36
 and Favourable number of cases are
 $\{(1, 4), (2, 3), (3, 2), (4, 1), (1, 5), (2, 4), (3, 3), (4, 2), (5, 1)\} = 9$
 Hence, the required probability = $\frac{9}{36} = \frac{1}{4}$.

5. Required probability = $\frac{15}{36} = \frac{5}{12}$

6. Prime numbers are $\{2, 3, 5, 7, 11\}$.
 Hence, required probability
 $= \frac{1+2+4+6+2}{36} = \frac{15}{36} = \frac{5}{12}$

7. $n(S) = 36$
 A: Event that product of numbers is even
 $n(A) = 27$

$$P(A) = \frac{27}{36} = \frac{3}{4}$$

8. Ways $\begin{array}{cccc} 9 & 10 & 11 & 12 \\ \downarrow & \downarrow & \downarrow & \downarrow \\ 4 & 3 & 2 & 1 \end{array}$

Hence, required probability = $\frac{10}{36} = \frac{5}{18}$

10. $n(S) = 6$
 $P(T) = P(R) = \frac{1}{6}$

$$\therefore P(T \text{ or } R) = P(T) + P(R) = \frac{1}{6} + \frac{1}{6} = \frac{1}{3}$$

11. Total number of ways = 2^n
 If head comes odd times, then favourable ways = 2^{n-1} .

$$\therefore \text{Required probability} = \frac{2^{n-1}}{2^n} = \frac{1}{2}$$

12. For m sided die, which is thrown n times, the probability that the number on the top is increasing is given by $\frac{{}^m C_n}{m^n}$

Here 6-faced die is thrown three times.

$$\therefore \text{Required probability} = \frac{{}^6 C_3}{6^3} = \frac{5}{54}$$



13. 3 coins are tossed
 $\therefore S = \{HHH, HHT, HTH, THH, TTH, THT, HTT, TTT\}$

A: Event of getting 2 heads
 $\Rightarrow A = \{HHT, HTH, THH\}$

$$\therefore n(A) = 3 \Rightarrow P(A) = \frac{3}{8}$$

14. $n(S) = 8$

$$P(2 \text{ tails}) = \frac{3}{8}$$

$$P(3 \text{ tails}) = \frac{1}{8}$$

$$\begin{aligned} P(\text{at least 2 tails}) &= P(2 \text{ tails}) + P(3 \text{ tails}) \\ &= \frac{3}{8} + \frac{1}{8} = \frac{1}{2} \end{aligned}$$

15. Three dice can be thrown in $6 \times 6 \times 6 = 216$ ways. A total 17 can be obtained as $\{(5, 6, 6), (6, 5, 6), (6, 6, 5)\}$. A total 18 can be obtained as $(6, 6, 6)$.

$$\text{Hence, the required probability} = \frac{4}{216} = \frac{1}{54}$$

16. Required combinations are $\{(2, 2, 1), (1, 2, 2), (2, 1, 2), (1, 3, 1), (3, 1, 1), (1, 1, 3)\}$

$$\therefore \text{Required probability} = \frac{6}{4^3} = \frac{6}{64} = \frac{3}{32}$$

17. Required probability = $\frac{2}{10} = \frac{1}{5}$

18. $n(S) = {}^4C_2$

$$\begin{aligned} P(\text{no black ball}) &= P(\text{red ball}) \\ &= \frac{{}^2C_2}{{}^4C_2} = \frac{1}{6} \end{aligned}$$

19. 3 batteries can be selected from 10 batteries in ${}^{10}C_3$ ways.

3 dead batteries can be selected from 4 dead batteries in 4C_3 ways.

\therefore Probability that the all 3 selected batteries are

$$\text{dead} = \frac{{}^4C_3}{{}^{10}C_3} = \frac{4 \times 3 \times 2}{10 \times 9 \times 8} = \frac{1}{30}$$

20. $n(S) = {}^{10}C_4$

A: Event of getting 2 red balls

$$n(A) = {}^4C_2 \cdot {}^6C_2$$

$$\therefore P(A) = \frac{{}^4C_2 \cdot {}^6C_2}{{}^{10}C_4} = \frac{9}{21}$$

21. STATISTICS \Rightarrow SSS TTT A II C

ASSISTANT \Rightarrow SSS TT AA I N

\therefore S, T, A and I are the common letters.

$$\therefore \text{Probability of choosing S} = \frac{{}^3C_1}{{}^{10}C_1} \times \frac{{}^3C_1}{{}^9C_1} = \frac{1}{10}$$

$$\text{Probability of choosing T} = \frac{{}^3C_1}{{}^{10}C_1} \times \frac{{}^2C_1}{{}^9C_1} = \frac{1}{15}$$

$$\text{Probability of choosing A} = \frac{{}^1C_1}{{}^{10}C_1} \times \frac{{}^2C_1}{{}^9C_1} = \frac{1}{45}$$

$$\text{Probability of choosing I} = \frac{{}^2C_1}{{}^{10}C_1} \times \frac{{}^1C_1}{{}^9C_1} = \frac{1}{45}$$

$$\begin{aligned} \therefore \text{Required probability} &= \frac{1}{10} + \frac{1}{15} + \frac{1}{45} + \frac{1}{45} \\ &= \frac{19}{90} \end{aligned}$$

22. $n(S) = {}^{12}C_3$

$$\begin{aligned} P(\text{not of same colour}) &= 1 - P(\text{Same colour}) \\ &= 1 - [P(\text{red ball}) + P(\text{black ball}) + P(\text{white ball})] \end{aligned}$$

$$= 1 - \left[\frac{{}^5C_3}{{}^{12}C_3} + \frac{{}^3C_3}{{}^{12}C_3} + \frac{{}^4C_3}{{}^{12}C_3} \right]$$

$$= 1 - \left(\frac{60 + 6 + 24}{1320} \right)$$

$$= \frac{41}{44}$$

23. Total rusted items = $3 + 5 = 8$;
 unrusted nails = 3.

$$\therefore \text{Required probability} = \frac{3+8}{6+10} = \frac{11}{16}$$

24. If both integers are even, then product is even.
 If both integers are odd, then product is odd.
 If one integer is odd and other is even, then product is even.

$$\therefore \text{Required probability} = \frac{2}{3}$$

25. Number which are cubes

$$1^3 = 1, 2^3 = 8, 3^3 = 27, 4^3 = 64$$

$$\therefore \text{Required probability} = \frac{4}{100} = \frac{1}{25}$$

26. $S = \{-18, -16, -14, \dots, 20\}$

$$n(S) = 20$$

A : no. divisible by both 4 and 6

$$A = \{-12, 0, 12\}$$

$$P(A) = \frac{n(A)}{n(S)} = \frac{3}{20}$$



27. In a non leap year, there are 365 days which has 52 weeks and 1 day.
 $\therefore P(53 \text{ Sundays}) = \frac{1}{7}$
28. Here, $n(S) = 36$
 Also, $n(F)$, where F is the set of favourable cases.
 $F = \{(6, 1), (5, 2), (4, 3)\}$
 where 1st number in ordered pair gives the number of black die and 2nd number gives the number on white die.
 \therefore required probability $= \frac{3}{36} = \frac{1}{12}$
29. Here, $n(S) = {}^{52}C_1 \times {}^{51}C_1 = 52 \times 51$
 A: Event that both cards chosen are Ace.
 $\therefore n(A) = {}^4C_1 \times {}^3C_1 = 12$
 $\therefore P(A) = \frac{12}{52 \times 51} = \frac{1}{221}$
30. There are 8 even numbers from 1 to 17
 \therefore Probability of selecting 1 even number $= \frac{8}{17}$
 Remaining number of tickets = 16
 There are 7 even numbers in the remaining tickets.
 \therefore Probability of selecting second even number $= \frac{7}{16}$
 \therefore Required probability $= \frac{8}{17} \times \frac{7}{16} = \frac{7}{34}$
31. Required probability $= \frac{10!}{\frac{2!}{11!}} = \frac{2}{11}$
 $\frac{2!}{2!2!}$
32. HULULULU \Rightarrow contains 4U, 3L, 1H
 Consider 3L together i.e. we have to arrange 6 units which contains 4U.
 Hence number of possible arrangements $= \frac{6!}{4!} = 6 \times 5 = 30$
 Number of ways of arranging all letters of given word $= \frac{8!}{3!4!} = \frac{8 \times 7 \times 6 \times 5}{3 \times 2} = 8 \times 7 \times 5$
 Hence required probability $= \frac{30}{8 \times 7 \times 5} = \frac{6}{8 \times 7} = \frac{3}{28}$
33. Let E be the event that the numbers are divisible by 4.
 $\therefore E = \{4, 8, 12, 16, 20, 24\}$
 $\therefore n(E) = 6$
 $\therefore n(\bar{E}) = 20$
 \therefore Required probability $= P(\bar{E}) = \frac{20}{26} = \frac{10}{13}$
34. $P(\text{at least 1H}) = 1 - P(\text{No head})$
 $= 1 - P(\text{four tail}) = 1 - \frac{1}{16} = \frac{15}{16}$
35. Required probability is $1 - P(\text{no die show up 1})$
 $= 1 - \left(\frac{5}{6}\right)^3 = \frac{216 - 125}{216} = \frac{91}{216}$
36. We have $P(\bar{A}) = 0.05 \Rightarrow P(A) = 0.95$
 Hence, the probability that the event will take place in 4 consecutive occasions $= \{P(A)\}^4 = (0.95)^4 = 0.81450625$
37. Probability that A does not solve the problem $= 1 - \frac{1}{2} = \frac{1}{2}$
 Probability that B does not solve the problem $= 1 - \frac{1}{3} = \frac{2}{3}$
 Probability that C does not solve the problem $= 1 - \frac{1}{5} = \frac{4}{5}$
 Probability that at least one of them solve problem $= 1 - \text{no one solves the problem}$
 $= 1 - \left(\frac{1}{2}\right)\left(\frac{2}{3}\right)\left(\frac{4}{5}\right)$
 $= 1 - \frac{4}{15} = \frac{11}{15}$
38. The probability of A, B, and C not finishing the game is, $1 - \frac{1}{2} = \frac{1}{2}$, $1 - \frac{1}{3} = \frac{2}{3}$ and $1 - \frac{1}{4} = \frac{3}{4}$ respectively.
 \therefore The probability that the game is not finished by any one of them $= \frac{1}{2} \times \frac{2}{3} \times \frac{3}{4} = \frac{1}{4}$
 \therefore The probability that the game is finished $= 1 - \frac{1}{4} = \frac{3}{4}$



39. Total balls = $5 + x$
Two balls are drawn.
 $\therefore n(S) = {}^{5+x}C_2$
Given, probability of red balls drawn = $\frac{5}{14}$
 $\therefore \frac{5}{14} = \frac{{}^5C_2}{{}^{5+x}C_2}$
 $\Rightarrow \frac{5}{14} = \frac{5!}{3!2!} \times \frac{(3+x)!2!}{(5+x)!}$
 $\Rightarrow \frac{5}{14} = \frac{20}{1} \times \frac{1}{(5+x)(4+x)}$
 $\Rightarrow (5+x)(4+x) = \frac{20 \times 14}{5}$
 $\Rightarrow (5+x)(4+x) = 56 \Rightarrow x = 3$
40. Number of ways in which two faulty machines may be detected (depending upon the test done to identify the faulty machines) = ${}^4C_2 = 6$
and Number of favourable cases = 1
[When faulty machines are identified in the first and the second test]
Hence, required probability = $\frac{1}{6}$.
41. Favorable number of cases = ${}^{20}C_1 = 20$
Sample space = ${}^{62}C_1 = 62$
 \therefore Required probability = $\frac{20}{62} = \frac{10}{31}$
42. The number of ways to arrange 7 white and 3 black balls in a row = $\frac{10!}{7!3!} = \frac{10 \cdot 9 \cdot 8}{1 \cdot 2 \cdot 3} = 120$
Numbers of blank places between 7 balls are 6. There is 1 place before first ball and 1 place after last ball. Hence, total number of places are 8.
Hence, 3 black balls are arranged on these 8 places so that no two black balls are together in number of ways
 $= {}^8C_3 = \frac{8 \times 7 \times 6}{1 \times 2 \times 3} = 56$
So required probability = $\frac{56}{120} = \frac{7}{15}$.
43. Since, we have
 $P(A + B) = P(A) + P(B) - P(AB)$
 $\Rightarrow 0.7 = 0.4 + P(B) - 0.2$
 $\Rightarrow P(B) = 0.5$.
44. $0.8 = 0.3 + x - 0.3x \Rightarrow x = \frac{5}{7}$.
45. Since events are mutually exclusive, therefore
 $P(A \cap B) = 0$ i.e., $P(A \cup B) = P(A) + P(B)$
 $\Rightarrow 0.7 = 0.4 + x \Rightarrow x = \frac{3}{10}$
46. Since, $P(A + B + C)$
 $= P(A) + P(B) + P(C)$
 $= \frac{2}{3} + \frac{1}{4} + \frac{1}{6} = \frac{13}{12}$, which is greater than 1.
Hence, the statement is wrong.
48. If $P(A) = P(B)$
As this gives,
 $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
or
 $P(A) = 2P(A) - P(A)$
 $\Rightarrow P(A \cup B) = P(A \cap B)$
49. A: Student who know lesson I
B: Student who know lesson II
 $P(A) = 0.6, P(B) = 0.4, P(A \cap B) = 0.2$
Required probability = $1 - P(A \cup B)$
 $= 1 - [P(A) + P(B) - P(A \cap B)]$
 $= 1 - (0.6 + 0.4 - 0.2)$
 $= 0.2$
 $= \frac{1}{5}$
50. Set of even numbers that can come up on die
 $= \{2, 4, 6\}$
 \therefore Probability of it being either 2 or 4
 $= \frac{1}{3} + \frac{1}{3} = \frac{2}{3}$
51. Here, $P(A) = \frac{3}{6} = \frac{1}{2}, P(B) = \frac{4}{6} = \frac{2}{3}$
and $P(A \cap B) =$ Probability of getting a number greater than 3 and less than 5
 $=$ Probability of getting 4 = $\frac{1}{6}$
 $\therefore P(A \cup B) = P(A) + P(B) - P(A \cap B)$
 $= \frac{1}{2} + \frac{2}{3} - \frac{1}{6} = 1$



52. $n(S) = {}^{10}C_3$
 A: event that minimum of chosen numbers is 3
 B: event that maximum of chosen number is 7.

$$P(A) = \frac{{}^7C_2}{{}^{10}C_3}, P(B) = \frac{{}^6C_2}{{}^{10}C_3}, P(A \cap B) = \frac{{}^3C_1}{{}^{10}C_3}$$

$$\begin{aligned} P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\ &= \frac{{}^7C_2}{{}^{10}C_3} + \frac{{}^6C_2}{{}^{10}C_3} - \frac{{}^3C_1}{{}^{10}C_3} \\ &= \frac{33}{120} \\ &= \frac{11}{40} \end{aligned}$$

53. Let R_1 be the event that the first ball drawn is red,
 B_1 be the event that the first ball drawn is black,
 R_2 be the event that the second ball drawn is red.

Required probability

$$\begin{aligned} &= P(R_1) \cdot P\left(\frac{R_2}{R_1}\right) + P(B_1) \cdot P\left(\frac{R_2}{B_1}\right) \\ &= \frac{4}{10} \times \frac{6}{12} + \frac{6}{10} \times \frac{4}{12} \\ &= \frac{2}{5} \end{aligned}$$

54. Given, $P(A \cup B) = 0.6$ and $P(A \cap B) = 0.2$
 We know that, if A and B are any two events,
 then $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
 $\Rightarrow 0.6 = 1 - P(\bar{A}) + 1 - P(\bar{B}) - 0.2$
 $\Rightarrow P(\bar{A}) + P(\bar{B}) = 2 - 0.8 = 1.2$

55. Given $P(A \cup B) = \frac{3}{5}$ and $P(A \cap B) = \frac{1}{5}$
 We know $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
 $\therefore \frac{3}{5} = 1 - P(\bar{A}) + 1 - P(\bar{B}) - \frac{1}{5}$
 $\therefore 2 - \frac{4}{5} = P(\bar{A}) + P(\bar{B})$
 $\Rightarrow P(\bar{A}) + P(\bar{B}) = \frac{6}{5}$

56. $P(A \cap B') = P(A) - P(A \cap B)$
 $= \frac{4}{5} - \frac{1}{2} = \frac{3}{10}$

57. $P(A \cap B') = P(A) - P(A \cap B)$
 $= 0.7 - 0.3 = 0.4 = \frac{2}{5}$

58. $P(\bar{A} \cap B) = P(B) - P(A \cap B) = y - z$.

59. $P(\bar{A} \cap \bar{B}) = P(A \cup B)'$
 $= 1 - P(A \cup B)$
 $= 1 - P(A) - P(B) + P(A \cap B)$
 $= 1 - 0.25 - 0.50 + 0.14 = 0.39$

60. $P(A' \cap B') = 1 - P(A \cup B)$
 $\Rightarrow P(A \cup B) = \frac{2}{3}$

Now $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

$$\Rightarrow \frac{2}{3} = x + x - \frac{1}{3} \Rightarrow x = \frac{1}{2}$$

61. Since A and B are mutually exclusive,
 $P(A \cup B) = P(A) + P(B)$
 $= \frac{3}{5} + \frac{1}{5} = \frac{4}{5} = 0.8$

62. Probability of getting head = $\frac{1}{2}$

Probability of die showing 3 = $\frac{1}{6}$

Since both events are independent, the
 required probability = $\frac{1}{2} \times \frac{1}{6} = \frac{1}{12}$

63. When two dice are thrown simultaneously,
 $n(S) = 36$
 A: Event that both the numbers on top are
 prime number
 $\therefore A = \{(2, 2), (2, 3), (2, 5), (3, 2), (3, 3), (3, 5),$
 $(5, 2), (5, 3), (5, 5)\}$
 $\therefore n(A) = 9$
 $\therefore P(A) = \frac{9}{36} = \frac{1}{4}$

When two coins are tossed simultaneously,
 $n(S) = 4$

B: Event that we get one head and one tail

$\therefore n(B) = 2$
 $\therefore P(B) = \frac{2}{4} = \frac{1}{2}$

Since both the events are independent of each
 other,

\therefore Required probability = $P(A) \cdot P(B) = \frac{1}{8}$



64. $P(\overline{A} \cap \overline{B}) = P(\overline{A \cup B}) = 1 - P(A \cup B)$
 Since A and B are mutually exclusive, so
 $P(A \cup B) = P(A) + P(B)$
 Hence, required probability = $1 - (0.5 + 0.3)$
 $= 0.2$.

65. Consider option (B)
 $P(A' \cap B') = [1 - P(A)][1 - P(B)]$
 $\Rightarrow P(A' \cap B') = P(A') \cdot P(B')$
 \therefore A and B are independent events.

66. $P(\text{neither A nor B}) = P(\overline{A} \cap \overline{B})$
 $= P(\overline{A}) \cdot P(\overline{B}) = 0.6 \times 0.5$
 $= 0.3$

67. $P(A' \cap B') = 1 - P(A \cup B)$
 $= 1 - [P(A) + P(B) - P(A \cap B)]$
 $= 1 - 1 = 0$

68. Here, $P(X) = 0.3$; $P(Y) = 0.2$
 Now $P(X \cup Y) = P(X) + P(Y) - P(X \cap Y)$
 Since, these are independent events
 $\therefore P(X \cap Y) = P(X) \cdot P(Y)$
 Thus, required probability
 $= 0.3 + 0.2 - 0.06 = 0.44$

69. Let A be the event that a man will live 10 more years.

$\therefore P(A) = \frac{1}{4}$

Let B be the event that his wife will live 10 more years.

$\therefore P(B) = \frac{1}{3}$

\therefore Required probability = $P(A' \cap B')$
 $= P(A') P(B')$
 $= \frac{3}{4} \times \frac{2}{3} = \frac{1}{2}$

70. $P(A) = \frac{3}{8}$ and $P(B) = \frac{1}{2}$

$\therefore P(A) P(B) = \frac{3}{8} \cdot \frac{1}{2} = \frac{3}{16}$
 and $P(A \cap B) = \frac{2}{8} = \frac{1}{4} \neq P(A) \cdot P(B)$
 \therefore A and B are dependent.

71. Since, A and B are independent events

$\therefore P(A \cap B) = P(A) \cdot P(B)$
 $= [1 - P(\overline{A})][1 - P(\overline{B})]$
 $= [1 - 2/3][1 - 2/7]$
 $= \frac{1}{3} \cdot \frac{5}{7} = \frac{5}{21}$

72. $P(A \cup \overline{B}) = 0.8$ and $P(B) = \frac{2}{7} \Rightarrow P(\overline{B}) = \frac{5}{7}$
 $\Rightarrow P(A) + P(\overline{B}) - P(A \cap \overline{B}) = 0.8$
 $\Rightarrow P(A) + \frac{5}{7} - \frac{5}{7} P(A) = 0.8$
 $\Rightarrow \frac{2}{7} P(A) = \frac{3}{35} \Rightarrow P(A) = 0.3$

73. Since $\overline{E_1} \cap \overline{E_2} = \overline{E_1 \cup E_2}$
 and $(E_1 \cup E_2) \cap (\overline{E_1 \cup E_2}) = \phi$
 $\therefore P\{(E_1 \cup E_2) \cap (\overline{E_1 \cup E_2})\} = P(\phi) = 0 < \frac{1}{4}$

74. $P(\overline{A \cup B}) = \frac{1}{6}$
 $\Rightarrow 1 - P(A \cup B) = \frac{1}{6}$
 $\Rightarrow P(A \cup B) = \frac{5}{6}$
 $\Rightarrow P(A) + P(B) - P(A \cap B) = \frac{5}{6}$
 $\Rightarrow \frac{3}{4} + P(B) - \frac{1}{4} = \frac{5}{6} \Rightarrow P(B) = \frac{1}{3}$
 Clearly, $P(A \cap B) = \frac{1}{4} = \frac{3}{4} \times \frac{1}{3} = P(A) P(B)$
 So, A and B are independent.
 Also, $P(A) \neq P(B)$. So, A and B are not equally likely.

75. $P(A \cap B) = \frac{1}{6}$ and $P(\overline{A} \cap \overline{B}) = \frac{1}{3}$
 $\Rightarrow P(A) P(B) = \frac{1}{6}$ and $P(\overline{A}) P(\overline{B}) = \frac{1}{3}$
 $\Rightarrow xy = \frac{1}{6}$ and $(1-x)(1-y) = \frac{1}{3}$,
 where $P(A) = x$, $P(B) = y$
 $\Rightarrow xy = \frac{1}{6}$ and $1 - x - y + \frac{1}{6} = \frac{1}{3}$



$$\Rightarrow xy = \frac{1}{6} \text{ and } x + y = \frac{5}{6}$$

$$\Rightarrow x = \frac{1}{2} \text{ and } y = \frac{1}{3} \text{ or } x = \frac{1}{3} \text{ and } y = \frac{1}{2}$$

76. Let A_i ($i = 1, 2$) denote the event that i^{th} plane hits the target.

Clearly, A_1 and A_2 are independent events.

$$\begin{aligned} \text{Required probability} &= P(\bar{A}_1 \cap A_2) \\ &= P(\bar{A}_1)P(A_2) \\ &= (1 - 0.3)(0.2) = 0.14 \end{aligned}$$

77. Total number of defective items
 $= \frac{2}{100} \times 2500 + \frac{3}{100} \times 3500 + \frac{5}{100} \times 4000$
 $= 355$

Number of defective items from machine C
 $= \frac{5}{100} \times 4000 = 200$

$$\therefore \text{Required probability} = \frac{200}{355} = \frac{40}{71}$$

78. $P[(A \cap (B \cup C))] = P[(A \cap B) \cup (A \cap C)]$
 $= P(A \cap B) + P(A \cap C) - P[(A \cap B) \cap (A \cap C)]$
 $= P(A \cap B) + P(A \cap C) - P(A \cap B \cap C)$

79. $P(B \cap C) = P(B) - [P(A \cap B \cap \bar{C}) + P(\bar{A} \cap B \cap \bar{C})]$
 $= \frac{3}{4} - \frac{2}{3} = \frac{1}{12}$

80. Let A_1 – student passes in Test - I
 A_2 – student passes in Test - II
 A_3 – student passes in Test - III
 A – student is successful
 $A = (A_1 \cap A_2 \cap A_3) \cup (A_1 \cap A_2 \cap \bar{A}_3) \cup (A_1 \cap \bar{A}_2 \cap A_3) \cup (A_1 \cap \bar{A}_2 \cap \bar{A}_3)$

$$\therefore P(A) = P(A_1) \cdot P(A_2) \cdot P(A_3) + P(A_1) \cdot P(A_2) \cdot P(\bar{A}_3) + P(A_1) \cdot P(\bar{A}_2) \cdot P(A_3) + P(A_1) \cdot P(\bar{A}_2) \cdot P(\bar{A}_3)$$

$$\therefore \frac{1}{2} = p \cdot q \cdot \frac{1}{2} + p \cdot (1 - q) \cdot \frac{1}{2} + p \cdot q \cdot \frac{1}{2}$$

$$\therefore p + pq = 1 \Rightarrow p(1 + q) = 1$$

81. Probability of first card to be a king = $\frac{4}{52}$

and probability of also second to be a king

$$= \frac{3}{51}$$

$$\text{Hence, required probability} = \frac{4}{52} \times \frac{3}{51} = \frac{1}{221}$$

82. Required probability = $P(\text{Diamond}) \cdot P(\text{king})$

$$= \frac{13}{52} \cdot \frac{4}{52} = \frac{1}{52}$$

83. Second white ball can draw in two ways.

i. First is white and second is white

$$\text{Probability} = \frac{4}{7} \times \frac{3}{6} = \frac{2}{7}$$

ii. First is black and second is white

$$\text{Probability} = \frac{3}{7} \times \frac{4}{6} = \frac{2}{7}$$

$$\text{Hence, required probability} = \frac{2}{7} + \frac{2}{7} = \frac{4}{7}$$

84. The sample space is [LWW, WLW]

$$\therefore P(LWW) + P(WLW)$$

= Probability that in 5 match series, it is India's second win

$$= P(L)P(W)P(W) + P(W)P(L)P(W)$$

$$= \frac{1}{8} + \frac{1}{8} = \frac{2}{8} = \frac{1}{4}$$

85. Here, $P(A) = \frac{3}{4}$, $P(B) = \frac{4}{5}$

$$\therefore P(\bar{A}) = \frac{1}{4} \text{ and } P(\bar{B}) = \frac{1}{5}$$

\therefore Required probability

$$= P(A) \cdot P(\bar{B}) + P(\bar{A}) \cdot P(B) = \frac{7}{20}$$

86. $P(A) = \frac{4}{5}$, $P(A') = \frac{1}{5}$

$$P(B) = \frac{3}{5}, P(B') = \frac{2}{5}$$

\therefore $P(\text{both are false}) = P(A') \cdot P(B')$

$$= \frac{1}{5} \cdot \frac{2}{5}$$

$$= \frac{2}{25}$$

\therefore $P(\text{at least one of them is true})$

$$= 1 - P(\text{both are false})$$

$$= 1 - \frac{2}{25} = \frac{23}{25}$$

87. Consider the following events:

$A = \text{'X' speaks truth}$, $B = \text{'Y' speaks truth}$.

$$\text{Then, } P(A) = \frac{60}{100} = \frac{3}{5} \text{ and } P(B) = \frac{50}{100} = \frac{1}{2}$$



$$\begin{aligned} \text{Required probability} &= P((A \cap \bar{B}) \cup (\bar{A} \cap B)) \\ &= P(A \cap \bar{B}) + P(\bar{A} \cap B) \\ &= \frac{3}{5} \times \frac{1}{2} + \frac{2}{5} \times \frac{1}{2} = \frac{1}{2} \end{aligned}$$

88. Consider the following events:

$X =$ 'A' speaks truth, $Y =$ 'B' speaks truth

$$\text{Then, } P(X) = \frac{70}{100} = \frac{7}{10} \text{ and } P(Y) = \frac{80}{100} = \frac{4}{5}$$

$$\begin{aligned} \text{Required probability} &= P[(X \cap \bar{Y}) \cup (\bar{X} \cap Y)] \\ &= \frac{7}{10} \times \frac{1}{5} + \frac{3}{10} \times \frac{4}{5} \\ &= \frac{19}{50} = 0.38 \end{aligned}$$

89. Consider the following events:

$A =$ family who owns a car,

$B =$ family who owns a house

$$\text{Required probability} = P(A \cup B) - P(A \cap B)$$

$$= \frac{60+30-20}{100} - \frac{20}{100} = \frac{70-20}{100} = 0.5$$

90. The probability of husband is not selected

$$= 1 - \frac{1}{7} = \frac{6}{7}$$

The probability that wife is not selected

$$= 1 - \frac{1}{5} = \frac{4}{5}$$

The probability that only husband selected

$$= \frac{1}{7} \times \frac{4}{5} = \frac{4}{35}$$

The probability that only wife selected

$$= \frac{1}{5} \times \frac{6}{7} = \frac{6}{35}$$

$$\begin{aligned} \text{Hence, required probability} &= \frac{6}{35} + \frac{4}{35} = \frac{10}{35} \\ &= \frac{2}{7} \end{aligned}$$

91. The probability of students not solving the

$$\text{problem are } 1 - \frac{1}{3} = \frac{2}{3}, 1 - \frac{1}{4} = \frac{3}{4} \text{ and } 1 - \frac{1}{5} = \frac{4}{5}$$

Therefore, the probability that the problem is

$$\text{not solved by any one of them} = \frac{2}{3} \times \frac{3}{4} \times \frac{4}{5} = \frac{2}{5}$$

Hence, the probability that problem is

$$\text{solved} = 1 - \frac{2}{5} = \frac{3}{5}.$$

92. i. This question can also be solved by one student

ii. This question can be solved by two students simultaneously

ii. This question can be solved by three students all together.

$$\text{We have, } P(A) = \frac{1}{2}, P(B) = \frac{1}{4}, P(C) = \frac{1}{6}$$

$$\begin{aligned} \therefore P(A \cup B \cup C) &= P(A) + P(B) + P(C) \\ &\quad - [P(A).P(B) + P(B).P(C) + P(C).P(A)] + \\ &\quad \quad \quad [P(A).P(B).P(C)] \end{aligned}$$

$$= \frac{1}{2} + \frac{1}{4} + \frac{1}{6} - \left[\frac{1}{2} \times \frac{1}{4} + \frac{1}{4} \times \frac{1}{6} + \frac{1}{6} \times \frac{1}{2} \right]$$

$$+ \left[\frac{1}{2} \times \frac{1}{4} \times \frac{1}{6} \right]$$

$$= \frac{33}{48}$$

$$93. P\left(\frac{B}{A}\right) = \frac{P(A \cap B)}{P(A)} = \frac{\left(\frac{1}{10}\right)}{\left(\frac{1}{4}\right)} = \frac{2}{5}.$$

94. For S and T as independent events,

$$P(S/T) = P(S). \text{ Thus, } P(S/T) = 0.3.$$

$$95. P(A/B) = \frac{P(A \cap B)}{P(B)} = \frac{\frac{7}{10}}{\frac{17}{20}} = \frac{7}{10} \times \frac{20}{17} = \frac{14}{17}$$

$$96. P(A \cap B) = P(A) P(B/A)$$

$$\therefore P(A \cap B) = \frac{1}{4} \times \frac{2}{3} = \frac{1}{6}$$

$$\text{Now, } P(A/B) = \frac{P(A \cap B)}{P(B)}$$

$$\Rightarrow \frac{1}{2} = \frac{1}{6} \times \frac{1}{P(B)}$$

$$\Rightarrow P(B) = \frac{1}{3}$$



$$\begin{aligned}
 97. \quad P(B / (A \cup B^c)) &= \frac{P(B \cap (A \cup B^c))}{P(A \cup B^c)} \\
 &= \frac{P(A \cap B)}{P(A) + P(B^c) - P(A \cap B^c)} \\
 &= \frac{P(A) - P(A \cap B^c)}{P(A) + P(B^c) - P(A \cap B^c)} \\
 &= \frac{0.7 - 0.5}{0.8} = \frac{1}{4}
 \end{aligned}$$

98. $S = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$

$\therefore n(E) = 4, n(F) = 4$ and $n(E \cap F) = 3$

$$\therefore P\left(\frac{E}{F}\right) = \frac{P(E \cap F)}{P(F)} = \frac{\frac{3}{8}}{\frac{4}{8}} = \frac{3}{4}$$

99. Event that at least one of them is a boy $\rightarrow A$,
Event that other is girl $\rightarrow B$,
So, required probability

$$P(B/A) = \frac{P(B \cap A)}{P(A)}$$

Now, total cases are 3 (BG, BB, GG)

$$\therefore \frac{P(B \cap A)}{P(A)} = \frac{\frac{1}{3}}{\frac{2}{3}} = \frac{1}{2}$$

....[$\because B \cap A = \{BG\}$ and $A = \{BG, BB\}$]

100. Consider the following events:
 $A =$ Sum of the digits on the selected tickets is 8.

$B =$ Product of the digits on the selected ticket is zero.

There are 14 tickets having product of digits appearing on them as zero. The numbers on such tickets are 00, 01, 02, 03, 04, 05, 06, 07, 08, 09, 10, 20, 30, 40.

$$\therefore P(B) = \frac{14}{50} \text{ and } P(A \cap B) = \frac{1}{50}$$

$$\begin{aligned}
 \therefore \text{Required probability} &= P(A/B) = \frac{P(A \cap B)}{P(B)} \\
 &= \frac{1}{14}
 \end{aligned}$$

101. M: student studying maths
S: student studying science

$$\therefore P(M \cap S) = 40\% = 0.4$$

$$P(M) = 60\% = 0.6$$

Probability of student studying science given the student is already studying maths

$$\begin{aligned}
 &= P(S/M) = P(M \cap S) / P(M) \\
 &= \frac{0.4}{0.6} = \frac{2}{3}
 \end{aligned}$$

$$102. P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\Rightarrow P(A \cap B) = P(A) + P(B) - P(A \cup B)$$

$$\{\because P(A \cap B) = P(A \cup B)\}$$

$$\Rightarrow 2 P(A \cap B) = P(A) + P(B)$$

$$\Rightarrow 2 P(A) \cdot \frac{P(A \cap B)}{P(A)} = P(A) + P(B)$$

$$\Rightarrow 2 P(A) \cdot P\left(\frac{B}{A}\right) = P(A) + P(B)$$

$$103. \text{ We know that } P(A / B) = \frac{P(A \cap B)}{P(B)}$$

Also we know that $P(A \cup B) \leq 1$

$$\Rightarrow P(A) + P(B) - P(A \cap B) \leq 1$$

$$\Rightarrow P(A \cap B) \geq P(A) + P(B) - 1$$

$$\Rightarrow \frac{P(A \cap B)}{P(B)} \geq \frac{P(A) + P(B) - 1}{P(B)}$$

$$\Rightarrow P(A / B) \geq \frac{P(A) + P(B) - 1}{P(B)}$$

$$104. P(E \cap F) = P(E) \cdot P(F)$$

$$\text{Now, } P(E \cap F^c) = P(E) - P(E \cap F)$$

$$= P(E)[1 - P(F)]$$

$$= P(E) \cdot P(F^c)$$

$$\text{and } P(E^c \cap F^c) = 1 - P(E \cup F)$$

$$= 1 - [P(E) + P(F) - P(E \cap F)]$$

$$= [1 - P(E)][1 - P(F)] = P(E^c)P(F^c)$$

$$\text{Also } P\left(\frac{E}{F}\right) = P(E) \text{ and } P\left(\frac{E^c}{F^c}\right) = P(E^c)$$

$$\Rightarrow P\left(\frac{E}{F}\right) + P\left(\frac{E^c}{F^c}\right) = 1.$$



$$105. P\left(\frac{B}{A}\right) = \frac{P(A \cap B)}{P(A)} \Rightarrow \frac{1}{2} = \frac{P(A \cap B)}{1/4}$$

$$\Rightarrow P(A \cap B) = \frac{1}{8}$$

Hence, events A and B are not mutually exclusive.

\therefore Statement II is incorrect.

$$\text{Now, } P\left(\frac{A}{B}\right) = \frac{P(A \cap B)}{P(B)} \Rightarrow P(B) = \frac{1}{2}$$

$$\dots \left[\because P(A \cap B) = \frac{1}{8} = P(A) \cdot P(B) \right]$$

\therefore events A and B are independent events.

$$\therefore P\left(\frac{A^c}{B^c}\right) = \frac{P(A^c \cap B^c)}{P(B^c)} = \frac{P(A^c)P(B^c)}{P(B^c)}$$

$$= \frac{3}{4} \cdot \frac{1}{2} = \frac{3}{8}$$

Hence, statement I is correct.

$$\text{Again } P\left(\frac{A}{B}\right) + P\left(\frac{A}{B^c}\right) = \frac{1}{4} + \frac{P(A \cap B^c)}{P(B^c)}$$

$$= \frac{1}{4} + \frac{P(A) - P(A \cap B)}{P(B^c)}$$

$$= \frac{1}{4} + \frac{\frac{1}{4} - \frac{1}{8}}{\frac{1}{2}}$$

$$= \frac{1}{4} + \frac{1}{4} = \frac{1}{2}$$

Hence, statement III is incorrect.

106. Consider the following events:

S = person is smoker,

NS = person is non smoker,

D = death due to lung cancer

$$P(D) = P(S) \cdot P\left(\frac{D}{S}\right) + P(NS) \cdot P\left(\frac{D}{NS}\right)$$

$$\Rightarrow 0.006 = \frac{20}{100} \times P\left(\frac{D}{S}\right) + \frac{80}{100} \times \frac{1}{10} \times P\left(\frac{D}{S}\right)$$

$$\Rightarrow P\left(\frac{D}{S}\right) = \frac{1000 \times 0.006}{280} = \frac{6}{280} = \frac{3}{140}$$

107. Let E denote the event that a five occurs and A be the event that the man reports it as '6'.

$$\text{Then, } P(E) = \frac{1}{6}, \quad P(E') = \frac{5}{6}$$

$$P(A/E) = \frac{2}{3}, \quad P(A/E') = \frac{1}{3}$$

From Baye's theorem,

$$P(E/A) = \frac{P(E) \cdot P(A/E)}{P(E) \cdot P(A/E) + P(E') \cdot P(A/E')}$$

$$= \frac{\frac{1}{6} \times \frac{2}{3}}{\frac{1}{6} \times \frac{2}{3} + \frac{5}{6} \times \frac{1}{3}}$$

$$= \frac{2}{7}$$

108. Let E_1 be the event that the ball is drawn from bag A, E_2 the event that it is drawn from bag B and E that the ball is red. We have to find $P(E_2/E)$.

Since both the bags are equally likely to be selected, we have $P(E_1) = P(E_2) = \frac{1}{2}$

$$\text{Also } P(E/E_1) = \frac{3}{5}, \quad P(E/E_2) = \frac{5}{9}$$

Hence by Baye's theorem, we have

$$P(E_2/E) = \frac{P(E_2)P(E/E_2)}{P(E_1)P(E/E_1) + P(E_2)P(E/E_2)}$$

$$= \frac{\frac{1}{2} \cdot \frac{5}{9}}{\frac{1}{2} \cdot \frac{3}{5} + \frac{1}{2} \cdot \frac{5}{9}} = \frac{25}{52}$$

109. Let A be the event of selecting bag X, B be the event of selecting bag Y and E be the event of drawing a white ball, the $P(A) = 1/2$, $P(B) = 1/2$, $P(E/A) = 2/5$, $P(E/B) = 4/6 = 2/3$

$\therefore P(E) = P(A)P(E/A) + P(B)P(E/B)$

$$= \frac{1}{2} \cdot \frac{2}{5} + \frac{1}{2} \cdot \frac{2}{3} = \frac{8}{15}$$



110. K = He knows the answers, NK = He randomly ticks the answers, C = He is correct

$$P\left(\frac{K}{C}\right) = \frac{P(K).P\left(\frac{C}{K}\right)}{P(K).P\left(\frac{C}{K}\right) + P(NK).P\left(\frac{C}{NK}\right)}$$

$$= \frac{p \times 1}{p \times 1 + (1-p) \times \frac{1}{5}} = \frac{5p}{4p+1}$$

111. Consider the following events:

$E_1 \rightarrow$ He knows the answer, $E_2 \rightarrow$ He guesses the answer

$A \rightarrow$ He gets the correct answer.

We have,

$$P(E_1) = \frac{90}{100} = \frac{9}{10}, P(E_2) = \frac{1}{10},$$

$$P(A/E_1) = 1, P(A/E_2) = \frac{1}{4}$$

\therefore Required probability = $P(E_2/A)$

$$= \frac{P(E_2)P(A/E_2)}{P(E_1)P(A/E_1) + P(E_2)P(A/E_2)}$$

$$= \frac{\frac{1}{10} \times \frac{1}{4}}{\frac{9}{10} \times 1 + \frac{1}{10} \times \frac{1}{4}} = \frac{1}{37}$$

112. Required probability

$$= \frac{\frac{1}{7} \times \frac{7}{9}}{\frac{1}{7} \times \frac{7}{9} + \frac{3}{7} \times \frac{8}{9} + \frac{2}{7} \times \frac{5}{9} + \frac{1}{7} \times \frac{8}{9}} = \frac{1}{7}$$

113. Required probability = $\frac{(21)!2!}{(22)!} = \frac{1}{11} = \frac{1}{1+10}$

\therefore Odds against = 10 : 1.

114. Required probability = $\frac{0.1}{0.1+0.32}$

$$= \frac{0.1}{0.42} = \frac{5}{21}$$

115. Probability [Person A will die in 30 years]

$$= \frac{8}{8+5}$$

$$\therefore P(A) = \frac{8}{13} \Rightarrow P(\bar{A}) = \frac{5}{13}$$

$$\text{Similarly, } P(B) = \frac{4}{7} \Rightarrow P(\bar{B}) = \frac{3}{7}$$

There are two ways in which one person is alive after 30 years. $\bar{A}B$ and $A\bar{B}$ are independent events.

So, required probability

$$= P(\bar{A}).P(B) + P(A).P(\bar{B})$$

$$= \frac{5}{13} \times \frac{4}{7} + \frac{8}{13} \times \frac{3}{7} = \frac{44}{91}$$

116. The probability of solving the question by

these three students are $\frac{1}{3}$, $\frac{2}{7}$ and $\frac{3}{8}$

respectively.

$$\therefore P(A) = \frac{1}{3}; P(B) = \frac{2}{7}; P(C) = \frac{3}{8}$$

Then, probability of question solved by only one student = $P(A\bar{B}\bar{C})$ or $\bar{A}B\bar{C}$ or $\bar{A}\bar{B}C$

$$= P(A)P(\bar{B})P(\bar{C}) + P(\bar{A})P(B)P(\bar{C})$$

$$+ P(\bar{A})P(\bar{B})P(C)$$

$$= \frac{1}{3} \cdot \frac{5}{7} \cdot \frac{5}{8} + \frac{2}{3} \cdot \frac{2}{7} \cdot \frac{5}{8} + \frac{2}{3} \cdot \frac{5}{7} \cdot \frac{3}{8}$$

$$= \frac{25+20+30}{168} = \frac{25}{56}$$

117. The quadratic equation $ax^2 + bx + c = 0$ has real roots when, $\Delta = b^2 - 4ac \geq 0$


Since a, b, c are chosen from the numbers 2, 3, 5.

6 different equations having distinct coefficients can be formed. Of these, only two equations having $b = 5$ will have real roots.

$$\therefore \text{Required probability} = \frac{2}{6} = \frac{1}{3}$$



118.



A is an event of (x, y) which satisfies $y^2 \leq x$

$$\therefore P(A) = \int_0^1 y \, dx = \int_0^1 \sqrt{x} \, dx = \frac{2}{3}$$

B is an event of (x, y) which satisfies $x^2 \leq y$

$$P(B) = \int_0^1 x \, dy = \int_0^1 x^2 \, dx = \frac{1}{3}$$

$$\therefore P(A \cap B) = \frac{2}{3} - \frac{1}{3} = \frac{1}{3}$$

119. Let, $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ be the given ellipse with eccentricity, $e = \frac{2\sqrt{2}}{3}$

$$\therefore e^2 = 1 - \frac{b^2}{a^2}$$


$$\therefore b^2 = a^2(1 - e^2) \quad \dots(i)$$

Area of ellipse = πab

$$= \pi a \cdot \sqrt{a^2(1 - e^2)} \quad \dots[\text{From (i)}]$$

$$= \pi a^2 \sqrt{1 - e^2}$$

$$= \pi a^2 \sqrt{1 - \frac{8}{9}} = \frac{\pi a^2}{3}$$



Also, radius of the circle = a

$$\therefore \text{Area of circle} = \pi a^2$$

\therefore Probability that point inside the circle lies

$$\text{outside the ellipse} = 1 - \frac{\frac{\pi a^2}{3}}{\pi a^2} = 1 - \frac{1}{3} = \frac{2}{3}$$



Evaluation Test

1. Out of 30 numbers from 1 to 30, three numbers can be chosen in ${}^{30}C_3$ ways.

So, total number of elementary events = ${}^{30}C_3$.

Three consecutive numbers can be chosen in one of the following ways:

(1, 2, 3), (2, 3, 4), ..., (28, 29, 30).

\therefore Number of elementary events in which three numbers are consecutive is 28.

\therefore Probability that the numbers are consecutive

$$= \frac{28}{{}^{30}C_3} = \frac{1}{145}$$

$$\therefore \text{required probability} = 1 - \frac{1}{145} = \frac{144}{145}$$

2. We have,

$$P(E) + P(F) - 2P(E \cap F) = \frac{11}{25} \text{ and}$$

$$P(\bar{E} \cap \bar{F}) = \frac{2}{25}$$

$$\Rightarrow P(E) + P(F) - 2P(E)P(F) = \frac{11}{25} \text{ and}$$

$$P(\bar{E})P(\bar{F}) = \frac{2}{25}$$

$$\Rightarrow x + y - 2xy = \frac{11}{25} \text{ and } 1 - x - y + xy = \frac{2}{25},$$

Where, $P(E) = x$ and $P(F) = y$

$$\Rightarrow x + y + 2 - 2x - 2y = \frac{11}{25} + 2 \times \frac{2}{25}$$

....[On eliminating xy]

$$\Rightarrow x + y = \frac{7}{5} \Rightarrow y = \frac{7}{5} - x$$

$$\text{Substituting } y = \frac{7}{5} - x \text{ in } 1 - x - y + xy = \frac{2}{25},$$

we get

$$1 - \frac{7}{5} + x\left(\frac{7}{5} - x\right) = \frac{2}{25}$$

$$\Rightarrow 25x^2 - 35x + 12 = 0$$

$$\Rightarrow x = \frac{3}{5}, \frac{4}{5}$$

When $x = \frac{3}{5}$, $y = \frac{4}{5}$ and $y = \frac{3}{5}$ for $x = \frac{4}{5}$

Hence, $P(E) = \frac{3}{5}$, $P(F) = \frac{4}{5}$ or $P(E) = \frac{4}{5}$,

$$P(F) = \frac{3}{5}$$



3. Let A denote the event that each American man is seated adjacent to his wife and B denote the event that Indian man is seated adjacent to his wife. Then,
required probability = $P(B/A)$

$$\begin{aligned} & \text{Number of ways in which Indian man} \\ & \text{sits adjacent to his wife when each} \\ & \text{man is sited adjacent to his wife} \\ & = \frac{\text{Number of ways in which each} \\ & \text{American man is seated} \\ & \text{adjacent to his wife}}{\text{Number of ways in which each} \\ & \text{American man is seated} \\ & \text{adjacent to his wife}} \\ & = \frac{(2!)^5 \times (5-1)!}{(2!)^4 (6-1)!} = \frac{2}{5} \end{aligned}$$

4. We have 13 denominations Ace, 2, 3, 4, ..., 10, J, Q, K. For selecting exactly one pair, we select first any 3 denominations, 2 cards from 1 and one each from the other two

$$\begin{aligned} \text{Thus, favourable ways} &= {}^{13}C_3 \cdot 3 \cdot {}^4C_2 \cdot {}^4C_1 \cdot {}^4C_1 \\ \text{Total ways} &= {}^{52}C_4 \end{aligned}$$

$$\begin{aligned} \therefore \text{required probability} &= \frac{13 \cdot 12 \cdot 11 \cdot 3 \cdot 6 \cdot 4 \cdot 4 \cdot 24}{6 \cdot 52 \cdot 51 \cdot 50 \cdot 49} \\ &= \frac{6336}{20825} = 0.3042 = 0.3 \end{aligned}$$

5. Let event A that minimum of the chosen number is 3 and B be the event that maximum of the chosen number is

$$\therefore P(A) = P(\text{choosing 3 and two other numbers from 4 to 10})$$

$$= \frac{{}^7C_2}{{}^{10}C_3} = \frac{7 \times 6 \times 3}{10 \times 9 \times 8} = \frac{7}{40}$$

$$P(B) = P(\text{choosing 7 and choosing two other numbers from 1 to 6})$$

$$= \frac{{}^6C_2}{{}^{10}C_3} = \frac{6 \times 5 \times 3}{10 \times 9 \times 7} = \frac{1}{8}$$

$$P(A \cap B) = P(\text{choosing 3 and 7 and one other from 4 to 6})$$

$$= \frac{3}{{}^{10}C_3} = \frac{3 \times 3 \times 2}{10 \times 9 \times 8} = \frac{1}{40}$$

$$\begin{aligned} \therefore P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\ &= \frac{7}{40} + \frac{1}{8} - \frac{1}{40} = \frac{11}{40} \end{aligned}$$

6. In the 22nd century there are 25 leap years viz. 2100, 2104, ..., 2196 and 75 non-leap years. Consider the following events:

E_1 = Selecting a leap year from 22nd century

E_2 = Selecting a non-leap year from 22nd century

A = There are 53 Sundays in a year of 22nd century

We have,

$$\begin{aligned} P(E_1) &= \frac{25}{100}, P(E_2) \\ &= \frac{75}{100} \end{aligned}$$

$$\begin{aligned} P(A/E_1) &= \frac{2}{7} \text{ and } P(A/E_2) \\ &= \frac{1}{7} \end{aligned}$$

$$\begin{aligned} \therefore \text{Required probability} &= P(A) \\ &= P((A \cap E_1) \cup (A \cap E_2)) \\ &= P(A \cap E_1) + P(A \cap E_2) \\ &= P(E_1) P(A/E_1) + P(E_2) P(A/E_2) \\ &= \frac{25}{100} \times \frac{2}{7} + \frac{75}{100} \times \frac{1}{7} \\ &= \frac{5}{28} \end{aligned}$$

7. We know that the probability of occurrence of an event is always less than or equal to 1 and it is given that $P(A \cup B \cup C) \geq 0.75$

$$\begin{aligned} \therefore 0.75 &\leq P(A \cup B \cup C) \leq 1 \\ &\Rightarrow 0.75 \leq P(A) + P(B) + P(C) - P(A \cap B) \\ &\quad - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C) \leq 1 \\ &\Rightarrow 0.75 \leq 0.3 + 0.4 + 0.8 - 0.08 - P(B \cap C) \\ &\quad - 0.28 + 0.09 \leq 1 \\ &\Rightarrow 0.75 \leq 1.23 - P(B \cap C) \leq 1 \\ &\Rightarrow -0.48 \leq -P(B \cap C) \leq -0.23 \\ &\Rightarrow 0.23 \leq P(B \cap C) \leq 0.48 \end{aligned}$$

8. From the tree diagram,


$$\begin{aligned} P(B_G) &= \frac{4}{5} \times \frac{3}{4} \times \frac{3}{4} + \frac{4}{5} \times \frac{1}{4} \times \frac{1}{4} + \frac{1}{5} \times \frac{3}{4} \times \frac{1}{4} + \frac{1}{5} \times \frac{1}{4} \times \frac{3}{4} \\ &= \frac{23}{40} \end{aligned}$$



$$P\left(\frac{B_G}{G}\right) = \frac{3}{4} \times \frac{3}{4} + \frac{1}{4} \times \frac{1}{4} = \frac{5}{8}$$

$$\therefore P(B_G \cap G) = P(G) P\left(\frac{B_G}{G}\right)$$

$$\Rightarrow P(B_G \cap G) = \frac{4}{5} \times \frac{5}{8} = \frac{1}{2}$$



$$\begin{aligned} \therefore \text{Required probability} &= P\left(\frac{G}{B_G}\right) = \frac{P(B_G \cap G)}{P(B_G)} \\ &= \frac{\frac{1}{2}}{\frac{23}{40}} = \frac{20}{23} \end{aligned}$$



Hints



Classical Thinking

- 'Bombay is the capital of India' is a statement. The other options are exclamatory and interrogative sentences.
- 'Two plus two is four' is a statement. The other options are imperative sentences.
- Even though $2 = 3$ is false, it is a statement in logic with truth value F.
- $\sim q$: Ram studies on holiday, 'and' is expressed by ' \wedge ' symbol
 \therefore Symbolic form is $p \wedge \sim q$.
- p : There are clouds in the sky, $\sim q$: It is not raining, 'and' is expressed by ' \wedge ' symbol.
 $\therefore p \wedge \sim q$
- $\sim p$: The sun has not set, $\sim q$: The moon has not risen, 'or' is expressed by ' \vee ' symbol.
 $\therefore \sim p \vee \sim q$
- $\sim p$: Rohit is short, 'or' is expressed by ' \vee ' symbol, 'and' is expressed by ' \wedge ' symbol.
- p : Candidates are present,
 q : Voters are ready to vote
 r : Ballot papers $\Rightarrow \sim r$: no Ballot papers
'and' and 'but' are represented by ' \wedge ' symbol.
- $\sim p$: She is not beautiful, ' \vee ' indicates 'or'.
- $\sim p$: Ram is not lazy, $\sim q$: Ram does not fail in the examination, ' \vee ' indicates 'or'.
- "Implies" is expressed as ' \rightarrow '.
 \therefore symbolic form is $p \rightarrow q$
- ($\sim d$: Driver is not drunk) implies ($\sim a$: He cannot meet with an accident).
- "if and only if" is expressed as ' \leftrightarrow '.
 \therefore symbolic form is $a \leftrightarrow b$.
- p : A, B, C, are distinct points
 q : Points are collinear
 r : Points form a triangle
 $\therefore p$ implies (q or r) i.e. $p \rightarrow (q \vee r)$
- ' $m \rightarrow n$ ' means 'If m then n ',
 \therefore option (C) is correct.

- Let p : x^2 is not even,
 q : x is not even
Converse of $p \rightarrow q$ is $q \rightarrow p$
i.e., If x is not even then x^2 is not even
- Converse of $p \rightarrow q$ is $q \rightarrow p$.
- Let p : $x > y$
 q : $x + a > y + a$
Converse of $p \rightarrow q$ is $q \rightarrow p$
i.e., If $x + a > y + a$, then $x > y$
- Let p : You access the internet
 q : You have to pay the charges
Given statement is written symbolically as,
 $p \rightarrow q$
Inverse of $p \rightarrow q$ is $\sim p \rightarrow \sim q$
i.e. If you do not access the internet then you do not have to pay the charges.
- Contrapositive of $p \rightarrow q$ is $\sim q \rightarrow \sim p$.
- $\sim p$: Sita does not get promotion and ' \leftrightarrow ' symbol indicates 'if and only if'.
- r : It is raining, c : I will go to college.
The given statement is $r \rightarrow c \equiv \sim c \rightarrow \sim r$

36.

p	q	$p \wedge q$	$(p \wedge q) \rightarrow p$
T	T	T	T
T	F	F	T
F	T	F	T
F	F	F	T

37.

p	q	$\sim q$	$p \wedge q$	$p \rightarrow \sim q$	$(p \wedge q) \wedge (p \rightarrow \sim q)$
T	T	F	T	F	F
T	F	T	F	T	F
F	T	F	F	T	F
F	F	T	F	T	F

38.

p	q	$\sim q$	$p \wedge \sim q$	$\sim(p \wedge \sim q)$	$p \rightarrow \sim(p \wedge \sim q)$
T	T	F	F	T	T
T	F	T	T	F	F
F	T	F	F	T	T
F	F	T	F	T	T



39.

p	q	$p \rightarrow q$	$\sim p$	$\sim q$	$\sim p \rightarrow \sim q$	$(p \rightarrow q) \leftrightarrow (\sim p \rightarrow \sim q)$
T	T	T	F	F	T	T
T	F	F	F	T	T	F
F	T	T	T	F	F	F
F	F	T	T	T	T	T

40. Option (C) is a true statement, since, $x = 3 \in \mathbb{N}$ satisfies $x + 5 = 8$.

41. Option (D) is the required true statement since $x = 6 \in \mathbb{W}$ satisfies $x^2 - 4 = 32$

43. p: Manoj has the job, q: he is not happy
Symbolic form is $p \wedge q$.
Its dual is $p \vee q$.
 \therefore Manoj has the job or he is not happy.

44. $\sim(p \wedge q) \equiv \sim p \vee \sim q$

45. $\sim[p \vee (\sim q)] \equiv \sim p \wedge \sim(\sim q) \equiv \sim p \wedge q$

46. p : I like Mathematics

q : I like English.

$\sim(p \wedge q) \equiv \sim p \vee \sim q$

\therefore Option (D) is correct.

47. We know that,

$p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$

$\therefore \sim(p \leftrightarrow q) \equiv \sim[(p \rightarrow q) \wedge (q \rightarrow p)]$
 $\equiv \sim(p \rightarrow q) \vee \sim(q \rightarrow p)$
.....[By Demorgan's Law]
 $\equiv (p \wedge \sim q) \vee (q \wedge \sim p)$
.....[$\because \sim(p \rightarrow q) = p \wedge \sim q$]

48. p : It is Sunday

q : It is a holiday

\therefore Symbolic form $p \rightarrow q$

$\sim(p \rightarrow q) \equiv p \wedge \sim q$

i.e. It is Sunday, but it is not a holiday

49. Given statement is ' $\forall x \in \mathbb{N}, x + 5 > 4$ '

$\therefore \sim[\forall x \in \mathbb{N}, x + 5 > 4]$
 $\equiv \exists x \in \mathbb{N}$, such that $x + 5 \leq 4$
i.e., there exists a natural number x , for which $x + 5 \leq 4$

51. Current will flow in the circuit if switch p and q are closed or switch r is closed.

It is represented by

$(p \wedge q) \vee r$.

\therefore option (A) is correct.



Critical Thinking

- 'Incorrect statement' means a statement in logic with truth value false.
Options (A) and (C) are not statements in logic.
Option (D) has truth value True.
Option (B) is a statement in logic with truth value false.
- p: One being lucky,
q: One should stop working
 \therefore Symbolic form: $(p \vee \sim p) \wedge \sim q$
- p: Physics is interesting.
q: Physics is difficult.
 \therefore Symbolic form: $\sim(\sim p \vee q)$
- p: Intelligent persons are polite.
q: Intelligent persons are helpful.
 \therefore Symbolic form: $\sim(\sim p \wedge \sim q)$
- $\sim p \wedge (q \vee \sim r)$ and $(p \rightarrow q) \wedge r$
 $\therefore \sim T \wedge (T \vee \sim F)$ and $(T \rightarrow T) \wedge F$
 $\Rightarrow F \wedge (T \vee T)$ and $(T \wedge F)$
 $\Rightarrow F \wedge T$ and $T \wedge F \Rightarrow F$ and F
- $(\sim p \vee q) \leftrightarrow \sim(p \wedge q)$ and $\sim p \leftrightarrow (p \rightarrow \sim q)$
 $\therefore (\sim F \vee F) \leftrightarrow \sim(F \wedge F)$ and $\sim F \leftrightarrow (F \rightarrow \sim F)$
 $\Rightarrow (T \vee F) \leftrightarrow \sim F$ and $T \leftrightarrow (F \rightarrow T)$
 $\Rightarrow T \leftrightarrow T$ and $T \leftrightarrow T$
 $\Rightarrow T$ and T
- $(p \rightarrow q) \leftrightarrow (\sim q \rightarrow \sim p)$ and $(\sim p \vee q) \wedge (\sim q \vee p)$
 $\therefore (T \rightarrow F) \leftrightarrow (\sim F \rightarrow \sim T)$ and $(\sim T \vee F) \wedge (\sim F \vee T)$
 $\Rightarrow F \leftrightarrow (T \rightarrow F)$ and $(F \vee F) \wedge (T \vee T)$
 $\Rightarrow F \leftrightarrow F$ and $F \wedge T \Rightarrow T$ and F
- $p \wedge q \equiv F \wedge T \equiv F$
 $p \vee \sim q \equiv F \vee \sim T \equiv F \vee F \equiv F$
 $q \rightarrow p \equiv T \rightarrow F \equiv F$
 $p \rightarrow q \equiv F \rightarrow T \equiv T$
- $\sim p \rightarrow \sim q \equiv \sim F \rightarrow \sim T \equiv T \rightarrow F \equiv F$
 $p \rightarrow (q \wedge p) \equiv F \rightarrow (T \wedge F) \equiv F \rightarrow F \equiv T$
 $p \rightarrow \sim q \equiv F \rightarrow \sim T \equiv F \rightarrow F \equiv T$
 $q \rightarrow \sim p \equiv T \rightarrow \sim F \equiv T \rightarrow T \equiv T$
- Consider option (C)
 $(p \vee q) \wedge (p \vee r) \equiv (T \vee T) \wedge (T \vee F)$
 $\equiv T \wedge T$
 $\equiv T$
 \therefore option (C) is correct.



11.

p	q	$\sim q$	$\sim q \vee p$	$p \leftrightarrow q$	$p \rightarrow q$	$q \rightarrow p$
T	T	F	T	T	T	T
T	F	T	T	F	F	T
F	T	F	F	F	T	F
F	F	T	T	T	T	T

Alternate Method:

$\sim q \vee p$: F

$\therefore \sim q$ is F, p is F

i.e., q is T, p is F

$\therefore p \rightarrow q \equiv F \rightarrow T \equiv T$

12. p: Seema solves a problem

q: She is happy

i. $p \rightarrow q$ ii. $\sim p \rightarrow \sim q$

iii. $\sim q \rightarrow \sim p$ iv. $q \rightarrow p$

(i) and (iii) have the same meaning,

(ii) and (iv) have the same meaning.

13. i. $b \rightarrow r$

ii. $\sim b \rightarrow \sim r$

iii. $r \rightarrow b$

iv. $\sim r \rightarrow \sim b$

(i) and (iv) are the same and (ii) and (iii) are the same.

14. $p \wedge (p \rightarrow q)$

$\equiv p \wedge (\sim p \vee q)$ [Conditional law]

$\equiv (p \wedge \sim p) \vee (p \wedge q)$ [Distributive law]

$\equiv F \vee (p \wedge q)$ [Complement law]

$\equiv p \wedge q$ [Identity law]

15. $\sim[p \rightarrow (p \vee \sim q)] \equiv \sim[\sim p \vee (p \vee \sim q)]$

....[$\because p \rightarrow q \equiv \sim p \vee q$]

$\equiv p \wedge \sim[p \vee (\sim q)]$

$\equiv p \wedge [\sim p \wedge \sim(\sim q)]$

$\equiv p \wedge (\sim p \wedge q)$

16. $(\sim q) \rightarrow (\sim p)$ is contrapositive of $p \rightarrow q$ and hence both are logically equivalent of each other.

17.

p	$\sim p$	$\sim(\sim p)$	$\sim(\sim p) \leftrightarrow p$
T	F	T	T
F	T	F	T

All the entries in the last column of the above truth table is T.

$\therefore \sim(\sim p) \leftrightarrow p$ is a tautology.

18.

p	q	r	$\sim p$	$\sim q$	$\sim p \wedge \sim q$	$q \wedge r$	$(\sim p \wedge \sim q) \wedge (q \wedge r)$
T	T	T	F	F	F	T	F
T	T	F	F	F	F	F	F
T	F	T	F	T	F	F	F
T	F	F	F	T	F	F	F
F	T	T	T	F	F	T	F
F	T	F	T	F	F	F	F
F	F	T	T	T	T	F	F
F	F	F	T	T	T	F	F

\therefore Given statement is contradiction.

19. Consider option (C)

p	q	$\sim q$	$p \rightarrow q$	$\sim(p \rightarrow q)$	$p \wedge \sim q$	$\sim(p \rightarrow q) \leftrightarrow (p \wedge \sim q)$
T	T	F	T	F	F	T
T	F	T	F	T	T	T
F	T	F	T	F	F	T
F	F	T	T	F	F	T

\therefore option (C) is correct.

20. consider option (B)

P	q	$\sim q$	$p \wedge \sim q$	$p \rightarrow q$	$(p \wedge \sim q) \wedge (p \rightarrow q)$
T	T	F	F	T	F
T	F	T	T	F	F
F	T	F	F	T	F
F	F	T	F	T	F

\therefore option (B) is correct.

21. Consider option (B)

p	q	$\sim p$	$\sim q$	$p \wedge q$	$\sim p \rightarrow \sim q$	$(p \wedge q) \leftrightarrow (\sim p \rightarrow \sim q)$
T	T	F	F	T	T	T
T	F	F	T	F	T	F
F	T	T	F	F	F	T
F	F	T	T	F	T	F

\therefore option (B) is correct.

22. Since, $x = 4, 5, 7, 9$ satisfies $x + 1 \leq 10$

\therefore option (B) is correct.

23. Option (A) is the true statement since square of every natural number is positive.

24. Option (C) is false, since for every natural number the statement $x - 1 \geq 0$ is always true.

25. Dual of $(p \vee q) \vee s$ is $(p \wedge q) \wedge s$.

27. Negation of $(p \vee q) \wedge (\sim q \wedge r)$ is

$\sim[(p \vee q) \wedge (\sim q \wedge r)]$

$\equiv \sim(p \vee q) \vee \sim(\sim q \wedge r)$

$\equiv (\sim p \wedge \sim q) \vee [\sim(\sim q) \vee \sim r]$

$\equiv (\sim p \wedge \sim q) \vee (q \vee \sim r)$



28. $\sim[p \vee (\sim q \wedge \sim p)]$
 $\equiv \sim p \wedge \sim(\sim q \wedge \sim p)$...[By De Morgan's law]
 $\equiv \sim p \wedge [\sim(\sim q) \vee \sim(\sim p)]$
 $\equiv \sim p \wedge (q \vee p)$
 $\equiv (\sim p \wedge q) \vee (\sim p \wedge p)$
 ...[Distributive property]
 $\equiv (\sim p \wedge q) \vee F$...[Complement law]
 $\equiv \sim p \wedge q$...[Identity law]
29. $\sim[p \rightarrow (p \vee \sim q)] \equiv p \wedge \sim[p \vee (\sim q)]$
 $\equiv p \wedge (\sim p \wedge q)$
30. $\sim[\exists x \in \mathbb{R}, \text{ such that } x^2 + 3 > 0]$
 $= \forall x \in \mathbb{R}, x^2 + 3 \leq 0$
31. p: Saral Mart does not reduce the prices.
 q: I will not shop there any more.
 Symbolic form is $p \rightarrow q$
 $\sim(p \rightarrow q) \equiv p \wedge \sim q$
 i.e. Saral Mart does not reduce the prices and still I will shop there.
36. The symbolic form of circuit is
 $(p \wedge q) \vee (\sim p \wedge q) \equiv (p \vee \sim p) \wedge q$
 $\equiv T \wedge q$
 $\equiv q$
37. The symbolic form of circuit is
 $[(\sim p \wedge \sim q) \vee p \vee q] \wedge r$
 $\equiv [\sim(p \vee q) \vee (p \vee q)] \wedge r$
 $\equiv T \wedge r$
 $\equiv r$

**Competitive Thinking**

2. Man is not rich : $\sim q$
 Man is not happy : $\sim p$
 \therefore The symbolic representation of the given statement is $\sim q \rightarrow \sim p$.
3. $\sim p$: Ram is not rich
 $\sim q$: Ram is not successful
 $\sim r$: Ram is not talented
 \therefore The symbolic form of the given statement is $\sim p \wedge \sim q \wedge \sim r$.
4. "Not a correct statement" means it is a statement whose truth value is false.
 Option (A) is not a statement.
 Options (C) and (D) are statements with truth value true.
 ' $\sqrt{3}$ is a prime' is false statement.
 Hence, option (B) is correct.
5. The symbol $p \wedge q$ means
 Mathematics is interesting and Mathematics is difficult.
6. p : roses are red
 q : The sun is a star
 $(\sim p) \vee q$: roses are not red or the sun is a star.
7. $\sim p$: Boys are not playing
 The symbol ' \vee ' means 'or'.
8. Consider option (C),
 $(p \wedge \sim q) \rightarrow q \equiv (T \wedge \sim T) \rightarrow T$
 $\equiv (T \wedge F) \rightarrow T$
 $\equiv F \rightarrow T$
 $\equiv T$
 \therefore option (C) is correct.
- 9.
- | p | q | $\sim p$ | $\sim p \vee q$ | $p \rightarrow (\sim p \vee q)$ |
|---|---|----------|-----------------|---------------------------------|
| T | T | F | T | T |
| T | F | F | F | F |
| F | T | T | T | T |
| F | F | T | T | T |
- \therefore From the table $p \rightarrow (\sim p \vee q)$ is false when p is true and q is false.
10. Since, $(p \wedge \sim q) \rightarrow (\sim p \vee r) \equiv F$
 $\Rightarrow p \wedge \sim q \equiv T$ and $\sim p \vee r \equiv F$
 $\Rightarrow p \equiv T, \sim q \equiv T$ and $\sim p \equiv F, r \equiv F$
 $\Rightarrow p \equiv T, q \equiv F, r \equiv F$
 \therefore The truth values of p, q and r are T, F, F respectively.
11. Since, both the given statements p and q have truth values T,
 $\therefore p \rightarrow q \equiv T \rightarrow T \equiv T$, and
 $p \leftrightarrow q \equiv T \leftrightarrow T \equiv T$
12. Contrapositive of $(p \vee q) \rightarrow r$ is
 $\sim r \rightarrow \sim(p \vee q)$ i.e. $\sim r \rightarrow \sim p \wedge \sim q$
13. Given $p \rightarrow q$
 Its contrapositive is $\sim q \rightarrow \sim p$
 and its converse is $\sim p \rightarrow \sim q$
14. Let p : Ram secures 100 marks in maths
 q : Ram will get a mobile
 Converse of $p \rightarrow q$ is $q \rightarrow p$
 i.e., If Ram will get a mobile, then he secures 100 marks in maths.



15. Inverse of $q \rightarrow p$ is $\sim q \rightarrow \sim p$
i.e., If a triangle is not equiangular then it is not equilateral.

16. Let p : It is raining
 q : I will not come
Contrapositive of $p \rightarrow q$ is $\sim q \rightarrow \sim p$
i.e., If I will come, then it is not raining.

17. Let $p = x$ is a prime number, $q = x$ is odd.
Contrapositive of $p \rightarrow q$ is $\sim q \rightarrow \sim p$

18. p : The weather is fine.
 q : My friends will come and we will go for a picnic.

\therefore Statement is $p \rightarrow q$
Contrapositive of $p \rightarrow q$ is $\sim q \rightarrow \sim p$
i.e., if my friends do not come or we do not go for a picnic then weather will not be fine.

19. Let p : x is prime number
 q : x is odd
 \therefore Statement is $p \rightarrow q$
Converse of $p \rightarrow q$ is $q \rightarrow p$
Contrapositive of $q \rightarrow p$ is $\sim p \rightarrow \sim q$.

21.

1	2	3	4	5	6
p	q	$\sim p$	$\sim p \wedge q$	$q \rightarrow p$	$\sim(q \rightarrow p)$
T	T	F	F	T	F
T	F	F	F	T	F
F	T	T	T	F	T
F	F	T	F	T	F

The entries in the columns 4 and 6 are identical.

$\therefore \sim p \wedge q \equiv \sim(q \rightarrow p)$

22. Consider option (B)
 $(p \vee q) \wedge \sim p \equiv (p \wedge \sim p) \vee (q \wedge \sim p)$
 $\equiv F \vee (q \wedge \sim p)$
 $\equiv q \wedge \sim p$
 $\equiv \sim p \wedge q$

23. $(p \wedge q) \vee (\sim q \wedge p) \equiv (p \wedge q) \vee (p \wedge \sim q)$
 $\equiv p \wedge (q \vee \sim q)$
 $\equiv p \wedge T \equiv p$

24. $\sim(p \vee q) \vee (\sim p \wedge q)$
 $\equiv (\sim p \wedge \sim q) \vee (\sim p \wedge q)$
 $\equiv \sim p \wedge (\sim q \vee q)$
 $\equiv \sim p \wedge T$
 $\equiv \sim p$

25. $(p \wedge \sim q) \vee q \vee (\sim p \wedge q)$
 $\equiv [(p \vee q) \wedge (\sim q \vee q)] \vee (\sim p \wedge q)$
 $\equiv [(p \vee q) \wedge T] \vee (\sim p \wedge q)$
 $\equiv (p \vee q) \vee (\sim p \wedge q)$
 $\equiv (p \vee q \vee \sim p) \wedge (p \vee q \vee q)$
 $\equiv (T \vee q) \wedge (p \vee q) \equiv T \wedge (p \vee q)$
 $\equiv p \vee q$

26.

1	2	3	4	5	6
p	q	$q \rightarrow p$	$p \rightarrow (q \rightarrow p)$	$q \vee p$	$p \rightarrow (q \vee p)$
T	T	T	T	T	T
T	F	T	T	T	T
F	T	F	T	T	T
F	F	T	T	F	T

The entries in the columns 4 and 6 are identical.

$\therefore p \rightarrow (q \rightarrow p) \equiv p \rightarrow (p \vee q)$

27. $p \rightarrow (\sim q) \equiv \sim p \vee \sim q$
 $\equiv \sim q \vee \sim p$

28. $\sim(p \vee q) \equiv \sim p \vee \sim q$ is not true as it contradicts De Morgan's law.
 \therefore option (D) is not true.

29. $p \wedge (\sim p \wedge q) \equiv (p \wedge \sim p) \wedge q$
 $\equiv F \wedge q$
 $\equiv F$

30.

p	$\sim p$	$p \rightarrow \sim p$	$\sim p \rightarrow p$	$(p \rightarrow \sim p) \wedge (\sim p \rightarrow p)$
T	F	F	T	F
F	T	T	F	F

31.

p	q	$\sim p$	$\sim q$	$(p \wedge \sim q)$	$(\sim p \wedge q)$	$(p \wedge \sim q) \wedge (\sim p \wedge q)$
T	T	F	F	F	F	F
T	F	F	T	T	F	F
F	T	T	F	F	T	F
F	F	T	T	F	F	F

\therefore Given statement is contradiction.

32. Since, $p \vee \sim p \equiv T$
 $\therefore (\sim q \wedge p) \vee (p \vee \sim p) \equiv (\sim q \wedge p) \vee T \equiv T$
 $\therefore (\sim q \wedge p) \vee (p \vee \sim p)$ is a tautology.

33. Consider option (C)

A	B	$A \rightarrow B$	$A \wedge (A \rightarrow B)$	$[A \wedge (A \rightarrow B)] \rightarrow B$
T	T	T	T	T
T	F	F	F	T
F	T	T	F	T
F	F	T	F	T

\therefore option (C) is correct.



34. Consider option (C)

p	q	$q \rightarrow p$	$\sim p$	$\sim p \leftrightarrow q$	$(q \rightarrow p) \vee (\sim p \leftrightarrow q)$
T	T	T	F	F	T
T	F	T	F	T	T
F	T	F	T	T	T
F	F	T	T	F	T

35. $p \rightarrow q$ is logically equivalent to $\sim q \rightarrow \sim p$
 $\therefore (p \rightarrow q) \leftrightarrow (\sim q \rightarrow \sim p)$ is tautology
 But, it is given contradiction.
 Hence, it is false statement.

36.

1	2	3	4	5	6
p	q	$\sim q$	$p \leftrightarrow \sim q$	$\sim(p \leftrightarrow \sim q)$	$p \leftrightarrow q$
T	T	F	F	T	T
T	F	T	T	F	F
F	T	F	T	F	F
F	F	T	F	T	T

The entries in the columns 5 and 6 are identical.

$\therefore \sim(p \leftrightarrow \sim q) \equiv p \leftrightarrow q$

37.

p	q	$\sim p$	$p \rightarrow q$	$\sim p \rightarrow q$	$(\sim p \rightarrow q) \rightarrow q$	$(p \rightarrow q) \rightarrow [(\sim p \rightarrow q) \rightarrow q]$
T	T	F	T	T	T	T
T	F	F	F	T	F	T
F	T	T	T	T	T	T
F	F	T	T	F	T	T

38. Option (C) is the correct answer since there exists a real number $x = 0$, such that $x^2 = 0$. Zero is neither positive nor negative.

39. Dual of $\sim p \wedge (q \vee c) = \sim p \vee (q \wedge t)$

40. Negation of $q \vee \sim(p \wedge r)$ is
 $\sim[q \vee \sim(p \wedge r)] \equiv \sim q \wedge \sim(\sim(p \wedge r))$
 $\equiv \sim q \wedge (p \wedge r)$

41. $\sim[(p \vee \sim q) \wedge q] \equiv \sim(p \vee \sim q) \vee \sim q$
[De Morgan's Law]
 $\equiv (\sim p) \wedge [\sim(\sim q)] \vee \sim q$
 $\equiv (\sim p \wedge q) \vee \sim q$

42. p : A is rich, q : A is silly
 $\therefore \sim(p \wedge q) \equiv \sim p \vee \sim q$

43. $\sim(p \wedge q) \equiv \sim p \vee \sim q$

44. p : 72 is divisible by 2.
 q : 72 is divisible by 3.
 \therefore Statement is $p \wedge q$
 $\sim(p \wedge q) \equiv \sim p \vee \sim q$

45. $\sim(p \vee q) \equiv (\sim p) \wedge (\sim q)$
 i.e., 7 is greater than 4 and Paris is not in France.

46. $\sim[\sim s \vee (\sim r \wedge s)]$
 $\equiv \sim(\sim s) \wedge \sim(\sim r \wedge s)$ [De Morgan's Law]
 $\equiv s \wedge (r \vee \sim s)$
 $\equiv (s \wedge r) \vee (s \wedge \sim s)$ [Distributive property]
 $\equiv (s \wedge r) \vee F$ [Complement law]
 $\equiv s \wedge r$ [Identity law]

47. $p \rightarrow q \equiv \sim p \vee q$
 $\therefore \sim(p \rightarrow q) \equiv p \wedge \sim q$

48. Since, $p \rightarrow q \equiv \sim p \vee q$
 $\therefore \sim p \rightarrow q \equiv p \vee q$
 $\therefore \sim(\sim p \rightarrow q) \equiv \sim(p \vee q)$
 $\equiv \sim p \wedge \sim q$

49. $\sim[p \rightarrow (\sim p \vee q)] \equiv p \wedge \sim(\sim p \vee q)$
 $\equiv p \wedge (p \wedge \sim q)$
 $\equiv (p \wedge p) \wedge \sim q$
 $\equiv p \wedge \sim q$

50. Since, $p \rightarrow q \equiv \sim p \vee q$
 $\therefore \sim[(p \wedge q) \rightarrow (\sim p \vee r)]$
 $\equiv \sim[\sim(p \wedge q) \vee (\sim p \vee r)]$
 $\equiv \sim[(\sim p \vee \sim q) \vee (\sim p \vee r)]$
 $\equiv \sim(\sim p \vee \sim q) \wedge \sim(\sim p \vee r)$
 $\equiv (p \wedge q) \wedge (p \wedge \sim r)$

52. Let p : 2 is prime, q : 3 is odd
 \therefore Symbolic form $p \rightarrow q$
 $\therefore \sim(p \rightarrow q) \equiv p \wedge \sim q$
 i.e., 2 is prime and 3 is not odd.

53. p : Hema gets admission in good college.
 q : Hema gets above 95% marks.
 \therefore Statement is $p \rightarrow q$
 $\sim(p \rightarrow q) \equiv p \wedge \sim q$

54. Given statement is
 $\exists x \in S$, such that $x > 0$
 $\therefore \sim(\exists x \in S$, such that $x > 0)$
 $\equiv \forall x \in S$, $x \leq 0$
 i.e., Every rational number $x \in S$ satisfies $x \leq 0$.

55. The current will flow through the circuit if p , q , r are closed or p , q' , r are closed.
 \therefore option (C) is the correct answer.



56. Let p : switch s_1 is closed.
 q : switch s_2 is closed.
 $\sim p$: switch s_1 is open
 $\sim q$: switch s_2 is open
 The current can flow in the circuit iff either s_1' and s_2 are closed or s_1 and s_2' are closed.
 It is represented by $(\sim p \wedge q) \vee (p \wedge \sim q)$.

58. The symbolic form of the given circuit is
 $(p \vee \sim p) \wedge q \equiv T \wedge q$
 $\equiv q$
59. Symbolic form of the circuit is
 $(p \wedge \sim q) \vee (\sim p \wedge q) \equiv (p \wedge \sim q) \vee (q \wedge \sim p)$
 $\equiv \sim (p \leftrightarrow q)$



Evaluation Test

1. $x + 3 = 10$ is an open sentence.
 \therefore It is not a statement.
 \therefore option (C) is correct.
2. Since $p \rightarrow q$ is false, when p is true and q is false.
 $p \rightarrow (q \vee r)$ is false,
 \therefore p is true and $q \vee r$ is false
 $\Rightarrow p$ is true and both q and r are false.
3. Since, contrapositive of $p \rightarrow q$ is $\sim q \rightarrow \sim p$.
 \therefore contrapositive of $(\sim p \wedge q) \rightarrow \sim r$ is
 $\sim(\sim r) \rightarrow \sim(\sim p \wedge q) \equiv r \rightarrow (p \vee \sim q)$
4. $\sim p$: Rohit is short.
 The given statement can be written symbolically as $p \vee (\sim p \wedge q)$.
5. Let p : \sqrt{x} is a complex number
 q : x is a negative number
 \therefore Logical statement is $p \rightarrow q$
 \therefore converse of $p \rightarrow q$ is $q \rightarrow p$
 \therefore option (B) is correct.
6. Consider option (C)

p	q	r	$\sim q$	$p \wedge \sim q$	$(p \wedge \sim q) \rightarrow r$
T	T	T	F	F	T
T	T	F	F	F	T
T	F	T	T	T	T
T	F	F	T	T	F
F	T	T	F	F	T
F	T	F	F	F	T
F	F	T	T	F	T
F	F	F	T	F	T

- $\therefore (p \wedge \sim q) \rightarrow r$ is a contingency
 \therefore option (C) is correct.
7. Consider option (A)

p	q	$p \wedge q$	$p \vee q$	$\sim(p \vee q)$	$(p \wedge q) \wedge (\sim(p \vee q))$
T	T	T	T	F	F
T	F	F	T	F	F
F	T	F	T	F	F
F	F	F	F	T	F

- $\therefore (p \wedge q) \wedge (\sim(p \vee q))$ is a contradiction.
 \therefore option (A) is correct.

8.

1	2	3	4	5	6
p	q	$\sim q$	$p \leftrightarrow q$	$p \leftrightarrow \sim q$	$\sim(p \leftrightarrow \sim q)$
T	T	F	T	F	T
T	F	T	F	T	F
F	T	F	F	T	F
F	F	T	T	F	T

- The entries in the columns 4 and 6 are identical.
 $\therefore \sim(p \leftrightarrow \sim q) \equiv p \leftrightarrow q$
 \therefore statement-1 is true.
 Also, all the entries in the last column of the above truth table are not T.
 $\therefore \sim(p \leftrightarrow \sim q)$ is not a tautology.
 \therefore statement-2 is false.
 \therefore option (B) is correct.
9. Consider option (C)
 $(p \vee q) \wedge (p \vee r) \equiv (T \vee T) \wedge (T \vee F)$
 $\equiv T \wedge T \equiv T$
 \therefore option (C) is correct.
11. The statement "Suman is brilliant and dishonest iff suman is rich" can be expressed as $Q \leftrightarrow (P \wedge \sim R)$
 The negation of this statement is
 $\sim(Q \leftrightarrow (P \wedge \sim R))$
12. $(\sim q) \rightarrow (\sim p)$ is contrapositive of $p \rightarrow q$.
 $\therefore p \rightarrow q \equiv (\sim q) \rightarrow (\sim p)$
 \therefore option (D) is true.
13. $(\sim p \wedge \sim q) \vee (p \wedge q) \vee (\sim p \wedge q)$
 $\equiv \sim p \wedge (\sim q \vee q) \vee (p \wedge q)$
 $\equiv (\sim p \wedge T) \vee (p \wedge q)$
 $\equiv \sim p \vee (p \wedge q)$
 $\equiv (\sim p \vee p) \wedge (\sim p \vee q)$
 $\equiv T \wedge (\sim p \vee q)$
 $\equiv \sim p \vee q$
 \therefore option (B) is correct.
14. Since, inverse of $p \rightarrow q$ is $\sim p \rightarrow \sim q$.
 \therefore inverse of $(p \wedge \sim q) \rightarrow r$ is $\sim(p \wedge \sim q) \rightarrow \sim r$
 i.e., $\sim p \vee q \rightarrow \sim r$

02 Matrices



Hints



Classical Thinking

$$1. \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} A$$

Applying $R_2 \rightarrow R_2 - 2R_1$,

$$\begin{bmatrix} 1 & -1 \\ 2-2(1) & 3-2(-1) \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0-2(1) & 1-2(0) \end{bmatrix} A$$

$$\begin{bmatrix} 1 & -1 \\ 0 & 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ -2 & 1 \end{bmatrix} A$$

$$2. A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & -2 & 5 \end{bmatrix}$$

Applying $R_1 \leftrightarrow R_2$,

$$A \sim \begin{bmatrix} 3 & -2 & 5 \\ 1 & 2 & -1 \end{bmatrix}$$

Applying $C_1 \rightarrow C_1 + 2C_3$,

$$A \sim \begin{bmatrix} 13 & -2 & 5 \\ -1 & 2 & -1 \end{bmatrix}$$

$$3. \text{ Let } A = \begin{bmatrix} 1 & -1 & 2 \\ 2 & 1 & 3 \\ 3 & 2 & 4 \end{bmatrix}$$

Applying $R_2 \rightarrow R_2 - 2R_1$ and $R_3 \rightarrow R_3 - 3R_1$,

$$A \sim \begin{bmatrix} 1 & -1 & 2 \\ 0 & 3 & -1 \\ 0 & 5 & -2 \end{bmatrix}$$

Applying $R_3 \rightarrow R_3 - \left(\frac{5}{3}\right)R_2$,

$$A \sim \begin{bmatrix} 1 & -1 & 2 \\ 0 & 3 & -1 \\ 0 & 0 & -\frac{1}{3} \end{bmatrix}$$

which is an upper triangular matrix.

4. If $|A| \neq 0$, then A^{-1} exists

$\therefore |A|$ is non zero

5. $M_{11} = \text{minor of } a_{11} = |a_{22}| = a_{22}$
....[By leaving first row and first column]

6. The minor of element $a_{21} = M_{21} = -1$
....[By leaving R_2 and C_1]

$$7. M_{31} = \begin{vmatrix} 2 & 3 \\ 4 & 2 \end{vmatrix} \quad \dots[\text{By leaving } R_3 \text{ and } C_1]$$

$$= -8$$

$$8. M_{23} = \begin{vmatrix} 1 & 1 \\ -1 & 2 \end{vmatrix} = 3$$

$$9. A_{12} = (-1)^{1+2} M_{12} = (-1)^3 (-3) = 3$$

$$10. A_{21} = (-1)^3 M_{21} = -(3) = -3$$

$$11. A_{32} = (-1)^{3+2} \cdot M_{32} = (-1)^5 \begin{vmatrix} 2 & 3 \\ 4 & 5 \end{vmatrix} = 2$$

$$12. A_{31} = (-1)^{3+1} \begin{vmatrix} 1 & 1 \\ 1 & -3 \end{vmatrix} = -3 - 1 = -4$$

$$A_{32} = -(-3 - 2) = -(-5) = 5$$

$$A_{33} = 1 - 2 = -1$$

\therefore Co-factors are $-4, 5, -1$

13. Matrix of co-factors

$$= [A_{ij}]_{2 \times 2} = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} = \begin{bmatrix} 2 & -(-3) \\ -5 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 3 \\ -5 & -1 \end{bmatrix}$$

$$\therefore \text{adj } A = [A_{ij}]_{2 \times 2}^T = \begin{bmatrix} 2 & 3 \\ -5 & -1 \end{bmatrix}^T = \begin{bmatrix} 2 & -5 \\ 3 & -1 \end{bmatrix}$$

$$14. \text{ Let } A = \begin{bmatrix} 1 & -4 \\ 3 & 2 \end{bmatrix}$$

Matrix of co-factors is:

$$[A_{ij}]_{2 \times 2} = \begin{bmatrix} 2 & -3 \\ 4 & 1 \end{bmatrix}$$

$$\therefore \text{adj } A = [A_{ij}]_{2 \times 2}^T = \begin{bmatrix} 2 & -3 \\ 4 & 1 \end{bmatrix}^T = \begin{bmatrix} 2 & 4 \\ -3 & 1 \end{bmatrix}$$



15. Matrix of co-factors is :

$$[A_{ij}]_{3 \times 3} = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{bmatrix} = \begin{bmatrix} -3 & 3 & -9 \\ 0 & -1 & -2 \\ 0 & 0 & 3 \end{bmatrix}$$

$$\therefore \text{adj } A = [A_{ij}]_{3 \times 3}^T = \begin{bmatrix} -3 & 3 & -9 \\ 0 & -1 & -2 \\ 0 & 0 & 3 \end{bmatrix}^T$$

$$\therefore \text{adj } A = \begin{bmatrix} -3 & 0 & 0 \\ 3 & -1 & 0 \\ -9 & -2 & 3 \end{bmatrix}$$

16. Matrix of co-factors is :

$$[A_{ij}]_{3 \times 3} = \begin{bmatrix} 3 & -9 & -5 \\ -4 & 1 & 3 \\ -5 & 4 & 1 \end{bmatrix}$$

$$\text{adj } A = [A_{ij}]_{3 \times 3}^T = \begin{bmatrix} 3 & -4 & -5 \\ -9 & 1 & 4 \\ -5 & 3 & 1 \end{bmatrix}$$

17. $|\text{adj}(\text{adj } A)| = |A| = 12 - 10 = 2$

18. $|A| = a^3$

$$|A| |\text{adj } A| = |A (\text{adj } A)| = |A| I$$

$$= \begin{vmatrix} |A| & 0 & 0 \\ 0 & |A| & 0 \\ 0 & 0 & |A| \end{vmatrix} = |A|^3 = (a^3)^3 = a^9$$

19. $|A| = \begin{vmatrix} 2 & 3 \\ 4 & 6 \end{vmatrix} = 12 - 12 = 0$

$\therefore A^{-1}$ does not exist.

20. Let $A = \begin{bmatrix} 1 & 3 \\ 3 & 10 \end{bmatrix}$

$$\therefore |A| = \begin{vmatrix} 1 & 3 \\ 3 & 10 \end{vmatrix} = 1$$

$$\text{adj } A = \begin{bmatrix} 10 & -3 \\ -3 & 1 \end{bmatrix}$$

$$A^{-1} = \frac{1}{|A|} \text{adj } A = \begin{bmatrix} 10 & -3 \\ -3 & 1 \end{bmatrix}$$

21. The multiplicative inverse of $A = A^{-1}$

$$|A| = \begin{vmatrix} 2 & 1 \\ 7 & 4 \end{vmatrix} = 1 \neq 0$$

$$\text{adj } A = \begin{bmatrix} 4 & -1 \\ -7 & 2 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \begin{bmatrix} 4 & -1 \\ -7 & 2 \end{bmatrix}$$

22. Let $A = \begin{bmatrix} 3 & -2 \\ 1 & 4 \end{bmatrix} \Rightarrow |A| = 14 \neq 0$

$$\text{adj } A = \begin{bmatrix} 4 & 2 \\ -1 & 3 \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} \frac{4}{14} & \frac{2}{14} \\ \frac{-1}{14} & \frac{3}{14} \end{bmatrix}$$

23. The inverse of the given diagonal matrix is

$$A^{-1} = \begin{bmatrix} \frac{1}{a} & 0 \\ 0 & b \end{bmatrix}$$

24. $|A| = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 0 & 2 \\ 3 & 1 & 1 \end{vmatrix} = 4 \neq 0$

$$\text{adj } A = \begin{bmatrix} -2 & 5 & 1 \\ 0 & -2 & 2 \\ 2 & -1 & -1 \end{bmatrix}^T = \begin{bmatrix} -2 & 0 & 2 \\ 5 & -2 & -1 \\ 1 & 2 & -1 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{4} \begin{bmatrix} -2 & 0 & 2 \\ 5 & -2 & -1 \\ 1 & 2 & -1 \end{bmatrix}$$

25. Let $A = \begin{bmatrix} 3 & -2 & -1 \\ -4 & 1 & -1 \\ 2 & 0 & 1 \end{bmatrix} \Rightarrow |A| = 1 \neq 0$

$$\text{adj } A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 7 \\ -2 & -4 & -5 \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 7 \\ -2 & -4 & -5 \end{bmatrix}$$

26. The inverse of the given diagonal matrix is,

$$A^{-1} = \begin{bmatrix} \frac{1}{a} & 0 & 0 \\ 0 & \frac{1}{b} & 0 \\ 0 & 0 & \frac{1}{c} \end{bmatrix}$$



$$27. |A| = \begin{vmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{vmatrix} = -1$$

$$\text{adj } A = \begin{bmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{bmatrix}^T = \begin{bmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

$$28. \text{ Given, } A^{-1} = \frac{1}{k} \text{adj } A$$

$$\therefore k = |A|$$

$$\therefore |A| = \begin{vmatrix} 3 & 2 & 4 \\ 1 & 2 & -1 \\ 0 & 1 & 1 \end{vmatrix}$$

$$= 3(2+1) - 2(1-0) + 4(1-0)$$

$$= 9 - 2 + 4 = 11$$

$$\Rightarrow k = 11$$

$$29. AB = AC$$

$$\Rightarrow A^{-1}AB = A^{-1}AC$$

$$\Rightarrow IB = IC$$

$$\Rightarrow B = C$$

$$\therefore \text{ For } B = C, A^{-1} \text{ must exist}$$

$$\Rightarrow A \text{ is non-singular}$$

$$30. \text{ Consider option (B),}$$

$$A^{-1} \text{ is a matrix and } |A|^{-1} \text{ is a number.}$$

$$\therefore \text{ option (B) is not true.}$$

$$32. (A^2 - 4A)A^{-1} = A \cdot A \cdot A^{-1} - 4A \cdot A^{-1}$$

$$= A - 4I$$

$$= \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} - \begin{bmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix}$$

$$= \begin{bmatrix} -3 & 2 & 2 \\ 2 & -3 & 2 \\ 2 & 2 & -3 \end{bmatrix}$$

33. The given system of equations can be written in matrix form $AX = B$, where

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix} \text{ and } B = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$$

$$\therefore \text{ Now, } \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$$

Applying $R_2 \rightarrow R_2 - 2R_1$,

$$\begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 \\ -2 \end{bmatrix}$$

$$\Rightarrow x + 2y = 3, \text{ and } \dots(i)$$

$$-y = -2 \dots(ii)$$

$$\Rightarrow y = 2$$

putting $y = 2$ in (i), we get

$$x + 2(2) = 3$$

$$\Rightarrow x = -1$$

Alternate method:

$$AX = B \Rightarrow X = A^{-1}B$$

$$|A| = -1 \neq 0$$

$$A^{-1} = \frac{1}{-1} \begin{bmatrix} 3 & -2 \\ -2 & 1 \end{bmatrix} = \begin{bmatrix} -3 & 2 \\ 2 & -1 \end{bmatrix}$$

$$X = \begin{bmatrix} -3 & 2 \\ 2 & -1 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

$$\therefore x = -1, y = 2$$

$$34. AX = B$$

$$\therefore \begin{bmatrix} 3 & -4 & 2 \\ 2 & 3 & 5 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -1 \\ 7 \\ 2 \end{bmatrix}$$

$$R_2 - 5R_3 \Rightarrow \begin{bmatrix} 3 & -4 & 2 \\ -3 & 3 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix}$$

$$R_1 - 2R_3 \Rightarrow \begin{bmatrix} 1 & -4 & 0 \\ -3 & 3 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -5 \\ -3 \\ 2 \end{bmatrix}$$

$$\Rightarrow x - 4y = -5, \text{ and } \dots(i)$$

$$-3x + 3y = -3 \dots(ii)$$

Solving (i) and (ii), we get $x = 3$

$$35. \begin{bmatrix} a & 1 & 1 \\ 1 & a & 1 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Applying $R_1 \rightarrow R_1 - R_3$,

$$\begin{bmatrix} a-1 & 0 & 0 \\ 1 & a & 1 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\therefore (a-1)x + 0 + 0 = 0$$

$$\Rightarrow a - 1 = 0$$

$$\Rightarrow a = 1$$



$$36. |A| = -\frac{1}{2} \neq 0$$

$$\text{adj } A = \begin{bmatrix} i & 0 \\ 2 & 0 \\ 0 & i \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{-\frac{1}{2}} \begin{bmatrix} i & 0 \\ 2 & 0 \\ 0 & i \end{bmatrix} = \begin{bmatrix} -i & 0 \\ 0 & -2i \end{bmatrix}$$

$$37. \therefore \text{adj } (AB) = \text{adj } (B) \text{adj } (A)$$



Critical Thinking

$$1. \begin{bmatrix} 1 & 3 & -2 \\ -3 & 0 & -5 \\ 2 & 5 & 0 \end{bmatrix} = A \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Applying $C_2 \rightarrow C_2 - 3C_1$ and $C_3 \rightarrow C_3 + 2C_1$,

$$\begin{bmatrix} 1 & 3-3 & -2+2 \\ -3 & 0+9 & -5-6 \\ 2 & 5-6 & 0+4 \end{bmatrix} = A \begin{bmatrix} 1 & 0-3 & 0+2 \\ 0 & 1-0 & 0+0 \\ 0 & 0-0 & 1+0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 & 0 & 0 \\ -3 & 9 & -11 \\ 2 & -1 & 4 \end{bmatrix} = A \begin{bmatrix} 1 & -3 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$2. A = \begin{bmatrix} 2 & -3 & 3 \\ 2 & 2 & 3 \\ 3 & -2 & 2 \end{bmatrix}$$

Applying $C_2 \rightarrow C_2 + 2C_1$,

$$\therefore A \sim \begin{bmatrix} 2 & 1 & 3 \\ 2 & 6 & 3 \\ 3 & 4 & 2 \end{bmatrix}$$

Applying $R_1 \rightarrow R_1 + R_3$,

$$A \sim \begin{bmatrix} 5 & 5 & 5 \\ 2 & 6 & 3 \\ 3 & 4 & 2 \end{bmatrix}$$

$$3. a_{11}A_{11} + a_{21}A_{21} + a_{31}A_{31} \\ = 1(4-3) + 3[-(4-1)] + 2(6-2) = 0 \\ \text{and } |A| = 1(4-3) - 2(6-6) + 1(3-4) = 0 \\ \therefore a_{11}A_{11} + a_{21}A_{21} + a_{31}A_{31} = |A|$$

4. Matrix of co-factors,

$$[A_{ij}]_{3 \times 3} = \begin{bmatrix} -4 & 1 & 4 \\ -3 & 0 & 4 \\ -3 & 1 & 3 \end{bmatrix}$$

$$\therefore \text{adj } N = [A_{ij}]_{3 \times 3}^T = \begin{bmatrix} -4 & -3 & -3 \\ 1 & 0 & 1 \\ 4 & 4 & 3 \end{bmatrix} = N$$

$$5. AB = \begin{bmatrix} 34 & 39 \\ 82 & 94 \end{bmatrix} \Rightarrow \text{adj } (AB) = \begin{bmatrix} 94 & -39 \\ -82 & 34 \end{bmatrix}$$

6. A is a 2×2 matrix

$$\therefore |\text{adj } A| = |A| = 10$$

7. A is a 3×3 Matrix

$$\therefore |\text{adj } A| = |A|^2 = (12)^2 = 144$$

8. $A(\text{Adj } A) = |A| \cdot (I_n)$

$$\therefore \begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix} = |A| \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix} = \begin{bmatrix} |A| & 0 \\ 0 & |A| \end{bmatrix}$$

$$\Rightarrow |A| = 10$$

9. $A(\text{adj } A) = |A| I$

$$\therefore |A(\text{adj } A)| = |A|^n \quad (\text{If } A \text{ is of order } n \times n)$$

$$\Rightarrow |A| |\text{adj } A| = |A|^n$$

$$\Rightarrow |\text{Adj } A| = |A|^{n-1}$$

Since, A is singular

$$\therefore |A| = 0$$

$$\Rightarrow |\text{Adj } A| = 0$$

Hence, adj A is a singular matrix.

10. A is a Singular matrix.

$$\therefore |A| = 0 \text{ and } A(\text{adj } A) = |A| \cdot I = 0 \cdot I = 0$$

$$\Rightarrow A(\text{adj } A) \text{ is a zero matrix.}$$

$$11. \text{adj } A = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

$$\therefore \text{adj } (\text{adj } A) = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = A$$

$$12. |A| = \begin{vmatrix} 2 & 0 & -1 \\ 5 & 1 & 0 \\ 0 & 1 & 3 \end{vmatrix} = 1 \neq 0$$

$$\text{adj } A = \begin{bmatrix} 3 & -1 & 1 \\ -15 & 6 & -5 \\ 5 & -2 & 2 \end{bmatrix}$$



$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A) = \begin{bmatrix} 3 & -1 & 1 \\ -15 & 6 & -5 \\ 5 & -2 & 2 \end{bmatrix}$$

$$13. |A| = \begin{vmatrix} 1 & 1 & 1 \\ 6 & 7 & 8 \\ 6 & 7 & -8 \end{vmatrix} = -16 \neq 0$$

$$\text{adj } A = \begin{bmatrix} -112 & 96 & 0 \\ 15 & -14 & -1 \\ 1 & -2 & 1 \end{bmatrix}^T$$

$$= \begin{bmatrix} -112 & 15 & 1 \\ 96 & -14 & -2 \\ 0 & -1 & 1 \end{bmatrix}$$

$$\therefore A^{-1} = -\frac{1}{16} \begin{bmatrix} -112 & 15 & 1 \\ 96 & -14 & -2 \\ 0 & -1 & 1 \end{bmatrix}$$

$$14. D = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{bmatrix}$$

The inverse of the given diagonal matrix is

$$D^{-1} = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{3} & 0 \\ 0 & 0 & \frac{1}{4} \end{bmatrix}$$

$$\therefore D^{-1} = \text{diag} \left[\frac{1}{2}, \frac{1}{3}, \frac{1}{4} \right]$$

$$15. \text{ If } AC = B, \text{ then } A = BC^{-1}$$

$$\begin{aligned} \therefore A &= \begin{bmatrix} 3 & -1 \\ 6 & 0 \end{bmatrix} \begin{bmatrix} 1 & 5 \\ 0 & 1 \end{bmatrix}^{-1} \\ &= \begin{bmatrix} 3 & -1 \\ 6 & 0 \end{bmatrix} \begin{bmatrix} 1 & -5 \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 3 & -16 \\ 6 & -30 \end{bmatrix} \end{aligned}$$

$$16. \text{ If } AB = C, \text{ then } B^{-1} A^{-1} = C^{-1}$$

$$\therefore A^{-1} = BC^{-1}$$

$$\text{Here, } A \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 1 & 1 \end{bmatrix}$$

$$\begin{aligned} \therefore A^{-1} &= \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 3 & 2 \\ 1 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 1 \\ 2 & -4 \end{bmatrix} \end{aligned}$$

$$17. \text{ If } AB = C, \text{ then } B^{-1} A^{-1} = C^{-1}$$

$$\therefore A^{-1} = BC^{-1}$$

$$\therefore A \begin{bmatrix} 0 & 1 \\ 2 & -4 \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 1 & 1 \end{bmatrix}$$

$$\begin{aligned} \Rightarrow A^{-1} &= \begin{bmatrix} 0 & 1 \\ 2 & -4 \end{bmatrix} \begin{bmatrix} 3 & 2 \\ 1 & 1 \end{bmatrix}^{-1} \\ &= \begin{bmatrix} 0 & 1 \\ 2 & -4 \end{bmatrix} \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix} \\ &= \begin{bmatrix} -1 & 3 \\ 6 & -16 \end{bmatrix} \end{aligned}$$

$$18. \text{ If } XAY = I, \text{ then } A = X^{-1} Y^{-1} = (YX)^{-1}$$

$$\text{Here, } YX = \begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 7 & 4 \end{bmatrix} = \begin{bmatrix} 8 & 5 \\ -11 & -7 \end{bmatrix}$$

$$\begin{aligned} \therefore A &= \begin{bmatrix} 8 & 5 \\ -11 & -7 \end{bmatrix}^{-1} \\ &= \begin{bmatrix} 7 & 5 \\ -11 & -8 \end{bmatrix} \end{aligned}$$

$$19. (BA)^{-1} = C$$

$$\Rightarrow A^{-1} B^{-1} = C \Rightarrow A^{-1} = CB$$

$$\begin{aligned} \therefore A^{-1} &= \begin{bmatrix} -1 & 0 & 1 \\ 1 & 1 & 3 \\ 2 & 0 & 2 \end{bmatrix} \begin{bmatrix} 2 & 6 & 4 \\ 1 & 0 & 1 \\ -1 & 1 & -1 \end{bmatrix} \\ &= \begin{bmatrix} -3 & -5 & -5 \\ 0 & 9 & 2 \\ 2 & 14 & 6 \end{bmatrix} \end{aligned}$$

$$20. \text{ Since, } PQ = -5I_3$$

$$\therefore (PQ)^{-1} = -\frac{1}{5} I_3$$

$$21. |A| = \begin{vmatrix} 2 & 3 \\ 1 & -2 \end{vmatrix} = -7 \neq 0$$

$$\text{adj } A = \begin{bmatrix} -2 & -3 \\ -1 & 2 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{-7} \begin{bmatrix} -2 & -3 \\ -1 & 2 \end{bmatrix} = \frac{1}{7} A$$

$$\therefore \alpha = \frac{1}{7}$$



$$22. |A| = \begin{vmatrix} 1 & 2 & -1 \\ 3 & 4 & 5 \\ 2 & 6 & 7 \end{vmatrix} = -34$$

Co-factor of element a_{23} of $A = A_{23}$

$$\therefore A_{23} = (-1)^{2+3} \begin{vmatrix} 1 & 2 \\ 2 & 6 \end{vmatrix} = 2$$

$$\therefore \text{Element } a_{32} \text{ of } A^{-1} = \frac{A_{23}}{|A|} = \frac{2}{-34} = \frac{-1}{17}$$

$$23. A^2 - 3A - 7I = 0$$

$$\Rightarrow A - 3I - 7A^{-1} = 0 \Rightarrow A^{-1} = \frac{1}{7}(A - 3I)$$

$$\therefore A^{-1} = \frac{1}{7} \left\{ \begin{bmatrix} 5 & 3 \\ -1 & -2 \end{bmatrix} - \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} \right\}$$

$$= \begin{bmatrix} \frac{2}{7} & \frac{3}{7} \\ -\frac{1}{7} & -\frac{5}{7} \end{bmatrix}$$

$$24. A^2 = \begin{bmatrix} x & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} x^2 + 1 & x \\ x & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow x = 0$$

$$\therefore A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$A^2 = I$$

$$\therefore A^{-1} A \cdot A = A^{-1} I$$

$$\Rightarrow I \cdot A = A^{-1} I$$

$$\Rightarrow A^{-1} = A$$

$$25. AB = 3I$$

$$\Rightarrow A^{-1} AB = 3 A^{-1} I$$

$$\Rightarrow B = 3A^{-1}$$

$$\therefore A^{-1} = \frac{1}{3} B$$

$$26. A^2 - A + 2I = 0$$

$$\Rightarrow A \cdot A - A + 2I = 0$$

$$\Rightarrow A^{-1} \cdot A \cdot A - A^{-1} \cdot A + 2 A^{-1} \cdot I = 0$$

$$\Rightarrow A - I + 2 A^{-1} = 0$$

$$\Rightarrow 2 A^{-1} = I - A$$

$$\Rightarrow A^{-1} = \frac{1}{2} (I - A)$$

$$27. A^2 + mA + nI = 0$$

$$\Rightarrow A \cdot A + mA + nI = 0$$

$$\Rightarrow A^{-1} \cdot A \cdot A + mA^{-1} \cdot A + nA^{-1} \cdot I = 0$$

$$\Rightarrow A + mA^{-1} + nA^{-1} = 0$$

$$\Rightarrow nA^{-1} = -A - mA^{-1}$$

$$\Rightarrow A^{-1} = \frac{-1}{n} (A + mA^{-1})$$

$$28. 4A^3 + 2A^2 + 7A + I = 0$$

$$\Rightarrow 4A^{-1}A^3 + 2A^{-1}A^2 + 7A^{-1}A + A^{-1}I = 0$$

$$\Rightarrow 4A^2 + 2A + 7I + A^{-1} = 0$$

$$\Rightarrow A^{-1} = -(4A^2 + 2A + 7I)$$

29. The given system of equations can be written in the matrix form as $AX = B$, where

$$A = \begin{bmatrix} 5 & -7 \\ 7 & -5 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix} \text{ and } B = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 5 & -7 \\ 7 & -5 \end{vmatrix} = 24 \neq 0$$

$$AX = B$$

$$\therefore \begin{bmatrix} 5 & -7 \\ 7 & -5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

$$R_1 \rightarrow R_1 - \frac{5}{7}R_2$$

$$\Rightarrow \begin{bmatrix} 0 & \frac{-24}{7} \\ 7 & -5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{-1}{7} \\ 3 \end{bmatrix}$$

$$\therefore \frac{-24}{7}y = \frac{-1}{7} \Rightarrow y = \frac{1}{24}$$

$$7x - 5y = 3 \Rightarrow x = \frac{11}{24}$$

Alternate method:

the given system of equations has a unique solution which is given by $X = A^{-1} B$.

$$\text{adj } A = \begin{bmatrix} -5 & 7 \\ -7 & 5 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{24} \begin{bmatrix} -5 & 7 \\ -7 & 5 \end{bmatrix}$$

$$\therefore X = A^{-1} B = \frac{1}{24} \begin{bmatrix} -5 & 7 \\ -7 & 5 \end{bmatrix} \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \frac{1}{24} \begin{bmatrix} 11 \\ 1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{11}{24} \\ \frac{1}{24} \end{bmatrix}$$

$$\Rightarrow x = \frac{11}{24}, y = \frac{1}{24}$$

$$30. AX = B$$

$$\begin{bmatrix} 1 & -1 & 2 \\ 2 & 0 & 1 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 4 \end{bmatrix}$$



$$R_1 \rightarrow 2R_1 + R_3$$

$$\begin{bmatrix} 5 & 0 & 5 \\ 2 & 0 & 1 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 10 \\ 1 \\ 4 \end{bmatrix}$$

$$R_1 \rightarrow R_1 - 5R_2$$

$$\begin{bmatrix} -5 & 0 & 0 \\ 2 & 0 & 1 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 5 \\ 1 \\ 4 \end{bmatrix}$$

$$\therefore -5x_1 = 5 \Rightarrow x_1 = -1$$

$$2x_1 + x_3 = 1 \Rightarrow x_3 = 3$$

$$3x_1 + 2x_2 + x_3 = 4 \Rightarrow x_2 = 2$$

$$\therefore X = \begin{bmatrix} -1 \\ 2 \\ 3 \end{bmatrix}$$

$$31. X = A^{-1} D$$

$$\Rightarrow AX = D$$

$$\begin{bmatrix} 1 & -1 & -2 \\ 2 & 1 & 1 \\ 4 & -1 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 5 \\ 11 \end{bmatrix}$$

$$R_1 \rightarrow R_1 + R_2, R_3 \rightarrow R_3 + R_2$$

$$\begin{bmatrix} 3 & 0 & -1 \\ 2 & 1 & 1 \\ 6 & 0 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 8 \\ 5 \\ 16 \end{bmatrix}$$

$$R_3 \rightarrow R_3 - R_1$$

$$\begin{bmatrix} 3 & 0 & -1 \\ 2 & 1 & 1 \\ 3 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 8 \\ 5 \\ 8 \end{bmatrix}$$

$$\therefore 3x = 8 \Rightarrow x = \frac{8}{3}$$

$$3x - z = 8 \Rightarrow z = 0$$

$$2x + y + z = 5 \Rightarrow y = \frac{-1}{3}$$

$$\therefore X = \begin{bmatrix} \frac{8}{3} \\ \frac{-1}{3} \\ 0 \end{bmatrix}$$

$$32. \Delta = \begin{vmatrix} 1 & \omega^n & \omega^{2n} \\ \omega^{2n} & 1 & \omega^n \\ \omega^n & \omega^{2n} & 1 \end{vmatrix}$$

Applying $C_1 \rightarrow C_1 + C_2 + C_3$, we get

$$\Delta = \begin{vmatrix} 1 + \omega^n + \omega^{2n} & \omega^n & \omega^{2n} \\ 1 + \omega^n + \omega^{2n} & 1 & \omega^n \\ 1 + \omega^n + \omega^{2n} & \omega^{2n} & 1 \end{vmatrix} = \begin{vmatrix} 0 & \omega^n & \omega^{2n} \\ 0 & 1 & \omega^n \\ 0 & \omega^{2n} & 1 \end{vmatrix}$$

....[: $1 + \omega^n + \omega^{2n} = 0$, if n is not multiple of 3]

$$\therefore \Delta = 0$$

$$33. |A| = \begin{vmatrix} 1 & \sin \alpha \\ -\sin \alpha & -1 \end{vmatrix} = -1 + \sin^2 \alpha \neq 0$$

$$\text{adj } A = \begin{bmatrix} -1 & -\sin \alpha \\ \sin \alpha & 1 \end{bmatrix}$$

$$\begin{aligned} \therefore A^{-1} &= \frac{1}{|A|} (\text{adj } A) = \frac{1}{-1 + \sin^2 \alpha} \begin{bmatrix} -1 & -\sin \alpha \\ \sin \alpha & 1 \end{bmatrix} \\ &= \frac{1}{\cos^2 \alpha} \begin{bmatrix} 1 & \sin \alpha \\ -\sin \alpha & -1 \end{bmatrix} = \sec^2 \alpha \begin{bmatrix} 1 & \sin \alpha \\ -\sin \alpha & -1 \end{bmatrix} \end{aligned}$$



Competitive Thinking

$$1. |A| = \begin{vmatrix} 2 & -2 \\ -2 & 2 \end{vmatrix} = 4 - 4 = 0$$

$$|B| = \begin{vmatrix} 1 & 1 \\ 1 & 1 \end{vmatrix} = 1 - 1 = 0$$

$\therefore A^{-1}$ and B^{-1} does not exist

$$2. \text{ The matrix is not invertible if } \begin{vmatrix} 1 & a & 2 \\ 1 & 2 & 5 \\ 2 & 1 & 1 \end{vmatrix} = 0$$

$$\Rightarrow 1(2 - 5) - a(1 - 10) + 2(1 - 4) = 0$$

$$\Rightarrow -3 + 9a - 6 = 0$$

$$\Rightarrow a = 1$$

3. $|A| = k^2 + 1$, which can be never zero. Hence matrix A is invertible for all real k .

4. The given matrix will be invertible, if

$$\begin{vmatrix} \lambda & -1 & 4 \\ -3 & 0 & 1 \\ -1 & 1 & 2 \end{vmatrix} \neq 0$$

$$\Rightarrow \lambda(0 - 1) + 1(-6 + 1) + 4(-3) \neq 0$$

$$\Rightarrow -\lambda - 5 - 12 \neq 0$$

$$\Rightarrow \lambda \neq -17$$



$$5. \text{ Let } A = \begin{bmatrix} \alpha & 14 & -1 \\ 2 & 3 & 1 \\ 6 & 2 & 3 \end{bmatrix} = 0$$

Since, A^{-1} does not exist

$$\therefore |A| = 0$$

$$\Rightarrow \begin{vmatrix} \alpha & 14 & -1 \\ 2 & 3 & 1 \\ 6 & 2 & 3 \end{vmatrix}$$

$$\Rightarrow \alpha(9-2) - 14(6-6) - 1(4-18) = 0$$

$$\Rightarrow 7\alpha = -14$$

$$\Rightarrow \alpha = -2$$

$$6. \quad a_{11} = 1, a_{12} = 1, a_{13} = 0$$

$$A_{21} = (-1)^{2+1} \begin{vmatrix} 1 & 0 \\ 2 & 1 \end{vmatrix} = -1$$

$$A_{22} = (-1)^{2+2} \begin{vmatrix} 1 & 0 \\ 1 & 1 \end{vmatrix} = 1$$

$$A_{23} = (-1)^{2+3} \begin{vmatrix} 1 & 1 \\ 1 & 2 \end{vmatrix} = -1$$

$$\therefore a_{11} \cdot A_{21} + a_{12} \cdot A_{22} + a_{13} \cdot A_{23} = 1 \times -1 + 1 \times 1 + 0 \times -1 = 0$$

$$7. \quad A = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 1 & 5 \\ 2 & 4 & 7 \end{bmatrix}$$

$$\begin{aligned} a_{31} A_{31} + a_{32} A_{32} + a_{33} A_{33} \\ = 2(10-3) + 4[-(5-3)] + 7(1-2) \\ = 14 - 8 - 7 = -1 \end{aligned}$$

$$8. \quad \text{Co-factor matrix of } X = \begin{bmatrix} t & -z \\ y & -x \end{bmatrix}$$

$$\therefore \text{Transpose of adj } X = \text{co-factor matrix of } X \\ = \begin{bmatrix} t & -z \\ y & -x \end{bmatrix}$$

9. Matrix of co-factors is

$$[A_{ij}]_{3 \times 3} = \begin{bmatrix} 2 & -5 & 32 \\ 0 & 1 & -6 \\ 0 & 0 & 2 \end{bmatrix}$$

$$\Rightarrow \text{adj } A = [A_{ij}]_{3 \times 3}^T = \begin{bmatrix} 2 & 0 & 0 \\ -5 & 1 & 0 \\ 32 & -6 & 2 \end{bmatrix}$$

$$10. \quad A = \begin{bmatrix} 2 & -3 \\ -4 & 1 \end{bmatrix}$$

$$\begin{aligned} 3A^2 &= 3 \begin{bmatrix} 2 & -3 \\ -4 & 1 \end{bmatrix} \begin{bmatrix} 2 & -3 \\ -4 & 1 \end{bmatrix} \\ &= 3 \begin{bmatrix} 16 & -9 \\ -12 & 13 \end{bmatrix} = \begin{bmatrix} 48 & -27 \\ -36 & 39 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} 3A^2 + 12A &= \begin{bmatrix} 48 & -27 \\ -36 & 39 \end{bmatrix} + \begin{bmatrix} 24 & -36 \\ -48 & 12 \end{bmatrix} \\ &= \begin{bmatrix} 72 & -63 \\ -84 & 51 \end{bmatrix} \end{aligned}$$

$$\therefore \text{adj}(3A^2 + 12A) = \begin{bmatrix} 51 & 63 \\ 84 & 72 \end{bmatrix}$$

$$11. \quad A = \begin{bmatrix} 1 & -1 & 1 \\ 0 & 2 & -3 \\ 2 & 1 & 0 \end{bmatrix}$$

$$B = \text{adj } A = \begin{bmatrix} 3 & 1 & 1 \\ -6 & -2 & 3 \\ -4 & -3 & 2 \end{bmatrix}$$

$$\therefore \text{adj } B = \begin{bmatrix} 5 & -5 & 5 \\ 0 & 10 & -15 \\ 10 & 5 & 0 \end{bmatrix} = 5A$$

$$\therefore \text{adj } B = C \quad \dots [\because C = 5A(\text{given})]$$

$$\Rightarrow |\text{adj } B| = |C|$$

$$\Rightarrow \frac{|\text{adj } B|}{|C|} = 1$$

$$12. \quad A(\text{adj } A) = |A| \cdot I_n$$

Where, n = order of the matrix

$$\therefore A(\text{adj } A) = \begin{bmatrix} 3 & 2 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix}$$

$$13. \quad |A| = \begin{vmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{vmatrix} = \cos^2 x + \sin^2 x = 1$$

$$\text{Since, } A(\text{adj } A) = |A| \cdot I$$

$$\therefore A(\text{adj } A) = 1 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$14. \quad \text{Since, } A(\text{adj } A) = |A| \cdot I$$

$$\Rightarrow \begin{bmatrix} k & 0 \\ 0 & k \end{bmatrix} = (\cos^2 \alpha + \sin^2 \alpha) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow k = 1$$



$$15. \text{ Let } I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \text{ then } kI = \begin{bmatrix} k & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & k \end{bmatrix}$$

$$\Rightarrow \text{adj}(kI) = \begin{bmatrix} k^2 & 0 & 0 \\ 0 & k^2 & 0 \\ 0 & 0 & k^2 \end{bmatrix} = k^2 I$$

$$16. \text{adj}(\lambda X) = \lambda^{3-1} (\text{adj } X) \\ \dots [\because \text{adj}(kA) = k^{n-1} (\text{adj } A)] \\ = \lambda^2 \text{adj } X$$

17. Given, A is a singular matrix.

$$\therefore |A| = 0$$

$$\text{Since, } |\text{adj } A| = |A|^{n-1}$$

$$\Rightarrow |\text{adj } A| = 0$$

\Rightarrow adj A is also singular.

$$18. |\text{Adj } A| = |A|^{n-1} = d^{n-1}$$

$$19. |A| = \begin{vmatrix} 4 & 2 \\ 3 & 4 \end{vmatrix} = 16 - 6 = 10$$

$$\therefore |\text{adj } A| = |A|^{n-1}$$

where n \Rightarrow order of matrix.

$$\therefore |\text{adj } A| = |A| = 10$$

$$20. A (\text{adj } A) = |A| I_n$$

$$\Rightarrow \begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix} = |A| I_n$$

$$\Rightarrow 10 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = |A| I_n$$

$$\Rightarrow 10 I_n = |A| I_n$$

$$\Rightarrow |A| = 10$$

21. Since, $A(\text{Adj } A) = |A| I$

$$\therefore |A| = 10$$

$$|\text{Adj } A| = |A|^{n-1}$$

$$\therefore |\text{Adj } A| = |A|^{3-1} = |A|^2 = 10^2 = 100$$

$$22. \text{adj } P = \begin{bmatrix} 1 & 4 & 4 \\ 2 & 1 & 7 \\ 1 & 1 & 3 \end{bmatrix}$$

$$|\text{adj } P| = |P|^2 \quad \dots [\because |\text{adj } A| = |A|^{n-1}]$$

$$\Rightarrow \begin{vmatrix} 1 & 4 & 4 \\ 2 & 1 & 7 \\ 1 & 1 & 3 \end{vmatrix} = |P|^2$$

$$\Rightarrow |P|^2 = 1(-4) - 4(-1) + 4(1)$$

$$\Rightarrow |P|^2 = 4 \Rightarrow |P| = \pm 2$$

$$23. |\text{adj } A| = |A|^{n-1} = |A|^{2-1} = |A| \\ \text{Adj}(\text{adj } A) = |A|^{n-2} A = |A|^0 A = A$$

\therefore option (B) is the correct answer.

$$24. \text{adj } AB - (\text{adj } B) (\text{adj } A) \\ = (\text{adj } B) (\text{adj } A) - (\text{adj } B) (\text{adj } A) \\ \dots [\because \text{adj } AB = (\text{adj } B) (\text{adj } A)]$$

$$= 0$$

25. Since, $AA^{-1} = I$

$$\therefore \begin{bmatrix} 2x & 0 \\ x & x \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 2x & 0 \\ 0 & 2x \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

By equality of matrices,

$$x = \frac{1}{2}$$

26. Since, $AA^{-1} = I$

$$\therefore \begin{bmatrix} x & -2 \\ 3 & 7 \end{bmatrix} \begin{bmatrix} \frac{7}{34} & \frac{1}{17} \\ -\frac{3}{34} & \frac{2}{17} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} \frac{7x+6}{34} & \frac{x-4}{17} \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

By equality of matrices,

$$\frac{x-4}{17} = 0 \Rightarrow x-4 = 0$$

$$\Rightarrow x = 4$$

$$27. 10 A^{-1} = \begin{bmatrix} 4 & 2 & 2 \\ -5 & 0 & \alpha \\ 1 & -2 & 3 \end{bmatrix} \quad \dots [\because B = A^{-1}]$$

$$\Rightarrow 10 A^{-1} A = \begin{bmatrix} 4 & 2 & 2 \\ -5 & 0 & \alpha \\ 1 & -2 & 3 \end{bmatrix} A$$

$$\Rightarrow 10 I = \begin{bmatrix} 4 & 2 & 2 \\ -5 & 0 & \alpha \\ 1 & -2 & 3 \end{bmatrix} \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & -3 \\ 1 & 1 & 1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 10 & 0 & 0 \\ 0 & 10 & 0 \\ 0 & 0 & 10 \end{bmatrix} = \begin{bmatrix} 10 & 0 & 0 \\ -5+\alpha & 5+\alpha & -5+\alpha \\ 0 & 0 & 10 \end{bmatrix}$$

$$\therefore -5 + \alpha = 0 \Rightarrow \alpha = 5$$



$$28. |A| = \begin{vmatrix} 5 & 4 \\ 3 & 2 \end{vmatrix} = -2$$

$$\text{adj } A = \begin{bmatrix} 2 & -4 \\ -3 & 5 \end{bmatrix}$$

$$A^{-1} = -\frac{1}{2} \begin{bmatrix} 2 & -4 \\ -3 & 5 \end{bmatrix}$$

$$29. |A| = \begin{vmatrix} 2 & -3 \\ -4 & 2 \end{vmatrix} = -8$$

$$\text{adj } A = \begin{bmatrix} 2 & 3 \\ 4 & 2 \end{bmatrix}$$

$$A^{-1} = -\frac{1}{8} \begin{bmatrix} 2 & 3 \\ 4 & 2 \end{bmatrix}$$

$$30. |U| = \begin{vmatrix} \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{vmatrix} = 1 \neq 0$$

$$\text{adj } U = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$\therefore U^{-1} = \frac{1}{|U|} (\text{adj } U) = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix} = U^T$$

$$31. |A| = \begin{vmatrix} a & c \\ d & b \end{vmatrix} = ab - cd$$

$$\text{adj } A = \begin{bmatrix} b & -c \\ -d & a \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{ab - cd} \begin{bmatrix} b & -c \\ -d & a \end{bmatrix}$$

$$32. \text{ The inverse of diagonal matrix } \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix} \text{ is}$$

$$\begin{bmatrix} \frac{1}{a} & 0 & 0 \\ 0 & \frac{1}{b} & 0 \\ 0 & 0 & \frac{1}{c} \end{bmatrix}$$

\therefore The inverse of the given diagonal matrix is

$$A^{-1} = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{3} & 0 \\ 0 & 0 & \frac{1}{4} \end{bmatrix}$$

$$33. \text{ If } B = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}, \text{ then } B^{-1} = \begin{bmatrix} \frac{1}{a} & 0 & 0 \\ 0 & \frac{1}{b} & 0 \\ 0 & 0 & \frac{1}{c} \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} \frac{2}{k} & 0 & 0 \\ 0 & \frac{3}{m} & 0 \\ 0 & 0 & \frac{4}{m} \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{3} & 0 \\ 0 & 0 & \frac{1}{4} \end{bmatrix}$$

$$\Rightarrow \frac{2}{k} = \frac{1}{2} \Rightarrow k = 4,$$

$$\frac{3}{m} = \frac{1}{3} \Rightarrow m = 9 \text{ and}$$

$$\frac{4}{m} = \frac{1}{4} \Rightarrow m = 16$$

$$\therefore k + m = 4 + 9 + 16 = 29$$

$$34. |A| = \begin{vmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{vmatrix} = -1 \neq 0$$

$$\text{adj } A = \begin{bmatrix} 0 & -1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} = A$$

$$35. |A| = \begin{vmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -1 \end{vmatrix} = -1 \neq 0$$

$$\text{adj } A = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



$$\begin{aligned} \therefore A^{-1} &= \frac{1}{|A|} (\text{adj } A) \\ &= \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{bmatrix} = A^T \end{aligned}$$

$$\begin{aligned} 36. \text{ Let } A &= \begin{bmatrix} 1 & 0 & 0 \\ 3 & 3 & 0 \\ 5 & 2 & -1 \end{bmatrix} \\ |A| &= -3 \neq 0 \\ \text{adj } A &= \begin{bmatrix} -3 & 0 & 0 \\ 3 & -1 & 0 \\ -9 & -2 & 3 \end{bmatrix} \\ A^{-1} &= \frac{1}{|A|} \text{adj } A \\ &= \frac{-1}{3} \begin{bmatrix} -3 & 0 & 0 \\ 3 & -1 & 0 \\ -9 & -2 & 3 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} 37. \quad |A| &= \begin{vmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{vmatrix} = -2 \neq 0 \\ \text{adj } A &= \begin{bmatrix} -1 & 1 & -1 \\ 8 & -6 & 2 \\ -5 & 3 & -1 \end{bmatrix} \\ \therefore A^{-1} &= \frac{1}{-2} \begin{bmatrix} -1 & 1 & -1 \\ 8 & -6 & 2 \\ -5 & 3 & -1 \end{bmatrix} \\ &= \begin{bmatrix} \frac{1}{2} & \frac{-1}{2} & \frac{1}{2} \\ -4 & 3 & -1 \\ \frac{5}{2} & \frac{-3}{2} & \frac{1}{2} \end{bmatrix} \end{aligned}$$

$$\begin{aligned} 38. \text{ Let } A &= \begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ b & c & 1 \end{bmatrix} \Rightarrow |A| = 1 \neq 0 \\ \text{adj } A &= \begin{bmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ ac-b & -c & 1 \end{bmatrix} \\ \therefore A^{-1} &= \begin{bmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ ac-b & -c & 1 \end{bmatrix} \end{aligned}$$

$$39. \quad |A| = \begin{vmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{vmatrix} = 1 \neq 0$$

$$\text{adj } A = \begin{bmatrix} 1 & -1 & 0 \\ -2 & 3 & -4 \\ -2 & 3 & -3 \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} 1 & -1 & 0 \\ -2 & 3 & -4 \\ -2 & 3 & -3 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 3 & -4 & 4 \\ 0 & -1 & 0 \\ -2 & 2 & -3 \end{bmatrix}$$

$$A^3 = A^2 \cdot A = \begin{bmatrix} 1 & -1 & 0 \\ -2 & 3 & -4 \\ -2 & 3 & -3 \end{bmatrix} = A^{-1}$$

$$40. \quad A = [a_{ij}]_{2 \times 2} \Rightarrow A = \begin{bmatrix} -1 & 3 \\ 3 & 0 \end{bmatrix}$$

$$\therefore |A| = -9$$

$$\text{adj } A = \begin{bmatrix} 0 & -3 \\ -3 & -1 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{-1}{9} \begin{bmatrix} 0 & -3 \\ -3 & -1 \end{bmatrix} = \frac{1}{9} \begin{bmatrix} 0 & 3 \\ 3 & 1 \end{bmatrix}$$

$$41. \text{ Let } A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix} \Rightarrow |A| = -2 \neq 0$$

Now, co-factor of element a_{32} of $A = A_{32}$

$$\therefore A_{32} = (-1)^{3+2} \begin{vmatrix} 1 & 1 \\ 2 & 0 \end{vmatrix} = 2$$

$$\therefore \text{Element } a_{23} \text{ of } A^{-1} = \frac{A_{32}}{|A|} = \frac{2}{-2} = -1$$

Alternate method:

$$|A| = -2 \neq 0$$

$$\text{adj } A = \begin{bmatrix} 1 & -2 & -1 \\ -2 & 2 & 2 \\ 1 & -2 & -3 \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} -\frac{1}{2} & 1 & \frac{1}{2} \\ 1 & -1 & -1 \\ -\frac{1}{2} & 1 & \frac{3}{2} \end{bmatrix}$$

$$\therefore \text{Element } a_{23} \text{ of } A^{-1} = -1.$$



$$42. \text{ Let } A = \begin{bmatrix} 1 & 2 & -3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \Rightarrow |A| = \begin{vmatrix} 1 & 2 & -3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{vmatrix} = 1 \neq 0$$

$$A_{31} = (-1)^{3+1} \begin{vmatrix} 2 & -3 \\ 1 & 2 \end{vmatrix} = 7$$

$$\therefore \text{ Element } a_{13} \text{ of } A^{-1} = \frac{A_{31}}{|A|} = 7$$

$$43. A = \begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} \frac{1}{2} & \frac{-1}{2} & \frac{1}{2} \\ -4 & 3 & -1 \\ \frac{5}{2} & \frac{-3}{2} & \frac{1}{2} \end{bmatrix}$$

$$\therefore \text{ sum of all the diagonal entries} = \frac{1}{2} + 3 + \frac{1}{2} = 4$$

$$44. A = \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix}$$

$$Ax = I \Rightarrow A^{-1}Ax = A^{-1}I$$

$$\Rightarrow x = A^{-1}$$

$$|A| = -5$$

$$\therefore A^{-1} = \frac{-1}{5} \begin{bmatrix} 3 & -2 \\ -4 & 1 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} -3 & 2 \\ 4 & -1 \end{bmatrix}$$

$$45. A \begin{bmatrix} 1 & 5 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & -1 \\ 6 & 0 \end{bmatrix}$$

$$\Rightarrow A = \begin{bmatrix} 3 & -1 \\ 6 & 0 \end{bmatrix} \begin{bmatrix} 1 & 5 \\ 0 & 1 \end{bmatrix}^{-1}$$

$$= \begin{bmatrix} 3 & -1 \\ 6 & 0 \end{bmatrix} \begin{bmatrix} 1 & -5 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & -16 \\ 6 & -30 \end{bmatrix}$$

$$46. |A| = \begin{vmatrix} 3 & 2 \\ 0 & 1 \end{vmatrix} = 3, \text{ adj } A = \begin{bmatrix} 1 & -2 \\ 0 & 3 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{3} \begin{bmatrix} 1 & -2 \\ 0 & 3 \end{bmatrix}$$

$$\begin{aligned} (A^{-1})^3 &= \frac{1}{27} \begin{bmatrix} 1 & -2 \\ 0 & 3 \end{bmatrix}^3 \\ &= \frac{1}{27} \begin{bmatrix} 1 & -2 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 & -2 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 & -2 \\ 0 & 3 \end{bmatrix} \\ &= \frac{1}{27} \begin{bmatrix} 1 & -26 \\ 0 & 27 \end{bmatrix} \end{aligned}$$

$$47. |A| = \begin{vmatrix} 0 & 3 \\ 2 & 0 \end{vmatrix} = -6 \neq 0$$

$$A^{-1} = \frac{1}{|A|} \text{adj } A = \lambda(\text{adj } A) \quad \dots[\text{Given}]$$

$$\therefore \lambda = \frac{1}{|A|} = -\frac{1}{6}$$

$$48. |A| = \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & -2 & 4 \end{vmatrix} = 6 \neq 0$$

$$\text{adj } A = \begin{bmatrix} 6 & 0 & 0 \\ 0 & 4 & -1 \\ 0 & 2 & 1 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{6} \begin{bmatrix} 6 & 0 & 0 \\ 0 & 4 & -1 \\ 0 & 2 & 1 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & -2 & 4 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & -2 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 5 \\ 0 & -10 & 14 \end{bmatrix}$$

$$\therefore A^2 + cA + dI$$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 5 \\ 0 & -10 & 14 \end{bmatrix} + \begin{bmatrix} c & 0 & 0 \\ 0 & c & c \\ 0 & -2c & 4c \end{bmatrix} + \begin{bmatrix} d & 0 & 0 \\ 0 & d & 0 \\ 0 & 0 & d \end{bmatrix}$$

$$= \begin{bmatrix} 1+c+d & 0 & 0 \\ 0 & -1+c+d & 5+c \\ 0 & -10-2c & 14+4c+d \end{bmatrix}$$

$$\text{Since, } 6A^{-1} = A^2 + cA + dI$$

$$\therefore \begin{bmatrix} 6 & 0 & 0 \\ 0 & 4 & -1 \\ 0 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 1+c+d & 0 & 0 \\ 0 & -1+c+d & 5+c \\ 0 & -10-2c & 14+4c+d \end{bmatrix}$$

$$\therefore \text{ by equality of matrices,}$$

$$1 + c + d = 6 \text{ and } 5 + c = -1,$$

$$\therefore c = -6 \text{ and } d = 11$$

$$49. \text{ By definition of inverse,}$$

$$I_3 I_3^{-1} = I_3$$

$$\Rightarrow I_3^{-1} = I_3$$

$$50. A^3 = I$$

$$\Rightarrow A^{-1}A^3 = A^{-1}I$$

$$\Rightarrow (A^{-1}A)A^2 = A^{-1}$$

$$\Rightarrow IA^2 = A^{-1} \Rightarrow A^2 = A^{-1}$$



51. $A^2 - A + I = 0$
 $\Rightarrow A \cdot A - A + I = 0$
 $\Rightarrow A^{-1} \cdot A \cdot A - A^{-1} \cdot A + A^{-1} \cdot I = 0$
 $\Rightarrow A - I + A^{-1} = 0$
 $\Rightarrow A^{-1} = I - A$
52. Given, $B = -A^{-1}BA$
 $\therefore AB = -AA^{-1}BA$
 $\Rightarrow AB = -I(BA) \Rightarrow AB = -BA$
 Now $(A + B)^2 = (A + B)(A + B)$
 $= A^2 + AB + BA + B^2$
 $= A^2 + B^2 \quad [\because BA = -AB]$
53. $(A^{-1}BA)^2 = (A^{-1}BA)(A^{-1}BA)$
 $= A^{-1}B(AA^{-1})BA$
 $= A^{-1}BIBA$
 $= A^{-1}B^2A$
 $(A^{-1}BA)^3 = (A^{-1}B^2A)(A^{-1}BA)$
 $= A^{-1}B^2(AA^{-1})BA$
 $= A^{-1}B^2IBA$
 $= A^{-1}B^3A$
- In general,
 $(A^{-1}BA)^n = A^{-1}B^nA$
54. $(M^{-1})^{-1} \neq (M^{-1})^1$
 $\therefore (M^{-1})^{-1} = (M^{-1})^1$ is not true
55. $(B^{-1}A^{-1})^{-1} = (A^{-1})^{-1} \cdot (B^{-1})^{-1} = A \cdot B$
 $\therefore A \cdot B = \begin{bmatrix} 2 & 2 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 2 & -2 \\ 2 & 3 \end{bmatrix}$
56. $(A^2 - 5A)A^{-1} = A \cdot A \cdot A^{-1} - 5A \cdot A^{-1}$
 $= A - 5I$
 $= \begin{bmatrix} 1 & 2 & 3 \\ -1 & 1 & 2 \\ 1 & 2 & 4 \end{bmatrix} - \begin{bmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$
 $= \begin{bmatrix} -4 & 2 & 3 \\ -1 & -4 & 2 \\ 1 & 2 & -1 \end{bmatrix}$
57. $\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$
 $\Rightarrow x + y = 2$ and $-x + y = 4$
 $\Rightarrow x = -1, y = 3$
58. $\begin{bmatrix} 1 & 2 & -3 \\ 0 & 4 & 5 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$
 $\Rightarrow z = 1$
 $4y + 5z = 1$
 $\Rightarrow y = -1$
 $x + 2y - 3z = 1$
 $\Rightarrow x = 6$

59. Let $A = \begin{bmatrix} 1 & 0 & 1 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ and $B = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$

Now $AX = B$

Applying $R_1 \rightarrow R_1 + R_2$,

$$\begin{bmatrix} 0 & 1 & 1 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 2 \end{bmatrix}$$

Applying $R_1 \rightarrow R_1 + R_3$,

$$\begin{bmatrix} 0 & 0 & 2 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 4 \\ 1 \\ 2 \end{bmatrix}$$

$$\begin{aligned} \therefore 2z &= 4 \Rightarrow z = 2 \\ -y + z &= 2 \Rightarrow y = 0 \\ -x + y &= 1 \Rightarrow x = -1 \\ \therefore (x, y, z) &= (-1, 0, 2) \end{aligned}$$

60. Applying $R_2 \rightarrow R_2 + 2R_1$,

$$\begin{bmatrix} 1 & 1 & 1 \\ 3 & 0 & 0 \\ 1 & 3 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \\ 4 \end{bmatrix}$$

Applying $R_1 \rightarrow R_1 - R_3$,

$$\begin{bmatrix} 0 & -2 & 0 \\ 3 & 0 & 0 \\ 1 & 3 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -4 \\ 3 \\ 4 \end{bmatrix}$$

$$\begin{aligned} \therefore -2y &= -4 \Rightarrow y = 2 \\ 3x &= 3 \Rightarrow x = 1 \\ x + 3y + z &= 4 \Rightarrow z = -3 \\ \therefore \begin{bmatrix} x \\ y \\ z \end{bmatrix} &= \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix} \end{aligned}$$

61. Applying $R_1 \rightarrow R_1 - R_3$

$$\begin{bmatrix} 0 & 0 & -1 \\ 1 & 4 & 4 \\ 1 & 3 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -1 \\ 15 \\ 13 \end{bmatrix}$$

Applying $R_2 \rightarrow R_2 - R_3$

$$\begin{bmatrix} 0 & 0 & -1 \\ 0 & 1 & 0 \\ 1 & 3 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ 13 \end{bmatrix}$$

$$\begin{aligned} \therefore -z &= -1 \Rightarrow z = 1 \\ y &= 2 \\ x + 3y + 4z &= 13 \Rightarrow x = 3 \\ \therefore (x, y, z) &= (3, 2, 1) \end{aligned}$$



62. Let $M = \begin{bmatrix} a & b & c \\ x & y & z \\ m & n & \end{bmatrix}$, then

$$M \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ 3 \end{bmatrix} \Rightarrow \begin{bmatrix} b \\ y \\ m \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ 3 \end{bmatrix}$$

∴ by the equality of matrices,
 $b = -1, y = 2, m = 3$

$$M \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} \Rightarrow \begin{bmatrix} a-b \\ x-y \\ -m \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$$

∴ by the equality of matrices,
 $a - b = 1, x - y = 1, -m = -1$
 $\Rightarrow a = 0, x = 3, m = 2$

$$M \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 12 \end{bmatrix} \Rightarrow \begin{bmatrix} a+b+c \\ x+y+z \\ +m+n \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 12 \end{bmatrix}$$

∴ by the equality of matrices,
 $a + b + c = 0, x + y + z = 0, +m + n = 12$
 $\Rightarrow c = 1, z = -5, n = 7$

∴ sum of diagonal elements of $M = a + y + n$
 $= 0 + 2 + 7 = 9$

63. Let $U_1 = \begin{bmatrix} a_1 \\ b_1 \\ c_1 \end{bmatrix}$, $U_2 = \begin{bmatrix} a_2 \\ b_2 \\ c_2 \end{bmatrix}$ and $U_3 = \begin{bmatrix} a_3 \\ b_3 \\ c_3 \end{bmatrix}$

$$AU_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$$\therefore \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ b_1 \\ c_1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$$\therefore \begin{bmatrix} a_1 \\ 2a_1 + b_1 \\ 3a_1 + 2b_1 + c_1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

∴ by the equality of matrices,
 $a_1 = 1, b_1 = -2$ and $c_1 = 1$
 Similarly $a_2 = 2, b_2 = -1$ and $c_2 = -4$
 $a_3 = 2, b_3 = -1$ and $c_3 = -3$

$$\therefore U = \begin{bmatrix} 1 & 2 & 2 \\ -2 & -1 & -1 \\ 1 & -4 & -3 \end{bmatrix}$$

$$|U| = \begin{vmatrix} 1 & 2 & 2 \\ -2 & -1 & -1 \\ 1 & -4 & -3 \end{vmatrix} = 3$$

∴ U^{-1} exists

$$\therefore U^{-1} = \begin{bmatrix} -\frac{1}{3} & -\frac{2}{3} & 0 \\ -\frac{7}{3} & -\frac{5}{3} & -1 \\ 3 & 2 & 1 \end{bmatrix}$$

∴ sum of elements of $U^{-1} = 0$

64. $a_{11}A_{11} + a_{12}A_{12} + a_{13}A_{13}$
 $= \cos\theta(\cos\theta - 0) + \sin\theta[-(-\sin\theta - 0)] + 0(0-0)$
 $= \cos^2\theta + \sin^2\theta = 1$

65. $|A| = 1 + \tan^2 \frac{\theta}{2} = \sec^2 \frac{\theta}{2}$

$$\text{adj}A = \begin{bmatrix} 1 & -\tan \frac{\theta}{2} \\ \tan \frac{\theta}{2} & 1 \end{bmatrix}$$

$$A^{-1} = \frac{1}{|A|} \text{adj}A = \frac{1}{\sec^2 \frac{\theta}{2}} \begin{bmatrix} 1 & -\tan \frac{\theta}{2} \\ \tan \frac{\theta}{2} & 1 \end{bmatrix}$$

$$AB = I \Rightarrow B = IA^{-1} \Rightarrow B = A^{-1}$$

$$\therefore B = \frac{1}{\sec^2 \frac{\theta}{2}} \begin{bmatrix} 1 & -\tan \frac{\theta}{2} \\ \tan \frac{\theta}{2} & 1 \end{bmatrix} = \cos^2 \frac{\theta}{2} \cdot A^T$$

66. $F(\alpha) \cdot F(-\alpha)$
 $= \begin{bmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \alpha & \sin \alpha & 0 \\ -\sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$
 $= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I$

∴ $[F(\alpha)]^{-1} = F(-\alpha)$

67. $I + A = \begin{bmatrix} 1 & -\tan \alpha \\ \tan \alpha & 1 \end{bmatrix}$

$$I - A = \begin{bmatrix} 1 & \tan \alpha \\ -\tan \alpha & 1 \end{bmatrix}$$

∴ $|I - A| = 1 + \tan^2 \alpha = \sec^2 \alpha \neq 0$



$$\text{adj}(I - A) = \begin{bmatrix} 1 & \tan \alpha \\ -\tan \alpha & 1 \end{bmatrix}^T = \begin{bmatrix} 1 & -\tan \alpha \\ \tan \alpha & 1 \end{bmatrix}$$

$$\Rightarrow (I - A)^{-1} = \frac{1}{|I - A|} [\text{adj}(I - A)]$$

$$= \begin{bmatrix} \cos^2 \alpha & -\sin \alpha \cos \alpha \\ \sin \alpha \cos \alpha & \cos^2 \alpha \end{bmatrix}$$

$$\therefore (I + A)(I - A)^{-1}$$

$$= \begin{bmatrix} 1 & -\tan \alpha \\ \tan \alpha & 1 \end{bmatrix} \begin{bmatrix} \cos^2 \alpha & -\sin \alpha \cos \alpha \\ \sin \alpha \cos \alpha & \cos^2 \alpha \end{bmatrix}$$

$$= \begin{bmatrix} \cos^2 \alpha - \sin^2 \alpha & -2 \sin \alpha \cos \alpha \\ 2 \sin \alpha \cos \alpha & \cos^2 \alpha - \sin^2 \alpha \end{bmatrix}$$

$$\therefore (I + A)(I - A)^{-1} = \begin{bmatrix} \cos 2\alpha & -\sin 2\alpha \\ \sin 2\alpha & \cos 2\alpha \end{bmatrix} \dots \text{(i)}$$

$$B(\alpha) = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

$$\therefore B(2\alpha) = \begin{bmatrix} \cos 2\alpha & -\sin 2\alpha \\ \sin 2\alpha & \cos 2\alpha \end{bmatrix} \dots \text{(ii)}$$

$$\therefore (I + A)(I - A)^{-1} = B(2\alpha) \dots [\text{From (i) and (ii)}]$$

$$68. \quad BB' = (A^{-1}A')(A^{-1}A')'$$

$$= (A^{-1}A')(A(A^{-1})')$$

$$= A^{-1}AA'(A^{-1})'$$

$$\dots [\because AA' = A'A \text{ (given)}]$$

$$= (A^{-1}A)(A'(A^{-1})')$$

$$= I(A^{-1}A)' = I.I = I^2 = I$$

$$69. \quad M^2N^2(M^T N)^{-1}(MN^{-1})^T$$

$$= M^2N^2N^{-1}(M^T)^{-1}(N^{-1})^T M^T$$

$$= M^2N(M^T)^{-1}(N^T)^{-1}M^T$$

$$= M^2N(-M)^{-1}(-N)^{-1}(-M)$$

$$\dots [\because \text{For skew-symmetric matrices } M \text{ and } N,$$

$$M^T = -M, N^T = -N]$$

$$= -M^2NM^{-1}N^{-1}M \quad \dots [\because (kA)^{-1} = \frac{1}{k}A^{-1}]$$

$$= -M(MN)(NM)^{-1}M$$

$$= -M(NM)(NM)^{-1}M$$

$$\dots [\because MN = NM \text{ (given)}]$$

$$= -M.I.M = -M^2$$



Evaluation Test

$$2. \quad |A| = \begin{vmatrix} 1 & 2 & 2 \\ -2 & -1 & -1 \\ 1 & -4 & -3 \end{vmatrix} = 1(-1) - 2(7) + 2(9)$$

$$= 3 \neq 0$$

$\therefore A^{-1}$ exists.

$$\therefore \text{adj } A = \begin{bmatrix} -1 & -7 & 9 \\ -2 & -5 & 6 \\ 0 & -3 & 3 \end{bmatrix}^T = \begin{bmatrix} -1 & -2 & 0 \\ -7 & -5 & -3 \\ 9 & 6 & 3 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{3} \begin{bmatrix} -1 & -2 & 0 \\ -7 & -5 & -3 \\ 9 & 6 & 3 \end{bmatrix}$$

$$\therefore \text{sum of the elements of } A^{-1}$$

$$= \frac{1}{3}(-1 - 2 + 0 - 7 - 5 - 3 + 9 + 6 + 3) = 0$$

$$3. \quad (\text{adj } A)A = |A|I_n$$

$$A = \begin{vmatrix} 2 & 0 & 3 \\ 1 & -1 & 2 \\ 3 & -2 & 0 \end{vmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= 11 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 11 & 0 & 0 \\ 0 & 11 & 0 \\ 0 & 0 & 11 \end{bmatrix}$$

$$4. \quad \text{Let } A = \begin{bmatrix} 1 & 2 & 4 \\ 3 & -19 & 7 \\ 2 & 4 & 8 \end{bmatrix}$$

$$\Rightarrow |A| = 0$$

$\therefore A^{-1}$ does not exist.

$$5. \quad \begin{bmatrix} 1 & \tan \theta \\ -\tan \theta & 1 \end{bmatrix}^{-1} = \frac{1}{\sec^2 \theta} \begin{bmatrix} 1 & \tan \theta \\ -\tan \theta & 1 \end{bmatrix}^T$$

$$= \frac{1}{\sec^2 \theta} \begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix}$$

$$\therefore \begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix} \cdot \frac{1}{\sec^2 \theta} \begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix}$$

$$= \frac{1}{\sec^2 \theta} \begin{bmatrix} 1 - \tan^2 \theta & -2 \tan \theta \\ 2 \tan \theta & 1 - \tan^2 \theta \end{bmatrix}$$

$$= \begin{bmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix}$$

By equality of matrices, we get

$$a = \cos 2\theta, b = \sin 2\theta$$



$$6. \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 1 \\ 3 & 2 & 9 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 6 \\ 7 \\ 14 \end{bmatrix}$$

$$R_2 \rightarrow R_2 - 2R_1, R_3 \rightarrow R_3 - 3R_1$$

$$\therefore \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & -5 \\ 0 & -4 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 6 \\ -5 \\ -4 \end{bmatrix}$$

$$\therefore -5z = -5 \Rightarrow z = 1$$

$$-4y = -4 \Rightarrow y = 1$$

$$x + 2y + 3z = 6 \Rightarrow x = 1$$

$$7. \text{ Let } A = \begin{bmatrix} 1 & -2 & -1 \\ 2 & \lambda & 3 \\ -1 & 0 & 3 \end{bmatrix}$$

Matrix will not be invertible if $|A| = 0$

$$\therefore \begin{vmatrix} 1 & -2 & -1 \\ 2 & \lambda & 3 \\ -1 & 0 & 3 \end{vmatrix} = 0$$

$$\Rightarrow 1(3\lambda) + 2(9) - 1(\lambda) = 0$$

$$\Rightarrow \lambda = -9$$

$$8. \text{ Given, } |A| \neq 0 \text{ and } |B| = 0$$

$$\therefore |AB| = |A| |B| = 0$$

$$\text{and } |A^{-1} B| = |A^{-1}| |B|$$

$$= \frac{1}{|A|} |B| \quad \dots \left[\because |A^{-1}| = \frac{1}{|A|} \right]$$

$$= 0$$

$$\therefore AB \text{ and } A^{-1} B \text{ are singular.}$$

$$9. (AB)^{-1} = B^{-1} A^{-1}$$

$$\therefore B^{-1} A^{-1} = \begin{bmatrix} \frac{-1}{2} & \frac{1}{2} \\ \frac{1}{4} & 0 \end{bmatrix}$$

$$10. (A^2 - 8A)A^{-1} = A.A.A^{-1} - 8A.A^{-1}$$

$$= A - 8I$$

$$= \begin{bmatrix} 1 & 4 & 4 \\ 4 & 1 & 4 \\ 4 & 4 & 1 \end{bmatrix} - \begin{bmatrix} 8 & 0 & 0 \\ 0 & 8 & 0 \\ 0 & 0 & 8 \end{bmatrix}$$

$$= \begin{bmatrix} -7 & 4 & 4 \\ 4 & -7 & 4 \\ 4 & 4 & -7 \end{bmatrix}$$

$$11. \det A = \begin{vmatrix} 1 & 2 & 0 \\ -1 & 1 & 2 \\ 2 & -1 & 1 \end{vmatrix}$$

$$= 13$$

$$\therefore \det(\text{adj}(\text{adj} A)) = (\det A)^{(3-1)^2}$$

$$\dots \left[\because |\text{adj}(\text{adj} A)| = |A|^{(n-1)^2} \right]$$

$$= (\det A)^4 = (13)^4$$

$$12. A. (\text{adj} A) = \begin{bmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix} \quad \dots(i)$$

$$= 4 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= 4.I$$

$$\text{Since, } A(\text{adj} A) = |A|.I$$

$$\therefore |A| = 4$$

$$\text{From (i), } |A| \cdot |\text{adj} A| = 64$$

$$\Rightarrow |\text{adj} A| = \frac{64}{4} = 16$$

$$\text{Also, } |\text{adj}(\text{adj} A)| = |A|^{(n-1)^2}$$

$$= |A|^{(3-1)^2}$$

$$= (4)^4 = 256$$

$$\therefore \frac{|\text{adj}(\text{adj} A)|}{|\text{adj} A|} = \frac{256}{16} = 16$$

$$13. \text{ Since, } A(\text{adj} A) = |A|.I$$

Replacing A by $\text{adj} A$, we get

$$\text{adj} A (\text{adj}(\text{adj} A)) = |\text{adj} A|.I$$

$$\Rightarrow A^{-1} \cdot |A| (\text{adj}(\text{adj} A)) = |\text{adj} A|.I$$

$$\dots \left[\because A^{-1} = \frac{1}{|A|} (\text{adj} A) \right]$$

$$\Rightarrow \alpha A^{-1} (\text{adj}(\text{adj} A)) = |A|^2.I$$

$$\dots \left[\because |\text{adj} A| = |A|^{n-1} \right]$$

$$\Rightarrow \alpha A^{-1} (\text{adj}(\text{adj} A)) = \alpha^2 I$$

$$\Rightarrow A^{-1} (\text{adj}(\text{adj} A)) = \alpha I$$

$$\text{Given, } A^{-1} (\text{adj}(\text{adj} A)) = kI$$

$$\therefore k = \alpha$$



Hints



Classical Thinking

2. $\tan \theta = \cot \alpha \Rightarrow \tan \theta = \tan \left(\frac{\pi}{2} - \alpha \right)$
 $\Rightarrow \theta = n\pi + \frac{\pi}{2} - \alpha$
[$\because \tan \theta = \tan \alpha \Rightarrow \theta = n\pi + \alpha$]
3. $\tan 3x = 1$
 $\therefore \tan 3x = \tan \frac{\pi}{4} \Rightarrow 3x = n\pi + \frac{\pi}{4}$
[$\because \tan \theta = \tan \alpha$
 $\Rightarrow \theta = n\pi + \alpha$]
- $\therefore x = \frac{n\pi}{3} + \frac{\pi}{12}, n \in I$
4. $\tan 3x = \cot x \Rightarrow \tan 3x = \tan \left(\frac{\pi}{2} - x \right)$
 $\therefore 3x = n\pi + \frac{\pi}{2} - x \Rightarrow 4x = n\pi + \frac{\pi}{2}$
 $\therefore x = \frac{n\pi}{4} + \frac{\pi}{8} = (2n+1) \frac{\pi}{8}$
5. $\sin^2 \theta + \sin \theta = 2$
 $\therefore (\sin \theta - 1)(\sin \theta + 2) = 0$
 $\therefore \sin \theta = 1, -2$
 Since, $\sin \theta \neq -2$
 $\therefore \sin \theta = 1 = \sin \left(\frac{\pi}{2} \right)$
 $\therefore \theta = n\pi + (-1)^n \frac{\pi}{2}, n \in I$
[$\because \sin \theta = \sin \alpha$
 $\Rightarrow \theta = n\pi + (-1)^n \alpha$]
6. $\cot \theta - \tan \theta = 2 \Rightarrow \frac{\cos \theta}{\sin \theta} - \frac{\sin \theta}{\cos \theta} = 2$
 $\therefore \cos^2 \theta - \sin^2 \theta = \sin 2\theta \Rightarrow \cos 2\theta = \sin 2\theta$
 $\therefore \tan 2\theta = \tan \frac{\pi}{4} \Rightarrow 2\theta = n\pi + \frac{\pi}{4}$
 $\therefore \theta = \frac{n\pi}{2} + \frac{\pi}{8}$

7. $\sin^2 \theta = \frac{1}{4} = \sin^2 \frac{\pi}{6} \Rightarrow \theta = n\pi \pm \frac{\pi}{6}$
[$\because \sin^2 \theta = \sin^2 \alpha \Rightarrow \theta = n\pi \pm \alpha$]
8. $4\cos^2 x + 6\sin^2 x = 5$
 $\therefore 4 + 2\sin^2 x = 5$
 $\therefore \sin^2 x = \frac{1}{2} = \sin^2 \frac{\pi}{4} \Rightarrow x = n\pi \pm \frac{\pi}{4}$
9. $\sec^2 \theta + \tan^2 \theta = \frac{5}{3}$ (i)
 $\therefore 1 + \tan^2 \theta + \tan^2 \theta = \frac{5}{3}$
 $\therefore 2 \tan^2 \theta = \frac{2}{3}$
 $\therefore \tan^2 \theta = \frac{1}{3} = \tan^2 \left(\frac{\pi}{6} \right) \Rightarrow \theta = n\pi \pm \frac{\pi}{6}$
[$\because \tan^2 \theta = \tan^2 \alpha \Rightarrow \theta = n\pi \pm \alpha$]
10. $\tan \theta + \tan 2\theta + \sqrt{3} \tan \theta \tan 2\theta = \sqrt{3}$
 $\therefore \tan \theta + \tan 2\theta = \sqrt{3} (1 - \tan \theta \tan 2\theta)$
 $\therefore \frac{\tan \theta + \tan 2\theta}{1 - \tan \theta \tan 2\theta} = \sqrt{3} \Rightarrow \tan 3\theta = \tan \frac{\pi}{3}$
 $\therefore 3\theta = n\pi + \frac{\pi}{3} \Rightarrow \theta = (3n+1) \frac{\pi}{9}$
11. By sine rule,
 $\frac{\sin A}{a} = \frac{\sin B}{b}$
 $\Rightarrow \frac{2/3}{2} = \frac{\sin B}{3}$
 $\Rightarrow \sin B = 1 = \sin 90^\circ \Rightarrow B = 90^\circ$
12. $\frac{\sin B}{\sin(A+B)} = \frac{\sin B}{\sin C} = \frac{b}{c}$
[$\because A+B+C = \pi, A+B = \pi - C$]
13. $2s = a + b + c = 16 + 24 + 20 = 60 \Rightarrow s = 30$
 $\therefore \cos \frac{B}{2} = \sqrt{\frac{s(s-b)}{ac}} = \sqrt{\frac{30 \times 6}{320}} = \frac{3}{4}$



14. Let $a = 4$ cm, $b = 5$ cm, $c = 6$ cm

$$s = \frac{a+b+c}{2} = \frac{4+5+6}{2} = \frac{15}{2}$$

$$A(\Delta ABC) = \sqrt{s(s-a)(s-b)(s-c)}$$

$$= \sqrt{\frac{15}{2} \left(\frac{15}{2} - 4 \right) \left(\frac{15}{2} - 5 \right) \left(\frac{15}{2} - 6 \right)} = \frac{15}{4} \sqrt{7}$$

15.
$$2ac \sin \frac{A-B+C}{2} = 2ac \sin \frac{\pi-2B}{2}$$

$$= 2ac \cos B$$

$$= 2ac \frac{c^2 + a^2 - b^2}{2ca}$$

....[By cosine rule]

$$= c^2 + a^2 - b^2$$

16. $s - a = 3 \Rightarrow b + c - a = 6$ (i)
 $s - c = 2 \Rightarrow a + b - c = 4$ (ii)
 Adding (i) and (ii), we get $b = 5$
 Since, $\angle B = 90^\circ$
 $\therefore b^2 = a^2 + c^2 \Rightarrow a^2 + c^2 = 25$ (iii)
 Solving, we get $a = 3, c = 4$

17. We know that,

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = k$$

$$\Rightarrow \frac{b}{1} = \frac{c}{\sqrt{2}} \Rightarrow c - \sqrt{2} b = 0$$
(i)
 By projection rule,
 $a = b \cos C + c \cos B$

$$\Rightarrow \sqrt{3} + 1 = \frac{b}{\sqrt{2}} + \frac{\sqrt{3}}{2} c$$

$$\Rightarrow 2(\sqrt{3} + 1) = \sqrt{2} b + \sqrt{3} c$$
(ii)
 From (i) and (ii), we get
 $2(\sqrt{3} + 1) = (\sqrt{3} + 1) c \Rightarrow c = 2$

18. $s = \frac{a+b+c}{2} = \frac{12}{2} = 6$

$$\sin \frac{B}{2} = \sqrt{\frac{(s-c)(s-a)}{ca}} = \sqrt{\frac{2 \times 3}{12}} = \sqrt{\frac{1}{2}}$$

$$\cos \frac{B}{2} = \sqrt{\frac{s(s-b)}{ca}} = \sqrt{\frac{6 \times 1}{12}} = \sqrt{\frac{1}{2}}$$

$$\therefore \sin \frac{B}{2} + \cos \frac{B}{2} = \frac{2}{\sqrt{2}} = \sqrt{2}$$

19.
$$\frac{\sin \frac{A}{2} \sin \frac{C}{2}}{\sin \frac{B}{2}} = \sqrt{\frac{ac(s-b)(s-c)(s-b)(s-a)}{(s-a)(s-c)bc \times ab}}$$

$$= \frac{s-b}{b}$$

But a, b and c are in A. P. $\Rightarrow 2b = a + c$
 $\Rightarrow 2b + b = a + b + c$
 $\Rightarrow 3b = 2s \Rightarrow s = \frac{3b}{2}$

$$\therefore \frac{s-b}{b} = \frac{\frac{3b}{2} - b}{b} = \frac{1}{2}$$

20. By Napier's analogy, we have

$$\tan \frac{B-C}{2} = \frac{b-c}{b+c} \cot \frac{A}{2} \Rightarrow x = \frac{b-c}{b+c}$$

21.
$$\tan \frac{A-B}{2} = \frac{a-b}{a+b} \cot \frac{C}{2}$$

$$= \frac{a-b}{a+b} \tan \left(\frac{A+B}{2} \right)$$

....[$\because A+B+C = \pi$]

$$\therefore \tan \frac{A-B}{2} \cot \frac{A+B}{2} = \frac{a-b}{a+b}$$

26. $\sin^{-1} \left(\frac{\sqrt{3}}{2} \right) - \sin^{-1} \left(\frac{1}{2} \right) = 60^\circ - 30^\circ = 30^\circ$

27. $\sin^{-1} \frac{1}{2} = \tan^{-1} x$
 $\Rightarrow \frac{\pi}{6} = \tan^{-1} x \Rightarrow \tan \frac{\pi}{6} = x$
 $\Rightarrow x = \frac{1}{\sqrt{3}}$

28. Let $\theta = \sin^{-1} \left(\frac{3}{5} \right)$

$$\therefore \sin \left(2 \sin^{-1} \left(\frac{3}{5} \right) \right) = \sin 2\theta$$

$$= 2 \sin \theta \cos \theta$$

$$= 2 \sin \left(\sin^{-1} \left(\frac{3}{5} \right) \right) \cos \left(\sin^{-1} \left(\frac{3}{5} \right) \right)$$

$$= 2 \times \frac{3}{5} \sqrt{1 - \left(\frac{3}{5} \right)^2}$$

....[$\because \cos(\sin^{-1} x) = \sqrt{1-x^2}$]

$$= 2 \times \frac{3}{5} \times \frac{4}{5} = \frac{24}{25}$$



$$29. \sin\left(3\sin^{-1}\left(\frac{2}{5}\right)\right) = \sin 3\theta,$$

$$\text{Where } \theta = \sin^{-1}\left(\frac{2}{5}\right)$$

$$\dots\left[\theta = \sin^{-1}\left(\frac{2}{5}\right), \sin \theta = \frac{2}{5}\right]$$

$$= 3\sin \theta - 4\sin^3 \theta$$

$$= 3\left(\frac{2}{5}\right) - 4\left(\frac{2}{5}\right)^3$$

$$\dots\left[\theta = \sin^{-1}\left(\frac{2}{5}\right), \sin \theta = \frac{2}{5}\right]$$

$$= \frac{6}{5} - \frac{32}{125} = \frac{118}{125}$$

$$30. \cos^{-1}(\cos 12) - \sin^{-1}(\sin 14) = 12 - 14 = -2$$

$$31. \tan^{-1}\left(\tan \frac{3\pi}{4}\right) = \tan^{-1}\left(\tan\left(\pi - \frac{\pi}{4}\right)\right)$$

$$= \tan^{-1}\left(-\tan \frac{\pi}{4}\right)$$

$$= -\tan^{-1}\left(\tan \frac{\pi}{4}\right) = -\frac{\pi}{4}$$

$$32. \text{ If } x = \sec \theta, \text{ then } \sqrt{x^2 - 1} = \sqrt{\sec^2 \theta - 1} = \tan \theta$$

$$\therefore \cot^{-1} \frac{1}{\sqrt{x^2 - 1}} = \cot^{-1}(\cot \theta) = \theta = \sec^{-1} x$$

$$33. \cot^{-1}(-\sqrt{3}) = \pi - \cot^{-1}(\sqrt{3})$$

$$= \pi - \frac{\pi}{6} = \frac{5\pi}{6}$$

$$34. \cos^{-1}(-1) = \pi - \cos^{-1} 1 = \pi - 0 = \pi$$

$$35. \sin\left(\frac{\pi}{3} - \sin^{-1}\left(-\frac{1}{2}\right)\right) = \sin\left(\frac{\pi}{3} + \sin^{-1}\left(\frac{1}{2}\right)\right)$$

$$= \sin\left(\frac{\pi}{3} + \frac{\pi}{6}\right) = \sin \frac{\pi}{2} = 1$$

$$37. \cos^{-1}\left[\cos \frac{5\pi}{3}\right] + \sin^{-1}\left[\cos \frac{5\pi}{3}\right] = \frac{\pi}{2}$$

$$\dots\left[\because \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}\right]$$

$$38. \cos\left\{\cos^{-1}\left(\frac{-1}{7}\right) + \sin^{-1}\left(\frac{-1}{7}\right)\right\} = \cos \frac{\pi}{2} = 0$$

$$39. \cot^{-1} x + \cot^{-1} y = \left(\frac{\pi}{2} - \tan^{-1} x\right) + \left(\frac{\pi}{2} - \tan^{-1} y\right)$$

$$\dots\left[\because \tan^{-1} x + \cot^{-1} x = \frac{\pi}{2}\right]$$

$$= \pi - (\tan^{-1} x + \tan^{-1} y)$$

$$= \pi - \frac{4\pi}{5} = \frac{\pi}{5}$$

$$40. \tan^{-1}(\sqrt{3}) - \cot^{-1}(-\sqrt{3})$$

$$= \tan^{-1} \sqrt{3} - [\pi - \cot^{-1} \sqrt{3}]$$

$$= \tan^{-1} \sqrt{3} + \cot^{-1} \sqrt{3} - \pi$$

$$= \frac{\pi}{2} - \pi = -\frac{\pi}{2}$$

$$41. \tan^{-1} \frac{1}{2} + \tan^{-1} \frac{2}{11} = \tan^{-1} \frac{\frac{1}{2} + \frac{2}{11}}{1 - \frac{1}{2} \times \frac{2}{11}}$$

$$= \tan^{-1} \frac{15}{20} = \tan^{-1} \frac{3}{4}$$

$$42. \tan^{-1} x - \tan^{-1} y = \tan^{-1} A$$

$$\Rightarrow \tan^{-1} \left(\frac{x-y}{1+xy}\right) = \tan^{-1} A$$

$$\Rightarrow A = \frac{x-y}{1+xy}$$

$$43. \sin^{-1}\left(\frac{3}{5}\right) + \sin^{-1}\left(\frac{8}{17}\right)$$

$$= \sin^{-1}\left(\frac{3}{5}\sqrt{1-\left(\frac{8}{17}\right)^2} + \frac{8}{17}\sqrt{1-\left(\frac{3}{5}\right)^2}\right)$$

$$\dots\left[\because \sin^{-1} x + \sin^{-1} y = \sin^{-1} x\sqrt{1-y^2} + y\sqrt{1-x^2}\right]$$

$$= \sin^{-1}\left(\frac{3}{5} \times \frac{15}{17} + \frac{8}{17} \times \frac{4}{5}\right) = \sin^{-1}\left(\frac{77}{85}\right)$$

$$44. \cos^{-1} \frac{3}{5} - \sin^{-1} \frac{4}{5} = \cos^{-1} x$$

$$\therefore \cos^{-1} \frac{3}{5} - \cos^{-1} \sqrt{1 - \frac{16}{25}} = \cos^{-1} x$$

$$\therefore \cos^{-1} \frac{3}{5} - \cos^{-1} \frac{3}{5} = \cos^{-1} x$$

$$\therefore \cos^{-1} x = 0 \Rightarrow x = 1$$



Critical Thinking

$$1. \quad \tan \theta + \frac{1}{\sqrt{3}} = 0 \Rightarrow \tan \theta = -\frac{1}{\sqrt{3}}$$

$$\Rightarrow \tan \theta = -\tan 30^\circ$$

$$\Rightarrow \tan \theta = \tan (180^\circ - 30^\circ) \text{ and}$$

$$\tan \theta = \tan (360^\circ - 30^\circ)$$

$$\Rightarrow \tan \theta = \tan 150^\circ \text{ and } \tan \theta = \tan 330^\circ$$

$$\Rightarrow \theta = 150^\circ \text{ and } 330^\circ$$

$$2. \quad \cos \theta = 1 - 2x^2$$

$$\therefore \cos \theta = 1 - 2 \cos^2 40^\circ \quad \dots [\because \cos 40^\circ = x]$$

$$= -(2 \cos^2 40^\circ - 1)$$

$$= -\cos (2 \times 40^\circ) = -\cos 80^\circ$$

$$\therefore \cos \theta = \cos (180^\circ + 80^\circ) = \cos 260^\circ$$

$$\text{and } \cos \theta = \cos (180^\circ - 80^\circ) = \cos 100^\circ$$

$$\therefore \theta = 100^\circ \text{ and } 260^\circ$$

$$3. \quad \tan \theta = \sqrt{3} = \tan \frac{\pi}{3} \Rightarrow \theta = n\pi + \frac{\pi}{3}$$

For $-\pi < \theta < 0$,

Put $n = -1$, we get $\theta = -\pi + \frac{\pi}{3} = \frac{-2\pi}{3} = \frac{-4\pi}{6}$

$$4. \quad \cot \theta + \tan \theta = 2 \operatorname{cosec} \theta \Rightarrow \frac{1}{\sin \theta \cos \theta} = \frac{2}{\sin \theta}$$

$$\Rightarrow \cos \theta = \frac{1}{2} = \cos \left(\frac{\pi}{3} \right) \Rightarrow \theta = 2n\pi \pm \frac{\pi}{3}$$

$$5. \quad \tan \theta + \tan \left(\frac{\pi}{2} - \theta \right) = 2$$

$$\Rightarrow \tan \theta + \frac{1}{\tan \theta} = 2 \Rightarrow \tan^2 \theta - 2 \tan \theta + 1 = 0$$

$$\Rightarrow \tan \theta = 1 = \tan \frac{\pi}{4} \Rightarrow \theta = n\pi + \frac{\pi}{4}$$

$$6. \quad \sin \theta = -\frac{1}{2} = -\sin \left(\frac{\pi}{6} \right) = \sin \left(\pi + \frac{\pi}{6} \right)$$

$$\tan \theta = \frac{1}{\sqrt{3}} = \tan \left(\frac{\pi}{6} \right) = \tan \left(\pi + \frac{\pi}{6} \right)$$

$$\Rightarrow \theta = \left(\pi + \frac{\pi}{6} \right)$$

Hence, general value of θ is $2n\pi + \frac{7\pi}{6}$.

$$7. \quad \cos x - \sin x = \frac{1}{\sqrt{2}}$$

Dividing both sides by $\sqrt{2}$, we get

$$\frac{1}{\sqrt{2}} \cos x - \frac{1}{\sqrt{2}} \sin x = \frac{1}{2}$$

$$\Rightarrow \cos \left(\frac{\pi}{4} + x \right) = \cos \frac{\pi}{3} \Rightarrow \frac{\pi}{4} + x = 2n\pi \pm \frac{\pi}{3}$$

$$\Rightarrow x = 2n\pi + \frac{\pi}{3} - \frac{\pi}{4} = 2n\pi + \frac{\pi}{12}$$

$$\text{or } x = 2n\pi - \frac{\pi}{3} - \frac{\pi}{4} = 2n\pi - \frac{7\pi}{12}$$

$$8. \quad 1 + \cot \theta = \operatorname{cosec} \theta$$

$$\Rightarrow \frac{1}{\sin \theta} = 1 + \frac{\cos \theta}{\sin \theta} \Rightarrow \sin \theta + \cos \theta = 1$$

Dividing both sides by $\sqrt{2}$, we get

$$\sin \theta \sin \frac{\pi}{4} + \cos \theta \cos \frac{\pi}{4} = \cos \frac{\pi}{4}$$

$$\Rightarrow \cos \left(\theta - \frac{\pi}{4} \right) = \cos \frac{\pi}{4} \Rightarrow \theta - \frac{\pi}{4} = 2n\pi \pm \frac{\pi}{4}$$

$$\Rightarrow \theta = 2n\pi \text{ or } \theta = 2n\pi + \frac{\pi}{2}$$

$$9. \quad \sin x - \cos x = \sqrt{2}$$

$$\Rightarrow \sin x \frac{1}{\sqrt{2}} - \cos x \frac{1}{\sqrt{2}} = 1$$

$$\Rightarrow \cos \left(x + \frac{\pi}{4} \right) = -1 = \cos \pi$$

$$\Rightarrow x + \frac{\pi}{4} = 2n\pi \pm \pi$$

$$\Rightarrow x = 2n\pi + \frac{3\pi}{4} \text{ or } 2n\pi - \frac{5\pi}{4}$$

$$10. \quad \cot \theta + \cot \left(\frac{\pi}{4} + \theta \right) = 2$$

$$\therefore \frac{\cos \theta}{\sin \theta} + \frac{\cos \left(\frac{\pi}{4} + \theta \right)}{\sin \left(\frac{\pi}{4} + \theta \right)} = 2$$

$$\therefore \sin \left(\frac{\pi}{4} + 2\theta \right) = 2 \sin \theta \sin \left(\frac{\pi}{4} + \theta \right)$$

$$= \cos \left(\theta - \frac{\pi}{4} - \theta \right) - \cos \left(\theta + \frac{\pi}{4} + \theta \right)$$



$$\begin{aligned} \therefore \sin\left(\frac{\pi}{4} + 2\theta\right) &= \cos\left(\frac{-\pi}{4}\right) - \cos\left(2\theta + \frac{\pi}{4}\right) \\ \Rightarrow \sin\left(\frac{\pi}{4} + 2\theta\right) + \cos\left(\frac{\pi}{4} + 2\theta\right) &= \frac{1}{\sqrt{2}} \\ \Rightarrow \left(\frac{1}{\sqrt{2}}\cos 2\theta + \frac{1}{\sqrt{2}}\sin 2\theta\right) \\ &+ \left(\frac{1}{\sqrt{2}}\cos 2\theta - \frac{1}{\sqrt{2}}\sin 2\theta\right) = \frac{1}{\sqrt{2}} \\ \Rightarrow \frac{2}{\sqrt{2}}\cos 2\theta &= \frac{1}{\sqrt{2}} \Rightarrow \cos 2\theta = \frac{1}{2} = \cos\left(\frac{\pi}{3}\right) \\ \Rightarrow 2\theta &= 2n\pi \pm \frac{\pi}{3} \Rightarrow \theta = n\pi \pm \frac{\pi}{6} \end{aligned}$$

$$\begin{aligned} 11. \quad \sin^2 x - 2\cos x + \frac{1}{4} &= 0 \\ \Rightarrow 1 - \cos^2 x - 2\cos x + \frac{1}{4} &= 0 \\ \text{Putting } \cos x = t, \text{ we get} \\ 1 - t^2 - 2t + \frac{1}{4} &= 0 \Rightarrow 4t^2 + 8t - 5 = 0 \end{aligned}$$

$$\therefore t = \frac{1}{2} \text{ or } t = -\frac{5}{2}$$

$$\text{Since, } \cos x \neq \frac{-5}{2}$$

$$\therefore \cos x = \frac{1}{2} = \cos \frac{\pi}{3} \Rightarrow x = 2n\pi \pm \frac{\pi}{3}$$

$$\begin{aligned} 12. \quad \text{We have, } \sec \theta + \tan \theta &= \sqrt{3} \quad \dots(i) \\ \Rightarrow \sec \theta - \tan \theta &= \frac{1}{\sqrt{3}} \quad \dots(ii) \\ \dots[\because \sec^2 \theta - \tan^2 \theta &= 1] \end{aligned}$$

By solving (i) and (ii), we get

$$\tan \theta = \frac{1}{2} \left(\sqrt{3} - \frac{1}{\sqrt{3}} \right) = \frac{1}{\sqrt{3}}$$

$$\therefore \tan \theta = \tan\left(\frac{\pi}{6}\right)$$

$$\Rightarrow \theta = n\pi + \frac{\pi}{6}$$

$$\therefore \theta = \frac{\pi}{6} \text{ and } \frac{7\pi}{6} \text{ in } [0, 2\pi]$$

Hence, there are two solutions.

$$\begin{aligned} 13. \quad r \sin \theta &= 3, \\ r &= 4(1 + \sin \theta) \\ \text{Eliminating } r, \text{ we get} \\ \frac{3}{\sin \theta} &= 4 + 4 \sin \theta \end{aligned}$$

$$\therefore \sin \theta = \frac{1}{2}, -\frac{3}{2}$$

$$\therefore \sin \theta = \frac{1}{2} \quad \dots \left[\because \sin \theta \neq \frac{-3}{2} \right]$$

$$\Rightarrow \theta = \frac{\pi}{6}, \pi - \frac{\pi}{6} = \frac{5\pi}{6} \text{ in } [0, 2\pi]$$

$$14. \quad 2\sin^2 \theta - 3\sin \theta - 2 = 0$$

$$\therefore \sin \theta = \frac{3 \pm \sqrt{9+16}}{4} = \frac{3 \pm 5}{4} = 2, -\frac{1}{2}$$

$$\therefore \sin \theta = -\frac{1}{2} \quad \dots [\because |\sin \theta| \leq 1]$$

$$\therefore \sin \theta = \sin\left(\frac{-\pi}{6}\right)$$

$$\therefore \theta = n\pi + (-1)^n \left(\frac{-\pi}{6}\right) = n\pi + (-1)^{n+1} \left(\frac{\pi}{6}\right)$$

$$15. \quad 2\cos^2 x + 3\sin x - 3 = 0$$

$$\Rightarrow 2 - 2\sin^2 x + 3\sin x - 3 = 0$$

$$\Rightarrow (2\sin x - 1)(\sin x - 1) = 0$$

$$\Rightarrow \sin x = \frac{1}{2} \text{ or } \sin x = 1$$

$$\Rightarrow x = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{\pi}{2} \text{ i.e., } 30^\circ, 150^\circ, 90^\circ.$$

$$16. \quad 4\sin^2 \theta + 2(\sqrt{3}+1)\cos \theta = 4 + \sqrt{3}$$

$$\Rightarrow 4 - 4\cos^2 \theta + 2(\sqrt{3}+1)\cos \theta = 4 + \sqrt{3}$$

$$\Rightarrow 4\cos^2 \theta - 2(\sqrt{3}+1)\cos \theta + \sqrt{3} = 0$$

$$\Rightarrow \cos \theta = \frac{2(\sqrt{3}+1) \pm \sqrt{4(\sqrt{3}+1)^2 - 16\sqrt{3}}}{8}$$

$$\Rightarrow \cos \theta = \frac{\sqrt{3}}{2} \text{ or } \frac{1}{2}$$

$$\Rightarrow \theta = 2n\pi \pm \frac{\pi}{6} \text{ or } 2n\pi \pm \frac{\pi}{3}$$

$$17. \quad \sin(A+B) = 1 \text{ and } \cos(A-B) = \frac{\sqrt{3}}{2}$$


$$\Rightarrow A+B = \frac{\pi}{2} \text{ and } A-B = \frac{\pi}{6}$$

$$\Rightarrow A = \frac{\pi}{3}, B = \frac{\pi}{6}$$



18. $\cos 7\theta = \cos \theta - \sin 4\theta$
 $\Rightarrow \sin 4\theta = \cos \theta - \cos 7\theta$
 $\Rightarrow \sin 4\theta = 2 \sin (4\theta) \sin (3\theta)$
 $\Rightarrow \sin 4\theta = 0 \Rightarrow 4\theta = n\pi$ or
 $\sin 3\theta = \frac{1}{2} = \sin \left(\frac{\pi}{6}\right)$
 $\Rightarrow 3\theta = n\pi + (-1)^n \frac{\pi}{6}$
 $\therefore \theta = \frac{n\pi}{4}, \frac{n\pi}{3} + (-1)^n \frac{\pi}{18}$
19. $\frac{1 - \tan^2 \theta}{\sec^2 \theta} = \frac{1}{2} \Rightarrow \cos^2 \theta - \sin^2 \theta = \frac{1}{2}$
 $\Rightarrow \cos 2\theta = \frac{1}{2} = \cos \left(\frac{\pi}{3}\right)$
 $\Rightarrow 2\theta = 2n\pi \pm \frac{\pi}{3} \Rightarrow \theta = n\pi \pm \frac{\pi}{6}$
20. $\sqrt{3} \tan 2\theta + \sqrt{3} \tan 3\theta + \tan 2\theta \tan 3\theta = 1$
 $\Rightarrow \frac{\tan 2\theta + \tan 3\theta}{1 - \tan 2\theta \tan 3\theta} = \frac{1}{\sqrt{3}} \Rightarrow \tan 5\theta = \tan \frac{\pi}{6}$
 $\Rightarrow 5\theta = n\pi + \frac{\pi}{6} \Rightarrow \theta = \left(n + \frac{1}{6}\right) \frac{\pi}{5}$
21. $\tan \theta + \tan 2\theta = \tan 3\theta (\tan \theta \tan 2\theta - 1)$
 $\Rightarrow \frac{\tan \theta + \tan 2\theta}{1 - \tan \theta \tan 2\theta} = -\tan 3\theta$
 $\Rightarrow 2 \tan 3\theta = 0 \Rightarrow 3\theta = n\pi$
 $\Rightarrow \theta = \frac{n\pi}{3}$
22. $2 \tan^2 \theta = \sec^2 \theta \Rightarrow 2 \tan^2 \theta = \tan^2 \theta + 1$
 $\Rightarrow \tan^2 \theta = 1 = \tan^2 \left(\frac{\pi}{4}\right) \Rightarrow \theta = n\pi \pm \frac{\pi}{4}$
23. $\tan \theta \tan 2\theta = 1$
 $\therefore \tan \theta \frac{2 \tan \theta}{1 - \tan^2 \theta} = 1$
 $\therefore 2 \tan^2 \theta = 1 - \tan^2 \theta$
 $\therefore 3 \tan^2 \theta = 1$
 $\therefore \tan^2 \theta = \frac{1}{3} = \tan^2 \left(\frac{\pi}{6}\right)$
 $\therefore \theta = n\pi \pm \frac{\pi}{6}$
24. $\sin 3\alpha = 4 \sin \alpha \sin (x + \alpha) \sin (x - \alpha)$
 $\therefore \sin 3\alpha = 4 \sin \alpha (\sin^2 x \cos^2 \alpha - \cos^2 x \sin^2 \alpha)$
 $\therefore 3 \sin \alpha - 4 \sin^3 \alpha = 4 \sin \alpha (\sin^2 x - \sin^2 \alpha)$

- $\therefore \sin^2 x = \left(\frac{3}{4}\right) \Rightarrow \sin^2 x = \sin^2 \frac{\pi}{3}$
 $\therefore x = n\pi \pm \frac{\pi}{3}$
25. $(\cos \theta + \cos 7\theta) + (\cos 3\theta + \cos 5\theta) = 0$
 $\Rightarrow 2 \cos 4\theta \cos 3\theta + 2 \cos 4\theta \cos \theta = 0$
 $\Rightarrow 2 \cos 4\theta (\cos 3\theta + \cos \theta) = 0$
 $\Rightarrow 4 \cos 4\theta \cos 2\theta \cos \theta = 0$
 $\Rightarrow 4 \frac{\sin 2^3 \theta}{2^3 \sin \theta} = 0$
 $\therefore \cos A \cos 2A \cos 2^2 A \cos 2^3 A \dots \cos 2^{n-1} A$
 $\dots \left[\begin{aligned} &= \frac{\sin 2^n A}{2^n \sin A} \end{aligned} \right]$
 $\Rightarrow \sin 8\theta = 0$
 $\Rightarrow 8\theta = n\pi$
 $\Rightarrow \theta = \frac{n\pi}{8}$
26. Given, $\frac{\cos A}{a} = \frac{\cos B}{b} = \frac{\cos C}{c}$ (i)
 By sine rule,
 $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$ (ii)
 From (i) and (ii), we get
 $\therefore \frac{\cos A}{\sin A} = \frac{\sin A}{\sin B}$
 $\therefore \sin (A - B) = 0 \Rightarrow A = B$
 Similarly, we get,
 $B = C$
 $\therefore A = B = C$
 Thus, ΔABC is an equilateral triangle.
27. We know that,
 $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = k$
 $\Rightarrow \frac{2}{\frac{3}{2}} = \frac{3}{\sin B} = \frac{c}{\sin C} = k$
 $\Rightarrow k = 3$
 $\therefore \frac{3}{\sin B} = 3$
 $\Rightarrow \sin B = 1$
 $\Rightarrow B = 90^\circ$
 Hence, the triangle is a right angled triangle.
 From the figure,
 $\cos C = \frac{BC}{AC} = \frac{2}{3}$





28. Since the angles are in A.P., therefore $B = 60^\circ$
By sine rule,

$$\frac{b}{c} = \frac{\sin B}{\sin C} \Rightarrow \frac{\sqrt{3}}{\sqrt{2}} = \frac{\sqrt{3}}{2\sin C} \Rightarrow C = 45^\circ$$

$$\therefore A = 180^\circ - 60^\circ - 45^\circ = 75^\circ$$

29. $B = 60^\circ, C = 75^\circ$
 $\Rightarrow A = 180^\circ - 60^\circ - 75^\circ = 45^\circ$

By sine rule,

$$\frac{b}{\sin B} = \frac{a}{\sin A} \Rightarrow \frac{b}{\sin 60^\circ} = \frac{2}{\sin 45^\circ} \Rightarrow b = \sqrt{6}$$

30. Let the angles of the triangle be $2x, 3x$ and $7x$.

$$\therefore 2x + 3x + 7x = 180^\circ \Rightarrow 12x = 180^\circ \Rightarrow x = 15^\circ$$

\therefore the angles are $30^\circ, 45^\circ$ and 105°

$$\therefore a : b : c = \sin 30^\circ : \sin 45^\circ : \sin 105^\circ$$

$$= \frac{1}{2} : \frac{1}{\sqrt{2}} : \frac{\sqrt{3}+1}{2\sqrt{2}}$$

$$= \sqrt{2} : 2 : (\sqrt{3} + 1)$$

$$\begin{aligned} 31. \quad \frac{b-c}{a} &= \frac{\sin B - \sin C}{\sin A} \\ &= \frac{2\sin \frac{B-C}{2} \cos \frac{B+C}{2}}{2\sin \frac{A}{2} \cos \frac{A}{2}} \\ &= \frac{\sin\left(\frac{B-C}{2}\right) \cos\left(\frac{B+C}{2}\right)}{\cos\left(\frac{B+C}{2}\right) \cos \frac{A}{2}} \\ &= \frac{\sin \frac{B-C}{2}}{\cos \frac{A}{2}} \end{aligned}$$

$$\Rightarrow (b-c) \cos \frac{A}{2} = a \sin \frac{B-C}{2}$$

$$32. \quad \frac{1 + \cos C \cos(A-B)}{1 + \cos(A-C) \cos B} = \frac{1 - \cos(A+B) \cos(A-B)}{1 - \cos(A-C) \cos(A+C)}$$

$$= \frac{1 - \frac{1}{2}(\cos 2A + \cos 2B)}{1 - \frac{1}{2}(\cos 2A + \cos 2C)}$$

$$= \frac{1 - \frac{1}{2}(1 - 2\sin^2 A + 1 - 2\sin^2 B)}{1 - \frac{1}{2}(1 - 2\sin^2 A + 1 - 2\sin^2 C)}$$

$$= \frac{\sin^2 A + \sin^2 B}{\sin^2 A + \sin^2 C} = \frac{a^2 + b^2}{a^2 + c^2}$$

$$\begin{aligned} 33. \quad \frac{\cos \frac{B-C}{2}}{\sin \frac{A}{2}} &= \frac{\sin \frac{B+C}{2} \cos \frac{B-C}{2}}{\sin \frac{B+C}{2} \sin \frac{A}{2}} \\ &= \frac{2\sin \frac{B+C}{2} \cos \frac{B-C}{2}}{2\sin\left(\frac{\pi}{2} - \frac{A}{2}\right) \sin \frac{A}{2}} \\ &= \frac{\sin B + \sin C}{\sin A} = \frac{b+c}{a} \end{aligned}$$

$$34. \quad \cos \theta = \frac{36 + 100 - (14)^2}{2.6.10}$$

$$\Rightarrow \theta = 120^\circ \Rightarrow \text{Obtuse angled triangle}$$

35. Since A, B and C are in A.P., therefore

$$B = 60^\circ \dots \left[\begin{array}{l} \because A + B + C = 180^\circ \\ \Rightarrow A + C = 2B \Rightarrow B = 60^\circ \end{array} \right]$$

Since sides a, b and c are in G.P., therefore $b^2 = ac$

$$\cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

$$\Rightarrow \frac{1}{2} = \frac{a^2 + c^2 - b^2}{2b^2}, \quad \dots [\because b^2 = ac]$$

$$\Rightarrow b^2 = a^2 + c^2 - b^2$$

$$\Rightarrow a^2 + c^2 = 2b^2$$

$\Rightarrow a^2, b^2, c^2$ are in A.P.

36. A, B, C are in A. P. then angle $B = 60^\circ$,

$$\dots \left[\begin{array}{l} \because A + B + C = 180^\circ \\ \Rightarrow A + C = 2B \Rightarrow B = 60^\circ \end{array} \right]$$

$$\therefore \cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

$$\Rightarrow \frac{1}{2} = \frac{a^2 + c^2 - b^2}{2ac} \Rightarrow a^2 + c^2 - b^2 = ac$$

$$\Rightarrow b^2 = a^2 + c^2 - ac$$

$$\begin{aligned} 37. \quad \frac{\cos A}{a} + \frac{\cos B}{b} + \frac{\cos C}{c} &= \frac{b^2 + c^2 - a^2 + a^2 + c^2 - b^2 + a^2 + b^2 - c^2}{2abc} \\ &= \frac{a^2 + b^2 + c^2}{2abc} \end{aligned}$$

38. We have, $a : b : c = 1 : \sqrt{3} : 2$

$$\text{i.e. } a = \lambda, b = \sqrt{3}\lambda, c = 2\lambda$$

$$\cos A = \frac{3\lambda^2 + 4\lambda^2 - \lambda^2}{2(\sqrt{3}\lambda)(2\lambda)} = \frac{6\lambda^2}{4\sqrt{3}\lambda^2} = \frac{\sqrt{3}}{2}$$

$$\Rightarrow A = 30^\circ$$



Similarly, $\cos B = \frac{1}{2} \Rightarrow B = 60^\circ$,

$\cos C = 0 \Rightarrow C = 90^\circ$.

Hence, $A : B : C = 1 : 2 : 3$

$$\begin{aligned}
 39. \quad & (a^2 + b^2 - 2ab) \cos^2 \frac{C}{2} + (a^2 + b^2 + 2ab) \sin^2 \frac{C}{2} \\
 &= (a^2 + b^2) \left(\cos^2 \frac{C}{2} + \sin^2 \frac{C}{2} \right) \\
 &\quad - 2ab \left(\cos^2 \frac{C}{2} - \sin^2 \frac{C}{2} \right) \\
 &= a^2 + b^2 - 2ab \cos C \\
 &= a^2 + b^2 - (a^2 + b^2 - c^2) = c^2
 \end{aligned}$$

$$\begin{aligned}
 40. \quad & \cos A = \frac{\sin B}{2 \sin C} \Rightarrow \frac{b^2 + c^2 - a^2}{2bc} = \frac{b}{2c} \\
 &\Rightarrow b^2 + c^2 - a^2 - b^2 = 0 \Rightarrow c^2 = a^2 \\
 &\Rightarrow c = a \Rightarrow \text{Triangle is isosceles}
 \end{aligned}$$

$$\begin{aligned}
 41. \quad & a = \sin \theta, b = \cos \theta \text{ and } c = \sqrt{1 + \sin \theta \cos \theta} \\
 & \text{Since } \sqrt{1 + \sin \theta \cos \theta} \text{ is greater than } \sin \theta \text{ and } \cos \theta. \\
 & \therefore C \text{ is the greatest angle,} \\
 & \therefore \cos C = \frac{a^2 + b^2 - c^2}{2ab} \\
 &= \frac{\sin^2 \theta + \cos^2 \theta - (1 + \sin \theta \cos \theta)}{2 \sin \theta \cos \theta} \\
 &= -\frac{1}{2} = \cos 120^\circ \\
 & \therefore C = 120^\circ
 \end{aligned}$$

$$\begin{aligned}
 42. \quad & 1 - \tan \frac{A}{2} \tan \frac{B}{2} = \frac{\cos \frac{A}{2} \cos \frac{B}{2} - \sin \frac{A}{2} \sin \frac{B}{2}}{\cos \frac{A}{2} \cos \frac{B}{2}} \\
 &= \frac{\cos \left(\frac{A}{2} + \frac{B}{2} \right)}{\cos \frac{A}{2} \cos \frac{B}{2}} \\
 &= \frac{\sin \frac{C}{2}}{\cos \frac{A}{2} \cos \frac{B}{2}} \\
 &= \left[\frac{(s-a)(s-b)bc \cdot ac}{ab \cdot s(s-a)s(s-b)} \right]^{1/2} \\
 &= \frac{c}{s} = \frac{2c}{a+b+c}
 \end{aligned}$$

$$\begin{aligned}
 43. \quad & \frac{s(s-a)}{bc} - \frac{(s-b)(s-c)}{bc} \\
 &= \cos^2 \frac{A}{2} - \sin^2 \frac{A}{2} = \cos \frac{2A}{2} = \cos A
 \end{aligned}$$

$$\begin{aligned}
 44. \quad & a \cos^2 \frac{C}{2} + c \cos^2 \frac{A}{2} = \frac{3b}{2} \\
 \therefore & a \frac{s(s-c)}{ab} + c \frac{s(s-a)}{bc} = \frac{3b}{2} \\
 \therefore & 2s(s-c + s-a) = 3b^2 \\
 \therefore & 2s(b) = 3b^2 \Rightarrow 2s = 3b \Rightarrow a + b + c = 3b \\
 \therefore & a + c = 2b \Rightarrow a, b, c \text{ are in A.P.}
 \end{aligned}$$

$$\begin{aligned}
 45. \quad & \tan \frac{C}{2} = \sqrt{\frac{(s-a)(s-b)}{s(s-c)}} = 1 \\
 \Rightarrow & \tan \frac{C}{2} = \tan 45^\circ \Rightarrow \frac{C}{2} = 45^\circ \\
 \Rightarrow & C = 90^\circ
 \end{aligned}$$

$$\begin{aligned}
 46. \quad & \frac{\tan \frac{A}{2} - \tan \frac{B}{2}}{\tan \frac{A}{2} + \tan \frac{B}{2}} = \frac{\sqrt{\frac{(s-b)(s-c)}{s(s-a)}} - \sqrt{\frac{(s-a)(s-c)}{s(s-b)}}}{\sqrt{\frac{(s-b)(s-c)}{s(s-a)}} + \sqrt{\frac{(s-a)(s-c)}{s(s-b)}}} \\
 &= \frac{(s-b)\sqrt{s(s-c)} - (s-a)\sqrt{s(s-c)}}{(s-b)\sqrt{s(s-c)} + (s-a)\sqrt{s(s-c)}} \\
 &= \frac{\sqrt{s(s-c)}(s-b-s+a)}{\sqrt{s(s-c)}(s-b+s-a)} = \frac{a-b}{c}
 \end{aligned}$$

$$\begin{aligned}
 47. \quad & \frac{1}{\sin^2 \frac{A}{2}}, \frac{1}{\sin^2 \frac{B}{2}}, \frac{1}{\sin^2 \frac{C}{2}} \text{ are in A. P.} \\
 \Rightarrow & \frac{1}{\sin^2 \frac{C}{2}} - \frac{1}{\sin^2 \frac{B}{2}} = \frac{1}{\sin^2 \frac{B}{2}} - \frac{1}{\sin^2 \frac{A}{2}} \\
 \Rightarrow & \frac{ab}{(s-a)(s-b)} - \frac{ac}{(s-a)(s-c)} \\
 &= \frac{ac}{(s-a)(s-c)} - \frac{bc}{(s-b)(s-c)} \\
 \Rightarrow & \left(\frac{a}{s-a} \right) \left(\frac{b(s-c) - c(s-b)}{(s-b)(s-c)} \right) \\
 &= \left(\frac{c}{s-c} \right) \left(\frac{a(s-b) - b(s-a)}{(s-a)(s-b)} \right) \\
 \Rightarrow & abs - abc - acs + abc = acs - abc - bcs + abc \\
 \Rightarrow & ab - ac = ac - bc \Rightarrow ab + bc = 2ac \\
 \Rightarrow & \frac{1}{c} + \frac{1}{a} = \frac{2}{b} \Rightarrow a, b, c \text{ are in H. P.}
 \end{aligned}$$



$$48. \text{ Let } t = \tan\left(\frac{A-B}{2}\right)$$

$$\cos(A-B) = \frac{1-t^2}{1+t^2} \Rightarrow \frac{4}{5} = \frac{1-t^2}{1+t^2} \Rightarrow t = \frac{1}{3}$$

$$\text{So, } \tan\left(\frac{A-B}{2}\right) = \frac{1}{3}$$

$$\text{Then, } \tan\left(\frac{A-B}{2}\right) = \frac{a-b}{a+b} \cot \frac{C}{2}$$

$$\Rightarrow \frac{1}{3} = \frac{6-3}{6+3} \cot \frac{C}{2} \Rightarrow C = 90^\circ$$

$$\therefore \Delta = \frac{1}{2} (6)(3) \sin 90^\circ = 9 \text{ square units.}$$

49. Let the common multiple be x .

\therefore the sides are $(2x)$, $(\sqrt{6}x)$, $(\sqrt{3}+1)x$

$\therefore (\sqrt{3}+1)x$ is the largest side.

If θ is the angle opposite to side $(\sqrt{3}+1)x$, then

$$\cos \theta = \frac{(2x)^2 + (\sqrt{6}x)^2 - [(\sqrt{3}+1)x]^2}{2 \times (2x) \times (\sqrt{6}x)}$$

$$= \frac{3-\sqrt{3}}{2\sqrt{6}}$$

$$\therefore \cos \theta = \frac{\sqrt{3}-1}{2\sqrt{2}} \Rightarrow \theta = 75^\circ$$

50. We have,

$$\tan\left(\frac{A-B}{2}\right) = \sqrt{\frac{1-\cos(A-B)}{1+\cos(A-B)}} = \sqrt{\frac{1-\left(\frac{31}{32}\right)}{1+\left(\frac{31}{32}\right)}}$$

$$\Rightarrow \frac{a-b}{a+b} \cot \frac{C}{2} = \frac{1}{\sqrt{63}}$$

$$\Rightarrow \frac{1}{9} \cot \frac{C}{2} = \frac{1}{\sqrt{63}}$$

$$\Rightarrow \tan \frac{C}{2} = \frac{\sqrt{7}}{3}$$

$$\text{Now, } \cos C = \frac{1-\tan^2\left(\frac{C}{2}\right)}{1+\tan^2\left(\frac{C}{2}\right)}$$

$$\Rightarrow \cos C = \frac{1-\left(\frac{7}{9}\right)}{1+\left(\frac{7}{9}\right)} = \frac{1}{8}$$

$$\therefore c^2 = a^2 + b^2 - 2ab \cos C$$

$$\Rightarrow c^2 = 25 + 16 - 40 \times \frac{1}{8} = 36 \Rightarrow c = 6$$

51. Since $\sin^{-1} x$ cannot be greater than $\frac{\pi}{2}$.

$$\therefore \sin^{-1} x = \sin^{-1} y = \sin^{-1} z = \frac{\pi}{2}$$

Therefore, $x = y = z = 1$

Putting these values in the expression, we get

$$1 + 1 + 1 - \frac{9}{1+1+1} = 0$$

$$52. A = \tan^{-1}\left(\frac{2}{3}\right) \Rightarrow \tan A = \frac{2}{3}$$

$$B = \operatorname{cosec}^{-1}\left(\frac{5}{3}\right) \Rightarrow \tan B = \frac{3}{4}$$

$$\cot(A+B) = \frac{1-\tan A \tan B}{\tan A + \tan B}$$

$$= \frac{1-\frac{2}{3} \cdot \frac{3}{4}}{\frac{2}{3} + \frac{3}{4}} = \frac{\frac{6}{12}}{\frac{17}{12}} = \frac{6}{17}$$

$$53. \sin^2\left(2 \tan^{-1} \sqrt{\frac{1+x}{1-x}}\right)$$

$$= \sin^2(2\theta), \text{ where } \theta = \tan^{-1} \sqrt{\frac{1+x}{1-x}}$$

$$= \left(\frac{2 \tan \theta}{1 + \tan^2 \theta}\right)^2, \text{ where } \tan \theta = \sqrt{\frac{1+x}{1-x}}$$

$$= \left(\frac{2\sqrt{1+x}}{\sqrt{1-x}}\right)^2 = \frac{4(1+x)(1-x)}{(1-x+1+x)^2} = 1-x^2$$

$$54. \text{ The principal value of } \sin^{-1}\left[\sin\left(\frac{2\pi}{3}\right)\right]$$

$$= \sin^{-1}\left[\sin\left(\pi - \frac{2\pi}{3}\right)\right] = \sin^{-1}\left[\sin\left(\frac{\pi}{3}\right)\right] = \frac{\pi}{3}$$

$$55. \text{ Let } \sin^{-1} \frac{5}{13} = x \Rightarrow \sin x = \frac{5}{13}$$

$$\Rightarrow \cos x = \sqrt{1 - \frac{25}{169}} = \frac{12}{13}$$

$$\Rightarrow \cos\left(\sin^{-1} \frac{5}{13}\right) = \cos\left(\cos^{-1} \frac{12}{13}\right) = \frac{12}{13}$$



$$\begin{aligned}
 56. \quad \theta &= \sin^{-1}[\sin(-600^\circ)] \\
 &\Rightarrow \theta = \sin^{-1}[-(\sin 240^\circ)] \\
 &\Rightarrow \theta = \sin^{-1}[-\sin(180^\circ + 60^\circ)] \\
 &\Rightarrow \theta = \sin^{-1}(\sin 60^\circ) = \sin^{-1}\left[\sin\left(\frac{\pi}{3}\right)\right] = \frac{\pi}{3}
 \end{aligned}$$

$$\begin{aligned}
 57. \quad \tan^{-1}\left(\frac{\cos x - \sin x}{\cos x + \sin x}\right) &= \tan^{-1}\left(\frac{1 - \tan x}{1 + \tan x}\right) \\
 &= \tan^{-1}\left(\tan\left(\frac{\pi}{4} - x\right)\right) \\
 &= \frac{\pi}{4} - x
 \end{aligned}$$

$$\begin{aligned}
 58. \quad \tan^{-1}\left[\frac{\frac{a \cos x - b \sin x}{b \cos x}}{\frac{b \cos x + a \sin x}{b \cos x}}\right] &= \tan^{-1}\left[\frac{\frac{a}{b} - \tan x}{1 + \frac{a}{b} \tan x}\right] \\
 &= \tan^{-1} \frac{a}{b} - \tan^{-1}(\tan x) \\
 &= \tan^{-1} \frac{a}{b} - x
 \end{aligned}$$

$$\begin{aligned}
 59. \quad \tan^{-1}\left(\frac{\cos x}{1 + \sin x}\right) &= \tan^{-1}\left[\frac{\sin\left(\frac{\pi}{2} - x\right)}{1 + \cos\left(\frac{\pi}{2} - x\right)}\right] \\
 &= \tan^{-1}\left[\frac{2 \sin\left(\frac{\pi}{4} - \frac{x}{2}\right) \cos\left(\frac{\pi}{4} - \frac{x}{2}\right)}{2 \cos^2\left(\frac{\pi}{4} - \frac{x}{2}\right)}\right] \\
 &= \tan^{-1}\left[\tan\left(\frac{\pi}{4} - \frac{x}{2}\right)\right] = \frac{\pi}{4} - \frac{x}{2}
 \end{aligned}$$

$$\begin{aligned}
 60. \quad \tan^{-1}\left\{\frac{3a^2x - x^3}{a(a^2 - 3x^2)}\right\} &= \tan^{-1}\left\{\frac{3a^2x - x^3}{a^3 - 3ax^2}\right\} \\
 &= \tan^{-1}\left\{\frac{3\left(\frac{x}{a}\right) - \left(\frac{x}{a}\right)^3}{1 - 3\left(\frac{x}{a}\right)^2}\right\}
 \end{aligned}$$

$$\text{Put } \frac{x}{a} = \tan \theta$$

\therefore The given expression becomes

$$\begin{aligned}
 \tan^{-1}\left(\frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta}\right) &= \tan^{-1}(\tan 3\theta) \\
 &= 3\theta = 3 \tan^{-1} \frac{x}{a}
 \end{aligned}$$

$$\begin{aligned}
 61. \quad 3 \sin^{-1} \frac{2x}{1+x^2} - 4 \cos^{-1} \frac{1-x^2}{1+x^2} \\
 + 2 \tan^{-1} \frac{2x}{1-x^2} = \frac{\pi}{3}
 \end{aligned}$$

Putting $x = \tan \theta$, we get

$$\begin{aligned}
 3 \sin^{-1}\left(\frac{2 \tan \theta}{1 + \tan^2 \theta}\right) - 4 \cos^{-1}\left(\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}\right) \\
 + 2 \tan^{-1}\left(\frac{2 \tan \theta}{1 - \tan^2 \theta}\right) = \frac{\pi}{3} \\
 \Rightarrow 3 \sin^{-1}(\sin 2\theta) - 4 \cos^{-1}(\cos 2\theta) \\
 + 2 \tan^{-1}(\tan 2\theta) = \frac{\pi}{3}
 \end{aligned}$$

$$\Rightarrow 3(2\theta) - 4(2\theta) + 2(2\theta) = \frac{\pi}{3}$$

$$\Rightarrow 6\theta - 8\theta + 4\theta = \frac{\pi}{3}$$

$$\Rightarrow \theta = \frac{\pi}{6} \Rightarrow \tan^{-1} x = \frac{\pi}{6}$$

$$\Rightarrow x = \tan \frac{\pi}{6} = \frac{1}{\sqrt{3}}$$

$$\begin{aligned}
 62. \quad \tan^{-1}\left(\frac{\sqrt{1+x^2}-1}{x}\right) \\
 = \tan^{-1}\left[\frac{\sqrt{1+\tan^2\theta}-1}{\tan\theta}\right] \\
 (\text{Putting } x = \tan \theta) \\
 = \tan^{-1}\left[\frac{\sec\theta-1}{\tan\theta}\right] = \tan^{-1}\left[\frac{1-\cos\theta}{\sin\theta}\right] \\
 = \tan^{-1}\left[\frac{2\sin^2\frac{\theta}{2}}{2\sin\frac{\theta}{2}\cos\frac{\theta}{2}}\right] \\
 = \tan^{-1}\left(\tan\frac{\theta}{2}\right) = \frac{\theta}{2} = \frac{1}{2} \tan^{-1} x
 \end{aligned}$$

$$\begin{aligned}
 63. \quad \text{Let } x = \sin \theta \text{ and } \sqrt{x} = \sin \phi \\
 \text{Hence}
 \end{aligned}$$

$$\begin{aligned}
 \therefore \quad \sin^{-1}(x\sqrt{1-x} - \sqrt{x}\sqrt{1-x^2}) \\
 = \sin^{-1}(\sin\theta\sqrt{1-\sin^2\phi} - \sin\phi\sqrt{1-\sin^2\theta}) \\
 = \sin^{-1}(\sin\theta\cos\phi - \sin\phi\cos\theta) \\
 = \sin^{-1}\sin(\theta - \phi) \\
 = \theta - \phi = \sin^{-1}(x) - \sin^{-1}(\sqrt{x})
 \end{aligned}$$



$$64. \quad \cos^{-1}\left(\frac{1}{x}\right) = \theta \Rightarrow \sec^{-1}x = \theta$$

$$\therefore x = \sec \theta$$

$$\begin{aligned} \therefore \tan \theta &= \sqrt{\sec^2 \theta - 1} \\ &= \sqrt{x^2 - 1} \end{aligned}$$

$$65. \quad \sin^{-1} \frac{1}{5} + \cos^{-1} x = \frac{\pi}{2}$$

$$\therefore \sin^{-1} \frac{1}{5} = \frac{\pi}{2} - \cos^{-1} x = \sin^{-1} x$$

$$\therefore x = \frac{1}{5}$$

$$\begin{aligned} 66. \quad \sin^{-1} x + \sin^{-1} \frac{1}{x} + \cos^{-1} x + \cos^{-1} \frac{1}{x} \\ &= \{\sin^{-1}(x) + \cos^{-1}(x)\} + \left\{\sin^{-1}\left(\frac{1}{x}\right) + \cos^{-1}\left(\frac{1}{x}\right)\right\} \\ &= \frac{\pi}{2} + \frac{\pi}{2} = \pi \end{aligned}$$

$$67. \quad \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$$

$$\Rightarrow \cos^{-1} x = \frac{\pi}{2} - \sin^{-1} x = \frac{\pi}{2} - \frac{\pi}{5} = \frac{3\pi}{10}$$

$$68. \quad \theta = \sin^{-1} x + \cos^{-1} x \quad \tan^{-1} x = \frac{\pi}{2} - \tan^{-1} x$$

$$\text{Since, } -\frac{\pi}{2} < \tan^{-1} x < \frac{\pi}{2}$$

$$\Rightarrow \frac{\pi}{2} > -\tan^{-1} x > -\frac{\pi}{2}$$

$$\Rightarrow 0 < \frac{\pi}{2} - \tan^{-1} x < \pi$$

$$69. \quad (\tan^{-1} x)^2 + (\cot^{-1} x)^2 = \frac{5\pi^2}{8}$$

$$\Rightarrow (\tan^{-1} x + \cot^{-1} x)^2$$

$$- 2\tan^{-1} x \left(\frac{\pi}{2} - \tan^{-1} x\right) = \frac{5\pi^2}{8}$$

$$\Rightarrow \frac{\pi^2}{4} - 2 \times \frac{\pi}{2} \tan^{-1} x + 2(\tan^{-1} x)^2 = \frac{5\pi^2}{8}$$

$$\Rightarrow 2(\tan^{-1} x)^2 - \pi \tan^{-1} x - \frac{3\pi^2}{8} = 0$$

$$\Rightarrow \tan^{-1} x = -\frac{\pi}{4}, \frac{3\pi}{4}$$

$$\Rightarrow \tan^{-1} x = -\frac{\pi}{4} \Rightarrow x = -1$$

$$\begin{aligned} 70. \quad \cos \left[\tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{2} \right] &= \cos \left[\tan^{-1} \left(\frac{\frac{1}{3} + \frac{1}{2}}{1 - \frac{1}{3} \times \frac{1}{2}} \right) \right] \\ &= \cos \{ \tan^{-1}(1) \} \\ &= \cos \frac{\pi}{4} = \frac{1}{\sqrt{2}} \end{aligned}$$

$$71. \quad \text{Let } \alpha = \cos^{-1} \left(\frac{4}{5} \right)$$

$$\therefore \cos \alpha = \left(\frac{4}{5} \right) \Rightarrow \tan \alpha = \left(\frac{3}{4} \right)$$

$$\therefore \alpha = \tan^{-1} \left(\frac{3}{4} \right)$$

$$\therefore \cos^{-1} \frac{4}{5} + \tan^{-1} \frac{3}{5} = \tan^{-1} \frac{3}{4} + \tan^{-1} \frac{3}{5}$$

$$= \tan^{-1} \left(\frac{\frac{3}{4} + \frac{3}{5}}{1 - \frac{3}{4} \times \frac{3}{5}} \right) = \tan^{-1} \left(\frac{27}{11} \right)$$

$$72. \quad \tan^{-1} \frac{x-1}{x+2} + \tan^{-1} \frac{x+1}{x+2} = \frac{\pi}{4}$$

$$\therefore \tan^{-1} \left[\frac{\frac{x-1}{x+2} + \frac{x+1}{x+2}}{1 - \left(\frac{x-1}{x+2} \right) \left(\frac{x+1}{x+2} \right)} \right] = \frac{\pi}{4}$$

$$\therefore \left[\frac{2x(x+2)}{x^2 + 4 + 4x - x^2 + 1} \right] = \tan \frac{\pi}{4}$$

$$\therefore \frac{2x(x+2)}{4x+5} = 1$$

$$\therefore 2x^2 + 4x = 4x + 5 \Rightarrow x = \pm \sqrt{\frac{5}{2}}$$

$$73. \quad \tan^{-1} \left[\frac{1}{\sqrt{\cos \alpha}} \right] - \tan^{-1} \left[\sqrt{\cos \alpha} \right] = x$$

$$\Rightarrow \tan^{-1} \left[\frac{1 - \sqrt{\cos \alpha}}{\sqrt{\cos \alpha} + \sqrt{\cos \alpha}} \right] = x$$

$$\Rightarrow \tan x = \frac{1 - \cos \alpha}{2\sqrt{\cos \alpha}}$$

$$\therefore \sin x = \frac{1 - \cos \alpha}{1 + \cos \alpha} = \frac{2 \sin^2 \frac{\alpha}{2}}{2 \cos^2 \frac{\alpha}{2}} = \tan^2 \left(\frac{\alpha}{2} \right)$$



$$74. \tan^{-1} \frac{a+x}{a} + \tan^{-1} \frac{a-x}{a} = \frac{\pi}{6}$$

$$\therefore \tan^{-1} \left(\frac{\frac{a+x}{a} + \frac{a-x}{a}}{1 - \frac{a+x}{a} \cdot \frac{a-x}{a}} \right) = \frac{\pi}{6}$$

$$\therefore \frac{2a^2}{x^2} = \tan \frac{\pi}{6} = \frac{1}{\sqrt{3}} \Rightarrow x^2 = 2\sqrt{3} a^2$$

$$75. \tan^{-1} \left(\frac{a}{b+c} \right) + \tan^{-1} \left(\frac{b}{c+a} \right)$$

$$= \tan^{-1} \left(\frac{ac+bc+a^2+b^2}{ac+bc+c^2} \right)$$

$$\dots \left[\because \tan^{-1} x + \tan^{-1} y = \tan^{-1} \left(\frac{x+y}{1-xy} \right) \right]$$

$$= \tan^{-1} (1) \quad \dots [\because c^2 = a^2 + b^2]$$

$$= \frac{\pi}{4}$$

$$76. \tan^{-1} \frac{3}{4} + \tan^{-1} \frac{3}{5} - \tan^{-1} \frac{8}{19}$$

$$= \tan^{-1} \left[\frac{\frac{3}{4} + \frac{3}{5}}{1 - \frac{3}{4} \times \frac{3}{5}} \right] - \tan^{-1} \frac{8}{19}$$

$$= \tan^{-1} \frac{27}{11} - \tan^{-1} \frac{8}{19} = \tan^{-1} \left[\frac{\frac{27}{11} - \frac{8}{19}}{1 + \frac{27}{11} \times \frac{8}{19}} \right]$$

$$= \tan^{-1} \left(\frac{425}{425} \right) = \tan^{-1} (1) = \frac{\pi}{4}$$

$$77. \tan^{-1} \left(\frac{xy}{zr} \right) + \tan^{-1} \left(\frac{yz}{xr} \right) + \tan^{-1} \left(\frac{xz}{yr} \right)$$

$$= \tan^{-1} \left[\frac{\frac{xy}{zr} + \frac{yz}{xr} + \frac{xz}{yr} - \frac{xyz}{r^3}}{1 - \left(\frac{x^2 + y^2 + z^2}{r^2} \right)} \right] = \tan^{-1} (\infty) = \frac{\pi}{2}$$

$$78. \tan^{-1} \left(\frac{c_1 x - y}{c_1 y + x} \right) + \tan^{-1} \left(\frac{c_2 - c_1}{1 + c_2 c_1} \right)$$

$$+ \tan^{-1} \left(\frac{c_3 - c_2}{1 + c_3 c_2} \right) + \dots + \tan^{-1} \frac{1}{c_n}$$

$$= \tan^{-1} \left(\frac{\frac{x}{y} - \frac{1}{c_1}}{1 + \frac{x}{y} \cdot \frac{1}{c_1}} \right) + \tan^{-1} \left(\frac{\frac{1}{c_1} - \frac{1}{c_2}}{1 + \frac{1}{c_1 c_2}} \right)$$

$$+ \tan^{-1} \left(\frac{\frac{1}{c_2} - \frac{1}{c_3}}{1 + \frac{1}{c_2 c_3}} \right) + \dots + \tan^{-1} \frac{1}{c_n}$$

$$= \tan^{-1} \left(\frac{x}{y} \right) - \tan^{-1} \left(\frac{1}{c_1} \right) + \tan^{-1} \left(\frac{1}{c_1} \right) - \tan^{-1} \left(\frac{1}{c_2} \right)$$

$$+ \tan^{-1} \left(\frac{1}{c_2} \right) - \tan^{-1} \left(\frac{1}{c_3} \right) + \dots + \tan^{-1} \left(\frac{1}{c_{n-1}} \right)$$

$$- \tan^{-1} \left(\frac{1}{c_n} \right) + \tan^{-1} \left(\frac{1}{c_n} \right)$$

$$= \tan^{-1} \left(\frac{x}{y} \right)$$

$$79. \tan^{-1} \left(\frac{d}{1 + a_1 a_2} \right) + \tan^{-1} \left(\frac{d}{1 + a_2 a_3} \right)$$

$$+ \dots + \tan^{-1} \left(\frac{d}{1 + a_{n-1} a_n} \right)$$

$$= \tan^{-1} \left(\frac{a_2 - a_1}{1 + a_1 a_2} \right) + \tan^{-1} \left(\frac{a_3 - a_2}{1 + a_2 a_3} \right)$$

$$+ \dots + \tan^{-1} \left(\frac{a_n - a_{n-1}}{1 + a_{n-1} a_n} \right)$$

$$= (\tan^{-1} a_2 - \tan^{-1} a_1) + (\tan^{-1} a_3 - \tan^{-1} a_2)$$

$$+ \dots + (\tan^{-1} a_n - \tan^{-1} a_{n-1})$$

$$= \tan^{-1} a_n - \tan^{-1} a_1 = \tan^{-1} \left(\frac{a_n - a_1}{1 + a_n a_1} \right)$$

$$= \tan^{-1} \left(\frac{(n-1)d}{1 + a_1 a_n} \right)$$

$$80. \tan \left[2 \tan^{-1} \left(\frac{1}{5} \right) - \frac{\pi}{4} \right]$$

$$= \tan \left[\tan^{-1} \frac{\frac{2}{5}}{1 - \frac{1}{25}} - \tan^{-1} (1) \right]$$

$$= \tan \left[\tan^{-1} \frac{5}{12} - \tan^{-1} (1) \right]$$

$$= \tan \left[\tan^{-1} \left(\frac{\frac{5}{12} - 1}{1 + \frac{5}{12}} \right) \right] = -\frac{7}{17}$$



$$\begin{aligned}
 81. \quad \sin \left[3 \sin^{-1} \left(\frac{1}{5} \right) \right] &= \sin \left[\sin^{-1} \left\{ 3 \left(\frac{1}{5} \right) - 4 \left(\frac{1}{5} \right)^3 \right\} \right] \\
 &= \sin \left[\sin^{-1} \left\{ \frac{3}{5} - \frac{4}{125} \right\} \right] = \sin \left[\sin^{-1} \left(\frac{75-4}{125} \right) \right] \\
 &= \sin \left[\sin^{-1} \frac{71}{125} \right] = \frac{71}{125}
 \end{aligned}$$

$$\begin{aligned}
 82. \quad \cot \left[\cos^{-1} \left(\frac{7}{25} \right) \right] &= \cot \left[\cot^{-1} \left(\frac{7}{24} \right) \right] = \frac{7}{24} \\
 &\dots \left[\because \cos^{-1} x = \cot^{-1} \frac{x}{\sqrt{1-x^2}} \right]
 \end{aligned}$$

$$\begin{aligned}
 83. \quad \text{Let } \sin^{-1} x = \theta &\Rightarrow x = \sin \theta \\
 \cos(2 \sin^{-1} x) &= \frac{1}{9} \quad \Rightarrow \cos 2\theta = \frac{1}{9} \\
 \Rightarrow 1 - 2 \sin^2 \theta &= \frac{1}{9} \quad \Rightarrow 1 - 2x^2 = \frac{1}{9} \\
 \Rightarrow 2x^2 = 1 - \frac{1}{9} &= \frac{8}{9} \quad \Rightarrow x^2 = \frac{4}{9} \\
 \Rightarrow x &= \pm \frac{2}{3}
 \end{aligned}$$

$$\begin{aligned}
 84. \quad \sin \left[2 \tan^{-1} \left(\frac{1}{3} \right) \right] + \cos [\tan^{-1} (2\sqrt{2})] \\
 &= \sin \left[\tan^{-1} \frac{2/3}{1-1/9} \right] + \cos [\tan^{-1} (2\sqrt{2})] \\
 &= \sin \left[\tan^{-1} \frac{3}{4} \right] + \cos [\tan^{-1} 2\sqrt{2}] \\
 &= \sin \left[\sin^{-1} \frac{\left(\frac{3}{4} \right)}{\sqrt{1+\left(\frac{3}{4} \right)^2}} \right] + \cos \left[\cos^{-1} \frac{1}{\sqrt{1+(2\sqrt{2})^2}} \right] \\
 &= \frac{3}{5} + \frac{1}{3} = \frac{14}{15}
 \end{aligned}$$

$$\begin{aligned}
 85. \quad \text{Given, } \tan^{-1} x &= \sin^{-1} \left[\frac{3}{\sqrt{10}} \right] \\
 \Rightarrow x &= \tan \left\{ \sin^{-1} \left[\frac{3}{\sqrt{10}} \right] \right\} = \tan \{ \tan^{-1} 3 \} \\
 \Rightarrow x &= 3
 \end{aligned}$$

$$\begin{aligned}
 86. \quad \tan \left(\cos^{-1} \frac{1}{5\sqrt{2}} - \sin^{-1} \frac{4}{\sqrt{17}} \right) \\
 &= \tan (\tan^{-1} 7 - \tan^{-1} 4) \\
 &= \tan \left[\tan^{-1} \left(\frac{7-4}{1+28} \right) \right] = \frac{3}{29}
 \end{aligned}$$

$$\begin{aligned}
 87. \quad \sin^{-1} \frac{1}{\sqrt{5}} + \cot^{-1} 3 &= \cot^{-1} \left(\frac{\sqrt{1-\frac{1}{5}}}{\frac{1}{\sqrt{5}}} \right) + \cot^{-1} 3 \\
 &= \cot^{-1}(2) + \cot^{-1}(3) \\
 &= \cot^{-1} \left(\frac{2 \times 3 - 1}{3 + 2} \right) \\
 &= \cot^{-1}(1) = \frac{\pi}{4}
 \end{aligned}$$

$$\begin{aligned}
 88. \quad \text{On expanding determinant,} \\
 \cos^2(A+B) + \sin^2(A+B) + \cos 2B &= 0 \\
 \therefore 1 + \cos 2B &= 0 \Rightarrow \cos 2B = \cos \pi \\
 \Rightarrow 2B &= 2n\pi + \pi \Rightarrow B = (2n+1) \frac{\pi}{2}, n \in \mathbb{Z}.
 \end{aligned}$$

**Competitive Thinking**

- $\tan^2 x = 1$
 $\Rightarrow \tan^2 x = \tan^2 \frac{\pi}{4} \Rightarrow x = n\pi \pm \frac{\pi}{4}$
- No solution as $|\sin x| \leq 1$, $|\cos x| \leq 1$ and both of them do not attain their maximum value for the same angle.
- $\cot \theta + \tan \theta = 2$
 $\therefore \frac{1}{\tan \theta} + \tan \theta = 2 \Rightarrow 1 + \tan^2 \theta = 2 \tan \theta$
 $\therefore \frac{2 \tan \theta}{1 + \tan^2 \theta} = 1 \Rightarrow \sin 2\theta = 1$
 $\Rightarrow 2\theta = n\pi + (-1)^n \frac{\pi}{2}$
 $\Rightarrow \theta = \frac{n\pi}{2} + (-1)^n \frac{\pi}{4}$
- $\tan 2\theta = 1$
 The value of $\tan \theta$ is positive if θ is in 1st and 3rd quadrant.
 \therefore Option (B) is the correct answer.
- The given equation is defined for $x \neq \frac{\pi}{2}, \frac{3\pi}{2}$.
 Now, $\sec x \cos 5x + 1 = 0$
 $\Rightarrow \sec x \cos 5x = -1$
 $\Rightarrow \cos 5x = -\cos x$
 $\Rightarrow \cos 5x + \cos x = 0$
 $\Rightarrow 2 \cos 3x \cdot \cos 2x = 0$
 $\Rightarrow \cos 3x = 0$ or $\cos 2x = 0$
 $\Rightarrow 3x = (2n+1) \frac{\pi}{2}$ or $2x = (2n+1) \frac{\pi}{2}$



- $\Rightarrow x = \frac{(2n+1)\pi}{6}$ or $x = \frac{(2n+1)\pi}{4}$
 $\Rightarrow x = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{11\pi}{6}, \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$ in $[0, 2\pi]$
 \therefore number of solutions = 8
6. $\cos\theta = \frac{-1}{2}$ and $0^\circ < \theta < 360^\circ$
 $\therefore \cos\theta = -\cos 60^\circ$
 $\therefore \cos\theta = \cos(180^\circ - 60^\circ)$ and $\cos\theta = \cos(180^\circ + 60^\circ)$
 $\Rightarrow \cos\theta = \cos 120^\circ$ and $\cos\theta = \cos 240^\circ$
 $\Rightarrow \theta = 120^\circ$ and 240°
7. $\cos\theta + \sqrt{3}\sin\theta = 2$
 $\Rightarrow \frac{1}{2}\cos\theta + \frac{\sqrt{3}}{2}\sin\theta = 1$
 $\Rightarrow \sin\left(\theta + \frac{\pi}{6}\right) = 1 = \sin\left(\frac{\pi}{2}\right) \Rightarrow \theta = \frac{\pi}{2} - \frac{\pi}{6} = \frac{\pi}{3}$
8. $\operatorname{cosec}\theta + 2 = 0$
 $\Rightarrow \sin\theta = -\frac{1}{2}$
 $\Rightarrow \sin\theta = -\sin 30^\circ$
 $\Rightarrow \sin\theta = \sin(180^\circ + 30^\circ)$ and $\sin\theta = \sin(360^\circ - 30^\circ)$
 $\Rightarrow \sin\theta = \sin 210^\circ$ and $\sin\theta = \sin 330^\circ$
 $\Rightarrow \theta = 210^\circ$ and $\theta = 330^\circ$
9. $\sin x + \sin y + \sin z = -3$ is satisfied only when $x = y = z = \frac{3\pi}{2}$, for $x, y, z \in [0, 2\pi]$.
 \therefore option (A) is the correct answer.
10. The given equation is defined for $x \neq \frac{\pi}{2}, \frac{3\pi}{2}$.
 Now, $\tan x + \sec x = 2 \cos x$
 $\Rightarrow \frac{\sin x}{\cos x} + \frac{1}{\cos x} = 2 \cos x$
 $\Rightarrow (\sin x + 1) = 2 \cos^2 x$
 $\Rightarrow (\sin x + 1) = 2(1 - \sin^2 x)$
 $\Rightarrow (\sin x + 1) = 2(1 - \sin x)(1 + \sin x)$
 $\Rightarrow (1 + \sin x)[2(1 - \sin x) - 1] = 0$
 $\Rightarrow 2(1 - \sin x) - 1 = 0$
 $\therefore \left[\begin{array}{l} \because \sin x \neq -1 \text{ otherwise } \cos x = 0 \text{ and} \\ \tan x, \sec x \text{ will be undefined} \end{array} \right]$
 $\Rightarrow \sin x = \frac{1}{2}$
 $\Rightarrow x = \frac{\pi}{6}, \frac{5\pi}{6}$ in $(0, 2\pi)$
 \therefore number of solutions = 2

11. $\tan(\pi \cos \theta) = \cot(\pi \sin \theta)$
 $\Rightarrow \tan(\pi \cos \theta) = \tan\left(\frac{\pi}{2} - \pi \sin \theta\right)$
 $\Rightarrow \pi \cos \theta = \frac{\pi}{2} - \pi \sin \theta$
 $\Rightarrow \sin \theta + \cos \theta = \frac{1}{2}$
 $\Rightarrow \frac{1}{\sqrt{2}} \sin \theta + \frac{1}{\sqrt{2}} \cos \theta = \frac{1}{2\sqrt{2}}$
 $\Rightarrow \cos \theta \cos \frac{\pi}{4} + \sin \theta \sin \frac{\pi}{4} = \frac{1}{2\sqrt{2}}$
 $\Rightarrow \cos\left(\theta - \frac{\pi}{4}\right) = \frac{1}{2\sqrt{2}}$
12. $\cos^2 \theta + \sin \theta + 1 = 0$
 $\Rightarrow 1 - \sin^2 \theta + \sin \theta + 1 = 0$
 $\Rightarrow \sin^2 \theta - \sin \theta - 2 = 0$
 $\Rightarrow (\sin \theta + 1)(\sin \theta - 2) = 0$
 $\Rightarrow \sin \theta = 2$, which is not possible and $\sin \theta = -1 = \sin \frac{3\pi}{2}$
 Therefore, solution of the given equation lies in the interval $\left(\frac{5\pi}{4}, \frac{7\pi}{4}\right)$.
13. $2 \sin^2 \theta = 4 + 3 \cos \theta$
 $\Rightarrow 2 - 2 \cos^2 \theta = 4 + 3 \cos \theta$
 $\Rightarrow 2 \cos^2 \theta + 3 \cos \theta + 2 = 0$
 $\Rightarrow \cos \theta = \frac{-3 \pm \sqrt{9 - 16}}{4}$,
 which are imaginary, hence no solution.
14. $\cos 2x + k \sin x = 2k - 7$
 $\Rightarrow 1 - 2 \sin^2 x + k \sin x - 2k + 7 = 0$
 $\Rightarrow 2 \sin^2 x - k \sin x + 2k - 8 = 0$
 $\Rightarrow \sin x = \frac{k \pm \sqrt{k^2 - 8(2k - 8)}}{4}$
 $\Rightarrow \sin x = \frac{k \pm (k - 8)}{4} = \frac{k - 4}{2}, 2$
 Since, $\sin x \neq 2$ and $-1 < \sin x < 1$
 $\therefore -1 < \frac{k - 4}{2} < 1 \Rightarrow -2 < k - 4 < 2$
 $\Rightarrow 2 < k < 6$
15. $(1 + \tan \theta)(1 + \tan \phi) = 2 \Rightarrow \frac{\tan \theta + \tan \phi}{1 - \tan \theta \tan \phi} = 1$
 $\Rightarrow \tan(\theta + \phi) = 1 \Rightarrow \theta + \phi = \frac{\pi}{4} = 45^\circ$



$$\begin{aligned}
 16. \quad & (1 + \tan \alpha)(1 + \tan 4\alpha) = 2 \\
 \therefore & 1 + \tan \alpha + \tan 4\alpha + \tan \alpha \cdot \tan 4\alpha = 2 \\
 \therefore & \tan \alpha + \tan 4\alpha = 1 - \tan \alpha \cdot \tan 4\alpha \\
 \therefore & \frac{\tan \alpha + \tan 4\alpha}{1 - \tan \alpha \cdot \tan 4\alpha} = 1 \\
 \therefore & \tan(\alpha + 4\alpha) = 1 \\
 \therefore & \tan 5\alpha = 1 \\
 \therefore & 5\alpha = \frac{\pi}{4} \quad \dots \left[\because \alpha \in \left(0, \frac{\pi}{16}\right) \right] \\
 \therefore & \alpha = \frac{\pi}{20}
 \end{aligned}$$

$$\begin{aligned}
 17. \quad & \cos x + \cos y = \frac{3}{2} \\
 \Rightarrow & 2 \cos \left(\frac{x+y}{2} \right) \cos \left(\frac{x-y}{2} \right) = \frac{3}{2} \\
 \Rightarrow & 2 \cos \frac{\pi}{3} \cos \left(\frac{x-y}{2} \right) = \frac{3}{2} \\
 & \dots \left[\because x+y = \frac{2\pi}{3} \text{ (given)} \right] \\
 \Rightarrow & \cos \left(\frac{x-y}{2} \right) = \frac{3}{2}, \text{ which is not possible} \\
 & \dots \left[\because \frac{3}{2} > 1 \right]
 \end{aligned}$$

Hence, the system of equations has no solution.

$$18. \quad 81^{\sin^2 x} + 81^{\cos^2 x} = 30 \quad \dots (i)$$

Check by options, put $x = \frac{\pi}{6}$ in (i),

$$81^{\sin^2 \frac{\pi}{6}} + 81^{\cos^2 \frac{\pi}{6}} = 30$$

$$\Rightarrow (81)^{\frac{1}{4}} + (81)^{\frac{3}{4}} = 30 \Rightarrow 30 = 30$$

\therefore option (A) is the correct answer.

$$\begin{aligned}
 19. \quad & 4 \sin^4 x + \cos^4 x = 1 \\
 \Rightarrow & 4 \sin^4 x = 1 - \cos^4 x \\
 \Rightarrow & 4 \sin^4 x = (1 - \cos^2 x)(1 + \cos^2 x) \\
 \Rightarrow & 4 \sin^4 x - (\sin^2 x)(1 + 1 - \sin^2 x) = 0 \\
 \Rightarrow & \sin^2 x [4 \sin^2 x - 2 + \sin^2 x] = 0 \\
 \Rightarrow & \sin^2 x (5 \sin^2 x - 2) = 0
 \end{aligned}$$

$$\Rightarrow \sin x = 0 \text{ or } \sin x = \pm \sqrt{\frac{2}{5}}$$

Hence $x = n\pi$ is the required answer.

$$\begin{aligned}
 20. \quad & 1 - \cos \theta = \sin \theta \cdot \sin \frac{\theta}{2} \\
 \Rightarrow & 2 \sin^2 \frac{\theta}{2} = 2 \sin \frac{\theta}{2} \cdot \cos \frac{\theta}{2} \cdot \sin \frac{\theta}{2}
 \end{aligned}$$

$$\Rightarrow 2 \sin^2 \frac{\theta}{2} - 2 \sin^2 \frac{\theta}{2} \cos \frac{\theta}{2} = 0$$

$$\Rightarrow 2 \sin^2 \frac{\theta}{2} \left(1 - \cos \frac{\theta}{2} \right) = 0$$

$$\Rightarrow 2 \sin^2 \frac{\theta}{2} \left(2 \sin^2 \frac{\theta}{4} \right) = 0$$

$$\Rightarrow 2 \sin^2 \frac{\theta}{2} = 0 \text{ or } 2 \sin^2 \frac{\theta}{4} = 0$$

$$\Rightarrow \sin \frac{\theta}{2} = 0 \text{ or } \sin \frac{\theta}{4} = 0$$

$$\Rightarrow \frac{\theta}{2} = k\pi \text{ or } \frac{\theta}{4} = k\pi$$

$$\Rightarrow \theta = 2k\pi \text{ or } \theta = 4k\pi, k \in I$$

\therefore option (B) is the correct answer.

$$21. \quad \sin 5x = \cos 2x$$

$$\Rightarrow \sin 5x = \sin \left(\frac{\pi}{2} - 2x \right)$$

$$\Rightarrow 5x = n\pi + (-1)^n \left(\frac{\pi}{2} - 2x \right)$$

$$\Rightarrow 5x + (-1)^n 2x = [2n + (-1)^n] \frac{\pi}{2}$$

$$\Rightarrow x [5 + 2(-1)^n] = [2n + (-1)^n] \frac{\pi}{2}$$

$$\Rightarrow x = \left[\frac{2n + (-1)^n}{5 + 2(-1)^n} \right] \frac{\pi}{2}$$

$$22. \quad \tan 5\theta = \cot 2\theta$$

$$\Rightarrow \tan 5\theta = \tan \left(\frac{\pi}{2} - 2\theta \right)$$

$$\Rightarrow 5\theta = n\pi + \frac{\pi}{2} - 2\theta$$

$$\Rightarrow 7\theta = n\pi + \frac{\pi}{2}$$

$$\Rightarrow \theta = \frac{n\pi}{7} + \frac{\pi}{14}$$

$$23. \quad \tan \theta = -1 \Rightarrow \tan \theta = \tan \left(2\pi - \frac{\pi}{4} \right)$$

$$\text{and } \cos \theta = \frac{1}{\sqrt{2}} \Rightarrow \cos \theta = \cos \left(2\pi - \frac{\pi}{4} \right)$$

$$\therefore \text{ general value is } 2n\pi + \left(2\pi - \frac{\pi}{4} \right) = 2n\pi + \frac{7\pi}{4}$$

$$\dots \left[\begin{array}{l} \text{If } \tan \theta = \tan \alpha \text{ and } \cos \theta = \cos \alpha \\ \Rightarrow \theta = 2n\pi + \alpha \end{array} \right]$$



$$24. \quad \tan \theta = -\frac{1}{\sqrt{3}} = \tan\left(\pi - \frac{\pi}{6}\right),$$

$$\sin \theta = \frac{1}{2} = \sin\left(\pi - \frac{\pi}{6}\right)$$

$$\text{and } \cos \theta = -\frac{\sqrt{3}}{2} = \cos\left(\pi - \frac{\pi}{6}\right)$$

$$\therefore \text{ principal value of } \theta = \pi - \frac{\pi}{6} = \frac{5\pi}{6}$$

$$25. \quad \cos p\theta = \cos q\theta \Rightarrow p\theta = 2n\pi \pm q\theta$$

$$\Rightarrow \theta = \frac{2n\pi}{p \pm q}$$

$$26. \quad (2 \cos x - 1)(3 + 2 \cos x) = 0$$

$$\Rightarrow \cos x = \frac{1}{2} \quad \dots \left[\because \cos x \neq \frac{-3}{2} \right]$$

$$\Rightarrow x = 2n\pi \pm \frac{\pi}{3}$$

$$\Rightarrow x = \frac{\pi}{3}, \frac{5\pi}{3} \text{ in } [0, 2\pi]$$

$$27. \quad \sin\left(\frac{\pi}{4} \cot \theta\right) = \cos\left(\frac{\pi}{4} \tan \theta\right)$$

$$\Rightarrow \sin\left(\frac{\pi}{4} \cot \theta\right) = \sin\left(\frac{\pi}{2} - \frac{\pi}{4} \tan \theta\right)$$

$$\Rightarrow \frac{\pi}{4} \cot \theta = \frac{\pi}{2} - \frac{\pi}{4} \tan \theta$$

$$\Rightarrow \tan \theta + \cot \theta = 2$$

$$\Rightarrow \frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} = 2$$

$$\Rightarrow \frac{1}{\sin \theta \cos \theta} = 2$$

$$\Rightarrow \sin 2\theta = 1 = \sin \frac{\pi}{2}$$

$$\Rightarrow 2\theta = (4n + 1) \frac{\pi}{2}$$

$$\Rightarrow \theta = n\pi + \frac{\pi}{4}$$

$$28. \quad \tan 2\theta \tan \theta = 1$$

$$\Rightarrow \frac{\tan 2\theta}{\cot \theta} = 1$$

$$\Rightarrow \tan 2\theta = \cot \theta$$

$$\Rightarrow \tan 2\theta = \tan\left(\frac{\pi}{2} - \theta\right)$$

$$\Rightarrow 2\theta = n\pi + \frac{\pi}{2} - \theta$$

$$\Rightarrow 3\theta = n\pi + \frac{\pi}{2}$$

$$\Rightarrow \theta = \frac{n\pi}{3} + \frac{\pi}{6} = \left(n + \frac{1}{2}\right) \frac{\pi}{3}$$

$$29. \quad \cos 2\theta = \sin \alpha$$

$$\Rightarrow \cos 2\theta = \cos\left(\frac{\pi}{2} - \alpha\right)$$

$$\Rightarrow 2\theta = 2n\pi \pm \left(\frac{\pi}{2} - \alpha\right)$$

$$\Rightarrow \theta = n\pi \pm \left(\frac{\pi}{4} - \frac{\alpha}{2}\right)$$

$$30. \quad \sin 6\theta + \sin 4\theta + \sin 2\theta = 0$$

$$\Rightarrow \sin 6\theta + \sin 2\theta + \sin 4\theta = 0$$

$$\Rightarrow 2 \sin 4\theta \cos 2\theta + \sin 4\theta = 0$$

$$\Rightarrow \sin 4\theta (2 \cos 2\theta + 1) = 0$$

$$\Rightarrow \sin 4\theta = 0 \quad \text{or} \quad 2 \cos 2\theta + 1 = 0$$

$$\Rightarrow 4\theta = n\pi \quad \text{or} \quad \cos 2\theta = \frac{-1}{2}$$

$$\Rightarrow \theta = \frac{n\pi}{4} \quad \text{or} \quad \cos 2\theta = -\cos \frac{\pi}{3}$$

$$\cos 2\theta = \cos\left(\pi - \frac{\pi}{3}\right)$$

$$\cos 2\theta = \cos \frac{2\pi}{3}$$

$$2\theta = 2n\pi \pm \frac{2\pi}{3}$$

$$\theta = n\pi \pm \frac{\pi}{3}$$

$$31. \quad \sin 5x + \sin 3x + \sin x = 0$$

$$\Rightarrow -\sin 3x = \sin 5x + \sin x = 2 \sin 3x \cos 2x$$

$$\Rightarrow \sin 3x = 0 \Rightarrow x = 0$$

$$\text{or } \cos 2x = -\frac{1}{2} = -\cos\left(\frac{\pi}{3}\right) = \cos\left(\pi - \frac{\pi}{3}\right)$$

$$\Rightarrow 2x = 2n\pi \pm \left(\pi - \frac{\pi}{3}\right) \Rightarrow x = n\pi \pm \left(\frac{\pi}{3}\right)$$

$$\Rightarrow x = \frac{\pi}{3} \quad \dots \left[\because 0 \leq x \leq \frac{\pi}{2} \right]$$

$$32. \quad \sin x + \sin 3x + \sin 5x = 0$$

$$\Rightarrow \sin 5x + \sin x + \sin 3x = 0$$

$$\Rightarrow 2 \sin 3x \cos 2x + \sin 3x = 0$$

$$\Rightarrow \sin 3x (2 \cos 2x + 1) = 0$$

$$\Rightarrow \sin 3x = 0 \quad \text{or} \quad 2 \cos 2x = -1$$



$$\begin{aligned} \Rightarrow 3x &= n\pi & \text{or} & & \cos 2x &= \frac{-1}{2} \\ \Rightarrow x &= \frac{n\pi}{3} & \text{or} & & \cos 2x &= -\cos \frac{\pi}{3} \\ & & & & \cos 2x &= \cos \left(\pi - \frac{\pi}{3} \right) \\ & & & & \cos 2x &= \cos \frac{2\pi}{3} \\ & & & & 2x &= 2n\pi \pm \frac{2\pi}{3} \\ & & & & x &= n\pi \pm \frac{\pi}{3} \\ \Rightarrow x &= \pi, \frac{2\pi}{3}, \frac{4\pi}{3} & \dots & \left[\because x \in \left[\frac{\pi}{2}, \frac{3\pi}{2} \right] \right] \end{aligned}$$

33. $\sin x - \sin 2x + \sin 3x = 2 \cos^2 x - \cos x$
 $\Rightarrow \sin x + \sin 3x - \sin 2x = \cos x (2 \cos x - 1)$
 $\Rightarrow 2 \sin 2x \cos x - \sin 2x = \cos x (2 \cos x - 1)$
 $\Rightarrow \sin 2x (2 \cos x - 1) = \cos x (2 \cos x - 1)$
 $\Rightarrow 2 \sin x \cos x = \cos x \text{ or } 2 \cos x - 1 = 0$
 $\Rightarrow \sin x = \frac{1}{2} \text{ or } \cos x = 0 \text{ or } \cos x = \frac{1}{2}$
 $\Rightarrow \sin x = \sin \frac{\pi}{6} \text{ or } \cos x = 0 \text{ or } \cos x = \cos \frac{\pi}{3}$
 $\Rightarrow x = n\pi + (-1)^n \frac{\pi}{6} \text{ or } x = (2n + 1) \frac{\pi}{2}$
 $\text{or } x = 2n\pi \pm \frac{\pi}{3}$
 $\Rightarrow x = \frac{\pi}{6}, \frac{\pi}{3}, \frac{\pi}{2}, \frac{5\pi}{6} \dots \left[\because x \in (0, \pi) \right]$

34. $\sin x \cos x = \frac{1}{4}$
 $\Rightarrow \sin 2x = \frac{1}{2} = \sin \frac{\pi}{6}$
 $\Rightarrow 2x = n\pi + (-1)^n \frac{\pi}{6}$
 $\Rightarrow x = \frac{n\pi}{2} + (-1)^n \frac{\pi}{12}$
 $\Rightarrow x = \frac{\pi}{12} \dots \left[\because x \in \left(0, \frac{\pi}{2} \right) \right]$

35. $\sin \theta + \cos \theta = 1$
Dividing both sides by $\sqrt{1^2 + 1^2} = \sqrt{2}$, we get
 $\frac{1}{\sqrt{2}} \sin \theta + \frac{1}{\sqrt{2}} \cos \theta = \frac{1}{\sqrt{2}}$
 $\Rightarrow \sin \theta \cos \frac{\pi}{4} + \cos \theta \sin \frac{\pi}{4} = \frac{1}{\sqrt{2}}$

$$\begin{aligned} \Rightarrow \sin \left(\theta + \frac{\pi}{4} \right) &= \frac{1}{\sqrt{2}} = \sin \frac{\pi}{4} \\ \dots \left[\because \sin (A+B) &= \sin A \cos B + \cos A \sin B \right] \\ \Rightarrow \theta + \frac{\pi}{4} &= n\pi + (-1)^n \frac{\pi}{4} \\ \Rightarrow \theta &= n\pi + (-1)^n \frac{\pi}{4} - \frac{\pi}{4} \end{aligned}$$

36. $\sqrt{2} \sec \theta + \tan \theta = 1 \Rightarrow \frac{\sqrt{2}}{\cos \theta} + \frac{\sin \theta}{\cos \theta} = 1$
 $\Rightarrow \sin \theta - \cos \theta = -\sqrt{2}$
 $\Rightarrow \frac{1}{\sqrt{2}} \cos \theta - \frac{1}{\sqrt{2}} \sin \theta = 1$
 $\Rightarrow \cos \left(\theta + \frac{\pi}{4} \right) = \cos (0)$
 $\Rightarrow \theta + \frac{\pi}{4} = 2n\pi \pm 0 \Rightarrow \theta = 2n\pi - \frac{\pi}{4}$

37. $\sqrt{3} \cos \theta + \sin \theta = \sqrt{2}$
Dividing both sides by $\sqrt{(\sqrt{3})^2 + 1^2} = 2$,
we get
 $\frac{\sqrt{3}}{2} \cos \theta + \frac{1}{2} \sin \theta = \frac{\sqrt{2}}{2}$
 $\Rightarrow \sin \theta \cos \frac{\pi}{3} + \cos \theta \sin \frac{\pi}{3} = \frac{1}{\sqrt{2}}$
 $\Rightarrow \sin \left(\theta + \frac{\pi}{3} \right) = \sin \left(\frac{\pi}{4} \right)$
 $\Rightarrow \theta + \frac{\pi}{3} = n\pi + (-1)^n \frac{\pi}{4}$
 $\Rightarrow \theta = n\pi + (-1)^n \frac{\pi}{4} - \frac{\pi}{3}$

38. $\sin 6\theta + \sin 4\theta + \sin 2\theta = 0$
 $\Rightarrow 2 \sin 4\theta \cos 2\theta + \sin 4\theta = 0$
 $\Rightarrow \sin 4\theta (2 \cos 2\theta + 1) = 0$
 $\Rightarrow \sin 4\theta = 0 \text{ or } 2 \cos 2\theta + 1 = 0$
Now, $\sin 4\theta = 0 \Rightarrow 4\theta = n\pi \Rightarrow \theta = \frac{n\pi}{4}$
and $2 \cos 2\theta = -1 \Rightarrow \cos 2\theta = -\frac{1}{2} = \cos \left(\frac{2\pi}{3} \right)$
 $\Rightarrow 2\theta = 2n\pi \pm \frac{2\pi}{3} \Rightarrow \theta = n\pi \pm \frac{\pi}{3}$
 $\therefore \theta = \frac{n\pi}{4} \text{ or } n\pi \pm \frac{\pi}{3}$



39. $\sin 7\theta = \sin 4\theta - \sin \theta$
 $\Rightarrow \sin 7\theta + \sin \theta - \sin 4\theta = 0$
 $\Rightarrow 2\sin 4\theta \cos 3\theta - \sin 4\theta = 0$
 $\Rightarrow \sin 4\theta (2\cos 3\theta - 1) = 0$
 $\Rightarrow \sin 4\theta = 0$ or $\cos 3\theta = \frac{1}{2}$
- Now, $\sin 4\theta = 0 \Rightarrow 4\theta = \pi \Rightarrow \theta = \frac{\pi}{4}$
- and $\cos 3\theta = \frac{1}{2} \Rightarrow 3\theta = \frac{\pi}{3} \Rightarrow \theta = \frac{\pi}{9}$
40. $\frac{\tan 3x - \tan 2x}{1 + \tan 3x \tan 2x} = 1$
 $\Rightarrow \tan(3x - 2x) = 1 \Rightarrow \tan x = 1$
 $\Rightarrow \tan x = \tan \frac{\pi}{4}$
 $\Rightarrow x = n\pi + \frac{\pi}{4}$
- But this value does not satisfy the given equation.
Hence, option (A) is the correct answer.
41. $\frac{\tan 3\theta - 1}{\tan 3\theta + 1} = \sqrt{3}$
 $\Rightarrow \frac{\tan 3\theta - \tan\left(\frac{\pi}{4}\right)}{1 + \tan 3\theta \tan\left(\frac{\pi}{4}\right)} = \sqrt{3}$
 $\Rightarrow \tan\left(3\theta - \frac{\pi}{4}\right) = \tan \frac{\pi}{3}$
 $\Rightarrow 3\theta - \frac{\pi}{4} = n\pi + \frac{\pi}{3} \Rightarrow 3\theta = n\pi + \frac{7\pi}{12}$
 $\Rightarrow \theta = \frac{n\pi}{3} + \frac{7\pi}{36}$
42. $\cos 3\theta = \sin 2\theta$
 $\Rightarrow \cos 3\theta = \cos\left(\frac{\pi}{2} - 2\theta\right)$
 $\Rightarrow 3\theta = 2n\pi \pm \left(\frac{\pi}{2} - 2\theta\right)$
 $\Rightarrow \theta = \frac{2n\pi}{3} \pm \left(\frac{\pi}{6} - \frac{2\theta}{3}\right)$
 $\Rightarrow \theta = \frac{2n\pi}{5} + \frac{\pi}{10}$ or $\theta = 2n\pi - \frac{\pi}{2}$
- Since, θ is acute
 $\Rightarrow \theta = \frac{\pi}{10} = 18^\circ$
 $\Rightarrow \sin \theta = \frac{\sqrt{5}-1}{4}$ $\left[\because \sin 18^\circ = \frac{\sqrt{5}-1}{4} \right]$

43. $\sin x - 3\sin 2x + \sin 3x$
 $= \cos x - 3\cos 2x + \cos 3x$
 $\Rightarrow (\sin x + \sin 3x) - 3\sin 2x - (\cos x + \cos 3x)$
 $+ 3\cos 2x = 0$
 $\Rightarrow 2\sin 2x \cos x - 3\sin 2x - 2\cos 2x \cos x$
 $+ 3\cos 2x = 0$
 $\Rightarrow \sin 2x(2\cos x - 3) - \cos 2x(2\cos x - 3) = 0$
 $\Rightarrow (\sin 2x - \cos 2x)(2\cos x - 3) = 0$
 $\Rightarrow \cos 2x = \sin 2x$ $\left[\because \cos x \neq \frac{3}{2} \right]$
- $\Rightarrow \cos 2x = \cos\left(\frac{\pi}{2} - 2x\right)$
 $\Rightarrow 2x = 2n\pi \pm \left(\frac{\pi}{2} - 2x\right)$
- Neglecting (-) sign, we get
 $x = \frac{n\pi}{2} + \frac{\pi}{8}$
44. $\cot(\alpha + \beta) = 0 \Rightarrow \cos(\alpha + \beta) = 0$
 $\Rightarrow \alpha + \beta = (2n + 1)\frac{\pi}{2}$
- $\therefore \sin(\alpha + 2\beta) = \sin(2\alpha + 2\beta - \alpha)$
 $= \sin[(2n + 1)\pi - \alpha]$
 $= \sin(2n\pi + \pi - \alpha)$
 $= \sin(\pi - \alpha) = \sin \alpha$
45. $\tan \theta + \tan 2\theta + \tan \theta \cdot \tan 2\theta = 1$
 $\Rightarrow \tan \theta + \tan 2\theta = 1 - \tan \theta \cdot \tan 2\theta$
 $\Rightarrow \frac{\tan \theta + \tan 2\theta}{1 - \tan \theta \cdot \tan 2\theta} = 1$
 $\Rightarrow \tan(\theta + 2\theta) = 1$
 $\Rightarrow \tan(3\theta) = 1 = \tan \frac{\pi}{4}$
 $\Rightarrow 3\theta = n\pi + \frac{\pi}{4} \Rightarrow \theta = \frac{n\pi}{3} + \frac{\pi}{12}$
46. $\sec 4\theta - \sec 2\theta = 2$
 $\Rightarrow \frac{1}{\cos 4\theta} - \frac{1}{\cos 2\theta} = 2$
 $\Rightarrow \cos 2\theta - \cos 4\theta = 2\cos 4\theta \cos 2\theta$
 $\Rightarrow \cos 2\theta - \cos 4\theta = \cos 6\theta + \cos 2\theta$
 $\dots \left[\because 2\cos A \cos B = \cos(A + B) + \cos(A - B) \right]$
 $\Rightarrow \cos 6\theta + \cos 4\theta = 0$
 $\Rightarrow 2\cos 5\theta \cos \theta = 0$
 $\dots \left[\because \cos C + \cos D = 2\cos\left(\frac{C+D}{2}\right)\cos\left(\frac{C-D}{2}\right) \right]$



$$\Rightarrow \cos \theta = 0 \text{ or } \cos 5\theta = 0$$

$$\Rightarrow \theta = (2n+1) \frac{\pi}{2} \text{ or } 5\theta = (2n+1) \frac{\pi}{2}$$

$$\Rightarrow \theta = n\pi + \frac{\pi}{2} \text{ or } \theta = \frac{n\pi}{5} + \frac{\pi}{10}$$

47. $\sin 2x + \sin 4x = 2 \sin 3x$
 $\Rightarrow 2 \sin 3x \cos x - 2 \sin 3x = 0$
 $\Rightarrow \sin 3x = 0 \text{ or } \cos x = 1 \Rightarrow 3x = n\pi \text{ or } x = 2n\pi$
 $\Rightarrow x = \frac{n\pi}{3} \text{ or } x = 2n\pi$

48. $a \sin x + b \cos x = c$
 $\Rightarrow \frac{a}{\sqrt{a^2+b^2}} \sin x + \frac{b}{\sqrt{a^2+b^2}} \cos x = \frac{c}{\sqrt{a^2+b^2}}$
 $\Rightarrow \cos \alpha \sin x + \sin \alpha \cos x = \frac{c}{\sqrt{a^2+b^2}}$
 $\Rightarrow \sin(x+\alpha) = \frac{c}{\sqrt{a^2+b^2}} > 1$, which is not possible.

\therefore there is no solution.

49. $\tan\left(\frac{p\pi}{4}\right) = \cot\left(\frac{q\pi}{4}\right)$
 $\Rightarrow \tan\left(\frac{p\pi}{4}\right) = \tan\left(\frac{\pi}{2} - \frac{q\pi}{4}\right)$
 $\Rightarrow \frac{p\pi}{4} = n\pi + \frac{\pi}{2} - \frac{q\pi}{4}$
 $\Rightarrow \frac{p}{4} = n + \frac{1}{2} - \frac{q}{4}$
 $\Rightarrow \frac{p+q}{4} = \frac{2n+1}{2}$
 $\Rightarrow p+q = 2(2n+1)$

50. $2\sin^2 x + \sin^2 2x = 2$
 $\Rightarrow (1 - \cos 2x) + (1 - \cos^2 2x) = 2$
 ...[$\because \sin^2 \theta + \cos^2 \theta = 1$ and $2\sin^2 \theta = 1 - \cos 2\theta$]
 $\Rightarrow \cos 2x (\cos 2x + 1) = 0$
 $\Rightarrow \cos 2x = 0 \text{ or } \cos 2x = -1$
 $\Rightarrow 2x = (2n+1) \frac{\pi}{2} \text{ or } (2n+1)\pi$
 $\Rightarrow x = (2n+1) \frac{\pi}{4} \text{ or } (2n+1) \frac{\pi}{2}$

Putting $n = -2, -1, 0, 1, 2$, we get

$$x = \frac{-3\pi}{4}, \frac{-\pi}{4}, \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}$$

and $\frac{-3\pi}{2}, \frac{-\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}$

Since, $-\pi < x < \pi$

$$\therefore x = \pm \frac{\pi}{4}, \pm \frac{\pi}{2}, \pm \frac{3\pi}{4}$$

\therefore option (B) is the correct answer.

51. $\tan(\cot x) = \cot(\tan x)$
 $\Rightarrow \tan(\cot x) = \tan\left(\frac{\pi}{2} - \tan x\right)$
 $\Rightarrow \cot x = n\pi + \frac{\pi}{2} - \tan x$
 $\Rightarrow \cot x + \tan x = n\pi + \frac{\pi}{2}$
 $\Rightarrow \frac{\cos x}{\sin x} + \frac{\sin x}{\cos x} = n\pi + \frac{\pi}{2}$
 $\Rightarrow \frac{2(\cos^2 x + \sin^2 x)}{2 \sin x \cos x} = n\pi + \frac{\pi}{2}$
 $\Rightarrow \frac{2}{\sin 2x} = n\pi + \frac{\pi}{2}$
 $\Rightarrow \sin 2x = \frac{2}{n\pi + \frac{\pi}{2}} = \frac{4}{(2n+1)\pi}$

52. Let $\sqrt{3} + 1 = r \cos \alpha$ and $\sqrt{3} - 1 = r \sin \alpha$.

$$\text{Then } r = \sqrt{(\sqrt{3}+1)^2 + (\sqrt{3}-1)^2} = 2\sqrt{2}$$

$$\tan \alpha = \frac{\sqrt{3}-1}{\sqrt{3}+1} = \frac{1 - \left(\frac{1}{\sqrt{3}}\right)}{1 + \left(\frac{1}{\sqrt{3}}\right)} = \tan\left(\frac{\pi}{4} - \frac{\pi}{6}\right)$$

$$\Rightarrow \alpha = \frac{\pi}{12}$$

The given equation reduces to

$$2\sqrt{2} \cos(\theta - \alpha) = 2$$

$$\Rightarrow \cos\left(\theta - \frac{\pi}{12}\right) = \cos \frac{\pi}{4}$$

$$\Rightarrow \theta - \frac{\pi}{12} = 2n\pi \pm \frac{\pi}{4} \Rightarrow \theta = 2n\pi \pm \frac{\pi}{4} + \frac{\pi}{12}$$

53. $\sec \theta - \operatorname{cosec} \theta = \frac{4}{3}$
 $\Rightarrow 3(\sin \theta - \cos \theta) = 4 \sin \theta \cos \theta$
 $\Rightarrow 3(\sin \theta - \cos \theta) = 2 \sin 2\theta$



Squaring on both sides, we get $9(1-s) = 4s^2$,
 where $s = \sin 2\theta$
 $\Rightarrow 4s^2 + 9s - 9 = 0$
 $\Rightarrow (s+3)(4s-3) = 0 \Rightarrow s = \frac{3}{4}$
 ...[$\because \sin 2\theta \neq -3$]

$$\Rightarrow \sin 2\theta = \frac{3}{4} = \sin \alpha$$

$$\Rightarrow 2\theta = n\pi + (-1)^n \alpha$$

$$\Rightarrow \theta = \frac{1}{2} \left[n\pi + (-1)^n \sin^{-1} \left(\frac{3}{4} \right) \right]$$

54. Using $\sec 2\theta = \frac{1}{\cos 2\theta} = \frac{1 + \tan^2 \theta}{1 - \tan^2 \theta}$, we can
 write the given equation as

$$\tan^2 \theta + \frac{1 + \tan^2 \theta}{1 - \tan^2 \theta} = 1$$

$$\Rightarrow \tan^2 \theta (1 - \tan^2 \theta) + 1 + \tan^2 \theta = 1 - \tan^2 \theta$$

$$\Rightarrow 3\tan^2 \theta - \tan^4 \theta = 0$$

$$\Rightarrow \tan^2 \theta (3 - \tan^2 \theta) = 0$$

$$\Rightarrow \tan \theta = 0 \text{ or } \tan^2 \theta = 3$$

$$\Rightarrow \tan \theta = 0 \text{ or } \tan^2 \theta = \tan^2 \frac{\pi}{3}$$

$$\Rightarrow \theta = m\pi \text{ or } \theta = n\pi \pm \frac{\pi}{3},$$

where m and n are integers.

55. $2\sqrt{3} \cos \theta = \tan \theta$

$$\Rightarrow 2\sqrt{3} \cos^2 \theta = \sin \theta$$

$$\Rightarrow 2\sqrt{3} \sin^2 \theta + \sin \theta - 2\sqrt{3} = 0$$

$$\Rightarrow \sin \theta = \frac{-1 \pm 7}{4\sqrt{3}} \Rightarrow \sin \theta = \frac{-8}{4\sqrt{3}},$$

which is not possible

$$\text{and } \sin \theta = \frac{6}{4\sqrt{3}} = \frac{\sqrt{3}}{2} = \sin \frac{\pi}{3}$$

$$\Rightarrow \theta = n\pi + (-1)^n \frac{\pi}{3}$$

56. $3 \sin^2 x - 7 \sin x + 2 = 0$

$$\Rightarrow 3\sin^2 x - 6 \sin x - \sin x + 2 = 0$$

$$\Rightarrow 3\sin x (\sin x - 2) - (\sin x - 2) = 0$$

$$\Rightarrow (3 \sin x - 1)(\sin x - 2) = 0$$

$$\Rightarrow \sin x = \frac{1}{3} \text{ or } 2$$

$$\Rightarrow \sin x = \frac{1}{3} \quad \dots[\because \sin x \neq 2]$$

Let $\sin^{-1} \frac{1}{3} = \alpha$, $0 < \alpha < \frac{\pi}{2}$ are the solutions

in $[0, 5\pi]$. Then, α , $\pi - \alpha$, $2\pi + \alpha$, $3\pi - \alpha$,
 $4\pi + \alpha$, $5\pi - \alpha$ are the solutions in $[0, 5\pi]$.

\therefore number of solutions = 6

57. $\sin 2x + \cos 2x = 0$

$$\Rightarrow (\sin 2x + \cos 2x)^2 = 0$$

$$\Rightarrow \sin^2 2x + \cos^2 2x + 2 \sin 2x \cos 2x = 0$$

$$\Rightarrow 1 + \sin 4x = 0 \Rightarrow \sin 4x = -1$$

$$\therefore 4x = n\pi + (-1)^n \left(\frac{-\pi}{2} \right)$$

$$\therefore 4x = n\pi + (-1)^{n+1} \frac{\pi}{2}$$

$$\therefore x = \frac{n\pi}{4} + (-1)^{n+1} \frac{\pi}{8}$$

\therefore For $\pi < x < 2\pi$, the values of x are $\frac{11\pi}{8}$, $\frac{15\pi}{8}$.

58. $2\sin^2 \theta = 3\cos \theta$

$$\Rightarrow 2 - 2\cos^2 \theta = 3 \cos \theta$$

$$\Rightarrow 2\cos^2 \theta + 3 \cos \theta - 2 = 0$$

$$\Rightarrow \cos \theta = \frac{-3 \pm \sqrt{9+16}}{4} = \frac{-3 \pm 5}{4}$$

Neglecting $(-)$ sign, we get

$$\cos \theta = \frac{1}{2} = \cos \left(\frac{\pi}{3} \right) \Rightarrow \theta = 2n\pi \pm \frac{\pi}{3}$$

The values of θ between 0 and 2π are $\frac{\pi}{3}$, $\frac{5\pi}{3}$.

59. $5 \cos 2\theta + 2\cos^2 \frac{\theta}{2} + 1 = 0$

$$\Rightarrow 5(2 \cos^2 \theta - 1) + (1 + \cos \theta) + 1 = 0$$

$$\Rightarrow 10 \cos^2 \theta + \cos \theta - 3 = 0$$

$$\Rightarrow (5 \cos \theta + 3)(2 \cos \theta - 1) = 0$$

$$\Rightarrow \cos \theta = \frac{1}{2}, \cos \theta = -\frac{3}{5}$$

$$\Rightarrow \theta = \frac{\pi}{3}, \pi - \cos^{-1} \left(\frac{3}{5} \right)$$

60. $2\sin^2 \theta + \sqrt{3} \cos \theta + 1 = 0$

$$\Rightarrow 2 - 2\cos^2 \theta + \sqrt{3} \cos \theta + 1 = 0$$

$$\Rightarrow 2 \cos^2 \theta - \sqrt{3} \cos \theta - 3 = 0$$

$$\Rightarrow \cos \theta = \frac{\sqrt{3} \pm \sqrt{3+24}}{4} = \frac{\sqrt{3}(1 \pm 3)}{4} = \sqrt{3} \left(\frac{-1}{2} \right)$$

$$\Rightarrow \theta = \frac{5\pi}{6}$$



61. $3\sin^2 x + 10 \cos x - 6 = 0$
 $\Rightarrow 3(1 - \cos^2 x) + 10 \cos x - 6 = 0$
 $\Rightarrow 3 - 3 \cos^2 x + 10 \cos x - 6 = 0$
 $\Rightarrow 3 \cos^2 x - 10 \cos x + 3 = 0$
 $\Rightarrow 3 \cos^2 x - 9 \cos x - \cos x + 3 = 0$
 $\Rightarrow 3 \cos x (\cos x - 3) - 1 (\cos x - 3) = 0$
 $\Rightarrow (\cos x - 3)(3 \cos x - 1) = 0$
 $\Rightarrow \cos x = 3, \text{ (which is not possible)}$
or $\cos x = \frac{1}{3}$
 $\Rightarrow \cos x = \frac{1}{3} = \cos \alpha \text{ (say)}$
 $\Rightarrow x = 2n\pi \pm \alpha$
 $\Rightarrow x = 2n\pi \pm \cos^{-1}\left(\frac{1}{3}\right)$

62. $\cos^2 x - 2 \cos x = 4 \sin x - \sin 2x$
 $\Rightarrow \cos^2 x - 2 \cos x = 4 \sin x - 2 \sin x \cos x$
 $\Rightarrow \cos x (\cos x - 2) = 2 \sin x (2 - \cos x)$
 $\Rightarrow \cos x (\cos x - 2) - 2 \sin x (2 - \cos x) = 0$
 $\Rightarrow \cos x (\cos x - 2) + 2 \sin x (\cos x - 2) = 0$
 $\Rightarrow (\cos x - 2)(\cos x + 2 \sin x) = 0$
 $\Rightarrow \cos x + 2 \sin x = 0 \quad \dots [\because \cos x \neq 2]$
 $\Rightarrow \cos x = -2 \sin x$
 $\Rightarrow \tan x = -\frac{1}{2} = \tan \alpha \text{ (say)}$
 $\Rightarrow x = n\pi + \alpha$
 $\Rightarrow x = n\pi + \tan^{-1}\left(-\frac{1}{2}\right), n \in \mathbb{I}$
Since, $0 \leq x \leq \pi$
 $\therefore x = \pi + \tan^{-1}\left(-\frac{1}{2}\right)$

63. $\cos 2\theta = (\sqrt{2} + 1) \left(\cos \theta - \frac{1}{\sqrt{2}} \right)$
 $\Rightarrow 2 \cos^2 \theta - 1 = \frac{\sqrt{2} + 1}{\sqrt{2}} (\sqrt{2} \cos \theta - 1)$
 $\Rightarrow 2 \cos^2 \theta - 1 - \frac{\sqrt{2} + 1}{\sqrt{2}} (\sqrt{2} \cos \theta - 1) = 0$
 $\Rightarrow (\sqrt{2} \cos \theta - 1) \left\{ (\sqrt{2} \cos \theta + 1) - \left(\frac{\sqrt{2} + 1}{\sqrt{2}} \right) \right\} = 0$
 $\Rightarrow \sqrt{2} \cos \theta - 1 = 0 \text{ or } \sqrt{2} \cos \theta + 1 = \frac{\sqrt{2} + 1}{\sqrt{2}}$
 $\Rightarrow \sqrt{2} \cos \theta = 1 \text{ or } \sqrt{2} \cos \theta = \frac{\sqrt{2} + 1 - \sqrt{2}}{\sqrt{2}}$

$$\Rightarrow \cos \theta = \frac{1}{\sqrt{2}} \text{ or } \sqrt{2} \cos \theta = \frac{1}{\sqrt{2}}$$

$$\Rightarrow \cos \theta = \frac{1}{\sqrt{2}} \text{ or } \cos \theta = \frac{1}{2}$$

$$\Rightarrow \cos \theta = \frac{1}{\sqrt{2}} \quad \dots \left[\because \cos \theta \neq \frac{1}{2} \right]$$

$$\Rightarrow \cos \theta = \cos \frac{\pi}{4}$$

$$\Rightarrow \theta = 2n\pi \pm \frac{\pi}{4}$$

64. $\cos 2\theta = \frac{1}{3}$
 $\Rightarrow \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} = \frac{1}{3}$
 $\Rightarrow 3 - 3 \tan^2 \theta = 1 + \tan^2 \theta \quad \Rightarrow 2 = 4 \tan^2 \theta$
 $\Rightarrow \tan^2 \theta = \frac{1}{2}$
 $\Rightarrow \tan^8 \theta = \frac{1}{16}$
Now, $32 \tan^8 \theta = 2 \cos^2 \alpha - 3 \cos \alpha$
 $\Rightarrow 32 \left(\frac{1}{16} \right) = 2 \cos^2 \alpha - 3 \cos \alpha$
 $\Rightarrow 2 \cos^2 \alpha - 3 \cos \alpha - 2 = 0$
 $\Rightarrow (2 \cos \alpha + 1)(\cos \alpha - 2) = 0$
But $\cos \alpha - 2 \neq 0$
 $\therefore 2 \cos \alpha + 1 = 0$
 $\Rightarrow \cos \alpha = -\frac{1}{2}$
 $\Rightarrow \cos \alpha = \cos \frac{2\pi}{3}$
 $\Rightarrow \alpha = 2n\pi \pm \frac{2\pi}{3}$

65. $\cos 2\theta = \sin \theta \Rightarrow 1 - 2 \sin^2 \theta = \sin \theta$
 $\Rightarrow 2 \sin^2 \theta + \sin \theta - 1 = 0$
 $\Rightarrow (2 \sin \theta - 1)(\sin \theta + 1) = 0$
 $\Rightarrow \sin \theta = \frac{1}{2} \text{ or } \sin \theta = -1$
 $\therefore \sin \theta = \frac{1}{2} = \sin \frac{\pi}{6} \Rightarrow \theta = n\pi + (-1)^n \frac{\pi}{6}$
and $\sin \theta = -1 = \sin \frac{3\pi}{2}$
 $\Rightarrow \theta = m\pi + (-1)^m \frac{3\pi}{2}$
 $\therefore \theta = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{3\pi}{2}$
 $\therefore \text{number of solutions} = 3$



$$\begin{aligned}
 66. \quad & \tan \theta = \cot 5\theta \\
 & \Rightarrow \tan \theta - \cot 5\theta = 0 \\
 & \Rightarrow \frac{\sin \theta}{\cos \theta} - \frac{\cos 5\theta}{\sin 5\theta} = 0 \\
 & \Rightarrow \cos 5\theta \cos \theta - \sin 5\theta \sin \theta = 0 \\
 & \Rightarrow \cos(5\theta + \theta) = 0 \\
 & \Rightarrow \cos 6\theta = 0 = \cos \frac{\pi}{2} \\
 & \Rightarrow 6\theta = 2n\pi \pm \frac{\pi}{2} \\
 & \Rightarrow 6\theta = \pm \frac{\pi}{2}, \pm \frac{3\pi}{2}, \pm \frac{5\pi}{2} \\
 & \Rightarrow \theta = \pm \frac{\pi}{12}, \pm \frac{\pi}{4}, \pm \frac{5\pi}{12} \\
 & \text{and } \sin 2\theta = \cos 4\theta \\
 & \Rightarrow \sin 2\theta = 1 - 2\sin^2 2\theta \\
 & \Rightarrow 2\sin^2 2\theta + \sin 2\theta - 1 = 0 \\
 & \Rightarrow (2\sin 2\theta - 1)(\sin 2\theta + 1) = 0 \\
 & \Rightarrow \sin 2\theta = \frac{1}{2} \text{ or } \sin 2\theta = -1 \\
 & \Rightarrow \sin 2\theta = \sin\left(\frac{\pi}{6}\right) \text{ or } \sin 2\theta = -1 \\
 & \Rightarrow 2\theta = n\pi + (-1)^n \frac{\pi}{6} \text{ or } 2\theta = (4n-1)\frac{\pi}{2} \\
 & \Rightarrow 2\theta = \frac{\pi}{6}, \frac{5\pi}{6} \text{ or } 2\theta = -\frac{\pi}{2} \\
 & \Rightarrow \theta = \frac{\pi}{12}, \frac{5\pi}{12} \text{ or } \theta = -\frac{\pi}{4}
 \end{aligned}$$

\therefore the common values of θ are $-\frac{\pi}{4}, \frac{\pi}{12}$ and $\frac{5\pi}{12}$.

Hence, there are 3 values of θ satisfying the given equation.

$$\begin{aligned}
 67. \quad & \cos^2\left(x + \frac{\pi}{6}\right) + \cos^2 x - 2\cos\left(x + \frac{\pi}{6}\right)\cos\left(\frac{\pi}{6}\right) \\
 & = \sin^2 \frac{\pi}{6} \\
 & \Rightarrow \cos^2\left(x + \frac{\pi}{6}\right) + \left(\cos^2 x - \sin^2 \frac{\pi}{6}\right) \\
 & \quad - 2\cos\left(x + \frac{\pi}{6}\right)\cos\left(\frac{\pi}{6}\right) = 0 \\
 & \Rightarrow \cos^2\left(x + \frac{\pi}{6}\right) + \cos\left(x + \frac{\pi}{6}\right)\cos\left(x - \frac{\pi}{6}\right) \\
 & \quad - 2\cos\left(x + \frac{\pi}{6}\right)\cos\frac{\pi}{6} = 0
 \end{aligned}$$

$$\begin{aligned}
 & \Rightarrow \cos\left(x + \frac{\pi}{6}\right)\left\{\cos\left(x + \frac{\pi}{6}\right) + \cos\left(x - \frac{\pi}{6}\right) - 2\cos\frac{\pi}{6}\right\} = 0 \\
 & \Rightarrow \cos\left(x + \frac{\pi}{6}\right)\left\{2\cos x \cos\frac{\pi}{6} - 2\cos\frac{\pi}{6}\right\} = 0 \\
 & \Rightarrow 2\cos\left(x + \frac{\pi}{6}\right)\cos\frac{\pi}{6}(\cos x - 1) = 0 \\
 & \Rightarrow \cos\left(x + \frac{\pi}{6}\right)(\cos x - 1) = 0 \\
 & \Rightarrow \cos\left(x + \frac{\pi}{6}\right) = 0 \text{ or } \cos x = 1 \\
 & \Rightarrow x + \frac{\pi}{6} = (2n+1)\frac{\pi}{2} \text{ or } x = 2n\pi \\
 & \Rightarrow x + \frac{\pi}{6} = \pm \frac{\pi}{2} \text{ or } x = 0 \\
 & \Rightarrow x = \frac{\pi}{3}, \frac{-2\pi}{3}, 0 \\
 & \Rightarrow x = 0, \frac{\pi}{3} \quad \dots \left[\because x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \right]
 \end{aligned}$$

\therefore number of solutions = 2.

$$\begin{aligned}
 68. \quad & 8\cos x \left[\cos\left(\frac{\pi}{6} + x\right) \cdot \cos\left(\frac{\pi}{6} - x\right) - \frac{1}{2} \right] = 1 \\
 & \Rightarrow 8\cos x \left(\cos^2 \frac{\pi}{6} - \sin^2 x - \frac{1}{2} \right) = 1 \\
 & \Rightarrow 8\cos x \left(\frac{3}{4} - \sin^2 x - \frac{1}{2} \right) = 1 \\
 & \Rightarrow 8\cos x \left(\frac{1}{4} - (1 - \cos^2 x) \right) = 1 \\
 & \Rightarrow 2(4\cos^3 x - 3\cos x) = 1 \\
 & \Rightarrow 2\cos 3x = 1 \\
 & \Rightarrow \cos 3x = \frac{1}{2} \\
 & \Rightarrow \cos 3x = \cos \frac{\pi}{3} \\
 & \Rightarrow x = \frac{2n\pi}{3} \pm \frac{\pi}{9} \\
 & \Rightarrow x = \frac{\pi}{9}, \frac{5\pi}{9}, \frac{7\pi}{9} \quad \dots \left[\because x \in [0, \pi] \right] \\
 & \text{Sum} = \frac{\pi}{9} + \frac{5\pi}{9} + \frac{7\pi}{9} = \frac{13\pi}{9} \\
 & \Rightarrow k = \frac{13}{9}
 \end{aligned}$$



$$69. \quad \sec^2 \theta = \frac{4}{3}$$

$$\Rightarrow \cos^2 \theta = \frac{3}{4} = \left(\frac{\sqrt{3}}{2}\right)^2$$

$$\Rightarrow \cos^2 \theta = \cos^2 \left(\frac{\pi}{6}\right)$$

$$\Rightarrow \theta = n\pi \pm \frac{\pi}{6}$$

....[$\because \cos^2 \theta = \cos^2 \alpha \Rightarrow \theta = n\pi \pm \alpha$]

$$70. \quad \cot \theta = \sin 2\theta, (\theta \neq n\pi)$$

$$\Rightarrow \frac{\cos \theta}{\sin \theta} = 2 \sin \theta \cos \theta$$

$$\Rightarrow 2 \sin^2 \theta \cos \theta = \cos \theta$$

$$\Rightarrow \cos \theta (2 \sin^2 \theta - 1) = 0$$

$$\Rightarrow \cos \theta = 0 \text{ or } \sin^2 \theta = \frac{1}{2}$$

$$\Rightarrow \cos \theta = 0 \text{ or } \sin^2 \theta = \sin^2 \left(\frac{\pi}{4}\right)$$

$$\Rightarrow \theta = (2n+1)\frac{\pi}{2} \text{ or } \theta = n\pi \pm \frac{\pi}{4}$$

$$\Rightarrow \theta = 90^\circ \text{ and } 45^\circ$$

....[\because at $\theta = 90^\circ$ and 45° ,
the given equation is satisfied.]

$$71. \quad \text{We have, } x - y = \frac{\pi}{4} \quad \dots\text{(i)}$$

and $\cot x + \cot y = 2$

$$\Rightarrow \frac{1}{\tan x} + \frac{1}{\tan y} = 2 \quad \dots\text{(ii)}$$

From (i) and (ii), we get

$$\frac{1}{\tan\left(y + \frac{\pi}{4}\right)} + \frac{1}{\tan y} = 2$$

$$\Rightarrow (1 - \tan y) \tan y + 1 + \tan y = 2 \tan y (1 + \tan y)$$

$$\Rightarrow 3 \tan^2 y = 1$$

$$\Rightarrow \tan^2 y = \frac{1}{3} = \tan^2 \frac{\pi}{6}$$

$$\Rightarrow y = \frac{\pi}{6} \quad \dots\text{[smallest +ve value]}$$

From (i),

$$x = \frac{\pi}{4} + y = \frac{\pi}{4} + \frac{\pi}{6} = \frac{5\pi}{12}$$

$$72. \quad \cos x + \cos 2x + \cos 3x + \cos 4x = 0$$

$$\therefore \cos x + \cos 4x + \cos 2x + \cos 3x = 0$$

$$\therefore 2 \cos \frac{5x}{2} \cdot \cos \frac{3x}{2} + 2 \cos \frac{5x}{2} \cdot \cos \frac{x}{2} = 0$$

$$\Rightarrow \cos \frac{5x}{2} \cdot \left[\cos \frac{3x}{2} + \cos \frac{x}{2} \right] = 0$$

$$\Rightarrow \cos \frac{5x}{2} \cdot 2 \cos x \cdot \cos \frac{x}{2} = 0$$

$$\Rightarrow x = (2n+1)\frac{\pi}{5}, (2k+1)\frac{\pi}{2} \text{ or } (2m+1)\pi$$

$$\Rightarrow x = \frac{\pi}{5}, \frac{3\pi}{5}, \pi, \frac{7\pi}{5}, \frac{9\pi}{5}, \frac{\pi}{2}, \frac{3\pi}{2} \text{ in } 0 \leq x < 2\pi$$

$$73. \quad \text{Let the angles of the triangle be } x, 2x \text{ and } 3x.$$

Then, $x + 2x + 3x = 180^\circ \Rightarrow x = 30^\circ$

$$\therefore \text{angles of the triangle are } 30^\circ, 60^\circ \text{ and } 90^\circ.$$

$$\therefore a : b : c = \sin 30^\circ : \sin 60^\circ : \sin 90^\circ$$

$$= \frac{1}{2} : \frac{\sqrt{3}}{2} : 1 = 1 : \sqrt{3} : 2$$

$$74. \quad \text{Let } x \text{ be the common multiple.}$$

$$\therefore A + B + C = 12x = 180^\circ \Rightarrow x = 15^\circ$$

$$\therefore A = 45^\circ, B = 75^\circ, C = 60^\circ$$

$$\frac{a}{\sin 45^\circ} = \frac{b}{\sin 75^\circ} = \frac{c}{\sin 60^\circ} = k$$

$$\therefore a = \frac{1}{\sqrt{2}}k, b = \frac{\sqrt{3}+1}{2\sqrt{2}}k, c = \frac{\sqrt{3}}{2}k$$

$$\therefore a + b + c\sqrt{2} = \frac{3+3\sqrt{3}}{2\sqrt{2}} = 3b$$

$$75. \quad \text{Let the angles of the triangle be } 4x, x \text{ and } x.$$

$$\therefore 4x + x + x = 180^\circ \Rightarrow 6x = 180^\circ \Rightarrow x = 30^\circ$$

$$\frac{\sin 120^\circ}{a} = \frac{\sin 30^\circ}{b} = \frac{\sin 30^\circ}{c}$$

$$\therefore a : (a + b + c) = (\sin 120^\circ) : (\sin 120^\circ + \sin 30^\circ + \sin 30^\circ)$$

$$= \frac{\sqrt{3}}{2} : \frac{\sqrt{3}+2}{2} = \sqrt{3} : \sqrt{3} + 2$$

$$76. \quad \text{Given, } \frac{\cos A}{a} = \frac{\cos B}{b} = \frac{\cos C}{c} \quad \dots\text{(i)}$$

By Sine rule,

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c} \quad \dots\text{(ii)}$$

From (i) and (ii), we get

$$\Rightarrow \frac{\cos A}{\sin A} = \frac{\cos B}{\sin B} = \frac{\cos C}{\sin C}$$

$$\Rightarrow \cot A = \cot B = \cot C$$


$$\Rightarrow A = B = C = 60^\circ$$

$\Rightarrow \Delta ABC$ is equilateral.

$$\therefore \Delta = \frac{\sqrt{3}}{4} a^2 = \frac{\sqrt{3}}{4} (2)^2 = \sqrt{3}$$



77. $\sin^2 A + \sin^2 B = \sin^2 C$
 $\Rightarrow \sin C = 1 \Rightarrow C = \frac{\pi}{2}$



$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$
 $\Rightarrow \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{10}{1}$
 $\Rightarrow a = 10\sin A, b = 10\sin B$
 $A(\Delta ABC) = \frac{1}{2}ab = \frac{1}{2}(10\sin A)(10\sin B)$
 $= \frac{1}{2} \times 100 \times \sin A \times \sin B$

Maximum value of $\sin A \sin B = \frac{1}{2}$

$\therefore A(\Delta ABC) = \frac{1}{2} \times 100 \times \frac{1}{2}$
 $= 25 \text{ sq. units}$

78. $\sin A \sin B = \frac{ab}{c^2}$
 $\Rightarrow \sin A \sin B = \frac{(k \sin A)(k \sin B)}{k^2 \sin^2 C}$
 $\Rightarrow \sin^2 C = 1 \Rightarrow \sin C = 1 \dots [\because \sin C \neq -1]$
 $\Rightarrow \angle C = 90^\circ$
 $\Rightarrow \Delta ABC$ is right angled.

79. According to the given condition,
 $a + b + c = \frac{6(\sin A + \sin B + \sin C)}{3}$
 $\Rightarrow k(\sin A + \sin B + \sin C) = 2(\sin A + \sin B + \sin C)$

where $k = \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$
 $\Rightarrow k = 2 \dots [\because \sin A + \sin B + \sin C \neq 0]$

$\therefore \frac{a}{\sin A} = 2 \Rightarrow \sin A = \frac{1}{2} \dots [\because a = 1]$

$\Rightarrow A = \frac{\pi}{6}$

80. $\frac{a+c}{b} = \frac{\sin A + \sin C}{\sin B}$
 $= \frac{2\sin\left(\frac{A+C}{2}\right)\cos\left(\frac{A-C}{2}\right)}{\sin B}$
 $= \frac{2\sin B}{\sin B}\cos\left(\frac{A-C}{2}\right) \dots [\because 2B = A + C]$
 $= 2\cos\left(\frac{A-C}{2}\right)$

81. $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$
 According to the given condition,
 In ΔABC , $a = 2b$ and
 $A - B = 60^\circ \Rightarrow A = 60^\circ + B$
 $\Rightarrow \frac{\sin(60^\circ + B)}{2b} = \frac{\sin B}{b}$
 $\Rightarrow \frac{\sin B}{\sin(B + 60^\circ)} = \frac{1}{2}$
 $\Rightarrow 2\sin B = \sin B \cos 60^\circ + \cos B \sin 60^\circ$
 $\Rightarrow \frac{3}{2}\sin B = \frac{\sqrt{3}}{2}\cos B$
 $\therefore \tan B = \frac{1}{\sqrt{3}} \Rightarrow B = 30^\circ$
 $\therefore A = 30^\circ + 60^\circ = 90^\circ$
 $\therefore \Delta ABC$ is right angled.

82. $\frac{\cos 2A}{a^2} - \frac{\cos 2B}{b^2}$
 $= \frac{1 - 2\sin^2 A}{a^2} - \frac{1 - 2\sin^2 B}{b^2}$
 $= \frac{1}{a^2} - \frac{1}{b^2} - \frac{2\sin^2 A}{a^2} + \frac{2\sin^2 B}{b^2}$
 $= \frac{1}{a^2} - \frac{1}{b^2} - 2\left(\frac{\sin^2 A}{a^2} - \frac{\sin^2 B}{b^2}\right)$
 $= \frac{1}{a^2} - \frac{1}{b^2} \dots \left[\text{By sine rule, } \frac{a}{\sin A} = \frac{b}{\sin B} \right]$

83. $\cos C = \frac{3^2 + 5^2 - 7^2}{2 \times 3 \times 5}$
 $\Rightarrow \cos C = -\frac{1}{2}$
 $\Rightarrow C = 120^\circ$
 \therefore option (B) is the correct answer.

84. $2\cos A = \frac{\sin B}{\sin C} \Rightarrow \frac{2(c^2 + b^2 - a^2)}{2bc} = \frac{b}{c}$
 $\Rightarrow c^2 = a^2 \Rightarrow c = a$

85. $\cos B = \frac{c^2 + a^2 - b^2}{2ac} \Rightarrow \cos B = \frac{1}{2} \Rightarrow B = \frac{\pi}{3}$

86. $(a + b + c)(a - b + c) = 3ac$
 $\Rightarrow a^2 + 2ac + c^2 - b^2 = 3ac$
 $\Rightarrow a^2 + c^2 - b^2 = ac$
 But $\cos B = \frac{a^2 + c^2 - b^2}{2ac} = \frac{1}{2} \Rightarrow B = 60^\circ$



87. $\cos A = \frac{8^2 + 10^2 - 6^2}{2 \cdot 8 \cdot 10} = \frac{128}{160} = \frac{4}{5}$
 $\therefore \sin A = \frac{3}{5}$
 $\therefore \sin 2A = 2 \sin A \cdot \cos A = 2 \cdot \frac{3}{5} \cdot \frac{4}{5} = \frac{24}{25}$

88. $\cos C = \frac{81 + 64 - x^2}{2 \cdot 9 \cdot 8} \Rightarrow \frac{2}{3} = \frac{145 - x^2}{144}$
 $\Rightarrow x^2 = 49 \Rightarrow x = 7$

89. $\cos A = \frac{b^2 + c^2 - a^2}{2bc} = \frac{9 + 16 - 4}{2 \times 3 \times 4} = \frac{7}{8}$
 $\Rightarrow A = \cos^{-1} \left(\frac{7}{8} \right)$

90. Let $a = 3, b = 5, c = 7$
 $\cos C = \frac{a^2 + b^2 - c^2}{2ab} = \frac{9 + 25 - 49}{2 \times 3 \times 5} = \frac{-15}{30} = -\frac{1}{2}$
 $\therefore \angle C = \frac{2\pi}{3}$

91. Since, $c = \sqrt{13}$ is the smallest side.
 $\therefore C$ is the smallest angle.
 $\therefore \cos C = \frac{b^2 + a^2 - c^2}{2ab} = \frac{48 + 49 - 13}{2 \times 7 \times 4\sqrt{3}}$
 $\Rightarrow \cos C = \frac{3}{2\sqrt{3}} = \frac{\sqrt{3}}{2}$
 $\Rightarrow C = 30^\circ$

92. $\frac{\sin(A-B)}{\sin(A+B)} = \frac{\sin A \cos B - \sin B \cos A}{\sin C}$
 $= \frac{a}{c} \cos B - \frac{b}{c} \cos A$
 But $\cos B = \frac{a^2 + c^2 - b^2}{2ac}, \cos A = \frac{b^2 + c^2 - a^2}{2bc}$
 $\therefore \frac{\sin(A-B)}{\sin(A+B)} = \frac{1}{2c^2} (a^2 + c^2 - b^2 - b^2 - c^2 + a^2)$
 $= \frac{a^2 - b^2}{c^2}$

93. $\frac{1}{a+c} + \frac{1}{b+c} = \frac{3}{a+b+c}$
 $\Rightarrow \frac{a+b+2c}{(a+c)(b+c)} = \frac{3}{a+b+c}$
 $\Rightarrow (a+b+2c)(a+b+c) = 3(a+c)(b+c)$
 $\Rightarrow a^2 + b^2 - c^2 = ab$
 $\therefore \cos C = \frac{a^2 + b^2 - c^2}{2ab} = \frac{ab}{2ab} = \frac{1}{2} = \cos 60^\circ$
 $\Rightarrow C = 60^\circ$

94. $\angle C = 60^\circ, \frac{1}{a+c} + \frac{1}{b+c} = \frac{3}{a+b+c}$
 $\therefore \frac{3}{a+b+c} - \frac{1}{a+c} = \frac{1}{b+c}$

95. $A + C = 2B \Rightarrow B = 60^\circ$
 $\cos B = \frac{a^2 + c^2 - b^2}{2ac}$
 $\Rightarrow \cos 60^\circ = \frac{a^2 + c^2 - b^2}{2ac}$
 $\Rightarrow b^2 = a^2 + c^2 - ac$
 $\therefore \frac{a+c}{\sqrt{a^2 - ac + c^2}} = \frac{a+c}{b} = \frac{\sin A + \sin C}{\sin B}$
 $= \frac{2 \sin \frac{A+C}{2} \cos \frac{A-C}{2}}{2 \sin \frac{B}{2} \sin \frac{A+C}{2}} = \frac{\cos \frac{A-C}{2}}{\sin \frac{B}{2}}$
 $= \frac{\cos \frac{A-C}{2}}{\sin 30^\circ} = 2 \cos \frac{A-C}{2}$

96. $ab^2 \cos A + ba^2 \cos B + ac^2 \cos A + ca^2 \cos C$
 $+ bc^2 \cos B + b^2c \cos C$
 $= ab(b \cos A + a \cos B) + ac(c \cos A + a \cos C)$
 $+ bc(c \cos B + b \cos C)$
 $= abc + abc + abc = 3abc$

97. Let $\frac{b+c}{11} = \frac{c+a}{12} = \frac{a+b}{13} = k$
 $\therefore b+c = 11k \dots(i)$
 $c+a = 12k \dots(ii)$
 and $a+b = 13k \dots(iii)$
 From (i) + (ii) + (iii), $2(a+b+c) = 36k$
 $\therefore a+b+c = 18k \dots(iv)$
 Now, (iv) - (i) gives, $a = 7k$
 (iv) - (ii) gives, $b = 6k$
 (iv) - (iii) gives, $c = 5k$
 Now,
 $\cos C = \frac{a^2 + b^2 - c^2}{2ab} = \frac{(7k)^2 + (6k)^2 - (5k)^2}{2 \times (7k) \times (6k)}$
 $= \frac{49k^2 + 36k^2 - 25k^2}{84k^2} = \frac{60k^2}{84k^2} = \frac{5}{7}$

98. $a(b \cos C - c \cos B)$
 $= a \left(b \frac{a^2 + b^2 - c^2}{2ab} - c \frac{c^2 + a^2 - b^2}{2ca} \right)$
 \dots [By cosine rule]
 $= a \left(\frac{a^2 + b^2 - c^2}{2a} - \frac{c^2 + a^2 - b^2}{2a} \right)$
 $= \frac{1}{2} \times 2(b^2 - c^2) = b^2 - c^2$



$$99. \quad a^2 \cos^2 A - b^2 - c^2 = 0$$

$$\Rightarrow \cos^2 A = \frac{b^2 + c^2}{a^2}$$

Since, $\cos^2 A \leq 1$ i.e., $\cos^2 A < 1$

$$\therefore \frac{b^2 + c^2}{a^2} < 1 \Rightarrow b^2 + c^2 - a^2 < 0$$

$$\therefore \frac{b^2 + c^2 - a^2}{2bc} < 0 \quad \dots [\because 2bc > 0]$$

$$\therefore \cos A < 0 \Rightarrow A \in \left(\frac{\pi}{2}, \pi \right)$$

$$100. \quad \text{Let } a = \alpha - \beta, b = \alpha + \beta, c = \sqrt{3\alpha^2 + \beta^2}$$

Since $\sqrt{3\alpha^2 + \beta^2}$ is the largest side, the largest angle is C.

$$\therefore \cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

$$\Rightarrow \cos C = \frac{\alpha^2 + \beta^2 - 2\alpha\beta + \alpha^2 + \beta^2 + 2\alpha\beta - 3\alpha^2 - \beta^2}{2(\alpha^2 - \beta^2)}$$

$$\Rightarrow \cos C = -\frac{(\alpha^2 - \beta^2)}{2(\alpha^2 - \beta^2)} = \cos\left(\frac{2\pi}{3}\right)$$

$$\Rightarrow C = \frac{2\pi}{3}$$

$$101. \quad \cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$\Rightarrow \cos 60^\circ = \frac{1}{2} = \frac{9 + c^2 - 16}{2 \times 3 \times c}$$

$$\Rightarrow 3c = c^2 - 7$$

$$\Rightarrow c^2 - 3c - 7 = 0$$

$$102. \quad \text{We have, } b = \sqrt{3}, c = 1, \angle A = 30^\circ$$

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$\Rightarrow \frac{\sqrt{3}}{2} = \frac{(\sqrt{3})^2 + 1^2 - a^2}{2 \cdot \sqrt{3} \cdot 1}$$

$$\Rightarrow a = 1, b = \sqrt{3}, c = 1$$

\therefore b is the largest side. Therefore, the largest angle B is given by

$$\cos B = \frac{a^2 + c^2 - b^2}{2ac} = \frac{1 + 1 - 3}{2 \cdot 1 \cdot 1} = -\frac{1}{2} = \cos 120^\circ$$

$$\Rightarrow B = 120^\circ$$

$$103. \quad a^4 + b^4 + c^4 = 2c^2(a^2 + b^2)$$

$$\Rightarrow a^4 + b^4 + c^4 - 2a^2c^2 - 2b^2c^2 + 2a^2b^2 = 2a^2b^2$$

$$\Rightarrow (a^2 + b^2 - c^2)^2 = 2a^2b^2$$

$$\Rightarrow a^2 + b^2 - c^2 = \pm \sqrt{2}ab$$

$$\Rightarrow \frac{a^2 + b^2 - c^2}{2ab} = \pm \frac{\sqrt{2}ab}{2ab} = \pm \frac{1}{\sqrt{2}}$$

$$\Rightarrow \cos C = \cos 45^\circ \text{ or } \cos 135^\circ$$

$$\Rightarrow C = 45^\circ \text{ or } 135^\circ$$

$$104. \quad \text{We have, } b + c = 2a \quad \dots (i)$$

$$\cos 60^\circ = \frac{b^2 + c^2 - a^2}{2bc} = \frac{(b+c)^2 - 2bc - a^2}{2bc}$$

$$\Rightarrow \frac{1}{2} = \frac{4a^2 - 2bc - a^2}{2bc} \Rightarrow \frac{1}{2} = \frac{3a^2}{2bc} - 1$$

$$\Rightarrow \frac{3}{2} = \frac{3a^2}{2bc}$$

$$\Rightarrow bc = a^2 \quad \dots (ii)$$

From (i) and (ii), we get

$$b + c = 2\sqrt{b}\sqrt{c}$$

$$\Rightarrow (\sqrt{b} - \sqrt{c})^2 = 0 \Rightarrow b = c$$

From (i), $a = b = c$

\therefore ΔABC is equilateral.

$$105. \quad \frac{\sin A}{\sin C} = \frac{\sin A \cos B - \cos A \sin B}{\sin B \cos C - \cos B \sin C}$$

$$\Rightarrow \frac{a}{c} = \frac{a \cos B - b \cos A}{b \cos C - c \cos B}$$

$$\Rightarrow ab \cos C - ac \cos B = ac \cos B - bc \cos A$$

$$\Rightarrow ab \cos C + bc \cos A = 2ac \cos B$$

$$\Rightarrow \frac{a^2 + b^2 - c^2}{2} + \frac{b^2 + c^2 - a^2}{2} = \frac{c^2 + a^2 - b^2}{1}$$

$$\Rightarrow b^2 = c^2 + a^2 - b^2 \Rightarrow b^2 = \frac{c^2 + a^2}{2}$$

$\Rightarrow a^2, b^2, c^2$ are in A.P.

$$106. \quad \frac{2 \cos A}{a} + \frac{\cos B}{b} + \frac{2 \cos C}{c} = \frac{a}{bc} + \frac{b}{ca}$$

$$\Rightarrow \frac{2(b^2 + c^2 - a^2)}{2abc} + \frac{a^2 + c^2 - b^2}{2abc} + \frac{2(a^2 + b^2 - c^2)}{2abc}$$

$$= \frac{a}{bc} + \frac{b}{ca}$$

$$\Rightarrow \frac{3b^2 + c^2 + a^2}{2abc} = \frac{a}{bc} + \frac{b}{ca}$$

$$\Rightarrow \frac{3b}{2ac} + \frac{c}{2ab} + \frac{a}{2bc} = \frac{a}{bc} + \frac{b}{ca}$$

$$\Rightarrow b^2 + c^2 = a^2$$

Hence, $\angle A = 90^\circ$




107. $4 \sin A = 4 \sin B = 3 \sin C$

$$\therefore 4a = 4b = 3c \text{ or } a = b$$

$$\begin{aligned} \therefore \cos C &= \frac{a^2 + b^2 - c^2}{2ab} = \frac{a^2 + a^2 - \left(\frac{4a}{3}\right)^2}{2 \times a \times a} \\ &= \frac{2a^2 - \frac{16a^2}{9}}{2a^2} = \frac{a^2 - \frac{8a^2}{9}}{a^2} = 1 - \frac{8}{9} = \frac{1}{9} \end{aligned}$$

108.



From the figure,

$$\begin{aligned} \cos 120^\circ &= \frac{x^2 + x^2 - AB^2}{2x^2} \\ \Rightarrow \frac{2x^2 - AB^2}{2x^2} &= \frac{-1}{2} \\ \Rightarrow 4x^2 - 2AB^2 &= -2x^2 \\ \Rightarrow 3x^2 &= AB^2 \Rightarrow AB = x\sqrt{3} \\ \Rightarrow a^2 : b^2 : c^2 &= (2x)^2 : x^2 : (x\sqrt{3})^2 \\ &= 4x^2 : x^2 : 3x^2 = 4 : 1 : 3. \end{aligned}$$

109. $\frac{\sqrt{3}}{2} < \frac{b}{a} < 1$

$$\Rightarrow \frac{b}{a} < 1$$

$$\Rightarrow b < a$$

$$\Rightarrow c < b < a$$

$$\Rightarrow B = 60^\circ \quad \dots[\because \text{Angles are in A.P.}]$$

$$\text{Consider } \frac{\sqrt{3}}{2} < \frac{b}{a} < 1$$

$$\Rightarrow \frac{\sqrt{3}}{2} < \frac{b}{a}$$

$$\Rightarrow \sqrt{3}a < 2b$$

$$\Rightarrow 3a^2 < 4b^2$$

$$\Rightarrow 4b^2 - 3a^2 > 0$$

$$\text{Now, } b^2 = a^2 + c^2 - 2ac \cos 60^\circ$$

$$\Rightarrow c^2 - ac + (a^2 - b^2) = 0$$

$$\therefore c = \frac{a \pm \sqrt{4b^2 - 3a^2}}{2}$$

110. $\cot A, \cot B$ and $\cot C$ are in A. P.

$$\Rightarrow \cot A + \cot C = 2 \cot B$$

$$\Rightarrow \frac{\cos A}{\sin A} + \frac{\cos C}{\sin C} = \frac{2 \cos B}{\sin B}$$

$$\Rightarrow \frac{b^2 + c^2 - a^2}{2bc(ka)} + \frac{a^2 + b^2 - c^2}{2ab(kc)} = 2 \frac{a^2 + c^2 - b^2}{2ac(kb)}$$

$$\Rightarrow a^2 + c^2 = 2b^2$$

 Hence, a^2, b^2, c^2 , are in A. P.

111. $\frac{\sin 3B}{\sin B} = \frac{3 \sin B - 4 \sin^3 B}{\sin B} = 3 - 4 \sin^2 B$

$$= 3 - 4 + 4 \cos^2 B$$

$$= -1 + \frac{4(a^2 + c^2 - b^2)^2}{4(ac)^2}$$

$$= -1 + \frac{\left(\frac{a^2 + c^2}{2}\right)^2}{(ac)^2} \dots[\because 2b^2 = a^2 + c^2]$$

$$= -1 + \frac{(a^2 + c^2)^2}{4(ac)^2}$$

$$= \frac{(a^2 + c^2)^2 - 4a^2c^2}{4(ac)^2} = \left(\frac{c^2 - a^2}{2ac}\right)^2$$

112. $\cot B + \cot C - \cot A = \frac{\cos B}{\sin B} + \frac{\cos C}{\sin C} - \cot A$

$$= \frac{\sin C \cos B + \cos C \sin B}{\sin B \sin C} - \cot A$$

$$= \frac{\sin(B+C)}{\sin B \sin C} - \frac{\cos A}{\sin A}$$

$$= \frac{\sin^2 A - \sin B \sin C \cos A}{\sin A \sin B \sin C} = \frac{a^2 - bc \cos A}{(abc)}$$

$$= \frac{a^2 - bc \frac{(b^2 + c^2 - a^2)}{2bc}}{(abc)}$$

$$= \frac{3a^2 - b^2 - c^2}{2(abc)}$$

$$= \frac{3a^2 - b^2 - c^2}{2(abc)} = \frac{3a^2 - (b^2 + c^2)}{2(abc)}$$

$$\therefore \cot B + \cot C - \cot A = \frac{3a^2 - 3a^2}{2(abc)} = 0$$

$$\dots[\because b^2 + c^2 = 3a^2]$$


113. Largest side is $\sqrt{p^2 + pq + q^2}$. If largest angle is θ , then

$$\cos \theta = \frac{p^2 + q^2 - p^2 - pq - q^2}{2pq} = -\frac{1}{2} = \cos\left(\frac{2\pi}{3}\right)$$

$$\Rightarrow \theta = \frac{2\pi}{3}$$



114.




$$\begin{aligned} (AB)^2 &= (a - c)^2 + (b - d)^2 \\ (OA)^2 &= (a - 0)^2 + (b - 0)^2 = a^2 + b^2 \\ \text{and } (OB)^2 &= c^2 + d^2 \end{aligned}$$

Now from triangle AOB,

$$\begin{aligned} \cos \theta &= \frac{(OA)^2 + (OB)^2 - (AB)^2}{2OA \cdot OB} \\ &= \frac{a^2 + b^2 + c^2 + d^2 - \{(a - c)^2 + (b - d)^2\}}{2\sqrt{a^2 + b^2} \cdot \sqrt{c^2 + d^2}} \\ &= \frac{ac + bd}{\sqrt{(a^2 + b^2)(c^2 + d^2)}} \end{aligned}$$

115.



$$\begin{aligned} \cos C &= \frac{b^2 + a^2 - c^2}{2ba} \\ \Rightarrow \cos \frac{\pi}{6} &= \frac{(x^2 - 1)^2 + (x^2 + x + 1)^2 - (2x + 1)^2}{2(x^2 - 1)(x^2 + x + 1)} \\ \Rightarrow \frac{\sqrt{3}}{2} &= \frac{(x^2 - 1)^2 + (x^2 + 3x + 2)(x^2 - x)}{2(x^2 + x + 1)(x^2 - 1)} \\ \Rightarrow \frac{\sqrt{3}}{2} &= \frac{(x^2 - 1)^2 + (x + 1)(x + 2)x(x - 1)}{2(x^2 + x + 1)(x^2 - 1)} \\ \Rightarrow \sqrt{3} &= \frac{(x^2 - 1)^2 + x(x^2 - 1)(x + 2)}{(x^2 + x + 1)(x^2 - 1)} \\ \Rightarrow \sqrt{3} &= \frac{x^2 - 1 + x(x + 2)}{x^2 + x + 1} \\ \Rightarrow \sqrt{3}(x^2 + x + 1) &= 2x^2 + 2x - 1 \\ \Rightarrow (\sqrt{3} - 2)x^2 + (\sqrt{3} - 2)x + (\sqrt{3} + 1) &= 0 \end{aligned}$$


On solving, $x^2 + x - (3\sqrt{3} + 5) = 0$

$$\Rightarrow x = \sqrt{3} + 1, -(2 + \sqrt{3})$$

Since, x cannot be negative.

$$\therefore x = 1 + \sqrt{3}$$

116. Let the fourth side be of length d .



From the figure,

In $\triangle ADC$,

$$AC^2 = CD^2 + DA^2 - 2 \cdot CD \cdot DA \cdot \cos 120^\circ$$

....[By Cosine rule]

In $\triangle BAC$,

$$AC^2 = AB^2 + BC^2 - 2 \cdot AB \cdot BC \cdot \cos 60^\circ$$

....[By Cosine rule]

$$\therefore 3^2 + d^2 - 2 \times 3 \times d \cos 120^\circ = 2^2 + 5^2 - 2 \times 2 \times 5 \cos 60^\circ$$

$$\Rightarrow d^2 + 3d - 10 = 0 \Rightarrow d = -5 \text{ or } d = 2$$

$$\therefore d = 2$$

117. By sine rule,

$$\begin{aligned} \frac{a}{\sin A} &= \frac{b}{\sin B} \\ \Rightarrow \frac{a}{\sin 2B} &= \frac{b}{\sin B} \\ \Rightarrow \sin 2B &= \frac{a}{b} \sin B \\ \Rightarrow \frac{a}{2 \cos B} &= b \\ \Rightarrow \frac{a}{2 \left(\frac{a^2 + c^2 - b^2}{2ac} \right)} &= b \\ \Rightarrow a^2 c &= b(a^2 + c^2 - b^2) \\ \Rightarrow a^2(b - c) - (b + c)(b - c) &= 0 \\ \Rightarrow a^2 - b(b + c) &= 0 \\ \Rightarrow a^2 &= b^2 + bc \end{aligned}$$

Now, $a = \alpha + 1, b = \alpha - 1, c = \alpha$

$$\begin{aligned} \therefore (\alpha + 1)^2 &= (\alpha - 1)^2 + \alpha(\alpha - 1) \\ \Rightarrow \alpha^2 + 2\alpha + 1 &= \alpha^2 - 2\alpha + 1 + \alpha^2 - \alpha \\ \Rightarrow \alpha^2 - 5\alpha &= 0 \Rightarrow \alpha(\alpha - 5) = 0 \\ \Rightarrow \alpha &= 0, 5 \end{aligned}$$

118. $(b + c) \cos A + (c + a) \cos B + (a + b) \cos C$

$$\begin{aligned} &= (b \cos C + c \cos B) + (c \cos A + a \cos C) + (a \cos B + b \cos A) \\ &= a + b + c \end{aligned}$$

....[By projection rule]



$$\begin{aligned}
 119. \quad & \frac{\sin B \cos C + \cos B \sin C}{\sin A} + \frac{\cos B \sin C}{\sin A} \\
 &= \left(\frac{b}{a} \cos C + \frac{c}{a} \cos B \right) \\
 &= 1 \quad \dots [\text{By projection rule}]
 \end{aligned}$$

$$\begin{aligned}
 120. \quad & \frac{\cos C + \cos A}{c + a} + \frac{\cos B}{b} \\
 &= \frac{(b \cos C + c \cos B) + (b \cos A + a \cos B)}{b(c + a)} \\
 &= \frac{a + c}{b(c + a)} \quad \dots [\text{By projection rule}] \\
 &= \frac{1}{b}
 \end{aligned}$$

121. Since, a, b, c are in A. P.,

$$\therefore 2b = a + c$$

$$\begin{aligned}
 & a \cos^2 \left(\frac{C}{2} \right) + c \cos^2 \left(\frac{A}{2} \right) \\
 &= \frac{a(1 + \cos C)}{2} + \frac{c(1 + \cos A)}{2} \\
 &= \frac{a + c + a \cos C + c \cos A}{2} \\
 &= \frac{a + c + b}{2} \\
 &= \frac{2b + b}{2} \\
 &= \frac{3b}{2}
 \end{aligned}$$

$$\begin{aligned}
 122. \quad & \sin \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{bc}} \\
 & \Rightarrow bc \sin^2 \frac{A}{2} = (s-b)(s-c) \\
 & \Rightarrow x = bc
 \end{aligned}$$

$$\begin{aligned}
 123. \quad & \sin \frac{A}{2} \cdot \sin \frac{C}{2} = \sin \frac{B}{2} \\
 \therefore & \sqrt{\frac{(s-b)(s-c)}{bc}} \times \sqrt{\frac{(s-a)(s-b)}{ab}} = \sqrt{\frac{(s-a)(s-c)}{ac}} \\
 & \Rightarrow \frac{(s-b)}{b} \sqrt{\frac{(s-a)(s-c)}{ac}} = \sqrt{\frac{(s-a)(s-c)}{ac}} \\
 & \Rightarrow s - b = b \Rightarrow s = 2b
 \end{aligned}$$

$$\begin{aligned}
 124. \quad & s = \frac{a + b + c}{2} = \frac{16 + 24 + 20}{2} = 30 \\
 & \cos \left(\frac{B}{2} \right) = \sqrt{\frac{s(s-b)}{ac}} \\
 &= \sqrt{\frac{30(30-24)}{16 \times 20}} = \sqrt{\frac{9}{16}} = \frac{3}{4}
 \end{aligned}$$

$$\begin{aligned}
 125. \quad & (a-b)^2 \cos^2 \frac{C}{2} + (a+b)^2 \sin^2 \frac{C}{2} \\
 &= (a^2 + b^2 - 2ab) \cos^2 \frac{C}{2} + (a^2 + b^2 + 2ab) \sin^2 \frac{C}{2} \\
 &= (a^2 + b^2) \left[\cos^2 \frac{C}{2} + \sin^2 \frac{C}{2} \right] \\
 & \quad - 2ab \left[\cos^2 \frac{C}{2} - \sin^2 \frac{C}{2} \right] \\
 &= a^2 + b^2 - 2ab \cos C \\
 &= c^2 \quad \dots [\text{By cosine rule}]
 \end{aligned}$$

$$\begin{aligned}
 126. \quad & a, b, c \text{ are in A.P.} \Rightarrow 2b = a + c \\
 & \Rightarrow 2s - 2b = 2s - (a + c) \\
 & \Rightarrow 2(s - b) = s - a + s - c \\
 & \Rightarrow 2s(s - b) = s(s - a) + s(s - c) \\
 & \Rightarrow 4s^2(s - b)^2 = s^2(s - a)^2 + s^2(s - c)^2 \\
 & \quad + 2s^2(s - a)(s - c) \\
 & \Rightarrow \frac{4s^2(s - b)^2}{s(s - a)(s - b)(s - c)} = \frac{s^2(s - a)^2}{s(s - a)(s - b)(s - c)} \\
 & \quad + \frac{s^2(s - c)^2}{s(s - a)(s - b)(s - c)} + \frac{2s^2(s - a)(s - c)}{s(s - a)(s - b)(s - c)}
 \end{aligned}$$

Taking square root on both sides, we get

$$\begin{aligned}
 2 \sqrt{\frac{s(s-b)}{(s-a)(s-c)}} &= \sqrt{\frac{s(s-a)}{(s-b)(s-c)}} \\
 & \quad + \sqrt{\frac{s(s-c)}{(s-b)(s-a)}} \\
 & \Rightarrow 2 \cot \frac{B}{2} = \cot \frac{A}{2} + \cot \frac{C}{2}
 \end{aligned}$$

$$\begin{aligned}
 127. \quad & \tan \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}, \tan \frac{B}{2} = \sqrt{\frac{(s-a)(s-c)}{s(s-b)}} \\
 & (a + b + c) \left(\tan \frac{A}{2} + \tan \frac{B}{2} \right) \\
 &= (a + b + c) \left[\sqrt{\frac{(s-b)(s-c)}{s(s-a)}} + \sqrt{\frac{(s-a)(s-c)}{s(s-b)}} \right] \\
 &= 2s \sqrt{\frac{s-c}{s}} \left[\sqrt{\frac{s-b}{s-a}} + \sqrt{\frac{s-a}{s-b}} \right]
 \end{aligned}$$



$$\begin{aligned}
 &= 2s \sqrt{\frac{s-c}{s}} \left[\frac{s-b+s-a}{\sqrt{(s-a)(s-b)}} \right] \\
 &= 2\sqrt{s(s-c)} \left[\frac{2s-a-b}{\sqrt{(s-a)(s-b)}} \right] \\
 &= 2c \sqrt{\frac{s(s-c)}{(s-a)(s-b)}} \quad \dots [\because 2s-a-b=c] \\
 &= 2c \cot \frac{C}{2}
 \end{aligned}$$

$$\therefore \tan \frac{A}{2} + \tan \frac{B}{2} = \frac{2c \cot \frac{C}{2}}{a+b+c}$$

$$\begin{aligned}
 128. \quad &\left\{ \cot \frac{A}{2} + \cot \frac{B}{2} \right\} \left\{ a \sin^2 \frac{B}{2} + b \sin^2 \frac{A}{2} \right\} \\
 &= \left\{ \frac{\cos \frac{C}{2}}{\sin \frac{A}{2} \sin \frac{B}{2}} \right\} \left\{ a \sin^2 \frac{B}{2} + b \sin^2 \frac{A}{2} \right\} \\
 &= \left\{ \cos \frac{C}{2} \right\} \left\{ a \frac{\sin \frac{B}{2}}{\sin \frac{A}{2}} + b \frac{\sin \frac{A}{2}}{\sin \frac{B}{2}} \right\} \\
 &= \sqrt{\frac{s(s-c)}{ab}} \left\{ a \frac{\sqrt{\frac{(s-a)(s-c)}{ac}}}{\sqrt{\frac{(s-b)(s-c)}{bc}}} + b \frac{\sqrt{\frac{(s-b)(s-c)}{bc}}}{\sqrt{\frac{(s-a)(s-c)}{ac}}} \right\} \\
 &= \sqrt{\frac{s(s-c)}{ab}} \left\{ \sqrt{\frac{(s-a)}{(s-b)}} ab + \sqrt{\frac{(s-b)}{(s-a)}} ab \right\} \\
 &= \sqrt{s(s-c)} \left\{ \frac{s-a+s-b}{\sqrt{(s-a)(s-b)}} \right\} \\
 &= \sqrt{s(s-c)} \left\{ \frac{2s-a-b}{\sqrt{(s-a)(s-b)}} \right\} \\
 &= c \sqrt{\frac{s(s-c)}{(s-a)(s-b)}} = c \cot \frac{C}{2}
 \end{aligned}$$

Alternate Method :

Let $a = 1$, $b = \sqrt{3}$, $c = 2$ and $A = 30^\circ$,
 $B = 60^\circ$, $C = 90^\circ$.

Hence, the given expression is equal to 2,
 which is given by option (D).

129. Let $\cot \frac{A}{2}$, $\cot \frac{B}{2}$ and $\cot \frac{C}{2}$ be in A.P.

$$\text{Then, } 2 \cot \frac{B}{2} = \cot \frac{C}{2} + \cot \frac{A}{2}$$

\therefore we need to prove that

$$2 \sqrt{\frac{s(s-b)}{(s-a)(s-c)}} = \sqrt{\frac{s(s-c)}{(s-a)(s-b)}} + \sqrt{\frac{s(s-a)}{(s-b)(s-c)}}$$

$$\begin{aligned}
 \text{R.H.S.} &= \sqrt{\frac{s}{(s-b)}} \left(\sqrt{\frac{(s-c)}{(s-a)}} + \sqrt{\frac{(s-a)}{(s-c)}} \right) \\
 &= \sqrt{\frac{s}{s-b}} \left(\frac{s-c+s-a}{\sqrt{(s-a)(s-c)}} \right) \\
 &= \sqrt{\frac{s}{s-b}} \left(\frac{2s-a-c}{\sqrt{(s-a)(s-c)}} \right) \\
 &= 2 \sqrt{\frac{s}{(s-b)}} \sqrt{\frac{(s-b)^2}{(s-a)(s-c)}} \\
 &\quad \dots \left[\begin{array}{l} \because 2b = a + c \\ \Rightarrow 2s - 2b = 2s - (a + c) \\ \Rightarrow 2(s-b) = 2s - a - c \end{array} \right] \\
 &= 2 \sqrt{\frac{s(s-b)}{(s-a)(s-c)}} = \text{L.H.S.}
 \end{aligned}$$

130. Δ is right angled, $\angle C = 90^\circ$

$$\therefore \Delta = \frac{1}{2} ab \sin 90^\circ = \frac{1}{2} ab$$

$$\therefore 4\Delta^2 = 4 \left(\frac{1}{2} ab \right)^2 = a^2 b^2$$

$$131. \Delta = \frac{1}{2} bc \sin A \Rightarrow 9 = \frac{1}{2} \cdot 36 \sin A$$

$$\Rightarrow \sin A = \frac{1}{2} \Rightarrow A = 30^\circ$$

132. We have, $a = 1$, $b = 2$, $\angle C = 60^\circ$

$$\begin{aligned}
 \text{Area of triangle} &= \frac{1}{2} ab \sin C \\
 &= \frac{1}{2} (1)(2) \sin 60^\circ = \frac{\sqrt{3}}{2}
 \end{aligned}$$

$$133. \Delta = \frac{1}{2} ab \sin C$$

$$= \frac{1}{2} \times 1 \times 2 \times \sin 60^\circ$$

$$= \frac{\sqrt{3}}{2}$$



$$\cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

$$\Rightarrow \cos 60^\circ = \frac{1 + 4 - c^2}{2(1)(2)}$$

$$\Rightarrow \frac{1}{2} = \frac{5 - c^2}{4}$$

$$\Rightarrow c^2 = 3$$

$$\begin{aligned} \therefore 4\Delta^2 + c^2 &= 4 \left(\frac{\sqrt{3}}{2} \right)^2 + 3 \\ &= 3 + 3 \\ &= 6 \end{aligned}$$

$$134. \quad a^2 \sin 2C + c^2 \sin 2A$$

$$= a^2(2 \sin C \cos C) + c^2(2 \sin A \cos A)$$

$$= 2a^2 \left(\frac{2\Delta}{ab} \cos C \right) + 2c^2 \left(\frac{2\Delta}{bc} \cos A \right)$$

$$\left[\begin{array}{l} \because \Delta = \frac{1}{2} ab \sin C = \frac{1}{2} bc \sin A, \\ \dots \\ \therefore \sin C = \frac{2\Delta}{ab}, \sin A = \frac{2\Delta}{bc} \end{array} \right]$$

$$= 4\Delta \left\{ \frac{a \cos C + c \cos A}{b} \right\} = 4\Delta \left(\frac{b}{b} \right) = 4\Delta$$

$$135. \quad \Delta = a^2 - (b - c)^2$$

$$= 2bc - (b^2 + c^2 - a^2)$$

$$= 2bc - 2bc \left(\frac{b^2 + c^2 - a^2}{2bc} \right) = 2bc(1 - \cos A)$$

$$\therefore \Delta = 2bc \cdot 2 \sin^2 \frac{A}{2} \quad \dots(i)$$

$$\text{Also, } \Delta = \frac{1}{2} bc \sin A$$

$$= \frac{1}{2} bc \cdot 2 \sin \frac{A}{2} \cos \frac{A}{2}$$

$$\therefore \Delta = bc \cdot \sin \frac{A}{2} \cos \frac{A}{2} \quad \dots(ii)$$


$$\therefore \tan \frac{A}{2} = \frac{1}{4} \quad \dots[\text{From (i) and (ii)}]$$

$$136. \quad a = 2 = QR,$$

$$b = \frac{7}{2} = PR,$$

$$c = \frac{5}{2} = PQ$$

$$s = \frac{a + b + c}{2} = \frac{8}{2} = 4$$



$$\frac{2 \sin P - 2 \sin P \cos P}{2 \sin P + 2 \sin P \cos P} = \frac{2 \sin P(1 - \cos P)}{2 \sin P(1 + \cos P)}$$

$$= \frac{1 - \cos P}{1 + \cos P} = \frac{2 \sin^2 \frac{P}{2}}{2 \cos^2 \frac{P}{2}} = \tan^2 \frac{P}{2}$$

$$= \frac{(s-b)(s-c)}{s(s-a)} = \frac{(s-b)^2(s-c)^2}{\Delta^2}$$

$$= \frac{\left(4 - \frac{7}{2}\right)^2 \left(4 - \frac{5}{2}\right)^2}{\Delta^2} = \left(\frac{3}{4\Delta}\right)^2$$

$$137. \quad \tan \left(\frac{B-C}{2} \right) = \frac{b-c}{b+c} \cot \frac{A}{2}$$

$$\Rightarrow \tan \left(\frac{90^\circ}{2} \right) = \frac{\sqrt{3}-1}{\sqrt{3}+1} \cot \frac{A}{2}$$

$$\Rightarrow \tan \left(\frac{A}{2} \right) = \frac{\sqrt{3}-1}{\sqrt{3}+1} = \frac{3+1-2\sqrt{3}}{2} = 2 - \sqrt{3}$$

$$\Rightarrow \frac{A}{2} = 15^\circ \Rightarrow A = 30^\circ$$

$$139. \quad \sin^{-1} \left(\frac{2x+1}{3} \right) \text{ is defined for}$$

$$-1 \leq \frac{2x+1}{3} \leq 1$$

$$\Rightarrow -3 \leq 2x+1 \leq 3 \Rightarrow -4 \leq 2x \leq 2$$

$$\Rightarrow -2 \leq x \leq 1$$

$$140. \quad \text{Given, } \sin^{-1} x = 2 \sin^{-1} 2a$$

$$\text{Since, } -\frac{\pi}{2} \leq \sin^{-1} x \leq \frac{\pi}{2}$$

$$\Rightarrow -\frac{\pi}{2} \leq 2 \sin^{-1} 2a \leq \frac{\pi}{2}$$

$$\Rightarrow -\frac{\pi}{4} \leq \sin^{-1} 2a \leq \frac{\pi}{4}$$

$$\Rightarrow \sin \left(-\frac{\pi}{4} \right) \leq \sin (\sin^{-1} 2a) \leq \sin \frac{\pi}{4}$$

$$\Rightarrow \frac{-1}{\sqrt{2}} \leq 2a \leq \frac{1}{\sqrt{2}}$$

$$\Rightarrow \frac{-1}{2\sqrt{2}} \leq a \leq \frac{1}{2\sqrt{2}} \quad \text{i.e., } |a| \leq \frac{1}{2\sqrt{2}}$$

$$141. \quad \text{Let } \cot^{-1} \left(\frac{1}{2} \right) = \theta$$

$$\Rightarrow \cot \theta = \frac{1}{2}$$

$$\therefore \sin \theta = \frac{2}{\sqrt{5}}$$



Let $\cos^{-1} x = \phi$

$$\Rightarrow x = \cos \phi$$

$$\text{Now, } \tan(\cos^{-1} x) = \sin\left(\cot^{-1}\left(\frac{1}{2}\right)\right)$$

$$\Rightarrow \tan \phi = \sin \theta$$

$$\Rightarrow \tan \phi = \frac{2}{\sqrt{5}}$$

$$\therefore x = \cos \phi = \frac{\sqrt{5}}{3}$$

142. Let $\tan^{-1} 2 = \alpha \Rightarrow \tan \alpha = 2$

and $\cot^{-1} 3 = \beta \Rightarrow \cot \beta = 3$

$$\begin{aligned} \therefore \sec^2(\tan^{-1} 2) + \operatorname{cosec}^2(\cot^{-1} 3) \\ = \sec^2 \alpha + \operatorname{cosec}^2 \beta = 1 + \tan^2 \alpha + 1 + \cot^2 \beta \\ = 2 + (2)^2 + (3)^2 = 15 \end{aligned}$$

143. $\sin^{-1} x + \sin^{-1} y + \sin^{-1} z = \frac{\pi}{2}$

Put $\sin^{-1} x = \alpha$, $\sin^{-1} y = \beta$, $\sin^{-1} z = \gamma$

$$\therefore \alpha + \beta + \gamma = \frac{\pi}{2}$$

$$\Rightarrow \alpha + \beta = \frac{\pi}{2} - \gamma \Rightarrow \cos(\alpha + \beta) = \cos\left(\frac{\pi}{2} - \gamma\right)$$

$$\Rightarrow \cos \alpha \cos \beta - \sin \alpha \sin \beta = \sin \gamma \quad \dots(i)$$

and, we have

$$\sin \alpha = x \Rightarrow \cos \alpha = \sqrt{1-x^2}$$

$$\text{Similarly, } \cos \beta = \sqrt{1-y^2}$$

\therefore From (i), we get

$$\sqrt{1-x^2} \cdot \sqrt{1-y^2} = xy + z$$

Squaring on both sides, we get

$$x^2 + y^2 + z^2 + 2xyz = 1$$

144. Let $\sin^{-1} a = A$,

$$\sin^{-1} b = B,$$

$$\sin^{-1} c = C$$

$$\therefore \sin A = a, \sin B = b, \sin C = c$$

and $A + B + C = \pi$ then

$$\sin 2A + \sin 2B + \sin 2C$$

$$= 4 \sin A \sin B \sin C$$

$$\Rightarrow \sin A \cos A + \sin B \cos B + \sin C \cos C$$

$$= 2 \sin A \sin B \sin C$$

$$\Rightarrow \sin A \sqrt{1-\sin^2 A} + \sin B \sqrt{1-\sin^2 B}$$

$$+ \sin C \sqrt{1-\sin^2 C} = 2 \sin A \sin B \sin C$$

$$\Rightarrow a \sqrt{1-a^2} + b \sqrt{1-b^2} + c \sqrt{1-c^2} = 2abc$$

145. Given $\cos^{-1} x + \cos^{-1} y + \cos^{-1} z = 3\pi$

$$\therefore 0 \leq \cos^{-1} x \leq \pi$$

$$\therefore 0 \leq \cos^{-1} y \leq \pi \text{ and } 0 \leq \cos^{-1} z \leq \pi$$

$$\text{Here, } \cos^{-1} x = \cos^{-1} y = \cos^{-1} z = \pi$$

$$\Rightarrow x = y = z = \cos \pi = -1$$

$$\begin{aligned} \therefore xy + yz + zx &= (-1)(-1) + (-1)(-1) + (-1)(-1) \\ &= 1 + 1 + 1 = 3 \end{aligned}$$

146. Let $\alpha = \cos^{-1} \sqrt{p}$, $\beta = \cos^{-1} \sqrt{1-p}$

and $\gamma = \cos^{-1} \sqrt{1-q}$ $\therefore \cos \alpha = \sqrt{p}$,

$$\cos \beta = \sqrt{1-p}$$

$$\text{and } \cos \gamma = \sqrt{1-q}$$

$$\therefore \sin \alpha = \sqrt{1-p}, \sin \beta = \sqrt{p} \text{ and } \sin \gamma = \sqrt{q}$$

The given equation can be written as

$$\alpha + \beta + \gamma = \frac{3\pi}{4} \Rightarrow \alpha + \beta = \frac{3\pi}{4} - \gamma$$

$$\Rightarrow \cos(\alpha + \beta) = \cos\left(\frac{3\pi}{4} - \gamma\right)$$

$$\Rightarrow \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$= \cos\left\{\pi - \left(\frac{\pi}{4} + \gamma\right)\right\} = -\cos\left(\frac{\pi}{4} + \gamma\right)$$

$$\Rightarrow \sqrt{p} \sqrt{1-p} - \sqrt{1-p} \sqrt{p}$$

$$= -\left(\frac{1}{\sqrt{2}} \sqrt{1-q} - \frac{1}{\sqrt{2}} \sqrt{q}\right)$$

$$\Rightarrow 0 = \sqrt{1-q} - \sqrt{q} \Rightarrow 1-q = q \Rightarrow q = \frac{1}{2}$$

147. $\tan^{-1} \sqrt{x(x+1)} + \sin^{-1} \sqrt{x^2+x+1} = \frac{\pi}{2}$

$\tan^{-1} \sqrt{x(x+1)}$ is defined when

$$x(x+1) \geq 0 \quad \dots(i)$$

$\sin^{-1} \sqrt{x^2+x+1}$ is defined when

$$x(x+1) + 1 \leq 1 \text{ or } x(x+1) \leq 0 \quad \dots(ii)$$

From (i) and (ii),

$$x(x+1) = 0 \text{ or } x = 0 \text{ and } -1.$$

Hence, number of solutions is 2.

148. Let $\cot^{-1} x = \theta \Rightarrow x = \cot \theta$

$$\text{Now } \operatorname{cosec} \theta = \sqrt{1+\cot^2 \theta} = \sqrt{1+x^2}$$

$$\therefore \sin \theta = \frac{1}{\operatorname{cosec} \theta} = \frac{1}{\sqrt{1+x^2}}$$



$$\Rightarrow \theta = \sin^{-1} \frac{1}{\sqrt{1+x^2}}$$

$$\begin{aligned} \therefore \sin(\cot^{-1} x) &= \sin\left(\sin^{-1} \frac{1}{\sqrt{1+x^2}}\right) \\ &= \frac{1}{\sqrt{1+x^2}} \\ &= (1+x^2)^{-\frac{1}{2}} \end{aligned}$$

$$149. \sin^{-1}\left(\frac{2a}{1+a^2}\right) + \sin^{-1}\left(\frac{2b}{1+b^2}\right) = 2\tan^{-1} x$$

Putting $a = \tan \theta$ and $b = \tan \phi$, we get

$$\sin^{-1}\left(\frac{2\tan\theta}{1+\tan^2\theta}\right) + \sin^{-1}\left(\frac{2\tan\phi}{1+\tan^2\phi}\right) = 2\tan^{-1} x$$

$$\Rightarrow \sin^{-1}[\sin(2\theta)] + \sin^{-1}[\sin(2\phi)] = 2\tan^{-1} x$$

$$\Rightarrow 2(\theta + \phi) = 2\tan^{-1} x$$

$$\Rightarrow x = \tan(\theta + \phi)$$

$$\Rightarrow x = \frac{\tan\theta + \tan\phi}{1 - \tan\theta \tan\phi}$$

Resubstituting the values of a and b , we get

$$x = \frac{a+b}{1-ab}$$

$$150. \cos(2\tan^{-1} x) = \frac{1}{2}$$

$$\Rightarrow 2\tan^{-1} x = \frac{\pi}{3}, \frac{-\pi}{3}$$

$$\Rightarrow \tan^{-1} x = \frac{\pi}{6}, \frac{-\pi}{6}$$

$$\Rightarrow x = \frac{1}{\sqrt{3}}, \frac{-1}{\sqrt{3}}$$

$$151. \cos\left[\cot^{-1}\left(\frac{1}{2}\right)\right] = \cos(\tan^{-1} 2)$$

$$= \cos\left[\cos^{-1}\left(\frac{1}{\sqrt{1+(2)^2}}\right)\right] = \frac{1}{\sqrt{5}}$$

$$\text{and } \cot(\cos^{-1} x) = \cot\left[\tan^{-1} \frac{\sqrt{1-x^2}}{x}\right]$$

$$= \cot\left(\cot^{-1} \frac{x}{\sqrt{1-x^2}}\right) = \frac{x}{\sqrt{1-x^2}}$$

$$\text{Given, } \cos\left[\cot^{-1}\left(\frac{1}{2}\right)\right] = \cot(\cos^{-1} x)$$

$$\Rightarrow \frac{1}{\sqrt{5}} = \frac{x}{\sqrt{1-x^2}} \Rightarrow \frac{1}{5} = \frac{x^2}{1-x^2}$$

$$\Rightarrow 6x^2 = 1$$

$$\Rightarrow x = \pm \frac{1}{\sqrt{6}}$$

$$152. \text{ Let } \cos^{-1} x = \theta \Rightarrow x = \cos \theta \Rightarrow \sec \theta = \frac{1}{x}$$

$$\Rightarrow \tan \theta = \sqrt{\sec^2 \theta - 1} = \sqrt{\frac{1}{x^2} - 1} = \frac{1}{x} \sqrt{1-x^2}$$

Now,

$$\sin[\cot^{-1}(\tan \theta)] = \sin\left[\cot^{-1}\left(\frac{1}{x} \sqrt{1-x^2}\right)\right]$$

Again, putting $x = \sin \theta$

$$\begin{aligned} \therefore \sin \cot^{-1}\left(\frac{1}{x} \sqrt{1-x^2}\right) &= \sin \cot^{-1}\left(\frac{\sqrt{1-\sin^2 \theta}}{\sin \theta}\right) \\ &= \sin \cot^{-1}(\cot \theta) \\ &= \sin \theta = x \end{aligned}$$

$$153. \cos^{-1} x + \cos^{-1}(2x) = -\pi$$

$$\Rightarrow \cos^{-1} 2x = -\pi - \cos^{-1} x$$

$$\Rightarrow 2x = \cos(\pi + \cos^{-1} x)$$

$$\Rightarrow 2x = (\cos \pi) \cos(\cos^{-1} x) - (\sin \pi) \sin(\cos^{-1} x)$$

$$\Rightarrow 2x = -x \Rightarrow x = 0$$

But $x = 0$ does not satisfy the given equation.

\therefore No solution will exist.

$$154. \cos \frac{7\pi}{6} = \cos\left(2\pi - \frac{7\pi}{6}\right) = \cos \frac{5\pi}{6}$$

$$\Rightarrow \cos \frac{7\pi}{6} = \cos \frac{5\pi}{6}$$

$$\therefore \cos^{-1}\left(\cos \frac{7\pi}{6}\right) = \cos^{-1}\left(\cos \frac{5\pi}{6}\right) = \frac{5\pi}{6}$$

$$155. \cos \frac{53\pi}{5} = \cos\left(\frac{50\pi}{5} + \frac{3\pi}{5}\right)$$

$$= \cos\left(10\pi + \frac{3\pi}{5}\right)$$

$$= \cos \frac{3\pi}{5}$$

$$= \sin\left(\frac{\pi}{2} - \frac{3\pi}{5}\right)$$

$$= \sin\left(\frac{-\pi}{10}\right)$$

$$\therefore \sin^{-1}\left(\cos \frac{53\pi}{5}\right) = \sin^{-1}\left(\sin \frac{-\pi}{10}\right) = \frac{-\pi}{10}$$



$$\begin{aligned}
 156. \quad & \tan^{-1}(\cot x) + \cot^{-1}(\tan x) \\
 &= \tan^{-1} \left[\tan \left(\frac{\pi}{2} - x \right) \right] + \cot^{-1} \left[\cot \left(\frac{\pi}{2} - x \right) \right] \\
 &= \frac{\pi}{2} - x + \frac{\pi}{2} - x \\
 &= \pi - 2x
 \end{aligned}$$

$$\begin{aligned}
 157. \quad & \text{Let } \theta = \cos^{-1} \frac{2}{\sqrt{5}} \Rightarrow \cos \theta = \frac{2}{\sqrt{5}} \\
 & \tan \left(\frac{1}{2} \cos^{-1} \frac{2}{\sqrt{5}} \right) = \tan \frac{\theta}{2} \\
 &= \frac{\sqrt{1 - \cos \theta}}{\sqrt{1 + \cos \theta}} \\
 &= \frac{\sqrt{1 - \frac{2}{\sqrt{5}}}}{\sqrt{1 + \frac{2}{\sqrt{5}}}} \\
 &= \frac{\sqrt{\sqrt{5} - 2}}{\sqrt{\sqrt{5} + 2}} \\
 &= \frac{(\sqrt{5} - 2)}{\sqrt{(\sqrt{5} + 2)(\sqrt{5} - 2)}} \\
 &= \frac{\sqrt{(\sqrt{5} - 2)^2}}{\sqrt{5 - 2}} \\
 &= \sqrt{5} - 2
 \end{aligned}$$

158. Putting

$a = \tan \theta$, $b = \tan \phi$ and $x = \tan \psi$ in the given expression, we get

$$\sin^{-1}(\sin 2\theta) - \cos^{-1}(\cos 2\phi) = \tan^{-1}(\tan 2\psi)$$

$$\Rightarrow 2\theta - 2\phi = 2\psi \Rightarrow \theta - \phi = \psi$$

Taking 'tan' on both sides, we get

$$\tan(\theta - \phi) = \tan \psi$$

$$\Rightarrow \frac{\tan \theta - \tan \phi}{1 + \tan \theta \tan \phi} = \tan \psi$$

$$\Rightarrow \frac{a - b}{1 + ab} = x$$

159. Putting $x = \tan \theta$, we get

$$\begin{aligned}
 & \sin \left[\tan^{-1} \left(\frac{1-x^2}{2x} \right) + \cos^{-1} \left(\frac{1-x^2}{1+x^2} \right) \right] \\
 &= \sin \left[\tan^{-1} \left(\frac{1-\tan^2 \theta}{2 \tan \theta} \right) + \cos^{-1} \left(\frac{1-\tan^2 \theta}{1+\tan^2 \theta} \right) \right] \\
 &= \sin [\tan^{-1}(\cot 2\theta) + \cos^{-1}(\cos 2\theta)]
 \end{aligned}$$

$$= \sin [\tan^{-1} \{ \tan \left(\frac{\pi}{2} - 2\theta \right) \} + \cos^{-1}(\cos 2\theta)]$$

$$= \sin \frac{\pi}{2} = 1$$

$$\begin{aligned}
 160. \quad & \cot^{-1} \left[\frac{\sqrt{1-\sin x} + \sqrt{1+\sin x}}{\sqrt{1-\sin x} - \sqrt{1+\sin x}} \right] \\
 &= \cot^{-1} \left[\frac{(\sqrt{1-\sin x} + \sqrt{1+\sin x})}{(\sqrt{1-\sin x} - \sqrt{1+\sin x})} \right. \\
 & \quad \left. \times \frac{(\sqrt{1-\sin x} + \sqrt{1+\sin x})}{(\sqrt{1-\sin x} + \sqrt{1+\sin x})} \right] \\
 &= \cot^{-1} \left[\frac{(1-\sin x) + (1+\sin x) + 2\sqrt{1-\sin^2 x}}{(1-\sin x) - (1+\sin x)} \right] \\
 &= \cot^{-1} \left[\frac{2(1+\cos x)}{-2\sin x} \right] \\
 &= \cot^{-1} \left[-\frac{2\cos^2 \left(\frac{x}{2} \right)}{2\sin \left(\frac{x}{2} \right) \cos \left(\frac{x}{2} \right)} \right] \\
 &= \cot^{-1} \left(-\cot \frac{x}{2} \right) = \cot^{-1} \left[\cot \left(\pi - \frac{x}{2} \right) \right] \\
 &= \pi - \frac{x}{2}
 \end{aligned}$$

161. Putting $x = \tan \theta$ in the given equation, we get

$$\begin{aligned}
 \cot^{-1} \left(\frac{\sqrt{1+x^2} - 1}{x} \right) &= \cot^{-1} \left(\frac{\sqrt{1+\tan^2 \theta} - 1}{\tan \theta} \right) \\
 &= \cot^{-1} \left(\frac{\sec \theta - 1}{\tan \theta} \right) \\
 &= \cot^{-1} \left(\frac{1 - \cos \theta}{\sin \theta} \right) \\
 &= \cot^{-1} \left(\frac{2\sin^2 \frac{\theta}{2}}{2\sin \frac{\theta}{2} \cos \frac{\theta}{2}} \right) \\
 &= \cot^{-1} \left(\tan \frac{\theta}{2} \right) \\
 &= \cot^{-1} \left[\cot \left(\frac{\pi}{2} - \frac{\theta}{2} \right) \right] \\
 &= \frac{\pi}{2} - \frac{\theta}{2} = \frac{\pi}{2} - \frac{\tan^{-1} x}{2}
 \end{aligned}$$



$$\begin{aligned}
 162. \quad \tan^{-1} \frac{1-x}{1+x} &= \frac{1}{2} \tan^{-1} x \\
 \Rightarrow \tan^{-1} \left[\frac{1-\tan \theta}{1+\tan \theta} \right] &= \frac{1}{2} \cdot \theta \quad \dots [\text{Put } x = \tan \theta] \\
 \Rightarrow \tan^{-1} \left[\frac{\tan \frac{\pi}{4} - \tan \theta}{1 + \tan \frac{\pi}{4} \tan \theta} \right] &= \frac{\theta}{2} \\
 \Rightarrow \tan^{-1} \tan \left(\frac{\pi}{4} - \theta \right) &= \frac{\theta}{2} \Rightarrow \frac{\pi}{4} - \theta = \frac{\theta}{2} \\
 \Rightarrow \theta = \frac{\pi}{6} \Rightarrow \tan^{-1} x &= \frac{\pi}{6} \Rightarrow x = \tan \frac{\pi}{6} = \frac{1}{\sqrt{3}}
 \end{aligned}$$

$$\begin{aligned}
 163. \quad \tan \left[\sin^{-1} \left\{ \frac{x}{\sqrt{2}} + \frac{\sqrt{1-x^2}}{\sqrt{2}} \right\} - \sin^{-1} x \right] \\
 = \tan \left[\sin^{-1} \left\{ \frac{x + \sqrt{1-x^2}}{\sqrt{2}} \right\} - \sin^{-1} x \right] \\
 = \tan \left[\sin^{-1} \left\{ \frac{\sin \theta + \cos \theta}{\sqrt{2}} \right\} - \theta \right] \\
 \dots \left[\begin{array}{l} \text{Put } \sin^{-1} x = \theta \\ \Rightarrow x = \sin \theta \end{array} \right] \\
 = \tan \left[\sin^{-1} \left[\sin \left(\theta + \frac{\pi}{4} \right) \right] - \theta \right] \\
 = \tan \left[\theta + \frac{\pi}{4} - \theta \right] = \tan \frac{\pi}{4} = 1
 \end{aligned}$$

$$\begin{aligned}
 164. \quad \sec^{-1} [\sec (-30^\circ)] \\
 = \sec^{-1} (\sec 30^\circ) \quad \dots [\because \sec (-\theta) = \sec \theta] \\
 = 30^\circ
 \end{aligned}$$

$$\begin{aligned}
 165. \quad \cos^{-1} \left[\cot \left(\frac{\pi}{2} \right) \right] + \cos^{-1} \left[\sin \left(\frac{2\pi}{3} \right) \right] \\
 = \cos^{-1} (0) + \cos^{-1} \left(\frac{\sqrt{3}}{2} \right) \\
 = \frac{\pi}{2} + \cos^{-1} \left(\cos \frac{\pi}{6} \right) \\
 = \frac{\pi}{2} + \frac{\pi}{6} = \frac{2\pi}{3}
 \end{aligned}$$

$$166. \quad \frac{\tan^{-1}(\sqrt{3}) - \sec^{-1}(-2)}{\operatorname{cosec}^{-1}(-\sqrt{2}) + \cos^{-1}\left(-\frac{1}{2}\right)} = \frac{\frac{\pi}{3} - \frac{2\pi}{3}}{\frac{-\pi}{4} + \frac{2\pi}{3}} = \frac{-4}{5}$$

$$\begin{aligned}
 167. \quad \text{Given,} \\
 \sec^{-1} x = \operatorname{cosec}^{-1} y \\
 \Rightarrow \cos^{-1} \left(\frac{1}{x} \right) = \sin^{-1} \left(\frac{1}{y} \right) \\
 \Rightarrow \cos^{-1} \left(\frac{1}{x} \right) = \frac{\pi}{2} - \cos^{-1} \left(\frac{1}{y} \right) \\
 \Rightarrow \cos^{-1} \left(\frac{1}{x} \right) + \cos^{-1} \left(\frac{1}{y} \right) = \frac{\pi}{2}
 \end{aligned}$$

$$\begin{aligned}
 168. \quad x^2 + 5|x| - 6 = 0 \\
 \Rightarrow |x|^2 + 5|x| - 6 = 0 \\
 \Rightarrow |x|^2 + 6|x| - |x| - 6 = 0 \\
 \Rightarrow (|x| + 6)(|x| - 1) = 0 \\
 \Rightarrow |x| = 1 \text{ or } |x| = -6 \\
 \text{But } |x| \text{ cannot be negative} \\
 \therefore |x| = 1 \quad \therefore x = \pm 1
 \end{aligned}$$

$$\begin{aligned}
 \alpha = 1, \beta = -1 \\
 \left| \tan^{-1} \alpha - \tan^{-1} \beta \right| &= \left| \tan^{-1} 1 - \tan^{-1}(-1) \right| \\
 &= \left| \frac{\pi}{4} - \left(-\frac{\pi}{4} \right) \right| \\
 &= \frac{\pi}{2}
 \end{aligned}$$

$$\begin{aligned}
 169. \quad 4 \sin^{-1} x + \cos^{-1} x = \pi \\
 \Rightarrow 3 \sin^{-1} x + \sin^{-1} x + \cos^{-1} x = \pi \\
 \Rightarrow 3 \sin^{-1} x = \pi - \frac{\pi}{2} \\
 \dots \left[\because \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2} \right] \\
 \Rightarrow 3 \sin^{-1} x = \frac{\pi}{2} \Rightarrow \sin^{-1} x = \frac{\pi}{6}
 \end{aligned}$$

$$\Rightarrow x = \sin \frac{\pi}{6} = \frac{1}{2}$$

$$\begin{aligned}
 170. \quad \cos \left(\cos^{-1} \frac{1}{5} + \sin^{-1} \frac{1}{5} + \cos^{-1} \frac{1}{5} \right) \\
 = \cos \left(\frac{\pi}{2} + \cos^{-1} \frac{1}{5} \right) = -\sin \left(\cos^{-1} \frac{1}{5} \right) \\
 = -\sin \left(\sin^{-1} \sqrt{\frac{24}{25}} \right) \\
 = -\frac{2\sqrt{6}}{5}
 \end{aligned}$$



$$\begin{aligned}
 171. \quad & \cos \left[2 \left(\tan^{-1} \frac{1}{5} + \tan^{-1} 5 \right) \right] \\
 &= \cos \left[2 \left(\cot^{-1} 5 + \tan^{-1} 5 \right) \right] \\
 &= \cos \left[2 \left(\frac{\pi}{2} \right) \right] \\
 &= \cos \pi \\
 &= -1
 \end{aligned}$$

$$\begin{aligned}
 172. \quad & \tan^{-1} (1+x) + \tan^{-1} (1-x) = \frac{\pi}{2} \\
 & \Rightarrow \tan^{-1} (1+x) = \frac{\pi}{2} - \tan^{-1} (1-x) \\
 & \Rightarrow \tan^{-1} (1+x) = \cot^{-1} (1-x) \\
 & \Rightarrow \tan^{-1} (1+x) = \tan^{-1} \left(\frac{1}{1-x} \right) \\
 & \Rightarrow 1+x = \frac{1}{1-x} \Rightarrow 1-x^2 = 1 \Rightarrow x = 0
 \end{aligned}$$

$$\begin{aligned}
 173. \quad & \text{The given equation can be written as} \\
 & \tan^{-1} x + \cot^{-1} x + \cot^{-1} x = \frac{2\pi}{3} \\
 & \Rightarrow \cot^{-1} x = \frac{2\pi}{3} - \frac{\pi}{2} \\
 & \quad \dots \left[\because \tan^{-1} x + \cot^{-1} x = \frac{\pi}{2} \right] \\
 & \Rightarrow \cot^{-1} x = \frac{\pi}{6} \Rightarrow x = \cot \frac{\pi}{6} \Rightarrow x = \sqrt{3}
 \end{aligned}$$

$$\begin{aligned}
 174. \quad & 3 \tan^{-1} x + \cot^{-1} x = \pi \\
 & \Rightarrow 2 \tan^{-1} x + \tan^{-1} x + \cot^{-1} x = \pi \\
 & \Rightarrow 2 \tan^{-1} x + \frac{\pi}{2} = \pi \\
 & \Rightarrow \tan^{-1} x = \frac{\pi}{4} \\
 & \Rightarrow \tan (\tan^{-1} x) = \tan \frac{\pi}{4} \\
 & \Rightarrow x = 1
 \end{aligned}$$

$$\begin{aligned}
 175. \quad & \tan^{-1} x + \tan^{-1} y = \frac{4\pi}{5} \\
 & \Rightarrow \frac{\pi}{2} - \cot^{-1} x + \frac{\pi}{2} - \cot^{-1} y = \frac{4\pi}{5} \\
 & \Rightarrow \cot^{-1} x + \cot^{-1} y = \frac{\pi}{5}
 \end{aligned}$$

$$\begin{aligned}
 176. \quad & \sin^{-1} \left(\frac{x}{13} \right) = \frac{\pi}{2} - \operatorname{cosec}^{-1} \left(\frac{13}{12} \right) \\
 & \quad \quad \quad = \sec^{-1} \left(\frac{13}{12} \right) = \cos^{-1} \left(\frac{12}{13} \right)
 \end{aligned}$$

$$\therefore \sin^{-1} \left(\frac{x}{13} \right) = \sin^{-1} \left(\frac{5}{13} \right)$$

$$\therefore x = 5$$

$$\begin{aligned}
 177. \quad & \cot^{-1} \alpha + \cot^{-1} \beta = \cot^{-1} x \\
 & \Rightarrow \cot^{-1} \left(\frac{\alpha\beta - 1}{\alpha + \beta} \right) = \cot^{-1} x \\
 & \quad \dots \left[\because \cot^{-1} x + \cot^{-1} y = \cot^{-1} \left(\frac{xy - 1}{x + y} \right) \right] \\
 & \Rightarrow x = \frac{\alpha\beta - 1}{\alpha + \beta}
 \end{aligned}$$

$$\begin{aligned}
 178. \quad & \tan^{-1} 2x + \tan^{-1} 3x = \frac{\pi}{4} \\
 & \Rightarrow \tan^{-1} \left(\frac{2x + 3x}{1 - (2x)(3x)} \right) = \frac{\pi}{4} \\
 & \Rightarrow \tan^{-1} \left(\frac{5x}{1 - 6x^2} \right) = \tan^{-1} (1) \\
 & \Rightarrow \frac{5x}{1 - 6x^2} = 1 \\
 & \Rightarrow 1 - 6x^2 = 5x \Rightarrow 6x^2 + 5x - 1 = 0 \\
 & \Rightarrow (x + 1) \left(x - \frac{1}{6} \right) = 0 \Rightarrow x = -1, \frac{1}{6}
 \end{aligned}$$

But $x = -1$ does not hold.


$$\therefore x = \frac{1}{6}$$

$$\begin{aligned}
 179. \quad & \tan^{-1} \left(\frac{x-1}{x-2} \right) + \tan^{-1} \left(\frac{x+1}{x+2} \right) = \frac{\pi}{4} \\
 & \Rightarrow \tan^{-1} \left[\frac{\frac{x-1}{x-2} + \frac{x+1}{x+2}}{1 - \left(\frac{x-1}{x-2} \right) \left(\frac{x+1}{x+2} \right)} \right] = \frac{\pi}{4} \\
 & \Rightarrow \frac{x-1}{x-2} + \frac{x+1}{x+2} = 1 - \left(\frac{x-1}{x-2} \right) \left(\frac{x+1}{x+2} \right) \\
 & \Rightarrow x^2 + x - 2 + x^2 - x - 2 = x^2 - 4 - x^2 + 1 \\
 & \Rightarrow 2x^2 = 1 \\
 & \Rightarrow x = \pm \frac{1}{\sqrt{2}}
 \end{aligned}$$



$$\begin{aligned}
 180. \quad & \tan^{-1} \frac{x}{y} - \tan^{-1} \left(\frac{x-y}{x+y} \right) \\
 &= \tan^{-1} \frac{x}{y} - \tan^{-1} \left(\frac{1-\frac{y}{x}}{1+\frac{y}{x}} \right) \\
 &= \tan^{-1} \frac{x}{y} - \left(\tan^{-1} 1 - \tan^{-1} \frac{y}{x} \right) \\
 &= \tan^{-1} \frac{x}{y} + \tan^{-1} \frac{y}{x} - \frac{\pi}{4} \\
 &= \tan^{-1} \frac{x}{y} + \cot^{-1} \frac{x}{y} - \frac{\pi}{4} = \frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4}
 \end{aligned}$$

$$\begin{aligned}
 181. \quad & A + B + C = \pi \\
 & \Rightarrow \tan^{-1} 2 + \tan^{-1} 3 + C = \pi \\
 & \Rightarrow \pi + \tan^{-1} \left(\frac{2+3}{1-2 \times 3} \right) + C = \pi \\
 & \Rightarrow \tan^{-1} \left(\frac{5}{-5} \right) + C = 0 \\
 & \Rightarrow -\tan^{-1} (1) + C = 0 \\
 & \Rightarrow -\frac{\pi}{4} + C = 0 \\
 & \Rightarrow C = \frac{\pi}{4}
 \end{aligned}$$

$$\begin{aligned}
 182. \quad & \angle A = 90^\circ \\
 & \tan^{-1} \left(\frac{c}{a+b} \right) + \tan^{-1} \left(\frac{b}{a+c} \right) \\
 &= \tan^{-1} \left[\frac{\frac{c}{a+b} + \frac{b}{a+c}}{1 - \left(\frac{c}{a+b} \right) \left(\frac{b}{a+c} \right)} \right] \\
 &= \tan^{-1} \left[\frac{ca + c^2 + ab + b^2}{a^2 + ab + ca + bc - bc} \right] \\
 &= \tan^{-1} \left[\frac{a^2 + ab + ca}{a^2 + ab + ca} \right] \quad \dots [\because b^2 + c^2 = a^2] \\
 &= \tan^{-1} (1) = \frac{\pi}{4}
 \end{aligned}$$


$$\begin{aligned}
 183. \quad & \text{Since, } \cot^{-1} x - \cot^{-1} y = \cot^{-1} \left(\frac{xy+1}{y-x} \right) \\
 \therefore & \cot^{-1} \frac{ab+1}{a-b} + \cot^{-1} \frac{bc+1}{b-c} + \cot^{-1} \frac{ca+1}{c-a} \\
 &= \cot^{-1} b - \cot^{-1} a + \cot^{-1} c - \cot^{-1} b \\
 & \quad \quad \quad + \cot^{-1} a - \cot^{-1} c \\
 &= 0
 \end{aligned}$$

$$\begin{aligned}
 184. \quad & \tan^{-1} x + \tan^{-1} y + \tan^{-1} z = \frac{\pi}{2} \\
 & \Rightarrow \tan^{-1} \left[\frac{x+y+z-xyz}{1-xy-yz-zx} \right] = \frac{\pi}{2} \\
 & \Rightarrow \left[\frac{x+y+z-xyz}{1-xy-yz-zx} \right] = \tan \frac{\pi}{2} \\
 & \Rightarrow xy + yz + zx - 1 = 0
 \end{aligned}$$

Alternate Method:

$$\text{Let } x = y = z = \frac{1}{\sqrt{3}}$$

$$\text{Then, } \tan^{-1} \frac{1}{\sqrt{3}} + \tan^{-1} \frac{1}{\sqrt{3}} + \tan^{-1} \frac{1}{\sqrt{3}} = \frac{\pi}{2}$$

Option (D) holds for these values of x, y, z

$$\begin{aligned}
 185. \quad & \text{Since, } 2 \tan^{-1} x = \tan^{-1} \frac{2x}{1-x^2} \\
 \therefore & 4 \tan^{-1} \frac{1}{5} = 2 \left[2 \tan^{-1} \frac{1}{5} \right] = 2 \tan^{-1} \frac{\frac{2}{5}}{1-\frac{1}{25}} \\
 &= 2 \tan^{-1} \frac{10}{24} = \tan^{-1} \frac{\frac{20}{24}}{1-\frac{100}{576}} = \tan^{-1} \frac{120}{119} \\
 \therefore & 4 \tan^{-1} \frac{1}{5} - \tan^{-1} \frac{1}{239} = \tan^{-1} \frac{120}{119} - \tan^{-1} \frac{1}{239} \\
 &= \tan^{-1} \frac{\frac{120}{119} - \frac{1}{239}}{1 + \frac{120}{119} \cdot \frac{1}{239}} = \tan^{-1} \frac{(120 \times 239) - 119}{(119 \times 239) + 120} \\
 &= \tan^{-1} 1 = \frac{\pi}{4}
 \end{aligned}$$

$$\begin{aligned}
 186. \quad & 2 \tan^{-1} (\cos x) = \tan^{-1} (2 \operatorname{cosec} x) \\
 \therefore & \tan^{-1} (\cos x) + \tan^{-1} (\cos x) = \tan^{-1} (2 \operatorname{cosec} x) \\
 \therefore & \tan [\tan^{-1} (\cos x) + \tan^{-1} (\cos x)] \\
 & \quad \quad \quad = \tan [\tan^{-1} (2 \operatorname{cosec} x)] \\
 & \Rightarrow \frac{\tan (\tan^{-1} \cos x) + \tan (\tan^{-1} \cos x)}{1 - \tan (\tan^{-1} \cos x) \cdot \tan (\tan^{-1} \cos x)} = 2 \operatorname{cosec} x \\
 & \Rightarrow \frac{\cos x + \cos x}{1 - \cos^2 x} = 2 \operatorname{cosec} x \\
 & \Rightarrow 2 \cos x = 2 \operatorname{cosec} x \cdot (1 - \cos^2 x) \\
 & \Rightarrow \cos x = \operatorname{cosec} x \cdot \sin^2 x \\
 & \Rightarrow \cos x = \sin x \\
 & \Rightarrow x = \frac{\pi}{4}
 \end{aligned}$$



$$\begin{aligned} \therefore \sin x + \cos x &= \sin \frac{\pi}{4} + \cos \frac{\pi}{4} \\ &= \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} = \sqrt{2} \end{aligned}$$

$$187. \theta = \tan^{-1} \sqrt{\frac{a(a+b+c)}{bc}} + \tan^{-1} \sqrt{\frac{b(a+b+c)}{ca}} + \tan^{-1} \sqrt{\frac{c(a+b+c)}{ab}}$$

$$\text{Let } s^2 = \frac{a+b+c}{abc}$$

$$\begin{aligned} \therefore \theta &= \tan^{-1} \sqrt{a^2 s^2} + \tan^{-1} \sqrt{b^2 s^2} + \tan^{-1} \sqrt{c^2 s^2} \\ &= \tan^{-1} (as) + \tan^{-1} (bs) + \tan^{-1} (cs) \\ &= \tan^{-1} \left[\frac{as + bs + cs - abcs^2}{1 - abs^2 - acs^2 - bcs^2} \right] \end{aligned}$$

$$\begin{aligned} \therefore \tan \theta &= \left[\frac{s[(a+b+c) - abcs^2]}{1 - (ab+bc+ca)s^2} \right] \\ &= \left[\frac{s[(a+b+c) - (a+b+c)]}{1 - s^2(ab+bc+ca)} \right] \\ &= 0 \end{aligned}$$

....[$\because s^2(abc) = (a+b+c)$]

Alternate Method :

Let $a = b = c = 1$. Then,

$$\begin{aligned} \theta &= \tan^{-1} \sqrt{3} + \tan^{-1} \sqrt{3} + \tan^{-1} \sqrt{3} = \pi \\ \Rightarrow \tan \theta &= 0 \end{aligned}$$

$$\begin{aligned} 188. 4 \tan^{-1} \frac{1}{5} - \tan^{-1} \frac{1}{70} + \tan^{-1} \frac{1}{99} \\ &= \tan^{-1} \frac{120}{119} + \tan^{-1} \frac{1}{99} - \tan^{-1} \frac{1}{70} \\ &= \tan^{-1} \left(\frac{120}{119} \right) + \tan^{-1} \left[\frac{\frac{1}{99} - \frac{1}{70}}{1 + \frac{1}{99} \cdot \frac{1}{70}} \right] \\ &= \tan^{-1} \left(\frac{120}{119} \right) + \tan^{-1} \left(\frac{-29}{6931} \right) \\ &= \tan^{-1} \frac{120}{119} - \tan^{-1} \frac{29}{6931} \\ &= \tan^{-1} \frac{120}{119} - \tan^{-1} \frac{1}{239} \\ &= \tan^{-1} 1 = \frac{\pi}{4} \end{aligned}$$

$$\begin{aligned} 189. 2 \tan^{-1} \frac{1}{5} + \sec^{-1} \frac{5\sqrt{2}}{7} + 2 \tan^{-1} \frac{1}{8} \\ &= 2 \left(\tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{8} \right) + \sec^{-1} \frac{5\sqrt{2}}{7} \\ &= 2 \tan^{-1} \left[\frac{\frac{1}{5} + \frac{1}{8}}{1 - \frac{1}{5} \left(\frac{1}{8} \right)} \right] + \tan^{-1} \sqrt{\left(\frac{5\sqrt{2}}{7} \right)^2 - 1} \\ &\quad \dots [\because \sec^{-1} x = \tan^{-1} \sqrt{x^2 - 1}] \\ &= 2 \tan^{-1} \frac{13}{39} + \tan^{-1} \frac{1}{7} \\ &= 2 \tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{7} \\ &= \tan^{-1} \left[\frac{2 \left(\frac{1}{3} \right)}{1 - \left(\frac{1}{3} \right)^2} \right] + \tan^{-1} \frac{1}{7} \\ &\quad \dots [\because 2 \tan^{-1} x = \tan^{-1} \frac{2x}{1-x^2}, \text{ if } -1 < x < 1] \\ &= \tan^{-1} \frac{3}{4} + \tan^{-1} \frac{1}{7} = \tan^{-1} \left[\frac{\frac{3}{4} + \frac{1}{7}}{1 - \frac{3}{4} \left(\frac{1}{7} \right)} \right] \end{aligned}$$

190. Consider option (A),

$$\sin(\cos^{-1} x) = \cos(\sin^{-1} x) = \sqrt{1-x^2}$$

$$\dots [\because \sin^{-1} x = \cos^{-1} \sqrt{1-x^2}, \cos^{-1} x = \sin^{-1} \sqrt{1-x^2}]$$

191. $\sin^{-1} [\cos(\sin^{-1} x)] + \cos^{-1} [\sin(\cos^{-1} x)]$

$$= \sin^{-1} \sqrt{1-x^2} + \cos^{-1} \sqrt{1-x^2}$$

$$\begin{aligned} \dots [\because \cos(\sin^{-1} x) = \sin(\cos^{-1} x) = \sqrt{1-x^2}] \\ = \frac{\pi}{2} \end{aligned}$$

192. $\sin^{-1} \frac{1}{3} + \sin^{-1} \frac{2}{3}$

$$= \sin^{-1} \left[\frac{1}{3} \sqrt{1-\frac{4}{9}} + \frac{2}{3} \sqrt{1-\frac{1}{9}} \right]$$

$$= \sin^{-1} \left[\frac{\sqrt{5+4\sqrt{2}}}{9} \right]$$

$$\therefore x = \frac{\sqrt{5+4\sqrt{2}}}{9}$$



$$\begin{aligned}
 193. \quad \sin^{-1}x + \cos^{-1}y &= \frac{2\pi}{5} \\
 \Rightarrow \frac{\pi}{2} - \cos^{-1}x + \frac{\pi}{2} - \sin^{-1}y &= \frac{2\pi}{5} \\
 \Rightarrow \pi - \cos^{-1}x - \sin^{-1}y &= \frac{2\pi}{5} \\
 \Rightarrow \cos^{-1}x + \sin^{-1}y &= \pi - \frac{2\pi}{5} \\
 &= \frac{3\pi}{5}
 \end{aligned}$$

$$\begin{aligned}
 194. \quad \text{Given, } \sin^{-1}x + \sin^{-1}y &= \frac{2\pi}{3} \\
 \therefore \cos^{-1}x + \cos^{-1}y &= \pi - \frac{2\pi}{3} = \frac{\pi}{3} \\
 &\dots \left[\begin{array}{l} \text{If } \sin^{-1}x + \sin^{-1}y = \theta, \\ \text{then } \cos^{-1}x + \cos^{-1}y = \pi - \theta \end{array} \right]
 \end{aligned}$$

$$\begin{aligned}
 195. \quad \sin^{-1}\left(\frac{2\sqrt{2}}{3}\right) + \sin^{-1}\left(\frac{1}{3}\right) \\
 = \sin^{-1}\left(\frac{2\sqrt{2}}{3}\right) + \cos^{-1}\sqrt{1 - \left(\frac{1}{3}\right)^2} \\
 \dots \left[\because \sin^{-1}x = \cos^{-1}\sqrt{1-x^2} \right] \\
 = \sin^{-1}\left(\frac{2\sqrt{2}}{3}\right) + \cos^{-1}\left(\frac{2\sqrt{2}}{3}\right) \\
 = \frac{\pi}{2} \quad \dots \left[\because \sin^{-1}x + \cos^{-1}x = \frac{\pi}{2} \right]
 \end{aligned}$$

$$\begin{aligned}
 196. \quad \left[\sin\left(\tan^{-1}\frac{3}{4}\right) \right]^2 \\
 = \left[\sin\left\{ \sin^{-1}\left(\frac{\frac{3}{4}}{\sqrt{1 + \left(\frac{3}{4}\right)^2}} \right) \right\} \right]^2 \\
 \dots \left[\because \tan^{-1}x = \sin^{-1}\left(\frac{x}{\sqrt{1+x^2}}\right) \right] \\
 = \left[\sin\left(\sin^{-1}\frac{3}{5}\right) \right]^2 = \left(\frac{3}{5}\right)^2 = \frac{9}{25}
 \end{aligned}$$

$$\begin{aligned}
 197. \quad \sin[\cot^{-1}(x+1)] &= \sin\left(\sin^{-1}\frac{1}{\sqrt{x^2+2x+2}}\right) \\
 &= \frac{1}{\sqrt{x^2+2x+2}}
 \end{aligned}$$

$$\begin{aligned}
 \text{And } \cos(\tan^{-1}x) &= \cos\left(\cos^{-1}\frac{1}{\sqrt{1+x^2}}\right) \\
 &= \frac{1}{\sqrt{1+x^2}}
 \end{aligned}$$

$$\begin{aligned}
 \text{Thus, } \frac{1}{\sqrt{x^2+2x+2}} &= \frac{1}{\sqrt{1+x^2}} \\
 \Rightarrow x^2+2x+2 &= 1+x^2 \Rightarrow x = -\frac{1}{2}
 \end{aligned}$$

$$\begin{aligned}
 198. \quad \cos(\tan^{-1}x) &= \cos\left[\cos^{-1}\frac{1}{\sqrt{1+x^2}}\right] \\
 &\dots \left[\because \tan^{-1}x = \cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right) \right] \\
 &= \frac{1}{\sqrt{1+x^2}}
 \end{aligned}$$

$$\begin{aligned}
 199. \quad \tan(\cos^{-1}x) \\
 = \tan\left[\tan^{-1}\frac{\sqrt{1-x^2}}{x}\right] \\
 \dots \left[\because \cos^{-1}x = \tan^{-1}\frac{\sqrt{1-x^2}}{x} \right] \\
 = \frac{\sqrt{1-x^2}}{x}
 \end{aligned}$$

$$\begin{aligned}
 200. \quad \cos\left[\tan^{-1}\left(\frac{3}{4}\right)\right] &= \cos\left[\cos^{-1}\frac{1}{\sqrt{1 + \left(\frac{3}{4}\right)^2}}\right] \\
 &= \cos\left[\cos^{-1}\left(\frac{4}{5}\right)\right] \\
 &= \frac{4}{5}
 \end{aligned}$$

$$\begin{aligned}
 201. \quad \text{Let } x = \cos\theta \Rightarrow \theta &= \cos^{-1}x \\
 \text{Now, } \cos^{-1}x + \cos^{-1}\left(\frac{x}{2} + \frac{1}{2}\sqrt{3-3x^2}\right) \\
 = \theta + \cos^{-1}\left(\frac{\cos\theta}{2} + \frac{\sqrt{3}}{2}\sqrt{1-\cos^2\theta}\right) \\
 = \theta + \cos^{-1}\left(\frac{1}{2}\cos\theta + \frac{\sqrt{3}}{2}\sin\theta\right)
 \end{aligned}$$



$$\begin{aligned}
 &= \theta + \cos^{-1} \left(\cos \frac{\pi}{3} \cdot \cos \theta + \sin \frac{\pi}{3} \cdot \sin \theta \right) \\
 &= \theta + \cos^{-1} \left[\cos \left(\frac{\pi}{3} - \theta \right) \right] = \theta + \frac{\pi}{3} - \theta \\
 &= \frac{\pi}{3}
 \end{aligned}$$

$$202. \tan^{-1} y = \tan^{-1} x + \tan^{-1} \left(\frac{2x}{1-x^2} \right)$$

$$= \tan^{-1} x + 2 \tan^{-1} x$$

$$\therefore \tan^{-1} y = 3 \tan^{-1} x$$

$$\text{Since, } 3 \tan^{-1} x = \tan^{-1} \left(\frac{3x-x^3}{1-3x^2} \right)$$

$$\therefore y = \frac{3x-x^3}{1-3x^2}$$

$$203. 2 \cot^{-1} \left(\frac{1}{2} \right) - \cot^{-1} \left(\frac{4}{3} \right)$$

$$= 2 \tan^{-1} (2) - \cot^{-1} \left(\frac{4}{3} \right)$$

$$\dots \left[\because \cot^{-1} (x) = \tan^{-1} \left(\frac{1}{x} \right), \text{ if } x > 0 \right]$$

$$= \pi + \tan^{-1} \left(\frac{4}{-3} \right) - \cot^{-1} \left(\frac{4}{3} \right)$$

$$\dots \left[\because 2 \tan^{-1} (x) = \pi + \tan^{-1} \left(\frac{2x}{1-x^2} \right), \text{ if } x > 1 \right]$$

$$= \pi - \tan^{-1} \left(\frac{4}{3} \right) - \cot^{-1} \left(\frac{4}{3} \right)$$

$$= \pi - \left[\tan^{-1} \left(\frac{4}{3} \right) + \cot^{-1} \left(\frac{4}{3} \right) \right]$$

$$= \pi - \frac{\pi}{2}$$

$$= \frac{\pi}{2}$$

$$204. \tan^{-1} \left(\frac{1}{2} \tan 2A \right) + \tan^{-1} (\cot A) + \tan^{-1} (\cot^3 A)$$

$$= \tan^{-1} \left[\frac{1}{2} \left(\frac{2 \tan A}{1 - \tan^2 A} \right) \right] + \pi + \tan^{-1} \left(\frac{\cot A + \cot^3 A}{1 - \cot^4 A} \right)$$

$$\left[\begin{array}{l}
 \because 0 \leq A \leq \frac{\pi}{4}, \\
 \dots \left[\tan^{-1} x + \tan^{-1} y = \pi + \tan^{-1} \left(\frac{x+y}{1-xy} \right), \right. \\
 \left. \text{for } x, y > 0 \text{ and } xy > 1 \right]
 \end{array} \right]$$

$$= \pi + \tan^{-1} \left(\frac{\tan A}{1 - \tan^2 A} \right) + \tan^{-1} \left[\frac{\cot A (1 + \cot^2 A)}{(1 + \cot^2 A)(1 - \cot^2 A)} \right]$$

$$= \pi + \tan^{-1} \left(\frac{\tan A}{1 - \tan^2 A} \right) + \tan^{-1} \left(\frac{\cot A}{1 - \cot^2 A} \right)$$

$$= \pi + \tan^{-1} \left(\frac{\tan A}{1 - \tan^2 A} \right) + \tan^{-1} \left(\frac{\tan A}{\tan^2 A - 1} \right)$$

$$= \pi + \tan^{-1} \left(\frac{\tan A}{1 - \tan^2 A} \right) + \tan^{-1} \left(\frac{-\tan A}{1 - \tan^2 A} \right)$$

$$= \pi + 0 \quad \dots [\tan^{-1}(-x) = -\tan^{-1} x]$$

$$= \pi$$

$$205. \cos^{-1} \left(\frac{15}{17} \right) + 2 \tan^{-1} \left(\frac{1}{5} \right)$$

$$= \cos^{-1} \left(\frac{15}{17} \right) + \cos^{-1} \left(\frac{1 - \frac{1}{25}}{1 + \frac{1}{25}} \right)$$

$$= \cos^{-1} \left(\frac{15}{17} \right) + \cos^{-1} \left(\frac{12}{13} \right)$$

$$= \cos^{-1} \left(\frac{15}{17} \times \frac{12}{13} - \sqrt{1 - \left(\frac{15}{17} \right)^2} \sqrt{1 - \left(\frac{12}{13} \right)^2} \right)$$

$$= \cos^{-1} \left(\frac{140}{221} \right)$$

$$206. \tan \left[\cos^{-1} \frac{4}{5} + \tan^{-1} \frac{2}{3} \right]$$

$$= \tan \left[\tan^{-1} \frac{\sqrt{1 - \frac{16}{25}}}{\frac{4}{5}} + \tan^{-1} \frac{2}{3} \right]$$

$$= \tan \left[\tan^{-1} \left(\frac{\frac{3}{4} + \frac{2}{3}}{1 - \frac{3}{4} \cdot \frac{2}{3}} \right) \right]$$

$$= \tan \left(\tan^{-1} \frac{17}{6} \right)$$

$$= \frac{17}{6}$$

$$207. \cot (\cos^{-1} x) = \sec \left[\tan^{-1} \left(\frac{a}{\sqrt{b^2 - a^2}} \right) \right]$$

$$\Rightarrow \cot \left[\cot^{-1} \left(\frac{x}{\sqrt{1 - x^2}} \right) \right] = \sec \left[\sec^{-1} \sqrt{1 + \frac{a^2}{b^2 - a^2}} \right]$$



$$\Rightarrow \frac{x}{\sqrt{1-x^2}} = \frac{b}{\sqrt{b^2-a^2}}$$

$$\Rightarrow \frac{x^2}{1-x^2} = \frac{b^2}{b^2-a^2}$$

$$\Rightarrow \frac{1-x^2}{x^2} = \frac{b^2-a^2}{b^2}$$

$$\Rightarrow \frac{1}{x^2} = \frac{2b^2-a^2}{b^2}$$

$$\Rightarrow x = \frac{b}{\sqrt{2b^2-a^2}}$$

$$\begin{aligned} 208. \quad & \tan^{-1} \left\{ \sin \left(\cos^{-1} \sqrt{\frac{2}{3}} \right) \right\} \\ &= \tan^{-1} \left\{ \sin \left(\sin^{-1} \sqrt{\frac{1}{3}} \right) \right\} \\ & \quad \dots \left[\because \cos^{-1} x = \sin^{-1} \sqrt{1-x^2} \right] \\ &= \tan^{-1} \left(\frac{1}{\sqrt{3}} \right) = \frac{\pi}{6} \end{aligned}$$

$$209. \quad \text{We know } \frac{5^x + 5^{-x}}{2} \geq 1 \quad \dots \left[\because \text{A.M.} \geq \text{G.M.} \right]$$

Since, $\cos(e^x) \leq 1$

So, there does not exist any solution.

210. Applying $R_1 \rightarrow R_1 - R_3$ and $R_2 \rightarrow R_2 - R_3$ in the given determinant, we get

$$\begin{vmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ \sin^2 \theta & \cos^2 \theta & 1+4\sin 4\theta \end{vmatrix} = 0$$

$$\Rightarrow 1 + 4\sin 4\theta + \cos^2 \theta + \sin^2 \theta = 0$$

$$\Rightarrow 4 \sin 4\theta = -2$$

$$\Rightarrow \sin 4\theta = \frac{-1}{2}$$

$$\Rightarrow 4\theta = \frac{7\pi}{6} \text{ or } \frac{11\pi}{6}$$

Since, $0 < \theta < \frac{\pi}{2}$

$$\Rightarrow 0 < 4\theta < 2\pi$$

$$\Rightarrow \theta = \frac{7\pi}{24} \text{ or } \frac{11\pi}{24}$$

$$211. \quad \frac{(x+1)^2}{x^3+x} = \frac{A}{x} + \frac{Bx+C}{x^2+1}$$

$$\therefore \frac{(x+1)^2}{x(x^2+1)} = \frac{Ax^2+A+Bx^2+Cx}{x(x^2+1)}$$

$$\therefore x^2+2x+1 = (A+B)x^2+Cx+A$$

Equating coefficients on both sides, we get

$$A+B=1, C=2, A=1$$

$$\Rightarrow B=0$$

$$\therefore \operatorname{cosec}^{-1} \left(\frac{1}{A} \right) + \cot^{-1} \left(\frac{1}{B} \right) + \sec^{-1} C$$

$$= \frac{\pi}{2} + 0 + \frac{\pi}{3} = \frac{5\pi}{6}$$

$$212. \quad 2y=1 \Rightarrow y = \frac{1}{2} \Rightarrow \sin x = \frac{1}{2}$$

$$\Rightarrow x = \frac{\pi}{6}, \frac{5\pi}{6}, -\frac{7\pi}{6}, -\frac{11\pi}{6} \text{ in } -2\pi \leq x \leq 2\pi$$

\therefore number of points of intersection = 4

$$213. \quad \frac{a}{\sin A} = \frac{b}{\sin B} \Rightarrow \frac{5}{\sin \left(\frac{\pi}{2} + B \right)} = \frac{4}{\sin B}$$

$$\Rightarrow \frac{5}{\cos B} = \frac{4}{\sin B} \Rightarrow \tan B = \frac{4}{5}$$

$$\text{Now, } \tan A = \tan \left(\frac{\pi}{2} + B \right) = -\cot B = \frac{-5}{4}$$

$$\tan C = \tan(\pi - (A+B))$$

$$= -\tan(A+B)$$

$$= -\frac{(\tan A + \tan B)}{1 - \tan A \cdot \tan B} = \frac{-\left(-\frac{5}{4} + \frac{4}{5} \right)}{1 - \left(-\frac{5}{4} \times \frac{4}{5} \right)} = \frac{9}{40}$$

$$\Rightarrow C = \tan^{-1} \left(\frac{2 \cdot \frac{1}{9}}{1 - \left(\frac{1}{9} \right)^2} \right)$$

$$\Rightarrow C = 2 \tan^{-1} \left(\frac{1}{9} \right)$$

$$214. \quad \text{Given, } x = \sin^{-1} K, y = \cos^{-1} K$$

$$\therefore \sin x = \cos y = K$$

$$\therefore \sin x = \sin \left(\frac{\pi}{2} - y \right)$$

$$\therefore x = \frac{\pi}{2} - y \Rightarrow x + y = \frac{\pi}{2}$$



$$\begin{aligned}
 215. \quad A - B &= \frac{1}{\pi} \begin{bmatrix} \sin^{-1}(\pi x) & \tan^{-1}\left(\frac{x}{\pi}\right) \\ \sin^{-1}\left(\frac{x}{\pi}\right) & \cot^{-1}(\pi x) \end{bmatrix} \\
 &\quad - \frac{1}{\pi} \begin{bmatrix} -\cos^{-1}(\pi x) & \tan^{-1}\left(\frac{x}{\pi}\right) \\ \sin^{-1}\left(\frac{x}{\pi}\right) & -\tan^{-1}(\pi x) \end{bmatrix} \\
 &= \frac{1}{\pi} \begin{bmatrix} \frac{\pi}{2} & 0 \\ 0 & \frac{\pi}{2} \end{bmatrix} \\
 &= \frac{1}{2} \mathbf{I}
 \end{aligned}$$

216. $\frac{1}{6} \sin \theta$, $\cos \theta$ and $\tan \theta$ are in G.P.

$$\therefore \cos^2 \theta = \frac{1}{6} \cdot \sin \theta \cdot \tan \theta$$

$$\Rightarrow 6 \cos^2 \theta = \sin \theta \cdot \frac{\sin \theta}{\cos \theta}$$

$$\Rightarrow 6 \cos^3 \theta = \sin^2 \theta$$

$$\Rightarrow 6 \cos^3 \theta + \cos^2 \theta - 1 = 0$$

Here, $\cos \theta = \frac{1}{2}$ is the only real root.

$$\therefore \theta = 2n\pi \pm \frac{\pi}{3}$$

$$217. \quad \frac{\sin B}{b} = \frac{\sin A}{a}$$

$$\Rightarrow \sin B = \frac{b \sin A}{a} = \frac{8 \sin 30^\circ}{6} = \frac{2}{3}$$

$$\Rightarrow \sin(\sin^{-1} x) = \frac{2}{3}$$

$$\Rightarrow x = \frac{2}{3}$$

$$\begin{aligned}
 218. \quad &\cot \left[\sum_{n=1}^{100} \cot^{-1} \left(1 + \sum_{k=1}^n 2k \right) \right] \\
 &= \cot \left[\sum_{n=1}^{100} \cot^{-1} (1 + 2 + 4 + 6 + \dots + 2n) \right] \\
 &= \cot \left\{ \sum_{n=1}^{100} \cot^{-1} [1 + n(n+1)] \right\} \\
 &= \cot \left\{ \sum_{n=1}^{100} \tan^{-1} \left[\frac{1}{1 + n(n+1)} \right] \right\} \\
 &= \cot \left\{ \sum_{n=1}^{100} \tan^{-1} \left[\frac{n+1-n}{1 + n(n+1)} \right] \right\} \\
 &= \cot \left\{ \sum_{n=1}^{100} [\tan^{-1}(n+1) - \tan^{-1} n] \right\} \\
 &= \cot [(\tan^{-1} 2 - \tan^{-1} 1) + (\tan^{-1} 3 - \tan^{-1} 2) + \dots \\
 &\quad + (\tan^{-1} 101 - \tan^{-1} 100)] \\
 &= \cot (\tan^{-1} 101 - \tan^{-1} 1) \\
 &= \cot \left[\tan^{-1} \left(\frac{101-1}{1+101} \right) \right] \\
 &= \cot \left[\tan^{-1} \left(\frac{100}{102} \right) \right] \\
 &= \cot \left[\tan^{-1} \left(\frac{50}{51} \right) \right] \\
 &= \cot \left[\cot^{-1} \left(\frac{51}{50} \right) \right] \\
 &= \frac{51}{50}
 \end{aligned}$$



Evaluation Test

1. $\sin 3x = 3 \sin x - 4 \sin^3 x$

$$\Rightarrow \sin^3 x = \frac{1}{4}(3 \sin x - \sin 3x)$$

$$\cos 3x = 4 \cos^3 x - 3 \cos x$$

$$\Rightarrow \cos^3 x = \frac{1}{4}(\cos 3x + 3 \cos x)$$

Given, $\cos 3x \cos^3 x + \sin 3x \sin^3 x = 0$

$$\Rightarrow \cos 3x \cdot \frac{1}{4}(\cos 3x + 3 \cos x)$$

$$+ \sin 3x \cdot \frac{1}{4}(3 \sin x - \sin 3x) = 0$$

$$\Rightarrow \frac{1}{4}[\cos^2 3x + 3 \cos x \cos 3x + 3 \sin x \sin 3x - \sin^2 3x] = 0$$

$$\Rightarrow \cos^2 3x - \sin^2 3x + 3(\cos 3x \cos x + \sin 3x \sin x) = 0$$

$$\Rightarrow \cos 6x + 3 \cos 2x = 0$$



$$\Rightarrow 4 \cos^3 2x - 3 \cos 2x + 3 \cos 2x = 0$$

$$\Rightarrow 4 \cos^3 2x = 0$$

$$\Rightarrow \cos 2x = 0$$

$$\Rightarrow 2x = (2n + 1) \frac{\pi}{2}$$

$$\Rightarrow x = (2n + 1) \frac{\pi}{4}$$

$$2. \quad \sin x \sqrt{8 \cos^2 x} = 1$$

$$\therefore \sin x |2\sqrt{2} \cos x| = 1 \quad \dots [\because \sqrt{8} = 2\sqrt{2}]$$

$$\therefore \sin x |\cos x| = \frac{1}{2\sqrt{2}}$$

Case I:

$$\text{If } \cos x > 0, \sin x \cos x = \frac{1}{2\sqrt{2}}$$

$$\therefore \frac{1}{2} \sin 2x = \frac{1}{2\sqrt{2}}$$

$$\therefore \sin 2x = \frac{1}{\sqrt{2}}$$

$$\therefore 2x = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{9\pi}{4}, \frac{11\pi}{4}$$

$$\dots [\because x \in (0, 2\pi), \therefore 2x \in (0, 4\pi)]$$

$$\therefore x = \frac{\pi}{8}, \frac{3\pi}{8}, \frac{9\pi}{8}, \frac{11\pi}{8}$$

But $\cos x > 0$ (x must be in 1st or 4th Quadrant)

$$\therefore \text{the possible values are } \frac{\pi}{8}, \frac{3\pi}{8}.$$

Case II:

If $\cos x < 0$,

$$\sin x (-\cos x) = \frac{1}{2\sqrt{2}} \Rightarrow \sin 2x = -\frac{1}{\sqrt{2}}$$

$$\therefore 2x = \frac{5\pi}{4}, \frac{7\pi}{4}$$

$$\therefore x = \frac{5\pi}{8}, \frac{7\pi}{8}$$

\therefore The values of x satisfying the given equation between 0 and 2π are $\frac{\pi}{8}, \frac{3\pi}{8}, \frac{5\pi}{8}, \frac{7\pi}{8}$.

These are in A.P. with common difference $\frac{\pi}{4}$.

$$3. \quad 16^{\sin^2 x} + 16^{\cos^2 x} = 10$$

$$\Rightarrow 16^{\sin^2 x} + 16^{1-\sin^2 x} = 10$$

$$\Rightarrow 16^{\sin^2 x} + \frac{16}{16^{\sin^2 x}} = 10$$

$$\text{Let } t = 16^{\sin^2 x}$$

$$\Rightarrow t + \frac{16}{t} = 10 \quad \Rightarrow t^2 + 16 = 10t$$

$$\Rightarrow t^2 - 10t + 16 = 0 \quad \Rightarrow (t-2)(t-8) = 0$$

$$\Rightarrow t = 2 \text{ or } t = 8$$

$$\Rightarrow 16^{\sin^2 x} = 2 \text{ or } 16^{\sin^2 x} = 8$$

$$\Rightarrow 2^{4\sin^2 x} = 2^1 \text{ or } 2^{4\sin^2 x} = 2^3$$

$$\Rightarrow 4 \sin^2 x = 1 \text{ or } 4 \sin^2 x = 3$$

$$\Rightarrow \sin^2 x = \frac{1}{4} \text{ or } \sin^2 x = \frac{3}{4}$$

$$\Rightarrow \sin^2 x = \sin^2\left(\frac{\pi}{6}\right) \text{ or } \sin^2 x = \sin^2\left(\frac{\pi}{3}\right)$$

$$\Rightarrow x = n\pi \pm \frac{\pi}{6} \text{ or } x = n\pi \pm \frac{\pi}{3}$$

$$\Rightarrow x = \frac{\pi}{6}, \frac{7\pi}{6}, \frac{5\pi}{6}, \frac{11\pi}{6} \text{ or } x = \frac{\pi}{3}, \frac{4\pi}{3}, \frac{2\pi}{3}, \frac{5\pi}{3}$$

\therefore There are 8 solutions in $[0, 2\pi]$.

4. The maximum value of $a \sin x + b \cos x$ is $\sqrt{a^2 + b^2}$.

\therefore Maximum value of $\sin x + \cos x$ is $\sqrt{2}$ and the maximum value of $1 + \sin 2x$ is 2.

\therefore The given equation will be true only when

$$\sin x + \cos x = \sqrt{2} \text{ and } 1 + \sin 2x = 2$$

$$\text{If } \sin x + \cos x = \sqrt{2}$$

$$\Rightarrow \cos x \cdot \frac{1}{\sqrt{2}} + \sin x \cdot \frac{1}{\sqrt{2}} = 1$$

$$\Rightarrow \cos x \cos \frac{\pi}{4} + \sin x \sin \frac{\pi}{4} = 1$$

$$\Rightarrow \cos\left(x - \frac{\pi}{4}\right) = 1$$

$$\Rightarrow x - \frac{\pi}{4} = 2n\pi,$$

$$\Rightarrow x = 2n\pi + \frac{\pi}{4} \quad \dots \text{(i)}$$

$$1 + \sin 2x = 2 \Rightarrow \sin 2x = 1$$

$$\Rightarrow \sin 2x = \sin \frac{\pi}{2}$$

$$\Rightarrow 2x = n\pi + (-1)^n \cdot \frac{\pi}{2}$$

$$\Rightarrow x = \frac{n\pi}{2} + (-1)^n \cdot \frac{\pi}{4} \quad \dots \text{(ii)}$$

The value of $x \in [-\pi, \pi]$ which satisfies both

(i) and (ii) is $\frac{\pi}{4}$.



$$\begin{aligned}
 5. \quad & \sin^4 x + \cos^4 x = \sin x \cos x \\
 & \Rightarrow (\sin^2 x + \cos^2 x)^2 - 2 \sin^2 x \cos^2 x = \sin x \cos x \\
 & \Rightarrow 1 - \frac{1}{2} (2 \sin x \cos x)^2 = \frac{1}{2} 2 \sin x \cos x \\
 & \Rightarrow 1 - \frac{1}{2} \sin^2 2x = \frac{1}{2} \sin 2x \\
 & \Rightarrow \sin^2 2x + \sin 2x - 2 = 0 \\
 & \Rightarrow (\sin 2x + 2)(\sin 2x - 1) = 0 \\
 & \Rightarrow \sin 2x = 1 \quad \dots [\because \sin 2x \neq -2] \\
 & \Rightarrow \sin 2x = \sin \frac{\pi}{2} \\
 & \Rightarrow 2x = n\pi + (-1)^n \frac{\pi}{2} \\
 & \Rightarrow x = \frac{n\pi}{2} + (-1)^n \frac{\pi}{4}
 \end{aligned}$$

\therefore The value of x in $[0, 2\pi]$ are $\frac{\pi}{4}$ and $\frac{5\pi}{4}$.

\therefore There are 2 solutions.

$$\begin{aligned}
 6. \quad & \tan^4 x - 2 \sec^2 x + a^2 = 0 \\
 & \Rightarrow \tan^4 x - 2(1 + \tan^2 x) + a^2 = 0 \\
 & \Rightarrow \tan^4 x - 2 \tan^2 x - 2 + a^2 = 0 \\
 & \Rightarrow \tan^4 x - 2 \tan^2 x + 1 - 3 + a^2 = 0 \\
 & \Rightarrow (\tan^2 x - 1)^2 = 3 - a^2 \\
 & \Rightarrow 3 - a^2 \geq 0 \\
 & \Rightarrow a^2 \leq 3 \\
 & \Rightarrow |a| \leq \sqrt{3}
 \end{aligned}$$

$$\begin{aligned}
 7. \quad & 3 \cos x + 4 \sin x = 5 \\
 \therefore & 3 \left(\frac{1 - \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} \right) + 4 \left(\frac{2 \tan \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} \right) = 5
 \end{aligned}$$

Let $\tan \frac{x}{2} = t$

$$\begin{aligned}
 \therefore \quad & 3 - 3t^2 + 8t = 5 + 5t^2 \quad \Rightarrow 8t^2 - 8t + 2 = 0 \\
 & \Rightarrow 4t^2 - 4t + 1 = 0 \quad \Rightarrow (2t - 1)^2 = 0 \\
 & \Rightarrow t = \frac{1}{2} \quad \Rightarrow \tan \frac{x}{2} = \tan \alpha \\
 & \Rightarrow \frac{x}{2} = n\pi + \alpha \quad \Rightarrow x = 2n\pi + 2\alpha
 \end{aligned}$$

$$\begin{aligned}
 8. \quad & \tan \theta + \tan \left(\theta + \frac{\pi}{3} \right) + \tan \left(\theta + \frac{2\pi}{3} \right) = 3 \\
 & \Rightarrow \tan \theta + \frac{\tan \theta + \sqrt{3}}{1 - \sqrt{3} \tan \theta} + \frac{\tan \theta - \sqrt{3}}{1 + \sqrt{3} \tan \theta} = 3 \\
 & \frac{\tan \theta (1 - 3 \tan^2 \theta) + (\tan \theta + \sqrt{3})(1 + \sqrt{3} \tan \theta) + (\tan \theta - \sqrt{3})(1 - \sqrt{3} \tan \theta)}{1 - 3 \tan^2 \theta} = 3
 \end{aligned}$$

$$\begin{aligned}
 & \Rightarrow \frac{9 \tan \theta - 3 \tan^3 \theta}{1 - 3 \tan^2 \theta} = 3 \\
 & \Rightarrow 3 \left(\frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta} \right) = 3 \\
 & \Rightarrow 3 \tan 3\theta = 3 \\
 & \Rightarrow \tan 3\theta = 1 = \tan \frac{\pi}{4} \\
 & \Rightarrow 3\theta = n\pi + \frac{\pi}{4} \\
 & \Rightarrow \theta = (4n + 1) \frac{\pi}{12}
 \end{aligned}$$

$$\begin{aligned}
 9. \quad & \frac{5}{6} > \frac{20}{37} \quad \therefore \tan \frac{A}{2} > \tan \frac{B}{2} \\
 \therefore & \frac{A}{2} > \frac{B}{2} \quad \therefore A > B
 \end{aligned}$$

$$\begin{aligned}
 \therefore \quad & \tan \left(\frac{A}{2} + \frac{B}{2} \right) = \frac{\tan \frac{A}{2} + \tan \frac{B}{2}}{1 - \tan \frac{A}{2} \tan \frac{B}{2}} \\
 & \Rightarrow \tan \left(\frac{A+B}{2} \right) = \frac{\frac{5}{6} + \frac{20}{37}}{1 - \frac{5}{6} \cdot \frac{20}{37}} \\
 & \Rightarrow \tan \left(\frac{\pi - C}{2} \right) = \frac{185 + 120}{222 - 100} \\
 & \Rightarrow \tan \left(\frac{\pi - C}{2} \right) = \frac{305}{122} \\
 & \Rightarrow \cot \frac{C}{2} = \frac{305}{122} \quad \Rightarrow \tan \frac{C}{2} = \frac{122}{305}
 \end{aligned}$$

Since, $\frac{20}{37} > \frac{122}{305}$


$$\begin{aligned}
 \therefore \quad & \tan \frac{B}{2} > \tan \frac{C}{2} \\
 & \Rightarrow \frac{B}{2} > \frac{C}{2} \quad \Rightarrow B > C \\
 & \Rightarrow A > B > C \quad \Rightarrow a > b > c
 \end{aligned}$$

$$\begin{aligned}
 10. \quad & A(\Delta ABC) = \frac{9\sqrt{3}}{2} \\
 & \Rightarrow \frac{1}{2} bc \sin A = \frac{9\sqrt{3}}{2} \\
 & \Rightarrow \frac{1}{2} \times bc \times \frac{\sqrt{3}}{2} = \frac{9\sqrt{3}}{2} \\
 & \dots \left[\because \sin A = \sin \frac{2\pi}{3} = \frac{\sqrt{3}}{2} \right]
 \end{aligned}$$



$$\begin{aligned} \Rightarrow bc &= 18 \\ \cos A &= \frac{b^2 + c^2 - a^2}{2bc} \\ \Rightarrow \cos \frac{2\pi}{3} &= \frac{(b-c)^2 + 2bc - a^2}{2bc} \\ \Rightarrow -\frac{1}{2} &= \frac{(3\sqrt{3})^2 + 2 \times 18 - a^2}{2 \times 18} \\ \Rightarrow -18 &= 27 + 36 - a^2 \\ \Rightarrow a^2 &= 27 + 36 + 18 = 81 \\ \Rightarrow a &= 9 \text{ cm} \end{aligned}$$

11.



Let $\angle B = 30^\circ$, $\angle C = 45^\circ \therefore \angle A = 105^\circ$

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$

$$\therefore \frac{\sin 105^\circ}{\sqrt{3}+1} = \frac{\sin 30^\circ}{b} = \frac{\sin 45^\circ}{c}$$

$$\therefore b = \frac{(\sqrt{3}+1)\sin 30^\circ}{\sin 105^\circ} = \frac{\sqrt{3}+1}{2\sin 105^\circ}$$

$$c = \frac{(\sqrt{3}+1)\sin 45^\circ}{\sin 105^\circ} = \frac{\sqrt{3}+1}{\sqrt{2}\sin 105^\circ}$$

$$\begin{aligned} A(\triangle ABC) &= \frac{1}{2} bc \sin A \\ &= \frac{1}{2} \times \frac{\sqrt{3}+1}{2\sin 105^\circ} \times \frac{\sqrt{3}+1}{\sqrt{2}\sin 105^\circ} \times \sin 105^\circ \\ &= \frac{(\sqrt{3}+1)^2}{4\sqrt{2}\sin(60^\circ+45^\circ)} \\ &= \frac{(\sqrt{3}+1)^2}{4\sqrt{2}\left(\frac{\sqrt{3}}{2} \cdot \frac{1}{\sqrt{2}} + \frac{1}{2} \cdot \frac{1}{\sqrt{2}}\right)} \\ &= \frac{(\sqrt{3}+1)^2}{4\sqrt{2}\left(\frac{\sqrt{3}+1}{2\sqrt{2}}\right)} \\ &= \frac{\sqrt{3}+1}{2} \end{aligned}$$

$$\begin{aligned} 12. \quad \text{Let } \frac{b+c}{11} = \frac{c+a}{12} = \frac{a+b}{13} = k &= \frac{2(a+b+c)}{36} \\ &= \frac{a+b+c}{18} \\ &\dots \text{(By property of equal ratio)} \end{aligned}$$


$$\therefore b+c = 11k, c+a = 12k, a+b = 13k,$$

$$\therefore a = 7k, b = 6k, c = 5k$$

$$\begin{aligned} \cos A &= \frac{b^2 + c^2 - a^2}{2bc} \\ &= \frac{36k^2 + 25k^2 - 49k^2}{2(6k)(5k)} \\ &= \frac{12k^2}{60k^2} = \frac{1}{5} \end{aligned}$$

$$\therefore \cos A = \frac{1}{5}$$

13.



Let $AC = n$, $AB = n+1$, $BC = n+2$

\therefore Largest angle is A and smallest angle is B.

$\therefore A = 2B$

Since, $A + B + C = 180^\circ$

$\therefore 3B + C = 180^\circ$

$$\Rightarrow C = 180^\circ - 3B$$

$$\Rightarrow \sin C = \sin(180^\circ - 3B) = \sin 3B$$

$$\Rightarrow \frac{\sin A}{n+2} = \frac{\sin B}{n} = \frac{\sin C}{n+1}$$

$$\Rightarrow \frac{\sin 2B}{n+2} = \frac{\sin B}{n} = \frac{\sin 3B}{n+1}$$

$$\Rightarrow \frac{2\sin B \cos B}{n+2} = \frac{\sin B}{n} = \frac{3\sin B - 4\sin^3 B}{n+1}$$

$$\Rightarrow \frac{2\cos B}{n+2} = \frac{1}{n} = \frac{3-4\sin^2 B}{n+1}$$

$$\therefore \cos B = \frac{n+2}{2n}, 3-4\sin^2 B = \frac{n+1}{n}$$

$$\therefore 3-4(1-\cos^2 B) = \frac{n+1}{n}$$

$$\therefore 3-4+4\left(\frac{n+2}{2n}\right)^2 = \frac{n+1}{n}$$

$$\Rightarrow -1 + \frac{n^2 + 4n + 4}{n^2} = \frac{n+1}{n}$$

$$\Rightarrow -n^2 + n^2 + 4n + 4 = n^2 + n$$



$$\Rightarrow n^2 - 3n - 4 = 0$$

$$\Rightarrow (n + 1)(n - 4) = 0$$


$$\Rightarrow n = -1 \text{ or } n = 4$$

But n cannot be negative.

$$\therefore n = 4$$

\therefore The sides of the Δ are 4, 5, 6.

14.



In ΔODC , $OD = OC = r$, $\angle DOC = \frac{360^\circ}{5} = 72^\circ$

$$\therefore A(\Delta ODC) = \frac{1}{2} r \cdot r \cdot \sin 72^\circ = \frac{1}{2} r^2 \sin 72^\circ$$

$$\therefore A_2 = \text{Area of pentagon} = \frac{5}{2} r^2 \sin 72^\circ$$

$A_1 = \text{Area of circle} = \pi r^2$

$$\therefore \frac{A_1}{A_2} = \frac{\pi r^2}{\frac{5}{2} r^2 \sin 72^\circ}$$

$$= \frac{2\pi}{5 \cos 18^\circ} = \frac{2\pi}{5} \sec 18^\circ = \frac{2\pi}{5} \sec \frac{\pi}{10}$$

15. Let $a = 4k$, $b = 5k$, $c = 6k$

$$\text{Now, } s = \frac{a + b + c}{2} = \frac{4k + 5k + 6k}{2} = \frac{15k}{2}$$

$$\Delta = \sqrt{s(s-a)(s-b)(s-c)}$$

$$= \sqrt{\frac{15k}{2} \left(\frac{15k}{2} - 4k \right) \left(\frac{15k}{2} - 5k \right) \left(\frac{15k}{2} - 6k \right)}$$

$$= \sqrt{\frac{15k}{2} \times \frac{7k}{2} \times \frac{5k}{2} \times \frac{3k}{2}} = \frac{15\sqrt{7}}{4} k^2$$

By sine Rule, $\frac{a}{\sin A} = 2R \therefore \sin A = \frac{a}{2R}$

$$\Delta = \frac{1}{2} bc \sin A$$

$$\Rightarrow \Delta = \frac{1}{2} bc \cdot \frac{a}{2R} = \frac{abc}{4R}$$

$$\therefore R = \frac{abc}{4\Delta} = \frac{4k \cdot 5k \cdot 6k}{15\sqrt{7}k^2} = \frac{8}{\sqrt{7}} k$$

Also $\Delta = rs$, where $r = \text{Radius of incircle of } \Delta ABC$

$$\therefore r = \frac{\Delta}{s} = \frac{\frac{15\sqrt{7}}{4} \cdot k^2}{\frac{15k}{2}} = \frac{\sqrt{7}}{2} k$$

$$\therefore \frac{R}{r} = \frac{8}{\sqrt{7}} k \times \frac{2}{\sqrt{7}k} = \frac{16}{7}$$

$$\therefore \frac{R}{r} = \frac{16}{7}$$

16. $\cos A = \frac{b^2 + c^2 - a^2}{2bc}$

$$\Rightarrow \cos 30^\circ = \frac{4 + 3 - a^2}{4\sqrt{3}}$$

$$\Rightarrow \frac{\sqrt{3}}{2} = \frac{7 - a^2}{4\sqrt{3}} \Rightarrow 7 - a^2 = 6$$

$$\Rightarrow a^2 = 1$$

$$\Rightarrow a = 1 \quad \dots [\because a \neq -1]$$

$$\Delta = \frac{1}{2} bc \sin A = \frac{1}{2} \times 2 \times \sqrt{3} \times \sin 30^\circ$$

$$= \sqrt{3} \times \frac{1}{2} = \frac{\sqrt{3}}{2}$$

$$s = \frac{a + b + c}{2} = \frac{1 + 2 + \sqrt{3}}{2} = \frac{3 + \sqrt{3}}{2}$$

$\Delta = rs$

$$\therefore r = \frac{\Delta}{s} = \frac{\frac{\sqrt{3}}{2} \times \frac{2}{3 + \sqrt{3}}}{\frac{3 + \sqrt{3}}{2}}$$

$$= \frac{\sqrt{3}(3 - \sqrt{3})}{9 - 3} = \frac{3\sqrt{3} - 3}{6} = \frac{\sqrt{3} - 1}{2}$$

17. $a^4 + b^4 + c^4 = 2a^2(b^2 + c^2)$

$$\therefore a^4 + b^4 + c^4 - 2a^2b^2 - 2a^2c^2 = 0$$

$$\therefore a^4 + b^4 + c^4 - 2a^2b^2 + 2b^2c^2 - 2a^2c^2 = 2b^2c^2$$


$$\therefore (b^2 + c^2 - a^2)^2 = (\sqrt{2}bc)^2$$

$$\therefore b^2 + c^2 - a^2 = \sqrt{2}bc$$

$$\therefore \cos A = \frac{b^2 + c^2 - a^2}{2bc} = \frac{\sqrt{2}bc}{2bc} = \frac{1}{\sqrt{2}}$$

$$\therefore \angle A = 45^\circ$$

18.





Let length of altitude = p

Since, $A + B + C = \pi$

$$\therefore A + \frac{\pi}{8} + \frac{5\pi}{8} = \pi$$

$$\therefore A = \pi - \frac{\pi}{8} - \frac{5\pi}{8} = \frac{\pi}{4}$$

$$\text{Area of } \Delta = \frac{1}{2}ap = \frac{1}{2}bc \sin A$$

$$\therefore ap = bc \sin \frac{\pi}{4}$$

$$\therefore ap = bc \times \frac{1}{\sqrt{2}}$$

$$\therefore p = \frac{bc}{\sqrt{2}a} \quad \dots(i)$$

By sine rule,

$$\frac{a}{\sin \frac{\pi}{4}} = \frac{b}{\sin \frac{\pi}{8}} = \frac{c}{\sin \frac{5\pi}{8}}$$

$$\therefore b = \frac{a \sin \frac{\pi}{8}}{\frac{1}{\sqrt{2}}} = \sqrt{2}a \sin \frac{\pi}{8}$$

$$c = \frac{a \sin \frac{5\pi}{8}}{\frac{1}{\sqrt{2}}} = \sqrt{2}a \sin \frac{5\pi}{8}$$

\therefore From (i),

$$p = \frac{\sqrt{2}a \sin \frac{\pi}{8} \cdot \sqrt{2}a \sin \frac{5\pi}{8}}{\sqrt{2}a} = \sqrt{2}a \sin \frac{5\pi}{8} \sin \frac{\pi}{8}$$

$$= \frac{\sqrt{2}a}{2} \left(2 \sin \frac{5\pi}{8} \sin \frac{\pi}{8} \right)$$

$$= \frac{a}{\sqrt{2}} \left[\cos \left(\frac{5\pi}{8} - \frac{\pi}{8} \right) - \cos \left(\frac{5\pi}{8} + \frac{\pi}{8} \right) \right]$$

$$= \frac{a}{\sqrt{2}} \left[\cos \frac{\pi}{2} - \cos \frac{3\pi}{4} \right]$$

$$= \frac{a}{\sqrt{2}} \left[0 - \left(-\frac{1}{\sqrt{2}} \right) \right]$$

$$\therefore p = \frac{a}{2}$$

19. $\tan \frac{A}{2}$ and $\tan \frac{B}{2}$ are the roots of the quadratic equation $6x^2 - 5x + 1 = 0$

$$\therefore \tan \frac{A}{2} + \tan \frac{B}{2} = \frac{5}{6}, \tan \frac{A}{2} \cdot \tan \frac{B}{2} = \frac{1}{6}$$

$$\therefore \tan \left(\frac{A}{2} + \frac{B}{2} \right) = \frac{\tan \frac{A}{2} + \tan \frac{B}{2}}{1 - \tan \frac{A}{2} \tan \frac{B}{2}} = \frac{\frac{5}{6}}{1 - \frac{1}{6}} = \frac{\frac{5}{6}}{\frac{5}{6}} = 1$$

$$\therefore \tan \left(\frac{A+B}{2} \right) = 1$$

$$\therefore \frac{A+B}{2} = \frac{\pi}{4}$$

$$\therefore A+B = \frac{\pi}{2} \quad \therefore \angle C = \frac{\pi}{2}$$

$\therefore \Delta ABC$ is a right angled triangle.

$$20. r = \frac{\Delta}{s} = \frac{\frac{1}{2}ac \sin B}{\frac{1}{2}(a+b+c)} = \frac{ac}{a+b+c}$$

$$\dots [\because \sin B = \sin 90^\circ = 1]$$

$$\begin{aligned} \therefore r &= \frac{ac}{a+c+b} \times \frac{a+c-b}{a+c-b} \\ &= \frac{ac(a+c-b)}{(a+c)^2 - b^2} = \frac{ac(a+c-b)}{a^2 + c^2 + 2ac - b^2} \\ &= \frac{a+c-b}{2} \quad \dots [\because a^2 + c^2 = b^2] \end{aligned}$$

$$\therefore \text{Diameter} = a + c - b$$

21. $\angle A = 55^\circ$, $\angle B = 15^\circ$, $\angle C = 110^\circ$

$$\therefore \frac{a}{\sin 55^\circ} = \frac{b}{\sin 15^\circ} = \frac{c}{\sin 110^\circ} = k$$

$$\therefore a = k \sin 55^\circ, b = k \sin 15^\circ, c = k \sin 110^\circ$$

$$\therefore c^2 - a^2 = k^2 \sin^2 110^\circ - k^2 \sin^2 55^\circ$$

$$= k^2 (\sin 110^\circ + \sin 55^\circ) (\sin 110^\circ - \sin 55^\circ)$$

$$= k^2 \left(2 \sin \frac{165^\circ}{2} \cos \frac{55^\circ}{2} \right) \left(2 \sin \frac{55^\circ}{2} \cos \frac{165^\circ}{2} \right)$$

$$= k^2 \sin 165^\circ \sin 55^\circ$$

$$= k^2 \sin 15^\circ \sin 55^\circ$$

$$= (k \sin 55^\circ) (k \sin 15^\circ)$$

$$= ab$$



22. A, B, C are in A.P.
 $\therefore A + C = 2B$
 Also, $A + B + C = 180^\circ$
 $\therefore \angle B = 60^\circ$
 $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c} = k$
 $\therefore \sin A = ak, \sin B = bk, \sin C = ck$
 $\therefore \frac{a}{c} \sin 2C + \frac{c}{a} \sin 2A$
 $= \frac{a}{c} (2 \sin C \cos C) + \frac{c}{a} (2 \sin A \cos A)$
 $= \frac{a}{c} (2 ck \cos C) + \frac{c}{a} (2ak \cos A)$
 $= 2ka \cos C + 2kc \cos A$
 $= 2k(a \cos C + c \cos A)$
 $= 2kb \quad \dots[\because b = a \cos C + c \cos A]$
 $= 2 \sin B$
 $= 2 \times \frac{\sqrt{3}}{2} \quad \dots[\because \angle B = 60^\circ]$
 $= \sqrt{3}$

23. $2 \cot^{-1} 3 = 2 \tan^{-1} \left(\frac{1}{3} \right) = \tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{3}$
 $= \tan^{-1} \left(\frac{\frac{1}{3} + \frac{1}{3}}{1 - \frac{1}{3} \cdot \frac{1}{3}} \right)$
 $= \tan^{-1} \left(\frac{3+3}{9-1} \right)$
 $= \tan^{-1} \left(\frac{6}{8} \right)$
 $= \tan^{-1} \frac{3}{4}$
 $\therefore \cot \left(\frac{\pi}{4} - 2 \cot^{-1} 3 \right) = \frac{1}{\tan \left(\frac{\pi}{4} - \tan^{-1} \frac{3}{4} \right)}$
 $= \frac{1 + \tan \frac{\pi}{4} \tan \left(\tan^{-1} \frac{3}{4} \right)}{\tan \frac{\pi}{4} - \tan \left(\tan^{-1} \frac{3}{4} \right)}$
 $= \frac{1 + 1 \cdot \frac{3}{4}}{1 - \frac{3}{4}} = \frac{4+3}{4-3} = 7$

24. Let $\frac{1}{2} \cos^{-1} \left(\frac{a}{b} \right) = \theta$
 $\therefore \cos^{-1} \left(\frac{a}{b} \right) = 2\theta$
 $\therefore \cos 2\theta = \frac{a}{b}$
 $\therefore \tan \left[\frac{\pi}{4} + \frac{1}{2} \cos^{-1} \left(\frac{a}{b} \right) \right] + \tan \left[\frac{\pi}{4} - \frac{1}{2} \cos^{-1} \left(\frac{a}{b} \right) \right]$
 $= \tan \left(\frac{\pi}{4} + \theta \right) + \tan \left(\frac{\pi}{4} - \theta \right)$
 $= \frac{1 + \tan \theta}{1 - \tan \theta} + \frac{1 - \tan \theta}{1 + \tan \theta}$
 $= \frac{(1 + \tan \theta)^2 + (1 - \tan \theta)^2}{1 - \tan^2 \theta}$
 $= \frac{2(1 + \tan^2 \theta)}{1 - \tan^2 \theta}$
 $= \frac{2}{\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}} = \frac{2}{\cos 2\theta} = \frac{2}{\frac{a}{b}} = \frac{2b}{a}$

25. $\cos^{-1} \alpha - \cos^{-1} \beta = \cos^{-1} \left[\alpha\beta + \sqrt{1 - \alpha^2} \sqrt{1 - \beta^2} \right]$

Given, $\cos^{-1} x - \cos^{-1} \frac{y}{2} = \alpha$

$\therefore \cos^{-1} \left[\frac{xy}{2} + \sqrt{(1 - x^2)} \left(1 - \frac{y^2}{4} \right) \right] = \alpha$

$\therefore \cos \alpha = \frac{xy}{2} + \sqrt{(1 - x^2)} \left(1 - \frac{y^2}{4} \right)$

$\therefore \sqrt{(1 - x^2)} \left(1 - \frac{y^2}{4} \right) = \cos \alpha - \frac{xy}{2}$

$\therefore 2 \sqrt{(1 - x^2)} \left(1 - \frac{y^2}{4} \right) = 2 \cos \alpha - xy$

Squaring on both sides, we get

$4(1 - x^2) \left(1 - \frac{y^2}{4} \right) = 4 \cos^2 \alpha - 4xy \cos \alpha + x^2 y^2$

$\therefore 4 - y^2 - 4x^2 + x^2 y^2 = 4 \cos^2 \alpha - 4xy \cos \alpha + x^2 y^2$

$\therefore 4x^2 + y^2 - 4xy \cos \alpha = 4 - 4 \cos^2 \alpha$

$\therefore 4x^2 + y^2 - 4xy \cos \alpha = 4 \sin^2 \alpha$



$$26. \quad \sin^{-1} x + \sin^{-1} 2x = \frac{\pi}{3}$$

$$\therefore \sin^{-1} 2x = \frac{\pi}{3} - \sin^{-1} x$$

$$\therefore 2x = \sin\left(\frac{\pi}{3} - \sin^{-1} x\right)$$

$$= \sin \frac{\pi}{3} \cos(\sin^{-1} x) - \cos \frac{\pi}{3} \sin(\sin^{-1} x)$$

$$\Rightarrow 2x = \frac{\sqrt{3}}{2} \cos(\sin^{-1} x) - \frac{1}{2} \cdot x \quad \dots (i)$$

$$\text{Let } \sin^{-1} x = \theta$$

$$\therefore \sin \theta = x$$

$$\cos \theta = \sqrt{1-x^2}$$

$$\therefore \cos(\sin^{-1} x) = \sqrt{1-x^2} \quad \dots (ii)$$

From (i) and (ii), we get

$$2x = \frac{\sqrt{3}}{2} \cdot \sqrt{1-x^2} - \frac{1}{2}x$$

$$\therefore 4x = \sqrt{3}\sqrt{1-x^2} - x$$

$$\therefore 5x = \sqrt{3}\sqrt{1-x^2}$$

$$\therefore 25x^2 = 3 - 3x^2 \quad (\text{squaring both sides})$$

$$\therefore 28x^2 = 3$$

$$\therefore x^2 = \frac{3}{28}$$

$$\therefore x = \sqrt{\frac{3}{28}} = \sqrt{\frac{1}{4} \cdot \frac{3}{7}} = \frac{1}{2} \sqrt{\frac{3}{7}}$$

(From the given relation it can be seen that x is positive)

$$27. \quad \text{L.H.S.} = \sin^{-1}\left(\sin \frac{33\pi}{7}\right) + \cos^{-1}\left(\cos \frac{46\pi}{7}\right)$$

$$+ \tan^{-1}\left(-\tan \frac{13\pi}{8}\right) + \cot^{-1}\left(-\cot \frac{19\pi}{8}\right)$$

$$= \sin^{-1}\left[\sin\left(5\pi - \frac{2\pi}{7}\right)\right] + \cos^{-1}\left[\cos\left(7\pi - \frac{3\pi}{7}\right)\right]$$

$$+ \tan^{-1}\left[-\tan\left(2\pi - \frac{3\pi}{8}\right)\right]$$

$$+ \cot^{-1}\left[-\cot\left(3\pi - \frac{5\pi}{8}\right)\right]$$

$$= \sin^{-1}\left(\sin \frac{2\pi}{7}\right) + \cos^{-1}\left(-\cos \frac{3\pi}{7}\right)$$

$$+ \tan^{-1}\left(\tan \frac{3\pi}{8}\right) + \cot^{-1}\left(\cot \frac{5\pi}{8}\right)$$

$$= \frac{2\pi}{7} + \pi - \frac{3\pi}{7} + \frac{3\pi}{8} + \frac{5\pi}{8}$$

$$\dots [\because \cos^{-1}(-x) = \pi - \cos^{-1}x]$$

$$= \pi - \frac{\pi}{7} + \pi = 2\pi - \frac{\pi}{7} = \frac{13\pi}{7}$$

$$\therefore \frac{13\pi}{7} = \frac{a\pi}{b}$$

$$\therefore a = 13, b = 7$$

$$\therefore a + b = 13 + 7 = 20$$

$$28. \quad \sin^{-1} \frac{4}{5} + \sin^{-1} \frac{5}{13} + \sin^{-1} \frac{16}{65}$$

$$= \sin^{-1} \left[\frac{4}{5} \sqrt{1 - \left(\frac{5}{13}\right)^2} + \frac{5}{13} \sqrt{1 - \left(\frac{4}{5}\right)^2} \right]$$

$$+ \sin^{-1} \frac{16}{65}$$

$$= \sin^{-1} \left[\frac{4}{5} \times \frac{12}{13} + \frac{5}{13} \times \frac{3}{5} \right] + \sin^{-1} \frac{16}{65}$$

$$= \sin^{-1} \left(\frac{48+15}{65} \right) + \sin^{-1} \left(\frac{16}{65} \right)$$

$$= \sin^{-1} \left(\frac{63}{65} \right) + \sin^{-1} \left(\frac{16}{65} \right)$$

$$= \cos^{-1} \left(\sqrt{1 - \left(\frac{63}{65}\right)^2} \right) + \sin^{-1} \left(\frac{16}{65} \right)$$

$$= \cos^{-1} \left(\frac{16}{65} \right) + \sin^{-1} \left(\frac{16}{65} \right)$$

$$= \frac{\pi}{2}$$

$$29. \quad \sqrt{2} = 1.414$$

$$\therefore 2\sqrt{2} - 1 = 2 \times 1.414 - 1 = 2.828 - 1 = 1.828$$

$$\therefore 2\sqrt{2} - 1 > \sqrt{3} \quad \dots [\because \sqrt{3} = 1.732]$$

$$\therefore \tan^{-1}(2\sqrt{2} - 1) > \tan^{-1}(\sqrt{3})$$

\dots [\because $\tan^{-1} x$ is an increasing function]

$$\therefore 2 \tan^{-1}(2\sqrt{2} - 1) > 2 \times \frac{\pi}{3}$$

$$\therefore A > \frac{2\pi}{3} \quad \dots (i)$$

$$\sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta$$

$$\therefore 3\theta = \sin^{-1}(3 \sin \theta - 4 \sin^3 \theta)$$



$$\text{Put } \sin \theta = \frac{1}{3}$$

$$\therefore \theta = \sin^{-1} \left(\frac{1}{3} \right)$$

$$\begin{aligned} \therefore 3 \sin^{-1} \left(\frac{1}{3} \right) &= \sin^{-1} \left[3 \cdot \frac{1}{3} - 4 \left(\frac{1}{3} \right)^3 \right] \\ &= \sin^{-1} \left(1 - \frac{4}{27} \right) \\ &= \sin^{-1} \left(\frac{23}{27} \right) = \sin^{-1} (0.852) \end{aligned}$$

$$\frac{\sqrt{3}}{2} = \frac{1.732}{2} = 0.866, 0.852 < 0.866$$

$$\therefore \sin^{-1} (0.852) < \sin^{-1} (0.866)$$

....[$\because \sin^{-1} x$ is also an increasing function]

$$\therefore 3 \sin^{-1} \left(\frac{1}{3} \right) < \sin^{-1} \left(\frac{\sqrt{3}}{2} \right)$$

$$\therefore 3 \sin^{-1} \left(\frac{1}{3} \right) < \frac{\pi}{3} \quad \dots(\text{ii})$$

$$\sin^{-1} \left(\frac{3}{5} \right) = \sin^{-1} (0.6) < \sin^{-1} \left(\frac{\sqrt{3}}{2} \right)$$

$$\therefore \sin^{-1} \left(\frac{3}{5} \right) < \frac{\pi}{3} \quad \dots(\text{iii})$$

From (ii) and (iii), we get

$$B = 3 \sin^{-1} \left(\frac{1}{3} \right) + \sin^{-1} \left(\frac{3}{5} \right) < \frac{\pi}{3} + \frac{\pi}{3} = \frac{2\pi}{3}$$

$$\therefore B < \frac{2\pi}{3} \quad \dots(\text{iv})$$

From (i) and (iv), $A > B$

$$30. \cot^{-1} x + \cot^{-1} y + \cot^{-1} z = \frac{\pi}{2}$$

$$\therefore \frac{\pi}{2} - \tan^{-1} x + \frac{\pi}{2} - \tan^{-1} y + \frac{\pi}{2} - \tan^{-1} z = \frac{\pi}{2}$$

$$\therefore \tan^{-1} x + \tan^{-1} y + \tan^{-1} z = \pi$$

$$\therefore \tan (\tan^{-1} x + \tan^{-1} y + \tan^{-1} z) = \tan \pi = 0$$

$$\text{Let } A = \tan^{-1} x, B = \tan^{-1} y, C = \tan^{-1} z$$

$$\therefore \tan (A + B + C) = \frac{\tan(A + B) + \tan C}{1 - \tan(A + B) \tan C}$$

$$= \frac{\frac{\tan A + \tan B}{1 - \tan A \tan B} + \tan C}{1 - \frac{\tan A + \tan B}{1 - \tan A \tan B} \cdot \tan C}$$

$$= \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan A \tan B - \tan B \tan C - \tan C \tan A}$$

$$\therefore \tan (A + B + C) = 0$$

$$\Rightarrow \tan A + \tan B + \tan C = \tan A \tan B \tan C$$

$$\therefore \tan (\tan^{-1} x) + \tan (\tan^{-1} y) + \tan (\tan^{-1} z)$$

$$= \tan (\tan^{-1} x) \tan (\tan^{-1} y) \tan (\tan^{-1} z)$$

$$\therefore x + y + z = xyz$$

$$31. \cos^{-1} \left\{ \frac{1}{\sqrt{2}} \left(\cos \frac{9\pi}{10} - \sin \frac{9\pi}{10} \right) \right\}$$

$$= \cos^{-1} \left[\cos \frac{\pi}{4} \cos \frac{9\pi}{10} - \sin \frac{\pi}{4} \sin \frac{9\pi}{10} \right]$$

$$= \cos^{-1} \left[\cos \left(\frac{\pi}{4} + \frac{9\pi}{10} \right) \right]$$

$$= \cos^{-1} \left[\cos \left(\frac{5\pi + 18\pi}{20} \right) \right]$$

$$= \cos^{-1} \left[\cos \left(\frac{23\pi}{20} \right) \right]$$

$$= \cos^{-1} \left[\cos \left(2\pi - \frac{23\pi}{20} \right) \right]$$

$$= \cos^{-1} \left[\cos \left(\frac{17\pi}{20} \right) \right] \text{ and } 0 \leq \frac{17\pi}{20} \leq \pi$$

$$= \frac{17\pi}{20}$$

$$\therefore \text{Principal value is } \frac{17\pi}{20}.$$

$$32. \tan^{-1} 2 + \tan^{-1} 3 = \pi + \tan^{-1} \left(\frac{2+3}{1-2 \times 3} \right)$$

....[$\because 2 \times 3 > 1$]

$$= \pi + \tan^{-1} (-1)$$

$$= \pi - \tan^{-1} 1$$

$$\therefore \tan^{-1} 1 + \tan^{-1} 2 + \tan^{-1} 3 = \pi$$



$$33. \tan^{-1} \frac{1}{1+2x} + \tan^{-1} \frac{1}{4x+1} = \tan^{-1} \frac{2}{x^2}$$

$$\therefore \tan^{-1} \left(\frac{\frac{1}{1+2x} + \frac{1}{4x+1}}{1 - \frac{1}{1+2x} \cdot \frac{1}{4x+1}} \right) = \tan^{-1} \frac{2}{x^2}$$

$$\therefore \frac{4x+1+2x+1}{(1+2x)(4x+1)-1} = \frac{2}{x^2}$$

$$\therefore \frac{6x+2}{4x+8x^2+1+2x-1} = \frac{2}{x^2}$$

$$\therefore x^2(6x+2) = 2(8x^2+6x)$$

$$\therefore 6x^3 + 2x^2 - 16x^2 - 12x = 0$$

$$\therefore 6x^3 - 14x^2 - 12x = 0$$

$$\therefore 3x^3 - 7x^2 - 6x = 0$$

$$\therefore x(3x^2 - 7x - 6) = 0$$

$$\therefore x(x-3)(3x+2) = 0$$

$$\therefore x = 0, 3, -\frac{2}{3}$$

$$\text{But } x > 0, \quad \therefore x = 3$$

$$34. \cot^{-1} x + \sin^{-1} \frac{1}{\sqrt{5}} = \frac{\pi}{4}$$

$$\therefore \tan^{-1} \frac{1}{x} + \tan^{-1} \frac{\frac{1}{\sqrt{5}}}{\sqrt{1-\frac{1}{5}}} = \frac{\pi}{4}$$

$$\dots \left[\because \sin^{-1} x = \tan^{-1} \frac{x}{\sqrt{1-x^2}} \right]$$

$$\therefore \tan^{-1} \frac{1}{x} + \tan^{-1} \frac{1}{2} = \frac{\pi}{4}$$

$$\therefore \tan^{-1} \left(\frac{\frac{1}{x} + \frac{1}{2}}{1 - \frac{1}{x} \cdot \frac{1}{2}} \right) = \frac{\pi}{4}$$

$$\therefore \frac{2+x}{2x-1} = \tan \frac{\pi}{4} = 1$$

$$\therefore 2+x = 2x-1$$

$$\therefore x = 3$$

04 Pair of Straight Lines



Hints



Classical Thinking

- Joint equation of pair of lines having slopes m_1 and m_2 and passing through the origin is $y^2 - (m_1 + m_2)xy + m_1m_2x^2 = 0$
 $\Rightarrow 3x^2 - 4xy + y^2 = 0$
Alternate method:
 Equations of the lines are $y = x$ and $y = 3x$ respectively.
 i.e. $y - x = 0$ and $y - 3x = 0$
 \therefore the combined equation of the pair of lines is $(y - x)(y - 3x) = 0$
 $\therefore y^2 - 3xy - xy + 3x^2 = 0 \Rightarrow 3x^2 - 4xy + y^2 = 0$
- The required equation is $y^2 - \left(\frac{8}{3}\right)xy - x^2 = 0$
 $\Rightarrow 3x^2 + 8xy - 3y^2 = 0$
- The required equation is $y^2 - \frac{3}{2}xy - x^2 = 0$
 $\Rightarrow 2x^2 + 3xy - 2y^2 = 0$
- $x^2 + xy - 12y^2 = 0$
 $\Rightarrow x^2 + 4xy - 3xy - 12y^2 = 0$
 $\Rightarrow x(x + 4y) - 3y(x + 4y) = 0$
 $\Rightarrow (x - 3y)(x + 4y) = 0$
 $\Rightarrow x - 3y = 0$ and $x + 4y = 0$
- It is a homogeneous equation of degree 2 in x and y .
 \therefore Correct option is (C).
- $3x^2 - 10xy - 8y^2 = 0$
 $\Rightarrow 3x^2 - 12xy + 2xy - 8y^2 = 0$
 $\Rightarrow 3x(x - 4y) + 2y(x - 4y) = 0$
 $\Rightarrow (3x + 2y)(x - 4y) = 0$
 $\Rightarrow 3x + 2y = 0$ and $x - 4y = 0$
- $6x^2 - 5xy + y^2 = 0$
 $\Rightarrow 6x^2 - 3xy - 2xy + y^2 = 0$
 $\Rightarrow 3x(2x - y) - y(2x - y) = 0$
 $\Rightarrow (2x - y)(3x - y) = 0$
 $\Rightarrow 3x - y = 0$ and $2x - y = 0$

- Equation of straight lines parallel to $ax^2 + 2hxy + by^2 = 0$ and passing through point (x_1, y_1) is found by shifting the origin to (x_1, y_1)
 \therefore The required equation is $a(x - x_1)^2 + 2h(x - x_1)(y - y_1) + b(y - y_1)^2 = 0$
- $L_1 = x^2 - y^2 = 0$ represents pair of straight lines passing through the origin
 To find equation of pair of straight lines parallel to L_1 and passing through $(3, 4)$, shift the origin to $(3, 4)$
 $\therefore (x - 3)^2 + (y - 4)^2 = 0$
 $\Rightarrow x^2 + y^2 - 6x - 8y + 25 = 0$
- $L_1: ax^2 + 2hxy + by^2 = 0$
 Equation of any line passing through origin and perpendicular to L_1 is given by $bx^2 - 2hxy + ay^2 = 0$
(interchanging coefficients of x^2 and y^2 and change of sign for xy term)
 \therefore The required equation is $ay^2 - 2hxy + bx^2 = 0$
- The required equation is $-3x^2 + 7xy + 5y^2 = 0$
 i.e. $3x^2 - 7xy - 5y^2 = 0$
- Comparing given equation with $ax^2 + 2hxy + by^2 = 0$, we get $h = \frac{-1}{2}$ and $b = -6$
 \therefore Sum of slopes $= m_1 + m_2 = \frac{-2h}{b}$
 $= \frac{-2\left(\frac{-1}{2}\right)}{-6} = \frac{-1}{-6} = \frac{1}{6}$
- Given equation of pair of lines is $ax^2 + 10xy + y^2 = 0$
 $\therefore A = a, H = 5, B = 1$
 Let the slopes of the lines given by be m_1 and m_2
 $m_1 + m_2 = \frac{-2H}{B}$ and $m_1m_2 = \frac{A}{B}$
 Given that $m_2 = 4m_1$
 $\therefore m_1 + 4m_1 = \frac{-2H}{B} = -10 \Rightarrow m_1 = -2$
 and $m_1 \times 4m_1 = \frac{A}{B} = a \Rightarrow 4m_1^2 = a \Rightarrow a = 16$



14. Given equation of pair of lines is
 $ax^2 + 4xy + y^2 = 0$

$$\therefore A = a, H = 2, B = 1$$

$$m_1 + m_2 = -4 \text{ and } m_1 m_2 = a$$

$$\text{Given that } m_1 = 3m_2$$

$$\therefore 3m_2 + m_2 = -4 \Rightarrow m_2 = -1$$

$$\text{Hence, } m_1 = -3$$

$$\therefore a = (-1)(-3) = 3$$

15. Given equation of pair of lines is
 $ax^2 + (3a + 1)xy + 3y^2 = 0$

$$\therefore A = a, H = \frac{3a+1}{2}, B = 3$$

$$\text{Given that } m_1 = \frac{1}{m_2} \Rightarrow m_1 m_2 = 1$$

$$\text{Now, } m_1 m_2 = \frac{a}{3} \Rightarrow \frac{a}{3} = 1 \Rightarrow a = 3$$

$$\text{Also, } m_1 + m_2 = -\left(\frac{3a+1}{3}\right) = \frac{-10}{3}$$

$$\Rightarrow m_1 + \frac{1}{m_1} = \frac{-10}{3} \Rightarrow 3m_1^2 + 10m_1 + 3 = 0$$

$$\therefore m_1 = \frac{-1}{3} \text{ or } -3.$$

16. Given equation of pair of lines is
 $6x^2 + 41xy - 7y^2 = 0$

$$\therefore a = 6, h = \frac{41}{2}, b = -7$$

α and β are angles made by the two lines with X-axis

\therefore their slopes m_1 and m_2 respectively are

$$m_1 = \tan \alpha \text{ and } m_2 = \tan \beta$$

$$\tan \alpha \cdot \tan \beta = m_1 m_2 = -\frac{6}{7}$$

17. Given equation of pair of lines is
 $6x^2 - xy - y^2 = 0$

$$\therefore a = 6, h = -\frac{1}{2}, b = -1$$

If θ is the acute angle between the pair of lines

$$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$

$$\Rightarrow \tan \theta = \left| \frac{2\sqrt{\frac{1}{4} + 6}}{5} \right| = 1$$

$$\Rightarrow \theta = \tan^{-1}(1) = 45^\circ$$

18. Given equation of pair of lines is
 $\sqrt{3}xy - y^2 = 0$

$$\therefore a = 0, h = \frac{\sqrt{3}}{2}, b = -1$$

$$\therefore \tan \theta = \left| \frac{2\sqrt{\frac{3}{4} - 0}}{0 - 1} \right| = \left| \frac{2 \times \frac{\sqrt{3}}{2}}{-1} \right| = \sqrt{3}$$

$$\Rightarrow \theta = \tan^{-1}(\sqrt{3}) = 60^\circ$$

19. Given equation of pair of lines is
 $11y^2 - 4xy + 4x^2 = 0$

$$\text{i.e. } 4x^2 - 4xy + 11y^2 = 0$$

$$\therefore a = 4, h = -2, b = 11$$

$$\therefore \tan \theta = \left| \frac{2\sqrt{144 - 44}}{4 + 11} \right| = \frac{2(10)}{15} = \frac{4}{3}$$

$$\Rightarrow \theta = \tan^{-1}\left(\frac{4}{3}\right)$$

20. Given equation of pair of lines is
 $2x^2 - 3xy + y^2 = 0$

$$\therefore a = 2, h = \frac{-3}{2}, b = 1$$

$$\therefore \tan \theta = \left| \frac{2\sqrt{\frac{9}{4} - 2}}{3} \right| = \left| \frac{\sqrt{9 - 8}}{3} \right| = \frac{1}{3}$$

$$\therefore \cot \theta = 3 \Rightarrow \theta = \cot^{-1}(3)$$

21. Given equation of pair of lines is
 $x^2(\cos \theta - \sin \theta) + 2xy \cos \theta$

$$+ y^2(\cos \theta + \sin \theta) = 0$$

$$\therefore a = \cos \theta - \sin \theta, h = \cos \theta, b = \cos \theta + \sin \theta$$

The acute angle α between the pair of lines is given by

$$\tan \alpha = \left| \frac{2\sqrt{\cos^2 \theta - (\cos^2 \theta - \sin^2 \theta)}}{2 \cos \theta} \right|$$

$$\Rightarrow \tan \alpha = \tan \theta \Rightarrow \alpha = \theta$$

22. Given equation of pair of lines is
 $x^2 - 4hxy + 3y^2 = 0$

$$\therefore A = 1, H = -2h, B = 3$$

$$\text{Now, } \theta = 60^\circ$$

$$\Rightarrow \tan \theta = \sqrt{3}$$

$$\therefore \tan \theta = \left| \frac{2\sqrt{H^2 - AB}}{A + B} \right|$$

$$\Rightarrow \sqrt{3} = \left| \frac{2\sqrt{4h^2 - 3}}{4} \right| \Rightarrow h = \pm \frac{\sqrt{15}}{2}$$



23. Given equation of pair of lines is

$$3x^2 + 18xy + by^2 = 0$$

$$\therefore a = 3, h = 9, b = b$$

$$\text{Now } \theta = \pi \Rightarrow \tan \theta = 0$$

$$\therefore \tan \theta = \left| \frac{2\sqrt{81 - 3b}}{3 + b} \right|$$

$$\Rightarrow 0 = \left| \frac{2\sqrt{81 - 3b}}{3 + b} \right|$$

$$\Rightarrow 81 = 3b \Rightarrow b = 27$$

24. Given equation of pair of lines is

$$3x^2 + 10xy + 8y^2 = 0$$

$$\therefore a = 3, h = 5, b = 8$$

$$\text{Now } \theta = \tan^{-1}(p) \Rightarrow \tan \theta = p$$

$$\therefore \tan \theta = \left| \frac{2\sqrt{25 - 24}}{11} \right|$$

$$\Rightarrow p = \left| \frac{2}{11} \right| = \frac{2}{11}$$

25. Given equation of pair of lines is

$$3x^2 + 2hxy + y^2 = 0$$

$$\therefore a = 3, h = h, b = 1$$

The two lines are real and coincident if $h^2 - ab = 0$

$$\therefore h^2 - ab = h^2 - 3$$

for these lines to be real and coincident, $h^2 - 3 \geq 0 \Rightarrow h^2 \geq 3$

26. Given equation of pair of lines is

$$9x^2 - 12xy + 4y^2 = 0$$

$$a = 9, h = -6, b = 4$$

$$\text{Now, } h^2 - ab = (-6)^2 - 9 \times 4 = 0$$

\therefore The lines are coincident.

27. The condition for a pair of straight lines to be real and coincident is $h^2 - ab = 0$

$$\text{Consider the equation } 4x^2 - 4xy + y^2 = 0$$

$$\therefore a = 4, h = -2, b = 1$$

$$h^2 - ab = (-2)^2 - (4)(1) = 0$$

\therefore Correct option is (A).

28. Given equation of pair of lines is

$$6x^2 + hxy + 12y^2 = 0$$

$$\therefore A = 6, H = \frac{h}{2}, B = 12$$

Since lines are parallel,

$$\therefore H^2 - AB = 0$$

$$\Rightarrow \frac{h^2}{4} = 6(12) \Rightarrow h^2 = (24)(12)$$

$$\Rightarrow h = \pm 12\sqrt{2}$$

29. Given equation of pair of lines is

$$4x^2 + hxy + y^2 = 0$$

The lines are coincident

$$\therefore H^2 = AB$$

$$\Rightarrow \frac{h^2}{4} = 4(1)$$

$$\Rightarrow h = \pm 4$$

30. Given equation of pair of lines is

$$x^2 + xy + y^2 = 0$$

$$\therefore a = 1, h = \frac{1}{2}, b = 1$$

$$\text{Here, } h^2 - ab = \frac{-3}{4} < 0$$

Hence, the lines are imaginary.

31. Given equation of pair of lines is

$$\lambda y^2 + (1 - \lambda^2)xy - \lambda x^2 = 0$$

$$\therefore a = -\lambda, b = \lambda$$

$$\text{Now } a + b = 0$$

\therefore the lines are perpendicular

\therefore Angle between the lines is 90° .

32. Given equation of pair of lines is

$$xy = 0$$

$$\therefore a = 0, h = \frac{1}{2}, b = 0$$

$$\text{Now, } a + b = 0$$

\therefore the lines are perpendicular to each other.

\therefore angle between the pair of line is 90° .

33. The condition for a pair of straight lines to be perpendicular is $a + b = 0$.

$$\text{Consider the equation } 2x^2 = 2y(2x + y)$$

$$\text{i.e. } 2x^2 - 4xy - 2y^2 = 0$$

$$\therefore a = 2, b = -2$$

$$\therefore a + b = 2 + (-2) = 0$$

\therefore Correct option is (A).

34. It is a homogeneous equation of degree 2 in x and y

Hence, it represents a pair of lines and

$$a + b = 0$$

\therefore lines are perpendicular

35. Given equation of pair of lines is

$$3y^2 + 9xy + kx^2 = 0$$

$$\text{i.e. } kx^2 + 9xy + 3y^2 = 0$$

$$\therefore a = k, b = 3$$

The lines are perpendicular

$$\therefore a + b = 0$$

$$\Rightarrow k + 3 = 0 \Rightarrow k = -3$$



36. Given equation of pair of lines is
 $a^2x^2 + bcy^2 = a(b + c)xy$
 $\therefore A = a^2, B = bc$
 Since the lines are mutually perpendicular,
 $\therefore A + B = 0$
 $\Rightarrow a^2 + bc = 0$
37. Consider $2x^2 + 3xy - 2y^2 + 5x + 5y + 3 = 0$
 Comparing the given equation with
 $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$, we get
 $a = 2, b = -2, c = 3, f = \frac{5}{2}, g = \frac{5}{2}, h = \frac{3}{2}$
 Condition for equation to represent pair of
 lines is $abc + 2fgh - af^2 - bg^2 - ch^2 = 0$
 $\therefore 2(-2)(3) + 2\left(\frac{5}{2}\right)\left(\frac{5}{2}\right)\left(\frac{3}{2}\right)$
 $\quad - 2\left(\frac{5}{2}\right)^2 - (-2)\left(\frac{5}{2}\right)^2 - 3\left(\frac{3}{2}\right)^2$
 $= -12 + \frac{75}{4} - \frac{50}{4} + \frac{50}{4} - \frac{27}{4} = 0$
 \therefore Condition is satisfied
 \therefore Correct answer is option (A).
38. Given equation of pair of lines is
 $y^2 + xy + px^2 - x - 2y = 0$
 $\therefore a = p, b = 1, c = 0, f = -1, g = \frac{-1}{2}, h = \frac{1}{2}$
 The given equation represents pair of straight
 lines if
 $abc + 2fgh - af^2 - bg^2 - ch^2 = 0$
 $\Rightarrow p(1)(0) + 2(-1)\left(\frac{-1}{2}\right)\left(\frac{1}{2}\right) - p(-1)^2$
 $\quad - 1\left(\frac{-1}{2}\right)^2 - 0 = 0$
 $\Rightarrow \frac{1}{2} - p - \frac{1}{4} = 0 \Rightarrow p = \frac{1}{4}$
39. Given equation of pair of lines is
 $6x^2 + 11xy - 10y^2 + x + 31y + k = 0$
 $\therefore a = 6, b = -10, c = k, f = \frac{31}{2}, g = \frac{1}{2}, h = \frac{11}{2}$
 Now, $abc + 2fgh - af^2 - bg^2 - ch^2 = 0$
 $\Rightarrow -6(10)k + 2\left(\frac{31}{2}\right)\left(\frac{1}{2}\right)\left(\frac{11}{2}\right) - 6\left(\frac{31}{2}\right)^2 + 10\left(\frac{1}{2}\right)^2$
 $\quad - k\left(\frac{11}{2}\right)^2 = 0$
 $\Rightarrow -k \frac{361}{4} = \frac{5415}{4} \Rightarrow k = -15$

40. Given equation of pair of lines is
 $x^2 - y^2 - x - \lambda y - 2 = 0$
 $\therefore a = 1, b = -1, c = -2, f = \frac{-\lambda}{2}, g = \frac{-1}{2}, h = 0$
 Now, $abc + 2fgh - af^2 - bg^2 - ch^2 = 0$
 $\therefore 2 - \frac{\lambda^2}{4} + \frac{1}{4} = 0 \Rightarrow \frac{\lambda^2}{4} = \frac{9}{4}$
 $\Rightarrow \lambda^2 = 9$
 $\Rightarrow \lambda = \pm 3$
41. Given equation of pair of lines is
 $3x^2 + 2hxy - 3y^2 - 40x + 30y - 75 = 0$
 $\therefore A = 3, B = -3, C = -75, F = 15, G = -20, H = h$
 Now $ABC + 2FGH - AF^2 - BG^2 - CH^2 = 0$
 $\Rightarrow (3)(-3)(-75) + 2(15)(-20)(h)$
 $\quad - 3(15)^2 - (-3)(-20)^2 - (-75)h^2 = 0$
 $\Rightarrow 675 - 600h - 675 + 1200 + 75h^2 = 0$
 $\Rightarrow h^2 - 8h + 16 = 0$
 $\Rightarrow (h - 4)^2 = 0$
 $\Rightarrow h = 4, 4$
42. Given equation of pair of lines is
 $2x^2 - 10xy + 12y^2 + 5x - 16y - 3 = 0$
 $\therefore a = 2, b = 12, c = -3, f = -8, g = \frac{5}{2}, h = -5$
 Equation of perpendicular drawn from origin
 on $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ is
 $bx^2 - 2hxy + ay^2 = 0$
 $\therefore 12x^2 + 10xy + 2y^2 = 0$
 i.e., $6x^2 + 5xy + y^2 = 0$
43. Given equation of pair of lines is
 $2x^2 - 5xy + 3y^2 + 8x - 9y + 6 = 0$
 $\therefore a = 2, b = 3, c = 6, f = -\frac{9}{2}, g = 4, h = \frac{-5}{2}$
 The point of intersection is given by
 $\left(\frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2}\right)$
 $= \left(\frac{\left(\frac{-5}{2}\right)\left(-\frac{9}{2}\right) - 3(4)}{2(3) - \left(\frac{5}{2}\right)^2}, \frac{4\left(\frac{-5}{2}\right) - 2\left(\frac{-9}{2}\right)}{2(3) - \left(\frac{5}{2}\right)^2}\right) = (3, 4)$
44. Given equation of pair of lines is
 $3x^2 + 10xy + 3y^2 - 15x - 21y + 18 = 0$
 $a = 3, b = 3, c = 18, f = \frac{-21}{2}, g = \frac{-15}{2}, h = 5$



The point of intersection is

$$\left(\frac{(5)\left(\frac{-21}{2}\right) - (3)\left(\frac{-15}{2}\right)}{(3)(3) - (5)^2}, \frac{\left(\frac{-15}{2}\right)(5) - (3)\left(\frac{-21}{2}\right)}{(3)(3) - (5)^2} \right)$$

$$\equiv \left(\frac{15}{8}, \frac{3}{8} \right)$$

45. Given equation of pair of lines is $6x^2 - xy - 12y^2 - 8x + 29y - 14 = 0$

$$a = 6, b = -12, h = \frac{-1}{2}$$

$$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right| = \frac{17}{6} \Rightarrow \theta = \tan^{-1} \left(\frac{17}{6} \right)$$

46. Given equation of pair of lines is

$$x^2 + 2\sqrt{3}xy + 3y^2 - 3x - 3\sqrt{3}y - 4 = 0$$

$$\therefore a = 1, h = \sqrt{3}, b = 3$$

$$\text{Now, } h^2 - ab = (\sqrt{3})^2 - (1)(3) = 0$$

\therefore the lines are parallel.

47. Given equation of pair of lines is

$$4x^2 + 2pxy + 25y^2 + 2x + 5y - 1 = 0$$

$$\therefore a = 4, b = 25, h = p$$

The lines are parallel

$$\therefore h^2 - ab = 0 \Rightarrow h^2 = ab$$

$$\Rightarrow p^2 = 4(25) = 100$$

$$\Rightarrow p = 10$$

48. Given equation of pair of lines is

$$3y^2 - 8xy - 3x^2 - 29x + 3y - 18 = 0$$

$$\therefore a = -3, b = 3$$

$$\text{Now, } a + b = -3 + 3 = 0,$$

\therefore The lines are perpendicular to each other.

49. Given equation of pair of lines is

$$x^2 - y^2 - 2y - 1 = 0$$

$$\therefore a = 1, b = -1$$

$$\text{Now, } a + b = 1 + (-1) = 0$$

\therefore The lines are perpendicular to each other.

50. Given equation of pair of lines is

$$3xy - 4y = 0$$

$$\therefore a = b = 0$$

$$\text{Now } a + b = 0$$

\therefore The lines are perpendicular to each other.

51. Given equation of pair of lines is

$$px^2 - 8xy + 3y^2 + 14x + 2y + q = 0$$

$$a = p, b = 3, c = q, f = 1, g = 7, h = -4$$

This lines are perpendicular if $a + b = 0$

$$\Rightarrow p + 3 = 0 \Rightarrow p = -3$$

Since the equation represents a pair of lines

$$\therefore abc + 2fgh - af^2 - bg^2 - ch^2 = 0$$

$$\Rightarrow -9q - 56 + 3 - 147 - 16q = 0$$

$$\Rightarrow -25q - 200 = 0 \Rightarrow q = -8$$

52. Given equation of pair of lines is

$$ax^2 + 6xy + by^2 - 10x + 10y - 6 = 0$$

$$A = a, B = b, C = -6, F = -5, G = 5, H = 3$$

The lines are perpendicular

$$\therefore a + b = 0 \Rightarrow a = -b$$

Also these lines satisfy the condition

$$ABC + 2FGH - AF^2 - BG^2 - CH^2 = 0$$

$$\Rightarrow 6a^2 + 2(-75) - 25a + 25a + 54 = 0$$

$$\Rightarrow 6a^2 - 96 = 0 \Rightarrow a^2 - 16 = 0 \Rightarrow a = \pm 4$$



Critical Thinking

1. The lines passing through origin and parallel to the given lines are $y = m_1x$ and $y = m_2x$,

\therefore the combined equation is $(y - m_1x)(y - m_2x) = 0$

$$\Rightarrow m_1m_2x^2 - (m_1 + m_2)xy + y^2 = 0$$

2. Given line $2x - y = 0 \Rightarrow$ Slope = 2

Let the slope of required line be m

$$\therefore \tan 30^\circ = \left| \frac{m - 2}{1 + 2m} \right|$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \left| \frac{m - 2}{1 + 2m} \right|$$

$$\Rightarrow m^2 + 16m - 11 = 0 \quad \dots(i)$$

Since, the line passes through origin, its equation is

$$y = mx \Rightarrow m = \frac{y}{x}$$

Substituting of m in equation (i), we get

$$\left(\frac{y}{x} \right)^2 + 16 \left(\frac{y}{x} \right) - 11 = 0$$

$$\Rightarrow 11x^2 - 16xy - y^2 = 0$$

3. From the diagram, the required lines are

$$y = \frac{x}{\sqrt{3}} \quad \text{i.e., } \sqrt{3}y - x = 0$$


and

$$y = \frac{-x}{\sqrt{3}} \quad \text{i.e., } \sqrt{3}y + x = 0$$

\therefore Combined equation is

$$(\sqrt{3}y - x)(\sqrt{3}y + x) = 0$$

$$\text{i.e., } 3y^2 - x^2 = 0$$





4. Let $y = mx$ be the equation of line.
Slope of the given line $y = -x - \sqrt{3}$ is -1
Since, the pair of straight lines and the given line form an equilateral triangle, angle between them is 60° .
- $$\therefore \tan \frac{\pi}{3} = \left| \frac{m+1}{1-m} \right| \Rightarrow \sqrt{3} = \left| \frac{m+1}{1-m} \right|$$
- $$\Rightarrow 3(1-m)^2 = (1+m)^2$$
- $$\Rightarrow 3(1+m^2-2m) = (1+m^2+2m)$$
- $$\therefore m^2 - 4m + 1 = 0 \quad \dots(i)$$
- The equation of line passing through origin is,
 $y = mx \Rightarrow m = \frac{y}{x}$
- Substituting the value of m in (i), we get
- $$\left(\frac{y}{x} \right)^2 - 4 \left(\frac{y}{x} \right) + 1 = 0 \Rightarrow x^2 - 4xy + y^2 = 0$$
5. $ax^2 + (a+b)xy + by^2 + x + y = 0$
 $\Rightarrow ax^2 + bxy + x + axy + by^2 + y = 0$
 $\Rightarrow x(ax + by + 1) + y(ax + by + 1) = 0$
 $\Rightarrow (x+y)(ax + by + 1) = 0$
6. $pq(x^2 - y^2) + (p^2 - q^2)xy = 0$
 $\Rightarrow pqx^2 - pqy^2 + p^2xy - q^2xy = 0$
 $\Rightarrow px(py + qx) - qy(py + qx) = 0$
 $\Rightarrow (px - qy)(py + qx) = 0$
 $\Rightarrow px - qy = 0$ and $py + qx = 0$
- \therefore
- Required equation of the line is
- $px - qy = 0$
7. $x^2 + 6xy = 0 \Rightarrow x(x + 6y) = 0$
 $\Rightarrow x = 0$ and $x + 6y = 0$ are two straight lines.
 $x = 0$ represents Y-axis.
8. $y^2 - x^2 + 2x - 1 = 0$
 $\Rightarrow y^2 - (x^2 - 2x + 1) = 0$
 $\Rightarrow (y - 0)^2 - (x - 1)^2 = 0$
This is equation of pair of straight lines passing through $(1, 0)$.
9. The given equation represents a pair of straight lines passing through $(5, 6)$.
10. The lines pass through $(-2, 2)$
Only $(-2, 2)$ satisfies the given equation.
11. Given equation of pair of lines is
 $ax^2 + xy - by^2 = 0$
Comparing the equations, with
 $Ax^2 + 2Hxy + By^2 = 0$
- $$\therefore A = a, H = \frac{1}{2} \text{ and } B = -a$$
- $$\therefore \text{the equation represents a pair of straight lines for all real values of 'a'.$$

12. The combined equation of pair of straight lines passing through origin and perpendicular to $3x^2 + xy - 2y^2 = 0$ is given by
 $-2x^2 - xy + 3y^2 = 0$
i.e. $2x^2 + xy - 3y^2 = 0$
Since the required lines pass through $(2, -3)$
- $$\therefore \text{By shifting the origin to } (2, -3), \text{ we get}$$
- $$2(x-2)^2 + (x-2)(y+3) - 3(y+3)^2 = 0$$
- $$\Rightarrow 2x^2 + xy - 3y^2 - 5x - 20y - 25 = 0$$
13. Separate equation of lines represented by
 $3x^2 - 8xy + 5y^2 = 0$ are
 $x - y = 0$ and
 $3x - 5y = 0$
Line perpendicular to $x - y = 0$ i.e. $y = x$ and passing through $(1, 2)$ is
 $(y - 2) = -1(x - 1)$
i.e. $x + y - 3 = 0 \quad \dots(i)$
Line perpendicular to $3x - 5y = 0$
i.e. $y = \frac{3}{5}x$ and passing through $(1, 2)$ is
 $(y - 2) = \frac{-5}{3}(x - 1)$
i.e. $5x + 3y - 11 = 0 \quad \dots(ii)$
- \therefore
- combined equation is
-
- $(x + y - 3)(5x + 3y - 11) = 0$
14. Slope of the line $4x + 3y = 0$ is $m = -\frac{4}{3}$
 $kx^2 - 5xy - 6y^2 = 0$
 $\Rightarrow -6m^2 - 5m + k = 0$
 $\Rightarrow -6\left(-\frac{4}{3}\right)^2 - 5\left(-\frac{4}{3}\right) + k = 0$
 $\Rightarrow k - \frac{32}{3} + \frac{20}{3} = 0$
 $\Rightarrow k = \frac{12}{3} \Rightarrow k = 4$
15. Let $y = mx$ be a line common to the given pair of lines,
It satisfies the given equations
- $$\therefore am^2 + 2m + 1 = 0 \text{ and } \dots(i)$$
- $$m^2 + 2m + a = 0 \quad \dots(ii)$$
- On solving (i) and (ii), we get
- $$\frac{m^2}{2(1-a)} = \frac{m}{a^2-1} = \frac{1}{2(1-a)}$$
- $$\therefore m^2 = 1 \text{ and } m = -\left(\frac{a+1}{2}\right)$$
- $$\therefore (a+1)^2 = 4 \Rightarrow a = 1 \text{ or } -3$$



But for $a = 1$ the two pair have both the lines common.

So $a = -3$ and the slope m of the line common to both the pairs is 1.

$$\text{Now } x^2 + 2xy + ay^2 = x^2 + 2xy - 3y^2 \\ = (x - y)(x + 3y)$$

$$\text{and } ax^2 + 2xy + y^2 = -3x^2 + 2xy + y^2 \\ = -(x - y)(3x + y)$$

Thus, required equation is $(x + 3y)(3x + y) = 0$
i.e., $3x^2 + 10xy + 3y^2 = 0$

16. The equation of given lines are

$$ax^2 + 2hxy + by^2 = 0 \quad \dots(i)$$

$$a'x^2 + 2h'xy + b'y^2 = 0 \quad \dots(ii)$$

Let the line common to both be $y = mx$.

It will satisfy both the above equations.

$$\text{Hence, } a + 2mh + bm^2 = 0 \quad \dots(iii)$$

$$\text{and } a' + 2mh' + b'm^2 = 0 \quad \dots(iv)$$

Now eliminating 'm' from the equations (iii) and (iv), we get

$$\frac{m^2}{2ha' - 2h'a} = \frac{-m}{ba' - b'a} = \frac{1}{2bh' - 2b'h}$$

$$\Rightarrow m^2 = \frac{ha' - h'a}{bh' - b'h} \quad \dots(v)$$

$$\text{and } m^2 = \frac{(ab' - ba')^2}{4(bh' - b'h)^2} \quad \dots(vi)$$

From (v) and (vi), we get the required condition.

17. Given equation of pair of lines

$$x^2 - 2xy \tan A - y^2 = 0$$

$$\therefore a = 1, h = -\tan A, b = -1$$

$$m_1 + m_2 = \frac{-2h}{b} \Rightarrow 4 = \frac{2 \tan A}{-1}$$

$$\Rightarrow \tan A = -2$$

$$\Rightarrow \angle A = \tan^{-1}(-2)$$

18. Given equation of pair of lines is

$$ax^2 + 2hxy + by^2 = 0$$

$$\text{Given that } m_1 = 5m_2$$

$$\therefore m_1 + m_2 = 5m_2 + m_2 = \frac{-2h}{b}$$

$$\Rightarrow m_2 = \frac{-h}{3b} \Rightarrow m_2^2 = \frac{h^2}{9b^2} \quad \dots(i)$$

$$m_1 m_2 = (5m_2)m_2 = \frac{a}{b}$$

$$\therefore m_2^2 = \frac{a}{5b} \quad \dots(ii)$$

$$\therefore \text{From (i) and (ii), we get } 5h^2 = 9ab$$

19. Let the gradient of one line be m .

\therefore the gradient of second line is $2m$

We know,

$$m + 2m = \frac{-2h}{b}$$

$$\therefore 3m = \frac{-2h}{b} \Rightarrow m = \frac{-2h}{3b} \quad \dots(i)$$

$$\text{Also, } m \times 2m = \frac{a}{b} \Rightarrow 2m^2 = \frac{a}{b} \quad \dots(ii)$$

\therefore from (i) and (ii), we get

$$2 \left(\frac{-2h}{3b} \right)^2 = \frac{a}{b} \Rightarrow \frac{8h^2}{9b^2} = \frac{a}{b} \Rightarrow ab = \frac{8h^2}{9}$$

20. Given equation of pair of lines is

$$ax^2 + 2hxy + by^2 = 0$$

given that $m_2 = \lambda m_1$

$$\text{Now, } m_1 + m_2 = m_1 + \lambda m_1 = \frac{-2h}{b}$$

$$\Rightarrow m_1 = \frac{-2h}{b(1 + \lambda)} \quad \dots(i)$$

$$m_1 \cdot m_2 = m_1 \cdot \lambda m_1 = \frac{a}{b} \Rightarrow m_1 = \sqrt{\frac{a}{b\lambda}} \quad \dots(ii)$$

\therefore from (i) and (ii), we get

$$\sqrt{\frac{a}{b\lambda}} = \frac{-2h}{b(1 + \lambda)}$$

Squaring both sides, we get

$$4\lambda h^2 = ab(1 + \lambda)^2$$

21. Given equation of pair of lines is

$$ax^2 + 2hxy + by^2 = 0$$

Given that, $m_1 = m_2^2$

$$m_1 m_2 = m_2^2 m_2 = \frac{a}{b}$$

$$\therefore m_2 = \left(\frac{a}{b} \right)^{\frac{1}{3}}$$

$$\text{Also, } m_1 + m_2 = m_2^2 + m_2 = \frac{-2h}{b}$$

$$\therefore \left\{ \left(\frac{a}{b} \right)^{\frac{1}{3}} \right\}^2 + \left(\frac{a}{b} \right)^{\frac{1}{3}} = \frac{-2h}{b}$$

Cubing both sides, we get

$$\left(\frac{a}{b} \right)^2 + \frac{a}{b} + 3 \left(\frac{a}{b} \right)^{\frac{2}{3}} \cdot \left(\frac{a}{b} \right)^{\frac{1}{3}} \cdot \left\{ \left(\frac{a}{b} \right)^{\frac{2}{3}} + \left(\frac{a}{b} \right)^{\frac{1}{3}} \right\} \\ = \frac{-8h^3}{b^3}$$



$$\therefore \left(\frac{a}{b}\right)^2 + \frac{a}{b} - \frac{6ah}{b^2} = \frac{-8h^3}{b^3}$$

$$\dots \left\{ \because \left(\frac{a}{b}\right)^{\frac{1}{3}} + \left(\frac{a}{b}\right)^{\frac{2}{3}} = \frac{-2h}{b} \right\}$$

$$\therefore ab(a+b) - 6abh + 8h^3 = 0$$

22. Given equation of pair of lines is
 $2x^2 - 5xy + 3y^2 = 0$

$$\therefore a = 2, h = \frac{-5}{2}, b = 3$$

$$\therefore m_1 + m_2 = \frac{5}{6} \text{ and } m_1 \cdot m_2 = \frac{2}{3} \quad \dots(i)$$

$$\text{Slopes of lines} = \frac{1}{m_1} \text{ and } \frac{1}{m_2}$$

\(\therefore\) Required equation of pair of lines is

$$y^2 - \left(\frac{1}{m_1} + \frac{1}{m_2}\right)xy + \frac{1}{m_1 m_2}x^2 = 0$$

$$\Rightarrow y^2 - \left(\frac{m_1 + m_2}{m_1 m_2}\right)xy + \frac{1}{m_1 m_2}x^2 = 0$$

$$\Rightarrow y^2 - \left(\frac{\frac{5}{6}}{\frac{2}{3}}\right)xy + \frac{1}{\left(\frac{2}{3}\right)}x^2 = 0$$

$$\Rightarrow 2y^2 - 5xy + 3x^2 = 0$$

23. Let the angle made by one of the lines with X-axis = θ

\(\therefore\) The angle made by other line with Y-axis = θ

$$\therefore m_1 = \tan \theta,$$

$$m_2 = \tan(90^\circ - \theta) = \cot \theta$$

$$\therefore m_1 m_2 = \frac{a}{b} = 1$$

$$\Rightarrow \frac{a}{b} = 1 \Rightarrow a = b$$

24. Given equation of pair of lines is
 $x^2(\sec^2 \theta - \sin^2 \theta) - 2xy \tan \theta + y^2 \sin^2 \theta = 0$

$$\therefore a = \sec^2 \theta - \sin^2 \theta, h = -\tan \theta, b = \sin^2 \theta$$

$$\text{Now, } m_1 + m_2 = \frac{2 \tan \theta}{\sin^2 \theta},$$

$$m_1 m_2 = \frac{\sec^2 \theta - \sin^2 \theta}{\sin^2 \theta}$$

$$\therefore m_1 - m_2 = \sqrt{(m_1 + m_2)^2 - 4m_1 m_2}$$

$$= \sqrt{\left(\frac{2 \tan \theta}{\sin^2 \theta}\right)^2 - 4\left(\frac{\sec^2 \theta - \sin^2 \theta}{\sin^2 \theta}\right)}$$

$$= \sqrt{\frac{4 \tan^2 \theta}{\sin^4 \theta} - 4(\sec^2 \theta \operatorname{cosec}^2 \theta - 1)}$$

$$= \sqrt{4 \sec^2 \theta \operatorname{cosec}^2 \theta - 4 \sec^2 \theta \operatorname{cosec}^2 \theta + 4}$$

$$= 2$$

25. Given equation of pair of lines

$$(\tan^2 \alpha + \cos^2 \alpha)x^2 - 2xy \tan \alpha + \sin^2 \alpha y^2 = 0$$

$$a = \tan^2 \alpha + \cos^2 \alpha, h = -\tan \alpha, b = \sin^2 \alpha$$

If θ_1 and θ_2 are the angles made by lines with X-axis, then $\tan \theta_1 = m_1$ and $\tan \theta_2 = m_2$

$$\text{Now, } m_1 + m_2 = \frac{2 \tan \alpha}{\sin^2 \alpha} = 2 \sec \alpha \operatorname{cosec} \alpha$$

$$m_1 m_2 = \frac{\tan^2 \alpha + \cos^2 \alpha}{\sin^2 \alpha} = \sec^2 \alpha + \cot^2 \alpha$$

$$\therefore m_1 - m_2 = \sqrt{4 \sec^2 \alpha \operatorname{cosec}^2 \alpha - 4(\sec^2 \alpha + \cot^2 \alpha)}$$

$$= \sqrt{4 \sec^2 \alpha (\operatorname{cosec}^2 \alpha - 1) - 4 \cot^2 \alpha}$$

$$= \sqrt{4 \cot^2 \alpha (\sec^2 \alpha - 1)}$$

$$= \sqrt{4 \cot^2 \alpha \tan^2 \alpha}$$

$$= 2$$

26. The equation of one of the lines passing through origin is $y = mx$.

The line makes an angle α with the line $y = x$

$$\therefore \tan \alpha = \pm \left\{ \frac{m_1 - m_2}{1 + m_1 m_2} \right\} = \pm \frac{(m-1)}{1+m}$$

$$\Rightarrow (1+m)^2 \tan^2 \alpha = (m-1)^2$$

$$\Rightarrow m^2 - 2m \left\{ \frac{1 + \tan^2 \alpha}{1 - \tan^2 \alpha} \right\} + 1 = 0$$

$$\Rightarrow m^2 - 2m \sec 2\alpha + 1 = 0$$

$$\dots \left\{ \because \frac{1 + \tan^2 \alpha}{1 - \tan^2 \alpha} = \sec 2\alpha \right\}$$

$$\text{But } m = \frac{y}{x}$$

On eliminating m we get

$$y^2 - 2xy \sec 2\alpha + x^2 = 0.$$

27. Let the equation of one of the line which bisects the angle between the co-ordinate axes be $y = x$

$$\therefore m_1 = \tan 45^\circ = 1$$

Let m_2 be the slope of the other line.

$$\text{Now, } m_1 m_2 = \frac{a}{b}$$



- Since $m_1 = 1$, we get $m_2 = \frac{a}{b}$;
- Also, $m_1 + m_2 = \frac{-2h}{b}$
- $$\Rightarrow 1 + \frac{a}{b} = \frac{-2h}{b}$$
- $$\Rightarrow a + b = -2h$$
28. Let the equation of one of the lines be $y = x$
- $$\therefore m_1 = \tan 45^\circ = 1$$
- Now, $m_1 m_2 = \frac{a}{c}$
- Since $m_1 = 1$, we get $m_2 = \frac{a}{c}$
- Also, $m_1 + m_2 = \frac{-b}{c}$
- $$\therefore 1 + \frac{a}{c} = \frac{-b}{c}$$
- $$\Rightarrow \frac{a+b+c}{c} = 0$$
- $$\Rightarrow a+b+c=0$$
29. Let the equation of one of the angle bisector of co-ordinate axes be $x + y = 0 \Rightarrow m_1 = -1$
- Now, $m_1 m_2 = \frac{a}{b}$
- $$\Rightarrow m_2 = -\frac{a}{b}$$
- Also, $m_1 + m_2 = \frac{-2h}{b}$
- $$\Rightarrow -1 - \frac{a}{b} = \frac{-2h}{b} \Rightarrow (a+b)^2 = 4h^2$$
30. The line makes angles α and β with X-axis
- $$\therefore m_1 = \tan \alpha \text{ and } m_2 = \tan \beta$$
- $$\Rightarrow \cot \alpha = \frac{1}{m_1} \text{ and } \cot \beta = \frac{1}{m_2}$$
- Given equation of pair of lines is
- $$2x^2 - 3xy + y^2 = 0$$
- $$\therefore a = 2, h = \frac{-3}{2}, b = 1$$
- Now, $m_1 + m_2 = 3$ and $m_1 m_2 = 2$
- $$\therefore \cot^2 \alpha + \cot^2 \beta = \frac{1}{m_1^2} + \frac{1}{m_2^2} = \frac{m_1^2 + m_2^2}{(m_1 m_2)^2}$$
- $$= \frac{(m_1 + m_2)^2 - 2m_1 m_2}{(m_1 m_2)^2}$$
- $$= \frac{(3)^2 - 2(2)}{(2)^2} = \frac{5}{4}$$

31. Given equation of pair of lines is
- $$ax^2 - bxy - y^2 = 0$$
- $$\therefore A = a, H = \frac{-b}{2}, B = -1$$
- Since lines make angles α and β with X-axis,
- $$\therefore m_1 = \tan \alpha \text{ and } m_2 = \tan \beta$$
- Now, $m_1 + m_2 = \frac{b}{-1} \Rightarrow \tan \alpha + \tan \beta = -b$
- and $m_1 m_2 = \frac{a}{-1} \Rightarrow \tan \alpha \tan \beta = -a$
- We know, $\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$
- $$= \frac{-b}{1 - (-a)} = \frac{-b}{1+a}$$
32. Given equation of pair of lines is
- $$ax^2 + 2hxy + by^2 = 0$$
- $$\therefore A = a, H = h, B = b$$
- $$\tan \theta = \left(\frac{2\sqrt{H^2 - AB}}{A + B} \right)$$
- $$= \left(\frac{\sqrt{4h^2 - 4ab}}{a + b} \right)$$
- $$= \left(\frac{\sqrt{3a^2 + 3b^2 + 10ab - 4ab}}{a + b} \right)$$
-[$\because 3a^2 + 3b^2 + 10ab = 4h^2$]
- $$\therefore \tan \theta = \left(\frac{\sqrt{3(a+b)^2}}{a+b} \right)$$
- $$\Rightarrow \theta = \tan^{-1}(\sqrt{3})$$
- $$= 60^\circ$$
33. Given equation of pair of lines is
- $$x^2 - 2pxy + y^2 = 0$$
- $$\therefore a = 1, h = -p, b = 1$$
- $$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$
- $$\Rightarrow \tan \theta = \frac{\pm 2\sqrt{p^2 - 1}}{1 + 1} = \pm \sqrt{p^2 - 1}$$
- $$\Rightarrow \tan^2 \theta = p^2 - 1$$
- $$\Rightarrow \sec^2 \theta - 1 = p^2 - 1$$
- $$\Rightarrow \theta = \sec^{-1} p$$



34. Given equation of pair of lines is

$$(x^2 + y^2) \sin \theta + 2xy = 0$$

$$\therefore a = b = \sin \theta, h = 1$$

$$\therefore \tan \theta = \left(\frac{2\sqrt{1 - \sin^2 \theta}}{2 \sin \theta} \right)$$

$$\Rightarrow \theta = \tan^{-1} \left(\frac{\cos \theta}{\sin \theta} \right) = \tan^{-1} (\cot \theta)$$

$$\Rightarrow \theta = \tan^{-1} \left\{ \tan \left(\frac{\pi}{2} - \theta \right) \right\} = \frac{\pi}{2} - \theta$$

35. Given equation of pair of lines is

$$ax^2 + xy + by^2 = 0$$

$$\therefore A = a, H = \frac{1}{2}, B = b$$

$$\text{Now, } \theta = 45^\circ \Rightarrow \tan \theta = 1$$

$$\therefore \tan 45^\circ = \left| \frac{2\sqrt{\frac{1}{4} - ab}}{a + b} \right|$$

$$\Rightarrow (a + b)^2 = (1 - 4ab)$$

$$\Rightarrow a^2 + b^2 + 6ab - 1 = 0$$

The above equation is satisfied by

$$a = 1 \text{ and } b = -6$$

36. Given equation of pair of lines is

$$\therefore a = -\tan^2 A, h = \frac{k}{2}, b = 1$$

$$\therefore \tan 2A = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$

$$\tan 2A = \frac{2\sqrt{\frac{k^2}{4} + \tan^2 A}}{1 - \tan^2 A}$$

$$\Rightarrow \frac{2 \tan A}{1 - \tan^2 A} = \frac{2\sqrt{\frac{k^2}{4} + \tan^2 A}}{1 - \tan^2 A}$$

$$\Rightarrow \frac{k^2}{4} + \tan^2 A = \tan^2 A \Rightarrow k = 0$$

37. Here, $a_1 = a, h_1 = h, b_1 = b,$

$$a_2 = 2, h_2 = \frac{-5}{2}, b_2 = 3$$

Given that $\theta_1 = \theta_2$

$$\Rightarrow \tan \theta_1 = \tan \theta_2$$

$$\Rightarrow \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right| = \left| \frac{2\sqrt{\frac{25}{4} - 6}}{5} \right|$$

$$\Rightarrow \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right| = \left| \frac{1}{5} \right|$$

Squaring both sides, we get

$$4 \times 25(h^2 - ab) = (a + b)^2$$

$$100(h^2 - ab) = (a + b)^2$$

Comparing with given condition,

$$k(h^2 - ab) = (a + b)^2, \text{ we get}$$

$$k = 100$$

38. Comparing the given equations with

$$ax^2 + 2hxy + by^2 = 0, \text{ we get,}$$

$$a_1 = 3, h_1 = \frac{-7}{2}, b_1 = 4$$

$$a_2 = 6, h_2 = \frac{-5}{2}, b_2 = 1$$

If θ_1 and θ_2 are acute angles between the two pairs of lines, then

$$\tan \theta_1 = \left(\frac{2\sqrt{\frac{49}{4} - 12}}{3 + 4} \right) = \frac{1}{7}$$

$$\Rightarrow \theta_1 = \tan^{-1} \left(\frac{1}{7} \right)$$

$$\tan \theta_2 = \left(\frac{2\sqrt{\frac{25}{4} - 6}}{6 + 1} \right) = \left(\frac{1}{7} \right)$$

$$\Rightarrow \theta_2 = \tan^{-1} \left(\frac{1}{7} \right)$$

Hence, $\theta_1 = \theta_2$.

39. Given equation of pair of lines is

$$a^2x^2 + bcy^2 = a(b + c)xy$$

$$\therefore A = a^2, H = \frac{-a(b + c)}{2}, B = bc$$

Since the lines are coincident

$$\therefore H^2 - AB = 0$$

$$\Rightarrow \left\{ \frac{-a(b + c)}{2} \right\}^2 - a^2(bc) = 0$$

$$\Rightarrow a^2(b - c)^2 = 0$$

$$\Rightarrow a = 0 \text{ or } b = c$$

40. Given equation of pair of lines is

$$(p - q)x^2 + 2(p + q)xy + (q - p)y^2 = 0$$

$$\therefore a = p - q, h = p + q, b = q - p$$

Since, the lines are mutually perpendicular

$$\therefore a + b = 0$$

$$\Rightarrow (p - q) + (q - p) = 0$$

The above equation is true for all values of p and q .



41. Given equation of pair of lines is $3ax^2 + 5xy + (a^2 - 2)y^2 = 0$
 $\therefore A = 3a, H = \frac{5}{2}, B = a^2 - 2$
 Since the lines are perpendicular
 $\therefore A + B = 0$
 $\Rightarrow 3a + (a^2 - 2) = 0$
 $\Rightarrow a^2 + 3a - 2 = 0$
 Since, the equation is a quadratic equation in 'a' and $B^2 - 4AC > 0$,
 The roots of 'a' are real and distinct.
 \therefore Lines are perpendicular to each other for two values of 'a'.
42. Given equation of pair of lines is $ay^2 + (-1 - \lambda^2)xy - ax^2 = 0$
 $\therefore A = -a, H = \frac{-1 - \lambda^2}{2}, B = a$
 $A + B = (-a) + a = 0$
 \Rightarrow Angle between the given lines is 90° .
 Now, consider $xy = 0$. Here, $A = B = 0$
 $\Rightarrow A + B = 0$
 \therefore the angle between the lines is 90°
 \therefore Correct option is (C).
43. Given equation of pair of lines is $x^2 + y^2 + 2gx + 2fy + 1 = 0$
 $A = 1, B = 1, C = 1, F = f, G = g, H = 0$
 The given equation represents a pair of lines
 $\therefore ABC + 2FGH - AF^2 - BG^2 - CH^2 = 0$
 $\Rightarrow (1)(1)(1) + 2fg(0) - (1)f^2 - 1(g)^2 - (1)(0)^2 = 0$
 $\Rightarrow f^2 + g^2 = 1$
44. Given equation of pair of lines is $ax^2 + by^2 + cx + cy = 0$
 $\therefore A = a, B = b, C = 0, F = \frac{c}{2}, G = \frac{c}{2}, H = 0$
 Now $ABC + 2FGH - AF^2 - BG^2 - CH^2 = 0$
 $\Rightarrow ab(0) + 2\left(\frac{c}{2}\right)\left(\frac{c}{2}\right)(0) - a\left(\frac{c}{2}\right)^2 - b\left(\frac{c}{2}\right)^2 - 0(0)^2 = 0$
 $\Rightarrow ac^2 + bc^2 = 0$
 $\Rightarrow c^2(a + b) = 0$
 $\Rightarrow c(a + b) = 0$

45. Given equation of pair of lines is $hxy + gx + fy + c = 0$
 $A = B = 0, C = c, F = \frac{f}{2}, G = \frac{g}{2}, H = \frac{h}{2}$
 Now, $ABC + 2FGH - AF^2 - BG^2 - CH^2 = 0$
 $\Rightarrow 0 + 2\left(\frac{f}{2}\right)\left(\frac{g}{2}\right)\left(\frac{h}{2}\right) - 0 - 0 - c\left(\frac{h}{2}\right)^2 = 0$
 $\Rightarrow \frac{fgh}{4} - \frac{ch^2}{4} = 0$
 $\Rightarrow fg = ch$
46. Given equation of pair of lines is $2x^2 + 5xy + 2y^2 + 3x + 3y + 1 = 0$
 $a = 2, b = 2, c = 1, f = \frac{3}{2}, g = \frac{3}{2}, h = \frac{5}{2}$
 $\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right| = \left| \frac{2\sqrt{\left(\frac{25}{4}\right) - 4}}{2 + 2} \right| = \frac{3}{4}$
 $\therefore \cos \theta = \frac{4}{5} \Rightarrow \theta = \cos^{-1}\left(\frac{4}{5}\right)$
47. Given equation of pair of lines is $x^2 - 3xy + \lambda y^2 + 3x - 5y + 2 = 0$
 $a = 1, b = \lambda, c = 2, f = \frac{-5}{2}, g = \frac{3}{2}, h = \frac{-3}{2}$
 $\theta = \tan^{-1}\left(\frac{1}{3}\right) \Rightarrow \tan \theta = \frac{1}{3}$
 Since, $\tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$
 $\Rightarrow \frac{1}{3} = \left| \frac{2\sqrt{\left(\frac{-3}{2}\right)^2 - \lambda}}{\lambda + 1} \right|$
 $\Rightarrow (\lambda + 1)^2 = 9(9 - 4\lambda) \Rightarrow \lambda^2 + 38\lambda - 80 = 0$
 $\Rightarrow (\lambda + 40)(\lambda - 2) = 0 \Rightarrow \lambda = -40, 2$
48. Given equation of pair of lines is $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$
 $\theta = \frac{\pi}{4} \Rightarrow \tan \theta = 1$
 $\therefore 1 = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$
 $\Rightarrow 4(h^2 - ab) = (a + b)^2$
 $\Rightarrow 4h^2 - 4ab = a^2 + 2ab + b^2$
 $\Rightarrow a^2 + 6ab + b^2 = 4h^2$



49. Given equation of pair of lines is
 $x^2 - 3xy + \lambda y^2 + 3x - 5y + 2 = 0$
 $a = 1, b = \lambda, c = 2, f = \frac{-5}{2}, g = \frac{3}{2}, h = \frac{-3}{2}$
 Now, $abc + 2 fgh - af^2 - bg^2 - ch^2 = 0$
 $\Rightarrow 2\lambda + 2\left(\frac{-5}{2}\right)\left(\frac{3}{2}\right)\left(\frac{-3}{2}\right) - \frac{25}{4} - \frac{9\lambda}{4} - \frac{18}{4} = 0$
 $\Rightarrow \lambda = 2$

$$\tan \theta = \frac{2\sqrt{\frac{9}{4} - 2}}{1 + 2} = \frac{1}{3}$$

$$\Rightarrow \cot \theta = 3$$

$$\therefore \operatorname{cosec}^2 \theta = 1 + \cot^2 \theta = 1 + 9 = 10$$

50. Given equation of pair of lines is
 $9x^2 + y^2 + 6xy - 4 = 0$

$$\therefore a = 9, b = 1, h = 3$$

$$h^2 - ab = 3^2 - 9(1) = 0$$

\therefore The lines are parallel

$$\text{Now, } 9x^2 + 6xy + y^2 = 4$$

$$\Rightarrow (3x + y)^2 = 4 \Rightarrow 3x + y = \pm 2$$

Hence, the lines are parallel and not coincident.

51. Given equation of pair of lines is
 $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$

$$\therefore A = a, B = b, H = h$$

The lines are parallel

$$\therefore H^2 = AB$$

$$\Rightarrow h = \sqrt{ab}$$

$$\text{Now } ABC + 2FGH - AF^2 - BG^2 - CH^2 = 0$$

$$\Rightarrow abc + 2fg\sqrt{ab} - af^2 - bg^2 - abc = 0$$

$$\Rightarrow (\sqrt{a}f - \sqrt{b}g)^2 = 0 \Rightarrow af^2 = bg^2$$

52. Given equation of pair of lines is
 $x^2 + k_1y^2 + 2k_2y = a^2$

$$a = 1, b = k_1, c = -a^2, f = k_2, g = 0, h = 0$$

The lines are perpendicular

$$\therefore a + b = 0 \Rightarrow k_1 = -1$$

Substituting value of k_1 in the given equation of lines, we get

$$x^2 - y^2 + 2k_2y - a^2 = 0$$

$$\Rightarrow a^2 - k_2^2 = 0 \Rightarrow k_2 = \pm a$$

53. $(x^2 + y^2)(h^2 + k^2 - a^2) = (hx + ky)^2$
 $\Rightarrow x^2(h^2 + k^2 - a^2) + y^2(h^2 + k^2 - a^2)$
 $\quad = h^2x^2 + k^2y^2 + 2hkxy$

$$\Rightarrow x^2(k^2 - a^2) + y^2(h^2 - a^2) - 2hkxy = 0$$

$$\therefore A = k^2 - a^2, B = h^2 - a^2$$

The lines are perpendicular

$$\therefore A + B = 0$$

$$\Rightarrow k^2 - a^2 + h^2 - a^2 = 0 \Rightarrow h^2 + k^2 = 2a^2$$

54. Given equation of pair of lines is

$$2x^2 - 4xy - py^2 + 4x + qy + 1 = 0$$

$$a = 2, b = -p, c = 1, f = \frac{q}{2}, g = 2, h = -2$$

The lines are perpendicular,

$$\therefore a + b = 0$$

$$\Rightarrow 2 - p = 0 \Rightarrow p = 2$$

The equations represents pair of lines

$$\therefore 2(-2)(1) + 2\left(\frac{q}{2}\right)(2)(2) - 2\left(\frac{q}{2}\right)^2 + 2(2)^2 - 1(2)^2 = 0$$

$$\Rightarrow q^2 - 8q = 0 \Rightarrow q = 0 \text{ or } 8$$

55. Given equation of pair of lines is

$$12x^2 + 7xy + by^2 + gx + 7y - 1 = 0$$

$$\therefore A = 12, B = b, C = -1, F = \frac{7}{2}, G = \frac{g}{2}, H = \frac{7}{2}$$

The lines are perpendicular

$$\therefore A + B = 0 \Rightarrow 12 + b = 0 \Rightarrow b = -12$$

$$\text{Also, } ABC + 2FGH - AF^2 - BG^2 - CH^2 = 0$$

$$\Rightarrow (12)(-12)(-1) + 2\left(\frac{7}{2}\right)\left(\frac{g}{2}\right)\left(\frac{7}{2}\right)$$

$$- (12)\left(\frac{7}{2}\right)^2 - (-12)\left(\frac{g}{2}\right)^2 - (-1)\left(\frac{7}{2}\right)^2 = 0$$

$$\Rightarrow 12g^2 + 49g + 37 = 0$$

$$\Rightarrow (g + 1)(12g + 37) = 0$$

$$\Rightarrow g = -1 \text{ or } -\frac{37}{12}$$

56. Given equation of pair of lines is

$$12x^2 + 7xy - py^2 - 18x + qy + 6 = 0$$

$$a = 12, b = -p, c = 6, f = \frac{q}{2}, g = -9, h = \frac{7}{2}$$

The lines are perpendicular

$$\therefore a + b = 0.$$

$$\Rightarrow 12 - p = 0 \Rightarrow p = 12$$

$$\text{Also, } abc + 2 fgh - af^2 - bg^2 - ch^2 = 0$$

$$\Rightarrow 12(-12)6 + 2\left(\frac{q}{2}\right)(-9)\left(\frac{7}{2}\right) - 12\left(\frac{q}{2}\right)^2$$

$$- (-12)(-9)^2 - 6\left(\frac{7}{2}\right)^2 = 0$$


$$\Rightarrow -864 - \frac{63q}{2} - 3q^2 + 972 - \frac{147}{2} = 0$$

$$\Rightarrow 23 - 21q - 2q^2 = 0$$

$$\Rightarrow (q - 1)(2q + 23) = 0 \Rightarrow q = 1 \text{ or } -\frac{23}{2}$$



57. The separate equations of lines represented by $x^2 - 7xy + 6y^2 = 0$ are $x - 6y = 0$ and $x - y = 0$. Let the 3 points be as shown in figure.



We know $\frac{0 + x_1 + x_2}{3} = 1$

$\Rightarrow x_1 + x_2 = 3$ (i)

and $y_1 + y_2 = 0$ (ii)

Also, $x_1 - 6y_1 = 0$ (iii)

$x_2 - y_2 = 0$ (iv)

[Since the points (x_1, y_1) and (x_2, y_2) lie on the lines AB and AC respectively]

On solving, we get the co-ordinates of B and C.

$\therefore B \equiv \left(\frac{18}{5}, \frac{3}{5}\right)$ and $C \equiv \left(\frac{-3}{5}, \frac{-3}{5}\right)$

Hence, the equation of third side i.e., BC is

$$\frac{y - \frac{3}{5}}{x - \frac{18}{5}} = \frac{\frac{-3}{5} - \frac{3}{5}}{\frac{-3}{5} - \frac{18}{5}}$$

$\Rightarrow 2x - 7y - 3 = 0$.

58. The given pair of lines can be separated as:

$L_1 = (+\sqrt{3}m)x + (m - \sqrt{3})y = 0$

$L_2 = (-\sqrt{3}m)x + (m + \sqrt{3})y = 0$

and $L_3 = x + my + n = 0$

- \therefore The slopes S_1, S_2 and S_3 of the three lines respectively are,

$S_1 = \frac{- (+\sqrt{3}m)}{(m - \sqrt{3})}, S_2 = \frac{- (-\sqrt{3}m)}{(m + \sqrt{3})}, S_3 = \frac{-}{m}$

Angle between L_1 and L_3 is

$$\theta_{13} = \tan^{-1} \left| \frac{S_1 - S_3}{1 + S_1 S_3} \right|$$

$$= \tan^{-1} \left| \frac{-\left(\frac{+\sqrt{3}m}{m - \sqrt{3}}\right) + \frac{-}{m}}{1 + \left(\frac{+\sqrt{3}m}{m - \sqrt{3}}\right)\left(\frac{-}{m}\right)} \right|$$

$$= \tan^{-1} \left| \frac{-\sqrt{3}m^2 - \sqrt{3}}{m^2 + m^2} \right| = \tan^{-1}(\sqrt{3}) = 60^\circ$$

Angle between L_2 and L_3 is

$$\theta_{23} = \tan^{-1} \left| \frac{S_2 - S_3}{1 + S_2 S_3} \right| = \tan^{-1} \left| \frac{-\left(\frac{-\sqrt{3}m}{m + \sqrt{3}}\right) + \frac{-}{m}}{1 + \left(\frac{-\sqrt{3}m}{m + \sqrt{3}}\right)\left(\frac{-}{m}\right)} \right|$$


$$= \tan^{-1} \left| \frac{\sqrt{3}m^2 + \sqrt{3}}{m^2 + m^2} \right| = \tan^{-1}(\sqrt{3}) = 60^\circ$$

- \therefore Angle between the lines L_1 and $L_2 = 60^\circ$
Hence, the triangle is equilateral.



Competitive Thinking

2.



Let OA and OB be the required lines.


- \therefore angles made by OA and OB with X-axis are 30° and 150° respectively.

- \therefore their equations are $y = \frac{1}{\sqrt{3}}x$ and $y = -\frac{1}{\sqrt{3}}x$

i.e., $x - \sqrt{3}y = 0$ and $x + \sqrt{3}y = 0$

- \therefore The joint equations of the lines is $(x - \sqrt{3}y)(x + \sqrt{3}y) = 0 \Rightarrow x^2 - 3y^2 = 0$

3. The lines trisecting the first quadrant are as shown in the figure.




- \therefore The joint equation of the lines is

$\left(y - \frac{1}{\sqrt{3}}x\right)(y - \sqrt{3}x) = 0$

$\Rightarrow (\sqrt{3}y - x)(y - \sqrt{3}x) = 0$

$\Rightarrow \sqrt{3}x^2 - 4xy + \sqrt{3}y^2 = 0$




The equations of bisectors are,
 $y - 3 = (1)(x - 5)$ and $y - 3 = (-1)(x - 5)$
 $\Rightarrow x - y - 2 = 0$ and $x + y - 8 = 0$

\therefore The joint equation of the bisectors is
 $(x - y - 2)(x + y - 8) = 0$
 $\Rightarrow x^2 - y^2 - 10x + 6y + 16 = 0$

5. Slope of QR = -2.
 Slope of PQ = m_1

$\therefore \tan 45^\circ = \left| \frac{m_1 + 2}{1 + m_1(-2)} \right|$
 $\Rightarrow 1 = \left| \frac{m_1 + 2}{1 - 2m_1} \right|$



$\Rightarrow m_1 = -\frac{1}{3}$

\therefore Equation of PQ passing through point P (2, 1) and having slope m_1 is

$y - 1 = -\frac{1}{3}(x - 2)$
 $\Rightarrow 3(y - 1) + (x - 2) = 0 \quad \dots \text{(i)}$

Slope of PR = $m_2 = 3 \quad \dots [\because PQ \perp PR]$

\therefore equation of PR is
 $y - 1 = 3(x - 2)$
 $\Rightarrow (y - 1) - 3(x - 2) = 0 \quad \dots \text{(ii)}$

\therefore The joint equation of the lines is
 $[3(y - 1) + (x - 2)][(y - 1) - 3(x - 2)] = 0$
 $\Rightarrow 3(y - 1)^2 - 8(y - 1)(x - 2) - 3(x - 2)^2 = 0$
 $\Rightarrow 3(x^2 - 4x + 4) + 8(xy - x - 2y + 2) - 3(y^2 - 2y + 1) = 0$
 $\Rightarrow 3x^2 - 3y^2 + 8xy - 20x - 10y + 25 = 0$

6. $x^2 - 7xy + 12y^2 = 0$
 $\Rightarrow (x - 3y)(x - 4y) = 0$
 Hence, the lines are intersecting and non-perpendicular.

7. $\sqrt{(x-2)^2 + y^2} + \sqrt{(x+2)^2 + y^2} = 4$
 i.e. $\sqrt{(x-2)^2 + y^2} = 4 - \sqrt{(x+2)^2 + y^2}$
 Squaring both sides, we get
 $(x-2)^2 + y^2 = 16 - 8\sqrt{(x+2)^2 + y^2} + (x+2)^2 + y^2$

$\Rightarrow x^2 - 4x + 4 + y^2 = 16 + x^2 + 4x + 4 + y^2 - 8\sqrt{(x+2)^2 + y^2}$

$\Rightarrow x + 2 = \sqrt{(x+2)^2 + y^2}$

Again squaring both sides, we get

$(x + 2)^2 = (x + 2)^2 + y^2$

$\Rightarrow y^2 = 0$

This is an equation of pair of two coincident straight lines.

8. The required lines are parallel to $x^2 - 4xy + 3y^2 = 0$, which pass through (3, -2).

\therefore the combined equation of lines is
 $(x - 3)^2 - 4(x - 3)(y + 2) + 3(y + 2)^2 = 0$
 $\Rightarrow x^2 - 6x + 9 - 4(xy + 2x - 3y - 6) + 3(y^2 + 4y + 4) = 0$
 $\Rightarrow x^2 - 6x + 9 - 4xy - 8x + 12y + 24 + 3y^2 + 12y + 12 = 0$
 $\Rightarrow x^2 - 4xy + 3y^2 - 14x + 24y + 45 = 0$

9. The required equation is $-2x^2 - 3xy + 5y^2 = 0$
 i.e., $2x^2 + 3xy - 5y^2 = 0$

10. Given equation of pair of lines is

$4xy + 2x + 6y + 3 = 0$
 $\Rightarrow 2x(2y + 1) + 3(2y + 1) = 0$
 $\Rightarrow (2y + 1)(2x + 3) = 0$

\therefore Separate equations of lines are $2x + 3 = 0$ and $2y + 1 = 0$

i.e. $x = -\frac{3}{2}$ and $y = -\frac{1}{2}$

The equation of line passing through (2, 1) and perpendicular to $x = -\frac{3}{2}$ is $y = 1$ i.e. $y - 1 = 0$


The equation of line passing through (2, 1) and perpendicular to $y = -\frac{1}{2}$ is $x = 2$ i.e. $x - 2 = 0$

\therefore Combined equation of pair of lines is
 $(x - 2)(y - 1) = 0$
 $\Rightarrow xy - x - 2y + 2 = 0$

11. OD is the median

$\therefore D \equiv \left(\frac{1+3}{2}, \frac{2+4}{2} \right)$

$\Rightarrow D \equiv (2, 3)$





Equation of OD is $y = mx$

$$\Rightarrow y = \frac{3}{2}x \Rightarrow 3x - 2y = 0$$

Slope of line AB = $\frac{2}{2} = 1$

Given, $OE \perp AB$

\therefore Slope of OE = -1

Equation of OE is $y = mx$

$$\Rightarrow y = -x \Rightarrow x + y = 0$$

\therefore Joint equation of median and altitude is

$$(3x - 2y)(x + y) = 0$$

$$\Rightarrow 3x^2 + xy - 2y^2 = 0$$

12. We have, $x^2 - 5x + 6 = 0$ and $y^2 - 6y + 5 = 0$


$$\Rightarrow (x - 3)(x - 2) = 0 \text{ and } (y - 1)(y - 5) = 0$$

\therefore One pair of opposite sides of parallelogram is

$$x - 3 = 0 \text{ and } x - 2 = 0 \text{ and the other pair is}$$

$$y - 1 = 0 \text{ and } y - 5 = 0$$

\therefore The vertices of the parallelogram are as shown in the figure below.



\therefore equation of diagonal d_1 is

$$y - 1 = \frac{5-1}{3-2}(x-2)$$

$$\Rightarrow y - 1 = 4(x-2) \Rightarrow y = 4x - 7$$

and equation of diagonal d_2 is


$$y - 1 = \frac{5-1}{2-3}(x-3)$$

$$\Rightarrow y - 1 = -4(x-3) \Rightarrow 4x + y = 13$$

\therefore the equations are $4x + y = 13$ and $y = 4x - 7$.

13. $2x^2 + 3xy - 2y^2 = 0$

$$\Rightarrow x + 2y = 0 \text{ and } 2x - y = 0$$



From the figure,

$$A\left(\frac{-2}{5}, \frac{1}{5}\right), B(0, 0), C\left(\frac{-1}{5}, \frac{-2}{5}\right)$$

Now, equation of side AD is

$$2x - y + c_1 = 0$$

Substituting $x = \frac{-2}{5}, y = \frac{1}{5}$ in above equation,

we get

$$c_1 = 1$$

\therefore equation of AD becomes $2x - y + 1 = 0$

Similarly equation of side DC is $x + 2y + c_2 = 0$

$$\text{i.e., } x + 2y + 1 = 0$$

$$\therefore D\left(\frac{-3}{5}, \frac{-1}{5}\right)$$

Now, equation of diagonal BD is

$$y - 0 = \frac{0 + \frac{1}{5}}{0 + \frac{3}{5}}(x - 0)$$

$$\Rightarrow \frac{3}{5}y = \frac{1}{5}x$$

$$\Rightarrow x - 3y = 0$$

14. Substituting the value of y in the equation

$$ax^2 + 2hxy + by^2 = 0.$$

$$\Rightarrow ax^2 + 2hx(mx) + b(mx)^2 = 0$$

$$\Rightarrow a + 2hm + bm^2 = 0$$

15. One of the lines is $3x + 4y = 0$

$$\text{i.e., } \frac{y}{x} = -\frac{3}{4}$$

The given joint equation is $6x^2 - xy + 4cy^2 = 0$

$$\Rightarrow 4c\left(\frac{y}{x}\right)^2 - \left(\frac{y}{x}\right) + 6 = 0 \quad \dots(i)$$

Substituting value of $\frac{y}{x}$ in equation (i), we get

$$4c\left(\frac{-3}{4}\right)^2 - \left(-\frac{3}{4}\right) + 6 = 0$$

$$\Rightarrow 4c \times \frac{9}{16} + \frac{3}{4} + 6 = 0$$

$$\Rightarrow \frac{9c}{4} + \frac{3+24}{4} = 0 \Rightarrow 9c + 27 = 0$$

$$\Rightarrow c = -3$$

16. Given equation of pair of lines is

$$kx^2 - 5xy - 3y^2 = 0$$

$$\Rightarrow k - 5\frac{y}{x} - 3\left(\frac{y}{x}\right)^2 = 0$$

$$\Rightarrow k - 5m - 3m^2 = 0 \quad \dots(i)$$

Now, slope of line $x - 2y + 3 = 0$ is $m_1 = \frac{1}{2}$.

\therefore slope of the line perpendicular to $x - 2y + 3 = 0$ is $m = -2$.

Substituting value of m in equation (i), we get

$$k - 5(-2) - 3(-2)^2 = 0$$

$$\Rightarrow k = -10 + 12 \Rightarrow k = 2$$



17. $6x^2 + xy - y^2 = 0$
 $\Rightarrow 6x^2 + 3xy - 2xy - y^2 = 0$
 $\Rightarrow 2x + y = 0$ and $3x - y = 0$
 let $a = \frac{1}{2}$
 \therefore equation $3x^2 - axy - y^2 = 0$ becomes
 $3x^2 - \frac{1}{2}xy - y^2 = 0$
 $\Rightarrow 6x^2 - xy - 2y^2 = 0$
 $\Rightarrow 3x - 2y = 0$ and $2x + y = 0$
 \therefore given pair of lines have common line $2x + y = 0$
 \therefore Option (A) is correct answer.

19. Given equation of pair of lines is
 $3x^2 + 5xy - 2y^2 = 0$

$$\therefore a = 3, h = \frac{5}{2}, b = -2$$

$$\text{Now, } m_1 + m_2 = \frac{-2h}{b} = \frac{5}{2}$$

20. Given equation of pair of lines is
 $4x^2 + 2hxy - 7y^2 = 0$

$$\therefore A = 4, H = h, B = -7$$

$$\text{Now, } m_1 + m_2 = -\frac{2H}{B} = \frac{2h}{7} \text{ and}$$

$$m_1 m_2 = \frac{A}{B} = \frac{4}{-7}$$

$$\text{Given that, } m_1 + m_2 = m_1 m_2$$

$$\Rightarrow \frac{2h}{7} = \frac{4}{-7} \Rightarrow h = -2$$

21. Given equation of pair of lines is
 $x^2 - 2cxy - 7y^2 = 0$

$$\therefore a = 1, h = -c, b = -7$$

$$\therefore m_1 + m_2 = \frac{-2c}{7} \text{ and } m_1 m_2 = \frac{-1}{7}$$

$$\text{Given that, } m_1 + m_2 = 4m_1 m_2$$

$$\Rightarrow \frac{-2c}{7} = \frac{-4}{7} \Rightarrow c = 2$$

22. Given equation of pair of lines is
 $ax^2 - 6xy + y^2 = 0$

$$\therefore A = a, H = -3, B = 1$$

$$\text{Given that, } m_1 = 2m_2$$

$$m_1 + m_2 = \frac{2(-3)}{1} = 6$$

$$\Rightarrow 2m_2 + m_2 = 6 \Rightarrow m_2 = 2 \Rightarrow m_1 = 4$$

$$\text{Now, } m_1 m_2 = \frac{a}{1} = a$$

$$\Rightarrow a = (4)(2) = 8$$

23. Given equation of pair of lines is
 $x^2 + hxy + 2y^2 = 0$

$$\therefore A = 1, H = \frac{h}{2}, B = 2$$

$$\text{Given that } m_1 = 2m_2$$

$$\text{Now, } m_1 + m_2 = \frac{-h}{2} \text{ and } m_1 m_2 = \frac{1}{2}$$

$$\therefore (2m_2)m_2 = \frac{1}{2} \Rightarrow 2(m_2)^2 = \frac{1}{2} \Rightarrow m_2 = \pm \frac{1}{2}$$

$$\text{Also, } 2m_2 + m_2 = \frac{-h}{2} \Rightarrow m_2 = \frac{-h}{6}$$

$$\Rightarrow \pm \frac{1}{2} = \frac{-h}{6} \Rightarrow h = \pm 3$$

24. Given equation of pairs of lines is
 $ax^2 + 2hxy + by^2 = 0$

$$\therefore m_1 + m_2 = \frac{-2h}{b} \text{ and } m_1 m_2 = \frac{a}{b}$$

$$\text{Given that, } m_1 = 2m_2$$

$$\therefore 2m_2 + m_2 = \frac{-2h}{b} \text{ and } 2m_2 m_2 = \frac{a}{b}$$

$$\therefore m_2 = \frac{-2h}{3b} \text{ and } m_2^2 = \frac{a}{2b}$$

$$\therefore \left(\frac{-2h}{3b}\right)^2 = \frac{a}{2b}$$

$$\therefore \frac{4h^2}{9b^2} = \frac{a}{2b}$$

$$\therefore 8h^2 = 9ab$$

25. Given equation of pairs of lines is
 $kx^2 + 5xy + y^2 = 0$

$$\therefore a = k, b = 1, h = \frac{5}{2}$$

$$\therefore m_1 + m_2 = \frac{-2h}{b} = -5$$

$$m_1 m_2 = \frac{a}{b} = k$$

$$\text{Given that, } m_1 - m_2 = 1$$

$$\text{Now, } (m_1 - m_2)^2 = (m_1 + m_2)^2 - 4m_1 m_2$$

$$\Rightarrow 1^2 = (-5)^2 - 4k \Rightarrow 4k = 24$$

$$\Rightarrow k = 6$$

26. If the gradients of two lines are in ratio 1 : n.

$$\text{then } \frac{h^2}{ab} = \frac{(n+1)^2}{4n} = \frac{(3+1)^2}{4 \cdot 3} = \frac{4}{3}$$

Alternate Method:

$$\text{Gradients } \frac{m_1}{m_2} = 1 : 3$$

$$\Rightarrow m_1 = m, m_2 = 3m$$



$$m_1 + m_2 = -\frac{2h}{b} \Rightarrow m + 3m = -\frac{2h}{b}$$

$$\Rightarrow m = \frac{-h}{2b}$$

$$m_1 \cdot m_2 = \frac{a}{b} \Rightarrow m \cdot 3m = \frac{a}{b}$$

$$\Rightarrow 3m^2 = \frac{a}{b} \Rightarrow 3 \cdot \frac{h^2}{4b^2} = \frac{a}{b} \Rightarrow \frac{h^2}{ab} = \frac{4}{3}$$

27. $m_1 : m_2 = 1 : 2$

$$\therefore \frac{h^2}{ab} = \frac{(2+1)^2}{4(2)} = \frac{9}{8}$$

$$\Rightarrow \frac{ab}{h^2} = \frac{8}{9}$$

28. Given equation of pair of lines is

$$x^2 + 4xy + y^2 = 0$$

$$\therefore a = 1, h = 2, b = 1$$

$$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a+b} \right| = \left| \frac{2\sqrt{(2)^2 - (1)(1)}}{1+1} \right| = \sqrt{3}$$

$$\Rightarrow \theta = \tan^{-1}(\sqrt{3}) = 60^\circ$$

29. Given equation of pair of lines is

$$(x^2 + y^2)\sqrt{3} = 4xy$$

$$\therefore a = \sqrt{3}, h = -2, b = \sqrt{3}$$

$$\therefore \tan \theta = \left| \frac{2\sqrt{4-3}}{2\sqrt{3}} \right| = \frac{1}{\sqrt{3}}$$

$$\Rightarrow \theta = \tan^{-1}\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6}$$

30. Given equation of pair of lines is

$$x^2 + 4y^2 - 7xy = 0$$

$$\therefore a = 1, h = -\frac{7}{2}, b = 4$$

$$\therefore \tan \theta = \left| \frac{2\sqrt{\left(\frac{-7}{2}\right)^2 - (1)(4)}}{1+(4)} \right|$$

$$= \left| \frac{2\sqrt{\frac{49}{4} - 4}}{5} \right| = \frac{\sqrt{33}}{5}$$

$$\Rightarrow \theta = \tan^{-1}\left(\frac{\sqrt{33}}{5}\right)$$

31. Given equation of pair of lines is

$$4x^2 - 24xy + 11y^2 = 0$$

$$\therefore a = 4, h = -12, b = 11$$

$$\therefore \tan \theta = \pm 2 \frac{\sqrt{h^2 - ab}}{a+b} = \pm 2 \frac{\sqrt{144 - 44}}{15} = \pm \frac{4}{3}$$

$$\Rightarrow \theta = \tan^{-1}\left(\pm \frac{4}{3}\right)$$

32. Given equation of pair of lines is

$$x^2 + 2xy \sec \theta + y^2 = 0$$

$$\therefore a = 1, h = \sec \theta, b = 1$$

Let ϕ be the angle between the lines.

$$\therefore \tan \phi = \left| \frac{2\sqrt{\sec^2 \theta - 1}}{2} \right|$$

$$\Rightarrow \tan \phi = \tan \theta \Rightarrow \phi = \theta$$

33. Let m_1 and m_2 be the slopes of the lines given

$$\text{by } x^2 + 4xy + y^2 = 0$$

$$\therefore m_1 + m_2 = -4 \Rightarrow m_2 = -4 - m_1$$

$$\text{and } m_1 \cdot m_2 = 1 \Rightarrow m_1(-4 - m_1) = 1$$

$$\Rightarrow m_1^2 + 4m_1 + 1 = 0$$

$$\therefore m_1, m_2 = -2 \pm \sqrt{3}$$

Slope of line $x - y = 4$ is

$$m_3 = 1$$

$$\therefore \text{Angle between first two lines,}$$

$$\tan^{-1} \theta_{12} = \left| \frac{m_1 - m_2}{1 + m_1 \cdot m_2} \right| = \left| \frac{(-2 + \sqrt{3}) - (-2 - \sqrt{3})}{1 + (-2 + \sqrt{3})(-2 - \sqrt{3})} \right|$$

$$\Rightarrow \theta_{12} = \tan^{-1}(\sqrt{3}) = 60^\circ$$

Angle between second and third line

$$\theta_{23} = \tan^{-1}\left(\frac{-2 - \sqrt{3} - 1}{1 + (-2 - \sqrt{3})1}\right) = \tan^{-1}(\sqrt{3}) = 60^\circ$$

Similarly, we have, $\theta_{31} = 60^\circ$

$$\therefore \text{The triangle formed by the lines is equilateral triangle.}$$

34. Let m_1 and m_2 be the slopes of the lines given by $23x^2 - 48xy + 3y^2 = 0$

$$\therefore m_1 + m_2 = \frac{48}{3} = 16 \Rightarrow m_2 = 16 - m_1$$

$$\text{and } m_1 m_2 = \frac{23}{3} \Rightarrow m_1(16 - m_1) = \frac{23}{3}$$

$$\Rightarrow -m_1^2 + 16m_1 - \frac{23}{3} = 0$$

$$\Rightarrow 3m_1^2 - 48m_1 + 23 = 0$$

$$\Rightarrow m_1, m_2 = \frac{24 \pm 13\sqrt{3}}{3}$$



slope of line is $2x + 3y + 4 = 0$ is

$$m_3 = \frac{-2}{3}$$

\therefore Angle between first two lines,

$$\begin{aligned} \tan^{-1} \theta_{12} &= \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right| \\ &= \left| \frac{\left(\frac{24 + 13\sqrt{3}}{3} \right) - \left(\frac{24 - 13\sqrt{3}}{3} \right)}{1 + \left(\frac{24 + 13\sqrt{3}}{3} \right) \left(\frac{24 - 13\sqrt{3}}{3} \right)} \right| \\ &= \left| \frac{\frac{26\sqrt{3}}{3}}{\frac{9 + 576 - 507}{9}} \right| = \left| \frac{\frac{26\sqrt{3}}{3}}{\frac{78}{9}} \right| \end{aligned}$$

$$\therefore \tan^{-1} \theta_{12} = \sqrt{3}$$

$$\Rightarrow \theta_{12} = \tan^{-1}(\sqrt{3}) = 60^\circ$$

Angle between second and third line

$$\begin{aligned} \theta_{23} &= \tan^{-1} \left(\frac{\frac{24 - 13\sqrt{3}}{3} - \left(-\frac{2}{3} \right)}{1 + \left(\frac{24 - 13\sqrt{3}}{3} \right) \left(-\frac{2}{3} \right)} \right) \\ &= \tan^{-1} \left(\frac{\frac{26 - 13\sqrt{3}}{3}}{\frac{9 - 48 + 26\sqrt{3}}{9}} \right) = \tan^{-1} \left(\frac{\frac{26 - 13\sqrt{3}}{3}}{\frac{-39 + 26\sqrt{3}}{9}} \right) \\ &= \tan^{-1} \left(\frac{26 - 13\sqrt{3}}{3} \times \frac{9}{-39 + 26\sqrt{3}} \right) \\ &= \tan^{-1} \left(\frac{13(2 - \sqrt{3}) \times 3}{13\sqrt{3}(2 - \sqrt{3})} \right) \\ &= \tan^{-1}(\sqrt{3}) = 60^\circ \end{aligned}$$

Similarly, we have, $\theta_{31} = 60^\circ$

\therefore The triangle formed by the lines is equilateral triangle.

35. Given equation of pair of lines is

$$4x^2 + 12xy + 9y^2 = 0$$

$$a = 4, h = 6, b = 9$$

Here,

$$h^2 - ab = (6)^2 - (4)(9) = 36 - 36 = 0$$

Hence, the lines are real and coincident.

36. Given equation of pair of lines is

$$x^2 + ky^2 + 4xy = 0$$

$$\therefore a = 1, h = \frac{k}{2}, b = 4$$

The pair of lines are coincident if $h^2 - ab = 0$

$$\Rightarrow h^2 = ab \Rightarrow \frac{k^2}{4} = 4(1)$$

$$\Rightarrow k = \pm 4$$

37. Given equation of pair of lines is

$$px^2 - qy^2 = 0$$

$$\therefore a = p, b = -q, c = 0$$

Since, the lines are real and distinct

$$\therefore h^2 - ab > 0$$

$$\Rightarrow 0 - p(-q) > 0$$

$$\Rightarrow pq > 0$$

38. Given equation of pair of lines is

$$y^2 \sin^2 \theta - xy \sin^2 \theta + x^2 (\cos^2 \theta - 1) = 0$$

$$\therefore a = \sin^2 \theta, b = \cos^2 \theta - 1 = -(1 - \cos^2 \theta) = -\sin^2 \theta$$

$$\text{Now, } a + b = \sin^2 \theta - \sin^2 \theta = 0$$

\therefore The lines are perpendicular.

$$\therefore \theta = \frac{\pi}{2}$$

39. Consider option (C)

$$\text{Given equation is } y^2 + x + 1 = 0$$

$$\therefore a = 0, b = 1, c = 0, f = 0, g = \frac{-1}{2}, h = 0$$

$$\text{Now, } abc + 2fgh - af^2 - bg^2 - ch^2$$

$$= 0 + 0 - 0 - \left(\frac{1}{4} \right) + 0 = \frac{-1}{4} \neq 0$$

\therefore The equation does not represent a pair of straight lines.

40. Given equation of pair of lines is

$$3x^2 + 7xy + 2y^2 + 5x + 5y + 2 = 0$$

$$a = 3, b = 2, c = 2, f = \frac{5}{2}, g = \frac{5}{2}, h = \frac{7}{2}$$

$$\text{Consider } abc + 2fgh - af^2 - bg^2 - ch^2$$

$$= (3)(2)(2) + 2 \left(\frac{5}{2} \right) \left(\frac{5}{2} \right) \left(\frac{7}{2} \right)$$

$$- 3 \left(\frac{5}{2} \right)^2 - 2 \left(\frac{5}{2} \right)^2 - 2 \left(\frac{7}{2} \right)^2 = 0$$

\therefore the given equation represents a pair of straight lines.



41. Given equation of pair of lines is
 $xy + a^2 = ax + ay$
 i.e. $ax + ay - xy - a^2 = 0$
- $\therefore A = 0, B = 0, C = -a^2, F = \frac{a}{2}, G = \frac{a}{2}, H = -\frac{1}{2}$
- Now, $ABC + 2FGH - AF^2 - BG^2 - CH^2$
 $= 0 - 2\left(\frac{a}{2}\right)\left(\frac{a}{2}\right)\left(\frac{-1}{2}\right) - (a^2)\left(\frac{-1}{2}\right)^2 = 0$
- \therefore the given equation represents a pair of straight lines.
42. Given equation of pair of lines is
 $ax^2 - y^2 + 4x - y = 0$
- $\therefore A = a, B = -1, C = 0, F = \frac{-1}{2}, G = 2, H = 0$
- The given equation represents a pair of straight lines,
- $\therefore ABC + 2FGH - AF^2 - BG^2 - CH^2 = 0$
 $\Rightarrow 0 - 0 - a\left(\frac{1}{4}\right) - (-1)(4) = 0$
 $\Rightarrow -\frac{a}{4} + 4 = 0 \Rightarrow a = 16$
43. Given equation of pair of lines is
 $kxy + 10x + 6y + 4 = 0$
- $\therefore a = b = 0, c = 4, f = 3, g = 5, h = \frac{k}{2}$
- Now, $abc + 2fgh - af^2 - bg^2 - ch^2 = 0$
 $\Rightarrow 0 + 2(3)(5)\left(\frac{k}{2}\right) - 0 - 0 - 4\left(\frac{k}{2}\right)^2 = 0$
 $\Rightarrow 15k - k^2 = 0 \Rightarrow k(15 - k) = 0$
 $\Rightarrow k = 0$ or $k = 15$
44. Given equation of pair of lines is
 $x^2 + kxy + y^2 - 5x - 7y + 6 = 0$
- $\therefore a = 1, b = 1, c = 6, f = \frac{-7}{2}, g = \frac{-5}{2}, h = \frac{k}{2}$
- Now, $abc + 2fgh - af^2 - bg^2 - ch^2 = 0$
 $\Rightarrow (1)(1)(6) + 2\left(\frac{-7}{2}\right)\left(\frac{-5}{2}\right)\left(\frac{k}{2}\right) - 1\left(\frac{-7}{2}\right)^2$
 $\quad - 1\left(\frac{-5}{2}\right)^2 - 6\left(\frac{k}{2}\right)^2 = 0$
 $\Rightarrow 6 + \frac{35k}{4} - \frac{49}{4} - \frac{25}{4} - \frac{6k^2}{4} = 0$
 $\Rightarrow -6k^2 + 35k - 50 = 0$
 $\Rightarrow (2k - 5)(3k - 10) = 0$
 $\Rightarrow k = \frac{5}{2}$ or $k = \frac{10}{3}$
45. Given equation of pair of lines is
 $x^2 - y^2 + x + 3y - 2 = 0$
- $\therefore a = 1, b = -1, g = \frac{1}{2}, f = \frac{3}{2}, c = -2$
- \therefore point of intersection of the lines is
 $\left(\frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2}\right) = \left(\frac{-1 \cdot \frac{3}{2}}{ab - h^2}, \frac{\frac{3}{2} - 1}{ab - h^2}\right) = \left(\frac{-1}{2}, \frac{3}{2}\right)$
46. Given equation of pair of lines is
 $2x^2 - 10xy + 2\lambda y^2 + 5x - 16y - 3 = 0$
- $\therefore a = 2, b = 2\lambda, c = -3, f = -8, g = \frac{5}{2}, h = -5$
- Since the equation represents pair of lines,
 $\therefore abc + 2fgh - af^2 - bg^2 - ch^2 = 0$
 $\Rightarrow 2(2\lambda)(-3) + 2(-8)\left(\frac{5}{2}\right)(-5) - 2(64)$
 $\quad - 2\lambda\left(\frac{25}{4}\right) + 3(25) = 0$
 $\Rightarrow \frac{49\lambda}{2} = 147 \Rightarrow \lambda = 6$
- \therefore Point of intersection of the lines is
 $\left(\frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2}\right)$
 $\equiv \left(\frac{(-5)(-8) - 2(6)\left(\frac{5}{2}\right)}{2(12) - (-5)^2}, \frac{\frac{5}{2}(-5) - 2(-8)}{2(12) - (-5)^2}\right)$
 $\equiv \left(-10, \frac{-7}{2}\right)$
47. Given equation of pair of lines is
 $2x^2 - 3xy - 2y^2 + 10x + 5y = 0$
- $\therefore a = 2, b = -2, c = 0, f = \frac{5}{2}, g = 5, h = \frac{-3}{2}$
- \therefore Point of intersection of the lines is
 $\left(\frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2}\right) \equiv (-1, 2)$
- Slope of line joining origin and $(-1, 2)$ $m = -2$
 Slope of $kx + y + 3 = 0$ is $-k$
 Now, $(-k)(-2) = -1 \Rightarrow k = \frac{-1}{2}$
48. The line $5x + y - 1 = 0$ is coincides
 $5x^2 + xy - kx - 2y + 2 = 0$
- $\therefore a = 5, b = 0, c = 2, f = -1, g = -\frac{k}{2}, h = \frac{1}{2}$
- $m_1 + m_2 = \frac{-2h}{b}$



As $b = 0$, this case is not defined

Slope of line $5x + y - 1 = 0$ is $m = -5$

\therefore Slope of another line must be infinite

\therefore equation of another line is $x = k_1$

\therefore Combine equation is $(5x + y - 1)(x - k_1) = 0$

$$\Rightarrow 5x^2 - 5xk_1 + xy - yk_1 - x + k_1 = 0$$

$$\Rightarrow 5x^2 + xy - (5k_1 + 1)x - yk_1 + k_1 = 0$$

Comparing this equation with the given equation, we get $k = 11$

49. Given equation of pair of lines is

$$3x^2 + 7xy + 2y^2 + 5x + 5y + 2 = 0$$

$$\therefore a = 3, b = 2, h = \frac{7}{2}$$

$$\tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right| = \left| \frac{2\sqrt{\frac{49}{4} - 6}}{3 + 2} \right| = \left| \frac{2\sqrt{\frac{25}{4}}}{5} \right|$$

$$\Rightarrow \tan \theta = 1$$

$$\Rightarrow \theta = \tan^{-1}(1) = \frac{\pi}{4}$$

50. Given equation of pair of lines is

$$x^2 - xy - 6y^2 - 7x + 31y - 18 = 0$$

$$\therefore a = 1, b = -6, h = -\frac{1}{2}$$

$$\therefore \tan \theta = \left| \frac{2\sqrt{\left(\frac{-1}{2}\right)^2 - 1(-6)}}{1 - 6} \right| = \left| \frac{2\sqrt{\frac{1}{4} + 6}}{-5} \right| = |-1| = 1$$

$$\Rightarrow \theta = \tan^{-1}(1) = 45^\circ$$

51. Given equation of pair of lines is

$$x^2 - 3xy + \lambda y^2 + 3x + 5y + 2 = 0$$

$$\therefore a = 1, b = \lambda, h = -\frac{3}{2}$$

$$\theta = \tan^{-1} 3 \Rightarrow \tan \theta = 3$$

$$\therefore \tan \theta = \left| \frac{2\sqrt{\left(-\frac{3}{2}\right)^2 - (1)\lambda}}{1 + \lambda} \right|$$

$$\Rightarrow 3 = \left| \frac{2\sqrt{\frac{9 - 4\lambda}{4}}}{1 + \lambda} \right| = \left| \frac{\sqrt{9 - 4\lambda}}{1 + \lambda} \right|$$

$$\Rightarrow \frac{9 - 4\lambda}{(1 + \lambda)^2} = 9$$

$$\Rightarrow 9 - 4\lambda = 9(1 + \lambda)^2$$

$$\Rightarrow 9\lambda^2 + 22\lambda = 0$$

$$\Rightarrow \lambda(9\lambda + 22) = 0$$

$$\Rightarrow \lambda = 0 \text{ or } \lambda = -\frac{22}{9}$$

But λ is non-negative

$$\therefore \lambda = 0$$

52. Given equation of pair of lines is

$$2x^2 + 5xy + 3y^2 + 6x + 7y + 4 = 0$$

$$a = 2, b = 3, h = \frac{5}{2}$$

$$\theta = \tan^{-1} m \Rightarrow \tan \theta = m$$

$$\tan \theta = \left| \frac{2\sqrt{\frac{25}{4} - 6}}{2 + 3} \right| \Rightarrow m = \frac{1}{5}$$

53. Given equation of pair of lines is

$$x^2 + y^2 - 2x - 1 = 0 \quad \dots(i)$$

$x + y = 1$ intersects the above pair of lines

\therefore It satisfies equation (i)

$$\therefore x^2 + y^2 - 2x(x + y) - (x + y)^2 = 0$$

$$\Rightarrow 2x^2 + 4xy = 0 \Rightarrow x^2 + 2xy = 0$$

$$\therefore a = 1, b = 0, h = 1$$

$$\therefore \tan \theta = \frac{2\sqrt{1^2 - 0}}{1}$$

$$\Rightarrow \theta = \tan^{-1}(2)$$

54. The joint equation of the pair of straight lines

joining the origin to the points of intersection

of the line $x + my + n = 0$ and

$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ is

$$ax^2 + 2hxy + by^2 + 2g\left(\frac{x + my}{-n}\right)x + 2f\left(\frac{x + my}{-n}\right)y + c\left(\frac{x + my}{-n}\right)^2 = 0$$

Here, $a = 2, m = 1, n = -1$ and

$$a = 3, b = 0, c = 1, f = 0, g = -2, h = 2$$

$$\therefore 3x^2 + 4xy - 4x(2x + y) + (2x + y)^2 = 0$$

$$\Rightarrow 3x^2 + 4xy - 8x^2 - 4xy + 4x^2 + y^2 + 4xy = 0$$

$$\Rightarrow x^2 - 4xy - y^2 = 0$$

$$\therefore A = 1, B = -1, H = -2$$

$$\therefore \tan \theta = \frac{2\sqrt{4 + 1}}{0} = \infty$$

$$\Rightarrow \theta = \frac{\pi}{2}$$

55. Given, $ax^2 + 2hxy + by^2 = -2gx$

$$a_1x^2 + 2h_1xy + b_1y^2 = -2g_1x$$

$$\therefore \frac{ax^2 + 2hxy + by^2}{a_1x^2 + 2h_1xy + b_1y^2} = \frac{g}{g_1}$$

We have,

$$(ag_1 - a_1g)x^2 + 2(hg_1 - h_1g)xy + (bg_1 - b_1g)y^2 = 0$$

$$\therefore A = (ag_1 - a_1g), B = (bg_1 - b_1g)$$

The lines are perpendicular

$$\therefore A + B = 0$$

$$\Rightarrow (ag_1 - a_1g) + (bg_1 - b_1g) = 0$$

$$\Rightarrow (a + b)g_1 = (a_1 + b_1)g$$




56. The equation of line is $y = 2\sqrt{2}x + c$
 $\Rightarrow \left(\frac{y - 2\sqrt{2}x}{c}\right) = 1 \quad \dots(i)$
 Given equation of circle is
 $x^2 + y^2 = 2(1)^2 \quad \dots(ii)$
 \therefore from (i) and (ii), we get
 $x^2 + y^2 = 2\left(\frac{y - 2\sqrt{2}x}{c}\right)^2$
 $\Rightarrow c^2(x^2 + y^2) = 2(y^2 - 4\sqrt{2}xy + 8x^2)$
 $\Rightarrow (c^2 - 16)x^2 + (c^2 - 2)y^2 + 8\sqrt{2}xy = 0$
 The lines are perpendicular if $A + B = 0$.
 $\therefore c^2 - 16 + c^2 - 2 = 0$
 $\Rightarrow 2c^2 - 18 = 0 \quad \Rightarrow c^2 - 9 = 0$
57. Lines represented by the equation
 $2y^2 - xy - 6x^2 = 0$ are
 $y = 2x$ and $y = -\frac{3}{2}x$
 The co-ordinates of the vertices of the triangle
 formed by above lines with $x + y = 1$ are
 $(0, 0)$, $\left(\frac{1}{3}, \frac{2}{3}\right)$ and $(-2, 3)$
 The altitude from vertex $(0, 0)$ on $x + y = 1$ is
 $y = x \quad \dots(i)$
 The altitude from vertex $\left(\frac{1}{3}, \frac{2}{3}\right)$ on $y = -\frac{3}{2}x$
 is $y - \frac{2}{3} = \frac{2}{3}\left(x - \frac{1}{3}\right)$
 $\Rightarrow 6x - 9y + 4 = 0 \quad \dots(ii)$
 Solving (i) and (ii), we get

$$x = \frac{4}{3} \text{ and } y = \frac{4}{3},$$

\therefore Orthocentre is $\left(\frac{4}{3}, \frac{4}{3}\right)$

58. Given equations of pair of lines are
 $xy + 4x - 3y - 12 = 0$ and
 $xy - 3x + 4y - 12 = 0$
 $\therefore x(y + 4) - 3(y + 4) = 0$ and $x(y - 3) + 4(y - 3) = 0$
 $\therefore (y + 4)(x - 3) = 0$ and $(x + 4)(y - 3) = 0$
 \therefore The vertices of the square are as shown in the figure



Equation of diagonal d_1 is

$$y + 4 = \frac{-4 - 3}{-4 - 3}(x + 4)$$

$$\Rightarrow y + 4 = x + 4$$

$$\Rightarrow x - y = 0$$

and equation of diagonal d_2 is

$$y + 4 = \frac{3 + 4}{-4 - 3}(x - 3)$$

$$\Rightarrow y + 4 = -1(x - 3)$$

$$\Rightarrow y + 4 = -x + 3$$

$$\Rightarrow x + y + 1 = 0$$

- \therefore Combined equation of diagonals d_1 and d_2 is
 $(x - y)(x + y + 1) = 0$
 $\Rightarrow x^2 - y^2 + x - y = 0$



Evaluation Test

1. $L_1: ax^2 + 2hxy + by^2 = 0$
 Equation of any line passing through origin
 and perpendicular to L_1 is given by
 $bx^2 - 2hxy + ay^2 = 0$
 \dots (interchanging coefficients of x^2 and y^2 and
 change of sign for xy term)
 \therefore The required equation of pair of lines is
 $-15x^2 + 7xy + 2y^2 = 0$
 i.e. $15x^2 - 7xy - 2y^2 = 0$
2. Here, $m_1 + m_2 = \frac{-2h}{b} \quad \dots(i)$
 and $m_1m_2 = \frac{a}{b}$

$$\begin{aligned} \therefore (m_1 - m_2)^2 &= (m_1 + m_2)^2 - 4m_1m_2 \\ &= \frac{4h^2 - 4ab}{b^2} \\ &= \frac{4h^2 - 3h^2}{b^2} \quad \dots[\because 4ab = 3h^2 \text{ (given)}] \\ &= \frac{h^2}{b^2} \end{aligned}$$

$$\therefore m_1 - m_2 = \frac{h}{b} \quad \dots(ii)$$

On solving (i) and (ii), we get

$$m_1 = \frac{-h}{2b} \text{ and } m_2 = \frac{-3h}{2b}$$

$$\therefore m_1 : m_2 = 1 : 3$$



3. The lines are parallel, if $af^2 = bg^2$
 $\therefore 4f^2 = 9g^2$
 $\Rightarrow f = \frac{3}{2}g$
 Let $g = 2$ and $f = 3$
 $\therefore abc + 2fgh - af^2 - bg^2 - ch^2$
 $= 4(9)(c) + 2(3)(2)(6) - 4(3)^2 - 9(2)^2 - c(6)^2 = 0$
 $\Rightarrow c$ is any number.
4. Given equation is $x^2 - y^2 - x - \lambda y - 2 = 0$.
 $\therefore a = 1, b = -1, c = -2, f = \frac{-\lambda}{2}, g = \frac{-1}{2}, h = 0$
 This equation represents a pair of straight lines, if $abc + 2fgh - af^2 - bg^2 - ch^2 = 0$
 $\Rightarrow 2 - \frac{\lambda^2}{4} + \frac{1}{4} = 0 \Rightarrow \frac{\lambda^2}{4} = \frac{9}{4} \Rightarrow \lambda^2 = 9 \Rightarrow \lambda = \pm 3$
5. The given equation of pair of lines is $x^2 + 2\sqrt{2}xy - y^2 = 0$
 $\therefore a = 1, b = -1, h = \sqrt{2}$
 Now, $a + b = 1 + (-1) = 0$
 \therefore The lines are perpendicular
6. The joint equation of the lines through the point (x_1, y_1) and at right angles to the lines $ax^2 + 2hxy + by^2 = 0$ is $b(x - x_1)^2 - 2h(x - x_1)(y - y_1) + a(y - y_1)^2 = 0$
 \therefore joint equation of pair of lines drawn through $(1, 1)$ and perpendicular to the pair of lines $3x^2 - 7xy + 2y^2 = 0$ is $2(x - 1)^2 + 7(x - 1)(y - 1) + 3(y - 1)^2 = 0$
7. The given equations are $x - y - 1 = 0$ and $2x + y - 6 = 0$
 \therefore The joint equation is given by $(x - y - 1)(2x + y - 6) = 0$
 $\Rightarrow 2x^2 + xy - 6x - 2xy - y^2 + 6y - 2x - y + 6 = 0$
 $\Rightarrow 2x^2 - y^2 - xy - 8x + 5y + 6 = 0$
8. Let the equation of one of the angle bisector of the co-ordinate axes be $x + y = 0 \Rightarrow m_1 = -1$
 Given equation of pair of lines is $2x^2 + 2hxy + 3y^2 = 0$
 $\therefore A = 2, H = h, B = 3$
 Now, $m_1 m_2 = \frac{a}{b} \Rightarrow m_2 = \frac{-2}{3}$
 Also $m_1 + m_2 = \frac{-2h}{b} \Rightarrow -1 - \frac{2}{3} = \frac{-2h}{3}$
 $\Rightarrow h = \frac{5}{2}$
9. The given equation of pair of lines is $3x^2 - 2y^2 + \lambda xy - x + 5y - 2 = 0$
 $\therefore a = 3, b = -2, c = -2, f = \frac{\lambda}{2}, g = \frac{-1}{2}, h = \frac{\lambda}{2}$

$$\text{Now } abc + 2fgh - af^2 - bg^2 - ch^2 = 0$$

$$12 - \frac{5\lambda}{4} - \frac{75}{4} + \frac{1}{2} + \frac{\lambda^2}{2} = 0$$

$$\Rightarrow 2\lambda^2 - 5\lambda - 25 = 0 \Rightarrow (\lambda - 5)(2\lambda + 5) = 0$$

$$\Rightarrow \lambda = 5 \text{ or } \frac{-5}{2}$$

10. Let $y = mx$ be the common line and let $y = m_1x$ and $y = m_2x$ be the other lines given by $2x^2 + axy + 3y^2 = 0$ and $2x^2 + bxy - 3y^2 = 0$ respectively. Then,

$$m + m_1 = -\frac{a}{3}, mm_1 = \frac{2}{3}, \text{ and}$$

$$m + m_2 = \frac{b}{3}, mm_2 = -\frac{2}{3}$$

$$\therefore (mm_1)(mm_2) = \frac{2}{3} \left(-\frac{2}{3} \right)$$

$$\Rightarrow m^2(m_1 m_2) = -\frac{4}{9}$$

$$\Rightarrow m^2 = \frac{4}{9} \quad \dots [\because m_1 m_2 = -1 \text{ (given)}]$$

$$\Rightarrow m = \pm \frac{2}{3}$$

$$\text{When } m = \frac{2}{3},$$

$$mm_1 = \frac{2}{3} \text{ and } mm_2 = -\frac{2}{3} \Rightarrow m_1 = 1 \text{ and } m_2 = -1$$

$$\therefore m + m_1 = -\frac{a}{3} \text{ and } m + m_2 = \frac{b}{3}$$

$$\Rightarrow a = -5 \text{ and } b = -1$$

$$\text{When } m = -\frac{2}{3},$$

$$mm_1 = \frac{2}{3} \text{ and } mm_2 = -\frac{2}{3} \Rightarrow m_1 = -1 \text{ and } m_2 = 1$$

$$\therefore m + m_1 = -\frac{a}{3} \text{ and } m + m_2 = \frac{b}{3}$$

$$\Rightarrow a = 5 \text{ and } b = 1$$

11. Given equation of pair of lines is $3x^2 - 48xy + 23y^2 = 0$

$$\therefore a = 3, h = -24, b = 23$$

$$\therefore \tan \theta = \left| \frac{2\sqrt{576 - 69}}{3 + 23} \right|$$

$$\Rightarrow \tan \theta = \left| \frac{2\sqrt{507}}{26} \right| = \left| \frac{2 \times 13\sqrt{3}}{26} \right| = \sqrt{3}$$

$$\Rightarrow \theta = \tan^{-1}(\sqrt{3}) = \frac{\pi}{3}$$

05 Vectors



Hints



Classical Thinking

1. Since the vectors are collinear,

$$\therefore \bar{b} = \lambda \bar{a}$$

$$\Rightarrow (-2\hat{i} + m\hat{j}) = \lambda(\hat{i} - \hat{j})$$

On comparing, we get

$$\lambda = -2 \text{ and } -\lambda = m$$

$$\Rightarrow m = 2$$

2. $\bar{c} = \lambda \bar{d}$

$$\Rightarrow (x-2)\bar{a} + \bar{b} = \lambda(2x+1)\bar{a} - \lambda\bar{b}$$

On comparing, we get

$$\lambda = -1 \text{ and}$$

$$(x-2) = \lambda(2x+1)$$

$$\Rightarrow x-2 = -2x-1$$

$$\Rightarrow x = \frac{1}{3}$$

3. Let $\bar{a} = 3\hat{i} - 2\hat{j} + 5\hat{k}$ and $\bar{b} = -2\hat{i} + p\hat{j} - q\hat{k}$

Two vector are collinear if

$$\frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3}$$

$$\Rightarrow \frac{3}{-2} = \frac{-2}{p} = \frac{5}{-q}$$

$$\Rightarrow p = \frac{4}{3}, q = \frac{10}{3}$$

4. For the points to be collinear,

$$\overline{AB} \times \overline{BC} = 0$$

$$\Rightarrow (\bar{b} - \bar{a}) \times (\bar{c} - \bar{b}) = \bar{0}$$

$$\Rightarrow \bar{b} \times \bar{c} - \bar{a} \times \bar{c} + \bar{a} \times \bar{b} = \bar{0}$$

$$\Rightarrow \bar{a} \times \bar{b} + \bar{b} \times \bar{c} + \bar{c} \times \bar{a} = \bar{0}$$

5. Here $\bar{a} = \hat{i} + \hat{j}$, $\bar{b} = 2\hat{i} - \hat{j}$ and $\bar{r} = 2\hat{i} - 4\hat{j}$

$$\text{Let } \bar{r} = t_1\bar{a} + t_2\bar{b}$$

$$\Rightarrow 2\hat{i} - 4\hat{j} = t_1(\hat{i} + \hat{j}) + t_2(2\hat{i} - \hat{j})$$

$$= (t_1 + 2t_2)\hat{i} + (t_1 - t_2)\hat{j}$$

Comparing the coefficients, we get

$$t_1 + 2t_2 = 2 \quad \dots\text{(i)}$$

$$t_1 - t_2 = -4 \quad \dots\text{(ii)}$$

On solving (i) and (ii), we get

$$t_1 = -2, t_2 = 2$$

6. Given, $3\bar{A} = 2\bar{B}$

$$\therefore 3(x+4y) = 2(y-2x+2)$$

$$\Rightarrow 7x + 10y = 4 \quad \dots\text{(i)}$$

$$\text{and } 3(2x+y+1) = 2(2x-3y-1)$$

$$\Rightarrow 2x + 9y = -5 \quad \dots\text{(ii)}$$

On solving (i) and (ii), we get

$$x = 2, y = -1$$

8. $1(\bar{a}) + 1(\bar{b}) = \bar{a} + \bar{b}$.

$$\therefore 1(\bar{a}) + 1(\bar{b}) - 1(\bar{a} + \bar{b}) = 0$$

\therefore The vectors are coplanar.

9. Let R(\bar{r}) be the point dividing PQ internally in the ratio 2 : 5

$$\therefore \bar{r} = \frac{5\bar{p} + 2\bar{q}}{7}$$

10. Let R(\bar{r}) divide line AB internally in the ratio 2 : 3

$$\therefore \bar{r} = \frac{2\bar{b} + 3\bar{a}}{2+3}$$

$$= \frac{2(3\hat{i} + \hat{j} + 4\hat{k}) + 3(2\hat{i} + 3\hat{j} - \hat{k})}{5}$$

$$= \frac{12\hat{i} + 11\hat{j} + 5\hat{k}}{5}$$

\therefore Co-ordinates of R are $\left(\frac{12}{5}, \frac{11}{5}, 1\right)$

11. $C \equiv \left(\frac{2-4}{2}, \frac{-1+3}{2}\right) \equiv (-1, 1)$

$$\therefore \overline{OC} = -\hat{i} + \hat{j}$$

12. If M(\bar{m}) is the mid-point of AB, then

$$\bar{m} = \frac{\bar{a} + \bar{b}}{2}$$

$$\Rightarrow \frac{\hat{i} + 3\hat{j} - \hat{k} + 3\hat{i} - \hat{j} - 3\hat{k}}{2} = 2\hat{i} + \hat{j} - 2\hat{k}$$

13. Let R(\bar{r}) divide AB externally in the ratio 5:2

$$\therefore \bar{r} = \frac{5(\hat{i} - \hat{j} + 2\hat{k}) - 2(2\hat{i} + \hat{j} - \hat{k})}{5-2} = \frac{\hat{i} - 7\hat{j} + 12\hat{k}}{3}$$



14. Let $R(\bar{r})$ divide PQ externally in the ratio 2 : 1

$$\begin{aligned}\therefore \bar{r} &= \frac{2\bar{q} - \bar{p}}{2-1} \\ &= \frac{2(3\hat{i} + 2\hat{j} + \hat{k}) - (2\hat{i} - \hat{j} + 4\hat{k})}{1} \\ &= 4\hat{i} + 5\hat{j} - 2\hat{k}\end{aligned}$$

\therefore Co-ordinates of R are (4, 5, -2)

15. Let P divide AB in the ratio $\lambda : 1$

$$\begin{aligned}\therefore \left(\frac{17}{4}, \frac{11}{4}, 0\right) &\equiv \left(\frac{2\lambda+5}{\lambda+1}, \frac{-7\lambda+a}{\lambda+1}, \frac{k\lambda-1}{\lambda+1}\right) \\ \Rightarrow \frac{17}{4} &= \frac{2\lambda+5}{\lambda+1} \Rightarrow \lambda = \frac{1}{3}\end{aligned}$$

16. If $A(\bar{a}), B(\bar{b}), C(\bar{c})$ are the vertices and $G(\bar{g})$ is the centroid of ΔABC , then

$$\bar{g} = \frac{\bar{a} + \bar{b} + \bar{c}}{3}$$

$$\begin{aligned}\therefore 3\hat{i} + 2\hat{j} + n\hat{k} &= \frac{(2\hat{i} + 3\hat{j} - 4\hat{k}) + (m\hat{i} + \hat{j} - \hat{k}) + (3\hat{i} + 2\hat{j} + 2\hat{k})}{3} \\ \Rightarrow 3(3\hat{i} + 2\hat{j} + n\hat{k}) &= (5+m)\hat{i} + 6\hat{j} + (-3)\hat{k} \\ \text{On comparing, we get} \\ 9 &= 5+m \Rightarrow m=4, \text{ and} \\ 3n &= -3 \Rightarrow n=-1\end{aligned}$$

$$\begin{aligned}17. G &\equiv \left(\frac{x_1+x_2+x_3}{3}, \frac{y_1+y_2+y_3}{3}, \frac{z_1+z_2+z_3}{3}\right) \\ \Rightarrow (2, 1, c) &\equiv \left(\frac{2a+1}{3}, \frac{4+b}{3}, \frac{1}{3}\right) \\ \Rightarrow 2 &= \frac{2a+1}{3}, 1 = \frac{4+b}{3}, c = \frac{1}{3} \\ \Rightarrow a &= \frac{5}{2}, b = -1, c = \frac{1}{3}\end{aligned}$$

$$18. [\hat{i} \hat{k} \hat{j}] = \hat{i} \cdot (\hat{k} \times \hat{j}) = \hat{i} \cdot (-\hat{i}) = -1.$$

$$\begin{aligned}19. 2\hat{i} \cdot [3\hat{j} \times (-5\hat{k})] &= -30 [\hat{i} \cdot (\hat{j} \times \hat{k})] \\ &= -30(\hat{i} \cdot \hat{i}) = -30(1) \\ &= -30\end{aligned}$$

$$\begin{aligned}20. (\hat{i} + \hat{j}) \cdot [(\hat{j} + \hat{k}) \times (\hat{k} + \hat{i})] \\ &= \begin{vmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{vmatrix} \\ &= 1(1) - 1(-1) = 2\end{aligned}$$

$$\begin{aligned}21. [\bar{a} \bar{b} \bar{c}] &= \begin{vmatrix} 1 & -1 & 1 \\ 1 & 1 & -4 \\ -1 & 2 & -1 \end{vmatrix} \\ &= 1(-1+8) + 1(-1-4) + 1(2+1) \\ &= 5\end{aligned}$$

$$\begin{aligned}22. \bar{a} \cdot (\bar{b} \times \bar{c}) &= \begin{vmatrix} 3 & -2 & 2 \\ 6 & 4 & -2 \\ 3 & -2 & -4 \end{vmatrix} \\ &= 3(-16-4) + 2(-24+6) + 2(-12-12) \\ &= -144\end{aligned}$$

$$24. \text{ Since } [\bar{a} \bar{b} \bar{c}] = [\bar{b} \bar{c} \bar{a}] = [\bar{c} \bar{a} \bar{b}] = -[\bar{b} \bar{a} \bar{c}]$$

$$\begin{aligned}25. [\hat{i} \hat{k} \hat{j}] + [\hat{k} \hat{j} \hat{i}] + [\hat{j} \hat{k} \hat{i}] &= [\hat{i} \hat{k} \hat{j}] + [\hat{i} \hat{k} \hat{j}] - [\hat{i} \hat{k} \hat{j}] \\ &= [\hat{i} \hat{k} \hat{j}] = -1\end{aligned}$$

$$\begin{aligned}27. [\bar{a} + 2\bar{b} \quad \bar{a} + \bar{c} \quad \bar{b}] \\ &= [\bar{a} \quad \bar{a} + \bar{c} \quad \bar{b}] + [2\bar{b} \quad \bar{a} + \bar{c} \quad \bar{b}] \\ &= [\bar{a} \bar{a} \bar{b}] + [\bar{a} \bar{c} \bar{b}] + [2\bar{b} \bar{a} \bar{b}] + [2\bar{b} \bar{c} \bar{b}] \\ &= 0 - [\bar{a} \bar{b} \bar{c}] + 2(0) + 2(0) \\ &= -[\bar{a} \bar{b} \bar{c}]\end{aligned}$$

$$28. \text{ Let } \bar{p} = \bar{a} - 2\bar{b} + 3\bar{c}, \bar{q} = 2\bar{a} + m\bar{b} - 4\bar{c} \text{ and } \bar{r} = -7\bar{b} + \bar{c}$$

Since the points are collinear.

$$\begin{aligned}\therefore [\bar{p} \quad \bar{q} \quad \bar{r}] &= 0 \\ \Rightarrow \begin{vmatrix} 1 & -2 & 3 \\ 2 & m & -4 \\ 0 & -7 & 10 \end{vmatrix} &= 0 \\ \Rightarrow 1(10m-28) + 2(20-0) + 3(-14-0) &= 0 \\ \Rightarrow 10m-30 &= 0 \Rightarrow m=3\end{aligned}$$

$$29. \text{ Let } \bar{a} = \hat{i} + 2\hat{j} + 3\hat{k}, \bar{b} = \lambda\hat{i} + 4\hat{j} + 7\hat{k}, \text{ and } \bar{c} = -3\hat{i} - 2\hat{j} - 5\hat{k}$$

Since the vectors are collinear,

$$\begin{aligned}\therefore \begin{vmatrix} 1 & 2 & 3 \\ \lambda & 4 & 7 \\ -3 & -2 & -5 \end{vmatrix} &= 0 \\ \Rightarrow -6 + 10\lambda - 42 - 6\lambda + 36 &= 0 \\ \Rightarrow \lambda &= 3\end{aligned}$$

30. We know that,

$$\begin{aligned}[\bar{a} - \bar{b} \quad \bar{b} - \bar{c} \quad \bar{c} - \bar{a}] &= 0 \\ \therefore \text{ Vectors } \bar{a} - \bar{b}, \bar{b} - \bar{c} \text{ and } \bar{c} - \bar{a} &\text{ are coplanar}\end{aligned}$$



31. Since, the vectors are coplanar,

$$\therefore [\bar{a} \ \bar{b} \ \bar{c}] = 0$$

$$\Rightarrow \begin{vmatrix} 1 & -1 & 1 \\ 1 & 2 & -1 \\ 3 & p & 5 \end{vmatrix} = 0$$

$$\Rightarrow 10 + p + 5 + 3 + p - 6 = 0$$

$$\Rightarrow p = -6$$

32. Let $\bar{a} = \hat{i} + 3\hat{j} + 2\hat{k}$, $\bar{b} = 2\hat{i} - \hat{j} + 4\hat{k}$, and $\bar{c} = 3\hat{i} + 2\hat{j} + x\hat{k}$

Since the vectors are coplanar,

$$\therefore \begin{vmatrix} 1 & 3 & -2 \\ 2 & -1 & 4 \\ 3 & 2 & x \end{vmatrix} = 0$$

$$\Rightarrow -x - 8 - 6x + 36 - 14 = 0$$

$$\Rightarrow x = 2$$

33. Let $\bar{a} = \hat{i} + \hat{j} + \hat{k}$, $\bar{b} = -\hat{i} + \hat{j}$ and $\bar{c} = \hat{i} + 2\hat{j} + a\hat{k}$

Since \bar{a} , \bar{b} and \bar{c} are coplanar.

$$\therefore \begin{vmatrix} 1 & 1 & 1 \\ -1 & 1 & 0 \\ 1 & 2 & a \end{vmatrix} = 0$$

$$\Rightarrow 1(a-0) - 1(-a-0) + 1(-2-1) = 0$$

$$\Rightarrow 2a = 3 \Rightarrow a = \frac{3}{2}$$

35. We have $[\bar{a} \ \bar{b} \ \bar{a} \times \bar{b}] = \bar{a} \cdot [\bar{b} \times (\bar{a} \times \bar{b})]$
 $= (\bar{a} \times \bar{b}) \cdot (\bar{a} \times \bar{b})$
 $= |\bar{a} \times \bar{b}|^2$

36. $[\bar{a} \ \bar{c} \ \bar{b}] = \bar{a} \cdot (\bar{c} \times \bar{b})$
 $= \bar{c} \cdot (\bar{b} \times \bar{a})$
 $= 0 \quad \dots [\because \bar{a} \text{ and } \bar{b} \text{ are parallel}]$

37. $\bar{a} \cdot (\bar{b} \times \bar{c}) = 0$ or $(\bar{a} \times \bar{b}) \cdot \bar{c} = 0$

38. Volume of parallelopiped = $[\bar{a} \ \bar{b} \ \bar{c}]$

$$= \left(\frac{11}{2}\right)(12) \left(\frac{13}{3}\right) [\hat{i} \ \hat{j} \ \hat{k}]$$

$$= 286 \text{ cu. unit.}$$

39. Let $\bar{a} = 2\hat{i} + \hat{j} - \hat{k}$, $\bar{b} = \hat{i} + 2\hat{j} + 3\hat{k}$ and $\bar{c} = -3\hat{i} - \hat{j} + \hat{k}$

$$\therefore \text{volume of parallelopiped} = \begin{vmatrix} 2 & 1 & -1 \\ 1 & 2 & 3 \\ -3 & -1 & 1 \end{vmatrix}$$

$$= 2(2+3) - 1(1+9) - 1(1-6)$$

$$= 5 \text{ cu. unit.}$$

$$40. \text{Volume of parallelopiped} = \begin{vmatrix} -3 & 7 & 5 \\ -3 & 7 & -3 \\ 7 & -5 & -3 \end{vmatrix}$$

$$= (-3)(-21-15) - 7(9+21) + 5(15-49)$$

$$= 108 - 210 - 170 = -272$$

But volume cannot be negative.

\therefore Volume of parallelopiped = 272 cu. unit.

41. A, B, C, D are vertices of tetrahedron.

$\therefore \overline{AB}$, \overline{AC} and \overline{AD} are its edges.

$$\text{Now, } \overline{AB} = -2\hat{i} - 2\hat{j} - 3\hat{k}$$

$$\overline{AC} = 4\hat{i} - 9\hat{k}$$

$$\overline{AD} = 6\hat{i} - 3\hat{j} - 3\hat{k}$$

$$\therefore \text{Volume of tetrahedron} = \frac{1}{6} [\overline{AB} \ \overline{AC} \ \overline{AD}]$$

$$= \frac{1}{6} \begin{vmatrix} -2 & -2 & -3 \\ 4 & 0 & -9 \\ 6 & -3 & -3 \end{vmatrix}$$

$$= \frac{1}{6} [-2(0-27) + 2(-12+54) - 3(-12-0)]$$

$$= \frac{1}{6} (174) = 29 \text{ cu. unit.}$$

42. Let A, B, C, D be the given points

$\therefore \overline{AB} = 2\hat{i} - 3\hat{j} + 6\hat{k}$, $\overline{AC} = -4\hat{i} - 5\hat{j} + 9\hat{k}$ and

$$\overline{AD} = -6\hat{i} - 2\hat{j} + 6\hat{k}$$

$$\therefore \text{Volume of tetrahedron} = \frac{1}{6} \begin{vmatrix} 2 & -3 & 6 \\ -4 & -5 & 9 \\ -6 & -2 & 6 \end{vmatrix}$$


$$= \frac{-66}{6} = -11$$

But volume cannot be negative

\therefore Volume of tetrahedron = 11 cu. unit.



43. Consider $\triangle ABC$,
AD, BE and CF are its medians.



$$\begin{aligned} \therefore \overline{AD} + \overline{BE} + \overline{CF} &= \overline{d} - \overline{a} + \overline{e} - \overline{b} + \overline{f} - \overline{c} \\ &= \frac{\overline{b} + \overline{c}}{2} - \overline{a} + \frac{\overline{a} + \overline{c}}{2} - \overline{b} + \frac{\overline{a} + \overline{b}}{2} - \overline{c} = 0 \end{aligned}$$

**Critical Thinking**

- Let $A(\overline{a})$, $B(\overline{b})$, $C(\overline{c})$ be the given points
 $\overline{a} = 60\hat{i} + 3\hat{j}$, $\overline{b} = 40\hat{i} - 8\hat{j}$, $\overline{c} = a\hat{i} - 52\hat{j}$
 $\therefore \overline{AB} = k(\overline{BC})$
 $\Rightarrow -20\hat{i} - 11\hat{j} = k[(a-40)\hat{i} - 44\hat{j}]$
On comparing, we get
 $-11 = -44k \Rightarrow k = \frac{1}{4}$
and $-20 = \frac{1}{4}(a-40) \Rightarrow a = -40$
- Let $\overline{a} = \hat{i} + 2\hat{k}$, $\overline{b} = \hat{j} + \hat{k}$ and $\overline{c} = \lambda\hat{i} + \mu\hat{j}$
 $\therefore \overline{AB} = m \cdot \overline{BC}$
 $\Rightarrow -\hat{i} + \hat{j} - \hat{k} = m[(\lambda\hat{i} + (\mu-1)\hat{j} - \hat{k})]$
On comparing, we get
 $-1 = -m \Rightarrow m = 1$,
 $-1 = \lambda \cdot m \Rightarrow \lambda = -1$,
and $1 = m(\mu-1) \Rightarrow \mu = 2$
- Let $\overline{a} = -\hat{i} + 3\hat{j} + 2\hat{k}$, $\overline{b} = -4\hat{i} + 2\hat{j} - 2\hat{k}$ and
 $\overline{c} = 5\hat{i} + \lambda\hat{j} + \mu\hat{k}$
 $\therefore \overline{AB} = m \cdot \overline{BC}$
 $\Rightarrow -3\hat{i} - \hat{j} - 4\hat{k} = m[9\hat{i} + (\lambda-2)\hat{j} + (\mu+2)\hat{k}]$
On comparing, we get
 $9m = -3 \Rightarrow m = \frac{-1}{3}$,
 $-1 = m(\lambda-2) \Rightarrow \lambda = 5$
and $-4 = m(\mu+2) \Rightarrow \mu = 10$
- Here $\overline{a} = \hat{i} + x\hat{j} + 3\hat{k}$, $\overline{b} = 3\hat{i} + 4\hat{j} + 7\hat{k}$, and
 $\overline{c} = y\hat{i} - 2\hat{j} - 5\hat{k}$
 $\therefore \overline{AB} = \lambda \overline{BC}$
 $\Rightarrow 2\hat{i} + (4-x)\hat{j} + 4\hat{k} = \lambda[(y-3)\hat{i} - 6\hat{j} - 12\hat{k}]$

On comparing, we get

$$4 = -12\lambda \Rightarrow \lambda = \frac{-1}{3},$$

$$4 - x = -6\lambda \Rightarrow x = 2, \text{ and}$$

$$2 = \lambda(y-3) \Rightarrow -6 = y-3 \Rightarrow y = -3$$

5. Here $\overline{a} = \hat{i} + \hat{j}$, $\overline{b} = \hat{i} - \hat{j}$, $\overline{c} = a\hat{i} + b\hat{j} + c\hat{k}$

The points are collinear

$$\therefore \overline{AB} = \lambda \overline{BC}$$

$$\Rightarrow -2\hat{j} = \lambda[(a-1)\hat{i} + (b+1)\hat{j} + c\hat{k}]$$

On comparing, we get

$$\lambda(a-1) = 0, \lambda(b+1) = -2, \lambda c = 0$$

Hence $a = 1$, $c = 0$ and b is arbitrary scalar.

6. Let A, B, C be the three collinear point.

$$\therefore \overline{AB} = \lambda \overline{BC}$$

$$\text{Here, } \overline{AB} = -2\overline{b}, \overline{BC} = (k+1)\overline{b}$$

$$\therefore \forall k \in \mathbb{R} \Rightarrow \overline{AB} = \lambda \overline{BC}$$

7. Since, $\overline{a} + 2\overline{b}$ is collinear with \overline{c} , and $\overline{b} + 3\overline{c}$ is collinear with \overline{a} .

$$\therefore \overline{a} + 2\overline{b} = x\overline{c} \text{ and } \overline{b} + 3\overline{c} = y\overline{a} \quad \forall x, y \in \mathbb{R}$$

$$\therefore \overline{a} + 2\overline{b} + 6\overline{c} = (x+6)\overline{c}$$

$$\text{Also, } \overline{a} + 2\overline{b} + 6\overline{c} = \overline{a} + 2(\overline{b} + 3\overline{c}) = (1+2y)\overline{a}$$

$$\therefore (x+6)\overline{c} = (1+2y)\overline{a}$$

Since, \overline{a} and \overline{c} are non-collinear.

$$\therefore x+6=0 \text{ and } 1+2y=0$$

$$\Rightarrow x = -6 \text{ and } y = -\frac{1}{2}$$

$$\text{Now, } \overline{a} + 2\overline{b} = x\overline{c}$$

$$\Rightarrow \overline{a} + 2\overline{b} + 6\overline{c} = \overline{0}$$

$$8. \overline{AB} = \overline{a} + \overline{b}$$

$$\overline{BD} = 3\overline{a} + 3\overline{b} = 3\overline{AB}$$

\therefore Points A, B, D are collinear.

$$9. \text{ Let } \overline{R} = x\overline{a} + y\overline{b} + z\overline{c}$$

$$\Rightarrow \overline{R} = x(2\overline{p} + 3\overline{q} - \overline{r}) + y(\overline{p} - 2\overline{q} + 2\overline{r})$$

$$+ z(-2\overline{p} + \overline{q} - 2\overline{r})$$

$$\Rightarrow 3\overline{p} - \overline{q} + 2\overline{r} = (2x+y-2z)\overline{p}$$

$$+ (3x-2y+z)\overline{q} + (-x+2y-2z)\overline{r}$$

On comparing, we get

$$2x+y-2z=3, \quad \dots(i)$$

$$3x-2y+z=-1, \quad \dots(ii)$$

$$-x+2y-2z=2 \quad \dots(iii)$$

Solving above equations, we get

$$x=2, y=5, z=3$$

$$\therefore \overline{R} = 2\overline{a} + 5\overline{b} + 3\overline{c}$$




10. $\vec{a} + \vec{b} + \vec{c} + \vec{d} = (1 + \lambda)\vec{d}$
 Also, $\vec{a} + \vec{b} + \vec{c} + \vec{d} = (1 + \mu)\vec{a}$
 $\Rightarrow (1 + \lambda)\vec{d} = (1 + \mu)\vec{a}$
 if $\lambda \neq -1$, then
 $\vec{d} = \left(\frac{1 + \mu}{1 + \lambda}\right)\vec{a}$
 Now, $\vec{a} + \vec{b} + \vec{c} + \vec{d} = (1 + \mu)\vec{a}$
 $\therefore \vec{a} + \vec{b} + \vec{c} + \left(\frac{1 + \mu}{1 + \lambda}\right)\vec{a} = (1 + \mu)\vec{a}$
 $\Rightarrow \left[1 + \left(\frac{1 + \mu}{1 + \lambda}\right) - (1 + \mu)\right]\vec{a} + \vec{b} + \vec{c} = 0$
 This contradicts the fact that $\vec{a}, \vec{b}, \vec{c}$ are non-coplanar
 $\Rightarrow \lambda = -1$
 $\therefore \vec{a} + \vec{b} + \vec{c} + \vec{d} = \vec{0}$

11. The position vector of A is $6\vec{b} - 2\vec{a}$ and the position vector of P is $\vec{a} - \vec{b}$
 Let the position vector of B be \vec{r}
 Since, P divides AB in the ratio 1 : 2
 $\therefore \vec{a} - \vec{b} = \frac{1(\vec{r}) + 2(6\vec{b} - 2\vec{a})}{3}$
 $\Rightarrow 3\vec{a} - 3\vec{b} - 12\vec{b} + 4\vec{a} = \vec{r}$
 $\Rightarrow \vec{r} = 7\vec{a} - 15\vec{b}$

12. $2\vec{a} + 3\vec{b} - 5\vec{c} = \vec{0}$
 $\Rightarrow 5\vec{c} = 2\vec{a} + 3\vec{b} \Rightarrow \vec{c} = \frac{3\vec{b} + 2\vec{a}}{3 + 2}$
 \therefore point C divides segment AB internally in the ratio 3:2.

13. $|\vec{OA}| = \sqrt{1+9+4} = \sqrt{14}$
 $|\vec{OB}| = \sqrt{9+1+4} = \sqrt{14}$
 $\therefore OA = OB$
 Let C be any point on angle bisector and on line AB
 $\therefore C$ is midpoint of AB



$\therefore \vec{c} = \frac{\vec{a} + \vec{b}}{2} = 2\hat{i} + 2\hat{j} - 2\hat{k}$

14. P(\vec{p}) divide AB internally in the ratio 3 : 1.
 $\therefore \vec{p} = \frac{3\vec{b} + \vec{a}}{4}$
 Q(\vec{q}) is midpoint of AP
 $\therefore \vec{q} = \frac{\vec{a} + \vec{p}}{2} = \frac{\vec{a} + \frac{3\vec{b} + \vec{a}}{4}}{2} = \frac{5\vec{a} + 3\vec{b}}{8}$

15. $2\vec{a} + \vec{b} = 3\vec{c}$
 $\Rightarrow 2\vec{a} = 3\vec{c} - \vec{b}$
 $\Rightarrow \vec{a} = \frac{3\vec{c} - \vec{b}}{2} = \frac{3\vec{c} - \vec{b}}{3 - 1}$
 \therefore A divides BC in the ratio 3 : 1 externally.

16. P(\vec{p}) is midpoint of BC
 $\therefore \vec{p} = \frac{\vec{b} + \vec{c}}{2}$
 $\Rightarrow 2\vec{p} = \vec{b} + \vec{c}$ (i)
 Q(\vec{q}) divides CA internally in the ratio 2:1
 $\therefore \vec{q} = \frac{2\vec{a} + \vec{c}}{3}$
 $\Rightarrow 3\vec{q} = 2\vec{a} + \vec{c}$ (ii)
 R(\vec{r}) divides AB externally in the ratio 1:2
 $\vec{r} = \frac{\vec{b} - 2\vec{a}}{1 - 2}$
 $= \frac{2\vec{p} - 3\vec{q}}{-1}$ [From (i) and (ii)]
 $\therefore \vec{r} = -2\vec{p} + 3\vec{q}$
 \therefore points P, Q and R are collinear.

17. $[\vec{a} \vec{b} \vec{c}] = \begin{vmatrix} 1 & 0 & -1 \\ x & 1 & 1-x \\ y & x & 1+x-y \end{vmatrix}$
 Applying, $C_3 \Rightarrow C_3 + C_1$
 $= \begin{vmatrix} 1 & 0 & 0 \\ x & 1 & 1 \\ y & x & 1+x \end{vmatrix} = 1(1 + x - x) = 1$

18. Let A \equiv (1, 1, 2), B \equiv (2, 1, p), C \equiv (1, 0, 3) and D \equiv (2, 2, 0).
 $\therefore \vec{AB} = \hat{i} + (p - 2)\hat{k}$
 $\vec{AC} = -\hat{j} + \hat{k}$, and
 $\vec{AD} = \hat{i} + \hat{j} - 2\hat{k}$
 The points are coplanar.
 $\therefore \vec{AB}, \vec{AC}$ and \vec{AD} are coplanar
 $[\vec{AB} \vec{AC} \vec{AD}] = 0$
 $\Rightarrow \begin{vmatrix} 1 & 0 & p-2 \\ 0 & -1 & 1 \\ 1 & 1 & -2 \end{vmatrix} = 0$
 $\Rightarrow 1(2 - 1) + (p - 2)(1) = 0$
 $\Rightarrow 1 + p - 2 = 0 \Rightarrow p = 1$



19. Since the points are coplanar,

$$\therefore \begin{vmatrix} 1 & 2 & 0 \\ 0 & 1 & 4 \\ \lambda - 1 & 2 & 3 \end{vmatrix} = 0$$

$$\Rightarrow 1(3 - 8) - 2[(0 - 4(\lambda - 1))] = 0$$

$$\Rightarrow -5 + 8\lambda - 8 = 0 \Rightarrow \lambda = \frac{13}{8}$$

20. Since, the given vectors are coplanar,

$$\therefore \begin{vmatrix} a & 1 & 1 \\ 1 & -b & 1 \\ 1 & 1 & -c \end{vmatrix} = 0$$

$$\Rightarrow a(bc - 1) - 1(-c - 1) + 1(1 + b) = 0$$

$$\Rightarrow abc - a + c + 1 + 1 + b = 0$$

$$\Rightarrow abc + 2 = a - b - c$$

21. Since the given vectors are coplanar,

$$\therefore \begin{vmatrix} -bc & b^2 + bc & c^2 + bc \\ a^2 + ac & -ac & c^2 + ac \\ a^2 + ab & b^2 + ab & -ab \end{vmatrix} = 0$$

$$\Rightarrow (ab + bc + ca)^3 = 0 \Rightarrow ab + bc + ca = 0.$$

22. Let $P(\vec{p})$, $Q(\vec{q})$, $R(\vec{r})$ be the three points.

$$\therefore \vec{p} = \vec{a} - \vec{b} + \vec{c}, \vec{q} = 4\vec{a} - 7\vec{b} - \vec{c} \text{ and}$$

$$\vec{r} = 3\vec{a} + 6\vec{b} + 6\vec{c}$$

\vec{PQ} is not scalar multiple of \vec{PR}

\therefore they are not collinear

$$\begin{bmatrix} \vec{p} & \vec{q} & \vec{r} \end{bmatrix} = \begin{vmatrix} 1 & -1 & 1 \\ 4 & -7 & -1 \\ 3 & 6 & 6 \end{vmatrix}$$

$$= 36 \neq 0$$

\therefore they are not coplanar.

$$23. \frac{(\vec{b} \times \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c})}{\lambda}$$

$$= \frac{(\vec{b} \times \vec{c}) \cdot \vec{a} + (\vec{b} \times \vec{c}) \cdot \vec{b} + (\vec{b} \times \vec{c}) \cdot \vec{c}}{\lambda}$$

$$= \frac{(\vec{b} \times \vec{c}) \cdot \vec{a} + \vec{0} + \vec{0}}{\lambda} = \frac{\vec{a} \cdot (\vec{b} \times \vec{c})}{\lambda} = \frac{\lambda}{\lambda} = 1$$

$$\begin{aligned} 24. & (\vec{a} - \vec{b}) \cdot [(\vec{b} + \vec{c}) \times (\vec{c} + \vec{a})] \\ &= (\vec{a} - \vec{b}) \cdot [\vec{b} \times \vec{c} + \vec{b} \times \vec{a} + \vec{c} \times \vec{c} + \vec{c} \times \vec{a}] \\ &= \vec{a} \cdot (\vec{b} \times \vec{c}) + \vec{a} \cdot (\vec{b} \times \vec{a}) + \vec{a} \cdot (\vec{c} \times \vec{a}) \\ &\quad - \vec{b} \cdot (\vec{b} \times \vec{c}) - \vec{b} \cdot (\vec{b} \times \vec{a}) - \vec{b} \cdot (\vec{c} \times \vec{a}) \\ &= [\vec{a} \vec{b} \vec{c}] - [\vec{a} \vec{b} \vec{c}] = 0 \end{aligned}$$

$$\begin{aligned} 25. & [\vec{a} + \vec{b} + \vec{c}] \cdot [(\vec{a} + \vec{b}) \times (\vec{a} + \vec{c})] \\ &= (\vec{a} + \vec{b}) \cdot [(\vec{a} + \vec{b}) \times (\vec{a} + \vec{c})] \\ &\quad + \vec{c} \cdot [(\vec{a} + \vec{b}) \times (\vec{a} + \vec{c})] \end{aligned}$$

$$= 0 + [\vec{c} \vec{a} + \vec{b} \vec{a} + \vec{c}]$$

$$= [\vec{c} \vec{a} \vec{a} + \vec{c}] + [\vec{c} \vec{b} \vec{a} + \vec{c}]$$

$$= [\vec{c} \vec{a} \vec{a}] + [\vec{c} \vec{a} \vec{c}] + [\vec{c} \vec{b} \vec{a}] + [\vec{c} \vec{b} \vec{c}]$$

$$= 0 + 0 + [\vec{c} \vec{b} \vec{a}] + 0 = -[\vec{a} \vec{b} \vec{c}]$$

$$26. \vec{r} = (\vec{b} \times \vec{c}) + m(\vec{c} \times \vec{a}) + n(\vec{a} \times \vec{b})$$

$$\therefore \vec{a} \cdot \vec{r} = \vec{a} \cdot (\vec{b} \times \vec{c}) + m\vec{a} \cdot (\vec{c} \times \vec{a}) + n\vec{a} \cdot (\vec{a} \times \vec{b})$$

$$= [\vec{a} \vec{b} \vec{c}] + 0 + 0$$

$$\vec{a} \cdot \vec{r} = 2 \quad \dots \left[\because [\vec{a} \vec{b} \vec{c}] = 2 \right] \dots (i)$$

Similarly,

$$\vec{b} \cdot \vec{r} = 2m, \quad \dots (ii)$$

$$\vec{c} \cdot \vec{r} = 2n \quad \dots (iii)$$

\therefore On adding equations (i), (ii) and (iii) we get

$$(\vec{a} + \vec{b} + \vec{c}) \cdot \vec{r} = 2(m + n)$$

$$\therefore m + n = \frac{1}{2}(\vec{a} + \vec{b} + \vec{c}) \cdot \vec{r}$$

27. Volume of parallelepiped

$$= \begin{vmatrix} 1 & 2 & -1 \\ 1 & -1 & 0 \\ 1 & -1 & -1 \end{vmatrix} [\vec{a} \vec{b} \vec{c}] = k[\vec{a} \vec{b} \vec{c}]$$

$$\Rightarrow 1(1 - 0) - 2(-1 - 0) - 1(-1 + 1) = k$$

$$\Rightarrow 1 + 2 - 0 = k \Rightarrow k = 3$$

$$28. \text{Volume of parallelepiped} = \begin{vmatrix} -p & 0 & 5 \\ 1 & -1 & q \\ 3 & -5 & 0 \end{vmatrix} = 8$$

$$\Rightarrow -p(0 + 5q) + 5(-5 + 3) = 8$$

$$\Rightarrow -5pq - 18 = 0$$

$$\Rightarrow 5pq + 18 = 0$$

29. Let $A \equiv (1, 2, 0)$, $B \equiv (2, 0, 4)$, $C \equiv (-1, 2, 0)$ and $D \equiv (-1, 1, \lambda)$ be the vertices of the tetrahedron

$$\therefore \vec{AB} = \hat{i} - 2\hat{j} + 4\hat{k}$$

$$\vec{AC} = -2\hat{i}$$

$$\vec{AD} = -2\hat{i} - \hat{j} + \lambda\hat{k}$$



$$\text{Volume of tetrahedron} = \frac{1}{6} [\overline{AB \ AC \ AD}]$$

$$\Rightarrow \frac{2}{3} = \frac{1}{6} \begin{vmatrix} 1 & -2 & 4 \\ -2 & 0 & 0 \\ -2 & -1 & \lambda \end{vmatrix}$$

$$\Rightarrow 2(-2\lambda) + 4(2) = 4$$

$$\Rightarrow \lambda = 1$$

$$\begin{aligned} 30. \quad \overline{AB} + \overline{BC} + \overline{AC} &= \overline{b} - \overline{a} + \overline{c} - \overline{b} + \overline{c} - \overline{a} \\ &= 2(\overline{c} - \overline{a}) \\ &= 2(\overline{c} - \overline{d} + \overline{d} - \overline{a}) \\ &= 2(\overline{DC} + \overline{AD}) \\ &= 2(\overline{DC} - \overline{BD}) \\ &\dots[\because D \text{ is mid-point of } AB] \end{aligned}$$


31. If AD is the median,

$$\begin{aligned} \therefore \overline{AD} &= \overline{d} - \overline{a} = \frac{\overline{b} + \overline{c}}{2} - \overline{a} \\ &= \frac{(\overline{b} - \overline{a}) + (\overline{c} - \overline{a})}{2} = \frac{1}{2} (\overline{AB} + \overline{AC}) \\ &= 4\hat{i} - \hat{j} + 4\hat{k} \end{aligned}$$

$$\therefore (AD) = \sqrt{16+1+16} = \sqrt{33}$$

$$\begin{aligned} 32. \quad \overline{AA'} + \overline{BB'} + \overline{CC'} &= \overline{a'} - \overline{a} + \overline{b'} - \overline{b} + \overline{c'} - \overline{c} \\ &= (\overline{a'} + \overline{b'} + \overline{c'}) - (\overline{a} + \overline{b} + \overline{c}) \\ &= 3\overline{g'} - 3\overline{g} \\ &\dots[\because G' \text{ and } G \text{ are centroids}] \\ &= 3\overline{GG'} \end{aligned}$$

33.



$$\overline{HA} = \overline{HO} + \overline{OA} \quad \dots(i)$$

$$\overline{HB} = \overline{HO} + \overline{OB} \quad \dots(ii)$$

$$\overline{HC} = \overline{HO} + \overline{OC} \quad \dots(iii)$$

\therefore Adding (i), (ii) and (iii), we get

$$\begin{aligned} \overline{HA} + \overline{HB} + \overline{HC} \\ = 3\overline{HO} + \overline{OA} + \overline{OB} + \overline{OC} \end{aligned}$$

$$\text{Since, } \overline{OA} + \overline{OB} + \overline{OC} = \overline{OH} = -\overline{HO}$$

$$\therefore \overline{HA} + \overline{HB} + \overline{HC} = 2\overline{HO}$$


34. Let \overline{a} , \overline{b} , \overline{c} , \overline{d} be the position vectors of A, B, C, D respectively

For parallelogram:

$$\overline{a} + \overline{c} = \overline{b} + \overline{d}$$

$$\Rightarrow \overline{d} = \overline{a} + \overline{c} - \overline{b} \Rightarrow \overline{d} = -\hat{i} + \hat{j} + \hat{k}$$

35.



$$\begin{aligned} \text{We have, } \overline{p} &= \overline{AC} + \overline{BD} \\ &= \overline{AC} + \overline{BC} + \overline{CD} \\ &= \overline{AC} + \lambda \overline{AD} + \overline{CD} \\ &= \lambda \overline{AD} + (\overline{AC} + \overline{CD}) \\ &= \lambda \overline{AD} + \overline{AD} \\ &= (\lambda + 1) \overline{AD} \end{aligned}$$

$$\text{Also, } \overline{p} = \mu \overline{AD}$$

$$\therefore \mu = \lambda + 1$$



Competitive Thinking

1. Here, $\overline{a} = \hat{i}$, $\overline{b} = \hat{j}$, $\overline{c} = x\hat{i} + 8\hat{j}$

$$\overline{AB} = -\hat{i} + \hat{j}, \overline{BC} = x\hat{i} + 7\hat{j}$$

Since the points are collinear,

$$\begin{aligned} \therefore \overline{AB} &= \lambda \overline{BC} \\ \Rightarrow -\hat{i} + \hat{j} &= \lambda (x\hat{i} + 7\hat{j}) \end{aligned}$$

On comparing, we get

$$7\lambda = 1 \Rightarrow \lambda = \frac{1}{7}$$

$$\lambda x = -1 \Rightarrow x = -7$$

2. Let A(\overline{a}), B(\overline{b}), C(\overline{c}) be the given points

$$\therefore \overline{a} = 20\hat{i} + p\hat{j}, \overline{b} = 5\hat{i} - \hat{j}, \overline{c} = 10\hat{i} - 13\hat{j}$$

$$\therefore \overline{AB} = k \overline{BC}$$

$$\Rightarrow -15\hat{i} - (p+1)\hat{j} = k(5\hat{i} - 12\hat{j})$$

On comparing, we get

$$-15 = 5k \Rightarrow k = -3 \text{ and}$$

$$-(p+1) = -12k$$

$$\Rightarrow -(p+1) = 36$$

$$\Rightarrow p = -37$$


3. $\overline{PQ} = k \overline{QR}$

$$\overline{a} + \overline{b} - \overline{c} = k(-2\overline{a} - 2\overline{b} + \overline{c})$$

On comparing, we get

$$1 = -2k \Rightarrow k = -\frac{1}{2} \text{ and } -1 = kt \Rightarrow t = 2$$



4. Here $\overline{AB} = \overline{b} - \overline{a}$ and
 $\overline{AC} = 2\overline{a} - 2\overline{b} = -2(\overline{b} - \overline{a})$
 $\therefore \overline{AC} = m\overline{AB}$
Hence A, B, C are collinear.
5. Since, $\overline{a} + 3\overline{b}$ is collinear with \overline{c} , and $\overline{b} + 2\overline{c}$ is collinear with \overline{a} ,
 $\therefore \overline{a} + 3\overline{b} = x\overline{c}$ and $\overline{b} + 2\overline{c} = y\overline{a} \quad \forall x, y \in \mathbb{R}$.
 $\therefore \overline{a} + 3\overline{b} + 6\overline{c} = (x+6)\overline{c}$
Also, $\overline{a} + 3\overline{b} + 6\overline{c} = \overline{a} + 3(\overline{b} + 2\overline{c}) = (1+3y)\overline{a}$
 $\therefore (x+6)\overline{c} = (1+3y)\overline{a}$
 $\Rightarrow (x+6)\overline{c} - (1+3y)\overline{a} = 0$
 $\therefore x+6=0$ and $1+3y=0$
 $\Rightarrow x = -6$ and $y = -\frac{1}{3}$
Now, $\overline{a} + 3\overline{b} = x\overline{c} \Rightarrow \overline{a} + 3\overline{b} + 6\overline{c} = \overline{0}$
6. Let $\overline{a} = 3\hat{i} + 2\hat{j} - \hat{k}$ and $\overline{b} = 6\hat{i} - 4x\hat{j} + y\hat{k}$
Since, \overline{a} and \overline{b} are parallel,
 $\frac{3}{6} = \frac{2}{-4x} = \frac{-1}{y}$
 $\Rightarrow x = -1$ and $y = -2$
7. The given vectors are collinear.
 $\therefore \frac{3}{a} = \frac{1}{b} = \frac{-5}{-15}$
 $\Rightarrow a = 9, b = 3$
8. $x = 0, y = 0$, otherwise one vector will be a scalar multiple of the other and hence collinear which is a contradiction.
11. $\overline{c} = m\overline{a} + n\overline{b}$
 $\therefore 3\hat{i} - \hat{k} = m(\hat{i} + \hat{j} - 2\hat{k}) + n(2\hat{i} - \hat{j} + \hat{k})$
Comparing the co-efficients of \hat{i} and \hat{j} , we get
 $3 = m + 2n$, and(i)
 $m = n$ (ii)
 \therefore Solving the above two equations, we get
 $m = n = 1$
 $\therefore m + n = 1 + 1 = 2$
12. Let P(\overline{p}) divide the line internally in the ratio 2 : 3
 $\therefore \overline{p} = \frac{3(2\overline{a} - 3\overline{b}) + 2(3\overline{a} - 2\overline{b})}{2+3} = \frac{12\overline{a} - 13\overline{b}}{5}$
13. $A \equiv (1, -1, 2), B \equiv (2, 3, -1)$
Point P divides AB internally in the ratio 2 : 3.
 $\therefore P \equiv \left[\frac{2(2) + 3(1)}{2+3}, \frac{2(3) + 3(-1)}{2+3}, \frac{2(-1) + 3(2)}{2+3} \right]$
 $\equiv \left(\frac{7}{5}, \frac{3}{5}, \frac{4}{5} \right)$
 \therefore the position vector of P is $\frac{1}{5}(7\hat{i} + 3\hat{j} + 4\hat{k})$
14. 
C divides AB internally in the ratio 1 : 2 and
D divides AB internally in the ratio 2 : 1.
 $\therefore z_1 + z_2 = \frac{1(6) + 2(4)}{1+2} + \frac{2(6) + 1(4)}{2+1}$
 $= \frac{14}{3} + \frac{16}{3} = \frac{30}{3}$
 $= 10$
15. Let position vector of B be \overline{r}
Since, \overline{a} divides AB in the ratio 2 : 3
 $\therefore \frac{2\overline{r} + 3(\overline{a} + 2\overline{b})}{2+3} = \overline{a}$
 $\Rightarrow 2\overline{r} = 5\overline{a} - 3\overline{a} - 6\overline{b} = 2\overline{a} - 6\overline{b}$
 $\Rightarrow \overline{r} = \overline{a} - 3\overline{b}$
16. We know that, centroid of a triangle divides the line segment joining the orthocentre and circumcentre in the ratio 2 : 1.
The co-ordinates of orthocentre and circumcentre are $(-1, 3, 2), (5, 3, 2)$ respectively.
 \therefore Co-ordinates of centroid
 $\equiv \left(\frac{2(5) + 1(-1)}{2+1}, \frac{2(3) + 1(3)}{2+1}, \frac{2(2) + 1(2)}{2+1} \right)$
 $\equiv (3, 3, 2)$
17. Let the co-ordinates of circumcentre be (x, y, z) .
Co-ordinates of orthocentre and centroid are $(-3, 5, 2)$ and $(3, 3, 4)$ respectively.
We know that, centroid of triangle divides the line segment joining its orthocentre and circumcentre in the ratio 2 : 1.
 $\therefore \left(\frac{2x - 3}{3}, \frac{2y + 5}{3}, \frac{2z + 2}{3} \right) \equiv (3, 3, 4)$
 $\Rightarrow \frac{2x - 3}{3} = 3, \frac{2y + 5}{3} = 3, \frac{2z + 2}{3} = 4$
 $\Rightarrow x = 6, y = 2, z = 5$



18. Let $N(\bar{n})$ divide line segment LM externally in the ratio $2 : 1$.

$$\begin{aligned}\therefore \bar{n} &= \frac{2(\bar{a} + 2\bar{b}) - (2\bar{a} - \bar{b})}{2 - 1} \\ &= \frac{2\bar{a} + 4\bar{b} - 2\bar{a} + \bar{b}}{1} = 5\bar{b}\end{aligned}$$

19. $R(\bar{r})$ divides PQ externally in the ratio $2 : 1$

$$\begin{aligned}\therefore \bar{r} &= \frac{2(-\hat{i} + \hat{j} - \hat{k}) - 1(\hat{i} + 2\hat{j} - \hat{k})}{2 - 1} \\ &= -2\hat{i} + 2\hat{j} - 2\hat{k} - \hat{i} - 2\hat{j} + \hat{k}\end{aligned}$$

$$\therefore \bar{r} = -3\hat{i} - \hat{k}$$

$$20. \quad 3\bar{P} + 2\bar{R} - 5\bar{Q} = \bar{0}$$

$$\Rightarrow 3\bar{P} + 2\bar{R} = 5\bar{Q}$$

$$\Rightarrow \bar{Q} = \frac{3\bar{P} + 2\bar{R}}{5}$$

- $\therefore \bar{Q}$ is the position vector of the point dividing P and R in the ratio $3 : 2$ internally.

Thus, P , Q and R are collinear.

21. Let the point B divide AC in the ratio $\lambda : 1$

$$\begin{aligned}\therefore 5\hat{i} - 2\hat{k} &= \frac{\lambda(11\hat{i} + 3\hat{j} + 7\hat{k}) + \hat{i} - 2\hat{j} - 8\hat{k}}{\lambda + 1} \\ \Rightarrow \lambda(5\hat{i} - 2\hat{k}) + (5\hat{i} - 2\hat{k}) &= \lambda(11\hat{i} + 3\hat{j} + 7\hat{k}) + (\hat{i} - 2\hat{j} - 8\hat{k})\end{aligned}$$

$$\Rightarrow -6\lambda\hat{i} - 3\lambda\hat{j} - 9\lambda\hat{k} = -4\hat{i} - 2\hat{j} - 6\hat{k}$$

$$\Rightarrow -6\lambda = -4$$

$$\Rightarrow \lambda = \frac{2}{3} \text{ i.e. ratio} = 2 : 3$$

22. M and N are the midpoints of sides PQ and RS

$$\therefore \bar{m} = \frac{\bar{p} + \bar{q}}{2} \text{ and } \bar{n} = \frac{\bar{r} + \bar{s}}{2}$$

$$\Rightarrow 2\bar{m} = \bar{p} + \bar{q} \text{ and } 2\bar{n} = \bar{r} + \bar{s}$$

$$\begin{aligned}\overline{PS} + \overline{QR} &= \bar{s} - \bar{p} + \bar{r} - \bar{q} \\ &= \bar{r} + \bar{s} - (\bar{p} + \bar{q}) \\ &= 2\bar{n} - 2\bar{m} \\ &= 2\overline{MN}\end{aligned}$$

23. Let the position vectors of A, B, C, L, M, N and K be $\bar{a}, \bar{b}, \bar{c}, \bar{l}, \bar{m}, \bar{n}$ and \bar{k} respectively.

$$\bar{l} = \frac{2\bar{b} + \bar{c}}{3}, \quad \bar{m} = \frac{2\bar{a} + 3\bar{c}}{5}, \quad \bar{n} = \frac{3\bar{b} + 5\bar{a}}{8},$$

$$\bar{k} = \frac{5\bar{b} + 3\bar{a}}{8}$$

$$\frac{|\overline{AL} + \overline{BM} + \overline{CN}|}{|\overline{CK}|}$$

$$= \frac{\left| \frac{2\bar{b} + \bar{c}}{3} - \bar{a} + \frac{2\bar{a} + 3\bar{c}}{5} - \bar{b} + \frac{3\bar{b} + 5\bar{a}}{8} - \bar{c} \right|}{\left| \frac{5\bar{b} + 3\bar{a}}{8} - \bar{c} \right|}$$

$$= \frac{\left| \frac{1}{120} [80\bar{b} + 40\bar{c} - 120\bar{a} + 48\bar{a} + 72\bar{c} - 120\bar{b} + 45\bar{b} + 75\bar{a} - 120\bar{c}] \right|}{\left| \frac{1}{8} [5\bar{b} + 3\bar{a} - 8\bar{c}] \right|}$$

$$= \frac{\left| \frac{1}{120} [3\bar{a} + 5\bar{b} - 8\bar{c}] \right|}{\left| \frac{1}{8} [3\bar{a} + 5\bar{b} - 8\bar{c}] \right|}$$

$$= \frac{1}{15}$$

24. G is the centroid.

$$\therefore \overline{OG} = \frac{\overline{OA} + \overline{OB} + \overline{OC}}{3}$$

$$\Rightarrow \overline{OA} + \overline{OB} + \overline{OC} = 3\overline{OG}$$

25. $\overline{GA} + \overline{GB} + \overline{GC} = (\bar{a} - \bar{g}) + (\bar{b} - \bar{g}) + (\bar{c} - \bar{g})$

$$= \bar{a} + \bar{b} + \bar{c} - 3\bar{g}$$

$$= \bar{a} + \bar{b} + \bar{c} - 3 \left(\frac{\bar{a} + \bar{b} + \bar{c}}{3} \right) = \bar{0}$$

$$\begin{aligned}26. \quad \bar{a} \cdot (\bar{b} \times \bar{c}) &= \begin{vmatrix} 2 & 1 & -1 \\ 1 & 2 & 1 \\ 1 & -1 & 2 \end{vmatrix} \\ &= 2(4 + 1) - 1(2 - 1) - 1(-1 - 2) \\ &= 12\end{aligned}$$

27. $\bar{a} \cdot (\bar{b} \times \bar{c}) = 10$

$$\therefore \begin{vmatrix} 1 & 1 & 1 \\ 2 & \lambda & 1 \\ 1 & -1 & 4 \end{vmatrix} = 10$$

$$\Rightarrow (4\lambda + 1) - (8 - 1) + (-2 - \lambda) = 10$$

$$\Rightarrow \lambda = 6$$



28. Let \hat{n} be the unit vector perpendicular to \vec{a} and \vec{b}

$$\begin{aligned} [\vec{a} \ \vec{b} \ \vec{c}] &= \vec{a} \cdot (\vec{b} \times \vec{c}) = \vec{a} \cdot (|\vec{b}| |\vec{c}| \sin \theta \hat{n}) \\ &= \vec{a} \cdot (3 \times 4 \sin \frac{2\pi}{3} \cdot \hat{n}) = \vec{a} \cdot \left(12 \times \frac{\sqrt{3}}{2} \hat{n} \right) \\ &= 6\sqrt{3} |\vec{a}| |\hat{n}| \cos 0 = 6\sqrt{3} \times 2 \times 1 \Rightarrow 12\sqrt{3}. \end{aligned}$$

29. $|\vec{a}| = 1, |\vec{b}| = 2, |\vec{c}| = 3$

$$\begin{aligned} &[\vec{a} + \vec{b} + \vec{c} \ \vec{b} - \vec{a} \ \vec{c}] \\ &= (\vec{a} + \vec{b} + \vec{c}) \cdot [(\vec{b} - \vec{a}) \times \vec{c}] \\ &= (\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{b} \times \vec{c} - \vec{a} \times \vec{c}) \\ &= [\vec{a} \ \vec{b} \ \vec{c}] - [\vec{b} \ \vec{a} \ \vec{c}] \\ &= 2[\vec{a} \ \vec{b} \ \vec{c}] \\ &= 2 \vec{a} \cdot (\vec{b} \times \vec{c}) \\ &= 2 |\vec{a}| \cdot |\vec{b} \times \vec{c}| \cos 0^\circ \\ &= 2 |\vec{a}| \cdot |\vec{b} \times \vec{c}| \\ &= 2 |\vec{a}| \cdot |\vec{b}| \cdot |\vec{c}| \sin 90^\circ \\ &= 2 \times 1 \times 2 \times 3 = 12 \end{aligned}$$

30. $[\vec{a} - \vec{b} \ \vec{b} - \vec{c} \ \vec{c} - \vec{a}] = [\vec{a} \ \vec{b} \ \vec{c}] - [\vec{b} \ \vec{c} \ \vec{a}]$
 $= [\vec{a} \ \vec{b} \ \vec{c}] - [\vec{a} \ \vec{b} \ \vec{c}] = 0$

31. $(\vec{a} - \vec{b}) \cdot [(\vec{b} - \vec{c}) \times (\vec{c} - \vec{a})]$
 $= [\vec{a} - \vec{b} \ \vec{b} - \vec{c} \ \vec{c} - \vec{a}] = 0$

32. Vector $\vec{\alpha}$ lies in the plane of $\vec{\beta}$ and $\vec{\gamma}$
 $\therefore \vec{\alpha}, \vec{\beta}, \vec{\gamma}$ are coplanar
 $\Rightarrow [\vec{\alpha} \ \vec{\beta} \ \vec{\gamma}] = 0$

33. $[\vec{a} + \vec{b} \ \vec{b} + \vec{c} \ \vec{c} + \vec{a}] = 2[\vec{a} \ \vec{b} \ \vec{c}]$
 $= 0$ [$\because \vec{a}, \vec{b}, \vec{c}$ are coplanar]

34. $[\vec{a} + \vec{b} \ \vec{b} + \vec{c} \ \vec{c} + \vec{a}] = 2[\vec{a} \ \vec{b} \ \vec{c}]$

35. Here $\vec{C} = C_1 \hat{i} - \hat{j} + \hat{k}$

To make three vectors coplanar $[\vec{A} \ \vec{B} \ \vec{C}] = 0$

$$\Rightarrow \begin{vmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ C_1 & -1 & 1 \end{vmatrix} = 0$$

$$\Rightarrow 1(0 - 0) - 1(1 - 0) + 1(-1 - 0) = 0$$

- \therefore The value of $[\vec{A} \ \vec{B} \ \vec{C}]$ is independent of C_1
Hence no value of C_1 can be found.

36. Let $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$, $\vec{b} = \hat{i} + 2\hat{j} - 3\hat{k}$ and
 $\vec{c} = 3\hat{i} + \lambda\hat{j} + 5\hat{k}$

Since, \vec{a}, \vec{b} and \vec{c} are coplanar,

$$\therefore [\vec{a} \ \vec{b} \ \vec{c}] = 0$$

$$\Rightarrow \begin{vmatrix} 2 & -1 & 1 \\ 1 & 2 & -3 \\ 3 & \lambda & 5 \end{vmatrix} = 0$$

$$\Rightarrow 2(10 + 3\lambda) + 1(5 + 9) + 1(\lambda - 6) = 0$$

$$\Rightarrow \lambda = -4$$

37. Let \vec{a}, \vec{b} and \vec{c} be the given vectors

The given vectors are coplanar

$$\therefore \begin{vmatrix} \lambda & 1 & 2 \\ 1 & \lambda & -1 \\ 2 & -1 & \lambda \end{vmatrix} = 0$$

$$\Rightarrow \lambda(\lambda^2 - 1) - (\lambda + 2) + 2(-1 - 2\lambda) = 0$$

$$\Rightarrow \lambda^3 - 6\lambda - 4 = 0$$

$$\Rightarrow (\lambda + 2)(\lambda^2 - 2\lambda - 2) = 0$$

$$\Rightarrow \lambda = -2 \text{ or } \lambda = \frac{2 \pm \sqrt{4 + 8}}{2} = 1 \pm \sqrt{3}$$

38. Let $\vec{a} = 4\hat{i} + 11\hat{j} + m\hat{k}$, $\vec{b} = 7\hat{i} + 2\hat{j} + 6\hat{k}$ and
 $\vec{c} = \hat{i} + 5\hat{j} + 4\hat{k}$.

Since \vec{a}, \vec{b} and \vec{c} are coplanar,

$$\therefore [\vec{a} \ \vec{b} \ \vec{c}] = 0$$

$$\Rightarrow \begin{vmatrix} 4 & 11 & m \\ 7 & 2 & 6 \\ 1 & 5 & 4 \end{vmatrix} = 0$$

$$\Rightarrow 4(8 - 30) - 11(28 - 6) + m(35 - 2) = 0$$

$$\Rightarrow -330 + 33m = 0$$

$$\Rightarrow m = 10$$

39. Here $\vec{a} = 2\hat{i} + 2\hat{j} + 6\hat{k}$, $\vec{b} = 2\hat{i} + \lambda\hat{j} + 6\hat{k}$,
 $\vec{c} = 2\hat{i} - 3\hat{j} + \hat{k}$

Since \vec{a}, \vec{b} and \vec{c} are coplanar,

$$\therefore \begin{vmatrix} 2 & 2 & 6 \\ 2 & \lambda & 6 \\ 2 & -3 & 1 \end{vmatrix} = 0$$

$$\Rightarrow 2(\lambda + 18) - 2(2 - 12) + 6(-6 - 2\lambda) = 0$$

$$\Rightarrow -10\lambda = -20$$

$$\Rightarrow \lambda = 2$$



40. Let $\vec{a} = -2\hat{i} + \hat{j} + \hat{k}$, $\vec{b} = \hat{i} + \hat{j} + \hat{k}$, $\vec{c} = \hat{j} - \hat{k}$ and $\vec{d} = \lambda\hat{j} + \hat{k}$

Since the given points are coplanar.

$$\therefore [\vec{AB} \vec{AC} \vec{AD}] = 0$$

$$\Rightarrow \begin{vmatrix} 3 & 0 & 0 \\ 2 & 0 & -2 \\ 2 & \lambda - 1 & 0 \end{vmatrix} = 0$$

$$\Rightarrow 3(2\lambda - 2) + 0 + 0 = 0$$

$$\Rightarrow 6\lambda - 6 = 0$$

$$\Rightarrow \lambda = 1$$

41. Since $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = \hat{i} - \hat{j} + 2\hat{k}$ and $\vec{c} = x\hat{i} + (x-2)\hat{j} - \hat{k}$ are coplanar vectors

$$\therefore [\vec{a} \vec{b} \vec{c}] = 0$$

$$\Rightarrow \begin{vmatrix} 1 & 1 & 1 \\ 1 & -1 & 2 \\ x & x-2 & -1 \end{vmatrix} = 0$$

$$\Rightarrow 1[1 - 2(x-2)] - 1(-1 - 2x) + 1(x-2+x) = 0$$

$$\Rightarrow 1 - 2x + 4 + 1 + 2x + 2x - 2 = 0$$

$$\Rightarrow 2x = -4$$

$$\Rightarrow x = -2$$

42. Let $\vec{a} = 3\hat{i} - 2\hat{j} - \hat{k}$, $\vec{b} = 2\hat{i} + 3\hat{j} - 4\hat{k}$, $\vec{c} = -\hat{i} + \hat{j} + 2\hat{k}$ and $\vec{d} = 4\hat{i} + 5\hat{j} + \lambda\hat{k}$

Since, the given points are coplanar,

$$\therefore [\vec{AB} \vec{AC} \vec{AD}] = 0$$

$$\Rightarrow \begin{vmatrix} -1 & 5 & -3 \\ -4 & 3 & 3 \\ 1 & 7 & \lambda + 1 \end{vmatrix} = 0$$

$$\Rightarrow -1(3\lambda + 3 - 21) - 5(-4\lambda - 4 - 3) - 3(-28 - 3) = 0$$

$$\Rightarrow -3\lambda + 18 + 20\lambda + 35 + 93 = 0$$

$$\Rightarrow 17\lambda = -146$$

$$\Rightarrow \lambda = \frac{-146}{17}$$

43. Let $\vec{s} = 2\vec{a} + 3\vec{b} - \vec{c}$, $\vec{t} = \vec{a} - 2\vec{b} + 3\vec{c}$,

$$\vec{u} = 3\vec{a} + 4\vec{b} - 2\vec{c}, \vec{v} = k\vec{a} - 6\vec{b} + 6\vec{c}$$

$$\therefore \vec{ST} = -\vec{a} - 5\vec{b} + 4\vec{c}, \vec{SU} = \vec{a} + \vec{b} - \vec{c}$$

$$\vec{SV} = (k-2)\vec{a} - 9\vec{b} + 7\vec{c}$$

Since, the given points are coplanar,

$$\therefore [\vec{ST} \vec{SU} \vec{SV}] = 0$$

$$\Rightarrow \begin{vmatrix} -1 & -5 & 4 \\ 1 & 1 & -1 \\ k-2 & -9 & 7 \end{vmatrix} = 0$$

$$\Rightarrow 2 + 5(7+k-2) + 4(-9-k+2) = 0$$

$$\Rightarrow 2 + 25 + 5k - 28 - 4k = 0$$

$$\Rightarrow -1 + k = 0$$

$$\Rightarrow k = 1$$

44. Since, $a\hat{i} + \hat{j} + \hat{k}$, $\hat{i} + b\hat{j} + \hat{k}$ and $\hat{i} + \hat{j} + c\hat{k}$ are coplanar,

$$\therefore \begin{vmatrix} a & 1 & 1 \\ 1 & b & 1 \\ 1 & 1 & c \end{vmatrix} = 0$$

$$\Rightarrow a(bc - 1) - 1(c - 1) + 1(1 - b) = 0$$

$$\Rightarrow abc - a - b - c + 2 = 0$$

$$\Rightarrow abc - (a + b + c) = -2$$

45. Let \vec{a}, \vec{b} and \vec{c} be the given vectors.

The vectors are coplanar

$$\therefore \begin{vmatrix} -\lambda^2 & 1 & 1 \\ 1 & -\lambda^2 & 1 \\ 1 & 1 & -\lambda^2 \end{vmatrix} = 0$$

$$\Rightarrow -\lambda^2(\lambda^4 - 1) - 1(-\lambda^2 - 1) + 1(1 + \lambda^2) = 0$$

$$\Rightarrow \lambda^6 - 3\lambda^2 - 2 = 0$$

$$\Rightarrow (1 + \lambda^2)^2 (\lambda^2 - 2) = 0$$

$$\Rightarrow \lambda = \pm \sqrt{2}$$

46. The given vectors are coplanar

$$\therefore \begin{vmatrix} \lambda^3 & 0 & 1 \\ 1 & -\lambda^3 & 0 \\ 1 & 2\lambda - \sin \lambda & -\lambda \end{vmatrix} = 0$$

$$\Rightarrow \lambda^3(\lambda^4 - 0) + 1(2\lambda - \sin \lambda + \lambda^3) = 0$$

$$\Rightarrow \lambda^7 + \lambda^3 + 2\lambda = \sin \lambda \quad \dots(i)$$

This is true for $\lambda = 0$.

For non-zero values of λ , equation (i) is

$$\lambda^6 + \lambda^2 + 2 = \frac{\sin \lambda}{\lambda} \quad \dots(ii)$$

We know that $\frac{\sin x}{x} < 1$ for all $x \neq 0$.

\therefore L.H.S. of (ii) is greater than 2 and R.H.S. is less than 1.

So, (ii) is not true for any non-zero λ .

Hence, there is only one value of λ .



47. Let $\vec{\alpha}$, $\vec{\beta}$ and $\vec{\gamma}$ be the given vectors

$\vec{\alpha}$, $\vec{\beta}$ and $\vec{\gamma}$ are coplanar

$$\therefore \begin{vmatrix} 1 & 2 & 3 \\ 0 & \lambda & 4 \\ 0 & 0 & (2\lambda - 1) \end{vmatrix} = 0$$

$$\Rightarrow \lambda(2\lambda - 1) = 0 \Rightarrow \lambda = 0, \frac{1}{2}$$

Hence, $\vec{\alpha}$, $\vec{\beta}$, $\vec{\gamma}$ are non-coplanar for all values of λ except 0 and $\frac{1}{2}$.

48. Since, O(0, 0, 0), P(2, 3, 4), Q(1, 2, 3), R(x, y, z) are co-planar

$$\therefore [\overline{OR} \quad \overline{OP} \quad \overline{OQ}] = 0$$

$$\Rightarrow \begin{vmatrix} x & y & z \\ 2 & 3 & 4 \\ 1 & 2 & 3 \end{vmatrix} = 0$$

$$\Rightarrow x(9 - 8) - y(6 - 4) + z(4 - 3) = 0$$

$$\Rightarrow x - 2y + z = 0$$

49. Let the vector be $a\hat{i} + b\hat{j} + c\hat{k}$.

It is perpendicular to $2\hat{i} + \hat{j} + \hat{k}$.

$$\therefore 2a + b + c = 0 \quad \dots(i)$$

The vector is coplanar with $\hat{i} + 2\hat{j} + \hat{k}$ and $\hat{i} + \hat{j} + 2\hat{k}$

$$\therefore \begin{vmatrix} a & b & c \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{vmatrix} = 0$$

$$\therefore 3a - b - c = 0 \quad \dots(ii)$$

On solving (i) and (ii), we get

$$a = 0, b = 5, c = -5$$

\therefore The required vector is $5(\hat{j} - \hat{k})$

$$50. [\vec{\alpha} \quad \vec{\beta} \quad \vec{\gamma}] = \begin{vmatrix} 5 & 6 & 7 \\ 7 & -8 & 9 \\ 3 & 20 & 5 \end{vmatrix}$$

$$= 5(-40 - 180) - 6(35 - 27) + 7(140 + 24) = 0$$

\therefore the given vectors are coplanar.

51. Since \vec{x} is a non-zero vector, the given conditions will be satisfied, if either

i. at least one of the vectors \vec{a} , \vec{b} , \vec{c} is zero or

ii. \vec{x} is perpendicular to all the vectors \vec{a} , \vec{b} , \vec{c}

In case (ii), \vec{a} , \vec{b} , \vec{c} are coplanar

$$\Rightarrow [\vec{a} \quad \vec{b} \quad \vec{c}] = 0$$

54. options (A), (B) and (D) = $[\vec{u} \quad \vec{v} \quad \vec{w}]$,

while option (C) = $-[\vec{u} \quad \vec{v} \quad \vec{w}]$

$$55. \vec{a} \cdot (\vec{a} \times \vec{b}) = (\vec{a} \times \vec{a}) \cdot \vec{b} = 0$$

$$56. \frac{\vec{a} \cdot \vec{b} \times \vec{c}}{\vec{c} \times \vec{a} \cdot \vec{b}} + \frac{\vec{b} \cdot \vec{a} \times \vec{c}}{\vec{c} \cdot \vec{a} \times \vec{b}} = \frac{\vec{a} \cdot \vec{b} \times \vec{c}}{\vec{c} \cdot \vec{a} \times \vec{b}} + \frac{\vec{b} \cdot \vec{a} \times \vec{c}}{\vec{c} \cdot \vec{a} \times \vec{b}}$$

$$= \frac{[\vec{a} \quad \vec{b} \quad \vec{c}]}{[\vec{c} \quad \vec{a} \quad \vec{b}]} + \frac{[\vec{b} \quad \vec{a} \quad \vec{c}]}{[\vec{c} \quad \vec{a} \quad \vec{b}]}$$

$$= \frac{[\vec{a} \quad \vec{b} \quad \vec{c}]}{[\vec{c} \quad \vec{a} \quad \vec{b}]} - \frac{[\vec{a} \quad \vec{b} \quad \vec{c}]}{[\vec{c} \quad \vec{a} \quad \vec{b}]} = 0$$

57. \vec{a} , \vec{b} and \vec{c} are non-coplanar.

$$\text{So, } [\vec{a} \quad \vec{b} \quad \vec{c}] \neq 0$$

$$\vec{a} \cdot \left\{ \frac{\vec{b} \times \vec{c}}{3\vec{b} \cdot (\vec{c} \times \vec{a})} \right\} - \vec{b} \cdot \left\{ \frac{\vec{c} \times \vec{a}}{2\vec{c} \cdot (\vec{a} \times \vec{b})} \right\}$$

$$= \frac{[\vec{a} \quad \vec{b} \quad \vec{c}]}{3[\vec{b} \quad \vec{c} \quad \vec{a}]} - \frac{[\vec{b} \quad \vec{c} \quad \vec{a}]}{2[\vec{c} \quad \vec{a} \quad \vec{b}]} = \frac{1}{3} - \frac{1}{2} = -\frac{1}{6}$$

58. $\vec{a} \cdot [(\vec{b} + \vec{c}) \times (\vec{a} + \vec{b} + \vec{c})]$

$$= \vec{a} \cdot [\vec{b} \times \vec{a} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} + \vec{c} \times \vec{b}]$$

$$\dots [\because \vec{b} \times \vec{b} = 0, \vec{c} \times \vec{c} = 0]$$

$$= [\vec{a} \quad \vec{b} \quad \vec{a}] + [\vec{a} \quad \vec{b} \quad \vec{c}] + [\vec{a} \quad \vec{c} \quad \vec{a}] + [\vec{a} \quad \vec{c} \quad \vec{b}]$$

$$= 0 + [\vec{a} \quad \vec{b} \quad \vec{c}] + 0 - [\vec{a} \quad \vec{b} \quad \vec{c}]$$

$$= 0$$

59. $(\vec{a} + \vec{b}) \cdot (\vec{b} + \vec{c}) \times (\vec{a} + \vec{b} + \vec{c})$

$$= (\vec{a} + \vec{b}) \cdot [\vec{b} \times \vec{a} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} + \vec{c} \times \vec{b}]$$

$$= [\vec{a} \quad \vec{b} \quad \vec{a}] + [\vec{a} \quad \vec{b} \quad \vec{c}] + [\vec{a} \quad \vec{c} \quad \vec{a}] + [\vec{a} \quad \vec{c} \quad \vec{b}]$$

$$+ [\vec{b} \quad \vec{b} \quad \vec{a}] + [\vec{b} \quad \vec{b} \quad \vec{c}] + [\vec{b} \quad \vec{c} \quad \vec{a}] + [\vec{b} \quad \vec{c} \quad \vec{b}]$$

$$= 0 + [\vec{a} \quad \vec{b} \quad \vec{c}] + 0 + [\vec{a} \quad \vec{c} \quad \vec{b}] + 0 + 0 + [\vec{b} \quad \vec{c} \quad \vec{a}] + 0$$

$$= [\vec{a} \quad \vec{b} \quad \vec{c}] - [\vec{a} \quad \vec{b} \quad \vec{c}] + [\vec{a} \quad \vec{b} \quad \vec{c}] = [\vec{a} \quad \vec{b} \quad \vec{c}]$$

60. Since, $\vec{a} \cdot \vec{b} = 0$

\therefore \vec{a} and \vec{b} are perpendicular unit vectors.

$$\text{Now, } (2\vec{a} - \vec{b}) \cdot \{(\vec{a} \times \vec{b}) \times (\vec{a} + 2\vec{b})\}$$

$$= [2\vec{a} - \vec{b} \quad \vec{a} \times \vec{b} \quad \vec{a} + 2\vec{b}]$$

$$= -[\vec{a} \times \vec{b} \quad 2\vec{a} - \vec{b} \quad \vec{a} + 2\vec{b}]$$

$$= -(\vec{a} \times \vec{b}) \cdot \{ (2\vec{a} - \vec{b}) \times (\vec{a} + 2\vec{b}) \}$$



$$\begin{aligned}
 &= -(\bar{a} \times \bar{b}) \cdot 5(\bar{a} \times \bar{b}) \\
 &= -5|\bar{a} \times \bar{b}| = -5|\bar{a}|^2|\bar{b}|^2 \quad \dots [\because \bar{a} \perp \bar{b}] \\
 &= -5 \quad \dots [\because |\bar{a}| = |\bar{b}| = 1]
 \end{aligned}$$

$$\begin{aligned}
 61. \quad \bar{p} + \bar{q} + \bar{r} &= \frac{\bar{b} \times \bar{c} + \bar{c} \times \bar{a} + \bar{a} \times \bar{b}}{[\bar{a} \ \bar{b} \ \bar{c}]} \\
 (\bar{a} + \bar{b} + \bar{c}) \cdot (\bar{p} + \bar{q} + \bar{r}) &= \frac{[\bar{a} \ \bar{b} \ \bar{c}] + [\bar{b} \ \bar{c} \ \bar{a}] + [\bar{c} \ \bar{a} \ \bar{b}]}{[\bar{a} \ \bar{b} \ \bar{c}]} \\
 &= 3
 \end{aligned}$$

$$\begin{aligned}
 62. \quad (\bar{u} + \bar{v} - \bar{w}) \cdot [(\bar{u} - \bar{v}) \times (\bar{v} - \bar{w})] \\
 &= \bar{u} \cdot (\bar{u} \times \bar{v}) - \bar{u} \cdot (\bar{u} \times \bar{w}) + \bar{u} \cdot (\bar{v} \times \bar{w}) + \bar{v} \cdot (\bar{u} \times \bar{v}) \\
 &\quad - \bar{v} \cdot (\bar{u} \times \bar{w}) + \bar{v} \cdot (\bar{v} \times \bar{w}) - \bar{w} \cdot (\bar{u} \times \bar{v}) \\
 &\quad + \bar{w} \cdot (\bar{u} \times \bar{w}) - \bar{w} \cdot (\bar{v} \times \bar{w}) \\
 &= [\bar{u} \ \bar{v} \ \bar{w}] - [\bar{v} \ \bar{u} \ \bar{w}] - [\bar{w} \ \bar{u} \ \bar{v}] \\
 &= [\bar{u} \ \bar{v} \ \bar{w}] + [\bar{u} \ \bar{v} \ \bar{w}] - [\bar{u} \ \bar{v} \ \bar{w}] = \bar{u} \cdot (\bar{v} \times \bar{w})
 \end{aligned}$$

$$\begin{aligned}
 63. \quad \bar{p} \cdot (\bar{a} + \bar{b}) &= \bar{p} \cdot \bar{a} + \bar{p} \cdot \bar{b} \\
 &= \frac{(\bar{b} \times \bar{c}) \cdot \bar{a}}{[\bar{a} \ \bar{b} \ \bar{c}]} + \frac{(\bar{b} \times \bar{c}) \cdot \bar{b}}{[\bar{a} \ \bar{b} \ \bar{c}]} \\
 &= \frac{[\bar{b} \ \bar{c} \ \bar{a}]}{[\bar{a} \ \bar{b} \ \bar{c}]} + \frac{[\bar{b} \ \bar{c} \ \bar{b}]}{[\bar{a} \ \bar{b} \ \bar{c}]} \\
 &= 1 + 0 \\
 &= 1
 \end{aligned}$$

Similarly, $\bar{q} \cdot (\bar{b} + \bar{c}) = 1$ and $\bar{r} \cdot (\bar{a} + \bar{c}) = 1$
 $(\bar{a} + \bar{b}) \cdot \bar{p} + (\bar{b} + \bar{c}) \cdot \bar{q} + (\bar{c} + \bar{a}) \cdot \bar{r}$
 $= 1 + 1 + 1 = 3$

$$\begin{aligned}
 64. \quad \text{Since } d &= \lambda a + \mu b + \nu c \\
 \therefore d \cdot (b \times c) &= \lambda a \cdot (b \times c) + \mu b \cdot (b \times c) + \nu c \cdot (b \times c) \\
 \Rightarrow d \cdot (b \times c) &= \lambda [a \ b \ c] \\
 \Rightarrow \lambda &= \frac{[d \ b \ c]}{[a \ b \ c]} = \frac{[b \ c \ d]}{[b \ c \ a]}
 \end{aligned}$$

$$\begin{aligned}
 65. \quad [3\bar{u} \ \bar{p} \ \bar{v} \ \bar{p} \ \bar{w}] - [p\bar{v} \ \bar{w} \ \bar{q} \ \bar{u}] - [2\bar{w} \ \bar{q} \ \bar{v} \ \bar{q} \ \bar{u}] &= 0 \\
 \Rightarrow 3p^2 [\bar{u} \ \bar{v} \ \bar{w}] - pq [\bar{v} \ \bar{w} \ \bar{u}] - 2q^2 [\bar{w} \ \bar{v} \ \bar{u}] &= 0 \\
 \Rightarrow 3p^2 [\bar{u} \ \bar{v} \ \bar{w}] - pq [\bar{u} \ \bar{v} \ \bar{w}] + 2q^2 [\bar{u} \ \bar{v} \ \bar{w}] &= 0 \\
 \Rightarrow (3p^2 - pq + 2q^2) [\bar{u} \ \bar{v} \ \bar{w}] &= 0
 \end{aligned}$$

But, $[\bar{u} \ \bar{v} \ \bar{w}] \neq 0$

$\therefore 3p^2 - pq + 2q^2 = 0$

$$\begin{aligned}
 \Rightarrow p^2 - \frac{1}{3}pq + \frac{2}{3}q^2 &= 0 \\
 \Rightarrow \left(p^2 - \frac{1}{3}pq + \frac{1}{36}q^2\right) - \frac{1}{36}q^2 + \frac{2}{3}q^2 &= 0 \\
 \Rightarrow \left(p - \frac{q}{6}\right)^2 + \frac{23}{36}q^2 &= 0
 \end{aligned}$$

$\Rightarrow p - \frac{q}{6} = 0, q = 0$

$\Rightarrow p = 0, q = 0$

Hence, there is exactly one value of (p, q).

$$\begin{aligned}
 66. \quad [\lambda(\bar{a} + \bar{b}) \ \lambda^2 \bar{b} \ \lambda \bar{c}] &= [\bar{a} \ \bar{b} + \bar{c} \ \bar{b}] \\
 \Rightarrow \lambda^4 [\bar{a} + \bar{b} \ \bar{b} \ \bar{c}] &= [\bar{a} \ \bar{b} + \bar{c} \ \bar{b}] \\
 \Rightarrow \lambda^4 \{[\bar{a} \ \bar{b} \ \bar{c}] + [\bar{b} \ \bar{b} \ \bar{c}]\} &= \{[\bar{a} \ \bar{b} \ \bar{b}] + [\bar{a} \ \bar{c} \ \bar{b}]\} \\
 \Rightarrow \lambda^4 [\bar{a} \ \bar{b} \ \bar{c}] &= -[\bar{a} \ \bar{b} \ \bar{c}] \\
 \Rightarrow (\lambda^4 + 1) [\bar{a} \ \bar{b} \ \bar{c}] &= 0
 \end{aligned}$$

But, $[\bar{a} \ \bar{b} \ \bar{c}] \neq 0$.

$\therefore \lambda^4 + 1 = 0$

This is not true for any real value of λ .

67. Let $\bar{a} = 2\hat{i} - 3\hat{j}$, $\bar{b} = \hat{i} + \hat{j} - \hat{k}$ and $\bar{c} = 3\hat{i} - \hat{k}$

Volume of parallelopiped = $[\bar{a} \ \bar{b} \ \bar{c}]$

$$\begin{vmatrix} 2 & -3 & 0 \\ 1 & 1 & -1 \\ 3 & 0 & -1 \end{vmatrix}$$

$= 2(-1) + 3(-1 + 3) = 4$ cu.unit.

$$68. \quad \text{Volume of parallelopiped} = \begin{vmatrix} 2 & -3 & 1 \\ 1 & -1 & 2 \\ 2 & 1 & -1 \end{vmatrix}$$

$= 2(1 - 2) + 3(-1 - 4) + 1(1 + 2) = -14$

But, volume cannot be negative.

\therefore Volume of parallelopiped = 14 cu. units.

$$69. \quad \text{Volume of tetrahedron} = \frac{1}{6} [\bar{a} \ \bar{b} \ \bar{c}]$$

$\Rightarrow 4 = \frac{1}{6} [\bar{a} \ \bar{b} \ \bar{c}] \Rightarrow [\bar{a} \ \bar{b} \ \bar{c}] = 24$

Edges of parallelopiped are $\bar{a} \times \bar{b}$, $\bar{b} \times \bar{c}$, $\bar{c} \times \bar{a}$

\therefore Volume of parallelopiped = $[\bar{a} \times \bar{b} \ \bar{b} \times \bar{c} \ \bar{c} \times \bar{a}]$
 $= [\bar{a} \ \bar{b} \ \bar{c}]^2$
 $= 24^2$
 $= 576$ sq. units



$$70. \text{ Volume of parallelepiped} = \begin{vmatrix} -12 & 0 & \alpha \\ 0 & 3 & -1 \\ 2 & 1 & -15 \end{vmatrix}$$

$$\Rightarrow 546 = -12(-45 + 1) + \alpha(0 - 6)$$

$$\Rightarrow \alpha = -3$$

$$71. \text{ Volume of parallelepiped} = [a - b \ b - c \ c - a]$$

$$= [\bar{a} \ \bar{b} \ \bar{c}] - [\bar{b} \ \bar{c} \ \bar{a}]$$

$$= [\bar{a} \ \bar{b} \ \bar{c}] - [\bar{a} \ \bar{b} \ \bar{c}]$$

$$= 0$$

$$72. \text{ Volume of parallelepiped}$$

$$= [\bar{a} + \bar{b} \ \bar{b} + \bar{c} \ \bar{c} + \bar{a}] = 2[\bar{a} \ \bar{b} \ \bar{c}]$$

$$= 2 \begin{vmatrix} 2 & -3 & 5 \\ 3 & -4 & 5 \\ 5 & -3 & -2 \end{vmatrix}$$

$$= 2[2(8 + 15) + 3(-6 - 25) + 5(-9 + 20)]$$

$$= 2[46 - 93 + 55]$$

$$= 16 \text{ cu. Unit}$$

$$73. \text{ Let A, B, C and D be the given points.}$$

$$\therefore \overline{AB} = -4\hat{i} - 6\hat{j}, \quad \overline{AC} = -\hat{i} + 4\hat{j} + 3\hat{k}, \quad \text{and}$$

$$\overline{AD} = -6\hat{i} - \hat{j} + 3\hat{k}$$

$$\text{Volume of tetrahedron} = \frac{1}{6} \begin{vmatrix} -4 & -6 & 0 \\ -1 & 4 & 3 \\ -6 & -1 & 3 \end{vmatrix}$$

$$= \frac{30}{6}$$

$$= 5$$

$$74. \text{ AD is the median}$$

$$\therefore \overline{AD} = \frac{\overline{AB} + \overline{AC}}{2} \quad \begin{matrix} 3\hat{i} + 5\hat{j} + 4\hat{k} & & 5\hat{i} - 5\hat{j} + 2\hat{k} \\ & \triangle & \\ & \text{A} & \\ & \text{B} \quad \text{D} \quad \text{C} & \end{matrix}$$

$$\Rightarrow \overline{AD} = \frac{(3+5)\hat{i} + (5-5)\hat{j} + (4+2)\hat{k}}{2}$$

$$= \frac{8\hat{i} + 6\hat{k}}{2}$$


$$= 4\hat{i} + 3\hat{k}$$

$$\therefore (\text{AD}) = |\overline{AD}| = \sqrt{16+9} = 5 \text{ units.}$$

$$75. \text{ Let AM be the angle bisector of } \angle \text{BAC}$$

$$|\overline{AB}| = \sqrt{16+25+9} = \sqrt{50} = 5\sqrt{2}$$

$$|\overline{AC}| = \sqrt{1+1+16} = \sqrt{18} = 3\sqrt{2}$$



$$\therefore \text{ M divides BC internally in the ratio } 5 : 3$$

$$\begin{aligned} \overline{M} &= \frac{5\bar{c} + 3\bar{b}}{8} = \frac{5(3\hat{i} + 2\hat{j} + \hat{k}) + 3(-2\hat{j} + 2\hat{k})}{8} \\ &= \frac{15\hat{i} + 4\hat{j} + 11\hat{k}}{8} \end{aligned}$$

$$\therefore \overline{M} = \left(\frac{15}{8}, \frac{4}{8}, \frac{11}{8} \right)$$

$$76. \text{ Let AM be the angle bisector of angle A}$$

$$|\overline{AB}| = 6 \text{ and } |\overline{AC}| = 3$$

$$\therefore \text{ M divides BC internally in the ratio } 2 : 1$$

$$\begin{aligned} \therefore \overline{M} &= \frac{2(2\hat{i} + 5\hat{j} + 7\hat{k}) + 1(2\hat{i} + 3\hat{j} + 4\hat{k})}{2+1} \\ &= \frac{6\hat{i} + 13\hat{j} + 18\hat{k}}{3} \end{aligned}$$

$$77. \overline{AB} = -6\hat{i} - 2\hat{j} + 3\hat{k}, \quad \overline{BC} = -2\hat{i} + 3\hat{j} - 6\hat{k}$$

$$\overline{CD} = 6\hat{i} + 2\hat{j} - 3\hat{k}, \quad \overline{DA} = 2\hat{i} - 3\hat{j} + 6\hat{k}$$

$$\overline{AC} = -8\hat{i} + \hat{j} - 3\hat{k} \text{ and } \overline{BD} = 4\hat{i} + 5\hat{j} - 9\hat{k}$$

$$\text{Here, } |\overline{AB}| = |\overline{BC}| = |\overline{CD}| = |\overline{DA}| = 7$$

$$\text{and } \overline{AC} \cdot \overline{BD} = 0 \Rightarrow \overline{AC} \perp \overline{BD}$$

Hence, ABCD is a rhombus.

$$78. \text{ In } \triangle ABC, \text{ hypotenuse } AB = p$$

$$\therefore \overline{AC} \perp \overline{CB}$$

$$\therefore \overline{AC} \cdot \overline{CB} = 0 \quad \dots(i)$$

$$\text{Now, } \overline{AB} \cdot \overline{AC} + \overline{BC} \cdot \overline{BA} + \overline{CA} \cdot \overline{CB}$$

$$= \overline{AB} \cdot \overline{AC} + \overline{BC} \cdot (-\overline{AB}) + (-\overline{AC}) \cdot \overline{CB}$$

$$= \overline{AB} \cdot \overline{AC} - \overline{BC} \cdot \overline{AB} - \overline{AC} \cdot \overline{CB}$$

$$= \overline{AB} \cdot (\overline{AC} - \overline{BC}) - 0 \quad \dots[\text{From (i)}]$$

$$= \overline{AB} \cdot (\overline{AC} + \overline{CB})$$

$$= \overline{AB} \cdot \overline{AB} \quad \dots[\because \overline{AC} + \overline{CB} = \overline{AB}]$$

$$= |\overline{AB}|^2 = p^2$$



79. Since, \vec{c} is coplanar with \vec{a} and \vec{b}
 $\therefore \vec{c} = x\vec{a} + y\vec{b}$
 $\Rightarrow \vec{c} = x(2\hat{i} + \hat{j} + \hat{k}) + y(\hat{i} + 2\hat{j} - \hat{k})$
 $\Rightarrow \vec{c} = (2x + y)\hat{i} + (x + 2y)\hat{j} + (x - y)\hat{k}$
 Also, $\vec{a} \cdot \vec{c} = 0$ [$\because \vec{c} \perp \vec{a}$]
 $\therefore 2(2x + y) + x + 2y + x - y = 0$
 $\Rightarrow y = -2x$

$\vec{c} = -3x\hat{j} + 3x\hat{k} = 3x(-\hat{j} + \hat{k})$
 Now, $|\vec{c}| = 1$
 $\therefore 9x^2 + 9x^2 = 1$
 $\Rightarrow x = \pm \frac{1}{3\sqrt{2}}$
 $\therefore \vec{c} = \frac{1}{\sqrt{2}}(-\hat{j} + \hat{k})$



Evaluation Test

1. Since, $\vec{a} + \vec{b}$ and $\vec{b} + \vec{c}$ are collinear with \vec{c} and \vec{a} respectively
 $\therefore \vec{a} + \vec{b} = t\vec{c}$... (i)
 $\vec{b} + \vec{c} = s\vec{a}$... (ii)
 From (i) and (ii), we get
 $\vec{a} - \vec{c} = t\vec{c} - s\vec{a} \Rightarrow \vec{a}(1 + s) = \vec{c}(1 + t)$
 But \vec{a} and \vec{c} are non-collinear
 $\therefore 1 + s = 0, 1 + t = 0 \Rightarrow s = -1, t = -1$
 Substituting value of t in (i) and value of s in (ii), we get
 $\vec{a} + \vec{b} = -\vec{c}$ and $\vec{b} + \vec{c} = -\vec{a}$
 Hence, $\vec{a} + \vec{b} + \vec{c} = 0$.

2. Given, $\vec{r} = \lambda_1\vec{r}_1 + \lambda_2\vec{r}_2 + \lambda_3\vec{r}_3$
 $\Rightarrow 2\vec{a} - 3\vec{b} + 4\vec{c} = (\lambda_1 - \lambda_2 + \lambda_3)\vec{a}$
 $\quad + (-\lambda_1 + \lambda_2 + \lambda_3)\vec{b} + (\lambda_1 + \lambda_2 + \lambda_3)\vec{c}$
 $\Rightarrow \lambda_1 - \lambda_2 + \lambda_3 = 2, -\lambda_1 + \lambda_2 + \lambda_3 = -3,$
 $\lambda_1 + \lambda_2 + \lambda_3 = 4$
 $\Rightarrow \lambda_1 = \frac{7}{2}, \lambda_2 = 1, \lambda_3 = -\frac{1}{2}$
 $\therefore \lambda_1 + \lambda_3 = 3$

3. Since, the given vectors are coplanar
 $\therefore \begin{vmatrix} a & a & c \\ 1 & 0 & 1 \\ c & c & b \end{vmatrix} = 0$

Applying $C_2 \rightarrow C_2 - C_1$,

$$\begin{vmatrix} a & 0 & c \\ 1 & -1 & 1 \\ c & 0 & b \end{vmatrix} = 0$$

$$\Rightarrow a(-b) + c(c) = 0 \Rightarrow c^2 = ab$$

Hence, c is the geometric mean of a and b .

4. $\hat{a} \cdot \hat{a} = 1, \hat{b} \cdot \hat{b} = 1, \hat{c} \cdot \hat{c} = 1,$
 $\hat{a} \cdot \hat{b} = \hat{b} \cdot \hat{c} = \hat{c} \cdot \hat{a} = \frac{1}{2}$

$$\therefore [\vec{abc}]^2 = \begin{vmatrix} \hat{a} \cdot \hat{a} & \hat{a} \cdot \hat{b} & \hat{a} \cdot \hat{c} \\ \hat{b} \cdot \hat{a} & \hat{b} \cdot \hat{b} & \hat{b} \cdot \hat{c} \\ \hat{c} \cdot \hat{a} & \hat{c} \cdot \hat{b} & \hat{c} \cdot \hat{c} \end{vmatrix}$$

$$= \begin{vmatrix} 1 & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & 1 & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & 1 \end{vmatrix} = \frac{1}{2}$$

$$\therefore [\vec{abc}] = \frac{1}{\sqrt{2}} \text{ cubic units}$$

5. Since, $\begin{vmatrix} a & 1 & 1 \\ 1 & b & 1 \\ 1 & 1 & c \end{vmatrix} = 0$

Applying $R_2 \rightarrow R_2 - R_1$ and $R_3 \rightarrow R_3 - R_1$, we get

$$\begin{vmatrix} a & 1 & 1 \\ 1-a & b-1 & 0 \\ 1-a & 0 & c-1 \end{vmatrix} = 0$$

$$\Rightarrow a(b-1)(c-1) - (1-a)(c-1) - (1-a)(b-1) = 0$$

Dividing by $(1-a)(1-b)(1-c)$, we get

$$\frac{a}{1-a} + \frac{1}{1-b} + \frac{1}{1-c} = 0 \quad \dots (i)$$

Consider, $\frac{1}{1-a} + \frac{1}{1-b} + \frac{1}{1-c}$

$$= \frac{1}{1-a} - \frac{a}{1-a}$$

$$= 1$$

....[From (i)]



6. Volume of the parallelepiped formed by vectors is

$$\text{i.e., } V = \begin{vmatrix} 1 & a & 1 \\ 0 & 1 & a \\ a & 0 & 1 \end{vmatrix} = 1 - a + a^3$$

$$\therefore \frac{dV}{da} = -1 + 3a^2, \quad \frac{d^2V}{da^2} = 6a$$

$$\text{For max. or min. of } V, \quad \frac{dV}{da} = 0$$

$$\therefore a^2 = \frac{1}{3} \quad \therefore a = \frac{1}{\sqrt{3}}$$

$$\frac{d^2V}{da^2} = 6a > 0 \text{ for } a = \frac{1}{\sqrt{3}}$$

$$\therefore V \text{ is minimum for } a = \frac{1}{\sqrt{3}}$$

7. Given, $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a} = 0$

The scalar triple product of three vectors is

$$[\vec{a} \vec{b} \vec{c}] = (\vec{a} \times \vec{b}) \cdot \vec{c}$$

$$\therefore \vec{a} \cdot \vec{b} = 0 \quad \therefore \vec{a} \perp \vec{b}$$

\therefore angle between \vec{a} and \vec{b} is $\theta = 90^\circ$

Similarly, $[\vec{a} \vec{b} \vec{c}] = |\vec{a}| |\vec{b}| |\hat{n}| |\vec{c}|$ where \hat{n} is a normal vector.

$\therefore \hat{n}$ and \vec{c} are parallel to each other

$$\therefore [\vec{a} \vec{b} \vec{c}] = |\vec{a}| |\vec{b}| |\hat{n}| |\vec{c}| = |\vec{a}| |\vec{b}| |\vec{c}|$$

8. Given, $\vec{r} \times \vec{b} = \vec{c} \times \vec{b}$

$$\Rightarrow (\vec{r} - \vec{c}) \times \vec{b} = \vec{0}$$

$\therefore \vec{r} - \vec{c}$ is parallel to \vec{b}

$$\Rightarrow \vec{r} - \vec{c} = \lambda \vec{b} \text{ for some scalar } \lambda$$

$$\Rightarrow \vec{r} = \vec{c} + \lambda \vec{b} \quad \dots (i)$$

$$\Rightarrow \vec{r} \cdot \vec{a} = \vec{c} \cdot \vec{a} + \lambda (\vec{b} \cdot \vec{a})$$

$$\Rightarrow 0 = \vec{c} \cdot \vec{a} + \lambda (\vec{b} \cdot \vec{a}) \quad \dots [\because \vec{r} \cdot \vec{a} = 0 (\text{given})]$$

$$\Rightarrow \lambda = - \frac{\vec{a} \cdot \vec{c}}{\vec{a} \cdot \vec{b}}$$

Substituting the value of λ in (i), we get

$$\vec{r} = \vec{c} - \frac{\vec{a} \cdot \vec{c}}{\vec{a} \cdot \vec{b}} \vec{b}$$

$$\Rightarrow \vec{r} \cdot \vec{b} = \vec{c} \cdot \vec{b} - \frac{\vec{a} \cdot \vec{c}}{\vec{a} \cdot \vec{b}} (\vec{b} \cdot \vec{b})$$

$$\Rightarrow \vec{r} \cdot \vec{b} = 1 - \frac{(-4)}{1} \times 2 = 9$$

9. Let $\vec{c} = 2\hat{i} + 3\hat{j} + 4\hat{k}$

$$\therefore \vec{a} \times \vec{c} = \vec{c} \times \vec{b}$$

$$\Rightarrow \vec{a} \times \vec{c} = -\vec{b} \times \vec{c} \Rightarrow (\vec{a} + \vec{b}) \times \vec{c} = \vec{0} \Rightarrow (\vec{a} + \vec{b}) \parallel \vec{c}$$

$$\text{Let } (\vec{a} + \vec{b}) = \lambda \vec{c}$$

$$\Rightarrow |\vec{a} + \vec{b}| = |\lambda| |\vec{c}| \Rightarrow \sqrt{29} = |\lambda| \cdot \sqrt{29} \Rightarrow \lambda = \pm 1$$

$$\therefore \vec{a} + \vec{b} = \pm(2\hat{i} + 3\hat{j} + 4\hat{k})$$

$$\text{Now, } (\vec{a} + \vec{b}) \cdot (-7\hat{i} + 2\hat{j} + 3\hat{k}) = \pm(-14 + 6 + 12) = \pm 4$$

10. Given,

$$[\vec{a} + m\vec{b} + n\vec{c} \quad \vec{b} + m\vec{c} + n\vec{a} \quad \vec{c} + m\vec{a} + n\vec{b}] = 0$$

$$\Rightarrow [\vec{a} + m\vec{b} + n\vec{c} \quad n\vec{a} + \vec{b} + m\vec{c} \quad m\vec{a} + n\vec{b} + \vec{c}] = 0$$

$$\Rightarrow \begin{vmatrix} m & n \\ n & m \end{vmatrix} [\vec{a} \vec{b} \vec{c}] = 0$$


$$\Rightarrow \begin{vmatrix} m & n \\ n & m \end{vmatrix} = 0 \quad \dots [\because [\vec{a} \vec{b} \vec{c}] \neq 0]$$

$$\Rightarrow m^2 + n^2 - 2mn = 0$$

$$\Rightarrow (m + n)^2 - (m - n)^2 = 0 \Rightarrow m + n = 0$$

$$\Rightarrow m + n = 0$$

11.



Let point O be the circumcentre of ΔABC .

Let $\vec{a}, \vec{b}, \vec{c}, \vec{p}, \vec{d}, \vec{h}, \vec{m}$ be the position vectors of the respective points.

Since, $\vec{h} = \vec{a} + \vec{b} + \vec{c} \quad \dots (\text{Standard formula})$

$$\therefore \vec{m} = \frac{\vec{p} + \vec{h}}{2} = \frac{\vec{p} + \vec{a} + \vec{b} + \vec{c}}{2}$$

$$\therefore \vec{DM} = \vec{m} - \vec{d} = \frac{\vec{p} + \vec{a} + \vec{b} + \vec{c}}{2} - \frac{\vec{b} + \vec{c}}{2} = \frac{\vec{p} + \vec{a}}{2}$$

$$\therefore \vec{DM} \cdot \vec{PA} = \left(\frac{\vec{p} + \vec{a}}{2} \right) \cdot (\vec{a} - \vec{p}) = \frac{1}{2} (a^2 - p^2) = 0$$

$\dots [\because O \text{ is circumcentre, } \therefore OA = OP \text{ i.e., } a = p]$

$\therefore DM$ is perpendicular to PA .



15. Let position vector of Q be \bar{r}
 Since, \bar{p} divides PQ in the ratio 3 : 4


$$\therefore \frac{3\bar{r} + 4(3\bar{p} + \bar{q})}{3+4} = \bar{p}$$

$$\Rightarrow 7\bar{p} = 3\bar{r} + 12\bar{p} + 4\bar{q}$$

$$\Rightarrow -5\bar{p} - 4\bar{q} = 3\bar{r}$$

$$\Rightarrow \bar{r} = \frac{-1}{3}(5\bar{p} + 4\bar{q})$$

16.



By distance formula,

$$AB = \sqrt{(5-3)^2 + (3-2)^2 + (2-0)^2}$$

$$= \sqrt{4+1+4} = \sqrt{9} = 3$$

$$AC = \sqrt{(3+9)^2 + (2-6)^2 + (0+3)^2}$$

$$= \sqrt{144+16+9} = \sqrt{169} = 13$$

\therefore Point D divides seg BC in the ratio of 3 : 13

\therefore By section formula,


$$D \equiv \left(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n}, \frac{mz_2 + nz_1}{m+n} \right)$$

$$\equiv \left(\frac{3(-9) + 13(5)}{3+13}, \frac{3(6) + 13(3)}{3+13}, \frac{3(-3) + 13(2)}{3+13} \right)$$

$$\equiv \left(\frac{-27 + 65}{16}, \frac{18 + 39}{16}, \frac{-9 + 26}{16} \right)$$

$$\equiv \left(\frac{38}{16}, \frac{57}{16}, \frac{17}{16} \right) \equiv \left(\frac{19}{8}, \frac{57}{16}, \frac{17}{16} \right)$$

17.



$$x_1 + x_2 = 2, x_2 + x_3 = 0, x_3 + x_1 = 0$$

On solving we get $x_1 = \frac{2}{3}, x_2 = -\frac{2}{3}, x_3 = -\frac{2}{3}$

$$y_1 + y_2 = 0, y_2 + y_3 = 2m, y_3 + y_1 = 0$$

On solving we get $y_1 = -m, y_2 = m, y_3 = m$

$$z_1 + z_2 = 0, z_2 + z_3 = 0, z_3 + z_1 = 2n$$

On solving we get $z_1 = n, z_3 = n, z_2 = -n$

$$\therefore A\left(\frac{2}{3}, -m, n\right), B\left(-\frac{2}{3}, m, -n\right), C\left(-\frac{2}{3}, m, n\right)$$

By distance formula,

$$AB^2 = \left(\frac{2}{3} + \frac{2}{3}\right)^2 + (-m - m)^2 + (n + n)^2 = 4m^2 + 4n^2$$

$$BC^2 = \left(\frac{2}{3} + \frac{2}{3}\right)^2 + (m - m)^2 + (-n - n)^2 = 4\left(\frac{2}{3}\right)^2 + 4n^2$$


$$CA^2 = \left(\frac{2}{3} + \frac{2}{3}\right)^2 + (-m - m)^2 + (n - n)^2 = 4\left(\frac{2}{3}\right)^2 + 4m^2$$

$$\therefore \frac{AB^2 + BC^2 + CA^2}{2 + m^2 + n^2}$$

$$= \frac{4m^2 + 4n^2 + 4\left(\frac{2}{3}\right)^2 + 4n^2 + 4\left(\frac{2}{3}\right)^2 + 4m^2}{2 + m^2 + n^2}$$

$$= 8 \frac{\left(\frac{2}{3} + m^2 + n^2\right)}{2 + m^2 + n^2} = 8$$

18.



Let D be the foot of perpendicular and let it divide BC in the ratio $\lambda : 1$ internally

$$\therefore D \equiv \left(\frac{3\lambda + 4}{\lambda + 1}, \frac{5\lambda + 7}{\lambda + 1}, \frac{3\lambda + 1}{\lambda + 1} \right)$$

$$\overline{AD} = \bar{d} - \bar{a}$$

$$= \left(\frac{3\lambda + 4}{\lambda + 1} - 1 \right) \hat{i} + \left(\frac{5\lambda + 7}{\lambda + 1} - 0 \right) \hat{j} + \left(\frac{3\lambda + 1}{\lambda + 1} - 3 \right) \hat{k} - \hat{i} - 3\hat{k}$$

$$= \left(\frac{2\lambda + 3}{\lambda + 1} \right) \hat{i} + \left(\frac{5\lambda + 7}{\lambda + 1} \right) \hat{j} - \left(\frac{2}{\lambda + 1} \right) \hat{k}$$

$$\overline{BC} = 3\hat{i} + 5\hat{j} + 3\hat{k} - (4\hat{i} + 7\hat{j} + \hat{k})$$

$$= -\hat{i} - 2\hat{j} + 2\hat{k}$$

Since, $\overline{AD} \perp \overline{BC}$.

$$\overline{AD} \cdot \overline{BC} = 0$$

$$\Rightarrow \left(\frac{2\lambda + 3}{\lambda + 1} \right)(-1) + \left(\frac{5\lambda + 7}{\lambda + 1} \right)(-2) + \left(\frac{-2}{\lambda + 1} \right)(2) = 0$$

$$\Rightarrow -2\lambda - 3 - 10\lambda - 14 - 4 = 0$$

$$\Rightarrow -12\lambda - 21 = 0 \Rightarrow \lambda = -\frac{7}{4}$$

$$\therefore D \equiv \left(\frac{3\left(-\frac{7}{4}\right) + 4}{-\frac{7}{4} + 1}, \frac{5\left(-\frac{7}{4}\right) + 7}{-\frac{7}{4} + 1}, \frac{3\left(-\frac{7}{4}\right) + 1}{-\frac{7}{4} + 1} \right)$$

$$\equiv \left(\frac{-21 + 16}{-7 + 4}, \frac{-35 + 28}{-7 + 4}, \frac{-21 + 4}{-7 + 4} \right) \equiv \left(\frac{5}{3}, \frac{7}{3}, \frac{17}{3} \right)$$

06 Three Dimensional Geometry



Hints



Classical Thinking

- For every point (x, y, z) on X-axis $y = 0, z = 0$
- Let the direction cosines of the line be l, m, n
 $\therefore l = \cos 45^\circ, m = \cos 60^\circ, n = \cos 60^\circ$
 $\Rightarrow l = \frac{1}{\sqrt{2}}, m = \frac{1}{2}$ and $n = \frac{1}{2}$
 \therefore d.c.s are $\frac{1}{\sqrt{2}}, \frac{1}{2}, \frac{1}{2}$.
- Let the d.c.s of the line be l, m, n
 $\therefore l = \cos 90^\circ, m = \cos 60^\circ, n = \cos 30^\circ$
 $\Rightarrow l = 0, m = \frac{1}{2}, n = \frac{\sqrt{3}}{2}$
 \therefore d.c.s are $0, \frac{1}{2}, \frac{\sqrt{3}}{2}$
- The d.c.s of Y-axis are $\cos 90^\circ, \cos 0^\circ, \cos 90^\circ$
 i.e. $0, 1, 0$
- The d.c.s of X-axis are $1, 0, 0$.
- For option (B),
 $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma \neq 1$
 \therefore option (B) is correct answer.
- Since, $l^2 + m^2 + n^2 = 1$
 $\therefore k^2 + \left(\frac{1}{2}\right)^2 + 0^2 = 1$
 $\Rightarrow k^2 = 1 - \frac{1}{4} = \frac{3}{4}$
 $\Rightarrow k = \pm \frac{\sqrt{3}}{2}$
- Since, $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$
 $\therefore \cos^2 45^\circ + \cos^2 60^\circ + \cos^2 \gamma = 1$
 $\Rightarrow \cos^2 \gamma = 1 - \frac{1}{2} - \frac{1}{4} = \frac{1}{4}$
 $\Rightarrow \cos \gamma = \pm \frac{1}{2}$
 $\Rightarrow \gamma = 60^\circ$ or 120°

- $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$
 $\Rightarrow \cos^2 \alpha + \cos^2(90^\circ - \alpha) + \cos^2 \gamma = 1$
 $\dots[\because \alpha + \beta = 90^\circ]$
 $\Rightarrow \cos^2 \alpha + \sin^2 \alpha + \cos^2 \gamma = 1$
 $\Rightarrow \cos^2 \gamma + 1 = 1$
 $\Rightarrow \cos^2 \gamma = 0$
 $\Rightarrow \gamma = 90^\circ$
- Let l, m, n be the d.c.s of the line.
 $\therefore l = \cos \alpha; m = \cos 60^\circ; n = \cos 45^\circ$
 Since, $\cos^2 \alpha + \cos^2 60^\circ + \cos^2 45^\circ = 1$
 $\Rightarrow \cos^2 \alpha = 1 - \frac{1}{2} - \frac{1}{4} = \frac{1}{4}$
 $\Rightarrow \cos \alpha = \pm \frac{1}{2}$
 \therefore the d.c.s are $\pm \frac{1}{2}, \frac{1}{2}, \frac{1}{\sqrt{2}}$
- Let $\vec{r} = 2\hat{i} + 2\hat{j} - \hat{k}$
 $|\vec{r}| = \sqrt{2^2 + 2^2 + (-1)^2} = 3$
 \therefore The d.c.s are $\frac{x}{|\vec{r}|}, \frac{y}{|\vec{r}|}, \frac{z}{|\vec{r}|}$
 i.e., $\frac{2}{3}, \frac{2}{3}, \frac{-1}{3}$
- Let $\vec{r} = 3\hat{i} + 4\hat{k}$.
 $|\vec{r}| = \sqrt{3^2 + 0^2 + 4^2} = 5$
 \therefore The d.c.s are $\frac{3}{5}, 0, \frac{4}{5}$
- D.c.s are $\frac{a}{|\vec{r}|}, \frac{b}{|\vec{r}|}, \frac{c}{|\vec{r}|}$
 i.e., $\frac{2}{7}, \frac{-3}{7}, \frac{6}{7}$
- $A \equiv (1, 2, 6)$ and $B \equiv (-4, 5, 0)$
 \therefore D.r.s of AB are $-4 - 1, 5 - 2, 0 - 6$
 i.e., $-5, 3, -6$
- On Y-axis, x and z co-ordinates are zero.
 Hence, (B) is the correct option.



18. Since $(-)^2 + (-m)^2 + (-n)^2 = 1$, we can say that $-$, $-m$, $-n$ are the direction cosines of the line.

$$\text{Also that } \frac{-}{m} = \frac{-m}{m} = \frac{-n}{n} = -1$$

Hence, we can say that $-$, $-m$, $-n$ are the d.r.s. of the line.

19. Let a , b , c be the d.r.s of the line.

\therefore The d.c.s are given by

$$\frac{a}{\sqrt{a^2 + b^2 + c^2}}, \frac{b}{\sqrt{a^2 + b^2 + c^2}}, \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

$$\text{i.e., } \frac{2}{\sqrt{2^2 + (-1)^2 + (-2)^2}}, \frac{-1}{\sqrt{2^2 + (-1)^2 + (-2)^2}},$$

$$\frac{-2}{\sqrt{2^2 + (-1)^2 + (-2)^2}}$$

$$\text{i.e., } \frac{2}{3}, \frac{-1}{3}, \frac{-2}{3}$$

20. The direction cosines are

$$\frac{\sqrt{2}}{\sqrt{2+5+2}}, \frac{-\sqrt{5}}{\sqrt{2+5+2}}, \frac{\sqrt{2}}{\sqrt{2+5+2}}$$

$$\text{i.e., } \frac{\sqrt{2}}{3}, \frac{-\sqrt{5}}{3}, \frac{\sqrt{2}}{3}$$

21. The d.r.s of line through $(1, 2, -3)$ and $(-2, 3, 1)$ are $-2-1, 3-2, 1-(-3)$

$$\text{i.e. } -3, 1, 4$$

\therefore d.c.s are

$$\frac{-3}{\sqrt{9+1+16}}, \frac{1}{\sqrt{9+1+16}}, \frac{4}{\sqrt{9+1+16}}$$

$$\text{i.e. } \frac{-3}{\sqrt{26}}, \frac{1}{\sqrt{26}}, \frac{4}{\sqrt{26}}$$

22. The d.r.s of AB are $2-14, -3-5, 1+3$

$$\text{i.e. } -12, -8, 4 \quad \text{i.e., } 3, 2, -1$$

\therefore The d.c.s are $\frac{3}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{-1}{\sqrt{14}}$

23. Let $O(0, 0, 0)$ and $P(1, 2, 3)$ be two points.

\therefore Then the d.r.s of OP are $1, 2, 3$

\therefore The d.c.s of OP are

$$\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}}$$

24. D.r.s. of line through $A(3, 1, 2)$, $B(4, \lambda, 0)$ are $4-3, \lambda-1, 0-2 \Rightarrow 1, \lambda-1, -2 \equiv a_1, b_1, c_1$

D.r.s. of line through $C(1, 2, 1)$, $D(2, 3, -1)$ are $2-1, 3-2, -1-1 \Rightarrow 1, 1, -2 \equiv a_2, b_2, c_2$

Since, $AB \parallel CD$,

$$\therefore \frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$$

$$\Rightarrow \frac{1}{1} = \frac{\lambda-1}{1} = \frac{-2}{-2}$$

$$\text{i.e., } \lambda-1=1 \Rightarrow \lambda=2$$

25. Let $A \equiv (5, 2, 4)$, $B \equiv (6, -1, 2)$ and $C \equiv (8, -7, k)$

\therefore The d.r.s of AB are $6-5, -1-2, 2-4$

$$\text{i.e., } 1, -3, -2, \text{ and}$$

The d.r.s of BC are $8-6, -7+1, k-2$

$$\text{i.e. } 2, -6, k-2$$

Since, the points A, B, C are collinear, $AB \parallel BC$

$$\therefore \frac{2}{1} = \frac{-6}{-3} = \frac{k-2}{-2}$$

$$\Rightarrow k-2 = -4$$

$$\Rightarrow k = 2-4 = -2$$

26. Let $A \equiv (-2, 4, \lambda)$, $B \equiv (3, -6, -8)$, $C \equiv (1, -2, -2)$

The d.r.s of AB are

$$-5, 10, \lambda+8, \text{ and}$$

The d.r.s of AC are

$$-3, 6, \lambda+2$$

Since, the points A, B, C are collinear, $AB \parallel AC$

$$\therefore \frac{-5}{-3} = \frac{10}{6} = \frac{\lambda+8}{\lambda+2}$$

$$\therefore 5(\lambda+2) = 3(\lambda+8)$$

$$\Rightarrow 5\lambda + 10 = 3\lambda + 24$$

$$\Rightarrow 2\lambda = 14$$

$$\Rightarrow \lambda = 7$$

27. Let, $n_1 = \frac{1}{\sqrt{6}}, m_1 = \frac{-1}{\sqrt{6}}, n_1 = \frac{2}{\sqrt{6}}$

$$\text{and } n_2 = \frac{2}{\sqrt{6}}, m_2 = \frac{1}{\sqrt{6}}, n_2 = \frac{-1}{\sqrt{6}}$$

\therefore angle between the lines is

$$\cos \theta = |n_1 m_2 + m_1 n_2 + n_1 n_2|$$

$$\Rightarrow \cos \theta = \left| \frac{1}{\sqrt{6}} \left(\frac{2}{\sqrt{6}} \right) + \left(\frac{-1}{\sqrt{6}} \right) \left(\frac{1}{\sqrt{6}} \right) + \left(\frac{2}{\sqrt{6}} \right) \left(\frac{-1}{\sqrt{6}} \right) \right|$$

$$\Rightarrow \cos \theta = \left| \frac{-1}{6} \right|$$

$$\therefore \theta = \cos^{-1} \left(\frac{1}{6} \right)$$



28. Let, $a_1, b_1, c_1 = 5, -12, 13$
and $a_2, b_2, c_2 = -3, 4, 5$

$$\begin{aligned}\therefore \cos\theta &= \left| \frac{a_1 a_2 + b_1 b_2 + c_1 c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \cdot \sqrt{a_2^2 + b_2^2 + c_2^2}} \right| \\ &= \left| \frac{5(-3) + (-12)4 + 13(5)}{\sqrt{5^2 + (-12)^2 + 13^2} \cdot \sqrt{(-3)^2 + 4^2 + 5^2}} \right| \\ &= \left| \frac{-15 - 48 + 65}{13\sqrt{2} \cdot 5\sqrt{2}} \right| \\ &= \frac{1}{65}\end{aligned}$$

$$\therefore \theta = \cos^{-1} \left(\frac{1}{65} \right)$$

29. Here, $A \equiv (1, 2, 3)$, $B \equiv (4, 5, 7)$,
 $C \equiv (-4, 3, -6)$ and $D \equiv (2, 9, 2)$

\therefore d.r.s of lines AB and CD are 3, 3, 4 and 6, 6, 8 respectively.

$$\begin{aligned}\therefore \theta &= \cos^{-1} \left[\frac{(3)(6) + (3)(6) + (4)(8)}{\sqrt{34} \cdot \sqrt{136}} \right] \\ &= \cos^{-1} \left[\frac{68}{2 \times 34} \right] = 0^\circ\end{aligned}$$

$$\begin{aligned}30. \cos 45^\circ &= \left| \frac{2a - 3 + 10}{\sqrt{2^2 + (-1)^2 + 2^2} \sqrt{a^2 + 3^2 + 5^2}} \right| \\ \Rightarrow \frac{1}{\sqrt{2}} &= \left| \frac{2a + 7}{3\sqrt{a^2 + 34}} \right| \\ \Rightarrow 9(a^2 + 34) &= 2(2a + 7)^2 \\ \Rightarrow 9a^2 + 306 &= 8a^2 + 56a + 98 \\ \Rightarrow a^2 - 56a + 208 &= 0 \\ \Rightarrow a &= 4\end{aligned}$$

31. Let $a_1, b_1, c_1 = 1, -2, 1$ and
 $a_2, b_2, c_2 = 2, 3, 4$

Consider,

$$a_1 a_2 + b_1 b_2 + c_1 c_2 = 1(2) + (-2)(3) + 1(4) = 0$$

$\therefore OP \perp OQ$.



Critical Thinking

1. If α, β, λ are direction angles of any vector \overline{OL} , then those of $\overline{OL'}$ are $\pi - \alpha, \pi - \beta, \pi - \gamma$ respectively
 \therefore correct answer is option (B).

2. We know that, $l^2 + m^2 + n^2 = 1$

Consider option (D)

$$\begin{aligned}\left(\frac{2}{\sqrt{25}} \right)^2 + \left(\frac{3}{\sqrt{25}} \right)^2 + \left(\frac{4}{\sqrt{25}} \right)^2 &= \frac{4+9+16}{25} \\ &= \frac{29}{25} \neq 1\end{aligned}$$

\therefore correct answer is option (D).

3. Consider option (B)

$$\begin{aligned}\therefore \cos^2 \frac{\pi}{4} + \cos^2 \frac{\pi}{3} + \cos^2 \frac{\pi}{3} \\ = \frac{1}{2} + \frac{1}{4} + \frac{1}{4} = 1\end{aligned}$$

\therefore correct answer is option (B).

4. $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$

$$\therefore \cos \gamma = \pm \sqrt{1 - \left(\frac{14}{15} \right)^2 - \left(\frac{1}{3} \right)^2} = \pm \sqrt{\frac{8}{9} - \left(\frac{196}{225} \right)} = \pm \frac{2}{15}$$

5. Since, $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$

$$\therefore \cos^2 \alpha + \cos^2 60^\circ + \cos^2 60^\circ = 1$$

$$\therefore \cos^2 \alpha = 1 - \frac{1}{4} - \frac{1}{4} = 1 - \frac{1}{2} = \frac{1}{2}$$

$$\Rightarrow \cos \alpha = \pm \frac{1}{\sqrt{2}}$$

$$\Rightarrow \alpha = 45^\circ \text{ or } \alpha = 135^\circ$$

6. Since, the line lies in ZOY plane, it makes an angle 90° with X-axis

Also, line makes angle 30° and $\pi - 30^\circ$ with positive Z-axis and 60° and $\pi - 60^\circ$ with positive Y-axis

\therefore d.c.s of the required line are

$$\pm \cos \alpha, \pm \cos \beta, \pm \cos \gamma$$

$$\text{i.e., } \pm \cos 60^\circ, \pm \cos 0^\circ, \pm \cos 30^\circ$$

$$\text{i.e. } \pm \frac{1}{2}, 0, \pm \frac{\sqrt{3}}{2}$$

7. $\cos \gamma = \sqrt{1 - \frac{3}{4} - \frac{1}{2}} = \sqrt{\frac{-1}{4}}$ which is not possible.

8. Let l, m, n be the d.c.s of \vec{r} .

$$l = m = n$$

$$\dots [\because \alpha = \beta = \gamma \Rightarrow \cos \alpha = \cos \beta = \cos \gamma]$$

$$\text{Now, } l^2 + m^2 + n^2 = 1$$

$$\Rightarrow l = \pm \frac{1}{\sqrt{3}}$$



9. Since, $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$
 $\Rightarrow \cos^2 \alpha + \cos^2 \alpha + \cos^2 \alpha = 1$ ($\because \alpha = \beta = \gamma$)

$$\Rightarrow \cos^2 \alpha = \frac{1}{3} \Rightarrow \cos \alpha = \frac{1}{\sqrt{3}}$$

Now, sum of d.c.s. = $+m+n$
 $= \cos \alpha + \cos \alpha + \cos \alpha$
 $= 3 \cos \alpha = \sqrt{3}$

10. $\cos 2\alpha + \cos 2\beta + \cos 2\gamma$
 $= 2\cos^2 \alpha - 1 + 2\cos^2 \beta - 1 + 2\cos^2 \gamma - 1$
 $= 2(\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma) - 3$
 $= 2(1) - 3 = -1$

11. $\sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma$
 $= (1 - \cos^2 \alpha) + (1 - \cos^2 \beta) + (1 - \cos^2 \gamma)$
 $= 3 - (\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma) = 3 - (1) = 2$

12. Let $\alpha = \frac{\pi}{6}$ and $\beta = \frac{\pi}{4}$

$$\therefore \cos \alpha = \frac{\sqrt{3}}{2} \text{ and } \cos \beta = \frac{1}{\sqrt{2}}$$

Since, $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$

$$\therefore \frac{3}{4} + \frac{1}{2} + \cos^2 \gamma = 1$$

$$\Rightarrow \cos^2 \gamma = -\frac{1}{4}$$

Square of a real number cannot be negative.

\therefore option (A) is the correct answer.

13. The line makes angle θ with X-axis and Z-axis and β with Y-axis.

$$\therefore \cos \theta, m = \cos \beta, n = \cos \theta$$

$$\cos^2 \theta + \cos^2 \beta + \cos^2 \theta = 1$$

$$\Rightarrow 2\cos^2 \theta = 1 - \cos^2 \beta$$

$$\Rightarrow 2 \cos^2 \theta = \sin^2 \beta \quad \dots(\text{i})$$

$$\text{But } \sin^2 \beta = 3\sin^2 \theta \quad \dots(\text{ii})$$

From (i) and (ii), we get

$$3\sin^2 \theta = 2\cos^2 \theta$$

$$\Rightarrow 3(1 - \cos^2 \theta) = 2\cos^2 \theta$$

$$\Rightarrow 3 = 5\cos^2 \theta \Rightarrow \cos^2 \theta = \frac{3}{5}$$

14. Let the length of the line segment be r and its d.c.s be l, m, n .

\therefore The projections on the co-ordinate axes are lr, mr, nr .

$$\therefore r = 4, mr = 6 \text{ and } nr = 12$$

$$\therefore l^2 r^2 + m^2 r^2 + n^2 r^2 = (4)^2 + (6)^2 + (12)^2$$

$$\Rightarrow r^2 (l^2 + m^2 + n^2) = 16 + 36 + 144$$

$$\Rightarrow r^2 = 196 \quad \dots[\because l^2 + m^2 + n^2 = 1]$$

$$\Rightarrow r = 14$$

$$\therefore \text{The d.c.s. of line are } \frac{4}{r}, \frac{6}{r}, \frac{12}{r}$$

$$\text{i.e., } \frac{2}{7}, \frac{3}{7}, \frac{6}{7}$$

15. Let α, β, γ be the angles which OP makes with the co-ordinates axes,

$$\therefore x = r \cos \alpha, y = r \cos \beta, z = r \cos \gamma$$

$$\therefore \cos \alpha = \frac{x}{r}; \cos \beta = \frac{y}{r}; \cos \gamma = \frac{z}{r}$$

So, the direction cosines are $\frac{x}{r}, \frac{y}{r}, \frac{z}{r}$.

16. We have $\cos 45^\circ = \frac{1}{\sqrt{2}}$,

$$m = \cos 60^\circ = \frac{1}{2} \text{ and } n = \cos \gamma$$

We know that $l^2 + m^2 + n^2 = 1$

$$\therefore \frac{1}{2} + \frac{1}{4} + n^2 = 1$$

$$\Rightarrow n^2 = 1 - \frac{3}{4} = \frac{1}{4} \Rightarrow n = \pm \frac{1}{2}$$

$$\Rightarrow \cos \gamma = \pm \frac{1}{2}$$

$$\vec{r} = r (\hat{i} + m\hat{j} + n\hat{k})$$

$$\Rightarrow \vec{r} = 12 \left(\frac{1}{\sqrt{2}} \hat{i} + \frac{1}{2} \hat{j} \pm \frac{1}{2} \hat{k} \right)$$

17. Since, $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$

$$\Rightarrow \cos^2 \alpha + \cos^2 \alpha + \cos^2 \alpha = 1$$
 ($\because \alpha = \beta = \gamma$)

$$\Rightarrow \cos^2 \alpha = \frac{1}{3} \Rightarrow \cos \alpha = \frac{1}{\sqrt{3}}$$

$$\therefore \text{The d.c.s are } \pm \frac{1}{\sqrt{3}}, \pm \frac{1}{\sqrt{3}}, \pm \frac{1}{\sqrt{3}}.$$

The magnitude of the given vector is 6.

$$\therefore \vec{r} = 6(\cos \alpha \hat{i} + \cos \beta \hat{j} + \cos \gamma \hat{k})$$

$$= \frac{\pm 6}{\sqrt{3}} (\hat{i} + \hat{j} + \hat{k}) = \pm 2\sqrt{3} (\hat{i} + \hat{j} + \hat{k})$$

18. For a line passing through origin, d.r.s are the co-ordinates of the point.

19. D.c.s. of the line are $\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}$

$$\therefore \cos \alpha = \frac{1}{\sqrt{3}}, \cos \beta = \frac{1}{\sqrt{3}}, \cos \gamma = \frac{1}{\sqrt{3}}$$

Hence, line is equally inclined to axes.



$$20. \quad \cos \gamma = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

$$= \frac{5}{\sqrt{9+16+25}} = \frac{5}{5\sqrt{2}} = \frac{1}{\sqrt{2}}$$

$$\Rightarrow \gamma = \frac{\pi}{4}$$

21. The d.r.s. of the given line are
 $2 - 6, -3 + 7, 1 + 1$
 i.e., $-2, 2, 1$. i.e., $2, -2, -1$

\therefore angle α is acute, $\cos \alpha > 0$

$$\Rightarrow \cos \alpha = \frac{2}{3}$$

Thus, required d.c.s are $\frac{2}{3}, \frac{-2}{3}, \frac{-1}{3}$

$$22. \quad x^2 + m^2 + n^2 = 1$$

$$\therefore \left(\frac{2}{7}\right)^2 + \left(\frac{-3}{7}\right)^2 + n^2 = 1$$

$$\therefore n^2 = 1 - \frac{13}{49} = \frac{36}{49}$$

Let a, b, c be the d.r.s. of the line.

$$\therefore a = 2, b = -3, c = z$$

$$\text{Since, } n = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

$$\therefore \frac{z}{\sqrt{4+9+z^2}} = \pm \frac{6}{7}$$

$$\Rightarrow \frac{z^2}{13+z^2} = \frac{36}{49}$$

$$\Rightarrow 49z^2 - 36z^2 = 13 \times 36$$

$$\Rightarrow z^2 = 36$$

$$\Rightarrow z = \pm 6$$

23. Let $A \equiv (2, a, -1), B \equiv (3, 4, b)$ and $C \equiv (1, -2, 3)$
 d.r.s of AB are $3 - 2, 4 - a, b - (-1)$

$$\text{i.e., } 1, 4 - a, b + 1$$

d.r.s of BC are $1 - 3, -2 - 4, 3 - b$

$$\text{i.e., } -2, -6, 3 - b$$

Since, the points A, B and C are collinear,
 $AB \parallel BC$

$$\therefore \frac{1}{-2} = \frac{4-a}{-6} = \frac{b+1}{3-b}$$

$$\Rightarrow \frac{4-a}{-6} = \frac{1}{-2}, \frac{b+1}{3-b} = \frac{1}{-2}$$

$$\Rightarrow 4 - a = 3, -2b - 2 = 3 - b$$

$$\Rightarrow a = 1, b = -5$$

24. Let $A \equiv (1, a, 1), B \equiv (3, -1, 2)$ and $C \equiv (1, a^2, 1)$
 d.r.s of AB are $3 - 1, -1 - a, 2 - 1$

$$\text{i.e. } 2, -1 - a, 1$$

d.r.s of BC are $1 - 3, a^2 + 1, 1 - 2$

$$\text{i.e. } -2, a^2 + 1, -1$$

Since, the points are collinear, $AB \parallel BC$

$$\therefore \frac{2}{-2} = \frac{-1-a}{a^2+1} = \frac{1}{-1}$$

$$\Rightarrow \frac{-1-a}{a^2+1} = -1$$

$$\Rightarrow a^2 + 1 = 1 + a$$

$$\Rightarrow a^2 - a = 0$$

$$\Rightarrow a(a - 1) = 0$$

$$\Rightarrow a = 0 \text{ or } a = 1$$

$$25. \quad \text{Here, } \frac{3 - (-2)}{1 - 3} = \frac{-6 - 4}{-2 - (-6)} = \frac{-8 - 7}{-2 - (-8)}$$

$$\Rightarrow -\frac{5}{2} = -\frac{5}{2} = -\frac{5}{2}$$

\therefore the given points are collinear.

26. Let, $a_1, m_1, n_1 = a, \frac{-2}{3}, \frac{1}{3}$ and

$$a_2, m_2, n_2 = \frac{2}{3}, \frac{1}{3}, \frac{-2}{3}$$

$$\cos \theta = \frac{a_1 a_2 + m_1 m_2 + n_1 n_2}{\sqrt{a_1^2 + m_1^2 + n_1^2} \sqrt{a_2^2 + m_2^2 + n_2^2}}$$

$$\therefore \cos \frac{\pi}{2} = \frac{a\left(\frac{2}{3}\right) + \left(\frac{-2}{3}\right)\left(\frac{1}{3}\right) + \frac{1}{3}\left(\frac{-2}{3}\right)}{\sqrt{a^2 + \frac{4}{9} + \frac{1}{9}} \sqrt{\frac{4}{9} + \frac{1}{9} + \frac{4}{9}}}$$

$$\Rightarrow 0 = \frac{2a}{3} - \frac{2}{9} - \frac{2}{9}$$

$$\Rightarrow \frac{2a}{3} = \frac{4}{9}$$

$$\Rightarrow a = \frac{2}{3}$$

27. Let, $a_1, b_1, c_1 = 5, 4, 1$

$$a_2, b_2, c_2 = -3, 2, 1$$

$$\cos \theta = \frac{a_1 a_2 + b_1 b_2 + c_1 c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

$$\therefore \cos \theta = \frac{5(-3) + 4(2) + 1(1)}{\sqrt{5^2 + 4^2 + 1^2} \sqrt{(-3)^2 + (2)^2 + (1)^2}}$$


$$= \frac{-15 + 8 + 1}{\sqrt{42} \sqrt{14}} = \frac{6}{14\sqrt{3}} = \frac{\sqrt{3}}{7}$$

$$\therefore \theta = \cos^{-1} \left(\frac{\sqrt{3}}{7} \right)$$



28. $\theta = \cos^{-1} \left| \frac{1(2) + 2(-3) + 1(4)}{\sqrt{1^2 + 2^2 + 1^2} \sqrt{2^2 + (-3)^2 + 4^2}} \right|$
 $\theta = \cos^{-1} (0) = \frac{\pi}{2}$
29. Given, $A \equiv (1, 2, -1)$, $B \equiv (2, 0, 3)$, $C \equiv (3, -1, 2)$
 The d.r.s of $AB = 1, -2, 4$ and d.r.s of $AC = 2, -3, 3$
 $\therefore \cos \theta = \left| \frac{1(2) + (-2)(-3) + 4(3)}{\sqrt{1+4+16} \sqrt{4+9+9}} \right|$
 $\Rightarrow \cos \theta = \frac{2+6+12}{\sqrt{21}\sqrt{22}} = \frac{20}{\sqrt{462}}$
 $\Rightarrow \sqrt{462} \cos \theta = 20$
30. $+m + n = 0$
 $\Rightarrow -(m+n)$ and $m = 0 \Rightarrow -(m+n)m = 0$
 $\Rightarrow m = 0$ or $m+n = 0 \Rightarrow m = 0$ or $m = -n$
 If $m = 0$, then $= -n$
 $\therefore \frac{-}{-1} = \frac{m}{0} = \frac{n}{1}$
 If $m = -n$, then $= 0$
 $\therefore \frac{-}{0} = \frac{m}{-1} = \frac{n}{1}$
 \therefore the d.r.s of the lines are proportional to $-1, 0, 1$ and $0, -1, 1$
 \therefore angle between them is
 $\cos \theta = \left| \frac{0+0+1}{\sqrt{1+0+1}\sqrt{0+1+1}} \right| = \frac{1}{2}$
 $\therefore \theta = \frac{\pi}{3}$
31. $+m - n = 0$ and $^2 + m^2 - n^2 = 0$
 $\Rightarrow +m = n$ and $^2 + m^2 = n^2$
 Putting $+m = n$ in $^2 + m^2 = n^2$, we get
 $^2 + m^2 = (+m)^2$
 $\Rightarrow 2m = 0 \Rightarrow = 0$ or $m = 0$
 If $= 0$, then $m = n$
 $\therefore \frac{-}{0} = \frac{m}{1} = \frac{n}{1}$
 If $m = 0$, then $= n$
 $\therefore \frac{-}{1} = \frac{m}{0} = \frac{n}{1}$
 \therefore the d.r.s of the lines are proportional to $0, 1, 1$ and $1, 0, 1$.
 $\therefore \cos \theta = \left| \frac{0(1) + 1(0) + 1(1)}{\sqrt{0+1+1}\sqrt{1+0+1}} \right| = \frac{1}{2}$
 $\Rightarrow \theta = \cos^{-1} \left(\frac{1}{2} \right) \Rightarrow \theta = \frac{\pi}{3}$

32. Since, the three lines are mutually perpendicular
 \therefore
 ${}_1{}_2 + m_1m_2 + n_1n_2 = 0$
 ${}_2{}_3 + m_2m_3 + n_2n_3 = 0$
 ${}_3{}_1 + m_3m_1 + n_3n_1 = 0$
 Also, ${}_1^2 + m_1^2 + n_1^2 = 1$,
 ${}_2^2 + m_2^2 + n_2^2 = 1$,
 ${}_3^2 + m_3^2 + n_3^2 = 1$
 Now, $({}_1 + {}_2 + {}_3)^2 + (m_1 + m_2 + m_3)^2 + (n_1 + n_2 + n_3)^2$
 $= ({}_1^2 + m_1^2 + n_1^2) + ({}_2^2 + m_2^2 + n_2^2) + ({}_3^2 + m_3^2 + n_3^2)$
 $+ 2({}_1{}_2 + m_1m_2 + n_1n_2) + 2({}_2{}_3 + m_2m_3 + n_2n_3)$
 $+ 2({}_3{}_1 + m_3m_1 + n_3n_1)$
 $= 3$
 $\Rightarrow ({}_1 + {}_2 + {}_3)^2 + (m_1 + m_2 + m_3)^2 + (n_1 + n_2 + n_3)^2 = 3$
 Hence, direction cosines of required line are :
 $\frac{{}_1 + {}_2 + {}_3}{\sqrt{3}}, \frac{m_1 + m_2 + m_3}{\sqrt{3}}, \frac{n_1 + n_2 + n_3}{\sqrt{3}}$



- The d.r.s of diagonal $EB = a, a, -a$
 The d.r.s of diagonal $AD = -a, a, a$
 \therefore Angle between EB and AD is
 $\cos \theta = \left| \frac{-a^2 + a^2 - a^2}{3a^2} \right|$

$\therefore \theta = \cos^{-1} \left(\frac{1}{3} \right)$

34. As d.r.s are proportional, the required lines are parallel to the given lines.
 \therefore (d.r.s.)₁ $\equiv 2, 3, -6$ and (d.r.s.)₂ $\equiv 3, -4, 5$
 $\therefore \cos \theta = \left| \frac{6-12-30}{\sqrt{49}\sqrt{50}} \right| = \frac{36}{7(5\sqrt{2})} = \frac{18\sqrt{2}}{7(5)}$
 $\therefore \theta = \cos^{-1} \left(\frac{18\sqrt{2}}{35} \right)$



35. Let $A \equiv (-2, 1, -8)$ and $B \equiv (a, b, c)$
 \therefore the d.r.s of the line AB are $a + 2, b - 1, c + 8$
 Since, AB is parallel to the line whose d.r.s are 6, 2, 3.

$$\therefore \frac{a+2}{6} = \frac{b-1}{2} = \frac{c+8}{3}$$

Only option (A) satisfies this condition.

36. The d.r.s of AB and CD are 1, 2, -2 and 2, 3, 4 respectively

$$\text{Now, } a_1a_2 + b_1b_2 + c_1c_2 = 1(2) + 2(3) + (-2)(4) = 0$$

- $\therefore AB \perp CD,$
 \therefore projection of AB on CD is 0.

37. As $\frac{a}{\left(\frac{1}{bc}\right)} = \frac{b}{\left(\frac{1}{ca}\right)} = \frac{c}{\left(\frac{1}{ab}\right)}$

the lines are parallel.

38.
$$\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & m_1 & n_1 \\ 2 & m_2 & n_2 \end{vmatrix}$$

$$= \hat{i}(m_1n_2 - m_2n_1) + \hat{j}(n_1 \cdot 2 - 1n_2) + \hat{k}(1m_2 - m_1 \cdot 2)$$

- \therefore The d.c.s are $m_1n_2 - m_2n_1, n_1 \cdot 2 - n_2 \cdot 1, 1m_2 - 2m_1$

39.
$$\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & -3 \\ 1 & -2 & 1 \end{vmatrix} = -5\hat{i} - 5\hat{j} - 5\hat{k}$$

- \therefore d.r.s of line are -5, -5, -5
 i.e. -1, -1, -1

\therefore the d.c.s are $\frac{-1}{\sqrt{3}}, \frac{-1}{\sqrt{3}}, \frac{-1}{\sqrt{3}}$

40. The vectors $4\hat{i} + \hat{j} + 3\hat{k}$ and $2\hat{i} - 3\hat{j} + \hat{k}$ will lie along the given lines.

The vector perpendicular to these vectors is given by

$$(4\hat{i} + \hat{j} + 3\hat{k}) \times (2\hat{i} - 3\hat{j} + \hat{k})$$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 4 & 1 & 3 \\ 2 & -3 & 1 \end{vmatrix} = 10\hat{i} + 2\hat{j} - 14\hat{k}$$

- \therefore The d.r.s of required line are 10, 2, -14.

\therefore The d.c.s are $\frac{1}{\sqrt{3}}, \frac{1}{5\sqrt{3}}, \frac{-7}{5\sqrt{3}}$



Competitive Thinking

2. $\cos 2\alpha + \cos 2\beta + \cos 2\gamma$
 $= 2 \cos^2 \alpha - 1 + 2 \cos^2 \beta - 1 + 2 \cos^2 \gamma - 1$
 $= 2(\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma) - 3 = 2(1) - 3 = -1$

3. $^2 + m^2 + n^2 = 1$
 $\therefore \left(\frac{1}{2}\right)^2 + \left(\frac{1}{3}\right)^2 + n^2 = 1$
 $\Rightarrow n^2 = \frac{23}{36} \Rightarrow n = \pm \frac{\sqrt{23}}{6}$

4. $^2 + m^2 + n^2 = 1$
 $\therefore \frac{1}{c^2} + \frac{1}{c^2} + \frac{1}{c^2} = 1$
 $\Rightarrow c^2 = 3 \Rightarrow c = \pm \sqrt{3}$

5. Since, $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$
 $\therefore \cos^2 45^\circ + \cos^2 120^\circ + \cos^2 \theta = 1$
 $\Rightarrow \cos^2 \theta = 1 - \frac{1}{2} - \frac{1}{4} = \frac{1}{4} \Rightarrow \cos \theta = \pm \frac{1}{2}$

Since, θ is an acute angle.

$\therefore \cos \theta = \frac{1}{2} \Rightarrow \theta = 60^\circ$

6. Since, $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$
 $\therefore \cos^2 120^\circ + \cos^2 \beta + \cos^2 60^\circ = 1$
 $\Rightarrow \left(\frac{-1}{2}\right)^2 + \cos^2 \beta + \left(\frac{1}{2}\right)^2 = 1$
 $\Rightarrow \cos^2 \beta = 1 - \frac{1}{4} - \frac{1}{4}$
 $\Rightarrow \cos^2 \beta = \frac{1}{2}$

$\Rightarrow \cos \beta = \pm \frac{1}{\sqrt{2}} \Rightarrow \beta = 45^\circ \text{ or } 135^\circ$

7. Since, $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$
 $\therefore \cos^2 \frac{\pi}{4} + \cos^2 \frac{\pi}{4} + \cos^2 \gamma = 1$

$\Rightarrow \cos^2 \gamma = 1 - \frac{1}{2} - \frac{1}{2} = 0$

$\Rightarrow \cos \gamma = 0 \Rightarrow \gamma = \frac{\pi}{2}$

8. Since, $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$
 $\therefore \cos^2 \left(\frac{\pi}{4}\right) + \cos^2 \left(\frac{5\pi}{4}\right) + \cos^2 \gamma = 1$

$\Rightarrow \cos^2 \gamma = 1 - \frac{1}{2} - \frac{1}{2}$

$\Rightarrow \cos^2 \gamma = 0 \Rightarrow \gamma = \frac{\pi}{2}$



9. $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$
 $\Rightarrow \cos^2 45^\circ + \cos^2 \beta + \cos^2 \beta = 1 \dots (\because \beta = \gamma)$
 $\Rightarrow 2\cos^2 \beta = 1 - \frac{1}{2} = \frac{1}{2}$
 $\Rightarrow \cos^2 \beta = \frac{1}{4}$
 $\therefore \beta = 60^\circ = \gamma$
 $\Rightarrow \alpha + \beta + \gamma = 165^\circ$

10. Since, $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$
 $\therefore \cos^2 \alpha + \cos^2 \alpha + \cos^2 \alpha = 1 \dots (\alpha = \beta = \gamma)$
 $\Rightarrow 3 \cos^2 \alpha = 1$
 $\Rightarrow \cos^2 \alpha = \frac{1}{3}$
 $\Rightarrow \cos \alpha = \pm \frac{1}{\sqrt{3}}$

Now, $l = m = n = \cos \alpha$

$\therefore l = m = n = \pm \frac{1}{\sqrt{3}}$

11. Since,
 $\alpha = \beta = \gamma \Rightarrow \cos^2 \alpha + \cos^2 \alpha + \cos^2 \alpha = 1$
 $\Rightarrow \cos \alpha = \left(\pm \frac{1}{\sqrt{3}} \right)$

So, there are four lines whose direction cosines are

$$\left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right), \left(\frac{-1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right), \left(\frac{1}{\sqrt{3}}, \frac{-1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right),$$

$$\left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{-1}{\sqrt{3}} \right).$$

12. Since, the vector is equally inclined to the co-ordinate axes,

$$l = m = n = \pm \frac{1}{\sqrt{3}}$$

13. $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$
 $\Rightarrow \frac{1 + \cos 2\alpha}{2} + \frac{1 + \cos 2\beta}{2} + \frac{1 + \cos 2\gamma}{2} = 1$
 $\Rightarrow \cos 2\alpha + \cos 2\beta + \cos 2\gamma + 3 = 2$
 $\Rightarrow \cos 2\alpha + \cos 2\beta + \cos 2\gamma + 1 = 0$

14. $\cos^2 \frac{\alpha}{2} + \cos^2 \frac{\beta}{2} + \cos^2 \frac{\gamma}{2} = 1$
 Now, $\cos \alpha + \cos \beta + \cos \gamma$
 $= 2\cos^2 \frac{\alpha}{2} - 1 + 2\cos^2 \frac{\beta}{2} - 1 + 2\cos^2 \frac{\gamma}{2} - 1$
 $= 2(1) - 3 = -1$

15. Let the length of the line segment be r and its direction cosines be l, m, n .

\therefore The projections on the co-ordinate axes are lr, mr, nr .

$\therefore r = 3, mr = 4$ and $nr = 5$

$\therefore l^2 r^2 + m^2 r^2 + n^2 r^2 = 3^2 + 4^2 + 5^2$

$\Rightarrow r^2 (l^2 + m^2 + n^2) = 9 + 16 + 25$

$\Rightarrow r^2 = 50 \dots [\because l^2 + m^2 + n^2 = 1]$

$\Rightarrow r = \sqrt{50} = 5\sqrt{2}$

16. The projections on the co-ordinate axes are lr, mr, nr .

$\therefore r = 2, mr = 3$ and $nr = 6$

$\therefore l^2 r^2 + m^2 r^2 + n^2 r^2 = 4 + 9 + 36$

$\Rightarrow r^2 (l^2 + m^2 + n^2) = 49$

$\Rightarrow r = 7$

17. d.r.s. of line are $-2 - 4, 1 - 3, -8 - (-5)$

i.e., $-6, -2, -3$

i.e. $6, 2, 3$

18. AD is the median

$\therefore D \equiv \left(\frac{\lambda - 1}{2}, \frac{5 + 3}{2}, \frac{\mu + 2}{2} \right) \equiv \left(\frac{\lambda - 1}{2}, 4, \frac{\mu + 2}{2} \right)$

\therefore d.r.s. of AD are $\frac{\lambda - 1}{2} - 2, 4 - 3, \frac{\mu + 2}{2} - 5$

i.e. $\frac{\lambda - 5}{2}, 1, \frac{\mu - 8}{2} \dots (i)$

Since AD is equally inclined to co-ordinate axes, its d.r.s. are $\pm 1, \pm 1, \pm 1$

Option (D) satisfies (i).

19. The d.c.s. are

$$\frac{1}{\sqrt{1+9+4}}, \frac{-3}{\sqrt{1+9+4}}, \frac{2}{\sqrt{1+9+4}}$$

$$\Rightarrow \frac{1}{\sqrt{14}}, \frac{-3}{\sqrt{14}}, \frac{2}{\sqrt{14}}$$

20. d.r.s. of line are $-2 - 4, 1 - 3, -8 + 5$

i.e., $-6, -2, -3$ i.e., $6, 2, 3$

\therefore The d.c.s. are $\frac{6}{7}, \frac{2}{7}, \frac{3}{7}$

21. The d.r.s of OP are $3, 12, 4$

\therefore The required d.c.s. are

i.e., $\frac{3}{13}, \frac{12}{13}, \frac{4}{13}$



22. Let the length of the line segment be r and its direction cosines be l, m, n .
 \therefore The projections on the co-ordinate axes are lr, mr, nr .
 $\therefore r = 6, mr = -3$ and $nr = 2$
 $\therefore l^2r^2 + m^2r^2 + n^2r^2 = (6)^2 + (-3)^2 + (2)^2$
 $\Rightarrow r^2(l^2 + m^2 + n^2) = 36 + 9 + 4$
 $\Rightarrow r^2 = 49 \quad \dots[\because l^2 + m^2 + n^2 = 1]$
 $\Rightarrow r = 7$
 Now, d.c.s. of line are $\frac{6}{r}, \frac{-3}{r}, \frac{2}{r}$
 i.e., $\frac{6}{7}, -\frac{3}{7}, \frac{2}{7}$.
23. Here, $\vec{a} = 3\hat{i} + 5\hat{j} - 2\hat{k}$, $\vec{b} = 6\hat{i} + 2\hat{j} + 3\hat{k}$
 \therefore Projection = $\frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = \frac{18 + 10 - 6}{7} = \frac{22}{7}$
24. For option (C), $\frac{4 - (-2)}{-3 - 4} \neq \frac{-3 - 4}{-2 - (-3)}$
 \therefore option (C) is the correct answer.
25. Let $A(5, -2, 7), B(2, 2, \beta), C(-1, 6, -1)$ be the given points
 d.r.s. of AB are $2 - 5, 2 + 2, \beta - 7$
 i.e., $-3, 4, \beta - 7$
 d.r.s. of BC are $-1 - 2, 6 - 2, -1 - \beta$
 i.e., $-3, 4, -1 - \beta$
 Since, the points are collinear
 $\therefore AB \parallel BC$
 $\therefore \frac{4}{-3} = \frac{\beta - 7}{-1 - \beta} \Rightarrow \beta - 7 = -1 - \beta \Rightarrow \beta = 3$
26. Let $A(-1, 2, -3), B(4, a, 1)$ and $C(b, 8, 5)$
 Since, the given points are collinear.
 $\therefore AB \parallel BC$
 $\therefore \frac{4 - (-1)}{b - 4} = \frac{a - 2}{8 - a} = \frac{1 - (-3)}{5 - 1}$
 $\Rightarrow \frac{5}{b - 4} = 1, \frac{a - 2}{8 - a} = 1$
 $\Rightarrow b = 9, a = 5$
27. $P(4, 5, x), Q(3, y, 4)$ and $R(5, 8, 0)$
 Since, the points are collinear
 $\therefore PQ \parallel QR$
 $\therefore \frac{-1}{2} = \frac{y - 5}{8 - y} = \frac{4 - x}{-4}$
 $\Rightarrow \frac{-1}{2} = \frac{y - 5}{8 - y}$ and $\frac{4 - x}{-4} = \frac{-1}{2}$
 $\Rightarrow y - 8 = 2y - 10$ and $8 - 2x = 4$
 $\Rightarrow y = 2$ and $x = 2$
 $\therefore x + y = 4$
28. d.r.s. of AB and BC are $(-2, 2, 2)$ and $(1, -1, -1)$ respectively.
 $\therefore \frac{-2}{1} = \frac{2}{-1} = \frac{2}{-1}$
 \therefore the given points are collinear.
29. The d.r.s. of the diagonal of the line joining the origin to the opposite corner of cube are $a - 0, a - 0, a - 0$ i.e. $1, 1, 1$.
30. Here, $a_1, b_1, c_1 = 1, 1, 2$ and
 $a_2, b_2, c_2 = \sqrt{3} - 1, -\sqrt{3} - 1, 4$
 $\therefore \cos \theta = \frac{1(\sqrt{3} - 1) + 1(-\sqrt{3} - 1) + 2(4)}{\sqrt{1 + 1 + 4} \sqrt{(\sqrt{3} - 1)^2 + (-\sqrt{3} - 1)^2 + 16}}$
 $= \frac{6}{\sqrt{6} \sqrt{4 + 4 + 16}} = \frac{6}{\sqrt{6} \sqrt{24}} = \frac{1}{2}$
 $\therefore \theta = 60^\circ$
31. D.r.s. are $2, 2, 1$ and
 $7 - 3, 2 - 1, 12 - 4 \equiv 4, 1, 8$
 $\therefore \cos \theta = \frac{2 \times 4 + 2 \times 1 + 1 \times 8}{\sqrt{2^2 + 2^2 + 1^2} \sqrt{4^2 + 1^2 + 8^2}} = \frac{18}{3 \times 9} = \frac{2}{3}$
 $\therefore \theta = \cos^{-1} \left(\frac{2}{3} \right)$
32. The direction ratios of AB = $1, 2, -2$ and the direction ratios of CD = $2, 3, 4$
 $a_1a_2 + b_1b_2 + c_1c_2 = (1)(2) + (2)(3) + (-2)(4) = 0$
 $\therefore AB \perp CD \quad \therefore \theta = \frac{\pi}{2}$
33. Putting $z = -m - n$ in $x^2 + y^2 + z^2 = m^2 + n^2$, we get
 $(-m - n)^2 = m^2 + n^2$
 $\Rightarrow mn = 0 \Rightarrow m = 0$ or $n = 0$
 If $m = 0$, then $z = -n$
 $\therefore \frac{x}{-1} = \frac{y}{0} = \frac{z}{1}$
 If $n = 0$, then $z = -m$
 $\therefore \frac{x}{-1} = \frac{y}{1} = \frac{z}{0}$
 $\therefore a_1, b_1, c_1 = -1, 0, 1$ and
 $a_2, b_2, c_2 = -1, 1, 0$
 \therefore The angle between the lines is given by
 $\cos \theta = \frac{1 + 0 + 0}{\sqrt{1 + 0 + 1} \sqrt{1 + 1 + 0}} = \frac{1}{2}$
 $\therefore \theta = \frac{\pi}{3}$



34. Let the direction ratios of the line perpendicular to both the lines be a, b, c .

The line is perpendicular to the lines with Direction ratios $-1, 2, 2$ and $0, 2, 1$

$$\therefore -a + 2b + 2c = 0 \quad \dots(i)$$

$$2b + c = 0 \quad \dots(ii)$$

Solving (i) and (ii), we get

$$\frac{a}{-2} = \frac{b}{1} = \frac{c}{-2}$$

\therefore The d.r.s. of the line are $2, -1, 2$.

\therefore The required d.c.s. of the line are $\frac{2}{3}, \frac{-1}{3}, \frac{2}{3}$.

35. The d.r.s. of the two lines are $1, -1, 2$ and $2, 1, -1$

Let d.r.s. of the line be a, b, c .

$$\therefore a - b + 2c = 0 \quad \dots(i)$$

$$\text{and } 2a + b - c = 0 \quad \dots(ii)$$

Solving (i) and (ii), we get

$$\frac{a}{-1} = \frac{b}{5} = \frac{c}{3}$$

\therefore d.r.s. of the line are $-1, 5, 3$.

\therefore the required d.c.s. are $\frac{-1}{\sqrt{35}}, \frac{5}{\sqrt{35}}, \frac{3}{\sqrt{35}}$.

36. If the straight line makes angles $\alpha, \beta, \gamma, \delta$ with diagonals of a cube, then

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma + \cos^2 \delta = \frac{4}{3}$$

$$\therefore \frac{1 + \cos 2\alpha}{2} + \frac{1 + \cos 2\beta}{2} + \frac{1 + \cos 2\gamma}{2} + \frac{1 + \cos 2\delta}{2} = \frac{4}{3}$$

$$\Rightarrow \cos 2\alpha + \cos 2\beta + \cos 2\gamma + \cos 2\delta = \frac{8}{3} - 4$$

$$\Rightarrow \cos 2\alpha + \cos 2\beta + \cos 2\gamma + \cos 2\delta = \frac{-4}{3}$$

$$37. \cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma + \cos^2 \delta = \frac{4}{3}$$

$$\therefore 1 - \sin^2 \alpha + 1 - \sin^2 \beta + 1 - \sin^2 \gamma + 1 - \sin^2 \delta = \frac{4}{3}$$

$$\Rightarrow \sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma + \sin^2 \delta = 4 - \frac{4}{3} = \frac{8}{3}$$



Evaluation Test

1. $\alpha = \beta = 2\gamma$

$$\Rightarrow \beta = \alpha, \gamma = \frac{\alpha}{2}$$

Since, $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$

$$\therefore \cos^2 \alpha + \cos^2 \alpha + \cos^2 \frac{\alpha}{2} = 1$$

$$\Rightarrow 2\cos^2 \alpha + \frac{1 + \cos \alpha}{2} = 1$$

$$\Rightarrow 4\cos^2 \alpha + \cos \alpha - 1 = 0$$

$$\Rightarrow \cos \alpha = \frac{-1 \pm \sqrt{1+16}}{2(4)} = \frac{-1 \pm \sqrt{17}}{8}$$

If α is acute, then $\cos \alpha$ is positive.

$$\therefore \cos \alpha = \frac{\sqrt{17}-1}{8}$$

2. $\cos \theta = l_1 l_2 + m_1 m_2 + n_1 n_2$

$$\sin^2 \theta = 1 - \cos^2 \theta$$

$$= 1 - (l_1 l_2 + m_1 m_2 + n_1 n_2)^2$$

$$= (l_1^2 + m_1^2 + n_1^2)(l_2^2 + m_2^2 + n_2^2)$$

$$- (l_1 l_2 + m_1 m_2 + n_1 n_2)^2$$

$$= l_1^2 l_2^2 + l_1^2 m_2^2 + l_1^2 n_2^2 + l_2^2 m_1^2 + m_1^2 m_2^2 + m_1^2 n_2^2 + l_2^2 n_1^2 + m_2^2 n_1^2 + n_1^2 n_2^2 - l_1^2 l_2^2 - m_1^2 m_2^2 - n_1^2 n_2^2 - 2l_1 l_2 m_1 m_2 - 2m_1 m_2 n_1 n_2 - 2n_1 n_2 l_1 l_2$$

$$= l_1^2 m_2^2 - 2l_1 l_2 m_1 m_2 + l_2^2 m_1^2 + m_1^2 n_2^2 - 2m_1 m_2 n_1 n_2 + m_2^2 n_1^2 + l_2^2 n_1^2 - 2l_1 l_2 n_1 n_2 + l_1^2 n_2^2$$

$$= (l_1 m_2 - l_2 m_1)^2 + (m_1 n_2 - m_2 n_1)^2 + (n_1 l_2 - n_2 l_1)^2$$

3. Given, $A(2, 3, 7), B(-1, 3, 2), C(p, 5, r)$

Let D be the midpoint of BC .

$$\therefore D \equiv \left(\frac{-1+p}{2}, \frac{3+5}{2}, \frac{2+r}{2} \right) \equiv \left(\frac{p-1}{2}, 4, \frac{r+2}{2} \right)$$

$$\therefore \text{d.r.s. of } AD \text{ are } \frac{p-1}{2} - 2, 4 - 3, \frac{r+2}{2} - 7$$

$$\text{i.e., } \frac{p-5}{2}, 1, \frac{r-12}{2}$$

Since, AD is equally inclined to the axes

$$\therefore \frac{p-5}{2} = 1 = \frac{r-12}{2}$$

$$\Rightarrow p = 7, r = 14$$



4. The d.r.s of AB are $3 - 1, 2 - 4, 6 - 5$
i.e. $2, -2, 1$
Let $a_1, b_1, c_1 = 2, -2, 1$
d.r.s. of BC are $1 - 3, 4 - 5, 5 - 3$
i.e., $-2, -1, 2$
Let $a_2, b_2, c_2 = -2, -1, 2$
 $\therefore a_1a_2 + b_1b_2 + c_1c_2 = 2(-2) + (-2)(-1) + 1(2)$
 $= -4 + 2 + 2 = 0$
 \Rightarrow AB and BC are perpendicular.
 $\therefore m\angle ABC = 90^\circ$
5. The given equations are
 $6mn - 2n + 5m = 0$, and(i)
 $3 + m + 5n = 0$
 $\Rightarrow m = -3 - 5n$ (ii)
Substituting value of m in equation (i),
we get
 $6(-3 - 5n)n - 2n + 5(-3 - 5n) = 0$
 $\Rightarrow -18n - 30n^2 - 2n - 15 - 25n = 0$
 $\Rightarrow 15^2 + 45n + 30n^2 = 0$
 $\Rightarrow n^2 + 3n + 2n^2 = 0$
 $\Rightarrow (n + 2)(n + 1) = 0$
 $\Rightarrow n = -2$ or $n = -1$
If $n = -2$, then $m = -3 - 5(-2)$
 $\Rightarrow m = -3 + 10 = 7$
 \therefore d.r.s. of the 1st line are $1, 2, -1$.
If $n = -1$, then $m = -3 - 5(-1)$
 $\Rightarrow m = -3 + 5 = 2$
 \therefore d.r.s. of the 2nd line are $-2, 1, 1$.
 $\therefore \cos \theta = \frac{1 \times (-2) + 2 \times 1 + (-1) \times 1}{\sqrt{1^2 + 2^2 + (-1)^2} \sqrt{(-2)^2 + 1^2 + 1^2}}$
 $= \frac{-2 + 2 - 1}{\sqrt{6} \sqrt{6}} = \frac{-1}{6}$
 $\therefore \theta = \cos^{-1}\left(\frac{-1}{6}\right)$
6. Since, $(-m)^2 \geq 0$
 $\therefore m^2 - 2m + m^2 \geq 0$
 $\therefore 2m^2 - 2m \geq 0$ (i)
Similarly, $m^2 + n^2 \geq 2mn$ (ii)
and $n^2 + m^2 \geq 2n$ (iii)

- Adding (i), (ii) and (iii), we get
 $2(m^2 + n^2) \geq 2(m + mn + n)$
 $\therefore m + mn + n \leq 1$
 \therefore The maximum value of $m + mn + n$ is 1.
7. Let $A = (a, 2, 3)$, $B = (3, b, 7)$ and
 $C = (-3, -2, -5)$
d.r.s of AB are $3 - a, b - 2, 4$
d.r.s of BC are $-6, -2 - b, -12$
Since the points are collinear
 $\therefore \frac{3 - a}{-6} = \frac{b - 2}{-2 - b} = \frac{4}{-12}$
 $\Rightarrow a = 2, b = 4$
8. Let the d.r.s of the line perpendicular to both
the lines be a, b, c .
d.r.s of lines is $1, -1, 0$ and $2, -1, 1$
 $\therefore a - b = 0$ (i)
 $2a - b + c = 0$ (ii)
On solving (i) and (ii), we get
 $\frac{a}{-1} = \frac{b}{-1} = \frac{c}{1}$
 \therefore d.r.s of the line are $-1, -1, 1$
 \therefore the required d.c.s are $\frac{-1}{\sqrt{3}}, \frac{-1}{\sqrt{3}}, \frac{1}{\sqrt{3}}$
9. Since $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$
 $\therefore \cos^2 \alpha + \cos^2 \alpha + \cos^2 \alpha = 1$ [$\because \alpha = \beta = \gamma$]
 $\Rightarrow 3 \cos^2 \alpha = 1$
 $\Rightarrow \cos \alpha = \pm \frac{1}{\sqrt{3}}$
 $\therefore m = n = \cos \alpha = \pm \frac{1}{\sqrt{3}}$

07 Line



Hints



Classical Thinking

- On X-axis, $y = 0$ and $z = 0$
- On Y-axis, the co-ordinates of x and $z = 0$
- Equation of X-axis is $y = 0, z = 0$.
Hence y and z remain fixed.
- Vector equation of line passing through \bar{a} and parallel to \bar{b} is
 $\bar{r} = \bar{a} + \lambda\bar{b}$
 $\therefore \bar{r} = (\hat{i} - \hat{j} + 3\hat{k}) + \lambda(3\hat{i} + 2\hat{j} - 5\hat{k})$
- Let $A \equiv (2, 1, -1)$
 $\therefore \bar{a} = 2\hat{i} + \hat{j} - \hat{k}$
 $\bar{b} = \hat{i} + 2\hat{j} + \hat{k}$
Now, $\bar{r} = (2\hat{i} + \hat{j} - \hat{k}) + \lambda(\hat{i} + 2\hat{j} + \hat{k})$
- The given line passes through $(3, -4, 6)$
The d.r.s. of line are $2, 5, 3$
 \therefore The given line is parallel to $2\hat{i} + 5\hat{j} + 3\hat{k}$
 \therefore The vector equation of the line is
 $\bar{r} = (3\hat{i} - 4\hat{j} + 6\hat{k}) + \lambda(2\hat{i} + 5\hat{j} + 3\hat{k})$
- The given line passes through $(5, -4, 6)$
The d.r.s. of line are $3, 7, 2$
 \therefore The given line is parallel to $3\hat{i} + 7\hat{j} + 2\hat{k}$
 \therefore The vector equation of the line is
 $\bar{r} = 5\hat{i} - 4\hat{j} + 6\hat{k} + \lambda(3\hat{i} + 7\hat{j} + 2\hat{k})$
- Given, cartesian equation of the line is
 $3x - 2 = 2y + 1 = 3z - 3$
 $\Rightarrow 3\left(x - \frac{2}{3}\right) = 2\left(y + \frac{1}{2}\right) = 3(z - 1)$
 $\Rightarrow \frac{x - \frac{2}{3}}{2} = \frac{y + \frac{1}{2}}{3} = \frac{z - 1}{2}$

- The given line passes through $\left(\frac{2}{3}, \frac{-1}{2}, 1\right)$,
and has direction ratios proportional to $2, 3, 2$.
- The vector equation is
 $\bar{r} = \left(\frac{2}{3}\hat{i} - \frac{1}{2}\hat{j} + \hat{k}\right) + \lambda(2\hat{i} + 3\hat{j} + 2\hat{k})$
- Given cartesian equation of the line is
 $6x - 2 = 3y + 1 = 1 - 2z$
 $\Rightarrow 6\left(x - \frac{1}{3}\right) = 3\left(y + \frac{1}{3}\right) = -2\left(z - \frac{1}{2}\right)$
 $\Rightarrow \frac{x - \frac{1}{3}}{1} = \frac{y + \frac{1}{3}}{2} = \frac{z - \frac{1}{2}}{-3}$
The given line passes through $\left(\frac{1}{3}, \frac{-1}{3}, \frac{1}{2}\right)$ and
the direction ratios are proportional to $1, 2, -3$
 \therefore The vector equation is
 $\bar{r} = \left(\frac{1}{3}\hat{i} - \frac{1}{3}\hat{j} + \frac{1}{2}\hat{k}\right) + \lambda(\hat{i} + 2\hat{j} - 3\hat{k})$
- The given vector equation is
 $\bar{r} = 3\hat{i} - 5\hat{j} + 7\hat{k} + \lambda(2\hat{i} + \hat{j} - 3\hat{k})$
 \therefore The line passes through $(3, -5, 7)$ and has
direction ratios $2, 1, -3$
 \therefore The equation of line is $\frac{x-3}{2} = \frac{y+5}{1} = \frac{z-7}{-3}$
- The required lines passes through $(2, -1, 1)$
and has d.r.s. proportional to $2, 7, -3$
 \therefore The equation of line is
 $\bar{r} = 2\hat{i} - \hat{j} + \hat{k} + \lambda(2\hat{i} + 7\hat{j} - 3\hat{k})$
- The line is parallel to $\frac{x-2}{3} = \frac{y-3}{-1} = \frac{z+1}{2}$
 \therefore d.r.s of line are $3, -1, 2$
also, the line passes through origin
 \therefore The equation of line is
 $\bar{r} = \lambda(3\hat{i} - \hat{j} + 2\hat{k})$



14. $\frac{2x-1}{2} = \frac{1-y}{1} = \frac{z}{3} \Rightarrow \frac{x-\frac{1}{2}}{1} = \frac{y-1}{-1} = \frac{z}{3}$
 \therefore The direction ratios of the required line are 1, -1, 3.
 Also line passes through (2, -1, 3)
 \therefore Equation of the line is $\frac{x-2}{1} = \frac{y+1}{-1} = \frac{z-3}{3}$
15. Let \vec{a} and \vec{b} be the position vectors of the points
 $\therefore \vec{a} = 3\hat{i} - 2\hat{j} - 5\hat{k}$ and $\vec{b} = 3\hat{i} - 2\hat{j} + 6\hat{k}$
 $\therefore \vec{b} - \vec{a} = 3\hat{i} - 2\hat{j} + 6\hat{k} - 3\hat{i} + 2\hat{j} + 5\hat{k}$
 $= 11\hat{k}$
 The vector equation of line is given by
 $\vec{r} = \vec{a} + \lambda(\vec{b} - \vec{a})$
 $\Rightarrow \vec{r} = 3\hat{i} - 2\hat{j} - 5\hat{k} + \lambda(11\hat{k})$
16. Let $\vec{a} = -2\hat{i} + \hat{j} + 3\hat{k}$ and $\vec{b} = \hat{i} - 2\hat{j} + 5\hat{k}$
 $\therefore \vec{b} - \vec{a} = 3\hat{i} - 3\hat{j} + 2\hat{k}$
 The vector equation of the line is
 $\vec{r} = \vec{a} + \lambda(\vec{b} - \vec{a})$
 $\vec{r} = -2\hat{i} + \hat{j} + 3\hat{k} + \lambda(3\hat{i} - 3\hat{j} + 2\hat{k})$
17. The equation of line passing through (x_1, y_1, z_1) and (x_2, y_2, z_2)
 $\frac{x-x_1}{x_2-x_1} = \frac{y-y_1}{y_2-y_1} = \frac{z-z_1}{z_2-z_1}$
 \therefore The equation of line passing through (4, -5, -2) and (-1, 5, 3) is
 $\frac{x-4}{-1-4} = \frac{y+5}{5+5} = \frac{z+2}{3+2}$
 $\Rightarrow \frac{x-4}{1} = \frac{y+5}{-2} = \frac{z+2}{-1}$
18. The required equation of line which passes through the points (1, 2, 3) and (0, 0, 0) is
 $\frac{x-1}{0-1} = \frac{y-2}{0-2} = \frac{z-3}{0-3}$
 $\Rightarrow \frac{x-1}{1} = \frac{y-2}{2} = \frac{z-3}{3}$
19. The equation of the line joining the points (-2, 4, 2) and (7, -2, 5) is
 $\frac{x+2}{7-(-2)} = \frac{y-4}{-2-4} = \frac{z-2}{5-2}$
 $\Rightarrow \frac{x+2}{3} = \frac{y-4}{-2} = \frac{z-2}{1}$

20. $2x + z - 4 = 0$
 $\Rightarrow 2x + z = 4$
 $\Rightarrow z = 4 - 2x$ (i)
 $2y + z = 0$
 $\Rightarrow z = -2y$ (ii)
 $\therefore 4 - 2x = -2y = z$ [From (i) and (ii)]
 $\Rightarrow -2(x-2) = -2y = z$
 $\Rightarrow x-2 = y = \frac{z}{-2}$
 $\Rightarrow x-2+2 = y+2 = \frac{z}{-2} + 2$
 $\Rightarrow \frac{x}{1} = \frac{y+2}{1} = \frac{z-4}{-2}$
21. $a_1, b_1, c_1 = 1, 2, 2$ and $a_2, b_2, c_2 = 3, 2, 6$
 $\therefore \cos \theta = \frac{|1 \times 3 + 2 \times 2 + 2 \times 6|}{\sqrt{1^2 + 2^2 + 2^2} \sqrt{3^2 + 2^2 + 6^2}}$
 $= \frac{19}{3 \times 7} = \frac{19}{21}$
22. $a_1, b_1, c_1 = 2, 2, -1$ and $a_2, b_2, c_2 = 1, 2, 2$
 $\cos \theta = \frac{|2 \times 1 + 2 \times 2 + (-1) \times 2|}{\sqrt{2^2 + 2^2 + (-1)^2} \sqrt{1^2 + 2^2 + 2^2}}$
 $= \frac{|2+4-2|}{3 \times 3} = \frac{4}{9}$
 $\Rightarrow \theta = \cos^{-1} \left(\frac{4}{9} \right)$
23. $a_1, b_1, c_1 = 3, -2, 0$ and $a_2, b_2, c_2 = 2, 3, 4$
 $\Rightarrow \cos \theta = \frac{|3 \times 2 + (-2) \times 3 + 0 \times 4|}{\sqrt{3^2 + (-2)^2 + 0} \sqrt{2^2 + 3^2 + 4^2}}$
 $\Rightarrow \cos \theta = 0$
 $\Rightarrow \theta = \frac{\pi}{2}$
24. $a_1, b_1, c_1 = 1, 2, 3$ and $a_2, b_2, c_2 = 2, 2, -2$
 $a_1 a_2 + b_1 b_2 + c_1 c_2 = 1(2) + 2(2) + 3(-2) = 0$
 \therefore The lines are at right angles.
25. $a_1, b_1, c_1 = 1, 2, 3$ and $a_2, b_2, c_2 = -5, 1, 1$
 $a_1 a_2 + b_1 b_2 + c_1 c_2 = (1)(-5) + (2)(1) + (3)(1) = 0$
 \therefore Lines are at right angle.
26. The given equation of line is,
 $\frac{x-2}{3} = \frac{y-3}{4}; z = 4$
 \therefore The line is perpendicular to Z-axis.
 Hence parallel to XY-plane.



27. Line $L_1: \vec{r} = (2\hat{j} - 3\hat{k}) + \lambda(\hat{i} + 2\hat{j} + 3\hat{k})$
 Line $L_2: \vec{r} = (2\hat{i} + 6\hat{j} + 3\hat{k}) + \mu(2\hat{i} + 3\hat{j} + 4\hat{k})$
 L_1 and L_2 can be written in cartesian form as
 $L_1: \frac{x}{1} = \frac{y-2}{2} = \frac{z+3}{3}$ and
 $L_2: \frac{x-2}{2} = \frac{y-6}{3} = \frac{z-3}{4}$
 The point (2, 6, 3) satisfies both the equations.
 \therefore it is the point of intersection.
Alternate method:
 $L_1: \frac{x}{1} = \frac{y-2}{2} = \frac{z+3}{3} = \lambda$
 $\Rightarrow x = \lambda, y = 2\lambda + 2, z = 3\lambda - 3.$
 $L_2: \frac{x-2}{2} = \frac{y-6}{3} = \frac{z-3}{4} = \mu$
 $\Rightarrow x = 2\mu + 2, y = 3\mu + 6, z = 4\mu + 3$
 Co-ordinates of a point on the line L_1 are
 ($\lambda, 2\lambda + 2, 3\lambda - 3$)
 Co-ordinates of a point on the line L_2 are
 ($2\mu + 2, 3\mu + 6, 4\mu + 3$)
 They intersect. Therefore, their co-ordinates must be same.
 $\therefore \lambda = 2\mu + 2, 2\lambda + 2 = 3\mu + 6, 3\lambda - 3 = 4\mu + 3$
 $\Rightarrow \lambda - 2\mu = 2 \quad \dots(i)$
 $2\lambda - 3\mu = 4 \quad \dots(ii)$
 $3\lambda - 4\mu = 6 \quad \dots(iii)$
 Solving equations (i) and (ii), we get
 $\lambda = 2, \mu = 0.$
 Equation (i) holds true for these values.
 \therefore Intersection is (2, 6, 3).
28. The point (-1, -1, -1) satisfies both the equations so it is the point of intersection
Alternate method:
 Let $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4} = \lambda$
 $\Rightarrow x = 1 + 2\lambda, y = 2 + 3\lambda, z = 3 + 4\lambda.$
 Let $\frac{x-4}{5} = \frac{y-1}{2} = \frac{z}{1} = \mu$
 $\Rightarrow x = 4 + 5\mu, y = 1 + 2\mu, z = \mu$
 Co-ordinates of a point on the first line are
 ($1 + 2\lambda, 2 + 3\lambda, 3 + 4\lambda$)
 Co-ordinates of a point on the second line are
 ($4 + 5\mu, 1 + 2\mu, \mu$)
 They intersect. Therefore, their co-ordinates must be same.
 $\therefore 1 + 2\lambda = 4 + 5\mu, 2 + 3\lambda = 1 + 2\mu, 3 + 4\lambda = \mu$
 $\Rightarrow 2\lambda - 5\mu = 3 \quad \dots(i)$
 $3\lambda - 2\mu = -1 \quad \dots(ii)$
 $4\lambda - \mu = -3 \quad \dots(iii)$

Solving equations (ii) and (iii), we get
 $\lambda = -1, \mu = -1.$

Equation (i) holds true for these values.

\therefore Intersection is (-1, -1, -1).

29. The point (4, 0, -1) satisfies both equations.

\therefore The two lines intersect at (4, 0, -1)

Alternate method:

$$\text{Let } \frac{x-1}{3} = \frac{y-1}{-1} = \lambda; z = -1$$

\Rightarrow general point on this line is
 ($3\lambda + 1, -\lambda + 1, -1$)

$$\text{Also, } \frac{x-4}{2} = \frac{z+1}{3} = \mu; y = 0$$

\Rightarrow general point on this line is
 ($2\mu + 4, 0, 3\mu - 1$)

For $\lambda = 1$ and $\mu = 0$, they have a common point on them. i.e., (4, 0, -1)

30. Co-ordinate of any point on Y-axis is
 $x = 0, z = 0$ i.e. (0, y, 0)

\therefore The foot of perpendicular from the point
 (α, β, γ) on Y-axis is (0, β , 0)

31. Any point on Z-axis is (0, 0, z)

\therefore The foot of perpendicular from the point
 (a, b, c) on Z-axis is (0, 0, c)

32. Distance from X-axis = $\sqrt{y^2 + z^2} = \sqrt{b^2 + c^2}$

33. Distance = $\sqrt{y^2 + z^2} = \sqrt{9 + 16} = 5$

34. Distance from Z-axis = $\sqrt{x^2 + y^2} = 5$

35. Distance from Y-axis = $\sqrt{1 + 9} = \sqrt{10}$

36. Let $p(\vec{\alpha}) = 2\hat{i} + \hat{j} + \hat{k}$

Comparing the equation of line with
 $\vec{r} = \vec{a} + \lambda\vec{b}$, we get

$$\vec{a} = -\hat{i} + 2\hat{j} + 2\hat{k}, \vec{b} = 3\hat{i} + \hat{k}$$

Now,

$$\vec{\alpha} - \vec{a} = 3\hat{i} - \hat{j} - \hat{k}$$

$$|\vec{\alpha} - \vec{a}| = \sqrt{3^2 + (-1)^2 + (-1)^2}$$

$$= \sqrt{11}$$

$$(\vec{\alpha} - \vec{a}) \cdot \vec{b} = (3\hat{i} - \hat{j} - \hat{k}) \cdot (3\hat{i} + \hat{k})$$

$$= 9 - 1$$

$$= 8$$



∴ The distance of point from the line is

$$d = \sqrt{|\bar{\alpha} - \bar{a}|^2 - \left[\frac{(\bar{\alpha} - \bar{a}) \cdot \bar{b}}{|\bar{b}|} \right]^2}$$

$$= \sqrt{11 - \frac{8 \times 8}{10}} = \sqrt{\frac{46}{10}} = \sqrt{\frac{23}{5}}$$

37. Let $A \equiv (2, 4, -1)$

$$\text{Let } \frac{x+5}{1} = \frac{y+3}{4} = \frac{z-6}{-9} = \lambda$$

Any point on the line is

$$P \equiv (\lambda - 5, 4\lambda - 3, -9\lambda + 6)$$

The d.r.s. of the line AP are

$$2 - \lambda + 5, 4 - 4\lambda + 3, -1 + 9\lambda - 6$$

Since, AP is perpendicular to the given line,

$$1(2 - \lambda + 5) + 4(4 - 4\lambda + 3) - 9(-1 + 9\lambda - 6) = 0$$

$$\therefore 2 - \lambda + 5 + 16 - 16\lambda + 12 + 9 - 81\lambda + 54 = 0$$

$$\therefore 98 - 98\lambda = 0 \Rightarrow \lambda = 1$$

The point P is $(1 - 5, 4 - 3, -9 + 6) \equiv (-4, 1, -3)$

$$AP = \sqrt{(2 - (-4))^2 + (4 - 1)^2 + (-1 + 3)^2}$$

$$= \sqrt{36 + 9 + 4} = 7$$

Alternate method:

Since the point is $(2, 4, -1)$

$$\therefore a = 2, b = 4, c = -1$$

Given equation of line is

$$\frac{x+5}{1} = \frac{y+3}{4} = \frac{z-6}{-9}$$

Comparing with

$$\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c},$$

$$x_1 = -5, y_1 = -3, z_1 = 6$$

d.r.s. are 1, 4, -9

$$\therefore \text{d.c.s. are } \frac{1}{\sqrt{98}}, \frac{4}{\sqrt{98}}, \frac{-9}{\sqrt{98}}$$

∴ Perpendicular distance of point from the line is

$$\sqrt{\frac{[(a-x_1)^2 + (b-y_1)^2 + (c-z_1)^2] - [(a-x_1) + (b-y_1)m + (c-z_1)n]^2}{(2+5)^2 + (4+3)^2 + (-1-6)^2}}$$

$$= \sqrt{\frac{-\left[(2+5)\frac{1}{\sqrt{98}} + (4+3)\frac{4}{\sqrt{98}} + (-1-6)\frac{-9}{\sqrt{98}}\right]^2}{49 + 49 + 49 - \frac{98 \times 98}{98}}}$$

$$= \sqrt{49}$$

$$= 7$$

$$38. \frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$$

Any point on the line is $P(\lambda, 2\lambda + 1, 3\lambda + 2)$

Given point is $A(1, 6, 3)$

∴ the d.r.s of the line AP are

$$\lambda - 1, 2\lambda + 1 - 6, 3\lambda + 2 - 3$$

$$\Rightarrow \lambda - 1, 2\lambda - 5, 3\lambda - 1$$

Since, AP is perpendicular to the given line,

$$(1)(\lambda - 1) + (2)(2\lambda - 5) + (3)(3\lambda - 1) = 0$$

$$\Rightarrow \lambda - 1 + 4\lambda - 10 + 9\lambda - 3 = 0$$

$$\Rightarrow 14\lambda - 14 = 0 \Rightarrow \lambda = 1$$

∴ $P \equiv (1, 3, 5)$

$$\therefore AP = \sqrt{(1-1)^2 + (6-3)^2 + (3-5)^2} = \sqrt{13}$$

40. First line passes through

$(x_1, y_1, z_1) = (4, -1, 0)$ and has d.r.s

$$a_1, b_1, c_1 = 1, 2, -3$$

Second line passes through

$(x_2, y_2, z_2) = (1, -1, 2)$ and has d.r.s

$$a_2, b_2, c_2 = 2, 4, -5$$

∴ Shortest distance between them is

$$d = \frac{\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix}}{\sqrt{(b_1 c_2 - b_2 c_1)^2 + (c_1 a_2 - c_2 a_1)^2 + (a_1 b_2 - a_2 b_1)^2}}$$

$$\Rightarrow d = \frac{\begin{vmatrix} 1-4 & -1+1 & 2-0 \\ 1 & 2 & -3 \\ 2 & 4 & -5 \end{vmatrix}}{\sqrt{(-10+12)^2 + (-6+5)^2 + (4-4)^2}}$$

$$= \frac{|-3(2) + 0 + 2(0)|}{\sqrt{5}} = \frac{6}{\sqrt{5}}$$

Alternate method:

Shortest distance between the lines

$\bar{r}_1 = \bar{a}_1 + \lambda \bar{b}_1$ and $\bar{r}_2 = \bar{a}_2 + \mu \bar{b}_2$ is given by

$$d = \frac{|(\bar{a}_2 - \bar{a}_1) \cdot (\bar{b}_1 \times \bar{b}_2)|}{|\bar{b}_1 \times \bar{b}_2|}$$

Here $\bar{a}_1 = 4\hat{i} - \hat{j}$, $\bar{a}_2 = \hat{i} - \hat{j} + 2\hat{k}$

$$\bar{b}_1 = \hat{i} + 2\hat{j} - 3\hat{k}, \bar{b}_2 = 2\hat{i} + 4\hat{j} - 5\hat{k}$$

Now $\bar{a}_2 - \bar{a}_1 = -3\hat{i} + 2\hat{k}$

$$\bar{b}_1 \times \bar{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & -3 \\ 2 & 4 & -5 \end{vmatrix} = 2\hat{i} - \hat{j}$$



$$\begin{aligned} \therefore \text{Shortest distance (d)} &= \left| \frac{(-3\hat{i} + 2\hat{k}) \cdot (2\hat{i} - \hat{j})}{\sqrt{4+1+0}} \right| \\ &= \left| -\frac{6}{\sqrt{5}} \right| \\ &= \frac{6}{\sqrt{5}} \end{aligned}$$

41. Here, $(x_1, y_1, z_1) = (1, -1, 0)$
 $(x_2, y_2, z_2) = (2, -1, 0)$
 $(a_1, b_1, c_1) = (2, 0, 1)$
 $(a_2, b_2, c_2) = (1, -1, -1)$

$$\begin{aligned} d &= \frac{\begin{vmatrix} 2-1 & -1+1 & 0-0 \\ 2 & 0 & 1 \\ 1 & -1 & -1 \end{vmatrix}}{\sqrt{(0+1)^2 + (1+2)^2 + (-2-0)^2}} \\ &= \frac{|1(0+1)|}{\sqrt{14}} \\ &= \frac{1}{\sqrt{14}} \end{aligned}$$

42. Here, $(x_1, y_1, z_1) = (3, 5, 7)$
 $(x_2, y_2, z_2) = (-1, -1, -1)$
 $(a_1, b_1, c_1) = (1, -2, 1)$
 $(a_2, b_2, c_2) = (7, -6, 1)$

$$\begin{aligned} d &= \frac{\begin{vmatrix} -4 & -6 & -8 \\ 1 & -2 & 1 \\ 7 & -6 & 1 \end{vmatrix}}{\sqrt{(-2+6)^2 + (7-1)^2 + (-6+14)^2}} \\ &= \frac{|-16 - 36 - 64|}{2\sqrt{29}} \\ &= \frac{116}{2\sqrt{29}} \\ &= 2\sqrt{29} \end{aligned}$$

43. Here, $(x_1, y_1, z_1) = (1, 2, 3)$
 $(x_2, y_2, z_2) = (2, 4, 5)$
 $(a_1, b_1, c_1) = (2, 3, 4)$

$$(a_2, b_2, c_2) = (3, 4, 5)$$

$$\begin{aligned} d &= \frac{\begin{vmatrix} 2-1 & 4-2 & 5-3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{vmatrix}}{\sqrt{(15-16)^2 + (12-10)^2 + (8-9)^2}} \\ &= \frac{|1(-1) - 2(-2) + 2(-1)|}{\sqrt{(-1)^2 + (2)^2 + (-1)^2}} \\ &= \frac{1}{\sqrt{6}} \end{aligned}$$

44. The given equation of lines are
 $x + a = 2y = -12z$ and $x + y + 2a = 6z - 6a$

$$\text{i.e., } \frac{x+a}{-12} = \frac{y}{-6} = \frac{z}{1} \text{ and } \frac{x}{6} = \frac{y+2a}{6} = \frac{z-a}{1}$$

$$\begin{aligned} d &= \frac{\begin{vmatrix} -a & 2a & -a \\ -12 & -6 & 1 \\ 6 & 6 & 1 \end{vmatrix}}{\sqrt{(-6-6)^2 + (6+12)^2 + (-72+36)^2}} \\ &= \frac{|-a(-12) - 2a(-12-6) - a(-72+36)|}{\sqrt{12^2 + 18^2 + 36^2}} \\ &= \frac{12a + 36a + 36a}{\sqrt{1764}} = \frac{84a}{42} = 2a \end{aligned}$$

45. Since, the line intersect each other,

$$\begin{aligned} \therefore \begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} \\ \Rightarrow \begin{vmatrix} 2-1 & 2-k & -1+1 \\ 3 & 6 & -2 \\ -1 & 4 & -1 \end{vmatrix} &= 0 \\ \Rightarrow 1(-6+8) - (2-k)(-3-2) + 0 &= 0 \\ \Rightarrow 2 + (2-k)5 &= 0 \\ \Rightarrow 12 - 5k &= 0 \\ \Rightarrow k &= \frac{12}{5} \end{aligned}$$

46. Comparing the given equations with
 $\vec{r} = \vec{a}_1 + \lambda \vec{b}_1$ and
 $\vec{r} = \vec{a}_2 + \lambda \vec{b}_2$ we get
 $\vec{a}_1 = -\hat{i} + 3\hat{j} + \hat{k}$, and $\vec{a}_2 = 3\hat{i} + \hat{j}$
 $\vec{b}_1 = \vec{b}_2 = \vec{b} = 5\hat{i} + \hat{j} + 4\hat{k}$



∴ The lines are parallel

$$\bar{a}_2 - \bar{a}_1 = 4\hat{i} - 2\hat{j} - \hat{k}$$

$$(\bar{a}_2 - \bar{a}_1) \times \bar{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 4 & -2 & -1 \\ 5 & 1 & 4 \end{vmatrix}$$

$$= \hat{i}(-8+1) - \hat{j}(16+5) + \hat{k}(4+10)$$

$$= -7\hat{i} - 21\hat{j} + 14\hat{k}$$

∴ The distance between the parallel lines is

$$d = \frac{|(\bar{a}_2 - \bar{a}_1) \times \bar{b}|}{|\bar{b}|}$$

$$\therefore d = \frac{|-7\hat{i} - 21\hat{j} + 14\hat{k}|}{\sqrt{25+1+16}}$$

$$= \frac{\sqrt{49+441+196}}{42} = \frac{7}{\sqrt{3}}$$



Critical Thinking

- The d.r.s. of line are 1, -2, 3 and it passes through point (1, 2, 3)
∴ the vector equation of the line is
 $\bar{r} = \hat{i} + 2\hat{j} + 3\hat{k} + \lambda(\hat{i} - 2\hat{j} + 3\hat{k})$
The cartesian equation of the line is
 $\frac{x-1}{1} = \frac{y-2}{-2} = \frac{z-3}{3}$
- The d.r.s. of line are 3, 2, -8 and it passes through (5, 2, -4)
∴ the vector equation of line is
 $\bar{r} = 5\hat{i} + 2\hat{j} - 4\hat{k} + \lambda(3\hat{i} + 2\hat{j} - 8\hat{k})$
The cartesian equation of the line is
 $\frac{x-5}{3} = \frac{y-2}{2} = \frac{z+4}{-8}$
- The line passes through (2, -3, 4) and has direction ratios proportional to 3, 4, -5.
∴ the cartesian equation of the line is
 $\frac{x-2}{3} = \frac{y+3}{4} = \frac{z-4}{-5}$
∴ $4x - 8 = 3y + 9$ and $-5y - 15 = 4z - 16$
i.e., $4x - 3y = 17$ and $5y + 4z = 1$
- Line || Z-axis
∴ d.r.s. are 0, 0, 1
∴ Required equation is
 $\bar{r} = 2\hat{i} + 3\hat{j} + 4\hat{k} + \lambda(0\hat{i} + 0\hat{j} + 1\hat{k})$
 $\Rightarrow \bar{r} = (2\hat{i} + 3\hat{j} + 4\hat{k}) + \lambda\hat{k}$

5. Let a, b, c, be the direction ratios of the required line.

Since, the line is perpendicular to the lines with d.r.s 3, -16, 7 and 3, 8, -5

$$\therefore 3a - 16b + 7c = 0 \quad \dots(i)$$

$$\text{and } 3a + 8b - 5c = 0 \quad \dots(ii)$$

$$\Rightarrow \frac{a}{2} = \frac{b}{3} = \frac{c}{6} \quad \dots[\text{From (i) and (ii)}]$$

∴ Equation of the required line is

$$\frac{x-1}{2} = \frac{y-2}{3} = \frac{z+4}{6}$$

6. Let A \equiv (-1, 3, -2) and B \equiv (-5, 3, -6)

Midpoint of AB = (-3, 3, -4)

Since the line is equally inclined to the axis

∴ d.r.s. are 1, 1, 1.

∴ equation of the line is

$$\frac{x+3}{1} = \frac{y-3}{1} = \frac{z+4}{1}$$

$$\Rightarrow x+3 = y-3 = z+4$$

7. Co-ordinates of G \equiv (1, 1, 1)

D.r.s of OG are 1, 1, 1 and it passes through (0, 0, 0)

∴ equation of line OG is

$$\frac{x-0}{1} = \frac{y-0}{1} = \frac{z-0}{1}$$

$$\Rightarrow x = y = z$$

8. $\bar{r} = (3\hat{i} + 4\hat{j} + \hat{k}) + t(2\hat{i} - 3\hat{j} + 5\hat{k})$

$$= (3+2t)\hat{i} + (4-3t)\hat{j} + (1+5t)\hat{k}$$

When the line crosses XY plane $\Rightarrow Z = 0$

$$\therefore 1 + 5t = 0 \Rightarrow t = \frac{-1}{5}$$

9. The equation of the line joining the points (-2, 1, -8) and (a, b, c) is

$$\frac{x-(-2)}{a+2} = \frac{y-1}{b-1} = \frac{z-(-8)}{c+8}$$

The above line is in the direction of vector

$$6\hat{i} + 2\hat{j} + 3\hat{k}$$

$$\therefore a+2 = 6, b-1 = 2, c+8 = 3$$

$$\Rightarrow a = 4, b = 3 \text{ and } c = -5$$

10. The equation of the line joining the points (2, 2, 1) and (5, 1, -2) is

$$\frac{x-2}{5-2} = \frac{y-2}{1-2} = \frac{z-1}{-2-1}$$

$$\Rightarrow \frac{x-2}{3} = \frac{y-2}{-1} = \frac{z-1}{-3} \quad \dots(i)$$



- Since, x co-ordinate is 4
 \therefore It satisfies (i)
 $\therefore \frac{4-2}{3} = \frac{y-2}{-1} = \frac{z-1}{-3}$
 $\therefore \frac{z-1}{-3} = \frac{2}{3}$
 $\therefore 3z - 3 = -6$
 $\therefore z = -1$
11. The equation of the line joining the points (3, 4, 1) and (5, 1, 6) is
 $\frac{x-3}{5-3} = \frac{y-4}{1-4} = \frac{z-1}{6-1}$
 $\Rightarrow \frac{x-3}{2} = \frac{y-4}{-3} = \frac{z-1}{5} \quad \dots(i)$
 Co-ordinate of any point on the XY-plane is $z = 0$
 $\therefore \frac{x-3}{2} = \frac{y-4}{-3} = \frac{0-1}{5}$
 $\therefore \frac{x-3}{2} = \frac{-1}{5}$
 $\Rightarrow x - 3 = -\frac{2}{5}$
 $\Rightarrow x = \frac{13}{5}$
 Also we have $\frac{y-4}{-3} = -\frac{1}{5}$
 $\Rightarrow y - 4 = \frac{3}{5} \Rightarrow y = \frac{23}{5}$
 \therefore The line meets the XY-plane at $\left(\frac{13}{5}, \frac{23}{5}, 0\right)$
12. Here, $(x_1, y_1, z_1) = (3, -6, 10)$ and $|r| = \sqrt{17}$
 $x_2 = x_1 + r = 3 - \frac{2}{\sqrt{17}} (\sqrt{17}) = 1$
 $y_2 = y_1 + mr = -6 + \frac{3}{\sqrt{17}} (\sqrt{17}) = -3$
 $z_2 = z_1 + nr = 10 - \frac{2}{\sqrt{17}} (\sqrt{17}) = 8$
13. The d.r.s. of the two lines are 2, -1, 1 and 4, -1, λ
 Since, the lines are perpendicular
 $a_1a_2 + b_1b_2 + c_1c_2 = 0$
 $\Rightarrow 2(4) + (-1)(-1) + (1)(\lambda) = 0$
 $\Rightarrow \lambda + 9 = 0$
 $\Rightarrow \lambda = -9$

14. $a_1, b_1, c_1 = 2, p, 5$ and
 $a_2, b_2, c_2 = 3, -p, p$
 Since, the given lines are perpendicular.
 $\therefore (2)(3) + p(-p) + (5)(p) = 0$
 $\Rightarrow 6 - p^2 + 5p = 0$
 $\Rightarrow p^2 - 5p - 6 = 0$
 $\Rightarrow (p - 6)(p + 1) = 0$
 $\Rightarrow p = 6$ or $p = -1$
15. $a_1, b_1, c_1 = 2, \lambda, 0$ and $a_2, b_2, c_2 = 1, 3, 1$
 Since, the lines are perpendicular.
 $a_1a_2 + b_1b_2 + c_1c_2 = 0$
 $\therefore 2(1) + \lambda(3) + 0(1) = 0$
 $\therefore 2 + 3\lambda = 0$
 $\therefore \lambda = \frac{-2}{3}$
16. Given lines pass through common point (1, 2, 3)
 Also, $a_1a_2 + b_1b_2 + c_1c_2 = 2(3) + 3(4) + 4(5) \neq 0$
 \therefore lines are intersecting
17. Let $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$, then
 $\vec{r} \times \vec{a} = \vec{b} \times \vec{a} \Rightarrow (\vec{r} - \vec{b}) \times \vec{a} = 0$
 $\therefore \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ x-2 & y & z+1 \\ 1 & 1 & 0 \end{vmatrix} = 0$
 $\Rightarrow (-z-1)\hat{i} - (-z-1)\hat{j} + (x-y-2)\hat{k} = 0$
 $\Rightarrow z = -1, x - y = 2 \quad \dots(i)$
 Now, $\vec{r} \times \vec{b} = \vec{a} \times \vec{b} \Rightarrow (\vec{r} - \vec{a}) \times \vec{b} = 0$
 $\therefore \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ x-1 & y-1 & z \\ 2 & 0 & -1 \end{vmatrix} = 0$
 $\Rightarrow (1-y)\hat{i} - (1-x-2z)\hat{j} + (2-2y)\hat{k} = 0$
 $\Rightarrow y = 1, x + 2z = 1 \quad \dots(ii)$
 Solving (i) and (ii), we get
 $x = 3, y = 1, z = -1$
18. Let P (x, y, z) be any point
 Now by the given condition, we get
 $\left[\sqrt{x^2 + y^2}\right]^2 + \left[\sqrt{y^2 + z^2}\right]^2 + \left[\sqrt{z^2 + x^2}\right]^2 = 36$
 i.e., $x^2 + y^2 + z^2 = 18$
 \therefore The distance from origin
 $= \sqrt{x^2 + y^2 + z^2} = \sqrt{18} = 3\sqrt{2}$




19. Let $\frac{x}{2} = \frac{y-1}{3} = \frac{z-1}{3} = \lambda$

∴ Any general point on this line is

Q $(2\lambda, 3\lambda+1, 3\lambda+1)$

Let P $\equiv (1, 2, 3)$.

∴ D.r.s. of PQ are $2\lambda - 1, 3\lambda - 1, 3\lambda - 2$



Since, PQ is perpendicular to given line

∴ $a_1a_2 + b_1b_2 + c_1c_2 = 0$.

∴ $(2\lambda - 1)2 + (3\lambda - 1)3 + (3\lambda - 2)3 = 0$

∴ $\lambda = \frac{1}{2}$

∴ Q $\equiv \left(1, \frac{5}{2}, \frac{5}{2}\right)$

20. Let $\frac{x}{2} = \frac{y-2}{3} = \frac{z-3}{4} = \lambda$

Any point on the line is

P $\equiv (2\lambda, 3\lambda + 2, 4\lambda + 3)$

Given point is A $(3, -1, 11)$

∴ The d.r.s. of AP are

$2\lambda - 3, 3\lambda + 3, 4\lambda - 8$

Since, the line AP is perpendicular to the given line

∴ $2(2\lambda - 3) + 3(3\lambda + 3) + 4(4\lambda - 8) = 0$

$\Rightarrow 29\lambda - 29 = 0$

$\Rightarrow \lambda = 1$

∴ P $\equiv (2, 5, 7)$

21. Let $\frac{x+3}{5} = \frac{y-1}{2} = \frac{z+4}{3} = \lambda$

∴ Any general point on this line is

Q $(5\lambda - 3, 2\lambda + 1, 3\lambda - 4)$

Let P $\equiv (0, 2, 3)$.

∴ The d.r.s. of PQ are $5\lambda - 3, 2\lambda - 1, 3\lambda - 7$

Since, PQ is perpendicular to given line

∴ $5(5\lambda - 3) + 2(2\lambda - 1) + 3(3\lambda - 7) = 0$

$\Rightarrow \lambda = 1$

∴ Q $\equiv (2, 3, -1)$

22. Distance of point P $(\vec{\alpha})$ from the

line $\vec{r} = \vec{a} + \lambda\vec{b}$ is

$$\sqrt{|\vec{\alpha} - \vec{a}|^2 - \left[\frac{(\vec{\alpha} - \vec{a}) \cdot \vec{b}}{|\vec{b}|} \right]^2}$$

Given, P $(\vec{\alpha}) \equiv (0, 0, 0)$ and

$\vec{t} = 4\hat{i} + 2\hat{j} + 4\hat{k} + \lambda(3\hat{i} + 4\hat{j} - 5\hat{k})$

∴ $\vec{a} = 4\hat{i} + 2\hat{j} + 4\hat{k}$ and

$\vec{b} = 3\hat{i} + 4\hat{j} - 5\hat{k}$


∴ Distance of point

$$\begin{aligned} &= \sqrt{[(-4)^2 + (-2)^2 + (-4)^2] - \left[\frac{-4(3) - 2(4) - 4(-5)}{\sqrt{3^2 + 4^2 + (-5)^2}} \right]^2} \\ &= \sqrt{16 + 4 + 16} \\ &= 6 \end{aligned}$$

Alternate method:

$\vec{AO} = 4\hat{i} + 2\hat{j} + 4\hat{k}$

∴ $OA = \sqrt{16 + 4 + 16} = 6$



AM = Projection of OA on AL

$$= \frac{12 + 8 - 20}{\sqrt{9 + 16 + 25}} = 0$$

In right angled ΔOAM , $d^2 = OA^2 - AM^2$

$\Rightarrow d^2 = 6^2 - 0 \Rightarrow d = 6$

23. Any point on the line $\frac{x-1}{2} = \frac{y}{9} = \frac{z}{5} = \lambda$ is

P $(2\lambda + 1, 9\lambda, 5\lambda)$

Let A $\equiv (5, 4, -1)$

The d.r.s. of the line AP are

$2\lambda + 1 - 5, 9\lambda - 4, 5\lambda - (-1)$

$\Rightarrow 2\lambda - 4, 9\lambda - 4, 5\lambda + 1$

Since, AP is perpendicular to the given line

∴ $2(2\lambda - 4) + 9(9\lambda - 4) + 5(5\lambda + 1) = 0$

$\Rightarrow 4\lambda - 8 + 81\lambda - 36 + 25\lambda + 5 = 0$

$\Rightarrow \lambda = \frac{39}{110}$



$$\therefore P \equiv \left(\frac{188}{110}, \frac{351}{110}, \frac{195}{110} \right)$$

$$\therefore AP = \sqrt{\left(5 - \frac{188}{110} \right)^2 + \left(4 - \frac{351}{110} \right)^2 + \left(-1 - \frac{195}{110} \right)^2}$$

$$= \frac{1}{\sqrt{110^2}} \sqrt{131044 + 7921 + 93025}$$

$$= \sqrt{\frac{2109}{110}}$$

24. Let $\frac{x-11}{10} = \frac{y+2}{-4} = \frac{z+8}{-11} = \lambda$
 Any point on the line is
 $P(10\lambda + 11, -4\lambda - 2, -11\lambda - 8)$
 Let $A \equiv (2, -1, 5)$
 The d.r.s. of the line AP are
 $10\lambda + 11 - 2, -4\lambda - 2 - (-1), -11\lambda - 8 - 5$
 i.e., $10\lambda + 9, -4\lambda - 1, -11\lambda - 13$
 Since, AP is perpendicular to the given line
 $\therefore 10(10\lambda + 9) - 4(-4\lambda - 1) - 11(-11\lambda - 13) = 0$
 $\Rightarrow 100\lambda + 90 + 16\lambda + 4 + 121\lambda + 143 = 0$
 $\Rightarrow 237\lambda + 237 = 0 \Rightarrow \lambda = -1$
 $\therefore P \equiv (1, 2, 3)$
 $\therefore AP = \sqrt{(2-1)^2 + (-1-2)^2 + (5-3)^2}$
 $= \sqrt{1+9+4} = \sqrt{14}$

25. The given equation of line is
 $\frac{x-1}{2} = \frac{y+1}{-3} = z$
 The co-ordinates of any point on the given line are $(2\lambda + 1, -3\lambda - 1, \lambda)$
 The distance of this point from the point $(1, -1, 0)$ is $4\sqrt{14}$.
 $\therefore (2\lambda)^2 + (-3\lambda)^2 + (\lambda)^2 = (4\sqrt{14})^2 \Rightarrow \lambda = \pm 4$
 \therefore The co-ordinates of the required point are $(9, -13, 4)$ or $(-7, 11, -4)$
 The point nearer to the origin is $(-7, 11, -4)$.

26. The equation of the line joining the points $A(2, -3, -1)$ and $B(8, -1, 2)$ is
 $\frac{x-2}{8-2} = \frac{y+3}{-1+3} = \frac{z+1}{2+1}$
 $\Rightarrow \frac{x-2}{6} = \frac{y+3}{2} = \frac{z+1}{3} = \lambda$
 Any point on the line is
 $(6\lambda + 2, 2\lambda - 3, 3\lambda - 1)$
 The distance of this point from the point $A(2, -3, -1)$ is 14 units.
 $\therefore (6\lambda)^2 + (2\lambda)^2 + (3\lambda)^2 = (14)^2$


$$\therefore 49\lambda^2 = 196$$

$$\therefore \lambda^2 = 4 \Rightarrow \lambda = \pm 2$$

$$\therefore$$
 The points are $(14, 1, 5)$ and $(-10, -7, -7)$

$$\therefore$$
 The point nearer to the origin is $(-10, -7, -7)$.

27. Any point on the line
 $\frac{x+5}{1} = \frac{y+3}{4} = \frac{z-6}{-9} = \lambda$ is given by
 $M \equiv (\lambda - 5, 4\lambda - 3, -9\lambda + 6)$.



The d.r.s. of PM are
 $\lambda - 7, 4\lambda - 7, -9\lambda + 7$
 Since, PM is perpendicular to AM,
 $\therefore 1(\lambda - 7) + 4(4\lambda - 7) - 9(-9\lambda + 7) = 0$
 $\Rightarrow 98\lambda - 98 = 0 \Rightarrow \lambda = 1$
 $\therefore M = (-4, 1, -3)$
 Now, Equation of perpendicular passing through $P(2, 4, -1)$ and $M(-4, 1, -3)$ is
 $\frac{x-2}{-4-2} = \frac{y-4}{1-4} = \frac{z+1}{-3+1}$
 $\Rightarrow \frac{x-2}{6} = \frac{y-4}{3} = \frac{z+1}{2}$

28. The direction ratios are same. Also both lines pass through origin.
 \therefore Given lines are coinciding lines.

29. The lines can be rewritten as
 $\vec{r} = (\hat{i} - 2\hat{j} + 3\hat{k}) + t(-\hat{i} + \hat{j} - \hat{k})$ and
 $\vec{r} = (\hat{i} - \hat{j} - \hat{k}) + s(\hat{i} + 2\hat{j} - 2\hat{k})$
 Here, $(x_1, y_1, z_1) = (1, -2, 3)$
 $(x_2, y_2, z_2) = (1, -1, -1)$
 $(a_1, b_1, c_1) = (-1, 1, -1)$
 $(a_2, b_2, c_2) = (1, 2, -2)$

\therefore Shortest distance (d)

$$d = \frac{\begin{vmatrix} 1-1 & -1+2 & -1-3 \\ -1 & 1 & -1 \\ 1 & 2 & -2 \end{vmatrix}}{\sqrt{(-2+2)^2 + (-1-2)^2 + (-2-1)^2}}$$

$$= \frac{|0-1(3)-4(-3)|}{3\sqrt{2}} = \frac{9}{3\sqrt{2}} = \frac{3}{\sqrt{2}}$$



30. The equations of the given lines are

$$\vec{r} = (-\hat{i} + \hat{j} - \hat{k}) + \lambda(\hat{i} + \hat{j} - \hat{k}) \text{ and}$$

$$\vec{r} = (\hat{i} - \hat{j} + 2\hat{k}) + \mu(-\hat{i} + 2\hat{j} + \hat{k})$$

Here, $(x_1, y_1, z_1) = (-1, 1, -1)$

$$(x_2, y_2, z_2) = (1, -1, 2)$$

$$(a_1, b_1, c_1) = (1, 1, -1)$$

$$(a_2, b_2, c_2) = (-1, 2, 1)$$

\therefore Shortest Distance (d)

$$= \frac{\begin{vmatrix} 1+1 & -1-1 & 2+1 \\ 1 & 1 & -1 \\ -1 & 2 & 1 \end{vmatrix}}{\sqrt{(1+2)^2 + (1-1)^2 + (2+1)^2}}$$

$$= \frac{|2(3) + 2(0) + 3(3)|}{3\sqrt{2}} = \frac{5}{\sqrt{2}}$$

31. The given equation of lines are

$$\frac{x-1}{k} = \frac{y+1}{3} = \frac{z-1}{4} \text{ and}$$

$$\frac{x-3}{1} = \frac{2y-9}{2k} = \frac{z}{1}$$

$$\text{i.e. } \frac{x-3}{1} = \frac{y-\frac{9}{2}}{\frac{1}{2}} = \frac{z}{1}$$

Since the line intersect,

$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

$$\therefore \begin{vmatrix} 2 & \frac{11}{2} & -1 \\ k & 3 & 4 \\ 1 & k & 1 \end{vmatrix} = 0$$

$$\therefore 2(3 - 4k) - \frac{11}{2}(k - 4) - 1(k^2 - 3) = 0$$

$$\therefore 6 - 8k - \frac{11}{2}k + 22 - k^2 + 3 = 0$$

$$\therefore 2k^2 + 27k - 62 = 0$$

$$\therefore 2k^2 - 4k + 31k - 62 = 0$$

$$\therefore 2k(k-2) + 31(k-2) = 0$$

$$\therefore k = 2 \text{ or } k = \frac{-31}{2}$$

$$32. \quad \vec{a}_2 - \vec{a}_1 = -3\hat{i} + 2\hat{k}$$

$$\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & -3 \\ 1 & 4 & -5 \end{vmatrix}$$

$$= \hat{i}(-10+12) - \hat{j}(-5+3) + \hat{k}(4-2)$$

$$= 2\hat{i} + 2\hat{j} + 2\hat{k}$$

$$\therefore (\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = (-3\hat{i} + 2\hat{k}) \cdot (2\hat{i} + 2\hat{j} + 2\hat{k})$$

$$= -6 + 4 = -2$$

33. Let the components of the line vector be a, b, c.

$$a^2 + b^2 + c^2 = (63)^2 \quad \dots(i)$$

$$\frac{a}{3} = \frac{b}{-2} = \frac{c}{6} = \lambda(\text{say})$$

$$\Rightarrow a = 3\lambda, b = -2\lambda, c = 6\lambda$$

$$\therefore 9\lambda^2 + 4\lambda^2 + 36\lambda^2 = (63)^2 \quad \dots[\text{From (i)}]$$

$$\therefore 49\lambda^2 = (63)^2 \Rightarrow \lambda = \pm \frac{63}{7} = \pm 9$$

Since, as the line makes an obtuse angle with X-axis, $a = 3\lambda < 0$, $\lambda = -9$

\therefore The required components are $-27, 18, -54$.



Competitive Thinking

1. The line passes through $(1, -2, -1)$

Let other point be (x_2, y_2, z_2)

Direction ratio are $0, 6, -1$

$$\therefore x_2 - 1 = 0 \Rightarrow x_2 = 1$$

$$y_2 - (-2) = 6 \Rightarrow y_2 = 4$$

$$z_2 - (-1) = -1 \Rightarrow z_2 = -2$$

2. The equation of line passing through (a, b, c)

and having d.r.s. $0, 0, 1$ is $\frac{x-a}{0} = \frac{y-b}{0} = \frac{z-c}{1}$

3. Let a, b, c be the d.r.s. of the required line

d.r.s. of the given lines are $2, -2, 1$ and $1, -2, 2$.

$$\therefore 2a - 2b + c = 0 \quad \dots(i)$$

$$a - 2b + 2c = 0 \quad \dots(ii)$$

$$\therefore \frac{a}{-4+2} = \frac{-b}{4-1} = \frac{c}{-4+2}$$

$$\Rightarrow \frac{a}{-2} = \frac{b}{-3} = \frac{c}{-2}$$

\therefore Equation of the required line is

$$\frac{x-3}{-2} = \frac{y+1}{-3} = \frac{z-2}{-2}$$

$$\Rightarrow \frac{x-3}{2} = \frac{y+1}{3} = \frac{z-2}{2}$$



4. $\frac{x+2}{2} = \frac{2y-5}{3}, z = -1$
 $\therefore \frac{x+2}{2} = \frac{y-\frac{5}{2}}{\frac{3}{2}}, z = -1$
 $\therefore \frac{x+2}{4} = \frac{y-\frac{5}{2}}{3}, z = -1$
 \therefore d.r.s of given line are 4, 3, 0
 \therefore d.c.s of the line are $\frac{4}{\sqrt{4^2+3^2}}, \frac{3}{\sqrt{4^2+3^2}}, 0 \Rightarrow \frac{4}{5}, \frac{3}{5}, 0$
5. d.r.s. of given line are 1, 1, 1
 \therefore d.c.s. are $\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}$
6. Given equation of line $x = 4z + 3, y = 2 - 3z$
 $\Rightarrow z = \frac{x-3}{4}, z = \frac{y-2}{-3}$
 \therefore Equation of line is $\frac{x-3}{4} = \frac{y-2}{-3} = \frac{z-0}{1}$
d.r.s of line are 4, -3, 1
 $\therefore \cos \alpha = \frac{4}{\sqrt{4^2 + (-3)^2 + 1^2}} = \frac{4}{\sqrt{26}}$,
 $\cos \beta = \frac{-3}{\sqrt{26}}, \cos \gamma = \frac{1}{\sqrt{26}}$
 $\therefore \cos \alpha + \cos \beta + \cos \gamma = \frac{4}{\sqrt{26}} - \frac{3}{\sqrt{26}} + \frac{1}{\sqrt{26}}$
 $= \frac{2}{\sqrt{26}}$
7. The given equation is $\frac{x-3}{3} = \frac{y-2}{1} = \frac{z-1}{0}$
The direction ratios of the above line are 3, 1, 0
 $\Rightarrow n = \cos \gamma = 0 \Rightarrow \gamma = 90^\circ$
 \therefore The given straight line is perpendicular to Z-axis.
8. Let a, b, c be the direction ratios of the line.
 $\therefore a - b + c = 0$ and(i)
 $a - 3b = 0$ (ii)
 $\therefore \frac{a}{3} = \frac{b}{1} = \frac{c}{-2}$
 \therefore the direction ratios of the line are 3, 1, -2.
9. If a line is equally inclined to axes, then
 $= m = n = \pm \frac{1}{\sqrt{3}}$
 \therefore d.r.s. of the line are 1, 1, 1

Given that the line passes through the point $(-3, 2, -5)$

$$\therefore \text{The equation of line is } \frac{x+3}{1} = \frac{y-2}{1} = \frac{z+5}{1}$$

10. Here, $(x_1, y_1, z_1) \equiv (a, b, c)$
and $(x_2, y_2, z_2) \equiv (a-b, b-c, c-a)$

Required equation of line is

$$\frac{x-a}{a-b-a} = \frac{y-b}{b-c-b} = \frac{z-c}{c-a-a}$$

$$\text{i.e., } \frac{x-a}{b} = \frac{y-b}{c} = \frac{z-c}{a}$$

11. Given equation is $\frac{x-1}{m} = \frac{y-2}{n} = \frac{z+1}{n}$

The equation of line passing through $(1, 2, -1)$ and $(-1, 0, 1)$ is

$$\frac{x-1}{-1-1} = \frac{y-2}{0-2} = \frac{z+1}{1+1}$$

$$\Rightarrow \frac{x-1}{-2} = \frac{y-2}{-2} = \frac{z+1}{2}$$

$$\Rightarrow \frac{x-1}{1} = \frac{y-2}{1} = \frac{z+1}{-1} \quad \dots\text{(i)}$$

Comparing (i) with given equation, we get
 $= 1, m = 1, n = -1$

12. Equation of line AB in vector form is

$$\bar{r} = 6\bar{a} - 4\bar{b} + 4\bar{c} + \lambda(-4\bar{c} - \{6\bar{a} - 4\bar{b} + 4\bar{c}\})$$

$$\Rightarrow \bar{r} = 6\bar{a} - 4\bar{b} + 4\bar{c} + \lambda(-6\bar{a} + 4\bar{b} - 8\bar{c}) \quad \dots\text{(i)}$$

Equation of line CD in vector form is

$$\bar{r}' = \bar{a} + 2\bar{b} - 5\bar{c} + \lambda'(-\bar{a} - 2\bar{b} - 3\bar{c} - \{\bar{a} + 2\bar{b} - 5\bar{c}\})$$

$$\Rightarrow \bar{r}' = \bar{a} + 2\bar{b} - 5\bar{c} + \lambda'(-2\bar{a} - 4\bar{b} + 2\bar{c}) \quad \dots\text{(ii)}$$

The point of intersection of AB and CD will satisfy

$$\bar{r} = \bar{r}'$$

$$\Rightarrow 6\bar{a} - 4\bar{b} + 4\bar{c} + \lambda(-6\bar{a} + 4\bar{b} - 8\bar{c})$$

$$= \bar{a} + 2\bar{b} - 5\bar{c} + \lambda'(-2\bar{a} - 4\bar{b} + 2\bar{c})$$

Comparing the coefficients of \bar{a} and \bar{b} , we get

$$6\lambda - 2\lambda' = 5 \quad \dots\text{(iii)}$$

$$2\lambda + 2\lambda' = 3 \quad \dots\text{(iv)}$$

$$\Rightarrow \lambda = 1 \text{ and } \lambda' = \frac{1}{2}$$

Substituting value of λ in equation (i), we get the point of intersection

- \therefore Point of intersection $\bar{r} = -4\bar{c}$ i.e. point B.



13. The equation of the line joining the points $(3, 5, -7)$ and $(-2, 1, 8)$ is

$$\frac{x-3}{-2-3} = \frac{y-5}{1-5} = \frac{z-(-7)}{8-(-7)}$$

$$\text{Let } \frac{x-3}{-5} = \frac{y-5}{-4} = \frac{z+7}{15} = \lambda$$

$$\Rightarrow x = 3 - 5\lambda, y = 5 - 4\lambda, z = -7 + 15\lambda$$

For YZ plane, $x = 0$

$$\therefore 3 - 5\lambda = 0 \Rightarrow \lambda = \frac{3}{5}$$

$$\text{Now, } y = 5 - 4\lambda = 5 - 4\left(\frac{3}{5}\right) = 5 - \frac{12}{5} = \frac{13}{5}$$

$$z = -7 + 15\lambda = -7 + 15\left(\frac{3}{5}\right) = 2$$

$$\therefore \text{The required point is } \left(0, \frac{13}{5}, 2\right)$$

14. Given equations of line are

$$\vec{r} = (\hat{i} + 2\hat{j} - \hat{k}) + \lambda(3\hat{i} - 4\hat{k}) \quad \dots(i)$$

$$\text{and } \vec{r} = (1-t)(4\hat{i} - \hat{j}) + t(2\hat{i} + \hat{j} - 3\hat{k})$$

$$\text{i.e., } \vec{r} = (4\hat{i} - \hat{j}) + t(-2\hat{i} + 2\hat{j} - 3\hat{k}) \quad \dots(ii)$$

Now, d.r.s. of line (i) and (ii) are

$$a_1, b_1, c_1 = 3, 0, -4$$

$$\text{and } a_2, b_2, c_2 = -2, 2, -3$$

$$\cos \theta = \left| \frac{3(-2) + 0(2) + (-4)(-3)}{\sqrt{9+0+16} \sqrt{4+4+9}} \right|$$

$$\Rightarrow \cos \theta = \frac{6}{5\sqrt{17}}$$

$$\Rightarrow \theta = \cos^{-1} \left(\frac{6}{5\sqrt{17}} \right)$$

15. The d.r.s. of the lines are 2, 5, -3 and -1, 8, 4

$$\therefore \cos \theta = \left| \frac{2(-1) + 5(8) + (-3)(4)}{\sqrt{2^2 + 5^2 + (-3)^2} \sqrt{(-1)^2 + 8^2 + 4^2}} \right|$$

$$\Rightarrow \cos \theta = \frac{26}{9\sqrt{38}}$$

$$\Rightarrow \theta = \cos^{-1} \left(\frac{26}{9\sqrt{38}} \right)$$

16. The d.r.s. of the lines are 1, 0, -1 and 3, 4, 5

$$\therefore \cos \theta = \left| \frac{1(3) + 0(4) + (-1)(5)}{\sqrt{1^2 + 0^2 + (-1)^2} \sqrt{3^2 + 4^2 + 5^2}} \right| = \left| \frac{-2}{10} \right|$$

$$\therefore \theta = \cos^{-1} \left(\frac{1}{5} \right)$$

$$17. \cos \theta = \left| \frac{2(1) + 2(2) + (-1)(2)}{\sqrt{4+4+1} \sqrt{1+4+4}} \right| = \left| \frac{4}{\sqrt{9} \cdot \sqrt{9}} \right| = \frac{4}{9}$$

$$\Rightarrow \theta = \cos^{-1} \left(\frac{4}{9} \right)$$

18. The d.r.s. of the line joining the points $(2, 1, -3)$ and $(-3, 1, 7)$ are -5, 0, 10

The d.r.s. of the line parallel to line

$$\frac{x-1}{3} = \frac{y}{4} = \frac{z+3}{5} \text{ are } 3, 4, 5$$

- \therefore The angle between the lines having d.r.s. -5, 0, 10 and 3, 4, 5 is

$$\cos \theta = \left| \frac{-5(3) + 0(4) + 10(5)}{\sqrt{25+0+100} \sqrt{9+16+25}} \right|$$

$$\Rightarrow \cos \theta = \frac{35}{25\sqrt{10}}$$

$$\Rightarrow \theta = \cos^{-1} \left(\frac{7}{5\sqrt{10}} \right)$$

19. $a_1a_2 + b_1b_2 + c_1c_2 = (2)(1) + (5)(2) + (4)(-3) = 0$

\therefore Lines are perpendicular

$$\therefore \theta = 90^\circ$$

20. The equation of given lines are

$$\frac{x}{3} = \frac{y}{2} = \frac{z}{-6} \text{ and } \frac{x}{2} = \frac{y}{-12} = \frac{z}{-3}$$

$$a_1a_2 + b_1b_2 + c_1c_2 = 3(2) + 2(-12) + (-6)(-3) = 0$$

\therefore Lines are perpendicular

$$\therefore \theta = 90^\circ$$

21. The first line is parallel to Z-axis and the second line is parallel to X-axis.

\therefore The angle between them is 90° .

22. Let the d.r.s of the given line be a, b, c

Then, according to given condition of perpendicularity,

$$-1.a + 2.b + 2.c = 0 \quad \dots(i)$$

$$0.a + 2.b + 1.c = 0 \quad \dots(ii)$$

On solving (i) and (ii), we get

$$a = 2, b = -1 \text{ and } c = 2$$

23. $a_1, b_1, c_1 = -3, 2k, 2$ and $a_2, b_2, c_2 = 3k, 1, -5$
Since, the lines are perpendicular to each other,

$$a_1a_2 + b_1b_2 + c_1c_2 = 0$$

$$\therefore (-3)(3k) + (2k)(1) + (2)(-5) = 0$$

$$\Rightarrow -9k + 2k - 10 = 0$$


$$\Rightarrow k = \frac{-10}{7}$$




24. Given equations of lines are
 $x = ay + b, z = cy + d$
 $\Rightarrow \frac{x-b}{a} = \frac{y}{1}, \frac{z-d}{c} = \frac{y}{1}$
 $\Rightarrow \frac{x-b}{a} = \frac{y}{1} = \frac{z-d}{c}$
 and $x = a'y + b', z = c'y + d'$
 $\Rightarrow \frac{x-b'}{a'} = \frac{y}{1}, \frac{z-d'}{c'} = \frac{y}{1}$
 $\Rightarrow \frac{x-b'}{a'} = \frac{y}{1} = \frac{z-d'}{c'}$
 Since, the lines are perpendicular to each other.
 $a_1a_2 + b_1b_2 + c_1c_2 = 0$
 $\Rightarrow aa' + 1(1) + cc' = 0$
 $\Rightarrow aa' + cc' = -1$
25. Equation of line BC is
 $\frac{x-0}{2-0} = \frac{y+11}{-3+11} = \frac{z-4}{1-4}$
 $\Rightarrow \frac{x}{2} = \frac{y+11}{8} = \frac{z-4}{-3}$
 Let $\frac{x}{2} = \frac{y+11}{8} = \frac{z-4}{-3} = \lambda$
 Any point D on the line is
 $\equiv (2\lambda, 8\lambda - 11, -3\lambda + 4)$
 Given point A $\equiv (1, 8, 4)$
 \therefore d.r.s of AD are $2\lambda - 1, 8\lambda - 11 - 8, -3\lambda + 4 - 4$
 $= 2\lambda - 1, 8\lambda - 19, -3\lambda$
 Since, $AD \perp BC$,
 $\therefore aa_1 + bb_1 + cc_1 = 0$
 $\Rightarrow 2(2\lambda - 1) + 8(8\lambda - 19) - 3(-3\lambda) = 0$
 $\Rightarrow 4\lambda - 2 + 64\lambda - 152 + 9\lambda = 0$
 $\Rightarrow 77\lambda = 154$
 $\Rightarrow \lambda = 2$
 $\therefore D \equiv (4, 5, -2)$
26. Given equation of line is
 $\vec{r} = (3+t)\hat{i} + (1-t)\hat{j} + (-2-2t)\hat{k}$
 $\Rightarrow \vec{r} = 3\hat{i} + \hat{j} - 2\hat{k} + (\hat{i} - \hat{j} - 2\hat{k})t$, where $t \in \mathbb{R}$
 \therefore The line passes through $(3, 1, -2)$ and is parallel to the vector $\hat{i} - \hat{j} - 2\hat{k}$.
 Equation of second line is
 $x = 4 + k, y = -k, z = -4 - 2k$,
 $\Rightarrow \frac{x-4}{1} = \frac{y}{-1} = \frac{z+4}{-2} = k$, where $k \in \mathbb{R}$
 \therefore d.r.s of the line are $1, -1, -2$. Also, it passes through $(3, 1, -2)$.
 \therefore Both lines are coincident.

27. Consider option (A)
 point $\left(21, \frac{5}{3}, \frac{10}{3}\right)$ satisfies both the equations of line
 \therefore option (A) is correct answer
Alternate method:
 Let $\frac{x-5}{3} = \frac{y-7}{-1} = \frac{z+2}{1} = \lambda$
 and $\frac{x+3}{-36} = \frac{y-3}{2} = \frac{z-6}{4} = \mu$
 $\therefore x = 3\lambda + 5, y = -\lambda + 7, z = \lambda - 2$ and
 $x = -36\mu - 3, y = 2\mu + 3, z = 4\mu + 6$
 On solving, we get $x = 21, y = \frac{5}{3}, z = \frac{10}{3}$

28. Consider option (B)
 Point $(-2, -4, -5)$ satisfies both the equations of the line.
 \therefore Option (B) is the correct answer.
29. Consider option (B)
 point $(-11, -4, 5)$ satisfies both the equations of line
 \therefore option (B) is correct answer
30. Consider option (B)
 point $(2, 3, 4)$ satisfies both the equations of line
 \therefore option (B) is correct answer
- 31.



- Let the two lines be AB and CD having equations $\frac{x}{1} = \frac{y+a}{1} = \frac{z}{1} = \lambda$ and
 $\frac{x+a}{2} = \frac{y}{1} = \frac{z}{1} = \mu$.
 Then, $P \equiv (\lambda, \lambda - a, \lambda)$ and $Q \equiv (2\mu - a, \mu, \mu)$
 According to the given condition,
 $\frac{\lambda - 2\mu + a}{2} = \frac{\lambda - a - \mu}{1} = \frac{\lambda - \mu}{2}$
 $\Rightarrow \mu = a$ and $\lambda = 3a$
 $\therefore P \equiv (3a, 2a, 3a)$ and $Q \equiv (a, a, a)$
32. d.r.s of the line joining $(0, -11, 4)$ and $(2, -3, 1)$ are $2, 8, -3$.
 \therefore Equation of line is $\frac{x}{2} = \frac{y+11}{8} = \frac{z-4}{-3}$



$$\text{Let } \frac{x}{2} = \frac{y+11}{8} = \frac{z-4}{-3} = \lambda$$

Any general point on this line is

$$M \equiv (2\lambda, 8\lambda - 11, -3\lambda + 4)$$

$$\text{Let } A \equiv (1, 8, 4)$$

d.r.s. of AM are $2\lambda - 1, 8\lambda - 19, -3\lambda$

Since, AM is perpendicular to the given line,


$$\therefore 2(2\lambda - 1) + 8(8\lambda - 19) - 3(-3\lambda) = 0$$

$$\Rightarrow 77\lambda = 154$$

$$\Rightarrow \lambda = 2$$

$$\therefore M \equiv (4, 5, -2)$$

$$33. \text{ Let } \frac{x+1}{3} = \frac{y-2}{-2} = \frac{z+1}{-1} = \lambda$$



Any general point on this line is

$$B(3\lambda - 1, -2\lambda + 2, -\lambda - 1)$$

$$\text{Let } A \equiv (1, 0, 2)$$

$$\therefore \text{d.r.s. of AB are } 3\lambda - 2, -2\lambda + 2, -\lambda - 3$$

Since, AB is perpendicular to the given line,

$$\therefore 3(3\lambda - 2) - 2(-2\lambda + 2) - 1(-\lambda - 3) = 0$$

$$\Rightarrow 14\lambda = 7$$

$$\Rightarrow \lambda = \frac{1}{2}$$

$$\therefore B \equiv \left(\frac{1}{2}, 1, \frac{-3}{2}\right)$$

34. Let M be the foot of perpendicular drawn from the point P(1, 2, 3) to the line

$$\text{and } \frac{x-6}{3} = \frac{y-7}{2} = \frac{z-7}{-2} = \lambda$$

\therefore The co-ordinates of any point on the line are

$$M \equiv (3\lambda + 6, 2\lambda + 7, -2\lambda + 7)$$

\therefore The d.r.s of PM are

$$3\lambda + 6 - 1, 2\lambda + 7 - 2, -2\lambda + 7 - 3$$

$$\text{i.e., } 3\lambda + 5, 2\lambda + 5, -2\lambda + 4.$$

Since, PM is perpendicular to the given line whose d.r.s. are 3, 2, -2,

$$\therefore 3(3\lambda + 5) + 2(2\lambda + 5) - 2(-2\lambda + 4) = 0$$

$$\Rightarrow 9\lambda + 15 + 4\lambda + 10 + 4\lambda - 8 = 0$$

$$\Rightarrow 17\lambda + 17 = 0$$

$$\Rightarrow \lambda = -1$$

$$\therefore M \equiv (3, 5, 9)$$

$$\therefore PM = \sqrt{(3-1)^2 + (5-2)^2 + (9-3)^2} \\ = \sqrt{4+9+36} = 7$$

35. Since the point is (-2, 4, -5),

$$\therefore a = -2, b = 4, c = -5$$

Given equation of line is

$$\frac{x+3}{3} = \frac{y-4}{5} = \frac{z+8}{6}$$

$$\therefore x_1 = -3, y_1 = 4, z_1 = -8$$

d.r.s of the line are 3, 5, 6

$$\therefore \text{d.c.s are } \frac{3}{\sqrt{70}}, \frac{5}{\sqrt{70}}, \frac{6}{\sqrt{70}}$$

Perpendicular distance of point from the line is

$$\frac{\sqrt{[(a-x_1)^2 + (b-y_1)^2 + (c-z_1)^2]}}{\sqrt{[(a-x_1) + (b-y_1)m + (c-z_1)n]^2}}$$

$$= \frac{\sqrt{1^2 + 0 + 3^2 - \left[\frac{3(1)}{\sqrt{70}} + \frac{0(5)}{\sqrt{70}} + \frac{3(6)}{\sqrt{70}}\right]^2}}{\sqrt{1+9 - \left(\frac{3}{\sqrt{70}} + \frac{18}{\sqrt{70}}\right)^2}}$$

$$= \frac{\sqrt{1+9 - \left(\frac{3}{\sqrt{70}} + \frac{18}{\sqrt{70}}\right)^2}}{\sqrt{10}}$$

$$= \frac{\sqrt{37}}{\sqrt{10}} \text{ units}$$

36. Let M be the foot of perpendicular drawn from the point P(2, 3, 4) to the line

$$\text{and } \frac{x-1}{-1} = \frac{y-0}{2} = \frac{z+1}{3} = \lambda$$

$$\therefore M \equiv (-\lambda + 1, 2\lambda, 3\lambda - 1).$$

The d.r.s of PM are $-\lambda - 1, 2\lambda - 3, 3\lambda - 5$

Since, PM is perpendicular to the given line,

$$-1(-\lambda - 1) + 2(2\lambda - 3) + 3(3\lambda - 5) = 0$$

$$\Rightarrow \lambda + 1 + 4\lambda - 6 + 9\lambda - 15 = 0$$

$$\Rightarrow 14\lambda = 20$$

$$\Rightarrow \lambda = \frac{10}{7}$$

$$\therefore M \equiv \left(\frac{-3}{7}, \frac{20}{7}, \frac{23}{7}\right)$$



$$\begin{aligned} \therefore PM &= \sqrt{\left(2 + \frac{3}{7}\right)^2 + \left(3 - \frac{20}{7}\right)^2 + \left(4 - \frac{23}{7}\right)^2} \\ &= \sqrt{\frac{289}{49} + \frac{1}{49} + \frac{25}{49}} \\ &= \frac{3}{7}\sqrt{35} \end{aligned}$$

37. The equation of the line joining the points $(-9, 4, 5)$ and $(11, 0, -1)$ is

$$\begin{aligned} \frac{x+9}{11+9} &= \frac{y-4}{0-4} = \frac{z-5}{-1-5} \\ \Rightarrow \frac{x+9}{20} &= \frac{y-4}{-4} = \frac{z-5}{-6} \\ \Rightarrow \frac{x+9}{10} &= \frac{y-4}{-2} = \frac{z-5}{-3} \end{aligned}$$

\therefore The d.r.s. of the given line are $10, -2, 3$

$$\text{Let } \frac{x+9}{10} = \frac{y-4}{-2} = \frac{z-5}{-3} = \lambda$$

\therefore Any point on the line is

$$P \equiv (10\lambda - 9, -2\lambda + 4, -3\lambda + 5)$$

\therefore The d.r.s. of OP are

$$10\lambda - 9, -2\lambda + 4, -3\lambda + 5$$

Since, the given line is perpendicular to OP,

$$10(10\lambda - 9) - 2(-2\lambda + 4) - 3(-3\lambda + 5) = 0$$


$$\Rightarrow 100\lambda - 90 + 4\lambda - 8 + 9\lambda - 15 = 0$$

$$\Rightarrow 113\lambda = 113$$

$$\Rightarrow \lambda = 1$$

$\therefore P \equiv (1, 2, 2)$

38. Let $\frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3} = \lambda$



Any general point on this line is

$$M \equiv (\lambda, 2\lambda + 1, 3\lambda + 2)$$

$$\text{Let } A \equiv (1, 6, 3)$$

$$\text{d.r.s. of AM are } \lambda - 1, 2\lambda - 5, 3\lambda - 1$$

Since, AM is perpendicular to the given line,

$$\therefore 1(\lambda - 1) + 2(2\lambda - 5) + 3(3\lambda - 1) = 0$$

$$\Rightarrow 14\lambda = 14$$

$$\Rightarrow \lambda = 1$$

$\therefore M \equiv (1, 3, 5)$

Now, M is the midpoint of AB.

$$\therefore \left(\frac{1+x_1}{2}, \frac{6+y_1}{2}, \frac{3+z_1}{2}\right) = (1, 3, 5)$$

$$\Rightarrow x_1 = 1, y_1 = 0, z_1 = 7$$

39. Let $\frac{x-1}{2} = \frac{y+1}{-2} = \frac{z+1}{1} = \lambda$

any point on the line is

$$P \equiv (2\lambda + 1, -2\lambda - 1, \lambda - 1)$$

$$\text{Let } A \equiv (1, -1, -1)$$

$$\text{Now, } PA = 3$$

$$\therefore \sqrt{(2\lambda + 1 - 1)^2 + (-2\lambda - 1 + 1)^2 + (\lambda - 1 + 1)^2} = 3$$

$$\Rightarrow \sqrt{4\lambda^2 + 4\lambda^2 + \lambda^2} = 3$$

$$\Rightarrow 9\lambda^2 = 9$$

$$\Rightarrow \lambda = \pm 1$$

$\therefore P \equiv (3, -3, 0)$ or $P \equiv (-1, 1, -2)$

40. First line passes through $(x_1, y_1, z_1) = (3, 8, 3)$

and has d.r.s. $(a_1, b_1, c_1) = (3, -1, 1)$

Second line passes through

$(x_2, y_2, z_2) = (-3, -7, 6)$ and has d.r.s.

$(a_2, b_2, c_2) = (-3, 2, 4)$

\therefore Shortest distance (d) between them is

$$d = \frac{\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix}}{\sqrt{(b_1c_2 - b_2c_1)^2 + (c_1a_2 - c_2a_1)^2 + (a_1b_2 - a_2b_1)^2}}$$

$$= \frac{\begin{vmatrix} -6 & -15 & 3 \\ 3 & -1 & 1 \\ -3 & 2 & 4 \end{vmatrix}}{\sqrt{(-4-2)^2 + (-3-12)^2 + (6-3)^2}}$$

$$= \frac{-6(-4-2) + 15(12+3) + 3(6-3)}{\sqrt{36 + 225 + 9}}$$

$$= \frac{270}{\sqrt{270}} = 3\sqrt{30}$$

41. Here, $(x_1, y_1, z_1) = (-1, -2, -1)$

$(x_2, y_2, z_2) = (2, -2, 3)$

$(a_1, b_1, c_1) = (3, 1, 2)$

$(a_2, b_2, c_2) = (1, 2, 3)$

$$d = \frac{\begin{vmatrix} 3 & 0 & 4 \\ 3 & 1 & 2 \\ 1 & 2 & 3 \end{vmatrix}}{\sqrt{(3-4)^2 + (2-9)^2 + (6-1)^2}}$$

$$= \frac{17}{\sqrt{75}} = \frac{17}{5\sqrt{3}}$$



42. Since, the given lines intersect each other,

$$\therefore \begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} 3-1 & k+1 & 0-1 \\ 2 & 3 & 4 \\ 1 & 2 & 1 \end{vmatrix} = 0$$

$$\Rightarrow 2(3-8) - (k+1)(2-4) - 1(4-3) = 0$$

$$\Rightarrow -10 + 2k + 2 - 1 = 0$$

$$\Rightarrow k = \frac{9}{2}$$

43. Since, the given lines intersect each other,

$$\therefore \begin{vmatrix} 2-1 & 3-2 & 1-3 \\ k & 2 & 3 \\ 3 & k & 2 \end{vmatrix} = 0$$

$$\Rightarrow 1(4-3k) - 1(2k-9) - 2(k^2-6) = 0$$

$$\Rightarrow 2k^2 + 5k - 25 = 0$$

$$\Rightarrow k = \frac{5}{2}, -5$$

44. Let the equation of a line passing through the

$$\text{origin be } \frac{x}{a} = \frac{y}{b} = \frac{z}{c}.$$

This meets the lines

$$\frac{x-2}{1} = \frac{y-1}{-2} = \frac{z+1}{1} \text{ and } \frac{x-\frac{8}{3}}{2} = \frac{y+3}{-1} = \frac{z-1}{1}$$

$$\therefore \begin{vmatrix} 2 & 1 & -1 \\ a & b & c \\ 1 & -2 & 1 \end{vmatrix} = 0 \text{ and } \begin{vmatrix} \frac{8}{3} & -3 & 1 \\ a & b & c \\ 2 & -1 & 1 \end{vmatrix} = 0$$

$$\Rightarrow a + 3b + 5c = 0 \text{ and } 3a + b - 5c = 0$$

$$\Rightarrow \frac{a}{5} = \frac{b}{-5} = \frac{c}{2}$$

Thus, the equation of the line through the origin intersecting the given lines is

$$\frac{x}{5} = \frac{y}{-5} = \frac{z}{2} = \lambda (\text{say})$$

The co-ordinates of any point on this line are $(5\lambda, -5\lambda, 2\lambda)$.

The co-ordinates of any point on

$$\frac{x-2}{1} = \frac{y-1}{-2} = \frac{z+1}{1} = \lambda_1 (\text{say}) \text{ are}$$

$$(\lambda_1 + 2, -2\lambda_1 + 1, \lambda_1 - 1).$$

If these two lines intersect, then

$$5\lambda = \lambda_1 + 2, -5\lambda = -2\lambda_1 + 1 \text{ and } 2\lambda = \lambda_1 - 1$$

$$\Rightarrow \lambda_1 = 3 \text{ and } \lambda = 1$$

So, the co-ordinates of P are $(5, -5, 2)$.

$$\text{Similarly, co-ordinates of Q are } \left(\frac{10}{3}, \frac{-10}{3}, \frac{8}{3} \right)$$

$$\therefore PQ^2 = \left(\frac{10}{3} - 5 \right)^2 + \left(\frac{-10}{3} + 5 \right)^2 + \left(\frac{8}{3} - 2 \right)^2 = 6$$

45. Lines L_1 and L_2 are parallel to the vectors $\bar{b}_1 = 3\hat{i} + \hat{j} + 2\hat{k}$ and $\bar{b}_2 = \hat{i} + 2\hat{j} + 3\hat{k}$ respectively.

\therefore The unit vector perpendicular to both L_1 and L_2 is

$$\hat{n} = \frac{\bar{b}_1 \times \bar{b}_2}{|\bar{b}_1 \times \bar{b}_2|}$$

$$\text{Now, } \bar{b}_1 \times \bar{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & 1 & 2 \\ 1 & 2 & 3 \end{vmatrix} = -\hat{i} - 7\hat{j} + 5\hat{k}$$

$$\therefore \hat{n} = \frac{1}{5\sqrt{3}} (-\hat{i} - 7\hat{j} + 5\hat{k})$$



Evaluation Test

1. Let $\frac{x+1}{-1} = \frac{y-12}{5} = \frac{z-7}{2} = r$

$$\therefore x = -r - 1, y = 5r + 12, z = 2r + 7$$

\therefore Co-ordinates of any point on the line are $(-r - 1, 5r + 12, 2r + 7)$.

This point lies on the curve $11x^2 - 5y^2 + z^2 = 0$

$$\therefore 11(-r-1)^2 - 5(5r+12)^2 + (2r+7)^2 = 0$$

$$\Rightarrow 11r^2 + 22r + 11 - 125r^2 - 600r - 720 + 4r^2 + 28r + 49 = 0$$

$$\Rightarrow -110r^2 - 550r - 660 = 0$$

$$\Rightarrow r^2 + 5r + 6 = 0$$

$$\Rightarrow (r+2)(r+3) = 0$$

$$\Rightarrow r = -2 \text{ or } r = -3$$

If $r = -2$, then the point is $(1, 2, 3)$


and if $r = -3$, then the point is $(2, -3, 1)$

\therefore option (A) is correct.



2. The given equation of line is
 $x = 4y + 5, z = 3y - 6$.
 It can be written as
 $\frac{x-5}{4} = y = \frac{z+6}{3} = r$, say
- \therefore co-ordinates of the any point on the line are
 $(4r + 5, r, 3r - 6)$.
 This point is at a distance of $3\sqrt{26}$ from the point $(5, 0, -6)$
- $\therefore (4r + 5 - 5)^2 + (r - 0)^2 + (3r - 6 + 6)^2 = (3\sqrt{26})^2$
 $\Rightarrow 16r^2 + r^2 + 9r^2 = 234$
 $\Rightarrow 26r^2 = 234$
 $\Rightarrow r^2 = 9$
 $\Rightarrow r = \pm 3$
 If $r = 3$, then the point is
 $(4 \times 3 + 5, 3, 3 \times 3 - 6) \equiv (17, 3, 3)$
3. Let the components of the line vector be a, b, c .
 $\therefore a^2 + b^2 + c^2 = (63)^2 \dots(i)$
 Also, $\frac{a}{3} = \frac{b}{-2} = \frac{c}{6} = k$, say
 $\therefore a = 3k, b = -2k, c = 6k$
 Substituting value of a, b and c in equation (i), we get
 $9k^2 + 4k^2 + 36k^2 = 63^2$
 $\therefore 49k^2 = 63 \times 63$
 $\therefore k^2 = \frac{63 \times 63}{49} = 81$
 $\therefore k = \pm 9$
 Since, the line makes obtuse angle with X-axis component along X-axis is negative.
 $\therefore k = -9$
 \therefore The components of the line vector are $3k, -2k, 6k$ i.e., $-27, 18, -54$
4. Let M be the foot of the perpendicular drawn from the point $P(3, -1, 11)$ to the given line.
 Let $\frac{x}{2} = \frac{y-2}{3} = \frac{z-3}{4} = \lambda$
 $\Rightarrow x = 2\lambda, y = 3\lambda + 2, z = 4\lambda + 3$
 $\therefore M \equiv (2\lambda, 3\lambda + 2, 4\lambda + 3)$
 d.r.s. of PM are $2\lambda - 3, 3\lambda + 3, 4\lambda - 8$
 Since, PM is perpendicular to the given line
 $\therefore (2\lambda - 3)(2) + (3\lambda + 3)(3) + (4\lambda - 8)(4) = 0$
 $\Rightarrow 4\lambda - 6 + 9\lambda + 9 + 16\lambda - 32 = 0$
 $\Rightarrow \lambda = 1$
 $\therefore M \equiv (2, 5, 7)$

- \therefore length of perpendicular (PM)
 $= \sqrt{(3-2)^2 + (-1-5)^2 + (11-7)^2}$
 $= \sqrt{1+36+16}$
 $= \sqrt{53}$
5. When square is folded co-ordinates will be
 $D(0, 0, a), C(a, 0, 0), A(-a, 0, 0), B(0, -a, 0)$.



Equation AB is, $\frac{x+a}{a} = \frac{y}{-a} = \frac{z}{0}$

and equation of DC is $\frac{x}{a} = \frac{y}{0} = \frac{z-a}{-a}$

- \therefore shortest distance

$$= \frac{\begin{vmatrix} -a & 0 & -a \\ a & -a & 0 \\ a & 0 & -a \end{vmatrix}}{\sqrt{(a^2 - 0)^2 + (0 + a^2)^2 + (0 + a^2)^2}}$$

$$= \frac{-a(a^2) - a(a^2)}{\sqrt{a^4 + a^4 + a^4}} = \frac{-2a^3}{\sqrt{3a^4}} = \frac{2a}{\sqrt{3}}$$

6. Given equation of motion of a rocket is
 $x = 2t, y = -4t, z = 4t$
 i.e., the equation of the path is $\frac{x}{2} = \frac{y}{-4} = \frac{z}{4}$
 i.e., $\frac{x}{1} = \frac{y}{-2} = \frac{z}{2}$
 Thus, the path of the rocket represents a straight line passing through the origin.
 For $t = 10$ sec.
 we have, $x = 20, y = -40, z = 40$
 Let $M(20, -40, 40)$
- $\therefore |\overline{OM}| = \sqrt{x^2 + y^2 + z^2}$
 $= \sqrt{400 + 1600 + 1600} = 60$ km
- \therefore Rocket will be at 60 km from the starting point $O(0, 0, 0)$ in 10 seconds.



7. d.r.s. of L_1 are 3, 1, 2 and d.r.s. of L_2 are 1, 2, 3


$$\therefore \text{vector perpendicular to } L_1 \text{ and } L_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & 1 & 2 \\ 1 & 2 & 3 \end{vmatrix}$$

$$= \hat{i}(3-4) - \hat{j}(9-2) + \hat{k}(6-1)$$

$$= -\hat{i} - 7\hat{j} + 5\hat{k}$$

$$\therefore \text{unit vector} = \frac{-\hat{i} - 7\hat{j} + 5\hat{k}}{\sqrt{1+49+25}} = \frac{-\hat{i} - 7\hat{j} + 5\hat{k}}{5\sqrt{3}}$$

8. Let S be the foot of perpendicular drawn from P(1, 0, 3) to the join of points A(4, 7, 1) and B(3, 5, 3)



Let S divide AB in the ratio $\lambda : 1$

$$\therefore S \equiv \left(\frac{3\lambda + 4}{\lambda + 1}, \frac{5\lambda + 7}{\lambda + 1}, \frac{3\lambda + 1}{\lambda + 1} \right) \dots (i)$$

Now, d.r.s. of PS are

$$\frac{3\lambda + 4}{\lambda + 1} - 1, \frac{5\lambda + 7}{\lambda + 1} - 0, \frac{3\lambda + 1}{\lambda + 1} - 3$$

$$\text{i.e., } \frac{2\lambda + 3}{\lambda + 1}, \frac{5\lambda + 7}{\lambda + 1}, \frac{-2}{\lambda + 1}$$

$$\text{i.e., } 2\lambda + 3, 5\lambda + 7, -2$$

Also, d.r.s. of AB are -1, -2, 2

Since, $PS \perp AB$

$$\therefore (2\lambda + 3)(-1) + (5\lambda + 7)(-2) + (-2)(2) = 0$$

$$\Rightarrow -2\lambda - 3 - 10\lambda - 14 - 4 = 0$$

$$\Rightarrow \lambda = -\frac{7}{4}$$

Substituting the value of λ in (i), we get

$$S = \left(\frac{5}{3}, \frac{7}{3}, \frac{17}{3} \right)$$

9. Equation of the line passing through the points (5, 1, a) and (3, b, 1) is

$$\frac{x-3}{5-3} = \frac{y-b}{1-b} = \frac{z-1}{a-1} \dots (i)$$

The line passes through the point $\left(0, \frac{17}{2}, \frac{-13}{2} \right)$

$$\therefore \frac{-3}{2} = \frac{\frac{17}{2} - b}{1-b} = \frac{\frac{-13}{2} - 1}{a-1} \dots [\text{From (i)}]$$

$$\therefore a - 1 = \frac{-15}{\frac{-3}{2}} = 5$$

$$\therefore a = 5 + 1 = 6$$

$$\text{and } -3 + 3b = 17 - 2b$$

$$\therefore 5b = 20 \Rightarrow b = 4$$

$$\therefore a = 6, b = 4$$

08 Plane



Hints



Classical Thinking

- Here, $\vec{n} = \hat{i} - 2\hat{j} + 3\hat{k}$ and $p = 1$

$$\hat{n} = \frac{\vec{n}}{|\vec{n}|} = \frac{\hat{i} - 2\hat{j} + 3\hat{k}}{\sqrt{1+4+9}} = \frac{\hat{i} - 2\hat{j} + 3\hat{k}}{\sqrt{14}}$$
 \therefore The vector equation of the plane is
 $\vec{r} \cdot \hat{n} = p$

$$\Rightarrow \vec{r} \cdot \left(\frac{\hat{i} - 2\hat{j} + 3\hat{k}}{\sqrt{14}} \right) = 1$$

$$\Rightarrow \vec{r} \cdot (\hat{i} - 2\hat{j} + 3\hat{k}) = \sqrt{14}$$
- The given vector equation is
 $\vec{r} \cdot (3\hat{i} - 2\hat{j} + 2\hat{k}) = 12 \quad \dots(i)$
 $\vec{r} \cdot \vec{n} = 12$, where $\vec{n} = 3\hat{i} - 2\hat{j} + 2\hat{k}$

$$\hat{n} = \frac{\vec{n}}{|\vec{n}|} = \frac{3\hat{i} - 2\hat{j} + 2\hat{k}}{\sqrt{9+4+4}} = \frac{3\hat{i} - 2\hat{j} + 2\hat{k}}{\sqrt{17}}$$

$$\Rightarrow \hat{n} = \frac{3}{\sqrt{17}}\hat{i} - \frac{2}{\sqrt{17}}\hat{j} + \frac{2}{\sqrt{17}}\hat{k}$$
 \therefore Normal form is

$$\vec{r} \cdot \left(\frac{3}{\sqrt{17}}\hat{i} - \frac{2}{\sqrt{17}}\hat{j} + \frac{2}{\sqrt{17}}\hat{k} \right) = \frac{12}{\sqrt{17}}$$
- Given equation of plane is
 $\vec{r} \cdot (2\hat{i} - 3\hat{j} + \hat{k}) + 9 = 0$

$$\Rightarrow \vec{r} \cdot (2\hat{i} - 3\hat{j} + \hat{k}) = -9 \quad \dots(i)$$

 $\vec{n} = 2\hat{i} - 3\hat{j} + \hat{k}$

$$\hat{n} = \frac{2\hat{i} - 3\hat{j} + \hat{k}}{\sqrt{4+9+1}} = \frac{2\hat{i} - 3\hat{j} + \hat{k}}{\sqrt{14}}$$
 \therefore The d.c.s. of normal to the plane are
 $\frac{2}{\sqrt{14}}, \frac{-3}{\sqrt{14}}, \frac{1}{\sqrt{14}}$
- Given that $x + my + nz = p$ is the equation of the plane in normal form.
 \therefore , m, n are the direction cosines.
 Also $2^2 + m^2 + n^2 = 1$,

- Since, p is the distance from the origin, p should be greater than zero.
- \therefore
- All the statements are true,
-
- \therefore
- correct answer is option (D)
- Equation of XY plane is $z = 0$,
 \therefore d.c.s. of its normal are 0, 0, 1
 - $$\frac{x}{7} + \frac{y}{7} + \frac{z}{7} = 1$$

 a
 For equal intercepts, $\frac{7}{a} = 7 \Rightarrow a = 1$
 - Equation of plane in intercept form is

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

 Here, $a = b = c$ and point $(1, -1, 2)$ lies in the plane,
 $\therefore \frac{1}{a} + \frac{-1}{a} + \frac{2}{a} = 1 \Rightarrow a = 2$
 \therefore the required equation of a plane is $x + y + z = 2$.
 - Here, $\vec{a} = \hat{i} + \hat{j} + 2\hat{k}$ and $\vec{n} = 3\hat{i} - 2\hat{j} + 3\hat{k}$
 The vector equation of the plane is
 $\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$

$$\Rightarrow \vec{r} \cdot (3\hat{i} - 2\hat{j} + 3\hat{k}) = (\hat{i} + \hat{j} + 2\hat{k}) \cdot (3\hat{i} - 2\hat{j} + 3\hat{k})$$

$$\Rightarrow \vec{r} \cdot (3\hat{i} - 2\hat{j} + 3\hat{k}) = 7$$
 - Let $\vec{a} = \hat{j} - 3\hat{k}$ and $\vec{n} = \hat{i} + 2\hat{j} + 4\hat{k}$
 The vector equation of plane is

$$\vec{r} \cdot (\hat{i} + 2\hat{j} + 4\hat{k}) = (\hat{j} - 3\hat{k}) \cdot (\hat{i} + 2\hat{j} + 4\hat{k})$$

$$\Rightarrow \vec{r} \cdot (\hat{i} + 2\hat{j} + 4\hat{k}) = -10$$
 - The plane passes through $(2, -1, 1)$
 This point satisfies the equation of plane in option (D)
 Also, it has d.r.s. 1, 1, -2.
 \therefore option (D) is correct answer.

**Alternate method:**Let $A \equiv (2, -1, 1)$ The d.r.s. of line joining the points $(2, 3, -1)$ and $(1, 2, 1)$ are $1, 1, -2$ \therefore the equation of the required plane is

$$1(x-2) + 1(y+1) - 2(z-1) = 0$$

$$\Rightarrow x + y - 2z + 1 = 0$$

12. The plane passes through $(3, 2, -1)$
This point satisfies the equation of plane in option (C).

Also, it has d.r.s. $2, 2, -3$ \therefore option (C) is correct answer.

13. The plane passes through $(-10, 5, 4)$
This point satisfies the equation of plane in option (B)

Also, it has d.r.s. $7, -3, -1$ \therefore option (B) is correct answer.

14. The plane passes through $(1, 2, -3)$
This point satisfies the equation of plane in option (A)

Also, it has d.r.s. $1, 2, -3$. \therefore option (A) is correct answer.**Alternate method:**Let $M(1, 2, -3)$ be the foot of perpendicular from the origin $O(0, 0, 0)$ to the plane
D. r. s of normal are $1, 2, -3$ \therefore the equation of the required plane is

$$1(x-1) + 2(y-2) - 3(z+3) = 0$$

$$\Rightarrow x + 2y - 3z = 14$$

15. The plane passes through $(2, 4, -3)$
This point satisfies the equation of plane in option (C)

Also, it has d.r.s. $2, 4, -3$. \therefore option (C) is correct answer.

16. The plane passes through $(1, -1, 1)$
This point satisfies the equation of plane in option (D)

$$\text{Also, it has d.r.s} = \vec{b} \times \vec{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & 1 \\ 0 & 1 & 2 \end{vmatrix}$$

$$= \hat{i}(2-1) - \hat{j}(4-0) + \hat{k}(2-0)$$

$$= \hat{i} - 4\hat{j} + 2\hat{k}$$

i.e., $1, -4, 2$ \therefore option (D) is correct answer.**Alternate Method**Let $\vec{a} = \hat{i} - \hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} + \hat{j} + \hat{k}$ and $\vec{c} = \hat{j} + 2\hat{k}$

Now, $\vec{b} \times \vec{c} = \hat{i} - 4\hat{j} + 2\hat{k}$

 \therefore the vector equation of required plane is

$$\vec{r} \cdot (\vec{b} \times \vec{c}) = \vec{a} \cdot (\vec{b} \times \vec{c})$$

$$\Rightarrow \vec{r} \cdot (\hat{i} - 4\hat{j} + 2\hat{k}) = (\hat{i} - \hat{j} + \hat{k}) \cdot (\hat{i} - 4\hat{j} + 2\hat{k})$$

$$\Rightarrow \vec{r} \cdot (\hat{i} - 4\hat{j} + 2\hat{k}) = 7$$

17. Let
- $(x_1, y_1, z_1) = (0, 1, 2)$
- ,

 $a_1, b_1, c_1 = 3, 1, 1$ and $a_2, b_2, c_2 = -1, 2, -5$ \therefore the equation of required plane is

$$\begin{vmatrix} x-x_1 & y-y_1 & z-z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} x-0 & y-1 & z-2 \\ 3 & 1 & 1 \\ -1 & 2 & -5 \end{vmatrix} = 0$$

$$\Rightarrow -7x + 14y - 14 + 7z - 14 = 0$$

$$\Rightarrow x - 2y - z + 4 = 0$$

18. Let
- $(x_1, y_1, z_1) = (1, 2, -1)$
- ,

 $a_1, b_1, c_1 = 2, 1, 3$ and $a_2, b_2, c_2 = 4, 1, 2$ \therefore the equation of required plane is

$$\begin{vmatrix} x-x_1 & y-y_1 & z-z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

$$\Rightarrow (x-1)(-2) + (y-2)(10) + (z+1)(-2) = 0$$

$$\Rightarrow -2x + 2 + 10y - 20 - 2z - 2 = 0$$

$$\Rightarrow x - 5y + z + 10 = 0$$

19. Required plane passes through point
- $(x_1, y_1, z_1) \equiv (1, -3, -2)$
- and is perpendicular to planes
- $x + 2y + 2z = 5$
- and
- $3x + 3y + 2z = 8$

 \therefore their normals are parallel to the required plane $\therefore a_1, b_1, c_1 = 1, 2, 2$ and $a_2, b_2, c_2 = 3, 3, 2$ \therefore the equation of required plane is

$$\begin{vmatrix} x-x_1 & y-y_1 & z-z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

$$\Rightarrow 2x - 4y + 3z - 8 = 0$$

20. The equation
- $\vec{r} = \vec{a} + \lambda\vec{b} + \mu\vec{c}$
- represents a plane passing through vector
- \vec{a}
- and parallel to
- \vec{b}
- and
- \vec{c}

 $\therefore \vec{a} = 3\hat{i} + \hat{j}$, $\vec{b} = -\hat{j} + \hat{k}$, $\vec{c} = \hat{i} + 2\hat{j} + 3\hat{k}$



$$\begin{aligned} \text{Now, } \vec{b} \times \vec{c} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & -1 & 1 \\ 1 & 2 & 3 \end{vmatrix} \\ &= -5\hat{i} + \hat{j} + \hat{k} \end{aligned}$$

- \therefore the equation of required plane is
 $\vec{r} \cdot (-5\hat{i} + \hat{j} + \hat{k}) = (3\hat{i} + \hat{j}) \cdot (-5\hat{i} + \hat{j} + \hat{k})$
 $\Rightarrow \vec{r} \cdot (-5\hat{i} + \hat{j} + \hat{k}) = -14$
21. Consider option (B)
 $\vec{r} \cdot (\hat{i} + 11\hat{j} + 3\hat{k}) = 14$
 Its Cartesian form is
 $x + 11y + 3z = 14$
 Since, the given points (1, 2, -3), (3, 1, 0) and (0, 1, 1) satisfy the above plane,
 \therefore correct answer is option (B)
Alternate method:
 Equation of a plane passing through three points is

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ x_3 - x_1 & y_3 - y_1 & z_3 - z_1 \end{vmatrix} = 0$$

 $\Rightarrow \begin{vmatrix} x - 1 & y - 2 & z + 3 \\ 2 & -1 & 3 \\ -1 & -1 & 4 \end{vmatrix} = 0$
 $\Rightarrow (x - 1)(-1) - (y - 2)(11) + (z + 3)(-3) = 0$
 $\Rightarrow -x - 11y - 3z + 14 = 0$
 $\Rightarrow x + 11y + 3z = 14$
 Its vector form is
 $\vec{r} \cdot (\hat{i} + 11\hat{j} + 3\hat{k}) = 14$
22. Consider option (B)
 $\vec{r} \cdot (3\hat{i} + \hat{j} - \hat{k}) + 4 = 0$
 Its Cartesian form is
 $3x + y - z = -4$
 Since the given points A(1, -2, 5), B(0, -5, -1) and C(-3, 5, 0) satisfy the above plane,
 \therefore correct answer is option (B).
23. Consider option (B)
 $\vec{r} \cdot (9\hat{i} + 3\hat{j} - \hat{k}) = 14$
 Its Cartesian form is
 $9x + 3y - z = 14$
 Since, given points (1, 1, -2), (2, -1, 1) and (1, 2, 1) satisfy the above plane,
 \therefore correct answer is option (B)

24. Consider option (D)
 $2x + 2y - 5z = 0$
 Since, the given points (4, 1, 2), (1, -1, 0) and (0, 0, 0) satisfy the above plane,
 \therefore correct answer is option (D)
25. Consider option (C)
 $3x - 4z + 1 = 0$
 Since, the given points (1, 1, 1), (1, -1, 1) and (-7, -3, -5) satisfy the above plane,
 \therefore correct answer is option (C)
26. Here $\vec{n}_1 = \hat{i} - \hat{j} + 2\hat{k}$ and $\vec{n}_2 = 3\hat{i} - \hat{j} - \hat{k}$
 The vector equation of plane passing through intersection of $\vec{r} \cdot \vec{n}_1 = p_1$ and $\vec{r} \cdot \vec{n}_2 = p_2$ is
 $\vec{r} \cdot (\vec{n}_1 + \lambda \vec{n}_2) = p_1 + \lambda p_2$
 $\Rightarrow \vec{r} \cdot [\hat{i} - \hat{j} + 2\hat{k} + \lambda(3\hat{i} - \hat{j} - \hat{k})] = 3 + \lambda(4)$
 $\Rightarrow \vec{r} \cdot [(1 + 3\lambda)\hat{i} - (1 + \lambda)\hat{j} + (2 - \lambda)\hat{k}] = 3 + 4\lambda$
27. Consider option (B)
 $\vec{r} \cdot (10\hat{i} + 11\hat{j} + 12\hat{k}) = 33$
 Its Cartesian form is
 $10x + 11y + 12z = 33$
 Since, the given point (1, 1, 1) satisfies the above plane
 \therefore correct answer is option (B)
Alternate method:
 The equation of plane through the intersection of given planes is
 $(x + y + z - 4) + \lambda(x + 2y + 3z + 3) = 0$
 Since, it passes through (1, 1, 1)
 $\therefore (1 + 1 + 1 - 4) + \lambda(1 + 2 + 3 + 3) = 0 \Rightarrow \lambda = \frac{1}{9}$
 $\Rightarrow (x + y + z - 4) + \frac{1}{9}(x + 2y + 3z + 3) = 0$
 $\Rightarrow 10x + 11y + 12z - 33 = 0$
 \therefore the equation of plane in vector form is
 $\vec{r} \cdot (10\hat{i} + 11\hat{j} + 12\hat{k}) = 33$
28. Consider option (D)
 $\vec{r} \cdot (11\hat{i} + 3\hat{j} - 5\hat{k}) = 22$
 Its Cartesian form is
 $11x + 3y - 5z = 22$
 Since, the given point (1, 2, -1) satisfies the above plane,
 \therefore correct answer is option (D)



29. Equation of plane passing through intersection of given planes is,
 $(x + y + z - 1) + \lambda(2x + 3y - z + 4) = 0$
 $(1 + 2\lambda)x + (1 + 3\lambda)y + (1 - \lambda)z + 4\lambda - 1 = 0$
 Since, the plane is parallel to X-axis,
 $\therefore (1 + 2\lambda) = 0 \Rightarrow \lambda = -\frac{1}{2}$
 Hence, the equation of required plane is
 $y - 3z + 6 = 0$
30. Plane passes through (1, 2, 3)
 The point (1, 2, 3) satisfies the equation of plane represented by option (B)
 \therefore option (B) is correct
Alternate method:
 Any plane parallel to $2x + 4y + 2z = 5$ is
 $2x + 4y + 2z = k$
 It passes through (1, 2, 3) $\Rightarrow k = 16$
 \therefore Equation of plane is $x + 2y + z = 8$
31. Plane passes through (0, 0, 0)
 The point (0, 0, 0) satisfies the equation of plane represented by option (A)
 \therefore option (A) is correct.
32. Equation of plane parallel to ZX-plane is $y = b$.
 It passes through (0, 2, 0)
 \therefore its equation is $y = 2$.
33. Equation of plane parallel to YZ-plane is $x = a$
 Since, it passes through (-1, 3, 4)
 \therefore equation of required plane is $x = -1$
 i.e., $x + 1 = 0$
34. Since, the plane is parallel to X-axis,
 \therefore the d.r.s. of the normal to the plane are 0, b, c
 \therefore The equation of required plane is $by + cz + d = 0$
35. Since, the plane is parallel to $ax + by + cz = 0$, their d.r.s will be same and
 It passes through (α, β, γ)
 \therefore The equation of the plane is
 $a(x - \alpha) + b(y - \beta) + c(z - \gamma) = 0$
 $\Rightarrow ax + by + cz = a\alpha + b\beta + c\gamma$
36. Equation of the plane through the origin is
 $ax + by + cz = 0$
 The required plane passes through the line
 $\frac{x-1}{5} = \frac{y-2}{4} = \frac{z-3}{5}$
 $\therefore 5a + 4b + 5c = 0 \quad \dots(i)$
 The plane passes through the point (1, 2, 3)
 $\therefore a + 2b + 3c = 0 \quad \dots(ii)$
- Solving (i) and (ii), we get
 $\therefore \frac{a}{12-10} = \frac{b}{5-15} = \frac{c}{10-4}$
 $\Rightarrow \frac{a}{1} = \frac{b}{-5} = \frac{c}{3}$
 \therefore The equation of the required plane is
 $x - 5y + 3z = 0$
37. Since, line is perpendicular to the plane
 \therefore d.r.s. of the line are a, b, c
 It passes through (α, β, γ)
 \therefore equation of perpendicular is
 $\frac{x-\alpha}{a} = \frac{y-\beta}{b} = \frac{z-\gamma}{c}$
38. Since, line is perpendicular to the plane
 \therefore d.r.s. of the line are 2, -3, 1
 It passes through (1, 1, 1)
 \therefore the equation of required line is
 $\frac{x-1}{2} = \frac{y-1}{-3} = \frac{z-1}{1}$
39. Since, line is perpendicular to the plane
 \therefore d.r.s. of the line are 1, -2, -3
 It passes through (1, 1, -1)
 \therefore the equation of required line is
 $\frac{x-1}{1} = \frac{y-1}{-2} = \frac{z+1}{-3}$
40. D.r.s of line perpendicular to YZ-plane are
 1, 0, 0
 It passes through (1, 2, 3)
 \therefore equation of required line is
 $\frac{x-1}{1} = \frac{y-2}{0} = \frac{z-3}{0}$
41. D.r.s of the normal to the XZ plane are a, 0, c
 The required line passes through (1, 2, 3)
 \therefore The equation of required line is
 $\frac{x-1}{a} = \frac{y-2}{0} = \frac{z-3}{c}$
42. Equation of line passing through point (1, 1, 1)
 is
 $\frac{x-1}{a} = \frac{y-1}{b} = \frac{z-1}{c}$
 Also, the line is parallel to the plane
 $2x + 3y + z + 5 = 0$
 $\therefore 2a + 3b + c = 0$
 The above equation is satisfied by -1, 1, -1
 \therefore correct answer is option (A)



43. The line $\frac{x-4}{1} = \frac{y-2}{1} = \frac{z-k}{2}$ lies in the plane $2x - 4y + z = 7$.

\therefore the point $(4, 2, k)$ lies on the line and hence lies in the plane

$$\therefore 2(4) - 4(2) + k = 7 \\ \Rightarrow k = 7$$

44. $\vec{n}_1 = 2\hat{i} - \hat{j} + \hat{k}$ and $\vec{n}_2 = \hat{i} + \hat{j} + 2\hat{k}$

$$\therefore \cos\theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| |\vec{n}_2|} \\ = \frac{|2(1) - 1(1) + 1(2)|}{\sqrt{4+1+1} \sqrt{1+1+4}} = \frac{1}{2} \\ \Rightarrow \theta = \frac{\pi}{3}$$

45. Let $a_1, b_1, c_1 = 1, 2, -3$ and $a_2, b_2, c_2 = 4, 1, 2$

\therefore The angle between the planes is

$$\cos\theta = \frac{|a_1 a_2 + b_1 b_2 + c_1 c_2|}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}} \\ = \frac{|1(4) + 2(1) + (-3)(2)|}{\sqrt{1+4+9} \sqrt{16+1+4}} = 0 \\ \Rightarrow \theta = \frac{\pi}{2}$$

48. The d.r.s. of normal to first plane are a, b, c and the d.r.s. of normal to second plane are a', b', c'

Since the two planes are perpendicular,

$$\therefore aa' + bb' + cc' = 0$$

49. The d.r.s. of the normal to the plane are $0, 2, 3$.

The d.r.s. of X axis are $1, 0, 0$

$$\text{Now, } a_1 a_2 + b_1 b_2 + c_1 c_2 = 0(1) + 2(0) + 3(0) \\ = 0$$

\therefore The plane $2y + 3z = 0$ passes through X-axis.

50. Comparing the equations of line and plane with $\vec{r} = \vec{a} + \lambda \vec{b}$ and $\vec{r} \cdot \vec{n} = p$, we get

$$\vec{b} = \hat{i} + 2\hat{j} - \hat{k} \text{ and } \vec{n} = 2\hat{i} - \hat{j} + \hat{k}$$

\therefore The angle between the line and plane is

$$\sin\theta = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| |\vec{n}|} \\ = \frac{|1(2) + 2(-1) - 1(1)|}{\sqrt{1+4+1} \sqrt{4+1+1}} = \frac{1}{6} \\ \Rightarrow \theta = \sin^{-1}\left(\frac{1}{6}\right)$$

51. Here, $\vec{b} = \hat{i} - \hat{j} + \hat{k}$ and $\vec{n} = 3\hat{i} - 4\hat{k}$

\therefore Angle between the line and plane is

$$\sin\theta = \frac{|(\hat{i} - \hat{j} + \hat{k}) \cdot (3\hat{i} - 4\hat{k})|}{|\sqrt{1+1+1}| \cdot |\sqrt{9+16}|} = \frac{|-1|}{|5\sqrt{3}|} \\ \Rightarrow \theta = \sin^{-1}\left(\frac{1}{5\sqrt{3}}\right)$$

52. Let $a, b, c = 2, 3, 4$ and $a_1, b_1, c_1 = 3, 2, -3$

$$\therefore \sin\theta = \frac{aa_1 + bb_1 + cc_1}{\sqrt{a^2 + b^2 + c^2} \sqrt{a_1^2 + b_1^2 + c_1^2}} \\ = \frac{2(3) + 3(2) + 4(-3)}{\sqrt{2^2 + 3^2 + 4^2} \sqrt{(3)^2 + (2)^2 + (-3)^2}} \\ \Rightarrow \sin\theta = 0 \\ \Rightarrow \theta = 0^\circ$$

53. Let $a, b, c = 3, 2, 4$ and $a_1, b_1, c_1 = 2, 1, -3$

$$\therefore \sin\theta = \frac{6 + 2 - 12}{\sqrt{9+4+16} \sqrt{4+1+9}} \\ \Rightarrow \sin\theta = \frac{-4}{\sqrt{29 \times 14}} = \frac{-4}{\sqrt{406}} \\ \Rightarrow \theta = \sin^{-1}\left(\frac{-4}{\sqrt{406}}\right)$$

54. The d.r.s. of line and plane are a, b, c

$$\therefore \sin\theta = \frac{aa + bb + cc}{\sqrt{a^2 + b^2 + c^2} \cdot \sqrt{a^2 + b^2 + c^2}} \\ = \frac{a^2 + b^2 + c^2}{a^2 + b^2 + c^2} = 1 \\ \Rightarrow \theta = 90^\circ$$

55. Given equation of line is $6x = 4y = 3z$

$$\text{i.e. } \frac{x}{2} = \frac{y}{3} = \frac{z}{4}$$

\therefore the d.r.s. of line are $2, 3, 4$

the d.r.s. of plane are $3, 2, -3$

$$\therefore \sin\theta = \frac{2(3) + 3(2) + 4(-3)}{\sqrt{4+9+16} \cdot \sqrt{9+4+9}} = 0 \\ \Rightarrow \theta = 90^\circ$$

57. Since the line $\vec{r} = \hat{i} + \lambda(2\hat{i} - m\hat{j} - 3\hat{k})$ is parallel to the plane $\vec{r} \cdot (m\hat{i} + 3\hat{j} + \hat{k}) = 0$

$$\therefore \vec{b} \cdot \vec{n} = 0 \\ \Rightarrow (2\hat{i} - m\hat{j} - 3\hat{k}) \cdot (m\hat{i} + 3\hat{j} + \hat{k}) = 0 \\ \Rightarrow 2(m) - m(3) - 3(1) = 0 \\ \Rightarrow -m = 3 \\ \Rightarrow m = -3$$



58. The line $\frac{x-1}{2} = \frac{y+1}{3} = \frac{z}{4}$ lie on the plane $4x + 4y - kz = 0$
Since the given line lies on the plane, it is parallel to the plane
 $\therefore aa_1 + bb_1 + cc_1 = 0$
 $\Rightarrow 4(2) + 4(3) - k(4) = 0$
 $\Rightarrow 4k = 20 \Rightarrow k = 5$

59. The equation of plane is $3x - 2y + 6z - 5 = 0$ and the point is $(2, 3, 4)$
 \therefore The distance of point from the plane is

$$d = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$$

$$= \frac{|2(3) + 3(-2) + 4(6) - 5|}{\sqrt{3^2 + 2^2 + 6^2}} = \frac{19}{7}$$

Alternate method:Let $A(\vec{a}) = (2, 3, 4)$

Given equation of plane is

$$\vec{r} \cdot (3\hat{i} - 2\hat{j} + 6\hat{k}) = 5$$

- $\therefore \vec{a} = 2\hat{i} + 3\hat{j} + 4\hat{k}$, and $\vec{n} = 3\hat{i} - 2\hat{j} + 6\hat{k}$

- \therefore The distance of point from plane is

$$d = \frac{|(\vec{a} \cdot \vec{n}) - p|}{|\vec{n}|} = \frac{|2(3) + 3(-2) + 4(6) - 5|}{\sqrt{3^2 + 2^2 + 6^2}} = \frac{19}{7}$$

60. Here, $a = 1, b = 1, c = 1, d = -3$ and $x = y = z = 0$

$$\therefore d = \frac{-3}{\sqrt{1^2 + 1^2 + 1^2}} = \sqrt{3}$$

61. Here, $a = 3, b = -6, c = 2, z = 11$ and $x = 2, y = 3, z = 4$

$$\therefore d = \frac{|3(2) + (-6)(3) + 2(4) + 11|}{\sqrt{3^2 + (-6)^2 + 2^2}} = 1$$

62. Let the intercepts made by the plane $a, b, c = 2, 1, -2$

- \therefore The distance of plane from origin is

$$d = \frac{1}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2}}} = \frac{1}{\sqrt{\frac{1}{4} + 1 + \frac{1}{4}}} = \frac{2}{\sqrt{6}}$$

Alternate method:

The equation of plane is

$$\frac{x}{2} + \frac{y}{1} + \frac{z}{-2} = 1$$

$$\text{i.e. } x + 2y - z - 2 = 0$$

- \therefore distance of plane from the origin is

$$d = \frac{|-2|}{\sqrt{1+4+1}}$$

$$d = \frac{2}{\sqrt{6}}$$

63. Let $a, b, c = -6, 3, 4$

- \therefore The length of perpendicular from origin is

$$d = \frac{1}{\sqrt{\frac{1}{(-6)^2} + \frac{1}{3^2} + \frac{1}{4^2}}} = \frac{1}{\sqrt{\frac{29}{144}}} = \frac{12}{\sqrt{29}}$$

64. The distance of $(1, 1, 1)$ from the origin is

$$d = \sqrt{(1)^2 + (1)^2 + (1)^2} = \sqrt{3}$$

Distance of $(1, 1, 1)$ from $x + y + z + k = 0$ is

$$d_1 = \frac{|(1) + (1) + (1) + k|}{\sqrt{(1)^2 + (1)^2 + (1)^2}} = \pm \frac{k+3}{\sqrt{3}}$$

$$\text{Now, } \sqrt{3} = \pm \frac{1}{2} \left(\frac{k+3}{\sqrt{3}} \right) \quad \dots(\text{given})$$

$$\Rightarrow 6 = \pm (k+3)$$

$$\Rightarrow k = 3, -9$$

65. Since, the points $(1, 1, k)$ and $(-3, 0, 1)$ are equidistance from the given plane

$$\therefore \frac{|3+4-12k+13|}{\sqrt{9+16+144}} = \frac{|-9-12+13|}{\sqrt{9+16+144}}$$

$$\Rightarrow |3+4-12k+13| = |-9-12+13|$$

$$\Rightarrow 20 - 12k = \pm 8 \Rightarrow k = 1, \frac{5}{7}$$

66. Given line passes through $(1, -2, 1)$ and the d.r.s. of normal to the plane are $2, 2, -1$

$$\therefore d = \frac{|2(1) + 2(-2) - 1(1) - 6|}{\sqrt{2^2 + 2^2 + (-1)^2}} = \frac{9}{\sqrt{9}} = 3$$

67. Given planes are parallel and can be written as

$$2x - 2y + z + 3 = 0 \text{ and } 2x - 2y + z + \frac{5}{2} = 0$$

- \therefore the distance between these planes is

$$d = \frac{|d_1 - d_2|}{\sqrt{a^2 + b^2 + c^2}}$$

$$= \frac{\left| 3 - \frac{5}{2} \right|}{\sqrt{4+4+1}} = \frac{\left(\frac{1}{2} \right)}{3} = \frac{1}{6}$$



68. Given planes are parallel, and can be written as
 $x + 2y + 3z + 7 = 0$ and $x + 2y + 3z + \frac{7}{2} = 0$

∴ the distance between these planes is

$$d = \left| \frac{7 - \frac{7}{2}}{\sqrt{1+4+9}} \right| = \frac{\sqrt{7}}{2\sqrt{2}}$$

69. The plane passes through points (1, -2, 3) and (4, 0, -1)

This points satisfies the equation of plane in option (A)

∴ option (A) is correct answer.

70. The plane passes through (1, 2, -1)

This point satisfies the equation of plane in option (A)

$$\begin{aligned} \text{Also, it has d.r.s } \vec{b} \times \vec{c} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & -1 \\ 1 & 1 & 3 \end{vmatrix} \\ &= 7\hat{i} - 4\hat{j} - \hat{k} \end{aligned}$$

i.e., 7, -4, -1

∴ option (A) is correct answer.

Alternate Method

Let $\vec{a} = \hat{i} - \hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} + \hat{j} + \hat{k}$ and $\vec{c} = \hat{j} + 2\hat{k}$

Now, $\vec{b} \times \vec{c} = \hat{i} - 4\hat{j} + 2\hat{k}$

∴ the vector equation of required plane is

$$\begin{aligned} \vec{r} \cdot (\vec{b} \times \vec{c}) &= \vec{a} \cdot (\vec{b} \times \vec{c}) \\ \Rightarrow \vec{r} \cdot (7\hat{i} - 4\hat{j} - \hat{k}) &= (\hat{i} - 2\hat{j} - \hat{k}) \cdot (7\hat{i} - 4\hat{j} - \hat{k}) \\ \Rightarrow \vec{r} \cdot (7\hat{i} - 4\hat{j} - \hat{k}) &= 0 \end{aligned}$$



Critical Thinking

1. Let x_1, y_1, z_1 be the intercepts made by the plane

∴ Equation of plane is

$$\frac{x}{x_1} + \frac{y}{y_1} + \frac{z}{z_1} = 1$$

Since it passes through (a, b, c),

$$\Rightarrow \frac{a}{x_1} + \frac{b}{y_1} + \frac{c}{z_1} = 1$$

∴ Locus of (x_1, y_1, z_1) is $\frac{a}{x} + \frac{b}{y} + \frac{c}{z} = 1$

2. Since, the plane contains the X-axis, it passes through the origin

∴ $d = 0$

∴ The equation of the plane is

$$ax + by + cz = 0 \quad \dots(i)$$

Also, plane passes through (1, 1, 1)

∴ $a + b + c = 0 \quad \dots(ii)$

The equation of the X-axis is $\frac{x}{1} = \frac{y}{0} = \frac{z}{0}$

As the plane contains the X-axis, the d.r.s of the normal to the plane are perpendicular to X-axis

∴ $a(1) + b(0) + c(0) = 0$

$\Rightarrow a = 0$

Substituting value of a in (ii) we get

$$b + c = 0 \Rightarrow b = -c$$

∴ The equation of the required plane is

$$by - bz = 0$$

$$\Rightarrow y - z = 0$$

3. The plane passes through (1, -1, 3) and (2, 3 -4)

The points satisfies the equation of plane in option (B)

∴ option (B) is correct answer.

Alternate method:

Let $ax + by + cz + d = 0$ be the equation of the required plane.

Since, the plane is parallel to X-axis,

∴ $a = 0$

The points (1, -1, 3) and (2, 3, -4) lie in the plane,

∴ $-b + 3c + d = 0$, and $\dots(i)$

$$3b - 4c + d = 0 \quad \dots(ii)$$


Solving the equations (i) and (ii), we get

$$\frac{b}{3 - (-4)} = \frac{c}{3 + 1} = \frac{d}{4 - 9}$$

$$\Rightarrow \frac{b}{7} = \frac{c}{4} = \frac{d}{-5}$$

∴ Equation of the required plane is $7y + 4z - 5 = 0$

4.





$$\begin{aligned}\therefore \vec{m} &= \frac{(1-1)\hat{i} + (2+2)\hat{j} + (4-6)\hat{k}}{2} \\ &= 2\hat{j} - \hat{k}\end{aligned}$$

\therefore equation of plane passing through the vector $2\hat{j} - \hat{k}$ and perpendicular to $\vec{AB} = -2\hat{i} - 10\hat{k}$ is

$$\vec{r} \cdot (-2\hat{i} - 10\hat{k}) = (2\hat{j} - \hat{k}) \cdot (-2\hat{i} - 10\hat{k})$$

$$\Rightarrow \vec{r} \cdot (\hat{i} + 5\hat{k}) = -10$$

5. P be the point (a, b, c).

\therefore The d.r.s of OP are a, b, c.

\therefore Equation of the plane passing through the point (a, b, c) is

$$a(x - a) + b(y - b) + c(z - c) = 0$$

$$\Rightarrow ax + by + cz = a^2 + b^2 + c^2$$

6. Mid-point of the line segment joining the points (-1, 2, 3) and (3, -5, 6) is

$$M \equiv \left(\frac{-1+3}{2}, \frac{2-5}{2}, \frac{3+6}{2} \right)$$

$$M \equiv \left(1, -\frac{3}{2}, \frac{9}{2} \right)$$

The plane passes through point M

It satisfies option (C)

Alternate method:

The required plane bisects the line segment perpendicularly.

\therefore the d.r.s of the normal to the plane are

$$3 - (-1), -5 - 2, 6 - 3$$

i.e. 4, -7, 3


Since, the mid-point $\left(1, -\frac{3}{2}, \frac{9}{2} \right)$ lies in the plane,

\therefore The equation of the plane is

$$4(x - 1) - 7\left(y + \frac{3}{2}\right) + 3\left(z - \frac{9}{2}\right) = 0$$

$$\Rightarrow 4x - 7y + 3z = 28$$

7.



Mid-point of line joining P(1, 2, 3) and Q(-3, 4, 5) is (-1, 3, 4)

It lies on the plane

The d.r.s. of normal to the plane are -4, 2, 2
i.e. -2, 1, 1

\therefore The equation of the plane is

$$-2(x + 1) + 1(y - 3) + 1(z - 4) = 0$$

$$\Rightarrow 2x - y - z = -9$$

$$\Rightarrow \frac{x}{-9} + \frac{y}{9} + \frac{z}{9} = 1$$

\therefore Intercepts are $\frac{-9}{2}, 9, 9$

8. (2, -1, 0) lies on the plane $9x - 2y - 3z = k$

$\therefore 9(2) - 2(-1) - 3(0) = k$

$$\Rightarrow k = 20$$

9. Since, the point (1, 0, z_1) lies on the plane

$$\vec{r} \cdot (-\hat{i} + 3\hat{k}) = 2$$

i.e. $-x + 3z = 2$

$$\Rightarrow z_1 = 1$$

10. (3, 2, -1) lies on the plane $5x + 3y - 2z = \lambda$

$\therefore 5(3) + 3(2) - 2(-1) = \lambda$

$$\Rightarrow \lambda = 23$$

11. The equation of the plane passing through the intersection of the planes $\vec{r} \cdot \vec{a} = p$ and

$$\vec{r} \cdot \vec{b} = q$$

$$\vec{r} \cdot (\vec{a} + \lambda \vec{b}) = p + \lambda q \quad \dots(i)$$

Since, the plane passes through the origin,

$$p + \lambda q = 0$$

$$\Rightarrow \lambda = \frac{-p}{q}$$

Substituting the value of λ in (i), we get

$$\vec{r} \cdot \left(\vec{a} - \frac{p}{q} \vec{b} \right) = p + \left(\frac{-p}{q} \right) (q)$$

$$\Rightarrow \vec{r} \cdot (\vec{a}q - \vec{b}p) = pq - pq$$

$$\Rightarrow \vec{r} \cdot (q\vec{a} - p\vec{b}) = 0$$

12. The line of intersection of the planes

$$\vec{r} \cdot (3\hat{i} - \hat{j} + \hat{k}) = 1 \text{ and } \vec{r} \cdot (\hat{i} + 4\hat{j} - 2\hat{k}) = 2$$

is perpendicular to each of the normal vectors

$$\vec{n}_1 = 3\hat{i} - \hat{j} + \hat{k} \text{ and } \vec{n}_2 = \hat{i} + 4\hat{j} - 2\hat{k}.$$

\therefore The line is parallel to the vector $\vec{n}_1 \times \vec{n}_2$

$$\begin{aligned}\therefore \vec{n}_1 \times \vec{n}_2 &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & -1 & 1 \\ 1 & 4 & -2 \end{vmatrix} \\ &= -2\hat{i} + 7\hat{j} + 13\hat{k}\end{aligned}$$



13. The equation of the required plane is
 $x + 2y + 3z - 4 + \lambda(2x + y - z + 5) = 0$
 $\Rightarrow (1 + 2\lambda)x + (2 + \lambda)y + (3 - \lambda)z - 4 + 5\lambda = 0$ (i)

Let a, b, c be the d.r.s of the required plane

- \therefore From equation (i), $a = 1 + 2\lambda$; $b = 2 + \lambda$;
 $c = 3 - \lambda$

The required plane is perpendicular to
 $5x + 3y - 6z + 8 = 0$

- \therefore $5a + 3b - 6c = 0$
 $\Rightarrow 5(1 + 2\lambda) + 3(2 + \lambda) - 6(3 - \lambda) = 0$
 $\Rightarrow 5 + 10\lambda + 6 + 3\lambda - 18 + 6\lambda = 0$
 $\Rightarrow -7 + 19\lambda = 0$

$$\Rightarrow \lambda = \frac{7}{19}$$

Substituting the value of λ in equation (i), we get

$$\left(1 + 2 \times \frac{7}{19}\right)x + \left(2 + \frac{7}{19}\right)y + \left(3 - \frac{7}{19}\right)z - 4 + 5\left(\frac{7}{19}\right) = 0$$

$$\Rightarrow \frac{33}{19}x + \frac{45}{19}y + \frac{50}{19}z - \frac{41}{19} = 0$$

$$\Rightarrow 33x + 45y + 50z - 41 = 0$$

14. The equation of the plane passing through the origin is $ax + by + cz = 0$.

The required plane is perpendicular to the line
 $x = 2y = 3z$

$$\text{i.e., } \frac{x}{6} = \frac{y}{3} = \frac{z}{2}$$

- \therefore the d.r.s. of the line are 6, 3, 2
 \therefore the d.r.s. of the normal to the plane are 6, 3 and 2.
 \therefore the equation of the required plane is
 $6x + 3y + 2z = 0$

15. Let a, b, c be the d.r.s. of the required plane.
 Since, the plane passes through Z-axis,

$$\therefore a(0) + b(0) + c(1) = 0$$


$$\Rightarrow c = 0$$

Given that the required plane is perpendicular

$$\text{to } \frac{x-1}{\cos\theta} = \frac{y+2}{\sin\theta} = \frac{z-3}{0}$$

- \therefore d.r.s of normal to plane are $\cos\theta, \sin\theta, 0$
 \therefore the equation of required plane is
 $x \cos\theta + y \sin\theta = 0$
 $\Rightarrow x + y \tan\theta = 0$

- 16.



$$\bar{n} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 3 & 2 \\ 2 & 7 & 5 \end{vmatrix} = \hat{i} - \hat{j} + \hat{k}$$

- \therefore the d.r.s of the normal to the plane are 1, -1, 1
 \therefore the equation of plane passing through the point (1, 2, 3)
 $1(x-1) - 1(y-2) + 1(z-3) = 0$
 $\Rightarrow x - y + z = 2$

17. Equation of any plane through (x_1, y_1, z_1) is
 $a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$ (i)
 it contains the line

$$\frac{x - x_2}{d_1} = \frac{y - y_2}{d_2} = \frac{z - z_2}{d_3} = 0$$

i.e. it passes through (x_2, y_2, z_2)

- $\therefore a(x_2 - x_1) + b(y_2 - y_1) + c(z_2 - z_1) = 0$ (ii)
 Also, $ad_1 + bd_2 + cd_3 = 0$ (iii)
 Eliminating a, b, c from (i), (ii), (iii), we get the equation of the required plane as

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ d_1 & d_2 & d_3 \end{vmatrix} = 0$$

18. Vector perpendicular to plane is
 $\bar{n} = 6\hat{i} - 3\hat{j} + 5\hat{k}$

Thus, the line perpendicular to the given line will be parallel to \bar{n}

- \therefore The equation of line which passes through
 $\bar{a} = 2\hat{i} - 3\hat{j} - 5\hat{k}$ and parallel to \bar{n} is
 $\bar{r} = \bar{a} + \lambda\bar{n}$
 $\Rightarrow \bar{r} = (2\hat{i} - 3\hat{j} - 5\hat{k}) + \lambda(6\hat{i} - 3\hat{j} + 5\hat{k})$

19. The d.r.s. of the line are 3, -4, 5 and it passes through is 3, -5, 7

- \therefore The equation of line is
 $\bar{r} = 3\hat{i} - 5\hat{j} + 7\hat{k} + \lambda(3\hat{i} - 4\hat{j} + 5\hat{k})$



20. The line is perpendicular to the plane

$$x + 2y - 5z + 9 = 0$$

∴ the d.r.s are 1, 2, -5

Also it passes through (1, 2, 3)

∴ The equation of line is $\frac{x-1}{1} = \frac{y-2}{2} = \frac{z-3}{-5}$

$$21. \quad \vec{n} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 2 \\ 3 & 1 & 1 \end{vmatrix} = -3\hat{i} + 5\hat{j} + 4\hat{k}$$

∴ the d.r.s. of line are -3, 5, 4

∴ The equation of the line passing through (1, 2, 3) and having d.r.s. -3, 5, 4 is

$$\frac{x-1}{-3} = \frac{y-2}{5} = \frac{z-3}{4}$$

22. Here, $\vec{n}_1 = (x\hat{i} + \hat{j} - \hat{k})$, and

$$\vec{n}_2 = (\hat{i} + x\hat{j} - \hat{k})$$

$$\therefore \cos \theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| |\vec{n}_2|}$$

$$\Rightarrow \cos \frac{\pi}{3} = \frac{|(x\hat{i} + \hat{j} - \hat{k}) \cdot (\hat{i} + x\hat{j} - \hat{k})|}{\sqrt{x^2 + 1 + 1} \cdot \sqrt{1 + x^2 + 1}}$$

$$\Rightarrow \frac{1}{2} = \pm \left(\frac{x + x - 1}{x^2 + 2} \right)$$

$$\Rightarrow \frac{2x-1}{x^2+2} = \frac{1}{2} \quad \dots(\text{considering positive value})$$

$$\Rightarrow x^2 + 2 - 4x + 2 = 0$$

$$\Rightarrow (x-2)^2 = 0$$

$$\Rightarrow x = 2$$

23. Consider plane OPQ

the equation of plane is

$$ax + by + cz = 0$$

The plane passes through P(1, 2, 1) and Q(2, 3, 0)

$$\therefore a + 2b + c = 0 \quad \dots(\text{i})$$

$$2a + 3b = 0 \quad \dots(\text{ii})$$

On solving (i) and (ii), we get

$$\frac{a}{-3} = \frac{b}{2} = \frac{c}{-1}$$

∴ The equation of plane OPQ is

$$-3x + 2y - z = 0 \quad \dots(\text{iii})$$

The equation of plane PQR is

$$a_1(x-1) + b_1(y-2) + c_1(z-1) = 0$$

On solving for a_1, b_1, c_1 , we get

$$a_1 = -3, b_1 = 3, c_1 = 0$$

∴ The equation of PQR is

$$x - y + 1 = 0 \quad \dots(\text{iv})$$

∴ The angle between the planes represented by equations (iii) and (iv) is

$$\begin{aligned} \cos \theta &= \frac{|(-3)(1) + 2(-1)|}{\sqrt{9+4+1} \cdot \sqrt{1+1}} \\ &= \frac{|-5|}{\sqrt{14} \cdot \sqrt{2}} \end{aligned}$$

$$\Rightarrow \theta = \cos^{-1} \left(\frac{5}{\sqrt{28}} \right)$$

24. The d.r.s. of normal to the plane are 2, 3, -1

The d.r.s. of X-axis are 1, 0, 0

∴ the angle between the plane and X-axis is

$$\sin \theta = \frac{aa_1 + bb_1 + cc_1}{\sqrt{a^2 + b^2 + c^2} \cdot \sqrt{a_1^2 + b_1^2 + c_1^2}}$$

$$\Rightarrow \sin \theta = \frac{2(1) + 0 + 0}{\sqrt{4+9+1} \cdot \sqrt{1}}$$

$$\Rightarrow \sin \theta = \frac{2}{\sqrt{14}}$$

$$\Rightarrow \theta = \sin^{-1} \left(\frac{2}{\sqrt{14}} \right)$$

25. Here $a = 1, b = k, c = 4$ and

$$a_1 = 1, b_1 = -3, c_1 = 2$$

The angle between the line and plane is

$$\sin \theta = \frac{aa_1 + bb_1 + cc_1}{\sqrt{a^2 + b^2 + c^2} \cdot \sqrt{a_1^2 + b_1^2 + c_1^2}}$$

$$\text{Now, } \theta = \sin^{-1} \left(\frac{3}{7\sqrt{6}} \right) \Rightarrow \sin \theta = \frac{3}{7\sqrt{6}}$$

$$\therefore \frac{3}{7\sqrt{6}} = \frac{1-3k+8}{\sqrt{1+k^2+16} \cdot \sqrt{1+9+4}}$$

$$\Rightarrow k^2 + 21k - 46 = 0$$

$$\Rightarrow k = 2 \text{ or } -23$$

26. Equation of the line L: $\frac{x-1}{2} = \frac{y}{3} = \frac{z-3}{2}$ and

equation of the plane P: $4x - 2y - z = 1$.

The d.r.s. of the line are 2, 3, 2, and

The d.r.s. of the normal to the plane are 4, -2, -1.

Now consider

$$a_1a_2 + b_1b_2 + c_1c_2 = 8 - 6 - 2 = 0$$

∴ Line L and plane P are parallel.

Since the point (1, 0, 3), which lies on the line L also satisfies the equation of the plane,

∴ The line L lies in the plane P.



27. Equation of the line
 $L: \frac{x+3}{2} = \frac{y-4}{3} = \frac{z+5}{1}$
 and equation of the plane
 $P: 2x - 3y + 5z = 1$.
 The d.r.s of the line are 2, 3, 1
 The d.r.s of the normal to the plane are 2, -3, 5.
 Now consider
 $a_1a_2 + b_1b_2 + c_1c_2 = 4 - 9 + 5 = 0$
 \therefore Line L is parallel to the plane P.

28. Since, the line $\frac{x-3}{2} = \frac{y-4}{3} = \frac{z-5}{4}$ lies in
 the plane $4x + 4y - cz - d = 0$,
 $\therefore aa_1 + bb_1 + cc_1 = 0$
 $\Rightarrow 2(4) + 3(4) + 4(-c) = 0$
 $\Rightarrow 20 - 4c = 0$
 $\Rightarrow c = 5$
 Also, the plane passes through (3, 4, 5)
 $\therefore 4(3) + 4(4) - 5(5) - d = 0$
 $\Rightarrow d = 3$

29. Given equation of plane
 $\frac{x-1}{2} = \frac{y-1}{3} = \frac{z+2}{2}$
 \therefore The line passes through (1, 1, -2)
 The above point lies on the plane
 $x + By - 3z + D = 0$
 $\therefore 1 + B + 6 + D = 0$
 $\Rightarrow B + D = -7 \quad \dots(i)$
 Also the given line is perpendicular to the
 normal to the plane
 $a_1a_2 + b_1b_2 + c_1c_2 = 0$
 $\Rightarrow 2(1) + 3(B) + 2(-3) = 0$
 $\Rightarrow B = \frac{4}{3}$
 Substituting value of B is equation (i), we get
 $D = \frac{-25}{3}$

30. Since both the given lines pass through the
 point with position vector $\hat{i} + \hat{j}$, the required
 plane also passes through $\hat{i} + \hat{j}$, and normal to
 the plane is perpendicular to the vectors
 $\hat{i} + 2\hat{j} - \hat{k}$ and $-\hat{i} + \hat{j} - 2\hat{k}$.
 Let a, b, c be the d.r.s. of the normal to the
 plane.

$$\therefore \bar{n} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & -1 \\ -1 & 1 & -2 \end{vmatrix}$$

$$\Rightarrow \bar{n} = -3\hat{i} + 3\hat{j} + 3\hat{k}$$

$$\text{i.e. } \bar{n} = -\hat{i} + \hat{j} + \hat{k}$$

\therefore Vector equation of the plane passing through
 $\hat{i} + \hat{j}$ and containing the given lines is
 $\bar{r} \cdot (-\hat{i} + \hat{j} + \hat{k}) = (\hat{i} + \hat{j}) \cdot (-\hat{i} + \hat{j} + \hat{k})$
 $\Rightarrow \bar{r} \cdot (-\hat{i} + \hat{j} + \hat{k}) = 0$

31. The plane passes through
 (0, 2, -3) and (2, 6, 3)
 The two points satisfy the equation of plane is
 option (A)

\therefore option (A) is correct.

Alternate Method:

The equation of the plane is

$$\begin{vmatrix} x-\alpha & y-\beta & z-\gamma \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} x & y-2 & z+3 \\ 1 & 2 & 3 \\ 2 & 3 & 4 \end{vmatrix} = 0$$

$$\Rightarrow -x - (y-2)(-2) + (z+3)(-1) = 0$$

$$\Rightarrow -x + 2y - 4 - z - 3 = 0$$

$$\Rightarrow x - 2y + z + 7 = 0$$

32. The plane passes through
 (5, 7, -3) and (8, 4, 5)
 The two points satisfy the equation of plane is
 option (A)

\therefore option (A) is correct.

33. Let a, b, c be the d.r.s of the normal to the
 plane

$$\therefore \bar{n} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & 5 & 7 \\ 1 & 4 & 7 \end{vmatrix} = \hat{i} - 2\hat{j} + \hat{k}$$

Since, the plane passes through (-1, -3, -5)

$$\therefore 1(x+1) - 2(y+3) + 1(z+5) = 0$$

$$\Rightarrow x - 2y + z = 0$$

From the given options only (0, 0, 0) satisfies
 the equation of the plane.

\therefore The plane passes through (0, 0, 0).

34. Here $x_1, y_1, z_1 = - , -3, -5$ and $x_2, y_2, z_2 = 2, 4, 6$
 $a_1, b_1, c_1 = 3, 5, 7$ and $a_2, b_2, c_2 = 1, 3, 5$

Since the given lines are coplanar

$$\therefore \begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$



$$\Rightarrow \begin{vmatrix} - & -2 & -3-4 & -5-6 \\ 3 & 5 & 7 & \\ 1 & 3 & 5 & \end{vmatrix} = 0$$

$$\Rightarrow (- -2)(25 - 21) - (-3-4)(15 - 7) + (-5-6)(9 - 5) = 0$$

$$\Rightarrow 12 = 4(+2)$$

$$\Rightarrow = 1.$$

35. The lines are coplanar

$$\therefore \begin{vmatrix} -1-2 & -3-4 & -5-6 \\ 1 & 4 & k \\ 3 & 5 & k \end{vmatrix} = 0$$

$$\Rightarrow -3(4k - 5k) + 7(k - 3k) - 11(-7) = 0$$

$$\Rightarrow k = 7$$

36. Since the given lines are coplanar, then

$$\therefore \begin{vmatrix} 3-1 & 1-2 & 3-1 \\ 1 & 2 & -\lambda \\ \lambda & 3 & 4 \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} 2 & -1 & 2 \\ 1 & 2 & -\lambda \\ \lambda & 3 & 4 \end{vmatrix} = 0$$

$$\Rightarrow \lambda^2 + 2\lambda + 26 = 0$$

$$\Delta = 4 - 4(1)(26) < 0$$

\(\therefore\) Roots are imaginary

So no real value of \(\lambda\) exists.

$$37. \frac{x}{1} = \frac{y-2}{2} = \frac{z+3}{3} \text{ and } \frac{x-2}{2} = \frac{y-6}{3} = \frac{z-3}{4}$$

The d.r.s. of the first line are 1, 2, 3 and

The d.r.s. of the second line are 2, 3, 4

Ratio of the d.r.s. are not same

$$\text{i.e. } \frac{2}{1} \neq \frac{3}{2} \neq \frac{4}{3}$$

\(\therefore\) The lines are not parallel.

Sum of the products of the d.r.s. is not equal to 0 i.e., $2(1) + 2(3) + 3(4) \neq 0$

\(\therefore\) The lines are not perpendicular.

$$\text{Consider } \begin{vmatrix} 0+2 & -2+6 & 3+3 \\ 1 & 2 & 3 \\ 2 & 3 & 4 \end{vmatrix} = \begin{vmatrix} 2 & 4 & 6 \\ 1 & 2 & 3 \\ 2 & 3 & 4 \end{vmatrix}$$

$$= 2 \begin{vmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \\ 2 & 3 & 4 \end{vmatrix} = 0 \quad (\because \text{the two rows are same})$$

\(\therefore\) The two lines are coplanar.

38. Let d_1 be the distance of the point $(1, 2, -1)$ from the plane $2x - 3y + z + k = 0$

$$\therefore d_1 = \frac{|2(1) - 3(2) + (-1) + k|}{\sqrt{2^2 + (-3)^2 + 1^2}} = \frac{|-5 + k|}{\sqrt{4 + 9 + 1}}$$

$$= \frac{|k - 5|}{\sqrt{14}}$$

Let d_2 be the distance of the point $(1, 2, -1)$ from the plane $x + 2y + 3z = 0$

$$\therefore d_2 = \frac{|(1) + 2(2) + 3(-1)|}{\sqrt{1^2 + 2^2 + 3^2}} = \frac{|2|}{\sqrt{14}}$$

Given that $d_1 \cdot d_2 = 1$.

$$\therefore \frac{|k-5|}{\sqrt{14}} \cdot \frac{|2|}{\sqrt{14}} = 1$$

$$\Rightarrow (k-5)2 = 14$$

$$\Rightarrow k-5 = 7$$

$$\Rightarrow k = 12$$

$$39. P_1 = \frac{|3(2) - 6(3) + 2(4) + 11|}{\sqrt{3^2 + (-6)^2 + (2)^2}} = 1$$

$$P_2 = \frac{|3(1) - 6(1) + 2(4) + 11|}{\sqrt{3^2 + (-6)^2 + (2)^2}} = \frac{16}{7}$$

the equation P_1 and P_2 satisfies

$$7P^2 - 23P + 16 = 0.$$

\(\therefore\) P_1 and P_2 are the roots of the equation (B).

40. Equation of plane parallel to $x - 2y + 2z = 5$ is $x - 2y + 2z + k = 0$... (i)

distance of the above plane from $(1, 2, 3)$ is 1.

$$\therefore \frac{|1 - 4 + 6 + k|}{\sqrt{9}} = 1$$

$$\text{i.e. } k + 3 = \pm 3$$

$$\Rightarrow k = 0 \text{ or } -6$$

41. Let x, y, z be any point

$$d_1^2 + d_2^2 + d_3^2 = 36$$

$$\therefore \left| \frac{x-z}{\sqrt{2}} \right|^2 + \left| \frac{x-2y+z}{\sqrt{6}} \right|^2 + \left| \frac{x+y+z}{\sqrt{3}} \right|^2 = 36$$

$$\Rightarrow \frac{1}{6} [3x^2 - 6xz + 3z^2 + x^2 + 4y^2 + z^2 - 4xy - 4yz + 2xz + 2x^2 + 2y^2 + 2z^2 + 4xy + 4yz + 4xz] = 36$$

$$\therefore \Rightarrow x^2 + y^2 + z^2 = 36$$



42. Since all the planes are parallel,

$$\therefore p_1 = \frac{|2-6|}{\sqrt{2^2+(-3)^2+4^2}} = \frac{4}{\sqrt{29}}$$

Equation of the plane $4x - 6y + 8z + 3 = 0$ can be written as $2x - 3y + 4z + \frac{3}{2} = 0$

$$\therefore p_2 = \frac{\left|2 - \frac{3}{2}\right|}{\sqrt{2^2+(-3)^2+4^2}} = \frac{1}{2\sqrt{29}}$$

$$\text{and } p_3 = \frac{|2+6|}{\sqrt{2^2+(-3)^2+4^2}} = \frac{8}{\sqrt{29}}$$

$$\begin{aligned} \text{Now consider } p_1 + 8p_2 - p_3 \\ = \frac{4}{\sqrt{29}} + \frac{4}{\sqrt{29}} - \frac{8}{\sqrt{29}} \\ = 0 \end{aligned}$$

43. Let $A \equiv (a, 0, 0)$, $B \equiv (0, b, 0)$ and $C \equiv (0, 0, c)$
The equation of the plane in intercept form is

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

Since, centroid is $(3, 3, 3)$

$$\therefore 3 = \frac{x_1 + x_2 + x_3}{3} = \frac{a+0+0}{3} = 3$$

$$\Rightarrow a = 9$$

$$\text{Similarly } \frac{0+b+0}{3} = 3 \Rightarrow b = 9, \text{ and}$$

$$\frac{0+0+c}{3} = 3 \Rightarrow c = 9$$

$$\therefore \text{The equation of plane is } \frac{x}{9} + \frac{y}{9} + \frac{z}{9} = 1$$

$$\Rightarrow x + y + z = 9$$

44. Let $A \equiv (a, 0, 0)$, $B \equiv (0, b, 0)$ and $C \equiv (0, 0, c)$.

Since, centroid is (α, β, γ)

$$\therefore a = 3\alpha, b = 3\beta, c = 3\gamma$$

$$\therefore \text{the equation of the plane is } \frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

$$\frac{x}{3\alpha} + \frac{y}{3\beta} + \frac{z}{3\gamma} = 1$$

$$\Rightarrow \frac{x}{\alpha} + \frac{y}{\beta} + \frac{z}{\gamma} = 3$$

45. The given equation of plane is $6x - 3y + 2z = 18$

$$\text{i.e. } \frac{x}{3} + \frac{y}{-6} + \frac{z}{9} = 1$$

If a, b, c are intercepts made by the plane, then

$$\text{Centroid} \equiv \left(\frac{a+0+0}{3}, \frac{0+b+0}{3}, \frac{0+0+c}{3} \right)$$

$$\begin{aligned} \therefore G &\equiv \left(\frac{3+0+0}{3}, \frac{0-6+0}{3}, \frac{9+0+0}{3} \right) \\ &\Rightarrow G \equiv (1, -2, 3) \end{aligned}$$

46. The given equations of plane is $ax + by + cz = 1$

$$\text{i.e. } \frac{x}{\frac{1}{a}} + \frac{y}{\frac{1}{b}} + \frac{z}{\frac{1}{c}} = 1$$

\therefore The intercepts made by the plane are $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}$

$$\therefore A \equiv \left(\frac{1}{a}, 0, 0 \right); B \equiv \left(0, \frac{1}{b}, 0 \right); C \equiv \left(0, 0, \frac{1}{c} \right)$$

$$\therefore \text{centroid} \equiv \left(\frac{\frac{1}{a}+0+0}{3}, \frac{0+\frac{1}{b}+0}{3}, \frac{0+0+\frac{1}{c}}{3} \right)$$

$$\Rightarrow G \equiv \left(\frac{1}{3a}, \frac{1}{3b}, \frac{1}{3c} \right)$$

47. Let equation of plane be $x + my + nz = p$

$$\text{i.e., } \frac{x}{\frac{p}{1}} + \frac{y}{\frac{p}{m}} + \frac{z}{\frac{p}{n}} = 1$$

$$\therefore A \equiv \left(\frac{p}{1}, 0, 0 \right), B \equiv \left(0, \frac{p}{m}, 0 \right), C \equiv \left(0, 0, \frac{p}{n} \right)$$

If centroid of ΔABC is (x_1, y_1, z_1) , then

$$x_1 = \frac{p}{3}, y_1 = \frac{p}{3m}, z_1 = \frac{p}{3n}$$

$$\text{Now, } p^2 + m^2 + n^2 = 1$$

$$\therefore \frac{p^2}{9x_1^2} + \frac{p^2}{9y_1^2} + \frac{p^2}{9z_1^2} = 1$$

$$\Rightarrow \frac{1}{x_1^2} + \frac{1}{y_1^2} + \frac{1}{z_1^2} = \frac{9}{p^2}$$

48. The equation of line perpendicular to given plane passing through $(2, 2, 2)$ is

$$\frac{x-2}{1} = \frac{y-2}{1} = \frac{z-2}{1} = \lambda \text{ (say)}$$

Any general point on it is $P \equiv (\lambda + 2, \lambda + 2, \lambda + 2)$

Since, P lies the plane $x + y + z = 0$

$$\therefore \lambda + 2 + \lambda + 2 + \lambda + 2 = 9 \Rightarrow \lambda = 1$$

\therefore The foot of perpendicular is $(3, 3, 3)$.

49. The required plane is perpendicular to the line

$$\frac{x-2}{1} = \frac{y-4}{2} = \frac{z-5}{2} = \lambda \text{ (say)}$$

the d.r.s of normal to the plane are proportional to $1, 2, 2$

\therefore Equation of the plane is

$$x + 2y + 2z + d = 0$$

....(i)



Since it passes through the point (5, 1, 2), we have

$$(5) + 2(1) + 2(2) + d = 0$$

$$\Rightarrow d = -11$$

\therefore The equation (i) becomes $x + 2y + 2z - 11 = 0$
Any general point on the given line is given by

$$\lambda + 2, 2\lambda + 4, 2\lambda + 5.$$

This point lies in the required plane

$$\therefore \lambda + 2 + 2(2\lambda + 4) + 2(2\lambda + 5) - 11 = 0$$

$$\Rightarrow \lambda + 2 + 4\lambda + 8 + 4\lambda + 10 - 11 = 0$$

$$\Rightarrow 9\lambda + 9 = 0 \Rightarrow \lambda = -1$$

\therefore The point of intersection is

$$[(-1) + 2, 2(-1) + 4, 2(-1) + 5]$$

$$\Rightarrow (1, 2, 3)$$

50. The equation of plane passing through the intersection of the given planes is

$$(2x - 5y + z - 3) + \lambda(x + y + 4z - 5) = 0$$

$$\Rightarrow (2 + \lambda)x + (-5 + \lambda)y$$

$$+ (1 + 4\lambda)z - 3 - 5\lambda = 0 \quad \dots(i)$$

This plane is parallel to the plane $x + 3y + 6z = 1$

$$\therefore \frac{2 + \lambda}{1} = \frac{-5 + \lambda}{3} = \frac{1 + 4\lambda}{6}$$

$$\Rightarrow \lambda = \frac{-11}{2}$$

\therefore Substituting value of λ in equation (i), we get

$$-\frac{7}{2}x - \frac{21}{2}y - \frac{42}{2}z + \frac{49}{2} = 0$$

$$\therefore x + 3y + 6z = 7$$

Comparing with $x + 3y + 6z = k$, we get

$$k = 7$$

51. The equation of the plane through the line of intersection of the planes,

$$4x + 7y + 4z + 81 = 0 \text{ and } 5x + 3y + 10z = 25$$

$$\text{is } (4x + 7y + 4z + 81) + \lambda(5x + 3y + 10z - 25) = 0$$

$$\Rightarrow (4 + 5\lambda)x + (7 + 3\lambda)y + (4 + 10\lambda)z + 81 - 25\lambda = 0 \quad \dots(i)$$

It is parallel to $x - 4y + 6z = k$,

$$\therefore \frac{4 + 5\lambda}{1} = \frac{7 + 3\lambda}{-4} = \frac{4 + 10\lambda}{6}$$

$$\Rightarrow \lambda = -1$$

Substituting value of λ in equation (i), we get

$$-x + 4y - 6z + 106 = 0$$

$$\Rightarrow x - 4y + 6z = 106$$

$$\text{Hence } k = 106$$

52. The equations of the planes bisecting the angle between the given planes are

$$\frac{a_1x + b_1y + c_1z + d_1}{\sqrt{a_1^2 + b_1^2 + c_1^2}} = \pm \frac{a_2x + b_2y + c_2z + d_2}{\sqrt{a_2^2 + b_2^2 + c_2^2}}$$

$$\Rightarrow \frac{2x - y + 2z + 3}{\sqrt{2^2 + (-1)^2 + 2^2}} = \pm \frac{3x - 2y + 6z + 8}{\sqrt{3^2 + (-2)^2 + 6^2}}$$

$$\Rightarrow 7(2x - y + 2z + 3) = \pm 3(3x - 2y + 6z + 8)$$

$$\Rightarrow 7(2x - y + 2z + 3) = 3(3x - 2y + 6z + 8)$$

$$\text{or } 7(2x - y + 2z + 3) = -3(3x - 2y + 6z + 8)$$

$$\Rightarrow 5x - y - 4z - 3 = 0 \text{ or } 23x - 13y + 32z + 45 = 0$$

53. The point (3, -2, 1) satisfies both the equations so it is the point of intersection

Alternate method:

$$\text{Line is } \frac{x+3}{3} = \frac{y-2}{-2} = \frac{z+1}{1} = \lambda \text{ (say)}$$

$$x = 3\lambda - 3; y = -2\lambda + 2; z = \lambda - 1$$

Line intersects plane,

$$4x + 5y + 3z - 5 = 0$$

$$\therefore 4(3\lambda - 3) + 5(-2\lambda + 2) + 3(\lambda - 1) - 5 = 0$$

$$\Rightarrow \lambda = 2.$$

\therefore The point of intersection is (3, -2, 1)

54. The point (1, -2, 7) satisfies the given equation of plane. So it is the point of intersection.

Alternate method:

The d.r.s ratios of the line joining the points (2, -3, 1) and (3, -4, -5) are 1, -1, -6

\therefore The equation of line is

$$\frac{x-2}{1} = \frac{y+3}{-1} = \frac{z-1}{-6} = \lambda(\text{say})$$

Any general point on the line is

$$(\lambda + 2, -\lambda - 3, -6\lambda + 1)$$

The above point lies on the plane

$$2x + y + z = 7$$

$$\therefore 2(\lambda + 2) + (-\lambda - 3) + (-6\lambda + 1) = 7$$

$$\Rightarrow -5\lambda + 2 = 7$$

$$\Rightarrow \lambda = -1$$

\therefore The point is (1, -2, 7)

55. The equations of line is

$$\frac{x-3}{1} = \frac{y-4}{2} = \frac{z-5}{2} = \lambda(\text{say})$$

Any point on the line is $(\lambda + 3, 2\lambda + 4, 2\lambda + 5)$

Since the point lies on the plane $x + y + z = 17$

$$\therefore \lambda + 3 + 2\lambda + 4 + 2\lambda + 5 = 17 \Rightarrow \lambda = 1$$

\therefore The point is (4, 6, 7).

Hence, the required distance is

$$\sqrt{(3-4)^2 + (4-6)^2 + (5-7)^2}$$


$$= \sqrt{1^2 + 2^2 + 2^2} = 3$$



56. The d.r.s ratios of the line $\frac{x}{2} = \frac{y}{3} = \frac{z}{-6}$ are
2, 3, -6.

∴ The d.r.s of any line parallel to it are also
2, 3, -6.

∴ The equation of the line passing through
P(1, -2, 3) and parallel to the given line is
 $\frac{x-1}{2} = \frac{y+2}{3} = \frac{z-3}{-6} = \lambda(\text{say}) \dots(i)$



Any point on the line is

$$Q \equiv (2\lambda + 1, 3\lambda - 2, -6\lambda + 3)$$

The point Q lies on the plane $x - y + z = 5$.

$$\therefore (2\lambda + 1) - (3\lambda - 2) + (-6\lambda + 3) = 5$$

$$\Rightarrow -7\lambda = -1 \Rightarrow \lambda = \frac{1}{7}$$

$$\therefore Q \equiv \left(\frac{9}{7}, \frac{-11}{7}, \frac{15}{7}\right)$$

∴ Required distance = (PQ) = d

$$\begin{aligned} \therefore d &= \sqrt{\left(\frac{9}{7}-1\right)^2 + \left(\frac{-11}{7}+2\right)^2 + \left(\frac{15}{7}-3\right)^2} \\ &= \sqrt{\left(\frac{2}{7}\right)^2 + \left(\frac{3}{7}\right)^2 + \left(\frac{-6}{7}\right)^2} \\ &= \sqrt{\frac{4}{49} + \frac{9}{49} + \frac{36}{49}} = \sqrt{\frac{49}{49}} = 1 \end{aligned}$$

57. Let $\left. \begin{array}{l} \pi_1 : x + 2y + 3z = 5 \\ \pi_2 : x + 2y + 3z = 7 \end{array} \right\}$ be two given planes

Any plane parallel to the given planes and
equidistant from these is given by

$$x + 2y + 3z = \frac{d_1 + d_2}{2} = \frac{5 + 7}{2}$$

$$\text{i.e. } x + 2y + 3z = 6$$

58. Given planes are parallel,

∴ the required plane is also parallel to them

Let $3x + 4y + 5z + \lambda = 0$ be the required plane

$$\lambda = \frac{d_1 + d_2}{2} = \frac{-6 + 6}{2} = 0$$

∴ the equation of required plane is

$$3x + 4y + 5z = 0$$



Competitive Thinking

$$1. \quad \bar{n} = 2\hat{i} - 3\hat{j} + \hat{k}$$

$$\Rightarrow \hat{n} = \frac{1}{\sqrt{14}}(2\hat{i} - 3\hat{j} + \hat{k})$$

The equation of required plane is

$$\bar{r} \cdot \hat{n} = d$$

$$\Rightarrow \bar{r} \cdot \frac{1}{\sqrt{14}}(2\hat{i} - 3\hat{j} + \hat{k}) = \frac{3}{\sqrt{14}}$$

$$\Rightarrow \bar{r} \cdot (2\hat{i} - 3\hat{j} + \hat{k}) = 3$$

$$2. \quad \text{Let } A \equiv (-1, 1, 2)$$

$$\therefore \bar{a} = -\hat{i} + \hat{j} + 2\hat{k}$$

$$\bar{n} = \hat{i} + \hat{j} + \hat{k}$$

$$\therefore \text{equation of plane is } \bar{r} \cdot \bar{n} = \bar{a} \cdot \bar{n}$$

$$\Rightarrow \bar{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = (-\hat{i} + \hat{j} + 2\hat{k}) \cdot (\hat{i} + \hat{j} + \hat{k})$$

$$\Rightarrow \bar{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 2$$

$$3. \quad \text{The d.r.s. of the normal to the plane are } 1, 2, -3$$

∴ the d.c.s. of the normal to the plane are

$$\frac{1}{\sqrt{1^2 + 2^2 + (-3)^2}}, \frac{2}{\sqrt{1^2 + 2^2 + (-3)^2}}, \frac{-3}{\sqrt{1^2 + 2^2 + (-3)^2}}$$

i.e., $\frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{-3}{\sqrt{14}}$.

$$4. \quad \text{d.c.s of normal to the plane are}$$

$$\cos \frac{\pi}{4}, \cos \frac{\pi}{4}, \cos \frac{\pi}{2} = \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0$$

Equation of the plane is $x + my + nz = p$

$$\Rightarrow \frac{x}{\sqrt{2}} + \frac{y}{\sqrt{2}} = \sqrt{2}$$

$$\Rightarrow x + y = 2$$

$$5. \quad \text{The equation of plane passing through } (1, 2, -3) \text{ and } (2, -2, 1) \text{ and parallel to X-axis is}$$

$$\begin{vmatrix} x-1 & y-2 & z+3 \\ 2-1 & -2-2 & 1+3 \\ 1 & 0 & 0 \end{vmatrix} = 0$$

$$\Rightarrow (y-2)(4) + (z+3)(4) = 0$$

$$\Rightarrow y + z + 1 = 0$$

$$6. \quad \text{The plane passes through } (2, 3, 4)$$

This point satisfies the equation of plane in
option (D)

Also, it has d.r.s. 1, 2, 4.

∴ option (D) is correct answer.

**Alternate method:**

The equation of the required plane parallel to the plane $x + 2y + 4z = 5$ is

$$x + 2y + 4z + k = 0$$

The plane passes through (2, 3, 4)

$$\therefore 2 + 2(3) + 4(4) + k = 0$$

$$\Rightarrow k = -24$$

\therefore the equation of the required plane is

$$x + 2y + 4z = 24$$

7. The plane passes through (2, 3, 4)

This point satisfies the equation of plane in option (B)

Also, it has d.r.s. 5, -6, 7.

\therefore option (B) is correct answer.

8. The plane passes through (1, 2, 3)

This point satisfies the equation of plane in option (D)

Also, it has d.r.s. 2, 3, -4.

\therefore option (D) is correct answer.

9. $5x - 3y + 6z = 60$

$$\Rightarrow \frac{5x}{60} - \frac{3y}{60} + \frac{6z}{60} = 1 \Rightarrow \frac{x}{12} + \frac{y}{-20} + \frac{z}{10} = 1$$

\therefore the intercepts are (12, -20, 10).

10. The plane $x - 3y + 5z = d$ passes through (1, 2, 4).

$\therefore d = 15$

\therefore the equation of plane becomes $x - 3y + 5z = 15$

$$\Rightarrow \frac{x}{15} + \frac{y}{-5} + \frac{z}{3} = 1$$

\therefore length of intercept cut by plane on the X, Y, Z axes are 15, -5, 3 respectively.

11. The plane π is parallel to Y-axis.

\therefore Y intercept is zero

\therefore the equation of plane is $\frac{x}{4} + \frac{z}{3} = 1$

$$\Rightarrow 3x + 4z = 12$$

12. Here, $a = b = c = 1$

\therefore the equation of the required plane is $\frac{x}{1} + \frac{y}{1} + \frac{z}{1} = 1$

$$\Rightarrow x + y + z = 1$$

13. The intercepts made by the plane are $a, b, c = \frac{1}{m}, \frac{1}{n}$

\therefore The distances of plane from origin is

$$d = \frac{1}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2}}}$$

$$\Rightarrow k = \frac{1}{\sqrt{\frac{1}{2^2} + \frac{1}{m^2} + \frac{1}{n^2}}} \Rightarrow \frac{1}{2} + \frac{1}{m^2} + \frac{1}{n^2} = \frac{1}{k^2}$$

14. Let $P \equiv (2, 3, 4)$ and $Q \equiv (6, 7, 8)$

If R is the mid-point of PQ,

$\therefore R \equiv (4, 5, 6)$

This point satisfies the equation of plane in option (D)

\therefore option (D) is correct answer

Alternate method:

$$\vec{n} = 4\hat{i} + 4\hat{j} + 4\hat{k}, \vec{a} = 4\hat{i} + 5\hat{j} + 6\hat{k}$$

\therefore equation of plane is

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$

$$\Rightarrow \vec{r} \cdot (4\hat{i} + 4\hat{j} + 4\hat{k}) = (4\hat{i} + 5\hat{j} + 6\hat{k}) \cdot (4\hat{i} + 4\hat{j} + 4\hat{k})$$

$$\Rightarrow 4x + 4y + 4z = 16 + 20 + 24$$

$$\Rightarrow x + y + z - 15 = 0$$

15. The plane passes through (1, 2, 2)

This point satisfies the equation of plane in option (B)

Also, it has d.r.s. 1, 2, 2.

\therefore option (B) is correct answer.

16. Let M (1, 2, 3) be the foot of perpendicular from the origin O(0, 0, 0) to the plane d.r.s. of normal are 1, 2, 3

\therefore the equation of the required plane is

$$1(x-1) + 2(y-2) + 3(z-3) = 0$$

$$\Rightarrow x - 1 + 2y - 4 + 3z - 9 = 0$$

$$\Rightarrow x + 2y + 3z - 14 = 0$$

Consider the option (B)

point (7, 2, 1) satisfies the above equation of plane.

\therefore option (B) is correct answer.

17. The plane is $y = \frac{-8}{5}$ which is parallel to XZ-plane

\therefore Foot of the perpendicular drawn from the

$$\text{origin} \equiv \left(0, \frac{-8}{5}, 0\right)$$

18. The plane passes through (2, 6, 3)

It satisfies option (D)

Alternate Method:

The d.r.s of OP are 2 - 0, 6 - 0, 3 - 0 i.e., 2, 6, 3

The plane passes through P(2, 6, 3).

\therefore the equation of the required plane is

$$2(x-2) + 6(y-6) + 3(z-3) = 0$$

$$\Rightarrow 2x + 6y + 3z = 49$$

19. The plane passes through (1, 1, 1) and

(1, -1, -1)

The above points satisfies the equation of plane in option (B)

\therefore option (B) is correct answer.



20. The plane passes through $A(-2, 2, 2)$ and $B(2, -2, -2)$
The above points satisfies the equation of plane in option (A)

\therefore option (A) is correct answer.

21. The plane passes through $(0, 1, 2)$ and $(-1, 0, 3)$
The above points satisfies the equation of plane in option (D)

\therefore option (D) is correct answer.

22. The plane passes through $(2, -3, 1)$
This point satisfies the equation of plane in option (A)

Also, it has d.r.s. $3 - 2, 4 + 1, -1 - 5$
i.e. $1, 5, -6$.

\therefore option (A) is correct answer.

Alternate method:

The d.r.s. of the line joining the points $(3, 4, -1)$ and $(2, -1, 5)$ are $1, 5, -6$.

The plane passes through $(2, -3, 1)$

- \therefore the equation of required plane is
 $1(x - 2) + 5(y + 3) - 6(z - 1) = 0$
 $\Rightarrow x + 5y - 6z + 19 = 0$

23. The d.r.s. of the line joining the points $(4, -1, 2)$ and $(-3, 2, 3)$ are $7, -3, -1$
The plane passes through $(-10, 5, 4)$

- \therefore The equation of required plane is
 $7(x + 10) - 3(y - 5) - 1(z - 4) = 0$
 $\Rightarrow 7x + 70 - 3y + 15 - z + 4 = 0$
 $\Rightarrow 7x - 3y - z + 89 = 0$

24. The equation of the plane is
 $b(x - 1) + c(y - 1) + a(z - 1) = 0$ (i)
Now, $2001 = 3 \times 23 \times 29$

Since, $a < b < c$

- \therefore $a = 3, b = 23$ and $c = 29$

Substituting the values of a, b, c in equation (i), we get

$$23x + 29y + 3z = 55$$

25. $\vec{r} = (1 - p - q)\vec{a} + p\vec{b} + q\vec{c}$
 $\Rightarrow \vec{r} = \vec{a} + p(\vec{b} - \vec{a}) + q(\vec{c} - \vec{a})$ (i)

Comparing with $\vec{r} = \vec{A} + \lambda\vec{B} + \mu\vec{C}$,

the equation (i) represents a plane passing through a point having position vector \vec{a} and parallel to the vectors $\vec{b} - \vec{a}$ and $\vec{c} - \vec{a}$.

26. Equation of plane passing through $(1, 0, 2)$, $(-1, 1, 2)$ and $(5, 0, 3)$ is

$$\begin{vmatrix} x-1 & y-0 & z-2 \\ -1-1 & 1-0 & 2-2 \\ 5-1 & 0-0 & 3-2 \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} x-1 & y & z-2 \\ -2 & 1 & 0 \\ 4 & 0 & 1 \end{vmatrix} = 0$$

$$\Rightarrow (x - 1) - y(-2) + (z - 2)(-4) = 0$$

$$\Rightarrow x - 1 + 2y - 4z + 8 = 0$$

$$\Rightarrow x + 2y - 4z + 7 = 0$$

27. Equation of plane passing through $(1, 2, 3)$, $(-1, 4, 2)$ and $(3, 1, 1)$ is

$$\begin{vmatrix} x-1 & y-2 & z-3 \\ -1-1 & 4-2 & 2-3 \\ 3-1 & 1-2 & 1-3 \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} x-1 & y-2 & z-3 \\ -2 & 2 & -1 \\ 2 & -1 & -2 \end{vmatrix} = 0$$

$$\Rightarrow (x - 1)(-4 - 1) - (y - 2)(4 + 2) + (z - 3)(2 - 4) = 0$$

$$\Rightarrow -5x + 5 - 6y + 12 - 2z + 6 = 0$$

$$\Rightarrow -5x - 6y - 2z + 23 = 0$$

$$\Rightarrow 5x + 6y + 2z = 23$$

28. Equation of plane passing through $(1, 2, 3)$, $(2, 3, 1)$ and $(3, 1, 2)$ is

$$\begin{vmatrix} x-1 & y-2 & z-3 \\ 2-1 & 3-2 & 1-3 \\ 3-1 & 1-2 & 2-3 \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} x-1 & y-2 & z-3 \\ 1 & 1 & -2 \\ 2 & -1 & -1 \end{vmatrix} = 0$$

$$\Rightarrow (x - 1)(-3) - (y - 2)(3) + (z - 3)(-3) = 0$$

$$\Rightarrow -3x + 3 - 3y + 6 - 3z + 9 = 0$$

$$\Rightarrow x + y + z = 6$$

Comparing the above equation with $ax + by + cz = d$, we get

$$a = 1, b = 1, c = 1$$

$$\text{Now, } a + 2b + 3c = (1) + 2(1) + 3(1) = 6$$

29. The equation of the required plane is
 $(x + 2y + 3z + 4) + \lambda(4x + 3y + 2z + 1) = 0$ (i)

The plane passes through origin i.e., $(0, 0, 0)$

- \therefore $4 + \lambda = 0 \Rightarrow \lambda = -4$



- Substituting value of λ in equation (i), we get
 $-15x - 10y - 5z = 0$
 $\Rightarrow 3x + 2y + z = 0$
30. The plane passes through $(2, 1, 0)$
 It satisfies option (C)
 The equation of the required plane is
 $(x - 2y + 3z - 4) + \lambda(x - y + z - 3) = 0$ (i)
 The plane passes through $(2, 1, 0)$.
 $\therefore (2 - 2 + 0 - 4) + \lambda(2 - 1 + 0 - 3) = 0$
 $\Rightarrow \lambda = -2$
 Substituting value of λ in (i), we get
 $-x + z + 2 = 0$
 $\Rightarrow x - z = 2$
31. The d.r.s. of the line are 1, 2, 3.
 The line is perpendicular to the plane
 \therefore The d.r.s. of plane are 1, 2, 3
 \therefore The equation of plane passing through $(2, 3, 4)$ is
 $a(x - 2) + b(y - 3) + c(z - 4) = 0$ (i)
 $\Rightarrow 1(x - 2) + 2(y - 3) + 3(z - 4) = 0$
 $\Rightarrow x + 2y + 3z = 20$
32. The plane passes through the line
 $\frac{x-3}{1} = \frac{y-6}{5} = \frac{z-4}{4}$ i.e. through $(3, 6, 4)$
 The points $(3, 2, 0)$ and $(3, 6, 4)$ satisfies
 option (A)
 \therefore option (A) is correct answer.
Alternate method:
 The equation of plane passing through $(3, 2, 0)$ is
 $a(x - 3) + b(y - 2) + c(z - 0) = 0$ (i)
 $\therefore a(3 - 3) + b(6 - 2) + c(4 - 0) = 0$
 $\Rightarrow 0 \cdot a + 4b + 4c = 0$ (ii)
 and $1 \cdot a + 5b + 4c = 0$ (iii)
 On solving (ii) and (iii), we get
 $a = -4, b = 4, c = -4$
 \therefore equation of required plane is $x - y + z = 1$
33. The equation of plane passing through $(2, -1, -3)$
 is
 $a(x - 2) + b(y + 1) + c(z + 3) = 0$
 Also, as the plane is parallel to the given two
 lines,
 $\therefore 3a + 2b - 4c = 0$ and $2a - 3b + 2c = 0$
 $\Rightarrow a = -8, b = -14, c = -13$
 \therefore The equation of the required plane is
 $-8(x - 2) - 14(y + 1) - 13(z + 3) = 0$
 $\Rightarrow 8x + 14y + 13z + 37 = 0$
34. Point $(2, 1, -2)$ lies in the plane
 $x + 3y - \alpha z + \beta = 0$
 $\therefore 2 + 3(1) - \alpha(-2) + \beta = 0$
 $\Rightarrow 2\alpha + \beta = -5$ (i)

Also, the d.r.s. of the normal are perpendicular
 to the given plane.

- $\therefore 3(1) + (-5)(3) + (2)(-\alpha) = 0$
 $\Rightarrow 3 - 15 - 2\alpha = 0$
 $\Rightarrow \alpha = -6$
 Substituting value of α in equation (i), we get
 $\beta = 7$
35. The d.r.s. of normal to the given planes are
 $1, 2, 2$ and $-5, 3, 4$
 $\therefore \cos \theta = \frac{(1)(-5) + (2)(3) + (2)(4)}{\sqrt{1^2 + 2^2 + 2^2} \sqrt{(-5)^2 + 3^2 + 4^2}} = \frac{3\sqrt{2}}{10}$
 $\Rightarrow \theta = \cos^{-1} \left(\frac{3\sqrt{2}}{10} \right)$
36. $\cos \theta = \frac{3(2) - 4(-1) + 5(-2)}{\sqrt{9 + 16 + 25} \sqrt{4 + 1 + 4}}$
 $\therefore \cos \theta = 0$
 $\Rightarrow \theta = \frac{\pi}{2}$
37. Given equation of locus $xy + yz = 0$
 $\Rightarrow y(x + z) = 0$
 $\Rightarrow y = 0$ or $x + z = 0$
 The planes $y = 0$ and $x + z = 0$ perpendicular
 to each other.
38. $\vec{r} \cdot (\hat{m}\hat{i} - \hat{j} + 2\hat{k}) + 3 = 0 \Rightarrow \vec{r} \cdot (\hat{m}\hat{i} - \hat{j} + 2\hat{k}) = -3$
 $\vec{r} \cdot (2\hat{i} - \hat{m}\hat{j} - \hat{k}) - 5 = 0 \Rightarrow \vec{r} \cdot (2\hat{i} - \hat{m}\hat{j} - \hat{k}) = 5$
 Here, $\vec{n}_1 = \hat{m}\hat{i} - \hat{j} + 2\hat{k}$ and $\vec{n}_2 = 2\hat{i} - \hat{m}\hat{j} - \hat{k}$
 $\therefore \cos \theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| |\vec{n}_2|}$
 $\Rightarrow \cos \frac{\pi}{3} = \frac{|(\hat{m}\hat{i} - \hat{j} + 2\hat{k}) \cdot (2\hat{i} - \hat{m}\hat{j} - \hat{k})|}{\sqrt{m^2 + 1 + 4} \sqrt{4 + m^2 + 1}}$
 $\Rightarrow \frac{1}{2} = \frac{2m + m - 2}{m^2 + 5}$
 ... (Considering positive value)
 $\Rightarrow m^2 + 5 = 6m - 4$
 $\Rightarrow m^2 - 6m + 9 = 0$
 $\Rightarrow (m - 3)^2 = 0$
 $\Rightarrow m = 3$



39. Here, $\vec{n}_1 = p\hat{i} - \hat{j} + 2\hat{k}$ and $\vec{n}_2 = 2\hat{i} - p\hat{j} - \hat{k}$

$$\begin{aligned} \therefore \cos\theta &= \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| |\vec{n}_2|} \\ \Rightarrow \cos\frac{\pi}{3} &= \frac{|(p\hat{i} - \hat{j} + 2\hat{k}) \cdot (2\hat{i} - p\hat{j} - \hat{k})|}{\sqrt{p^2 + 1 + 4} \sqrt{4 + p^2 + 1}} \\ \Rightarrow \frac{1}{2} &= \pm \left(\frac{2p + p - 2}{p^2 + 5} \right) \\ \Rightarrow \frac{1}{2} &= \frac{3p - 2}{p^2 + 5} \end{aligned}$$

..... (considering positive value)

$$\begin{aligned} \Rightarrow p^2 + 5 &= 6p - 4 \\ \Rightarrow p^2 - 6p + 9 &= 0 \\ \Rightarrow (p - 3)^2 &= 0 \\ \Rightarrow p &= 3 \end{aligned}$$

40. Let the d.r.s of the normal to the plane be proportional to a, b, c.

It passes through (1, 0, 0)

\therefore the equation of the plane is
 $a(x - 1) + b(y - 0) + c(z - 0) = 0$ (i)

Also, the plane passes through (0, 1, 0).

$\therefore a(-1) + b(1) + c(0) = 0$
 $\Rightarrow a = b$ (ii)

Now, the angle between the required plane and the plane $x + y = 3$ is $\frac{\pi}{4}$.

$$\begin{aligned} \therefore \cos\frac{\pi}{4} &= \frac{a(1) + b(1) + c(0)}{\sqrt{a^2 + b^2 + c^2} \sqrt{1 + 1}} \\ \Rightarrow \frac{1}{\sqrt{2}} &= \frac{a + b}{\sqrt{a^2 + b^2 + c^2} \sqrt{2}} \end{aligned}$$

Squaring both sides, we get
 $\Rightarrow a^2 + b^2 + c^2 = a^2 + b^2 + 2ab$
 $\Rightarrow c^2 = 2ab$ (iii)

From (ii) and (iii), we get
 $a : b : c = a : a : \sqrt{2}a = 1 : 1 : \sqrt{2}$

41. For perpendicular planes, $a_1a_2 + b_1b_2 + c_1c_2 = 0$
 $\Rightarrow 2(1) + 1(2) - 2(k) = 0$
 $\Rightarrow k = 2$

42. Since the planes are perpendicular,
 $\therefore (3)(2) + (-6)(1) + (-2)(-k) = 0$
 $\Rightarrow k = 0$

43. Since, the planes are perpendicular to each other.
 $\therefore 3(4) + (-2)(3) + 2 \times (-k) = 0$
 $\Rightarrow k = 3$

44. The equation of plane passing through (4, 4, 0) is $a(x - 4) + b(y - 4) + c(z - 0) = 0$

$$\Rightarrow a(x - 4) + b(y - 4) + cz = 0 \quad \dots(i)$$

Since, plane (i) is perpendicular to the planes $2x + y + 2z + 3 = 0$ and $3x + 3y + 2z - 8 = 0$

$$\therefore 2a + b + 2c = 0, \text{ and} \quad \dots(ii)$$

$$3a + 3b + 2c = 0 \quad \dots(iii)$$

On solving (i) and (ii), we get

$$a = -4, b = 2, c = 3$$

Substituting the values of a, b, c in (i), we get

$$-4(x - 4) + 2(y - 4) + 3z = 0$$

$$\Rightarrow -4x + 16 + 2y - 8 + 3z = 0$$

$$\Rightarrow 4x - 2y - 3z = 8$$

45. Comparing with $\vec{r} = \vec{a} + \lambda\vec{b}$ and $\vec{r} \cdot \vec{n} = p$, we get

$$\vec{b} = -\hat{i} + \hat{j} + \hat{k} \text{ and } \vec{n} = 3\hat{i} + 2\hat{j} - \hat{k}$$

\therefore Angle between the line and plane is given by

$$\begin{aligned} \sin\theta &= \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| |\vec{n}|} \\ &= \frac{(-1)(3) + (1)(2) + (1)(-1)}{\sqrt{3} \sqrt{14}} = \frac{-2}{\sqrt{42}} \\ \Rightarrow \theta &= \sin^{-1} \left(\frac{-2}{\sqrt{42}} \right) \end{aligned}$$

46. The d.r.s. of line are 3, 4, 5 and

the d.r.s. of normal to the plane are 2, -2, 1

\therefore The angle between line and plane is

$$\begin{aligned} \sin\theta &= \frac{aa_1 + bb_1 + cc_1}{\sqrt{a^2 + b^2 + c^2} \cdot \sqrt{a_1^2 + b_1^2 + c_1^2}} \\ &= \frac{(2)(3) + (-2)(4) + (1)(5)}{\sqrt{2^2 + (-2)^2 + (1)^2} \cdot \sqrt{3^2 + 4^2 + 5^2}} \\ &= \frac{3}{\sqrt{9} \sqrt{50}} = \frac{1}{5\sqrt{2}} = \frac{\sqrt{2}}{10} \end{aligned}$$

47. The d.r.s. of line are 1, 2, 2 and

the d.r.s. of normal to the plane are 2, -1, $\sqrt{\lambda}$

$$\therefore \sin\theta = \frac{1(2) + 2(-1) + 2(\sqrt{\lambda})}{\sqrt{1 + 4 + 4} \cdot \sqrt{4 + 1 + \lambda}}$$

$$\Rightarrow \frac{1}{3} = \frac{2\sqrt{\lambda}}{3\sqrt{5 + \lambda}}$$

$$\Rightarrow 2\sqrt{\lambda} = \sqrt{5 + \lambda}$$

$$\Rightarrow 4\lambda = 5 + \lambda$$

$$\Rightarrow \lambda = \frac{5}{3}$$



48. d.r.s. of normal to the plane are 2, -3, 6
d.r.s. of X-axis are 1, 0, 0.
∴ The angle between the plane and X-axis is

$$\sin \theta = \frac{aa_1 + bb_1 + cc_1}{\sqrt{a^2 + b^2 + c^2} \sqrt{a_1^2 + b_1^2 + c_1^2}}$$

$$= \frac{2(1) - 3(0) + 6(0)}{\sqrt{4 + 9 + 36} \cdot \sqrt{1}}$$

$$= \frac{2}{7}$$
∴ $\theta = \sin^{-1} \left(\frac{2}{7} \right)$
But $\theta = \sin^{-1} \alpha$
∴ $\alpha = \frac{2}{7}$
49. The d.r.s. of line are 1, 2, λ and
The d.r.s. of normal to the plane are 1, 2, 3.
∴ $\sin \theta = \frac{1(1) + 2(2) + \lambda(3)}{\sqrt{1+4+9} \sqrt{1+4+\lambda^2}}$

$$\Rightarrow \sin \theta = \frac{5+3\lambda}{\sqrt{14} \sqrt{5+\lambda^2}}$$

$$\Rightarrow \sin^2 \theta = \frac{(5+3\lambda)^2}{14(5+\lambda^2)}$$

$$\Rightarrow 1 - \frac{5}{14} = \frac{(5+3\lambda)^2}{14(5+\lambda^2)}$$

$$\dots \left[\because \cos \theta = \sqrt{\frac{5}{14}} \text{ (given)} \right]$$

$$\Rightarrow \frac{9}{14} = \frac{25+30\lambda+9\lambda^2}{14(5+\lambda^2)}$$
On solving, we get
 $\lambda = \frac{2}{3}$
50. Let a, b, c = 3, 2 + λ , -1 and $a_1, b_1, c_1 = 1, -2, 0$
Since, the line lies on the plane
∴ $aa_1 + bb_1 + cc_1 = 0$
 $\Rightarrow 3(1) + (2 + \lambda)(-2) + (-1)(0) = 0$
 $\Rightarrow \lambda = \frac{-1}{2}$
51. The line is parallel to the plane if
 $aa_1 + bb_1 + cc_1 = 0$
Consider option (B), $2(3) + 1(4) - 2(5) = 0$
∴ $2x + y - 2z = 0$ is the required plane.
52. The equation of the plane is $ax + by + cz + d = 0$
∴ the d.r.s. of the normal to the plane are a, b, c
Since the given line is parallel to the plane,
∴ $a + bm + cn = 0$
53. line $\frac{x-4}{1} = \frac{y-2}{1} = \frac{z-k}{2}$ lies on the plane
 $2x - 4y + z = 7$.
∴ Point (4, 2, k) lies on the plane $2x - 4y + z = 7$
∴ $2(4) - 4(2) + k = 7$
 $\Rightarrow k = 7$
54. Line is perpendicular to normal of plane
 $\Rightarrow (2\hat{i} - \hat{j} + 3\hat{k}) \cdot (\hat{i} + m\hat{j} - \hat{k}) = 0$
 $2 - m - 3 = 0 \quad \dots(i)$
(3, -2, -4) lies on the plane $x + my - z = 9$
∴ $3 - 2m + 4 = 9$
 $\Rightarrow 3 - 2m = 5 \quad \dots(ii)$
Solving (i) and (ii)
 $= 1, m = -1$
 $^2 + m^2 = 2$
55. The d.r.s. of the XY-plane are 0, 0, 1
the d.r.s. of the given line are , m, n
Since, the line is parallel to the plane
∴ $aa_1 + bb_1 + cc_1 = 0$
 $\Rightarrow (0) + m(0) + n(1) = 0$
 $\Rightarrow n = 0$
56. Let the position vector of Q be
 $(\hat{i} - \hat{j} + 2\hat{k}) + \mu(-3\hat{i} + \hat{j} + 5\hat{k})$
 $= (-3\mu + 1)\hat{i} + (\mu - 1)\hat{j} + (5\mu + 2)\hat{k}$
∴ $\overline{PQ} = (-3\mu - 2)\hat{i} + (\mu - 3)\hat{j} + (5\mu - 4)\hat{k}$
Since, \overline{PQ} is parallel to the plane
∴ $(-3\mu - 2)(1) + (\mu - 3)(-4) + (5\mu - 4)(3) = 0$
 $\Rightarrow \mu = \frac{1}{4}$
57. The plane passes through points (-3, 0, 2) and
(3, 2, 6)
This points satisfies the equation of plane in
option (D)
∴ option (D) is correct answer.
58. Lines are coplanar if

$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$
∴
$$\begin{vmatrix} 1-2 & 4-3 & 5-4 \\ 1 & 1 & -k \\ k & 2 & 1 \end{vmatrix} = 0$$



$$\Rightarrow \begin{vmatrix} -1 & 1 & 1 \\ 1 & 1 & -k \\ k & 2 & 1 \end{vmatrix} = 0$$

$$\Rightarrow -1(1 + 2k) - 1(1 + k^2) + 1(2 - k) = 0$$

$$\Rightarrow k^2 + 3k = 0$$

$$\Rightarrow k = 0, -3$$

59. The planes are concurrent,

$$\therefore \begin{vmatrix} 1 & -c & -b \\ c & -1 & a \\ b & a & -1 \end{vmatrix} = 0$$

$$\Rightarrow a^2 + b^2 + c^2 + 2abc - 1 = 0$$

$$\Rightarrow a^2 + b^2 + c^2 = 1 - 2abc$$

60. The equation of the plane is $\frac{x}{8} + \frac{y}{4} + \frac{z}{4} = 1$

$$\text{i.e., } x + 2y + 2z = 8$$

\therefore The length of the perpendicular from origin to the plane is

$$d = \left| \frac{-8}{\sqrt{1+4+4}} \right| = \frac{8}{3}$$

61. The equations of the plane with reference to the two systems of rectangular axes are

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1 \quad \dots(i)$$

$$\text{and } \frac{X}{a'} + \frac{Y}{b'} + \frac{Z}{c'} = 1 \quad \dots(ii)$$

Since the origin of axes is same.

\therefore Length of the perpendicular from (0, 0, 0) on plane (i)


= Length of the perpendicular from (0, 0, 0) on plane (ii)

$$\Rightarrow \left| \frac{-1}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2}}} \right| = \left| \frac{-1}{\sqrt{\frac{1}{a'^2} + \frac{1}{b'^2} + \frac{1}{c'^2}}} \right|$$

$$\Rightarrow \frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} - \frac{1}{a'^2} - \frac{1}{b'^2} - \frac{1}{c'^2} = 0$$

62. Since the line is parallel to XY-plane, the distance of the point P (6, 7, 8) from this plane is equal to its Z co-ordinate i.e. 8 units.

63.



Distance of point P from the given plane is given by

$$d = \left| \frac{3(-6) - 6(2) + 2(3) + 10}{\sqrt{(3)^2 + (-6)^2 + (2)^2}} \right|$$

$$= \left| \frac{-18 - 12 + 6 + 10}{\sqrt{9 + 36 + 4}} \right|$$

$$= \left| \frac{-14}{\sqrt{49}} \right| = \frac{14}{7}$$

$$\therefore d = 2$$

64. Given equation of plane is $\vec{r} \cdot (3\hat{i} + 2\hat{j} + 6\hat{k}) = 13$

The vector form of the equation is

$$3x + 2y + 6z = 13$$

$$\Rightarrow 3x + 2y + 6z - 13 = 0$$

Given point $\equiv (2, 3, \lambda)$

\therefore Distance of the point from the plane

$$= \left| \frac{ax_1 + by_1 + cz_1 + d}{\sqrt{a^2 + b^2 + c^2}} \right|$$

$$\Rightarrow 5 = \left| \frac{3(2) + 2(3) + 6(\lambda) - 13}{\sqrt{9 + 4 + 36}} \right|$$

$$\Rightarrow 5 = \left| \frac{6\lambda - 1}{7} \right|$$

$$\Rightarrow 6\lambda - 1 = \pm 35$$

$$\Rightarrow \lambda = 6, \frac{-17}{3}$$

65. Here, $a = 2, b = 1, c = 2, d = 5, x = 2, y = 1, z = 0$

$$\therefore d = \left| \frac{2(2) + 1(1) + 2(0) + 5}{\sqrt{2^2 + 1^2 + 2^2}} \right|$$

$$= \left| \frac{10}{\sqrt{9}} \right| = \frac{10}{3}$$

$$66. \text{ Normal vector } \hat{n} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -2 & 3 \\ 2 & -1 & -1 \end{vmatrix}$$

$$= \hat{i}(2+3) - \hat{j}(-1-6) + \hat{k}(-1+4)$$

$$= 5\hat{i} + 7\hat{j} + 3\hat{k}$$

Let $A \equiv (1, -1, -1)$

$$\therefore \vec{a} = \hat{i} - \hat{j} - \hat{k}$$

\therefore Equation of the plane is

$$5(x - 1) + 7(y + 1) + 3(z + 1) = 0$$

$$\Rightarrow 5x + 7y + 3z + 5 = 0$$



Distance of $(1, 3, -7)$ from the above plane is

$$d = \left| \frac{5(1) + 7(3) + 3(-7) + 5}{\sqrt{25 + 49 + 9}} \right|$$

$$= \frac{10}{\sqrt{83}} \text{ units}$$

67. $\vec{n} = 2\hat{i} + \hat{j} + 2\hat{k}$ and $p = 5$

$$\hat{n} = \frac{\vec{n}}{|\vec{n}|} = \frac{2\hat{i} + \hat{j} + 2\hat{k}}{\sqrt{4+1+4}} = \frac{2\hat{i} + \hat{j} + 2\hat{k}}{3}$$

\therefore The vector equation of the plane is $\vec{r} \cdot \hat{n} = p$

$$\Rightarrow \vec{r} \cdot \left(\frac{2\hat{i} + \hat{j} + 2\hat{k}}{3} \right) = 5$$

$$\Rightarrow \vec{r} \cdot (2\hat{i} + \hat{j} + 2\hat{k}) = 15$$

68. Let $a, b, c = -3, 2, 6$

\therefore the equation of plane is

$$\therefore -3x + 2y + 6z + d = 0 \quad \dots(i)$$

Now, the perpendicular distance (D) from origin is

$$D = \left| \frac{d}{\sqrt{(-3)^2 + 2^2 + 6^2}} \right|$$

$$\Rightarrow 7 = \frac{|d|}{7} \Rightarrow d = \pm 49$$

\therefore The equation of plane is

$$-3x + 2y + 6z + 49 = 0$$

$$\text{or } -3x + 2y + 6z - 49 = 0$$

69. The equation of a plane passing through $(1, -2, 1)$ is

$$a(x-1) + b(y+2) + c(z-1) = 0 \quad \dots(i)$$

Plane (i) is perpendicular to planes

$$2x - 2y + z = 0 \text{ and } x - y + 2z = 4.$$

$$\therefore 2a - 2b + c = 0, \text{ and} \quad \dots(ii)$$

$$a - b + 2c = 0 \quad \dots(iii)$$

Solving (ii) and (iii), we get

$$a = -3, b = -3, c = 0$$

Substituting the values of a, b, c in equation (i), we get

$$x + y + 1 = 0$$

\therefore The distance of this plane from $(1, 2, 2)$ is

$$d = \left| \frac{1+2+1}{\sqrt{1+1}} \right| = 2\sqrt{2}$$

70. Equation of L_1 i.e., the line of intersection of the first two given planes is

$$(2x - 2y + 3z - 2) + \lambda(x - y + z + 1) = 0$$

$$\Rightarrow (\lambda + 2)x - (2 + \lambda)y + (\lambda + 3)z + (\lambda - 2) = 0 \quad \dots(i)$$

Equation of L_2 i.e., the line of intersection of the next two given planes is

$$(1 + 3\mu)x + (2 - \mu)y + (2\mu - 1)z - (\mu + 3) = 0 \quad \dots(ii)$$

Since, equations (i) and (ii) represent the same plane.

\therefore by comparing, we get

$$\frac{2 + \lambda}{1 + 3\mu} = \frac{-(2 + \lambda)}{2 - \mu}$$

$$\Rightarrow 1 + 3\mu = \mu - 2 \quad \Rightarrow \mu = -\frac{3}{2}$$

Substituting $\mu = -\frac{3}{2}$ in (ii), we get

$$7x - 7y + 8z + 3 = 0$$

Perpendicular distance from the origin $(0, 0, 0)$

$$= \left| \frac{7(0) - 7(0) + 8(0) + 3}{\sqrt{7^2 + 7^2 + 8^2}} \right| = \frac{3}{\sqrt{162}}$$

$$= \frac{3}{9\sqrt{2}} = \frac{1}{3\sqrt{2}}$$

71. The equation of a plane passing through the line of intersection of the planes

$$x + 2y + 3z = 2 \text{ and } x - y + z = 3 \text{ is}$$

$$(x + 2y + 3z - 2) + \lambda(x - y + z - 3) = 0$$

$$\Rightarrow x(1 + \lambda) + y(2 - \lambda) + z(3 + \lambda) - 2 - 3\lambda = 0 \quad \dots(i)$$

This plane is at a distance of $\frac{2}{\sqrt{3}}$ units from

$$(3, 1, -1).$$

$$\therefore \frac{|3(1 + \lambda) + 1(2 - \lambda) - 1(3 + \lambda) - 2 - 3\lambda|}{\sqrt{(1 + \lambda)^2 + (2 - \lambda)^2 + (3 + \lambda)^2}} = \frac{2}{\sqrt{3}}$$

$$\Rightarrow \frac{|-2\lambda|}{\sqrt{3\lambda^2 + 4\lambda + 14}} = \frac{2}{\sqrt{3}}$$

Squaring both sides, we get

$$3\lambda^2 + 4\lambda + 14 = 3\lambda^2 \Rightarrow 4\lambda = -14 \Rightarrow \lambda = \frac{-7}{2}$$

Substituting value of λ in equation (i), we get

$$-\frac{5}{2}x + \frac{11}{2}y - \frac{z}{2} + \frac{17}{2} = 0$$

$$\Rightarrow 5x - 11y + z - 17 = 0$$

72. The equation of the plane passing through $(-1, 3, 0)$ is

$$a(x + 1) + b(y - 3) + c(z - 0) = 0 \quad \dots(i)$$

Also, the plane passes through the points $(2, 2, 1)$ and $(1, 1, 3)$.

$$\therefore 3a - b + c = 0 \quad \dots(ii)$$

$$2a - 2b + 3c = 0 \quad \dots(iii)$$



Solving (ii) and (iii), we get

$$a = -1, b = -7, c = -4$$

Substituting the values of a, b, c in equation (i), we get

$$-1(x+1) - 7(y-3) - 4(z) = 0$$

$$\Rightarrow x + 7y + 4z - 20 = 0$$

\(\therefore\) The distance of this plane from the point (5, 7, 8) is

$$d = \frac{|1(5) + 7(7) + 4(8) - 20|}{\sqrt{1^2 + 7^2 + 4^2}} = \frac{66}{\sqrt{66}} = \sqrt{66}$$

73. Given planes are

$$2x + y + 2z - 8 = 0$$

$$4x + 2y + 4z - 16 = 0 \quad \dots(i)$$

$$\text{and } 4x + 2y + 4z + 5 = 0 \quad \dots(ii)$$

The distance between two parallel planes is

$$d = \frac{|d_1 - d_2|}{\sqrt{a^2 + b^2 + c^2}} = \frac{|-16 - 5|}{\sqrt{4^2 + 2^2 + 4^2}} = \frac{21}{6} = \frac{7}{2}$$

74. $x^2 - 5x + 6 = 0$

\(\Rightarrow\) $(x - 2) = 0$ or $(x - 3) = 0$, which represents a plane.

75. Here, the co-ordinates of A, B, C are $(3a, 0, 0)$, $(0, 3b, 0)$ and $(0, 0, 3c)$ respectively.

\(\therefore\) The centroid is (a, b, c) .

76. Let $A \equiv (a, 0, 0)$, $B \equiv (0, b, 0)$ and $C \equiv (0, 0, c)$
The equation of the plane in intercept form is

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

Since, centroid is $(6, 6, 3)$

$$\therefore 6 = \frac{x_1 + x_2 + x_3}{3}$$

$$\Rightarrow 6 = \frac{a + 0 + 0}{3} \Rightarrow a = 18$$

$$\text{Similarly } \frac{0 + b + 0}{3} = 6 \Rightarrow b = 18$$

$$\frac{0 + 0 + c}{3} = 3 \Rightarrow c = 9$$

\(\therefore\) The equation of plane is $\frac{x}{18} + \frac{y}{18} + \frac{z}{9} = 1$

$$\Rightarrow x + y + 2z - 18 = 0$$

77. Given equation of plane is $ax + by + cz = 1$

$$\therefore A \equiv \left(\frac{1}{a}, 0, 0\right), B \equiv \left(0, \frac{1}{b}, 0\right) \text{ and}$$

$$C \equiv \left(0, 0, \frac{1}{c}\right)$$

$$\therefore \text{Centroid} \equiv \left(\frac{1}{3a}, \frac{1}{3b}, \frac{1}{3c}\right) = \left(\frac{1}{6}, \frac{-1}{3}, 1\right)$$

$$\therefore 3a = 6 \Rightarrow a = 2$$

$$3b = -3 \Rightarrow b = -1$$

$$3c = 1 \Rightarrow c = \frac{1}{3}$$

$$\therefore a + b + 3c = 2 - 1 + 3\left(\frac{1}{3}\right) = 2$$

$$78. [a \ b \ c] \begin{bmatrix} 1 & 9 & 7 \\ 8 & 2 & 7 \\ 7 & 3 & 7 \end{bmatrix} = [0 \ 0 \ 0]$$

$$\Rightarrow a + 8b + 7c = 0, 9a + 2b + 3c = 0,$$

$$7a + 7b + 7c = 0$$

$$\Rightarrow a = 1, b = 6, c = -7$$

$P(a, b, c)$ lies on the plane $2x + y + z = 1$.

$$\therefore 7a + b + c = 7 + 6 - 7 = 6$$

79. The equation of the required plane is

$$(2x - 5y + z - 3) + \lambda(x + y + 4z - 5) = 0 \quad \dots(i)$$

$$\therefore (2 + \lambda)x + (-5 + \lambda)y + (1 + 4\lambda)z + (-3 - 5\lambda) = 0$$

Since, this plane is parallel to $x + 3y + 6z = 1$

$$\therefore \frac{2 + \lambda}{1} = \frac{-5 + \lambda}{3} = \frac{1 + 4\lambda}{6}$$

On solving, we get

$$\lambda = -\frac{11}{2}$$

Substituting the value of λ in equation (i), we get

$$(2x - 5y + z - 3) - \frac{11}{2}(x + y + 4z - 5) = 0$$

$$\Rightarrow -7x - 21y - 42z + 49 = 0$$

$$\Rightarrow x + 3y + 6z - 7 = 0$$

80. The point $\left(\frac{-1}{11}, \frac{9}{11}, \frac{-25}{11}\right)$ satisfies both the equations

\(\therefore\) it is the point of intersection.

Alternate method:

$$\text{Let } \frac{x}{1} = \frac{y-1}{2} = \frac{z+2}{3} = \lambda (\text{say})$$

Any general point on the line is $(\lambda, 2\lambda+1, 3\lambda-2)$

This point lies on the plane $2x + 3y + z = 0$

$$\therefore 2\lambda + 3(2\lambda + 1) + (3\lambda - 2) = 0$$

$$\Rightarrow \lambda = \frac{-1}{11}$$

$$\therefore x = \frac{-1}{11}, y = \frac{9}{11}, z = \frac{-25}{11}$$

\(\therefore\) The point is $\left(\frac{-1}{11}, \frac{9}{11}, \frac{-25}{11}\right)$



81. The point $(5, -1, 1)$ satisfies both the equations

\therefore it is the point of intersection

\therefore option (D) is correct

82. The point $(10, 10, 3)$ satisfies both the equations.

\therefore it is the point of intersection.

\therefore option (B) is correct

83. The point $(-4, -3, 0)$ satisfies the given equations

\therefore correct answer is option (D).

84. Let $\frac{x-2}{3} = \frac{y+1}{4} = \frac{z-2}{12} = \lambda$

\therefore the co-ordinates of any point on the line are

$$P \equiv (3\lambda + 2, 4\lambda - 1, 12\lambda + 2)$$

This point lies on the plane $x - y + z = 16$

$\therefore 3\lambda + 2 - 4\lambda + 1 + 12\lambda + 2 = 16$

$$\Rightarrow 11\lambda = 11 \Rightarrow \lambda = 1$$


$\therefore P \equiv (5, 3, 14)$

\therefore Let $Q \equiv (1, 0, 2)$

\therefore distance PQ is given by

$$d = \sqrt{(5-1)^2 + (3-0)^2 + (14-2)^2} = 13$$

85.



Distance of point $P(1, -2, 1)$ from the $x + 2y - 2z = \alpha$ plane is 5

$$\therefore \frac{|1 - 4 - 2 - \alpha|}{\sqrt{1 + 4 + 4}} = 5$$

$$\Rightarrow |\alpha + 5| = 15$$

$$\Rightarrow \alpha + 5 = \pm 15$$

$$\Rightarrow \alpha = 10, -20$$

$$\Rightarrow \alpha = 10 \quad \dots (\because \alpha > 0)$$

The equation of line PM whose d.r.s. are $1, 2, -2$ is

$$\frac{x-1}{1} = \frac{y+2}{2} = \frac{z-1}{-2} = \lambda(\text{say})$$

The co-ordinates of M are $(\lambda + 1, 2\lambda - 2, -2\lambda + 1)$

Since, M lies on the plane

$$x + 2y - 2z = 10$$

$$\therefore \lambda + 1 + 4\lambda - 4 + 4\lambda - 2 = 10$$

$$\Rightarrow 9\lambda = 15 \Rightarrow \lambda = \frac{5}{3}$$


Hence, the co-ordinates of M are $\left(\frac{8}{3}, \frac{4}{3}, \frac{-7}{3}\right)$.

86. The d.r.s. of the line $\frac{x}{1} = \frac{y}{4} = \frac{z}{5}$ are $1, 4, 5$

\therefore The d.r.s. of any line parallel to it are also $1, 4, 5$

The equation of the line passing through $Q(1, -2, 3)$

$$\frac{x-1}{1} = \frac{y+2}{4} = \frac{z-3}{5} = \lambda(\text{say}) \quad \dots(i)$$



Any point on the line is

$$P \equiv (\lambda + 1, 4\lambda - 2, 5\lambda + 3)$$

The point P lies on the plane

$$2x + 3y - 4z + 22 = 0$$

$$\therefore 2(\lambda + 1) + 3(4\lambda - 2) - 4(5\lambda + 3) + 22 = 0$$

$$\Rightarrow 6\lambda = 6 \quad \Rightarrow \lambda = 1$$

$$\therefore P = (2, 2, 8)$$

$$\therefore \text{Required distance} = (PQ) = d$$

$$d = \sqrt{(2-1)^2 + (2+2)^2 + (8-3)^2}$$

$$= \sqrt{1 + 16 + 25}$$

$$\therefore d = \sqrt{42} \text{ units}$$

87. Since line PQ is parallel to line $\frac{x}{1} = \frac{y}{4} = \frac{z}{5}$

\therefore d.r.s. of PQ are $1, 4, 5$

\therefore Equation of line PQ passing through $P(1, -2, 3)$ is

$$\frac{x-1}{1} = \frac{y+2}{4} = \frac{z-3}{5}$$

$$\text{Let } \frac{x-1}{1} = \frac{y+2}{4} = \frac{z-3}{5} = \lambda$$

Any point R on PQ $\equiv (\lambda + 1, 4\lambda - 2, 5\lambda + 3)$



Since point R lies in the plane

$$2x + 3y - 4z + 22 = 0$$

$$\therefore 2(\lambda + 1) + 3(4\lambda - 2) - 4(5\lambda + 3) + 22 = 0$$

$$\Rightarrow -6\lambda + 6 = 0$$

$$\Rightarrow \lambda = 1$$

$$\therefore R \equiv (2, 2, 8)$$


$$PQ = 2PR$$

$$= 2\sqrt{(2-1)^2 + (2+2)^2 + (8-3)^2}$$

$$= 2\sqrt{42} \text{ units}$$

88. Let $A = (5, -1, 4)$, $B = (4, -1, 3)$

$$\overline{AB} = -\hat{i} - \hat{k} \Rightarrow |\overline{AB}| = \sqrt{2}$$



Projection of \overline{AB} in the plane $x + y + z = 7$

$$\text{is } |\overline{AB}| \cos \theta = |\overline{A'B'}| \cos \theta$$

Direction ratios of normal to the given plane is 1, 1, 1.

$$\cos(90^\circ - \theta) = \frac{|1(-1) + 1(0) + 1(-1)|}{\sqrt{1^2 + 1^2 + 1^2} \sqrt{1^2 + 0^2 + 1^2}}$$

$$\Rightarrow \sin \theta = \frac{2}{\sqrt{6}} \Rightarrow \cos \theta = \sqrt{1 - \frac{4}{6}} = \sqrt{\frac{1}{3}}$$

$$\text{Required projection} = |\overline{AB}| \cos \theta$$

$$= \sqrt{2} \times \frac{1}{\sqrt{3}} = \sqrt{\frac{2}{3}}$$

89. The line of intersection of first two planes is

$$\frac{x-5}{0} = \frac{y}{-3} = \frac{z+\frac{8}{-5a}}{-5a}$$

It must lie on third plane.

$$\therefore 3b(0) + (-3)(1) + (-3)(-5a) = 0$$

$$\text{and } 3b(5) + 0(1) + (-3)\left(\frac{-8}{3}\right) = 0$$

$$\Rightarrow a = \frac{1}{5} \text{ and } 15b + 8 = 0$$

$$\Rightarrow a = \frac{1}{5} \text{ and } b = -\frac{8}{15}$$



Evaluation Test

1. Given planes are

$$x - cy - bz = 0 \quad \dots(i)$$

$$cx - y + az = 0 \quad \dots(ii)$$

$$bx + ay - z = 0 \quad \dots(iii)$$

Equation of a plane passing through the line of intersection of planes (i) and (ii) is

$$x - cy - bz + k(cx - y + az) = 0$$

$$\Rightarrow (1 + ck)x - (c + k)y - (b - ak)z = 0 \quad \dots(iv)$$

Now, planes (iii) and (iv) are same for some value of k,

$$\therefore \frac{1 + ck}{b} = -\frac{c + k}{a} = \frac{-(b - ak)}{-1}$$

$$\Rightarrow \frac{1 + ck}{b} = -\frac{c + k}{a}$$

$$\Rightarrow a + ack = -bc - bk$$

$$\Rightarrow k(b + ac) = -(a + bc)$$

$$\Rightarrow k = -\left(\frac{a + bc}{b + ac}\right)$$

$$\text{Also, } -\frac{c + k}{a} = b - ak$$

$$\Rightarrow -\left(\frac{c - \frac{a + bc}{b + ac}}{a}\right) = b + a\left(\frac{a + bc}{b + ac}\right)$$

$$\Rightarrow \frac{-bc - ac^2 + a + bc}{a} = b^2 + abc + a^2 + abc$$

$$\Rightarrow 1 - c^2 = a^2 + b^2 + 2abc$$

$$\Rightarrow a^2 + b^2 + c^2 + 2abc = 1$$

2. Let a, b, c be the intercepts form by the plane on co-ordinate axes.

$$\text{Since, } \frac{1}{a} + \frac{1}{b} + \frac{1}{c} = \frac{1}{2}$$

$$\therefore \frac{2}{a} + \frac{2}{b} + \frac{2}{c} = 1$$



\therefore The point $(2, 2, 2)$ satisfies the equation of the plane $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$.

\therefore the required point is $(2, 2, 2)$.

3. Given equation of line and plane are

$$\vec{r} = \hat{i} + \hat{j} + \lambda(2\hat{i} + \hat{j} + 4\hat{k}), \text{ and}$$

$$\vec{r} \cdot (\hat{i} + 2\hat{j} - \hat{k}) = 3$$

$\therefore \vec{b} = 2\hat{i} + \hat{j} + 4\hat{k}$ and

$$\vec{n} = \hat{i} + 2\hat{j} - \hat{k}$$

$$\begin{aligned} \text{Consider } \vec{b} \cdot \vec{n} &= (2\hat{i} + \hat{j} + 4\hat{k}) \cdot (\hat{i} + 2\hat{j} - \hat{k}) \\ &= 2 + 2 - 4 \\ &= 0 \end{aligned}$$

\therefore the line lies in the plane.

4. The equation of the given line is

$$x = 2 + t, y = 1 + t, z = -\frac{1}{2} - \frac{1}{2}t$$

$$\Rightarrow \frac{x-2}{1} = \frac{y-1}{1} = \frac{z+\frac{1}{2}}{-\frac{1}{2}}$$

\therefore The given line passes through the point $(2, 1, -\frac{1}{2})$ and its d. r.s are $1, 1, -\frac{1}{2}$

The equation of the given plane is $x + 2y + 6z = 10$

\therefore d.r.s of the normal to the plane are $1, 2, 6$

$$\begin{aligned} \therefore p &= \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}} \\ &= \frac{|1(2) + 2(1) + 6\left(-\frac{1}{2}\right) - 10|}{\sqrt{1^2 + 2^2 + 6^2}} \\ &= \frac{|2 + 2 - 3 - 10|}{\sqrt{1 + 4 + 36}} = \frac{9}{\sqrt{41}} \end{aligned}$$

$$\therefore \frac{\lambda}{\sqrt{\mu}} = \frac{9}{\sqrt{41}}$$

$$\Rightarrow \lambda = 9, \mu = 41$$

$$\therefore 5\lambda - \mu = 5(9) - 41 = 45 - 41 = 4$$

5. Let \vec{a} be the vector along the line of intersection of the planes $3x - 7y - 5z = 1$ and $8x - 11y + 2z = 0$. the d.r.s of the normals to the planes are $3, -7, -5$ and $8, -11, 2$.

$$\begin{aligned} \therefore \vec{a} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & -7 & -5 \\ 8 & -11 & 2 \end{vmatrix} \\ &= \hat{i}(-14 - 55) - \hat{j}(6 + 40) + \hat{k}(-33 + 56) \\ &= -69\hat{i} - 46\hat{j} + 23\hat{k} \end{aligned}$$

Similarly, let \vec{b} the vector along the line of intersection of the planes $5x - 13y + 3z + 2 = 0$ and $8x - 11y + 2z = 0$

the d.r.s of the normals to the planes are $5, -13, 3$ and $8, -11, 2$

$$\begin{aligned} \therefore \vec{b} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 5 & -13 & 3 \\ 8 & -11 & 2 \end{vmatrix} \\ &= \hat{i}(-26 + 33) - \hat{j}(10 - 24) + \hat{k}(-55 + 104) \\ &= 7\hat{i} + 14\hat{j} + 49\hat{k} \end{aligned}$$

Consider,

$$\begin{aligned} \vec{a} \cdot \vec{b} &= (-69\hat{i} - 46\hat{j} + 23\hat{k}) \cdot (7\hat{i} + 14\hat{j} + 49\hat{k}) \\ &= -69 \times 7 + (-46) \times 14 + 23 \times 49 \\ &= -483 - 644 + 1127 \\ &= -1127 + 1127 \\ &= 0 \end{aligned}$$

$\therefore \vec{a}$ and \vec{b} are perpendicular

$$\Rightarrow \theta = 90^\circ$$


$\therefore \sin \theta = \sin 90^\circ = 1$

6. The equation of the given plane is

$$\begin{aligned} 2\lambda x - (1 + \lambda)y + 3z &= 0 \\ \Rightarrow 2\lambda x - y - \lambda y + 3z &= 0 \\ \Rightarrow (2x - y)\lambda - (y - 3z) &= 0 \\ \Rightarrow (2x - y) - \frac{1}{\lambda}(y - 3z) &= 0 \end{aligned}$$

\therefore The plane passes through the point of intersection of the planes $2x - y = 0$ and $y - 3z = 0$

7. $A(2, -1, 3)$



Let $A \equiv (2, -1, 3)$, AM be \perp to the given plane and let $B \equiv (x, y, z)$ be the image of A in the Plane.

the d.r.s of the normal to the plane are $3, -2, -1$



∴ The equation of the line AM is $\frac{x-2}{3} = \frac{y+1}{-2} = \frac{z-3}{-1} = k$, say

∴ $x = 3k + 2, y = -2k - 1, z = -k + 3$
 Let $M \equiv (3k + 2, -2k - 1, -k + 3)$

∴ equation of plane becomes $3(3k + 2) - 2(-2k - 1) - (-k + 3) = 9$

∴ $k = \frac{2}{7}$

∴ $M \equiv \left(\frac{6}{7} + 2, -\frac{4}{7} - 1, -\frac{2}{7} + 3\right) \equiv \left(\frac{20}{7}, -\frac{11}{7}, \frac{19}{7}\right)$

Since, M is the mid point of AB.

∴ $\frac{x_1 + 2}{2} = \frac{20}{7}, \frac{y_1 - 1}{2} = -\frac{11}{7}, \frac{z_1 + 3}{2} = \frac{19}{7}$

∴ $x_1 = \frac{26}{7}, y_1 = -\frac{15}{7}, z_1 = \frac{17}{7}$

Image of A is $B\left(\frac{26}{7}, -\frac{15}{7}, \frac{17}{7}\right)$

8. Since, \vec{a} and \vec{b} are coplanar, $\vec{a} \times \vec{b}$ is a vector perpendicular to the plane containing \vec{a} and \vec{b} . Similarly, $\vec{c} \times \vec{d}$ is a vector perpendicular to the plane containing \vec{c} and \vec{d} . The two planes will be parallel, if their normals $\vec{a} \times \vec{b}$ and $\vec{c} \times \vec{d}$ are parallel.

∴ $(\vec{a} \times \vec{b}) \times (\vec{c} \times \vec{d}) = \vec{0}$

9. Equation of the plane containing the given lines is

$$\begin{vmatrix} x-1 & y-2 & z-3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{vmatrix} = 0$$

$\Rightarrow (x-1)(15-16) - (y-2)(10-12) + (z-3)(8-9) = 0$

$\Rightarrow (x-1)(-1) - (y-2)(-2) + (z-3)(-1) = 0$

$\Rightarrow -x + 1 + 2y - 4 - z + 3 = 0$

$\Rightarrow -x + 2y - z = 0$

$\Rightarrow x - 2y + z = 0 \quad \dots(i)$

Given equation of plane is $Ax - 2y + z = d \quad \dots(ii)$

The planes given by equation (i) and (ii) are parallel.


∴ $A = 1$
 distance between the planes (D) is

$$D = \left| \frac{d}{\sqrt{1^2 + (-2)^2 + 1^2}} \right| = \left| \frac{d}{\sqrt{6}} \right|$$

∴ $\left| \frac{d}{\sqrt{6}} \right| = \sqrt{6}$

$\Rightarrow |d| = 6$

10.



Since, direction cosines of PQ are equal and positive

∴ the d.r.s. of PQ are $\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}$

∴ The equation of the line PQ is $\frac{x-2}{1/\sqrt{3}} = \frac{y+1}{1/\sqrt{3}} = \frac{z-2}{1/\sqrt{3}}$

$\Rightarrow x - 2 = y + 1 = z - 2 = k$, say

∴ Co-ordinate of the point Q are $(k + 2, k - 1, k + 2)$

The point Q lies on the plane $2x + y + z = 9$

∴ $2(k + 2) + k - 1 + k + 2 = 9$

$\Rightarrow 4k + 5 = 9 \quad \Rightarrow k = 1$

∴ $Q \equiv (3, 0, 3)$

∴ $PQ = \sqrt{(3-2)^2 + (0+1)^2 + (3-2)^2}$

$= \sqrt{1+1+1} = \sqrt{3}$

11. Let $A \equiv (a, 0, 0), B \equiv (0, b, 0), C \equiv (0, 0, c)$

∴ $G \equiv (x, y, z) \equiv \left(\frac{a}{3}, \frac{b}{3}, \frac{c}{3}\right)$

$\Rightarrow \frac{a}{3} = x, \frac{b}{3} = y, \frac{c}{3} = z$

$\Rightarrow a = 3x, b = 3y, c = 3z \quad \dots(i)$



The equation of the plane is

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

Since, this plane is at a distance of 1 unit from the origin,

$$\left| \frac{-1}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2}}} \right| = 1$$

$$\Rightarrow \frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} = 1$$

$$\Rightarrow \frac{1}{9x^2} + \frac{1}{9y^2} + \frac{1}{9z^2} = 1 \dots [\text{From (i)}]$$

$$\Rightarrow \frac{1}{x^2} + \frac{1}{y^2} + \frac{1}{z^2} = 9$$

$$\Rightarrow k = 9$$

12. Let the equation of the plane OAB be $ax + by + cz = d$

This plane passes through the points A(1, 2, 1) and B(2, 1, 3)

$$\therefore a + 2b + c = 0, \quad \dots \text{(i)}$$

$$\text{and } 2a + b + 3c = 0 \quad \dots \text{(ii)}$$

\therefore on solving (i) and (ii), we get

$$\frac{a}{5} = \frac{b}{-1} = \frac{c}{-3}$$

Similarly, let the equation of the plane ABC be

$$a'(x+1) + b'(y-1) + c'(z-2) = 0$$

Substituting the co-ordinates of A and B, we get

$$2a' + b' - c' = 0,$$

$$\text{and } 3a' + c' = 0$$

$$\therefore \frac{a'}{1} = \frac{b'}{-5} = \frac{c'}{-3}$$

If θ is the angle between two planes, then it is the angle between their normals.

$$\therefore \cos \theta = \frac{5 \times 1 + (-1) \times (-5) + (-3) \times (-3)}{\sqrt{25+1+9} \sqrt{1+25+9}}$$

$$= \frac{5+5+9}{\sqrt{35} \sqrt{35}}$$

$$= \frac{19}{35}$$

$$\Rightarrow \theta = \cos^{-1} \left(\frac{19}{35} \right)$$

13. The equation of the given plane can be written

$$\text{as } \frac{x}{20} + \frac{y}{15} + \frac{z}{-12} = 1$$

Let the plane intersects the x , y and z axes in the points A(20, 0, 0), B(0, 15, 0), C(0, 0, -12)

$$\therefore \bar{a} = 20\hat{i}, \bar{b} = 15\hat{j}, \text{ and } \bar{c} = -12\hat{k}$$

$$\therefore \text{Volume of tetrahedron} = \frac{1}{6} [\bar{a} \ \bar{b} \ \bar{c}]$$

$$= \frac{1}{6} \begin{vmatrix} 20 & 0 & 0 \\ 0 & 15 & 0 \\ 0 & 0 & -12 \end{vmatrix} = |-600| = 600$$

14. Given lines are coplanar.

$$\therefore \begin{vmatrix} 1-2 & 4-3 & 5-4 \\ 1 & 1 & -k \\ k & 2 & 1 \end{vmatrix} = 0$$

$$\Rightarrow -1(1+2k) - 1(1+k^2) + 1(2-k) = 0$$

$$\Rightarrow -1 - 2k - 1 - k^2 + 2 - k = 0$$

$$\Rightarrow -k^2 - 3k = 0 \Rightarrow k(k+3) = 0$$

$$\Rightarrow k = 0 \text{ or } k = -3$$

09 Linear Programming



Hints



Classical Thinking

3. Option D is the only option which is non-linear.
4. 'p' is a linear inequality and 'q' is a non-linear inequality
5. Since the profit should be maximum, the objective function is
Maximum profit, $z = 40x + 25y$.
9. Let x = number of table clothes produced in a day, and
 y = number of curtains produced in a day
 $\therefore x \geq 0, y \geq 0$...[\because both items cannot be negative]

Representing the given information in tabular form, we get

	Table cloth (x)	Curtain (y)	Total availability
Money earned (₹)	50	250	500
Hours of work	1	3	z

- $\therefore 50x + 250y \geq 500$
- \therefore total hours = $z = x + 3y$
- \therefore Required LPP is formulated as
Minimize, $z = x + 3y$, subject to $50x + 250y \geq 500, x \geq 0, y \geq 0$
15. At (800, 400), $P = 12(800) + 6(400) = 12000$
At (1050, 150), $P = 12(1050) + 6(150) = 13500$
At (600, 0), $P = 12(600) + 6(0) = 7200$
 \therefore Maximum value of P is 13500.
16. The corner points of feasible region are O(0, 0), A(7, 0), B(3, 4) and D(0, 2)
At A(7, 0), $z = 5(7) + 7(0) = 35$
At B(3, 4), $z = 5(3) + 7(4) = 43$
At C(0, 2), $z = 5(0) + 7(2) = 14$
 \therefore Maximum value of z is 43.
17. The corners of feasible region are O(0, 0), A(25, 0), B(16, 16) and C(0, 24)
At O(0, 0), $z = 0$
At A(25, 0), $z = 4(25) + 3(0) = 100$
At B(16, 16), $z = 4(16) + 3(16) = 112$
At C(0, 24), $z = 4(0) + 3(24) = 72$
 \therefore Maximum value of z is 112.
18. The corners of feasible region are O(0,0), A(52, 0), E(44, 16) and D(0, 38).
 \therefore At A(52, 0), $z = 3(52) + 4(0) = 156$
At E(44, 16), $z = 3(44) + 4(16) = 196$
At D(0, 38), $z = 3(0) + 4(38) = 152$
 \therefore Maximum value of z is 196



19. At A (50, 50), $P = \frac{5}{2}(50) + \frac{3}{2}(50) + 410 = 610$

At B (10, 50), $P = \frac{5}{2}(10) + \frac{3}{2}(50) + 410 = 510$

At C (60, 0), $P = \frac{5}{2}(60) + \frac{3}{2}(0) + 410 = 560$

At D (60, 40), $P = \frac{5}{2}(60) + \frac{3}{2}(40) + 410 = 620$

∴ Minimum value of P is 510 at B (10, 50)

20. The corners of given feasible region are A(12, 0), B(4, 2), C(1, 5) and D(0, 10)

At A(12, 0), $z = 3(12) + 2(0) = 36$

At B(4, 2), $z = 3(4) + 2(2) = 16$

At C(1, 5), $z = 3(1) + 2(5) = 13$

At D(0, 10), $z = 3(0) + 2(10) = 20$

Minimum value of z is 13

21. The corner points of feasible region are (0, 3), (0, 5) and (3, 2)

∴ At (0, 3), $z = 11(0) + 7(3) = 21$

At (0, 5), $z = 11(0) + 7(5) = 35$

At (3, 2), $z = 11(3) + 7(2) = 47$

∴ Minimum value of z is 21

22. At P $\left(\frac{3}{13}, \frac{24}{13}\right)$, $z = \frac{3}{13} + 2\left(\frac{24}{13}\right) = \frac{51}{13} = 3.923$

At Q $\left(\frac{3}{2}, \frac{15}{4}\right)$, $z = \frac{3}{2} + 2\left(\frac{15}{4}\right) = 9$

At R $\left(\frac{7}{2}, \frac{3}{4}\right)$, $z = \frac{7}{2} + 2\left(\frac{3}{4}\right) = 5$

At S $\left(\frac{18}{7}, \frac{2}{7}\right)$, $z = \frac{18}{7} + 2\left(\frac{2}{7}\right) = \frac{22}{7} = 3.143$

∴ Maximum value of z is 9, and

Minimum value of z is $\frac{22}{7}$.

23. Assume that x and y take arbitrary large values. So the objective function can be made as large as we want. Hence the problem has unbounded solution.

24. The feasible region is unbounded. x and y can take arbitrary large values. Hence the problem has unbounded solution.

25. Since there are two disjoint feasible regions, the LPP has no solution.

26. The feasible region is disjoint.

∴ There is no solution.



Critical Thinking

1. From the given table the constraints are $2x + 3y \leq 36$; $5x + 2y \leq 50$; $2x + 6y \leq 60$
Also $x \geq 0, y \geq 0$ [\because number of magazines cannot be negative]

\therefore The number of constraints are 5.

2. Representing the given information in table form, we get

	Shirt (x)	Pants (y)	Total availability
Work time on machine (hours)	2	3	70
Man labour (hours)	3	2	75

Linear constraints are $2x + 3y \leq 70, 3x + 2y \leq 75$.

Also, $x \geq 0, y \geq 0$ [\because number of shirts and pants cannot be negative]

3. Let the factory owner purchase x units of machine A and y units of machine B for his factory.

$\therefore x \geq 0, y \geq 0$ [\because number of machines cannot be negative]

Representing the given information in tabular form, we get

	Machine A(x)	Machine B(y)	Total Availability
Machine Area (m^2)	1000	1200	7600
Skilled men	12	8	72
Daily output (no. of units)	50	40	z

$\therefore 1000x + 1200y \leq 7600$

$$12x + 8y \leq 72$$

4. Let, x = number of necklaces, and y = number of bracelets

$\therefore x \geq 0, y \geq 0$ [\because number of necklaces and bracelets cannot be negative]

Representing the given information in tabular form, we get

	Necklace (x)	Bracelet (y)	Total availability
Time required (hrs)	$\frac{1}{2}$	1	16
Profit (₹)	100	200	z

$\therefore \frac{1}{2}x + y \leq 16 \Rightarrow x + 2y \leq 32$

$$x + y \leq 24$$

$$\text{total profit } z = 100x + 300y$$

\therefore Required LPP is formulated as

Maximize $z = 100x + 300y$, subject to

$$x + y \leq 24, x + 2y \leq 32, x \geq 0, y \geq 0$$

5. Let the consumption per day be, x grams of food X and Y grams of food Y.

$\therefore x \geq 0$ and $y \geq 0$ [\because the quantities cannot be negative]

Representing the given information in table form, we get

Type of food	Food X (x)	Food Y (y)	Minimum requirement
Vitamin A per gram (units)	4	6	90
Vitamin B per gram (units)	7	11	130
Cost per gram (paise)	15	22	z

$\therefore 4x + 6y \geq 90,$

$$7x + 11y \geq 130, \text{ and } z = 15x + 22y$$

\therefore Required LLP is formulated as,

Minimize $z = 15x + 22y$, subject to constraints

$$4x + 6y \geq 90, 7x + 11y \geq 130, x \geq 0, y \geq 0$$




6. Suppose x kg of food A and y kg of food B are consumed to form a weekly diet.
 $\therefore x \geq 0, y \geq 0$[Since quantity of food cannot be negative]

Representing the given information in table form, we get

	Food A (x)	Food B (y)	Minimum requirement
Fats (units)	4	12	18
Carbohydrates (units)	16	4	24
Protein (units)	8	6	16
Cost (₹)	6	5	z

- \therefore Required LPP is formulated as
 Minimize, $z = 6x + 5y$ subject to constraints,
 $4x + 12y \geq 18, 16x + 4y \geq 24, 8x + 6y \geq 24, x \geq 0, y \geq 0$

7. Converting the given inequalities into equations, we get $x + y = 4$
 The equation intersects the axes at $(4, 0)$ and $(0, 4)$
 The feasible region lies on origin side of lines $y = 5$ and $x + y = 4$ and in first quadrant.
 It is bounded in first quadrant.



8. Converting given inequalities into equations, we get

$$y - x = 1 \quad \text{i.e.} \quad \frac{x}{(-1)} + \frac{y}{1} = 1 \quad \dots\text{(i)}$$

$$2x - 6y = 3 \quad \text{i.e.} \quad \frac{x}{\frac{3}{2}} + \frac{y}{\left(\frac{-1}{2}\right)} = 1 \quad \dots\text{(ii)}$$


$$x = 0, y = 0$$

- \therefore Equation (i) intersects the axes at $(-1, 0)$ and $(0, 1)$


Equation (ii) intersects the axes at $\left(\frac{3}{2}, 0\right)$ and $\left(0, \frac{-1}{2}\right)$


Substituting $x = 0, y = 0$ in given inequalities, we get
 $(0) - (0) = 0 \leq 1$, and $2(0) - 6(0) = 0 \leq 3$

- \therefore Feasible region lies on the origin side of both the lines, in first quadrant.
 It is unbounded and convex.



9. The feasible region lies on origin side of line $2x + 3y - 5 = 0$ and non-origin side of line $4x - 3y + 2 = 0$. However, it is not bounded by any axes.




10. 

The feasible region lies on origin side of the lines $-x + 3y = 9$ and $-x + y = 1$, and in first quadrant.
 It is unbounded.




11. Feasible region lies on origin side of line $2x - 3y = 5$.
 \therefore O lies inside the region
 Substituting P (2, -2) in given inequality, we get
 $2(2) - 3(-2) = 10 \not\leq 5$
 \therefore P lies outside the region.




12. It is clear from the graph that origin is not there in the feasible region. Out of the 4 options, only option (B) satisfies this condition i.e., $4(0) - 2(0) = 0 \leq -3$ is correct.

13. The shaded region lies;
 On origin side of line $x + 2y = 8 \Rightarrow x + 2y \leq 8$,
 On non-origin side of line $2x + y = 2 \Rightarrow 2x + y \geq 2$,
 On origin side of line $x - y = 1 \Rightarrow x - y \leq 1$
 and in first quadrant $\Rightarrow x \geq 0, y \geq 0$.

14. The feasible region lies on non-origin side of line $2x + y = 2$ and origin side of line $x - y = 3$ as shown in the figure.
 By solving the two equations, we get the point of intersection $(\frac{5}{3}, \frac{-4}{3})$, which is the vertex of the common graph.




15. Feasible region lies on origin side of line $x + y = 6$, non-origin side of line $3x + 2y = 6$ and in the first quadrant.
 \therefore Vertices of the feasible region are (0, 6), (0, 3), (2, 0) and (6, 0)





16. Converting the given inequalities into equations, we get $x = 5$, $x = 10$, $y = 5$ and $y = 10$

The feasible region is as shown in the figure



∴ The vertices of the feasible region are (5, 5), (10, 5), (10, 10) and (5, 10)

17. Converting the given inequations into equations, we get

$$2x + 3y = 6 \text{ i.e. } \frac{x}{3} + \frac{y}{2} = 1 \quad \dots(i)$$

$$5x + 3y = 15 \text{ i.e. } \frac{x}{3} + \frac{y}{5} = 1 \quad \dots(ii)$$

∴ Equation (i) intersects the axes at points (3, 0) and (0, 2)

Equation (ii), intersects at points (3, 0) and (0, 5).


Also substituting origin (0, 0) in both in equalities we get,

$$2(0) + 3(0) = 0 \leq 6 \text{ and } 5(0) + 3(0) = 0 \leq 15$$

∴ Feasible region lies on origin side of both the lines as shown in the graph

∴ the vertices of feasible region are (0, 2), (0, 0) and (3, 0)

∴ (0, 5) is not a vertex of feasible region.



18. Using two point form we have, equation of line AB : $x + 2y = 8$ and equation of line CD : $3x + 2y = 12$

Since, the shaded region lies on, origin side of line AB, non-origin side of line CD and above X- axis.

∴ $x + 2y \leq 8$, $3x + 2y \geq 12$ and $y \geq 0$

19. Take a test point (1, 1) that lies within the feasible region. Since $(1) + (1) = 2 \leq 5$, is true we have $x + y \leq 5$.

Since $1 \leq 4$ and $1 \leq 3$ are true, we have $x \leq 4$ and $y \leq 3$. Since $4(1) + 1 = 5 \geq 4$, we have $4x + y \geq 4$

20. The feasible region lies on the origin side of $2x + y = 30$ and $x + 2y = 24$, in the first quadrant.


The corners of the feasible region are O (0, 0), A (15, 0), B (0, 12) and C (12, 6)

At A(15, 0), $z = 90$

At B(0, 12), $z = 96$

At C(12, 6), $z = 120$

∴ Maximum value of z is 120.



21. The feasible region lies on origin side of lines $x + y = 5$ and $3x + y = 9$, in first quadrant.


∴ The corners of feasible region are

O (0, 0), A (0, 5), B (2, 3) and C (3, 0)

∴ Maximum value of objective function

$$z = 12x + 3y \text{ is at } C (3, 0)$$

∴ $z = 12(3) + 3(0) = 36$






22. The feasible region lies on the origin side of $3x + 5y = 15$ and $5x + 2y = 10$, in first quadrant.

The corners of the feasible region are

O (0, 0), B (0, 3), E $\left(\frac{20}{19}, \frac{45}{19}\right)$ and C (2, 0)

The maximum value of $z = 5x + 3y$ is at E $\left(\frac{20}{19}, \frac{45}{19}\right)$

\therefore Maximum $z = 5 \left(\frac{20}{19}\right) + 3 \left(\frac{45}{19}\right) = \frac{235}{19}$



23. Feasible region lies on the origin side of $x + 5y = 200$ and $2x + 3y = 134$, in first quadrant.


The corner points of the feasible region are O (0, 0), A (67, 0), B (10, 38) and C (0, 40)

At A (67, 0), $z = 268$

At A (10, 38), $z = 382$

At A (0, 40), $z = 360$

\therefore Maximum value of z is at B (10, 38)



24. $z = px + qy$

At (15, 15), $z = 15p + 15q$

At (0, 20), $z = 0 + 20q = 20q$

\therefore Maximum z occurs at both the points,

$$\Rightarrow 15p + 15q = 20q \Rightarrow 15p = 5q \Rightarrow 3p = q$$

25. Suppose that the manufacturer produces x soaps of type I and y soaps of type II.

$\therefore x \geq 0; y \geq 0; 2x + 3y \leq 480$ and $3x + 5y \leq 480$

Feasible region lies on origin side on both inequalities, in first quadrant.

The corners of the feasible region are


O (0, 0), A (0, 96) and B (160, 0)

Maximum profit, $P = 0.25x + 0.5y$

\therefore At A (0, 96), $P = 0.25(0) + 0.5(96) = 48$

At B (160, 0), $P = 0.25(160) + 0.5(0) = 40$

For maximum profit of ₹ 48, 96 soaps of type II must be manufactured.



26. The feasible region lies on the origin side of both the lines.

The corner points of feasible region are

O (0, 0), A (30, 0), B (0, 40) and P (30, 40)


\therefore At O (0, 0), $z = 4(0) + 5(0) = 0$

At A (30, 0), $z = 4(30) + 5(0) = 120$

At B (0, 40), $z = 4(0) + 5(40) = 200$


At P (30, 40), $z = 4(30) + 5(40) = 320$

\therefore The minimum value of z is 0





27. The feasible region lies on origin side of line $2x + 3y = 6$ and non-origin side of line $x + y = 1$
 The corners of feasible region are
 A (3, 0), B (0, 2), C (1, 0) and D (0, 1)
 $\therefore z = 3x + y$ will be minimum at C or D.
 \therefore At C (1, 0), $z = 3(1) + (0) = 3$
 At D (0, 1), $z = 3(0) + 1 = 1$
 \therefore Minimum value of z is 1



28. Feasible region lies on origin side of lines $5x + 8y = 40$ and $3x + y = 6$ and above line $y = 2$, in first quadrant.

The corner points of the feasible region

A(0, 2), B($\frac{4}{3}$, 2), C($\frac{8}{19}$, $\frac{90}{19}$) and D(0, 5)


At A (0, 2), $z = 14$

At B ($\frac{4}{3}$, 2), $z = 22$


At C ($\frac{8}{19}$, $\frac{90}{19}$), $z = \frac{678}{19}$

At D (0, 5), $z = 35$


- \therefore Minimum value of z is 14




30. The corner points of feasible region are
 A(1, 0), B(10, 0), C (2, 4), D(0, 4) and E (0, 1)
 At A (1, 0), $z = 1 + 0 = 1 = 1$
 At B (10, 0), $z = 10 + 0 = 10$
 At C (2, 4), $z = 2 + 4 = 6$
 At D (0, 4), $z = 0 + 4 = 4$
 At E (0, 1), $z = 0 + 1 = 1$
 z has minimum value at both A (1, 0) and E (0, 1).
 $\therefore z$ has infinite solutions on seg AE.



31. Feasible region lies on origin side of line $x_1 + x_2 = 1$ and non-origin side of line $3x_1 + x_2 = 3$ in first quadrant.
 \therefore there is no feasible region.





The feasible regions are disjoint. Hence there is no point in common.

∴ There is no optimum value of the objective function.



Competitive Thinking


5. Condition (i),
 $i = 1, x_{11} + x_{12} + x_{13} + \dots + x_{1n}$
 $i = 2, x_{21} + x_{22} + x_{23} + \dots + x_{2n}$
 $i = 3, x_{31} + x_{32} + x_{33} + \dots + x_{3n}$

 $i = m, x_{m1} + x_{m2} + x_{m3} + \dots + x_{mn} \rightarrow m \text{ constraints}$
 Condition (ii),
 $j = 1, x_{11} + x_{21} + x_{31} + \dots + x_{m1}$
 $j = 2, x_{12} + x_{22} + x_{32} + \dots + x_{m2}$

 $j = n, x_{1n} + x_{2n} + x_{3n} + \dots + x_{mn} \rightarrow n \text{ constraints}$


∴ Total constraints = $m + n$

7. In linear programming problem, concave region is not used. Convex region is used in linear programming.




Feasible region is on non-origin side of $3y + x = 3$ and in first quadrant. Hence, it is unbounded.

9. The feasible region lies on origin side of the lines $-x_1 + x_2 = 1$ and $-x_1 + 3x_2 = 9$, in first quadrant. It is unbounded.





10. Feasible region lies on non-origin side of both lines and is true for positive values of x and both positive and negative values of y .



11. Since shaded region lies on origin side of lines $x + y = 20$ and $2x + 5y = 80$ and is in first quadrant
 $\therefore x + y \leq 20$, $2x + 5y \leq 80$, $x \geq 0$, $y \geq 0$
12. Shaded region lies on origin side of $x + 2y = 8$ and $x - y = 1$, and on non-origin side of $2x + y = 2$.
 $\therefore x + 2y \leq 8$, $x - y \leq 1$, $2x + y \geq 2$
13. Take a test point $(2, 1)$ which lies within the feasible region.
 Since, $2 - 1 = 1 \geq 0$, $2 \leq 5$, $1 \leq 3$ and $2, 1 \geq 0$
 $\therefore x \geq 0$, $x - y \geq 0$, $x \leq 5$, $y \leq 3$.
14. Since shaded region lies on non-origin side of $5x + 4y = 20$, and on origin side of the lines $x = 6$ and $y = 3$
 $\therefore 5x + 4y \geq 20$, $x \leq 6$, $y \leq 3$, $x \geq 0$, $y \geq 0$

17. The feasible region lies on the origin side of $x + y = 40$ and $x + 2y = 6$, in first quadrant.

The corners of feasible region are


$O(0, 0)$, $A(0, 30)$, $B(20, 20)$ and $C(40, 0)$

\therefore At $A(0, 30)$, $P = 0 + 4(30) = 120$

At $B(20, 20)$, $P = 3(20) + 4(20) = 140$

At $C(40, 0)$, $P = 3(40) + 0 = 120$

\therefore Maximum value of P is 140.




18. The feasible region lies on origin side of $4x + 5y = 20$, non-origin side of $x + y = 3$ and in first quadrant.

\therefore The corners of feasible region are

$A(5, 0)$, $B(0, 4)$, $C(3, 0)$ and $D(0, 3)$

\therefore Maximum $2x + 3y$ is at $B(0, 4)$

\therefore Maximum $2x + 3y = 2(0) + 4(3) = 12$






19. Feasible region lies on origin side of $x + y = 7$ and $x + 2y = 10$, and in first quadrant.

The corners of feasible region are $O(0, 0)$, $A(7, 0)$, $E(4, 3)$ and $D(0, 5)$

- \therefore Maximum $z = 5x + 2y$ is at $A(7, 0)$
 \therefore Maximum, $z = 5(7) + 2(0) = 35$



20. Corner points of the feasible region are

$(0, 0)$, $(6, 0)$, $(\frac{9}{2}, \frac{5}{2})$ and $(0, \frac{26}{5})$


At $(0, 0)$, $z = 2(0) + 0 = 0$

At $(6, 0)$, $z = 2(6) + 0 = 12$

At $(\frac{9}{2}, \frac{5}{2})$, $z = 2(\frac{9}{2}) + \frac{5}{2} = 11.5$

At $(0, \frac{26}{5})$, $z = 2(0) + \frac{26}{5} = 5.2$

- \therefore Maximum value of z is 12 at $(6, 0)$.



21. The feasible region lies on origin side of all the inequalities and in first quadrant

The corners of feasible region are $(0, 0)$, $(4, 0)$, $(4, 3)$, $(2, 6)$ and $(0, 6)$.


At $(4, 0)$, $z = 3(4) + 0 = 12$

At $(4, 3)$, $z = 3(4) + 5(3) = 27$

At $(2, 6)$, $z = 3(2) + 5(6) = 36$

At $(0, 6)$, $z = 0 + 5(6) = 30$

- \therefore Maximum value of z is 36 at $(2, 6)$




22. The feasible region lies on the origin side of the lines $x + y = 20$, $x + 2y = 35$ and $x - 3y = 12$

The corners of the feasible region are

$O(0, 0)$, $E(12, 0)$, $H(18, 2)$, $G(5, 15)$, $D(0, \frac{35}{2})$

- \therefore The maximum value of $4x + 5y$ is at $G(5, 15)$

- \therefore Maximum $4x + 5y = 4(5) + 5(15) = 95$






23. The feasible region lies on origin side of all the lines and in first quadrant.

∴ The corners of feasible region are

O (0, 0), A (2, 0), B (2, 1), C $\left(\frac{2}{3}, \frac{7}{3}\right)$ and D (0, 1)

Maximum value of $z = 3x + 2y$ is at B (2, 1)

∴ Maximum $z = 3(2) + 2(1) = 8$




24. The feasible region lies on origin side of $x + y = 2$

The corners of feasible region are

A (2, 0), B (0, 2) and O (0, 0).

At A (2, 0), the value of z is maximum = 6



25. The feasible region lies on the origin side of the lines


$6x + 4y = 120$ and $3x + 10y = 180$

The corners of feasible region are

O (0, 0), A (20, 0), E (10, 15) and D (0, 18)

∴ The maximum value of $45x + 55y$ is at E (10, 15)

Max $(45x + 55y) = 45(10) + 55(15)$
 $= 1275$



26. Feasible region lies on origin side of all lines and in first quadrant.


The corners of feasible region are

O (0, 0), A (0, 4), B $\left(1, \frac{7}{2}\right)$, C $\left(\frac{5}{2}, 2\right)$, D $\left(\frac{7}{2}, 0\right)$

Substituting the above points in $P = 2x + 3y$, we get

Max $P = 12.5$ at B $\left(1, \frac{7}{2}\right)$

∴ B $\equiv (1, 3.5)$



27. At (15, 15), $z = 15p + 15q$

At (0, 20), $z = 20q$

Since, maximum occurs at (15, 15) and (0, 20),

∴ $z_{\max} = 15p + 15q = 20q$

$\Rightarrow 15p + 15q = 20q$

$\Rightarrow 15p = 5q \Rightarrow 3p = q$

28. $z = px + qy$

At (25, 20), $z = 25p + 20q$

At (0, 30), $z = 0 + 30q = 30q$

Since maximum z occurs at both the points,

$25p + 20q = 30q$


$\Rightarrow 25p = 10q \Rightarrow 5p = 2q$



29. At (5, 5), $z = 3(5) + 9(5) = 60$
 At (0, 10), $z = 3(0) + 9(10) = 90$
 At (0, 20), $z = 3(0) + 9(20) = 180$
 At (15, 15), $z = 3(15) + 9(15) = 180$
 \therefore Minimum value of z is 60 at (5, 5).

30. The feasible region lies on non-origin side of all the lines, X_2 in first quadrant
 The corners of feasible region are A(11, 0), B(4, 2), C(1, 5) and D(0, 10).


- \therefore At A(11,0), $z = 2(11) + 0 = 22$
 At B(4, 2), $z = 2(4) + 3(2) = 14$
 At C(1, 5), $z = 2(1) + 3(5) = 17$
 At D(0, 1), $z = 0 + 3(10) = 30$
 \therefore Maximum value of z is 14



31. The feasible region is unbounded whose vertex is $(\frac{5}{4}, \frac{5}{4})$.

- \therefore Minimum $z = 2x + 10y$ is at $(\frac{5}{4}, \frac{5}{4})$

- $\therefore z = 2(\frac{5}{4}) + 10(\frac{5}{4}) = 15$




32. The feasible region region lies on the non-origin side of $2x + 3y = 6$ and $y = 1$ and on origin side of $x + y = 8$
 The corners of feasible region are

A $(\frac{3}{2}, 1)$, B(0, 2), C(7, 1) and D(0, 8).

Substituting above points in $z = 4x + 6y$, we get


Min. $z = 12$ at A $(\frac{3}{2}, 1)$ and B (0, 2).



33. The feasible region lies on origin side of line $x + y - 20 = 0$ and above the line $y = 5$.

The corners of feasible region are B (0, 20), C (0, 5) and D (15, 5)

- \therefore The minimum value of $z = 7x - 8y$ is at B (0, 20)
 $z = 7(0) - 8(20) = -160$





34. Corner points of the feasible region are (60, 0), (120, 0), (60, 30), (40, 20).


At (60, 0), $z = 5(60) + 10(0) = 300$

At (120, 0), $z = 5(120) + 10(0) = 600$

At (60, 30), $z = 5(60) + 10(30) = 600$

At (40, 20), $z = 5(40) + 10(20) = 400$

∴ Minimum value of z is 300 at (60, 0).



35. The corner points of the feasible region are

A(3.5, 0), B(7.5, 0), C(3, 3) and D(2, 3)


At A(3.5, 0), $z = 4(3.5) + 5(0) = 14$

At B(7.5, 0), $z = 4(7.5) + 5(0) = 30$

At C(3, 3), $z = 4(3) + 5(3) = 27$

At D(2, 3), $z = 4(2) + 3(3) = 17$

∴ z is minimum at A(3.5, 0).



36. The corner points of feasible region are

A (6, 0), B (6, 4), C (3, 7) and D (0, 5)

∴ At A (6, 0), $z = 6 + 0 = 6$


At B (6, 4), $z = 6 + 4 = 10$

At C (3, 7), $z = 3 + 7 = 10$

At D (0, 5), $z = 0 + 5 = 5$

∴ z is maximum at B (6, 4) and C (3, 7)

∴ Infinite number of solutions exists along BC. $-2x + 3y = 15$



37. The corners of feasible region are

A(8,0), B(0, 8), F(0, 3), G(1, 3/2) and C(4, 0)

At F(0,3), $z = 30(0) + 20(3) = 60$

At G(1,3/2), $z = 30(1) + 20(3/2) = 30 + 30 = 60$


At A (8, 0), $z = 30(8) + 0 = 240$

At A (0, 8), $z = 0 + 20(8) = 160$

At C (4, 0), $z = 30(4) + 0 = 120$


∴ z has minimum value at F (0, 3) and G (1, 3/2)

∴ z has infinite solution on seg FG.







38. The feasible region is unbounded.
 \therefore its maximum value does not exist.



39. The feasible region lies on the origin side of the line $x + 2y = 2$ and on non-origin side of $x + 2y = 8$.
 \therefore There is no feasible solution.




40. The feasible region is disjoint.
 \therefore there is no point common to all inequations.
 \therefore There is no maximum value of z .




Evaluation Test

1. Let no. of model $M_1 = x$ and no. of model $M_2 = y$
 $\therefore x \geq 0, y \geq 0$
 Constraints are $4x + 2y \leq 80 \Rightarrow 2x + y \leq 40, 2x + 5y \leq 180$
 Maximize $z = 3x + 4y$
 The corners of feasible region are
 $O(0, 0), A(20, 0), B(2.5, 35), C(0, 36)$
 \therefore At A (20, 0), $z = 3(20) + 0 = 60$
 At B (2.5, 35), $z = 3(2.5) + 4(35) = 147.5$
 At C (0, 36), $z = 0 + 3(36) = 108$
 $\therefore z$ is maximum at B(2.5, 35).






3. Objective function $P = 2x + 3y$
 The corner points of feasible region are
 $B(12, 12)$, $C(3, 3)$, $D(20, 3)$, $E(20, 10)$, $F(18, 12)$
 At $B = P_B = 2(12) + 3(12) = 60$
 At $C = P_C = 2(3) + 3(3) = 15$
 At $D = P_D = 2(20) + 3(3) = 49$
 At $E = P_E = 2(20) + 3(10) = 70$
 At $F = P_F = 2(18) + 3(12) = 72$
 $\therefore P$ is maximum at $F(18, 12)$.



4. For $(1, 3)$, $3x + 2y = 3 + 6 > 0$,
 for $(5, 0)$, $3 \times 5 + 0 > 0$,
 and for $(-1, 2)$, $-3 + 4 > 0$
 Similarly, other inequalities satisfies the given points.
 \therefore Option (D) is the correct answer.



- OABCD is the feasible region
 $\therefore O(0, 0)$, $A(0, 600)$, $B(800, 600)$, $C(1000, 500)$, $D(1500, 0)$
 $z = x_1 + x_2$
 At point C and D, z is maximum. $\text{Max } z = 1500$
 \therefore Infinite optimal solutions exist along CD.

6. Consider option (C)
 $3 + 2(4) \geq 11$
 $3(3) + 4(4) \leq 30$
 $2(3) + 5(4) \leq 30$
 \therefore All the above three in-equalities hold for point $(3, 4)$.
 \therefore Option (C) is the correct answer.


7. Let the manufacturer produce x and y bottles of medicines A and B.
 He must have $\frac{3x}{1000} + \frac{y}{1000} \leq 66$, $x + y \leq 45000$, $x \leq 20000$, $y \leq 40,000$, $x \geq 0$, $y \geq 0$.
 \therefore the number of constraints is 6.



8. Let the company produce x telephones of A type and y telephones of B type.

\therefore Constraints are $2x + 4y \leq 800 \Rightarrow x + 2y \leq 400$, $x + y \leq 300$

Maximize $z = 300x + 400y$



\therefore the feasible region of the LPP is bounded.

9. Given that $4x + 2y \leq 8$, $2x + 5y \leq 10$

\therefore the feasible region lies on origin side of $4x + 2y = 8$ and $2x + 5y = 10$.

Also, $x, y \geq 0$

\therefore the feasible region lies in first quadrant.

\therefore option (C) is correct.


10. Objective function $z = x_1 + x_2$

The corner points of feasible region are

$O(0, 0)$, $A(2, 0)$, $B(2, 1)$, $C\left(\frac{2}{3}, \frac{7}{3}\right)$ and $D(0, 1)$

At $B(2, 1)$ and $C\left(\frac{2}{3}, \frac{7}{3}\right)$, z is maximum. $\text{Max } z = 3$

\therefore Infinite number of solutions exists along BC.



11. Objective function $z = 3x + 2y$

The corner points of feasible region are

$A\left(\frac{1}{4}, \frac{5}{4}\right)$, $B\left(\frac{1}{6}, \frac{5}{6}\right)$, $C(1, 0)$, $D(3, 0)$, $E(3, 3)$, $F\left(\frac{5}{2}, \frac{7}{2}\right)$

At A = $z_A = 3\left(\frac{1}{4}\right) + 2\left(\frac{5}{4}\right) = 3.25$

At B = $z_B = 3\left(\frac{1}{6}\right) + 2\left(\frac{5}{6}\right) = 2.167$


At C = $z_C = 3(1) + 2(0) = 3$

At D = $z_D = 3(3) + 2(0) = 9$

At E = $z_E = 3(3) + 2(3) = 15$

At F = $z_F = 3\left(\frac{5}{2}\right) + 2\left(\frac{7}{2}\right) = 14.5$

\therefore Maximum value of z at $(3, 3)$ is 15.



01 Continuity



Hints



Classical Thinking

$$1. \quad \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \left(\frac{\sin x}{x} + \cos x \right) \\ = 1 + 1 = 2 = f(0)$$

$\therefore f(x)$ is continuous at $x = 0$.

$$2. \quad \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{1}{2} \sin^2 x = 0 = f(0)$$

$\therefore f(x)$ is continuous at $x = 0$.

$$3. \quad \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \left(1 + \frac{4}{5} \right)^{\frac{1}{x}} \\ = \left[\lim_{x \rightarrow 0} \left(1 + \frac{4}{5} \right)^{\frac{5}{4}x} \right]^{\frac{4}{5}} = e^4 = f(0)$$

$\therefore f(x)$ is continuous at $x = 0$.

4. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x) \\ = \lim_{x \rightarrow 0} (\sin x - \cos x) \\ = \sin 0 - \cos 0 = -1$$

5. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x) \\ = \lim_{x \rightarrow 0} \frac{2 + \tan x}{x} \\ = \lim_{x \rightarrow 0} \left(2 + \frac{\tan x}{x} \right) = 2 + 1 = 3$$

6. Since, $f(x)$ is continuous at $x = 1$.

$$\therefore f(1) = \lim_{x \rightarrow 1} f(x) \quad \Rightarrow k = \lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 1} \\ \Rightarrow k = \lim_{x \rightarrow 1} (x + 1) \quad \Rightarrow k = 2$$

7. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x) \\ \Rightarrow \frac{k}{2} = \lim_{x \rightarrow 0} \frac{\sin 3x}{x} = \lim_{x \rightarrow 0} \frac{\sin 3x}{3} \cdot 3 \\ \Rightarrow \frac{k}{2} = 3 \quad \Rightarrow k = 6$$

8. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x) \\ \Rightarrow k = \lim_{x \rightarrow 0} \frac{\sin \pi x}{5} \\ \Rightarrow k = \lim_{x \rightarrow 0} \left(\frac{\sin \pi x}{\pi x} \right) \cdot \frac{\pi}{5} \\ \Rightarrow k = (1) \cdot \frac{\pi}{5}$$

$$\Rightarrow k = \frac{\pi}{5}$$

9. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{(e^3 - 1) \sin x}{2} \\ = \lim_{x \rightarrow 0} \frac{e^3 - 1}{3} \times 3 \times \frac{\sin x}{x} = 1 \times 3 \times 1$$

$\therefore f(0) = 3$

$$10. \quad \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (2x + 1) = 3 \neq f(1)$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (x^2 + 1) = 2 = f(1)$$

$$11. \quad f\left(\frac{1}{2}\right) = 1 - \frac{1}{2} = \frac{1}{2}$$

$$\lim_{x \rightarrow \frac{1}{2}^-} f(x) = \lim_{x \rightarrow \frac{1}{2}^-} (x) = \frac{1}{2}$$

$$\lim_{x \rightarrow \frac{1}{2}^+} f(x) = \lim_{x \rightarrow \frac{1}{2}^+} (1 - x) = 1 - \frac{1}{2} = \frac{1}{2}$$

$$\therefore \lim_{x \rightarrow \frac{1}{2}^-} f(x) = \lim_{x \rightarrow \frac{1}{2}^+} f(x) = f\left(\frac{1}{2}\right)$$

$\therefore f(x)$ is continuous at $x = \frac{1}{2}$.

12. Since, $f(x)$ is continuous at $x = 1$.

$$\therefore \lim_{x \rightarrow 1} f(x) = f(1) \\ \Rightarrow \lim_{x \rightarrow 1} (8x - 1) = k \\ \Rightarrow k = 7$$



13. Since, $f(x)$ is continuous at $x = 2$.
 $\therefore f(2) = \lim_{x \rightarrow 2^-} f(x)$
 $\Rightarrow 3 = \lim_{x \rightarrow 2^-} (kx - 1)$
 $\Rightarrow 3 = 2k - 1$
 $\Rightarrow k = 2$
14. Since, $f(x)$ is continuous at $x = 1$.
 $\therefore \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x)$
 $\Rightarrow 2 = \lim_{x \rightarrow 1} (cx - 2)$
 $\Rightarrow 2 = c - 2$
 $\Rightarrow c = 4$
15. Since, $f(x)$ is continuous at $x = 0$.
 $\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x)$
 $\Rightarrow \lim_{x \rightarrow 0^-} (-x^2 - k) = \lim_{x \rightarrow 0^+} (x^2 + k)$
 $\Rightarrow -k = k$
 $\Rightarrow k = 0$
16. Since, $f(x)$ is continuous at $x = 1$.
 $\therefore \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x)$
 $\Rightarrow \lim_{x \rightarrow 1^-} (2x + 1) = \lim_{x \rightarrow 1^+} (3 - kx^2)$
 $\Rightarrow 2 + 1 = 3 - k(1)^2$
 $\Rightarrow k = 0$
17. Since, $f(x)$ is continuous at $x = 3$.
 $\therefore f(3) = \lim_{x \rightarrow 3^-} f(x)$
 $\Rightarrow 4 = \lim_{h \rightarrow 0} f(3 - h) \Rightarrow 4 = \lim_{h \rightarrow 0} (3 - h + \lambda)$
 $\Rightarrow 3 + \lambda = 4 \Rightarrow \lambda = 1$
18. Since, $f(x)$ is continuous at $x = 2$.
 $\therefore f(2) = \lim_{x \rightarrow 2^-} f(x)$
 $\Rightarrow f(2) = \lim_{x \rightarrow 2^-} \left(\frac{x^2 - 4}{-2} + a \right) \Rightarrow 8 = 4 + a$
 $\Rightarrow a = 4$
 Also, $f(2) = \lim_{x \rightarrow 2^+} f(x)$
 $\Rightarrow f(2) = \lim_{x \rightarrow 2^+} (x + b + 4) \Rightarrow 8 = 6 + b$
 $\Rightarrow b = 2$
19. Since, $f(x)$ is continuous at $x = \frac{\pi}{2}$.
 $\therefore \lim_{x \rightarrow \frac{\pi}{2}^-} f(x) = \lim_{x \rightarrow \frac{\pi}{2}^+} f(x)$
 $\Rightarrow \lim_{x \rightarrow \frac{\pi}{2}^-} (a + 1) = \lim_{x \rightarrow \frac{\pi}{2}^+} (\sin x + b)$
 $\Rightarrow a \cdot \frac{\pi}{2} + 1 = 1 + b \Rightarrow b = \frac{a\pi}{2}$
20. $\lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 1} x^2 = 1$ and $f(1) = 2$
 $\therefore f(x)$ is discontinuous at $x = 1$.
21. $\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} x^2 = 1$
 $\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (x + 5) = 6$
 $\therefore f(x)$ is discontinuous at $x = 1$.
22. $\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} x = 1$
 $\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (x + 1) = 1 + 1 = 2$
 $\therefore \lim_{x \rightarrow 1^-} f(x) \neq \lim_{x \rightarrow 1^+} f(x)$
 $\therefore f(x)$ is discontinuous at $x = 1$.
23. $\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (x - 1) = -1$
 $\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} x^2 = 0$
 $\therefore \lim_{x \rightarrow 0^-} f(x) \neq \lim_{x \rightarrow 0^+} f(x)$
 $\therefore f(x)$ is discontinuous at $x = 0$.
24. $\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} \left(\frac{5}{2} - x \right) = \frac{1}{2}$
 $\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} \left(-\frac{3}{2} \right) = \frac{1}{2}$ and $f(2) = 1$
 $\therefore \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x) \neq f(2)$
 $\therefore f(x)$ is discontinuous at $x = 2$.
25. $\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (1 - x) = 0$
 $\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (1 + x^2) = 1 + 1^2 = 2$
 $\therefore \lim_{x \rightarrow 1^+} f(x) \neq \lim_{x \rightarrow 1^-} f(x)$
 $\therefore f(x)$ is discontinuous at $x = 1$.
26. $\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (x^2 - x - 1) = 4 - 2 - 1 = 1$
 $\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} (4x + 1) = 8 + 1 = 9$
 $\therefore \lim_{x \rightarrow 2^-} f(x) \neq \lim_{x \rightarrow 2^+} f(x)$
 $\therefore f(x)$ is discontinuous at $x = 2$.
28. $\lim_{x \rightarrow 3} f(x) = \lim_{x \rightarrow 3} \sqrt{x - 2} = 1$
 $f(3) = \sqrt{3 - 2} = 1$
 $\therefore \lim_{x \rightarrow 3} f(x) = f(3)$
 $\therefore f(x)$ is continuous at $x = 3$.
 Since, $3 \in (2, 4)$
 $\therefore f(x)$ is continuous in $(2, 4)$.



29. For $x > 0$, $f(x) = x^2$
 Since f is a polynomial function, it is continuous for all $x > 0$.
 For $x < 0$, $f(x) = x^2$
 Since f is a polynomial function, it is continuous for all $x < 0$.
 $\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} x^2 = 0$
 $\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} x^2 = 0$
 $f(0) = 0$
 $\therefore f(x)$ is continuous at $x = 0$.
 $\therefore f(x)$ is continuous on \mathbb{R} .
30. $f(x)$ being a rational function, is continuous in $[0, 1]$ except at those points where the denominator $(x-2)(x-5) = 0$
 i.e., when $x = 2$ or $x = 5$
 Since $2, 5 \notin [0, 1]$
 $\therefore f(x)$ is continuous in $[0, 1]$.
31. For $x < 2$, $f(x) = x - 1$
 Since f is a polynomial function, it is continuous for all $x < 2$.
 For $x > 2$, $f(x) = 2x - 3$
 Since f is a polynomial function, it is continuous for all $x > 2$.
 $\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (x - 1) = 1$
 $\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} (2x - 3) = 1$
 $f(2) = 1$
 $\therefore f(x)$ is continuous for all real values of x .
32. Since, $f(x)$ is continuous in $[0, 3]$.
 \therefore it is continuous at $x = 2$.
 $\therefore \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x)$
 $\Rightarrow \lim_{x \rightarrow 2^-} (3x - 4) = \lim_{x \rightarrow 2^+} (2x + k)$
 $\Rightarrow 3(2) - 4 = 2(2) + k$
 $\Rightarrow 2 = 4 + k \Rightarrow k = -2$
33. Since, $f(x)$ is continuous in $[-2, 2]$.
 \therefore it is continuous at $x = 0$ and $x = 1$.
 $\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x)$
 $\Rightarrow \lim_{x \rightarrow 0^-} (x + a) = \lim_{x \rightarrow 0^+} (x + a)$
 $\Rightarrow a = 0$
 Also, $\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x)$
 $\Rightarrow \lim_{x \rightarrow 1^-} (x + a) = \lim_{x \rightarrow 1^+} (bx - 1)$
 $\Rightarrow 1 = b - 1$
 $\Rightarrow b = 2$

34. Since, $f(x)$ is continuous on $[-4, 2]$.
 \therefore it is continuous at $x = -2$.
 $\therefore \lim_{x \rightarrow -2^-} f(x) = \lim_{x \rightarrow -2^+} f(x)$
 $\Rightarrow \lim_{x \rightarrow -2^-} (6b - 3a) = \lim_{x \rightarrow -2^+} (4x + 1)$
 $\Rightarrow 6b - 3a(-2) = 4(-2) + 1$
 $\Rightarrow 6b + 6a = -7$
 $\Rightarrow a + b = -\frac{7}{6}$

**Critical Thinking**

1. Since, $f(x)$ is continuous at $x = 5$.
 $\therefore f(5) = \lim_{x \rightarrow 5} f(x)$
 $= \lim_{x \rightarrow 5} \frac{x^2 - 10x + 25}{x^2 - 7x + 10}$
 $= \lim_{x \rightarrow 5} \frac{(x-5)^2}{(x-2)(x-5)}$
 $= \frac{5-5}{5-2} = 0$
2. Since, $f(x)$ is continuous at $x = \frac{1}{2}$.
 $\therefore f\left(\frac{1}{2}\right) = \lim_{x \rightarrow \frac{1}{2}} f(x)$
 $= \lim_{x \rightarrow \frac{1}{2}} \frac{6x^5 - 1}{3x^3 - \frac{1}{8}}$
 $\Rightarrow k = \lim_{x \rightarrow \frac{1}{2}} \frac{6x^5 - 1}{3x^3 - \frac{1}{8}}$
 Applying L'Hospital rule on R.H.S., we get
 $k = \lim_{x \rightarrow \frac{1}{2}} \frac{6x^5}{3x^3} = \lim_{x \rightarrow \frac{1}{2}} 2x^2 = 2\left(\frac{1}{2}\right)^2 = \frac{1}{2}$
3. $\lim_{x \rightarrow 0} f(x) = \sin^{-1}(0) = 0 = f(0)$
 $\therefore f(x)$ is continuous at $x = 0$.
4. $\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} 2x \sin \frac{1}{x}$, but $-1 \leq \sin \frac{1}{x} \leq 1$ and $x \rightarrow 0$
 $\therefore \lim_{x \rightarrow 0} f(x) = 0 = \lim_{x \rightarrow 0} f(x) = f(0)$
 $\therefore f(x)$ is continuous at $x = 0$.
5. Since, $f(x)$ is continuous at $x = 0$.
 $\therefore \lim_{x \rightarrow 0} f(x) = f(0)$
 $\therefore \lim_{x \rightarrow 0} x^a \sin \frac{1}{x} = 0$, if $a > 0$



6. $\lim_{x \rightarrow 0^-} f(x) = \lim_{h \rightarrow 0} f(0-h)$

$$= \lim_{h \rightarrow 0} \frac{-h}{e^h + 1} = \lim_{h \rightarrow 0} \frac{-h}{1 + \frac{1}{e^h}} = 0$$

 $\lim_{x \rightarrow 0^+} f(x) = \lim_{h \rightarrow 0} f(0+h) = \lim_{h \rightarrow 0} \frac{h}{\frac{1}{e^h} + 1} = 0$
 $\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$
 $\therefore f(x)$ is continuous at $x = 0$.
7. $\lim_{x \rightarrow 0} \frac{\sin 2x}{x} = 2 \neq f(0)$
 $\lim_{x \rightarrow 0} (1+x)^{\frac{1}{x}} = e \neq f(0)$
 $\lim_{x \rightarrow 0} e^{\frac{1}{x}} = \lim_{x \rightarrow 0} \frac{1}{\frac{1}{e^x}} = \frac{1}{e^\infty} = \frac{1}{\infty} = 0 \neq f(0)$
 $\lim_{x \rightarrow 0} \left(\frac{3}{x} + \frac{4 \tan x}{x} \right) = 3 + 4 = 7 = f(0)$
 $\therefore f(x)$ is continuous at $x = 0$.
8. $\lim_{x \rightarrow 0^-} f(x) = \lim_{h \rightarrow 0^-} 5^{\frac{1}{h}} = \lim_{h \rightarrow 0^-} 5^{-\frac{1}{h}} = 0$
 $\lim_{x \rightarrow 0^+} f(x) = \lim_{h \rightarrow 0^+} \lambda \left[\frac{1}{h} \right] = 0$, for all $\lambda \in \mathbb{R}$
 $f(0) = \lambda(0) = 0$
 $\therefore f$ is continuous at $x = 0$, whatever λ may be.
9. Since, $f(x)$ is continuous at $x = a$.
 $\therefore f(a) = \lim_{x \rightarrow a} f(x)$

$$= \lim_{x \rightarrow a} \frac{-a}{\sqrt{x} - \sqrt{a}} \times \frac{\sqrt{x} + \sqrt{a}}{\sqrt{x} + \sqrt{a}}$$

$$= \lim_{x \rightarrow a} (\sqrt{x} + \sqrt{a}) = \sqrt{a} + \sqrt{a} = 2\sqrt{a}$$
10. Since, $f(x)$ is continuous at $x = 1$.
 $\therefore f(1) = \lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 1} \frac{\sqrt{x+3} - 2}{x^3 - 1}$

$$= \lim_{x \rightarrow 1} \frac{\sqrt{x+3} - 2}{x^3 - 1^3} \times \frac{\sqrt{x+3} + 2}{\sqrt{x+3} + 2}$$

$$= \lim_{x \rightarrow 1} \frac{-1}{(x-1)(x^2 + x + 1)(\sqrt{x+3} + 2)} = \frac{1}{3(4)} = \frac{1}{12}$$

11. $\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0} \frac{\sqrt{1+k} - \sqrt{1-k}}{x}$
 By rationalising, we get

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0} \frac{2k}{(\sqrt{1+k} + \sqrt{1-k})x}$$

$$= 2 \lim_{x \rightarrow 0} \frac{k}{\sqrt{1+k} + \sqrt{1-k}} = k$$

 $\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0} (2^x + 3^{-x} - 2) = -2$
 Since, $f(x)$ is continuous at $x = 0$.
 $\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x)$
 $\Rightarrow k = -2$
12. Since, $f(x)$ is continuous at $x = 4$.
 $\therefore f(4) = \lim_{x \rightarrow 4} f(x)$

$$= \lim_{x \rightarrow 4} \frac{x^4 - 64}{\sqrt{x^2 + 9} - 5}$$

$$= \lim_{x \rightarrow 4} \frac{(x^3 - 64)(\sqrt{x^2 + 9} + 5)}{(x^2 + 9) - 25}$$

$$= \left(\lim_{x \rightarrow 4} \frac{x^3 - 64}{x^2 - 4^2} \right) \left[\lim_{x \rightarrow 4} (\sqrt{x^2 + 9} + 5) \right]$$

$$= \frac{3}{2}(4) \left[4(\sqrt{16+9} + 5) \right]$$

$$= 240$$
13. Since, $f(x)$ is continuous at $x = 0$.
 $\therefore f(0) = \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{\tan(x^2 - 1)}{x - 1}$

$$= \lim_{x \rightarrow 0} \frac{\tan[(x-1)]}{(x-1)} \times (x-1) = 1 \times (-1) = -1$$
14. Since, $f(x)$ is continuous at $x = 0$.
 $\therefore f(0) = \lim_{x \rightarrow 0} f(x)$
 $\Rightarrow k = \lim_{x \rightarrow 0} \frac{1 - \cos 4x}{8x^2} = \lim_{x \rightarrow 0} \frac{2 \sin^2 2x}{8x^2}$
 $\Rightarrow k = \lim_{x \rightarrow 0} \frac{\sin^2 2x}{4x^2} = 1$
15. Since, $f(x)$ is continuous at $x = 0$.
 $\therefore f(0) = \lim_{x \rightarrow 0} f(x)$
 $\Rightarrow a = \lim_{x \rightarrow 0} \frac{1 - \cos 4x}{2}$

$$= \lim_{x \rightarrow 0} \frac{2 \sin^2 2x}{2}$$

$$= 2 \lim_{x \rightarrow 0} \frac{\sin^2 2x}{(2x)^2} \times 4 = 2 \times 4 = 8$$



16. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x)$$

$$\Rightarrow k = \lim_{x \rightarrow 0} \frac{1 - \cos 3x}{\tan x}$$

$$\Rightarrow k = \lim_{x \rightarrow 0} \frac{1 - \cos 3x}{x^2} \times \frac{1}{\frac{\tan x}{x}}$$

$$\Rightarrow k = \frac{3^2}{2} \times 1 \quad \dots \left[\because \lim_{x \rightarrow 0} \left(\frac{1 - \cos kx}{x^2} \right) = \frac{k^2}{2} \right]$$

$$\Rightarrow k = \frac{9}{2}$$

17. Since, $f(x)$ is continuous at $x = \frac{\pi}{2}$.

$$\therefore f\left(\frac{\pi}{2}\right) = \lim_{x \rightarrow \frac{\pi}{2}} f(x)$$

$$\Rightarrow 3 = \lim_{x \rightarrow \frac{\pi}{2}} \left(\frac{k \cos x}{\pi - 2x} \right)$$

Applying L'Hospital rule on R.H.S., we get

$$3 = \lim_{x \rightarrow \frac{\pi}{2}} \frac{k(-\sin x)}{-2}$$

$$\Rightarrow 3 = \frac{k}{2} \Rightarrow k = 6$$

18. Since, $f(x)$ is continuous at $x = \frac{\pi}{4}$.

$$\therefore f\left(\frac{\pi}{4}\right) = \lim_{x \rightarrow \frac{\pi}{4}} f(x)$$

$$\Rightarrow k = \lim_{x \rightarrow \frac{\pi}{4}} \frac{\cos x - \sin x}{\cos 2x}$$

$$\Rightarrow k = \lim_{x \rightarrow \frac{\pi}{4}} \frac{\cos x - \sin x}{\cos^2 x - \sin^2 x}$$

$$\Rightarrow k = \lim_{x \rightarrow \frac{\pi}{4}} \frac{\cos x - \sin x}{(\cos x - \sin x)(\cos x + \sin x)}$$

$$\Rightarrow k = \lim_{x \rightarrow \frac{\pi}{4}} \frac{1}{\cos x + \sin x} = \frac{1}{\sqrt{2}}$$

19. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x)$$

$$\Rightarrow \lambda = \lim_{x \rightarrow 0} \frac{\cos 3x - \cos x}{x^2}$$

Applying L'Hospital rule on R.H.S., we get

$$\lambda = \lim_{x \rightarrow 0} \frac{-3\sin 3x + \sin x}{2x}$$

Applying L'Hospital rule on R.H.S., we get

$$\lambda = \lim_{x \rightarrow 0} \frac{-9\cos 3x + \cos x}{2} \Rightarrow \lambda = \frac{-9+1}{2} = -4$$

20. Since, $f(x)$ is continuous at $x = \frac{\pi}{4}$.

$$\therefore f\left(\frac{\pi}{4}\right) = \lim_{x \rightarrow \frac{\pi}{4}} f(x)$$

$$\Rightarrow k = \lim_{x \rightarrow \frac{\pi}{4}} \frac{1 - \tan x}{1 - \sqrt{2} \sin x}$$

Applying L'Hospital rule on R.H.S., we get

$$k = \lim_{x \rightarrow \frac{\pi}{4}} \frac{-\sec^2 x}{-\sqrt{2} \cos x} = \frac{-2}{-1} = 2$$

21. Since, $f(x)$ is continuous at $x = \frac{\pi}{6}$,

$$\therefore \lim_{x \rightarrow \frac{\pi}{6}} f(x) = f\left(\frac{\pi}{6}\right)$$

$$\Rightarrow \lim_{x \rightarrow \frac{\pi}{6}} \frac{3\sin x - \sqrt{3}\cos x}{6 - \pi} = a$$

Applying L'Hospital rule to L.H.S, we get

$$\lim_{x \rightarrow \frac{\pi}{6}} \frac{3\cos x + \sqrt{3}\sin x}{6} = a$$

$$\Rightarrow \frac{3\left(\frac{\sqrt{3}}{2}\right) + \sqrt{3}\left(\frac{1}{2}\right)}{6} = a$$

$$\Rightarrow \frac{4\sqrt{3}}{12} = a \Rightarrow a = \frac{1}{\sqrt{3}}$$

22. Since, $f(x)$ is continuous at $x = \frac{\pi}{2}$.

$$\therefore f\left(\frac{\pi}{2}\right) = \lim_{x \rightarrow \frac{\pi}{2}} f(x)$$

$$\Rightarrow \lambda = \lim_{x \rightarrow \frac{\pi}{2}} \frac{1 - \sin x}{(\pi - 2x)^2}$$

Applying L'Hospital rule on R.H.S., we get

$$\lambda = \lim_{x \rightarrow \frac{\pi}{2}} \frac{-\cos x}{-4(\pi - 2x)}$$

Applying L'Hospital rule on R.H.S., we get

$$\lambda = \lim_{x \rightarrow \frac{\pi}{2}} \frac{\sin x}{-4(-2)} \Rightarrow \lambda = \frac{1}{8}$$



23. For $f(x)$ to be continuous at $x = 0$,

$$f(0) = \lim_{x \rightarrow 0} f(x)$$

$$\Rightarrow f(0) = \lim_{x \rightarrow 0} \frac{(a+x)^2 \sin(a+x) - a^2 \sin a}{1}$$

Applying L'Hospital rule on R.H.S., we get

$$f(0) = \lim_{x \rightarrow 0} \frac{2(a+x) \sin(a+x) + (a+x)^2 \cos(a+x)}{1}$$

$$\Rightarrow f(0) = 2a \sin a + a^2 \cos a$$

24. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{2 - \sqrt{x+4}}{\sin 2x}$$

Applying L'Hospital rule on R.H.S., we get

$$f(0) = \lim_{x \rightarrow 0} \frac{\left(-\frac{1}{2\sqrt{x+4}} \right)}{2 \cos 2x} = -\frac{1}{8}$$

25. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{(27-2x)^{\frac{1}{3}} - 3}{9 - 3(243+5x)^{\frac{1}{5}}}$$

Applying L'Hospital rule on R.H.S., we get

$$f(0) = \lim_{x \rightarrow 0} \frac{\frac{1}{3}(27-2x)^{-\frac{2}{3}}(-2)}{-\frac{3}{5}(243+5x)^{-\frac{4}{5}}(5)} = 2$$

26. For $f(x)$ to be continuous at $x = \frac{\pi}{2}$,

$$f\left(\frac{\pi}{2}\right) = \lim_{x \rightarrow \frac{\pi}{2}} f(x)$$

$$= \lim_{x \rightarrow \frac{\pi}{2}} \frac{\sqrt{2} - \sqrt{1+\sin x}}{\cos^2 x}$$

$$= \lim_{x \rightarrow \frac{\pi}{2}} \frac{2 - (1 + \sin x)}{(1 - \sin^2 x)(\sqrt{2} + \sqrt{1 + \sin x})}$$

$$= \lim_{x \rightarrow \frac{\pi}{2}} \frac{1 - \sin x}{(1 - \sin x)(1 + \sin x)(\sqrt{2} + \sqrt{1 + \sin x})}$$

$$= \lim_{x \rightarrow \frac{\pi}{2}} \frac{1}{(1 + \sin x)(\sqrt{2} + \sqrt{1 + \sin x})}$$

$$= \frac{1}{(1+1)(\sqrt{2} + \sqrt{1+1})} = \frac{1}{4\sqrt{2}}$$

27. For $f(x)$ to be continuous at $x = 0$,

$$f(0) = \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{\sqrt{1+\sin x} - \sqrt{1-\sin x}}{1}$$

Applying L'Hospital rule on R.H.S., we get

$$f(0) = \lim_{x \rightarrow 0} \frac{\frac{\cos x}{2\sqrt{1+\sin x}} + \frac{\cos x}{2\sqrt{1-\sin x}}}{1}$$

$$= \frac{1}{2}(1+1) = 1$$

28. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x)$$

$$\Rightarrow f(0) = \lim_{x \rightarrow 0} \frac{\cos^2 x - \sin^2 x - 1}{\sqrt{x^2+1} - 1}$$

$$= \lim_{x \rightarrow 0} \frac{(\cos^2 x - 1) - \sin^2 x}{\sqrt{x^2+1} - 1} \times \frac{\sqrt{x^2+1} + 1}{\sqrt{x^2+1} + 1}$$

$$= \lim_{x \rightarrow 0} \frac{(-\sin^2 x - \sin^2 x)(\sqrt{x^2+1} + 1)}{x^2 + 1 - 1}$$

$$= \lim_{x \rightarrow 0} \frac{-2\sin^2 x}{2} \times (\sqrt{x^2+1} + 1)$$

$$= -2(1)^2 (\sqrt{0^2+1} + 1) = -4$$

29. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \left(\frac{2 - 2^{-x}}{1} \right)$$

Applying L'Hospital rule on R.H.S., we get

$$f(0) = \lim_{x \rightarrow 0} \left[\frac{(2 + 2^{-x}) \log_e 2}{1} \right]$$

$$= (2^0 + 2^0) \log_e 2$$

$$\therefore f(0) = 2 \log_e 2 = \log_e 4$$

30. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{3 + 3^{-x} - 2}{2}$$

$$= \lim_{x \rightarrow 0} \frac{(3 - 1)^2}{2}$$

$$= \frac{(\log 3)^2}{1} = (\log 3)^2$$

31. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x)$$

$$\Rightarrow 2 = \lim_{x \rightarrow 0} \frac{8 - 2}{k - 1}$$

$$\Rightarrow 2 = \lim_{x \rightarrow 0} \frac{2 \left(\frac{4 - 1}{k - 1} \right)}{k - 1} \Rightarrow 2 = \frac{2^0 \log 4}{\log k}$$

$$\Rightarrow 2 \log k = \log 4$$

$$\Rightarrow 2 \log k = 2 \log 2$$

$$\Rightarrow k = 2$$



32. Since, $f(x)$ is continuous at $x = 0$.

$$\begin{aligned}\therefore f(0) &= \lim_{x \rightarrow 0} f(x) \\ &= \lim_{x \rightarrow 0} \frac{(e^3 - 1)\sin \frac{\pi}{2}}{2} \\ &= \lim_{x \rightarrow 0} \frac{e^3 - 1}{3} \times 3 \times \frac{\sin \frac{\pi}{180}}{\frac{\pi}{180}} \times \frac{\pi}{180} \\ &= 1 \times 3 \times 1 \times \frac{\pi}{180} = \frac{\pi}{60}\end{aligned}$$

33. Since, $f(x)$ is continuous at $x = 0$.

$$\begin{aligned}\therefore f(0) &= \lim_{x \rightarrow 0} f(x) \\ \Rightarrow \frac{k}{2} &= \lim_{x \rightarrow 0} \frac{e^5 - e^2}{\sin 3}\end{aligned}$$

Applying L'Hospital rule on R.H.S., we get

$$\begin{aligned}\frac{k}{2} &= \lim_{x \rightarrow 0} \frac{5e^5 - 2e^2}{3 \cos 3} \\ \Rightarrow \frac{k}{2} &= \frac{5e^0 - 2e^0}{3 \cos 0} = \frac{5 - 2}{3} = 1 \\ \Rightarrow k &= 2\end{aligned}$$

34. For $f(x)$ to be continuous at $x = 0$,

$$\begin{aligned}f(0) &= \lim_{x \rightarrow 0} f(x) \\ &= \lim_{x \rightarrow 0} \frac{e - e^{\sin x}}{2(1 - \sin x)} \\ &= \frac{1}{2} \lim_{x \rightarrow 0} e^{\sin x} \left(\frac{e^{-\sin x} - 1}{-\sin x} \right) \\ &= \frac{1}{2} \times e^0 \times 1 \quad \dots \left[\because \lim_{x \rightarrow 0} \frac{e^{-x} - 1}{-x} = 1 \right] \\ &= \frac{1}{2}\end{aligned}$$

35. For $f(x)$ to be continuous at $x = 0$,

$$\begin{aligned}f(0) &= \lim_{x \rightarrow 0} f(x) \\ &= \lim_{x \rightarrow 0} (1 + x)^{\cot x} \\ &= \lim_{x \rightarrow 0} \left\{ (1 + x)^{\frac{1}{x}} \right\}^{\cot x} \\ &= \lim_{x \rightarrow 0} \left\{ (1 + x)^{\frac{1}{x}} \right\}^{\lim_{x \rightarrow 0} \left(\frac{1}{\tan x} \right)} \\ &= e^1 = e\end{aligned}$$

36. Since, $f(x)$ is continuous at $x = 0$.

$$\begin{aligned}\therefore f(0) &= \lim_{x \rightarrow 0} f(x) \\ &= \lim_{x \rightarrow 0} \left(\frac{4 + x}{1 - 4} \right)^{\frac{1}{x}} \\ &= \lim_{x \rightarrow 0} \frac{\left[(1 + 4)^{\frac{1}{4}} \right]^4}{\left[(1 - 4)^{-\frac{1}{4}} \right]^{-4}} \\ &= \frac{e^4}{e^{-4}} = e^8\end{aligned}$$

37. Since, $f(x)$ is continuous at $x = 0$.

$$\begin{aligned}\therefore f(0) &= \lim_{x \rightarrow 0} f(x) \\ \Rightarrow k &= \lim_{x \rightarrow 0} (\sec^2 x)^{\cot^2 x} \\ \Rightarrow k &= \lim_{x \rightarrow 0} (1 + \tan^2 x)^{\frac{1}{\tan^2 x}} \\ \Rightarrow k &= e\end{aligned}$$

38. Since, $f(x)$ is continuous at $x = \frac{\pi}{2}$,

$$\begin{aligned}\therefore f\left(\frac{\pi}{2}\right) &= \lim_{x \rightarrow \frac{\pi}{2}} f(x) \\ &= \lim_{x \rightarrow \frac{\pi}{2}} (\sin x)^{\frac{1}{\pi - 2}} \\ &= \lim_{x \rightarrow \frac{\pi}{2}} [1 + (\sin x - 1)]^{\frac{1}{\pi - 2}} \\ &= e^{\lim_{x \rightarrow \frac{\pi}{2}} \left(\frac{\sin x - 1}{\pi - 2} \right)} \\ &= e^{-\frac{1}{2} \lim_{x \rightarrow \frac{\pi}{2}} \frac{1 - \cos\left(\frac{\pi}{2} - x\right)}{\left(\frac{\pi}{2} - x\right)}} \\ &= e^0 \\ &= 1\end{aligned}$$

39. Since, $f(x)$ is continuous at $x = 0$.

$$\begin{aligned}\therefore f(0) &= \lim_{x \rightarrow 0} f(x) \\ \Rightarrow 5 &= \lim_{x \rightarrow 0} \frac{\log(1 + k)}{\sin x} \\ &= \lim_{x \rightarrow 0} \frac{\log(1 + k)}{k} \times k \\ \Rightarrow 5 &= \lim_{x \rightarrow 0} \frac{k}{\sin x} \\ \Rightarrow 5 &= \frac{1 \times k}{1} \Rightarrow k = 5\end{aligned}$$



40. Since, $f(x)$ is continuous at $x = 7$.

$$\therefore f(7) = \lim_{x \rightarrow 7} f(x)$$

$$\Rightarrow k = \lim_{x \rightarrow 7} \frac{\log x - \log 7}{x - 7}$$

Applying L'Hospital rule on R.H.S., we get

$$k = \lim_{x \rightarrow 7} \frac{1}{x} = \frac{1}{7}$$

41. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x)$$

$$\Rightarrow k = \lim_{x \rightarrow 0} \frac{\log(1+2a^x) - \log(1-b^x)}{x}$$

$$\Rightarrow k = \lim_{x \rightarrow 0} \left[\frac{\log(1+2a^x)}{2a^x} \times 2a^x + \frac{\log(1-b^x)}{-b^x} \times b^x \right]$$

$$\Rightarrow k = 2a + b$$

42. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x)$$

$$\Rightarrow k = \lim_{x \rightarrow 0} \frac{(3^{\sin x} - 1)^2}{\log(1+x)}$$

$$\Rightarrow k = \lim_{x \rightarrow 0} \frac{\left(\frac{3^{\sin x} - 1}{\sin x}\right)^2 \cdot \left(\frac{\sin x}{x}\right)^2}{\log(1+x)}$$

$$\Rightarrow k = \frac{(\log 3)^2 \times (1)^2}{1} = (\log 3)^2$$

43. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x)$$

$$= \lim_{x \rightarrow 0} \frac{\log(\sec^2 x)}{\sin x}$$

$$= \lim_{x \rightarrow 0} \frac{\log(1 + \tan^2 x)}{\tan^2 x} \times \frac{\tan^2 x}{\sin x}$$

$$= \lim_{x \rightarrow 0} \frac{\log(1 + \tan^2 x)}{\tan^2 x} \times \frac{\tan^2 x}{\sin^2 x}$$

$$= 1 \times \frac{1^2}{1} = 1$$

44. Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x)$$

$$\Rightarrow 12(\log 4)^3 = \lim_{x \rightarrow 0} \frac{(4-x)^3}{\sin x - \log\left(1 + \frac{x^2}{3}\right)}$$

$$\Rightarrow 12(\log 4)^3$$

$$= \lim_{x \rightarrow 0} \left(\frac{4-x}{3} \right)^3 \times \frac{\left(\frac{-x}{p}\right)}{\left(\frac{\sin x}{p}\right)} \times \frac{p}{\frac{\log\left(1 + \frac{x^2}{3}\right)}{\frac{x^2}{3} \times 3}}$$

$$\Rightarrow 12(\log 4)^3 = (\log 4)^3 (1) \left(\frac{3p}{1}\right)$$

$$\Rightarrow p = 4$$

45. Since, $f(x)$ is continuous at $x = 3$.

$$\therefore \lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^+} f(x) = f(3)$$

$$\lim_{x \rightarrow 3^-} f(x) = f(3)$$

$$\Rightarrow \lim_{x \rightarrow 3^-} (2^{x^2} + 3^x + b) = 5$$

$$\Rightarrow 2(3)^2 + 3(3) + b = 5$$

$$\Rightarrow b = -22$$

Also, $\lim_{x \rightarrow 3^+} f(x) = f(3)$

$$\Rightarrow \lim_{x \rightarrow 3^+} \left(\frac{x^2 - 9}{-3} + a \right) = 5$$

$$\Rightarrow (3 + 3 + a) = 5$$

$$\Rightarrow a = -1$$

46. $f\left(\frac{1}{2}\right) = \left(\frac{1}{2}\right)^2 + a$

$$\Rightarrow 2 = \frac{1}{4} + a \Rightarrow a = \frac{7}{4} \quad \dots(i)$$

Since, $f(x)$ is continuous at $x = 0$.

$$\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x)$$

$$\Rightarrow \lim_{x \rightarrow 0^-} (2\sqrt{x^2 + 1} + b) = \lim_{x \rightarrow 0^+} (x^2 + a)$$

$$\Rightarrow 2\sqrt{0+1} + b = 0 + a$$

$$\Rightarrow 2 + b = \frac{7}{4} \quad \dots[\text{From (i)}]$$

$$\Rightarrow b = -\frac{1}{4}$$

47. $\lim_{h \rightarrow 0} f(x) = \lim_{h \rightarrow 0} f(4-h)$

$$= \lim_{h \rightarrow 0} \frac{4-h-4}{|4-h-4|} + a$$

$$= \lim_{h \rightarrow 0} \left(-\frac{h}{h} + a \right) = a - 1$$



- $$\lim_{h \rightarrow 0^+} f(4+h) = \lim_{h \rightarrow 0^+} f(4+h)$$

$$= \lim_{h \rightarrow 0^+} \frac{4+h-4}{|4+h-4|} + b = b + 1$$
 and $f(4) = a + b$
 Since, $f(x)$ is continuous at $x = 4$.
 $\therefore \lim_{h \rightarrow 0^+} f(4+h) = f(4) = \lim_{h \rightarrow 0^+} f(4+h)$
 $\Rightarrow a - 1 = a + b = b + 1$
 $\Rightarrow b = -1$ and $a = 1$
48.
$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} \frac{\sin(a+1) + \sin x}{(a+1) + \sin x}$$

$$= \lim_{x \rightarrow 0^-} \left[\frac{\sin(a+1)}{(a+1)} \times (a+1) + \frac{\sin x}{\sin x} \right]$$

$$= a + 1 + 1$$

$$= a + 2$$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} \frac{\sqrt{1+b^2} - \sqrt{1+b^2-x}}{b\sqrt{1+b^2-x}}$$

$$= \lim_{x \rightarrow 0^+} \frac{\sqrt{1+b^2} - \sqrt{1+b^2-x}}{b\sqrt{1+b^2-x}}$$

$$= \lim_{x \rightarrow 0^+} \frac{\sqrt{1+b^2} - 1}{b} = \frac{0}{b} = 0, \text{ if } b \neq 0$$
 Since, $f(x)$ is continuous at $x = 0$.
 $\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$
 $\Rightarrow a + 2 = 0 = c$
 $\Rightarrow a = -2, c = 0$
 $\therefore a = -2, b \neq 0$ and $c = 0$
49.
$$\lim_{h \rightarrow 0^-} f(x) = \lim_{h \rightarrow 0^-} e^{-1/h} = 0$$

$$\lim_{h \rightarrow 0^+} f(x) = \lim_{h \rightarrow 0^+} e^{1/h} = \infty$$
 $\therefore \lim_{h \rightarrow 0^-} f(x) \neq \lim_{h \rightarrow 0^+} f(x)$
 $\therefore f(x)$ is discontinuous at $x = 0$.
50.
$$\lim_{x \rightarrow 2} f(x) = \lim_{x \rightarrow 2} \frac{x^4 - 16}{x - 2}$$

$$= \lim_{x \rightarrow 2} \frac{(x-2)(x+2)(x^2+4)}{x-2}$$

$$= \lim_{x \rightarrow 2} (x+2)(x^2+4) = 32 \text{ and } f(2) = 16$$
 $\therefore \lim_{x \rightarrow 2} f(x) \neq f(2)$
 $\therefore f(x)$ is discontinuous at $x = 2$.

51. As $f(x) = \frac{|x-4|}{3-x}$ is discontinuous at $x = 0$.
 $\therefore \frac{|3-4|}{3-4}$ is discontinuous at $x = 3-4 = 0$.
 $\therefore \lim_{x \rightarrow 0} f(x) = \frac{4}{3}$
52.
$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} |x| = \lim_{x \rightarrow 0^-} (-x) = 0$$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} |x| = 0$$

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} |x-1| = 1$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} |x-1| = 1$$
53. When $x < 0, |x| = -x$
 $\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} \frac{-x}{-x} = \lim_{x \rightarrow 0^-} (-1) = -1$
 When $x > 0, |x| = x$
 $\therefore \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} \frac{x}{x} = \lim_{x \rightarrow 0^+} (1) = 1$
 $\therefore \lim_{x \rightarrow 0^-} f(x) \neq \lim_{x \rightarrow 0^+} f(x)$
 $\therefore f(x)$ is discontinuous at $x = 0$.
54.
$$\lim_{x \rightarrow 0^+} \frac{\sin x}{|x|} = \lim_{x \rightarrow 0^+} \frac{\sin x}{x} = 1$$
 and
$$\lim_{x \rightarrow 0^-} \frac{\sin x}{|x|} = \lim_{x \rightarrow 0^-} \frac{\sin x}{-x} = -1$$
 \therefore the given function is discontinuous at $x = 0$.
55.
$$\lim_{x \rightarrow 0^-} f(x) = 1 + 1 = 2$$

$$\lim_{x \rightarrow 0^+} f(x) = 0$$
 $\therefore f(x)$ is discontinuous at $x = 0$.
56.
$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} \frac{|x|}{x^2 + 2}$$

$$= \lim_{x \rightarrow 0^-} \frac{-x}{x^2 + 2}$$

$$= -\frac{1}{2}$$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} \frac{|x|}{x^2 + 2}$$

$$= \lim_{x \rightarrow 0^+} \frac{x}{x^2 + 2} = \frac{1}{2}$$
 $\therefore \lim_{x \rightarrow 0} f(x)$ does not exist.



$$57. \lim_{h \rightarrow 0^-} f(h) = \lim_{h \rightarrow 0^-} \frac{e^{-\frac{1}{h}} - 1}{e^{-\frac{1}{h}} + 1} = \lim_{h \rightarrow 0^-} \frac{\frac{1}{e^{\frac{1}{h}}} - 1}{\frac{1}{e^{\frac{1}{h}}} + 1} = \frac{0 - 1}{0 + 1} = -1$$

$$\lim_{h \rightarrow 0^+} f(h) = \lim_{h \rightarrow 0^+} \frac{e^{\frac{1}{h}} - 1}{e^{\frac{1}{h}} + 1} = \lim_{h \rightarrow 0^+} \frac{1 - \frac{1}{e^{\frac{1}{h}}}}{1 + \frac{1}{e^{\frac{1}{h}}}} = \frac{1 - 0}{1 + 0} = 1$$

$$\therefore \lim_{h \rightarrow 0^-} f(h) \neq \lim_{h \rightarrow 0^+} f(h)$$

$\therefore f(x)$ is not continuous at $x = 0$.

58. $f(x)$ is discontinuous, when $x^2 - 3x + 2 = 0$
i.e., $(x - 1)(x - 2) = 0 \Rightarrow x = 1, x = 2$

59. $f(x) = \frac{x + 1}{(x - 3)(x + 4)}$

$\therefore f(x)$ is discontinuous at $x = 3, -4$.

60. $f(x) = \frac{4 - x^2}{(4 - x^2)} = \frac{4 - x^2}{(2 + x)(2 - x)}$

Since, $f(x)$ does not exist at $x = 0, 2, -2$.

\therefore there are three points of discontinuity.

61. $\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (4 - 3x) = 4 - 6 = -2$

$$\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} (2 - 6x) = 4 - 6 = -2$$

$$f(2) = 4 - 3(2) = -2$$

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^-} (2 - 6x) = 6 - 6 = 0$$

$$\lim_{x \rightarrow 3^+} f(x) = \lim_{x \rightarrow 3^+} (x + 5) = 3 + 5 = 8$$

$$\therefore \lim_{x \rightarrow 3^-} f(x) \neq \lim_{x \rightarrow 3^+} f(x)$$

$\therefore f(x)$ is continuous at $x = 2$ and discontinuous at $x = 3$.

62. $\lim_{x \rightarrow \frac{\pi}{2}^-} f(x) = \lim_{x \rightarrow \frac{\pi}{2}^-} \sin x = \frac{\pi}{2}$

$$\lim_{x \rightarrow \frac{\pi}{2}^+} f(x) = \lim_{x \rightarrow \frac{\pi}{2}^+} \frac{\pi}{2} \sin(\pi + x) = \frac{-\pi}{2}$$

$\therefore f(x)$ is discontinuous at $x = \frac{\pi}{2}$.

63. $\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{(e^2 - 1)}{2} \times 2 \times \frac{\sin x}{x}$

$$= \log e \times 2 \times 1 = 2$$

$$\text{and } f(0) = 4$$

$\therefore f(x)$ is discontinuous at $x = 0$.

64. $\lim_{a \rightarrow 0} f(a) = \lim_{a \rightarrow 0} \frac{\sin^2 a}{(a)^2} \cdot a^2 = a^2$ and $f(0) = 1$.

$\therefore f(x)$ is discontinuous at $x = 0$,
when $a \neq \pm 1$

65. Let $f(x) = \tan x$

The point of discontinuity of $f(x)$ are those points where $\tan x$ is infinite.

i.e., $\tan x = \infty$

$$\Rightarrow x = (2n + 1) \frac{\pi}{2}, n \in \mathbb{I}$$

66. $\lim_{x \rightarrow \frac{\pi}{2}^-} f(x) = \lim_{x \rightarrow \frac{\pi}{2}^-} \frac{\sin 2x}{\sqrt{1 - \cos 2x}} = \frac{\sin \pi}{\sqrt{1 - \cos \pi}} = 0$

$$\lim_{x \rightarrow \frac{\pi}{2}^+} f(x) = \lim_{x \rightarrow \frac{\pi}{2}^+} \frac{\cos x}{\pi - 2x}$$

$$= \lim_{h \rightarrow 0} \frac{\cos\left(\frac{\pi}{2} + h\right)}{\pi - 2\left(\frac{\pi}{2} + h\right)}$$

$$= \lim_{h \rightarrow 0} \frac{-\sin h}{-2h}$$

$$= \frac{1}{2}(1) = \frac{1}{2}$$

$$\therefore \lim_{x \rightarrow \frac{\pi}{2}^-} f(x) \neq \lim_{x \rightarrow \frac{\pi}{2}^+} f(x)$$

$\therefore f(x)$ is discontinuous at $x = \frac{\pi}{2}$.

67. $\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{\cos x - 3 \tan x}{x^2 + 2 \sin x}$

$$= \lim_{x \rightarrow 0} \frac{\cos x - 3 \tan x}{x^2 + 2 \sin x}$$

$$= \lim_{x \rightarrow 0} \frac{\cos x - \frac{3 \tan x}{x}}{x + 2 \sin x}$$

$$= \frac{1 - 3}{0 + 2} = -1$$

$$\therefore \lim_{x \rightarrow 0} f(x) \neq f(0)$$

$\therefore f(x)$ is discontinuous at $x = 0$.



$$\begin{aligned}
 68. \quad \lim_{x \rightarrow 0} f(x) &= \lim_{x \rightarrow 0} \frac{5^x - e^x}{\sin 2x} = \lim_{x \rightarrow 0} \frac{5^{-1+1} - e^{-1+1}}{\sin 2} \\
 &= \lim_{x \rightarrow 0} \frac{5^{-1} - e^{-1}}{\frac{\sin 2}{2} \times 2} \\
 &= \frac{\log 5 - \log e}{2} \\
 &= \frac{1}{2} (\log 5 - 1)
 \end{aligned}$$

$$\therefore \lim_{x \rightarrow 0} f(x) \neq f(0)$$

$\therefore f(x)$ is discontinuous at $x = 0$.

69. Applying L'Hospital rule, we get

$$\begin{aligned}
 \lim_{x \rightarrow \frac{\pi}{2}} \frac{5^{\cos x} - 1}{\frac{\pi}{2} - x} &= \lim_{x \rightarrow \frac{\pi}{2}} \frac{5^{\cos x} \cdot \log 5 (-\sin x)}{-1} \\
 &= 5^{\cos \frac{\pi}{2}} \cdot \log 5 \sin \frac{\pi}{2} \\
 &= \log 5
 \end{aligned}$$

$$\text{and } f\left(\frac{\pi}{2}\right) = 2 \log 5$$

$\therefore f(x)$ is discontinuous at $x = \frac{\pi}{2}$.

Here, $\lim_{x \rightarrow \frac{\pi}{2}} f(x)$ exists but not equal to $f\left(\frac{\pi}{2}\right)$.

\therefore the discontinuity at $x = \frac{\pi}{2}$ is removable.

$$\begin{aligned}
 70. \quad \lim_{x \rightarrow 0} f(x) &= \lim_{x \rightarrow 0} \frac{(2^x - 1)^2}{\tan x \cdot \log(1+x)} \\
 &= \lim_{x \rightarrow 0} \frac{(2^x - 1)^2}{x^2} \times \frac{1}{\tan x} \times \frac{1}{\log(1+x)} \\
 &= (\log 2)^2 \times 1 = (\log 2)^2
 \end{aligned}$$

$$\text{and } f(0) = \log 4$$

$\therefore f(x)$ is discontinuous at $x = 0$.

Here, $\lim_{x \rightarrow 0} f(x)$ exists but not equal to $f(0)$.

\therefore the discontinuity at $x = 0$ is removable.

71. Since, $f(x)$ is continuous in $\left(0, \frac{\pi}{2}\right)$.

$\therefore f(x)$ is continuous at $x = \frac{\pi}{4}$.

$$\therefore f\left(\frac{\pi}{4}\right) = \lim_{x \rightarrow \frac{\pi}{4}} f(x)$$

$$\Rightarrow f\left(\frac{\pi}{4}\right) = \lim_{x \rightarrow \frac{\pi}{4}} \frac{\tan\left(\frac{\pi}{4} - x\right)}{\cot 2x}$$

Applying L'Hospital rule on R.H.S., we get

$$f\left(\frac{\pi}{4}\right) = \lim_{x \rightarrow \frac{\pi}{4}} \frac{-\sec^2\left(\frac{\pi}{4} - x\right)}{-2 \operatorname{cosec}^2 2x} = \frac{-1}{-2} = \frac{1}{2}$$

72. Since, $f(x)$ is continuous in $[-1, 1]$.

\therefore it is continuous at $x = 0$.

$$\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x)$$

$$\Rightarrow \lim_{x \rightarrow 0} \frac{\sqrt{1+p} - \sqrt{1-p}}{x} = \lim_{x \rightarrow 0} \frac{2 + 1}{-2}$$

$$\Rightarrow \lim_{x \rightarrow 0} \frac{(1+p) - (1-p)}{(\sqrt{1+p} + \sqrt{1-p})} = \frac{-1}{2}$$

$$\Rightarrow p = \frac{-1}{2}$$

73. For all $x \in \mathbb{R}$, $-1 \leq \sin x \leq 1$

$\therefore f(x)$ is continuous for all real values of x .

74. Since, $f(x)$ is continuous in $[0, 8]$.

\therefore it is continuous at $x = 2$ and $x = 4$.

$$\therefore \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x)$$

$$\Rightarrow \lim_{x \rightarrow 2^-} (x^2 + a + 6) = \lim_{x \rightarrow 2^+} (3x + 2)$$

$$\Rightarrow (2)^2 + 2a + 6 = 3(2) + 2$$

$$\Rightarrow 10 + 2a = 8$$

$$\Rightarrow a = -1 \quad \dots \text{(i)}$$

$$\text{Also, } \lim_{x \rightarrow 4^-} f(x) = \lim_{x \rightarrow 4^+} f(x)$$

$$\Rightarrow \lim_{x \rightarrow 4^-} (3x + 2) = \lim_{x \rightarrow 4^+} (2a + 5b)$$

$$\Rightarrow 3(4) + 2 = 2a(4) + 5b$$

$$\Rightarrow 14 = 8a + 5b$$

$$\Rightarrow b = \frac{22}{5} \quad \dots \text{[From (i)]}$$

75. Since, $f(x)$ is continuous in $[-2, 2]$.

\therefore it is continuous at $x = 0$ and $x = 1$.

$$\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x)$$

$$\Rightarrow \lim_{x \rightarrow 0^-} \left(\frac{\sin a}{x} - 2 \right) = \lim_{x \rightarrow 0^+} (2 + 1)$$

$$\Rightarrow a - 2 = 0 + 1 \Rightarrow a = 3$$



Also, $\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x)$

$\Rightarrow \lim_{x \rightarrow 1^-} (2x + 1) = \lim_{x \rightarrow 1^+} (2b\sqrt{x^2 + 3} - 1)$

$\Rightarrow 2(1) + 1 = 2b\sqrt{1+3} - 1$

$\Rightarrow 3 = 4b - 1$

$\Rightarrow b = 1$

$\therefore a + b = 3 + 1 = 4$

76. Since, $f(x)$ is continuous on its domain.

\therefore it is continuous at $x = 2$ and $x = 9$.

$\therefore \lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^-} f(x)$

$\Rightarrow \lim_{x \rightarrow 2^+} (a + b) = 7$

$\Rightarrow 2a + b = 7 \quad \dots(i)$

Also, $\lim_{x \rightarrow 9^-} f(x) = \lim_{x \rightarrow 9^+} f(x)$

$\Rightarrow \lim_{x \rightarrow 9^-} (a + b) = 21$

$\Rightarrow 9a + b = 21 \quad \dots(ii)$

Solving (i) and (ii), we get $a = 2, b = 3$

77. $f(x)$ is continuous in $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ except at $x = 0$.

For $f(x)$ to be continuous in $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$,

$f(0) = \lim_{x \rightarrow 0} f(x)$

$\Rightarrow f(0) = \lim_{x \rightarrow 0} \frac{e^x + e^{-x} - 2}{\sin x}$

Applying L'Hospital rule on R.H.S., we get

$f(0) = \lim_{x \rightarrow 0} \frac{e^x - e^{-x}}{\cos x + \sin x}$

Applying L'Hospital rule on R.H.S., we get

$f(0) = \lim_{x \rightarrow 0} \frac{e^x + e^{-x}}{\sin x + \cos x + \cos x}$
 $= \frac{e^0 + e^0}{0 + 2\cos 0} = \frac{1+1}{2(1)} = 1$

78. $f(x) = \frac{(x-1)(x+1)(x-2)(x+2)}{|x-1||x-2|}$

Since, $\lim_{x \rightarrow 1} \frac{-1}{|x-1|}$ does not exist.

Also, $\lim_{x \rightarrow 2} \frac{-2}{|x-2|}$ does not exist

$\therefore f(x)$ is discontinuous at $x = 1, 2$.

For any $x \neq 1, 2, f(x)$ is the quotient of two polynomials and a polynomial is everywhere continuous. Therefore, $f(x)$ is continuous for all $x \neq 1, 2$.

$\therefore f(x)$ is continuous on $\mathbb{R} - \{1, 2\}$.

79. Since, $f(x)$ is continuous in $[0, \pi]$.

\therefore it is continuous at $x = \frac{\pi}{4}$ and $x = \frac{\pi}{2}$.

$\therefore \lim_{x \rightarrow (\frac{\pi}{4})^-} f(x) = \lim_{x \rightarrow (\frac{\pi}{4})^+} f(x)$

$\Rightarrow \lim_{x \rightarrow (\frac{\pi}{4})^-} (x + a\sqrt{2} \sin x) = \lim_{x \rightarrow (\frac{\pi}{4})^+} (2 \cot x + b)$

$\Rightarrow \frac{\pi}{4} + a\sqrt{2} \left(\frac{1}{\sqrt{2}}\right) = 2 \left(\frac{\pi}{4}\right) + b$

$\Rightarrow \frac{\pi}{4} + a = \frac{\pi}{2} + b$

$\Rightarrow a - b = \frac{\pi}{4} \quad \dots(i)$

Also, $\lim_{x \rightarrow (\frac{\pi}{2})^-} f(x) = \lim_{x \rightarrow (\frac{\pi}{2})^+} f(x)$

$\Rightarrow \lim_{x \rightarrow (\frac{\pi}{2})^-} (2 \cot x + b) = \lim_{x \rightarrow (\frac{\pi}{2})^+} (a \cos 2x - b \sin x)$

$\Rightarrow 2 \left(\frac{\pi}{2}\right) + b = a(-1) - b(1)$

$\Rightarrow b = -a - b$

$\Rightarrow a + 2b = 0 \quad \dots(ii)$

From (i) and (ii), we get

$a = \frac{\pi}{6}$ and $b = \frac{-\pi}{12}$

80. Since, $f(x)$ is continuous in $(-\infty, 6)$.

\therefore it is continuous at $x = 1$ and $x = 3$.

$\therefore \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x)$

$\Rightarrow \lim_{x \rightarrow 1^-} \left(1 + \sin \frac{\pi}{2}\right) = \lim_{x \rightarrow 1^+} (a + b)$

$\Rightarrow 1 + \sin \frac{\pi}{2} = a + b$

$\Rightarrow a + b = 2 \quad \dots(i)$

Also, $\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^+} f(x)$

$\Rightarrow \lim_{x \rightarrow 3^-} (a + b) = \lim_{x \rightarrow 3^+} \left(6 \tan \frac{\pi}{12}\right)$

$\Rightarrow 3a + b = 6 \tan \frac{3\pi}{12}$

$\Rightarrow 3a + b = 6 \quad \dots(ii)$

From (i) and (ii), we get $a = 2, b = 0$



Competitive Thinking

- $f(2) = k(2)^2 = 4k$
 $\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} 3 = 3$
 Since the function is continuous at $x = 2$,
 $\lim_{x \rightarrow 2^+} f(x) = f(2)$
 $\Rightarrow 4k = 3$
 $\Rightarrow k = \frac{3}{4}$
- Since, $f(x)$ is continuous at $x = a$.
 $\therefore f(a) = \lim_{x \rightarrow a} f(x)$
 $\Rightarrow b = \lim_{x \rightarrow a} \frac{x^3 - a^3}{x - a}$
 $\Rightarrow b = 3a^{3-1} = 3a^2$
- Since, $f(x)$ is continuous at $x = 2$.
 $\therefore \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x) = f(2)$
 $\Rightarrow \lim_{x \rightarrow 2^-} (x^2 - 1) = \lim_{x \rightarrow 2^+} (2x - 1) = k$
 $\Rightarrow 3 = 3 = k \Rightarrow k = 3$
- $\lim_{x \rightarrow 5^-} f(x) = \lim_{x \rightarrow 5} (3x - 8) = 7$
 $\lim_{x \rightarrow 5^+} f(x) = \lim_{x \rightarrow 5} 2k = 2k$
 Since, $f(x)$ is continuous at $x = 5$.
 $\therefore \lim_{x \rightarrow 5^-} f(x) = \lim_{x \rightarrow 5^+} f(x)$
 $\Rightarrow 7 = 2k \Rightarrow k = \frac{7}{2}$
- Since, $f(x)$ is continuous at $x = 0$,
 $\therefore f(0) = \lim_{x \rightarrow 0^-} f(x)$
 $\Rightarrow 0^2 + \alpha = \lim_{x \rightarrow 0} 2\sqrt{x^2 + 1} + \beta$
 $\Rightarrow \alpha = 2 + \beta$
 $\Rightarrow \beta = \alpha - 2$
 Also, $f\left(\frac{1}{2}\right) = 2$
 $\Rightarrow \left(\frac{1}{2}\right)^2 + \alpha = 2$
 $\Rightarrow \frac{1}{4} + \alpha = 2 \Rightarrow \alpha = \frac{7}{4}$
 $\beta = \alpha - 2 = \frac{7}{4} - 2 = -\frac{1}{4}$
 $\therefore \alpha^2 + \beta^2 = \left(\frac{7}{4}\right)^2 + \left(-\frac{1}{4}\right)^2 = \frac{50}{16} = \frac{25}{8}$

- For $f(x)$ to be continuous at $x = 0$,
 $f(0) = \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} (1 + x)^{\frac{1}{x}} = e$
- Since, $f(x)$ is continuous at $x = -5$.
 $\therefore f(-5) = \lim_{x \rightarrow -5} f(x)$
 $\Rightarrow a = \lim_{x \rightarrow -5} \frac{x^2 + 3x - 10}{x^2 + 2x - 15}$
 $\Rightarrow a = \lim_{x \rightarrow -5} \frac{(x - 2)(x + 5)}{(x + 5)(x - 3)}$
 $\Rightarrow a = \lim_{x \rightarrow -5} \frac{-2}{-3} = \frac{2}{3}$
- Since, $f(x)$ is continuous at $x = 3$.
 $\therefore f(3) = \lim_{x \rightarrow 3} f(x)$
 $\Rightarrow 2(3) + k = \lim_{x \rightarrow 3} \frac{x^2 - 9}{x - 3}$
 $\Rightarrow 6 + k = \lim_{x \rightarrow 3} \frac{(x + 3)(x - 3)}{x - 3}$
 $\Rightarrow 6 + k = \lim_{x \rightarrow 3} (x + 3)$
 $\Rightarrow 6 + k = 6 \Rightarrow k = 0$
- Since, $f(x)$ is continuous at $x = 0$,
 $\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x)$
 $\Rightarrow \lim_{x \rightarrow 0} \frac{\sqrt{1+k} - \sqrt{1-k}}{x} = \lim_{x \rightarrow 0} \frac{2}{x-1}$
 Applying L'Hospital rule on L.H.S, we get
 $\lim_{x \rightarrow 0} \frac{\frac{k}{2\sqrt{1+k}} - \frac{(-k)}{2\sqrt{1-k}}}{1} = -1$
 $\Rightarrow \frac{k}{2} + \frac{k}{2} = -1 \Rightarrow k = -1$
- Since, $f(x)$ is continuous at $x = 0$.
 $\therefore f(0) = \lim_{x \rightarrow 0} f(x)$
 $\Rightarrow k = \lim_{x \rightarrow 0} \frac{\sin 2x}{5}$
 $\Rightarrow k = \lim_{x \rightarrow 0} \frac{\sin 2x}{2} \times \frac{2}{5}$
 $\Rightarrow k = \frac{2}{5}$
- Since, $f(x)$ is continuous at $x = 0$.
 $\therefore f(0) = \lim_{x \rightarrow 0} f(x)$
 $\Rightarrow 2k = \lim_{x \rightarrow 0} \frac{3 \sin \pi x}{5} = \lim_{x \rightarrow 0} \frac{3 \sin \pi x}{5(\pi x)} \times \pi = \frac{3\pi}{5}$
 $\therefore k = \frac{3\pi}{10}$



12. Here, $f(2) = 0$
 $\lim_{h \rightarrow 2^-} f(h) = \lim_{h \rightarrow 2^-} f(2-h) = \lim_{h \rightarrow 0} |2-h-2| = 0$

$$\lim_{h \rightarrow 2^+} f(h) = \lim_{h \rightarrow 2^+} f(2+h) = \lim_{h \rightarrow 0} |2+h-2| = 0$$

$\therefore f(h)$ is continuous at $h = 2$.

13. Here, $f(b) = 0$

$$\lim_{h \rightarrow b^-} f(h) = \lim_{h \rightarrow 0} f(b-h) = \lim_{h \rightarrow 0} |b-h-b| = 0$$

$$\lim_{h \rightarrow b^+} f(h) = \lim_{h \rightarrow 0} f(b+h) = \lim_{h \rightarrow 0} |b+h-b| = 0$$

$\therefore f(h)$ is continuous at $h = b$.

14. Here, $f\left(\frac{3\pi}{4}\right) = 1$ and $\lim_{h \rightarrow \frac{3\pi}{4}} f(h) = 1$

$$\lim_{h \rightarrow \frac{3\pi}{4}} f(h) = \lim_{h \rightarrow 0} 2 \sin \frac{2}{9} \left(\frac{3\pi}{4} + h \right)$$

$$= 2 \sin \frac{\pi}{6} = 1$$

$\therefore f(h)$ is continuous at $h = \frac{3\pi}{4}$.

15. Since, $f(h)$ is continuous at $h = 0$.

$$\therefore f(0) = \lim_{h \rightarrow 0} f(h)$$

$$= \lim_{h \rightarrow 0} \frac{1 - \cos h}{h^2} = \lim_{h \rightarrow 0} \frac{2 \sin^2 \frac{h}{2}}{h^2} \times \frac{1}{4}$$

$$\therefore f(0) = 2(1)(0) = 0$$

Alternate method:

Since, $f(h)$ is continuous at $h = 0$.

$$\therefore f(0) = \lim_{h \rightarrow 0} f(h)$$

$$\Rightarrow f(0) = \lim_{h \rightarrow 0} \frac{1 - \cos h}{h^2}$$

Applying L'Hospital rule on R.H.S., we get

$$f(0) = \lim_{h \rightarrow 0} \sin h = 0$$

16. Since, $f(h)$ is continuous at $h = 0$.

$$\therefore f(0) = \lim_{h \rightarrow 0} f(h)$$

$$\Rightarrow k = \lim_{h \rightarrow 0} (\cos h)^{\frac{1}{h}}$$

$$\Rightarrow \log k = \lim_{h \rightarrow 0} \frac{1}{h} \log(\cos h)$$

Applying L'Hospital rule on R.H.S., we get

$$\log k = \lim_{h \rightarrow 0} \frac{-\frac{\sin h}{\cos h}}{1}$$

$$\Rightarrow \log k = 0$$

$$\Rightarrow k = e^0 = 1$$

17. Since, $f(h)$ is continuous at $h = 1$.

$$\therefore f(1) = \lim_{h \rightarrow 1} f(h)$$

$$\Rightarrow k = \lim_{h \rightarrow 1} \frac{\log h}{h-1}$$

Applying L'Hospital rule on R.H.S., we get

$$\frac{1}{h}$$

$$k = \lim_{h \rightarrow 1} \frac{1}{1} = 1$$

18. For $f(h)$ to be continuous at $h = \pi$,

$$f(\pi) = \lim_{h \rightarrow \pi} f(h) = \lim_{h \rightarrow \pi} \frac{1 - \sin h + \cos h}{1 + \sin h + \cos h}$$

Applying L'Hospital rule on R.H.S., we get

$$f(\pi) = \lim_{h \rightarrow \pi} \frac{-\cos h - \sin h}{\cos h - \sin h}$$

$$\Rightarrow f(\pi) = -1$$

19. Since, $f(h)$ is continuous at $h = 0$.

$$\therefore f(0) = \lim_{h \rightarrow 0} f(h)$$

$$\Rightarrow 4 = \lim_{h \rightarrow 0} \frac{(e^k - 1)^2 \sin h}{h^3}$$

$$\Rightarrow 4 = \lim_{h \rightarrow 0} \left[k^2 \times \frac{(e^k - 1)^2}{k^2} \cdot \frac{\sin h}{h} \right]$$

$$\Rightarrow 4 = k^2$$

$$\Rightarrow k = \pm 2$$

20. For $f(h)$ to be continuous at $h = 0$,

$$f(0) = \lim_{h \rightarrow 0} f(h)$$

$$= \lim_{h \rightarrow 0} \frac{e^{2h} - \cos h}{h^2}$$

$$= \lim_{h \rightarrow 0} \frac{e^{2h} - 1 + 1 - \cos h}{h^2}$$

$$= \lim_{h \rightarrow 0} \frac{e^{2h} - 1}{h^2} + \lim_{h \rightarrow 0} \frac{1 - \cos h}{h^2}$$

$$= 1 + \frac{1}{2} = \frac{3}{2}$$

21. For $f(h)$ to be continuous at $h = 0$,

$$f(0) = \lim_{h \rightarrow 0} f(h)$$

$$\Rightarrow k = \lim_{h \rightarrow 0} \frac{\log(1+2^h) \sin \frac{\pi}{2h}}{h^2}$$

$$\Rightarrow k = \lim_{h \rightarrow 0} \frac{\log(1+2^h)}{h^2} \times 2 \times \frac{\sin \frac{\pi}{180}}{\frac{\pi}{180}} \times \frac{\pi}{180}$$

$$\Rightarrow k = 1 \times 2 \times 1 \times \frac{\pi}{180} = \frac{\pi}{90}$$



22. Since, $f(x)$ is continuous at $x = 0$.

$$\begin{aligned}\therefore f(0) &= \lim_{x \rightarrow 0} f(x) \\ \Rightarrow k &= \lim_{x \rightarrow 0} \log_{(1-3)}(1+3) \\ \Rightarrow k &= \lim_{x \rightarrow 0} \frac{\log(1+3)}{\log(1-3)} \\ \Rightarrow k &= \frac{\lim_{x \rightarrow 0} \frac{\log(1+3)}{3} \times 3}{\lim_{x \rightarrow 0} \frac{\log(1-3)}{-3} \times -3} \\ \Rightarrow k &= -1\end{aligned}$$

23. Since the function is continuous at $x = 0$,
 $\lim_{x \rightarrow 0} f(x) = f(0)$

$$\begin{aligned}\therefore k &= \lim_{x \rightarrow 0} \log(\sec^2 x)^{\cot^2 x} \\ &= \lim_{x \rightarrow 0} \cot^2 x \log \sec^2 x \\ &= \lim_{x \rightarrow 0} \frac{\log(1 + \tan^2 x)}{\tan^2 x} \\ &= 1\end{aligned}$$

24. Since, $f(x)$ is continuous at $x = 0$

$$\begin{aligned}\therefore f(0) &= \lim_{x \rightarrow 0} f(x) \\ \Rightarrow k &= \lim_{x \rightarrow 0} \left[\tan\left(\frac{\pi}{4} + x\right) \right]^{\frac{1}{\tan x}} \\ &= \lim_{x \rightarrow 0} \left(\frac{1 + \tan x}{1 - \tan x} \right)^{\frac{1}{\tan x}} \\ &= \lim_{x \rightarrow 0} \frac{\left[(1 + \tan x)^{\frac{1}{\tan x}} \right]^{\tan x}}{\left[(1 - \tan x)^{\frac{1}{-\tan x}} \right]^{\tan x}} \\ &= \frac{e^1}{e^{-1}} = e^2\end{aligned}$$

$$\begin{aligned}25. \quad \lim_{x \rightarrow 1} (\log_2 2)^{\log_2 8} \\ &= \lim_{x \rightarrow 1} [\log_2 2 + \log_2 2]^{\log_2 2^3} \\ &= \lim_{x \rightarrow 1} [1 + \log_2 2]^{3 \log_2 2} \\ &= \lim_{x \rightarrow 1} [1 + \log_2 2]^{\frac{3}{\log_2 2}} \\ &= e^{\lim_{x \rightarrow 1} \log_2 2 \times \frac{3}{\log_2 2}} \\ &= e^3\end{aligned}$$

Since the function is continuous at $x = 1$,

$$\begin{aligned}\therefore \lim_{x \rightarrow 1} f(x) &= f(1) \\ \Rightarrow e^3 &= (k-1)^3 \\ \Rightarrow e &= k-1 \\ \Rightarrow k &= e+1\end{aligned}$$

26. For $f(x)$ to be continuous at $x = 0$,

$$\begin{aligned}f(0) &= \lim_{x \rightarrow 0} f(x) \\ &= \lim_{x \rightarrow 0} \frac{\log_e(1+x) - \log_e(1-x)}{x}\end{aligned}$$

Applying L'Hospital rule on R.H.S., we get

$$\begin{aligned}f(0) &= \lim_{x \rightarrow 0} \frac{\frac{1}{1+x} + \frac{1}{1-x}}{1} \\ \Rightarrow f(0) &= 2\end{aligned}$$

27. For $f(x)$ to be continuous at $x = 0$,

$$\begin{aligned}f(0) &= \lim_{x \rightarrow 0} f(x) \\ \Rightarrow f(0) &= \lim_{x \rightarrow 0} \frac{\log(1+a) - \log(1-b)}{x}\end{aligned}$$

Applying L'Hospital rule on R.H.S., we get

$$\begin{aligned}f(0) &= \lim_{x \rightarrow 0} \frac{\frac{a}{1+a} + \frac{b}{1-b}}{1} \\ \Rightarrow f(0) &= a+b\end{aligned}$$

28. Since, $f(x)$ is continuous at $x = 2$.

$$\begin{aligned}\therefore f(2) &= \lim_{x \rightarrow 2} f(x) \\ \Rightarrow 2 &= \lim_{x \rightarrow 2} \frac{x^2 - (A+2)x + A}{x-2} \\ \Rightarrow 2 &= \lim_{x \rightarrow 2} \frac{(x-2) - A(x-1)}{x-2},\end{aligned}$$

which is true if $A = 0$

29. If $x \rightarrow 0$, then the value of $\sin^{-1} \frac{1}{x}$ passes through $[-1, 1]$ infinitely many ways, therefore limit of the function does not exist at $x = 0$. Hence, there is no value of k for which the function is continuous at $x = 0$.

30. $\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \sin^{-1} \frac{1}{x}$, but $-1 \leq \sin^{-1} \frac{1}{x} \leq 1$ and $x \rightarrow 0$

$$\therefore \lim_{x \rightarrow 0} f(x) = 0$$

Since, $f(x)$ is continuous at $x = 0$.

$$\begin{aligned}\therefore f(0) &= \lim_{x \rightarrow 0} f(x) \\ \Rightarrow k &= 0\end{aligned}$$



31. Since, $f(x)$ is continuous at $x = 1$.

$$\begin{aligned}\therefore f(1) &= \lim_{x \rightarrow 1^-} f(x) \\ &\Rightarrow 2 = \lim_{x \rightarrow 1} (a^2 - b)\end{aligned}$$

$$\Rightarrow 2 = a - b$$

The values of a and b in options (A), (B) and (C) satisfies this relation.

\therefore option (D) is the correct answer.

32. $f(x) = \sin x$

$$\therefore f(0) = \sin 0 = 0$$

$$\lim_{x \rightarrow 0^+} x^2 + a^2 = 0^2 + a^2 = a^2$$

Since the function is continuous at $x = 0$,

$$\lim_{x \rightarrow 0^+} f(x) = f(0)$$

$$\Rightarrow 0 = a^2$$

$$\Rightarrow a = 0$$

$$\lim_{x \rightarrow 1^-} x^2 + a^2 = 1^2 + a^2$$

$$= 1$$

$$f(1) = b + 2$$

$$\therefore f(1) = b + 2$$

$$\text{Now, } \lim_{x \rightarrow 1^-} f(x) = f(1)$$

$$\Rightarrow 1 = b + 2$$

$$\Rightarrow b = -1$$

$$a + b + ab = 0 - 1 + 0(-1) = -1$$

33. Since, $f(x)$ is continuous at $x = \frac{\pi}{2}$.

$$\therefore f\left(\frac{\pi}{2}\right) = \lim_{x \rightarrow \frac{\pi}{2}} f(x)$$

$$\Rightarrow \lambda = \lim_{x \rightarrow \frac{\pi}{2}} \frac{1 - \sin x}{\pi - 2x}$$

Applying L'Hospital rule on R.H.S., we get

$$\lambda = \lim_{x \rightarrow \frac{\pi}{2}} \frac{-\cos x}{-2}$$

$$\Rightarrow \lambda = \lim_{x \rightarrow \frac{\pi}{2}} \frac{\cos x}{2} = \frac{\cos \frac{\pi}{2}}{2} = 0$$

34. Since, $f(x)$ is continuous at $x = \frac{\pi}{4}$,

$$\therefore f\left(\frac{\pi}{4}\right) = \lim_{x \rightarrow \frac{\pi}{4}} f(x)$$

$$\Rightarrow k = \lim_{x \rightarrow \frac{\pi}{4}} \frac{1 - \sqrt{2} \sin x}{\pi - 4x}$$

Applying L'Hospital rule on R.H.S., we get

$$k = \lim_{x \rightarrow \frac{\pi}{4}} \frac{-\sqrt{2} \cos x}{-4} \Rightarrow k = \frac{1}{4}$$

35. Since, $f(x)$ is continuous at $x = \frac{\pi}{4}$.

$$\therefore f\left(\frac{\pi}{4}\right) = \lim_{x \rightarrow \frac{\pi}{4}} f(x)$$

$$\Rightarrow a = \lim_{x \rightarrow \frac{\pi}{4}} \frac{\tan x - \cot x}{-\frac{\pi}{4}}$$

Applying L'Hospital rule on R.H.S., we get

$$a = \lim_{x \rightarrow \frac{\pi}{4}} \frac{\sec^2 x + \operatorname{cosec}^2 x}{1}$$

$$\Rightarrow a = (\sqrt{2})^2 + (\sqrt{2})^2 = 4$$

36. For $f(x)$ to be continuous at $x = 0$,

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$$

$$\Rightarrow f(0) = \lim_{x \rightarrow 0^-} f(x)$$

$$\Rightarrow k = \lim_{h \rightarrow 0} f(0 - h) = \lim_{h \rightarrow 0} \frac{\cos \frac{\pi}{2} [0 - h]}{[0 - h]}$$

$$\Rightarrow k = \lim_{h \rightarrow 0} \frac{\cos \frac{\pi}{2} [-h]}{[-h]}$$

$$\Rightarrow k = \lim_{h \rightarrow 0} \frac{\cos\left(-\frac{\pi}{2}\right)}{-1}$$

$$\Rightarrow k = 0$$

37. If $f(x)$ is continuous from right at $x = 2$, then

$$f(2) = \lim_{x \rightarrow 2^+} f(x)$$

$$\Rightarrow k = \lim_{h \rightarrow 0} f(2 + h)$$

$$\Rightarrow k = \lim_{h \rightarrow 0} \left[(2 + h)^2 + e^{\frac{1}{2 - (2 + h)}} \right]^{-1}$$

$$\Rightarrow k = \lim_{h \rightarrow 0} \left[4 + h^2 + 4h + e^{\frac{-1}{h}} \right]^{-1}$$

$$\Rightarrow k = (4 + 0 + 0 + e^{-\infty})^{-1}$$

$$\Rightarrow k = \frac{1}{4}$$



38. Since, $f(x)$ is continuous at $x = 0$.

$$\begin{aligned}\therefore f(0) &= \lim_{x \rightarrow 0} f(x) \\ &= \lim_{x \rightarrow 0} \frac{\log_e(1 + \tan^2 x)}{\sin^3 x} \\ &= \lim_{x \rightarrow 0} \left(\frac{\log_e(1 + \tan^2 x)}{\tan^2 x} \cdot \frac{\tan^2 x}{\sin^3 x} \right) \\ &= \lim_{x \rightarrow 0} \left(\frac{\log_e(1 + \tan^2 x)}{\tan^2 x} \cdot \frac{\tan^3 x}{\sin^3 x} \cdot \frac{\tan x}{\tan x} \right)\end{aligned}$$

$$\therefore f(0) = 1$$

39. Since, $f(x)$ is continuous at $x = 0$.

$$\begin{aligned}\therefore f(0) &= \lim_{x \rightarrow 0} f(x) \\ &= \lim_{x \rightarrow 0} \left(\frac{1}{x} - \frac{2}{e^x - 1} \right) \\ &= \lim_{x \rightarrow 0} \frac{e^x - 1 - 2x}{x(e^x - 1)}\end{aligned}$$

Applying L'Hospital rule on R.H.S., we get

$$f(0) = \lim_{x \rightarrow 0} \frac{2e^x - 2}{(2e^x + 1)(e^x - 1)}$$

Applying L'Hospital rule on R.H.S., we get

$$f(0) = \lim_{x \rightarrow 0} \frac{4e^x}{2(2e^x) + e^x(2) + 2e^x}$$

$$\Rightarrow f(0) = \frac{4}{2+2} = 1$$

40. Since, $f(x)$ is continuous at $x = 0$.

$$\begin{aligned}\therefore f(0) &= \lim_{x \rightarrow 0} f(x) \\ &\Rightarrow \left(\frac{k}{16} \right) \log \left(\frac{10}{3} \right) \cdot \log 2 = \lim_{x \rightarrow 0} \frac{20 + 3x - 6x - 10x}{1 - \cos 8x} \\ &= \lim_{x \rightarrow 0} \frac{(10 - 3x)(2 - x)}{2 \sin^2 4x} \\ &= \lim_{x \rightarrow 0} \frac{\left(\frac{10 - 1x}{2} - \frac{3 - 1x}{2} \right) \cdot \left(\frac{2 - 1x}{2} \right)}{2 \times \frac{\sin^2 4}{16} \times 16} \\ &= \frac{(\log 10 - \log 3)(\log 2)}{32}\end{aligned}$$

$$\therefore \left(\frac{k}{16} \right) \log \left(\frac{10}{3} \right) \cdot \log 2 = \frac{1}{32} \log \left(\frac{10}{3} \right) \log 2$$

$$\Rightarrow \frac{k}{16} = \frac{1}{32}$$

$$\Rightarrow k = \frac{1}{2}$$

$$\Rightarrow k = 3^{\log_3 \left(\frac{1}{2} \right)} \quad \dots \left[\because a^{\log_a x} = x \right]$$

$$\begin{aligned}41. \quad \lim_{x \rightarrow 2^-} f(x) &= \lim_{h \rightarrow 0} f(2 - h) \\ &= \lim_{h \rightarrow 0} \frac{2 - h - 2}{|2 - h - 2|} + a \\ &= \lim_{h \rightarrow 0} \left(-\frac{h}{h} + a \right) = a - 1 \\ \lim_{x \rightarrow 2^+} f(x) &= \lim_{h \rightarrow 0} f(2 + h) \\ &= \lim_{h \rightarrow 0} \frac{2 + h - 2}{|2 + h - 2|} + b = b + 1\end{aligned}$$

$$\text{and } f(2) = a + b$$

Since, $f(x)$ is continuous at $x = 2$,

$$\begin{aligned}\therefore \lim_{x \rightarrow 2^-} f(x) &= f(2) = \lim_{x \rightarrow 2^+} f(x) \\ \Rightarrow a - 1 &= a + b = b + 1 \\ \Rightarrow b &= -1 \text{ and } a = 1\end{aligned}$$

42. Given, $f(x) = |x| + |x - 1|$

$$\therefore f(x) = \begin{cases} -x - (x - 1), & \text{if } x < 0 \\ -(x - 1), & \text{if } 0 \leq x < 1 \\ +(x - 1), & \text{if } x \geq 1 \end{cases}$$

$$\therefore f(x) = \begin{cases} -2x + 1, & \text{if } x < 0 \\ 1, & \text{if } 0 \leq x < 1 \\ 2x - 1, & \text{if } x \geq 1 \end{cases}$$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (-2x + 1) = 1$$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} 1 = 1$$

$$f(0) = 1$$

$$\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$$

$\therefore f(x)$ is continuous at $x = 0$.

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} 1 = 1$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (2x - 1) = 1$$

$$f(1) = 2(1) - 1 = 1$$

$$\therefore \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = f(1)$$

$\therefore f(x)$ is continuous at $x = 1$.



$$43. \lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 1} \frac{x^2 - 4x + 3}{x^2 - 1} = \lim_{x \rightarrow 1} \frac{(x-3)(x+1)}{(x-1)(x+1)} = -1$$

$$f(1) = 2$$

$$\therefore \lim_{x \rightarrow 1} f(x) \neq f(1)$$

$\therefore f(x)$ is discontinuous at $x = 1$.

$$44. \lim_{x \rightarrow a^-} f(x) = -1$$

$$\lim_{x \rightarrow a^+} f(x) = 1$$

$\therefore f(x)$ is discontinuous at $x = a$.

45. $|x|$ is continuous at $x = 0$ and $\frac{1}{|x|}$ is discontinuous at $x = 0$.

$\therefore f(x) = |x| + \frac{1}{|x|}$ is discontinuous at $x = 0$.

$$46. f(x) = \frac{x^2 + 7}{x^2(x+3) - 1(x+3)} = \frac{x^2 + 7}{(x^2 - 1)(x+3)}$$

$$= \frac{x^2 + 7}{(x-1)(x+1)(x+3)}$$

\therefore the points of discontinuity are $x = 1$, $x = -1$ and $x = -3$ only.

$$47. \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1} 2 - x^2 = 1$$

$$f(1) = 1 - 1 = 0$$

$$\lim_{x \rightarrow 1} f(x) \neq f(1)$$

\therefore The function is discontinuous at $x = 1$

$$\lim_{x \rightarrow 5^-} f(x) = \lim_{x \rightarrow 5} -10 = -5$$

$$f(5) = 2(5) = 10$$

$$\lim_{x \rightarrow 5} f(x) \neq f(5)$$

\therefore The function is discontinuous at $x = 5$

$$\lim_{x \rightarrow 3^+} f(x) = -10 = -7$$

$$f(3) = 2 - 3^2 = -7$$

$$\lim_{x \rightarrow 3} f(x) = f(3)$$

\therefore The function is continuous at $x = 3$

48. Since, $f(x)$ is not defined at $x = 0, 1, -1$ and at all other points $f(x)$ is continuous.

\therefore the given function is discontinuous at 3 points.

49. Given, $f(x) = [x]$, $x \in (-3.5, 100)$

As we know greatest integer function is discontinuous on integer values.

In given interval, the integer values are $(-3, -2, -1, 0, \dots, 99)$.

\therefore the total number of integers are 103.

50. Since, $f(x)$ is continuous at every point of its domain.

\therefore it is continuous at $x = 1$.

$$\therefore \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x)$$

$$\Rightarrow \lim_{x \rightarrow 1} (5x - 4) = \lim_{x \rightarrow 1} (4x^2 + 3bx)$$

$$\Rightarrow 1 = 4 + 3b$$

$$\Rightarrow b = -1$$

51. Since, $f(x)$ is continuous for all x .

$\therefore f(x)$ is continuous at $x = 2$.

$$\therefore f(2) = \lim_{x \rightarrow 2} f(x)$$

$$\Rightarrow k = \lim_{x \rightarrow 2} \frac{x^3 + x^2 - 16x + 20}{(x-2)^2}$$

$$= \lim_{x \rightarrow 2} \frac{(x-2)(x^2 + 3x - 10)}{(x-2)^2}$$

$$= \lim_{x \rightarrow 2} \frac{(x-2)^2(x+5)}{(x-2)^2}$$

$$= 7$$

52. Since, $f(x) = [x]$ is continuous at every non integer points.

\therefore option (C) is the correct answer.

53. Let $g(x) = |x|$ and $h(x) = \sin x$.

Then, $f(x) = (hog)(x)$ for all $x \in \mathbb{R}$.

As both g and h are continuous functions on \mathbb{R} .

$\therefore f(x)$ is also continuous for all $x \in \mathbb{R}$.

54. Since, $f(x)$ is continuous in $\left[0, \frac{\pi}{2}\right]$.

\therefore it is continuous at $x = \frac{\pi}{4}$.

$$\therefore f\left(\frac{\pi}{4}\right) = \lim_{x \rightarrow \frac{\pi}{4}} f(x) = \lim_{x \rightarrow \frac{\pi}{4}} \frac{1 - \tan x}{4 - \pi}$$



Applying L'Hospital rule on R.H.S., we get

$$f\left(\frac{\pi}{4}\right) = \lim_{x \rightarrow \frac{\pi}{4}} \frac{-\sec^2}{4}$$

$$\Rightarrow f\left(\frac{\pi}{4}\right) = \frac{-2}{4} = \frac{-1}{2}$$

55. Since, $f(x)$ is continuous at each point of its domain.

\therefore it is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x)$$

$$= \lim_{x \rightarrow 0} \left(\frac{2 - \sin^{-1} x}{2 + \tan^{-1} x} \right)$$

Applying L'Hospital rule on R.H.S., we get

$$f(0) = \lim_{x \rightarrow 0} \frac{\left(2 - \frac{1}{\sqrt{1-x^2}} \right)}{\left(2 + \frac{1}{1+x^2} \right)}$$

$$= \frac{2-1}{2+1} = \frac{1}{3}$$

56. The given function is defined only in the interval $[1, \infty)$. For $x > 2$, $y = 3x - 2$ which is a straight line, hence continuous. Also, the given function is continuous at $x = 2$.

\therefore option (C) is the correct answer.

57. $\lim_{x \rightarrow 1^-} f(x) = 0$, $\lim_{x \rightarrow 1^+} f(x) = 0$ and

$$f(1) = 0$$

$\therefore \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = f(1)$

$\therefore f(x)$ is continuous at $x = 1$.

$$\lim_{x \rightarrow 2^-} f(x) = 0 \text{ and } \lim_{x \rightarrow 2^+} f(x) = 1$$

$\therefore \lim_{x \rightarrow 2^-} f(x) \neq \lim_{x \rightarrow 2^+} f(x)$

$\therefore f(x)$ is not continuous at $x = 2$.

58. Since, $f(x)$ is continuous over $[-\pi, \pi]$.

\therefore it is continuous at $x = -\frac{\pi}{2}$ and $x = \frac{\pi}{2}$.

$$\therefore \lim_{x \rightarrow -\frac{\pi}{2}} f(x) = \lim_{x \rightarrow -\frac{\pi}{2}} f(x)$$

$$\Rightarrow \lim_{x \rightarrow -\frac{\pi}{2}} (-2 \sin x) = \lim_{x \rightarrow -\frac{\pi}{2}} (\alpha \sin x + \beta)$$

$$\Rightarrow -2(-1) = \alpha(-1) + \beta$$

$$\Rightarrow -\alpha + \beta = 2 \quad \dots(i)$$

$$\text{Also, } \lim_{x \rightarrow \frac{\pi}{2}} f(x) = \lim_{x \rightarrow \frac{\pi}{2}} f(x)$$

$$\Rightarrow \lim_{x \rightarrow \frac{\pi}{2}} (\alpha \sin x + \beta) = \lim_{x \rightarrow \frac{\pi}{2}} (\cos x)$$

$$\Rightarrow \alpha(1) + \beta = 0$$

$$\Rightarrow \alpha + \beta = 0 \quad \dots(ii)$$

From (i) and (ii), we get

$$\alpha = -1, \beta = 1$$

59. For $f(x)$ to be continuous at $x = \frac{-\pi}{2}$,

$$\lim_{x \rightarrow \frac{-\pi}{2}^+} f(x) = \lim_{x \rightarrow \frac{-\pi}{2}^-} f(x) = f\left(\frac{-\pi}{2}\right)$$

$$\therefore \lim_{x \rightarrow \frac{-\pi}{2}} A \sin x + B = -2 \sin\left(\frac{-\pi}{2}\right)$$

$$\Rightarrow -A + B = 2$$

$$\Rightarrow A - B = -2 \quad \dots(i)$$

For $f(x)$ to be continuous at $x = \frac{\pi}{2}$,

$$\lim_{x \rightarrow \frac{\pi}{2}^+} f(x) = \lim_{x \rightarrow \frac{\pi}{2}^-} f(x) = f\left(\frac{\pi}{2}\right)$$

$$\therefore \lim_{x \rightarrow \frac{\pi}{2}} A \sin x + B = \cos \frac{\pi}{2}$$

$$\Rightarrow A + B = 0$$

$$\dots(ii)$$

On solving (i) and (ii), we get

$$A = -1, B = 1$$

60. Since, $f(x)$ is continuous for all x in \mathbb{R} .

$\therefore f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0^-} f(x)$$

$$\Rightarrow q = \lim_{x \rightarrow 0} \frac{\sin(p+1) + \sin x}{x}$$

$$\Rightarrow q = \lim_{x \rightarrow 0} \left[(p+1) \times \frac{\sin(p+1)}{(p+1)} + \frac{\sin x}{x} \right]$$

$$\Rightarrow q = (p+1) + 1$$

$$\Rightarrow q = p + 2$$

The values of p and q in option (C) satisfies this condition.



61. Since, f is continuous at every point in \mathbb{R} .

$\therefore f$ is continuous at $x = 2n$.

$$\therefore \lim_{x \rightarrow (2n)^-} f(x) = \lim_{x \rightarrow (2n)^+} f(x) = f(2n)$$

$$\Rightarrow \lim_{x \rightarrow (2n)^-} (b_n + \cos \pi) = \lim_{x \rightarrow (2n)^+} (a_n + \sin \pi)$$

$$\Rightarrow b_n + \cos 2n\pi = a_n + \sin 2n\pi$$

$$\Rightarrow b_n + 1 = a_n \Rightarrow a_n - b_n = 1$$

So, option (C) is correct.

Also, f is continuous at $x = 2n + 1$.

$$\lim_{x \rightarrow (2n+1)^-} f(x) = \lim_{x \rightarrow (2n+1)^+} f(x) = f(2n + 1)$$

$$\Rightarrow \lim_{x \rightarrow (2n+1)^-} (a_n + \sin \pi) = \lim_{x \rightarrow (2n+1)^+} (b_{n+1} + \cos \pi)$$

$$\Rightarrow a_n + \sin(2n + 1)\pi = b_{n+1} + \cos(2n + 1)\pi$$

$$\dots[\because f(x) = b_{n+1} + \cos \pi, x \in (2n + 1, 2n + 2)]$$

$$\Rightarrow a_n = b_{n+1} - 1$$

$$\Rightarrow a_n - b_{n+1} = -1$$

Replacing n by $n - 1$, we get

$$a_{n-1} - b_n = -1$$

So, options (A) and (D) are correct.

Hence, option (B) does not hold.

63. For $f(x)$ to be continuous at $x = 0$,

$$f(0) = \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \left[(1+x)^{\frac{1}{2}} - (1+x)^{\frac{1}{3}} \right]$$

$$= \lim_{x \rightarrow 0} \left[\left(1 + \frac{1}{2}x - \frac{1}{8}x^2 + \dots \right) - \left(1 + \frac{1}{3}x - \frac{1}{9}x^2 + \dots \right) \right]$$

$$= \lim_{x \rightarrow 0} \left[\left(\frac{1}{2} - \frac{1}{3} \right)x + \left(\frac{1}{9} - \frac{1}{8} \right)x^2 + \dots \right]$$

$$= \lim_{x \rightarrow 0} \left[\left(\frac{1}{2} - \frac{1}{3} \right)x + \left(\frac{1}{9} - \frac{1}{8} \right)x^2 + \dots \right]$$

$$\therefore f(0) = \frac{1}{6}$$



Evaluation Test

1. $f(0) = \lim_{x \rightarrow 0} \frac{1 - \cos(1 - \cos x)}{4}$

$$= \lim_{x \rightarrow 0} \frac{2 \sin^2 \left[\frac{2 \sin^2 \left(\frac{x}{2} \right)}{2} \right]}{4}$$

$$= \lim_{x \rightarrow 0} \frac{2 \sin^2 \left[\sin^2 \left(\frac{x}{2} \right) \right] \left[\sin^2 \left(\frac{x}{2} \right) \right]^2}{4 \left[\sin^2 \left(\frac{x}{2} \right) \right]^2}$$

$$= 2 \lim_{x \rightarrow 0} \frac{\sin^4 \left(\frac{x}{2} \right)}{\left(\frac{x}{2} \right)^4} = \frac{1}{2^3} = \frac{1}{8}$$

2. f is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0} \left[\frac{\log(1+x^2) - \log(1-x^2)}{\sec x - \cos x} \right]$$

$$= \lim_{x \rightarrow 0} \left[\frac{\log(1+x^2) - \log(1-x^2)}{\left[\frac{1 - \cos^2 x}{\cos x} \right]} \right]$$

$$= \lim_{x \rightarrow 0} \left[\frac{\cos x \left[\log(1+x^2) - \log(1-x^2) \right]}{\sin^2 x} \right]$$

$$= \lim_{x \rightarrow 0} \left[\frac{\cos x \left[\frac{\log(1+x^2)}{2} + \frac{\log(1-x^2)}{-2} \right]}{\left(\frac{\sin^2 x}{2} \right)} \right]$$

$$= (\cos 0) \left[\frac{1+1}{1^2} \right] = 2$$



3. For $f(x)$ to be continuous at $x = 0$, we must have $f(0) = \lim_{x \rightarrow 0} f(x)$

$$\begin{aligned} &= \lim_{x \rightarrow 0} \frac{\sqrt{a^2 - a + 2} - \sqrt{a^2 + a + 2}}{\sqrt{a+} - \sqrt{a-}} \\ &= \lim_{x \rightarrow 0} \frac{\sqrt{a^2 - a + 2} - \sqrt{a^2 + a + 2}}{\sqrt{a+} - \sqrt{a-}} \\ &\quad \times \frac{\sqrt{a+} + \sqrt{a-}}{\sqrt{a+} + \sqrt{a-}} \times \frac{\sqrt{a^2 - a + 2} + \sqrt{a^2 + a + 2}}{\sqrt{a^2 - a + 2} + \sqrt{a^2 + a + 2}} \\ &= \lim_{x \rightarrow 0} \frac{[(a^2 - a + 2) - (a^2 + a + 2)][\sqrt{a+} + \sqrt{a-}]}{[(a+)(a-)][\sqrt{a^2 - a + 2} + \sqrt{a^2 + a + 2}]} \\ &= \lim_{x \rightarrow 0} \frac{-2a(\sqrt{a+} + \sqrt{a-})}{2(\sqrt{a^2 - a + 2} + \sqrt{a^2 + a + 2})} \\ &= \frac{-a(\sqrt{a+} + \sqrt{a-})}{\sqrt{a^2} + \sqrt{a^2}} \end{aligned}$$

$$\therefore f(0) = -\sqrt{a}$$

$$\begin{aligned} 4. \quad &\lim_{x \rightarrow 0} \frac{5 \cdot 2 + 7 - 7 \cdot 2 - 5}{2 \sin^2 \frac{x}{2}} \\ &= \lim_{x \rightarrow 0} \frac{5(2-1) - 7(2-1)}{2 \sin^2 \frac{x}{2}} \\ &= \lim_{x \rightarrow 0} \frac{1}{2} \left(\frac{2-1}{\sin^2 \frac{x}{2}} \right) \left(\frac{5-1-7-1}{\frac{x}{2} \times \frac{1}{4}} \right) \\ &= 2(\log 2) \left(\log \frac{5}{7} \right) \end{aligned}$$

It is discontinuous at $x = 0$ and it is removable.

$$\begin{aligned} 5. \quad a &= \lim_{x \rightarrow 0} \frac{\sin^3 \sqrt{x} \log(1+3x)}{(\tan^{-1} \sqrt{x})^2 (e^{5\sqrt{x}} - 1)} \\ &= \lim_{x \rightarrow 0} \frac{\frac{\sin^3 \sqrt{x}}{(\sqrt{x})^3} \cdot (\sqrt{x})^3 \cdot \frac{\log(1+3x)}{3} \cdot 3}{\frac{(\tan^{-1} \sqrt{x})^2}{(\sqrt{x})^2} \cdot (\sqrt{x})^2 \cdot \frac{e^{5\sqrt{x}} - 1}{5\sqrt{x}} \cdot 5\sqrt{x}} \end{aligned}$$

$$\begin{aligned} &= \frac{(1)^3 (\sqrt{x})^3 \cdot (1)3}{(1)^2 (\sqrt{x})^2 (1)5\sqrt{x}} \\ &= \frac{3}{5} \end{aligned}$$

6. For f to be continuous at $x = 2$,

$$\begin{aligned} f(2) &= \lim_{x \rightarrow 0} (x-1)^{\frac{1}{2-x}} \\ &= \lim_{x \rightarrow 0} (1+(x-2))^{-\frac{1}{2-x}} = e^{-1} \end{aligned}$$

7. Given function is continuous at $(-\infty, 6)$.

\therefore at $x = 1$ and $x = 3$, function is continuous.

If the function $f(x)$ is continuous at $x = 1$, then

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x)$$

$$\Rightarrow 1 + \sin \frac{\pi}{2} = a + b$$

$$\therefore a + b = 2 \quad \dots (i)$$

If the function is continuous at $x = 3$, then

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^+} f(x)$$

$$\Rightarrow 3a + b = 6 \tan \frac{3\pi}{12}$$

$$\therefore 3a + b = 6 \quad \dots (ii)$$

From (i) and (ii), $a = 2$, $b = 0$

8. Since, $\sin x$ and $|x|$ are continuous for all x .

$\therefore \sin x + |x|$ is continuous for $x \in (-\infty, \infty)$.

9. For $f(x)$ to be continuous at $x = 0$, we must have

$$\lim_{x \rightarrow 0^-} f(x) = f(0) = \lim_{x \rightarrow 0^+} f(x)$$

$$\begin{aligned} \lim_{x \rightarrow 0^+} f(x) &= \lim_{x \rightarrow 0^+} e^{\tan 2x / \tan 3x} \\ &= \lim_{x \rightarrow 0^+} e^{\left(\frac{\tan 2x}{2} \times 2 \right) / \left(\frac{\tan 3x}{3} \times 3 \right)} \\ &= e^{\frac{2}{3}} \end{aligned}$$

$$f(0) = \lim_{x \rightarrow 0^+} f(x)$$

$$\Rightarrow b = e^{\frac{2}{3}}$$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (1 + |\sin x|)^{a/|\sin x|}$$

$$= e^{\lim_{x \rightarrow 0} \left(|\sin x| \times \frac{a}{|\sin x|} \right)} = e^a$$

$$f(0) = \lim_{x \rightarrow 0^-} f(x)$$

$$\Rightarrow b = e^a \Rightarrow e^{\frac{2}{3}} = e^a$$

$$\Rightarrow a = \frac{2}{3}$$



10. Given, $f(x) = [x]^2 - [x^2]$
- $-1 < x < 0$, $f(x) = (-1)^2 - 0 = 1$
 $= 0$, $f(x) = 0^2 - 0 = 0$
- $0 < x < 1$, $f(x) = 0^2 - 0 = 0$
 $= 1$, $f(x) = 1^2 - 1 = 0$
- $1 < x < \sqrt{2}$, $f(x) = 1^2 - 1 = 0$
 $= \sqrt{2}$, $f(x) = 1^2 - 2 = -1$
- $\sqrt{2} < x < \sqrt{3}$, $f(x) = 1^2 - 2 = -1$
 $= \sqrt{3}$, $f(x) = 1^2 - 3 = -2$
- $\sqrt{3} < x < 2$, $f(x) = 1^2 - 3 = -2$
 $= 2$, $f(x) = 2^2 - 4 = 0$
- $2 < x < \sqrt{5}$, $f(x) = 2^2 - 4 = 0$
 $= \sqrt{5}$, $f(x) = 2^2 - 5 = -1$

Hence, the given function is discontinuous at all integers except 1.

02 Differentiation



Hints



Classical Thinking

- $$f'(2^-) = \lim_{h \rightarrow 0^-} \frac{f(2+h) - f(2)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(2+h+1) - 3}{h}$$

$$= \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

$$f'(2^+) = \lim_{h \rightarrow 0^+} \frac{f(2+h) - f(2)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2(2+h) - 1 - 3}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2h}{h} = 2$$

$\therefore f'(2^-) \neq f'(2^+)$
 $\therefore f'(2)$ does not exist.
- $$f'(3^-) = \lim_{h \rightarrow 0^-} \frac{f(3+h) - f(3)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(3+h+2) - 5}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

$$f'(3^+) = \lim_{h \rightarrow 0^+} \frac{f(3+h) - f(3)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{8 - (3+h) - 5}{h} = \lim_{h \rightarrow 0} \frac{-h}{h} = -1$$

$\therefore f'(3^-) \neq f'(3^+)$
 $\therefore f'(3)$ does not exist.
- $$f'(0^+) = \lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0^+} \frac{|h| - 0}{h}$$

$$= \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

$$f'(0^-) = \lim_{h \rightarrow 0^-} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0^-} \frac{|h| - 0}{h}$$

$$= \lim_{h \rightarrow 0} \frac{-h}{h} = -1$$

$\therefore f'(0^+) \neq f'(0^-)$
 $\therefore f'(0)$ does not exist.

- $$f'(1^-) = \lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{x - 1}$$

$$= \lim_{x \rightarrow 1} \frac{p^2 + 1 - p - 1}{-1} = \lim_{x \rightarrow 1} \frac{p^2 - 1}{-1}$$

$$= p \lim_{x \rightarrow 1} (x + 1)$$

$$= 2p$$

$$f'(1^+) = \lim_{x \rightarrow 1^+} \frac{f(x) - f(1)}{x - 1} = \lim_{x \rightarrow 1} \frac{+p - p - 1}{-1}$$

$$= \lim_{x \rightarrow 1} \frac{-1}{-1}$$

$$= 1$$

Since, $f(x)$ is differentiable at $x = 1$.

$\therefore 2p = 1 \Rightarrow p = \frac{1}{2}$
- $$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1} = 1$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1} (2x - 1) = 1$$

$$f(1) = 1$$

$\therefore f(x)$ is continuous at $x = 1$.

$$f'(1^-) = \lim_{h \rightarrow 0^-} \frac{f(1+h) - f(1)}{h} = \lim_{h \rightarrow 0} \frac{1+h-1}{h}$$

$$= \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

$$f'(1^+) = \lim_{h \rightarrow 0^+} \frac{f(1+h) - f(1)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2(1+h) - 1 - 1}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2h}{h} = 2$$

$\therefore f'(1^-) \neq f'(1^+)$
 $\therefore f(x)$ is not differentiable at $x = 1$.
- $$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2} (x + 1) = 3$$

$$\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2} (5 - x) = 3$$

$$f(2) = 1 + 2 = 3$$

$\therefore f(x)$ is continuous at $x = 2$.



$$f'(2^-) = \lim_{h \rightarrow 0^-} \frac{f(2+h) - f(2)}{h}$$

$$= \lim_{h \rightarrow 0^-} \frac{1 + (2+h) - 3}{h}$$

$$= \lim_{h \rightarrow 0^-} \frac{h}{h} = 1$$

$$f'(2^+) = \lim_{h \rightarrow 0^+} \frac{f(2+h) - f(2)}{h}$$

$$= \lim_{h \rightarrow 0^+} \frac{5 - (2+h) - 3}{h}$$

$$= \lim_{h \rightarrow 0^+} \frac{-h}{h} = -1$$

$$\therefore f'(2^-) \neq f'(2^+)$$

$\therefore f(x)$ is not differentiable at $x = 2$.

7. $f(0) = 0$

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \sin \frac{1}{x} = 0$$

$\therefore f(x)$ is continuous at $x = 0$.

$$f(0^-) = \lim_{h \rightarrow 0^-} \frac{f(x) - f(0)}{x - 0}$$

$$= \lim_{h \rightarrow 0^-} \frac{f(0-h) - f(0)}{0-h-0}$$

$$= \lim_{h \rightarrow 0^-} \frac{-h \sin\left(\frac{1}{-h}\right) - 0}{-h}$$

$$= -\lim_{h \rightarrow 0^-} \sin\left(\frac{1}{h}\right)$$

= (a number which oscillates between -1 and 1)

$\therefore f(0^-)$ does not exist.

$\therefore f(x)$ is not differentiable at $x = 0$.

8. $\frac{d}{dx} [\sin(2x+3)] = \cos(2x+3) \cdot \frac{d}{dx} (2x+3)$
 $= 2 \cos(2x+3)$

9. $y = e^{\sqrt{x}} \Rightarrow \frac{dy}{dx} = e^{\sqrt{x}} \cdot \frac{d}{dx}(\sqrt{x}) \Rightarrow \frac{dy}{dx} = \frac{e^{\sqrt{x}}}{2\sqrt{x}}$

10. $\frac{d}{dx}(e^3) = e^3 \cdot \frac{d}{dx}(3) = 3^2 \cdot e^3$

11. Let $y = (\log x)^4$

$$\therefore \frac{dy}{dx} = 4(\log x)^3 \cdot \frac{d}{dx}(\log x)$$

$$= \underline{4(\log x)^3}$$

12. $\frac{d}{dx} [\log(\log x)] = \frac{1}{\log x} \cdot \frac{d}{dx}(\log x)$
 $= \frac{1}{\log x} \cdot \frac{1}{x} = (\log x)^{-1}$

13. $= \log_{10} |x| = \frac{\log_e |x|}{\log_e 10}$

$$\therefore \frac{d}{dx} = \frac{1}{\log_e 10} \cdot \frac{1}{|x|} \cdot \frac{d}{dx} |x| = \frac{1}{\log_e 10}$$

14. $y = f(a^2 + b)$

$$\therefore \frac{dy}{dx} = f'(a^2 + b) \cdot \frac{d}{dx}(a^2 + b) = 2a \cdot f'(a^2 + b)$$

15. $y = (4^3 - 5^2 + 1)^4$

$$\therefore \frac{dy}{dx} = 4(4^3 - 5^2 + 1)^3 \cdot \frac{d}{dx}(4^3 - 5^2 + 1)$$

$$= 4(4^3 - 5^2 + 1)^3 (12^2 - 10)$$

16. $\frac{d}{dx} (x^2 + \cos x)^4 = 4(x^2 + \cos x)^3 \cdot \frac{d}{dx} (x^2 + \cos x)$
 $= 4(x^2 + \cos x)^3 (2x - \sin x)$

17. $\frac{d}{dx} = \frac{d}{du} \cdot \frac{du}{dx}$
 $= \frac{2}{(u+1)^2} \cdot \frac{1}{2\sqrt{u}}$
 $= \frac{1}{(\sqrt{u}+1)^2} \cdot \frac{1}{\sqrt{u}}$
 $= \frac{1}{\sqrt{u}(1+\sqrt{u})^2}$

18. $y = \log(\tan \sqrt{x})$

$$\therefore \frac{dy}{dx} = \frac{1}{\tan \sqrt{x}} \cdot \frac{d}{dx}(\tan \sqrt{x})$$

$$= \frac{1}{\tan \sqrt{x}} \cdot \sec^2 \sqrt{x} \cdot \frac{1}{2\sqrt{x}}$$

$$= \frac{\sec^2 \sqrt{x}}{2\sqrt{x} \tan \sqrt{x}}$$

19. $y = \log(\sec x + \tan x)$

$$\therefore \frac{dy}{dx} = \frac{1}{\sec x + \tan x} \cdot \frac{d}{dx}(\sec x + \tan x)$$

$$= \frac{\sec x \tan x + \sec^2 x}{\sec x + \tan x}$$

$$= \sec x$$



$$\begin{aligned}
 20. \quad &= \log(\log(\log^3)) \\
 \therefore \frac{d}{d} &= \frac{1}{\log(\log^3)} \cdot \frac{d}{d} [\log(\log^3)] \\
 &= \frac{1}{\log(\log^3)} \cdot \frac{1}{\log^3} \cdot \frac{d}{d} (\log^3) \\
 &= \frac{1}{\log(\log^3)} \cdot \frac{1}{3\log} \cdot \frac{1}{3} \cdot 3^2 \\
 &= \frac{1}{\log \log(\log^3)}
 \end{aligned}$$

21. Derivative exists if $1 - x^2 > 0$ i.e., $1 > x^2$
i.e., $x^2 < 1$ i.e., $|x| < 1$ i.e., $-1 < x < 1$

$$\begin{aligned}
 22. \quad \frac{d}{d} [\tan^{-1}(\sqrt{x})] &= \frac{1}{1+(\sqrt{x})^2} \cdot \frac{1}{2\sqrt{x}} \\
 &= \frac{1}{2\sqrt{x}(1+x)}
 \end{aligned}$$

$$\begin{aligned}
 23. \quad &= \cos^{-1}\left(\frac{1}{3}\right) \\
 \therefore &= \sec^{-1}(3) \\
 \therefore \frac{d}{d} &= \frac{1}{3\sqrt{(3)^2-1}} \cdot 3^2 = \frac{3}{\sqrt{6}-1}
 \end{aligned}$$

$$\begin{aligned}
 24. \quad \text{Put } x &= \sin \theta \Rightarrow \theta = \sin^{-1} x \\
 \therefore &= \tan^{-1}\left(\frac{\sin \theta}{\sqrt{1-\sin^2 \theta}}\right) = \tan^{-1}\left(\frac{\sin \theta}{\cos \theta}\right) \\
 &= \tan^{-1}(\tan \theta) = \theta = \sin^{-1} x \\
 \therefore \frac{d}{d} &= \frac{1}{\sqrt{1-x^2}}
 \end{aligned}$$

$$\begin{aligned}
 25. \quad \text{Let } x &= \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right) \\
 \text{Put } x &= \tan \theta \Rightarrow \theta = \tan^{-1} x \\
 \therefore &= \cos^{-1}\left(\frac{1-\tan^2 \theta}{1+\tan^2 \theta}\right) \\
 \Rightarrow &= \cos^{-1}(\cos 2\theta) = 2\theta = 2 \tan^{-1} x \\
 \therefore \frac{d}{d} &= \frac{2}{1+x^2}
 \end{aligned}$$

$$\begin{aligned}
 26. \quad \text{Let } x &= \operatorname{cosec}^{-1}\left(\frac{1+x^2}{2}\right) \\
 &= \sin^{-1}\left(\frac{2}{1+x^2}\right)
 \end{aligned}$$

$$\begin{aligned}
 \text{Put } x &= \tan \theta \Rightarrow \theta = \tan^{-1} x \\
 \therefore &= \sin^{-1}\left(\frac{2 \tan \theta}{1+\tan^2 \theta}\right) \\
 &= \sin^{-1}(\sin 2\theta) \\
 &= 2\theta = 2 \tan^{-1} x \\
 \therefore \frac{d}{d} &= \frac{2}{1+x^2}
 \end{aligned}$$

$$\begin{aligned}
 27. \quad \text{Put } x &= \tan \theta \Rightarrow \theta = \tan^{-1} x \\
 \therefore &= \sin^{-1}\left(\frac{1-\tan^2 \theta}{1+\tan^2 \theta}\right) = \sin^{-1}(\cos 2\theta) \\
 &= \sin^{-1}\left(\sin\left(\frac{\pi}{2}-2\theta\right)\right) \\
 &= \frac{\pi}{2} - 2\theta = \frac{\pi}{2} - 2 \tan^{-1} x \\
 \therefore \frac{d}{d} &= -\frac{2}{1+x^2}
 \end{aligned}$$

$$\begin{aligned}
 28. \quad \text{Put } x &= \cos \theta \Rightarrow \theta = \cos^{-1} x \\
 \therefore &= \sec^{-1}\left(\frac{1}{2\cos^2 \theta - 1}\right) \\
 &= \sec^{-1}\left(\frac{1}{\cos 2\theta}\right) \\
 &= \sec^{-1}(\sec 2\theta) \\
 &= 2\theta = 2 \cos^{-1} x \\
 \therefore \frac{d}{d} &= -\frac{2}{\sqrt{1-x^2}}, \quad x \neq \pm 1
 \end{aligned}$$

$$\begin{aligned}
 29. \quad \text{Let } x &= e^{\sin \theta} \\
 \text{Taking logarithm on both sides, we get} \\
 \log x &= \sin \theta \\
 \text{Differentiating both sides w.r.t. } \theta, \text{ we get} \\
 \frac{1}{x} \cdot \frac{d}{d} x &= \cos \theta
 \end{aligned}$$

$$\therefore \frac{d}{d} x = e^{\sin \theta} (\sin \theta + \cos \theta)$$

$$\begin{aligned}
 30. \quad \text{Let } x &= e^{\log \theta} \\
 \text{Taking logarithm on both sides, we get} \\
 \log x &= \log \theta \\
 \text{Differentiating both sides w.r.t. } \theta, \text{ we get} \\
 \frac{1}{x} \cdot \frac{d}{d} x &= \frac{1}{\theta} + \log \theta \cdot 1
 \end{aligned}$$

$$\begin{aligned}
 \therefore \frac{d}{d} x &= (1 + \log \theta) x \\
 &= (\log e + \log \theta) x = \log(e \theta) x
 \end{aligned}$$



$$31. \quad = \log$$

Taking logarithm on both sides, we get

$$\log = \log \log$$

$$= (\log)^2$$

Differentiating both sides w.r.t. , we get

$$\frac{1}{d} \cdot \frac{d}{d} = 2 \log \cdot \frac{1}{d}$$

$$\Rightarrow \frac{d}{d} = \frac{2}{d} \log$$

$$\Rightarrow \frac{d}{d} = 2 \frac{\log}{d} \cdot \log = 2 \log^{-1} \cdot \log$$

$$32. \quad = 2 + \log$$

$$\therefore \frac{d}{d} = 2 + \frac{d}{d} (\log)$$

$$\Rightarrow \frac{d}{d} = 2 + \frac{2}{d} \log (\log)$$

$$33. \quad \frac{2}{3} + \frac{2}{3} = a^{\frac{2}{3}}$$

Differentiating both sides w.r.t. , we get

$$\frac{2}{3} \frac{2}{3}^{-1} + \frac{2}{3} \frac{2}{3}^{-1} \cdot \frac{d}{d} = 0$$

$$\Rightarrow \frac{2}{3} \frac{-1}{3} + \frac{2}{3} \frac{-1}{3} \cdot \frac{d}{d} = 0$$

$$\Rightarrow \frac{-1}{3} \cdot \frac{d}{d} = -\frac{-1}{3} \Rightarrow \frac{d}{d} = -\left(-\frac{1}{3}\right)$$

$$34. \quad 3^2 + 3^2 - 3a = 0$$

Differentiating w.r.t. , we get

$$3^2 + 3^2 \cdot \frac{d}{d} - 3a \left(\frac{d}{d} + \right) = 0$$

$$\Rightarrow 3(2 - a) + 3 \frac{d}{d} (2 - a) = 0$$

$$\Rightarrow \frac{d}{d} = \frac{a - 2}{2 - a}$$

$$35. \quad 3^2 + 8 + 3^2 = 64$$

Differentiating both sides w.r.t. , we get

$$3^2 + 8 \left(\frac{d}{d} + \frac{d}{d} \right) + 3^2 \frac{d}{d} = 0$$

$$\Rightarrow \frac{d}{d} = -\frac{3^2 + 8}{8 + 3^2}$$

$$36. \quad = \cos (+)$$

$$\therefore \frac{d}{d} = -\sin (+) \cdot \left(1 + \frac{d}{d} \right)$$

$$\Rightarrow \frac{d}{d} [1 + \sin (+)] = -\sin (+)$$

$$\Rightarrow \frac{d}{d} = \frac{-\sin (+)}{1 + \sin (+)}$$

$$37. \quad \sin^2 + 2 \cos + = 0$$

Differentiating w.r.t. , we get

$$2 \sin \cos - 2 \sin \frac{d}{d} + + \frac{d}{d} = 0$$

$$\Rightarrow \frac{d}{d} (-2 \sin) = - - \sin 2$$

$$\Rightarrow \frac{d}{d} = \frac{+ \sin 2}{2 \sin -}$$

$$38. \quad a^2 + 2h + b^2 + 2g + 2f + c = 0$$

Differentiating w.r.t. , we get

$$2a + 2h \left(\frac{d}{d} + \frac{d}{d} \right) + 2b \frac{d}{d} + 2g + 2f \frac{d}{d} = 0$$

$$\Rightarrow \frac{d}{d} (2h + 2b + 2f) = -(2a + 2h + 2g)$$

$$\Rightarrow \frac{d}{d} = -\frac{a + h + g}{h + b + f}$$

$$39. \quad \sqrt{ } + \sqrt{ } = 1$$

Differentiating both sides w.r.t. , we get

$$\frac{d}{d} = -\frac{\sqrt{ }}{\sqrt{ }}$$

$$\therefore \left(\frac{d}{d} \right)_{\left(\frac{1}{4} \frac{1}{4} \right)} = -1$$

$$40. \quad = a \cos \theta \text{ and } = b \sin \theta$$

$$\therefore \frac{d}{d\theta} = -a \sin \theta \text{ and } \frac{d}{d\theta} = b \cos \theta$$

$$\therefore \frac{d}{d} = \frac{\frac{d}{d\theta}}{\frac{d}{d\theta}} = \frac{b \cos \theta}{-a \sin \theta} = \left(-\frac{b}{a} \right) \cot \theta$$

$$41. \quad \text{Let } = 5 \text{ and } z = \log_5$$

$$\therefore \frac{d}{d} = 5 \log 5 \text{ and } \frac{dz}{d} = \frac{1}{\log 5}$$

$$\therefore \frac{d}{dz} = \frac{\frac{d}{d}}{\frac{dz}{d}} = \frac{5 \log 5}{\frac{1}{\log 5}} = .5 (\log 5)^2$$



$$42. \quad = \frac{1}{1-t^2} \text{ and } = 1+t^2$$

$$\therefore \frac{d}{dt} = \frac{2t}{(1-t^2)^2} \text{ and } \frac{d}{dt} = 2t$$

$$\therefore \frac{d}{d} = \frac{\frac{dt}{d}}{\frac{dt}{d}} = \frac{2t}{2t} = (1-t^2)^2$$

$$43. \text{ Let } = \sin^2 \text{ and } z = 2$$

$$\therefore \frac{d}{d} = \cos^2 \cdot (2) = 2 \cos^2$$

$$\text{and } \frac{dz}{d} = 2$$

$$\therefore \frac{d}{dz} = \frac{\frac{d}{d}}{\frac{dz}{d}} = \cos^2$$

$$44. \text{ Let } = e^3 \text{ and } z = \log$$

$$\therefore \frac{d}{d} = e^3 \cdot 3^2 = 3^2 e^3 \text{ and } \frac{dz}{d} = \frac{1}{-}$$

$$\therefore \frac{d}{dz} = \frac{\frac{d}{d}}{\frac{dz}{d}} = \frac{3^2 e^3}{\left(\frac{1}{-}\right)} = 3^3 e^3$$

$$45. \text{ Let } = a^{\sin^{-1}} \text{ and } z = \sin^{-1}$$

$$\therefore = a^z$$

$$\therefore \frac{d}{dz} = a^z \log a = a^{\sin^{-1}} \log a$$

$$46. \quad = a \sec^2 \theta \text{ and } = b \tan^2 \theta$$

$$\therefore \frac{d}{d\theta} = 2a \sec^2 \theta \cdot \tan \theta$$

$$\text{and } \frac{d}{d\theta} = 2b \tan \theta \cdot \sec^2 \theta$$

$$\therefore \frac{d}{d} = \frac{\frac{d\theta}{d}}{\frac{d\theta}{d}} = \frac{b}{a}$$

$$47. \quad = a^2 (\sin \theta + \operatorname{cosec} \theta) \quad \dots(i)$$

$$= a^2 (\sin \theta - \operatorname{cosec} \theta) \quad \dots(ii)$$

Squaring (i) and (ii) and subtracting, we get

$$^2 - ^2 = 4a^4$$

Differentiating w.r.t. , we get

$$2 - 2 \frac{d}{d} = 0 \quad \Rightarrow \frac{d}{d} = -$$

$$48. \quad = \log(a + b)$$

$$\therefore \frac{d}{d} = \frac{1}{a+b} \times a$$

$$\therefore \frac{d^2}{d^2} = \frac{-a^2}{(a+b)^2}$$

$$49. \quad = \log(\sin)$$

$$\therefore \frac{d}{d} = \frac{1}{\sin} \cdot \cos = \cot$$

$$\therefore \frac{d^2}{d^2} = -\operatorname{cosec}^2$$

$$50. \quad \sqrt{\quad} = 1$$

$$\Rightarrow = 1$$

$$\Rightarrow = \frac{1}{-}$$

$$\therefore \frac{d}{d} = \frac{-1}{2}$$

$$\therefore \frac{d^2}{d^2} = \frac{2}{3}$$

$$51. \quad = \sin m \quad \dots(i)$$

$$\therefore \frac{d}{d} = m \cos m$$

$$\therefore \frac{d^2}{d^2} = -m^2 \sin m$$

$$\Rightarrow \frac{d^2}{d^2} + m^2 = 0 \quad \dots[\text{From (i)}]$$

$$52. \quad = 2 \sin + 3 \cos$$

$$\therefore \frac{d}{d} = 2 \cos - 3 \sin$$

$$\therefore \frac{d^2}{d^2} = -2 \sin - 3 \cos$$

$$\therefore + \frac{d^2}{d^2} = 2 \sin + 3 \cos - 2 \sin - 3 \cos$$

$$\Rightarrow + \frac{d^2}{d^2} = 0$$

$$53. \quad = a \cos nt - b \sin nt \quad \dots(i)$$

$$\therefore \frac{d}{dt} = -na \sin nt - nb \cos nt$$

$$\therefore \frac{d^2}{dt^2} = -n^2 a \cos nt + n^2 b \sin nt$$

$$= -n^2 (a \cos nt - b \sin nt)$$

$$= -n^2 \quad \dots[\text{From (i)}]$$



$$54. \quad a \sin(m) + b \cos(m) \quad \dots(i)$$

$$\therefore \frac{d}{d} = am \cos(m) - bm \sin(m)$$

$$\therefore \frac{d^2}{d^2} = -am^2 \sin(m) - bm^2 \cos(m)$$

$$= -m^2 [a \sin(m) + b \cos(m)]$$

$$= -m^2 \quad \dots[\text{From (i)}]$$

$$55. \quad = a + b^2$$

$$\therefore \frac{d}{d} = 2b \quad \dots(i)$$

$$\therefore \frac{d^2}{d^2} = 2b$$

$$\Rightarrow \frac{d^2}{d^2} = 2b = \frac{d}{d} \quad \dots[\text{From (i)}]$$

$$56. \quad f(x) = be^a + ae^b$$

$$\therefore f'(x) = abe^a + abe^b$$

$$\therefore f''(x) = a^2be^a + ab^2e^b$$

$$\therefore f''(0) = a^2b + ab^2 = ab(a + b)$$



Critical Thinking

$$1. \quad f(x) = \begin{cases} -(x-3), & x < 3 \\ (x-3), & x \geq 3 \end{cases}$$

$$f'(3^-) = -1 \text{ and } f'(3^+) = 1$$

$$\therefore f'(3^-) \neq f'(3^+)$$

$$\therefore f'(3) \text{ does not exist.}$$

$$2. \quad \lim_{x \rightarrow 2^-} f(x) = \lim_{h \rightarrow 0} f(2-h) = \lim_{h \rightarrow 0} |2-h-2| = 0$$

$$\lim_{x \rightarrow 2^+} f(x) = \lim_{h \rightarrow 0} f(2+h) = \lim_{h \rightarrow 0} |2+h-2| = 0$$

$$\therefore f(2) = 0$$

$$\therefore f(x) \text{ is continuous at } x = 2.$$

$$f'(2^-) = \lim_{h \rightarrow 0^-} \frac{f(2+h) - f(2)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{-(2+h-2) - 0}{h}$$

$$= -1$$

$$f'(2^+) = \lim_{h \rightarrow 0^+} \frac{f(2+h) - f(2)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2+h-2-0}{h} = 1$$

$$\therefore f'(2^-) \neq f'(2^+)$$

$$\therefore f(x) \text{ is not differentiable at } x = 2.$$

$$3. \quad f'(1^-) = \lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{-1}$$

$$= \lim_{x \rightarrow 1^-} \frac{2^2 + 3x + 4 - 9}{-1}$$

$$= \lim_{x \rightarrow 1^-} \frac{(x-1)(2x+5)}{-1}$$

$$= \lim_{x \rightarrow 1^-} (2x+5)$$

$$= 7$$

$$f'(1^+) = \lim_{x \rightarrow 1^+} \frac{f(x) - f(1)}{-1}$$

$$= \lim_{x \rightarrow 1^+} \frac{kx + 9 - k - 9}{-1}$$

$$= \lim_{x \rightarrow 1^+} \frac{k(x-1)}{-1}$$

$$= -k$$

Since, $f(x)$ is differentiable at $x = 1$.

$$\therefore -k = 7 \Rightarrow k = -7$$

$$4. \quad f(x) = \frac{1}{1+|x|}$$

$$\therefore f(x) = \begin{cases} \frac{1}{1-x}, & x < 0 \\ \frac{1}{1+x}, & x \geq 0 \end{cases}$$

$$\therefore f'(x) = \begin{cases} \frac{1}{(1-x)^2}, & x < 0 \\ \frac{1}{(1+x)^2}, & x \geq 0 \end{cases}$$

$f(x)$ is differentiable at $(-\infty, \infty)$.

$$5. \quad \text{Applying L'Hospital rule, we get}$$

$$\lim_{x \rightarrow 2} \frac{2^2 - 4f'(x)}{-2} = \lim_{x \rightarrow 2} \frac{4 - 4f''(x)}{1}$$

$$= 8 - 4f''(2) = 8 - 4(1) = 4$$

$$6. \quad \text{Since, } f'(a) \text{ exists.}$$

$$\therefore \lim_{x \rightarrow a} \frac{f(x) - f(a)}{-a} = f'(a) \quad \dots(i)$$

$$\text{Now, } \lim_{x \rightarrow a} \frac{f(a) - af(x)}{-a}$$

$$= \lim_{x \rightarrow a} \frac{(x-a)f(a) - a(f(x) - f(a))}{-a}$$

$$= \lim_{x \rightarrow a} f(a) - a \lim_{x \rightarrow a} \left\{ \frac{f(x) - f(a)}{-a} \right\}$$

$$= f(a) - af'(a) \quad \dots[\text{From (i)}]$$

7. If a function $f(x)$ is continuous at $x = a$, then it may or may not be differentiable at $x = a$.

\therefore Option (B) is not true.



$$\begin{aligned}
 8. \quad f'(0^-) &= \lim_{h \rightarrow 0^-} \frac{f(0-h) - f(0)}{-h} \\
 &= \lim_{h \rightarrow 0^-} \frac{f(0-h) - f(0)}{-h} \\
 &= \lim_{h \rightarrow 0^-} \frac{h^2 \sin\left(\frac{1}{-h}\right) - 0}{-h} \\
 &= 0
 \end{aligned}$$

$$\begin{aligned}
 f'(0^+) &= \lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h} \\
 &= \lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h} \\
 &= \lim_{h \rightarrow 0^+} \frac{h^2 \sin\frac{1}{h} - 0}{h} \\
 &= 0
 \end{aligned}$$

$$\therefore f'(0^-) = f'(0^+)$$

$\therefore f(x)$ is derivable at $x = 0$.

$$\begin{aligned}
 9. \quad f'(0^-) &= \lim_{h \rightarrow 0^-} \frac{f(0-h) - f(0)}{-h} \\
 &= \lim_{h \rightarrow 0^-} \frac{-h - 0}{-h} \\
 &= -1
 \end{aligned}$$

$$\begin{aligned}
 f'(0^+) &= \lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h} \\
 &= \lim_{h \rightarrow 0^+} \frac{h^2 - 0}{h} \\
 &= 0
 \end{aligned}$$

$$\therefore f'(0^-) \neq f'(0^+)$$

$$\begin{aligned}
 f'(1^-) &= \lim_{h \rightarrow 1^-} \frac{f(1-h) - f(1)}{-h} \\
 &= \lim_{h \rightarrow 1^-} \frac{h^2 - 1}{-h} \\
 &= 2
 \end{aligned}$$

$$\begin{aligned}
 f'(1^+) &= \lim_{h \rightarrow 1^+} \frac{f(1+h) - f(1)}{h} \\
 &= \lim_{h \rightarrow 1^+} \frac{h^3 - 1 + 1 - 1}{h} \\
 &= \lim_{h \rightarrow 1^+} \frac{(h^2 - 1)}{h} \\
 &= 2
 \end{aligned}$$

$$\therefore f'(1^-) = f'(1^+)$$

$\therefore f(x)$ is differentiable at $x = 1$.

10. $f(x)$ is continuous at $x = 1$.

$$\begin{aligned}
 \therefore f(1) &= \lim_{x \rightarrow 1^+} f(x) \\
 &\Rightarrow a + b = b + a + c \\
 &\Rightarrow c = 0
 \end{aligned}$$

Also, $f(x)$ is differentiable at $x = 1$.

$$\begin{aligned}
 \therefore Lf'(1) &= Rf'(1) \\
 &\Rightarrow \left[\frac{d}{dx}(a^2 + b) \right]_{x=1} = \left[\frac{d}{dx}(b^2 + a + c) \right]_{x=1} \\
 &\Rightarrow [2a]_{x=1} = [2b + a]_{x=1} \\
 &\Rightarrow 2a = 2b + a \\
 &\Rightarrow a = 2b
 \end{aligned}$$

$$\begin{aligned}
 11. \quad f'(0) &= \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{h^p \cos\frac{1}{h} - 0}{h} \\
 &= \lim_{h \rightarrow 0} h^{p-1} \cos\frac{1}{h} = 0, \text{ if } p - 1 > 0
 \end{aligned}$$

i.e., if $p > 1$

$\therefore f(x)$ will be differentiable at $x = 0$, if $p > 1$

$$12. \quad f(x) = \begin{cases} e^{-x}, & x \geq 0 \\ e^x, & x < 0 \end{cases}$$

Clearly, $f(x)$ is continuous and differentiable for all non-zero x .

$$\text{Now, } \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} e^x = 1$$

$$\text{and } \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} e^{-x} = 1$$

$$\text{Also at } x = 0, f(0) = e^0 = 1$$

So, $f(x)$ is continuous for all values of x .

$$Lf'(0) = \lim_{h \rightarrow 0^-} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0^-} \frac{e^h - 1}{h} = 1$$

$$Rf'(0) = \lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0^+} \frac{e^{-h} - 1}{h} = -1$$

So, $f(x)$ is not differentiable at $x = 0$.

$\therefore f(x)$ is continuous every where but not differentiable at $x = 0$.

$$13. \quad f(x) = \begin{cases} e^{-\left(\frac{1}{x} + 1\right)}, & x < 0 \\ e^{-2/x}, & x > 0 \\ 0, & x = 0 \end{cases}$$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} 0 = 0$$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} e^{-2/x} = 0 \text{ and } f(0) = 0$$

$$\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$$



So, $f(x)$ is continuous at $x = 0$

Also,

$$Lf'(0) = 1 \text{ and}$$

$$Rf'(0) = \lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h}$$

$$= \lim_{h \rightarrow 0^+} \frac{he^{-2/h} - 0}{h} = \lim_{h \rightarrow 0^+} e^{-2/h} = 0$$

$\therefore f$ is not differentiable at $x = 0$.

Thus, $f(x)$ is everywhere continuous but not differentiable at $x = 0$.

$$14. f(x) = \begin{cases} x^2 = x^2, & x > 0 \\ 0, & x = 0 \\ -x^2 = -x^2, & x < 0 \end{cases}$$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (-x^2) = 0, \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} x^2 = 0$$

and $f(0) = 0$.

So, $f(x)$ is continuous at $x = 0$.

Also, $f(x)$ is continuous for all other values of x .

Hence, $f(x)$ is continuous everywhere.

Here, $Lf'(0) = -1$ and $Rf'(0) = 1$.

$\therefore f(x)$ is not differentiable at $x = 0$.

$$15. Lf'(0) = \lim_{x \rightarrow 0^-} \frac{f(x) - f(0)}{x - 0} = \lim_{x \rightarrow 0^-} \frac{e^x + a - b}{x}$$

$$Rf'(0) = \lim_{x \rightarrow 0^+} \frac{f(x) - f(0)}{x - 0}$$

$$= \lim_{x \rightarrow 0^+} \frac{b(x - 1)^2 - b}{x}$$

$$= \lim_{x \rightarrow 0^+} \frac{b(x^2 - 2x + 1 - 1)}{x}$$

$$= \lim_{x \rightarrow 0^+} b(x - 2) = -2b$$

Since $f'(0)$ exists.

$\therefore Lf'(0)$ must exist.

$$\therefore 1 - b = 0 \Rightarrow b = 1$$

$$Lf'(0) = Rf'(0) = -2$$

$$\text{and } Lf'(0) = \lim_{x \rightarrow 0^-} \frac{e^x + a - 1}{x}$$

$$= \lim_{x \rightarrow 0^-} \left(\frac{e^x - 1}{x} + a \right) = 1 + a$$

$$\therefore 1 + a = -2 \Rightarrow a = -3$$

$$\therefore (a, b) = (-3, 1)$$

$$16. Lf'(1) = \lim_{h \rightarrow 0} \frac{f(1-h) - f(1)}{-h}$$

$$= \lim_{h \rightarrow 0} \frac{m(1-h)^2 - m}{-h} = \lim_{h \rightarrow 0} \frac{m(1+h^2 - 2h - 1)}{-h}$$

$$= \lim_{h \rightarrow 0} m(2-h) = 2m$$

$$Rf'(1) = \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h} = \lim_{h \rightarrow 0} \frac{2(1+h) - m}{h}$$

For differentiability, $Lf'(1) = Rf'(1)$.

But for any value of m , $Lf'(1) = Rf'(1)$ is not possible.

$$17. Lf'(0) = \lim_{x \rightarrow 0^-} \frac{f(x) - f(0)}{x - 0}$$

$$= \lim_{x \rightarrow 0^-} \left(\frac{-a \sin x + be^{-x} - b}{x} \right)$$

Applying L' Hospital rule, we get

$$Lf'(0) = \lim_{x \rightarrow 0^-} \left(\frac{-a \cos x - be^{-x}}{1} \right) = -(a + b)$$

$$Rf'(0) = \lim_{x \rightarrow 0^+} \frac{f(x) - f(0)}{x - 0}$$

$$= \lim_{x \rightarrow 0^+} \left(\frac{a \sin x + be^{-x} - b}{x} \right)$$

Applying L' Hospital rule, we get

$$Rf'(0) = \lim_{x \rightarrow 0^+} \left(\frac{a \cos x + be^{-x}}{1} \right) = a + b$$

Since, $f(x)$ is differentiable at $x = 0$.

$$\therefore Lf'(0) = Rf'(0)$$

$$\Rightarrow -(a + b) = a + b$$

$$\Rightarrow a + b = 0$$

$$18. \text{ Let } y = \sqrt{\sqrt{x} + 1}$$

$$\therefore \frac{dy}{dx} = \frac{1}{2\sqrt{\sqrt{x} + 1}} \cdot \frac{d}{dx} (\sqrt{x} + 1)$$

$$= \frac{1}{4\sqrt{x} \cdot \sqrt{\sqrt{x} + 1}}$$

$$= \frac{1}{4\sqrt{x} (\sqrt{x} + 1)}$$

$$19. \text{ As } 1^\circ = \frac{\pi}{180} \text{ radian.}$$

$$\therefore \frac{d}{dx} = \frac{\pi}{180} \sec^2 \theta \tan \theta$$



$$\begin{aligned}
 20. \quad & 10^{-\tan} \left[\frac{d}{d} (10^{\tan}) \right] \\
 & = 10^{-\tan} \cdot 10^{\tan} \cdot \log 10 \cdot \frac{d}{d} (\tan) \\
 & = \log 10 (\tan + \sec^2)
 \end{aligned}$$

$$\begin{aligned}
 21. \quad & = e^{\frac{2}{1+2}} \\
 \therefore \frac{d}{d} & = e^{\frac{2}{1+2}} \cdot \frac{d}{d} \left(\frac{2}{1+2} \right) \\
 & = e^{\frac{2}{1+2}} \cdot \left[\frac{(1+2) \cdot (2) - 2 \cdot (0+2)}{(1+2)^2} \right] \\
 & = \frac{2 e^{\frac{2}{1+2}}}{(1+2)^2}
 \end{aligned}$$

$$\begin{aligned}
 22. \quad \frac{d}{d} & = \frac{d}{du} \times \frac{du}{dv} \times \frac{dv}{d} \\
 & = \frac{1}{2\sqrt{u}} \times (3-4v) \times 2 \\
 & = \frac{1}{\sqrt{(3-2v)v}} \times (3-4v) \times \\
 & = \frac{1}{\sqrt{(3-2^2)^2}} \times (3-4^2) \times \\
 & = \frac{3-4^2}{\sqrt{3-2^2}}
 \end{aligned}$$

$$\begin{aligned}
 23. \quad & = (\cos^2)^2 \\
 \therefore \frac{d}{d} & = 2 \cos^2 \cdot \frac{d}{d} (\cos^2) \\
 & = 2 \cos^2 \cdot (-\sin^2) \cdot \frac{d}{d} (2) \\
 & = 2 \cos^2 \cdot (-\sin^2) \cdot 2 \\
 & = -2 (2 \sin^2 \cos^2) = -2 \sin 2^2
 \end{aligned}$$

$$\begin{aligned}
 24. \quad & = \frac{\tan + \cot}{\tan - \cot} = \frac{\tan + \frac{1}{\tan}}{\tan - \frac{1}{\tan}} \\
 & = \frac{1 + \tan^2}{1 - \tan^2} = -\sec 2 \\
 \therefore \frac{d}{d} & = -\sec 2 \tan 2 \cdot \frac{d}{d} (2) \\
 & = -2 \sec 2 \tan 2
 \end{aligned}$$

$$\begin{aligned}
 25. \quad & = \log (\sqrt{+} + \sqrt{-a}) \\
 \therefore \frac{d}{d} & = \frac{1}{\sqrt{+} + \sqrt{-a}} \cdot \frac{d}{d} (\sqrt{+} + \sqrt{-a}) \\
 & = \frac{1}{\sqrt{+} + \sqrt{-a}} \left(\frac{1}{2\sqrt{+}} + \frac{1}{2\sqrt{-a}} \right) \\
 & = \frac{1}{2(\sqrt{+} + \sqrt{-a})} \left(\frac{1}{\sqrt{+}} + \frac{1}{\sqrt{-a}} \right) \\
 & = \frac{1}{2(\sqrt{+} + \sqrt{-a})} \left(\frac{\sqrt{-a} + \sqrt{+}}{\sqrt{+}\sqrt{-a}} \right) \\
 & = \frac{1}{2\sqrt{+}\sqrt{-a}}
 \end{aligned}$$

$$\begin{aligned}
 26. \quad \frac{d}{d} & \left[\log (\sqrt{\sin \sqrt{e}}) \right] = \frac{d}{d} \left[\frac{1}{2} \log (\sin \sqrt{e}) \right] \\
 & = \frac{1}{2} \cdot \frac{1}{\sin \sqrt{e}} \cdot \frac{d}{d} (\sin \sqrt{e}) \\
 & = \frac{1}{2} \cdot \frac{1}{\sin \sqrt{e}} \cdot \cos \sqrt{e} \cdot \frac{d}{d} (\sqrt{e}) \\
 & = \frac{1}{2} \cot \sqrt{e} \cdot \frac{1}{2\sqrt{e}} \cdot e = \frac{1}{4} e^{\frac{1}{2}} \cot (e^{\frac{1}{2}})
 \end{aligned}$$

$$\begin{aligned}
 27. \quad \frac{d}{d} & \left(\frac{e^a}{\sin(b+c)} \right) \\
 & = \frac{\sin(b+c) \cdot e^a \cdot \frac{d}{d} (a) - e^a \cdot \cos(b+c) \cdot \frac{d}{d} (b+c)}{\{\sin(b+c)\}^2} \\
 & = \frac{\sin(b+c) \cdot e^a \cdot a - e^a \cos(b+c) \cdot b}{\{\sin(b+c)\}^2} \\
 & = \frac{e^a [a \sin(b+c) - b \cos(b+c)]}{\sin^2(b+c)}
 \end{aligned}$$

$$\begin{aligned}
 28. \quad & = \sin(\sqrt{\sin + \cos}) \\
 \therefore \frac{d}{d} & = \cos(\sqrt{\sin + \cos}) \cdot \frac{d}{d} (\sqrt{\sin + \cos}) \\
 & = \cos(\sqrt{\sin + \cos}) \cdot \frac{1}{2\sqrt{\sin + \cos}} \cdot \frac{d}{d} (\sin + \cos) \\
 & = \frac{\cos(\sqrt{\sin + \cos})}{2\sqrt{\sin + \cos}} \cdot (\cos - \sin)
 \end{aligned}$$



$$\begin{aligned}
 29. \quad \frac{d}{d}(\sqrt{\sec^2 + \operatorname{cosec}^2}) &= \frac{d}{d} \left[\sqrt{\left(\frac{1}{\cos^2} + \frac{1}{\sin^2}\right)} \right] \\
 &= \frac{d}{d} \left(\sqrt{\frac{1}{\cos^2 \sin^2}} \right) = \frac{d}{d} \left(\sqrt{\frac{4}{\sin^2 2}} \right) \\
 &= \frac{d}{d} (2 \operatorname{cosec} 2) = -2 \operatorname{cosec} 2 \cot 2 \cdot \frac{d}{d}(2) \\
 &= -4 \operatorname{cosec} 2 \cot 2
 \end{aligned}$$

$$\begin{aligned}
 30. \quad &= (\cot^3)^{\frac{3}{2}} \\
 \therefore \frac{d}{d} &= \frac{3}{2} (\cot^3)^{\frac{1}{2}} \cdot \frac{d}{d}(\cot^3) \\
 &= \frac{3}{2} (\cot^3)^{\frac{1}{2}} \left[\cot^3 \cdot 1 + 3 \cot^2 \cdot \frac{d}{d}(\cot) \right] \\
 &= \frac{3}{2} (\cot^3)^{\frac{1}{2}} [\cot^3 + 3 \cot^2 (-\operatorname{cosec}^2)] \\
 &= \frac{3}{2} (\cot^3)^{\frac{1}{2}} (\cot^3 - 3 \cot^2 \operatorname{cosec}^2)
 \end{aligned}$$

$$\begin{aligned}
 31. \quad &= \sqrt{\frac{1+\tan}{1-\tan}} = \sqrt{\tan\left(\frac{\pi}{4} + \right)} \\
 \therefore \frac{d}{d} &= \frac{1}{2\sqrt{\tan\left(\frac{\pi}{4} + \right)}} \cdot \frac{d}{d} \left[\tan\left(\frac{\pi}{4} + \right) \right] \\
 &= \frac{1}{2\sqrt{1+\tan}} \cdot \sec^2\left(\frac{\pi}{4} + \right)
 \end{aligned}$$

$$\begin{aligned}
 32. \quad &= \log \left[\tan\left(\frac{\pi}{4} + \frac{1}{2}\right) \right] \\
 \therefore \frac{d}{d} &= \frac{1}{\tan\left(\frac{\pi}{4} + \frac{1}{2}\right)} \cdot \frac{d}{d} \left[\tan\left(\frac{\pi}{4} + \frac{1}{2}\right) \right] \\
 &= \frac{1}{\tan\left(\frac{\pi}{4} + \frac{1}{2}\right)} \cdot \sec^2\left(\frac{\pi}{4} + \frac{1}{2}\right) \cdot \frac{1}{2} \\
 &= \frac{1}{2 \sin\left(\frac{\pi}{4} + \frac{1}{2}\right) \cos\left(\frac{\pi}{4} + \frac{1}{2}\right)} = \frac{1}{\sin\left(\frac{\pi}{2} + \right)} \\
 &= \frac{1}{\cos} = \sec
 \end{aligned}$$

$$\begin{aligned}
 33. \quad \log \left(\sqrt{\frac{1-\cos}{1+\cos}} \right) &= \left(\log \sqrt{\frac{2 \sin^2\left(\frac{1}{2}\right)}{2 \cos^2\left(\frac{1}{2}\right)}} \right) \\
 &= \log \left(\tan \frac{1}{2} \right)
 \end{aligned}$$

$$\begin{aligned}
 \therefore \frac{d}{d} &= \frac{1}{\tan \frac{1}{2}} \cdot \frac{d}{d} \left(\tan \frac{1}{2} \right) \\
 &= \frac{1}{\tan \frac{1}{2}} \cdot \sec^2 \frac{1}{2} \cdot \frac{1}{2} \\
 &= \frac{1}{2 \sin \frac{1}{2} \cos \frac{1}{2}} \\
 &= \frac{1}{\sin} \\
 &= \operatorname{cosec}
 \end{aligned}$$

$$\begin{aligned}
 34. \quad \text{Let } &= \left[\log \left\{ e \left(\frac{-2}{+2} \right)^{\frac{3}{4}} \right\} \right] \\
 \Rightarrow &= \log e + \log \left(\frac{-2}{+2} \right)^{\frac{3}{4}} \\
 \Rightarrow &= + \frac{3}{4} [\log(-2) - \log(+2)] \\
 \therefore \frac{d}{d} &= 1 + \frac{3}{4} \left(\frac{1}{-2} - \frac{1}{+2} \right) \\
 &= 1 + \frac{3}{2-4} = \frac{2-1}{2-4}
 \end{aligned}$$

$$\begin{aligned}
 35. \quad f() &= \frac{1}{\sqrt{2+a^2} + \sqrt{2+b^2}} \\
 &= \frac{1}{\sqrt{2+a^2} + \sqrt{2+b^2}} \times \frac{\sqrt{2+a^2} - \sqrt{2+b^2}}{\sqrt{2+a^2} - \sqrt{2+b^2}} \\
 &= \frac{1}{a^2 - b^2} \left[\sqrt{2+a^2} - \sqrt{2+b^2} \right] \\
 \therefore f'() &= \frac{1}{a^2 - b^2} \left[\frac{1}{2\sqrt{2+a^2}} \cdot 2 - \frac{1}{2\sqrt{2+b^2}} \cdot 2 \right] \\
 &= \frac{1}{a^2 - b^2} \left[\frac{1}{\sqrt{2+a^2}} - \frac{1}{\sqrt{2+b^2}} \right]
 \end{aligned}$$



$$\begin{aligned}
 36. \quad &= \log\left(\sqrt{\frac{1+\sin}{1-\sin}}\right) = \frac{1}{2}\log\left(\frac{1+\sin}{1-\sin}\right) \\
 &= \frac{1}{2}\log(1+\sin) - \frac{1}{2}\log(1-\sin) \\
 \therefore \frac{d}{d} &= \frac{1}{2} \cdot \frac{1}{1+\sin} \cdot \cos - \frac{1}{2} \cdot \frac{1}{1-\sin} \cdot (-\cos) \\
 &= \frac{1}{2}\cos\left(\frac{1}{1+\sin} + \frac{1}{1-\sin}\right) \\
 &= \frac{1}{2}\cos\left(\frac{2}{1-\sin^2}\right) = \frac{2\cos}{2\cos^2} \\
 &= \frac{1}{\cos} = \sec
 \end{aligned}$$

$$\begin{aligned}
 37. \quad &= \frac{1}{2}\sqrt{a^2 + 2} + \frac{a^2}{2}\log\left(\frac{1+\sqrt{2+a^2}}{1-\sqrt{2+a^2}}\right) \\
 \therefore \frac{d}{d} &= \frac{1}{2}\left[\frac{1}{\sqrt{a^2+2}} + \frac{1}{2\sqrt{a^2+2}} \cdot 2\right] \\
 &\quad + \frac{a^2}{2} \cdot \frac{1}{1-\sqrt{2+a^2}} \left[1 + \frac{1}{2\sqrt{2+a^2}} \cdot 2\right] \\
 &= \frac{1}{2(\sqrt{a^2+2})} (a^2 + 2 + 2) \\
 &\quad + \frac{a^2}{2} \cdot \frac{1}{1-\sqrt{2+a^2}} \cdot \frac{\sqrt{2+a^2} + 1}{\sqrt{2+a^2}} \\
 &= \frac{1}{2(\sqrt{a^2+2})} (a^2 + 2 + a^2) \\
 &= \frac{2(a^2+2)}{2\sqrt{a^2+2}} = \sqrt{a^2+2}
 \end{aligned}$$

$$\begin{aligned}
 38. \quad &f(x) = \cos(\sin^2 x) \\
 \therefore f'(x) &= -\sin(\sin^2 x) \cdot \frac{d}{d}(\sin^2 x) \\
 &= -\sin(\sin^2 x) \cdot (\cos^2 x) \cdot (2x) \\
 \therefore f'\left(\sqrt{\frac{\pi}{2}}\right) &= -2\sqrt{\frac{\pi}{2}} \sin\left(\sin^2 \frac{\pi}{2}\right) \cos \frac{\pi}{2} \\
 &= 0 \quad \dots \left[\because \cos \frac{\pi}{2} = 0 \right]
 \end{aligned}$$

$$\begin{aligned}
 39. \quad \text{Let } &= \frac{d}{d}[\log f(e+2)] \\
 &= \frac{1}{f(e+2)} \cdot \frac{d}{d}[f(e+2)] \\
 &= \frac{1}{f(e+2)} \cdot f'(e+2) \cdot \frac{d}{d}(e+2) \\
 &= \frac{f'(e+2)(e+2)}{f(e+2)}
 \end{aligned}$$

$$\therefore ()_{(=0)} = \frac{f'(1) \cdot 3}{f(1)} = \frac{2 \cdot 3}{3} = 2$$

$$\begin{aligned}
 40. \quad &= \sin^{-1}\left(\frac{19}{20}\right) + \cos^{-1}\left(\frac{19}{20}\right) \\
 &= \frac{\pi}{2} \quad \dots \left[\because \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2} \right]
 \end{aligned}$$

$$\therefore \frac{d}{d} = 0$$

$$\begin{aligned}
 41. \quad &= \sec^{-1}\left(\frac{-1}{-1}\right) + \sin^{-1}\left(\frac{-1}{+1}\right) \\
 &= \cos^{-1}\left(\frac{-1}{+1}\right) + \sin^{-1}\left(\frac{-1}{+1}\right) \\
 \therefore &= \frac{\pi}{2} \quad \dots \left[\because \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2} \right]
 \end{aligned}$$

$$\therefore \frac{d}{d} = 0$$

$$42. \quad = \cos^{-1}\left(\frac{\sqrt{-1}}{\sqrt{+1}}\right) + \sin^{-1}\left(\frac{\sqrt{-1}}{\sqrt{+1}}\right) = \pi/2$$

$$\therefore \frac{d}{d} = 0$$

$$\begin{aligned}
 43. \quad &\sin^{-1} x + \sin^{-1} \sqrt{1-x^2} \\
 &= \sin^{-1} x + \cos^{-1} x \quad \dots \left[\because \cos^{-1} x = \sin^{-1} \sqrt{1-x^2} \right] \\
 &= \frac{\pi}{2}
 \end{aligned}$$

$$\therefore \frac{d}{d} (\sin^{-1} x + \sin^{-1} \sqrt{1-x^2}) = \frac{d}{d} \left(\frac{\pi}{2}\right) = 0$$

$$\begin{aligned}
 44. \quad \text{Let } &= \tan^{-1}(\cot x) + \cot^{-1}(\tan x) \\
 &= \tan^{-1}\left[\tan\left(\frac{\pi}{2} - x\right)\right] + \cot^{-1}\left[\cot\left(\frac{\pi}{2} - x\right)\right] \\
 &= \pi - 2x
 \end{aligned}$$

$$\therefore \frac{d}{d} = -2$$



$$\begin{aligned}
 45. \quad & \frac{d}{d} \{ \sin (2 \cos^{-1} (\sin \frac{\pi}{2})) \} \\
 &= \frac{d}{d} \left\{ \sin \left(2 \cos^{-1} \left(\cos \left(\frac{\pi}{2} - \frac{\pi}{2} \right) \right) \right) \right\} \\
 &= \frac{d}{d} \left\{ \sin \left(2 \left(\frac{\pi}{2} - \frac{\pi}{2} \right) \right) \right\} \\
 &= \frac{d}{d} \{ \sin (\pi - 2) \} \\
 &= -2 \cdot \cos (\pi - 2) \\
 &= 2 \cos 2
 \end{aligned}$$

$$\begin{aligned}
 46. \quad &= \tan^{-1} \left(\frac{\frac{1}{3} + a^{\frac{1}{3}}}{1 - \frac{1}{3} a^{\frac{1}{3}}} \right) = \tan^{-1} \left(\frac{1}{3} \right) + \tan^{-1} \left(a^{\frac{1}{3}} \right) \\
 \therefore \quad & \frac{d}{d} = \frac{1}{1 + \left(\frac{1}{3} \right)^2} \cdot \frac{1}{3} \cdot \frac{-2}{3} = \frac{1}{3^{\frac{2}{3}} \left(1 + \frac{2}{3} \right)}
 \end{aligned}$$

$$\begin{aligned}
 47. \quad &= \tan^{-1} \left(\frac{\frac{6}{5} + \tan \frac{\pi}{5}}{1 - \frac{6}{5} \tan \frac{\pi}{5}} \right) \\
 &= \tan^{-1} \left(\frac{6}{5} \right) + \tan^{-1} (\tan \frac{\pi}{5}) \\
 \therefore \quad &= \tan^{-1} \left(\frac{6}{5} \right) + \\
 \therefore \quad & \frac{d}{d} = 0 + 1 = 1
 \end{aligned}$$

$$\begin{aligned}
 48. \quad &= \tan^{-1} \left(\frac{5 - \frac{2}{3}}{1 + 5 \cdot \frac{2}{3}} \right) + \tan^{-1} \left(\frac{\frac{2}{3} + \frac{2}{3}}{1 - \frac{2}{3} \cdot \frac{2}{3}} \right) \\
 &= \tan^{-1} 5 - \tan^{-1} \frac{2}{3} + \tan^{-1} \frac{2}{3} + \tan^{-1} \frac{2}{3} \\
 &= \tan^{-1} 5 + \tan^{-1} \frac{2}{3} \\
 \therefore \quad & \frac{d}{d} = \frac{1}{1 + (5)^2} \cdot 5 = \frac{5}{1 + 25^2}
 \end{aligned}$$

$$\begin{aligned}
 49. \quad & \frac{d}{d} \left[\tan^{-1} \left(\frac{\cos \frac{\pi}{2}}{1 + \sin \frac{\pi}{2}} \right) \right] \\
 &= \frac{d}{d} \left[\tan^{-1} \left(\frac{\cos^2 \frac{\pi}{2} - \sin^2 \frac{\pi}{2}}{\cos^2 \frac{\pi}{2} + \sin^2 \frac{\pi}{2} + 2 \sin \frac{\pi}{2} \cos \frac{\pi}{2}} \right) \right]
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{d}{d} \left[\tan^{-1} \left(\frac{\cos^2 \frac{\pi}{2} - \sin^2 \frac{\pi}{2}}{\left(\cos \frac{\pi}{2} + \sin \frac{\pi}{2} \right)^2} \right) \right] \\
 &= \frac{d}{d} \left[\tan^{-1} \left(\frac{\cos \frac{\pi}{2} - \sin \frac{\pi}{2}}{\cos \frac{\pi}{2} + \sin \frac{\pi}{2}} \right) \right] \\
 &= \frac{d}{d} \left[\tan^{-1} \left(\frac{1 - \tan \frac{\pi}{2}}{1 + \tan \frac{\pi}{2}} \right) \right] \\
 &= \frac{d}{d} \left[\tan^{-1} \tan \left(\frac{\pi}{4} - \frac{\pi}{2} \right) \right] \\
 &= \frac{d}{d} \left(\frac{\pi}{4} - \frac{\pi}{2} \right) = -\frac{1}{2}
 \end{aligned}$$

Alternate Method:

$$\begin{aligned}
 \text{Let } &= \tan^{-1} \left(\frac{\cos \frac{\pi}{2}}{1 + \sin \frac{\pi}{2}} \right) \\
 &= \tan^{-1} \left(\frac{\sin \left(\frac{\pi}{2} - \frac{\pi}{2} \right)}{1 + \cos \left(\frac{\pi}{2} - \frac{\pi}{2} \right)} \right) \\
 &= \tan^{-1} \left(\frac{2 \sin \left(\frac{\pi}{4} - \frac{\pi}{2} \right) \cos \left(\frac{\pi}{4} - \frac{\pi}{2} \right)}{2 \cos^2 \left(\frac{\pi}{4} - \frac{\pi}{2} \right)} \right) \\
 &= \tan^{-1} \left(\tan \left(\frac{\pi}{4} - \frac{\pi}{2} \right) \right) = \frac{\pi}{4} - \frac{\pi}{2}
 \end{aligned}$$

$$\therefore \quad \frac{d}{d} = \frac{-1}{2}$$

$$\begin{aligned}
 50. \quad &= \tan^{-1} (\sec \frac{\pi}{2} - \tan \frac{\pi}{2}) \\
 \therefore \quad & \frac{d}{d} = \frac{d}{d} \left[\tan^{-1} \left(\frac{1 - \sin \frac{\pi}{2}}{\cos \frac{\pi}{2}} \right) \right]
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{d}{d} \left[\tan^{-1} \left(\frac{\cos \frac{\pi}{2} - \sin \frac{\pi}{2}}{\cos \frac{\pi}{2} + \sin \frac{\pi}{2}} \right) \right] \\
 &= \frac{d}{d} \left[\tan^{-1} \left(\frac{1 - \tan \frac{\pi}{2}}{1 + \tan \frac{\pi}{2}} \right) \right]
 \end{aligned}$$



$$= \frac{d}{d} \left[\tan^{-1} \left(\tan \left(\frac{\pi}{4} - \frac{1}{2} \right) \right) \right]$$

$$= \frac{d}{d} \left(\frac{\pi}{4} - \frac{1}{2} \right) = -\frac{1}{2}$$

$$51. \quad \frac{d}{d} \left(\tan^{-1} \sqrt{\frac{1+\cos}{1-\cos}} \right)$$

$$= \frac{d}{d} \left(\tan^{-1} \sqrt{\frac{2\cos^2 \frac{1}{2}}{2\sin^2 \frac{1}{2}}} \right)$$

$$= \frac{d}{d} \left(\tan^{-1} \left(\cot \frac{1}{2} \right) \right)$$

$$= \frac{d}{d} \left(\tan^{-1} \left(\tan \left(\frac{\pi}{2} - \frac{1}{2} \right) \right) \right)$$

$$= \frac{d}{d} \left(\frac{\pi}{2} - \frac{1}{2} \right)$$

$$= -\frac{1}{2}$$

$$52. \quad = \cot^{-1} \left[\frac{\sqrt{1+\sin} + \sqrt{1-\sin}}{\sqrt{1+\sin} - \sqrt{1-\sin}} \right]$$

By rationalizing the denominator, we get

$$= \cot^{-1} \left[\frac{2+2\cos}{2\sin} \right]$$

$$= \cot^{-1} \left[\frac{1+\cos}{\sin} \right]$$

$$= \cot^{-1} \left[\cot \frac{1}{2} \right] = \frac{1}{2}$$

$$\therefore \frac{d}{d} = \frac{1}{2}$$

$$53. \quad \text{Put } \cos \alpha = \frac{5}{\sqrt{41}}, \sin \alpha = \frac{4}{\sqrt{41}}$$

$$\therefore = \sin^{-1} [\sin(\frac{1}{2} + \alpha)] = \frac{1}{2} + \alpha$$

$$\therefore \frac{d}{d} = 1$$

$$54. \quad \text{Put } = \tan \theta \Rightarrow \theta = \tan^{-1}$$

$$\therefore = \tan^{-1} \left(\frac{1+\tan \theta}{1-\tan \theta} \right)$$

$$= \tan^{-1} \left[\tan \left(\frac{\pi}{4} + \theta \right) \right] = \frac{\pi}{4} + \theta$$

$$= \frac{\pi}{4} + \tan^{-1}$$

$$\therefore \frac{d}{d} = \frac{1}{1+^2}$$

$$55. \quad \text{Let } = \tan^{-1} \left(\frac{2}{1-^2} \right) = \tan^{-1} \left(\frac{2}{1-^2} \right)$$

$$\text{Put } = \tan \theta \Rightarrow \theta = \tan^{-1}$$

$$\therefore = \tan^{-1} \left(\frac{2 \tan \theta}{1-\tan^2 \theta} \right)$$

$$= \tan^{-1}(\tan 2\theta) = 2\theta = 2 \tan^{-1}$$

$$\therefore \frac{d}{d} = \frac{2}{1+^2}$$

$$56. \quad \text{Let } = \cos^{-1} \left(\frac{-^2-1}{+^2-1} \right) = \cos^{-1} \left(\frac{^2-1}{^2+1} \right)$$

$$\text{Put } = \cot \theta \Rightarrow \theta = \cot^{-1}$$

$$\therefore = \cos^{-1} \left(\frac{\cot^2 \theta - 1}{\cot^2 \theta + 1} \right) = \cos^{-1} \left(\frac{1-\tan^2 \theta}{1+\tan^2 \theta} \right)$$

$$= \cos^{-1}(\cos 2\theta) = 2\theta = 2 \cot^{-1}$$

$$\therefore \frac{d}{d} = \frac{-2}{1+^2}$$

$$57. \quad \text{Let } = \tan^{-1} \left(\frac{1}{\sqrt{a^2-^2}} \right)$$

$$\text{Put } = a \sin \theta \Rightarrow \theta = \sin^{-1} \left(\frac{1}{a} \right)$$

$$\therefore = \tan^{-1} \left(\frac{a \sin \theta}{\sqrt{a^2-a^2 \sin^2 \theta}} \right)$$

$$= \tan^{-1} \left(\frac{a \sin \theta}{a \cos \theta} \right) = \tan^{-1}(\tan \theta) = \theta$$

$$= \sin^{-1} \left(\frac{1}{a} \right)$$

$$\therefore \frac{d}{d} = \frac{1}{a} \cdot \frac{1}{\sqrt{1-\left(\frac{1}{a}\right)^2}} = \frac{1}{\sqrt{a^2-^2}}$$



$$\begin{aligned}
 58. \quad \text{Put } &= \tan \theta \Rightarrow \theta = \tan^{-1} \\
 \therefore &= \sin^{-1} \left(\frac{2 \tan \theta}{1 + \tan^2 \theta} \right) + \sec^{-1} \left(\frac{1 + \tan^2 \theta}{1 - \tan^2 \theta} \right) \\
 &= \sin^{-1}(\sin 2\theta) + \sec^{-1}(\sec 2\theta) \\
 &= 2\theta + 2\theta = 4\theta = 4 \tan^{-1} \\
 \therefore \frac{d}{d} &= \frac{4}{1 + ^2}
 \end{aligned}$$

$$\begin{aligned}
 59. \quad \text{Put } &= \cos 2\theta \Rightarrow \theta = \frac{1}{2} \cos^{-1} \\
 \therefore \sin^{-1} \left(\sqrt{\frac{1 - }{2}} \right) &= \sin^{-1} \left(\sqrt{\frac{1 - \cos 2\theta}{2}} \right) \\
 &= \sin^{-1} \left(\sqrt{\sin^2 \theta} \right) = \theta \\
 &= \frac{1}{2} \cos^{-1} \\
 \therefore \frac{d}{d} \left[\sin^{-1} \left(\sqrt{\frac{1 - }{2}} \right) \right] &= \frac{d}{d} \left(\frac{1}{2} \cos^{-1} \right) \\
 &= \frac{-1}{2\sqrt{1 - ^2}}
 \end{aligned}$$

$$\begin{aligned}
 60. \quad \text{Put } e^2 &= \cot \theta \Rightarrow \theta = \cot^{-1}(e^2) \\
 \therefore &= \tan^{-1} \left(\frac{\cot \theta + 1}{\cot \theta - 1} \right) = \tan^{-1} \left(\frac{1 + \tan \theta}{1 - \tan \theta} \right) \\
 &= \tan^{-1} \left[\tan \left(\frac{\pi}{4} + \theta \right) \right] \\
 &= \frac{\pi}{4} + \theta = \frac{\pi}{4} + \cot^{-1}(e^2) \\
 \therefore \frac{d}{d} &= 0 - \frac{1}{1 + (e^2)^2} \cdot e^2 \cdot 2 \\
 \therefore \frac{d}{d} &= -\frac{2e^2}{1 + e^4}
 \end{aligned}$$

$$\begin{aligned}
 61. \quad \text{Let } &= \sin^{-1} \left(\frac{\sqrt{1 + } - \sqrt{1 - }}{2} \right) \\
 \text{Put } &= \cos 2\theta \Rightarrow \theta = \frac{1}{2} \cos^{-1} \\
 \therefore &= \sin^{-1} \left[\frac{\sqrt{2}}{2} \cos \theta - \frac{\sqrt{2}}{2} \sin \theta \right] \\
 &= \sin^{-1} \left[\frac{1}{\sqrt{2}} \cos \theta - \frac{1}{\sqrt{2}} \sin \theta \right]
 \end{aligned}$$

$$\begin{aligned}
 &= \sin^{-1} \left(\sin \frac{\pi}{4} \cos \theta - \cos \frac{\pi}{4} \sin \theta \right) \\
 &= \sin^{-1} \left(\sin \left(\frac{\pi}{4} - \theta \right) \right) = \frac{\pi}{4} - \frac{1}{2} \cos^{-1} \\
 \therefore \frac{d}{d} &= \frac{1}{2\sqrt{1 - ^2}}
 \end{aligned}$$

$$\begin{aligned}
 62. \quad \text{Let } &= \tan^{-1} \left(\frac{\sqrt{1 + } - \sqrt{1 - }}{\sqrt{1 + } + \sqrt{1 - }} \right) \\
 \text{Put } &= \cos 2\theta \Rightarrow \theta = \frac{1}{2} \cos^{-1} \\
 \therefore &= \tan^{-1} \left(\frac{\sqrt{1 + \cos 2\theta} - \sqrt{1 - \cos 2\theta}}{\sqrt{1 + \cos 2\theta} + \sqrt{1 - \cos 2\theta}} \right) \\
 &= \tan^{-1} \left(\frac{\sqrt{2 \cos^2 \theta} - \sqrt{2 \sin^2 \theta}}{\sqrt{2 \cos^2 \theta} + \sqrt{2 \sin^2 \theta}} \right) \\
 &= \tan^{-1} \left(\frac{\cos \theta - \sin \theta}{\cos \theta + \sin \theta} \right) = \tan^{-1} \left(\frac{1 - \tan \theta}{1 + \tan \theta} \right) \\
 &= \tan^{-1} \left(\tan \left(\frac{\pi}{4} - \theta \right) \right) \\
 \therefore &= \frac{\pi}{4} - \theta = \frac{\pi}{4} - \frac{1}{2} \cos^{-1} \\
 \therefore \frac{d}{d} &= \frac{1}{2} \left(\frac{1}{\sqrt{1 - ^2}} \right)
 \end{aligned}$$

$$\begin{aligned}
 63. \quad \text{Let } &= \sin^2 \left\{ \cot^{-1} \left(\sqrt{\frac{1 - }{1 + }} \right) \right\} \\
 \text{Put } &= \cos \theta \\
 \therefore &= \sin^2 \left\{ \cot^{-1} \left(\sqrt{\frac{1 - \cos \theta}{1 + \cos \theta}} \right) \right\} \\
 &= \sin^2 \left\{ \cot^{-1} \left(\sqrt{\frac{2 \sin^2 \frac{\theta}{2}}{2 \cos^2 \frac{\theta}{2}}} \right) \right\} \\
 &= \sin^2 \left\{ \cot^{-1} \left(\tan \frac{\theta}{2} \right) \right\} \\
 &= \sin^2 \left\{ \cot^{-1} \left(\cot \left(\frac{\pi}{2} - \frac{\theta}{2} \right) \right) \right\} \\
 &= \sin^2 \left(\frac{\pi}{2} - \frac{\theta}{2} \right) \\
 \therefore &= \cos^2 \frac{\theta}{2} = \frac{1 + \cos \theta}{2} = \frac{1 + }{2} \\
 \therefore \frac{d}{d} &= \frac{1}{2}
 \end{aligned}$$



64. $f(x) = \cot^{-1}(\cos 2x)^{1/2}$
 $\therefore f(x) = \cot^{-1}(\sqrt{\cos 2x})$
 $\therefore f'(x) = \frac{-1}{1 + \cos 2x} \left[\frac{-2 \sin 2x}{2\sqrt{\cos 2x}} \right]$
 $= \frac{\sin 2x}{(1 + \cos 2x)\sqrt{\cos 2x}}$
 $\therefore f'\left(\frac{\pi}{6}\right) = \frac{\frac{\sqrt{3}}{2}}{\left(1 + \frac{1}{2}\right)\left(\sqrt{\frac{1}{2}}\right)} = \sqrt{\frac{2}{3}}$
65. Since, $1 + \sin \theta = \left(\sin \frac{\theta}{2} + \cos \frac{\theta}{2}\right)^2$
and $1 - \sin \theta = \left(\sin \frac{\theta}{2} - \cos \frac{\theta}{2}\right)^2$
 $\therefore f(x) = \tan^{-1} \left(\frac{\cos \frac{x}{2} + \sin \frac{x}{2}}{\cos \frac{x}{2} - \sin \frac{x}{2}} \right) = \tan^{-1} \left(\frac{1 + \tan \frac{x}{2}}{1 - \tan \frac{x}{2}} \right)$
 $= \tan^{-1} \left[\tan \left(\frac{\pi}{4} + \frac{x}{2} \right) \right]$
 $\therefore f(x) = \frac{\pi}{4} + \frac{x}{2} \quad \therefore f'(x) = \frac{1}{2}$
 $\therefore f'\left(\frac{\pi}{6}\right) = \frac{1}{2}$
66. Put $\log x = \tan \theta \Rightarrow \theta = \tan^{-1}(\log x)$
 $\therefore f(x) = \cos^{-1} \left(\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right)$
 $= \cos^{-1}(\cos 2\theta)$
 $= 2\theta = 2 \tan^{-1}(\log x)$
 $\therefore f'(x) = 2 \cdot \frac{1}{1 + (\log x)^2} \cdot \frac{1}{x}$
 $\therefore f'(e) = \frac{2}{1 + (\log e)^2} \cdot \frac{1}{e} = \frac{2}{1 + 1^2} \cdot \frac{1}{e} = \frac{1}{e}$
67. $y = (x^2 + 2 \log x)^2$
Taking logarithm on both sides, we get
 $\log y = \log (x^2 + 2 \log x)^2$
Differentiating both sides w.r.t. x , we get
 $\frac{1}{y} \cdot \frac{dy}{dx} = 2 \cdot \frac{1}{x^2 + 2 \log x} \cdot (2x + 2 \cdot \frac{1}{x})$
 $\Rightarrow \frac{dy}{dx} = (x^2 + 2 \log x)^2 \cdot \frac{2(2x + \frac{2}{x})}{x^2 + 2 \log x}$

68. $y = x^2 + 2 \log x$
Taking logarithm on both sides, we get
 $\log y = \log (x^2 + 2 \log x)$
Differentiating both sides w.r.t. x , we get
 $\frac{1}{y} \cdot \frac{dy}{dx} = 2 \cdot \frac{1}{x} + \frac{2}{x^2 + 2 \log x} \cdot \frac{1}{x}$ (2)
 $\therefore \frac{dy}{dx} = (x^2 + 2 \log x) \left(\frac{2}{x} + \frac{2}{x(x^2 + 2 \log x)} \right)$
 $= (x^2 + 2 \log x) \cdot \frac{2(x^2 + 2 \log x) + 2}{x(x^2 + 2 \log x)}$
69. Let $y = 4^{3x^2}$
Taking logarithm on both sides, we get
 $\log y = 3x^2 \cdot \log 4$
Differentiating both sides w.r.t. x , we get
 $\frac{1}{y} \cdot \frac{dy}{dx} = 4^{3x^2} + 12x \log 4$
 $\therefore \frac{dy}{dx} = 4^{3x^2} \cdot 4^{3x^2} (1 + 3 \log 4)$
 $= 4^{4^{3x^2} + 1} (1 + 3 \log 4)$
70. $y = \sqrt{\frac{1+x}{1-x}}$
Taking logarithm on both sides, we get
 $\log y = \frac{1}{2} \log(1+x) - \frac{1}{2} \log(1-x)$
Differentiating w.r.t. x , we get
 $\frac{1}{y} \cdot \frac{dy}{dx} = \frac{1}{2} \cdot \frac{1}{1+x} - \frac{1}{2} \cdot \frac{1}{1-x} \cdot (-1)$
 $\Rightarrow \frac{dy}{dx} = \sqrt{\frac{1+x}{1-x}} \cdot \frac{1}{(1+x)(1-x)}$
 $= \frac{1}{(1+x)^{\frac{1}{2}}(1-x)^{\frac{3}{2}}}$
71. $y = \frac{2(-\sin x)^{\frac{3}{2}}}{\sqrt{x}}$
Taking logarithm on both sides, we get
 $\log y = \log 2 + \frac{3}{2} \log(-\sin x) - \frac{1}{2} \log x$
Differentiating w.r.t. x , we get
 $\frac{1}{y} \cdot \frac{dy}{dx} = 0 + \frac{3}{2} \cdot \frac{1}{-\sin x} \cdot (-\cos x) - \frac{1}{2} \cdot \frac{1}{x}$
 $\Rightarrow \frac{dy}{dx} = \frac{2(-\sin x)^{\frac{3}{2}}}{\sqrt{x}} \left(\frac{3}{2} \cdot \frac{1 - \cos x}{-\sin x} - \frac{1}{2x} \right)$



$$72. \quad = \frac{e \log}{2}$$

Taking logarithm on both sides, we get

$$\log = + \log (\log) - 2 \log$$

Differentiating w.r.t. , we get

$$\frac{1}{d} \cdot \frac{d}{d} = 1 + \frac{1}{\log} - \frac{2}{d}$$

$$\Rightarrow \frac{d}{d} = \frac{e \log}{2} \left(\frac{\log + 1 - 2 \log}{\log} \right)$$

$$= \frac{e [(-2) \log + 1]}{3}$$

$$73. \quad \text{Let } = (\sin)^{\log}$$

Taking logarithm on both sides, we get

$$\log = \log \log (\sin)$$

Differentiating both sides w.r.t. , we get

$$\frac{1}{d} \cdot \frac{d}{d} = \log \cdot \frac{1}{\sin} \cdot \cos + \log (\sin) \cdot \frac{1}{d}$$

$$\Rightarrow \frac{d}{d} = (\sin)^{\log} \left(\frac{1}{\sin} \log \sin + \cot \log \right)$$

$$74. \quad = (\tan)^{\sin}$$

Taking logarithm on both sides, we get

$$\log = \sin \log (\tan)$$

Differentiating both sides w.r.t. , we get

$$\frac{1}{d} \cdot \frac{d}{d} = \sin \cdot \frac{1}{\tan} \cdot \sec^2 + \log (\tan) \cdot \cos$$

$$\Rightarrow \frac{1}{d} \cdot \frac{d}{d} = \sin \cdot \frac{\cos}{\sin} \cdot \frac{1}{\cos^2} + \cos \log (\tan)$$

$$\Rightarrow \frac{d}{d} = (\tan)^{\sin} [\sec + \cos \log (\tan)]$$

$$75. \quad 2e + 2e + 13 = 0$$

Differentiating w.r.t. , we get

$$2e + 2e \frac{d}{d} + 2 \left(e + e \frac{d}{d} + e \right) = 0$$

$$\Rightarrow \frac{d}{d} = -\frac{2e + 2(e + e)}{(e + 2e)}$$

$$\Rightarrow \frac{d}{d} = -\frac{2e^{-} + 2(+1)}{(e^{-} + 2)}$$

$$76. \quad \sec \left(\frac{+}{-} \right) = a \Rightarrow \frac{+}{-} = \sec^{-1} a$$

Differentiating both sides w.r.t. , we get

$$\frac{(-)}{(-)} \left(1 + \frac{d}{d} \right) - \left(\frac{+}{+} \right) \left(1 - \frac{d}{d} \right) = 0$$

$$\Rightarrow (-) + (-) + (-) + (-) \frac{d}{d} = 0$$

$$\Rightarrow 2 \frac{d}{d} = 2$$

$$\Rightarrow \frac{d}{d} = 1$$

$$77. \quad \cos(+) = \sin$$

Differentiating both sides w.r.t. , we get

$$-\sin(+) \cdot \left(1 + \frac{d}{d} \right) = \cos + \sin \frac{d}{d}$$

$$\Rightarrow \frac{d}{d} = -\frac{\cos + \sin(+)}{\sin(+)+\sin}$$

$$78. \quad \sin(+) + \cos(+)=1$$

Differentiating both sides w.r.t. , we get

$$\cos(+). \left(1 + \frac{d}{d} \right) - \sin(+). \left(1 + \frac{d}{d} \right) = 0$$

$$\Rightarrow \frac{d}{d} [\cos(+)-\sin(+)]$$

$$= -\cos(+)+\sin(+)$$

$$\Rightarrow \frac{d}{d} = \frac{\sin(+)-\cos(+)}{\cos(+)-\sin(+)}$$

$$\Rightarrow \frac{d}{d} = -1$$

$$79. \quad \sin(+)=\log(+)$$

Differentiating both sides w.r.t. , we get

$$\cos(+)\left[1+\frac{d}{d}\right]=\frac{1}{+}\left(1+\frac{d}{d}\right)$$

$$\Rightarrow \cos(+)\frac{d}{d}-\frac{1}{+}\frac{d}{d}=\frac{1}{+}-\cos(+)$$

$$\Rightarrow -\frac{d}{d}\left[\frac{1}{+}-\cos(+)\right]=\frac{1}{+}-\cos(+)$$

$$\Rightarrow \frac{d}{d} = -1$$

$$80. \quad 3\sin(+)+4\cos(+)=5$$

Differentiating both sides w.r.t. , we get

$$3\cos(+)\left[+\frac{d}{d}\right]-4\sin(+)\left[+\frac{d}{d}\right]=0$$

$$\Rightarrow \frac{d}{d} = \frac{4\sin(+)-3\cos(+)}{3\cos(+)-4\sin(+)}$$

$$= \frac{[4\sin(+)-3\cos(+)]}{- [4\sin(+)-3\cos(+)]} = -1$$



$$81. \quad = \sqrt{1-x^2}$$

Differentiating both sides w.r.t. x , we get

$$1 = \frac{d}{dx} \sqrt{1-x^2} + \frac{1}{2\sqrt{1-x^2}} \cdot (-2x) \cdot \frac{d}{dx} x$$

$$\Rightarrow 1 = \frac{d}{dx} \sqrt{1-x^2} - \frac{x}{\sqrt{1-x^2}} \cdot \frac{d}{dx} x$$

$$\Rightarrow 1 = \frac{d}{dx} \left(\frac{1-x^2-x^2}{\sqrt{1-x^2}} \right)$$

$$\Rightarrow 1 = \frac{d}{dx} \left(\frac{1-2x^2}{\sqrt{1-x^2}} \right)$$

$$\Rightarrow \frac{d}{dx} = \frac{\sqrt{1-x^2}}{1-2x^2}$$

$$82. \quad \sqrt{1+x} + \sqrt{1+x} = 0$$

$$\Rightarrow (1+x)^2 = (1+x)^2$$

$$\Rightarrow (1-x)(1+x) = 0$$

$$\Rightarrow 1+x = 0 \quad \dots \text{(i) } [\because x \neq -1]$$

Differentiating w.r.t. x , we get

$$1 + \frac{d}{dx} + \frac{d}{dx} + 1 = 0$$

$$\Rightarrow (1+x) \frac{d}{dx} = -(1+x)$$

$$\Rightarrow (1+x)^2 \frac{d}{dx} = -(1+x)(1+x)$$

$$\Rightarrow \frac{d}{dx} = \frac{-(1+x)(1+x)}{(1+x)^2}$$

$$\Rightarrow \frac{d}{dx} = \frac{-1}{(1+x)^2} \quad \dots \text{[From (i)]}$$

$$83. \quad \text{If } y = \sqrt{f(x)} + \sqrt{f(x)} + \sqrt{f(x)} + \dots \infty,$$

then $\frac{d}{dx} = \frac{f'(x)}{2-1} \quad \therefore \frac{d}{dx} = \frac{1}{(2-1)}$

$$84. \quad \text{If } y = \sqrt{f(x)} + \dots, \text{ then}$$

$$\frac{d}{dx} = \frac{f'(x)}{2-1} \quad \therefore \frac{d}{dx} = \frac{\cos}{2-1}$$

$$85. \quad \text{If } y = \sqrt{f(x)} + \sqrt{f(x)} + \sqrt{f(x)} + \dots \infty, \text{ then}$$

$$\frac{d}{dx} = \frac{f'(x)}{2-1}$$

$$\therefore \frac{d}{dx} = \frac{-\sin}{2-1} = \frac{\sin}{1-2}$$

$$86. \quad y = e^{+e^{+e^{+\dots \infty}}}$$

$$\Rightarrow y = e^y$$

Taking logarithm on both sides, we get

$$\log y = (y) \log e$$

Differentiating both sides w.r.t. x , we get

$$\frac{1}{y} \cdot \frac{d}{dx} y = 1 + \frac{d}{dx} y \Rightarrow \frac{d}{dx} y = \frac{y}{1-y}$$

$$87. \quad y = \sqrt{x} + \sqrt{x} + \dots$$

$$\Rightarrow (2-x) = \sqrt{2}$$

$$\Rightarrow (2-x)^2 = 2$$

Differentiating both sides w.r.t. x , we get

$$2(2-x) \left(2 \frac{d}{dx} - 1 \right) = 2 \frac{d}{dx}$$

$$\therefore \frac{d}{dx} = \frac{(2-x)}{2^3 - 2 - 1}$$

$$88. \quad \text{If } y = f(x)^{f(x)^{f(x)^{\dots \infty}}}, \text{ then}$$

$$\frac{d}{dx} = \frac{2f'(x)}{f(x)[1 - \log f(x)]}$$

$$\therefore \frac{d}{dx} = \frac{2}{(1 - \log y)}$$

$$\Rightarrow (1 - \log y) \frac{d}{dx} = 2$$

$$89. \quad \text{If } y = f(x)^{f(x)^{f(x)^{\dots \infty}}}, \text{ then}$$

$$\frac{d}{dx} = \frac{2f'(x)}{f(x)[1 - \log f(x)]}$$

$$\therefore \frac{d}{dx} = \frac{2 \cdot \frac{1}{2\sqrt{y}}}{\sqrt{y}(1 - \log \sqrt{y})} = \frac{2}{2 \left(1 - \frac{1}{2} \log y \right)}$$

$$\Rightarrow (2 - \log y) \frac{d}{dx} = 2$$

$$90. \quad \text{If } y = f(x)^{f(x)^{f(x)^{\dots \infty}}}, \text{ then}$$

$$\frac{d}{dx} = \frac{2f'(x)}{f(x)[1 - \log f(x)]}$$

$$\therefore \frac{d}{dx} = \frac{2 \cos}{\sin(1 - \log \sin)}$$

$$= \frac{2 \cot}{1 - \log \sin}$$



91. $y = 2 + \frac{1}{x}$

If $y = f(x) + \frac{1}{x}$, then $\frac{dy}{dx} = \frac{f'(x)}{2 - f(x)}$

$\therefore \frac{dy}{dx} = \frac{2}{2 - 2}$

92. $y = e^x$

Taking logarithm on both sides, we get

$\log y = \log e^x + \log e$

$\Rightarrow \log y = \log e + x$

Differentiating both sides w.r.t. x we get

$\frac{1}{y} \cdot \frac{dy}{dx} = \frac{1}{e} + \frac{dx}{dx}$

$\Rightarrow \frac{dy}{dx} \left(\frac{1}{e} \right) = \frac{1}{e} + 1$

$\Rightarrow \frac{dy}{dx} = \frac{(1 + e)}{(1 - e)}$

93. $y = e^{\log x}$

Taking logarithm on both sides, we get

$\log_e y = \log_e e^x$

Differentiating both sides w.r.t. x , we get

$\log_e \frac{dy}{dx} + x = \log_e e + \frac{1}{e} \cdot \frac{dx}{dx}$

$\Rightarrow \frac{dy}{dx} \left(\frac{\log_e y}{e} \right) = \frac{\log_e e}{e}$

$\Rightarrow \frac{dy}{dx} = \frac{(\log_e y - x)}{(\log_e y - e)}$

94. $y = 1$

Taking logarithm on both sides, we get

$\log y + \log x = 0$

Differentiating w.r.t. x , we get

$\log \frac{dy}{dx} + \frac{1}{x} + \frac{1}{x} \cdot \frac{dx}{dx} + \log x \cdot 1 = 0$

$\Rightarrow \frac{dy}{dx} \left(\log \frac{dy}{dx} + \frac{1}{x} \right) + \frac{1}{x} + \log x = 0$

$\Rightarrow \frac{dy}{dx} = - \frac{\left(\frac{1}{x} + \log x \right)}{\log \frac{dy}{dx} + \frac{1}{x}}$

$\Rightarrow \frac{dy}{dx} = - \frac{\left(\frac{1}{x} + \log x \right)}{\left(\frac{1}{x} + \log x \right)}$

95. $m^n = 2(x + y)^{m+n}$

Taking logarithm on both sides, we get

$m \log x + n \log y = \log 2 + (m+n) \log(x+y)$

Differentiating both sides w.r.t. x we get

$\frac{m}{x} + \frac{n}{y} \cdot \frac{dy}{dx} = \frac{m+n}{x+y} \left(1 + \frac{dy}{dx} \right)$

$\Rightarrow \left(\frac{m+n}{x+y} - \frac{n}{y} \right) \frac{dy}{dx} = \frac{m}{x} - \frac{m+n}{x+y}$

$\Rightarrow \frac{dy}{dx} = \dots$

96. $y = 2^{-x}$

Taking logarithm on both sides, we get

$\log y = (-x) \log 2$

Differentiating both sides w.r.t. x , we get

$\frac{1}{y} \cdot \frac{dy}{dx} = \log 2 \left(1 - \frac{dx}{dx} \right)$

$\Rightarrow (\log y + \log 2) \frac{dy}{dx} = \log 2 - \dots$

$\Rightarrow \left[\log(2^{-x}) \right] \frac{dy}{dx} = \frac{\log 2 - \dots}{\dots}$

$\Rightarrow \frac{dy}{dx} = \frac{\log 2 - \dots}{\log(2^{-x})}$

97. $y = a^x$

$\therefore \log y = x \log a$

$\therefore \log(\log y) = \log x + \log(\log a)$

Differentiating both sides w.r.t. x , we get

$\frac{1}{\log y} \cdot \frac{1}{y} \cdot \frac{dy}{dx} = \frac{1}{x} + \frac{dx}{dx} \log a$

$\Rightarrow \left(\frac{1}{\log y} - \log a \right) \frac{dy}{dx} = \dots$

$\Rightarrow (1 - \log y \log a) \frac{dy}{dx} = x^2 \log a$

98. $\log(x+y) = 2 \dots (i)$

Differentiating both sides w.r.t. x , we get

$\left(\frac{1}{x+y} \right) \left(1 + \frac{dy}{dx} \right) = 2 \left(\frac{dx}{dx} + \dots \right)$

$\Rightarrow \frac{dy}{dx} = \frac{1-2}{2^2+2} - \frac{2^2}{-1}$

Putting $y = 0$ in (i) we get

$y = 1$

$\therefore 'y'(0) = \frac{1-0-2}{0+0-1} = 1$



99. Let $y = e^{\cos \theta}$ and $z = e^{-\sin \theta}$

$$\therefore \frac{d}{d\theta} = e^{\cos \theta} (-\sin \theta) \text{ and}$$

$$\frac{dz}{d\theta} = e^{-\sin \theta} (-\cos \theta)$$

$$\therefore \frac{d}{dz} = \frac{\frac{d}{d\theta}}{\frac{dz}{d\theta}} = e^2$$

100. Let $y = \cos^{-1}(\sqrt{x})$ and $z = \sqrt{1-x}$

$$\therefore \frac{d}{dx} = \frac{-1}{\sqrt{1-x}} \cdot \frac{1}{2\sqrt{x}} \text{ and } \frac{dz}{dx} = \frac{-1}{2\sqrt{1-x}}$$

$$\therefore \frac{d}{dz} = \frac{\frac{d}{dx}}{\frac{dz}{dx}} = \frac{1}{\sqrt{x}}$$

101. $y = \frac{e^t + e^{-t}}{2}$ and $z = \frac{e^t - e^{-t}}{2}$

$$\therefore \frac{d}{dt} = \frac{e^t - e^{-t}}{2} \text{ and } \frac{dz}{dt} = \frac{e^t + e^{-t}}{2}$$

$$\therefore \frac{d}{dz} = \frac{\frac{d}{dt}}{\frac{dz}{dt}} = \frac{\frac{e^t - e^{-t}}{2}}{\frac{e^t + e^{-t}}{2}} = -$$

102. $y = a(t \cos t - \sin t)$ and $z = a(t \sin t + \cos t)$

$$\therefore \frac{d}{dt} = a(-t \sin t + \cos t - \cos t) = -at \sin t$$

$$\text{and } \frac{dz}{dt} = a(t \cos t + \sin t - \sin t) = at \cos t$$

$$\therefore \frac{d}{dz} = \frac{\frac{d}{dt}}{\frac{dz}{dt}} = \frac{-at \sin t}{at \cos t} = -\cot t$$

103. $y = a \cos^3 \theta$ and $z = a \sin^3 \theta$

$$\therefore \frac{d}{d\theta} = -3a \cos^2 \theta \cdot \sin \theta$$

$$\text{and } \frac{dz}{d\theta} = 3a \sin^2 \theta \cdot \cos \theta$$

$$\therefore \frac{d}{dz} = \frac{\frac{d}{d\theta}}{\frac{dz}{d\theta}} = -\tan \theta$$

$$\therefore \sqrt{1 + \left(\frac{d}{dz}\right)^2} = \sqrt{1 + \tan^2 \theta} = \sqrt{\sec^2 \theta} = |\sec \theta|$$

104. $y = \log(1 + \theta)$, $z = \sin^{-1} \theta$

$$\therefore \frac{d}{d\theta} = \frac{1}{1 + \theta}, \frac{dz}{d\theta} = \frac{1}{\sqrt{1 - \theta^2}}$$

$$\begin{aligned} \therefore \frac{d}{dz} &= \frac{\frac{d}{d\theta}}{\frac{dz}{d\theta}} = \frac{\sqrt{1 - \theta^2}}{1 + \theta} \\ &= \frac{\sqrt{(1 + \theta)(1 - \theta)}}{(1 + \theta)^2} = \sqrt{\frac{1 - \theta}{1 + \theta}} \end{aligned}$$

105. Let $y = \sin^{-1} \sqrt{x}$ and $z = \cos^{-1}(\sqrt{1-x^2})$

$$\therefore \frac{d}{dx} = \frac{1}{\sqrt{1-x^2}}$$

$$z = \cos^{-1}(\sqrt{1-x^2}) = \sin^{-1} x$$

$$\therefore \frac{dz}{dx} = \frac{1}{\sqrt{1-x^2}}$$

$$\therefore \frac{d}{dz} = \frac{\frac{d}{dx}}{\frac{dz}{dx}} = 1$$

106. Let $y = \sin^{-1}\left(\frac{1-x}{1+x}\right)$ and $z = \sqrt{x}$

$$\begin{aligned} \therefore \frac{d}{dx} &= \frac{1}{\sqrt{1 - \left(\frac{1-x}{1+x}\right)^2}} \cdot \frac{(1+x)(-1) - (1-x)(1)}{(1+x)^2} \\ &= \frac{-1}{\sqrt{x}(1+x)} \end{aligned}$$

$$\text{and } \frac{dz}{dx} = \frac{1}{2\sqrt{x}}$$

$$\therefore \frac{d}{dz} = \frac{\frac{d}{dx}}{\frac{dz}{dx}} = \frac{-2}{1+x}$$



107. Let $y = a^{\sec t}$ and $z = a^{\tan t}$

$$\therefore \frac{dy}{dt} = a^{\sec t} \log a \sec t$$

$$\text{and } \frac{dz}{dt} = a^{\tan t} \log a \sec^2 t$$

$$\begin{aligned} \therefore \frac{dy}{dz} &= \frac{\frac{dy}{dt}}{\frac{dz}{dt}} = \frac{a^{\sec t} \log a \sec t}{a^{\tan t} \log a \sec^2 t} \\ &= a^{\sec t - \tan t} \cdot \frac{\sin t}{\cos t} = \sin t \cdot a^{\sec t - \tan t} \end{aligned}$$

108. $y = e^{\theta} \left(\theta + \frac{1}{\theta} \right)$

$$\therefore \frac{dy}{d\theta} = e^{\theta} \left(1 - \frac{1}{\theta^2} \right) + e^{\theta} \left(\theta + \frac{1}{\theta} \right)$$

$$= e^{\theta} \left(1 + \theta + \frac{1}{\theta} - \frac{1}{\theta^2} \right)$$

$$= e^{\theta} \left(\frac{\theta^2 + \theta^3 + \theta - 1}{\theta^2} \right)$$

$$= e^{-\theta} \left(\theta - \frac{1}{\theta} \right)$$

$$\therefore \frac{dy}{d\theta} = e^{-\theta} \left(1 + \frac{1}{\theta^2} \right) - e^{-\theta} \left(\theta - \frac{1}{\theta} \right)$$

$$= e^{-\theta} \left(1 + \frac{1}{\theta^2} - \theta + \frac{1}{\theta} \right)$$

$$= e^{-\theta} \left(\frac{\theta^2 + 1 - \theta^3 + \theta}{\theta^2} \right)$$

$$\therefore \frac{dy}{d\theta} = \frac{\frac{dy}{d\theta}}{\frac{d\theta}{d\theta}} = \frac{e^{-2\theta} (1 + \theta^2 - \theta^3 + \theta)}{\theta^2 - 1 + \theta^3 + \theta}$$

109. $\frac{dy}{dt} = a \left(-\sin t + \frac{1}{\tan \frac{t}{2}} \cdot \sec^2 \frac{t}{2} \cdot \frac{1}{2} \right)$

$$= a \left(-\sin t + \frac{1}{2 \sin \frac{t}{2} \cos \frac{t}{2}} \right)$$

$$= a \left(-\sin t + \frac{1}{\sin t} \right) = a \left(\frac{\cos^2 t}{\sin t} \right)$$

$$= a \cos t \cot t$$

$$\text{and } \frac{dz}{dt} = a \cos t$$

$$\therefore \frac{dy}{dz} = \frac{\frac{dy}{dt}}{\frac{dz}{dt}} = \frac{1}{\cot t} = \tan t$$

110. $y = a(\sin 2\theta + \frac{1}{2} \sin 4\theta)$,
 $= b \left[\cos 2\theta - \frac{1}{2}(1 + \cos 4\theta) \right]$

$$\therefore \frac{dy}{d\theta} = 2a(\cos 2\theta + \cos 4\theta) = 2a(2\cos 3\theta \cos \theta)$$

$$\text{and } \frac{dz}{d\theta} = 2b(\sin 4\theta - \sin 2\theta) = 2b(2\cos 3\theta \sin \theta)$$

$$\therefore \frac{dy}{dz} = \frac{\frac{dy}{d\theta}}{\frac{dz}{d\theta}} = \frac{b}{a} \tan \theta$$

111. $y = t \log t$ and $z = t^t$

$$\therefore \frac{dy}{dz} = \log t^t = \log z$$

Differentiating both sides w.r.t. z , we get

$$1 = \frac{1}{z} \cdot \frac{dz}{dz}$$

$$\Rightarrow \frac{dy}{dz} = \frac{1}{z} = \frac{1}{t^t}$$

Since, $y = t \log t$

$$\therefore \frac{dy}{dz} = \log t^t$$

$$\Rightarrow e^y = t^t$$

$$\therefore \frac{dy}{dz} = e^{-y}$$

112. Let $x = \tan^{-1} \left(\frac{2}{1-t^2} \right)$ and $z = \sin^{-1} \left(\frac{2}{1+t^2} \right)$

Put $x = \tan \theta$

$$\therefore \theta = \tan^{-1} \left(\frac{2 \tan \theta}{1 - \tan^2 \theta} \right) = \tan^{-1} (\tan 2\theta) = 2\theta$$

$$\text{and } z = \sin^{-1} \left(\frac{2 \tan \theta}{1 + \tan^2 \theta} \right) = \sin^{-1} (\sin 2\theta) = 2\theta$$

$$\therefore x = z$$

$$\therefore \frac{dx}{dz} = 1$$



$$113. \text{ Let } x = \sin^{-1} \left(\frac{2}{1+x^2} \right) \text{ and } z = \cos^{-1} \left(\frac{1-x^2}{1+x^2} \right)$$

$$\therefore x = 2 \tan^{-1} x \text{ and } z = 2 \tan^{-1} x$$

$$\therefore \frac{dx}{dz} = \frac{\frac{d}{dx}}{\frac{dz}{dx}} = 1$$

$$114. \text{ Let } x = \cos^{-1} \left(\frac{1-x^2}{1+x^2} \right) \text{ and } z = \cot^{-1} \left(\frac{1-3x^2}{3-x^3} \right)$$

$$\therefore x = 2 \tan^{-1} x \text{ and } z = 3 \tan^{-1} x$$

$$\therefore \frac{dx}{dz} = \frac{\frac{d}{dx}}{\frac{dz}{dx}} = \frac{2}{3} \cdot \frac{1+x^2}{3} = \frac{2}{3}$$

$$115. \text{ Put } t = \sin \theta$$

$$\therefore x = \sin^{-1} (3 \sin \theta - 4 \sin^3 \theta)$$

$$= \sin^{-1} (\sin 3\theta) = 3\theta$$

$$= \cos^{-1} (\sqrt{1-\sin^2 \theta}) = \cos^{-1} (\cos \theta) = \theta$$

$$\therefore x = 3\theta$$

$$\Rightarrow \frac{dx}{d\theta} = \frac{1}{3}$$

$$\therefore \frac{dx}{dt} = \frac{1}{3}$$

$$116. \sin x = \frac{t}{\sqrt{1+t^2}}$$

$$\text{Put } t = \tan \theta$$

$$\therefore \sin x = \frac{\tan \theta}{\sec \theta} = \sin \theta$$

$$\therefore x = \theta$$

$$\therefore \frac{dx}{d\theta} = 1$$

$$\cos x = \frac{1}{\sqrt{1+t^2}} = \frac{1}{\sec \theta} = \cos \theta$$

$$\therefore x = \theta \quad \therefore \frac{dx}{d\theta} = 1$$

$$\therefore \frac{dx}{dt} = \frac{\frac{d}{d\theta}}{\frac{d}{d\theta}} = 1$$

$$117. \text{ Let } x = \tan^{-1} \left(\sqrt{\frac{1-x^2}{1+x^2}} \right) \text{ and } z = \cos^{-1}(x^2)$$

$$\text{Put } x^2 = \cos 2\theta \Rightarrow \theta = \frac{1}{2} \cos^{-1}(x^2)$$

$$\therefore x = \tan^{-1} \left(\sqrt{\frac{2 \sin^2 \theta}{2 \cos^2 \theta}} \right) = \tan^{-1}(\tan \theta) = \theta$$

$$\Rightarrow x = \frac{1}{2} \cos^{-1}(x^2)$$

$$\Rightarrow x = \frac{1}{2} z$$

$$\therefore \frac{dx}{dz} = \frac{1}{2}$$

$$118. \text{ Putting } t = \tan \theta \text{ in the given equations, we get}$$

$$= \frac{1-\tan^2 \theta}{1+\tan^2 \theta} = \cos 2\theta \text{ and}$$

$$= \frac{2 \tan \theta}{1+\tan^2 \theta} = \sin 2\theta$$

$$\therefore \frac{dx}{d\theta} = -2 \sin 2\theta \text{ and } \frac{dy}{d\theta} = 2 \cos 2\theta$$

$$\therefore \frac{dx}{dy} = \frac{\frac{dx}{d\theta}}{\frac{dy}{d\theta}} = \frac{-2 \sin 2\theta}{2 \cos 2\theta} = -\tan 2\theta$$

$$119. \text{ Put } x = \sin \theta \Rightarrow 2 \sin^{-1} x = 2\theta$$

$$\Rightarrow \sin(2 \sin^{-1} x) = \sin 2\theta \Rightarrow x = \sin 2\theta$$

$$\therefore \frac{dx}{d\theta} = \cos \theta \text{ and } \frac{dy}{d\theta} = 2 \cos 2\theta$$

$$\therefore \frac{dx}{dy} = \frac{\frac{dx}{d\theta}}{\frac{dy}{d\theta}} = \frac{\cos \theta}{2 \cos 2\theta} = \frac{2(1-2 \sin^2 \theta)}{\sqrt{1-\sin^2 \theta}} = \frac{2-4x^2}{\sqrt{1-x^2}}$$

$$120. \text{ Let } x = \tan^{-1} \left[\frac{\sin \theta}{1+\cos \theta} \right] \text{ and } z = \tan^{-1} \left[\frac{\cos \theta}{1+\sin \theta} \right]$$

$$= \tan^{-1} \left[\frac{2 \sin(\theta/2) \cos(\theta/2)}{2 \cos^2(\theta/2)} \right]$$

$$= \tan^{-1} \left[\tan \left(\frac{\theta}{2} \right) \right] = \frac{\theta}{2}$$

$$\therefore \frac{dx}{dz} = \frac{1}{2}$$



$$\begin{aligned}
 z &= \tan^{-1} \left[\frac{\cos}{1 + \sin} \right] \\
 &= \tan^{-1} \left[\frac{(\cos^2 / 2 - \sin^2 / 2)}{(\sin / 2 + \cos / 2)^2} \right] \\
 &= \tan^{-1} \left[\frac{(\cos / 2 - \sin / 2)}{(\cos / 2 + \sin / 2)} \right] \\
 &= \tan^{-1} \left(\frac{1 - \tan / 2}{1 + \tan / 2} \right) \\
 &= \tan^{-1} [\tan(\pi/4 - / 2)] = \pi/4 - / 2
 \end{aligned}$$

$$\therefore \frac{dz}{d} = -\frac{1}{2}$$

$$\therefore \frac{d}{dz} = \frac{d}{\frac{dz}{d}} = -1$$

$$121. \quad = a \cos^4 \theta \text{ and } = a \sin^4 \theta$$

$$\therefore \frac{d}{d\theta} = -4a \cos^3 \theta \sin \theta$$

$$\text{and } \frac{d}{d\theta} = 4a \sin^3 \theta \cos \theta$$

$$\therefore \frac{d}{d} = \frac{d\theta}{d} = \frac{-\sin^2 \theta}{\cos^2 \theta} = -\tan^2 \theta$$

$$\therefore \left(\frac{d}{d} \right)_{\left(\theta = \frac{3\pi}{4} \right)} = -\tan^2 \left(\frac{3\pi}{4} \right) = -(-1)^2 = -1$$

$$122. \quad \text{Let } = \sec^{-1} \left(\frac{1}{2^2 - 1} \right) \text{ and } z = \sqrt{1+3}$$

$$\therefore = \cos^{-1} (2^2 - 1) = 2\cos^{-1}$$

$$\therefore \frac{d}{d} = \frac{-2}{\sqrt{1-^2}} \text{ and } \frac{dz}{d} = \frac{3}{2\sqrt{1+3}}$$

$$\therefore \frac{d}{dz} = \frac{d}{\frac{dz}{d}} = \frac{-2}{\sqrt{1-^2}} \times \frac{2\sqrt{1+3}}{3}$$

$$\therefore \left(\frac{d}{dz} \right)_{\left(\frac{-1}{3} \right)} = 0$$

$$123. \quad = \sin t \cos 2t \text{ and } = \cos t \sin 2t$$

$$\therefore \frac{d}{dt} = \cos t \cos 2t - 2\sin t \sin 2t$$

$$\text{and } \frac{d}{dt} = 2\cos t \cos 2t - \sin t \sin 2t$$

$$\therefore \frac{d}{d} = \frac{d}{\frac{d}{dt}} = \frac{2\cos t \cos 2t - \sin t \sin 2t}{\cos t \cos 2t - 2\sin t \sin 2t}$$

$$\therefore \left(\frac{d}{d} \right)_{\left(t = \frac{\pi}{4} \right)} = \frac{0 - \frac{1}{\sqrt{2}}}{0 - 2 \times \frac{1}{\sqrt{2}}} = \frac{1}{2}$$

$$124. \quad \text{Let } = \tan^{-1} \left(\frac{\sqrt{1+^2} - 1}{2} \right) \text{ and}$$

$$z = \tan^{-1} \left(\frac{2\sqrt{1-^2}}{1-2^2} \right)$$

$$\text{Put } = \tan \theta \Rightarrow \theta = \tan^{-1}$$

$$\therefore = \tan^{-1} \left(\frac{\sec \theta - 1}{\tan \theta} \right) = \tan^{-1} \left(\tan \frac{\theta}{2} \right)$$

$$= \frac{\theta}{2} = \frac{1}{2} \tan^{-1}$$

$$\therefore \frac{d}{d} = \frac{1}{2(1+^2)}$$

$$z = \tan^{-1} \left(\frac{2\sqrt{1-^2}}{1-2^2} \right)$$

$$\text{Put } = \sin \theta \Rightarrow \theta = \sin^{-1}$$

$$\therefore z = \tan^{-1} \left(\frac{2\sin \theta \cos \theta}{1-2\sin^2 \theta} \right) = \tan^{-1} \left(\frac{\sin 2\theta}{\cos 2\theta} \right)$$

$$= \tan^{-1}(\tan 2\theta) = 2\theta$$

$$= 2 \sin^{-1}$$

$$\therefore \frac{dz}{d} = \frac{2}{\sqrt{1-^2}}$$

$$\therefore \frac{d}{dz} = \frac{d}{\frac{dz}{d}} = \frac{\sqrt{1-^2}}{4(1+^2)}$$

$$\therefore \left(\frac{d}{dz} \right)_{=0} = \frac{1}{4}$$

$$125. \quad = \cos^2 \frac{3}{2} - \sin^2 \frac{3}{2}$$

$$\Rightarrow = \cos 3 \quad \dots(i)$$

$$\therefore \frac{d}{d} = -3 \sin 3$$

$$\therefore \frac{d^2}{d^2} = -9 \cos 3$$

$$\Rightarrow \frac{d^2}{d^2} = -9 \quad \dots[\text{From (i)}]$$



$$126. \quad = t^2 \text{ and } = t^3 + 1$$

$$\therefore \frac{d}{dt} = 2t \text{ and } \frac{d}{dt} = 3t^2$$

$$\therefore \frac{d}{d} = \frac{\frac{d}{dt}}{\frac{d}{dt}} = \frac{3t}{2}$$

$$\therefore \frac{d^2}{d^2} = \frac{3}{2} \cdot \frac{dt}{d} = \frac{3}{2} \cdot \frac{1}{2t} = \frac{3}{4t}$$

$$127. \quad \frac{d}{dt} = 10t^9 \text{ and } \frac{d}{dt} = 8t^7$$

$$\therefore \frac{d}{d} = \frac{\frac{d}{dt}}{\frac{d}{dt}} = \frac{5t^2}{4}$$

$$\therefore \frac{d^2}{d^2} = \frac{5}{4} \cdot 2t \cdot \frac{dt}{d} = \frac{5t}{2} \cdot \frac{1}{8t^7} = \frac{5}{16t^6}$$

$$128. \quad = \log t \text{ and } = \frac{1}{t}$$

$$\therefore \frac{d}{dt} = \frac{1}{t} \text{ and } \frac{d}{dt} = -\frac{1}{t^2}$$

$$\therefore \frac{d}{d} = \frac{\frac{d}{dt}}{\frac{d}{dt}} = -\frac{1}{t} \quad \dots(i)$$

$$\therefore \frac{d^2}{d^2} = -\left(-\frac{1}{t^2}\right) \frac{dt}{d}$$

$$= \frac{1}{t^2} \cdot \frac{1}{\frac{d}{dt}} = \frac{1}{t^2} \cdot \frac{1}{\frac{1}{t}} = \frac{1}{t}$$

$$\therefore \frac{d^2}{d^2} = -\frac{d}{d} \quad \dots[\text{From (i)}]$$

$$129. \quad = 1 - \frac{1}{2!} + \frac{1}{3!} - \frac{1}{4!} + \dots$$

$$\Rightarrow = e^{-1} \quad \dots(i)$$

$$\therefore \frac{d}{d} = e^{-1}(-1)$$

$$\therefore \frac{d^2}{d^2} = (-1)\{e^{-1}(-1)\} = e^{-1} = \dots[\text{From (i)}]$$

130. Consider option (C),

$$f(x) = \sin x$$

$$\Rightarrow f(0) = 0 \text{ and}$$

$$f'(x) = \cos x$$

$$\Rightarrow f'(0) = 1$$

$$\text{Also, } f''(x) = -\sin x = -f(x)$$

\(\therefore\) option (C) is the correct answer.

$$131. \quad e^{-(x+1)} = 1 \quad \Rightarrow e^{-x} = \frac{1}{+1}$$

$$\Rightarrow = \log\left(\frac{1}{+1}\right)$$

$$\Rightarrow = -\log(+1)$$

$$\therefore \frac{d}{d} = -\frac{1}{+1} \quad \dots(i)$$

$$\therefore \frac{d^2}{d^2} = \frac{1}{(+1)^2} = \left(\frac{-1}{+1}\right)^2$$

$$= \left(\frac{d}{d}\right)^2 \quad \dots[\text{From (i)}]$$

$$132. \quad = a^5 + \frac{b}{4} \quad \dots(i)$$

$$\therefore \frac{d}{d} = 5a^4 - \frac{4b}{5}$$

$$\therefore \frac{d^2}{d^2} = 20a^3 + \frac{20b}{6}$$

$$= \frac{20}{2} \left(a^5 + \frac{b}{4}\right) = \frac{20}{2} \quad \dots[\text{From (i)}]$$

$$133. \quad = a^{n+1} + b^{-n} \quad \dots(i)$$

$$\therefore \frac{d}{d} = (n+1)a^n - nb^{-n-1}$$

$$\therefore \frac{d^2}{d^2} = n(n+1)a^{n-1} + n(n+1)b^{-n-2}$$

$$\Rightarrow \frac{d^2}{d^2} = \frac{n(n+1)}{2}(a^{n+1} + b^{-n})$$

$$\Rightarrow 2 \frac{d^2}{d^2} = n(n+1) \quad \dots[\text{From (i)}]$$

$$134. \quad = a \cos(\log x) + b \sin(\log x) \quad \dots(i)$$

$$\therefore ' = \frac{-a \sin(\log x)}{x} + \frac{b \cos(\log x)}{x}$$

$$\Rightarrow ' = -a \sin(\log x) + b \cos(\log x)$$

Differentiating both sides w.r.t. x , we get

$$'' + ' = \frac{-a \cos(\log x)}{x} - \frac{b \sin(\log x)}{x}$$

$$\Rightarrow 2'' + ' = -[a \cos(\log x) + b \sin(\log x)]$$

$$\Rightarrow 2'' + ' = - \dots[\text{From (i)}]$$

$$135. \quad = a \cdot b^{2^{-1}} \quad \dots(i)$$

$$\therefore \frac{d}{d} = b^{2^{-1}} \cdot a \log a + a \cdot 2b^{2^{-1}} \log b$$

$$= a b^{2^{-1}} (\log a + 2 \log b)$$

$$\therefore \frac{d^2}{d^2} = a b^{2^{-1}} (\log a + 2 \log b)^2$$

$$= a b^{2^{-1}} (\log ab^2)^2$$

$$= (\log ab^2)^2 \quad \dots[\text{From (i)}]$$



$$136. \quad = \log \left(+ \sqrt{^2 + a^2} \right)$$

$$\therefore \frac{d}{d} = \frac{1}{+\sqrt{^2 + a^2}} \cdot \left(1 + \frac{1}{2\sqrt{^2 + a^2}} \cdot 2 \right)$$

$$\Rightarrow \frac{d}{d} = \frac{1}{+\sqrt{^2 + a^2}} \times \frac{\sqrt{^2 + a^2} +}{\sqrt{^2 + a^2}}$$

$$\Rightarrow \frac{d}{d} = \frac{1}{\sqrt{^2 + a^2}}$$

$$\therefore \frac{d^2}{d^2} = \frac{-1}{2} \left(^2 + a^2 \right)^{\frac{3}{2}} \cdot 2 = \frac{-}{\left(^2 + a^2 \right)^{\frac{3}{2}}}$$

$$137. \quad = ^2 + 2 + 3$$

$$\therefore \frac{d}{d} = 2 + 2$$

$$\Rightarrow \frac{d}{d} = \frac{1}{2 + 2}$$

$$\therefore \frac{d^2}{d^2} = \frac{-1}{2(+1)^2} \cdot \frac{d}{d} = \frac{-1}{4(+1)^3}$$

$$138. \quad = + e$$

$$\therefore \frac{d}{d} = 1 + e \quad \dots(i)$$

$$\therefore \frac{d}{d} = \frac{1}{1+e} = (1+e)^{-1}$$

$$\therefore \frac{d^2}{d^2} = -(1+e)^{-2} \cdot \frac{d}{d} (1+e)$$

$$= -(1+e)^{-2} \cdot e \cdot \frac{d}{d}$$

$$= -\frac{e}{(1+e)^2} \cdot \frac{1}{1+e} \quad \dots[\text{From (i)}]$$

$$= -\frac{e}{(1+e)^3}$$

$$139. \quad = \sin + e$$

$$\therefore \frac{d}{d} = \cos + e$$

$$\Rightarrow \frac{d}{d} = (\cos + e)^{-1} \quad \dots(i)$$

$$\therefore \frac{d^2}{d^2} = -(\cos + e)^{-2} \cdot (-\sin + e) \cdot \frac{d}{d}$$

$$= \frac{(\sin - e)}{(\cos + e)^2} \cdot (\cos + e)^{-1} \quad \dots[\text{From (i)}]$$

$$= \frac{\sin - e}{(\cos + e)^3}$$

$$140. \quad = e^2$$

$$\therefore \frac{d}{d} = 2e^2 \quad \therefore \frac{d^2}{d^2} = 4e^2$$

Now, $= e^2$

$$\therefore \log = 2$$

$$\therefore = \frac{1}{2} \log \quad \therefore \frac{d}{d} = \frac{1}{2}$$

$$\therefore \frac{d^2}{d^2} = \frac{-1}{2^2} = \frac{-1}{2(e^2)^2}$$

$$\therefore \frac{d^2}{d^2} \times \frac{d^2}{d^2} = \frac{-2}{e^2} = -2e^{-2}$$

$$141. \quad \text{Let } = 2 \cos \cos 3$$

$$\Rightarrow = \cos 4 + \cos 2$$

$$\therefore \frac{d}{d} = -4 \sin 4 - 2 \sin 2$$

$$\therefore \frac{d^2}{d^2} = -16 \cos 4 - 4 \cos 2$$

$$= -4(\cos 2 + 4 \cos 4)$$

$$= -2^2(\cos 2 + 2^2 \cos 4)$$

$$142. \quad \frac{d}{d} = \left(\frac{d}{d} \right)^{-1}$$

$$\therefore \frac{d}{d} \left(\frac{d}{d} \right) = \frac{d}{d} \left\{ \left(\frac{d}{d} \right)^{-1} \right\}$$

$$\Rightarrow \frac{d^2}{d^2} = \frac{d}{d} \left\{ \left(\frac{d}{d} \right)^{-1} \right\} \frac{d}{d}$$

$$\Rightarrow \frac{d^2}{d^2} = -\left(\frac{d}{d} \right)^{-2} \cdot \frac{d}{d} \left(\frac{d}{d} \right) \cdot \frac{d}{d}$$

$$\Rightarrow \frac{d^2}{d^2} = -\left(\frac{d}{d} \right)^{-3} \left(\frac{d^2}{d^2} \right)$$

$$143. \quad \frac{^2}{a^2} + \frac{^2}{b^2} = 1 \quad \dots(i)$$

Differentiating both sides w.r.t. , we get

$$\frac{2}{a^2} + \frac{2}{b^2} \cdot \frac{d}{d} = 0$$

$$\therefore \frac{d}{d} = -\frac{b^2}{a^2} \quad \dots(ii)$$



$$\begin{aligned} \therefore \frac{d^2}{d^2} &= -\frac{b^2}{a^2} \left(-\frac{d}{d} \right) \\ &= -\frac{b^2}{a^2} \left(-\frac{d}{d} \right) \\ &= -\frac{b^2}{a^2} \left(+\frac{b^2}{a^2} \right) \quad \dots[\text{From (ii)}] \\ &= -\frac{b^2}{a^2} \cdot \frac{b^2}{a^2} \left(\frac{b^2}{b^2} + \frac{a^2}{a^2} \right) = \frac{-b^4}{a^2 \cdot a^2} \dots[\text{From (i)}] \end{aligned}$$

$$144. \quad = f(t) \text{ and } = g(t)$$

$$\therefore \frac{d}{dt} = f'(t) \text{ and } \frac{d}{dt} = g'(t)$$

$$\therefore \frac{d}{d} = \frac{\frac{d}{dt}}{\frac{d}{dt}} = \frac{g'(t)}{f'(t)}$$

$$\begin{aligned} \therefore \frac{d^2}{d^2} &= \frac{f'(t) \cdot g''(t) - g'(t) \cdot f''(t)}{[f'(t)]^2} \cdot \frac{dt}{d} \\ &= \frac{f'(t) \cdot g''(t) - g'(t) \cdot f''(t)}{[f'(t)]^3} \end{aligned}$$

$$145. \quad = \frac{\cos - \sin}{\cos + \sin} = \frac{1 - \tan}{1 + \tan}$$

$$\Rightarrow = \tan \left(\frac{\pi}{4} - \right) \quad \dots(i)$$

$$\therefore \frac{d}{d} = -\sec^2 \left(\frac{\pi}{4} - \right)$$

$$\therefore \frac{d^2}{d^2} = 2 \sec^2 \left(\frac{\pi}{4} - \right) \cdot \tan \left(\frac{\pi}{4} - \right)$$

$$\therefore \frac{\frac{d^2}{d^2}}{\frac{d}{d}} = -2 \tan \left(\frac{\pi}{4} - \right) = -2 \quad \dots[\text{From (i)}]$$

$$146. \quad = \cos(\log) \quad \dots(i)$$

$$\therefore \frac{d}{d} = -\sin(\log) \cdot \frac{1}{d}$$

$$\Rightarrow \frac{d}{d} = -\sin(\log)$$

Differentiating both sides w.r.t. , we get

$$\frac{d^2}{d^2} + \frac{d}{d} \cdot 1 = -\cos(\log) \cdot \frac{1}{d}$$

$$\Rightarrow \frac{d^2}{d^2} + \frac{d}{d} = - \quad \dots[\text{From (i)}]$$

$$\Rightarrow \frac{d^2}{d^2} + \frac{d}{d} + = 0$$

$$147. \quad = e^{\tan}$$

$$\Rightarrow \log = \tan$$

Differentiating both sides w.r.t. , we get

$$\frac{1}{d} \cdot \frac{d}{d} = \sec^2 \Rightarrow \frac{d}{d} = \frac{1}{\cos^2}$$

$$\Rightarrow \cos^2 \frac{d}{d} =$$

Differentiating both sides w.r.t. , we get

$$\cos^2 \frac{d^2}{d^2} - 2 \cos \sin \frac{d}{d} = \frac{d}{d}$$

$$\Rightarrow \cos^2 \frac{d^2}{d^2} = (1 + \sin 2) \frac{d}{d}$$

$$148. \quad = e^{m \cos^{-1}} \quad \dots(i)$$

$$\therefore \frac{d}{d} = e^{m \cos^{-1}} \cdot m \cdot \frac{-1}{\sqrt{1 - ^2}}$$

$$\Rightarrow \sqrt{1 - ^2} \frac{d}{d} = -m \quad \dots[\text{From (i)}]$$

$$\Rightarrow (1 - ^2) \left(\frac{d}{d} \right)^2 = m^2$$

Differentiating both sides w.r.t. , we get

$$(1 - ^2) \cdot 2 \frac{d}{d} \cdot \frac{d^2}{d^2} + \left(\frac{d}{d} \right)^2 \cdot (0 - 2) = 2m^2 \frac{d}{d}$$

$$\Rightarrow (1 - ^2) \frac{d^2}{d^2} - \frac{d}{d} = m^2$$

$$\Rightarrow (1 - ^2) \frac{d^2}{d^2} - \frac{d}{d} - m^2 = 0$$

$$149. \quad \frac{d}{d} = \frac{2 \sin^{-1}}{\sqrt{1 - ^2}} - \frac{2 \cos^{-1}}{\sqrt{1 - ^2}}$$

$$\Rightarrow \frac{d}{d} = \frac{2(\sin^{-1} - \cos^{-1})}{\sqrt{1 - ^2}}$$

$$\Rightarrow \sqrt{1 - ^2} \frac{d}{d} = 2(\sin^{-1} - \cos^{-1})$$

Differentiating both sides w.r.t. , we get

$$\sqrt{1 - ^2} \cdot \frac{d^2}{d^2} + \frac{d}{d} \cdot \frac{1}{2\sqrt{1 - ^2}} \cdot (-2)$$

$$= 2 \left(\frac{1}{\sqrt{1 - ^2}} - \frac{(-1)}{\sqrt{1 - ^2}} \right) = \frac{4}{\sqrt{1 - ^2}}$$

$$\therefore (1 - ^2) \frac{d^2}{d^2} - \frac{d}{d} = 4$$



150. $y = \cos(m \sin^{-1} x) \dots (i)$
 $\therefore \frac{dy}{dx} = -\sin(m \sin^{-1} x) \cdot \frac{m}{\sqrt{1-x^2}}$
 $\Rightarrow \sqrt{1-x^2} \frac{dy}{dx} = -m \sin(m \sin^{-1} x)$
 Differentiating both sides w.r.t. x , we get
 $\sqrt{1-x^2} \frac{d^2y}{dx^2} - \frac{1}{\sqrt{1-x^2}} \frac{dy}{dx} = -m \cos(m \sin^{-1} x) \cdot \frac{m}{\sqrt{1-x^2}}$
 $\Rightarrow (1-x^2) \frac{d^2y}{dx^2} - \frac{dy}{dx} = -m^2 \dots [From (i)]$
 $\Rightarrow (1-x^2) \frac{d^2y}{dx^2} - \frac{dy}{dx} + m^2 = 0$

151. $y^2 = a^2 + b^2 + c$
 Differentiating both sides w.r.t. x , we get
 $2y \frac{dy}{dx} = 2a^2 + b^2$
 Differentiating both sides w.r.t. x , we get
 $2y \frac{d^2y}{dx^2} + \frac{dy}{dx} \cdot 2 \frac{dy}{dx} = 2a^2$
 Multiplying both the sides by y^2 , we get
 $3y^3 \frac{d^2y}{dx^2} = a^2 - \left(\frac{dy}{dx}\right)^2$
 $= a(a^2 + b^2 + c) - \left(a + \frac{b}{2}\right)^2$
 $= a^2 + ab + ac - a^2 - \frac{b^2}{4} - ab$
 $= ac - \frac{b^2}{4} = a \text{ constant}$

152. $y = \tan^{-1} \left[\frac{\log e}{\log \frac{e}{x}} \right] + \tan^{-1} \left[\frac{8 - \log x}{1 + 8 \log x} \right]$
 $\Rightarrow y = \tan^{-1} \left[\frac{1 + \log x}{1 - \log x} \right] + \tan^{-1} \left[\frac{8 - \log x}{1 + 8 \log x} \right]$
 $\Rightarrow y = \tan^{-1} 1 + \tan^{-1}(\log x) + \tan^{-1} 8 - \tan^{-1}(\log x)$
 $\Rightarrow y = \tan^{-1} 1 + \tan^{-1} 8$
 $\therefore \frac{dy}{dx} = 0, \quad \therefore \frac{d^2y}{dx^2} = 0$

153. $y = \sin t$ and $x = \sin^3 t$
 $\therefore \frac{dy}{dx} = 3x^2$
 $\therefore \frac{d^2y}{dx^2} = 6$
 At $t = \frac{\pi}{2}$, $y = \sin \frac{\pi}{2} = 1$
 $\therefore \left(\frac{d^2y}{dx^2}\right)_{t=\frac{\pi}{2}} = \left(\frac{d^2y}{dx^2}\right)_{y=1} = 6(1) = 6$

154. $y = a(1 - \cos \theta)$ and $x = a(\theta + \sin \theta)$
 $\therefore \frac{dy}{d\theta} = a \sin \theta$ and $\frac{dx}{d\theta} = a(1 + \cos \theta)$
 $\therefore \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{a \sin \theta}{a(1 + \cos \theta)} = \frac{2 \cos^2 \frac{\theta}{2}}{2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}} = \cot \frac{\theta}{2}$
 $\therefore \frac{d^2y}{dx^2} = -\operatorname{cosec}^2 \frac{\theta}{2} \cdot \frac{1}{2} \cdot \frac{d\theta}{dx}$
 $= -\frac{1}{2} \operatorname{cosec}^2 \frac{\theta}{2} \cdot \frac{1}{a \sin \theta}$
 $\therefore \left(\frac{d^2y}{dx^2}\right)_{\theta=\frac{\pi}{2}} = -\frac{1}{2} (\sqrt{2})^2 \cdot \frac{1}{a(1)} = -\frac{1}{a}$

155. Let $y = a \sin^3 t$ and $x = a \cos^3 t$
 $\therefore \frac{dy}{dt} = 3a \sin^2 t \cos t$
 and $\frac{dx}{dt} = -3a \cos^2 t \sin t \dots (i)$
 $\therefore \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = -\tan t$
 $\therefore \frac{d^2y}{dx^2} = -\sec^2 t \cdot \frac{dt}{dx}$
 $= -\sec^2 t \cdot \frac{1}{-3a \cos^2 t \sin t} \dots [From (i)]$
 $= \frac{1}{3a \cos^4 t \sin t}$
 $\therefore \left(\frac{d^2y}{dx^2}\right)_{t=\frac{\pi}{4}} = \frac{1}{3a \cos^4 \left(\frac{\pi}{4}\right) \sin \left(\frac{\pi}{4}\right)}$
 $= \frac{1}{3a \left(\frac{1}{\sqrt{2}}\right)^5} = \frac{4\sqrt{2}}{3a}$

156. $e^y + x = e$
 Differentiating both sides w.r.t. x , we get
 $e \frac{dy}{dx} + 1 = 0 \dots (i)$
 Again, differentiating both sides w.r.t. x , we get
 $e \frac{d^2y}{dx^2} + e \left(\frac{dy}{dx}\right)^2 + 2 \frac{dy}{dx} + \frac{d^2x}{dx^2} = 0 \dots (ii)$
 Putting $\frac{dy}{dx} = 0$ in $e \frac{dy}{dx} + 1 = 0$, we get $e = 1$
 Putting $e = 1$, $\frac{dy}{dx} = 0$ in (ii), we get
 $\frac{d^2y}{dx^2} = -\frac{1}{e}$



Putting $x = 0$, $y = 1$, $\frac{dy}{dx} = -\frac{1}{e}$ in (ii), we get

$$e \frac{d^2 y}{dx^2} + e \cdot \frac{1}{e^2} - \frac{2}{e} + 0 = 0 \Rightarrow \frac{d^2 y}{dx^2} = \frac{1}{e^2}$$

158. $f(-x) = -f(x)$... [$\because f(x)$ is an odd function]

$$\therefore f(-x) = -f(x)$$

Differentiating w.r.t. x , we get

$$f'(x) = -[-f'(-x)]$$

$$\Rightarrow f'(x) = f'(-x)$$

$$\Rightarrow f'(3) = f'(-3)$$

$$\Rightarrow f'(-3) = 2$$

$$159. \left(\frac{1}{x} \right) + \left(\frac{1}{x} \right) = 2$$

$$\Rightarrow \frac{1}{x^2} + \frac{1}{x^2} = 2$$

$$\Rightarrow \left(\frac{1}{x} \right)^2 = 0$$

$$\Rightarrow \frac{1}{x} = 0$$

$$\Rightarrow \frac{d}{dx} = 1$$

$$\therefore \frac{d}{dx} = 1$$

$$160. y = e \cdot e^2 \cdot e^3 \dots e^n$$

$$\Rightarrow y = e^{(1+2+3+\dots+n)}$$

$$\Rightarrow y = e^{\left[\frac{n(n+1)}{2} \right]}$$

$$\Rightarrow \log y = \left[\frac{n(n+1)}{2} \right]$$

Differentiating both sides w.r.t. x , we get

$$\frac{1}{y} \frac{dy}{dx} = \frac{n(n+1)}{2}$$

$$\Rightarrow \frac{dy}{dx} = \frac{n(n+1)}{2} y$$

$$161. y = \sqrt{\frac{1+e}{1-e}} \Rightarrow y^2 = \frac{1+e}{1-e}$$

Differentiating both sides w.r.t. x , we get

$$2 \frac{dy}{dx} = \frac{(1-e)e + (1+e)e}{(1-e)^2} = \frac{2e}{(1-e)^2}$$

$$\therefore \frac{dy}{dx} = \frac{e}{(1-e)^2} \sqrt{\frac{1-e}{1+e}}$$

$$= \frac{e}{(1-e)^2} \sqrt{\left(\frac{1-e}{1+e} \right) \left(\frac{1-e}{1-e} \right)}$$

$$= \frac{e}{(1-e) \sqrt{1-e^2}}$$



Competitive Thinking

$$1. Lf'(0) = \lim_{h \rightarrow 0^-} \frac{f(0+h) - f(0)}{h}$$

$$= \lim_{h \rightarrow 0^-} \frac{1-1}{h} = 0$$

$$Rf'(0) = \lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h}$$

$$= \lim_{h \rightarrow 0^+} \frac{1 + \sinh - 1}{h}$$

$$= \lim_{h \rightarrow 0^+} \frac{\sinh}{h} = 1$$

$$\therefore Lf'(0) \neq Rf'(0)$$

$$\therefore f'(0) \text{ does not exist.}$$

$$2. f(x) = \begin{cases} \frac{1}{x-1}, & \text{if } x \neq 1, 2 \\ 2, & \text{if } x = 1 \\ 1, & \text{if } x = 2 \end{cases}$$

$$\therefore \lim_{x \rightarrow 2} \frac{f(x) - f(2)}{x - 2} = \lim_{x \rightarrow 2} \frac{\frac{1}{x-1} - 1}{x - 2}$$

$$= -\lim_{x \rightarrow 2} \frac{-2}{(x-1)(x-2)}$$

$$= -\lim_{x \rightarrow 2} \frac{1}{x-1}$$

$$= -1$$

$$3. f(x) = \begin{cases} \frac{1}{2x-5}, & \text{for } x \neq 1 \\ -\frac{1}{3}, & \text{for } x = 1 \end{cases}$$

$$\therefore f'(1) = \lim_{x \rightarrow 1} \frac{f(x) - f(1)}{x - 1}$$

$$= \lim_{x \rightarrow 1} \frac{\frac{1}{2x-5} - \left(-\frac{1}{3}\right)}{x - 1}$$

$$= \lim_{x \rightarrow 1} \frac{2 - 2}{3(2x-5)(x-1)}$$

$$= \frac{2}{3} \lim_{x \rightarrow 1} \frac{-1}{(2x-5)(x-1)}$$

$$= \frac{2}{3} \lim_{x \rightarrow 1} \frac{1}{2x-5} = -\frac{2}{9}$$



$$\begin{aligned}
 4. \quad f'(0^-) &= \lim_{h \rightarrow 0^-} \frac{f(0-h) - f(0)}{-h} \\
 &= \lim_{h \rightarrow 0^-} \frac{f(0-h) - f(0)}{-h} \\
 &= \lim_{h \rightarrow 0^-} \frac{h^2 \log(\cosh) - 0}{\log(1+h^2)} \\
 &= \lim_{h \rightarrow 0^-} \frac{-\log(\cosh)}{h} \lim_{h \rightarrow 0^-} \frac{1}{\frac{\log(1+h^2)}{h^2}} \\
 &= \lim_{h \rightarrow 0^-} \frac{-\log(\cosh)}{h} \cdot (1)
 \end{aligned}$$

Applying L'Hospital rule, we get

$$\begin{aligned}
 &= \lim_{h \rightarrow 0^-} \frac{\sinh}{\cosh} \\
 &= 0
 \end{aligned}$$


$$\begin{aligned}
 f(0^+) &= \lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h} \\
 &= \lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h} \\
 &= \lim_{h \rightarrow 0^+} \frac{h^2 \log(\cosh) - 0}{\log(1+h^2)} \\
 &= 0
 \end{aligned}$$

- ∴ $f'(0^-) = f'(0^+)$
- ∴ $f(x)$ is differentiable at zero.

$$\begin{aligned}
 5. \quad Lf'(1) &= \lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{x - 1} \\
 &= \lim_{x \rightarrow 1^-} \frac{\frac{x^2}{4} - \frac{3}{2} + \frac{13}{4} - 2}{x - 1} \\
 &= \lim_{x \rightarrow 1^-} \frac{x^2 - 6x + 5}{4(x - 1)} \\
 &= \lim_{x \rightarrow 1^-} \frac{(x - 5)(x - 1)}{4(x - 1)} \\
 &= \frac{1}{4} \lim_{x \rightarrow 1^-} (x - 5) = -1 \\
 Rf'(1) &= \lim_{x \rightarrow 1^+} \frac{f(x) - f(1)}{x - 1} = \lim_{x \rightarrow 1^+} \frac{|x - 3| - 2}{x - 1} \\
 &= \lim_{x \rightarrow 1^+} \frac{3 - x - 2}{x - 1} = -1
 \end{aligned}$$

- ∴ $f'(1) = -1$
- ∴ $f(x)$ is differentiable at $x = 1$.
- If $f(x)$ is differentiable, it has to be continuous.
- ∴ $f(x)$ is continuous and differentiable at $x = 1$.

- 6. $\lim_{x \rightarrow 0^+} f(x) = 0$
- $\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} 0 = 0$
- $\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$
- ∴ The function is continuous at $x = 0$



- Since the function has a sharp edge at $x = 0$,
- ∴ The function is not differentiable.

- 7. $\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x - 1) = 1 - 1 = 0$
- $\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (x^3 - 1) = 1 - 1 = 0$
- $f(1) = 0$
- ∴ $f(x)$ is continuous at $x = 1$.
- $Lf'(1) = \lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{x - 1} = \lim_{x \rightarrow 1^-} \frac{-1 - 0}{-1} = 1$

$$\begin{aligned}
 Rf'(1) &= \lim_{x \rightarrow 1^+} \frac{f(x) - f(1)}{x - 1} \\
 &= \lim_{x \rightarrow 1^+} \frac{x^3 - 1}{x - 1} = \lim_{x \rightarrow 1^+} \frac{x^2 + x + 1}{1} = 3
 \end{aligned}$$

- ∴ $Lf'(1) \neq Rf'(1)$
- ∴ $f(x)$ is not differentiable at $x = 1$.

- 8. Since $f(x)$ is differentiable at $x = 1$.
- ∴ $Lf'(1) = Rf'(1)$

$$\begin{aligned}
 &\Rightarrow \left[\frac{d}{dx} (x^2 + bx + c) \right]_{x=1} = \left[\frac{d}{dx} (x) \right]_{x=1} \\
 &\Rightarrow [2x + b]_{x=1} = 1 \\
 &\Rightarrow 2 + b = 1 \\
 &\Rightarrow b = -1 \quad \dots(i) \\
 &f(x) \text{ is differentiable at } x = 1. \\
 &\Rightarrow f(x) \text{ is continuous at } x = 1.
 \end{aligned}$$



$$\begin{aligned} \therefore f(1) &= \lim_{x \rightarrow 1^-} f(x) \\ \Rightarrow 1 &= \lim_{x \rightarrow 1} (x^2 + b + c) \\ \Rightarrow 1 &= 1 + b + c \\ \Rightarrow b + c &= 0 \\ \Rightarrow c &= 1 \quad \dots[\text{From (i)}] \end{aligned}$$

$$\therefore b - c = -1 - 1 = -2$$

9. $\lim_{x \rightarrow 1} \frac{x^2 f(1) - f(x)}{-1}$
Applying L' Hospital rule, we get
 $\lim_{x \rightarrow 1} 2x f(1) - f'(x) = 2f(1) - f'(1)$

10. Applying L'Hospital rule, we get
 $\lim_{x \rightarrow 2} \frac{f(2) - 2f(x)}{-2} = \lim_{x \rightarrow 2} \frac{f(2) - 2f'(x)}{1}$
 $= f(2) - 2f'(2)$
 $= 4 - 2(1) = 2$

11. Since, $f(x)$ is differentiable at $x = a$.
 $\therefore f'(x)$ exists
Let $\lim_{x \rightarrow a} \frac{f(x) - f(a)}{-a} = f'(a) \quad \dots(i)$


Now, $\lim_{x \rightarrow a} \frac{x^2 f(a) - a^2 f(x)}{-a}$
 $= \lim_{x \rightarrow a} \frac{x^2 f(a) - a^2 f(a) + a^2 f(a) - a^2 f(x)}{-a}$
 $= \lim_{x \rightarrow a} \frac{(x^2 - a^2)f(a) - a^2 \{f(x) - f(a)\}}{-a}$
 $= \lim_{x \rightarrow a} \left[\frac{(x^2 - a^2)f(a)}{-a} - a^2 \left\{ \frac{f(x) - f(a)}{-a} \right\} \right]$
 $= \lim_{x \rightarrow a} (x + a) f(a) - a^2 \lim_{x \rightarrow a} \frac{f(x) - f(a)}{-a}$
 $= 2a f(a) - a^2 f'(a) \quad \dots[\text{From (i)}]$

12. Since, $f(x)$ is differentiable for all x . So, it is everywhere continuous.

$$\begin{aligned} \therefore \lim_{x \rightarrow 1} f(x) &= f(1) \\ \Rightarrow \lim_{h \rightarrow 0} f(1+h) &= f(1) \\ \Rightarrow \lim_{h \rightarrow 0} \frac{f(1+h)}{h} \times h &= f(1) \\ \Rightarrow \lim_{h \rightarrow 0} \frac{f(1+h)}{h} \times \lim_{h \rightarrow 0} h &= f(1) \\ \Rightarrow 5 \times 0 &= f(1) \\ \Rightarrow f(1) &= 0 \end{aligned}$$

Now, $f'(1) = \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h}$
 $\Rightarrow f'(1) = \lim_{h \rightarrow 0} \frac{f(1+h) - 0}{h}$
 $\Rightarrow f'(1) = 5$

13. The continuous line shown in the figure below represents the graph of $f(x)$.



$$\Rightarrow f(x) = \begin{cases} x, & x \leq -1 \\ 3x, & -1 < x \leq 0 \\ x, & 0 < x < 1 \\ 3x, & 1 \leq x \end{cases}$$

Clearly, $f(x)$ is not differentiable at $x = -1, 0, 1$.

14. Let $f(x) = |x - 1| = \begin{cases} -1, & x \geq 1 \\ 1 - x, & x < 1 \end{cases}$
 $\therefore p =$ left hand derivative of $f(x)$ at $x = 1$
 $\Rightarrow p = \lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{-1} = \lim_{x \rightarrow 1^-} \frac{1 - x - 0}{-1} = -1$

Now, $\lim_{x \rightarrow 1^+} g(x) = p$
 $\Rightarrow \lim_{h \rightarrow 0} g(1+h) = -1$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{h^n}{\log \cos^m h} = -1$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{h^n}{m \log \cos h} = -1$$

Applying L'Hospital rule on L.H.S., we get

$$\frac{1}{m} \lim_{h \rightarrow 0} \frac{n h^{n-1}}{-\tan h} = -1$$

$$\Rightarrow \frac{n}{m} \lim_{h \rightarrow 0} \frac{h^{n-2}}{\left(\frac{\tan h}{h}\right)} = 1$$

$$\Rightarrow n = 2 \text{ and } \frac{n}{m} = 1$$

$$\Rightarrow m = n = 2$$



$$15. \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{2 \left| \cos \frac{\pi}{x} \right| - 0}{x}$$

$$= \lim_{x \rightarrow 0} \left| \cos \frac{\pi}{x} \right| = 0$$

So, $f(x)$ is differentiable at $x = 0$.

$$\text{Now, } Rf'(2) = \lim_{h \rightarrow 0^+} \frac{f(2+h) - f(2)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(2+h)^2 \cos\left(\frac{\pi}{2+h}\right) - 0}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(2+h)^2 \sin\left(\frac{\pi}{2} - \frac{\pi}{2+h}\right)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(2+h)^2 \sin\left\{\frac{\pi h}{2(2+h)}\right\}}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\sin\left\{\frac{\pi h}{2(2+h)}\right\}}{\frac{\pi h}{2(2+h)}} \times \frac{(2+h)\pi}{2} = \pi$$

Similarly, $Lf'(2) = -\pi$

$\therefore Lf'(2) \neq Rf'(2)$

So, $f(x)$ is not differentiable at $x = 2$.

$$16. Lg'(3) = \lim_{x \rightarrow 3^-} \frac{g(x) - g(3)}{x - 3} = \lim_{x \rightarrow 3^-} \frac{k\sqrt{x+1} - 2k}{x - 3}$$

$$= \lim_{x \rightarrow 3^-} k \left[\frac{x+1-4}{(x-3)(\sqrt{x+1}+2)} \right]$$

$$= \lim_{x \rightarrow 3^-} \frac{k}{\sqrt{x+1}+2} = \frac{k}{4}$$

$$Rg'(3) = \lim_{x \rightarrow 3^+} \frac{g(x) - g(3)}{x - 3} = \lim_{x \rightarrow 3^+} \frac{m + 2 - 2k}{x - 3}$$

Applying L'Hospital rule, we get

$$Rg'(3) = m$$

Since, $g'(3)$ exists.

$\therefore Rg'(3)$ must exist.

$$\therefore 3m + 2 - 2k = 0 \quad \dots(i)$$

Since, $g(x)$ is differentiable.

$\therefore Lg'(3) = Rg'(3)$

$$\therefore \frac{k}{4} = m \Rightarrow k = 4m \quad \dots(ii)$$

Solving (i) and (ii), we get

$$m = \frac{2}{5} \text{ and } k = \frac{8}{5}$$

$$\therefore k + m = \frac{8}{5} + \frac{2}{5} = 2$$

17. **Differentiability at $x = \pi$:**

$$Ls'(\pi)$$

$$= \lim_{h \rightarrow 0} \frac{|\pi - h - \pi| \left(e^{|\pi-h|} - 1 \right) \sin|\pi - h| - 0}{-h}$$

$$= 0$$

$$Rs'(\pi)$$

$$= \lim_{h \rightarrow 0} \frac{|\pi + h - \pi| \left(e^{|\pi+h|} - 1 \right) \sin|\pi + h| - 0}{h}$$

$$= 0$$

Differentiability at $x = 0$:

$$Ls'(0) = \lim_{h \rightarrow 0} \frac{|-h - \pi| \left(e^{|-h|} - 1 \right) \sin|-h| - 0}{-h}$$

$$= 0$$

$$Rs'(0) = \lim_{h \rightarrow 0} \frac{|h - \pi| \left(e^{|h|} - 1 \right) \sin|h| - 0}{h}$$

$$= 0$$

The function $f(x)$ is differentiable at $x = 0, \pi$.
 \Rightarrow Set S is an empty set.

$$18. \quad = \cos(2 + 45)$$

$$\therefore \frac{d}{dx} = -\sin(2 + 45) \cdot \frac{d}{dx}(2 + 45)$$

$$= -2 \sin(2 + 45)$$

$$19. \quad = \sqrt{\sin \sqrt{x}}$$

$$\therefore \frac{d}{dx} = \frac{1}{2\sqrt{\sin \sqrt{x}}} \cdot \frac{d}{dx}(\sin \sqrt{x})$$

$$= \frac{1}{2\sqrt{\sin \sqrt{x}}} \cdot \cos \sqrt{x} \cdot \frac{1}{2\sqrt{x}}$$

$$= \frac{\cos \sqrt{x}}{4\sqrt{x} \sqrt{\sin \sqrt{x}}}$$

$$20. \quad \frac{d}{dx} \log_{|e|} |e| = \frac{d}{dx} \left(\frac{1}{|\log ||} \right)$$

$$= \frac{-1}{\log^2 ||} \times \frac{1}{|} = \frac{-1}{(\log ||)^2}$$



$$\begin{aligned}
 21. \quad f(x) &= \log(\log x) \\
 \therefore f[\log x] &= \log(\log(\log x)) \\
 \therefore f'[\log x] &= \frac{1}{\log x} \cdot \frac{d}{dx}(\log x) \\
 &= \frac{1}{\log x}
 \end{aligned}$$

$$\begin{aligned}
 22. \quad f(x) &= \log_2(\log_2 x) \\
 &= \frac{\log\left(\frac{\log x}{\log 2}\right)}{\log 2} \\
 &= \frac{\log(\log x) - \log(\log 2)}{\log 2}
 \end{aligned}$$

$$\begin{aligned}
 \therefore \frac{d}{dx} &= \frac{1}{\log 2} \left(\frac{1}{\log x} \cdot \frac{1}{x} - 0 \right) \\
 &= \frac{1}{(x \log x) \log 2}
 \end{aligned}$$

$$\begin{aligned}
 23. \quad f(x) &= \log\left(\frac{1-x^2}{1+x^2}\right) \\
 \therefore \frac{d}{dx} &= \frac{1}{\left(\frac{1-x^2}{1+x^2}\right)} \cdot \frac{(1+x^2)(0-2x) - (1-x^2)(0+2x)}{(1+x^2)^2} \\
 &= \frac{1}{(1-x^2)} \cdot \frac{-2x - 2x^3 - 2x + 2x^3}{(1+x^2)^2}
 \end{aligned}$$

$$\therefore \frac{d}{dx} = \frac{-4x}{1-x^4}$$

$$\begin{aligned}
 24. \quad \frac{d}{dx}[\cos(1-x^2)^2] &= -\sin(1-x^2)^2 \cdot \frac{d}{dx}[(1-x^2)^2] \\
 &= -\sin(1-x^2)^2 \cdot 2(1-x^2) \cdot \frac{d}{dx}(1-x^2) \\
 &= -\sin(1-x^2)^2 \cdot 2(1-x^2) \cdot (-2x) \\
 &= 4x(1-x^2)\sin(1-x^2)^2
 \end{aligned}$$

$$\begin{aligned}
 25. \quad \frac{d}{dx}[e^{\log(1+x^2)}] &= \log(1+x^2) \cdot e + e \cdot \frac{1}{1+x^2} \cdot \frac{d}{dx}(1+x^2) \\
 &= e \log(1+x^2) + \frac{e}{1+x^2} \cdot 2x \\
 &= e \left[\log(1+x^2) + \frac{2x}{1+x^2} \right]
 \end{aligned}$$

$$\begin{aligned}
 26. \quad \frac{d}{dx}(e^{\log \sin 2x}) &= \log \sin 2x \cdot e + e \cdot \frac{1}{\sin 2x} \cdot \frac{d}{dx}(\sin 2x) \\
 &= e \log \sin 2x + e \cdot \frac{1}{\sin 2x} \cdot \cos 2x \cdot \frac{d}{dx}(2x) \\
 &= e \log \sin 2x + e \cot 2x \cdot 2 \\
 &= e (\log \sin 2x + 2 \cot 2x)
 \end{aligned}$$

$$\begin{aligned}
 27. \quad \frac{d}{dx}(e^{\sqrt{1-x^2}} \cdot \tan^{-1} x) &= e^{\sqrt{1-x^2}} \cdot \sec^2 x + \tan^{-1} x \cdot e^{\sqrt{1-x^2}} \cdot \frac{d}{dx}(\sqrt{1-x^2}) \\
 &= e^{\sqrt{1-x^2}} \cdot \sec^2 x + \tan^{-1} x \cdot e^{\sqrt{1-x^2}} \cdot \frac{1}{2\sqrt{1-x^2}} \cdot (-2x) \\
 &= e^{\sqrt{1-x^2}} \left[\sec^2 x - \frac{x \tan^{-1} x}{\sqrt{1-x^2}} \right]
 \end{aligned}$$

$$\begin{aligned}
 28. \quad f(x) &= \frac{e^x + e^{-x}}{e^x - e^{-x}} \\
 \therefore \frac{d}{dx} &= \frac{1}{(e^x - e^{-x})^2} \left[(e^x - e^{-x}) \cdot 2(e^x - e^{-x}) - (e^x + e^{-x}) \cdot 2(e^x + e^{-x}) \right] \\
 &= \frac{-8}{(e^x - e^{-x})^2}
 \end{aligned}$$

$$\begin{aligned}
 29. \quad f(x) &= \log e^{(\tan^{-1} x)^2} \\
 \therefore \frac{d}{dx} &= e^{(\tan^{-1} x)^2} \cdot \frac{1}{e^{(\tan^{-1} x)^2}} + \log e^{(\tan^{-1} x)^2} \cdot \frac{d}{dx}(\tan^{-1} x)^2 \\
 &= e^{(\tan^{-1} x)^2} \cdot \frac{1}{e^{(\tan^{-1} x)^2}} + \log e^{(\tan^{-1} x)^2} (\sec^2 x + 2) \\
 &= e^{(\tan^{-1} x)^2} \left[\frac{1}{e^{(\tan^{-1} x)^2}} + (\sec^2 x + 2) \log e \right]
 \end{aligned}$$

$$\begin{aligned}
 30. \quad H(x) &= G[F(x)] \\
 &= e^{-e^x} \\
 \therefore H'(x) &= -e^{-e^x} \cdot e^x \\
 \therefore H'(0) &= -e^0 \cdot e^{-e^0} \\
 &= -e^{-1} \\
 &= -\frac{1}{e}
 \end{aligned}$$



31. $h(x) = f(g(x))$
 $\Rightarrow h(x) = f(\sin^{-1} x) = e^{\sin^{-1} x}$... (i)
 $\therefore h'(x) = e^{\sin^{-1} x} \cdot \frac{d}{dx}(\sin^{-1} x) = e^{\sin^{-1} x} \cdot \frac{1}{\sqrt{1-x^2}}$
 $\therefore \frac{h'(x)}{h(x)} = \frac{1}{\sqrt{1-x^2}}$... [From (i)]
32. At $x = 1$, $f(x)$ is not defined.
 For $x \in \mathbb{R} - \{1\}$,
 $g(x) = f[f\{f(x)\}] = f\left[f\left(\frac{1}{1-x}\right)\right] = f\left(\frac{1}{1-\frac{1}{1-x}}\right)$
 $= f\left(\frac{-1}{1-x-1}\right) = \frac{1}{1-x-1} =$
 $\therefore g'(x) = 1$ for all $x \in \mathbb{R} - \{1\}$
33. Let $t = \frac{2-x-1}{x^2+1}$. Then, $y = f(t)$
 $\therefore \frac{dy}{dx} = f'(t) \cdot \frac{dt}{dx} = \sin t^2 \cdot \frac{d}{dx}\left(\frac{2-x-1}{x^2+1}\right)$
 ... [$\because f'(x) = \sin^2(x)$ (given)]
 $= \sin^2 t^2 \left[\frac{(x^2+1)(2-x-1) - (2-x-1)(2x+2)}{(x^2+1)^2} \right]$
 $= \frac{-2x^2+2x+2}{(x^2+1)^2} \cdot \sin^2\left(\frac{2-x-1}{x^2+1}\right)$
34. $f^{-1}(x) = g(x)$
 $\Rightarrow x = f[g(x)]$
 Differentiating w.r.t. x , we get
 $f'[g(x)] \cdot g'(x) = 1$
 $\Rightarrow \frac{1}{1+[g(x)]^4} \cdot g'(x) = 1$... [$\because f'(x) = \frac{1}{1+x^4}$]
 $\Rightarrow g'(x) = 1 + [g(x)]^4$
35. $g(x) = [f(2f(x)+2)]^2$
 $\therefore g'(x) = 2[f(2f(x)+2)] \cdot [f(2f(x)+2)]'$
 $= 2[(2f(x)+2)] f'[2f(x)+2] \cdot 2f'(x)$
 $\therefore g'(0) = 2[f(-2+2)] f'[2f(0)+2] \cdot 2(1)$
 $= 2[f(0)](1) \cdot 2$
 $= 4(-1)$
 $= -4$

36. $= 5(1-x)^{-\frac{2}{3}} + \cos^2(2x+1)$
 $\therefore \frac{d}{dx} = 5 \cdot \frac{-2}{3}(1-x)^{-\frac{5}{3}} \cdot \frac{d}{dx}(1-x) + 5(1-x)^{-\frac{2}{3}}$
 $+ 2\cos(2x+1) \cdot \frac{d}{dx}[\cos(2x+1)]$
 $= \frac{10}{3(1-x)^{\frac{5}{3}}} + \frac{5}{(1-x)^{\frac{2}{3}}}$
 $- 2[2\cos(2x+1)\sin(2x+1)]$
 $= \frac{5}{(1-x)^{\frac{2}{3}}} \left[\frac{2}{3(1-x)} + 1 \right] - 2\sin(4x+2)$
 ... [$\because 2\sin\theta\cos\theta = \sin 2\theta$]
 $= \frac{5(3-x)}{3(1-x)^{\frac{5}{3}}} - 2\sin(4x+2)$
37. $y = f(x^2+2)$
 $\therefore \frac{dy}{dx} = f'(x^2+2) \cdot (2x)$
 $\therefore \left(\frac{dy}{dx}\right)_{x=1} = f'(1^2+2) \cdot (2 \times 1)$
 $= f'(3) \cdot 2 = 5 \cdot 2 = 10$
38. $f(x) = \log(\log x) = \frac{\log(\log x)}{\log}$
 $\therefore f'(x) = \frac{\log \cdot \frac{1}{\log} \cdot \frac{d}{dx}(\log x) - \log(\log x) \cdot \frac{1}{\log}}{(\log x)^2}$
 $= \frac{\frac{1}{\log} - \frac{1}{\log} \log(\log x)}{(\log x)^2}$
 $\therefore f'(e) = \frac{\frac{1}{e} - 0}{(1)^2} = \frac{1}{e}$
39. $f(x) = \sqrt{1+\cos^2(x^2)}$
 $\therefore f'(x) = \frac{1}{2\sqrt{1+\cos^2(x^2)}} \cdot (2\cos^2(x^2)) \cdot (-\sin^2(x^2)) \cdot (2x)$
 $\therefore f'(x) = \frac{-\sin^2(x^2)}{\sqrt{1+\cos^2(x^2)}}$
 $\therefore f'\left(\frac{\sqrt{\pi}}{2}\right) = \frac{-\frac{\sqrt{\pi}}{2} \cdot \sin \frac{2\pi}{4}}{\sqrt{1+\cos^2 \frac{\pi}{4}}} = \frac{-\frac{\sqrt{\pi}}{2} \cdot 1}{\sqrt{\frac{3}{2}}} = -\sqrt{\frac{\pi}{6}}$



$$\begin{aligned}
 40. \quad f(x) &= \frac{\sin^2}{1 + \cot} + \frac{\cos^2}{1 + \tan} \\
 &= \frac{\sin^2 (\sin)}{\sin + \cos} + \frac{\cos^2 (\cos)}{\cos + \sin} \\
 &= \frac{\sin^3 + \cos^3}{\sin + \cos} \\
 &= \sin^2 - \sin \cos + \cos^2 \\
 &\quad \dots [\because a^3 + b^3 = (a + b)(a^2 - ab + b^2)]
 \end{aligned}$$

$$\begin{aligned}
 &= (\sin^2 + \cos^2) - \frac{1}{2}(2 \sin \cos) \\
 &= 1 - \frac{1}{2} \sin 2
 \end{aligned}$$

$$\therefore f'(x) = -\cos 2 \Rightarrow f'\left(\frac{\pi}{4}\right) = -\cos\left(\frac{\pi}{2}\right) = 0$$

$$\begin{aligned}
 41. \quad \frac{d}{dx} \tan^{-1}\left(\frac{1-x}{1+x}\right) &= \frac{d}{dx} [\tan^{-1}(1) - \tan^{-1}(x)] \\
 &= 0 - \frac{1}{1+x^2} = \frac{-1}{1+x^2}
 \end{aligned}$$

$$\begin{aligned}
 42. \quad &= \tan^{-1}\left(\frac{\sqrt{a}-\sqrt{a}}{1+\sqrt{a}}\right) = \tan^{-1} \sqrt{a} - \tan^{-1} \sqrt{a} \\
 \therefore \frac{d}{dx} &= 0 - \frac{1}{1+(\sqrt{a})^2} \cdot \frac{d}{dx}(\sqrt{a}) = -\frac{1}{(1+a)} \cdot \frac{1}{2\sqrt{a}}
 \end{aligned}$$

$$\begin{aligned}
 43. \quad &= \tan^{-1}\left(\frac{\sin + \cos}{\cos - \sin}\right) = \tan^{-1}\left(\frac{1 + \tan}{1 - \tan}\right) \\
 &= \tan^{-1}\left[\tan\left(\frac{\pi}{4} + \right)\right] = \frac{\pi}{4} +
 \end{aligned}$$

$$\therefore \frac{d}{dx} = 1$$

$$\begin{aligned}
 44. \quad &= \tan^{-1}\left(\frac{a \cos - b \sin}{b \cos + a \sin}\right) \\
 &= \tan^{-1}\left(\frac{\frac{a}{b} - \tan}{1 + \frac{a}{b} \tan}\right) \\
 &= \tan^{-1}\left(\frac{a}{b}\right) - \tan^{-1}(\tan)
 \end{aligned}$$

$$\therefore = \tan^{-1}\left(\frac{a}{b}\right) - \quad \therefore \frac{d}{dx} = -1$$

$$\begin{aligned}
 45. \quad &= \sec(\tan^{-1} x) \\
 \therefore \frac{d}{dx} &= \sec(\tan^{-1} x) \tan(\tan^{-1} x) \cdot \frac{1}{1+x^2} \\
 &= \sqrt{1+x^2} \cdot \frac{1}{1+x^2} \\
 &\quad \dots [\because \tan^{-1} x = \sec^{-1} \sqrt{1+x^2}] \\
 &= \frac{1}{\sqrt{1+x^2}}
 \end{aligned}$$

$$\begin{aligned}
 46. \quad \text{Let } x &= \tan^{-1}\left(\frac{6\sqrt{x}}{1-9x^3}\right) = \tan^{-1}\left[\frac{6x^{\frac{3}{2}}}{1-\left(3x^{\frac{3}{2}}\right)^2}\right] \\
 &= \tan^{-1}\left[\frac{2 \times 3x^{\frac{3}{2}}}{1-\left(3x^{\frac{3}{2}}\right)^2}\right] \\
 &= 2 \tan^{-1} 3x^{\frac{3}{2}}
 \end{aligned}$$

$$\therefore \frac{d}{dx} = \frac{2}{1+\left(3x^{\frac{3}{2}}\right)^2} \cdot 3 \times \frac{3}{2} x^{\frac{1}{2}} = \frac{9}{1+9x^3} \sqrt{x}$$

Comparing with $\sqrt{x} g(x)$, we get

$$g(x) = \frac{9}{1+9x^3}$$

$$47. \quad = e^{m \sin^{-1} x} \quad \dots (i)$$

$$\therefore \frac{d}{dx} = e^{m \sin^{-1} x} \cdot \frac{m}{\sqrt{1-x^2}}$$

$$\Rightarrow \sqrt{1-x^2} \frac{d}{dx} = m \quad \dots [\text{From (i)}]$$

$$\Rightarrow (1-x^2) \left(\frac{d}{dx}\right)^2 = m^2 - 2$$

$$\therefore A = m^2$$

$$\begin{aligned}
 48. \quad \text{Putting } x &= \sin A \text{ and } \sqrt{x} = \sin B, \text{ we get} \\
 &= \sin^{-1}(\sin A \sqrt{1-\sin^2 B} + \sin B \sqrt{1-\sin^2 A}) \\
 &= \sin^{-1}(\sin A \cos B + \sin B \cos A) \\
 &= \sin^{-1}[\sin(A+B)] = A + B = \sin^{-1} x + \sin^{-1} \sqrt{x}
 \end{aligned}$$



$$\begin{aligned} \therefore \frac{d}{d} &= \frac{1}{\sqrt{1-x^2}} + \frac{1}{\sqrt{1-(\sqrt{x})^2}} \cdot \frac{1}{2\sqrt{x}} \\ &= \frac{1}{\sqrt{1-x^2}} + \frac{1}{2\sqrt{x-x^2}} \end{aligned}$$

49. $= \tan^{-1} \left(\frac{1}{1+\sqrt{1-x^2}} \right) + \sin \left(2 \tan^{-1} \sqrt{\frac{1-x}{1+x}} \right)$

Put $= \cos \theta \Rightarrow \theta = \cos^{-1}$

$$\begin{aligned} \therefore &= \tan^{-1} \left(\frac{\cos \theta}{1+\sin \theta} \right) + \sin \left(2 \tan^{-1} \sqrt{\frac{1-\cos \theta}{1+\cos \theta}} \right) \\ &= \tan^{-1} \left(\frac{1-\tan \frac{\theta}{2}}{1+\tan \frac{\theta}{2}} \right) + \sin \left(2 \tan^{-1} \sqrt{\frac{2 \sin^2 \frac{\theta}{2}}{2 \cos^2 \frac{\theta}{2}}} \right) \\ &= \tan^{-1} \left[\tan \left(\frac{\pi}{4} - \frac{\theta}{2} \right) \right] + \sin \left[2 \tan^{-1} \left(\tan \frac{\theta}{2} \right) \right] \\ &= \frac{\pi}{4} - \frac{\theta}{2} + \sin \theta \\ &= \frac{\pi}{4} - \frac{\cos^{-1}}{2} + \sin(\cos^{-1}) \\ &= \frac{\pi}{4} - \frac{\cos^{-1}}{2} + \sin \left(\sin^{-1} \sqrt{1-x^2} \right) \\ &= \frac{\pi}{4} - \frac{\cos^{-1}}{2} + \sqrt{1-x^2} \end{aligned}$$

$$\begin{aligned} \therefore \frac{d}{d} &= \frac{1}{2\sqrt{1-x^2}} + \frac{(-2)}{2\sqrt{1-x^2}} \\ &= \frac{1-2}{2\sqrt{1-x^2}} \end{aligned}$$

50. Put $= \tan \theta \Rightarrow \theta = \tan^{-1} ()$

$$\begin{aligned} \therefore f() &= \cot^{-1} \left(\frac{\tan^2 \theta - 1}{2 \tan \theta} \right) \\ &= \cot^{-1} (-\cot 2\theta) \\ &= \pi - \cot^{-1}(\cot 2\theta) \\ \therefore f() &= \pi - 2\theta = \pi - 2 \tan^{-1}() \\ \therefore f'() &= \frac{-2}{1+ } \cdot (1 + \log) \\ \therefore f'(1) &= \frac{-2}{1+1^2} \cdot 1(1+0) = -1 \end{aligned}$$

51. $= \tan^{-1} \left(\frac{\sqrt{x-x^3}}{1+\frac{x}{2}} \right) = \tan^{-1}(\sqrt{x}) - \tan^{-1} ()$

$$\therefore ' = \frac{1}{1+ } \cdot \frac{1}{2\sqrt{x}} - \frac{1}{1+x^2}$$

$$\therefore '(1) = \frac{1}{2} \cdot \frac{1}{2} - \frac{1}{2} = \frac{-1}{4}$$

52. $= \tan^{-1} \left(\frac{\sqrt{1+x^2}-1}{ } \right)$

Put $= \tan \theta \Rightarrow \theta = \tan^{-1}$

$$\therefore = \tan^{-1} \left(\frac{\sqrt{1+\tan^2 \theta} - 1}{\tan \theta} \right) = \tan^{-1} \left(\frac{\sec \theta - 1}{\tan \theta} \right)$$

$$= \tan^{-1} \left(\frac{1-\cos \theta}{\sin \theta} \right)$$

$$= \tan^{-1} \left(\frac{2 \sin^2 \frac{\theta}{2}}{2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}} \right)$$

$$= \tan^{-1} \left(\tan \frac{\theta}{2} \right) = \frac{\theta}{2} = \frac{1}{2} \tan^{-1}$$

$$\therefore ' = \frac{1}{2(1+x^2)}$$

$$\therefore '(1) = \frac{1}{2(1+1^2)} = \frac{1}{4}$$

53. $= \left(1 + \frac{1}{ } \right)$

Taking logarithm on both sides, we get

$$\log = \log \left(1 + \frac{1}{ } \right)$$

Differentiating w.r.t. , we get

$$\frac{1}{ } \cdot \frac{d}{d} = \log \left(1 + \frac{1}{ } \right) + \frac{1}{1+ } \cdot \left(-\frac{1}{2} \right)$$

$$\Rightarrow \frac{1}{ } \cdot \frac{d}{d} = \log \left(1 + \frac{1}{ } \right) - \frac{1}{1+ }$$

$$\Rightarrow \frac{d}{d} = \left(1 + \frac{1}{ } \right) \left[\log \left(1 + \frac{1}{ } \right) - \frac{1}{1+ } \right]$$



$$54. \quad = (\sin^{-1} x)^{\tan^{-1} x}$$

Taking logarithm on both sides, we get
 $\log = \tan^{-1} x \cdot \log(\sin^{-1} x)$

Differentiating w.r.t. x , we get

$$\frac{1}{x} \cdot \frac{d}{dx} = \tan^{-1} x \cdot \cot^{-1} x + \log(\sin^{-1} x) \cdot \sec^2 x$$

$$\Rightarrow \frac{d}{dx} = (\sin^{-1} x)^{\tan^{-1} x} [1 + \sec^2 x \log(\sin^{-1} x)]$$

$$55. \quad = \frac{e^2 \cos x}{\sin x}$$

Taking logarithm on both sides, we get

$$\log = 2 + \log(\cos x) - \log \sin x$$

Differentiating w.r.t. x , we get

$$\frac{1}{x} \cdot \frac{d}{dx} = 2 + \left(\frac{-\sin x}{\cos x} \right) - \frac{1}{\sin x} \cdot \cos x$$

$$\Rightarrow \frac{d}{dx} = \frac{e^2 \cos x}{\sin x} \left(2 - \frac{\sin x}{\cos x} - \frac{1}{\sin x} \cdot \cos x \right)$$

$$= e^2 \left(2 \cot x - \frac{1}{\sin x} - \frac{1}{2} \cot x - \frac{\cot^2 x}{2} \right)$$

$$= \frac{e^2}{2} [2 \cot x - \cot x - (1 + \cot^2 x)]$$

$$= \frac{e^2}{2} [(2 - 1) \cot x - \operatorname{cosec}^2 x]$$

$$56. \quad = \{f(x)\}^{\phi(x)}$$

Taking logarithm on both sides, we get

$$\log = \phi(x) \log \{f(x)\}$$

$$\Rightarrow = e^{\phi(x) \log f(x)}$$

$$\therefore \frac{d}{dx} = e^{\phi(x) \log f(x)} \cdot \frac{d}{dx} [\phi(x) \log f(x)]$$

$$= e^{\phi(x) \log f(x)} \left\{ \phi(x) \cdot \frac{f'(x)}{f(x)} + \log f(x) \cdot \phi'(x) \right\}$$

$$57. \quad = (\log x)^{\log(\log x)}$$

Taking logarithm on both sides, we get

$$\log = \log(\log x) [\log x + \log(\log x)]$$

Differentiating w.r.t. x , we get

$$\frac{1}{x} \cdot \frac{d}{dx} = \frac{1}{\log x} [\log x + \log(\log x)]$$

$$+ \log(\log x) \left(\frac{1}{x} + \frac{1}{\log x} \right)$$

$$\Rightarrow \frac{d}{dx} = (\log x)^{\log(\log x)} \left\{ \frac{1}{\log x} [\log x + \log(\log x)] \right.$$

$$\left. + \log(\log x) \left(\frac{1}{x} + \frac{1}{\log x} \right) \right\}$$

$$58. \quad = [(\tan^{-1} x)^{\tan^{-1} x}]^{\tan^{-1} x}$$

Taking logarithm on both sides, we get

$$\log = \tan^{-1} x \log(\tan^{-1} x)^{\tan^{-1} x}$$

$$\Rightarrow \log = (\tan^{-1} x)^2 \log(\tan^{-1} x)$$

Differentiating w.r.t. x , we get

$$\frac{1}{x} \cdot \frac{d}{dx} = (\tan^{-1} x)^2 \cdot \frac{1}{\tan x} \cdot \sec^2 x$$

$$+ \log(\tan^{-1} x) \cdot 2 \tan^{-1} x \cdot \sec^2 x$$

$$\Rightarrow \frac{d}{dx} = [(\tan^{-1} x)^{\tan^{-1} x}]^{\tan^{-1} x} \cdot \tan^{-1} x \sec^2 x [1 + 2 \log(\tan^{-1} x)]$$

$$\therefore \left(\frac{d}{dx} \right)_{\left(x = \frac{\pi}{4} \right)} = 1.1 \cdot (\sqrt{2})^2 (1 + 0) = 2$$

$$59. \quad = 1 + e^{-x} \quad \dots (i)$$

$$\therefore \frac{d}{dx} = e^{-x} \cdot 1 + (-e^{-x}) \cdot \frac{d}{dx} x$$

$$\Rightarrow (1 - e^{-x}) \frac{d}{dx} = e^{-x}$$

$$\Rightarrow (2 - e^{-x}) \frac{d}{dx} = e^{-x} \quad \dots [\text{From (i)}]$$

$$\Rightarrow \frac{d}{dx} = \frac{e^{-x}}{2 - e^{-x}}$$

$$60. \quad = 1 + \log x$$

Differentiating both sides w.r.t. x , we get

$$\frac{d}{dx} + \frac{1}{x} = \frac{1}{x} \cdot \frac{d}{dx} x$$

$$\Rightarrow (x - 1) \frac{d}{dx} + 1 = 0$$

$$\therefore k = -1$$

$$61. \quad \tan^{-1} (x^2 + 2) = \alpha$$

$$\Rightarrow x^2 + 2 = \tan \alpha$$

Differentiating both sides w.r.t. x , we get

$$2x + 2 \frac{d}{dx} = 0 \Rightarrow \frac{d}{dx} = -\frac{x}{x+1}$$

$$62. \quad e^{\sin^{-1}(t^2-1)} \text{ and } = e^{\sec^{-1}\left(\frac{1}{t^2-1}\right)} = e^{\cos^{-1}(t^2-1)}$$

$$\therefore = e^{\frac{\pi}{2}} \quad \dots \left[\because \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2} \right]$$

Differentiating both sides w.r.t. t , we get

$$\frac{d}{dt} + \frac{1}{t} = 0$$

$$\Rightarrow \frac{d}{dt} = -\frac{1}{t}$$



63. $2^2 - 3^2 + 2^2 + 2^2 - 8 = 0$
 Differentiating w.r.t. x , we get
 $4 \cdot 2 \left(\frac{d}{dx} \right) + 2 \left(\frac{d}{dx} \right) + 1 + 2 \frac{d}{dx} = 0$
 $\Rightarrow (-3 + 2 + 2) \frac{d}{dx} + 4 - 3 + 1 = 0$
 $\Rightarrow \frac{d}{dx} = \frac{3 - 4 - 1}{2 - 3 + 2}$

64. $\sec^2 x + \tan^2 x = 0$
 Differentiating w.r.t. x , we get
 $\sec^2 x \frac{d}{dx} + \sec^2 x \tan x + \sec^2 x + 2 \tan x \frac{d}{dx} = 0$
 $\Rightarrow \frac{d}{dx} = -\frac{2 \tan x + \sec^2 x + \sec^2 x \tan x}{2 + \sec^2 x}$

65. If $y = \sqrt{f(x)} + \sqrt{f(x)} + \sqrt{f(x)} + \dots \infty$,
 then $\frac{dy}{dx} = \frac{f'(x)}{2\sqrt{f(x)}}$
 $\therefore \frac{dy}{dx} = \frac{1}{2\sqrt{f(x)}}$

66. $y = e^{-x}$
 Taking logarithm on both sides, we get
 $\log y = -x$
 $\Rightarrow \frac{1}{y} \frac{dy}{dx} = -1$
 $\therefore \frac{dy}{dx} = \frac{(1 + \log y) \cdot 1 - \left(0 + \frac{1}{y}\right)}{(1 + \log y)^2}$
 $= \frac{\log y}{(1 + \log y)^2}$

67. $y = x^p + x^q = (x^p + x^q)^{p+q}$
 Taking logarithm on both sides, we get
 $p \log x + q \log x = (p + q) \log(x^p + x^q)$
 Differentiating both sides w.r.t. x , we get
 $\frac{p}{x} + \frac{q}{x} \cdot \frac{d}{dx} = \frac{p+q}{x^p + x^q} \left(1 + \frac{d}{dx}\right)$
 $\Rightarrow \frac{d}{dx} = -$

68. $y = \sin^{-1} x$
 Taking logarithm on both sides, we get
 $\log y = \log \sin^{-1} x + \log(\sin^{-1} x)$
 Differentiating both sides w.r.t. x , we get
 $\frac{1}{y} \cdot \frac{d}{dx} + \log \frac{d}{dx} = \frac{1}{\sin^{-1} x} \cdot \cos x \cdot \frac{d}{dx}$
 $\Rightarrow \frac{d}{dx} (1 + \log \sin^{-1} x - \cot x) = \frac{1}{\sin^{-1} x}$
 $\Rightarrow \frac{d}{dx} = \frac{1}{(1 + \log \sin^{-1} x - \cot x)}$

69. $\log_{10} \left(\frac{2^x - 2^{-x}}{2^x + 2^{-x}} \right) = 2$
 $\Rightarrow \frac{2^x - 2^{-x}}{2^x + 2^{-x}} = 10^2$
 $\Rightarrow 2^x - 2^{-x} = 100 \cdot 2^x + 100 \cdot 2^{-x}$
 $\Rightarrow 99 \cdot 2^x + 101 \cdot 2^{-x} = 0$
 Differentiating w.r.t. x , we get
 $99(2^x) + 101 \left(2 \frac{d}{dx} \right) = 0$
 $\Rightarrow \frac{d}{dx} = -\frac{99}{101}$

70. $\log_{10} \left(\frac{3^x - 3^{-x}}{3^x + 3^{-x}} \right) = 2$
 $\Rightarrow \frac{3^x - 3^{-x}}{3^x + 3^{-x}} = 10^2$
 $\Rightarrow 3^x - 3^{-x} = 100 \cdot 3^x + 100 \cdot 3^{-x}$
 $\Rightarrow 99 \cdot 3^x = -101 \cdot 3^{-x} \dots (i)$
 Differentiating w.r.t. x , we get
 $99(3^x) = -101(3^{-x}) \frac{d}{dx}$
 $\Rightarrow \frac{d}{dx} = \frac{-99 \cdot 3^x}{101 \cdot 3^{-x}}$
 $\Rightarrow \frac{d}{dx} = \left(\frac{101 \cdot 3^x}{101 \cdot 3^{-x}} \right) \times \frac{1}{101 \cdot 3^{-x}} \dots [\text{From (i)}]$
 $\Rightarrow \frac{d}{dx} = -$

71. $\cos^{-1} \left(\frac{2^x - 2^{-x}}{2^x + 2^{-x}} \right) = \log a$
 $\therefore \frac{2^x - 2^{-x}}{2^x + 2^{-x}} = \cos(\log a)$



Differentiating both sides w.r.t. x , we get

$$\frac{(x^2 + 2) \left(2 - 2 \frac{d}{dx} \right) - (x^2 - 2) \left(2 + 2 \frac{d}{dx} \right)}{(x^2 + 2)^2} = 0$$

$$\Rightarrow (x^2 + 2) \left(2 - 2 \frac{d}{dx} \right) - (x^2 - 2) \left(2 + 2 \frac{d}{dx} \right) = 0$$

$$\Rightarrow 4x^2 - 4x^2 \frac{d}{dx} = 0$$

$$\Rightarrow 4x^2 = 4x^2 \frac{d}{dx}$$

$$\Rightarrow \frac{d}{dx} = 1$$

72. $\sin x = \sin(a + x)$

$$\Rightarrow \frac{d}{dx} \sin x = \frac{d}{dx} \sin(a + x)$$

Differentiating both sides w.r.t. x , we get

$$1 = \frac{\sin(a + x) \cdot \cos x \frac{d}{dx} - \sin x \cdot \cos(a + x) \frac{d}{dx}}{\sin^2(a + x)}$$

$$\Rightarrow 1 = \frac{\frac{d}{dx} \cdot \sin(a + x)}{\sin^2(a + x)}$$

$$\Rightarrow \frac{d}{dx} = \frac{\sin^2(a + x)}{\sin a}$$

73. $\cos x = \cos(a + x)$

$$\Rightarrow \frac{d}{dx} \cos x = \frac{d}{dx} \cos(a + x)$$

Differentiating both sides w.r.t. x , we get

$$1 = \frac{-\cos(a + x) \sin x \frac{d}{dx} + \cos x \sin(a + x) \frac{d}{dx}}{\cos^2(a + x)}$$

$$\Rightarrow 1 = \frac{\frac{d}{dx} \sin(a + x)}{\cos^2(a + x)}$$

$$\Rightarrow \frac{d}{dx} = \frac{\cos^2(a + x)}{\sin a}$$

74. $\sin(x) + \cos(x) = 2 - x$

Differentiating both sides w.r.t. x , we get

$$\cos(x) \left[1 + \frac{d}{dx} \right] + \left(-\frac{1}{2} \right) \frac{d}{dx} + 1 = 2 - \frac{d}{dx}$$

$$\Rightarrow \left[\cos(x) \left(1 + \frac{d}{dx} \right) - \frac{1}{2} \frac{d}{dx} + 1 \right] \frac{d}{dx} = 2 - \frac{d}{dx} - \cos(x)$$

$$\Rightarrow \frac{d}{dx} = \frac{[2 - \cos(x) - 1] \frac{d}{dx}}{2 \cos(x) + \frac{d}{dx} - 1}$$

75. $\sqrt{x^2 + 1} = \log(\sqrt{x^2 + 1} - x)$

Differentiating both sides w.r.t. x , we get

$$\frac{d}{dx} \sqrt{x^2 + 1} + \frac{1}{2\sqrt{x^2 + 1}} \cdot 2x = \frac{1}{\sqrt{x^2 + 1} - x} \times \left(\frac{1}{2\sqrt{x^2 + 1}} \cdot 2x - 1 \right)$$

$$\Rightarrow \sqrt{x^2 + 1} \cdot \frac{d}{dx} + \frac{x}{\sqrt{x^2 + 1}}$$

$$= \frac{1}{\sqrt{x^2 + 1} - x} \times \frac{-\sqrt{x^2 + 1}}{\sqrt{x^2 + 1}}$$

$$\Rightarrow (x^2 + 1) \frac{d}{dx} + \frac{x}{\sqrt{x^2 + 1}} = \sqrt{x^2 + 1} \cdot \frac{-1}{\sqrt{x^2 + 1}}$$

$$\Rightarrow (x^2 + 1) \frac{d}{dx} + \frac{x}{\sqrt{x^2 + 1}} + 1 = 0$$

76. $e^x = x + \sin^2 x$... (i)

When $x = 0$, $e^0 = 0 + \sin^2 0$

Differentiating (i) w.r.t. x , we get

$$e^x + e^x \left(\frac{d}{dx} x + \frac{d}{dx} \sin^2 x \right) = \frac{d}{dx} x + 2 \sin x \cos x$$

Putting $x = 0$, $e^0 = 0$, we get

$$\frac{d}{dx} = 1$$

77. $2^x + 2 = 2^{x+1}$

Differentiating both sides w.r.t. x , we get

$$2(\log 2) + 2(\log 2) \frac{d}{dx} = 2^{(x+1)} (\log 2) \left(1 + \frac{d}{dx} \right)$$

$$\Rightarrow 2 + 2 \frac{d}{dx} = 2^{x+1} + 2^{x+1} \left(\frac{d}{dx} \right)$$

$$\Rightarrow \frac{d}{dx} (2 - 2^{x+1}) = 2^{x+1} - 2$$

$$\Rightarrow \frac{d}{dx} = \frac{2^{x+1} - 2}{2 - 2^{x+1}}$$

$$\therefore \left(\frac{d}{dx} \right)_{x=1} = \frac{2^2 - 2}{2 - 2^2} = \frac{2}{-2} = -1$$



$$78. \quad \sin^{-1} e^{-\cos x} = e$$

Differentiating both sides w.r.t. x , we get

$$\cos x \frac{d}{dx} + e^{-\cos x} \left\{ (-1) \left(-\sin x \frac{d}{dx} \right) + \cos x (-1) \right\} = 0$$

$$\Rightarrow \cos x \frac{d}{dx} + \sin x e^{-\cos x} \frac{d}{dx} - \cos x e^{-\cos x} = 0$$

$$\Rightarrow \frac{d}{dx} = \frac{\cos x e^{-\cos x}}{\cos x + \sin x e^{-\cos x}}$$

$$\therefore \left(\frac{d}{dx} \right)_{(1, \pi)} = \frac{\cos \pi e^{-\cos \pi}}{\cos \pi + \sin \pi e^{-\cos \pi}} = \frac{(-1)e}{-1+0} = e$$

$$79. \quad 2^x - 2 \cot x - 1 = 0 \quad \dots(i)$$

Putting $x = 1$ in (i), we get

$$1 - 2 \cot x - 1 = 0 \Rightarrow \cot x = 0 \Rightarrow x = \frac{\pi}{2}$$

Differentiating (i) w.r.t. x , we get

$$2^x (1 + \log 2) - 2 (1 + \log 2) \cot x + 2 \operatorname{cosec}^2 x \cdot \frac{d}{dx} = 0$$

Putting $x = 1$ and $\frac{d}{dx} = \frac{\pi}{2}$, we get

$$2 - 0 + 2 \frac{d}{dx} = 0 \Rightarrow \frac{d}{dx} = -1$$

$$80. \quad \text{Let } y = 6^x \text{ and } z = 3^x$$

$$\therefore \frac{dy}{dx} = 6^x \ln 6 \text{ and } \frac{dz}{dx} = 3^x \ln 3$$

$$\therefore \frac{dy}{dz} = \frac{\frac{dy}{dx}}{\frac{dz}{dx}} = \frac{6^x \ln 6}{3^x \ln 3} = 2^x$$

$$81. \quad \text{Let } y = \sin x \text{ and } z = \cos x$$

$$\therefore \frac{dy}{dx} = \cos x \text{ and } \frac{dz}{dx} = -\sin x$$

$$\therefore \frac{dy}{dz} = \frac{\frac{dy}{dx}}{\frac{dz}{dx}} = \frac{\cos x}{-\sin x} = -\cot x$$

$$82. \quad \text{Let } y = \sin^2 x \text{ and } z = \cos^2 x$$

$$\therefore \frac{dy}{dx} = 2 \sin x \cos x \text{ and } \frac{dz}{dx} = -2 \cos x \sin x$$

$$\therefore \frac{dy}{dz} = \frac{\frac{dy}{dx}}{\frac{dz}{dx}} = -1$$

$$83. \quad \text{Let } y = \cos^3 x \text{ and } z = \sin^3 x$$

$$\therefore \frac{dy}{dx} = -3 \cos^2 x \sin x \text{ and } \frac{dz}{dx} = 3 \sin^2 x \cos x$$

$$\therefore \frac{dy}{dz} = \frac{\frac{dy}{dx}}{\frac{dz}{dx}} = \frac{-\cos x}{\sin x} = -\cot x$$

$$84. \quad \text{Let } y = \log_{10} x \text{ and } z = x^2$$

$$\therefore \frac{dy}{dx} = \frac{1}{x \log_e 10} \text{ and } \frac{dz}{dx} = 2x$$

$$\therefore \frac{dy}{dz} = \frac{\frac{dy}{dx}}{\frac{dz}{dx}} = \frac{1}{2x^2 \log_e 10} = \frac{1}{2x^2} \log_{10} e$$

$$85. \quad \text{Let } y = \log_{10} x \text{ and } z = \log x$$

$$\therefore \frac{dy}{dx} = \frac{1}{x \log 10}$$

$$\text{and } \frac{dz}{dx} = \log 10 \cdot \left[-\frac{1}{(\log x)^2} \cdot \frac{1}{x} \right] = -\frac{\log 10}{(x \log x)^2}$$

$$\therefore \frac{dy}{dz} = \frac{\frac{dy}{dx}}{\frac{dz}{dx}} = \frac{\frac{1}{x \log 10}}{-\frac{\log 10}{(x \log x)^2}} = -(\log_{10} x)^2$$

$$86. \quad y = a \cos^3 \theta \text{ and } z = a \sin^3 \theta$$

$$\therefore \frac{dy}{d\theta} = -3a \cos^2 \theta \sin \theta$$

$$\text{and } \frac{dz}{d\theta} = 3a \sin^2 \theta \cos \theta$$

$$\therefore \frac{dy}{dz} = -\tan \theta$$

$$\therefore 1 + \left(\frac{dy}{dz} \right)^2 = 1 + \tan^2 \theta = \sec^2 \theta$$

$$87. \quad y = \log(1+t^2) \text{ and } z = t - \tan^{-1} t$$

$$\therefore \frac{dy}{dt} = \frac{2t}{1+t^2} \text{ and } \frac{dz}{dt} = 1 - \frac{1}{1+t^2} = \frac{t^2}{1+t^2}$$

$$\therefore \frac{dy}{dz} = \frac{\frac{dy}{dt}}{\frac{dz}{dt}} = \frac{t}{t^2} = \frac{1}{t}$$

Since, $y = \log(1+t^2)$

$$\therefore t = (e^y - 1)^{1/2}$$

$$\therefore \frac{dy}{dz} = \frac{(e^y - 1)^{1/2}}{2}$$



88. $y = a(t - \sin t)$ and $x = a(1 - \cos t)$
 $\therefore \frac{dy}{dt} = a(1 - \cos t)$ and $\frac{dx}{dt} = a \sin t$
 $\therefore \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{a \sin t}{a(1 - \cos t)} = \frac{2a \sin \frac{t}{2} \cos \frac{t}{2}}{2a \sin^2 \frac{t}{2}}$
 $= \cot \frac{t}{2}$

89. $y = 2 \cos \theta - \cos 2\theta$ and $x = 2 \sin \theta - \sin 2\theta$
 $\therefore \frac{dy}{d\theta} = -2 \sin \theta + 2 \sin 2\theta$ and
 $\frac{dx}{d\theta} = 2 \cos \theta - 2 \cos 2\theta$
 $\therefore \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{\cos \theta - \cos 2\theta}{\sin 2\theta - \sin \theta}$
 $= \frac{2 \sin \frac{3\theta}{2} \sin \frac{\theta}{2}}{2 \cos \frac{3\theta}{2} \sin \frac{\theta}{2}}$
 $= \tan \frac{3\theta}{2}$

90. $\sin \theta = \frac{2t}{1+t^2}$, $\tan \theta = \frac{2t}{1-t^2}$
 Putting $t = \tan \theta$ in both equations, we get
 $\sin \theta = \frac{2 \tan \theta}{1 + \tan^2 \theta}$
 $\Rightarrow \sin \theta = \sin 2\theta$
 $\Rightarrow \theta = 2\theta$
 $\therefore \frac{d\theta}{d\theta} = 2$
 $\tan \theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}$
 $\Rightarrow \tan \theta = \tan 2\theta$
 $\Rightarrow \theta = 2\theta$
 $\therefore \frac{d\theta}{d\theta} = 2$
 $\frac{d}{dx} = \left(\frac{d}{d\theta} \right) = 1$

91. Let $y = (\log x)$ and $z = \log(\log x)$
 $\therefore \log x = \log(\log x)$
 Differentiating both sides w.r.t. x , we get
 $\frac{1}{x} \cdot \frac{dy}{dx} = \log(\log x) + \frac{1}{\log x}$
 $\Rightarrow \frac{dy}{dx} = (\log x) \left[\log(\log x) + \frac{1}{\log x} \right]$
 $z = \log(\log x)$
 $\therefore \frac{dz}{dx} = \frac{1}{\log x}$
 $\therefore \frac{dy}{dz} = \frac{\frac{dy}{dx}}{\frac{dz}{dx}} = (\log x) \left[\log(\log x) + \frac{1}{\log x} \right]$

92. Let $y = \tan^{-1} \left(\frac{\sin \theta}{1 + \sqrt{1 - \sin^2 \theta}} \right)$ and $z = \sin^{-1} \theta$
 Put $y = \sin \theta \Rightarrow \theta = \sin^{-1} y$
 $\therefore y = \tan^{-1} \left(\frac{\sin \theta}{1 + \cos \theta} \right)$
 $= \tan^{-1} \left(\tan \frac{\theta}{2} \right) = \frac{\theta}{2}$
 $= \frac{\sin^{-1} y}{2}$
 $\therefore \frac{dy}{dx} = \frac{1}{2\sqrt{1-x^2}}$ and $\frac{dz}{dx} = \frac{1}{\sqrt{1-x^2}}$
 $\therefore \frac{dy}{dz} = \frac{\frac{dy}{dx}}{\frac{dz}{dx}} = \frac{1}{2}$

93. Let $u = \cos^{-1}(2x^2 - 1)$ and $v = \cos^{-1} x$
 Putting $x = \cos \theta$ in both equations, we get
 $u = \cos^{-1}(2 \cos^2 \theta - 1)$
 $u = \cos^{-1}(\cos 2\theta)$
 $= 2\theta$
 $v = \cos^{-1}(\cos \theta)$
 $= \theta$
 $\therefore \frac{du}{d\theta} = 2$ and $\frac{dv}{d\theta} = 1$
 $\frac{du}{dv} = \left(\frac{du}{d\theta} \right) = 2$



94. Let $y = \tan^{-1}\left(\frac{\sqrt{1+z^2}-1}{z}\right)$ and $z = \tan^{-1} \theta$

Put $y = \tan \theta \Rightarrow \theta = \tan^{-1} z$

$\therefore y = \tan^{-1}\left(\frac{\sec \theta - 1}{\tan \theta}\right)$

$= \tan^{-1}\left(\frac{1 - \cos \theta}{\sin \theta}\right)$

$= \tan^{-1}\left(\tan \frac{\theta}{2}\right) = \frac{\theta}{2} = \frac{\tan^{-1} z}{2}$

$\therefore \frac{dy}{dz} = \frac{1}{2(1+z^2)}$ and $\frac{dz}{d\theta} = \frac{1}{1+\theta^2}$

$\therefore \frac{dy}{d\theta} = \frac{dy}{dz} \cdot \frac{dz}{d\theta} = \frac{1}{2}$

95. Let $y = \sin^{-1}(2\sqrt{1-z^2})$

and $z = \sin^{-1}(3-4z^3)$

Put $y = \sin \theta \Rightarrow \theta = \sin^{-1} z$

$\therefore y = \sin^{-1}(2\sin \theta \sqrt{1-\sin^2 \theta})$ and

$z = \sin^{-1}(3\sin \theta - 4\sin^3 \theta)$

$\Rightarrow y = \sin^{-1}(\sin 2\theta)$ and $z = \sin^{-1}(\sin 3\theta)$

$\Rightarrow y = 2\theta = 2\sin^{-1} z$ and $z = 3\theta = 3\sin^{-1} z$

$\therefore \frac{dy}{dz} = \frac{2}{\sqrt{1-z^2}}$ and $\frac{dz}{d\theta} = \frac{3}{\sqrt{1-z^2}}$

$\therefore \frac{dy}{d\theta} = \frac{dy}{dz} \cdot \frac{dz}{d\theta} = \frac{2}{3}$

96. Let $y = \tan^{-1}\left(\frac{z}{\sqrt{1-z^2}}\right)$

and $z = \sin^{-1}(3-4z^3)$

Put $y = \sin \theta \Rightarrow \theta = \sin^{-1} z$

$\therefore y = \tan^{-1}\left(\frac{\sin \theta}{\sqrt{1-\sin^2 \theta}}\right)$

$= \tan^{-1}(\tan \theta) = \theta = \sin^{-1} z$ and

$z = \sin^{-1}(3\sin \theta - 4\sin^3 \theta)$

$= \sin^{-1}(\sin 3\theta) = 3\theta = 3\sin^{-1} z$

$\therefore \frac{dy}{dz} = \frac{dy}{d\theta} \cdot \frac{d\theta}{dz} = \frac{1}{3} \cdot \frac{1}{\sqrt{1-z^2}} = \frac{1}{3\sqrt{1-z^2}}$

97. $f(x) = \tan^{-1} \log x$

$\therefore \log f(x) = \tan^{-1} \log x$

$\therefore \frac{1}{f(x)} f'(x) = \frac{\log x}{1+x^2} + \frac{\tan^{-1} x}{x}$

$\Rightarrow f'(x) = \tan^{-1} x \left[\frac{\log x}{1+x^2} + \frac{\tan^{-1} x}{x} \right]$

$g(x) = \sec^{-1}\left(\frac{1}{2-x^2}\right)$

$\therefore g(x) = \cos^{-1}(2-x^2)$

Put $y = \cos \theta \Rightarrow \theta = \cos^{-1} x$

$\therefore g(x) = \cos^{-1}(2\cos^2 \theta - 1)$

$= \cos^{-1}(\cos 2\theta)$

$= 2\theta$

$\therefore g(x) = 2\cos^{-1} x$

$\therefore g'(x) = \frac{-2}{\sqrt{1-x^2}}$

Now,

$\frac{f'(x)}{g'(x)} = \frac{\tan^{-1} x \left[\frac{\log x}{1+x^2} + \frac{\tan^{-1} x}{x} \right]}{-2}$

$= -\frac{1}{2} \sqrt{1-x^2} \tan^{-1} x \left[\frac{\log x}{1+x^2} + \frac{\tan^{-1} x}{x} \right]$

98. $y = ct$ and $x = \frac{c}{t}$

$\therefore \frac{dy}{dt} = c$ and $\frac{dx}{dt} = \frac{-c}{t^2}$

$\therefore \frac{dy}{dx} = \frac{c}{\frac{-c}{t^2}} = \frac{-1}{t^2}$

$\therefore \left(\frac{dy}{dx}\right)_{(t=2)} = \frac{-1}{2^2} = \frac{-1}{4}$

99. $y = a \sin^3 \theta$ and $x = a \cos^3 \theta$

$\therefore \frac{dy}{d\theta} = 3a \sin^2 \theta \cos \theta$ and $\frac{dx}{d\theta} = -3a \cos^2 \theta \sin \theta$

$\therefore \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{-\sin \theta}{\cos \theta} = -\tan \theta$

$\therefore \left(\frac{dy}{dx}\right)_{\theta=\frac{\pi}{3}} = -\tan \frac{\pi}{3} = -\sqrt{3}$



$$\begin{aligned}
 100. \quad &= e^\theta(\sin\theta - \cos\theta) \\
 \therefore \quad &\frac{d}{d\theta} = e^\theta(\cos\theta + \sin\theta) + e^\theta(\sin\theta - \cos\theta) \\
 &= 2e^\theta \sin\theta \\
 &= e^\theta(\sin\theta + \cos\theta) \\
 \therefore \quad &\frac{d}{d\theta} = e^\theta(\cos\theta - \sin\theta) + e^\theta(\sin\theta + \cos\theta) \\
 &= 2e^\theta \cos\theta \\
 \frac{d}{d} &= \frac{\frac{d}{d\theta}}{\frac{d}{d}} = \frac{2e^\theta \cos\theta}{2e^\theta \sin\theta} = \cot\theta \\
 \therefore \quad &\frac{d}{d}\left(\theta = \frac{\pi}{4}\right) = 1
 \end{aligned}$$

$$\begin{aligned}
 101. \quad &\text{Let } = \log(\sec\theta + \tan\theta) \text{ and } z = \sec\theta \\
 \therefore \quad &\frac{d}{d\theta} = \frac{1}{\sec\theta + \tan\theta} \cdot (\sec\theta \tan\theta + \sec^2\theta) = \sec\theta \\
 \text{and } \frac{dz}{d\theta} &= \sec\theta \tan\theta \\
 \therefore \quad &\frac{d}{dz} = \frac{\frac{d}{d\theta}}{\frac{dz}{d\theta}} = \frac{\sec\theta}{\sec\theta \tan\theta} = \frac{1}{\tan\theta} = \cot\theta \\
 \therefore \quad &\left(\frac{d}{dz}\right)_{\theta = \frac{\pi}{4}} = \cot\frac{\pi}{4} = 1
 \end{aligned}$$

$$\begin{aligned}
 102. \quad &\text{Let } = \sec^{-1}\left(\frac{1}{2^2 - 1}\right) \text{ and } z = \sqrt{1 - ^2} \\
 \therefore \quad &= \cos^{-1}(2^2 - 1) \\
 \text{Put } &= \cos\theta \Rightarrow \theta = \cos^{-1} \\
 \therefore \quad &= \cos^{-1}(2 \cos^2\theta - 1) \\
 &= \cos^{-1}(\cos 2\theta) = 2\theta = 2\cos^{-1} \\
 \therefore \quad &\frac{d}{d} = \frac{-2}{\sqrt{1 - ^2}} \\
 \text{and } \frac{dz}{d} &= \frac{-2}{2\sqrt{1 - ^2}} = \frac{-}{\sqrt{1 - ^2}} \\
 \therefore \quad &\frac{d}{dz} = \frac{\frac{d}{d}}{\frac{dz}{d}} = 2 \\
 \therefore \quad &\left(\frac{d}{dz}\right)_{\left(-\frac{1}{2}\right)} = 4
 \end{aligned}$$

$$\begin{aligned}
 103. \quad &\text{Let } = f(\tan) \text{ and } z = g(\sec) \\
 \therefore \quad &\frac{d}{d} = f'(\tan) \cdot \sec^2 \\
 \text{and } \frac{dz}{d} &= g'(\sec) \cdot \sec \tan \\
 \therefore \quad &\frac{d}{dz} = \frac{\frac{d}{d}}{\frac{dz}{d}} = \frac{f'(\tan)}{g'(\sec)} \cdot \text{cosec} \\
 \therefore \quad &\left(\frac{d}{dz}\right)_{\left(-\frac{\pi}{4}\right)} = \frac{f'(1)}{g'(\sqrt{2})} \cdot \sqrt{2} = \frac{2\sqrt{2}}{4} = \frac{1}{\sqrt{2}}
 \end{aligned}$$

$$\begin{aligned}
 104. \quad &= A \sin 5 \quad \dots(i) \\
 \therefore \quad &\frac{d}{d} = 5 A \cos 5 \\
 \therefore \quad &\frac{d^2}{d^2} = -25 A \sin 5 \\
 \Rightarrow \frac{d^2}{d^2} &= -25 \quad \dots[\text{From (i)}]
 \end{aligned}$$

$$\begin{aligned}
 105. \quad &= A \cos 4t + B \sin 4t \\
 \therefore \quad &\frac{d}{dt} = -4A \sin 4t + 4B \cos 4t \\
 \therefore \quad &\frac{d^2}{dt^2} = -16A \cos 4t - 16B \sin 4t \\
 &= -16(A \cos 4t + B \sin 4t) \\
 &= -16
 \end{aligned}$$

$$\begin{aligned}
 106. \quad &\frac{^2}{a^2} + \frac{^2}{b^2} = 1 \\
 \Rightarrow b^2 ^2 + a^2 ^2 &= a^2 b^2 \\
 \text{Differentiating w.r.t } , \text{ we get} \\
 2b^2 + 2a^2 \frac{d}{d} &= 0 \\
 \Rightarrow 2a^2 \frac{d}{d} &= -2b^2 \\
 \Rightarrow \frac{d}{d} &= \frac{-b^2}{a^2} \left(-\right)
 \end{aligned}$$



$$\begin{aligned} \Rightarrow \frac{d^2}{d^2} &= \frac{-b^2}{a^2} \left[\frac{-\frac{d}{d}}{\frac{d}{d}} \right] \\ &= \frac{-b^2}{a^2} \left[- \left(\frac{-b^2}{a^2} \right) \right] \\ &= \frac{-b^2}{a^2} \left[\frac{a^2 + b^2}{a^2} \right] \\ &= \frac{-b^2}{a^2} \times \frac{a^2 + b^2}{a^2} \\ &= \frac{-b^4}{a^2} \end{aligned}$$

$$107. \quad = \log(\log)$$

$$\therefore \frac{d}{d} = \frac{1}{\log}$$

$$\therefore \frac{d^2}{d} = \frac{-1}{(\log)^2} [1 + \log]$$

$$108. \quad \text{Let } = \frac{e+1}{e} = 1 + \frac{1}{e} = 1 + e^{-}$$

$$\therefore \frac{d}{d} = -e^{-}$$

$$\therefore \frac{d^2}{d^2} = e^{-} = \frac{1}{e}$$

$$109. \quad = (\tan^{-1})^2$$

$$\therefore \frac{d}{d} = \frac{2 \tan^{-1}}{1 + ^2}$$

$$\Rightarrow \frac{d}{d} (1 + ^2) = 2 \tan^{-1}$$

$$\therefore \frac{d}{d} (2) + (1 + ^2) \frac{d^2}{d^2} = \frac{2}{1 + ^2}$$

$$\Rightarrow (2 + 1)^2 \frac{d^2}{d^2} + 2 (2 + 1) \frac{d}{d} = 2$$

$$110. \quad = (\sin^{-1})^2$$

$$\therefore \frac{d}{d} = \frac{2 \sin^{-1}}{\sqrt{1 - ^2}} \quad \dots(i)$$

$$\therefore \frac{d^2}{d^2} = 2 \left[\frac{1 + \sin^{-1} \cdot (1 - ^2)^{-1/2}}{1 - ^2} \right]$$

$$\Rightarrow (1 - ^2) \frac{d^2}{d^2} = 2 [1 + \sin^{-1} \cdot (1 - ^2)^{-1/2}]$$

$$\Rightarrow (1 - ^2) \frac{d^2}{d^2} - \frac{d}{d} = 2 \quad \dots[\text{From (i)}]$$

$$111. \quad = \frac{(\sin^{-1})^2}{2} \quad \therefore \frac{d}{d} = \frac{\sin^{-1}}{\sqrt{1 - ^2}}$$

$$\Rightarrow \sqrt{1 - ^2} \frac{d}{d} = \sin^{-1}$$

$$\Rightarrow \sqrt{1 - ^2} \frac{d^2}{d^2} + \frac{d}{d} \left(\frac{-}{\sqrt{1 - ^2}} \right) = \frac{1}{\sqrt{1 - ^2}}$$

$$\Rightarrow (1 - ^2) \frac{d^2}{d^2} - \frac{d}{d} = 1$$

$$112. \quad \sqrt{r} = \cos^{-1} \Rightarrow = (\cos^{-1})^2$$

$$\therefore \frac{d}{d} = -\frac{2 \cos^{-1}}{\sqrt{1 - ^2}}$$

$$\Rightarrow \frac{d^2}{d^2} = \frac{2 - \frac{2 \cos^{-1}}{\sqrt{1 - ^2}}}{1 - ^2}$$

$$\Rightarrow \frac{d^2}{d^2} = \frac{2 + \frac{d}{d}}{1 - ^2}$$

$$\Rightarrow (1 - ^2) \frac{d^2}{d^2} - \frac{d}{d} = 2$$

$$113. \quad \sqrt{r} = a \cdot e^{\theta(\cot \alpha)} \Rightarrow r = a^2 \cdot e^{2\theta(\cot \alpha)}$$

$$\therefore \frac{dr}{d\theta} = a^2 \cdot e^{2\theta(\cot \alpha)} \cdot 2 \cot \alpha$$

$$\Rightarrow \frac{dr}{d\theta} = 2a^2 \cot \alpha \cdot e^{2\theta(\cot \alpha)}$$

$$\therefore \frac{d^2 r}{d\theta^2} = 4a^2 \cot^2 \alpha \cdot e^{2\theta(\cot \alpha)}$$

$$\therefore \frac{d^2 r}{d\theta^2} - 4r \cot^2 \alpha$$

$$= 4a^2 \cot^2 \alpha \cdot e^{2\theta(\cot \alpha)} - 4a^2 \cot^2 \alpha \cdot e^{2\theta(\cot \alpha)} = 0$$

$$114. \quad = \frac{2}{\sqrt{a^2 - b^2}} \tan^{-1} \left(\sqrt{\frac{a-b}{a+b}} \tan \frac{r}{2} \right)$$

$$\therefore \frac{d}{d} = \frac{2}{\sqrt{a^2 - b^2}} \frac{1}{1 + \left(\frac{a-b}{a+b} \right) \tan^2 \frac{r}{2}} \times \sqrt{\frac{a-b}{a+b}} \sec^2 \frac{r}{2} \left(\frac{1}{2} \right)$$

$$= \frac{1}{a+b} \times \frac{\sec^2 \frac{r}{2}}{1 + \left(\frac{a-b}{a+b} \right) \tan^2 \frac{r}{2}}$$



$$\Rightarrow \frac{d}{d} = \frac{\sec^2 \frac{\pi}{2}}{(a+b) + (a-b)\tan^2 \frac{\pi}{2}}$$

$$\left[(a+b) + (a-b)\tan^2 \frac{\pi}{2} \right] \left(\sec \frac{\pi}{2} \sec \frac{\pi}{2} \tan \frac{\pi}{2} \right)$$

$$\therefore \frac{d^2}{d^2} = \frac{-\sec^2 \frac{\pi}{2} \left[(a-b)\tan \frac{\pi}{2} \sec^2 \frac{\pi}{2} \right]}{\left[(a+b) + (a-b)\tan^2 \frac{\pi}{2} \right]^2}$$

$$(a+b+a-b)(\sqrt{2} \times \sqrt{2} \times 1)$$

$$\left(\frac{d^2}{d^2} \right)_{\left(\theta = \frac{\pi}{4} \right)} = \frac{-(\sqrt{2})^2 \left[(a-b)(\sqrt{2})^2 \right]}{(a+b+a-b)^2}$$

$$= \frac{4a - 4(a-b)}{4a^2}$$

$$= \frac{4b}{4a^2} = \frac{b}{a^2}$$

115. Here, $\frac{d}{ds} = 1, \frac{d}{ds} = 2$ (i)

and $\frac{d^2}{ds^2} = 0, \frac{d^2}{ds^2} = 0$ (ii)

Now, $u = s^2 + \frac{1}{s^2}$

$$\therefore \frac{du}{ds} = 2 \cdot \frac{d}{ds} + 2 \cdot \frac{d}{ds}$$

$$\therefore \frac{d^2 u}{ds^2} = 2 \left(\frac{d}{ds} \right)^2 + 2 \left(\frac{d^2}{ds^2} \right) + 2 \left(\frac{d}{ds} \right)^2 + 2 \left(\frac{d^2}{ds^2} \right)$$

From (i) and (ii), we get

$$\frac{d^2 u}{ds^2} = 2(1) + 0 + 2(4) + 0 = 10$$

116. $y = at^2$

$$\therefore \frac{dy}{dt} = 2at$$

$$= 2at$$

$$\therefore \frac{d}{dt} = 2a \quad \dots(i)$$

$$\therefore \frac{d}{d} = \frac{2at}{2a} \Rightarrow \frac{d}{d} = t$$

$$\therefore \frac{d}{d} \left(\frac{d}{d} \right) = \frac{d}{d} (t)$$

$$\therefore \frac{d^2}{d^2} = \frac{dt}{d}$$

$$\therefore \frac{d^2}{d^2} = \frac{1}{2a} \quad \dots[\text{From (i)}]$$

117. $y = f(t)$ and $y = g(t)$

$$\therefore \frac{dy}{dt} = f'(t) \text{ and } \frac{dy}{dt} = g'(t)$$

$$\therefore \frac{d}{d} = \frac{\frac{dy}{dt}}{\frac{dy}{dt}} = \frac{g'(t)}{f'(t)}$$

$$\therefore \frac{d^2}{d^2} = \frac{f'(t) \cdot g''(t) - g'(t) \cdot f''(t)}{[f'(t)]^2} \cdot \frac{dt}{d}$$

$$= \frac{f'(t) \cdot g''(t) - g'(t) \cdot f''(t)}{[f'(t)]^2} \cdot \frac{1}{f'(t)}$$

$$= \frac{f'(t) \cdot g''(t) - g'(t) \cdot f''(t)}{[f'(t)]^3}$$

118. $y = (1 + \sqrt{1+x^2})^n$ (i)

$$\therefore \frac{dy}{dx} = n(1 + \sqrt{1+x^2})^{n-1} \cdot \left(1 + \frac{x}{\sqrt{1+x^2}} \right)$$

$$\Rightarrow \frac{dy}{dx} = \frac{n(1 + \sqrt{1+x^2})^n}{\sqrt{1+x^2}}$$

$$\Rightarrow (\sqrt{1+x^2}) \frac{dy}{dx} = n(1 + \sqrt{1+x^2})^n$$

Again, differentiating both sides w.r.t. x , we get

$$\sqrt{1+x^2} \cdot \frac{d^2 y}{dx^2} + \frac{dy}{dx} \cdot \left(\frac{x}{\sqrt{1+x^2}} \right)$$

$$= n^2(1 + \sqrt{1+x^2})^{n-1} \left(1 + \frac{x}{\sqrt{1+x^2}} \right)$$

$$\Rightarrow (1+x^2) \frac{d^2 y}{dx^2} + \frac{dy}{dx} = n^2(1 + \sqrt{1+x^2})^n$$

$$\Rightarrow (1+x^2) \frac{d^2 y}{dx^2} + \frac{dy}{dx} = n^2 \quad \dots[\text{From (i)}]$$

119. $y = (x^2 + 3)^5$

Taking logarithm on both sides, we get

$$2 \log x + 3 \log y = 5 \log (x^2 + 3)$$

Differentiating both sides w.r.t. x , we get

$$\frac{2}{x} + \frac{3}{y} \cdot \frac{dy}{dx} = \frac{5}{x^2 + 3} \left(1 + \frac{d}{dx} \right)$$

$$\Rightarrow \frac{dy}{dx} \left(\frac{3}{y} - \frac{5}{x^2 + 3} \right) = \frac{5}{x^2 + 3} - \frac{2}{x}$$

$$\Rightarrow \frac{dy}{dx} = \dots \quad \dots(i)$$



$$\begin{aligned} \therefore \frac{d^2}{d^2} &= \frac{d}{d^2} \\ \Rightarrow \frac{d^2}{d^2} &= \frac{(-)}{2} \dots [\text{From (i)}] \\ \Rightarrow \frac{d^2}{d^2} &= 0 \end{aligned}$$

120. $y = \sin t$ and $x = \sin pt$

$$\therefore \frac{dy}{dt} = \cos t$$

and $\frac{dx}{dt} = p \cos pt$

$$\therefore \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{\cos t}{p \cos pt}$$

$$\therefore \frac{dy}{dx} = \frac{p\sqrt{1-t^2}}{\sqrt{1-t^2}} \dots (i)$$

Again, differentiating w.r.t. x , we get

$$\frac{d^2 y}{dx^2} = \frac{p\sqrt{1-t^2} \left(\frac{-2}{2\sqrt{1-t^2}} \right) \frac{d}{dx} - p\sqrt{1-t^2} \left(\frac{-2}{2\sqrt{1-t^2}} \right)}{(\sqrt{1-t^2})^2}$$

$$\Rightarrow (1-t^2) \frac{d^2 y}{dx^2} = -p \frac{\sqrt{1-t^2}}{\sqrt{1-t^2}} \cdot \frac{d}{dx} + p \frac{\sqrt{1-t^2}}{\sqrt{1-t^2}}$$

$$\Rightarrow (1-t^2) \frac{d^2 y}{dx^2} = -p^2 + \frac{d}{dx} \dots [\text{From (i)}]$$

$$\Rightarrow (1-t^2) \frac{d^2 y}{dx^2} - \frac{d}{dx} + p^2 = 0$$

121. $y = \cos \theta$ and $x = \sin 5\theta$

$$\therefore \frac{dy}{d\theta} = -\sin \theta \text{ and } \frac{dx}{d\theta} = 5 \cos 5\theta$$

$$\therefore \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = -\frac{\sin \theta}{5 \cos 5\theta}$$

$$\Rightarrow \frac{dy}{dx} = -\frac{5\sqrt{1-t^2}}{\sqrt{1-t^2}} \dots (i)$$

$$\therefore \frac{d^2 y}{dx^2} = \frac{-5\sqrt{1-t^2} \left(\frac{-2}{2\sqrt{1-t^2}} \right) \frac{d}{dx} + 5\sqrt{1-t^2} \left(\frac{-2}{2\sqrt{1-t^2}} \right)}{(\sqrt{1-t^2})^2}$$

$$\Rightarrow (1-t^2) \frac{d^2 y}{dx^2} = \frac{5\sqrt{1-t^2}}{\sqrt{1-t^2}} \cdot \frac{d}{dx} - \frac{5\sqrt{1-t^2}}{\sqrt{1-t^2}}$$

$$\Rightarrow (1-t^2) \frac{d^2 y}{dx^2} = -25 + \frac{d}{dx} \dots [\text{From (i)}]$$

$$\Rightarrow (1-t^2) \frac{d^2 y}{dx^2} - \frac{d}{dx} = -25$$

122. $y = e^{\sqrt{x}} + e^{-\sqrt{x}}$

$$\therefore \frac{dy}{dx} = \frac{e^{\sqrt{x}}}{2\sqrt{x}} - \frac{e^{-\sqrt{x}}}{2\sqrt{x}}$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{2\sqrt{x}} (e^{\sqrt{x}} - e^{-\sqrt{x}})$$

$$\therefore \frac{d^2 y}{dx^2} = \frac{1}{2\sqrt{x}} \left(\frac{e^{\sqrt{x}}}{2\sqrt{x}} + \frac{e^{-\sqrt{x}}}{2\sqrt{x}} \right) + \frac{(e^{\sqrt{x}} - e^{-\sqrt{x}})}{2} \left(\frac{-1}{2x^{3/2}} \right)$$

$$\Rightarrow \frac{d^2 y}{dx^2} = \frac{e^{\sqrt{x}} + e^{-\sqrt{x}}}{4} - \frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{4\sqrt{x}}$$

$$\therefore \frac{d^2 y}{dx^2} + \frac{1}{2} \cdot \frac{dy}{dx} = \left(\frac{e^{\sqrt{x}} + e^{-\sqrt{x}}}{4} - \frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{4\sqrt{x}} \right) + \frac{1}{2} \left(\frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{2\sqrt{x}} \right)$$

$$= \frac{e^{\sqrt{x}} + e^{-\sqrt{x}}}{4} - \frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{4\sqrt{x}} + \frac{e^{\sqrt{x}} - e^{-\sqrt{x}}}{4\sqrt{x}} = \frac{e^{\sqrt{x}} + e^{-\sqrt{x}}}{4}$$

$$\therefore \frac{d^2 y}{dx^2} + \frac{1}{2} \cdot \frac{dy}{dx} = \frac{1}{4} \dots [\text{From (i)}]$$

123. $y = 2at^2$ and $x = at^4$

$$\therefore \frac{dy}{dt} = 4at \text{ and } \frac{dx}{dt} = 4at^3$$

$$\therefore \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = t^2$$

$$\therefore \frac{d^2 y}{dx^2} = 2t \cdot \frac{dt}{dx} = 2t \cdot \frac{1}{4at} = \frac{1}{2a}$$

$$\therefore \left(\frac{d^2 y}{dx^2} \right)_{(t=2)} = \frac{1}{2a}$$



$$124. \quad x = a \sin \theta \text{ and } y = b \cos \theta$$

$$\therefore \frac{dx}{d\theta} = a \cos \theta \text{ and } \frac{dy}{d\theta} = -b \sin \theta$$

$$\therefore \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{-b}{a} \tan \theta$$

$$\therefore \frac{d^2y}{dx^2} = \frac{-b}{a} \sec^2 \theta \cdot \frac{d\theta}{dx} = \frac{-b}{a^2} \sec^3 \theta$$

$$\begin{aligned} \therefore \left(\frac{d^2y}{dx^2} \right)_{\left(\theta = \frac{\pi}{4} \right)} &= \frac{-b}{a^2} \sec^3 \frac{\pi}{4} \\ &= -2\sqrt{2} \frac{b}{a^2} \end{aligned}$$

$$125. \quad y = x^3 \log \log_e(1+x)$$

$$\therefore y' = 3x^2 \log \log_e(1+x) + \frac{x^3}{\log_e(1+x)} \cdot \frac{1}{1+x}$$

$$\therefore y'' = 6x \log \log_e(1+x) + \frac{3x^2}{\log_e(1+x)} \cdot \frac{1}{(1+x)}$$

$$+ \left[\frac{(1+x) \log_e(1+x) \cdot 3x^2 - x^3 \left[\frac{1}{1+x} + \log_e(1+x) \right]}{(1+x)^2 [\log_e(1+x)]^2} \right]$$

$$\therefore y''(0) = 0$$

$$126. \quad \text{At } (1, 1), x = e^t \sin t \text{ and } y = e^t \cos t$$

$$\therefore \tan t = 1 \Rightarrow t = \frac{\pi}{4}$$

$$\text{Now, } \frac{dx}{dt} = e^t \sin t \text{ and } \frac{dy}{dt} = e^t \cos t$$

$$\therefore \frac{dy}{dx} = e^t(\sin t + \cos t) \text{ and } \frac{d^2y}{dx^2} = e^t(\cos t - \sin t)$$

$$\therefore \frac{d^2y}{dx^2} = \frac{\frac{d}{dt} \left(\frac{dy}{dx} \right)}{\frac{dx}{dt}} = \frac{\cos t - \sin t}{\cos t + \sin t}$$

$$\begin{aligned} \therefore \frac{d^2y}{dx^2} &= \frac{d}{dt} \left(\frac{\cos t - \sin t}{\cos t + \sin t} \right) \frac{dt}{dx} \\ &= \left[\frac{(\cos t + \sin t)(-\sin t - \cos t) - (\cos t - \sin t)(-\sin t + \cos t)}{(\cos t + \sin t)^2} \right] \frac{dt}{dx} \\ &= \frac{-2}{(\cos t + \sin t)^2} \cdot \frac{1}{e^t(\sin t + \cos t)} \end{aligned}$$

$$\begin{aligned} &= \frac{-2}{(e^t \sin t + e^t \cos t)^2} \cdot \frac{1}{(\cos t + \sin t)^2} \\ &= \frac{-2}{+} \cdot \frac{1}{(\cos t + \sin t)^2} \end{aligned}$$

$$\therefore \left(\frac{d^2y}{dx^2} \right)_{(1,1)} = \frac{-2}{1+1} \cdot \frac{1}{\left(\cos \frac{\pi}{4} + \sin \frac{\pi}{4} \right)^2} = \frac{-1}{2}$$

$$127. \quad \text{At } \left(\frac{3}{2}\sqrt{2}, 2\sqrt{2} \right)$$

$$\cos t = \frac{1}{\sqrt{2}} \text{ and } \sin t = \frac{1}{\sqrt{2}}$$

$$\therefore \tan t = 1 \Rightarrow t = \frac{\pi}{4}$$

$$\text{Now, } \frac{dx}{dt} = 3 \cos t \text{ and } \frac{dy}{dt} = 4 \sin t$$

$$\therefore \frac{dy}{dx} = -3 \sin t \text{ and } \frac{d^2y}{dx^2} = 4 \cos t$$

$$\therefore \frac{d^2y}{dx^2} = \frac{\frac{d}{dt} \left(\frac{dy}{dx} \right)}{\frac{dx}{dt}} = -\frac{4}{3} \cot t$$

$$\therefore \frac{d^2y}{dx^2} = \frac{4}{3} \operatorname{cosec}^2 t \frac{dt}{dx} = \frac{4}{3} \operatorname{cosec}^2 t \times \frac{1}{3 \sin t}$$

$$\begin{aligned} \therefore \left(\frac{d^2y}{dx^2} \right)_{\left(\frac{3}{2}\sqrt{2}, 2\sqrt{2} \right)} &= \frac{4}{3} \operatorname{cosec}^2(\pi/4) \times \frac{-1}{3 \sin(\pi/4)} \\ &= \frac{-8\sqrt{2}}{9} \end{aligned}$$

$$128. \quad f(x) = \frac{x^2 - a + 1}{x^2 + a + 1}$$

$$\therefore f'(x) = \frac{(x^2 + a + 1)(2x) - (x^2 - a + 1)(2x + a)}{(x^2 + a + 1)^2}$$

$$\Rightarrow f'(x) = \frac{2a(x^2 - 1)}{(x^2 + a + 1)^2}$$

$$\therefore f''(x)$$

$$= \frac{4a(x^2 + a + 1)^2 - 4a(x^2 - 1)(2x + a)(x^2 + a + 1)}{(x^2 + a + 1)^4}$$

$$\Rightarrow f''(x) = \frac{4a[(x^2 + a + 1) - (x^2 - 1)(2x + a)]}{(x^2 + a + 1)^3}$$

$$\therefore f'(1) = 0, f''(1) = \frac{4a}{(2+a)^2} \text{ and } f''(-1) = -\frac{4a}{(2-a)^2}$$

$$\therefore (2+a)^2 f''(1) + (2-a)^2 f''(-1) = 0$$



$$129. \quad = \frac{a^{\cos^{-1}}}{1+a^{\cos^{-1}}} \text{ and } z = a^{\cos^{-1}} \Rightarrow = \frac{z}{1+z}$$

$$\therefore \frac{d}{dz} = \frac{(1+z)1-z(1)}{(1+z)^2} = \frac{1}{(1+z)^2} = \frac{1}{(1+a^{\cos^{-1}})^2}$$

$$130. \text{ Let } f(x) = \cos^{-1}\left(\sin\sqrt{\frac{1+x}{2}}\right) +$$

$$= \cos^{-1}\left[\cos\left(\frac{\pi}{2} - \sqrt{\frac{1+x}{2}}\right)\right] +$$

$$= \frac{\pi}{2} - \sqrt{\frac{1+x}{2}} +$$

$$\therefore f'(x) = -\frac{1}{\sqrt{2}} \cdot \frac{1}{2\sqrt{1+x}} + (1 + \log)$$

$$\therefore f'(1) = -\frac{1}{4} + 1 = \frac{3}{4}$$

$$131. \text{ Since, } g(x) \text{ is the inverse of } f(x).$$

$$\therefore fog(x) =$$

$$\Rightarrow \frac{d}{dx} [fog(x)] = \frac{d}{dx} (x)$$

$$\Rightarrow f'[g(x)] \cdot g'(x) = 1$$

$$\Rightarrow \frac{1}{1+[g(x)]^3} \cdot g'(x) = 1$$

$$\dots \left[\because f'(x) = \frac{1}{1+x^3} \text{ (given)} \right]$$

$$\Rightarrow g'(x) = 1 + [g(x)]^3$$

$$132. f(x) = \tan^{-1}$$

$$\therefore f'(x) = \frac{1}{1+x^2}$$

$$\therefore f''(x) = \frac{-1}{(1+x^2)^2} \cdot 2x$$

Since, $f'(x) + f''(x) = 0$

$$\therefore \frac{1}{1+x^2} - \frac{2x}{(1+x^2)^2} = 0$$

$$\Rightarrow 1+x^2 - 2x = 0$$

$$\Rightarrow x = 1$$

$$133. \quad = a\left(t - \frac{1}{t}\right) \dots(i)$$

$$\text{and } = a\left(t + \frac{1}{t}\right) \dots(ii)$$

Squaring (i) and (ii) and subtracting, we get

$$2 - 2 = a^2(-4) \Rightarrow 2 - 2 = 4a^2$$

Differentiating both sides w.r.t. t , we get

$$2 \frac{d}{dt} - 2 = 0 \Rightarrow \frac{d}{dt} = -$$

$$134. \quad 2 = \sin^{-1}(x + 5)$$

$$\Rightarrow \sin 2 = x + 5$$

Differentiating both sides w.r.t. x , we get

$$2 \cos 2 \left(\frac{d}{dx}\right) = 1 + 5\left(\frac{d}{dx}\right)$$

$$\Rightarrow \frac{d}{dx} (2 \cos 2 - 5) = 1$$

$$\Rightarrow \frac{d}{dx} = \frac{1}{2 \cos 2 - 5}$$

$$\text{Now, } \frac{d}{dx} = \frac{1}{(d/dx)}$$

$$\Rightarrow \frac{d}{dx} = 2 \cos 2 - 5$$

$$135. f(x+y) = f(x) + f(y) \text{ for all } x, y \in \mathbb{R}$$

Putting $x = 0$ and $y = 0$, we get

$$f(0) = f(0) + f(0) \Rightarrow f(0) = 0$$

$$\text{Now, } f'(0) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h}$$

$$\Rightarrow f'(0) = \lim_{h \rightarrow 0} \frac{f(h)}{h} \dots(i)$$

$$\therefore f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{f(x) + f(h) - f(x)}{h}$$

$$\Rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{f(h)}{h} = f'(0) \dots[\text{From (i)}]$$

$$\Rightarrow f(x) = f'(0) + c$$



But, $f(0) = 0$

$$\therefore c = 0$$

Hence, $f(x) = f'(0)$ for all $x \in \mathbb{R}$

Clearly, $f(x)$ is everywhere continuous and differentiable and $f'(x)$ is constant for all $x \in \mathbb{R}$.

Hence, option (D) is incorrect.

$$136. \quad x^2 + x^2 = t + \frac{2}{t}$$

Squaring on both sides, we get

$$4x^4 + 4x^4 + 2x^2 \cdot 2 = t^2 + \frac{4}{t^2} + 4$$

$$\Rightarrow \left(t^2 + \frac{4}{t^2} \right) + 2x^2 \cdot 2 = t^2 + \frac{4}{t^2} + 4$$

$$\Rightarrow 2x^2 = 2 \quad \dots(i)$$

Differentiating both sides w.r.t. x , we get

$$2 \cdot 2x \frac{d}{dx} + 2 \cdot 2 = 0$$

$$\Rightarrow 2x \frac{d}{dx} = -2$$

$$\Rightarrow 3x \frac{d}{dx} = -2x^2$$

$$\Rightarrow 3x \frac{d}{dx} = -2 \quad \dots[\text{From (i)}]$$

$$137. \quad \frac{d}{dx} f_n(x) = \frac{d}{dx} e^{f_{n-1}(x)}$$

Let $n = 3$

$$\begin{aligned} \therefore \frac{d}{dx} f_3(x) &= \frac{d}{dx} e^{f_2(x)} \\ &= e^{f_2(x)} \frac{d}{dx} f_2(x) \\ &= e^{f_2(x)} \frac{d}{dx} e^{f_1(x)} \\ &= e^{f_2(x)} e^{f_1(x)} \frac{d}{dx} f_1(x) \\ &= e^{f_2(x)} e^{f_1(x)} \frac{d}{dx} e \\ &= e^{f_2(x)} e^{f_1(x)} e \end{aligned}$$

$$\frac{d}{dx} f_3(x) = f_3(x) f_2(x) f_1(x)$$

Similarly,

$$\frac{d}{dx} f_n(x) = f_n(x) f_{n-1}(x) \dots f_1(x)$$

$$138. \quad f(x) = f(-x) \quad \dots[\because f(x) \text{ is an even function}]$$

$$\therefore f'(x) = -f'(-x)$$

$$\therefore f'(0) = -f'(0)$$

$$\therefore 2f'(0) = 0$$

$$\therefore f'(0) = 0$$



Evaluation Test

$$\begin{aligned} 1. \quad &= \frac{-1}{4} + \frac{(-1)^3}{12} + \frac{(-1)^5}{20} + \frac{(-1)^7}{28} + \dots \\ &= \frac{1}{4} \left[(-1) + \frac{(-1)^3}{3} + \frac{(-1)^5}{5} + \frac{(-1)^7}{7} + \dots \right] \end{aligned}$$

$$\text{Now, } \log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

$$\therefore \log(1-x) = -x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} - \dots$$

$$\begin{aligned} \therefore \log\left(\frac{1+x}{1-x}\right) &= \log(1+x) - \log(1-x) \\ &= 2 \left[x + \frac{x^3}{3} + \frac{x^5}{5} + \dots \right] \end{aligned}$$

$$\therefore -1 + \frac{(-1)^3}{3} + \frac{(-1)^5}{5} + \frac{(-1)^7}{7} + \dots$$

$$= \frac{1}{2} \log \left[\frac{1+(-1)}{1-(-1)} \right]$$

$$= \frac{1}{2} \log \left(\frac{0}{2} \right)$$

$$\therefore = \frac{1}{8} \log \left(\frac{0}{2} \right)$$



$$\begin{aligned} \therefore \frac{d}{d} &= \frac{1}{8} \left(\frac{2-}{ } \right) \left[\frac{(2-)(1) - (-1)}{(2-)^2} \right] \\ &= \frac{1}{8} \left(\frac{2-}{ } \right) \left[\frac{2- +}{(2-)^2} \right] = \frac{1}{4(2-)} \end{aligned}$$

$$2. = (\cos + i \sin) (\cos 3 + i \sin 3) \dots (\cos(2n-1) + i \sin(2n-1))$$

Since, $\cos \theta + i \sin \theta = e^{i\theta}$

$$\begin{aligned} \therefore &= e^i \cdot e^{i3} \cdot e^{i5} \dots e^{i(2n-1)} \\ &= e^{i[1+3+5+\dots+(2n-1)]} \\ &= e^{in^2} \end{aligned}$$

$$\therefore \frac{d}{d} = in^2 e^{in^2}$$

$$\therefore \frac{d^2}{d^2} = i^2 n^4 e^{in^2} = -n^4$$

$$3. = f\left(\frac{3 + \pi}{5 + 4}\right)$$

$$\begin{aligned} \therefore \frac{d}{d} &= f' \left(\frac{3 + \pi}{5 + 4} \right) \cdot \frac{d}{d} \left(\frac{3 + \pi}{5 + 4} \right) \\ &= f' \left(\frac{3 + \pi}{5 + 4} \right) \left[\frac{(5 + 4)3 - 5(3 + \pi)}{(5 + 4)^2} \right] \end{aligned}$$

$$\begin{aligned} \therefore \left(\frac{d}{d} \right)_{=0} &= f' \left(\frac{\pi}{4} \right) \left[\frac{12 - 5\pi}{16} \right] \\ &= \tan^2 \left(\frac{\pi}{4} \right) \left(\frac{12 - 5\pi}{16} \right) \\ &= (1)^2 \left(\frac{12 - 5\pi}{16} \right) \\ &= \frac{12 - 5\pi}{16} \end{aligned}$$

$$4. = |\cos | + |\sin |$$

Since, $\frac{d}{d} | | = \frac{| |}{| |}$

$$\begin{aligned} \therefore \frac{d}{d} &= \frac{|\cos |}{\cos } \cdot \frac{d}{d} (\cos) + \frac{|\sin |}{\sin } \cdot \frac{d}{d} (\sin) \\ &= \frac{|\cos |}{\cos } (-\sin) + \frac{|\sin |}{\sin } \cos \end{aligned}$$

$$\text{When } = \frac{2\pi}{3}, \cos = \cos \frac{2\pi}{3} = \frac{-1}{2}, |\cos | = \frac{1}{2}$$

$$\text{and } \sin = \sin \frac{2\pi}{3} = \frac{\sqrt{3}}{2}, |\sin | = \frac{\sqrt{3}}{2}$$

$$\begin{aligned} \therefore \left(\frac{d}{d} \right)_{=\frac{2\pi}{3}} &= -1 \left(\frac{-\sqrt{3}}{2} \right) + 1 \left(\frac{-1}{2} \right) \\ &= \frac{\sqrt{3} - 1}{2} \end{aligned}$$

$$5. = \left(1 + \frac{1}{ } \right) \left(1 + \frac{2}{ } \right) \dots \left(1 + \frac{n}{ } \right)$$

$$\begin{aligned} \therefore \frac{d}{d} &= \left(-\frac{1}{2} \right) \left(1 + \frac{2}{ } \right) \left(1 + \frac{3}{ } \right) \dots \left(1 + \frac{n}{ } \right) \\ &\quad + \left(1 + \frac{1}{ } \right) \left(-\frac{2}{2} \right) \left(1 + \frac{3}{ } \right) \dots \left(1 + \frac{n}{ } \right) \\ &\quad + \left(1 + \frac{1}{ } \right) \left(1 + \frac{2}{ } \right) \left(-\frac{3}{2} \right) \dots \left(1 + \frac{n}{ } \right) + \dots \end{aligned}$$

$$\text{When } = -1, 1 + \frac{1}{-1} = 1 + \frac{1}{(-1)} = 1 - 1 = 0$$

\therefore Except 1st term all terms are 0.

$$\begin{aligned} \therefore \left(\frac{d}{d} \right)_{(=-1)} &= (-1)(-1)(-2) \dots (1-n) \\ &= (-1)^n (n-1)! \end{aligned}$$

$$6. f() = \begin{cases} \frac{1+}{ }, & \geq 0 \\ \frac{1-}{ }, & < 0 \end{cases}$$

$$\therefore Lf'(0) = \lim_{\rightarrow 0^-} \frac{f() - f(0)}{-0} = \lim_{\rightarrow 0} \frac{1- - 0}{-0} = 1$$

$$Rf'(0) = \lim_{\rightarrow 0^+} \frac{1+ - 0}{-0} = 1$$

$\therefore f()$ is differentiable at $= 0$ and $f'(0) = 1$.

$$7. f() = \sin(\log)$$

$$\begin{aligned} \therefore f'() &= \cos(\log) \cdot \frac{1}{ } \\ &= f\left(\frac{2 + 3}{3 - 2}\right) \end{aligned}$$



$$\begin{aligned} \therefore \frac{d}{d} &= f' \left(\frac{2+3}{3-2} \right) \cdot \frac{d}{d} \left(\frac{2+3}{3-2} \right) \\ &= \cos \left(\log \left(\frac{2+3}{3-2} \right) \right) \\ &\quad \cdot \left[\frac{(3-2)(2) - (-2)(2+3)}{(3-2)^2} \right] \cdot \left(\frac{3-2}{2+3} \right) \\ &= \cos \left(\log \left(\frac{2+3}{3-2} \right) \right) \left[\frac{6-4+4+6}{3-2} \right] \\ &\quad \cdot \frac{1}{2+3} \\ &= \frac{12}{9-4^2} \cos \left\{ \log \left(\frac{2+3}{3-2} \right) \right\} \end{aligned}$$

$$8. \quad \frac{d}{d} \left[a \tan^{-1} + b \log \left(\frac{-1}{+1} \right) \right] = \frac{1}{4-1}$$

$$\begin{aligned} \therefore a \tan^{-1} + b \log \left(\frac{-1}{+1} \right) \\ &= \int \frac{1}{4-1} \\ &= \int \frac{1}{(x^2-1)(x^2+1)} \\ &= \frac{1}{2} \int \left(\frac{1}{x^2-1} - \frac{1}{x^2+1} \right) dx \\ &= \frac{1}{2} \cdot \frac{1}{2} \log \left(\frac{-1}{+1} \right) - \frac{1}{2} \tan^{-1} \end{aligned}$$

$$\therefore a = -\frac{1}{2}, b = \frac{1}{4}$$

$$\therefore a - 2b = -\frac{1}{2} - 2 \left(\frac{1}{4} \right) = -\frac{1}{2} - \frac{1}{2} = -1$$

$$\begin{aligned} 9. \quad f(x) &= \cos x \cos 2x \cos 4x \cos 8x \cos 16x \\ &= \frac{1}{32} \times \frac{16}{\sin x} (2 \sin x \cos x \cos 2x \cos 4x \\ &\quad \cos 8x \cos 16x) \\ &= \frac{1}{32} \times \frac{16}{\sin x} (\sin 2x \cos 2x \cos 4x \cos 8x \\ &\quad \cos 16x) \\ &= \frac{1}{32} \times \frac{8}{\sin x} (\sin 4x \cos 4x \cos 8x \\ &\quad \cos 16x) \\ &= \frac{1}{32} \times \frac{4}{\sin x} (\sin 8x \cos 8x \cos 16x) \end{aligned}$$

$$\begin{aligned} &= \frac{1}{32} \times \frac{2}{\sin x} \sin 16x \cos 16x \\ &= \frac{\sin 32x}{32 \sin x} \end{aligned}$$

$$\therefore f'(x) = \frac{1}{32} \left[\frac{\sin x \cdot 32 \cos 32x - \sin 32x \cos x}{\sin^2 x} \right]$$

$$\begin{aligned} \therefore f' \left(\frac{\pi}{4} \right) &= \frac{1}{32} \frac{\left[\frac{1}{\sqrt{2}} \cdot 32(1) - 0 \right]}{\left(\frac{1}{\sqrt{2}} \right)^2} \\ &= \frac{1}{32} \times \frac{1}{\sqrt{2}} \times 32 \times 2 = \frac{2}{\sqrt{2}} = \sqrt{2} \end{aligned}$$

$$\begin{aligned} 10. \quad 1 + x^4 + x^8 &= 1 + 2x^4 + x^8 - x^4 \\ &= (1 + x^4)^2 - x^4 \\ &= (1 + x^4 + x^2)(1 + x^4 - x^2) \end{aligned}$$

$$\therefore \frac{1 + x^4 + x^8}{1 + x^2 + x^4} = 1 - x^2 + x^4$$

$$\begin{aligned} \therefore \frac{d}{d} \left(\frac{1 + x^4 + x^8}{1 + x^2 + x^4} \right) &= \frac{d}{d} (1 - x^2 + x^4) \\ &= 4x^3 - 2x = ax^3 + bx \end{aligned}$$

$$\therefore a = 4, b = -2$$

$$11. \quad 2 = \frac{1}{5} + \frac{-1}{5}$$

$$\text{Let } \frac{1}{5} = a$$

$$\therefore \frac{-1}{5} = \frac{1}{a}, \quad \therefore a + \frac{1}{a} = 2$$

$$\therefore a^2 - 2a + 1 = 0$$

$$\therefore a = \frac{2 \pm \sqrt{4 - 2 - 4}}{2}$$

$$\therefore \frac{1}{5} = +\sqrt{x^2-1}$$

$$\therefore = \left(+\sqrt{x^2-1} \right)^5$$

$$\therefore \frac{d}{d} = 5 \left(+\sqrt{x^2-1} \right)^4 \left(1 + \frac{1}{2\sqrt{x^2-1}} \cdot 2 \right)$$

$$\begin{aligned} \therefore \sqrt{x^2-1} \frac{d}{d} &= 5 \left(+\sqrt{x^2-1} \right)^4 \left(+\sqrt{x^2-1} \right) \\ &= 5 \end{aligned}$$

$$\therefore (x^2-1) \left(\frac{d}{d} \right)^2 = 25x^2$$

$$\therefore (x^2-1) \times \frac{2d}{d} \cdot \frac{d^2}{d^2} + \left(\frac{d}{d} \right)^2 (2x) = 25 \times 2 \frac{d}{d}$$



Dividing both sides by $2\frac{d}{d}$, we get

$$(d^2 - 1)\frac{d^2}{d^2} + \frac{d}{d} = 25$$

$$\therefore k = 25$$

12. $\sqrt{x^2 + 2} = ae^{\tan^{-1}(-)} \dots(i)$

Diff. w.r.t. x , we get

$$\frac{1}{2\sqrt{x^2 + 2}} \left(2x + 2 \frac{d}{d} \right)$$

$$= ae^{\tan^{-1}(-)} \cdot \frac{1}{1 + \frac{d}{d}} \left(\frac{\frac{d}{d}}{\frac{d}{d}} \right)$$

$$\therefore \frac{1}{\sqrt{x^2 + 2}} \left(x + \frac{d}{d} \right) = \sqrt{x^2 + 2} \left(\frac{\frac{d}{d}}{\frac{d}{d}} \right)$$

$$\dots \left[\because ae^{\tan^{-1}(-)} = \sqrt{x^2 + 2} \right]$$

$$\therefore x + \frac{d}{d} = \frac{d}{d} \dots(ii)$$

Diff. w.r.t. x , we get

$$1 + \frac{d^2}{d^2} + \left(\frac{d}{d} \right)^2 = \frac{d^2}{d^2} + \frac{d}{d} - \frac{d}{d}$$

$$\therefore 1 + \frac{d^2}{d^2} + \left(\frac{d}{d} \right)^2 = \frac{d^2}{d^2}$$

$$\therefore \left(\frac{d}{d} \right)^2 = - \left(1 + \left(\frac{d}{d} \right)^2 \right)$$

From (i), when $x = 0$, $y = ae^{\frac{\pi}{2}}$

From (ii), when $x = 0$, $\frac{d}{d} = -1$

$$\therefore ae^{\frac{\pi}{2}} \frac{d^2}{d^2} = -2$$

$$\therefore \left(\frac{d^2}{d^2} \right)_{=0} = -\frac{2}{a} e^{-\frac{\pi}{2}}$$

13. $f(x) = \begin{vmatrix} f & g & h \\ f' & g' & h' \\ f'' & g'' & h'' \end{vmatrix}$

$$\therefore f'(x) = \begin{vmatrix} f' & g' & h' \\ f'' & g'' & h'' \\ f''' & g''' & h''' \end{vmatrix} + \begin{vmatrix} f & g & h \\ f'' & g'' & h'' \\ f''' & g''' & h''' \end{vmatrix} + \begin{vmatrix} f & g & h \\ f' & g' & h' \\ f''' & g''' & h''' \end{vmatrix}$$

$$= 0 + 0 + 0$$

$\dots[\because f, g, h$ are polynomials of 2nd degree, $f''' = g''' = h''' = 0]$

$$= 0$$

14. $\begin{vmatrix} 1 & 2 \\ 3 & 4 & 5 \\ 6 & 7 & 8 \end{vmatrix} = \begin{vmatrix} \cos a & -a \sin a & -a^2 \cos a \\ a^3 \sin a & a^4 \cos a & -a^5 \sin a \\ -a^6 \cos a & a^7 \sin a & a^8 \cos a \end{vmatrix}$

$$= -a^2 \times 0 \dots[\because C_1 \equiv C_3]$$

$$= 0$$

15. $= \sin[\cos^{-1}\{\sin(\cos^{-1} x)\}]$

$$= \sin\left[\cos^{-1}\left\{\sin\left(\frac{\pi}{2} - \sin^{-1} x\right)\right\}\right]$$

$$= \sin[\cos^{-1}(\cos(\sin^{-1} x))]$$

$$= \sin(\sin^{-1} x) = x$$

$$\therefore \frac{d}{d} = 1$$

$$\therefore \left(\frac{d}{d} \right)_{=\frac{1}{2}} = 1$$

16. $8f(x) + 6f\left(\frac{1}{x}\right) = x + 5 \dots(i)$

Replacing x by $\frac{1}{x}$, we get

$$8f\left(\frac{1}{x}\right) + 6f(x) = \frac{1}{x} + 5$$

$$\therefore 6f(x) + 8f\left(\frac{1}{x}\right) = \frac{1}{x} + 5 \dots(ii)$$

(i) $\times 8$ - (ii) $\times 6$ gives

$$64f(x) - 36f\left(\frac{1}{x}\right) = 8x + 40 - \frac{6}{x} - 30$$

$$\therefore 28f(x) = 8x - \frac{6}{x} + 10$$



$$\text{Given, } = {}^2f(x) = \frac{x^2}{28} \left(8x^3 - 6x + 10 \right)$$

$$\therefore = \frac{1}{28} (8x^3 - 6x + 10)$$

$$\therefore \frac{d}{dx} = \frac{1}{28} (24x^2 - 6 + 20)$$

$$\therefore \left(\frac{d}{dx} \right)_{x=1} = \frac{1}{28} (24 - 6 + 20) = -\frac{2}{28} = -\frac{1}{14}$$

17. $f(x^3) = x^5$
Diff. w.r.t. x , we get
 $f'(x^3) \cdot 3x^2 = 5x^4$

$$\therefore f'(x^3) = \frac{5x^2}{3}$$

$$\therefore f'(27) = f'(3^3) = \frac{5}{3} (3)^2 = 15$$

18. Since, $g(x)$ is the inverse of $f(x)$.

$$\therefore f[g(x)] = x$$

$$\Rightarrow f'(g(x))g'(x) = 1$$

$$\Rightarrow f'(g(1))g'(1) = 1$$

$$\Rightarrow g'(1) = \frac{1}{f'(g(1))} \quad \dots(i)$$

$$f(x) = x^3 + e^{1/2}$$

$$\therefore f(0) = 1$$

$$\Rightarrow 0 = f^{-1}(1)$$

$$\Rightarrow g(1) = 0 \quad \dots[\because g(x) = f^{-1}(x)(\text{given})]$$

From (i), we get

$$g'(1) = \frac{1}{f'(0)}$$

$$\text{Now, } f(x) = x^3 + e^{1/2}$$

$$\Rightarrow f'(x) = 3x^2 + \frac{1}{2}e^{1/2}$$

$$\Rightarrow f'(0) = \frac{1}{2}$$

$$\therefore g'(1) = \frac{1}{1/2} = 2$$

19. $y = f(x^3)$

$$\therefore \frac{dy}{dx} = f'(x^3) \cdot 3x^2 = 3x^2 \tan(x^3)$$

$$z = g(x^5)$$

$$\therefore \frac{dz}{dx} = g'(x^5) \cdot 5x^4 = 5x^4 \sec(x^5)$$

$$\therefore \frac{dz}{dy} = \frac{\frac{dz}{dx}}{\frac{dy}{dx}} = \frac{5x^4 \sec(x^5)}{3x^2 \tan(x^3)} = \frac{5 \tan^3(x^3)}{5^2 \sec^5(x^5)}$$

20. $\sqrt{1-x^6} + \sqrt{1-x^6} = a^3(x^3 - x^3)$

$$\text{Put } x^3 = \sin \alpha \text{ and } x^3 = \sin \beta$$

$$\therefore \sqrt{1-\sin^2 \alpha} + \sqrt{1-\sin^2 \beta} = a^3(\sin \alpha - \sin \beta)$$

$$\therefore \cos \alpha + \cos \beta = a^3(\sin \alpha - \sin \beta)$$

$$\therefore 2 \cos\left(\frac{\alpha+\beta}{2}\right) \cos\left(\frac{\alpha-\beta}{2}\right)$$

$$= a^3 \cdot 2 \sin\left(\frac{\alpha-\beta}{2}\right) \cos\left(\frac{\alpha+\beta}{2}\right)$$

$$\therefore \cot\left(\frac{\alpha-\beta}{2}\right) = a^3$$

$$\therefore \alpha - \beta = 2 \cot^{-1} a^3$$

$$\therefore \sin^{-1} x^3 - \sin^{-1} x^3 = \text{constant}$$

Diff. w.r.t. x , we get

$$\frac{3x^2}{\sqrt{1-x^6}} - \frac{3x^2}{\sqrt{1-x^6}} \cdot \frac{d}{dx} = 0$$

$$\therefore \frac{d}{dx} = \frac{2}{2} \sqrt{\frac{1-x^6}{1-x^6}}$$

21. Let $f(x) = p^2 + q + r$

$$\therefore f(1) = f(-1) \Rightarrow p + q + r = p - q + r \Rightarrow q = 0$$

$$\therefore f(x) = p^2 + r$$

$$\Rightarrow f'(x) = 2p$$

$$\Rightarrow f'(a) = 2ap, f'(b) = 2bp \text{ and } f'(c) = 2cp$$

Since, a, b, c are in A.P.

$$\therefore 2ap, 2bp, 2cp \text{ are in A.P.}$$

$$\Rightarrow f'(a), f'(b), f'(c) \text{ are in A.P.}$$

22. $\frac{d}{d\theta} = \sec \theta \tan \theta + \sin \theta$

$$\text{and } \frac{d}{d\theta} = n \sec^{n-1} \theta \cdot \sec \theta \tan \theta - n \cos^{n-1} \theta \cdot (-\sin \theta)$$

$$= n \sec^n \theta \tan \theta + n \cos^{n-1} \theta \sin \theta$$

$$\therefore \frac{d}{dx} = \frac{\frac{d}{d\theta}}{\frac{d}{d\theta}} = \frac{n \sec^n \theta \tan \theta + n \cos^{n-1} \theta \sin \theta}{\sec \theta \tan \theta + \sin \theta}$$

Dividing N^r and D^r by $\tan \theta$, we get

$$\frac{d}{dx} = \frac{n(\sec^n \theta + \cos^n \theta)}{\sec \theta + \cos \theta}$$



$$\Rightarrow \left(\frac{d}{d}\right)^2 = \frac{n^2(\sec^n \theta + \cos^n \theta)^2}{(\sec \theta + \cos \theta)^2}$$

$$= \frac{n^2[(\sec^n \theta - \cos^n \theta)^2 + 4\sec^n \theta \cos^n \theta]}{(\sec \theta - \cos \theta)^2 + 4\sec \theta \cos \theta}$$

$$= \frac{n^2(x^2 + 4)}{x^2 + 4}$$

$$\therefore (x^2 + 4)\left(\frac{d}{d}\right)^2 = n^2(x^2 + 4)$$

23. $f(x) = \begin{vmatrix} x^2 & \sin x & \cos x \\ 2 & \tan x & -3 \\ 2 & \sin 2 & 5 \end{vmatrix}$

$$\therefore f'(x) = \begin{vmatrix} 1 & \sin x & \cos x \\ 2 & \tan x & -3 \\ 2 & \sin 2 & 5 \end{vmatrix}$$

$$+ \begin{vmatrix} x^2 & \cos x & \cos x \\ 2 & \sec^2 x & -3 \\ 2 & 2\cos 2 & 5 \end{vmatrix} + \begin{vmatrix} x^2 & \sin x & -\sin x \\ 2 & \tan x & -3 \\ 2 & \sin 2 & 5 \end{vmatrix}$$

$$\therefore \frac{f'(x)}{x^2} = \begin{vmatrix} 1 & \sin x & \cos x \\ 2 & \tan x & -3 \\ 2 & \sin 2 & 5 \end{vmatrix}$$

$$+ \begin{vmatrix} 1 & \cos x & \cos x \\ 2 & \sec^2 x & -3 \\ 2 & 2\cos 2 & 5 \end{vmatrix} + \begin{vmatrix} 1 & \sin x & -\sin x \\ 2 & \tan x & -3 \\ 2 & \sin 2 & 5 \end{vmatrix}$$

$$\therefore \lim_{x \rightarrow 0} \frac{f'(x)}{x^2} = \begin{vmatrix} 1 & 0 & 1 \\ 2 & 1 & 0 \\ 2 & 0 & 0 \end{vmatrix} + \begin{vmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 2 & 2 & 0 \end{vmatrix} + \begin{vmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 2 & 0 & 5 \end{vmatrix}$$

$$= -2 - 2 + 0 = -4$$

24. Since, g is the inverse of f.

$$\therefore f[g(x)] = x$$

Diff. w.r.t. x, we get

$$f'(g(x)) g'(x) = 1$$

$$\therefore g'(x) = \frac{1}{f'(g(x))} = 1 + [g(x)]^5$$

25.
$$= \frac{\sin x}{\sin x \sin 2} + \frac{\sin x}{\sin 2 \sin 3} + \dots + \frac{\sin x}{\sin n \sin(n+1)}$$

$$= \frac{\sin(2-x)}{\sin x \sin 2} + \frac{\sin(3-x)}{\sin 2 \sin 3} + \dots + \frac{\sin((n+1)-x)}{\sin n \sin(n+1)}$$

$$= \frac{\sin 2 \cos x}{\sin x \sin 2} - \frac{\cos 2 \sin x}{\sin x \sin 2} + \frac{\sin 3 \cos x}{\sin 2 \sin 3}$$

$$- \frac{\cos 3 \sin x}{\sin 2 \sin 3} + \dots + \frac{\sin(n+1) \cos x}{\sin n \sin(n+1)} - \frac{\cos(n+1) \sin x}{\sin n \sin(n+1)}$$

$$= \cot x - \cot 2 + \cot 2 - \cot 3$$

$$+ \dots + \cot n - \cot(n+1)$$

$$\therefore = \cot x - \cot(n+1)$$

$$\therefore \frac{d}{dx} = -\operatorname{cosec}^2 x - [-\operatorname{cosec}^2(n+1)](n+1)$$

$$= (n+1) \operatorname{cosec}^2(n+1) - \operatorname{cosec}^2 x$$

26. If $|r| < 1$, $a + ar + ar^2 + \dots + \infty = \frac{a}{1-r}$

$$\therefore \sin^2 x + \sin^4 x + \sin^6 x + \dots = \frac{\sin^2 x}{1-\sin^2 x}$$

$$= \frac{\sin^2 x}{\cos^2 x} = \tan^2 x$$

$$\therefore = e^{\tan^2 x}$$

$$\therefore \frac{d}{dx} = e^{\tan^2 x} \cdot 2 \tan x \sec^2 x = 2e^{\tan^2 x} \tan x \sec^2 x$$

27.
$$= \tan^{-1} \frac{1}{1+x^2} + \tan^{-1} \frac{1}{x^2+3} + \frac{1}{x^2+3} + \frac{1}{x^2+5} + \dots$$
 to n terms

$$= \tan^{-1} \frac{1}{1+(1+x^2)} + \tan^{-1} \frac{1}{1+(x^2+2)(x^2+1)}$$

$$+ \tan^{-1} \frac{1}{1+(x^2+3)(x^2+2)} + \dots$$
 to n terms

$$= \tan^{-1} \left[\frac{(x^2+1) - (x^2+2)}{1+(x^2+1)(x^2+2)} \right] + \tan^{-1} \left[\frac{(x^2+2) - (x^2+1)}{1+(x^2+2)(x^2+1)} \right]$$

$$+ \tan^{-1} \left[\frac{(x^2+3) - (x^2+2)}{1+(x^2+3)(x^2+2)} \right] + \dots$$
 to n terms

$$= \tan^{-1}(x^2+1) - \tan^{-1}(x^2+2) + \tan^{-1}(x^2+2)$$

$$- \tan^{-1}(x^2+1) + \tan^{-1}(x^2+3) - \tan^{-1}(x^2+2)$$

$$+ \dots + \tan^{-1}(x^2+n) - \tan^{-1}(x^2+(n-1))$$

$$\therefore = \tan^{-1}(x^2+n) - \tan^{-1}(x^2+1)$$

$$\therefore \frac{d}{dx} = \frac{1}{1+(x^2+n)^2} - \frac{1}{1+x^2}$$

$$\therefore \left(\frac{d}{dx}\right)_{x=0} = \frac{1}{1+n^2} - 1 = \frac{1-1-n^2}{1+n^2} = -\frac{n^2}{1+n^2}$$



$$\begin{aligned}
 28. \quad &= a \sin(b + c) \\
 \therefore \quad &_1 = a \cos(b + c).b = ab \sin\left(\frac{\pi}{2} + b + c\right) \\
 \therefore \quad &_2 = -ab \sin(b + c).b = ab^2 \sin(\pi + b + c) \\
 &_3 = -ab^2 \cos(b + c).b = ab^3 \sin\left(\frac{3\pi}{2} + b + c\right) \\
 &_4 = -ab^3(-\sin(b + c).b) = ab^4 \sin(2\pi + b + c) \\
 &= ab^4 \sin\left(\frac{4\pi}{2} + b + c\right)
 \end{aligned}$$

$$\text{In general, } _n = ab^n \sin\left(\frac{n\pi}{2} + b + c\right)$$

$$\begin{aligned}
 29. \quad &f(x) = x^n \\
 &f'(x) = n x^{n-1} \\
 &f''(x) = n(n-1) x^{n-2} \\
 &f'''(x) = n(n-1)(n-2) x^{n-3} \\
 \therefore \quad &f(1) = 1^n = 1 = {}^n C_0 \\
 &\frac{f'(1)}{1!} = \frac{n(1)^{n-1}}{1} = n = {}^n C_1 \\
 &\frac{f''(1)}{2!} = \frac{n(n-1)(1)^{n-2}}{2!} = \frac{n(n-1)}{2!} = {}^n C_2 \\
 &\frac{f'''(1)}{3!} = \frac{n(n-1)(n-2)(1)^{n-3}}{3!} = \frac{n(n-1)(n-2)}{3!} \\
 &= {}^n C_3 \\
 \therefore \quad &f(1) - \frac{f'(1)}{1!} + \frac{f''(1)}{2!} - \frac{f'''(1)}{3!} + \dots + (-1)^n \frac{f^n(1)}{n!} \\
 &= {}^n C_0 - {}^n C_1 + {}^n C_2 - {}^n C_3 + \dots + (-1)^n {}^n C_n \\
 &= 0
 \end{aligned}$$

$$\begin{aligned}
 30. \quad &p = a^2 \cos^2 \theta + b^2 \sin^2 \theta \\
 \therefore \quad &\frac{dp}{d\theta} = a^2 \cdot 2 \cos \theta (-\sin \theta) + b^2 \cdot 2 \sin \theta \cos \theta \\
 &= (b^2 - a^2) \sin 2\theta \\
 \therefore \quad &\frac{d^2 p}{d\theta^2} = 2(b^2 - a^2) \cos 2\theta \\
 &= 2(b^2 - a^2) (\cos^2 \theta - \sin^2 \theta) \\
 \therefore \quad &4p + \frac{d^2 p}{d\theta^2} = 4a^2 \cos^2 \theta + 4b^2 \sin^2 \theta \\
 &\quad + 2(b^2 - a^2) (\cos^2 \theta - \sin^2 \theta) \\
 &= \cos^2 \theta (4a^2 + 2b^2 - 2a^2) \\
 &\quad + \sin^2 \theta (4b^2 - 2b^2 + 2a^2)
 \end{aligned}$$

$$\begin{aligned}
 &= \cos^2 \theta (2a^2 + 2b^2) + \sin^2 \theta (2a^2 + 2b^2) \\
 &= (2a^2 + 2b^2) (\cos^2 \theta + \sin^2 \theta) \\
 &= 2a^2 + 2b^2 \\
 &= 2(a^2 + b^2) \\
 &= 2c^2 \quad \dots [\because a^2 + b^2 = c^2 \text{ (given)}]
 \end{aligned}$$



Hints



Classical Thinking

1. $y = 3t^2 + 1, \quad x = t^3 - 1$

$$\therefore \frac{dy}{dt} = 6t, \quad \frac{dx}{dt} = 3t^2$$

$$\therefore \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{6t}{3t^2} = \frac{2}{t}$$

$$\therefore \left(\frac{dy}{dx}\right)_{t=1} = \frac{1}{2}$$

2. $y = 3x^2 - 1$

$$\therefore \frac{dy}{dx} = 6x - 1$$

$$\therefore \left(\frac{dy}{dx}\right)_{x=2} = 3(2)^2 - 1 = 11$$

$$\therefore \text{slope of normal at } x = 2 = -\frac{1}{\left(\frac{dy}{dx}\right)_{x=2}} = -\frac{1}{11}$$

3. If the tangent is perpendicular to X-axis, then $\theta = 90^\circ$

$$\therefore \cot \theta = 0$$

$$\Rightarrow \frac{1}{\tan \theta} = 0 \Rightarrow \frac{dy}{dx} = 0$$

4. $y = 3x^3 - 3x^2 - 9x + 5 \Rightarrow \frac{dy}{dx} = 9x^2 - 6x - 9$

Since, the tangent is parallel to X-axis.

$$\therefore \frac{dy}{dx} = 0$$

$$\Rightarrow 9x^2 - 6x - 9 = 0 \Rightarrow x = -1, 3$$

5. $y^2 = -4x$

Differentiating both sides w.r.t. x , we get

$$2y \frac{dy}{dx} = -4$$

$$\Rightarrow \frac{dy}{dx} = \frac{-2}{y}$$

$\therefore m = \text{Slope of the tangent at } (-4, -4)$

$$= \left(\frac{dy}{dx}\right)_{(-4,-4)} = 2$$

\therefore equation of the tangent at $(-4, -4)$ is

$$y - y_1 = m(x - x_1) \\ \Rightarrow y + 4 = 2(x + 4) \\ \Rightarrow 2x - y + 4 = 0$$

6. $\sqrt{x} + \sqrt{y} = a$

Differentiating both sides w.r.t. x , we get

$$\frac{1}{2\sqrt{x}} + \frac{1}{2\sqrt{y}} \cdot \frac{dy}{dx} = 0$$

$$\Rightarrow \frac{dy}{dx} = -\frac{\sqrt{y}}{\sqrt{x}}$$

$$\text{At } \left(\frac{a^2}{4}, \frac{a^2}{4}\right), \frac{dy}{dx} = -\sqrt{\frac{\frac{a^2}{4}}{\frac{a^2}{4}}} = -1$$

\therefore Equation of the tangent at $\left(\frac{a^2}{4}, \frac{a^2}{4}\right)$ is

$$y - \frac{a^2}{4} = -1 \left(x - \frac{a^2}{4}\right)$$

$$\Rightarrow x + y = \frac{a^2}{2}$$

7. $y = x^2 - 2x + 1$

$$\therefore \frac{dy}{dx} = 2x - 2$$

$\therefore m = \text{slope of the normal at } (0, 1)$

$$= \frac{-1}{\left(\frac{dy}{dx}\right)_{(0,1)}} = \frac{-1}{2(0) - 2} = \frac{1}{2}$$

\therefore Equation of the normal at $(0, 1)$ is

$$y - 1 = m(x - 0)$$

$$\Rightarrow y - 1 = \frac{1}{2}(x - 0)$$

$$\Rightarrow x - 2y + 2 = 0$$

8. $y = \sin \frac{\pi}{2} \Rightarrow \frac{dy}{dx} = \frac{\pi}{2} \cos \frac{\pi}{2} \Rightarrow \left(\frac{dy}{dx}\right)_{(1,1)} = 0$

\therefore Equation of the normal at $(1, 1)$ is $x = 1$



$$9. \quad \text{At } t = \frac{\pi}{4}, \quad \frac{d}{dt} = \frac{2}{\sqrt{2}} = \sqrt{2}$$

$$= 2 \sin$$

$$\therefore \frac{d}{dt} = 2 \cos$$

$$\therefore \left(\frac{d}{dt} \right)_{t=\frac{\pi}{4}} = \sqrt{2}$$

\therefore Equation of the tangent at $\left(\frac{\pi}{4}, \sqrt{2} \right)$ is

$$-\sqrt{2} = \frac{1}{\sqrt{2}} \left(-\frac{\pi}{4} \right)$$

$$10. \quad \text{At } t = \frac{\pi}{2},$$

$$= 4 + \cos^2 \frac{\pi}{2} = 4$$

$$= 4 + \cos^2$$

$$\therefore \frac{d}{dt} = 2 \cos (-\sin)$$

$$\therefore \left(\frac{d}{dt} \right)_{t=\frac{\pi}{2}} = 2 \cos \frac{\pi}{2} \left(-\sin \frac{\pi}{2} \right) = 0$$

\therefore Equation of the tangent at $\left(\frac{\pi}{2}, 4 \right)$ is

$$-4 = 0 \left(-\frac{\pi}{2} \right)$$

$$\therefore -4 = 0 \Rightarrow = 4$$

$$11. \quad \text{At } t = \frac{\pi}{2},$$

$$= \frac{\pi}{2} - \sin \frac{\pi}{2} \cos \frac{\pi}{2} = \frac{\pi}{2}$$

$$= -\sin \cos$$

$$\therefore \frac{d}{dt} = 1 - \cos \cos - \sin (-\sin)$$

$$= 1 - \cos^2 + \sin^2$$

$$\therefore \left(\frac{d}{dt} \right)_{t=\frac{\pi}{2}} = 1 - \cos^2 \frac{\pi}{2} + \sin^2 \frac{\pi}{2} = 2$$

\therefore Equation of the tangent at $\left(\frac{\pi}{2}, \frac{\pi}{2} \right)$ is

$$-\frac{\pi}{2} = 2 \left(-\frac{\pi}{2} \right)$$

$$\Rightarrow = 2 - \frac{\pi}{2}$$

$$12. \quad \text{At } t = \frac{\pi}{3}, \quad \frac{d}{dt} = 2 \sin \frac{\pi}{3} + \sin \frac{2\pi}{3} = \frac{3\sqrt{3}}{2}$$

$$= 2 \sin + \sin 2$$

$$\therefore \frac{d}{dt} = 2 \cos + 2 \cos 2$$

$$\therefore \left(\frac{d}{dt} \right)_{t=\frac{\pi}{3}} = 2 \cos \frac{\pi}{3} + 2 \cos \frac{2\pi}{3} = 0$$

\therefore Equation of the tangent at $\left(\frac{\pi}{3}, \frac{3\sqrt{3}}{2} \right)$ is

$$-\frac{3\sqrt{3}}{2} = 0 \left(-\frac{\pi}{3} \right) \Rightarrow 2 = 3\sqrt{3}$$

$$13. \quad \text{At } t = \frac{\pi}{4}, \quad \frac{d}{dt} = \frac{2}{\sqrt{2}} = \sqrt{2}$$

$$= 2 \cos$$

$$\therefore \frac{d}{dt} = -2 \sin$$

$$\therefore \left(\frac{d}{dt} \right)_{t=\frac{\pi}{4}} = -\sqrt{2}$$

\therefore Equation of the normal at $\left(\frac{\pi}{4}, \sqrt{2} \right)$ is

$$-\sqrt{2} = \frac{1}{\sqrt{2}} \left(-\frac{\pi}{4} \right)$$

$$14. \quad s = 3t^2 + 2t - 5$$

$$\therefore \frac{ds}{dt} = 6t + 2$$

$$\therefore \text{Acceleration} = \frac{d^2s}{dt^2} = 6$$

$$15. \quad s = 2t^2 - 3t + 1$$

$$\therefore v = \frac{ds}{dt} = 4t - 3 \quad \therefore \frac{d^2s}{dt^2} = 4$$

$$16. \quad \frac{ds}{dt} = \text{velocity} = 45 + 22t - 3t^2$$

When particle will come to rest, then $v = 0$

$$\Rightarrow 3t^2 - 22t - 45 = 0 \Rightarrow t = 9 \quad \dots \left[\because t \neq -\frac{5}{3} \right]$$

$$17. \quad \text{Given, } s = a \sin t + b \cos 2t$$

$$\therefore \frac{ds}{dt} = a \cos t - 2b \sin 2t$$

$$\therefore \frac{d^2s}{dt^2} = -a \sin t - 4b \cos 2t$$

$$\text{At } t = 0, \quad \frac{d^2s}{dt^2} = -a \sin 0^\circ - 4b \cos 0^\circ = -4b$$




18. $s = 2t^3 - 9t^2 + 12t$
 $\Rightarrow \frac{ds}{dt} = 6t^2 - 18t + 12$
 $\Rightarrow \frac{d^2s}{dt^2} = 12t - 18 = \text{acceleration}$
 When acceleration of the particle will be zero,
 $12t - 18 = 0$
 $\Rightarrow t = \frac{3}{2} \text{ sec}$
 Hence, the acceleration of the particle will be zero after $\frac{3}{2}$ sec.
19. $s = \frac{1}{2}gt^2 \Rightarrow \frac{ds}{dt} = gt \Rightarrow \frac{d^2s}{dt^2} = g$
 \therefore the acceleration of the stone is uniform.
20. $\frac{dr}{dt} = 3$
 $A = \pi r^2 \Rightarrow \frac{dA}{dt} = 2\pi r \frac{dr}{dt}$
 $\Rightarrow \left(\frac{dA}{dt}\right)_{r=10} = 2\pi \times 10 \times 3 = 60\pi \text{ cm}^2/\text{sec}$
21. $A = s^2$
 $\therefore \frac{dA}{dt} = 2s \frac{ds}{dt}$
 $\therefore \frac{dA}{dt} = 2 \times 10 \times 0.5 = 2 \times 5 = 10 \text{ cm}^2/\text{sec}$
22. $V = 5 - \frac{2}{6}$
 $\Rightarrow \frac{dV}{dt} = 5 \frac{d}{dt} - \frac{2}{3} \cdot \frac{d}{dt}$
 $\Rightarrow \frac{d}{dt} = \frac{\frac{dV}{dt}}{\left(5 - \frac{2}{3}\right)}$
 $\Rightarrow \left(\frac{d}{dt}\right)_{=2} = \frac{5}{5 - \frac{2}{3}} = \frac{15}{13} \text{ cm/sec}$
23. Let $f(x) = \sqrt{x}$
 $\therefore f'(x) = \frac{1}{2\sqrt{x}}$
 Here, $a = 25$ and $h = 0.2$
 $\therefore f(a) = f(25) = \sqrt{25} = 5$
 and $f'(a) = f'(25) = \frac{1}{2\sqrt{25}} = \frac{1}{10}$


- $\therefore f(a+h) \approx f(a) + hf'(a)$
 $\approx 5 + (0.2)\left(\frac{1}{10}\right)$
 $\approx 5 + 0.02$
 $\therefore \sqrt{25.2} \approx 5.02$
24. Let $f(x) = x^{\frac{1}{3}}$
 $\therefore f'(x) = \frac{1}{3}x^{-\frac{2}{3}} = \frac{1}{3x^{\frac{2}{3}}}$
 Here, $a = 27$ and $h = 2$
 $\therefore f(a+h) \approx f(a) + hf'(a)$
 $\approx (27)^{\frac{1}{3}} + 2\left[\frac{1}{3(27)^{\frac{2}{3}}}\right]$
 $\approx 3 + 2\left(\frac{1}{27}\right)$
 $\approx 3 + 0.07407$
 $\therefore (29)^{\frac{1}{3}} \approx 3.07407$
25. If Rolle's theorem is true for any function $f(x)$ in $[a, b]$.
 Then $f(a) = f(b)$
 Only option (B) satisfies this condition.
26. According to Lagrange's mean value theorem, in interval $[a, b]$ for $f(x)$,
 $f'(c) = \frac{f(b) - f(a)}{b - a}$, where $a < c < b$
 $\therefore a < c < b$
27. $f(x) = 2 - 3x$
 $\therefore f'(x) = -3 < 0$
 $\therefore f(x)$ is a decreasing function.
28. $f(x) = x^2 \Rightarrow f'(x) = 2x$
 For increasing function,
 $f'(x) > 0$
 $\Rightarrow 2x > 0$
 $\Rightarrow x \in (0, \infty)$
29. $f(x) = ax + b$
 $\therefore f'(x) = a$
 For $f(x)$ to be decreasing,
 $f'(x) < 0$
 $\Rightarrow a < 0$
30. $f(x) = 5^{-x}$
 $\therefore f'(x) = -5^{-x} \log_e 5 = -\frac{\log_e 5}{5}$
 $\Rightarrow f'(x) < 0$ for all x
 i.e., $f(x)$ is decreasing for all x .




32. Let $f(x) = 4^x - 4 \Rightarrow f'(x) = 4^x - 4$
 For $f(x)$ to be decreasing, $f'(x) < 0$
 $\Rightarrow 4^x - 4 < 0 \Rightarrow 4^x < 4$
 $\Rightarrow x \in (-\infty, 1)$

33. $f(x) = 4^{4x} - 2 + 1$
 $\therefore f'(x) = 16^{3x} - 2$
 For $f(x)$ to be increasing,
 $f'(x) > 0$
 $\Rightarrow 16^{3x} - 2 > 0$
 $\Rightarrow 4^{3x} > \frac{1}{8}$
 $\Rightarrow 3x > \frac{1}{2}$

34. $f(x) = 2^{3x} + 9^{2x} + 12 + 20$
 $\therefore f'(x) = 6^{2x} + 18^{2x} + 12$ 
 For $f(x)$ to be increasing,
 $f'(x) > 0$
 $\Rightarrow 2^{2x} + 3^{2x} + 2 > 0$
 $\Rightarrow (x + 2)(x + 1) > 0$
 $\Rightarrow x \in (-\infty, -2) \cup (-1, \infty)$

35. $f(x) = 2^{3x} - 3^{2x} - 12 + 12$
 $\therefore f'(x) = 6^{2x} - 6^{2x} - 12$
 For $f(x)$ to be increasing,
 $f'(x) > 0$
 $\Rightarrow 2^{2x} - 3^{2x} > 0$ 
 $\Rightarrow (x - 2)(x + 1) > 0$
 $\Rightarrow x \in (-\infty, -1) \cup (2, \infty)$

36. $f(x) = 3^{3x} - 6^{2x} + 9^{2x} + 3$
 $\therefore f'(x) = 3^{2x} - 12^{2x} + 9^{2x}$ 
 For $f(x)$ to be decreasing,
 $f'(x) < 0$
 $\Rightarrow 3(2^{2x} - 4^{2x} + 3) < 0$
 $\Rightarrow (x - 3)(x - 1) < 0$
 $\Rightarrow x \in (1, 3)$

37. Let $f(x) = 2^{3x} - 6^{2x} + 5$
 $\therefore f'(x) = 6^{2x} - 6^{2x}$
 For $f(x)$ to be increasing, $f'(x) > 0$
 $\Rightarrow 6^{2x} - 6^{2x} > 0 \Rightarrow (x - 1)(x + 1) > 0$
 $\Rightarrow x > 1$ or $x < -1$

38. Let $f(x) = \frac{1}{1+x^2}$
 $\therefore f'(x) = -\frac{2x}{(1+x^2)^2}$
 For $f(x)$ to be decreasing,
 $f'(x) < 0 \Rightarrow -\frac{2x}{(1+x^2)^2} < 0$
 $\Rightarrow x > 0 \Rightarrow x \in (0, \infty)$

39. Let $f(x) = \log(\sin x) \Rightarrow f'(x) = \cot x$
 \therefore the given function is increasing in the interval
 $\left(0, \frac{\pi}{2}\right)$.

40. $f(x) = 2^{3x} - 3^{2x} - 36 + 7$
 $\therefore f'(x) = 6^{2x} - 6^{2x} - 36$
 For decreasing function, $f'(x) < 0$
 $\Rightarrow 2^{2x} - 3^{2x} - 6 < 0$
 $\Rightarrow (x - 3)(x + 2) < 0$
 $\Rightarrow x \in (-2, 3)$

41. $f(x) = 2^{3x} - 3^{2x} - 12 + 5$
 $\therefore f'(x) = 6^{2x} - 6^{2x} - 12$
 For maximum or minimum,
 $f'(x) = 0$
 $\Rightarrow 2^{2x} - 3^{2x} - 2 = 0$
 $\Rightarrow (x - 2)(x + 1) = 0$
 $\Rightarrow x = 2, -1$
 Now, $f''(x) = 12^{2x} - 6$
 $\therefore f''(2) = 18 > 0$
 $\therefore f(x)$ is minimum at $x = 2$.

43. $f(x) = 7 - 20x + 11x^2$
 $\therefore f'(x) = -20 + 22x$
 For maximum or minimum,
 $f'(x) = 0 \Rightarrow -20 + 22x = 0$
 $\Rightarrow x = 10/11$
 Now, $f''(x) = 22 > 0$
 $\therefore f(x)$ is minimum at $x = \frac{10}{11}$.

$\therefore [f(x)]_{\min} = f\left(\frac{10}{11}\right)$
 $= 7 - \frac{200}{11} + \frac{100 \times 11}{121} = -\frac{23}{11}$

44. Let $f(x) = 2^{2x} + x - 1$
 $\therefore f'(x) = 4^{2x} + 1$
 For maximum or minimum,
 $f'(x) = 0 \Rightarrow 4^{2x} = -1$
 Now, $f''(x) = 4^{2x} > 0$
 $\therefore f(x)$ is minimum at $x = \frac{-1}{4}$.

$\therefore [f(x)]_{\min} = \left[f\left(-\frac{1}{4}\right) \right] = \frac{2}{16} - \frac{1}{4} - 1 = \frac{-9}{8}$



45. $f(x) = 2x^3 - 3x^2 - 12x + 4$
 $\therefore f'(x) = 6x^2 - 6x - 12$
 For maximum or minimum,
 $f'(x) = 0 \Rightarrow 2x^2 - x - 2 = 0 \Rightarrow x = 2, -1$
 Now, $f''(x) = 12x - 6$
 $\therefore f''(2) = 18 > 0$ and $f''(-1) = -18 < 0$
 \therefore the given function has one maximum and one minimum.

46. $y = 1 - \cos x$
 $\therefore y' = \sin x$
 For maximum or minimum,
 $y' = 0 \Rightarrow \sin x = 0 \Rightarrow x = 0, \pi$
 Now, $y'' = \cos x$
 $\Rightarrow y''(0) = 1 > 0$ and $y''(\pi) = -1 < 0$
 $\therefore y$ is maximum when $x = \pi$.



Critical Thinking

1. $y = 15x$
 $\therefore y' = 15$
 $\therefore y' = -\frac{15}{2}$
 At $(3, 5)$, $y' = -\frac{15}{9}$
 \therefore Slope of normal at $(3, 5) = \frac{9}{15}$
 $\therefore \theta = \tan^{-1}\left(\frac{9}{15}\right)$
 2. $y^2 = 2x$
 Differentiating both sides w.r.t. x , we get
 $2y \frac{dy}{dx} = 2$
 $\Rightarrow \frac{dy}{dx} = \frac{1}{y}$
 $\therefore \left(\frac{dy}{dx}\right)_{(1, \frac{1}{2})} = 1$
 $\Rightarrow \tan \theta = 1$
 $\Rightarrow \theta = 45^\circ$ [$\because \tan 45^\circ = 1$]
 3. $x^3 - 8a^2 = 0$ (i)
 Differentiating w.r.t. x , we get
 $3x^2 - 8a^2 \frac{dx}{dx} = 0$
 $\Rightarrow \frac{dx}{dx} = \frac{3x^2}{8a^2}$

\therefore Slope of the normal $= -\frac{1}{\frac{dx}{dy}} = -\frac{1}{\frac{1}{8a^2}} = -\frac{8a^2}{1}$
 According to the given condition,
 $-\frac{8a^2}{3^2} = \frac{-2}{3}$
 $\Rightarrow 4a^2 = 2$
 $\Rightarrow a = 2$
 From (i), $8a^3 - 8a^2 = 0 \Rightarrow a = a$
 \therefore the required point is $(2a, a)$.
 4. $y^2 = 3x - 2$ (i)
 Differentiating both sides w.r.t. x , we get
 $2y \frac{dy}{dx} = -2 \frac{dx}{dx} \Rightarrow \frac{dy}{dx} = -\frac{1}{y}$
 \therefore Slope of the tangent $= -\frac{1}{y}$
 Slope of the given line is -1 .
 Since, the tangent is parallel to the given line.
 $-\frac{1}{y} = -1 \Rightarrow y = 1$
 From (i), $x = 1$
 \therefore the required point is $(1, 1)$.
 5. $y = 6x - 2$ (i)
 $\therefore \frac{dy}{dx} = 6 - 2$
 Slope of the given line is 2.
 Since, the tangent is parallel to the given line.
 $6 - 2 = 2 \Rightarrow x = 2$
 From (i), $y = 8$
 \therefore the point of tangency will be $(2, 8)$.
 6. Let the coordinates of P be (x_1, y_1) .
 Then, $y_1 = 2x_1^2 - x_1 + 1$ (i)
 Now, $y = 2x^2 - x + 1$
 $\therefore \frac{dy}{dx} = 4x - 1$
 $\therefore \left(\frac{dy}{dx}\right)_{(x_1, y_1)} = 4x_1 - 1$
 Slope of the given line is 3.
 Since, the tangent is parallel to the given line.
 \therefore slope of the tangent $= 3$
 $\Rightarrow 4x_1 - 1 = 3$
 $\Rightarrow x_1 = 1$
 From (i), $y_1 = 2$
 \therefore the coordinates of P are $(1, 2)$.



$$7. \quad = \log \quad \dots(i)$$

$$\therefore \frac{d}{d} = 1 + \log$$

$$\text{Slope of the normal} = -\frac{1}{\left(\frac{d}{d}\right)} = \frac{-1}{1 + \log}$$

Slope of the given line is 1.

Since, the normal is parallel to the given line.

$$\therefore \frac{-1}{1 + \log} = 1$$

$$\Rightarrow \log = -2$$

$$\Rightarrow = e^{-2}$$

$$\text{From (i), } = -2e^{-2}$$

\(\therefore\) Co-ordinates of the point are \((e^{-2}, -2e^{-2})\).

$$8. \quad = (-3)^2$$

$$\therefore ' = 2(-3)$$

Since, the tangent is parallel to the line joining \((3, 0)\) and \((4, 1)\).

$$\therefore 2(-3) = \frac{1-0}{4-3}$$

$$\Rightarrow 2 - 6 = 1 \Rightarrow = \frac{7}{2}$$

$$\text{When } = \frac{7}{2},$$

$$= \left(\frac{7}{2} - 3\right)^2 = \frac{1}{4}$$

\(\therefore\) the required point is \(\left(\frac{7}{2}, \frac{1}{4}\right)\).

$$9. \quad = x^2 - 4x + 5 \quad \dots(i)$$

$$\therefore \frac{d}{d} = 2x - 4$$

$$\text{Slope of the given line} = -\frac{1}{2}$$

Since, the tangent is perpendicular to the given line.

$$\therefore (2x - 4) \left(-\frac{1}{2}\right) = -1$$

$$\Rightarrow 2x - 4 = 2$$

$$\Rightarrow x = 3$$

$$\text{From (i), } = 2$$

\(\therefore\) the required point is \((3, 2)\).

$$10. \quad x^2 + y^2 - 2x - 3 = 0 \quad \dots(i)$$

Differentiating w.r.t. , we get

$$2x + 2y \frac{d}{d} - 2 = 0$$

$$\therefore \frac{d}{d} = \frac{1-}{y}$$

Since, the tangent is parallel to X-axis.

$$\therefore \frac{d}{d} = 0$$

$$\Rightarrow \frac{1-}{y} = 0 \Rightarrow = 1$$

From (i),

$$= \pm 2$$

$$11. \quad x^3 + 3x^2 - 12 = 0 \quad \dots(i)$$

Differentiating w.r.t. , we get

$$\frac{d}{d} = -\frac{6}{3x^2 - 12}$$

Since, the tangent is parallel to Y-axis.

$$\therefore \frac{d}{d} = 0$$

$$\Rightarrow 3x^2 - 12 = 0$$

$$\Rightarrow x^2 = 4 \Rightarrow x = \pm 2$$

$$\therefore x = 2 \quad \dots[\because x \neq -2]$$

From (i),

$$= \pm \frac{4}{\sqrt{3}}$$

$$12. \quad y = a^2 + b$$

$$\therefore \frac{d}{d} = 2a + b \Rightarrow \left(\frac{d}{d}\right)_{(2,-8)} = 4a + b$$

Since, the tangent is parallel to X-axis.

$$\therefore \left(\frac{d}{d}\right)_{(2,-8)} = 0 \Rightarrow b = -4a \quad \dots(i)$$

Also, the point \((2, -8)\) lies on the curve

$$y = a^2 + b$$

$$\therefore -8 = 4a + 2b \quad \dots(ii)$$

From (i) and (ii), we get \(a = 2, b = -8\)

$$13. \quad y = a^2 - 6x + b$$

$$\therefore \frac{d}{d} = 2a - 6$$

$$\therefore \left(\frac{d}{d}\right)_{\left(\frac{3}{2}\right)} = 3a - 6$$

Since, the tangent is parallel to X-axis at \(\left(\frac{3}{2}, \right)\).

$$\therefore \left(\frac{d}{d}\right)_{\left(\frac{3}{2}\right)} = 0$$

$$\Rightarrow 3a - 6 = 0 \Rightarrow a = 2$$

Now, the given curve passes through \((0, 2)\).

$$\therefore 2 = 0 - 0 + b$$

$$\Rightarrow b = 2$$



14. At $t = 2$, $y = \frac{1}{2}$
 and $x = 2 - \frac{1}{2} = \frac{3}{2}$
 Now, $\frac{dy}{dx} = \frac{\frac{d}{dt}}{\frac{dt}{dx}} = \frac{1 + \frac{1}{t^2}}{\frac{-1}{t^2}} = \frac{t^2 + 1}{-1}$
 $\therefore \left(\frac{d}{dx}\right)_{(t=2)} = -5$
 \therefore Equation of the normal at $\left(\frac{1}{2}, \frac{3}{2}\right)$ is
 $-\frac{3}{2} = \frac{1}{5}\left(x - \frac{1}{2}\right)$
 $\therefore -5x + 7 = 0$
15. At $\theta = \frac{\pi}{6}$,
 $y = a \sec \frac{\pi}{6} = \frac{2a}{\sqrt{3}}$ and $x = a \tan \frac{\pi}{6} = \frac{a}{\sqrt{3}}$
 $\frac{dy}{dx} = \frac{\frac{d}{d\theta}}{\frac{dx}{d\theta}} = \frac{a \sec^2 \theta}{a \sec \theta \tan \theta} = \frac{1}{\sin \theta} = \operatorname{cosec} \theta$
 $\therefore \left(\frac{d}{dx}\right)_{\theta=\frac{\pi}{6}} = \operatorname{cosec} \frac{\pi}{6} = 2$
 \therefore Equation of the tangent at $\left(\frac{2a}{\sqrt{3}}, \frac{a}{\sqrt{3}}\right)$ is
 $-\frac{a}{\sqrt{3}} = 2\left(x - \frac{2a}{\sqrt{3}}\right)$
 $\Rightarrow 2x - \frac{4a}{\sqrt{3}} = \frac{a}{\sqrt{3}}$
16. $y = x^3 + 2x^2 - 4x - 43$
 $\therefore \frac{dy}{dx} = 3x^2 + 4x - 4$
 $\therefore \left(\frac{d}{dx}\right)_{(-2,5)} = 3(-2)^2 + 4(-2) - 4 = 0$
 \therefore equation of the tangent at $(-2, 5)$ is
 $y - 5 = 0 \cdot (x + 2)$
 i.e., $y = 5$ (parallel to X-axis)
 Normal is perpendicular to X-axis and passes through $(-2, 5)$.
 \therefore equation of the normal is
 $x + 2 = 0$, i.e., $x + 2 = 0$

17. $x^2 = 5 - y$
 $\therefore \frac{dy}{dx} = \frac{5}{2}$
 $\therefore \left(\frac{d}{dx}\right)_{(1,-2)} = \frac{-5}{4}$
 \therefore Equation of the normal at $(1, -2)$ is
 $-(-2) = \frac{4}{5}(x - 1)$
 $\therefore 4x - 5y - 14 = 0 \dots(i)$
 As the normal is of the form $ax - 5y + b = 0$, comparing this with (i), we get
 $a = 4$ and $b = -14$
18. $\frac{2}{3} + \frac{2}{3} = a^{\frac{2}{3}}$
 Differentiating both sides w.r.t. x , we get
 $\frac{2}{3} \cdot \frac{-1}{3} + \frac{2}{3} \cdot \frac{-1}{3} \frac{d}{dx} = 0$
 $\Rightarrow \frac{d}{dx} = \frac{-\frac{1}{3}}{\frac{1}{3}}$
 At $(a \sin^3 \theta, a \cos^3 \theta)$,
 $\frac{dy}{dx} = -\frac{\cos \theta}{\sin \theta} = -\cot \theta$
 \therefore slope of the normal is $\tan \theta$.
 \therefore equation of the normal at $(a \sin^3 \theta, a \cos^3 \theta)$ is
 $-a \cos^3 \theta = \tan \theta (x - a \sin^3 \theta)$
 $\Rightarrow \cos \theta - a \cos^4 \theta = \sin \theta - a \sin^4 \theta$
 $\Rightarrow \sin \theta - \cos \theta = a \sin^4 \theta - a \cos^4 \theta$
19. Let (x_1, y_1) be a point on the curve $y = x + \frac{4}{2}$.
 Since, the tangent is parallel to X-axis.
 $\therefore \left(\frac{d}{dx}\right)_{(x_1, y_1)} = 0 \Rightarrow 1 - \frac{8}{x_1} = 0 \Rightarrow x_1 = 2$
 Now, $y_1 = x_1 + \frac{4}{2}$
 $\Rightarrow y_1 = 2 + \frac{4}{2}$
 $\Rightarrow y_1 = 3$
 \therefore equation of the tangent at $(2, 3)$ is
 $y - 3 = 0 \Rightarrow y = 3$
20. Since, the given curve crosses the X-axis,
 $y = 0$
 $\therefore 0 = 2 - x \Rightarrow x = 2$
 \therefore the given curve crosses the X-axis at $(2, 0)$.
 Now, $(1 + x^2) = 2 -$



Differentiating both sides w.r.t. x , we get

$$(1 + x^2) \frac{d}{dx} + 2x = -1$$

$$\therefore \frac{d}{dx} = \frac{-1-2x}{1+x^2}$$

$$\therefore \left(\frac{d}{dx}\right)_{(2,0)} = -\frac{1}{5}$$

\therefore equation of the tangent at $(2, 0)$ is

$$y - 0 = -\frac{1}{5}(x - 2)$$

$$\Rightarrow x + 5y = 2$$

21. Since, the given curve crosses the Y-axis, $x = 0$

$$\therefore y = be^0 \Rightarrow y = b$$

\therefore the given curve crosses the Y-axis at $(0, b)$.

$$\text{Now, } y = be^{-\frac{x}{a}}$$

$$\therefore \frac{d}{dx} = -\frac{b}{a} e^{-\frac{x}{a}}$$

$$\therefore \left(\frac{d}{dx}\right)_{(0,b)} = -\frac{b}{a}$$

\therefore the equation of the tangent at $(0, b)$ is

$$-b = -\frac{b}{a}(x - 0)$$

$$\Rightarrow \frac{x}{a} + \frac{y}{b} = 1$$

22. $y = e^{2x}$

$$\therefore \frac{d}{dx} = 2e^{2x}$$

$$\therefore \left(\frac{d}{dx}\right)_{(0,1)} = 2$$

\therefore equation of the tangent at $(0, 1)$ is

$$y - 1 = 2(x - 0)$$

$$\Rightarrow y = 2x + 1$$

This tangent meets X-axis, $\therefore y = 0$

$$\therefore 0 = 2x + 1 \Rightarrow x = -\frac{1}{2}$$

\therefore the required point is $\left(-\frac{1}{2}, 0\right)$.

23. Let the required point be (x_1, y_1) .

$$\therefore y_1 = be^{-\frac{x_1}{a}} \quad \dots(i)$$

$$\text{Now, } y = be^{-\frac{x}{a}}$$

$$\therefore \frac{d}{dx} = -\frac{b}{a} \cdot e^{-\frac{x}{a}}$$

$$\therefore \left(\frac{d}{dx}\right)_{(x_1, y_1)} = -\frac{b}{a} \cdot e^{-\frac{x_1}{a}} = -\frac{y_1}{a} \quad \dots[\text{From (i)}]$$

\therefore equation of the tangent at (x_1, y_1) is

$$y - y_1 = -\frac{y_1}{a}(x - x_1)$$

$$\Rightarrow \frac{x}{a} + \frac{y}{y_1} = \frac{x_1}{a} + 1$$

Comparing this equation with $\frac{x}{a} + \frac{y}{b} = 1$, we get

$$y_1 = b \text{ and } 1 + \frac{x_1}{a} = 1 \Rightarrow x_1 = 0$$

\therefore the required point is $(0, b)$.

24. When $x = 0$,

$$y = (1 + 0) + \sin^{-1}(0) \Rightarrow y = 1$$

$$\text{Now, } y = (1 + x) + \sin^{-1}(\sin^2 x)$$

$$\therefore \frac{d}{dx} = (1 + x) \left\{ \frac{d}{dx} \log(1 + x) + \frac{1}{1 + x} \right\} + \frac{\sin 2x}{\sqrt{1 - \sin^4 x}}$$

$$\therefore \left(\frac{d}{dx}\right)_{(0,1)} = 1$$

\therefore the equation of the normal at $(0, 1)$ is

$$y - 1 = -1(x - 0) \Rightarrow x + y = 1$$

25. Let (x_1, y_1) be the point on the curve $y = 2x^2 + 7$, where the tangent is parallel to the line $4x - y + 3 = 0$.

$$\text{Then, } y_1 = 2x_1^2 + 7 \quad \dots(i)$$

$$\text{Now, } y = 2x^2 + 7$$

$$\therefore \frac{d}{dx} = 4x$$

$$\therefore \left(\frac{d}{dx}\right)_{(x_1, y_1)} = 4x_1$$

Slope of the given line is 4.

Since, the tangent is parallel to the given line.

\therefore slope of the tangent = 4

$$\Rightarrow 4x_1 = 4$$

$$\Rightarrow x_1 = 1$$

$$\text{From (i), } y_1 = 9$$

\therefore the coordinates of the point are $(1, 9)$.

\therefore Equation of the tangent at $(1, 9)$ is

$$y - 9 = 4(x - 1)$$

$$\Rightarrow 4x - y + 5 = 0$$



26. $8 = (-2)^2$
Differentiating both sides w.r.t. x , we get

$$\frac{d}{dx} = \frac{-2}{4}$$

$$\therefore \left(\frac{d}{dx}\right)_{(-6,8)} = \frac{-6-2}{4} = -2 \quad \dots(i)$$

$$= +\frac{3}{-}$$

$$\therefore \frac{d}{dx} = 1 - \frac{3}{2}$$

$$\therefore \left(\frac{d}{dx}\right)_{(1,4)} = 1 - \frac{3}{1^2} = -2 \quad \dots(ii)$$

From (i) and (ii),
 T_1 is parallel to T_2

27. $\frac{d}{dx} = 1$

$$\therefore \frac{d}{dx} = \frac{1}{x}$$

$$\therefore \frac{d}{dx} = \frac{-1}{2}$$

\therefore Slope of the normal = $-\frac{1}{2}$

Slope of the line $ax + by + c = 0$ is $-\frac{a}{b}$.

Since, the line $ax + by + c = 0$ is a normal to the curve $y = 1/x$.

$$\therefore \frac{1}{2} = -\frac{a}{b}$$

For this condition to hold true, either $a < 0, b > 0$ or $b < 0, a > 0$

28. $\frac{d}{dx} = 1 - 2x + 3x^2$

$$\therefore \frac{d}{dx} = 3x^2 - 2x + 1 = 3\left(x^2 - \frac{2}{3}x + \frac{1}{3}\right)$$

$$= 3\left(x^2 - \frac{2}{3}x + \frac{1}{9} - \frac{1}{9} + \frac{1}{3}\right)$$

$$= 3\left[\left(x - \frac{1}{3}\right)^2 + \frac{2}{9}\right]$$

$$= 3\left[\left(x - \frac{1}{3}\right)^2 + \frac{2}{9}\right] > 0$$

Slope of the given line is $-\frac{l}{m}$.

The slope will be positive only if l and m have opposite signs.

\therefore option (D) is the correct answer.

29. $\left(\frac{1}{a}\right)^n + \left(\frac{1}{b}\right)^n = 2$

Differentiating both sides w.r.t. x , we get

$$\frac{1}{a^n} n^{-n-1} + \frac{1}{b^n} n^{-n-1} \cdot \frac{dn}{dx} = 0$$

$$\therefore \frac{dn}{dx} = \frac{-b^n}{a^n} \cdot \frac{n-1}{n-1}$$

At (a, b) ,

$$\frac{dn}{dx} = \frac{-b^n}{a^n} \cdot \frac{a^{n-1}}{b^{n-1}} = \frac{-b}{a}, \text{ which is}$$

independent of n .

30. Since, $(1, 1)$ lies on the curve $y = a^2/x$.

$$\therefore \frac{1}{1} = \frac{a^2}{1} \quad \dots(i)$$

Now, $y = a^2/x$

Differentiating w.r.t. x , we get

$$\frac{dy}{dx} + \frac{y}{x} = 0$$

$$\Rightarrow \frac{dy}{dx} = -\frac{y}{x}$$

$$\therefore \left(\frac{dy}{dx}\right)_{(1,1)} = -\frac{1}{1}$$

\therefore equation of the tangent at $(1, 1)$ is

$$y - 1 = -\frac{1}{1}(x - 1)$$

$$\Rightarrow y + 1 = 2 - x$$

$$\Rightarrow y + 1 = 2a^2 - x \quad \dots[\text{From (i)}]$$

This tangent meets the coordinate axes at

$$\left(\frac{2a^2}{1}, 0\right) \text{ and } \left(0, \frac{2a^2}{1}\right).$$

$$\therefore \text{required area} = \frac{1}{2} \left(\frac{2a^2}{1}\right) \left(\frac{2a^2}{1}\right)$$

$$= \frac{2a^4}{1}$$

$$= 2a^2 \quad \dots[\text{From (i)}]$$

31. $y = x^2 \Rightarrow \frac{dy}{dx} = 2x \Rightarrow \left(\frac{dy}{dx}\right)_{(1,1)} = 2 = m_1(\text{say})$

$$6 = 7 - x^3 \Rightarrow 6 \cdot \frac{dx}{dx} = -3x^2$$

$$\Rightarrow \left(\frac{dx}{dy}\right)_{(1,1)} = -\frac{1}{2} = m_2(\text{say})$$

Since, $m_1 m_2 = -1$

\therefore the angle of intersection is $\frac{\pi}{2}$.



32. $\frac{d}{dt} = 2$
 $\therefore \frac{d}{dt} = 2$
 $\therefore \left(\frac{d}{dt}\right)_{(1,1)} = 2 = m_1$ (say)
 and $\frac{d}{dt} = 2$
 $\therefore 1 = 2 \frac{d}{dt}$
 $\Rightarrow \frac{d}{dt} = \frac{1}{2}$
 $\therefore \left(\frac{d}{dt}\right)_{(1,1)} = \frac{1}{2} = m_2$ (say)
 \therefore angle of intersection is $\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$
 $= \left| \frac{2 - \frac{1}{2}}{1 + 2 \left(\frac{1}{2}\right)} \right| = \frac{3}{4}$
 $\therefore \theta = \tan^{-1} \left(\frac{3}{4} \right)$
33. The point of intersection of the given curves is (0, 1).
 Now, $\frac{d}{dt} = a$
 $\therefore \frac{d}{dt} = a \log a$
 $\therefore \left(\frac{d}{dt}\right)_{(0,1)} = \log a = m_1$ (say)
 Also, $\frac{d}{dt} = b$
 $\therefore \frac{d}{dt} = b \log b$
 $\therefore \left(\frac{d}{dt}\right)_{(0,1)} = \log b = m_2$ (say)
 $\therefore \tan \alpha = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right| = \frac{\log a - \log b}{1 + \log a \log b}$
34. $s = ae^t + \frac{b}{e^t}$
 $\therefore \frac{ds}{dt} = \text{velocity} = ae^t - \frac{b}{e^t}$
 $\therefore \frac{d^2s}{dt^2} = \text{acceleration} = ae^t + \frac{b}{e^t} = s$
35. $\frac{dS}{dt} = \text{velocity} = 15 + 12t - 3t^2$
 When particle comes to rest, $v = 0$
 $\Rightarrow 3t^2 - 12t - 15 = 0$
 $\Rightarrow t = 5$ sec

36. $s = \sqrt{t} \Rightarrow \frac{ds}{dt} = \frac{1}{2\sqrt{t}}$
 and $\frac{d^2s}{dt^2} = -\frac{1}{4t^{\frac{3}{2}}}$
 $= -\frac{1}{4} \left(\frac{2ds}{dt} \right)^3 = -2 \left(\frac{ds}{dt} \right)^3$

Hence, acceleration \propto (velocity)³.

37. $s = \sqrt{at^2 + bt + c}$
 $\therefore v = \frac{ds}{dt} = \frac{1}{2} \frac{2at + b}{\sqrt{at^2 + bt + c}}$
 $= \frac{2at + b}{2s}$
 acceleration $= \frac{d^2s}{dt^2} = \frac{dv}{dt}$
 $= \frac{2s(2a) - (2at + b) \cdot 2 \frac{ds}{dt}}{4s^2}$
 $= \frac{4as - 2(2at + b) \left(\frac{2at + b}{2s} \right)}{4s^2}$
 $= \frac{4as^2 - (2at + b)^2}{4s^3}$
 $= \frac{4a(at^2 + bt + c) - (4a^2t^2 + 4abt + b^2)}{4s^3}$
 $= \frac{4ac - b^2}{4s^3}$

\therefore acceleration varies as $\frac{1}{s^3}$

38. Area of a circle is $A = \pi R^2$ and $\frac{dR}{dt} = 0.2$

$\therefore \frac{dA}{dt} = 2\pi R \frac{dR}{dt} = 1.2\pi \text{ cm}^2$

39. Let a be each side and A be the area of the square at any time t . Then,

$$A = a^2$$

$$\Rightarrow \frac{dA}{dt} = 2a \frac{da}{dt}$$

$$= 2(2)(4)$$

$$\dots \left[\because \frac{da}{dt} = 4 \text{ cm/sec and } a = 2 \text{ cm (given)} \right]$$

$$= 16 \text{ cm}^2/\text{sec}$$



40. Radius of balloon = $r = \frac{3}{4}(2 + 3)$

$$\Rightarrow \frac{dr}{dt} = \frac{3}{2}$$

$$V = \frac{4}{3} \pi r^3$$

$$\begin{aligned} \therefore \frac{dV}{dt} &= 4\pi \left(\frac{3}{4}\right)^2 (2 + 3)^2 \cdot \frac{3}{2} \\ &= \frac{27\pi}{8} (2 + 3)^2 \end{aligned}$$

41. Given, $\frac{dr}{dt} = 2$ cm/sec, where r be the radius of circle and t be the time.

Now, area of circle is given by $A = \pi r^2$

$$\therefore \frac{dA}{dt} = 2\pi r \frac{dr}{dt}$$

$$\Rightarrow \frac{dA}{dt} = 2\pi \cdot 20 \cdot 2$$

$$\therefore \frac{dA}{dt} = 80\pi \text{ cm}^2/\text{sec}$$

\therefore the rate of change of area of circle with respect to time is $80\pi \text{ cm}^2/\text{sec}$.

42. Let r be the radius and V be the volume of the spherical balloon at any time t . Then,

$$V = \frac{4}{3} \pi r^3$$

$$\Rightarrow \frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

$$\therefore \left(\frac{dV}{dt}\right)_{(r=15)} = 4\pi \times (15)^2 \times \left(\frac{dr}{dt}\right)_{(r=15)}$$

$$\begin{aligned} \Rightarrow 30 &= 900\pi \left(\frac{dr}{dt}\right)_{(r=15)} \\ &\dots \left[\because \frac{dV}{dt} = 30 \text{ ft}^3 / \text{min (given)} \right] \end{aligned}$$

$$\Rightarrow \left(\frac{dr}{dt}\right)_{(r=15)} = \frac{1}{30\pi} \text{ ft / min}$$

43. Let velocity $V = 5$ cm/sec
(Increasing the rate/sec is called the velocity)

$$\frac{da}{dt} = 5 \quad \dots(i)$$

But if a is edge of a cube, then $V = a^3$

$$\therefore \frac{dV}{dt} = 3a^2 \frac{da}{dt} = 3a^2 \cdot 5$$

$$\begin{aligned} &= 15a^2 = 15 \times (12)^2 \dots [\because \text{edge } a = 12 \text{ cm}] \\ &= 2160 \text{ cm}^3/\text{sec} \end{aligned}$$

44. $\frac{da}{dt} = 60$ cm/sec where a is edge and t is time.

$$V = a^3$$

$$\begin{aligned} \therefore \frac{dV}{dt} &= 3a^2 \frac{da}{dt} \\ &= 3a^2 \times 60 = 180a^2 \\ &= 180 \times (90)^2 \\ &= 1458000 \text{ cm}^3/\text{sec}. \end{aligned}$$

45. $V = \frac{4}{3} \pi (r + 10)^3$, where r is thickness of ice.

$$\therefore \frac{dV}{dt} = 4\pi (r + 10)^2 \frac{dr}{dt}$$

$$\text{But, } \frac{dV}{dt} = 50$$

$$\therefore 50 = 4\pi (10 + r)^2 \frac{dr}{dt}$$

$$\text{At } r = 5, \frac{dr}{dt} = \frac{50}{4\pi(10+5)^2}$$

$$= \frac{50}{4\pi(225)}$$


$$= \frac{1}{18\pi} \text{ cm / min}$$

46. $\frac{d}{dt} = 0.5$ cm/sec

$$\therefore \text{Area} = \frac{a^2}{2}$$

$$\therefore \frac{dA}{dt} = \frac{2}{2} \cdot \frac{da}{dt}$$


$$\begin{aligned} \therefore \left[\frac{dA}{dt}\right]_{A=400} &= \frac{1}{2} \sqrt{800} \quad \dots \left[\because A = 400 \text{ cm}^2 \right] \\ &= 10\sqrt{2} \text{ cm}^2/\text{sec} \quad \dots \left[\therefore a = \sqrt{800} \text{ cm} \right] \end{aligned}$$



47. From the figure,

$$\frac{6}{2} = \frac{+}{6}$$


$$\Rightarrow 4 = 2 \Rightarrow \frac{d}{dt} = \frac{1}{2}$$




$$\therefore \frac{d}{dt} = \frac{1}{2} \frac{d}{dt} = \frac{5}{2} \text{ metre/hour}$$



48. When $a = 8$, $b = 15$, $a^2 + b^2 = 17^2$
 Differentiating both sides w.r.t. t , we get
 $2a \frac{da}{dt} + 2b \frac{db}{dt} = 0$
- $\therefore 8(1) + 15 \frac{db}{dt} = 0$
- $\therefore \frac{db}{dt} = \frac{-8}{15} \text{ m/sec}$
- \therefore the upper end is coming down at the rate of $\frac{8}{15} \text{ m/sec}$.




49. Let the position of the kite at time t be at C .
 $\therefore BC = 151.5 \text{ m}$
 Let AD be the boy who is flying the kite.
 $\therefore CE = BC - BE = 151.5 - 1.5 = 150 \text{ m}$



- In right angled ΔCDE ,
 $2^2 = 2^2 + (150)^2$
 Differentiating both sides w.r.t. t , we get
 $2 \frac{d}{dt} = 2 \frac{d}{dt}$
 $\Rightarrow \frac{dy}{dt} = - \dots \left[\because \frac{d}{dt} = 10 \right]$
 $= \frac{10\sqrt{2^2 - (150)^2}}{250}$
 $= \frac{10\sqrt{(250)^2 - (150)^2}}{250} = \frac{10 \times 200}{250} = 8 \text{ m/s}$

50.



Let OC be the wall. Let AB be the position of the ladder at any time t such that $OA =$ and $OB =$.

Length of the ladder $AB = 20 \text{ ft}$.

In right angled ΔAOB ,
 $2^2 + 2^2 = (20)^2$

Differentiating both sides w.r.t. t , we get

$$2 \frac{d}{dt} + 2 \frac{d}{dt} = 0$$

$$\therefore \frac{d}{dt} = - \frac{d}{dt} = \frac{-}{\sqrt{400 - 2^2}} \cdot \frac{d}{dt}$$

$$\therefore \left(\frac{d}{dt} \right)_{=16} = - \frac{16}{\sqrt{400 - (16)^2}} \cdot \frac{d}{dt} = - \frac{4}{3} \cdot \frac{d}{dt}$$

Negative sign indicates, that when increases with time, decreases.

Hence, the upper end is moving $\frac{4}{3}$ times as fast as the lower end.

51. $f(x) = x^3 - 3x + 5$
 $\therefore f'(x) = 3x^2 - 3$
 Here, $a = 2$ and $h = -0.01$
 $\therefore f(a + h) \approx f(a) + hf'(a)$
 $\approx 7 + (-0.01)(9)$
 $\therefore f(1.99) \approx 7 - 0.09 \approx 6.91$

52. Let $f(x) = \frac{1}{\sqrt{x}}$

$$\therefore f'(x) = \frac{-1}{2} x^{-\frac{3}{2}} = \frac{-1}{2x^{\frac{3}{2}}}$$

Here, $a = 25$ and $h = 0.1$

$$\therefore f(a + h) \approx f(a) + hf'(a)$$

$$\approx \frac{1}{5} - \frac{(0.1)}{2 \times 125} \approx \frac{1}{5} - \frac{0.1}{250}$$

$$\approx \frac{1}{5} \left(1 - \frac{1}{500} \right) \approx \frac{1}{5} \left(\frac{499}{500} \right) \approx \frac{1}{5} \left(\frac{998}{1000} \right)$$

$$\therefore \frac{1}{\sqrt{25.1}} \approx \frac{1}{5} \times 0.998 \approx 0.1996$$

53. Let $f(x) = \frac{1}{2x}$

$$\therefore f'(x) = -2^{-3} = \frac{-2}{3}$$

Here, $a = 2$ and $h = 0.002$

$$\therefore f(a + h) \approx f(a) + hf'(a)$$

$$\approx \frac{1}{4} + (0.002) \left(\frac{-2}{8} \right) \approx \frac{1}{4} - \frac{0.002}{4}$$

$$\therefore \frac{1}{(2.002)^2} \approx \frac{0.998}{4} \approx 0.2495$$



54. Let $f(x) = \frac{1}{4}$
 $\therefore f'(x) = \frac{1}{4} \cdot \frac{-3}{4} = -\frac{3}{16}$
 Here, $a = 81$ and $h = -1$
 $\therefore f(a+h) \approx f(a) + hf'(a)$
 $\approx (81)^{\frac{1}{4}} + (-1) \left[\frac{1}{4(81)^{\frac{3}{4}}} \right]$
 $\approx 3 - \frac{1}{108}$
 $\approx 3 - 0.009259$
 $\therefore (80)^{\frac{1}{4}} \approx 2.9907$
55. Let $f(x) = \cot^{-1}$
 $\therefore f'(x) = \frac{-1}{1+x^2}$
 Here, $a = 1$ and $h = 0.001$
 $\therefore f(a+h) \approx f(a) + hf'(a)$
 $\approx \frac{\pi}{4} + 0.001 \left(\frac{-1}{2} \right)$
 $\approx \frac{3.14}{4} - 0.0005$
 $\therefore \cot^{-1}(1.001) \approx 0.785 - 0.0005 \approx 0.7845$
56. Let $f(x) = \tan^{-1}$
 $\therefore f'(x) = \frac{1}{1+x^2}$
 Here, $a = 1$ and $h = -0.001$
 $\therefore f(a+h) \approx f(a) + hf'(a)$
 $\therefore \tan^{-1}(0.999) \approx \frac{\pi}{4} + \frac{1}{1+1} (-0.001)$
 $\approx \frac{\pi}{4} - \frac{0.001}{2}$
 $\approx \frac{\pi}{4} - 0.0005$
57. Let $f(x) = \cos$
 $\therefore f'(x) = -\sin$
 Here, $a = 90^\circ$
 and $h = 30' = \left(\frac{1}{2}\right)^\circ = \left(\frac{1}{2} \times 0.0175\right)^\circ$
 $= 0.00875$
 $f(a) = f(90^\circ) = \cos 90^\circ = 0$
 $f'(a) = f'(90^\circ) = -\sin 90^\circ = -1$
 $\therefore f(a+h) \approx f(a) + hf'(a)$
 $\therefore \cos(90^\circ 30') \approx 0 + (0.00875) \times (-1) \approx -0.00875$

58. Let $f(x) = \sin$
 $\therefore f'(x) = \cos$
 Here, $a = 30^\circ$ and
 $h = 1^\circ = 0.0175^\circ$
 $\therefore f(a+h) \approx f(a) + hf'(a)$
 $\approx \frac{1}{2} + 0.0175 \times 0.8660$
 $\approx 0.5 + 0.01515$
 $\therefore \sin(31^\circ) \approx 0.51515$
 ≈ 0.5152
59. Let $f(x) = \tan$
 $\therefore f'(x) = \sec^2$
 Here, $a = 45^\circ = \left(\frac{\pi}{4}\right)^\circ$ and $h = 1^\circ = 0.0175^\circ$
 $f(a+h) \approx f(a) + hf'(a)$
 $\approx \tan(a) + h \sec^2 a$
 $\approx \tan(a) + h \frac{1}{\cos^2 a}$
 $\approx \tan\left(\frac{\pi}{4}\right) + (0.0175) \frac{1}{(1/\sqrt{2})^2}$
 $\approx 1 + 0.035$
 $\therefore \tan 46^\circ \approx 1.035$
60. Let $f(x) = \log_e$
 $\therefore f'(x) = \frac{1}{x}$
 Here, $a = 9$ and $h = 0.01$
 $\therefore f(a+h) \approx f(a) + hf'(a)$
 $\approx f(9) + (0.01) f'(9)$
 $\approx \log_e 3^2 + \frac{0.01}{9}$
 $\approx 2 \log_e 3 + \frac{0.01}{9}$
 $\approx 2.1972 + 0.0011$
 ≈ 2.1983
61. Consider option (B),
 $f(x) = x^2$ is a polynomial function.
 $\therefore f(x)$ is continuous and differentiable in the given interval.
 Also, $f(1) = f(-1) = 1$
 So, Rolle's theorem is applicable to
 $f(x) = x^2$ on $[-1, 1]$.



$$62. f(x) = |x| = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$$

$$f'(0^-) = \lim_{h \rightarrow 0} \frac{f(0-h) - f(0)}{-h}$$

$$= \lim_{h \rightarrow 0} \frac{|-h|}{-h} = \lim_{h \rightarrow 0} \frac{h}{-h} = -1$$

$$f'(0^+) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{|h|}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

$\therefore f(x)$ is not differentiable at $x = 0$.

$$63. f(x) = e^{-2x} \sin 2x$$

$$\Rightarrow f'(x) = 2e^{-2x} (\cos 2x - \sin 2x)$$

Now, $f'(c) = 0$

$$\Rightarrow \cos 2c - \sin 2c = 0$$

$$\Rightarrow \tan 2c = 1 \Rightarrow 2c = \frac{\pi}{4} \Rightarrow c = \frac{\pi}{8}$$

$$64. f(x) = x^3 - 6x^2 + ax + b$$

$$\Rightarrow f'(x) = 3x^2 - 12x + a$$

Now, $f'(c) = 0 \Rightarrow f'\left(2 + \frac{1}{\sqrt{3}}\right) = 0$

$$\Rightarrow 3\left(2 + \frac{1}{\sqrt{3}}\right)^2 - 12\left(2 + \frac{1}{\sqrt{3}}\right) + a = 0$$

$$\Rightarrow 3\left(4 + \frac{4}{3} + \frac{4}{\sqrt{3}}\right) - 12\left(2 + \frac{1}{\sqrt{3}}\right) + a = 0$$

$$\Rightarrow 12 + 4 + 4\sqrt{3} - 24 - 4\sqrt{3} + a = 0$$

$$\Rightarrow a = 11$$

$$65. f(x) = (x+3)e^{-\frac{1}{2}x}$$

$$\therefore f'(x) = (x+3)e^{-\frac{1}{2}x} \cdot \left(-\frac{1}{2}\right) + (x+3)e^{-\frac{1}{2}x}$$

$$= e^{-\frac{1}{2}x} \left\{ -\frac{1}{2}(x+3) + (x+3) \right\}$$

$$= -\frac{1}{2}e^{-\frac{1}{2}x}(x-3)$$

Since, $f(x)$ satisfies all the conditions of Rolle's theorem. So, there exists $c \in (-3, 0)$ such that

$$f'(c) = 0$$

$$\Rightarrow c^2 - c - 6 = 0$$

$$\Rightarrow c = 3, -2$$

But $c = -2 \in [-3, 0]$

$\therefore c = -2$

66. Here, $f(x)$ is continuous and differentiable on $(0, 1)$ for $\alpha > 0$

Also, $f(0) = f(1) = 0$

For $f(x)$ to be continuous at $x = 0$,

$$\lim_{x \rightarrow 0^+} f(x) = f(0) \Rightarrow \lim_{x \rightarrow 0^+} x^\alpha \log x = 0$$

$$\Rightarrow \lim_{x \rightarrow 0^+} \frac{\log x}{-x^{-\alpha}} = 0$$

Applying L'Hospital rule, we get

$$\lim_{x \rightarrow 0^+} \frac{1}{-\frac{\alpha}{x^{\alpha+1}}} = 0 \Rightarrow \lim_{x \rightarrow 0^+} -\frac{x^{\alpha+1}}{\alpha} = 0,$$

which is possible only when $\alpha > 0$

\therefore option (D) is the correct answer.

67. $f(x) = \log_e x$

$\therefore f(1) = \log_e 1 = 0,$

$$f(3) = \log_e 3 \text{ and } f'(x) = \frac{1}{x}$$

By Lagrange's mean value theorem,

$$f'(c) = \frac{f(3) - f(1)}{3 - 1}$$

$$\Rightarrow \frac{1}{c} = \frac{\log_e 3 - 0}{2} \Rightarrow c = \frac{2}{\log_e 3} \Rightarrow c = 2 \log_3 e$$

68. $f(x) = x + \frac{1}{x}$

$\therefore f(3) = \frac{10}{3}, f(1) = 2$ and $f'(x) = 1 - \frac{1}{x^2}$

By Lagrange's mean value theorem,

$$f'(c) = \frac{f(3) - f(1)}{3 - 1}$$

$$\Rightarrow 1 - \frac{1}{c^2} = \frac{\frac{10}{3} - 2}{2} \Rightarrow 1 - \frac{1}{c^2} = \frac{2}{3}$$

$$\Rightarrow c^2 = 3 \Rightarrow c = \sqrt{3}$$

69. $f(x) = \frac{1}{x}$

$$\Rightarrow f(a) = \frac{1}{a}, f(b) = \frac{1}{b} \text{ and } f'(x) = -\frac{1}{x^2}$$

Given, $f(b) - f(a) = (b-a)f'(c)$

$$\Rightarrow \frac{1}{b} - \frac{1}{a} = (b-a) \left(-\frac{1}{c^2} \right)$$

$$\Rightarrow \frac{a-b}{ab} = \frac{(a-b)}{c^2}$$

$$\Rightarrow \frac{1}{c^2} = \frac{1}{ab} \Rightarrow c = \sqrt{ab}$$



70. $f(x) = (x - 1)^2 = x^2 - 2x + 1$
 $\therefore f(0) = 0, f(2) = 2$ and $f'(x) = 2x - 2$

By mean value theorem,

$$f'(c) = \frac{f(2) - f(0)}{2 - 0}$$

$$\Rightarrow 3c^2 - 4c + 1 = \frac{2 - 0}{2 - 0} = 1$$

$$\Rightarrow 3c^2 - 4c = 0$$

$$\Rightarrow c(3c - 4) = 0 \Rightarrow c = 0, c = \frac{4}{3}$$

71. $f(x) = (x - 1)(x - 2)$
 $\therefore f(a) = f(0) = 0, f(b) = f\left(\frac{1}{2}\right) = \frac{3}{8}$ and

$$f'(x) = (x - 1)(x - 2) + (x - 2) + (x - 1)$$

$$\Rightarrow f'(c) = (c - 1)(c - 2) + c(c - 2) + c(c - 1)$$

$$\Rightarrow f'(c) = c^2 - 3c + 2 + c^2 - 2c + c^2 - c$$

$$\Rightarrow f'(c) = 3c^2 - 6c + 2$$

Given, $f'(c) = \frac{f(b) - f(a)}{b - a}$


$$\Rightarrow 3c^2 - 6c + 2 = \frac{\frac{3}{8} - 0}{\frac{1}{2} - 0} = \frac{3}{4}$$

$$\Rightarrow 3c^2 - 6c + \frac{5}{4} = 0$$

$$\therefore c = \frac{6 \pm \sqrt{36 - 15}}{2 \times 3} = \frac{6 \pm \sqrt{21}}{6} = 1 \pm \frac{\sqrt{21}}{6}$$

72. $f(x) = 1 - x^3 - x^5 \Rightarrow f'(x) = -3x^2 - 5x^4$
 $\Rightarrow f'(x) < 0$ for all values of x .

73. $f(x) = 2x^3 + 3x^2 - 12x + 5$
 $\therefore f'(x) = 6x^2 + 6x - 12$
 $= 6(x^2 + x - 2)$
 $= 6(x + 2)(x - 1)$



Increasing at $(-\infty, -2) \cup (1, \infty) = I_1$
 Decreasing at $(-2, 1) = I_2$

74. $f(x) = \frac{1}{1 + |x|}$

$$\therefore f'(x) = \frac{1 + |x| - |x|}{(1 + |x|)^2} = \frac{1}{(1 + |x|)^2} > 0$$

\therefore the given function is increasing.

75. $f(x) = \frac{\log x}{x}$

$$\therefore f'(x) = \frac{1 - \log x}{x^2} < 0$$

$$\Rightarrow 1 - \log x < 0$$

$$\Rightarrow 1 < \log x$$

$$\Rightarrow \log x > 1$$

$$\Rightarrow x > e$$

76. $\frac{d}{dx}(f(x)) = \frac{-2 - x}{(x + 1)^2}$

For $x > 0$,

$$\frac{d}{dx}(f(x)) < 0$$

77. $f'(x) = 3x^2 + 3x + 3 = 3(x^2 + x + 1)$

$$= 3 \left[\left(x + \frac{1}{2}\right)^2 + \frac{3}{4} \right] \geq \frac{9}{4} > 0$$

$\therefore f(x)$ is an increasing function.

78. $f'(x) = 2e^{-x} - 2e^{-x} = e^{-x}(2 - x)$

Since, f is increasing, $f'(x) > 0$

$$\Rightarrow e^{-x}(2 - x) > 0 \Rightarrow (2 - x) > 0$$

$$\Rightarrow x < 2, 2 - x > 0 \text{ or } x < 0, 2 - x < 0$$

$$\Rightarrow x > 0, 2 > x \text{ or } x < 0, 2 < x$$

$$\Rightarrow 0 < x < 2 \text{ or } 2 < x < 0 \text{ (Not possible)}$$

$$\Rightarrow 0 < x < 2 \Rightarrow x \in (0, 2)$$

79. $f(x) = e^a + e^{-a}$

$$\therefore f'(x) = a(e^a - e^{-a}) < 0$$

But, $a < 0$

$$\therefore e^a - e^{-a} > 0$$

$$\Rightarrow e^a > e^{-a}$$

$$\Rightarrow a > -a$$

$$\Rightarrow 2a > 0$$

$\therefore a > 0$, then $x < 0 \dots [\because a < 0]$

80. $f(x) = 3kx^2 - 18x + 9$
 $= 3(kx^2 - 6x + 3)$

Since, $f(x)$ is increasing on $\mathbb{R} \therefore f'(x) > 0$

$$\therefore kx^2 - 6x + 3 > 0 \forall x \in \mathbb{R}$$

$$\Rightarrow k > 0 \text{ and } 36 - 12k < 0$$

$$\Rightarrow k > 3$$

$\dots [\because a^2 + b^2 + c > 0 \forall x \in \mathbb{R}$
 $\Rightarrow a > 0 \text{ and } b^2 - 4ac < 0]$

Hence, $f(x)$ is increasing on \mathbb{R} if $k > 3$.



81. Since, $f(x)$ is increasing for all x .


$\therefore f'(x) > 0$ for all x

$$\Rightarrow \frac{K-2}{(\sin x + \cos x)^2} > 0 \text{ for all } x$$

$$\Rightarrow K-2 > 0 \Rightarrow K > 2$$

82. The graph of $\operatorname{cosec} x$ is opposite in interval

$$\left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$$



At the point $x = \pi$, $\operatorname{cosec} x$ is not defined and

$$x \in \left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$$

\therefore equation is neither increasing nor decreasing.

Also $\frac{d}{dx}(\tan x) = \sec^2 x > 0$ which is an increasing function.

Also $y = x^2$ is a parabola, which is increasing

Also $y = |x - 1|$ is a V-shaped upward curve, which is always increasing.

\therefore option (A) is the correct answer.

83. Let $f(x) = x + \frac{1}{x}$

$$\therefore f'(x) = 1 - \frac{1}{x^2} \leq 0 \Rightarrow 1 \leq \frac{1}{x^2} \Rightarrow x^2 \leq 1$$

$$\Rightarrow x \in [-1, 1]$$

84. Since $f'(x) = \frac{3}{(x+1)^2}$ is greater than '0' in

interval $(-\infty, \infty)$, therefore $f(x) = \frac{-2}{x+1}$ is

increasing in interval $(-\infty, \infty)$ or \mathbb{R} .

85. Let $f(x) = \sin x - b + c$

$$\therefore f'(x) = \cos x - b > 0 \Rightarrow \cos x > b \Rightarrow b < -1$$

$$86. f(x) = 4x^3 - \frac{x^3}{3} \Rightarrow f'(x) = 4^3 - x^2$$

$$\text{For } f(x) \text{ to be increasing, } 4^3 - x^2 > 0$$

$$\Rightarrow x^2(4 - x) > 0$$

\therefore the function is increasing for $x > \frac{1}{4}$

Similarly, decreasing for $x < \frac{1}{4}$

$$87. f(x) = 2x^3 - 15x^2 + 36x + 1$$

$$\therefore f'(x) = 6x^2 - 30x + 36 = 6(x^2 - 5x + 6)$$

$$= 6(x-2)(x-3)$$

To be monotonic decreasing, $f'(x) < 0$

$$\Rightarrow (x-2)(x-3) < 0 \Rightarrow x \in (2, 3)$$

$$88. \text{ As } f(x) = \sin 2x \Rightarrow f'(x) = 2 \cos 2x$$

$$\text{Here, } f'(x) > 0 \text{ in } \left(0, \frac{\pi}{4}\right) \text{ and } f'(x) < 0 \text{ in } \left(\frac{\pi}{4}, \frac{\pi}{2}\right)$$

$$89. f(x) = x + \cos x \Rightarrow f'(x) = 1 - \sin x$$

$f'(x) > 0$ for all values of x .

$\therefore f(x)$ is always increasing.

$$90. f(x) = \frac{\log x}{x}$$

$$\therefore f'(x) = \frac{1}{x} - \frac{\log x}{x^2} = \frac{1 - \log x}{x^2}$$

For $f(x)$ to be increasing, $f'(x) > 0$

$$\Rightarrow 1 - \log x > 0 \Rightarrow 1 > \log x \Rightarrow x < e$$

$\therefore f(x)$ is increasing in the interval $(0, e)$.

$$91. f(x) = 1 - e^{-\frac{x^2}{2}}$$

$$\Rightarrow f'(x) = -e^{-\frac{x^2}{2}} \left(-x\right) = x e^{-\frac{x^2}{2}}$$

For $f(x)$ to be increasing, $f'(x) > 0$

$$\Rightarrow x e^{-\frac{x^2}{2}} > 0$$

$\Rightarrow f(x)$ is decreasing for $x < 0$ and increasing for $x > 0$.

$$92. f(x) = \frac{a \sin x + b \cos x}{c \sin x + d \cos x}$$

$f(x)$ will be decreasing, if $f'(x) < 0$

$$\therefore \frac{1}{(c \sin x + d \cos x)^2} [(c \sin x + d \cos x)(a \cos x - b \sin x)$$

$$- (a \sin x + b \cos x)(c \cos x - d \sin x)] < 0$$

$$\Rightarrow a c \sin x \cos x - b c \sin^2 x + a d \cos^2 x$$

$$- b d \sin x \cos x - a c \sin x \cos x + a d \sin^2 x$$

$$- b c \cos^2 x + b d \sin x \cos x < 0$$

$$\Rightarrow a d (\sin^2 x + \cos^2 x) - b c (\sin^2 x + \cos^2 x) < 0$$

$$\Rightarrow a d - b c < 0$$



93. $f(x) = x^4 - 62x^2 + a + 9$ (i)
 $\therefore f'(x) = 4x^3 - 124x + a$
 For maximum or minimum, $f'(x) = 0$
 $\Rightarrow 4x^3 - 124x + a = 0$
 Since, $x = 1$ is a root of (i).
 $\therefore f'(1) = 4 - 124 + a = 0 \therefore a = 120$
94. $f(x) = (x - \alpha)(x - \beta) = x^2 - (\alpha + \beta)x + \alpha\beta$
 $\therefore f'(x) = 2x - (\alpha + \beta)$
 For maximum or minimum, $f'(x) = 0$
 $\Rightarrow x = \frac{\alpha + \beta}{2}$
 Now $f''(x) = 2 > 0$
 $\therefore f(x) = f\left(\frac{\alpha + \beta}{2}\right)$
 $= \left(\frac{\alpha + \beta}{2} - \alpha\right)\left(\frac{\alpha + \beta}{2} - \beta\right)$
 $= \left(\frac{\beta - \alpha}{2}\right)\left(\frac{\alpha - \beta}{2}\right) = -\frac{(\alpha - \beta)^2}{4}$
95. $y = e^{-x}$
 $\therefore y' = -e^{-x} = -e^{-x}$
 $\therefore y'' = e^{-x}$
 At $x = -1$,
 $y'' = -e^{-1} + e^{-1} + e^{-1} = \frac{1}{e} > 0$
 \therefore Minimum at $x = -1$.
96. $f(x) = 5x^5 - 5x^4 + 5x^3 - 10x^2$
 $\therefore f'(x) = 5x^4 - 20x^3 + 15x^2$
 For maximum or minimum, $f'(x) = 0$
 $\Rightarrow 5x^2(x^2 - 4x + 3) = 0$
 $\Rightarrow x^2(x - 3)(x - 1) = 0$
 $\Rightarrow x = 0, x = 3, x = 1$
 $f''(x) = 20x^3 - 60x^2 + 30x = 10x(2x^2 - 6x + 3)$
 $f''(0) = 0$
 $f''(3) = \text{Positive (Minima)}$
 $f''(1) = \text{Negative (Maxima)}$
 $\therefore (p, q) = (1, 3)$
97. $\frac{d}{dx} = a + 2b + 1 \Rightarrow \left(\frac{d}{dx}\right)_{x=1} = a + 2b + 1 = 0$
 $\Rightarrow a = -2b - 1$ and $\left(\frac{d}{dx}\right)_{x=2} = \frac{a}{2} + 4b + 1 = 0$
 $\Rightarrow \frac{-2b - 1}{2} + 4b + 1 = 0$
 $\Rightarrow -b + 4b + \frac{1}{2} = 0$
 $\Rightarrow 3b = \frac{-1}{2} \Rightarrow b = \frac{-1}{6}$ and $a = \frac{1}{3} - 1 = \frac{-2}{3}$

98. $f(x) = 3x^4 - 4x^3$
 $\therefore f'(x) = 12x^3 - 12x^2$
 $\therefore f'(x) = 0 \Rightarrow x = 1, 0$
 Now $f''(x) = 36x^2 - 24x$
 $f''(1) = 12 > 0$
 $f''(0) = 0$
 $f(1) = 3 - 4 = -1$
 $f(-1) = 3 + 4 = 7$
 $f(2) = 48 - 32 = 16$
 \therefore Maximum at 2 and Minimum at 1 and
 Maximum value is 16 and Minimum value is -1.
99. Let $f(x) = x^3 - 12x^2 + 36x + 17$
 $\therefore f'(x) = 3x^2 - 24x + 36 = 0$ at $x = 2, 6$
 Again $f''(x) = 6x - 24$ is -ve at $x = 2$
 So that $f(6) = 17$, $f(2) = 49$
 At the end points, $f(1) = 42$, $f(10) = 177$
 So that $f(x)$ has its maximum value as 177
100. $x + y = 16 \Rightarrow y = 16 - x$
 $\Rightarrow z = x^2 + y^2 = x^2 + (16 - x)^2$
 Let $z = x^2 + (16 - x)^2$
 $\Rightarrow z' = 4x - 32$
 To be minimum of z , $z' > 0$,
 Therefore $4x - 32 = 0 \Rightarrow x = 8, y = 8$
101. $f(x) = (x - 1)^{\frac{1}{3}}(x - 2)$
 $\therefore f'(x) = (x - 1)^{\frac{1}{3}} \cdot 1 + (x - 2) \cdot \frac{1}{3}(x - 1)^{-\frac{2}{3}}$
 $= \frac{4x - 5}{3(x - 1)^{\frac{2}{3}}}$
 For maxima or minima, $f'(x) = 0$
 $\therefore \frac{4x - 5}{3(x - 1)^{\frac{2}{3}}} = 0$
 $\therefore x = \frac{5}{4}$
 $\therefore f(1) = (1 - 1)^{\frac{1}{3}}(1 - 2) = 0$
 $f\left(\frac{5}{4}\right) = \left(\frac{5}{4} - 1\right)^{\frac{1}{3}}\left(\frac{5}{4} - 2\right) = \frac{-3}{4^{\frac{2}{3}}}$, $f(9) = 14$
 \therefore absolute minimum occurs at $x = \frac{5}{4}$ and min.
 value = $\frac{-3}{4^{\frac{2}{3}}}$
 Absolute maximum occurs at $x = 9$ and max.
 value = 14.



102. Let $f(x) = \sqrt{1-x^2}$
 $\Rightarrow f'(x) = \frac{1-2x^2}{\sqrt{1-x^2}} = 0 \Rightarrow x = \pm \frac{1}{\sqrt{2}}$
 But as $x > 0$, we have $x = \frac{1}{\sqrt{2}}$
 Now,

$$f''(x) = \frac{\sqrt{1-x^2}(-4x) - (1-2x^2)\frac{-x}{\sqrt{1-x^2}}}{(1-x^2)^2}$$

$$= \frac{2x^3 - 3}{(1-x^2)^{3/2}}$$
 $\Rightarrow f''\left(\frac{1}{\sqrt{2}}\right) = -ve$
 $\therefore f(x)$ is maximum at $x = \frac{1}{\sqrt{2}}$
103. Let x and y be two natural numbers such that
 $x + y = 10$ and the product is P .
 $P = x(10-x) = 10x - x^2 = f(x)$
 $\therefore f'(x) = 10 - 2x$
 $\therefore f''(x) = -2$
 Roots of $f'(x) = 0$,
 i.e., $10 - 2x = 0$ i.e., $x = 5$
 $f'(5) = 10 - 10 = 0$
 $\therefore f$ is maximum when $x = 5$, $y = 5$
 \therefore The product is maximum if $x = 5$, $y = 5$
104. $2(x + y) = 24$
 $\therefore x + y = 12$
 $\therefore y = 12 - x$
 $f(x) = xy = x(12-x) = 12x - x^2$
 $\therefore f'(x) = 12 - 2x = 0$
 $\therefore x = 6$ At $x = 6$, $y = 6$
 \therefore max area is 36 m^2 .
105. Let $x + y = 3$
 According to the given condition,
 $f(x) = x^2 \times (3-x) = 3x^2 - x^3 \dots(i)$
 $\therefore f'(x) = 6x - 3x^2 = 0$
 $\therefore 3x(2-x) = 0$
 $\therefore x = 0$, $x = 2$
 Now $f''(x)$ is max at $x = 2$
 Its maximum value is 4 \dots [From (i)]
106. Let one number be $(100 - x)$ and then another is x . Therefore $f(x) = 2(100 - x) + x^2$
 $= 200 - 2x + x^2$
 $\therefore f'(x) = 0 \Rightarrow 2 - 2 = 0 \Rightarrow x = 1$
 Here $f''(x) = 2 > 0$
 Therefore function is minimum at $x = 1$.
 So the numbers are 99 and 1.

107. According to the given condition,
 $2x + 2y = 100 \Rightarrow x + y = 50 \dots(i)$
 Let the area of rectangle be A .
 $\therefore A = xy \Rightarrow y = \frac{A}{x}$
 Put in (i), we have $x + \frac{A}{x} = 50 \Rightarrow A = 50x - x^2$
 $\Rightarrow \frac{dA}{dx} = 50 - 2x$
 For maximum area, $\frac{dA}{dx} = 0$
 $\therefore 50 - 2x = 0 \Rightarrow x = 25$ and $y = 25$
 Hence, adjacent sides are 25 and 25 cm.
108. Let the number be x , then the function
 $f(x) = \frac{x^2}{x^2 + 16}$
 On differentiating with respect to x , we get
 $\Rightarrow f'(x) = \frac{(x^2 + 16) \cdot 1 - (x^2) \cdot (2x)}{(x^2 + 16)^2}$
 $= \frac{x^2 + 16 - 2x^2}{(x^2 + 16)^2}$
 $= \frac{16 - x^2}{(x^2 + 16)^2}$
 Put $f'(x) = 0$ for maxima or minima
 $f'(x) = 0 \Rightarrow 16 - x^2 = 0 \Rightarrow x = 4, -4$
 Again differentiating
 $f''(x) = \frac{(x^2 + 16)^2(-2x) - (16 - x^2)2(x^2 + 16)2x}{(x^2 + 16)^4}$
 At $x = 4$, $f''(x) < 0$ and at $x = -4$, $f''(x) > 0$
 \therefore Least value of $f(x) = \frac{-4}{16 + 16} = -\frac{1}{8}$
109. Let $y = x^2 \Rightarrow \log y = 2 \log x$, ($x > 0$)
 Differentiating, $\frac{dy}{dx} = 2x^2(1 + \log x)$;
 $\therefore \frac{dy}{dx} = 0$
 $\Rightarrow \log x = -1 \Rightarrow x = e^{-1} = \frac{1}{e}$
 \therefore Stationary point is $\left(\frac{1}{e}, \frac{1}{e}\right)$
110. $x + y = 8$ $\therefore y = 8 - x$
 Now $f(x) = xy = x(8 - x) = 8x - x^2$
 $\therefore f'(x) = 8 - 2x$
 For maximum value of $f(x)$, $f'(x) = 0$
 $\therefore 8 - 2x = 0$ and $x = 4$
 So, maximum value of $f(x) = 4 \times 4 = 16$



111. $f(x) = 2x^3 - 21x^2 + 36x - 30$
 $\Rightarrow f'(x) = 6x^2 - 42x + 36$
 $\therefore f'(x) = 0 \Rightarrow x = 6, 1$ and $f''(x) = 12x - 42$
 Here $f''(1) = -30 < 0$ and $f''(6) = 30 > 0$
 $\therefore f(x)$ has maxima at $x = 1$ and minima at $x = 6$.

112. $f(x) = \cos x + \cos(\sqrt{2}x)$
 $\therefore f'(x) = -\sin x - \sqrt{2}\sin(\sqrt{2}x) = 0$
 $\therefore x = 0$ is the only solution.
 $f''(x) = -\cos x - 2\cos(\sqrt{2}x) < 0$ at $x = 0$
 Hence, maxima occurs at $x = 0$.

113. Let $f(x) = x^3 - 18x^2 + 96x$
 $\therefore f'(x) = 3x^2 - 36x + 96$
 For maximum or minimum, $f'(x) = 0$
 $\Rightarrow x^2 - 12x + 32 = 0 \Rightarrow (x-4)(x-8) = 0$
 $\Rightarrow x = 4, 8$
 Now, $f''(x) = 6x - 36$
 At $x = 4$, $f''(x) = 24 - 36 = -12 < 0$
 At $x = 4$, $f(x)$ will be maximum
 and $[f(4)]_{\max} = 64 - 288 + 384 = 160$
 At $x = 8$, $\frac{d^2}{dx^2} = 48 - 36 = 12 > 0$
 At $x = 8$, $f(x)$ will be minimum and
 $[f(8)]_{\min} = 128$

114. Let $PQ = a$ and $PR = b$, then $\Delta = \frac{1}{2}ab \sin \theta$
 $\therefore -1 \leq \sin \theta \leq 1$
 Since, area is maximum when $\sin \theta = 1$
 $\Rightarrow \theta = \frac{\pi}{2}$

115. Here $f(x) = |\sin 4x + 3|$
 We know that minimum value of $\sin x$ is -1
 and maximum is 1 .
 Hence minimum $|\sin 4x + 3| = |-1 + 3| = 2$ and
 maximum $|\sin 4x + 3| = |1 + 3| = 4$

116. $f(x) = |px - 9| + |r|$, $x \in (-\infty, \infty)$
 Where $p > 0$, $q > 0$ and $r > 0$ can assume its
 minimum value only at one point, if $p = q = r$.

117. $f(x) = 3x^4 - 8x^3 + 12x^2 - 48x + 25$
 $\therefore f'(x) = 12x^3 - 24x^2 + 24x - 48$
 $= 12(x^3 - 2x^2 + 2x - 4) = 12[(x-2)(x^2+2)]$
 For maximum or minimum of $f(x)$, $f'(x) = 0$
 $\Rightarrow 12[(x-2)(x^2+2)] = 0$
 $\Rightarrow x = 2$.
 Now, $f''(x) = 12(3x^2 - 4x + 2)$
 $\therefore f''(2) = 12(12 - 8 + 2) = 72 > 0$

- $\therefore f$ has minimum at $x = 2$ and the minimum
 value of f at $x = 2$ is
 $f(2) = 48 - 64 + 48 - 96 + 25 = -39$

118. $f(x) = (x - \alpha)(x - \beta) = x^2 - (\alpha + \beta)x + \alpha\beta$
 $\therefore f'(x) = 2x - (\alpha + \beta)$
 For maximum or minimum of $f(x)$, $f'(x) = 0$
 $\Rightarrow 2x - (\alpha + \beta) = 0$
 Now, $f''(x) = 2 > 0$

- $\therefore f$ has minimum at $x = \frac{\alpha + \beta}{2}$

and the minimum value of f at $x = \frac{\alpha + \beta}{2}$ is

$$\left(\frac{\alpha + \beta}{2} - \alpha\right)\left(\frac{\alpha + \beta}{2} - \beta\right)$$

$$= \left(\frac{\beta - \alpha}{2}\right)\left(\frac{\alpha - \beta}{2}\right) = -\frac{(\alpha - \beta)^2}{4}$$

119. Let $x + y = 20 \Rightarrow x = 20 - y$ (i)
 and $x^3 + y^3 = z$
 $\Rightarrow z = (20 - y)^3 + y^3 \Rightarrow z = 400^3 + y^3 - 40^4$

- $\therefore \frac{dz}{dy} = 1200y^2 + 5y^4 - 160^3$

For maximum or minimum,

$$\frac{dz}{dy} = 0$$

$$\Rightarrow 1200y^2 + 5y^4 - 160^3 = 0$$

$$\Rightarrow y = 12, 20$$

$$\frac{d^2z}{dy^2} = 2400y + 20y^3 - 480^2$$

- $\therefore \left(\frac{d^2z}{dy^2}\right)_{y=12} = -5760 < 0$

- $\therefore z$ is maximum at $y = 12$.

$$\text{From (i), } x = 20 - 12 = 8$$

- \therefore the parts of 20 are 12 and 8.

120. Let $x = \sin^p \theta \cdot \cos^q \theta$

- $\therefore \frac{dx}{d\theta} = p \sin^{p-1} \theta \cdot \cos \theta \cdot \cos^q \theta - q \cos^{q-1} \theta \cdot \sin^p \theta$

$$= (p \sin^{p-1} \theta \cdot \cos^q \theta - q \cos^{q-1} \theta \cdot \sin^p \theta)$$

- $\therefore \frac{dx}{d\theta} = p \sin^{p-1} \theta \cdot \cos^q \theta - q \cos^{q-1} \theta \cdot \sin^p \theta$

For maximum or minimum,

$$\frac{dx}{d\theta} = 0$$


- $\therefore \tan^2 \theta = \frac{p}{q} \Rightarrow \tan \theta = \pm \sqrt{\frac{p}{q}}$

- \therefore Point of maxima $x = \tan^{-1} \sqrt{\frac{p}{q}}$



121. $4 + 2\pi r = a$
 $A = 2 + \pi r^2 = \frac{1}{16} (a - 2\pi r)^2 + \pi r^2$
 $\therefore \frac{dA}{dr} = 0$ gives $r = \frac{a}{2(\pi + 4)}$, thus $\frac{d^2A}{dr^2} > 0$
 and hence minimum,
 $\therefore 4 = a - 2\pi r = a - \frac{a\pi}{\pi + 4} = \frac{4a}{\pi + 4}$
 $\therefore a = \frac{4(\pi + 4)}{\pi + 4} = 4$
 $\therefore A = 2 + \pi r^2 = \frac{a^2}{4(\pi + 4)}$

122.



Let OAB be a given sector of a circle of a radius r cm such that arc $AB = l$ cm, and $\angle AOB = \theta$ radians.

Then

$2r + l = 20$ (i)

$\frac{l}{r} = \theta$ (ii)

$A = \frac{1}{2} r^2 \theta$ (iii)

From (i), (ii), (iii), we get

$A = \frac{1}{2} r^2 \times \frac{l}{r} = \frac{1}{2} r l = \frac{1}{2} r(20 - 2r)$

$\Rightarrow A = 10r - r^2$ (iv)

Now, $\frac{dA}{dr} = 10 - 2r = 0 \Rightarrow r = 5$

$\therefore \frac{d^2A}{dr^2} = -2 < 0$

$\therefore A$ is maximum at $r = 5$
 Hence the maximum area
 $= 10 \times 5 - 25 = 25 \text{ cm}^2$ [From (iv)]

123. $2l + 2\pi R = 440$

$\therefore l + \pi R = 220$ (i)



Now $f(R) = l(2R) = (220 - \pi R)(2R)$

$\therefore f(R) = 440R - 2\pi R^2$

$\therefore f'(R) = 440 - 4\pi R = 0$

$\therefore 0 = 110 - \pi R$

$\therefore 110 = \frac{22}{7} R$

$\therefore R = 35$ $\therefore 2R = 70$


From (i),

$l + \frac{22}{7} \times 35 = 220$

$\therefore l + 110 = 220$

$\therefore l = 110$

124. Let the length of side of each square cut out be x cm. Then, each side of base of the box is $(12 - 2x)$ cm and x cm will be height of box.



$V =$ Volume of box

$= (12 - 2x)^2 \times x = 4(36 - 24x + 12x^2 - x^3)$
 $= 4(36 - 24x + 12x^2 - x^3)$

$\Rightarrow \frac{dV}{dx} = 4(36 - 24 + 24x - 3x^2)$
 $= 12(12 - 8x + 3x^2)$

and $\frac{d^2V}{dx^2} = 4(6 - 6x)$

Now, $\frac{dV}{dx} = 0 \Rightarrow 12 - 8x + 3x^2 = 0$

$\Rightarrow (3x - 2)(x - 6) = 0 \Rightarrow x = 2$ or $x = 6$

But $x < 6$

$\therefore x = 2$

For $x = 2$, $\frac{d^2V}{dx^2} = 4(12 - 24) = -48 < 0$

\therefore Volume is maximum when each square of 2 cm length is cut out from each corner.

125. Given equation is $10s = 10ut - 4.9t^2$

$\Rightarrow s = ut - 4.9t^2$

$\Rightarrow \frac{ds}{dt} = u - 9.8t = v$

When stone reaches the maximum height, then $v = 0$


$\Rightarrow u - 9.8t = 0 \Rightarrow u = 9.8t$

But time $t = 5$ sec

So the value of $u = 9.8 \times 5 = 49.0$ m/sec



126. Let L be the lamp and PQ be the man and OQ = metre be his shadow and let MQ = metre.



$\frac{d}{dt}$ = speed of the man = 3 m/s (given)

Since, ΔOPQ and ΔOLM are similar.

$$\therefore \frac{OM}{OQ} = \frac{LM}{PQ} \Rightarrow \frac{OQ + MQ}{OQ} = \frac{5}{2}$$

$$\Rightarrow \frac{3}{2} = \frac{OQ + MQ}{OQ}$$

$$\therefore \frac{d}{dt} = \frac{3}{2} \cdot \frac{d}{dt}$$

$$\Rightarrow 3 = \frac{3}{2} \cdot \frac{d}{dt}$$

$$\Rightarrow \frac{d}{dt} = 2 \text{ m/s.}$$

127. Let A, P and x be the area, perimeter and length of the side of the square respectively at time t seconds. Then, $A = x^2$ and $P = 4x$

$$\therefore P = 4\sqrt{A}$$

$$\therefore \frac{dP}{dt} = 4 \cdot \frac{1}{2\sqrt{A}} \cdot \frac{dA}{dt}$$

$$= \frac{2}{\sqrt{A}} \cdot \frac{dA}{dt} = \frac{2}{16} \cdot 2 = \frac{1}{4} \text{ cm/sec.}$$

128. Let

$$f(A) = \cos A \cos B = \cos A \cos\left(\frac{\pi}{2} - A\right) = \cos A \sin A$$

$$\therefore f'(A) = \cos^2 A - \sin^2 A = \cos 2A$$

For maximum or minimum,

$$f'(A) = 0 \Rightarrow \cos 2A = 0$$

$$\Rightarrow 2A = \frac{\pi}{2} \Rightarrow A = \frac{\pi}{4}$$

Now, $f''(A) = -2 \sin 2A$

$$= -2 \sin \frac{\pi}{2} = -2 < 0$$

$$\therefore f(A) \text{ is maximum at } A = \frac{\pi}{4}.$$

$$\therefore \text{Maximum value} = \cos \frac{\pi}{4} \sin \frac{\pi}{4} = \frac{1}{2}$$

129. Since, $f(x)$ satisfies all the conditions of Rolle's theorem.

$$\therefore f(3) = f(5) = 0$$

$\Rightarrow x = 3$ and $x = 5$ are the roots of $f(x)$.

$$\Rightarrow f(x) = (x - 3)(x - 5) = x^2 - 8x + 15$$

$$\therefore \int_3^5 f(x) dx = \int_3^5 (x^2 - 8x + 15) dx$$

$$= \left[\frac{x^3}{3} - 4x^2 + 15x \right]_3^5$$

$$= \frac{1}{3}(125 - 27) - 4(25 - 9) + 15(5 - 3)$$

$$= -\frac{4}{3}$$



Competitive Thinking

1. $y = x^2 - \frac{1}{2}$

$$\therefore \frac{dy}{dx} = 2x + \frac{2}{3}$$

$$\therefore \left(\frac{dy}{dx}\right)_{(-1,0)} = 2(-1) + \frac{2}{(-1)^3} = -4$$

$$\therefore \text{Slope of normal at } (-1, 0) = -\frac{1}{\left(\frac{dy}{dx}\right)_{(-1,0)}} = \frac{1}{4}$$

2. For the point $(2, -1)$ on the curve $y = t^2 + 3t - 8$, $x = 2t^2 - 2t - 5$, we have

$$t^2 + 3t - 8 = 2 \text{ and } 2t^2 - 2t - 5 = -1$$

$$\Rightarrow (t + 5)(t - 2) = 0 \text{ and } (t - 2)(t + 1) = 0$$

$$\Rightarrow t = 2$$

$$\text{Now, } \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{4t - 2}{2t + 3}$$

$$\therefore \left(\frac{dy}{dx}\right)_{(t=2)} = \frac{4(2) - 2}{2(2) + 3} = \frac{6}{7}$$

3. Slope of the normal = $-\frac{1}{\frac{dy}{dx}}$

$$\Rightarrow \tan \frac{3\pi}{4} = \frac{-1}{\left(\frac{dy}{dx}\right)_{(3,4)}}$$

$$\Rightarrow \left(\frac{dy}{dx}\right)_{(3,4)} = 1 \Rightarrow f'(3) = 1$$



$$4. \quad = a^3 + b + 4$$

$$\therefore \frac{d}{d} = 3a^2 + b$$

$$\text{Slope of tangent at } (2, 14) = \left(\frac{d}{d}\right)_{(2,14)}$$

$$\Rightarrow 21 = 3a(2)^2 + b$$

$$\Rightarrow 21 = 12a + b \quad \dots(i)$$

$$= a^3 + b + 4$$

$$\therefore 14 = a(8) + b(2) + 4$$

$$\Rightarrow 8a + 2b = 10 \quad \dots(ii)$$

On solving (i) and (ii), we get

$$a = 2, b = -3$$

$$5. \quad = t^2 - 1, \quad = t^2 - t$$

$$\therefore \frac{d}{d} = \frac{\frac{d}{dt}}{\frac{d}{dt}} = \frac{2t-1}{2t}$$

Since, the tangent is perpendicular to X-axis.

$$\therefore \frac{d}{d} = 0 \Rightarrow \frac{2t}{2t-1} = 0 \Rightarrow t = 0$$

$$6. \quad = 3 \Rightarrow \frac{d}{d} = 3^2$$

According to the given condition,

$$3^2 =$$

$$\Rightarrow 3^2 = 3 \quad \dots[\because = 3]$$

$$\Rightarrow = 0, 3$$

Thus, the two points are (0, 0) and (3, 27).

$$7. \quad = 2 - 3 + 2 \Rightarrow \frac{d}{d} = 2 - 3$$

Slope of the given line = 1

Since, the tangent is perpendicular to the given line.

$$\therefore (2 - 3)(1) = -1$$

$$\Rightarrow = 1$$

$$\text{At } = 1, \quad = 0$$

\therefore the required point is (1, 0).

$$8. \quad \text{Given equation of curve is } = \sqrt{-1}$$

Slope of tangent to the curve is

$$\frac{d}{d} = \frac{1}{2\sqrt{-1}}$$

Slope of line $2 + - 5 = 0$ is -2

Since the tangent is perpendicular to the given line,

$$\left(\frac{1}{2\sqrt{-1}}\right)(-2) = -1$$

$$\Rightarrow \sqrt{-1} = 1 \quad \Rightarrow = 2$$

$$= \sqrt{-1} = \sqrt{2-1} = 1$$

$$\therefore (,) = (2, 1)$$

$$9. \quad = p^3 + q \quad \dots(i)$$

Differentiating both sides w.r.t. , we get

$$2 \cdot \frac{d}{d} = 3p^2$$

$$\Rightarrow \frac{d}{d} = \frac{3p}{2} \left(\frac{2}{2}\right)$$

$$\therefore \left(\frac{d}{d}\right)_{(2,3)} = \frac{3p}{2} \times \frac{4}{3} = 2p$$

Since the line touches the curve, their slopes are equal.

$$\therefore 2p = 4 \Rightarrow p = 2$$

Since, (2,3) lies on $= p^3 + q$.

$$\therefore 9 = 2 \times 8 + q \Rightarrow q = -7$$

$$10. \quad = a^3 + b \quad \dots(i)$$

Differentiating both sides w.r.t. , we get

$$2 \cdot \frac{d}{d} = 3a^2$$

$$\Rightarrow \frac{d}{d} = \frac{3a}{2} \left(\frac{2}{2}\right)$$

$$\therefore \left(\frac{d}{d}\right)_{(2,3)} = \frac{3a}{2} \times \frac{4}{3} = 2a$$

Since, the line touches the curve, their slopes are equal.

$$\therefore 2a = 4 \Rightarrow a = 2$$

Since, (2,3) lies on $= a^3 + b$.

$$\therefore 9 = 2 \times 8 + b \Rightarrow b = -7$$

$$\text{Now, } 7a + 2b = 7(2) + 2(-7) = 0$$

$$11. \quad = \frac{1}{2} \quad \dots(i)$$

$$\therefore \frac{d}{d} = \frac{-1}{2}$$

$$\therefore \text{Slope of tangent to the curve} = \frac{-1}{2}$$

Slope of $= -4 + b$ is -4 .

$$\therefore \frac{-1}{2} = -4 \quad \Rightarrow = \pm \frac{1}{2}$$

From (i),

$$= \pm 2$$

Putting the values of and in $= -4 + b$, we get

$$b = \pm 4$$



12. $y^2 = 2(x - 3)$ (i)
 Differentiating w.r.t. x , we get
 $2 \frac{dy}{dx} = 2 \Rightarrow \frac{dy}{dx} = 1$
- \therefore Slope of the normal $= \frac{-1}{\frac{dy}{dx}} = -1$
- Slope of the given line $= 2$.
 Since, the normal is parallel to the given line.
 \therefore Slope $= -2$
 From (i), $y = 5$
 \therefore the required point is $(5, -2)$.
13. Given equation of curve is
 $x^2 - 4y^2 = 1$ (i)
 Slope of tangent to the curve is
 $\frac{dy}{dx} = \frac{1}{4}$
- Slope of line is $y = 2$ is $\frac{1}{2}$
 Since, the tangent is parallel to the given line,
 $\frac{1}{4} = \frac{1}{2}$
 \therefore Slope $= 2$
 Substituting $y = 2$ in equation (i), we get
 $(2x)^2 - 4(2)^2 = 1$
 \therefore tangent is parallel to curve at zero point.
14. $x = a(1 + \cos \theta)$ and $y = a \sin \theta$
 $\therefore \frac{dx}{d\theta} = -a \sin \theta$ and $\frac{dy}{d\theta} = a \cos \theta$
- $\therefore \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = -\cot \theta$
- \therefore slope of the normal $= -\frac{1}{\frac{dy}{dx}} = \frac{-1}{-\cot \theta} = \tan \theta$
- \therefore equation of the normal at θ is
 $-a \sin \theta = \tan \theta [x - a(1 + \cos \theta)]$
 Clearly, this line passes through $(a, 0)$.
15. $y^2 = 12$ (i)
 Differentiating w.r.t. x , we get
 $2 \frac{dy}{dx} = 12 \Rightarrow \frac{dy}{dx} = 6$
- \therefore slope of the normal $= -\frac{1}{\frac{dy}{dx}} = -\frac{1}{6}$

- Slope of the line $x + y = k$ is -1 .
 $\therefore -\frac{1}{6} = -1 \Rightarrow k = 6$
 From (i), $y = 3$
 Putting the values of x and y in $x + y = k$, we get $k = 9$
16. Slope of given line $= -\frac{a}{b}$
 $= -\frac{4}{2} \Rightarrow \frac{dy}{dx} = -\frac{4}{2}$
 $\Rightarrow -\frac{a}{b} = -\frac{4}{2}$
 $\Rightarrow \frac{a}{b} = \frac{4}{2} > 0$
 $\Rightarrow a < 0, b < 0$
17. $y^2 = 4ax$
 $\therefore 2 \frac{dy}{dx} = 4a$
 $\Rightarrow \frac{dy}{dx} = \frac{2a}{y}$
 $\Rightarrow \left(\frac{dy}{dx}\right)_{(at^2, 2at)} = \frac{2a}{2at} = \frac{1}{t}$
- \therefore Slope of tangent (m_1) $= \frac{1}{t}$
 $y^2 - x^2 = a^2$
 $\Rightarrow 2y \frac{dy}{dx} - 2x = 0$
 $\Rightarrow \frac{dy}{dx} = \frac{x}{y}$
 $\Rightarrow \left(\frac{dy}{dx}\right)_{(a \sec \theta, a \tan \theta)} = \frac{a \sec \theta}{a \tan \theta} = \operatorname{cosec} \theta$
- \therefore Slope of normal (m_2) $= \operatorname{cosec} \theta$
 Now, $m_1 \cdot m_2 = -1$
 $\Rightarrow \left(\frac{1}{t}\right) (\operatorname{cosec} \theta) = -1$
 $\Rightarrow t = -\operatorname{cosec} \theta$
18. $9x^2 = 3$ (i)
 Differentiating w.r.t. x , we get
 $18 \frac{dx}{dx} = 3 \cdot 2$
 $\Rightarrow \frac{dx}{dx} = \frac{2}{6}$



$$\therefore \text{slope of the normal} = -\frac{6}{2}$$

Since, the normal to the given curve makes equal intercepts with the axis.

$$\therefore -\frac{6}{2} = \pm 1$$

$$\Rightarrow = -\frac{2}{6} \text{ or } \frac{2}{6}$$

Putting these values in (i), we get

$$9\left(\frac{4}{36}\right) = 3 \Rightarrow = 0 \text{ or } = 4$$

$$\therefore = 0 \text{ or } = -\frac{16}{6} \text{ or } \frac{16}{6} = -\frac{8}{3} \text{ or } \frac{8}{3}$$

$$\therefore \text{the required points are } \left(4, \frac{8}{3}\right) \text{ or } \left(4, -\frac{8}{3}\right).$$

$$19. = \frac{2}{3} x^3 + \frac{1}{2} x^2 \quad \dots(i)$$

$$\therefore \frac{d}{d} = 2x^2 +$$

Since, the tangent makes equal angles with the axis.

$$\therefore \frac{d}{d} = \pm 1$$

$$\Rightarrow 2x^2 + = \pm 1$$

$$\Rightarrow 2x^2 + = 1 \quad (\text{taking +ve sign})$$

$$\Rightarrow 2x^2 + - 1 = 0$$

$$\Rightarrow (2x - 1)(x + 1) = 0$$

$$\Rightarrow = \frac{1}{2}, -1$$

From (i),

$$\text{when } = \frac{1}{2}, = \frac{2}{3} \cdot \frac{1}{8} + \frac{1}{2} \cdot \frac{1}{4} = \frac{5}{24}$$

$$\text{and when } = -1, = \frac{2}{3}(-1) + \frac{1}{2} \cdot 1 = -\frac{1}{6}$$

$$\therefore \text{the required points are } \left(\frac{1}{2}, \frac{5}{24}\right) \text{ and } \left(-1, -\frac{1}{6}\right).$$

$$20. \text{ At } = 4, \\ 4^2 = 8 \Rightarrow = 2$$

$$\text{Now, } ^2 = 8$$

Differentiating w.r.t. , we get

$$2 = 8 \frac{d}{d} \Rightarrow \frac{d}{d} = \frac{1}{4}$$

$$\therefore \left(\frac{d}{d}\right)_{(4,2)} = 1$$

$$\therefore \text{equation of the normal at } (4, 2) \text{ is} \\ -2 = -1(x - 4) \Rightarrow + = 6$$

$$21. \text{ At } t = 1, = (1)^2 = 1 \text{ and } = 2(1) = 2$$

$$\frac{d}{d} = \frac{\frac{d}{dt}}{\frac{d}{dt}} = \frac{2}{2t} = \frac{1}{t}$$

$$\therefore \left(\frac{d}{d}\right)_{t=1} = 1$$

\therefore Equation of the normal at (1, 2) is

$$-2 = -1(x - 1) \Rightarrow + - 3 = 0$$

22. Centre of circle is (1, -2) and point A(2,1) lie on circle.

$$\therefore \text{Equation of normal is } + 2 = \frac{1+2}{2-1}(x - 1)$$

$$\Rightarrow + 2 = 3(x - 1) \Rightarrow = 3 - 5$$

$$23. \left(\frac{a}{a}\right)^n + \left(\frac{b}{b}\right)^n = 2$$

Differentiating w.r.t. , we get

$$n \left(\frac{a}{a}\right)^{n-1} \left(\frac{1}{a}\right) + n \left(\frac{b}{b}\right)^{n-1} \left(\frac{1}{b}\right) \left(\frac{d}{d}\right) = 0$$

$$\Rightarrow \frac{n}{b} \left(\frac{a}{b}\right)^{n-1} \frac{d}{d} = \frac{-n}{a} \left(\frac{a}{a}\right)^{n-1}$$

$$\Rightarrow \frac{d}{d} = \frac{-b}{a} \left(\frac{a}{a}\right)^{n-1} \left(\frac{b}{b}\right)^{n-1}$$

$$\text{Slope of tangent at } (a, b) = \left(\frac{d}{d}\right)_{(a,b)}$$

$$= \frac{-b}{a} \left(\frac{a}{a}\right)^{n-1} \left(\frac{b}{b}\right)^{n-1}$$

$$= \frac{-b}{a}$$

$$\text{Equation of tangent is } -b = \frac{-b}{a}(x - a)$$

$$\Rightarrow a - ab = -b + ab$$

$$\Rightarrow a + b = 2ab$$

$$\Rightarrow \frac{1}{a} + \frac{1}{b} = 2$$

$$24. \text{ At } \theta = \frac{\pi}{4},$$

$$= 2 \cos^3 \frac{\pi}{4} = \frac{1}{\sqrt{2}} \text{ and } = 3 \sin^3 \frac{\pi}{4} = \frac{3}{2\sqrt{2}}$$

$$= 2 \cos^3 \theta \text{ and } = 3 \sin^3 \theta$$

$$\therefore \frac{d}{d\theta} = -6 \cos^2 \theta \sin \theta \text{ and } \frac{d}{d\theta} = 9 \sin^2 \theta \cos \theta$$



$$\therefore \frac{d}{d} = \frac{d\theta}{d} = -\frac{3}{2} \tan \theta$$

$$\therefore \left(\frac{d}{d}\right)_{\left(\theta=\frac{\pi}{4}\right)} = -\frac{3}{2}$$

\therefore equation of the tangent at $\left(\frac{1}{\sqrt{2}}, \frac{3}{2\sqrt{2}}\right)$ is

$$-\frac{3}{2\sqrt{2}} = -\frac{3}{2} \left(-\frac{1}{\sqrt{2}}\right)$$

$$\Rightarrow 3 + 2 = 3\sqrt{2}$$

25. At $t = 0$, $y = e^0 + 0 = 1$
 $\quad \quad \quad = e^2 + 2$

$$\therefore \frac{d}{d} = 2e^2 + 2$$

$$\therefore \left(\frac{d}{d}\right)_{(0,1)} = 2$$

Also, $-\left(\frac{d}{d}\right)_{(0,1)} = -\frac{1}{2}$

Equation of normal at (0, 1) is

$$(y - 1) = \frac{-1}{2}(x - 0)$$

$$\Rightarrow 2y - 2 = -x \Rightarrow x + 2y - 2 = 0$$

\therefore distance between origin and normal

$$= \frac{|0 + 0 - 2|}{\sqrt{1 + 4}} = \frac{2}{\sqrt{5}}$$

26. $x^2 + y^2 - 13 = 0$

$$\therefore 2x + 2 \frac{dy}{dx} = 0$$

$$\Rightarrow \frac{dy}{dx} = -x$$

$$\therefore \text{Slope of tangent at } (2, 3) = \left(\frac{dy}{dx}\right)_{(2,3)}$$

$$\Rightarrow m = -\frac{2}{3}$$

Given equation of circle is $x^2 + y^2 = 13$

\therefore Centre of circle $O = (0, 0)$, radius = $\sqrt{13}$ units

$$\text{Given point } M \left(m, \frac{-1}{m}\right) = \left(\frac{-2}{3}, \frac{3}{2}\right)$$

$$= (-0.67, 1.5)$$

$OM < \text{radius}$

\therefore The point lies inside the circle

27. $y = a(\sin \theta - \theta \cos \theta)$, $x = a(\cos \theta + \theta \sin \theta)$

$$\therefore \frac{dy}{dx} = \frac{a(\cos \theta - \cos \theta + \theta \sin \theta)}{a(\cos \theta + \theta \sin \theta)} = \frac{\theta \sin \theta}{\cos \theta + \theta \sin \theta}$$

and $\frac{dy}{d\theta} = a(-\sin \theta + \sin \theta + \theta \cos \theta)$

$$= a \theta \cos \theta$$

$$\therefore \frac{dy}{dx} = \frac{d/d\theta}{d/d\theta} = \frac{a\theta \sin \theta}{a\theta \cos \theta} = \tan \theta$$

$$\therefore \text{Slope of the normal} = \frac{-1}{\tan \theta} = -\cot \theta$$

\therefore Equation of the normal is

$$-a \sin \theta + a \theta \cos \theta$$

$$= -\frac{\cos \theta}{\sin \theta} (-a \cos \theta - a \theta \sin \theta)$$

$$\Rightarrow \sin \theta - a \sin^2 \theta + a \theta \sin \theta \cos \theta$$

$$= -\cos \theta + a \cos^2 \theta + a \theta \sin \theta \cos \theta$$

$$\Rightarrow \cos \theta + \sin \theta = a(\sin^2 \theta + \cos^2 \theta)$$

$$\Rightarrow \cos \theta + \sin \theta = a$$

$$\therefore \text{Distance from origin} = \frac{|-a|}{\sqrt{\sin^2 \theta + \cos^2 \theta}} = a = \text{constant}$$

28. $y = x^2 - x + 1$

$$\Rightarrow \frac{dy}{dx} = 2x - 1$$

$$\left(\frac{dy}{dx}\right)_{(=0)} = -1, \left(\frac{dy}{dx}\right)_{(=-1)} = -3, \left(\frac{dy}{dx}\right)_{\left(=\frac{5}{2}\right)} = 4$$

equation of normal at (0, 1) and having slope 1

$$\text{is } y - 1 = -1(x - 0)$$

$$\Rightarrow -x + 1 = 0 \quad \dots(i)$$

Equation of normal at (-1, 3) and having slope

$$\frac{1}{3} \text{ is}$$

$$-3 = \frac{1}{3}(x + 1)$$

$$\Rightarrow -3x + 10 = 0 \quad \dots(ii)$$

Equation of normal at $\left(\frac{5}{2}, \frac{19}{4}\right)$ and having

$$\text{slope } \frac{-1}{4} \text{ is}$$

$$-\frac{19}{4} = \frac{-1}{4} \left(x - \frac{5}{2}\right) \Rightarrow 4x - 19 = -x + \frac{5}{2}$$

$$\Rightarrow 2x + 8 - 43 = 0 \quad \dots(iii)$$

Equation (i), (ii) and (iii) are passes through

$$\text{point } \left(\frac{7}{2}, \frac{9}{2}\right).$$

\therefore they are concurrent



29. Given, $x^2 + 2x - 3 = 0$ (i)

Differentiating w.r.t. x , we get

$$2x + 2 \left(\frac{d}{dx} \right) - 6 \frac{d}{dx} = 0$$

$$\Rightarrow \frac{d}{dx} = \frac{-2}{3-2} \Rightarrow \left(\frac{d}{dx} \right)_{(1,1)} = 1$$

\therefore equation of the normal at $(1, 1)$ is

$$y - 1 = -1(x - 1)$$

$$\Rightarrow y = 2 - x$$

Putting $y = 2 - x$ in (i), we get

$$x^2 + 2(2 - x) - 3(2 - x)^2 = 0$$

$$\Rightarrow x^2 - 4x + 3 = 0$$

$$\Rightarrow x = 1, 3$$

\therefore the points of intersection are $(1, 1)$ and $(3, -1)$.

\therefore the normal at $(1, 1)$ meets the curve again at $(3, -1)$ which lies in the fourth quadrant.

30. $\sqrt{x} + \sqrt{y} = \sqrt{a}$

$$\therefore \frac{1}{2\sqrt{x}} + \frac{1}{2\sqrt{y}} \cdot \frac{dy}{dx} = 0$$

$$\therefore \frac{dy}{dx} = -\frac{\sqrt{y}}{\sqrt{x}}$$

\therefore Equation of the tangent at (x, y) is

$$Y - y = -\frac{\sqrt{y}}{\sqrt{x}}(X - x)$$

$$\therefore X\sqrt{y} + Y\sqrt{x} = \sqrt{x}(\sqrt{y} + \sqrt{x})$$

$$\therefore X\sqrt{y} + Y\sqrt{x} = \sqrt{x} \cdot \sqrt{a}$$

$$\therefore \frac{X}{\sqrt{a}\sqrt{x}} + \frac{Y}{\sqrt{a}\sqrt{y}} = 1$$

Clearly its intercepts on the axes are $\sqrt{a}\sqrt{x}$ and $\sqrt{a}\sqrt{y}$.

Sum of the intercepts

$$= \sqrt{a}(\sqrt{x} + \sqrt{y}) = \sqrt{a} \cdot \sqrt{a} = a$$

31. Let the coordinates of P be (x, y) .

$$\therefore \frac{x^2}{1} + \frac{y^2}{3} = a^2 \quad \dots(i)$$

$$\text{Now, } \frac{x^2}{3} + \frac{y^2}{3} = a^2$$

Differentiating w.r.t. x , we get

$$\frac{2}{3} \cdot \frac{1}{3} + \frac{2}{3} \cdot \frac{1}{3} \cdot \frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = -\frac{1}{3}$$

$$\therefore \left(\frac{dy}{dx} \right)_{(1, 1)} = -\frac{1}{3}$$

\therefore equation of the tangent at $(1, 1)$ is

$$y - 1 = -\frac{1}{3}(x - 1)$$

$$\Rightarrow \frac{y-1}{\frac{1}{3}} = -\frac{x-1}{\frac{1}{3}}$$

$$\Rightarrow \frac{y-1}{\frac{1}{3}} + \frac{x-1}{\frac{1}{3}} = \frac{2}{3} + \frac{2}{3}$$

$$\Rightarrow \frac{y-1}{\frac{1}{3}} + \frac{x-1}{\frac{1}{3}} = a^2 \quad \dots[\text{From (i)}]$$

This tangent meets the coordinate axes at

$$A\left(\frac{2}{a^2}, \frac{1}{3}, 0\right) \text{ and } B\left(0, a^2, \frac{1}{3}\right).$$

$$\therefore AB = \sqrt{a^2 \frac{4}{1} + a^2 \frac{4}{1}} = \sqrt{a^2 \left(\frac{2}{1} + \frac{2}{1} \right)}$$

$$= \sqrt{a^2 \cdot a^2} \quad \dots[\text{From (i)}]$$

$$= a$$

32. $y = x^2 - 5x + 6$

$$\therefore \frac{dy}{dx} = 2x - 5$$

$$\therefore \left(\frac{dy}{dx} \right)_{(2,0)} = 2(2) - 5 = -1 = m_1 \text{ (say)}$$

$$\text{and } \left(\frac{dy}{dx} \right)_{(3,0)} = 2(3) - 5 = 1 = m_2 \text{ (say)}$$

Here, $m_1 m_2 = -1$

\therefore the required angle is $\frac{\pi}{2}$.

33. If $\sin \theta = \cos \phi$, then $\theta = \frac{\pi}{4}$

Now, $\theta = \sin \phi$

$$\therefore \frac{d\theta}{d\phi} = \cos \phi$$

$$\therefore \left(\frac{d\theta}{d\phi} \right)_{\left(\frac{\pi}{4}\right)} = \frac{1}{\sqrt{2}} = m_1 \text{ (say)}$$

Also, $\theta = \cos \phi$

$$\therefore \frac{d\theta}{d\phi} = -\sin \phi$$

$$\therefore \left(\frac{d\theta}{d\phi} \right)_{\left(\frac{\pi}{4}\right)} = -\frac{1}{\sqrt{2}} = m_2 \text{ (say)}$$



∴ angle between the curves is

$$\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right| = \left| \frac{\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}}{1 + \left(\frac{1}{\sqrt{2}}\right)\left(-\frac{1}{\sqrt{2}}\right)} \right|$$

$$\Rightarrow \tan \theta = 2\sqrt{2}$$

$$\Rightarrow \theta = \tan^{-1}(2\sqrt{2})$$

34. $y = e^2$ (i)

$y = e^2 \sin$ (ii)

From (i) and (ii), we get

$$e^2 = e^2 \sin$$

∴ $\sin = 1 \Rightarrow = \frac{\pi}{2}$

Slope of tangent to (i) at $= \frac{\pi}{2}$ is given by

$$\left(\frac{d}{d} \right)_{=\frac{\pi}{2}} = \left[2 e^2 \right]_{=\frac{\pi}{2}} = \pi e^{\frac{\pi^2}{4}}$$

Slope of tangent to (ii) at $= \frac{\pi}{2}$ is given by

$$\left(\frac{d}{d} \right)_{=\frac{\pi}{2}} = \left[2 e^2 \sin + e^2 \cos \right]_{=\frac{\pi}{2}} = \pi e^{\frac{\pi^2}{4}}$$

Since both tangents have equal slopes, the angle between them is zero.

35. Let the given curves intersect each other at P(1, 1).
 $2^2 = 6$

Differentiating w.r.t. , we get

$$2 \frac{d}{d} = 6 \Rightarrow \left(\frac{d}{d} \right)_P = \frac{3}{1}$$

$$9^2 + b^2 = 16$$

Differentiating w.r.t. , we get

$$18 + 2b \frac{d}{d} = 0$$

$$\Rightarrow \left(\frac{d}{d} \right)_P = -\frac{9}{b_1}$$

Since, the given curves intersect each other at right angles.

$$\Rightarrow \left(\frac{3}{1} \right) \left(\frac{-9}{b_1} \right) = -1$$

$$\Rightarrow \frac{27}{b_1^2} = 1$$

$$\Rightarrow b = \frac{9}{2} \quad \dots \left[1^2 = 6 \right]$$

36. Acceleration, $\frac{dv}{dt} = 2t$, then acceleration after 3 second = $2 \times 3 = 6 \text{ cm / sec}^2$.

37. Motion of a particle $s = 15t - 2t^2$

∴ velocity = $\frac{ds}{dt} = 15 - 4t$

$$\Rightarrow \left(\frac{ds}{dt} \right)_{t=0} = 15 \text{ and } \left(\frac{ds}{dt} \right)_{t=3} = 3$$

∴ average velocity = $\frac{15+3}{2} = 9 \text{ units}$

38. Velocity, $v^2 = 2 - 3$

Differentiating both sides w.r.t.t, we get

$$2v \frac{dv}{dt} = -3 \frac{d}{dt}$$

$$\Rightarrow 2v \frac{dv}{dt} = -3v$$

$$\Rightarrow \frac{dv}{dt} = -\frac{3}{2}$$

Hence, the acceleration is uniform.

39. $y = At^2 + Bt + C$

∴ $v = 2At + B \Rightarrow v^2 = 4A^2t^2 + 4ABt + B^2$

and $4A = 4A^2t^2 + 4ABt + 4AC$

$$\Rightarrow v^2 - 4A = B^2 - 4AC$$

$$\Rightarrow 4A - v^2 = 4AC - B^2$$

40. $t = \frac{v^2}{2} \Rightarrow v^2 = 2t$

Differentiating both sides w.r.t.t., we get

$$2v \frac{dv}{dt} = 2$$

$$\Rightarrow \frac{dv}{dt} = \frac{1}{v} = f$$

$$\Rightarrow \frac{df}{dt} = -\frac{1}{v^2} \cdot \frac{dv}{dt} = -\frac{1}{v^2} \times \frac{1}{v}$$

$$\Rightarrow -\frac{df}{dt} = \frac{1}{v^3} = f^3$$

41. $\frac{d^2t}{d^2} = \frac{d}{d} \left(\frac{dt}{d} \right) = \frac{d}{d} \left(\frac{1}{v} \right) = -\frac{1}{v^2} \cdot \frac{dv}{d}$

Since, $v \frac{dv}{d} = f \Rightarrow \frac{dv}{d} = \frac{f}{v}$

∴ $\frac{d^2t}{d^2} = -\frac{1}{v^2} \cdot \frac{f}{v} \Rightarrow -v^3 \frac{d^2t}{d^2} = f$




42. $s = \frac{1}{2}vt \Rightarrow 2s = vt \Rightarrow 2 \frac{ds}{dt} = v + t \cdot \frac{dv}{dt}$
 $\Rightarrow 2 \frac{d^2s}{dt^2} = \frac{dv}{dt} + t \cdot \frac{d^2v}{dt^2} + \frac{dv}{dt}$
 But $\frac{dv}{dt} = \text{acceleration (a)}$
 $\Rightarrow 2a = a + t \cdot \frac{da}{dt} + a \Rightarrow \frac{da}{dt} = 0$ or $t = 0$
 But $t = 0$ is impossible
 $\therefore \frac{da}{dt} = 0$ i.e., a is constant.

44. $s = 6 + 48t - t^3$
 $\therefore v = \frac{ds}{dt} = 0 + 48 - 3t^2$
 When direction of motion reverses, $v = 0$
 $\Rightarrow 48 - 3t^2 = 0 \Rightarrow t = -4, 4$
 $\therefore (s)_4 = 6 + 192 - 64 = 134$

45. $a + bv^2 = c$
 Differentiating both sides w.r.t.t, we get
 $0 + b \left(2v \cdot \frac{dv}{dt} \right) = 2 \frac{d}{dt} c$
 $\Rightarrow v \cdot b \frac{dv}{dt} = \frac{d}{dt} c \Rightarrow \frac{dv}{dt} = \frac{d}{dt} c \cdot \left[\because \frac{d}{dt} c = v \right]$


46. $\frac{d}{dt} = 1.2$.
 From the figure,
 $\frac{2}{3} \Rightarrow \frac{d}{dt} = \frac{2}{3} \cdot \frac{d}{dt}$
 \therefore Required rate of length of shadow
 $= \frac{d}{dt} = 0.8 \text{ m/s}$



47. From the figure, $x^2 + y^2 = 100$ (i)
 $\Rightarrow 2 \frac{dx}{dt} + 2 \frac{dy}{dt} = 0$ (ii)

From (i) and (ii),
 $\frac{dx}{dt} = -\frac{16}{6} = -\frac{8}{3} \text{ cm/sec}$

The rate at which the end B is moving is $\frac{8}{3} \text{ cm/sec}$.



48. According to the figure, $x^2 + y^2 = 25$ (i)
 Differentiate (i) w.r.t. t, we get

$$2 \frac{dx}{dt} + 2 \frac{dy}{dt} = 0$$


Here $x = 4$ and $\frac{dy}{dt} = 1.5$

From (i), $4^2 + y^2 = 25 \Rightarrow y = 3$

From (ii), $2(4)\left(\frac{dx}{dt}\right) + 2(3)(1.5) = 0$

$$\Rightarrow \frac{dx}{dt} = -2 \text{ m/sec}$$

Hence, length of the highest point decreases at the rate of 2 m/sec.



49. According to the figure, $x^2 + y^2 = 400$ (i)
 Differentiate (i) w.r.t. t, we get

$$2 \frac{dx}{dt} + 2 \frac{dy}{dt} = 0$$

Here $x = 12$ and $\frac{dy}{dt} = 2$

From (i), $12^2 + y^2 = 400$

$$\Rightarrow y = 16$$

From (ii), $2(12)\left(\frac{dx}{dt}\right) + 2(16)(2) = 0$

$$\Rightarrow \frac{dx}{dt} = -\frac{8}{3}$$

50. Surface area, $S = 4\pi r^2$ and $\frac{dr}{dt} = 2$

$$\therefore \frac{dS}{dt} = 4\pi \times 2r \frac{dr}{dt} = 8\pi r \times 2 = 16\pi r$$

$$\Rightarrow \frac{dS}{dt} \propto r$$

51. Given the rate of increasing the radius

$$= \frac{dr}{dt} = 3.5 \text{ cm/sec and } r = 10 \text{ cm}$$

$$\text{Area of circle} = A = \pi r^2$$

$$\therefore \frac{dA}{dt} = 2\pi r \cdot \frac{dr}{dt}$$

$$\Rightarrow \frac{dA}{dt} = 2\pi \times 10 \times 3.5 \Rightarrow \frac{dA}{dt} = 220 \text{ cm}^2/\text{sec}$$

52. If s is the length of each side of an equilateral triangle and A is its area, then

$$A = \frac{\sqrt{3}}{4} s^2 \Rightarrow \frac{dA}{dt} = \frac{\sqrt{3}}{4} \cdot 2 \frac{ds}{dt}$$

Here, $s = 10 \text{ cm}$ and $\frac{ds}{dt} = 2 \text{ cm/sec}$

$$\therefore \frac{dA}{dt} = 10\sqrt{3} \text{ sq. unit/sec}$$



$$53. \quad A_1 = 2^2, \text{ and } A_2 = 2^2 \\ \Rightarrow \frac{dA_1}{dt} = 2 \frac{d}{dt}, \text{ and } \frac{dA_2}{dt} = 2 \frac{d}{dt}$$

$$\therefore \frac{dA_2}{dA_1} = \frac{\frac{dA_2}{dt}}{\frac{dA_1}{dt}} = \frac{2 \frac{d}{dt}}{2 \frac{d}{dt}} = - \left(\frac{d}{d} \right)$$

$$\text{Given, } = + 2^2$$

$$\therefore \frac{d}{d} = 1 + 2$$

$$\therefore \frac{dA_2}{dA_1} = - (1 + 2)$$

$$= - \frac{+ 2^2}{(1 + 2)}$$

$$= (1 + 2)(1 + 2) = 2^2 + 3 + 1$$

$$54. \quad h = 6 \text{ m, } r = 4 \text{ m} = \frac{2}{3} h$$

$$V = \frac{1}{3} \pi r^2 h$$

$$\Rightarrow V = \frac{1}{3} \times \frac{4}{9} \times \pi h^3$$

$$\Rightarrow \frac{dV}{dt} = \frac{4}{9} \pi h^2 \frac{dh}{dt}$$

$$\text{But } \frac{dV}{dt} = 3 \text{ m}^3/\text{min} \text{ and } h = 3 \text{ m}$$

$$\therefore 3 = \frac{4}{9} \pi \times 9 \times \frac{dh}{dt}$$

$$\Rightarrow \frac{dh}{dt} = \frac{3}{4\pi} \text{ m/min}$$

$$55. \quad V = \frac{4}{3} \pi r^3$$

$$\therefore \frac{dV}{dt} = 4\pi r^2 \cdot \frac{dr}{dt} \Rightarrow \frac{dr}{dt} = \frac{1}{4\pi r^2} \cdot \frac{dV}{dt}$$

$$\Rightarrow \frac{dr}{dt} = \frac{1}{4 \times \pi \times 15 \times 15} \times 900$$

$$\Rightarrow \frac{dr}{dt} = \frac{1}{\pi} = \frac{7}{22}$$

$$56. \quad \text{Here, } V = \frac{4}{3} \pi r^3 \text{ and } S = 4\pi r^2$$

$$\Rightarrow \frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt} \Rightarrow \frac{dr}{dt} = \frac{40}{4\pi r^2} = \frac{5}{32\pi}$$

$$\therefore \frac{dS}{dt} = 8\pi r \frac{dr}{dt}$$

$$= 8\pi \times 8 \times \frac{5}{32\pi} = 10 \text{ cm}^2/\text{min}$$

$$57. \quad V = \frac{4}{3} \pi r^3$$

$$\Rightarrow 288 \pi = \frac{4}{3} \pi r^3$$

$$\Rightarrow r = 6 \text{ cm}$$

$$V = \frac{4}{3} \pi r^3$$

$$\therefore \frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

$$\Rightarrow 4\pi = 4\pi r^2 \frac{dr}{dt}$$

$$\Rightarrow \frac{dr}{dt} = \frac{1}{r^2}$$

$$\text{Now, } A = 4\pi r^2$$

$$\therefore \frac{dA}{dt} = 8\pi r \frac{dr}{dt}$$

$$= 8\pi r \times \frac{1}{r^2}$$

$$= \frac{8\pi}{r} = \frac{8\pi}{6} = \frac{4\pi}{3} \text{ cm}^2/\text{sec}$$

$$58. \quad \text{Volume} = V = \frac{4}{3} \pi r^3$$

$$\Rightarrow \frac{dV}{dt} = 4\pi r^2 \cdot \frac{dr}{dt}, \text{ at } r = 7 \text{ cm}$$

$$\Rightarrow 35 = 4\pi(7)^2 \frac{dr}{dt} \Rightarrow \frac{dr}{dt} = \frac{5}{28\pi}$$

$$\text{Surface area, } S = 4\pi r^2$$

$$\therefore \frac{dS}{dt} = 8\pi r \frac{dr}{dt} = 8\pi(7) \left(\frac{5}{28\pi} \right) = 10 \text{ cm}^2/\text{min}$$

$$59. \quad V = \frac{4}{3} \pi r^3$$

$$\therefore \frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt} \quad \dots(i)$$

$$\text{After 49 min, } (4500 - 49 \times 72)\pi = 972 \pi \text{ m}^3$$

$$\therefore 972 \pi = \frac{4}{3} \pi r^3$$

$$\therefore r^3 = 3 \times 243 = 3 \times 3^5$$

$$\therefore r = 9$$

$$\text{Given, } \frac{dV}{dt} = 72\pi$$

$$\therefore 72\pi = 4\pi \times 9 \times 9 \left(\frac{dr}{dt} \right) \quad \dots[\text{From (i)}]$$

$$\therefore \frac{dr}{dt} = \frac{2}{9}$$



60. Volume of sphere (V) = $\frac{4}{3}\pi r^3$
 Surface area of sphere (A) = $4\pi r^2$
 $\frac{dV}{dr} = 4\pi r^2$ and $\frac{dA}{dr} = 8\pi r$
 $\therefore \left(\frac{dV}{dA}\right) = \left(\frac{\frac{dV}{dr}}{\frac{dA}{dr}}\right) = \frac{4\pi r^2}{8\pi r} = \frac{r}{2}$
 $\therefore \left(\frac{dV}{dA}\right)_{r=4} = \frac{4}{2} = 2 \text{ cm}^3/\text{cm}^2$
61. $W = nw$, $n = 2t^2 + 3$ and $w = t^2 - t + 2$
 $\therefore \frac{dW}{dt} = w \frac{dn}{dt} + n \frac{dw}{dt}$, $\frac{dW}{dt} = 4t$, $\frac{dW}{dt} = 2t - 1$
 At $t = 1$,
 $n = 5$, $w = 2$, $\frac{dn}{dt} = 4$, $\frac{dW}{dt} = 1$
 $\therefore \left(\frac{dW}{dt}\right)_{(t=1)} = 2(4) + 5(1) = 13$
62. According to the given condition,
 $\frac{d}{dt} = 8 \frac{d}{dt}$ (i)
 Given, $6 = 3 + 2$ (ii)
 $\Rightarrow 6 \left(\frac{d}{d}\right) = 3^2 \frac{d}{dt}$
 $\Rightarrow 6 \left(\frac{8d}{dt}\right) = 3^2 \frac{d}{dt}$ [From (i)]
 $\Rightarrow 3^2 = 48 \Rightarrow 2 = 16 \Rightarrow = \pm 4$
 Putting $= 4$ in (ii), we get
 $6 = (4)^3 + 2 = 64 + 2$
 $= 11$
 Putting $= -4$ in (ii), we get
 $= -64 + 2$
 $= \frac{-62}{6} = \frac{-31}{3}$
 \therefore the required points on the curve are (4, 11) and $\left(-4, \frac{-31}{3}\right)$.
63. $f(x) = x^3 + 5x^2 - 7x + 9$
 $\therefore f'(x) = 3x^2 + 10x - 7$
 Here, $a = 1$ and $h = 0.1$
 $\therefore f(a) = f(1) = 1^3 + 5(1)^2 - 7(1) + 9 = 8$
 and $f'(a) = f'(1) = 3(1)^2 + 10(1) - 7 = 6$
 $\therefore f(a+h) \approx f(a) + hf'(a)$
 $\approx 8 + 0.1(6)$
 $\approx 8 + 0.6 \approx 8.6$

64. Let $f(x) = \sqrt[3]{x} = x^{\frac{1}{3}}$
 $\therefore f'(x) = \frac{1}{3} x^{-\frac{2}{3}} = \frac{1}{3x^{\frac{2}{3}}}$
 Here, $a = -1$, and $h = 0.01$
 $f(a+h) \approx f(a) + hf'(a)$
 $\approx (-1)^{\frac{1}{3}} + 0.01 \times \frac{1}{3(-1)^{\frac{2}{3}}}$
 $\approx -1 + 0.0033$
 ≈ -0.9967
65. Let $f(x) = \sqrt[5]{x} = x^{1/5}$
 $\therefore f'(x) = \frac{1}{5} x^{-4/5} = \frac{1}{5x^{4/5}}$
 Here, $a = 243$ and $h = -0.001$
 $f(a+h) \approx f(a) + hf'(a)$
 $= (243)^{1/5} - 0.001 \times \frac{1}{5(243)^{4/5}}$
 $= 3 - \frac{0.001}{5 \times 81}$
 $= 3 - \frac{1}{405000}$
 $\therefore f(242.999) = \frac{1214999}{405000}$
66. Let $f(x) = \cos x$
 $\therefore f'(x) = -\sin x$
 Here, $a = 30^\circ$ and $h = 1^\circ = 0.0174$
 $\therefore f(a+h) \approx f(a) + hf'(a)$
 $\approx \frac{\sqrt{3}}{2} + 0.0174 \left(\frac{-1}{2}\right)$
 $\approx \frac{1.73}{2} - \frac{0.0174}{2}$
 ≈ 0.8563
67. $f(x) = e^x (\sin x - \cos x)$
 $\therefore f'(x) = e^x (\sin x - \cos x) + e^x (\cos x + \sin x)$
 $\therefore f'(x) = 2e^x \sin x$
 Now, $f'(c) = 0$
 $\Rightarrow 2e^c \sin c = 0$
 $\Rightarrow \sin c = 0 = \sin \pi$
 $\Rightarrow c = \pi$
68. Here, $f\left(\frac{\pi}{2}\right) = e^0 = 1$ and $f\left(\frac{3\pi}{2}\right) = e^0 = 1$
 $\therefore f\left(\frac{\pi}{2}\right) = f\left(\frac{3\pi}{2}\right)$
 \therefore Third condition of Rolle's theorem is satisfied by option (A) only.



69. (A) $f(x) = |x|$ is not differentiable at $x = 0$.
 (B) $f(x) = \tan x$ is discontinuous at $x = \frac{\pi}{2}$.
 (C) $f(x) = 1 + (x-2)^{\frac{2}{3}}$ is not differentiable at $x = 2$.
 (D) $f(x) = (x-2)^2$ is a polynomial function.
 $\therefore f(x)$ is continuous on $[0, 2]$ and differentiable on $(0, 2)$.
 Also, $f(0) = f(2)$
 Hence, Rolle's theorem is applicable.

70. $f(x) = e^x$
 $\therefore f(0) = e^0 = 1, f(1) = e$ and $f'(x) = e^x$
 By mean value theorem,

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$\Rightarrow f'(c) = \frac{e^b - e^a}{b - a}$$

$$\Rightarrow e^c = \frac{e - 1}{1 - 0}$$

$$\Rightarrow c = \log(e - 1)$$

71. $f(x) = x^2$
 $f(2) = 4, f(4) = 16$
 $f'(x) = 2x$
 \therefore By Lagrange's mean value theorem,

$$f'(c) = \frac{f(4) - f(2)}{4 - 2}$$

$$\Rightarrow 2c = \frac{16 - 4}{2}$$

$$\Rightarrow c = 3$$

72. $f(x) = \sqrt{x}$
 $\therefore f(a) = f(4) = \sqrt{4} = 2, f(b) = f(9) = \sqrt{9} = 3$ and
 $f'(x) = \frac{1}{2\sqrt{x}}$
 Given, $f'(c) = \frac{f(b) - f(a)}{b - a} = \frac{3 - 2}{9 - 4} = \frac{1}{5}$
 $\therefore \frac{1}{2\sqrt{c}} = \frac{1}{5} \Rightarrow c = \frac{25}{4} = 6.25$

73. $f(x) = (x-1)(x-2)$
 $\Rightarrow f(x) = x^2 - 3x + 2$
 $f(0) = 2$
 $f\left(\frac{1}{2}\right) = \frac{3}{4}$
 $f'(x) = 2x - 3$
 By Lagrange's mean value theorem,

$$f'(c) = \frac{f\left(\frac{1}{2}\right) - f(0)}{\frac{1}{2} - 0}$$

$$\Rightarrow 2c - 3 = \frac{\frac{3}{4} - 2}{\frac{1}{2}}$$

$$\Rightarrow 2c = \frac{-5}{2} + 3 \Rightarrow c = \frac{1}{4}$$

74. $f(x) = \frac{2x + 3}{4x - 1}$
 $f(1) = \frac{5}{3}, f(2) = 1$
 $f'(x) = \frac{(4x - 1)(2) - (2x + 3)(4)}{(4x - 1)^2} = \frac{-14}{(4x - 1)^2}$
 \therefore By Lagrange's mean value theorem,

$$f'(c) = \frac{f(2) - f(1)}{2 - 1}$$

$$\Rightarrow \frac{-14}{(4c - 1)^2} = \frac{1 - \frac{5}{3}}{1}$$

$$\Rightarrow (4c - 1)^2 = \frac{-14}{\frac{-2}{3}}$$

$$\Rightarrow 16c^2 - 8c + 1 = 21$$

$$\Rightarrow 4c^2 - 2c - 5 = 0$$

$$\Rightarrow c = \frac{1 + \sqrt{21}}{4}$$

75. $f(x) = \cos x$
 $\therefore f(0) = 1, f\left(\frac{\pi}{2}\right) = 0$ and $f'(x) = -\sin x$
 By mean value theorem,

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$\Rightarrow -\sin c = \frac{f\left(\frac{\pi}{2}\right) - f(0)}{\frac{\pi}{2} - 0}$$

$$\Rightarrow -\sin c = \frac{0 - 1}{\frac{\pi}{2}} = -\frac{2}{\pi}$$

$$\Rightarrow \sin c = \frac{2}{\pi} \Rightarrow c = \sin^{-1}\left(\frac{2}{\pi}\right)$$




76. $g(x) = \frac{f(x)}{x+1}$
 $\therefore g(0) = \frac{f(0)}{0+1} = 12$ and $f(6) = \frac{f(6)}{6+1} = \frac{-4}{7}$
 By mean value theorem,
 $g'(c) = \frac{g(6) - g(0)}{6 - 0}$
 $= \frac{\frac{-4}{7} - 12}{6}$
 $= \frac{-4 - 84}{7 \times 6} = -\frac{44}{21}$

77. Consider option (A),
 $Lf'\left(\frac{1}{2}\right) = -1$ and $Rf'\left(\frac{1}{2}\right) = 0$
 So, it is not differentiable at $x = \frac{1}{2} \in (0, 1)$.
 Hence, Lagrange's mean value theorem is not applicable.

78. $f(x) = x^3 = f(x)$
 $\therefore f(2) = 8, f(-2) = -8$ and $f'(x) = 3x^2$
 By mean value theorem,
 $f'(c) = \frac{f(2) - f(-2)}{2 - (-2)}$
 $\Rightarrow 3c^2 = \frac{8 - (-8)}{4}$
 $\Rightarrow c^2 = \frac{4}{3}$
 $\Rightarrow c = \pm \frac{2}{\sqrt{3}}$

79. $f(b) = f(2) = 8 - 24a + 10 = 18 - 24a$
 $f(a) = f(1) = 1 - 6a + 5 = 6 - 6a$
 $f'(x) = 3x^2 - 12a + 5$
 By Lagrange's mean value theorem,
 $f'(c) = \frac{f(b) - f(a)}{b - a} = \frac{18 - 24a - 6 + 6a}{2 - 1}$
 $\therefore f'(c) = 12 - 18a$
 $\therefore 3c^2 - 12a + 5 = 12 - 18a$
 At $c = \frac{7}{4}$,
 $3\left(\frac{49}{16}\right) - 12a\left(\frac{7}{4}\right) + 5 = 12 - 18a$
 $\Rightarrow 3a = \frac{147}{16} - 7 \Rightarrow 3a = \frac{35}{16} \Rightarrow a = \frac{35}{48}$

80. Since, $f(x) = x^3 \Rightarrow f'(x) = 3x^2$, which is non-negative for all real values of x .
 \therefore Option (C) is the correct answer.
81. $f(x) = ax + b \Rightarrow f'(x) = a$
 \therefore For strictly increasing, $f'(x) > 0$
 $\Rightarrow a > 0$ for all real x .
82. $f(x) = (x - 1)^2 - 1$. Hence decreasing in $x < 1$.



Alternate Method:

$f'(x) = 2(x - 1) = 2(x - 1)$
 To be decreasing, $2(x - 1) < 0 \Rightarrow (x - 1) < 0$
 $\Rightarrow x < 1$

83. $\frac{d}{dx} \tan^{-1} x = \frac{1}{1+x^2} \Rightarrow \frac{d}{dx} \tan^{-1} x = \frac{1}{1+x^2} \geq 0$

84. $f(x) = \begin{cases} 0 & , x \leq 0 \\ -3 & , x > 0 \end{cases}$

$\therefore f'(x) = \begin{cases} 0 & , x \leq 0 \\ 1 & , x > 0 \end{cases}$

\therefore It is strictly increasing when $x > 0$.

85. $f(x) = \sin(x - 2p) < 0$
 $\Rightarrow f'(x) = -\sin(x - 2p) < 0$
 $\Rightarrow \frac{1}{2} \sin(x - 2p) + p > 0$
 $\Rightarrow p > \frac{1}{2} \dots [\because -1 \leq \sin(x - 2p) \leq 1]$

86. Function is monotonically decreasing, when $f'(x) < 0$
 $\Rightarrow 6x^2 - 18x + 12 < 0$
 $\Rightarrow x^2 - 3x + 2 < 0$
 $\Rightarrow (x - 2)(x - 1) < 0$
 $\Rightarrow 1 < x < 2$

87. $f(x) = x^2 + 2x - 5$
 $f'(x) = 2x + 2 = 2(x + 1)$
 For increasing function, $f'(x) > 0$
 $\Rightarrow 2(x + 1) > 0$
 $\Rightarrow x > -1$
 $\Rightarrow x \in (-1, \infty)$



88. $f(x) = x^3 - 3x^2 - 24x + 5$
 For $f(x)$ to be increasing, $f'(x) > 0$
 $\Rightarrow 3x^2 - 6x - 24 > 0$
 $\Rightarrow x^2 - 2x - 8 > 0$
 $\Rightarrow x^2 - 4x + 2x - 8 > 0$
 $\Rightarrow (x + 2)(x - 4) > 0$
 $\Rightarrow x \in (-\infty, -2) \cup (4, \infty)$

89. $f(x) = -2x^3 - 9x^2 - 12x + 1$
 $\Rightarrow f'(x) = -6x^2 - 18x - 12$
 For $f(x)$ to be decreasing, $f'(x) < 0$
 $\Rightarrow -6x^2 - 18x - 12 < 0$
 $\Rightarrow x^2 + 3x + 2 > 0 \Rightarrow (x + 2)(x + 1) > 0$
 $\Rightarrow x < -2$ or $x > -1$
 $\Rightarrow x \in (-1, \infty)$ or $(-\infty, -2)$

90. $f(x) = x + \sqrt{1-x}$
 $\therefore f'(x) = 1 - \frac{1}{2\sqrt{1-x}}$
 For $f(x)$ to be decreasing $f'(x) < 0$
 $\Rightarrow 1 - \frac{1}{2\sqrt{1-x}} < 0$
 $\Rightarrow 1 < \frac{1}{2\sqrt{1-x}}$
 $\Rightarrow 2\sqrt{1-x} < 1$
 $\Rightarrow 4(1-x) < 1$
 $\Rightarrow 1-x < \frac{1}{4}$
 $\Rightarrow \frac{3}{4} < x$
 $\therefore x \in \left(\frac{3}{4}, 1\right)$

91. $f(x) = \sin^4 x + \cos^4 x$
 $\therefore f'(x) = -\sin 4x$
 $\therefore f'(x) > 0$
 $\therefore -\sin 4x > 0$
 $\therefore \sin 4x < 0$
 $\therefore (2n+1)\pi < 4x < (2n+2)\pi$
 $\Rightarrow \frac{(2n+1)\pi}{4} < x < \frac{(n+1)\pi}{2}$
 For $n = 0, \frac{\pi}{4} < x < \frac{\pi}{2}$
 Now, $\frac{\pi}{2} = \frac{4\pi}{8} > \frac{3\pi}{8}$
 $\therefore f(x)$ is increasing in $\left(\frac{\pi}{4}, \frac{3\pi}{8}\right)$.

92. If $f(x) = (a+2)^3 - 3a^2 + 9a - 1$
 decreases monotonically for all $x \in \mathbb{R}$,
 then $f'(x) \leq 0$ for all $x \in \mathbb{R}$
 $\Rightarrow 3(a+2)^2 - 6a + 9a \leq 0$ for all $x \in \mathbb{R}$
 $\Rightarrow (a+2)^2 - 2a + 3a \leq 0$ for all $x \in \mathbb{R}$
 $\Rightarrow a+2 < 0$ and Discriminant ≤ 0
 $\Rightarrow a < -2, -8a^2 - 24a \leq 0$
 $\Rightarrow a < -2$ and $a(a+3) \geq 0$
 $\Rightarrow a < -2, a \leq -3$ or $a \geq 0$
 $\Rightarrow a \leq -3 \Rightarrow -\infty < a \leq -3$

93. $f(x) = \frac{1}{x^2+1}$
 $\therefore f'(x) = \frac{(x^2+1)(1) - (2x)}{(x^2+1)^2} = \frac{1-x^2}{(x^2+1)^2}$
 For $f(x)$ to be increasing
 $f'(x) > 0 \Rightarrow \frac{1-x^2}{(x^2+1)^2} > 0$
 $x^2+1 \neq 0 \Rightarrow x^2 \neq -1$
 $1-x^2 > 0$
 $\Rightarrow x^2 < 1$
 $\Rightarrow x \in (-1, 1)$

94. $f(x) = \log(1+x) - \frac{2}{2+x}$
 $\Rightarrow f'(x) = \frac{1}{1+x} - \frac{(2+x) \cdot (2) - 2(1)}{(2+x)^2}$
 $\Rightarrow f'(x) = \frac{x^2}{(x+1)(x+2)^2}$
 $\therefore f'(x) > 0$ for all $x > 0$
 Hence, $f(x)$ is increasing on $(0, \infty)$.

95. $f(x) = (x+2)e^{-x}$
 $\therefore f'(x) = e^{-x} - e^{-x}(x+2) = -e^{-x}(x+1)$
 For $f(x)$ to be increasing,
 $-e^{-x}(x+1) > 0 \Rightarrow e^{-x}(x+1) < 0$
 $\Rightarrow (x+1) < 0$
 $\Rightarrow x < -1$
 $\therefore x \in (-\infty, -1)$
 \therefore the function is increasing in $(-\infty, -1)$.
 \therefore For $f(x)$ to be decreasing, $-e^{-x}(x+1) < 0$
 $\Rightarrow e^{-x}(x+1) > 0$
 $\Rightarrow x+1 > 0$
 $\Rightarrow x > -1$
 $\Rightarrow x \in (-1, \infty)$
 \therefore the function is decreasing in $(-1, \infty)$.



$$96. f(x) = 3x^2 - 2x + 1, \Rightarrow f'(x) = 6x - 2 \geq 0 \Rightarrow x \geq \frac{1}{3}$$

Option (A) is incorrectly matched.

$$97. \text{ Let } f(x) = \frac{\ln(\pi + x)}{\ln(e + x)}$$

$$\begin{aligned} \therefore f'(x) &= \frac{\ln(e + x) \times \frac{1}{\pi + x} - \ln(\pi + x) \times \frac{1}{e + x}}{[\ln(e + x)]^2} \\ &= \frac{(e + x)\ln(e + x) - (\pi + x)\ln(\pi + x)}{[\ln(e + x)]^2 \times (e + x)(\pi + x)} \end{aligned}$$

$$\Rightarrow f'(x) < 0 \text{ for all } x \geq 0 \quad \dots [\because \pi > e]$$

$\therefore f(x)$ is decreasing on $[0, \infty)$.

$$98. f(x) = x^3 - 10x^2 + 200x - 10$$

$$\Rightarrow f'(x) = 3x^2 - 20x + 200$$

For $f(x)$ to be increasing $f'(x) > 0$

$$\Rightarrow 3x^2 - 20x + 200 > 0$$

$$\Rightarrow 3 \left(x^2 - \frac{20}{3}x + \frac{200}{3} + \frac{100}{9} - \frac{100}{9} \right) > 0$$

$$\Rightarrow 3 \left[\left(x - \frac{10}{3} \right)^2 + \frac{500}{9} \right] > 0$$

$$\Rightarrow 3 \left(x - \frac{10}{3} \right)^2 + \frac{500}{9} > 0$$

Always increasing throughout real line.

$$99. f(x) = \frac{3}{2}(3x - 10), \quad x \geq 0$$

$$\begin{aligned} \therefore f'(x) &= \frac{3}{2} \cdot \frac{1}{2}(3x - 10) + \frac{3}{2}(3) \\ &= \frac{15}{2} \cdot \frac{1}{2}(x - 2) \end{aligned}$$

For $f(x)$ to be increasing,

$$f'(x) \geq 0 \Rightarrow \frac{15}{2} \cdot \frac{1}{2}(x - 2) \geq 0$$

$$\Rightarrow x \geq 2 \Rightarrow x \in [2, \infty)$$

$$100. f(x) = \tan^{-1}(\sin x + \cos x)$$

$$\begin{aligned} \Rightarrow f'(x) &= \frac{1}{1 + (\sin x + \cos x)^2} \times (\cos x - \sin x) \\ &= \frac{\sqrt{2} \cos \left(x + \frac{\pi}{4} \right)}{1 + (\sin x + \cos x)^2} \end{aligned}$$

For $f(x)$ to be increasing, $f'(x) > 0$

$$\Rightarrow \sqrt{2} \cos \left(x + \frac{\pi}{4} \right) > 0$$

$$\Rightarrow \cos \left(x + \frac{\pi}{4} \right) > 0$$

$$\Rightarrow -\frac{\pi}{2} < x + \frac{\pi}{4} < \frac{\pi}{2}$$

$$\Rightarrow -\frac{3\pi}{4} < x < \frac{\pi}{4}$$

$\therefore f(x)$ is an increasing function in $\left(-\frac{\pi}{2}, \frac{\pi}{4} \right)$.

$$101. f(x) = \log(\sin x + \cos x)$$

$$\Rightarrow f'(x) = \frac{\cos x - \sin x}{\sin x + \cos x} = \frac{1 - \tan x}{1 + \tan x} = \tan \left(\frac{\pi}{4} - x \right)$$

For $f(x)$ to be increasing,

$$f'(x) > 0$$

$$\Rightarrow \tan \left(\frac{\pi}{4} - x \right) > 0$$

$$\Rightarrow 0 < \frac{\pi}{4} - x < \frac{\pi}{2}$$

$$\Rightarrow -\frac{\pi}{4} < -x < \frac{\pi}{4}$$

$$\Rightarrow -\frac{\pi}{4} < x < \frac{\pi}{4}$$

$$\Rightarrow x \in \left(-\frac{\pi}{4}, \frac{\pi}{4} \right)$$

$$102. f(x) = \int e^{-(x-1)}(x-2) dx$$

$$\Rightarrow f'(x) = e^{-(x-1)}(x-2)$$

For $f(x)$ to be decreasing, $f'(x) < 0$

$$\Rightarrow e^{-(x-1)}(x-2) < 0$$

$$\Rightarrow (x-1)(x-2) < 0$$

$$\Rightarrow x \in (1, 2)$$

$$103. f(x) = \frac{1}{\sin x}$$

$$\Rightarrow f'(x) = \frac{\sin x - \cos x}{\sin^2 x} = \frac{\cos x (\tan x - 1)}{\sin^2 x}$$

$\therefore f'(x) > 0$ for $0 < x \leq 1$

$\Rightarrow f(x)$ is an increasing function.

$$\text{Now, } g(x) = \frac{1}{\sin x}$$

$$\Rightarrow g'(x) = \frac{\tan x - \sec^2 x}{\tan^2 x}$$

$$= \frac{\sin x \cos x - 1}{\sin^2 x}$$

$$= \frac{\sin 2x - 2}{2\sin^2 x}$$

$\therefore g'(x) < 0$ for $0 < x \leq 1$.

$\Rightarrow g(x)$ is a decreasing function.



104. $f(x) = \sin x - \cos x$
 $\Rightarrow f'(x) = \cos x + \sin x = \sqrt{2} \left[\cos \left(x - \frac{\pi}{4} \right) \right]$
 For $f(x)$ to be decreasing, $f'(x) < 0$
 $\Rightarrow \sqrt{2} \cos \left(x - \frac{\pi}{4} \right) < 0$
 $\Rightarrow \cos \left(x - \frac{\pi}{4} \right) < 0$
 $\Rightarrow \frac{\pi}{2} < x - \frac{\pi}{4} < \frac{3\pi}{2} \Rightarrow \frac{3\pi}{4} < x < \frac{7\pi}{4}$
105. $h(x) = f(x) - (f(x))^2 + (f(x))^3$
 $\therefore h'(x) = f'(x) - 2f(x)f'(x) + 3(f(x))^2 f'(x)$
 $= f'(x) [1 - 2f(x) + 3(f(x))^2]$
 Here, $1 - 2f(x) + 3(f(x))^2 > 0$ for all $f(x)$
 $\Rightarrow h'(x) > 0$, if $f'(x) > 0$ and $h'(x) < 0$, if $f'(x) < 0$
 $\Rightarrow h$ is increasing whenever f is increasing and h is decreasing whenever f is decreasing.
106. $f(x) = [x(x-2)]^2$
 $\Rightarrow f(x) = x^2(x-2)^2$
 $\Rightarrow f'(x) = 2x(x-2) + x^2(2(x-2))$
 $= 2x(x-2) \{ 1 + (x-2) \}$
 $= 4x(x-2)(x-1)$
 For $f(x)$ to be increasing, $f'(x) > 0$
 $\Rightarrow 4x(x-1)(x-2) > 0$
 $\Rightarrow (x-1)(x-2) > 0 \Rightarrow x \in (0, 1) \cup (2, \infty)$
107. $y = \{x(x-3)\}^2$
 $\Rightarrow y = x^2(x-3)^2$
 $\therefore \frac{dy}{dx} = 2x(x-3)^2 + 2x(x-3) \cdot 2(x-3)$
 $= 2x(x-3)[x-3+2x]$
 $= 2x(x-3)(3x-3)$
 For y to be increasing, $\frac{dy}{dx} > 0$
 $\Rightarrow 2x(x-3)(3x-3) > 0$
 $\Rightarrow (x-3)(2x-3) > 0 \Rightarrow x \in \left(0, \frac{3}{2}\right)$
108. $f(a) = 2a^2 - 3a + 10$
 $\Rightarrow f'(a) = 4a - 3 \Rightarrow f''(a) = 4 > 0$
 For minimum value of $f(a)$,
 $f'(a) = 0 \Rightarrow a = \frac{3}{4}$
 $\therefore f(a)$ is minimum at $a = \frac{3}{4}$
 $\therefore [f(a)]_{\min} = f\left(\frac{3}{4}\right) = 2\left(\frac{3}{4}\right)^2 - 3\left(\frac{3}{4}\right) + 10 = \frac{71}{8}$

109. $f(x) = a \sin x + \frac{1}{3} \sin 3x$
 $\Rightarrow f'(x) = a \cos x + \frac{1}{3} \cdot 3 \cos 3x$
 $\Rightarrow f'(x) = a \cos x + \cos 3x$
 Now, $f'\left(\frac{\pi}{3}\right) = 0$
 $\Rightarrow a \cos \frac{\pi}{3} + \cos \pi = \frac{a}{2} - 1 = 0$
 $\Rightarrow a = 2$
110. Clearly, it has a maximum at $x = 1$.
112. $y = x^3 - 3x^2 + 5$
 $f(x) = x^3 - 3x^2 + 5$
 $f'(x) = 3x^2 - 6$
 $f''(x) = 6x - 6$
 $f'(x) = 0$ at $x = 0, x = 2$
 $f''(0) < 0, f''(2) > 0$
 $\therefore f(x)$ is maximum at $x = 2$
113. Let $f(x) = 2x^3 - 15x^2 + 36x + 4$
 $\therefore f'(x) = 6x^2 - 30x + 36 = 0$ at $x = 3, 2$
 $\therefore f''(x) = 12x - 30$ is -ve at $x = 2$
 \therefore maximum value of $f(x)$ attained at $x = 2$
114. $f'(x) = 6x^2 - 6x - 12$
 $f'(x) = 0 \Rightarrow (x-2)(x+1) = 0 \Rightarrow x = -1, 2$
 Here $f(4) = 128 - 48 - 48 + 5 = 37$
 $f(-1) = -2 - 3 + 12 + 5 = 12$
 $f(2) = 16 - 12 - 24 + 5 = -15$
 $f(-2) = -16 - 12 + 24 + 5 = 1$
 \therefore the maximum value of function is 37 at $x = 4$.
115. Given $f(x) = (1-x)^2, f(x) = x^3 - 2x^2 + 1$
 $\therefore f'(x) = 3x^2 - 4x + 1$
 Put $f'(x) = 0$ i.e. $3x^2 - 4x + 1 = 0$
 $\Rightarrow 3x^2 - 3x - x + 1 = 0 \Rightarrow x = 1, 1/3$
 $f''(x) = 6x - 4$
 $\therefore f''(1) = 2 > 0$ and $f''(1/3) = -2 < 0$
 $\therefore f(x)$ is maximum at $x = \frac{1}{3}$
 \therefore Maximum value = $f\left(\frac{1}{3}\right) = \frac{4}{27}$
116. Let $f(x) = x^2 + \frac{250}{x}$
 $\Rightarrow f'(x) = 2x - \frac{250}{x^2}$
 $\Rightarrow f''(x) = 2 + \frac{500}{x^3}$



For maximum or minimum of $f(x)$,

$$f'(x) = 0 \Rightarrow 2^3 - 250 = 0$$

$$\Rightarrow 2^3 = 250 \Rightarrow x = 5$$

$$\therefore f''(5) = 2 + \frac{500}{125} = 6 > 0$$

$\therefore f$ has minimum at $x = 5$ and minimum value of f at $x = 5$ is $f(5) = 25 + 50 = 75$

$$117. f(x) = \log x$$

$$f'(x) = 1 + \log x$$

$$\text{for minimum, } f'(x) = 0 \Rightarrow \log x = -1 \Rightarrow x = \frac{1}{e}$$

$$f''(x) = \frac{1}{x} \quad f''\left(\frac{1}{e}\right) = \frac{1}{\frac{1}{e}} = e > 0$$

$\therefore f(x)$ is minimum at $x = \frac{1}{e}$

$$\therefore f\left(\frac{1}{e}\right) = \frac{1}{e} \log\left(\frac{1}{e}\right) = -\frac{1}{e}$$

$$118. \text{ Let } f(x) = \frac{\log x}{x} \Rightarrow f'(x) = \frac{1}{x^2} - \frac{\log x}{x^2}$$

For maximum or minimum value of $f(x)$,

$$f'(x) = 0$$

$$\Rightarrow \frac{1 - \log x}{x^2} = 0$$

$\therefore \log x = 1$ or $x = e$, which lie in $(0, \infty)$.

For $x = e$, $\frac{d^2}{dx^2} = -\frac{1}{e^3}$, which is -ve.

$\therefore x = e$ is maximum at $x = e$

$$\text{and its maximum value} = \frac{\log e}{e} = \frac{1}{e}$$

$$119. x + y = 32 \Rightarrow y = 32 - x$$

$$\Rightarrow x^2 + y^2 = x^2 + (32 - x)^2$$

$$\text{Let } z = x^2 + (32 - x)^2$$

$$\Rightarrow z' = 2x + 2(32 - x)(-1) = 4x - 64$$

$$\text{Now, } z'' = 4 > 0$$

\therefore at $x = 16$ and $y = 32 - 16 = 16$

$$x^2 + y^2 = 32 \text{ have minimum value}$$

\therefore minimum value $= x^2 + y^2 = (16)^2 + (16)^2 = 512$

$$120. \text{ Let } f(x) = 25(1 - x)^{75}$$

$$\therefore f'(x) = 25(75)(1 - x)^{74}(-1) + 25^{-24}(1 - x)^{75}$$

For maximum value of $f(x)$, $f'(x) = 0$

$$\Rightarrow -75 \cdot 25(1 - x)^{74} + 25^{-24}(1 - x)^{75} = 0$$

$$\Rightarrow 25^{-24}(1 - x)^{74}(1 - 4x) = 0$$

$$\Rightarrow x = 0 \text{ or } x = 1 \text{ or } x = \frac{1}{4}$$

$$\text{At } x = \frac{1}{4}, f'\left(\frac{1}{4} - h\right) > 0 \text{ and } f'\left(\frac{1}{4} + h\right) < 0$$

$\therefore f(x)$ has maximum value at $x = \frac{1}{4}$.

$$21. h(x) = \frac{x^2 + \frac{1}{2}}{-\frac{1}{x}}$$

$$= \frac{\left(-\frac{1}{x}\right)^2 + 2}{\left(-\frac{1}{x}\right)}$$

$$= \left(-\frac{1}{x}\right) + \frac{2}{\left(-\frac{1}{x}\right)}$$

$$\text{When } -\frac{1}{x} < 0, \left(-\frac{1}{x}\right) + \frac{2}{\left(-\frac{1}{x}\right)} \leq -2\sqrt{2}$$

$$\text{When } -\frac{1}{x} > 0, \left(-\frac{1}{x}\right) + \frac{2}{\left(-\frac{1}{x}\right)} \geq 2\sqrt{2}$$

The local minimum value of $h(x)$ is $2\sqrt{2}$.

$$122. f(x) = 2x^3 - 9ax^2 + 12a^2x + 1$$

$$\Rightarrow f'(x) = 6x^2 - 18ax + 12a^2$$

$$\Rightarrow f''(x) = 12x - 18a$$

For maximum or minimum of $f(x)$, $f'(x) = 0$

$$\Rightarrow 6x^2 - 18ax + 12a^2 = 0$$

$$\Rightarrow x^2 - 3ax + 2a^2 = 0$$

$$\Rightarrow x = a \text{ or } x = 2a$$

At $x = a$, f has maximum $(5a^3 + 1)$

and at $x = 2a$, f has minimum $(4a^3 + 1)$

Since, $p^3 = q$

$$\therefore a^3 = 2a \Rightarrow a = \sqrt{2} \text{ or } a = 0$$

But $a > 0$

$$\therefore a = \sqrt{2}$$

$$123. f(x) = x^2 + 2bx + 2c^2$$

$$f'(x) = 2x + 2b = 0, \text{ at } x = -b$$

$$f''(x) = 2 > 0$$

$\therefore f(x)$ is minimum at $x = -b$

$$\therefore f(-b) = b^2 - 2b^2 + 2c^2 = 2c^2 - b^2$$

$$g(x) = -x^2 - 2cx + b^2$$

$$g'(x) = -2x - 2c = 0 \text{ at } x = -c$$

$$g''(x) = -2 < 0$$

$\therefore g(x)$ is maximum at $x = -c$

$$\therefore g(-c) = -c^2 + 2c^2 + b^2 = b^2 + c^2$$

Given, minimum value of $f(x) >$ maximum of

$$g(x)$$

$$\Rightarrow 2c^2 - b^2 > b^2 + c^2$$

$$\Rightarrow c^2 > 2b^2$$



124. $f(x) = x^2 + e$
 $f'(x) = 2x + e$
 $f''(x) = 2 + e$
 $f'''(x) = e$
 $f^{(4)}(x) = e$
 $\Rightarrow f_3 = f_4 \Rightarrow n = 3$

125. Let a and b be the lengths of two adjacent sides of the rectangle.

Then, its perimeter is $2(a + b) = 36$
 $\Rightarrow a + b = 18 \Rightarrow b = 18 - a$ (i)

Area of rectangle,

$A = ab = a(18 - a) = 18a - a^2$

$\therefore \frac{dA}{da} = 18 - 2a$

For maximum or minimum,


$\frac{dA}{da} = 0 \Rightarrow 18 - 2a = 0 \Rightarrow a = 9$

From (i), $b = 18 - 9 = 9$

126. Total length of wire = $r + r + r\theta$

$\Rightarrow 20 = 2r + r\theta$

$\Rightarrow \theta = \frac{20 - 2r}{r}$



$A = \frac{1}{2}r^2\theta$

$= \frac{1}{2}r^2\left(\frac{20 - 2r}{r}\right) = 10r - r^2$

$\therefore \frac{dA}{dr} = 10 - 2r$

For maximum area, $\frac{dA}{dr} = 0$

$\Rightarrow 0 = 10 - 2r \Rightarrow 10 = 2r \Rightarrow r = 5$ m

\therefore Area = $\frac{1}{2}r(20 - 2r)$

$= \frac{1}{2} \times 5 \times (20 - 10) = 25$ sq.m.

127. Let $\frac{1}{x} + \frac{1}{y} = 4 \Rightarrow \frac{1}{y} = 4 - \frac{1}{x}$

$\frac{1}{x} + \frac{1}{4 - \frac{1}{x}} = \frac{1}{2}$

$f(x) = \frac{4}{x} = \frac{4}{(4 - \frac{1}{x})} = \frac{4}{4 - \frac{1}{x}}$

$\therefore f'(x) = \frac{-4}{(4 - \frac{1}{x})^2} \cdot (-\frac{1}{x^2})$

For maximum or minimum of $f(x)$,

$f'(x) = 0 \Rightarrow 4 - \frac{1}{x} = 0$

$\therefore x = 2$ and $y = 2$

$\therefore \min\left(\frac{1}{x} + \frac{1}{y}\right) = \frac{1}{2} + \frac{1}{2} = 1$

128. Let a and b be the lengths of two adjacent sides of the rectangle.

Then, its perimeter is $P = 2(a + b)$ (i)

$\Rightarrow a = \frac{P - 2b}{2}$

Area of rectangle, $A = ab$

$= \left(\frac{P - 2b}{2}\right)b = \frac{Pb - 2b^2}{2}$

$\therefore \frac{dA}{db} = \frac{P - 4b}{2}$ and $\frac{d^2A}{db^2} = -2$

For maximum or minimum,

$\frac{dA}{db} = 0$

$\Rightarrow \frac{P - 4b}{2} = 0$

$\Rightarrow P = 4b$

$\Rightarrow 2a + 2b = 4b$

....[From (i)]

$\Rightarrow a = b$

$\therefore \left(\frac{d^2A}{db^2}\right)_b = -2 < 0$

Hence, the area of a rectangle will be maximum when rectangle is a square.

129. $p(t) = 1000 + \frac{1000t}{100 + t^2}$

$\therefore \frac{dp}{dt} = \frac{(100 + t^2)1000 - 1000t \cdot 2t}{(100 + t^2)^2}$

$= \frac{1000(100 - t^2)}{(100 + t^2)^2}$

For extremum,

$\frac{dp}{dt} = 0 \Rightarrow t = 10$

Now $\left.\frac{dp}{dt}\right|_{t < 10} > 0$ and $\left.\frac{dp}{dt}\right|_{t > 10} < 0$

\therefore At $t = 10$, $\frac{dp}{dt}$ change from positive to negative.

It is a critical point.

\therefore p is maximum at $t = 10$.

$\therefore p_{\max} = p(10)$

$= 1000 + \frac{1000 \cdot 10}{100 + 10^2} = 1050$

130. $f(x) = x + \frac{1}{x} \Rightarrow f'(x) = 1 - \frac{1}{x^2}$

$\Rightarrow f'(x) = 0 \Rightarrow x^2 - 1 = 0 \Rightarrow x = 1, -1$

But it is given that x is positive

\therefore at $x = 1$, $f(x) = 1 + \frac{1}{1} = 2$



131. $f(x) = x + \sin x \Rightarrow f'(x) = 1 + \cos x$
 Now, $f'(x) = 0 \Rightarrow 1 + \cos x = 0 \Rightarrow \cos x = -1$
 $\Rightarrow x = \pi$
 Now, $f''(x) = -\sin x$, $f''(\pi) = 0$
 $f'''(x) = -\cos x$
 $\therefore f'''(\pi) = 1 \neq 0$
 \therefore Neither maximum nor minimum.

132. Let α, β be the roots of the equation
 $x^2 - (a-2)x - a + 1 = 0$,
 then $\alpha + \beta = a - 2$, $\alpha\beta = -a + 1$
 $\therefore z = \alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$
 $= (a-2)^2 + 2(a-1) = a^2 - 2a + 2$
 $\therefore \frac{dz}{da} = 2a - 2 = 0 \Rightarrow a = 1$
 $\frac{d^2z}{da^2} = 2 > 0$, so z has minima at $a = 1$
 So $\alpha^2 + \beta^2$ has least value for $a = 1$. This is because we have only one stationary value at which we have minima. Hence $a = 1$.

133. $f(x) = \frac{x^2 - 1}{x^2 + 1} = \frac{x^2 + 1 - 2}{x^2 + 1} = 1 - \frac{2}{x^2 + 1}$
 $\therefore f(x) < 1 \forall x$ and $f(x) \geq -1$ as $-\frac{2}{x^2 + 1} \leq 2$
 $\therefore -1 \leq f(x) < 1$
 $\therefore f(x)$ has minimum value -1 .

134. Let $f(x) = \frac{x^2 - 1}{x^2 + 1}$
 $\Rightarrow \frac{d}{dx} = \frac{(x^2 + 1)(2x) - (x^2 - 1)(2x)}{(x^2 + 1)^2}$
 $\Rightarrow \frac{d}{dx} = \frac{2x^2 - 2}{(x^2 + 1)^2} = 0$
 $\Rightarrow 2x^2 - 2 = 0 \Rightarrow x = -1, +1$
 $\frac{d^2}{dx^2} = \frac{4(-3x + 1)}{(x^2 + 1)^2}$
 At $x = -1$, $\frac{d^2}{dx^2} < 0$ the function will occupy maximum value,
 $\therefore f(-1) = 3$ and at $x = 1$, $\frac{d^2}{dx^2} > 0$ the function will occupy minimum value.
 $\therefore f(1) = \frac{1}{3}$

135. Let $f(x) = \exp(2 + \sqrt{3} \cos x + \sin x)$
 $\Rightarrow f'(x) = \exp(2 + \sqrt{3} \cos x + \sin x)$
 $\times (-\sqrt{3} \sin x + \cos x)$

For maximum or minimum of $f(x)$, $f'(x) = 0$
 $\Rightarrow \exp(2 + \sqrt{3} \cos x + \sin x) (-\sqrt{3} \sin x + \cos x) = 0$
 $\Rightarrow -\sqrt{3} \sin x + \cos x = 0$
 $\Rightarrow \frac{\sqrt{3}}{2} \sin x - \frac{1}{2} \cos x = 0$
 $\Rightarrow \sin\left(x - \frac{\pi}{6}\right) = 0$
 $\Rightarrow x = \frac{\pi}{6}$

At $x = \frac{\pi}{6}$, $f''(x)$ is negative

- $\therefore f$ has maximum at $x = \frac{\pi}{6}$ and maximum value of f at $x = \frac{\pi}{6}$ is

$$f\left(\frac{\pi}{6}\right) = \exp\left(2 + \sqrt{3}\left(\frac{\sqrt{3}}{2}\right) + \frac{1}{2}\right) = \exp(4)$$

136. Let $y = x \Rightarrow \log y = x \cdot \log x$, ($x > 0$)
 Differentiating, $\frac{dy}{dx} = x(1 + \log x)$;

$\therefore \frac{dy}{dx} = 0$

$$\Rightarrow \log x = -1 \Rightarrow x = e^{-1} = \frac{1}{e}$$

- \therefore Stationary point is $x = \frac{1}{e}$

137. Let $f(x) = \left(\frac{1}{e}\right)^x$

$$\Rightarrow f(x) = e^{-x}$$

$$\Rightarrow f'(x) = -e^{-x} (1 + \log x)$$

$$\Rightarrow f''(x) = -e^{-x} (1 + \log x)^2 - e^{-x-1}$$

For maximum or minimum of $f(x)$, $f'(x) = 0$

$$\Rightarrow -e^{-x} (1 + \log x) = 0$$

$$\Rightarrow 1 + \log x = 0$$

$$\Rightarrow \log x = -1 = \log \frac{1}{e}$$


$$\Rightarrow x = \frac{1}{e}$$

- $\therefore f''\left(\frac{1}{e}\right) = \left(\frac{1}{e}\right)^{-\frac{1}{e}} \left(1 + \log \frac{1}{e}\right)^2 - \left(\frac{1}{e}\right)^{-\frac{1}{e}-1}$
 $= e^{\frac{1}{e}} (1-1)^2 - e^{\frac{1}{e}+1}$
 $= -e^{\frac{1}{e}+1} < 0$



$\therefore f$ has maximum at $= \frac{1}{e}$ and maximum value of f at $= \frac{1}{e}$ is $f\left(\frac{1}{e}\right) = (e)^{1/e}$

138. Let r be the radius and h be the height, then from the figure, $r^2 + \left(\frac{h}{2}\right)^2 = 3^2$




$\therefore h^2 = 4(9 - r^2)$
 $\Rightarrow r^2 = 36 - h^2$
 Now, $V = \pi r^2 h$
 $\Rightarrow V = \pi(36 - h^2)h$

$\therefore \frac{dV}{dh} = \pi(36 - 3h^2)$

for max or min, $\frac{dV}{dh} = 0$

$\Rightarrow \pi(36 - 3h^2) = 0 \Rightarrow h^2 = 12 \Rightarrow h = 2\sqrt{3}$

139. Let r be the radius and h be the height, then from the figure, $r^2 + \left(\frac{h}{2}\right)^2 = R^2$



$\therefore h^2 = 4(R^2 - r^2)$

Now, $V = \pi r^2 h = 2\pi r^2 \sqrt{R^2 - r^2}$

$\therefore \frac{dV}{dr} = 4\pi r \sqrt{R^2 - r^2} + 2\pi r^2 \cdot \frac{(-2r)}{2\sqrt{R^2 - r^2}}$


For max. or min., $\frac{dV}{dr} = 0$

$\Rightarrow 4\pi r \sqrt{R^2 - r^2} = \frac{2\pi r^3}{\sqrt{R^2 - r^2}} \Rightarrow 2(R^2 - r^2) = r^2$

$\Rightarrow 2R^2 = 3r^2 \Rightarrow r = \sqrt{\frac{2}{3}} R \Rightarrow \frac{d^2V}{dr^2} = -ve$

$\therefore V$ is max., when $r = \sqrt{\frac{2}{3}} R$.

140. Let diameter of sphere be $AE = 2r$
 Let radius of cone be and height be .



$\therefore AD =$
 Since, $BD^2 = AD \cdot DE$
 $\Rightarrow r^2 = (2r -)$

Volume of cone $V = \frac{1}{3} \pi r^2 h = \frac{1}{3} \pi (2r -)$
 $= \frac{1}{3} \pi (2r^2 - 3)$

$\Rightarrow \frac{dV}{d} = \frac{1}{3} \pi (4r - 3)$

Now $\frac{dV}{d} = 0$

$\Rightarrow \frac{1}{3} \pi (4r - 3) = 0 \Rightarrow (4r - 3) = 0$

$\Rightarrow = \frac{4}{3} r, 0$


Now $\frac{d^2V}{d^2} = \frac{1}{3} \pi (4r - 6)$

$\Rightarrow \left(\frac{d^2V}{d^2}\right)_{\frac{4}{3}r} = \frac{1}{3} \pi \left(4r - 6 \times \frac{4}{3} r\right) < 0$

So, volume of cone is maximum at $= \frac{4}{3} r$.

$\therefore \frac{\text{Height of Cone}}{\text{Diameter of Sphere}} = \frac{2}{3}$

141.



Area of rectangle ABCD

$= (2a \cos \theta) (2b \sin \theta) = 2ab \sin 2\theta$


Hence, area of greatest rectangle is equal to $2ab$, when $\sin 2\theta = 1$.



142. Let $f(x) = 3x^3 - px + q$. Then,
 $f'(x) = 3x^2 - p$

$$= 3 \left(-\sqrt{\frac{p}{3}} \right) \left(+\sqrt{\frac{p}{3}} \right)$$

The signs of $f'(x)$ for different values of x are as shown below:



Since, $f'(x)$ changes its sign from positive to negative in the neighbourhood of $-\sqrt{\frac{p}{3}}$.

So, $-\sqrt{\frac{p}{3}}$ is a point of local maximum.

Similarly, $+\sqrt{\frac{p}{3}}$ is a point of local minimum.

143. For any $x \in [0, 1]$, we have
 $x^2 \leq e^x \leq 1$

$$\Rightarrow x^2 e^{-x} \leq e^{-x} \leq e^{-x}$$

$$\Rightarrow e^{-x} + x^2 e^{-x} \leq e^{-x} + e^{-x} \leq e^{-x} + e^{-x}$$

$$\Rightarrow h(x) \leq g(x) \leq f(x)$$

Now, $f(x) = e^x + e^{-x}$
 $\Rightarrow f'(x) = 2(e^x - e^{-x}) > 0$ for all $x \in (0, 1]$

$\Rightarrow f(x)$ is increasing on $(0, 1]$
 $\Rightarrow f(1)$ is the maximum value of $f(x)$ on $[0, 1]$
 $\Rightarrow a = e + e^{-1}$

Also, $f(1) = g(1) = h(1) = e + e^{-1}$


$\therefore a = b = c = e + e^{-1}$

144. If $f(x)$ has a local minimum at $x = -1$, then

$$\lim_{x \rightarrow -1^+} f(x) = \lim_{x \rightarrow -1^-} f(x)$$

$$\Rightarrow \lim_{x \rightarrow -1^+} (2x + 3) = \lim_{x \rightarrow -1^-} (kx - 2)$$

$$\Rightarrow -2 + 3 = k + 2 \Rightarrow k = -1$$




145. $f(x) = \int_0^x \sqrt{t} \sin t dt \Rightarrow f'(x) = \sqrt{x} \sin x$

For local maximum or minimum of $f(x)$,

$$f'(x) = 0 \Rightarrow \sqrt{x} \sin x = 0$$

$$\Rightarrow x = \pi, 2\pi \dots \left[\because x \in \left(0, \frac{5\pi}{2}\right) \right]$$

The changes in signs of $f'(x)$ in the neighbourhoods of π and 2π are as shown below:




Clearly, $f'(x)$ changes its sign from positive to negative in the neighbourhood of $x = \pi$ and negative to positive in the neighbourhood of $x = 2\pi$. Thus, $f(x)$ has a local maximum at $x = \pi$ and a local minimum at $x = 2\pi$.

146. $f(x) = |x+1| + |x^2-1| = \begin{cases} -x^2 - 1, & x \leq -1 \\ -x^2 + 1, & -1 < x < 0 \\ -x^2 + 1, & 0 \leq x < 1 \\ x^2 - 1, & x \geq 1 \end{cases}$

$$\therefore f'(x) = \begin{cases} -2x - 1, & x < -1 \\ -2x - 1, & -1 < x < 0 \\ -2x + 1, & 0 < x < 1 \\ 2x - 1, & x > 1 \end{cases}$$

Here, $f(x)$ is not differentiable at $x = -1, 0, 1$.

The changes in signs of $f'(x)$ for different values of x are as shown below:



So, $f'(x)$ changes its sign at 5 points.

Hence, total number of points of local maximum or local minimum of $f(x)$ is 5.

147. $f(x) = \begin{cases} e^x, & 0 \leq x \leq 1 \\ 2 - e^{-x}, & 1 < x \leq 2 \\ -e^x, & 2 < x \leq 3 \end{cases}$

and $g(x) = \int_0^x f(t) dt, x \in [1, 3]$

$$\therefore g'(x) = f(x) = \begin{cases} 2 - e^{-x}, & 1 < x \leq 2 \\ -e^x, & 2 < x \leq 3 \end{cases}$$

Now, $g'(x) = 0 \Rightarrow x = 1 + \log_e 2$ and $x = e$

Also, $g'(x) > 0$ for $x \in (1, 1 + \log_e 2)$

and $g'(x) < 0$ for $x \in (1 + \log_e 2, 2)$.



So, $g(x)$ attains a local maximum at $x = 1 + \log_e 2$.
 Similarly,
 $g'(x) < 0$ for $2 < x < e$
 and $g'(x) > 0$ for $e < x < 3$
 So, $g(x)$ attains a local minimum at $x = e$.
 We have,

$$f'(x) = \begin{cases} e & , 0 < x < 1 \\ -e^{-1} & , 1 < x < 2 \\ 1 & , 2 < x < 3 \end{cases}$$

Clearly, $f'(x) > 0$ for $x \in (0, 1)$
 $f'(x) < 0$ for $x \in (1, 2)$
 $f'(x) > 0$ for $x \in (2, 3)$
 So, $f(x)$ attains local maximum at $x = 1$ and local minimum at $x = 2$.
 Hence, option (C) is incorrect.

148. $f(x) = \begin{cases} (2+x)^3 & , -3 < x \leq -1 \\ x^{2/3} & , -1 < x < 2 \end{cases}$
 $\Rightarrow f'(x) = \begin{cases} 3(2+x)^2 & , -3 < x < -1 \\ \frac{2}{3}x^{-1/3} & , -1 < x < 2 \end{cases}$

Clearly, $f'(x)$ changes its sign from positive to negative as x passes through $x = -1$ from left to right.
 So, $f(x)$ attains a local maximum at $x = -1$.
 Here, $f'(x) > 0$ for all $x \in (-3, -1)$ and $f'(x) < 0$ for $x \in (-1, 0)$.
 Also, $f'(x) > 0$ for $x \in (0, 2)$.
 But, $f'(0)$ does not exist.
 So, $f(x)$ attains a local minimum at $x = 0$.
 Hence, the total number of local maxima and local minima is 2.

149. $f(x) = (1 + b^2)x^2 + 2bx + 1$
 $\Rightarrow f'(x) = 2(1 + b^2)x + 2b$
 $\Rightarrow f''(x) = 2(1 + b^2) > 0$
 For minimum value of $f(x)$,
 $f'(x) = 0$
 $\Rightarrow 2(1 + b^2)x + 2b = 0$
 $\Rightarrow x = -\frac{b}{1 + b^2}$
 $\therefore f(x)$ is minimum at $x = -\frac{b}{1 + b^2}$
 \therefore Minimum value of $f(x) = \frac{1}{1 + b^2}$
 $\therefore m(b) = \frac{1}{1 + b^2}$

Since, $\frac{1}{1 + b^2} \leq 1$ and $\frac{1}{1 + b^2} > 0 \forall b \in \mathbb{R}$

$\therefore 0 < m(b) \leq 1$
 \therefore range of $m(b)$ is $(0, 1]$.
 150. $P(x) = x^4 + ax^3 + bx^2 + cx + d$
 $\Rightarrow P'(x) = 4x^3 + 3ax^2 + 2bx + c \dots(i)$
 Since, $x = 0$ is the only real root of $P'(x) = 0$.
 $\therefore P'(0) = 0 \Rightarrow c = 0$
 Putting $c = 0$ in (i), we get
 $P'(x) = (4x^2 + 3ax + 2b)$
 Since, $x = 0$ is the only real root of $P'(x) = 0$.
 $\therefore 4x^2 + 3ax + 2b = 0$ has no real root.
 $\Rightarrow 9a^2 - 32b < 0$
 Given, $P(-1) < P(1)$
 $\Rightarrow 1 - a + b - c + d < 1 + a + b + c + d$
 $\Rightarrow a > 0$
 But, $9a^2 - 32b < 0$. Therefore, $b > 0$
 $\therefore P'(x) = (4x^2 + 3ax + 2b) > 0$ for all $x \in (0, 1]$
 $\Rightarrow P(x)$ is increasing in $(0, 1]$
 $\Rightarrow P(1)$ is the maximum value of $P(x)$.
 Also, $P'(x) = (4x^2 + 3ax + 2b) < 0$ for all $x \in [-1, 0)$
 $\dots[\because 4x^2 + 3ax + 2b > 0$ for all $x \in [-1, 0)$
 $\Rightarrow P(x)$ is decreasing in $[-1, 0)$.
 $\Rightarrow P(-1)$ is not the minimum value of P .

151. $f(x) = \ln\{g(x)\}$
 $\Rightarrow g(x) = e^{f(x)}$
 $\Rightarrow g'(x) = e^{f(x)} f'(x)$
 For local maximum of $g(x)$,
 $g'(x) = 0$
 $\Rightarrow e^{f(x)} f'(x) = 0$
 $\Rightarrow f'(x) = 0$
 $\Rightarrow 2010(x - 2009)(x - 2010)^2(x - 2011)^3 \times (x - 2012)^4 = 0$
 $\Rightarrow x = 2009, 2010, 2011, 2012$
 $\therefore f'(x)$ changes its sign from positive to negative in the neighbourhood of $x = 2009$.
 $\therefore g(x)$ has a local maximum at $x = 2009$ only.

152. According to the given condition,
 $\frac{d}{dx} = 0$
 $\Rightarrow 12 - 3x^2 = 0$
 $\Rightarrow x = \pm 2$
 When $x = 2$, $y = 16$
 When $x = -2$, $y = -16$
 \therefore the required points are $(2, 16)$ and $(-2, -16)$.



153. $v = \frac{d}{dt} = 4t^3 - 3kt^2$

$\therefore \frac{dv}{dt} = 12t^2 - 6kt$

At $t = 2$, $\frac{dv}{dt} = 0$


$\Rightarrow 48 - 12k = 0 \Rightarrow k = 4$

154. Since, $f(x)$ satisfies the conditions of Rolle's theorem.

$\therefore f(2) = f(1)$

Now, $\int_1^2 f'(x) dx = [f(x)]_1^2 = f(2) - f(1) = 0$

155. It is always increasing.



156. $f(x) = x^3 + bx^2 + cx + d$

$\therefore f'(x) = 3x^2 + 2bx + c$

Now its discriminant $= 4(b^2 - 3c)$

$\Rightarrow 4(b^2 - c) - 8c < 0$, as $b^2 < c$ and $c > 0$

$\Rightarrow f'(x) > 0$ for all $x \in \mathbb{R}$

$\Rightarrow f$ is strictly increasing on \mathbb{R} .

157. Since $x = 1$ and $x = 3$ are extreme points of $p(x)$.

$\therefore p'(1) = 0$ and $p'(3) = 0$

$\therefore (x - 1)$ and $(x - 3)$ are the factors of $p'(x)$.

$\therefore p'(x) = k(x - 1)(x - 3) = k(x^2 - 4x + 3)$

$\Rightarrow p(x) = k\left(\frac{x^3}{3} - 2x^2 + 3x\right) + c$

Given, $p(1) = 6$ and $p(3) = 2$

$\Rightarrow 6 = k\left(\frac{1}{3} - 2 + 3\right) + c$ and 2

$= k(9 - 18 + 9) + c$

$\Rightarrow 6 = \frac{4k}{3} + c$ and $c = 2 \Rightarrow k = 3$

$\therefore p'(x) = 3(x^2 - 4x + 3)$

$\therefore p'(0) = 9$

158. Let $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4$

Given, $\lim_{x \rightarrow 0} \left[1 + \frac{f(x)}{2}\right] = 3$

$\Rightarrow \lim_{x \rightarrow 0} \frac{f(x)}{2} = 3 - 1 = 2$

$\Rightarrow \lim_{x \rightarrow 0} \frac{a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4}{2} = 2$

$\therefore a_0 = 0, a_1 = 0, a_2 = 2$

$\therefore f(x) = 2x^2 + a_3x^3 + a_4x^4$

$\therefore f'(x) = 4x + 3a_3x^2 + 4a_4x^3 = (4 + 3a_3x + 4a_4x^2)$

Given, $f'(1) = 0$ and $f'(2) = 0$

$\Rightarrow 4 + 3a_3 + 4a_4 = 0 \dots(i)$

and $4 + 6a_3 + 16a_4 = 0 \dots(ii)$

Solving (i) and (ii), we get

$a_4 = \frac{1}{2}, a_3 = -2$

$\therefore f(x) = 2x^2 - 2x^3 + \frac{x^4}{2}$

$\therefore f(2) = 8 - 16 + 8 = 0$

159. $\tan A \cdot \tan B$ is maximum if $A = B = \frac{\pi}{6}$

\therefore Maximum of $\tan A \cdot \tan B = \frac{1}{3}$

160. According to the given condition,

$4 + 2\pi r = 2$

$\Rightarrow 2 + \pi r = 1 \dots(i)$

$A = x^2 + \pi r^2 = \left(\frac{1 - \pi r}{2}\right)^2 + \pi r^2$

$\therefore \frac{dA}{dr} = 2\left(\frac{1 - \pi r}{2}\right)\left(-\frac{\pi}{2}\right) + 2\pi r$

For maximum or minimum, $\frac{dA}{dr} = 0$

$\Rightarrow \pi(1 - \pi r) = 4\pi r$

$\Rightarrow 1 = 4r + \pi r \dots(ii)$

From (i) and (ii), we get

$2 + \pi r = 4r + \pi r$

$\Rightarrow r = 2$

161. $f(x) = \tan^{-1} \sqrt{\frac{1 + \sin x}{1 - \sin x}}$

$= \tan^{-1} \sqrt{\frac{\left(\cos \frac{x}{2} + \sin \frac{x}{2}\right)^2}{\left(\cos \frac{x}{2} - \sin \frac{x}{2}\right)^2}}$

$= \tan^{-1} \left(\tan\left(\frac{\pi}{4} + \frac{x}{2}\right)\right) = \frac{\pi}{4} + \frac{x}{2}$

$\Rightarrow f'(x) = \frac{1}{2}$ and at $x = \frac{\pi}{6}, f(x) = \frac{\pi}{3}$

\therefore equation of the normal at $\left(\frac{\pi}{6}, \frac{\pi}{3}\right)$ is

$-\frac{\pi}{3} = -2 \left(-\frac{\pi}{6}\right) \Rightarrow -\frac{\pi}{3} + 2 = \frac{2\pi}{3}$

Only option (A) satisfies this equation.



Evaluation Test

- Let $f(x) = ax^4 + bx^3 + cx^2 + dx$
 $\therefore f(0) = 0$
 and $f(3) = a \cdot 3^4 + b \cdot 3^3 + c \cdot 3^2 + d \cdot 3$
 $= 81a + 27b + 9c + 3d$
 $= 3(27a + 9b + 3c + d)$
 $= 3 \times 0$
 $\therefore f(0) = f(3) = 0$
 $f(x)$ is a polynomial function, it is continuous and differentiable.
 Now, $f'(x) = 4ax^3 + 3bx^2 + 2cx + d$
 By Rolle's theorem, there exist at least 1 root of the equation $f'(x) = 0$ in between 0 and 3.
- The equation of the curve is $y = x^2 + bx + c$.
 $\therefore \frac{dy}{dx} = 2x + b \dots (i)$
 Since, the curve touches the line $y = 1$ at (1,1).
 $\therefore [2x + b]_{(1,1)} = 1$
 $\therefore 2(1) + b = 1$
 $\Rightarrow b = -1$
 Substituting the value of b in equation (i), we get
 $\frac{dy}{dx} = 2x - 1$
 Since, gradient is negative.
 $\therefore \frac{dy}{dx} < 0$
 $\Rightarrow 2x - 1 < 0$
 $\Rightarrow 2x < 1$
 $\Rightarrow x < \frac{1}{2}$
- The equation of the parabola is $y^2 = 8x$.
 $\therefore 2 \frac{dy}{dx} = 8$
 $\therefore \frac{dy}{dx} = \frac{8}{2} = 4 = m_1$
 Slope of given line, $m_2 = 3$
 Since, $\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$
 $\therefore \tan \frac{\pi}{4} = \left| \frac{4 - 3}{1 + 4 \cdot 3} \right|$
 $\Rightarrow 1 = \left| \frac{4 - 3}{1 + 12} \right|$
 $\therefore \frac{4 - 3}{1 + 12} = -2 \text{ or } \frac{4 - 3}{1 + 12} = 8$

- Putting $x = -2$ in the equation of the curve, we get
 $y = \frac{1}{2}$
 \therefore the point of contact is $\left(\frac{1}{2}, -2\right)$.
- $f(x) = \tan^{-1} x - \frac{1}{2} \log x$
 $\therefore f'(x) = \frac{1}{1+x^2} - \frac{1}{2x} = -\frac{(x-1)^2}{2(1+x^2)}$
 Now, $f'(x) = 0 \Rightarrow x = 1$
 $f(1) = \tan^{-1} 1 - \frac{1}{2} \log 1 = \frac{\pi}{4} = \frac{3.14}{4} = 0.785$
 Since, we are finding maxima on an interval $\left(\frac{1}{\sqrt{3}}, \sqrt{3}\right)$. We have to find the value of $f(x)$ at $\left(\frac{1}{\sqrt{3}}\right)$ and $(\sqrt{3})$
 $f\left(\frac{1}{\sqrt{3}}\right) = \tan^{-1} \frac{1}{\sqrt{3}} + \frac{1}{4} \log 3 = \frac{\pi}{6} + \frac{1}{4} \log 3$
 $= \frac{3.14}{6} + \frac{1}{4} \log 3 = 0.52 + \frac{1}{4} \times 1.0986$
 $= 0.52 + 0.2746 = 0.7946$
 $f(\sqrt{3}) = \tan^{-1}(\sqrt{3}) - \frac{1}{4} \log 3 = \frac{\pi}{3} - \frac{1}{4} \log 3$
 $= \frac{3.14}{3} - 0.2746$
 $= 1.04 - 0.2746$
 $= 0.7654$
 \therefore the greatest value of $f(x)$ is $\frac{\pi}{6} + \frac{1}{4} \log 3$.
 - $\alpha + \beta = \frac{\pi}{2}$
 $\therefore \cos \beta = \cos\left(\frac{\pi}{2} - \alpha\right) = \sin \alpha$
 Let $y = \cos \alpha \cos \beta = \cos \alpha \sin \alpha = \frac{1}{2} \sin 2\alpha$
 $\therefore \frac{dy}{d\alpha} = \frac{1}{2} \cos 2\alpha \cdot 2 = \cos 2\alpha$
 Now, $\frac{dy}{d\alpha} = 0 \Rightarrow \cos 2\alpha = 0 \Rightarrow 2\alpha = \frac{\pi}{2}$
 $\Rightarrow \alpha = \frac{\pi}{4}$



$$\text{Also, } \frac{d^2}{d\alpha^2} = -2 \sin 2\alpha = -2 < 0$$

$$\therefore \text{ is maximum when } \alpha = \frac{\pi}{4}$$

$$\therefore \text{ it is maximum at } \beta = \frac{\pi}{4}$$

6. Let $P(x_1, y_1)$ be the point on the curve at which tangent is drawn.

$$\text{The equation of the curve is } x^2 + y^2 = c^2.$$

$$\therefore \frac{d}{dx} + (1) = 0$$

$$\therefore \left(\frac{dy}{dx}\right)_{(x_1, y_1)} = -\frac{x_1}{y_1}$$

\therefore The equation of the tangent is

$$-y_1 = -\frac{x_1}{y_1}(x - x_1)$$

$$\therefore x_1 - x_1 y_1 = -x_1 + x_1 y_1$$

$$\therefore x_1 + x_1 = 2 x_1 y_1$$

$$\therefore \frac{x_1}{2 y_1} + \frac{x_1}{2 y_1} = 1$$

\therefore The tangent meets the X-axis in the point $A(2x_1, 0)$ and the Y-axis in the point $B(0, 2y_1)$

\therefore P is the mid point of AB

\therefore The ratio is 1 : 1

$$7. \text{ Let } f(x) = \frac{a^3}{3} + \frac{b^2}{2} + c + d$$

$$\therefore f'(x) = a^2 + b + c$$

$$\text{Now, } f(1) = \frac{a}{3} + \frac{b}{2} + c + d = \frac{2a + 3b + 6c + 6d}{6}$$

$$\Rightarrow f(1) = \frac{0 + 6d}{6} = d \quad \dots [\because 2a + 3b + 6c = 0]$$

$$\text{Also, } f(0) = d$$

$$\therefore f(0) = f(1)$$

$f(x)$ is a polynomial function, it is continuous and differentiable.

\therefore There exists at least one value of x in $(0, 1)$ at which $f'(x) = a^2 + b + c = 0$

\therefore one root of the equation $a^2 + b + c = 0$ has value between 0 and 1.

$$8. f(x) = \sin(1 + \cos x)$$

$$= \sin x + \sin x \cos x$$

$$\therefore f(x) = \sin x + \frac{1}{2} \sin 2x$$

$$\therefore f'(x) = \cos x + \cos 2x = 2 \cos \frac{x}{2} \cos \frac{3x}{2}$$

$$\therefore f'(x) = 0 \Rightarrow \cos \frac{x}{2} = 0 \text{ or } \cos \frac{3x}{2} = 0$$

$$\therefore \frac{x}{2} = \frac{\pi}{2} \text{ or } \frac{3x}{2} = \frac{\pi}{2}$$

$$\therefore x = \pi \text{ or } x = \frac{\pi}{3}$$

$$f''(x) = -\sin x - 2 \sin 2x < 0, \text{ only when } x = \frac{\pi}{3}$$

\therefore The maximum value of function is at $\frac{\pi}{3}$

$$\therefore f\left(\frac{\pi}{3}\right) = \frac{\sqrt{3}}{2} \left(1 + \frac{1}{2}\right) = \frac{\sqrt{3}}{2} \times \frac{3}{2} = \frac{3\sqrt{3}}{4}$$

$$9. f(x) = \sin^4 x + \cos^4 x$$

$$= (\sin^2 x + \cos^2 x)^2 - 2 \sin^2 x \cos^2 x$$

$$\therefore f(x) = 1 - \frac{1}{2} \sin^2 2x$$

$$\therefore f'(x) = -\frac{1}{2} (2 \sin 2x \cos 2x) \times 2$$

$$f'(x) = -2 \sin 2x \cos 2x$$

$$\text{Now, } f'(x) = 0$$

$$\Rightarrow \sin 2x = 0 \text{ or } \cos 2x = 0$$

$$\Rightarrow x = 0 \text{ or } x = \frac{\pi}{4}$$

$$\text{Since, } f'(x) = -2 \sin 2x \cos 2x$$

$$\Rightarrow f'(x) = -\sin 4x$$

$$\therefore f''(x) = -4 \cos 4x$$

$$\text{For } x = 0, f''(x) = -4 < 0$$

$$\text{For } x = \frac{\pi}{4}, f''(x) = 4 > 0$$

\therefore At $x = \frac{\pi}{4}$, $f(x)$ is minimum

$$\therefore \text{ Minimum value of } f(x) = 1 - \frac{1}{2}(1) = 1 - \frac{1}{2} = \frac{1}{2}$$

10. $2^{(2-3)^3+27}$ is minimum when $(2-3)^3+27$ is minimum.

$$\text{Since, } (2-3)^3+27$$

$$= 6 - 9^4 + 27^2$$

$$= 2(4^4 - 9^2 + 27)$$

$$= 2 \left[\left(2 - \frac{9}{2}\right)^2 + \frac{27}{4} \right] \geq 0, \text{ for all } x$$

\therefore Minimum value of $(2-3)^3+27$ is 0.

\therefore Minimum value of $2^{(2-3)^3+27} = 2^0 = 1$



11. $f(x) = 3 \cos |x| - 6a + b$
 $= 3 \cos - 6a + b$
 ...[$\because \cos(-x) = \cos x$]
 $\therefore f'(x) = -3 \sin - 6a$
 The function $f(x)$ is increasing for all $x \in \mathbb{R}$.
 $\therefore f'(x) > 0$
 $\therefore -3 \sin - 6a > 0$
 $\Rightarrow 6a < -3 \sin$
 $\Rightarrow a < -\frac{1}{2} \sin$
 $\Rightarrow a < -\frac{1}{2}$

12. Let $f(x) = a^2 \sec^2 x + b^2 \operatorname{cosec}^2 x$
 $\therefore f'(x) = a^2 \cdot 2 \sec x \cdot \sec x \tan x + b^2 \cdot 2 \operatorname{cosec} x (-\operatorname{cosec} x \cot x)$
 $= 2a^2 \sec^2 x \tan x - 2b^2 \operatorname{cosec}^2 x \cot x$
 Now, $f'(x) = 0$
 $\Rightarrow 2a^2 \sec^2 x \tan x - 2b^2 \operatorname{cosec}^2 x \cot x = 0$
 $2a^2 \cdot \frac{1}{\cos^2 x} \cdot \frac{\sin x}{\cos x} = 2b^2 \cdot \frac{1}{\sin^2 x} \cdot \frac{\cos x}{\sin x}$
 $\therefore \frac{\sin^4 x}{\cos^4 x} = \frac{b^2}{a^2}$
 $\therefore \tan^4 x = \frac{b^2}{a^2}$
 $\therefore \tan^2 x = \frac{b}{a}$ and $\cot^2 x = \frac{a}{b}$

Also,
 $f''(x) = 2a^2 [\sec^2 x \cdot \sec^2 x + \tan x \cdot 2 \sec x \cdot \sec x \tan x]$
 $- 2b^2 [\operatorname{cosec}^2 x (-\operatorname{cosec}^2 x) + \cot x \cdot 2 \operatorname{cosec} x (-\operatorname{cosec} x \cot x)]$
 $= 2a^2 [\sec^4 x + 2 \sec^2 x \tan^2 x]$
 $+ 2b^2 [\operatorname{cosec}^4 x + 2 \operatorname{cosec}^2 x \cot^2 x]$
 $\therefore f''(x) > 0$ for all x .
 $\therefore f(x)$ is minimum when $\tan^2 x = \frac{b}{a}$
 \therefore Minimum value of $f(x) = a^2(1 + \tan^2 x) + b^2(1 + \cot^2 x)$
 $= a^2 \left(1 + \frac{b}{a}\right) + b^2 \left(1 + \frac{a}{b}\right)$
 $= a^2 \left(\frac{a+b}{a}\right) + b^2 \left(\frac{a+b}{b}\right)$
 $= a(a+b) + b(a+b) = (a+b)^2$

13. $= \frac{a+b}{(x-4)(x-1)} = \frac{a+b}{x^2-5x+4}$
 $\therefore \frac{d}{dx} = \frac{(x^2-5x+4)a - (a+b)(2x-5)}{(x^2-5x+4)^2}$
 For extreme (i.e., maximum or minimum)
 $\frac{d}{dx} = 0$
 $\therefore a(x^2-5x+4) - (a+b)(2x-5) = 0$
 Since, x has an extreme at $P(2, -1)$
 $\therefore x = 2$ satisfies above equation
 $a(4-10+4) - (2a+b)(-1) = 0$
 $\Rightarrow -2a+2a+b = 0$
 $\Rightarrow b = 0$
 $x = -1$ satisfies the equation of the curve
 $\therefore -1 = \frac{a(2)+b}{4-10+4}$
 $\therefore -1 = \frac{2a+0}{-2} = -a$
 $\therefore a = 1$ $\therefore a = 1, b = 0$

14. Let $f(x) = \tan x$
 $\therefore f'(x) = \sec^2 x + \tan x$
 $\therefore f'(x) > 0$ for $x \in \left(0, \frac{\pi}{2}\right)$
 $\therefore f(x)$ is increasing in the interval $\left(0, \frac{\pi}{2}\right)$
 Since, $0 < \alpha < \beta < \frac{\pi}{2}$
 $\therefore f(\alpha) < f(\beta)$
 $\therefore \alpha \tan \alpha < \beta \tan \beta$
 $\Rightarrow \frac{\alpha}{\beta} < \frac{\tan \beta}{\tan \alpha}$

15. The point of intersection of the given curves is $(0, 1)$.
 Now, $m = 3$
 $\therefore \frac{d}{dx} = 3 \log 3$
 $\therefore \left(\frac{d}{dx}\right)_{(0,1)} = \log 3 = m_1$ (say)
 Also, $m = 5$
 $\therefore \frac{d}{dx} = 5 \log 5$
 $\therefore \left(\frac{d}{dx}\right)_{(0,1)} = \log 5 = m_2$ (say)
 $\therefore \tan \alpha = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right| = \frac{\log 3 - \log 5}{1 + \log 3 \log 5}$



16. Let $f(x) = ax^2 + bx + c$
 $\therefore f'(x) = 2ax + b$
 since, α and β are roots of the equation
 $ax^2 + bx + c = 0$
 $\therefore f(\alpha) = f(\beta) = 0$
 $\therefore f(x)$ being a polynomial function in x ,
 it is continuous and differentiable.
 \therefore There exists k in (α, β) such that $f'(k) = 0$
 $\therefore 2ak + b = 0, \therefore k = -\frac{b}{2a}$
 But $k \in [\alpha, \beta]$
 $\therefore \alpha < k < \beta$
 $\therefore \alpha < -\frac{b}{2a} < \beta$

17. $f(x) = \tan^{-1}(\sin x + \cos x)$
 $\therefore f'(x) = \frac{1}{1+(\sin x + \cos x)^2} \times (\cos x - \sin x)$

$$= \frac{\sqrt{2} \cos\left(x + \frac{\pi}{4}\right)}{1+(\sin x + \cos x)^2}$$

For $f(x)$ to be increasing, $f'(x) > 0$

$$\Rightarrow \sqrt{2} \cos\left(x + \frac{\pi}{4}\right) > 0$$

$$\Rightarrow \cos\left(x + \frac{\pi}{4}\right) > 0$$

$$\Rightarrow -\frac{\pi}{2} < x + \frac{\pi}{4} < \frac{\pi}{2} \Rightarrow -\frac{3\pi}{4} < x < \frac{\pi}{4}$$

- $\therefore f(x)$ is an increasing function in $\left(-\frac{\pi}{2}, \frac{\pi}{4}\right)$.

18. $f(x) = 3x^3 - 12ax^2 + 36a^2x - 4$
 Diff. w.r.t. x , we get
 $f'(x) = 3x^2 - 12a(2x) + 36a^2(1)$
 $= 3x^2 - 24ax + 36a^2$
 Now, $f'(x) = 0 \Rightarrow 3x^2 - 24ax + 36a^2 = 0$
 $\therefore x^2 - 8ax + 12a^2 = 0$
 $\therefore (x - 2a)(x - 6a) = 0$
 $\therefore x = 2a$ or $x = 6a$
 Also, $f''(x) = 6x - 24a$
 $[f''(x)]_{x=2a} = 12a - 24a = -12a < 0$
 $[f''(x)]_{x=6a} = 36a - 24a = 12a > 0$
 \therefore Maxima at $p = 2a$ and minima at $q = 6a$
 $3p = q^2 \dots(\text{given})$
 $\therefore 3 \times 2a = (6a)^2$
 $\therefore 6a = 36a^2$
 $\therefore a = \frac{1}{6}$

19. The functions e^{-x} , $\sin x$, $\cos x$ are continuous and differentiable in their respective domains.
 $\therefore f(x)$ is continuous and differentiable

$$\text{Also } f\left(\frac{\pi}{4}\right) = 0 = f\left(\frac{5\pi}{4}\right)$$

Now,

$$\begin{aligned} f'(x) &= -e^{-x}(\sin x - \cos x) + e^{-x}(\cos x + \sin x) \\ &= e^{-x}(-\sin x + \cos x + \cos x + \sin x) \\ &= 2e^{-x} \cos x \end{aligned}$$

$$\text{Also, } f'(x) = 0 \Rightarrow \cos x = 0$$

$$\therefore x = \frac{\pi}{2} \in \left[\frac{\pi}{4}, \frac{5\pi}{4}\right]$$

20. $a^2 = 3x^3 \dots(i)$

Diff. w.r.t. x , we get

$$2a \frac{d}{dx} = 3x^2$$

$$\Rightarrow \frac{d}{dx} = \frac{3x^2}{2a}$$

$$\therefore \text{slope of the normal} = -\frac{2a}{3x^2}$$

Since, the normal to the given curve makes equal intercepts with the axis.

$$\therefore -\frac{2a}{3x^2} = -1$$

$$\Rightarrow x = \frac{3x^2}{2a}$$

Substituting $x = \frac{3x^2}{2a}$ in (i) and solving, we get

$$\text{the point } \left(\frac{4a}{9}, \frac{8a}{27}\right).$$

04 Integration



Hints



Classical Thinking

$$1. \int 7e^{7+5} d = 7 \cdot \frac{e^{7+5}}{7} + c = e^{7+5} + c$$

$$2. \int (a - a^2) d = \frac{a}{\log a} - \frac{a^2}{\log a} \cdot \frac{1}{2} + c = \frac{1}{\log a} \left(a - \frac{a^2}{2} \right) + c$$

$$3. \int \frac{2+3}{5} d = \int \left(\frac{2}{5} + \frac{3}{5} \right) d = \int \left(\left(\frac{2}{5} \right) + \left(\frac{3}{5} \right) \right) d = \frac{\left(\frac{2}{5} \right)}{\log \left(\frac{2}{5} \right)} + \frac{\left(\frac{3}{5} \right)}{\log \left(\frac{3}{5} \right)} + c$$

$$4. \int \frac{1}{(-5)^2} d = \frac{(-5)^{-2+1}}{-2+1} + c = \frac{(-5)^{-1}}{-1} + c = -\frac{1}{(-5)} + c$$

$$5. \int \frac{d}{\sqrt{1-d}} = \int (1-d)^{-1/2} d = \frac{(1-d)^{-1/2+1}}{(-1)\left(-\frac{1}{2}+1\right)} + c = \frac{-2\sqrt{1-d}}{1} + c$$

$$6. \int \frac{2-1}{3} d = \int \frac{1}{3} - \int \frac{-3}{3} d = \log \frac{1}{3} + \frac{1}{2} + c$$

$$7. \int \frac{3^3 - 2\sqrt{d}}{d} = 3 \int \frac{2}{d} - 2 \int \frac{1}{2} d = 3 \log 2 - 4\sqrt{d} + c$$

$$8. \int \frac{a^{-2} + b^{-1} + c}{-3} d = \int (a^{-2} + b^{-1} + c^{-3}) d = \frac{1}{2} a^{-2+1} + \frac{1}{3} b^{-1+1} + \frac{1}{4} c^{-3+1} + c$$

$$9. \int \left(\frac{1}{d} + \frac{1}{d^3} \right) d = \int \left(\frac{1}{d} + \frac{1}{3} \frac{1}{d^3} + \frac{1}{3} \frac{1}{d^3} \right) d = \frac{1}{d} + \frac{1}{2} \frac{1}{d^2} + 3 \log \frac{1}{d^2} + c$$

$$10. \int \frac{1}{2} (2+d)^3 d = \int \frac{(8+12d+6d^2+d^3)}{2} d = \int \left(4 + 6d + 3d^2 + \frac{1}{2}d^3 \right) d = 4d + 3d^2 + 6 \log \frac{1}{2} + c$$

$$11. \int \frac{3^3 - 2\sqrt{d}}{d} = 3 \int \frac{2}{d} - 2 \int \frac{1}{2} d = 3 \log 2 - 4\sqrt{d} + c$$

$$12. \int \frac{(\sqrt{d} + \sqrt[3]{d^2})^2}{d} d = \int \left(\frac{1}{2} + \frac{2}{3} \right) d = \int \left(1 + 2 \frac{1}{6} + \frac{4}{3} \right) d = \frac{7}{6} d + \frac{4}{3} d + c = \frac{12}{7} \frac{7}{6} + \frac{3}{4} \frac{4}{3} + c$$

$$13. \frac{(1+d)^2}{(1+d^2)} = \frac{(1+d^2)+2d}{(1+d^2)} = \frac{1}{1+d^2} + 2 \cdot \frac{d}{1+d^2}$$

$$\therefore \int \frac{(1+d)^2}{(1+d^2)} d = \int \frac{1}{1+d^2} d + \int \frac{2d}{1+d^2} d = \log \frac{1}{1+d^2} + 2 \tan^{-1} d + c$$

$$15. \int \frac{\sin^2 d - \cos^2 d}{\sin^2 d \cos^2 d} d = \int \sec^2 d - \int \operatorname{cosec}^2 d = \tan d + \cot d + c$$



$$\begin{aligned}
 16. \quad & \int \sqrt{1 + \cos} \, d \\
 &= \int \sqrt{2\cos^2 \frac{1}{2}} \, d \\
 &= \sqrt{2} \int \cos\left(\frac{1}{2}\right) d \\
 &= 2\sqrt{2} \sin\left(\frac{1}{2}\right) + c
 \end{aligned}$$

$$\begin{aligned}
 17. \quad & \int \left(\cos \frac{1}{2} - \sin \frac{1}{2}\right)^2 d \\
 &= \int \left(\cos^2 \frac{1}{2} + \sin^2 \frac{1}{2} - 2\sin \frac{1}{2} \cos \frac{1}{2}\right) d \\
 &= \int (1 - \sin) d = + \cos + c
 \end{aligned}$$

$$\begin{aligned}
 18. \quad & f(1) = \int f'(1) d \\
 &= \int (1^2 + 5) d \\
 &= \frac{1^3}{3} + 5 + c
 \end{aligned}$$

$$\therefore f(0) = \frac{0}{3} + 0 + c$$

$$\Rightarrow c = -1$$

$$\therefore f(1) = \frac{1}{3} + 5 - 1$$

$$\begin{aligned}
 19. \quad & f(1) = \int f'(1) d = \int \left(\frac{1}{1+x}\right) d \\
 &= \log \frac{1}{2} + \frac{1^2}{2} + c
 \end{aligned}$$

$$\therefore f(1) = \log 1 + \frac{1^2}{2} + c$$

$$\Rightarrow \frac{5}{2} = 0 + \frac{1}{2} + c \Rightarrow c = 2$$

$$\therefore f(1) = \log \frac{1}{2} + \frac{1^2}{2} + 2$$

$$20. \quad \text{Put } \cos = t \Rightarrow -\sin d = dt$$

$$\begin{aligned}
 \therefore \int \sin \cos^4 d &= \int t^4 (-dt) = -\frac{t^5}{5} + c \\
 &= -\frac{\cos^5}{5} + c
 \end{aligned}$$

$$21. \quad \text{Put } (1 + \log) = t \Rightarrow \frac{1}{t} d = dt$$

$$\begin{aligned}
 \therefore \int \frac{(1 + \log)^2}{3} d &= \int t^2 dt = \frac{t^3}{3} + c \\
 &= \frac{(1 + \log)^3}{3} + c
 \end{aligned}$$

$$22. \quad \text{Put } 1 + x^2 = t \Rightarrow d = \frac{dt}{2}$$

$$\begin{aligned}
 \therefore \int \sqrt{1 + x^2} d &= \frac{1}{2} \int t^{1/2} dt \\
 &= \frac{1}{2} \times \frac{t^{3/2}}{3/2} = \frac{1}{3} (1 + x^2)^{3/2} + c
 \end{aligned}$$

$$23. \quad \text{Put } t = \tan^{-1} \Rightarrow dt = \frac{1}{1+x^2} d$$

$$\therefore \int \frac{e^{\tan^{-1}}}{1+x^2} d = \int e^t dt = e^t + c = e^{\tan^{-1}} + c$$

$$24. \quad \text{Put } t = 1 + \tan \Rightarrow dt = \sec^2 d$$

$$\begin{aligned}
 \therefore \int \frac{\sec^2}{1 + \tan} d &= \int \frac{1}{t} dt = \log |t| + c \\
 &= \log |1 + \tan| + c
 \end{aligned}$$

$$25. \quad \text{Put } \log \sin = t \Rightarrow \cot d = dt$$

$$\begin{aligned}
 \therefore \int \frac{\cot}{\log \sin} d &= \int \frac{dt}{t} = \log t + c \\
 &= \log(\log \sin) + c
 \end{aligned}$$

$$26. \quad \text{Put } (1 + \sin^2) = t \Rightarrow \sin 2 d = dt$$

$$\begin{aligned}
 \therefore \int \frac{\sin 2}{1 + \sin^2} d &= \int \frac{1}{t} dt = \log t + c \\
 &= \log(1 + \sin^2) + c
 \end{aligned}$$

$$27. \quad \text{Let } I = \int \frac{e - e^{-}}{e + e^{-}} d$$

$$\text{Put } e + e^{-} = t \Rightarrow (e - e^{-}) d = dt$$

$$\therefore I = \int \frac{dt}{t} = \log |t| + c = \log |e + e^{-}| + c$$

$$28. \quad \text{Put } \cos^{-1} = t \Rightarrow -\frac{1}{\sqrt{1-x^2}} d = dt$$

$$\begin{aligned}
 \therefore \int \frac{1}{\cos^{-1} \sqrt{1-x^2}} d &= -\int \frac{1}{t} dt = -\log |t| + c \\
 &= -\log |\cos^{-1}| + c
 \end{aligned}$$

$$29. \quad \text{Put } 1 + \cos^2 = t$$

$$\Rightarrow [1 + 2 \cos (-\sin)] d = dt$$

$$\Rightarrow (1 - \sin 2) d = dt$$

$$\begin{aligned}
 \therefore \int \frac{1 - \sin 2}{1 + \cos^2} d &= \int \frac{dt}{t} = \log |t| + c \\
 &= \log |1 + \cos^2| + c
 \end{aligned}$$

$$30. \quad \text{Put } \sqrt{x} = t \Rightarrow \frac{1}{2\sqrt{x}} d = dt$$

$$\begin{aligned}
 \therefore \int \frac{\cos \sqrt{x}}{\sqrt{x}} d &= 2 \int \cos t dt \\
 &= 2 \sin t + c \\
 &= 2 \sin \sqrt{x} + c
 \end{aligned}$$



$$31. \text{ Put } e^{\sqrt{t}} = t \Rightarrow \frac{e^{\sqrt{t}}}{\sqrt{t}} dt = 2 dt$$

$$\begin{aligned} \therefore \int \frac{e^{\sqrt{t}} \cos(e^{\sqrt{t}})}{\sqrt{t}} dt &= 2 \int \cos t dt \\ &= 2 \sin t + c = 2 \sin(e^{\sqrt{t}}) + c \end{aligned}$$

$$32. \text{ Put } a^t = t \Rightarrow a^t dt = \frac{dt}{\log a}$$

$$\begin{aligned} \therefore \int \frac{1}{\log a} (a^t \cos a^t) dt &= \int \frac{1}{(\log a)^2} \cos t dt \\ &= \frac{1}{(\log a)^2} \sin t + c \\ &= \frac{1}{(\log a)^2} \sin a^t + c \end{aligned}$$

$$33. \text{ Put } \log t = t \Rightarrow \frac{1}{t} dt = dt$$

$$\begin{aligned} \therefore \int \frac{\tan(\log t)}{t} dt &= \int \tan(t) dt = \log |\sec(t)| + c \\ &= \log |\sec(\log t)| + c \end{aligned}$$

$$34. \text{ Put } \log t = t \\ \Rightarrow \frac{1}{t} dt = dt$$

$$\begin{aligned} \therefore \int \frac{\sec^2(\log t)}{t} dt &= \int \sec^2 t dt = \tan t + c \\ &= \tan(\log t) + c \end{aligned}$$

$$35. \text{ Let } I = \int \frac{\log(\log t)}{\log t} dt$$

$$\text{Put } \log(\log t) = t \Rightarrow \frac{1}{\log t} dt = dt$$

$$\therefore I = \int t dt = \frac{t^2}{2} + c = \frac{[\log(\log t)]^2}{2} + c$$

$$36. \text{ Let } I = \int \sec t \cdot \log(\sec t + \tan t) dt$$

$$\text{Put } \log(\sec t + \tan t) = t \\ \Rightarrow \sec t dt = dt$$

$$\therefore I = \int t dt = \frac{t^2}{2} + c = \frac{1}{2} [\log(\sec t + \tan t)]^2 + c$$

$$37. \int \frac{d}{\sqrt{1-16t^2}} = \int \frac{d}{\sqrt{1-(4t)^2}} \\ = \frac{1}{4} \sin^{-1}(4t) + c$$

$$38. \int \frac{d}{\sqrt{2-t^2}} = \int \frac{d}{\sqrt{1-(t/\sqrt{2})^2}} \\ = \sin^{-1}\left(\frac{t}{\sqrt{2}}\right) + c$$

$$39. \int \frac{d}{t^2-2t+2} \\ = \int \frac{d}{t^2-2t+1+2-1} \\ = \int \frac{d}{(t-1)^2+1} \\ = \tan^{-1}\left(\frac{t-1}{1}\right) + c = \tan^{-1}(t-1) + c$$

$$40. \int \frac{d}{t^2+4t+13} = \int \frac{d}{(t+2)^2+3^2} \\ = \frac{1}{3} \tan^{-1}\left(\frac{t+2}{3}\right) + c$$

$$41. \text{ Let } I = \int \frac{1}{9t^2-25} dt = \frac{1}{9} \int \frac{d}{t^2 - \left(\frac{5}{3}\right)^2}$$

$$= \frac{1}{9} \cdot \frac{1}{2\left(\frac{5}{3}\right)} \cdot \log \left| \frac{t - \frac{5}{3}}{t + \frac{5}{3}} \right| + c$$

$$= \frac{1}{30} \log \left| \frac{3t-5}{3t+5} \right| + c$$

42. Integrating by parts, taking $\cos d$ as the first function.

$$\begin{aligned} \therefore \int \cos d &= \int \cos d - \int \left[\frac{d}{d} (\cos d) \int \cos d \right] d \\ &= \sin d - \int 1 \cdot \sin d = \sin d - \int \sin d \\ &= \sin d + \cos d + c \end{aligned}$$

$$43. \int e^d dt = \int e^d - \int \left[\frac{d}{d} (e^d) \int e^d \right] dt \\ = e^d - \int 1 \cdot e^d dt \\ = e^d - e^d + c = e^d (-1) + c$$

$$44. \int \frac{d}{\cos^2 d} = \int \sec^2 d \\ = \tan d - \int 1 \cdot \tan d \\ = \tan d + \log |\cos d| + c$$



$$\begin{aligned}
45. \quad \int \sin 2x dx &= \int \sin 2x dx - \int \left[\frac{d}{dx} (\sin 2x) \int \sin 2x dx \right] dx \\
&= \left(-\frac{\cos 2x}{2} \right) - \int 1 \cdot \left(-\frac{\cos 2x}{2} \right) dx \\
&= -\frac{\cos 2x}{2} + \frac{1}{2} \cdot \frac{\sin 2x}{2} + c \\
&= \frac{\sin 2x}{4} - \frac{\cos 2x}{2} + c
\end{aligned}$$

$$\begin{aligned}
46. \quad \int x^2 \log x dx &= \log x \cdot \frac{x^3}{3} - \int \frac{1}{3} \cdot \frac{3x^2}{3} dx \\
&= \frac{1}{3} x^3 \cdot \log x - \frac{1}{3} \left[\frac{x^3}{3} \right] + c \\
&= \frac{1}{3} x^3 \log x - \frac{1}{9} x^3 + c
\end{aligned}$$

$$\begin{aligned}
47. \quad \int (-1)e^{-x} dx &= \frac{(-1)e^{-x}}{-1} - \int \frac{e^{-x}}{-1} dx \\
&= -e^{-x} + e^{-x} - e^{-x} + c \\
&= -e^{-x} + c
\end{aligned}$$

$$\begin{aligned}
48. \quad \text{Let } I &= \int e^x \sin x dx \\
&= \sin x \cdot e^x - \int \cos x \cdot e^x dx \\
&= e^x \sin x - \cos x \cdot e^x + \int (-\sin x) \cdot e^x dx \\
&= e^x \sin x - e^x \cos x - \int e^x \sin x dx
\end{aligned}$$

$$\therefore I = e^x \sin x - e^x \cos x - I$$

$$\therefore 2I = e^x (\sin x - \cos x) + c$$

$$\therefore I = \frac{e^x}{2} (\sin x - \cos x) + c$$

$$\begin{aligned}
49. \quad \int e^x (\sin x + \cos x) dx &= e^x \sin x + c \\
&\dots [\because \int e^x [f(x) + f'(x)] dx = e^x f(x) + c]
\end{aligned}$$

$$\begin{aligned}
50. \quad \int e^x (\sec x + \sec x \tan x) dx &= e^x \sec x + c \\
&\dots [\because \int e^x [f(x) + f'(x)] dx = e^x f(x) + c]
\end{aligned}$$

$$\begin{aligned}
51. \quad \int \frac{dx}{x^2-1} &= \int \left(\frac{1}{x-1} + \frac{1}{1-x} \right) dx \\
&= \log |x-1| - \log |1-x| + c
\end{aligned}$$

$$\begin{aligned}
52. \quad \int \frac{dx}{(x+1)(x+2)} &= \int \left(\frac{1}{x+1} - \frac{1}{x+2} \right) dx \\
&= \log |x+1| - \log |x+2| + c \\
&= \log \left| \frac{x+1}{x+2} \right| + c
\end{aligned}$$

$$\begin{aligned}
53. \quad \text{Let } \frac{-1}{(x-3)(x-2)} &= \frac{A}{x-3} + \frac{B}{x-2} \\
\therefore -1 &= A(x-2) + B(x-3) \quad \dots (i) \\
\text{Putting } x=2 \text{ in (i), we get} & \\
B &= -1 \\
\text{Putting } x=3 \text{ in (i), we get} & \\
A &= 2
\end{aligned}$$

$$\begin{aligned}
\therefore \int \frac{-1}{(x-3)(x-2)} dx &= \int \left(\frac{2}{x-3} - \frac{1}{x-2} \right) dx \\
&= 2 \log |x-3| - \log |x-2| + c \\
&= \log |(x-3)^2| - \log |x-2| + c
\end{aligned}$$

$$\begin{aligned}
54. \quad \int \frac{dx}{(x-2)(x-1)} &= -\int \frac{1}{x-1} dx + \int \frac{2}{x-2} dx \\
&= -\log |x-1| + 2 \log |x-2| + p \\
&= \log \left| \frac{(x-2)^2}{x-1} \right| + p
\end{aligned}$$

$$\begin{aligned}
55. \quad \int \frac{dx}{x^4+5x^2+4} &= \int \frac{dx}{(x^2+1)(x^2+4)} \\
&= \frac{1}{3} \int \left[\frac{1}{x^2+1} - \frac{1}{x^2+4} \right] dx \\
&= \frac{1}{3} \tan^{-1} x - \frac{1}{6} \tan^{-1} \left(\frac{x}{2} \right) + c
\end{aligned}$$

**Critical Thinking**

$$\begin{aligned}
1. \quad \int \frac{(x^3+3x^2+3x+1)}{(x+1)^5} dx &= \int \frac{(x+1)^3}{(x+1)^5} dx \\
&= \int \frac{1}{(x+1)^2} dx = \int (x+1)^{-2} dx \\
&= -\frac{1}{x+1} + c
\end{aligned}$$

$$\begin{aligned}
2. \quad \int (1+2x+3x^2+4x^3+\dots) dx & \\
&= \int (1-x)^{-2} dx \\
&= (1-x)^{-1} + c
\end{aligned}$$

$$3. \quad \int \left(1 + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \right) dx = \int e^x dx = e^x + c$$



4. Rationalizing the denominator, we get

$$\begin{aligned} & \int \frac{d}{\sqrt{x+3}-\sqrt{x+2}} \\ &= \int \frac{\sqrt{x+3}+\sqrt{x+2}}{(\sqrt{x+3}-\sqrt{x+2})(\sqrt{x+3}+\sqrt{x+2})} d \\ &= \int \frac{\sqrt{x+3}+\sqrt{x+2}}{(x+3)-(x+2)} d = \int \frac{\sqrt{x+3}+\sqrt{x+2}}{x+3-x-2} d \\ &= \int \left\{ (x+3)^{\frac{1}{2}} + (x+2)^{\frac{1}{2}} \right\} d \\ &= \frac{(x+3)^{\frac{3}{2}}}{\frac{3}{2}} + \frac{(x+2)^{\frac{3}{2}}}{\frac{3}{2}} + c \\ &= \frac{2}{3} \left[(x+3)^{\frac{3}{2}} + (x+2)^{\frac{3}{2}} \right] + c \end{aligned}$$

5.
$$\begin{aligned} \int \frac{-1}{(x+1)^2} dx &= \int \frac{x+1-2}{(x+1)^2} dx \\ &= \int \frac{1}{x+1} dx - \int \frac{2}{(x+1)^2} dx \\ &= \log|x+1| + \frac{2}{(x+1)} + c \end{aligned}$$

6. Since, the degree of the N^f is more than degree of the D^f , divide the N^f by D^f .

$$\begin{array}{r} \overline{2x^2-1} \\ 2x^2+1 \\ \hline -2+1 \\ -2-1 \\ \hline + \\ 2 \end{array}$$

$$\begin{aligned} \therefore \int \frac{x^4+1}{x^2+1} dx &= \int \left(x^2-1 + \frac{2}{x^2+1} \right) dx \\ &= \frac{x^3}{3} - x + 2 \tan^{-1} x + c \end{aligned}$$

7.
$$\begin{aligned} \int \frac{5(x^6+1)}{x^2+1} dx &= \int \frac{5(x^2+1)(x^4-x^2+1)}{(x^2+1)} dx \\ &= 5 \int (x^4-x^2+1) dx \\ &= 5 \left[\frac{x^5}{5} - \frac{x^3}{3} + x \right] + c \end{aligned}$$

8.
$$\begin{aligned} \int 2 \cdot 3^{x+1} \cdot 4^{x+2} dx &= 16 \times 3 \int 2 \cdot 3^x \cdot 4^x dx \\ &= 48 \int (24)^x dx = \frac{48(24)^x}{\log 24} + c \\ &= \frac{2 \cdot 3^{x+1} \cdot 4^{x+2}}{\log 2 + \log 4 + \log 3} + c \end{aligned}$$

9.
$$f(x) = \frac{1}{1-x}$$

$$\begin{aligned} \therefore f(f(f(x))) &= f\left(f\left(\frac{1}{1-x}\right)\right) \\ &= f\left(\frac{1}{1-\left(\frac{1}{1-x}\right)}\right) \\ &= f\left(\frac{1-x}{-}\right) \\ &= \frac{1}{1-\left(\frac{1-x}{-}\right)} = \frac{1}{1+\frac{1-x}{-}} \\ &= \frac{2}{-} + c \end{aligned}$$

∴ Required integral = $\frac{2}{-} + c$

10. Since, $a^{\log_a m} = m$

$$\begin{aligned} \therefore \int 9^{\log_3(\sec x)} dx &= \int \sec^2 x dx \quad \dots \left[\begin{array}{l} \because 3^{2\log_3(\sec x)} \\ = 3^{\log_3(\sec x)^2} \\ = (\sec x)^2 \end{array} \right] \\ &= \tan x + c \end{aligned}$$

11.
$$\begin{aligned} \int (e^{a \log x} + e^{\log_a x}) dx &= \int (e^{\log_e x^a} + e^{\log_e x^a}) dx \\ &= \int (x^a + x^a) dx \\ &= \frac{x^{a+1}}{a+1} + \frac{x^a}{\log a} + c \end{aligned}$$

12. Since, $\sec^2 x \cdot \operatorname{cosec}^2 x = \sec^2 x + \operatorname{cosec}^2 x$

$$\therefore \int \sec^2 x \cdot \operatorname{cosec}^2 x dx = \int \sec^2 x dx + \int \operatorname{cosec}^2 x dx = \tan x - \cot x + c$$

13.
$$\begin{aligned} \int (\sin^{-1} x + \cos^{-1} x) dx &= \int \left(\frac{\pi}{2}\right) dx \\ &\dots \left[\because \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2} \right] \\ &= \frac{\pi}{2} x + c = (\cos^{-1} x + \sin^{-1} x) + c \end{aligned}$$



$$14. \int \sin^{-1}(\cos x) dx = \int \left\{ \frac{\pi}{2} - \cos^{-1}(\cos x) \right\} dx$$

$$= \frac{\pi}{2}x - \frac{x^2}{2} = \frac{\pi x - x^2}{2}$$

$$15. \int (\cos x - \sin x) dx$$

$$= \sin x + \cos x + c$$

$$= \sqrt{2} \left(\frac{\sin x}{\sqrt{2}} + \frac{\cos x}{\sqrt{2}} \right) + c$$

$$= \sqrt{2} \sin \left(x + \frac{\pi}{4} \right) + c$$

$$\therefore \alpha = \frac{\pi}{4}$$

$$16. \int \sin 3x \cos 4x dx$$

$$= \frac{1}{2} \int 2 \sin 3x \cos 4x dx$$

$$= \frac{1}{2} \int [\sin (3x + 4x) + \sin (3x - 4x)] dx$$

$$[\because 2 \sin A \cos B = \sin (A + B) + \sin (A - B)]$$

$$= \frac{1}{2} (\int \sin 7x dx - \int \sin x dx)$$

$$= \frac{1}{2} \left(\frac{-\cos 7x}{7} + \cos x \right) + c$$

$$= \frac{-\cos 7x}{14} + \frac{\cos x}{2} + c$$

$$17. \int \frac{dx}{\tan x + \cot x} = \int \frac{dx}{\frac{\sin x}{\cos x} + \frac{\cos x}{\sin x}}$$

$$= \frac{1}{2} \int 2 \sin x \cos x dx = \frac{-\cos 2x}{4} + c$$

$$18. \int 2 \sin x \cos x dx$$

$$= \int \sin 2x dx = -\frac{\cos 2x}{2} + c_1$$

$$= -\frac{(1 - 2 \sin^2 x)}{2} + c_1$$

$$= -\frac{1}{2} + \sin^2 x + c_1$$

$$= \sin^2 x + c, \text{ where } c = \frac{-1}{2} + c_1$$

$$19. \int \sqrt{1 + \sin 2x} dx$$

$$= \int \sqrt{\sin^2 x + \cos^2 x + 2 \sin x \cos x} dx$$

$$\dots [\because \sin^2 x + \cos^2 x = 1]$$

$$= \int \sqrt{(\cos x + \sin x)^2} dx$$

$$= \int (\cos x + \sin x) dx$$

$$= \int \cos x dx + \int \sin x dx$$

$$= \sin x - \cos x + c$$

$$20. \int \frac{\sin x + \cos x}{\sqrt{1 + \sin 2x}} dx = \int \frac{\sin x + \cos x}{\sqrt{(\sin x + \cos x)^2}} dx$$

$$= \int \frac{dx}{1} = x + c$$

$$21. \sin^2 2x = (2 \sin x \cos x)^2$$

$$= 4 \sin^2 x \cos^2 x$$

$$\therefore 4 \int \frac{\sin^3 x + \cos^3 x}{\sin^2 2x} dx$$

$$= 4 \int \frac{\sin^3 x + \cos^3 x}{4 \sin^2 x \cos^2 x} dx$$

$$= \int \frac{\sin x}{\cos^2 x} dx + \int \frac{\cos x}{\sin^2 x} dx$$

$$= \int \tan x \sec x dx + \int \cot x \operatorname{cosec} x dx$$

$$= \sec x - \operatorname{cosec} x + c$$

$$22. \int \frac{dx}{\cos 2x + \sin^2 x} = \int \frac{dx}{\cos^2 x - \sin^2 x + \sin^2 x}$$

$$= \int \frac{dx}{\cos^2 x} = \int \sec^2 x dx = \tan x + c$$

$$23. \int \frac{\cos 2x - \cos 2\alpha}{\cos x - \cos \alpha} dx = \int \frac{2(\cos^2 x - \cos^2 \alpha)}{\cos x - \cos \alpha} dx$$

$$= 2 \int (\cos x + \cos \alpha) dx$$

$$= 2(\sin x + \cos \alpha x) + c$$

$$24. \text{ Since, } 1 + \cos 2x = 2 \cos^2 x$$

$$\therefore \int \sqrt{2 + \sqrt{2 + \sqrt{2 + 2 \cos 8x}}} dx$$

$$= \int \sqrt{2 + \sqrt{2 + 2 \cos 4x}} dx$$

$$= \int \sqrt{2 + 2 \cos 2x} dx$$

$$= \int 2 \cos x dx = 2 \sin x + c$$

$$25. \sin^4 x - \cos^4 x = (\sin^2 x - \cos^2 x)(\sin^2 x + \cos^2 x)$$

$$= -(\cos^2 x - \sin^2 x)(1)$$

$$= -\cos 2x$$

$$\therefore \int (\sin^4 x - \cos^4 x) dx = -\int \cos 2x dx$$

$$= -\frac{\sin 2x}{2} + c$$

$$= -\sin x \cos x + c$$



$$\begin{aligned}
 26. \quad & \int \frac{\sin^8 - \cos^8}{1 - 2\sin^2 \cos^2} d \\
 &= \int \frac{(\sin^4 + \cos^4)(\sin^4 - \cos^4)}{(\sin^2 + \cos^2)^2 - 2\sin^2 \cos^2} d \\
 &= \int (\sin^4 - \cos^4) d \\
 &= \int (\sin^2 + \cos^2)(\sin^2 - \cos^2) d \\
 &= \int (\sin^2 - \cos^2) d = \int -\cos 2 d \\
 &= -\frac{\sin 2}{2} + c
 \end{aligned}$$

$$\begin{aligned}
 27. \quad & \int \tan^{-1} \left(\sqrt{\frac{1-\cos 2}{1+\cos 2}} \right) d = \int \tan^{-1} \left(\sqrt{\frac{2\sin^2}{2\cos^2}} \right) d \\
 &= \int \tan^{-1} (\tan \frac{d}{2}) d \\
 &= \int \frac{d}{2} = \frac{d^2}{2} + c
 \end{aligned}$$

$$\begin{aligned}
 28. \quad & \int \tan^{-1} \left(\frac{\sin}{1+\cos} \right) d \\
 &= \int \tan^{-1} \left(\frac{2\sin \frac{d}{2} \cos \frac{d}{2}}{2\cos^2 \frac{d}{2}} \right) d \\
 &= \int \tan^{-1} \left(\tan \frac{d}{2} \right) d \\
 &= \int \frac{d}{2} \\
 &= \frac{1}{2} \cdot \frac{d^2}{2} + c \\
 &= \frac{d^2}{4} + c
 \end{aligned}$$

$$\begin{aligned}
 29. \quad & \int \frac{\cos 4 + 1}{\cot - \tan} d = \int \frac{(\cos 4 + 1)(\sin \cos)}{\cos^2 - \sin^2} d \\
 &= \int \frac{2\cos^2 2 (\sin \cos)}{\cos 2} d \\
 &= \int \cos 2 (2 \sin \cos) d \\
 &= \int \cos 2 \sin 2 d \\
 &= \frac{1}{2} \int \sin 4 d \\
 &= -\frac{1}{8} \cos 4 + c
 \end{aligned}$$

$$\therefore A = -\frac{1}{8}$$

$$\begin{aligned}
 30. \quad & \int \cos \frac{d}{16} \cos \frac{d}{8} \cos \frac{d}{4} \sin \frac{d}{16} d \\
 &= \int \cos \frac{d}{16} \sin \frac{d}{16} \cos \frac{d}{8} \cos \frac{d}{4} d \\
 &= \frac{1}{2} \int \sin \frac{d}{8} \cos \frac{d}{8} \cos \frac{d}{4} d \\
 &\quad \dots [\because \sin 2\theta = 2 \sin \theta \cos \theta] \\
 &= \frac{1}{4} \int \sin \frac{d}{4} \cos \frac{d}{4} d \\
 &= \frac{1}{8} \int \sin \frac{d}{2} d \\
 &= \left(\frac{2}{8} \right) \left(-\cos \frac{d}{2} \right) + c \\
 &= -\frac{1}{4} \cos \frac{d}{2} + c
 \end{aligned}$$

$$\begin{aligned}
 31. \quad & \int \frac{\sin^6 + \cos^6}{\sin^2 \cos^2} d \\
 &= \int \frac{(\sin^2 + \cos^2)^3 - 3\sin^2 \cos^2 (\sin^2 + \cos^2)}{\sin^2 \cos^2} d \\
 &\quad \dots \left[\begin{array}{l} \because a^3 + b^3 = (a+b)^3 - 3ab(a+b) \\ \text{and } \sin^6 = (\sin^2)^3 \end{array} \right] \\
 &= \int \frac{1 - 3\sin^2 \cos^2}{\sin^2 \cos^2} d \\
 &= \int \left(\frac{1}{\sin^2 \cos^2} - 3 \right) d \\
 &= \int \left(\frac{\sin^2 + \cos^2}{\sin^2 \cos^2} - 3 \right) d \\
 &= \int \left(\frac{1}{\cos^2} + \frac{1}{\sin^2} - 3 \right) d \\
 &= \int \sec^2 d + \int \operatorname{cosec}^2 d - 3 \int d \\
 &= \tan - \cot - 3d + c
 \end{aligned}$$

$$\begin{aligned}
 32. \quad & \int \frac{\cot \tan}{\sec^2 - 1} d = \int \frac{1}{\tan^2} d \\
 &= \int \cot^2 d \\
 &= \int (\operatorname{cosec}^2 - 1) d \\
 &= -\cot - d + c
 \end{aligned}$$

$$\begin{aligned}
 33. \quad & \int \frac{1+\cos^2}{\sin^2} d = \int (\operatorname{cosec}^2 + \cot^2) d \\
 &= \int (2\operatorname{cosec}^2 - 1) d \\
 &\quad \dots [\because 1 + \cot^2 \theta = \operatorname{cosec}^2 \theta] \\
 &= -2 \cot - d + c
 \end{aligned}$$



$$\begin{aligned}
 34. \quad 2 \int \frac{1 + \cos 4}{1 - \cos 4} d &= 2 \int \frac{\cos^2 2}{\sin^2 2} d \\
 &= 2 \int \cot^2 2 d \\
 &= 2 \int (\operatorname{cosec}^2 2 - 1) d \\
 &= 2 \left(-\frac{\cot 2}{2} \right) - 2 + c \\
 &= -\cot 2 - 2 + c
 \end{aligned}$$

$$\begin{aligned}
 35. \quad \int \left(\frac{1 + \tan}{1 - \tan} \right)^2 d &= \int \left(\tan \left(\frac{\pi}{4} + \right) \right)^2 d \\
 &= \int \left(\sec^2 \left(\frac{\pi}{4} + \right) - 1 \right) d \\
 &= \tan \left(\frac{\pi}{4} + \right) - + c
 \end{aligned}$$

$$\begin{aligned}
 36. \quad \int (\sec + \tan)^2 d &= \int (\sec^2 + \tan^2 + 2 \sec \tan) d \\
 &= \int (2 \sec^2 - 1 + 2 \sec \tan) d \\
 &= 2 \tan + 2 \sec - + c \\
 &= 2 (\sec + \tan) - + c
 \end{aligned}$$

$$\begin{aligned}
 37. \quad \int \frac{\tan}{(\sec + \tan)} d &= \int \frac{\tan (\sec - \tan)}{(\sec + \tan)(\sec - \tan)} d \\
 &= \int \frac{\tan (\sec - \tan)}{(\sec^2 - \tan^2)} d \\
 &= \int (\sec \tan - \tan^2) d \\
 &= \int \sec \tan d - \int (\sec^2 - 1) d \\
 &= \int \sec \tan d - \int \sec^2 d + \int 1 d \\
 &= \sec - \tan + + c
 \end{aligned}$$

$$\begin{aligned}
 38. \quad \text{Put } t = 3 - 5 \Rightarrow dt = 3d \\
 \therefore \int \tan(3 - 5) \sec(3 - 5) d &= \frac{1}{3} \int \tan t \cdot \sec t dt \\
 &= \frac{\sec t}{3} + c = \frac{\sec(3 - 5)}{3} + c
 \end{aligned}$$

$$\begin{aligned}
 39. \quad \text{Put } f(\) = t \Rightarrow f'(\) d &= dt \\
 \therefore \int \frac{f'(\)}{[f(\)]^2} d &= \int \frac{1}{t^2} dt = -\frac{1}{t} + c = -\frac{1}{f(\)} + c
 \end{aligned}$$

$$\begin{aligned}
 40. \quad \text{Put } 10 + 10 = t \Rightarrow (10^9 + 10 \log_e 10) d &= dt \\
 \therefore \int \frac{10^9 + 10 \log_e 10}{10 + 10} d &= \int \frac{1}{t} dt = \log t + c \\
 &= \log(10 + 10) + c
 \end{aligned}$$

$$\begin{aligned}
 41. \quad \text{Put } t^2 - 4 + 3 = t \\
 \Rightarrow (2t - 4) dt = dt \Rightarrow (2t - 4) dt = \frac{1}{2} dt \\
 \therefore \int \frac{-2}{t^2 - 4 + 3} dt = \frac{1}{2} \int \frac{dt}{t} = \frac{1}{2} \log t + c \\
 = \frac{1}{2} \log(t^2 - 4 + 3) + c \\
 = \log(\sqrt{t^2 - 4 + 3}) + c
 \end{aligned}$$

$$\begin{aligned}
 42. \quad \text{Put } 5^7 = t \\
 \Rightarrow 35^6 d = dt \\
 \Rightarrow 6 d = \frac{dt}{35} \\
 \therefore \int 6 \sin(5^7) d = \int \sin t \cdot \frac{dt}{35} \\
 = -\frac{\cos t}{35} = \frac{-\cos(5^7)}{35}
 \end{aligned}$$

$$\therefore k = -\frac{1}{7}$$

$$\begin{aligned}
 43. \quad \int \frac{\sin}{\sin(-\alpha)} d &= \int \frac{\sin(-\alpha + \alpha)}{\sin(-\alpha)} d \\
 &= \int \frac{\{\sin(-\alpha)\cos\alpha + \cos(-\alpha)\sin\alpha\}}{\sin(-\alpha)} d \\
 &= \int \cos \alpha d + \int \sin \alpha \cot(-\alpha) d \\
 &= \cos \alpha + \sin \alpha \cdot \log |\sin(-\alpha)| + c
 \end{aligned}$$

$$\begin{aligned}
 44. \quad \int \frac{\cos(\alpha + \alpha)}{\cos} d &= \int \left[\frac{\cos \cos \alpha - \sin \sin \alpha}{\cos} \right] d \\
 &= \int (\cos \alpha - \sin \alpha \tan) d \\
 &= (\cos \alpha) - \sin \alpha \log |\sec| + c
 \end{aligned}$$

$$\begin{aligned}
 45. \quad \int \frac{1}{\sqrt{1 + \cos}} d &= \int \frac{d}{\sqrt{2 \cos^2 \left(\frac{\ }{2} \right)}} = \frac{1}{\sqrt{2}} \int \sec \frac{d}{2} \\
 &= \frac{1}{\sqrt{2}} \left\{ \log \left| \sec \frac{d}{2} + \tan \frac{d}{2} \right| \right\} \cdot \frac{1}{1/2} + c \\
 &= \sqrt{2} \log \left| \sec \frac{d}{2} + \tan \frac{d}{2} \right| + c
 \end{aligned}$$

$$\begin{aligned}
 46. \quad \int \frac{d}{4 \cos^3 2 - 3 \cos 2} &= \int \frac{d}{\cos 6} = \int \sec 6 d \\
 &= \frac{1}{6} \log |\sec 6 + \tan 6| + c
 \end{aligned}$$



$$\begin{aligned}
 47. \int \frac{d}{\sin \frac{d}{2} + \sqrt{3} \cos \frac{d}{2}} &= \frac{1}{2} \int \frac{d}{\frac{\sin \frac{d}{2}}{2} + \frac{\sqrt{3}}{2} \cos \frac{d}{2}} \\
 &= \frac{1}{2} \int \frac{d}{\sin \frac{d}{2} \cos \frac{\pi}{3} + \sin \frac{\pi}{3} \cos \frac{d}{2}} \\
 &= \frac{1}{2} \int \frac{d}{\sin \left(\frac{d}{2} + \frac{\pi}{3} \right)} \\
 &= \frac{1}{2} \int \operatorname{cosec} \left(\frac{d}{2} + \frac{\pi}{3} \right) d \\
 &= \frac{1}{2} \log \left| \tan \left(\frac{d}{2} + \frac{\pi}{6} \right) \right| + c
 \end{aligned}$$

$$\begin{aligned}
 48. \int \frac{\sin 2}{\sin 3 \sin 5} d &= \int \frac{\sin(5-3)}{\sin 3 \sin 5} d \\
 &= \int \frac{\sin 5 \cos 3 - \cos 5 \sin 3}{\sin 3 \sin 5} d \\
 &= \int \cot 3 d - \int \cot 5 d \\
 &= \frac{1}{3} \log |\sin 3| - \frac{1}{5} \log |\sin 5| + c
 \end{aligned}$$

$$\begin{aligned}
 49. \text{ Let } I &= \int \sqrt{\frac{1+t}{1-t}} d = \int \frac{1+t}{\sqrt{1-t^2}} d \\
 &= \int \frac{d}{\sqrt{1-t^2}} + \int \frac{t}{\sqrt{1-t^2}} d \\
 \text{ In 2}^{\text{nd}} \text{ integral, put } 1-t^2 &= t \Rightarrow -2 dt = dt \\
 \therefore I &= \int \frac{d}{\sqrt{1-t^2}} - \frac{1}{2} \int \frac{dt}{t^2} \\
 &= \sin^{-1} t - \sqrt{t} + c \\
 &= \sin^{-1} \sqrt{1-t^2} + c
 \end{aligned}$$

$$\begin{aligned}
 50. \text{ Put } t &= 1 + \log x \Rightarrow dt = \left(1 + \frac{1}{x} \right) dx \\
 \therefore \int \frac{(1+\log x)^2}{x} dx &= \int t^2 dt = \frac{t^3}{3} + c \\
 &= \frac{1}{3} (1 + \log x)^3 + c
 \end{aligned}$$

$$\begin{aligned}
 51. \text{ Put } \log(\log x) &= t \\
 \Rightarrow \frac{1}{\log x} dx &= dt \\
 \therefore \int \frac{dx}{\log x \log(\log x)} &= \int \frac{dt}{t} = \log |t| + c \\
 &= \log |\log(\log x)| + c
 \end{aligned}$$

$$\begin{aligned}
 52. \text{ Let } I &= \int \frac{1}{3} [\log x]^2 dx = \int \frac{1}{3} [\log x]^2 dx \\
 &= \int \frac{1}{2} (\log x)^2 dx
 \end{aligned}$$

$$\text{ Put } \log x = t \Rightarrow \frac{1}{x} dx = dt$$

$$\therefore I = \int t^2 dt = \frac{t^3}{3} + c = \frac{(\log x)^3}{3} + c$$

$$53. \frac{dI}{dx} = 3^{\cos x} \sin x$$

$$\therefore I = \int 3^{\cos x} \sin x \cdot dx$$

$$\text{ Put } \cos x = t$$

$$\Rightarrow -\sin x dx = dt \Rightarrow \sin x dx = -dt$$

$$\therefore I = -\int 3^t dt = \frac{-3^t}{\log 3} + c = \frac{-3^{\cos x}}{\log 3} + c$$

$$54. \text{ Put } a^x = t$$

$$\Rightarrow a \log a dx = dt \Rightarrow a dx = \frac{1}{\log a} dt$$

$$\begin{aligned}
 \therefore \int a^x \cdot a dx &= \frac{1}{\log a} \int a^t dt = \frac{1}{\log a} \cdot a^t \cdot \frac{1}{\log a} + c \\
 &= \frac{a^x}{(\log a)^2} + c
 \end{aligned}$$

$$55. \text{ Put } 2e^{-x} + 5 = t \Rightarrow -2e^{-x} dx = dt$$

$$\begin{aligned}
 \therefore \int e^{-x} \operatorname{cosec}^2(2e^{-x} + 5) dx &= -\frac{1}{2} \int \operatorname{cosec}^2 t dt \\
 &= \frac{1}{2} \cot t + c \\
 &= \frac{1}{2} \cot(2e^{-x} + 5) + c
 \end{aligned}$$

$$56. \text{ Let } I = \int \frac{dx}{1+e^x} = \int \frac{e^{-x}}{1+e^{-x}} dx$$

$$\text{ Put } 1+e^{-x} = t \Rightarrow e^{-x} dx = -dt$$

$$\therefore I = -\int \frac{dt}{t} = -\log |t| + c = -\log |1+e^{-x}| + c$$

$$\begin{aligned}
 57. \text{ Let } I &= \int \frac{1}{(e^x + e^{-x})^2} dx \\
 &= \int \frac{e^{2x}}{(e^4 + 1)^2} dx
 \end{aligned}$$

$$\text{ Put } e^4 + 1 = t \Rightarrow 4e^4 dx = dt$$

$$\begin{aligned}
 \therefore I &= \frac{1}{4} \int \frac{1}{t^2} dt \\
 &= \frac{1}{4} \left(\frac{-1}{t} \right) + c = \frac{-1}{4(e^4 + 1)} + c
 \end{aligned}$$



58. Put $\log t = t \Rightarrow t = e^t \Rightarrow dt = e^t dt$
 $\therefore \int \frac{\log t}{(1 + \log t)^2} dt = \int \frac{t}{(1+t)^2} e^t dt$
 $= \int e^t \left[\frac{t+1-1}{(1+t)^2} \right] dt$
 $= \int e^t \left[\frac{1}{1+t} - \frac{1}{(1+t)^2} \right] dt$
 $= \frac{e^t}{1+t} + c$
 $= \frac{e^{\log t}}{1 + \log t} + c$

59. Put $e^t = t$
 $\Rightarrow (e^t + e^t) dt = dt \Rightarrow e^t (1 + t) dt = dt$
 $\therefore \int \frac{e^t (1+t)}{\sin(e^t)} dt = \int \frac{dt}{\sin t} = \int \operatorname{cosec} t dt$
 $= \log \left| \tan \left(\frac{t}{2} \right) \right| + c$
 $= \log \left| \tan \left(\frac{e^t}{2} \right) \right| + c$

60. Put $\log \left(\tan \frac{t}{2} \right) = t$
 $\Rightarrow \frac{1}{\tan \frac{t}{2}} \cdot \frac{1}{2} \sec^2 \frac{t}{2} dt = dt \Rightarrow \operatorname{cosec} \frac{t}{2} dt = dt$
 $\therefore \int \frac{\operatorname{cosec} \frac{t}{2}}{\log \left(\tan \frac{t}{2} \right)} dt = \int \frac{1}{t} dt = \log |t| + c$
 $= \log \left| \log \left(\tan \frac{t}{2} \right) \right| + c$

61. Put $\tan^{-1} (t^3) = t$
 $\Rightarrow \frac{1}{1+(t^3)^2} \cdot 3t^2 dt = dt \Rightarrow \frac{3t^2}{1+t^6} dt = \frac{dt}{3}$
 $\therefore \int \frac{3t^2 \tan^{-1} (t^3)}{1+t^6} dt = \frac{1}{3} \int t dt = \frac{1}{3} \cdot \frac{t^2}{2} + c$
 $= \frac{(\tan^{-1} t^3)^2}{6} + c$

62. Put $\tan \sqrt{t} = t \Rightarrow \frac{\sec^2 \sqrt{t}}{2\sqrt{t}} dt = dt$
 $\therefore \int \frac{1}{\sqrt{t}} \tan^4 \sqrt{t} \cdot \sec^2 \sqrt{t} dt = 2 \int t^4 dt$
 $= \frac{2t^5}{5} + c = \frac{2}{5} \tan^5 \sqrt{t} + c$

63. Put $e^t = t \Rightarrow e dt = dt$
 $\therefore \int e \tan^2(e) dt = \int \tan^2 t dt = \int (\sec^2 t - 1) dt$
 $= \tan t - t + c$
 $= \tan(e) - e + c$

64. Let $I = \int \frac{t^2+1}{(t^2-1)} dt = \int \frac{1+\frac{1}{2}}{\frac{1}{2}} dt$
Put $\frac{1}{2} = t \Rightarrow \left(1 + \frac{1}{2}\right) dt = dt$
 $\therefore I = \int \frac{dt}{t} = \log t + c = \log \left(\frac{1}{2} \right) + c$
 $= \log \left(\frac{t^2-1}{2} \right) + c$

65. Let $I = \int \frac{(t^4-1)^{\frac{1}{4}} dt}{5}$
 $= \int \frac{\left(1 - \frac{1}{3}\right)^{\frac{1}{4}} dt}{5} = \int \frac{\left(1 - \frac{1}{3}\right)^{\frac{1}{4}} dt}{4}$
Put $1 - \frac{1}{3} = t \Rightarrow \frac{3}{4} dt = dt$
 $\therefore I = \int t^{\frac{1}{4}} \cdot \frac{dt}{3} = \frac{t^{\frac{5}{4}}}{\frac{5}{4}} \cdot \frac{1}{3} + c$
 $= \frac{4}{15} t^{\frac{5}{4}} + c = \frac{4}{15} \left(1 - \frac{1}{3}\right)^{\frac{5}{4}} + c$

66. Let $I = \int \frac{d}{(d^7+1)} = \int \frac{d}{8 \left(1 + \frac{1}{7}\right)}$
Put $1 + \frac{1}{7} = t \Rightarrow \frac{-7}{8} dt = dt$
 $\therefore I = \frac{-1}{7} \int \frac{dt}{t} = \frac{-1}{7} \log |t| + c$
 $= \frac{-1}{7} \log \left| \frac{d^7+1}{7} \right| + c$
 $= \frac{1}{7} \log \left| \frac{d^7}{d^7+1} \right| + c$



$$\begin{aligned}
 67. \quad \int \frac{1}{(x^n + 1)} dx &= \int \frac{dx}{x^{n+1} \left(1 + \frac{1}{x^n}\right)} \\
 &= \frac{-1}{n} \int \frac{x^{-n-1}}{(1 + x^{-n})} dx \\
 &= \frac{-1}{n} \log \left| 1 + \frac{1}{x^n} \right| + c \\
 &= \frac{-1}{n} \log \left| \frac{x^n + 1}{x^n} \right| + c \\
 &= \frac{1}{n} \log \left| \frac{x^n}{x^n + 1} \right| + c
 \end{aligned}$$

$$\begin{aligned}
 68. \quad \text{Let } I &= \int \frac{1}{\cos^2 (1 - \tan^2)^2} dx = \int \frac{\sec^2 dx}{(\tan^2 - 1)^2} \\
 \text{Put } \tan^2 - 1 &= t \Rightarrow \sec^2 dx = dt \\
 \therefore I &= \int \frac{1}{t^2} dt = -\frac{1}{t} + c = \frac{-1}{\tan^2 - 1} + c = \frac{1}{1 - \tan^2} + c
 \end{aligned}$$

$$\begin{aligned}
 69. \quad \text{Put } x &= \tan \theta \Rightarrow dx = \sec^2 \theta d\theta \\
 \therefore \int \frac{1}{\sqrt{1+x^2}} dx &= \int \frac{\sec^2 \theta d\theta}{\tan^2 \theta \sec \theta} \\
 &= \int \operatorname{cosec} \theta \cot \theta d\theta = -\operatorname{cosec} \theta + c \\
 &= \frac{-\sqrt{\tan^2 \theta + 1}}{\tan \theta} + c = \frac{-\sqrt{1+x^2}}{x} + c
 \end{aligned}$$

$$\begin{aligned}
 70. \quad \int \frac{x^3}{\sqrt{1+x^4}} dx &= \frac{1}{4} \int \frac{4x^3}{\sqrt{1+x^4}} dx \\
 &= \frac{1}{4} \int \frac{dt}{t^{1/2}} \\
 &\dots [\text{Put } 1+x^4 = t \Rightarrow 4x^3 dx = dt] \\
 &= \frac{1}{4} \cdot \frac{t^{-1/2+1}}{-1/2+1} + c \\
 &= \frac{1}{2} \sqrt{t} + c = \frac{1}{2} \sqrt{1+x^4} + c
 \end{aligned}$$

$$\begin{aligned}
 71. \quad \text{Let } I &= \int \frac{\sin 2x}{\sin^2 x + 2\cos^2 x} dx = \int \frac{2\sin x \cos x}{1 + \cos^2 x} dx \\
 \text{Put } 1 + \cos^2 x &= t \Rightarrow -2\sin x \cos x dx = dt \\
 \therefore I &= \int -\left(\frac{dt}{t}\right) = -\log |t| + c \\
 &= -\log |1 + \cos^2 x| + c
 \end{aligned}$$

$$\begin{aligned}
 72. \quad \text{Let } I &= \int \frac{\cos 2x}{(\cos^2 x + \sin^2 x)^2} dx \\
 &= \int \frac{(\cos x - \sin x)(\cos x + \sin x) dx}{(\cos^2 x + \sin^2 x)^2} \\
 &= \int \frac{\cos x - \sin x}{\cos^2 x + \sin^2 x} dx \\
 \text{Put } t &= \sin x + \cos x \\
 \Rightarrow dt &= (\cos x - \sin x) dx \\
 \therefore I &= \int \frac{1}{t} dt = \log |t| + c = \log |\sin x + \cos x| + c
 \end{aligned}$$

$$\begin{aligned}
 73. \quad \text{Put } 3\sin^2 x + 5\cos^2 x &= t \\
 \Rightarrow (3 \times 2 \sin x \cos x - 5 \times 2 \sin x \cos x) dx &= dt \\
 \Rightarrow -4 \sin x \cos x dx &= dt \\
 \Rightarrow \sin x \cos x dx &= \frac{dt}{-4} \\
 \therefore \int \frac{\sin x \cos x}{3\sin^2 x + 5\cos^2 x} dx &= \int \frac{dt}{(-4)t} \\
 &= -\frac{1}{4} \int \frac{1}{t} dt = -\frac{1}{4} \log |t| + c \\
 &= -\frac{1}{4} \log |3\sin^2 x + 5\cos^2 x| + c
 \end{aligned}$$

$$\begin{aligned}
 74. \quad \text{Let } I &= \int \frac{\cos^2 x + \sin^2 x}{2 + \cos x} dx = \int \frac{\cos^2 x + \sin^2 x}{1 + \frac{\cos x}{2}} dx \\
 \text{Put } 1 + \frac{\cos x}{2} &= t \\
 \Rightarrow \frac{-(\sin x + \cos x)}{2} dx &= dt
 \end{aligned}$$

$$\begin{aligned}
 \therefore I &= -\int \frac{dt}{t} = -\log |t| + c \\
 &= -\log \left| \frac{\cos x + 2}{2} \right| + c \\
 &= \log \left| \frac{2}{\cos x + 2} \right| + c
 \end{aligned}$$

$$\begin{aligned}
 75. \quad \text{Let } I &= \int \frac{\log \left(\frac{x + \sqrt{1+x^2}}{\sqrt{1+x^2}} \right)}{\sqrt{1+x^2}} dx \\
 \text{Put } \log \left(\frac{x + \sqrt{1+x^2}}{\sqrt{1+x^2}} \right) &= t \\
 \Rightarrow \frac{1 + \frac{2}{2\sqrt{1+x^2}}}{x + \sqrt{1+x^2}} dx &= dt \Rightarrow \frac{dx}{\sqrt{1+x^2}} = dt \\
 \therefore I &= \int t dt = \frac{t^2}{2} + c = \frac{\left[\log \left(\frac{x + \sqrt{1+x^2}}{\sqrt{1+x^2}} \right) \right]^2}{2} + c
 \end{aligned}$$



$$\begin{aligned}
76. \quad & \text{Put } (\) = t \Rightarrow \log(\) = \log t \\
& \Rightarrow {}^2 \log = \log t \\
& \Rightarrow (2 \log +)d = \frac{1}{t} \cdot dt \\
& \Rightarrow (2 \log + 1) (\) d = dt \\
\therefore & \int (\) (2 \log + 1) d = \int dt = t + c = (\) + c
\end{aligned}$$

$$\begin{aligned}
77. \quad & 1 + 2 \tan (\sec + \tan) \\
& = 1 + 2 \tan \cdot \sec + 2 \tan^2 \\
& = (1 + \tan^2) + 2 \sec \cdot \tan + \tan^2 \\
& = \sec^2 + 2 \sec \cdot \tan + \tan^2 \\
& = (\sec + \tan)^2 \\
\therefore & \int \sqrt{1 + 2 \tan (\sec + \tan)} d \\
& = \int (\sec + \tan) d \\
& = \int \frac{1 + \sin}{\cos} d \\
& = \int \frac{1 - \sin^2}{\cos (1 - \sin)} d \\
& = - \int \frac{(-\cos)}{1 - \sin} d = - \log |1 - \sin| + c
\end{aligned}$$

$$\begin{aligned}
78. \quad & \text{Put } \log = t \Rightarrow \frac{1}{\ } d = dt \\
\therefore & \int \frac{d}{\sqrt{1 - (\log)^2}} = \int \frac{dt}{\sqrt{1 - t^2}} = \sin^{-1} t + c \\
& = \sin^{-1} (\log) + c
\end{aligned}$$

$$\begin{aligned}
79. \quad & \text{Put } t = \cos \Rightarrow dt = -\sin \ d \\
\therefore & \int \frac{\sin}{\sqrt{4 - \cos^2}} d = - \int \frac{dt}{\sqrt{2^2 - t^2}} \\
& = - \sin^{-1} \left(\frac{t}{2} \right) + c = - \sin^{-1} \left(\frac{\cos}{2} \right) + c
\end{aligned}$$

$$\begin{aligned}
80. \quad & \int \frac{\sec d}{\sqrt{\cos 2}} = \int \frac{\sec}{\sqrt{\cos^2 - \sin^2}} d \\
& = \int \frac{\sec^2 d}{\sqrt{1 - \tan^2}} \\
& \dots [\text{Multiplying } N^r \text{ and } D^r \text{ by } \sec \]
\end{aligned}$$

$$\begin{aligned}
& \text{Put } \tan = t \Rightarrow \sec^2 d = dt \\
\therefore & \int \frac{\sec d}{\sqrt{\cos 2}} = \int \frac{dt}{\sqrt{1 - t^2}} = \sin^{-1} t + c \\
& = \sin^{-1} (\tan) + c
\end{aligned}$$

$$\begin{aligned}
81. \quad & \text{Put } 2 = \sin \theta \Rightarrow 2d = \cos \theta d \theta \\
\therefore & \int \frac{2d}{\sqrt{1 - 4^2}} = \int \frac{\cos \theta d \theta}{\sqrt{1 - \sin^2 \theta}} = \int \frac{\cos \theta}{\cos \theta} d \theta \\
& = \int d \theta = \theta + c \\
& = \sin^{-1} (2) + c
\end{aligned}$$

$$\begin{aligned}
82. \quad & \int \frac{d}{\sqrt{2 - 3 -^2}} = \int \frac{d}{\sqrt{\left(\frac{17}{4}\right) - \left(+\frac{3}{2}\right)^2}} \\
& = \int \frac{d}{\sqrt{\left(\frac{\sqrt{17}}{2}\right)^2 - \left(+\frac{3}{2}\right)^2}} \\
& = \sin^{-1} \left[\frac{\left(+\frac{3}{2}\right)}{\left(\frac{\sqrt{17}}{2}\right)} \right] + c \\
& = \sin^{-1} \left(\frac{2 + 3}{\sqrt{17}} \right) + c
\end{aligned}$$

$$\begin{aligned}
83. \quad & \text{Put } \sin = t \Rightarrow \cos d = dt \\
\therefore & \int \cos \sqrt{4 - \sin^2} d = \int \sqrt{4 - t^2} dt \\
& = \int \sqrt{(2)^2 - t^2} dt = \frac{t}{2} \sqrt{4 - t^2} + \frac{4}{2} \sin^{-1} \left(\frac{t}{2} \right) + c \\
& = \frac{1}{2} \sin \sqrt{4 - \sin^2} + 2 \sin^{-1} \left(\frac{1}{2} \sin \right) + c
\end{aligned}$$

$$84. \quad \text{Let } I = \int \frac{3^2}{\sqrt{9 - 16^6}} d = \int \frac{3^2}{\sqrt{(3)^2 - (4^3)^2}} d$$

$$\begin{aligned}
& \text{Put } 4^3 = t \\
& \Rightarrow 12^2 d = dt
\end{aligned}$$

$$\begin{aligned}
\therefore & I = \frac{1}{4} \int \frac{dt}{\sqrt{(3)^2 - t^2}} \\
& = \frac{1}{4} \sin^{-1} \left(\frac{t}{3} \right) + c \\
& = \frac{1}{4} \sin^{-1} \left(\frac{4^3}{3} \right) + c
\end{aligned}$$

$$\begin{aligned}
85. \quad & \int \sqrt{\frac{a-}{a+}} d = \int \frac{a-}{\sqrt{a^2 -^2}} d \\
& = \int \left(\frac{a}{\sqrt{a^2 -^2}} - \frac{1}{\sqrt{a^2 -^2}} \right) d \\
& = a \int \frac{1}{\sqrt{a^2 -^2}} d + \frac{1}{2} \int \frac{-2}{\sqrt{a^2 -^2}} d \\
& = a \cdot \sin^{-1} \left(\frac{-}{a} \right) + \frac{1}{2} \cdot 2 \sqrt{a^2 -^2} + c \\
& = a \sin^{-1} \left(\frac{-}{a} \right) + \sqrt{a^2 -^2} + c
\end{aligned}$$



$$86. \text{ Let } I = \int \frac{2}{\sqrt{1-(2)^2}} d$$

$$\text{Put } 2 = t \Rightarrow 2 \, d = \frac{dt}{\log 2}$$

$$\begin{aligned} \therefore I &= \frac{1}{\log 2} \int \frac{dt}{\sqrt{1-t^2}} = \frac{1}{\log 2} \sin^{-1} t + c \\ &= \frac{1}{\log 2} \sin^{-1} 2 + c \end{aligned}$$

$$\therefore K = \frac{1}{\log 2}$$

$$\begin{aligned} 87. \int \frac{\sqrt{\quad}}{1+\quad} d &= \int \frac{\sqrt{\quad} \cdot \sqrt{\quad}}{\sqrt{\quad}(1+\quad)} d \\ &= \int \frac{+1}{\sqrt{\quad}(+\quad)} d - \int \frac{1}{\sqrt{\quad}(+\quad)} d \\ &= \int \frac{1}{\sqrt{\quad}} d - \int \frac{1}{\sqrt{\quad}[1+(\sqrt{\quad})^2]} d \\ &= 2\sqrt{\quad} - 2\tan^{-1}\sqrt{\quad} + c \\ &= 2(\sqrt{\quad} - \tan^{-1}\sqrt{\quad}) + c \end{aligned}$$

$$88. \text{ Let } I = \int \frac{e^{\log\left(1+\frac{1}{2}\right)}}{2+\frac{1}{2}} d = \int \frac{1+\frac{1}{2}}{\left(\frac{-1}{-}\right)^2+2} d$$

.... $[\because e^{\log a} = a]$

$$\text{Put } 1 + \frac{1}{2} = t$$

$$\Rightarrow \left(1 + \frac{1}{2}\right) d = dt$$

$$\begin{aligned} \therefore I &= \int \frac{dt}{t^2+2} = \frac{1}{\sqrt{2}} \tan^{-1}\left(\frac{t}{\sqrt{2}}\right) + c \\ &= \frac{1}{\sqrt{2}} \tan^{-1}\left(\frac{1+\frac{1}{2}}{\sqrt{2}}\right) + c \\ &= \frac{1}{\sqrt{2}} \tan^{-1}\left(\frac{2+1}{\sqrt{2}}\right) + c \end{aligned}$$

$$89. \text{ Put } t^2 = t \Rightarrow d = \frac{1}{2} dt$$

$$\begin{aligned} \therefore \int \frac{1}{t^4 + t^2 + 1} d &= \frac{1}{2} \int \frac{dt}{t^2 + t + 1} \\ &= \frac{1}{2} \int \frac{dt}{t^2 + t + \frac{1}{4} + \frac{3}{4}} \\ &= \frac{1}{2} \int \frac{dt}{(t+1/2)^2 + \left(\frac{\sqrt{3}}{2}\right)^2} \\ &= \frac{1}{2} \cdot \frac{1}{(\sqrt{3}/2)} \tan^{-1}\left(\frac{t+1/2}{\sqrt{3}/2}\right) + c \\ &= \frac{1}{\sqrt{3}} \tan^{-1}\left(\frac{2t+1}{\sqrt{3}}\right) + c \\ &= \frac{1}{\sqrt{3}} \tan^{-1}\left(\frac{2t^2+1}{\sqrt{3}}\right) + c \end{aligned}$$

$$\begin{aligned} 90. \text{ Let } I &= \int \frac{1}{1+\sin^2} d = \int \frac{d}{2\sin^2 + \cos^2} \\ &= \int \frac{\sec^2 d}{2\tan^2 + 1} \\ &= \frac{1}{2} \int \frac{\sec^2 d}{\tan^2 + \frac{1}{2}} \end{aligned}$$

$$\text{Put } \tan = t \Rightarrow \sec^2 d = dt$$

$$\begin{aligned} \therefore I &= \frac{1}{2} \int \frac{dt}{t^2 + \frac{1}{2}} = \frac{1}{2} \cdot \frac{1}{\sqrt{2}} \tan^{-1} \frac{t}{\frac{1}{\sqrt{2}}} + c \\ &= \frac{1}{\sqrt{2}} \tan^{-1}(\sqrt{2} \tan) + c \end{aligned}$$

$$91. \int \frac{1}{1+\cos^2} d = \int \frac{\sec^2}{\sec^2 + 1} d = \int \frac{\sec^2}{\tan^2 + 2} d$$

$$\text{Put } \tan = t \Rightarrow \sec^2 d = dt$$

$$\begin{aligned} \therefore \int \frac{1}{1+\cos^2} d &= \int \frac{dt}{t^2+2} = \frac{1}{\sqrt{2}} \tan^{-1}\left(\frac{t}{\sqrt{2}}\right) + c \\ &= \frac{1}{\sqrt{2}} \tan^{-1}\left(\frac{1}{\sqrt{2}} \tan\right) + c \end{aligned}$$



92. Put $\cos = t$
 $\Rightarrow -\sin d = dt$
 $\therefore \int \frac{\sin}{3+4\cos^2} d = \int \frac{-dt}{3+4t^2} = \frac{-1}{4} \int \frac{dt}{t^2 + \left(\frac{\sqrt{3}}{2}\right)^2}$
 $= -\frac{1}{4 \cdot \frac{\sqrt{3}}{2}} \cdot \tan^{-1} \frac{t}{\left(\frac{\sqrt{3}}{2}\right)} + c$
 $= \frac{-1}{2\sqrt{3}} \tan^{-1} \left(\frac{2t}{\sqrt{3}}\right) + c$
 $= \frac{-1}{2\sqrt{3}} \tan^{-1} \left(\frac{2\cos}{\sqrt{3}}\right) + c$

93. Let $I = \int \frac{d}{a^2 \sin^2 + b^2 \cos^2}$
 Dividing N^r and D^r by \cos^2 , we get
 $I = \int \frac{\sec^2}{a^2 \tan^2 + b^2} d$
 Put $\tan = t \Rightarrow \sec^2 d = dt$
 $\therefore I = \int \frac{dt}{a^2 t^2 + b^2} = \frac{1}{a^2} \int \frac{dt}{t^2 + \frac{b^2}{a^2}}$
 $= \frac{1}{a^2} \cdot \frac{1}{\left(\frac{b}{a}\right)} \tan^{-1} \left(\frac{t}{\left(\frac{b}{a}\right)}\right) + c$
 $= \frac{1}{ab} \tan^{-1} \left(\frac{a}{b} \tan\right) + c$

94. Let $I = \int \frac{d}{4\sin^2 + 5\cos^2}$
 Dividing N^r and D^r by \cos^2 , we get
 $I = \int \frac{\sec^2 d}{4\tan^2 + 5}$
 $= \frac{1}{4} \int \frac{\sec^2 d}{\tan^2 + \frac{5}{4}}$
 Put $\tan = t$
 $\Rightarrow \sec^2 d = dt$
 $\therefore I = \frac{1}{4} \int \frac{dt}{t^2 + \left(\frac{\sqrt{5}}{2}\right)^2} = \frac{1}{4} \cdot \frac{1}{\frac{\sqrt{5}}{2}} \tan^{-1} \left(\frac{2t}{\sqrt{5}}\right) + c$
 $= \frac{1}{2\sqrt{5}} \tan^{-1} \left(\frac{2\tan}{\sqrt{5}}\right) + c$

95. Let $I = \int \frac{d}{2\sin^2 - 3\cos^2 + 7}$
 Dividing N^r and D^r by \cos^2 , we get

$$I = \int \frac{\sec^2 d}{2\tan^2 - 3 + 7\sec^2}$$

$$= \int \frac{\sec^2 d}{2\tan^2 - 3 + 7(1 + \tan^2)}$$

$$= \int \frac{\sec^2 d}{4 + 9\tan^2}$$

Put $t = \tan$
 $\Rightarrow dt = \sec^2 d$

$$\therefore I = \int \frac{dt}{2^2 + (3t)^2} = \frac{1}{6} \tan^{-1} \left(\frac{3t}{2}\right) + c$$

$$= \frac{1}{6} \tan^{-1} \left(\frac{3\tan}{2}\right) + c$$

96. $\int \frac{d}{2 + \cos}$
 $= \int \frac{d}{2\sin^2\left(\frac{d}{2}\right) + 2\cos^2\left(\frac{d}{2}\right) + \cos^2\left(\frac{d}{2}\right) - \sin^2\left(\frac{d}{2}\right)}$
 $= \int \frac{d}{\sin^2\left(\frac{d}{2}\right) + 3\cos^2\left(\frac{d}{2}\right)} = \int \frac{\sec^2\left(\frac{d}{2}\right) d}{\tan^2\left(\frac{d}{2}\right) + 3}$

Put $\tan\left(\frac{d}{2}\right) = t$

$$\Rightarrow \sec^2\left(\frac{d}{2}\right) d = 2dt$$

$$\therefore \int \frac{d}{2 + \cos} = 2 \int \frac{dt}{t^2 + 3} = \frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{t}{\sqrt{3}}\right) + c$$

$$= \frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{\tan\left(\frac{d}{2}\right)}{\sqrt{3}}\right) + c$$

97. Let $I = \int \frac{d}{2\sin + \cos + 3}$

Put $t = \tan\left(\frac{d}{2}\right)$

$$\therefore d = \frac{2dt}{1+t^2} \text{ and } \cos = \frac{1-t^2}{1+t^2}, \sin = \frac{2t}{1+t^2}$$



$$\begin{aligned} \therefore I &= \int \frac{\frac{2dt}{1+t^2}}{2\left(\frac{2t}{1+t^2}\right) + \left(\frac{1-t^2}{1+t^2}\right) + 3} \\ &= 2 \int \frac{dt}{4t+1-t^2+3+3t^2} \\ &= 2 \int \frac{dt}{2t^2+4t+4} = \int \frac{dt}{t^2+2t+2} \\ &= \int \frac{dt}{t^2+2t+1+1} = \int \frac{dt}{(t+1)^2+1^2} \\ &= \tan^{-1}\left(\frac{t+1}{1}\right) + c = \tan^{-1}\left(\tan\left(\frac{t}{2}\right)+1\right) + c \end{aligned}$$

98. Put $t^2 = t$
 $\Rightarrow 2t \, dt = dt \quad \Rightarrow d = \frac{dt}{2} = \frac{dt}{2\sqrt{t}}$

$$\begin{aligned} \therefore \int \frac{d}{\sqrt{t^4-1}} &= \int \frac{dt}{2t\sqrt{t^2-1}} = \frac{1}{2} \sec^{-1} t + c \\ &= \frac{1}{2} \sec^{-1} t^2 + c \end{aligned}$$

99. Let $I = \int \frac{t^2-1}{(t^2+1)\sqrt{t^4+1}} dt$
 Dividing N^r and D^r by t^2 , we get

$$= \int \frac{\left(1 - \frac{1}{t^2}\right)}{\left(t + \frac{1}{t}\right)\sqrt{t^2 + \frac{1}{t^2}}} dt$$

$$= \int \frac{\left(1 - \frac{1}{t^2}\right)}{\left(t + \frac{1}{t}\right)\sqrt{\left(t + \frac{1}{t}\right)^2 - 2}} dt$$

Put $t + \frac{1}{t} = t \Rightarrow \left(1 - \frac{1}{t^2}\right) dt = dt$

$$\begin{aligned} \therefore I &= \int \frac{dt}{t\sqrt{t^2-2}} = \frac{1}{\sqrt{2}} \sec^{-1}\left(\frac{t}{\sqrt{2}}\right) + c \\ &= \frac{1}{\sqrt{2}} \sec^{-1}\left[\frac{\left(t + \frac{1}{t}\right)}{\sqrt{2}}\right] + c \\ &= \frac{1}{\sqrt{2}} \sec^{-1}\left(\frac{t^2+1}{\sqrt{2}t}\right) + c \end{aligned}$$

100. Put $t = \tan \theta \Rightarrow dt = \sec^2 \theta \, d\theta$
 $\therefore \int \frac{\sec^2 \theta \, d\theta}{\sqrt{\tan^2 \theta + 4}} = \int \frac{1}{\sqrt{t^2+2^2}} dt$
 $= \log|t + \sqrt{t^2+4}| + c$
 $= \log|\tan \theta + \sqrt{\tan^2 \theta + 4}| + c$

101. $\int \sqrt{t^2-8} + 7 \, dt = \int \sqrt{(t-4)^2 - (3)^2} \, dt$
 $= \frac{(t-4)}{2} \sqrt{t^2-8} + 7$
 $- \frac{9}{2} \log| -4 + \sqrt{t^2-8} + 7| + c$

102. Put $t^2 = t$
 $\Rightarrow 2t \, dt = dt$
 $\Rightarrow d = \frac{dt}{2}$

$$\begin{aligned} \therefore \int \frac{d}{\sqrt{t^4-4}} &= \frac{1}{2} \int \frac{dt}{\sqrt{t^2-2^2}} \\ &= \frac{1}{2} \log|t + \sqrt{t^2-4}| + c \\ &= \frac{1}{2} \log|t^2 + \sqrt{t^4-4}| + c \end{aligned}$$

103. Let $I = \int \frac{e}{\sqrt{e^2+4e+13}} \, dt$
 Put $e = t \Rightarrow e \, dt = dt$
 $\therefore I = \int \frac{dt}{\sqrt{t^2+4t+13}}$
 $= \int \frac{dt}{\sqrt{(t+2)^2+3^2}}$
 $= \log|t+2 + \sqrt{(t+2)^2+3^2}| + c$
 $= \log|e+2 + \sqrt{e^2+4e+13}| + c$

104. Let $I = \int \frac{1}{\sqrt{\operatorname{cosec}^2 t + \cot^2 t}} \, dt$
 $= \int \frac{\sin t}{\sqrt{1+\cos^2 t}} \, dt$
 Put $t = \cos \theta \Rightarrow dt = -\sin \theta \, d\theta$
 $\therefore I = - \int \frac{dt}{\sqrt{1+t^2}} = -\log(t + \sqrt{1+t^2}) + c$
 $= -\log(\cos \theta + \sqrt{1+\cos^2 \theta}) + c$



$$\begin{aligned}
 105. \int \frac{d}{2^2 + -1} &= \frac{1}{2} \int \frac{d}{2 + \frac{1}{2} - \frac{1}{2}} \\
 &= \frac{1}{2} \int \frac{d}{\left(\frac{1}{4}\right)^2 - \left(\frac{3}{4}\right)^2} \\
 &= \frac{1}{2} \cdot \frac{1}{2 \cdot \left(\frac{3}{4}\right)} \log \left| \frac{\left(\frac{1}{4}\right) - \frac{3}{4}}{\left(\frac{1}{4}\right) + \frac{3}{4}} \right| + c \\
 &= \frac{1}{3} \log \left| \frac{2 - 1}{2(+ 1)} \right| + c
 \end{aligned}$$

$$\begin{aligned}
 106. \text{ Put } \log &= t \\
 \Rightarrow \frac{1}{d} &= dt
 \end{aligned}$$

$$\begin{aligned}
 \therefore \int \frac{d}{[(\log)^2 + 4\log - 1]} &= \int \frac{dt}{t^2 + 4t - 1} \\
 &= \int \frac{dt}{(t+2)^2 - (\sqrt{5})^2} \\
 &= \frac{1}{2\sqrt{5}} \log \left| \frac{t+2-\sqrt{5}}{t+2+\sqrt{5}} \right| + c \\
 &= \frac{1}{2\sqrt{5}} \log \left| \frac{\log + 2 - \sqrt{5}}{\log + 2 + \sqrt{5}} \right| + c
 \end{aligned}$$

$$\begin{aligned}
 107. \text{ Let } I &= \int \frac{1}{(t^2 - 1)\sqrt{t^2 + 1}} dt \\
 \text{Put } &= \frac{1}{t} \Rightarrow dt = -\frac{1}{t^2} dt
 \end{aligned}$$

$$\begin{aligned}
 \therefore I &= \int \frac{-\frac{1}{t^2}}{\left(\frac{1}{t^2} - 1\right)\sqrt{\frac{1}{t^2} + 1}} dt \\
 &= - \int \frac{t}{(1-t^2)\sqrt{1+t^2}} dt
 \end{aligned}$$

$$\text{Put } \sqrt{1+t^2} = u \Rightarrow 1+t^2 = u^2 \Rightarrow tdt = udu$$

$$\begin{aligned}
 \therefore I &= - \int \frac{u}{[1-(u^2-1)]u} du \\
 &= - \int \frac{du}{2-u^2} = \int \frac{du}{u^2 - (\sqrt{2})^2} \\
 &= \frac{1}{2\sqrt{2}} \log \left| \frac{u-\sqrt{2}}{u+\sqrt{2}} \right| + c
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{1+t^2} - \sqrt{2}}{\sqrt{1+t^2} + \sqrt{2}} \right| + c \\
 &= \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{1+\left(\frac{1}{t}\right)^2} - \sqrt{2}}{\sqrt{1+\left(\frac{1}{t}\right)^2} + \sqrt{2}} \right| + c \\
 &= \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{t^2+1} - \sqrt{2}}{\sqrt{t^2+1} + \sqrt{2}} \right| + c
 \end{aligned}$$

$$\begin{aligned}
 108. \text{ Let } I &= \int \frac{2 + 1}{4 + 2^3 + t^2 - 1} dt \\
 &= \int \frac{2 + 1}{[(t + 1)]^2 - 1} dt
 \end{aligned}$$

$$\begin{aligned}
 \text{Put } t &= (+ 1) \\
 \Rightarrow dt &= (2 + 1) dt
 \end{aligned}$$

$$\begin{aligned}
 \therefore I &= \int \frac{dt}{t^2 - 1} \\
 &= \frac{1}{2} \log \left| \frac{t-1}{t+1} \right| + c \\
 &= \frac{1}{2} \log \left| \frac{t^2 + -1}{t^2 + +1} \right| + c \\
 &= -\frac{1}{2} \log \left| \frac{t^2 + +1}{t^2 + -1} \right| + c
 \end{aligned}$$

$$\therefore A = -\frac{1}{2}$$

$$\begin{aligned}
 109. \int \frac{1}{\sin \sqrt{\sin} \cdot \cos} dt &= \int \frac{1}{\sin \sqrt{\sin^2} \cdot \frac{\cos}{\sin}} dt \\
 &= \int \frac{1}{\sin \cdot \sin \sqrt{\frac{\cos}{\sin}}} dt \\
 &= \int \frac{1}{\sqrt{\cot}} \times \operatorname{cosec}^2 dt
 \end{aligned}$$

$$\begin{aligned}
 \text{Put } t &= \cot \\
 \Rightarrow -dt &= \operatorname{cosec}^2 dt
 \end{aligned}$$

$$\begin{aligned}
 \therefore \int \frac{1}{\sin \sqrt{\sin} \cdot \cos} dt &= \int \frac{-dt}{\sqrt{t}} = -\int t^{-\frac{1}{2}} dt \\
 &= -2t^{\frac{1}{2}} + c \\
 &= -2\sqrt{\cot} + c
 \end{aligned}$$



$$\begin{aligned}
 110. \text{ Let } I &= \int \sin^3 \sqrt{\cos} \, d \\
 &= \int (1 - \cos^2) \sqrt{\cos} \sin \, d \\
 \text{Put } t &= \cos \\
 \Rightarrow dt &= -\sin \, d \\
 \Rightarrow -dt &= \sin \, d \\
 \therefore I &= \int (1 - t^2) \sqrt{t} (-dt) \\
 &= -\int \sqrt{t} \, dt + \int t^{5/2} \, dt = -\frac{t^{3/2}}{3/2} + \frac{t^{7/2}}{7/2} + c \\
 &= \frac{2}{7} (\sqrt{\cos})^7 - \frac{2}{3} (\sqrt{\cos})^3 + c
 \end{aligned}$$

$$\begin{aligned}
 111. \text{ Let } I &= \int \frac{d}{\sqrt{t^3+3}\sqrt{t}} \\
 \text{Put } \frac{1}{6} &= t \\
 \Rightarrow &= t^6 \\
 \Rightarrow d &= 6t^5 dt \\
 \therefore I &= \int \frac{6t^5}{t^3+t^2} dt \\
 &= \int \frac{6t^5}{t^2(t+1)} dt = 6 \int \frac{t^3}{t+1} dt \\
 &= 6 \int \frac{t^3+1-1}{t+1} dt = 6 \int \left(t^2 - t + 1 - \frac{1}{t+1} \right) dt \\
 &= 6 \left[\frac{t^3}{3} - \frac{t^2}{2} + t - \log|t+1| \right] + c \\
 &= 2\sqrt{} - 3\sqrt{} + 6\sqrt{} - 6 \log|\sqrt{} + 1| + c
 \end{aligned}$$

$$\begin{aligned}
 112. \text{ Let } I &= \int \frac{5d}{\sqrt{1+d^3}} = \int \frac{3 \cdot 2}{\sqrt{1+d^3}} d \\
 \text{Put } 1 + d^3 &= t^2 \\
 \Rightarrow 3 \cdot 2 d &= 2t dt \\
 \therefore I &= \int \frac{(t^2-1)}{t} \left(\frac{2}{3} \right) t dt \\
 &= \frac{2}{3} \int (t^2-1) dt = \frac{2}{3} \left(\frac{t^3}{3} - t \right) + c \\
 &= \frac{2}{9} t(t^2-3) + c \\
 &= \frac{2}{9} \sqrt{1+d^3} (1+d^3-3) + c \\
 &= \frac{2}{9} \sqrt{1+d^3} (d^3-2) + c
 \end{aligned}$$

$$\begin{aligned}
 113. \text{ Let } I &= \int \sec^6 d = \int \sec^4 \cdot \sec^2 d \\
 &= \int (1 + \tan^2)^2 \sec^2 d \\
 \text{Put } t &= \tan \\
 \Rightarrow dt &= \sec^2 d \\
 \therefore I &= \int (1+t^2)^2 dt = \int (1+2t^2+t^4) dt \\
 &= t + \frac{2t^3}{3} + \frac{t^5}{5} + c \\
 &= \tan + \frac{2}{3} \tan^3 + \frac{1}{5} \tan^5 + c
 \end{aligned}$$

$$\begin{aligned}
 114. \text{ Let } I &= \int \sec^{\frac{2}{3}} \operatorname{cosec}^{\frac{4}{3}} d = \int \frac{1}{\cos^{\frac{2}{3}} \sin^{\frac{4}{3}}} d \\
 \text{Dividing } N^r \text{ and } D^r &\text{ by } \cos^{\frac{4}{3}}, \text{ we get} \\
 I &= \int \frac{\sec^2}{\tan^3} d \\
 \text{Put } \tan &= t \\
 \Rightarrow \sec^2 d &= dt \\
 \therefore I &= \int \frac{dt}{t^{4/3}} = \int t^{-4/3} dt = \frac{t^{-1/3}}{-1/3} + c \\
 &= -3 (\tan)^{-1/3} + c
 \end{aligned}$$

$$\begin{aligned}
 115. \text{ Let } I &= \int \tan^4 d = \int \tan^2 (\sec^2 - 1) d \\
 &= \int (\tan^2 \sec^2 - \tan^2) d \\
 &= \int (\tan^2 \cdot \sec^2 - \sec^2 + 1) d \\
 &= \int (\tan^2 - 1) \sec^2 d + \int 1 d
 \end{aligned}$$

In 1st integral,
 Put $t = \tan$
 $\Rightarrow dt = \sec^2 d$

$$\begin{aligned}
 \therefore I &= \frac{t^3}{3} - t + c \\
 \therefore I &= \frac{\tan^3}{3} - \tan + c \\
 \therefore A &= \frac{1}{3}, B = -1, f(\) = + c
 \end{aligned}$$

$$\begin{aligned}
 116. \text{ Let } I &= \int \frac{\sin^3 2}{\cos^5 2} d \\
 &= \int \frac{\sin^3 2}{\cos^3 2} \cdot \frac{1}{\cos^2 2} d \\
 &= \int \tan^3 2 \cdot \sec^2 2 d
 \end{aligned}$$



$$\text{Put } \tan 2 = t \Rightarrow 2 \sec^2 2 \, d = dt$$

$$\begin{aligned} \therefore I &= \int t^3 \cdot \frac{dt}{2} = \frac{1}{2} \cdot \frac{t^4}{4} + c \\ &= \frac{1}{8} (\tan^4 2) + c \end{aligned}$$

$$\begin{aligned} 117. \int \log x \, dx &= \int \log x \cdot 1 \, dx \\ &= \log x \cdot x - \int 1 \cdot x \, dx \\ &= \log x \cdot x - \frac{x^2}{2} + c = (x \log x - \frac{x^2}{2}) + c \\ &= (x \log x - \log e) + c = \log \left(\frac{x}{e} \right) + c \end{aligned}$$

$$\begin{aligned} 118. \int e^{2x} \, dx &= \frac{e^{2x}}{2} - \int 1 \cdot \frac{e^{2x}}{2} \, dx \\ &= \frac{e^{2x}}{2} - \frac{e^{2x}}{4} + c \\ &= e^{2x} \left(\frac{2-1}{4} \right) + c \end{aligned}$$

$$\therefore f(x) = \frac{2-1}{4}$$

$$\begin{aligned} 119. \int x^2 e^{3x} \, dx &= \frac{x^2 \cdot e^{3x}}{3} - \int 2x \cdot \frac{e^{3x}}{3} \, dx \\ &= \frac{2e^{3x}}{3} - \frac{2}{3} \left[\frac{e^{3x}}{3} - \int 1 \cdot \frac{e^{3x}}{3} \, dx \right] \\ &= \frac{2e^{3x}}{3} - \frac{2}{3} \left(\frac{e^{3x}}{3} - \frac{e^{3x}}{9} \right) + c \\ &= \frac{1}{3} x^2 e^{3x} - \frac{2}{9} e^{3x} + \frac{2}{27} e^{3x} + c \end{aligned}$$

$$\begin{aligned} 120. \int x^3 \log x \, dx &= \log x \cdot \frac{x^4}{4} - \int 1 \cdot \frac{x^4}{4} \, dx \\ &= \frac{x^4}{4} \log x - \int \frac{x^3}{4} \, dx \\ &= \frac{x^4}{4} \log x - \frac{x^4}{16} + c \\ &= \frac{1}{16} (4^4 \log x - x^4) + c \end{aligned}$$

$$\begin{aligned} 121. \int \frac{\log x}{x^3} \, dx &= \int \log x \cdot x^{-3} \, dx \\ &= \log x \cdot \frac{x^{-2}}{-2} - \int \left(\frac{1}{x} \cdot \frac{x^{-2}}{-2} \right) dx \\ &= -\frac{\log x}{2x^2} + \frac{1}{2} \int x^{-3} \, dx \end{aligned}$$

$$\begin{aligned} &= -\frac{\log x}{2x^2} + \frac{1}{2} \cdot \frac{x^{-2}}{-2} + c \\ &= -\frac{\log x}{2x^2} - \frac{1}{4x^2} + c \\ &= -\frac{1}{4x^2} (2 \log x + 1) + c \end{aligned}$$

$$\begin{aligned} 122. \int x^n \log x \, dx &= \log x \cdot \frac{x^{n+1}}{n+1} - \int 1 \cdot \frac{x^{n+1}}{n+1} \, dx \\ &= \frac{x^{n+1}}{n+1} \log x - \frac{x^{n+1}}{(n+1)^2} + c \\ &= \frac{x^{n+1}}{n+1} \left(\log x - \frac{1}{n+1} \right) + c \end{aligned}$$

$$\begin{aligned} 123. \int [f(x) + f'(x)] \, dx \\ &= \int f(x) \, dx + \int f'(x) \, dx \\ &= \int f(x) \, dx + x \cdot f(x) - \int f(x) \, dx + c \\ &= x \cdot f(x) + c \end{aligned}$$

$$\begin{aligned} 124. \int [f(x)g'(x) - f'(x)g(x)] \, dx \\ &= f(x)g'(x) - \int f'(x)g'(x) \, dx - g(x)f'(x) \\ &\quad + \int f'(x)g'(x) \, dx \\ &= f(x)g'(x) - g(x)f'(x) + c \end{aligned}$$

$$\begin{aligned} 125. I_5 + 5I_4 &= \int x^5 e^x \, dx + 5 \int x^4 \cdot e^x \, dx \\ &= x^5 e^x - 5 \int x^4 e^x \, dx + 5 \int x^4 \cdot e^x \, dx + c \\ &= x^5 e^x + c \end{aligned}$$

$$\begin{aligned} 126. \text{ Let } I &= \int \tan^{-1} x \cdot 1 \, dx \\ &= \tan^{-1} x - \int \frac{1}{1+x^2} \cdot dx \\ &= \tan^{-1} x - \frac{1}{2} \int \frac{2}{1+x^2} \, dx \\ &= \tan^{-1} x - \frac{1}{2} \log |1+x^2| + c \end{aligned}$$

$$\begin{aligned} 127. \int \tan^{-1} x \, dx &= (\tan^{-1} x) \left(\frac{x^2}{2} \right) - \int \frac{1}{1+x^2} \left(\frac{x^2}{2} \right) dx \\ &= \frac{1}{2} x^2 \tan^{-1} x - \frac{1}{2} \int \frac{x^2+1-1}{1+x^2} \, dx \\ &= \frac{1}{2} x^2 \tan^{-1} x - \frac{1}{2} \int \left(1 - \frac{1}{x^2+1} \right) dx \\ &= \frac{1}{2} x^2 \tan^{-1} x - \frac{1}{2} (x - \tan^{-1} x) + c \\ &= \frac{1}{2} (x^2+1) \tan^{-1} x - \frac{1}{2} x + c \end{aligned}$$



$$128. I = \int \tan^{-1} \left(\frac{2}{1-x^2} \right) dx$$

$$\therefore I = 2 \int \tan^{-1} x \, dx$$

$$= 2 \left(\tan^{-1} x - \int \frac{1}{1+x^2} \cdot dx \right)$$

$$= 2 \left(\tan^{-1} x - \frac{1}{2} \int \frac{2}{1+x^2} dx \right)$$

$$\therefore I = 2 \tan^{-1} x - \log(1+x^2) + c$$

$$\therefore I - 2 \tan^{-1} x = -\log(1+x^2) + c$$

$$129. \int x^3 \tan^{-1} x \, dx = \tan^{-1} x \cdot \frac{x^4}{4} - \frac{1}{4} \int \frac{x^4}{x^2+1} dx$$

$$= \frac{x^4}{4} \tan^{-1} x - \frac{1}{4} \int \frac{x^4 - 1 + 1}{x^2 + 1} dx$$

$$= \frac{x^4}{4} \tan^{-1} x - \frac{1}{4} \int \left[(x^2 - 1) + \frac{1}{x^2 + 1} \right] dx$$

$$= \frac{x^4}{4} \tan^{-1} x - \frac{1}{4} \left[\frac{x^3}{3} - x + \tan^{-1} x \right] + c$$

$$= \frac{1}{4} \left[(x^4 - 1) \tan^{-1} x - \frac{x^3}{3} + x \right] + c$$

$$130. \int \log \left(1 + \frac{1}{x} \right) dx$$

$$= \log \left(1 + \frac{1}{x} \right) \cdot \frac{x^2}{2} + \frac{1}{2} \int \frac{-1}{x^2+1} dx$$

$$= \frac{x^2}{2} \log \left(1 + \frac{1}{x} \right) + \frac{1}{2} \left[-\frac{1}{2} \log(x+1) \right] + c$$

$$= \left(\frac{x^2-1}{2} \right) \log(x+1) - \frac{x^2}{2} \log x + \frac{1}{2} + c$$

$$131. \int \log(x^2 + 1) dx = \int \log[(x+1)(x-1)] dx$$

$$= \int \log(x-1) dx + \int \log(x+1) dx$$

$$= \log(x-1) \cdot x - \int \frac{1}{x-1} dx + \log(x+1) \cdot x - \int \frac{1}{x+1} dx$$

$$= x \log(x-1) - \int \frac{1}{x} dx + x \log(x+1) - \int \left(\frac{x+1-1}{x+1} \right) dx$$

$$= x \log(x-1) - x + x \log(x+1) - \int \left(1 - \frac{1}{x+1} \right) dx$$

$$= x \log(x-1) - x + x \log(x+1) - x + \log|x+1| + c$$

$$= [x \log(x-1) + x \log(x+1)] - 2x + \log|x+1| + c$$

$$= \log(x^2 - 1) - 2x + \log|x+1| + c$$

$$\therefore A = -2x + \log|x+1| + c$$

$$132. \text{ Put } \sin^{-1} x = t \Rightarrow x = \sin t \Rightarrow dx = \cos t \, dt$$

$$\therefore \int \sin^{-1} x \, dx = \int t \cos t \, dt$$

$$= t \sin t - \int 1 \cdot \sin t \, dt$$

$$= t \sin t + \cos t + c$$

$$= \sin^{-1} x + \sqrt{1-x^2} + c$$

$$133. \text{ Put } x = t^2 \Rightarrow dx = 2t \, dt$$

$$\therefore \int \sin \sqrt{x} \, dx = \int \sin t (2t) \, dt$$

$$= 2 \int t \sin t \, dt$$

$$= 2 \left[t(-\cos t) - \int (1)(-\cos t) dt \right]$$

$$= 2 \left(-t \cos t + \int \cos t \, dt \right)$$

$$= -2t \cos t + 2 \sin t + c$$

$$= -2\sqrt{x} \cos \sqrt{x} + 2 \sin \sqrt{x} + c$$

$$134. \text{ Put } x = t^2 \Rightarrow dx = 2t \, dt$$

$$\therefore \int \sqrt{x} \cdot e^{\sqrt{x}} \, dx = 2 \int t^2 \cdot e^t \, dt$$

$$= 2(t^2 \cdot e^t - 2te^t + 2e^t) + c$$

$$= 2 \left(x \cdot e^{\sqrt{x}} - 2\sqrt{x} \cdot e^{\sqrt{x}} + 2e^{\sqrt{x}} \right) + c$$

$$= e^{\sqrt{x}} (2x - 4\sqrt{x} + 4) + c$$

$$135. \text{ Put } x^2 = t \Rightarrow 2x \, dx = dt$$

$$\therefore \int x^5 e^{x^2} \, dx = \frac{1}{2} \int t^2 e^t \, dt$$

$$= \frac{1}{2} \left[t^2 e^t - 2 \int t e^t \, dt \right]$$

$$= \frac{t^2 e^t}{2} - (t e^t - e^t) + c$$

$$= \frac{1}{2} x^4 e^{x^2} - x^2 e^{x^2} + e^{x^2} + c$$

$$136. \text{ Let } I = \int \sin(\log x) \, dx$$

$$\text{Put } \log x = t \Rightarrow x = e^t \Rightarrow dx = e^t \, dt$$

$$\therefore I = \int \sin t \cdot e^t \, dt = \sin t \cdot e^t - \int \cos t \cdot e^t \, dt$$

$$= \sin t \cdot e^t - \left[\cos t \cdot e^t + \int \sin t \cdot e^t \, dt \right]$$

$$\therefore I = \sin t \cdot e^t - \cos t \cdot e^t - I + c_1$$

$$\Rightarrow 2I = \sin t \cdot e^t - \cos t \cdot e^t + c_1$$

$$\Rightarrow I = \frac{1}{2} [\sin(\log x) - \cos(\log x)] + c,$$

where $c = \frac{c_1}{2}$



$$\begin{aligned}
 137. \int \sin \log(\sec \theta + \tan \theta) d\theta & \\
 &= \log(\sec \theta + \tan \theta) \cdot (-\cos \theta) - \int \sec \theta \cdot (-\cos \theta) d\theta \\
 &\quad \dots \left[\because \frac{d}{d\theta} [\log(\sec \theta + \tan \theta)] = \sec \theta \right] \\
 &= -\cos \theta \log(\sec \theta + \tan \theta) + \int 1 d\theta \\
 &= -\cos \theta \log(\sec \theta + \tan \theta) + c
 \end{aligned}$$

$$\begin{aligned}
 138. \text{ Put } \theta &= \sin \theta \\
 \Rightarrow d\theta &= \cos \theta d\theta \\
 \therefore \int \sin^{-1}(3 - 4\theta^3) d\theta & \\
 &= \int \sin^{-1}(\sin 3\theta) \cos \theta d\theta \\
 &= \int 3\theta \cos \theta d\theta = 3(\theta \sin \theta - \int \sin \theta d\theta) \\
 &= 3(\theta \sin \theta + \cos \theta) + c \\
 &= 3\left(\sin^{-1} \sqrt{1 - \theta^2} + \sqrt{1 - \theta^2}\right) + c
 \end{aligned}$$

$$\begin{aligned}
 139. \text{ Put } \sin^{-1} \theta &= t \Rightarrow \frac{1}{\sqrt{1 - \theta^2}} d\theta = dt \\
 \therefore \int \frac{\sin^{-1} \theta}{(1 - \theta^2)^{3/2}} d\theta &= \int t \sec^2 t dt \\
 &= t \tan t - \int 1 \cdot \tan t dt \\
 &= t \tan t + \log(\cos t) + c \\
 &= t \cdot \frac{\sin t}{\sqrt{1 - \sin^2 t}} + \log(\sqrt{1 - \sin^2 t}) + c \\
 &= \frac{\sin^{-1} \theta}{\sqrt{1 - \theta^2}} + \frac{1}{2} \log(1 - \theta^2) + c
 \end{aligned}$$

$$\begin{aligned}
 140. \text{ Put } \tan^{-1} \theta &= \theta \Rightarrow \theta = \tan \theta \Rightarrow d\theta = \sec^2 \theta d\theta \\
 \therefore \int \frac{\tan^{-1} \theta}{(1 + \theta^2)^{3/2}} d\theta &= \int \frac{\theta \tan \theta \sec^2 \theta}{(1 + \tan^2 \theta)^{3/2}} d\theta \\
 &= \int \theta \sin \theta d\theta = -\theta \cos \theta + \sin \theta + c \\
 &= \frac{\tan \theta}{\sqrt{1 + \tan^2 \theta}} - \theta \cdot \frac{1}{\sqrt{1 + \tan^2 \theta}} + c \\
 &= \frac{\tan \theta}{\sqrt{1 + \theta^2}} - \tan^{-1} \theta \cdot \frac{1}{\sqrt{1 + \theta^2}} + c \\
 &= \frac{-\tan^{-1} \theta}{\sqrt{1 + \theta^2}} + c
 \end{aligned}$$

$$\begin{aligned}
 141. \int \sec^3 \theta d\theta &= \int \sec \theta \cdot \sec^2 \theta d\theta \\
 &= \sec \theta \tan \theta - \int (\sec \theta \tan \theta \cdot \tan \theta) d\theta \\
 &= \sec \theta \tan \theta - \int \sec \theta \tan^2 \theta d\theta \\
 &= \sec \theta \tan \theta - \int \sec \theta (\sec^2 \theta - 1) d\theta \\
 &= \sec \theta \tan \theta - \int (\sec^3 \theta - \sec \theta) d\theta \\
 &= \sec \theta \tan \theta - \int \sec^3 \theta d\theta + \int \sec \theta d\theta \\
 \therefore 2 \int \sec^3 \theta d\theta &= \sec \theta \tan \theta + \int \sec \theta d\theta \\
 &= \sec \theta \tan \theta + \log |\sec \theta + \tan \theta| + c \\
 \therefore \int \sec^3 \theta d\theta &= \frac{1}{2} [\sec \theta \tan \theta + \log |\sec \theta + \tan \theta|] + c
 \end{aligned}$$

$$\begin{aligned}
 142. \int \frac{e^{(x-1)}}{2} dx &= \int e^{(x-1)} dx = \frac{e^x}{2} + c \\
 \dots \left[\because \int e^{[f(x) + f'(x)]} dx &= e^{f(x)} + c \right]
 \end{aligned}$$

$$\begin{aligned}
 143. \int e^{(x^5 + 5x^4 + 1)} dx & \\
 &= \int e^{(x^5 + 5x^4)} dx + \int e dx \\
 &= e \cdot x^5 + e + c
 \end{aligned}$$

$$\begin{aligned}
 144. \int e^{[\tan^{-1} x - \log(\cos x)]} dx & \\
 &= \int e^{[\tan^{-1} x + \log(\sec x)]} dx \\
 &= e \log(\sec x) + c \\
 \dots \left[\because \int e^{[f(x) + f'(x)]} dx &= e^{f(x)} + c \right]
 \end{aligned}$$

$$\begin{aligned}
 145. \int e^{(1 + \tan x + \tan^2 x)} dx &= \int e^{(\tan x + \sec^2 x)} dx \\
 &= e \tan x + c \\
 \dots \left[\because \int e^{[f(x) + f'(x)]} dx &= e^{f(x)} + c \right]
 \end{aligned}$$

$$\begin{aligned}
 146. \int e^{(1 - \cot x + \cot^2 x)} dx & \\
 &= \int e^{(-\cot x + \operatorname{cosec}^2 x)} dx \\
 &= e^{(-\cot x)} + c = -e \cot x + c
 \end{aligned}$$

$$\begin{aligned}
 147. \int \frac{e}{(x+1)^2} dx &= \int e \frac{(x+1-1)}{(x+1)^2} dx \\
 &= \int e \left[\frac{1}{x+1} - \frac{1}{(x+1)^2} \right] dx \\
 &= \frac{e}{x+1} + c \\
 \dots \left[\because \int e^{[f(x) + f'(x)]} dx &= e^{f(x)} + c \right]
 \end{aligned}$$



148.
$$\int \frac{(x+3)e^x}{(x+4)^2} dx = \int \frac{(x+4-1)e^x}{(x+4)^2} dx$$

$$= \int e^x \left(\frac{1}{x+4} - \frac{1}{(x+4)^2} \right) dx$$

$$= \frac{e^x}{x+4} + c$$

149.
$$\int \frac{e^{(x^2+1)}}{(x^2+1)^2} dx = \int \frac{e^{(x^2-1+2)}}{(x^2+1)^2} dx$$

$$= \int e^x \left[\frac{-1}{x+1} + \frac{2}{(x^2+1)^2} \right] dx$$

$$= e^x \left(\frac{-1}{x+1} \right) + c$$

150.
$$\int \left(\frac{2+\sin 2x}{1+\cos 2x} \right) e^x dx$$

$$= \int \left(\frac{2+2\sin x \cos x}{2\cos^2 x} \right) e^x dx$$

$$= \int (\sec^2 x + \tan x) e^x dx$$

$$= e^x \tan x + c$$

151.
$$\int e^x \left(\frac{1-\sin x}{1-\cos x} \right) dx = \int e^x \left[\frac{1-2\sin \frac{x}{2} \cos \frac{x}{2}}{2\sin^2 \left(\frac{x}{2} \right)} \right] dx$$

$$= \int e^x \left[\frac{1}{2} \operatorname{cosec}^2 \left(\frac{x}{2} \right) - \cot \left(\frac{x}{2} \right) \right] dx$$

$$= -e^x \cot \left(\frac{x}{2} \right) + c$$

152.
$$\int e^{2x} (2 \cos x - \sin x) dx = e^{2x} \cos x + c$$

.... $\left[\int e^{mx} [mf(x) + f'(x)] dx = e^{mx} f(x) + c \right]$

153. Let $I = \int \log(\log x + 2) dx$
 Put $\log x = t \Rightarrow x = e^t \Rightarrow dx = e^t dt$
 $\therefore I = \int t(t+2)e^t dt = \int e^t (t^2 + 2t) dt$
 $= e^t \cdot t^2 + c = (\log x)^2 + c$

154. Put $\log x = t \Rightarrow x = e^t \Rightarrow dx = e^t dt$
 $\therefore \int \left[\frac{1}{\log x} - \frac{1}{(\log x)^2} \right] dx = \int \left(\frac{1}{t} - \frac{1}{t^2} \right) e^t dt$
 $= \frac{e^t}{t} + c$
 $= \frac{x}{\log x} + c$

155. Put $\log x = t \Rightarrow x = e^t \Rightarrow dx = e^t dt$
 $\therefore \int \frac{\log x}{(1+\log x)^2} dx = \int \frac{t}{(1+t)^2} e^t dt$
 $= \int e^t \left[\frac{t+1-1}{(1+t)^2} \right] dt$
 $= \int e^t \left[\frac{1}{1+t} - \frac{1}{(1+t)^2} \right] dt$
 $= \frac{e^t}{1+t} + c$
 $= \frac{x}{1+\log x} + c$

156. Let $I = \int \left\{ \log(\log x) + \frac{1}{(\log x)^2} \right\} dx$
 Put $\log x = t \Rightarrow x = e^t \Rightarrow dx = e^t dt$
 $\therefore I = \int e^t \left(\log t + \frac{1}{t^2} \right) dt$
 $= \int e^t \left(\log t + \frac{1}{t} \right) dt + \int e^t \left(-\frac{1}{t} + \frac{1}{t^2} \right) dt$
 $= e^t \log t + e^t \left(-\frac{1}{t} \right) + c = \left(\log(\log x) - \frac{1}{\log x} \right) + c$
 $\therefore f(x) = \log(\log x)$ and $g(x) = \frac{1}{\log x}$

157.
$$\int \frac{dx}{x^2-3} = \int \frac{(1-x)dx}{x^2(1-x)} + \int \frac{dx}{x^2(1-x)}$$

$$= \int \frac{1}{x} dx + \int \frac{dx}{(1-x)}$$

$$= -\frac{1}{x} + \int \frac{dx}{1-x} + \int \frac{dx}{1-x}$$

$$= -\frac{1}{x} + \log|x| - \log|1-x| + c$$

$$= \log \left| \frac{x}{1-x} \right| - \frac{1}{x} + c$$

158.
$$\int \frac{x^2 + x - 1}{x^2 + x - 6} dx = \int \left[1 + \frac{5}{x^2 + x - 6} \right] dx$$

$$= \int \left[1 + \frac{5}{(x+3)(x-2)} \right] dx$$

$$= \int dx + \int \frac{dx}{x-2} - \int \frac{dx}{x+3}$$

$$= x + \log|x-2| - \log|x+3| + c$$



$$\begin{aligned}
 159. \int \frac{d}{x^4-1} &= \int \frac{d}{(x^2-1)(x^2+1)} \\
 &= \frac{1}{2} \int \left(\frac{1}{x^2-1} - \frac{1}{x^2+1} \right) d \\
 &= \frac{1}{4} \int \left(\frac{2}{x^2-1} - \frac{2}{x^2+1} \right) d \\
 &= \frac{1}{4} \log |x^2-1| - \frac{1}{4} \log |x^2+1| + c \\
 &= \frac{1}{4} \log \left| \frac{x^2-1}{x^2+1} \right| + c
 \end{aligned}$$

$$\begin{aligned}
 160. \int \frac{x^2}{(x^2+2)(x^2+3)} dx &= \int \left[\frac{3}{x^2+3} - \frac{2}{x^2+2} \right] dx \\
 &= \frac{3}{\sqrt{3}} \tan^{-1} \left(\frac{x}{\sqrt{3}} \right) - \frac{2}{\sqrt{2}} \tan^{-1} \left(\frac{x}{\sqrt{2}} \right) + c \\
 &= \sqrt{3} \tan^{-1} \left(\frac{x}{\sqrt{3}} \right) - \sqrt{2} \tan^{-1} \left(\frac{x}{\sqrt{2}} \right) + c
 \end{aligned}$$

$$\begin{aligned}
 161. \int \frac{dx}{(x^2-1)(1-2x)} &= \int \frac{-1}{(1-x)(1+x)(1-2x)} dx \\
 \text{Let } \frac{-1}{(1-x)(1+x)(1-2x)} &= \frac{A}{1-x} + \frac{B}{1+x} + \frac{C}{1-2x} \\
 \Rightarrow -1 &= A(1+x)(1-2x) + B(1-x)(1-2x) \\
 &\quad + C(1-x)(1+x) \dots (i) \\
 \text{Putting } x &= -1 \text{ in (i), we get } B = -\frac{1}{6} \\
 \text{Putting } x &= 1 \text{ in (i), we get } A = \frac{1}{2} \\
 \text{Putting } x &= \frac{1}{2} \text{ in (i), we get } C = -\frac{4}{3}
 \end{aligned}$$

$$\begin{aligned}
 \therefore \int \frac{dx}{(x^2-1)(1-2x)} &= \frac{1}{2} \int \frac{1}{1-x} dx - \frac{1}{6} \int \frac{1}{1+x} dx - \frac{4}{3} \int \frac{1}{1-2x} dx \\
 &= -\frac{1}{2} \log |1-x| - \frac{1}{6} \log |1+x| + \frac{2}{3} \log |1-2x| + c
 \end{aligned}$$

$$\begin{aligned}
 162. \int \frac{1}{x^3} dx &= \int \frac{1}{(1+x)(1-x)} dx \\
 &= \frac{1}{2} \int \left(\frac{2}{1+x} - \frac{1}{1-x} \right) dx \\
 &= \frac{1}{2} (2 \log |1+x| - \log |1-x|) + c \\
 &= \frac{1}{2} (\log |x^2| - \log |1-x^2|) + c = \frac{1}{2} \log \left| \frac{x^2}{1-x^2} \right| + c
 \end{aligned}$$

$$163. \text{ Put } e = t \Rightarrow e dx = dt$$

$$\begin{aligned}
 \therefore \int \frac{e}{(1+e)(2+e)} dx &= \int \frac{dt}{(1+t)(2+t)} \\
 &= \int \left(\frac{1}{1+t} - \frac{1}{2+t} \right) dt \\
 &= \log |1+t| - \log |2+t| + c \\
 &= \log |1+e| - \log |2+e| + c \\
 &= \log \left| \frac{1+e}{2+e} \right| + c
 \end{aligned}$$

$$164. \int \frac{dx}{e^x+1-2e^{-x}} = \int \frac{e^x}{e^{2x}+e-2} dx$$

$$\text{Put } e = t \Rightarrow e dx = dt$$

$$\begin{aligned}
 \therefore \int \frac{dx}{e^x+1-2e^{-x}} &= \int \frac{dt}{t^2+t-2} \\
 &= \int \frac{dt}{(t+2)(t-1)} = \frac{1}{3} \int \left(\frac{1}{t-1} - \frac{1}{t+2} \right) dt \\
 &= \frac{1}{3} \log |t-1| - \frac{1}{3} \log |t+2| + c \\
 &= \frac{1}{3} \log |e^x-1| - \frac{1}{3} \log |e^x+2| + c
 \end{aligned}$$

$$165. \text{ Let } I = \int \frac{a}{b+ce^x} dx = \int \frac{ae^x}{be^x+ce^{2x}} dx$$

$$\text{Put } e = t \Rightarrow e dx = dt$$

$$\begin{aligned}
 \therefore I &= a \int \frac{dt}{bt+ct^2} \\
 &= a \int \frac{dt}{t(ct+b)} \\
 &= -\frac{a}{b} \int \left(\frac{c}{ct+b} - \frac{1}{t} \right) dt \\
 &= -\frac{a}{b} \log |ct+b| + \frac{a}{b} \log |t| + c \\
 &= \frac{a}{b} \log \left| \frac{t}{ct+b} \right| + c \\
 &= \frac{a}{b} \log \left| \frac{e}{b+ce^x} \right| + c
 \end{aligned}$$

$$166. \text{ Put } \sin x = t \\ \Rightarrow \cos x dx = dt$$

$$\begin{aligned}
 \therefore \int \frac{\cos x}{(1+\sin x)(2+\sin x)} dx &= \int \frac{dt}{(t+1)(t+2)} \\
 &= \int \frac{1}{t+1} dt - \int \frac{1}{t+2} dt
 \end{aligned}$$



$$\begin{aligned}
 &= \log|t+1| - \log|t+2| + c \\
 &= \log \left| \frac{t+1}{t+2} \right| + c \\
 &= \log \left| \frac{\sin^{-1} t + 1}{\sin^{-1} t + 2} \right| + c
 \end{aligned}$$

167. $\int \frac{t^3-1}{t^3+1} dt = \int \frac{t^3}{t^3+1} dt - \int \frac{1}{t^3+1} dt$

$$\begin{aligned}
 &= \int \frac{t^2}{t^2+1} dt - \int \left(\frac{1}{t^2+1} - \frac{t^2}{t^2+1} \right) dt \\
 &= \int \left(1 - \frac{1}{t^2+1} \right) dt - \int \frac{1}{t^2+1} dt + \frac{1}{2} \int \frac{2}{t^2+1} dt \\
 &= -\tan^{-1} t - \log|t| + \frac{1}{2} \log|t^2+1| + c \\
 &= -\tan^{-1} t - \log|t| + \log|\sqrt{t^2+1}| + c
 \end{aligned}$$

168. Let $\frac{2t+7}{(t-4)^2} = \frac{A}{t-4} + \frac{B}{(t-4)^2}$

$$\begin{aligned}
 \Rightarrow 2t+7 &= A(t-4) + B = At + (-4A+B) \\
 \therefore A &= 2 \text{ and } -4A+B = 7 \\
 \therefore B &= 7+4A = 7+8 = 15 \\
 \therefore \int \frac{2t+7}{(t-4)^2} dt &= \int \left(\frac{2}{t-4} + \frac{15}{(t-4)^2} \right) dt \\
 &= 2 \log|t-4| - \frac{15}{t-4} + c
 \end{aligned}$$

169. Let $\frac{t^2+1}{(t-2)^2(t+3)} = \frac{A}{t-2} + \frac{B}{(t-2)^2} + \frac{C}{t+3}$

$$\begin{aligned}
 \Rightarrow t^2+1 &= A(t-2)(t+3) + B(t+3) + C(t-2)^2 \quad \dots(i) \\
 \text{Putting } t &= 2 \text{ in (i), we get } B = 1 \\
 \text{Putting } t &= -3 \text{ in (i), we get } C = \frac{2}{5} \\
 \text{Putting } t &= 3 \text{ in (i), we get} \\
 6A + 6B + C &= 10 \Rightarrow A = \frac{3}{5} \\
 \therefore \int \frac{t^2+1}{(t-2)^2(t+3)} dt &= \frac{3}{5} \int \frac{1}{t-2} dt + \int \frac{1}{(t-2)^2} dt + \frac{2}{5} \int \frac{1}{t+3} dt \\
 &= \frac{3}{5} \log|t-2| - \frac{1}{t-2} + \frac{2}{5} \log|t+3| + c
 \end{aligned}$$

170. Let $\frac{1}{(t-1)(t^2+1)} = \frac{A}{t-1} + \frac{B+C}{t^2+1}$

$$\therefore 1 = A(t^2+1) + (B+C)(t-1) \quad \dots(i)$$

Putting $t = 1$ in (i), we get

$$A = \frac{1}{2}$$

Putting $t = 0$ in (i), we get

$$A - C = 1 \Rightarrow C = -\frac{1}{2}$$

Comparing the coefficient of t^2 , we get

$$A + B = 0 \Rightarrow B = -\frac{1}{2}$$

$$\begin{aligned}
 \therefore \int \frac{1}{(t-1)(t^2+1)} dt &= \int \left[\frac{1}{2(t-1)} - \frac{1}{2(t^2+1)} \right] dt \\
 &= \frac{1}{2} \int \frac{1}{t-1} dt - \frac{1}{4} \int \frac{2}{t^2+1} dt - \frac{1}{2} \int \frac{1}{t^2+1} dt \\
 &= \frac{1}{2} \log|t-1| - \frac{1}{4} \log|t^2+1| - \frac{1}{2} \tan^{-1} t + c
 \end{aligned}$$

171. $\int \frac{dx}{1+x^2+x^3} = \int \frac{dx}{(1+x)(1+x^2)}$

$$\begin{aligned}
 &= \frac{1}{2} \int \frac{1}{1+x^2} dx + \frac{1}{2} \int \frac{1}{1+x} dx - \frac{1}{2} \int \frac{1}{1+x^2} dx \\
 &= \frac{1}{2} \tan^{-1} x + \log|\sqrt{1+x}| - \frac{1}{2} \log|\sqrt{1+x^2}| + c
 \end{aligned}$$

172. $\int \frac{t^4}{(t-1)(t^2+1)} dt$

$$\begin{aligned}
 &= \int \frac{t^4-1}{(t-1)(t^2+1)} dt + \int \frac{1}{(t-1)(t^2+1)} dt \\
 &= \int \frac{(t+1)(t-1)(t^2+1)}{(t-1)(t^2+1)} dt + \int \frac{1}{(t-1)(t^2+1)} dt \\
 &= \int (t+1) dt + \int \left[\frac{1}{2(t-1)} - \frac{1}{2(t^2+1)} \right] dt \\
 &= \int t dt + \int dt + \frac{1}{2} \int \frac{1}{t-1} dt \\
 &\quad - \frac{1}{4} \int \frac{2}{t^2+1} dt - \frac{1}{2} \int \frac{1}{t^2+1} dt \\
 &= \frac{t^2}{2} + \frac{1}{2} \log|t-1| - \frac{1}{4} \log|t^2+1| - \frac{1}{2} \tan^{-1} t + c
 \end{aligned}$$



$$173. \int \frac{d}{f(x)} = \log [f(x)]^2 + c$$

Differentiating on both sides, we get

$$\frac{1}{f(x)} = \frac{2f(x)f'(x)}{[f(x)]^2}$$

$$\Rightarrow f'(x) = \frac{1}{2}$$

$$\therefore f(x) = \int f'(x) dx = \frac{x}{2} + \alpha$$

$$174. |A| = \begin{vmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{vmatrix} = abc$$

$$\text{Let } I = \int |A| dx = \int 7^{7^x} 7^{7^x} dx$$

$$\text{Put } 7^{7^x} = t$$

$$\Rightarrow 7^{7^x} (\log 7)^3 7^x dx = dt$$

$$\Rightarrow 7^x dx = \frac{dt}{7^{7^x} (\log 7)^3} = \frac{dt}{t(\log 7)^3}$$

$$\therefore I = \frac{1}{(\log 7)^3} \int dt = \frac{t}{(\log 7)^3} + c$$

$$= \frac{7^{7^x}}{(\log 7)^3} + c$$

$$175. \text{ Put } x^2 = t \Rightarrow 2x dx = dt$$

$$\therefore \int x^3 \cos^2 x dx$$

$$= \frac{1}{2} \int t \cos t dt = \frac{1}{2} (t \sin t - \int \sin t dt)$$

$$= \frac{1}{2} (t \sin t + \cos t) + c$$

$$= \frac{1}{2} (x^2 \sin x^2 + \cos x^2) + c$$

$$176. \text{ Put } x = \cos \theta \Rightarrow \theta = \cos^{-1} x$$

$$\therefore \tan^{-1} \sqrt{\frac{1-x}{1+x}} = \tan^{-1} \left(\sqrt{\frac{1-\cos \theta}{1+\cos \theta}} \right)$$

$$= \tan^{-1} \left(\sqrt{\frac{2 \sin^2 \frac{\theta}{2}}{2 \cos^2 \frac{\theta}{2}}} \right)$$

$$= \tan^{-1} \left(\tan \frac{\theta}{2} \right) = \frac{\theta}{2}$$

$$= \frac{1}{2} \cos^{-1} x$$

$$\therefore \int \tan^{-1} \sqrt{\frac{1-x}{1+x}} dx = \frac{1}{2} \int (\cos^{-1} x) dx$$

$$= \frac{1}{2} \left[\cos^{-1} x + \int \frac{1}{\sqrt{1-x^2}} dx \right]$$

$$= \frac{1}{2} (\cos^{-1} x - \sqrt{1-x^2}) + c$$

$$177. \text{ Let } I = \int \sin x \sec^3 x dx$$

$$= \int \tan x \sec^2 x dx$$

$$\text{Put } \tan x = t \Rightarrow \sec^2 x dx = dt$$

$$\therefore I = \int \tan^{-1} t \cdot t dt$$

$$= \tan^{-1} t \cdot \frac{t^2}{2} - \int \frac{1}{1+t^2} \cdot \frac{t^2}{2} dt$$

$$= \frac{t^2 \tan^{-1} t}{2} - \frac{1}{2} \int \left(\frac{t^2+1-1}{1+t^2} \right) dt$$

$$= \frac{t^2 \tan^{-1} t}{2} - \frac{1}{2} \int \left(1 - \frac{1}{1+t^2} \right) dt$$

$$= \frac{t^2 \tan^{-1} t}{2} - \frac{1}{2} t + \frac{1}{2} \tan^{-1} t + c$$

$$= \frac{\tan^2 x}{2} - \frac{1}{2} \tan x + \frac{1}{2} + c$$

$$= \frac{(\sec^2 x - 1)}{2} - \frac{1}{2} \tan x + \frac{1}{2} + c$$

$$= \frac{1}{2} (\sec^2 x - \tan x) + c$$

$$178. \int \log(x+1) dx = \int \log(x+1) \cdot 1 dx$$

$$= \log(x+1) \cdot x - \int \frac{x}{x+1} dx$$

$$= \log(x+1) \cdot x - \int \frac{x+1-1}{x+1} dx$$

$$= \log(x+1) \cdot x - \int \left(1 - \frac{1}{x+1} \right) dx$$

$$= \log(x+1) \cdot x - x + \log(x+1) + c$$

$$= (x+1) \log(x+1) - x + c$$

$$179. \int \frac{1}{\cos x (1+\cos x)} dx$$

$$= \int \frac{1+\cos x - \cos x}{\cos x (1+\cos x)} dx$$

$$= \int \frac{dx}{\cos x} - \int \frac{dx}{1+\cos x}$$



$$\begin{aligned}
 &= \int \sec d - \int \frac{d}{2\cos^2 \frac{d}{2}} \\
 &= \int \sec d - \frac{1}{2} \int \sec^2 \frac{d}{2} \\
 &= \log|\sec + \tan| - \frac{1}{2} \left(\tan \frac{d}{2} \right) \cdot 2 + c \\
 &= \log|\sec + \tan| - \tan \frac{d}{2} + c
 \end{aligned}$$



Competitive Thinking

1. Rationalizing the denominator, we get

$$\begin{aligned}
 \int \frac{d}{\sqrt{d} + \sqrt{d-2}} &= \frac{1}{2} \int (\sqrt{d} - \sqrt{d-2}) d \\
 &= \frac{1}{2} \left[\frac{3}{2} d - \frac{(d-2)^{3/2}}{3/2} \right] + c \\
 &= \frac{1}{3} \left\{ \frac{3}{2} d - (d-2)^{3/2} \right\} + c
 \end{aligned}$$

2. Let $f(d) = e^d$

$$\therefore \int [f(d)]^2 d = \int (e^d)^2 d = \frac{e^{2d}}{2} = \frac{1}{2} [f(d)]^2$$

$$\begin{aligned}
 3. \int e^{\log a} \cdot e^d d &= \int e^{\log a} \cdot e^d d = \int a e^d d \\
 &= \int (ae) d = \frac{(ae)}{\log(ae)} + c
 \end{aligned}$$

$$\begin{aligned}
 4. \int \frac{e^{5\log d} - e^{4\log d}}{e^{3\log d} - e^{2\log d}} d &= \int \frac{d^5 - d^4}{d^3 - d^2} d \\
 &= \int \frac{d^4(d-1)}{d^2(d-1)} d = \int d^2 d = \frac{d^3}{3} + c
 \end{aligned}$$

$$\begin{aligned}
 5. \int \frac{e^{6\log d} - e^{5\log d}}{e^{4\log d} - e^{3\log d}} d &= \int \frac{d^6 - d^5}{d^4 - d^3} d \\
 &= \int \frac{d^5(d-1)}{d^3(d-1)} d \\
 &= \int d^2 d = \frac{d^3}{3} + c
 \end{aligned}$$

$$\begin{aligned}
 6. \int {}^{51}(\tan^{-1} + \cot^{-1}) d &= \int {}^{51} \frac{\pi}{2} d \\
 &\dots \left[\because \tan^{-1} + \cot^{-1} = \frac{\pi}{2} \right] \\
 &= \frac{\pi}{2} \cdot \frac{52}{2} + c \\
 &= \frac{52}{2} (\tan^{-1} + \cot^{-1}) + c
 \end{aligned}$$

$$\begin{aligned}
 7. \int \frac{d+1}{\sqrt{d}} d &= \int \left(\frac{1}{\sqrt{d}} + \frac{1}{\sqrt{d}} \right) d \\
 &= \int \left(d^{-\frac{1}{2}} + d^{-\frac{1}{2}} \right) d \\
 &= \frac{d^{\frac{1}{2}}}{\frac{1}{2}} + \frac{d^{\frac{1}{2}}}{\frac{1}{2}} + c \\
 &= \frac{2}{3} d^{\frac{3}{2}} + 2 d^{\frac{1}{2}} + c
 \end{aligned}$$

$$\begin{aligned}
 8. \int \frac{d + \sqrt{1-d^2}}{\sqrt{1-d^2}} d &= \int \frac{1}{\sqrt{1-d^2}} d + \int \frac{1}{\sqrt{1-d^2}} d \\
 &= \sin^{-1} d + \log|d + \sqrt{1-d^2}| + c
 \end{aligned}$$

$$\begin{aligned}
 9. \int \frac{\sin^2 d}{1 + \cos d} d &= \int \frac{1 - \cos^2 d}{1 + \cos d} d \\
 &= \int (1 - \cos d) d \\
 &= d - \sin d + C
 \end{aligned}$$

$$\begin{aligned}
 10. \int \frac{d}{1 - \sin d} &= \int \frac{(1 + \sin d) d}{1 - \sin^2 d} = \int \frac{1 + \sin d}{\cos^2 d} d \\
 &= \int \sec^2 d + \int \tan d \cdot \sec d = \tan d + \sec d + c
 \end{aligned}$$

$$\begin{aligned}
 11. \int \frac{d}{\sin^2 d \cos^2 d} &= \int \frac{(\cos^2 d + \sin^2 d) d}{\cos^2 d \sin^2 d} \\
 &= \int \left(\frac{1}{\sin^2 d} + \frac{1}{\cos^2 d} \right) d \\
 &= \int \operatorname{cosec}^2 d + \int \sec^2 d \\
 &= -\cot d + \tan d + c
 \end{aligned}$$

$$\begin{aligned}
 12. \int \frac{d}{\sin d \cos^2 d} &= \int \frac{\sin^2 d + \cos^2 d}{\sin d \cos^2 d} d \\
 &= \int \frac{\sin d}{\cos^2 d} d + \int \frac{1}{\sin d} d \\
 &= \int \sec d \tan d + \int \operatorname{cosec} d \\
 &= \sec d + \log|\operatorname{cosec} d - \cot d| + c
 \end{aligned}$$

$$\begin{aligned}
 13. \int \frac{1}{1 + \cos 8} d &= \int \frac{1}{2\cos^2 4} d \\
 &= \frac{1}{2} \int \sec^2 4 d \\
 &= \frac{\tan 4}{8} + c
 \end{aligned}$$



$$\begin{aligned}
 14. \quad \text{Let } I &= \int \frac{\sin 2}{\sin 5 \sin 3} d \\
 &= \int \frac{\sin(5-3)}{\sin 5 \sin 3} d \\
 &= \int \frac{\sin 5 \cos 3 - \cos 5 \sin 3}{\sin 5 \sin 3} d \\
 &= \int (\cot 3 - \cot 5) d \\
 &= \frac{1}{3} \log |\sin 3| - \frac{1}{5} \log |\sin 5| + c
 \end{aligned}$$

$$\begin{aligned}
 15. \quad \int \frac{\cos 2 - \cos 2\theta}{\cos - \cos \theta} d &= \int \frac{2(\cos^2 - \cos^2 \theta)}{\cos - \cos \theta} d \\
 &= 2 \int (\cos + \cos \theta) d \\
 &= 2(\sin + \cos \theta) + c
 \end{aligned}$$

$$\begin{aligned}
 16. \quad \int \frac{\cos - 1}{\cos + 1} d &= \int \frac{2\sin^2 \frac{-}{2}}{2\cos^2 \frac{-}{2}} d \\
 &= -\int \tan^2 \frac{-}{2} d = -\int \left(\sec^2 \frac{-}{2} - 1 \right) d \\
 &= \int \left(1 - \sec^2 \frac{-}{2} \right) d = -2 \tan \frac{-}{2} + c
 \end{aligned}$$

$$\begin{aligned}
 17. \quad \int \frac{+\sin}{1+\cos} d &= \int \frac{+2\sin \frac{-}{2} \cos \frac{-}{2}}{2\cos^2 \frac{-}{2}} d \\
 &= \frac{1}{2} \int \sec^2 \frac{-}{2} d + \int \tan \frac{-}{2} d \\
 &= \frac{1}{2} \cdot \frac{\tan \frac{-}{2}}{\frac{1}{2}} - \frac{1}{2} \int \frac{\tan \frac{-}{2}}{\frac{1}{2}} d + \int \tan \frac{-}{2} d \\
 &= \tan \frac{-}{2} + c
 \end{aligned}$$

$$\begin{aligned}
 18. \quad \int (1 - \cos) \operatorname{cosec}^2 d & \\
 &= \int \operatorname{cosec}^2 d - \int \cot \operatorname{cosec} d \\
 &= -\cot + \operatorname{cosec} + c \\
 &= \frac{1 - \cos}{\sin} + c \\
 &= \frac{2\sin^2 \frac{-}{2}}{2\sin \frac{-}{2} \cos \frac{-}{2}} + c \\
 &= \tan \left(\frac{-}{2} \right) + c
 \end{aligned}$$

$$\begin{aligned}
 19. \quad \int \sqrt{1 - \sin 2} d & \\
 &= \int \sqrt{\cos^2 + \sin^2 - 2\sin \cos} d \\
 &= \int \sqrt{(\cos - \sin)^2} d \\
 &= \int (\cos - \sin) d = \sin + \cos + c
 \end{aligned}$$

$$\begin{aligned}
 20. \quad \int \sqrt{1 + \sin \frac{-}{2}} d & \\
 &= \int \sqrt{\left(\sin^2 \frac{-}{4} + \cos^2 \frac{-}{4} + 2\sin \frac{-}{4} \cos \frac{-}{4} \right)} d \\
 &= \int \left(\sin \frac{-}{4} + \cos \frac{-}{4} \right) d = 4 \left(\sin \frac{-}{4} - \cos \frac{-}{4} \right) + c
 \end{aligned}$$

$$\begin{aligned}
 21. \quad \int \sqrt{2} \sqrt{1 + \sin} d &= \sqrt{2} \int \left(\sin \frac{-}{2} + \cos \frac{-}{2} \right) d \\
 &= 2 \int \sin \left(\frac{-}{2} + \frac{\pi}{4} \right) d \\
 &= -4 \cos \left(\frac{-}{2} + \frac{\pi}{4} \right) + c
 \end{aligned}$$

$$\therefore a = \frac{1}{2}, b = \frac{\pi}{4}$$

$$\begin{aligned}
 22. \quad \int (\sin 2 - \cos 2) d &= \frac{1}{\sqrt{2}} \sin(2 - a) + b \\
 \Rightarrow -\frac{1}{2} (\sin 2 + \cos 2) &= \frac{1}{\sqrt{2}} \sin(2 - a) + b \\
 \Rightarrow -\left[\frac{1}{\sqrt{2}} \sin 2 + \frac{1}{\sqrt{2}} \cos 2 \right] &= \sin(2 - a) + b\sqrt{2} \\
 \Rightarrow -\sin \left(2 + \frac{\pi}{4} \right) &= \sin(2 - a) + b\sqrt{2} \\
 \Rightarrow \sin \left(2 + \frac{5\pi}{4} \right) &= \sin(2 - a) + b\sqrt{2}
 \end{aligned}$$

$$\therefore b \text{ is any constant and } a = \frac{-5\pi}{4}$$

$$\begin{aligned}
 23. \quad \int \frac{d}{1 + \sin} &= \int \frac{d}{1 + \cos \left(\frac{\pi}{2} - \right)} \\
 &= \int \frac{d}{2\cos^2 \left(\frac{\pi}{4} - \frac{-}{2} \right)} \\
 &= \frac{1}{2} \int \sec^2 \left(\frac{-}{2} - \frac{\pi}{4} \right) d
 \end{aligned}$$



$$= \frac{1}{2} \cdot \frac{\tan\left(\frac{\alpha}{2} - \frac{\pi}{4}\right)}{\frac{1}{2}} + c$$

$$= \tan\left(\frac{\alpha}{2} - \frac{\pi}{4}\right) + c$$

∴ $a = \frac{-\pi}{4}$ and $b =$ arbitrary constant

24.
$$\int \frac{\cos \alpha}{\sin \cos(\alpha -)} d = \int \frac{\cos[(\alpha -) +]}{\sin \cos(\alpha -)} d$$

$$= \int \frac{\cos(\alpha -)\cos - \sin(\alpha -)\sin}{\sin \cos(\alpha -)} d$$

$$= \int [\cot - \tan(\alpha -)] d$$

$$= \log |\sin | - \log |\cos(\alpha -)| + c_1$$

$$= -\log \left| \frac{\cos(\alpha -)}{\sin} \right| + c_1$$

$$= -\log \left| \frac{\cos \alpha \cos + \sin \alpha \sin}{\sin} \right| + c_1$$

$$= -\log |\cos \alpha (\cot + \tan \alpha)| + c_1$$

$$= -\log |\cot + \tan \alpha| - \log |\cos \alpha| + c_1$$

$$= -\log |\cot + \tan \alpha| + c$$

25.
$$\int \sec^4 \tan d = \int \sec^3 \sec \tan d$$

Put $t = \sec \Rightarrow dt = \sec \tan d$

∴
$$\int \sec^4 \tan d = \int t^3 dt = \frac{t^4}{4} + c = \frac{1}{4} \sec^4 + c$$

26.
$$I_4 = \int \tan^4 d, I_6 = \int \tan^6 d$$

$$I_4 + I_6 = \int (\tan^4 + \tan^6) d$$

$$= \int \tan^4 (1 + \tan^2) d$$

$$= \int \tan^4 \cdot \sec^2 d$$

$$= \frac{1}{5} \tan^5 + c$$

Comparing with $a \tan^5 + b^{-5} + c$, we get

∴ $a = \frac{1}{5}, b = 0$

27.
$$I_4 - \frac{2}{3} I_2 = \int \left(\sec^4 - \frac{2}{3} \sec^2 \right) d$$

$$= \int \sec^2 \left(\sec^2 - \frac{2}{3} \right) d$$

$$= \int \sec^2 \left(\frac{3 \sec^2 - 2}{3} \right) d$$

$$= \frac{1}{3} \int \sec^2 (3 \tan^2 + 1) d$$

Put $\tan = t \Rightarrow \sec^2 d = dt$

∴
$$I_4 - \frac{2}{3} I_2 = \frac{1}{3} \int (3t^2 + 1) dt$$

$$= \frac{1}{3} (t^3 + t) + c$$

$$= \frac{1}{3} [t(t^2 + 1)] + c$$

$$= \frac{1}{3} [\tan (\tan^2 + 1)] + c$$

$$= \frac{1}{3} \sec^2 \tan + c$$

28. Put $\log = t \Rightarrow \frac{1}{d} = dt$

∴
$$\int \frac{\cos(\log)}{d} = \int \cos t dt$$

$$= \sin t + c = \sin(\log) + c$$

29. Put $2 = t \Rightarrow 2 d = dt$

$$\int e^{2 \log^2} e^2 d = \frac{1}{2} \int e^t 2 dt$$

$$= \frac{1}{2} \int (2e)^t dt$$

$$= \frac{(2e)^t}{2 \log(2e)} + c$$

$$= \frac{2^2 e^2}{2(\log 2 + 1)} + c$$

30. Put $t = \tan^{-1} 2 \Rightarrow dt = \frac{1}{1 + 4} \cdot 2 d$

∴
$$\int \frac{2 \tan^{-1} 2}{1 + 4} d = \int t dt = \frac{t^2}{2} + c$$

$$= \frac{1}{2} (\tan^{-1} 2)^2 + c$$

31. Let $I = \int (e + e^{-})^2 (e - e^{-}) d$

Put $e + e^{-} = t \Rightarrow (e - e^{-}) d = dt$

∴
$$I = \int t^2 \cdot dt = \frac{t^3}{3} + c = \frac{1}{3} (e + e^{-})^3 + c$$

32. Put $t = + \log \sec$

$\Rightarrow dt = (1 + \tan) d$

∴
$$\int \frac{1 + \tan}{+ \log \sec} d = \int \frac{1}{t} dt = \log t + c$$

$= \log(+ \log \sec) + c$



$$33. \text{ Put } 1 + \log x = t \Rightarrow \frac{1}{x} dx = dt$$

$$\therefore \int \frac{1 + \log x}{x} dx = \int t dt = \frac{t^2}{2} + c = \frac{(1 + \log x)^2}{2} + c$$

$$34. \text{ Put } a^2 + b^2 \sin^2 x = t$$

$$\Rightarrow b^2 \sin 2x dx = dt$$

$$\therefore \int \frac{\sin 2x}{a^2 + b^2 \sin^2 x} dx = \frac{1}{b^2} \int \frac{dt}{t}$$

$$= \frac{1}{b^2} \log t + c$$

$$= \frac{1}{b^2} \log(a^2 + b^2 \sin^2 x) + c$$

$$35. \text{ Put } x^3 = t \Rightarrow 3x^2 dx = dt$$

$$\therefore \int x^2 \sec^3 x^3 dx = \frac{1}{3} \int \sec t dt$$

$$= \frac{1}{3} \log(\sec t + \tan t) + c$$

$$= \frac{1}{3} \log(\sec^3 x + \tan^3 x) + c$$

$$36. \int \frac{e^x (e^x + 1)}{\cos^2(e^x)} dx = \int e^x (e^x + 1) \sec^2(e^x) dx$$

$$\text{Put } e^x = t \Rightarrow (e^x + 1) e^x dx = dt$$

$$\therefore \int \frac{e^x (e^x + 1)}{\cos^2(e^x)} dx = \int \sec^2 t dt = \tan t + c$$

$$= \tan(e^x) + c$$

$$37. \text{ Let } I = \int \frac{1 + e^x}{1 + e^{-x}} dx = \int \frac{e^x (1 + e^x)}{e^x + 1} dx$$

$$\text{Put } e^x + 1 = t \Rightarrow e^x (1 + e^x) dx = dt$$

$$\therefore I = \int \frac{dt}{t} = \log |t| + c = \log |1 + e^x| + c$$

$$38. \text{ Put } x + \tan^{-1} x = t$$

$$\Rightarrow \left(1 + \frac{1}{1+x^2}\right) dx = dt \Rightarrow \frac{2 + x^2}{1+x^2} dx = dt$$

$$\therefore \int \left[\frac{(x^2 + 2)a^{(x + \tan^{-1} x)}}{1+x^2} \right] dx = \int a^t dt$$

$$= \frac{a^t}{\log a} + c$$

$$= \frac{a^{x + \tan^{-1} x}}{\log a} + c$$

$$39. \text{ Let } I = \int \frac{1}{4\sqrt{x} + x} dx = \int \frac{1}{\sqrt{x}(4 + \sqrt{x})} dx$$

$$\text{Put } 4 + \sqrt{x} = t \Rightarrow \frac{1}{2\sqrt{x}} dx = dt$$

$$\therefore I = \int \frac{2dt}{t} = 2 \log t + c$$

$$= 2 \log(\sqrt{x} + 4) + c$$

$$40. \text{ Let } I = \int \frac{\sin x}{\cos x - \sin x - 1} dx$$

$$\text{Put } \cos x - \sin x - 1 = t$$

$$\Rightarrow \sin x dx = -dt$$

$$\therefore I = -\int \frac{dt}{t} = -\log |t| + c$$

$$= -\log |\cos x - \sin x - 1| + c$$

$$41. \text{ Let } I = \int \frac{dx}{\sin x \cos x + 3 \cos^2 x} = \int \frac{\sec^2 x}{\tan x + 3} dx$$

$$\text{Put } \tan x + 3 = t$$

$$\Rightarrow \sec^2 x dx = dt$$

$$\therefore I = \int \frac{dt}{t} = \log |t| + c = \log |\tan x + 3| + c$$

$$42. \int \frac{\cos x}{\sqrt{1 + \sin x}} dx = 2\sqrt{1 + \sin x} + c$$

$$\dots \left[\because \int \frac{f'(x)}{\sqrt{f(x)}} dx = 2\sqrt{f(x)} + c \right]$$

$$= 2\sqrt{\left(\sin \frac{x}{2} + \cos \frac{x}{2}\right)^2} + c$$

$$= 2 \left[\sin \left(\frac{x}{2}\right) + \cos \left(\frac{x}{2}\right) \right] + c$$

$$43. \text{ Put } 1 + \log \tan \frac{x}{2} = t$$

$$\Rightarrow \left(\frac{1}{\tan \left(\frac{x}{2}\right)} \cdot \sec^2 \frac{x}{2} \cdot \frac{1}{2} \right) dx = dt$$

$$\Rightarrow \operatorname{cosec} x dx = dt$$

$$\therefore \int \frac{\operatorname{cosec} x}{\cos^2 \left(1 + \log \tan \frac{x}{2}\right)} dx = \int \frac{dt}{\cos^2 t}$$

$$= \int \sec^2 t dt$$

$$= \tan t + c$$

$$= \tan \left(1 + \log \tan \frac{x}{2}\right) + c$$



44. Put $x = t^2 \Rightarrow dx = 2t dt$
 $\therefore \int \frac{\log \sqrt{x}}{3} dx = \int \frac{\log t}{3t^2} (2t dt)$
 $= \frac{2}{3} \int \frac{\log t}{t} dt$
 $= \frac{2}{3} \cdot \frac{(\log t)^2}{2} + c$
 $= \frac{(\log \sqrt{x})^2}{3} + c$
45. Let $I = \int \left[\frac{\log^{-1}}{1 + (\log^{-1})^2} \right] dx$
 Put $\log^{-1} = t$
 $\therefore dx = e^t \Rightarrow dx = e^t dt$
 $\therefore I = \int \left(\frac{t-1}{1+t^2} \right) e^t dt$
 $= \int \left[\frac{1+t^2-2t}{(1+t^2)^2} \right] e^t dt$
 $= \int e^t \left[\frac{1}{1+t^2} + \frac{(-2t)}{(1+t^2)^2} \right] dt$
 $= e^t \left(\frac{1}{1+t^2} \right) + c$
 $\dots \left[\because \int e^{f(x)} [f'(x) + f(x)] dx = e^{f(x)} f(x) + c \right]$
 $= \frac{e^t}{1 + (\log^{-1})^2} + c$
46. $\int \frac{\sqrt{\tan x}}{\sin x \cos x} dx = \int \frac{\tan x}{\sqrt{\tan x} \sin x \cos x} dx$
 $= \int \frac{\sin x \sec x}{\sqrt{\tan x} \sin x \cos x} dx$
 $= \int \frac{\sec^2 x}{\sqrt{\tan x}} dx$
 Put $t = \tan x \Rightarrow dt = \sec^2 x dx$
 $\therefore \int \frac{\sqrt{\tan x}}{\sin x \cos x} dx = \int \frac{1}{\sqrt{t}} dt = 2t^{1/2} + c = 2\sqrt{\tan x} + c$
47. Let $I = \int \frac{\sin^{-1} \left[\tan^{-1} \left(\frac{x}{4} \right) \right]}{1 + x^8} dx$
 Put $x^4 = t \Rightarrow 4x^3 dx = dt$
 $\therefore I = \frac{1}{4} \int \frac{\sin^{-1}(\tan^{-1} t)}{1+t^2} dt$

- Put $\tan^{-1} t = z \Rightarrow \frac{1}{1+t^2} dt = dz$
 $\therefore I = \frac{1}{4} \int \sin z dz = \frac{1}{4} (-\cos z) + c$
 $= -\frac{1}{4} \cos(\tan^{-1} t) + c = -\frac{1}{4} \cos[\tan^{-1} \left(\frac{x}{4} \right)] + c$
48. Let $I = \int \operatorname{cosec}^4 x dx = \int \operatorname{cosec}^2 x \cdot \operatorname{cosec}^2 x dx$
 $= \int \operatorname{cosec}^2 x (1 + \cot^2 x) dx$
 $= \int \operatorname{cosec}^2 x dx + \int \cot^2 x \cdot \operatorname{cosec}^2 x dx$
 In 2nd integral, put $\cot x = t \Rightarrow -\operatorname{cosec}^2 x dx = dt$
 $\therefore I = \int \operatorname{cosec}^2 x dx - \int t^2 dt$
 $= -\cot x - \frac{t^3}{3} + c = -\cot x - \frac{\cot^3 x}{3} + c$
49. Let $I = \int (x+1)(x+2)^7(x+3) dx$
 Put $x+2 = t \Rightarrow dx = dt$
 $\therefore I = \int (t-1)t^7(t+1) dt = \int (t^2-1)t^7 dt$
 $= \int (t^9 - t^7) dt$
 $= \frac{t^{10}}{10} - \frac{t^8}{8} + c$
 $= \frac{(x+2)^{10}}{10} - \frac{(x+2)^8}{8} + c$
50. $\int \sec x dx = \log(\sec x + \tan x) + c$
 $= \log \left(\frac{\sec^2 x - \tan^2 x}{\sec x - \tan x} \right) + c$
 $= \log \left(\frac{1}{\sec x - \tan x} \right) + c$
 $= -\log(\sec x - \tan x) + c$
51. $\int \frac{1 - \tan x}{1 + \tan x} dx = \int \tan \left(\frac{\pi}{4} - x \right) dx$
 $= \frac{-\log \cos \left(\frac{\pi}{4} - x \right)}{-1} + c$
 $= \log \cos \left(\frac{\pi}{4} - x \right) + c$
 $= \log \sin \left[\frac{\pi}{2} - \left(\frac{\pi}{4} - x \right) \right] + c$
 $= \log \sin \left(\frac{\pi}{4} + x \right) + c$



$$\begin{aligned}
 52. \quad \int \frac{d}{\sin \theta + \cos \theta} &= \frac{1}{\sqrt{2}} \int \frac{d}{\sin \theta \cos \frac{\pi}{4} + \cos \theta \sin \frac{\pi}{4}} \\
 &= \frac{1}{\sqrt{2}} \int \operatorname{cosec} \left(\theta + \frac{\pi}{4} \right) d\theta \\
 &= \frac{1}{\sqrt{2}} \log \tan \left(\frac{\theta}{2} + \frac{\pi}{8} \right) + c
 \end{aligned}$$

$$\begin{aligned}
 53. \quad \int \frac{1}{\sqrt{1 + \sin \theta}} d\theta &= \int \frac{1}{\sin \frac{\theta}{2} + \cos \frac{\theta}{2}} d\theta \\
 &= \int \frac{1}{\sqrt{2} \sin \left(\frac{\theta}{2} + \frac{\pi}{4} \right)} d\theta \\
 &= \frac{1}{\sqrt{2}} \int \operatorname{cosec} \left(\frac{\theta}{2} + \frac{\pi}{4} \right) d\theta \\
 &= \sqrt{2} \log \tan \left(\frac{\theta}{8} + \frac{\pi}{4} \right) + c
 \end{aligned}$$

$$\begin{aligned}
 54. \quad \text{Let } I &= \sqrt{2} \int \frac{\sin \theta}{\sin \left(\theta - \frac{\pi}{4} \right)} d\theta \\
 \text{Put } \theta - \frac{\pi}{4} &= t \Rightarrow d\theta = dt \\
 \therefore I &= \sqrt{2} \int \frac{\sin \left(\frac{\pi}{4} + t \right)}{\sin t} dt = \int \frac{\cos t + \sin t}{\sin t} dt \\
 &= \int \cot t dt + \int dt = \log |\sin t| + t + c_1 \\
 &= -\frac{\pi}{4} + \log \left| \sin \left(\theta - \frac{\pi}{4} \right) \right| + c_1 \\
 &= \log \left| \sin \left(\theta - \frac{\pi}{4} \right) \right| + c, \text{ where } c = c_1 - \frac{\pi}{4}
 \end{aligned}$$

$$\begin{aligned}
 55. \quad \int (1 + 2 \tan^2 \theta + 2 \tan \theta \sec \theta)^{1/2} d\theta \\
 &= \int (\sec^2 \theta + \tan^2 \theta + 2 \tan \theta \sec \theta)^{1/2} d\theta \\
 &= \int (\sec \theta + \tan \theta) d\theta \\
 &= \log(\sec \theta + \tan \theta) + \log \sec \theta + c \\
 &= \log \sec \theta (\sec \theta + \tan \theta) + c
 \end{aligned}$$

$$\begin{aligned}
 56. \quad \int \frac{d}{e + e^{-\theta}} &= \int \frac{e}{e^2 + 1} d\theta \\
 &= \int \frac{dt}{t^2 + 1} \\
 &\quad \dots [\text{Put } e^\theta = t \Rightarrow e d\theta = dt] \\
 &= \tan^{-1}(t) + c \\
 &= \tan^{-1}(e) + c
 \end{aligned}$$

$$\begin{aligned}
 57. \quad \text{Let } I &= \int \frac{1}{1 + e^{-\theta}} d\theta = \int \frac{1}{1 + \frac{1}{e^\theta}} d\theta \\
 &= \int \frac{e^\theta}{1 + e^\theta} d\theta
 \end{aligned}$$

$$\text{Put } e^\theta = t \Rightarrow e^\theta d\theta = dt$$

$$\begin{aligned}
 \therefore I &= -\int \frac{1}{1+t} dt \\
 &= -\log(1+t) + c \\
 &= -\log(1+e^\theta) + c \\
 &= -\log \left(\frac{1+e}{e} \right) + c \\
 &= \log \left(\frac{e}{1+e} \right) + c
 \end{aligned}$$

$$\begin{aligned}
 58. \quad \text{Let } I &= \int \frac{d}{e + e^{-\theta} + 2} \\
 &= \int \frac{e d\theta}{e^2 + 2e + 1}
 \end{aligned}$$

$$\text{Put } e^\theta = t \Rightarrow e d\theta = dt$$

$$\begin{aligned}
 \therefore I &= \int \frac{dt}{t^2 + 2t + 1} = \int \frac{dt}{(t+1)^2} \\
 &= \frac{-1}{t+1} + c = \frac{-1}{e+1} + c
 \end{aligned}$$

$$59. \quad \text{Put } t^2 = \theta \Rightarrow d\theta = \frac{dt}{2}$$

$$\begin{aligned}
 \therefore \int \frac{d\theta}{1 + \theta^2} &= \frac{1}{2} \int \frac{dt}{1+t^2} \\
 &= \frac{1}{2} \tan^{-1} t + c \\
 &= \frac{1}{2} \tan^{-1}(\sqrt{\theta}) + c
 \end{aligned}$$

$$60. \quad \int \frac{1}{(1 + \sqrt{\theta}) \sqrt{\theta}} d\theta = \int \frac{1}{\left[1 + (\sqrt{\theta})^2 \right] \sqrt{\theta}} d\theta$$

$$\text{Put } \sqrt{\theta} = t \Rightarrow \frac{1}{2\sqrt{\theta}} d\theta = dt$$

$$\begin{aligned}
 \therefore \int \frac{1}{(1 + \sqrt{\theta}) \sqrt{\theta}} d\theta &= \int \frac{2}{1+t^2} dt \\
 &= 2 \tan^{-1} t + A
 \end{aligned}$$

$$\therefore \int \frac{1}{(1 + \sqrt{\theta}) \sqrt{\theta}} d\theta = 2 \tan^{-1} \sqrt{\theta} + A$$

$$\therefore f(\theta) = 2 \tan^{-1} \sqrt{\theta}$$



$$61. \text{ Let } I = \int \frac{2}{1 + (\sqrt[3]{x})^2} dx$$

$$\text{Put } \sqrt[3]{x} = t \Rightarrow 3^{-2} dx = dt$$

$$\begin{aligned} \therefore I &= \frac{1}{3} \int \frac{dt}{1+t^2} \\ &= \frac{1}{3} \tan^{-1} t + c \\ &= \frac{1}{3} \tan^{-1} \sqrt[3]{x} + c \end{aligned}$$

$$62. \text{ Put } x^4 = t \Rightarrow 4x^3 dx = dt$$

$$\begin{aligned} \therefore \int \frac{x^3 dx}{1+x^8} &= \frac{1}{4} \int \frac{dt}{1+t^2} = \frac{1}{4} \tan^{-1} t + c \\ &= \frac{1}{4} \tan^{-1} (x^4) + c \end{aligned}$$

$$\begin{aligned} 63. \int \frac{dx}{16x^2+9} &= \int \frac{dx}{(4x)^2+3^2} \\ &= \frac{1}{12} \tan^{-1} \left(\frac{4x}{3} \right) + c \end{aligned}$$

$$\begin{aligned} 64. \text{ Let } I &= \int \frac{1}{\sqrt{9-16x^2}} dx \\ &= \int \frac{1}{\sqrt{3^2-(4x)^2}} dx \\ &= \frac{1}{4} \sin^{-1} \left(\frac{4x}{3} \right) + c \end{aligned}$$

Comparing with $\alpha \sin^{-1}(\beta x) + c$, we get

$$\alpha = \frac{1}{4}, \beta = \frac{4}{3}$$

$$\therefore \alpha + \frac{1}{\beta} = \frac{1}{4} + \frac{3}{4} = 1$$

$$\begin{aligned} 65. \text{ Let } I &= \int \frac{dx}{\sqrt{16-9x^2}} \\ &= \int \frac{1}{\sqrt{4^2-(3x)^2}} dx \\ &= \frac{1}{3} \sin^{-1} \frac{3x}{4} + C \end{aligned}$$

Comparing with $A \sin^{-1}(Bx) + C$, we get

$$A = \frac{1}{3}, B = \frac{3}{4}$$

$$\therefore A + B = \frac{1}{3} + \frac{3}{4} = \frac{13}{12}$$

$$\begin{aligned} 66. \text{ Let } I &= \int \frac{dx}{\sqrt{10-x^2}} \\ &= \int \frac{dx}{\sqrt{8-x^2}} \\ &= \int \frac{x^3 dx}{4\sqrt{8-x^2}} \end{aligned}$$

$$\text{Put } x^4 = t \Rightarrow 4x^3 dx = dt$$

$$\begin{aligned} \therefore I &= \frac{1}{4} \int \frac{dt}{t\sqrt{t^2-1}} \\ &= \frac{1}{4} \sec^{-1} t + c \\ &= \frac{1}{4} \sec^{-1} (x^4) + c \end{aligned}$$

$$\begin{aligned} 67. \text{ Let } I &= \int \frac{\sin 2x}{\sin^4 x + \cos^4 x} dx \\ &= \int \frac{2 \sin x \cos x}{\sin^4 x + \cos^4 x} dx \\ &= \int \frac{2 \tan x \sec^2 x}{1 + \tan^4 x} dx \end{aligned}$$

$$\text{Put } \tan^2 x = t \Rightarrow 2 \tan x \sec^2 x dx = dt$$

$$\therefore I = \int \frac{dt}{1+t^2} = \tan^{-1} t + c = \tan^{-1}(\tan^2 x) + c$$

$$\begin{aligned} 68. \text{ Let } I &= \int \frac{\sin 2x}{\sin^4 x + \cos^4 x} dx \\ &= \int \frac{2 \sin x \cos x}{\sin^4 x + \cos^4 x} dx \\ &= \int \frac{2 \tan x \sec^2 x}{1 + \tan^4 x} dx \end{aligned}$$

$$\text{Put } \tan^2 x = t \Rightarrow 2 \tan x \sec^2 x dx = dt$$

$$\therefore I = \int \frac{dt}{1+t^2} = \tan^{-1} t + c = \tan^{-1}(\tan^2 x) + c$$

Comparing with $\tan^{-1}[f(x)] + c$, we get
 $f(x) = \tan^2 x$

$$\therefore f\left(\frac{\pi}{3}\right) = \tan^2 \frac{\pi}{3} = (\sqrt{3})^2 = 3$$

$$\begin{aligned} 69. \text{ Let } I &= \int \frac{\sin^2 x \cos^2 x}{(\sin^5 x + \cos^3 x \sin^2 x + \sin^3 x \cos^2 x + \cos^5 x)^2} dx \\ &= \int \frac{\sin^2 x \cos^2 x}{\left[(\sin^2 x + \cos^2 x)(\sin^3 x + \cos^3 x) \right]^2} dx \\ &= \int \frac{\sin^2 x \cos^2 x}{(\sin^3 x + \cos^3 x)^2} dx \end{aligned}$$



Dividing numerator and denominator by \cos^6 , we get

$$I = \int \frac{\tan^2 \cdot \sec^2}{(1 + \tan^3)^2} dx$$

$$\text{Put } 1 + \tan^3 = t \Rightarrow 3 \tan^2 \sec^2 dx = dt$$

$$\therefore I = \frac{1}{3} \int \frac{dt}{t^2} = -\frac{1}{3t} + c = \frac{-1}{3(1 + \tan^3)} + c$$

70. Put $a = t \Rightarrow a \log_e a dx = dt$

$$\begin{aligned} \therefore \int \frac{a}{\sqrt{1-a^2}} dx &= \frac{1}{\log_e a} \int \frac{dt}{\sqrt{1-t^2}} \\ &= \frac{1}{\log_e a} \sin^{-1}(t) + c = \frac{\sin^{-1}(a)}{\log_e a} + c \end{aligned}$$

71. $\int \frac{1}{\sqrt{8+2x-x^2}} dx = \int \frac{1}{\sqrt{8+1-(x^2-2x+1)}} dx$

$$= \int \frac{1}{\sqrt{3^2 - (x-1)^2}} dx$$

$$= \sin^{-1} \left(\frac{x-1}{3} \right) + c$$

72. $\int \frac{1}{\sqrt{3-6x-9x^2}} dx = \int \frac{1}{\sqrt{3-(9x^2+6x)}} dx$

$$= \int \frac{1}{\sqrt{4-(9x^2+6x+1)}} dx$$

$$= \int \frac{1}{\sqrt{2^2 - (3x+1)^2}} dx$$

$$= \frac{1}{3} \sin^{-1} \left(\frac{3x+1}{2} \right) + c$$

73. $I = \int \frac{dx}{\sqrt{(1-x)(x-2)}} = \int \frac{dx}{\sqrt{-2+3x-x^2}}$

$$= \int \frac{dx}{\sqrt{-2+\frac{9}{4}-\left(x-\frac{3}{2}\right)^2}}$$

$$= \int \frac{dx}{\sqrt{\left(\frac{1}{2}\right)^2 - \left(x-\frac{3}{2}\right)^2}}$$

$$= \sin^{-1} \left(\frac{x-\frac{3}{2}}{\frac{1}{2}} \right) + C$$

$\therefore I = \sin^{-1}(2x-3) + C$

74. Let $I = \int \sqrt{\frac{-5}{-7}} dx$

$$= \int \frac{-5}{\sqrt{(x-7)(x-5)}} dx$$

$$= \int \frac{-5}{\sqrt{x^2-12x+35}} dx$$

$$= \frac{1}{2} \int \frac{2x-10}{\sqrt{x^2-12x+35}} dx$$

$$= \frac{1}{2} \int \frac{2x-12+2}{\sqrt{x^2-12x+35}} dx$$

$$= \frac{1}{2} \int \frac{2x-12}{\sqrt{x^2-12x+35}} dx + \frac{2}{2} \int \frac{dx}{\sqrt{x^2-12x+36-1}}$$

$$= \frac{1}{2} \times 2\sqrt{x^2-12x+35} + \int \frac{dx}{\sqrt{(x-6)^2-1}}$$

$$= \sqrt{x^2-12x+35} + \log \left| (x-6) + \sqrt{x^2-12x+35} \right| + c$$

Comparing with $A\sqrt{x^2-12x+35}$

$$+ \log \left| (x-6) + \sqrt{x^2-12x+35} \right| + c, \text{ we get}$$

$$A = 1$$

75. Let $I = \int \sqrt{x^2+2x+5} dx$

$$= \int \sqrt{(x+1)^2+2^2} dx$$

$$= \frac{+1}{2} \sqrt{x^2+2x+5} + 2 \log \left| x+1 + \sqrt{x^2+2x+5} \right| + c$$

76. Let $I = \int \frac{\sec^8}{\operatorname{cosec}} dx$

$$= \int \frac{\sin}{\cos^8} dx$$

$$= \int \tan^7 \cdot \sec^7 dx$$

$$= \int \sec^6 \sec \tan dx$$

Put $\sec = t \Rightarrow \sec \tan dx = dt$

$\therefore I = \int t^6 dt$

$$= \frac{t^7}{7} + c$$

$$= \frac{\sec^7}{7} + c$$



$$77. \int \frac{d}{5+4\cos} = \int \frac{d}{5+4 \left[\frac{1-\tan^2 \frac{\theta}{2}}{1+\tan^2 \frac{\theta}{2}} \right]}$$

$$= \int \frac{\sec^2 \frac{\theta}{2} d}{9+\tan^2 \frac{\theta}{2}}$$

$$\text{Put } \tan \frac{\theta}{2} = t \Rightarrow \frac{1}{2} \sec^2 \frac{\theta}{2} d = dt$$

$$\therefore \int \frac{d}{5+4\cos} = 2 \int \frac{dt}{3^2+t^2}$$

$$= \frac{2}{3} \tan^{-1} \left(\frac{t}{3} \right) + c$$

$$= \frac{2}{3} \tan^{-1} \left(\frac{1}{3} \tan \frac{\theta}{2} \right) + c$$

$$78. \text{ Let } I = \int \frac{d}{7+5\cos} = \int \frac{d}{7+5 \left[\frac{1-\tan^2 \left(\frac{\theta}{2} \right)}{1+\tan^2 \left(\frac{\theta}{2} \right)} \right]}$$

$$= \int \frac{\sec^2 \left(\frac{\theta}{2} \right) d}{12+2\tan^2 \left(\frac{\theta}{2} \right)} = \int \frac{\frac{1}{2} \sec^2 \left(\frac{\theta}{2} \right) d}{6+\tan^2 \left(\frac{\theta}{2} \right)}$$

$$\text{Put } \tan \frac{\theta}{2} = t \Rightarrow \frac{1}{2} \sec^2 \frac{\theta}{2} d = dt$$

$$\therefore I = \int \frac{dt}{t^2+(\sqrt{6})^2} = \frac{1}{\sqrt{6}} \tan^{-1} \left(\frac{t}{\sqrt{6}} \right) + c$$

$$= \frac{1}{\sqrt{6}} \tan^{-1} \left(\frac{\tan \frac{\theta}{2}}{\sqrt{6}} \right) + c$$

$$79. \int \frac{d}{1+3\sin^2} = \int \frac{d}{\sin^2 + \cos^2 + 3\sin^2}$$

$$= \int \frac{d}{4\sin^2 + \cos^2}$$

$$= \int \frac{\sec^2 d}{4\tan^2 + 1}$$

$$= \frac{1}{4} \int \frac{\sec^2 d}{\tan^2 + \frac{1}{4}}$$

$$\text{Put } t = \tan \theta \Rightarrow dt = \sec^2 \theta d\theta$$

$$\therefore \int \frac{d}{1+3\sin^2} = \frac{1}{4} \int \frac{dt}{t^2 + \left(\frac{1}{2} \right)^2} = \frac{1}{4} \cdot 2 \tan^{-1} (2t) + c$$

$$= \frac{1}{2} \tan^{-1} (2t) + c$$

$$= \frac{1}{2} \tan^{-1} (2 \tan \theta) + c$$

$$80. \int \frac{5 \tan \theta}{\tan \theta - 2} d\theta = \int \frac{5 \sin \theta}{\sin \theta - 2 \cos \theta} d\theta$$

$$= \int \frac{(\sin \theta - 2 \cos \theta) + 2(\cos \theta + 2 \sin \theta)}{\sin \theta - 2 \cos \theta} d\theta$$

$$= \int d\theta + 2 \int \frac{\cos \theta + 2 \sin \theta}{\sin \theta - 2 \cos \theta} d\theta$$

$$= \theta + 2 \log |\sin \theta - 2 \cos \theta| + k$$

$$\therefore a = 2$$

$$81. \int \frac{\sin \theta d\theta}{\sin \theta - \cos \theta} = \frac{1}{2} \int \frac{2 \sin \theta}{\sin \theta - \cos \theta} d\theta$$

$$= \frac{1}{2} \int \frac{(\sin \theta - \cos \theta) + (\sin \theta + \cos \theta)}{\sin \theta - \cos \theta} d\theta$$

$$= \frac{1}{2} \int \left(1 + \frac{\sin \theta + \cos \theta}{\sin \theta - \cos \theta} \right) d\theta$$

$$= \frac{1}{2} [\theta + \log |\sin \theta - \cos \theta|] + c$$

$$82. \int \frac{4e^{-\theta} - 25}{2e^{-\theta} - 5} d\theta = \int \frac{5(2e^{-\theta} - 5) - 3(2e^{-\theta})}{2e^{-\theta} - 5} d\theta$$

$$= 5 \int d\theta - 3 \int \frac{2e^{-\theta}}{2e^{-\theta} - 5} d\theta$$

$$= 5\theta - 3 \log |2e^{-\theta} - 5| + c$$

$$\therefore A = 5 \text{ and } B = -3$$

$$83. \int \frac{d\theta}{\sin(a-\theta)\sin(b-\theta)}$$

$$= \frac{1}{\sin(a-b)} \int \frac{\sin\{(\theta-b) - (\theta-a)\}}{\sin(a-\theta)\sin(b-\theta)} d\theta$$

$$= \frac{1}{\sin(a-b)} \int \frac{\sin(b-a)\cos(\theta-b)\cos(\theta-a) - \cos(b-a)\sin(\theta-b)\sin(\theta-a)}{\sin(a-\theta)\sin(b-\theta)} d\theta$$

$$= \frac{1}{\sin(a-b)} \left[\int \cot(\theta-a) d\theta - \int \cot(\theta-b) d\theta \right]$$

$$= \frac{1}{\sin(a-b)} \left[\log |\sin(\theta-a)| - \log |\sin(\theta-b)| \right] + c$$

$$= \frac{1}{\sin(a-b)} \log \left| \frac{\sin(\theta-a)}{\sin(\theta-b)} \right| + c$$



84. Let $I = \int \sqrt{e^{-1}} dt$

Put $e^{-1} = t^2$

$\Rightarrow e^{-1} dt = 2t dt$

$\Rightarrow dt = \frac{2t}{t^2+1} dt$

$\therefore I = \int t \cdot \frac{2t}{t^2+1} dt = \int \frac{2t^2}{t^2+1} dt$

$= \int \frac{2(t^2+1)-2}{t^2+1} dt$

$= 2 \int dt - 2 \int \frac{dt}{t^2+1}$

$= 2t - 2 \tan^{-1} t + c$

$= 2(\sqrt{e^{-1}} - \tan^{-1} \sqrt{e^{-1}}) + c$

85. Let $I = \int \frac{1}{(e^x + e^{-x})^2} dx = \int \frac{e^{2x}}{(e^{2x} + 1)^2} dx$

Put $e^{2x} + 1 = t \Rightarrow 2e^{2x} dx = dt$

$\therefore I = \frac{1}{2} \int \frac{1}{t^2} dt = -\frac{1}{2} \cdot \frac{1}{t} + c$

$= \frac{-1}{2(e^x + 1)} + c$

86. $\int \frac{1}{\sqrt{1-e^2}} dx = \int \frac{e^{-x}}{\sqrt{e^{-2x}-1}} dx$

Put $e^{-x} = t \Rightarrow -e^{-x} dx = dt$

$\therefore \int \frac{1}{\sqrt{1-e^2}} dx = -\int \frac{1}{\sqrt{t^2-1}} dt$

$= -\log[t + \sqrt{t^2-1}] + c$

$= -\log[e^{-x} + \sqrt{e^{-2x}-1}] + c$

$= -\log\left[\frac{1}{e} + \frac{\sqrt{1-e^2}}{e}\right] + c$

$= -\log[1 + \sqrt{1-e^2}] + \log e + c$

$= -\log[1 + \sqrt{1-e^2}] + c$

87. Put $x = \sin \theta \Rightarrow dx = \cos \theta d\theta$

$\therefore \int \frac{1+x^2}{\sqrt{1-x^2}} dx = \int \frac{1+\sin^2 \theta}{\sqrt{1-\sin^2 \theta}} \cdot \cos \theta d\theta$

$= \int (1 + \sin^2 \theta) d\theta$

$= \int d\theta + \int \left(\frac{1-\cos 2\theta}{2}\right) d\theta$

$= \theta + \frac{1}{2} \left(\theta - \frac{\sin 2\theta}{2}\right) + c$

$= \frac{3}{2}\theta - \frac{\sin \theta \cos \theta}{2} + c$

$= \frac{3}{2}\theta - \frac{\sin \theta \sqrt{1-\sin^2 \theta}}{2} + c$

$= \frac{3}{2} \sin^{-1} x - \frac{1}{2} \sqrt{1-x^2} + c$

88. Let $I = \int \frac{1}{2(x^4+1)^{\frac{3}{4}}} dx = \int \frac{dx}{5\left(1+\frac{1}{4}\right)^{\frac{3}{4}}}$

Put $1 + \frac{1}{4} = t \Rightarrow \frac{-4}{5} dx = dt$

$\therefore I = -\frac{1}{4} \int \frac{dt}{t^{\frac{3}{4}}} = -\frac{1}{4} \times 4t^{\frac{1}{4}} + c = -t^{\frac{1}{4}} + c$

$= -\left(1 + \frac{1}{4}\right)^{\frac{1}{4}} + c = -\left(\frac{4+1}{4}\right)^{\frac{1}{4}} + c$

89. $\int \frac{2^{12} + 5^9}{(x^5 + x^3 + 1)^3} dx$

$= \int \frac{15\left(\frac{2}{3} + \frac{5}{6}\right)}{15\left(1 + \frac{1}{2} + \frac{1}{5}\right)^3} dx$

Put $1 + \frac{1}{2} + \frac{1}{5} = t \Rightarrow \left(\frac{-2}{3} - \frac{5}{6}\right) dx = dt$

$\therefore \int \frac{2^{12} + 5^9}{(x^5 + x^3 + 1)^3} dx = -\int \frac{dt}{t^3}$

$= \frac{1}{2t^2} + C$

$= \frac{1}{2\left(1 + \frac{1}{2} + \frac{1}{5}\right)^2} + C$

$= \frac{10}{2\left(x^5 + x^3 + 1\right)^2} + C$



$$90. \text{ Put } \frac{-1}{+2} = t \Rightarrow \frac{1}{(+2)^2} d \frac{1}{3} dt$$

$$\begin{aligned} \therefore \int \frac{1}{\left[(-1)^3(+2)^5\right]^{1/4}} d &= \int \frac{1}{(-1)^{3/4} (+2)^{-3/4} (+2)^2} d \\ &= \frac{1}{3} \int t^{-3/4} dt = \frac{1}{3} \cdot \frac{t^4}{\frac{4}{4}} + c \\ &= \frac{4}{3} \left(\frac{-1}{+2} \right)^{1/4} + c \end{aligned}$$

$$\begin{aligned} 91. \text{ Let } I &= \int \frac{(-2)d}{\left\{(-2)^2(+3)\right\}^{1/3}} \\ &= \int \frac{d}{(-2)^{-1/3} (+3)^{7/3}} \\ &= \int \frac{d}{(-2)^{-1/3} \cdot (-2)^{7/3} \left(\frac{+3}{-2}\right)^{7/3}} \end{aligned}$$

$$\therefore I = \int \frac{d}{(-2)^2 \left(\frac{+3}{-2}\right)^{7/3}}$$

$$\text{Put } \frac{+3}{-2} = t \Rightarrow \frac{-5}{(-2)^2} d = dt$$

$$\Rightarrow \frac{d}{(-2)^2} = \frac{-1}{5} dt$$

$$\begin{aligned} \therefore I &= \frac{-1}{5} \int \frac{dt}{t^{7/3}} = \frac{-1}{5} \cdot \frac{t^{-4/3}}{\left(\frac{-4}{3}\right)} + c \\ &= \frac{3}{20} \left(\frac{-2}{+3}\right)^{4/3} + c \end{aligned}$$

$$92. I = \int \frac{\sin 2}{(3+4\cos)^3} d$$

$$\Rightarrow I = \int \frac{2\sin \cos}{(3+4\cos)^3} d$$

$$\text{Put } 3+4\cos = t \Rightarrow \cos = \frac{t-3}{4}$$

$$\Rightarrow \sin d = \frac{dt}{(-4)}$$

$$\begin{aligned} \therefore I &= \int \frac{2 \left(\frac{dt}{-4}\right) \cdot \left(\frac{t-3}{4}\right)}{(t)^3} \\ &= \frac{-1}{8} \int \frac{t-3}{t^3} dt \\ &= \frac{-1}{8} \left(\int \frac{dt}{t^2} - 3 \int \frac{dt}{t^3} \right) \\ &= \frac{-1}{8} \left(\frac{-1}{t} + \frac{3}{2t^2} \right) + C \\ &= \left(\frac{1}{8t} - \frac{3}{16t^2} \right) + C \\ &= \frac{2t-3}{16t^2} + C = \frac{2(3+4\cos) - 3}{16(3+4\cos)^2} + C \end{aligned}$$

$$\therefore I = \frac{3+8\cos}{16(3+4\cos)^2} + C$$

$$\begin{aligned} 93. \text{ Let } I &= \int \frac{d}{\cos \sqrt{1+\cos 2} + \sin 2} \\ &= \int \frac{d}{\cos \sqrt{2\cos^2 + 2\sin \cos}} \\ &= \int \frac{d}{\cos^2 \sqrt{2+2\tan}} \\ &= \int \frac{\sec^2 d}{\sqrt{2+2\tan}} \end{aligned}$$

$$\text{Put } 2+2\tan = t \Rightarrow 2\sec^2 d = dt$$

$$\begin{aligned} \therefore I &= \frac{1}{2} \int \frac{dt}{\sqrt{t}} \\ &= \frac{1}{2} (2)\sqrt{t} + c \\ &= \sqrt{t} + c = \sqrt{2+2\tan} + c \end{aligned}$$

$$\begin{aligned} 94. \text{ Let } I &= \int \frac{\sin \theta + \cos \theta}{\sqrt{2\sin \theta \cos \theta}} d\theta \\ &= \int \frac{\sin \theta + \cos \theta}{\sqrt{1-(1-2\sin \theta \cos \theta)}} d\theta \\ &= \int \frac{(\sin \theta + \cos \theta) d\theta}{\sqrt{1-(\sin^2 \theta + \cos^2 \theta - 2\sin \theta \cos \theta)}} \\ &= \int \frac{\sin \theta + \cos \theta}{\sqrt{1-(\sin \theta - \cos \theta)^2}} d\theta \end{aligned}$$

$$\begin{aligned} \text{Put } (\sin \theta - \cos \theta) &= t \\ \Rightarrow (\cos \theta + \sin \theta) d\theta &= dt \end{aligned}$$

$$\therefore I = \int \frac{dt}{\sqrt{1-t^2}} = \sin^{-1}(t) + c = \sin^{-1}(\sin \theta - \cos \theta) + c$$



$$\begin{aligned}
 95. \quad \text{Let } I &= \int \frac{d}{(a^2 + x^2)^{\frac{3}{2}}} \\
 \text{Put } x &= a \tan \theta \Rightarrow dx = a \sec^2 \theta d\theta \\
 \therefore I &= \int \frac{a \sec^2 \theta}{(a^2 + a^2 \tan^2 \theta)^{\frac{3}{2}}} d\theta = \int \frac{a \sec^2 \theta}{a^3 (\sec^2 \theta)^{\frac{3}{2}}} d\theta \\
 &= \frac{1}{a^2} \int \frac{d\theta}{\sec \theta} = \frac{1}{a^2} \int \cos \theta d\theta = \frac{1}{a^2} \sin \theta + c \\
 &= \frac{1}{a^2} \cdot \frac{\tan \theta}{\sqrt{1 + \tan^2 \theta}} + c \\
 &= \frac{x}{a^2 (x^2 + a^2)^{\frac{1}{2}}} + c
 \end{aligned}$$

$$\begin{aligned}
 96. \quad \text{Let } I &= \int \frac{1}{a + be^{-x}} dx = \int \frac{e^{-x}}{ae^{-x} + b} dx \\
 \text{Put } ae^{-x} + b &= t \Rightarrow -ae^{-x} dx = dt \\
 \therefore I &= -\frac{1}{a} \int \frac{dt}{t} = -\frac{1}{a} \log |t| + c \\
 &= -\frac{1}{a} \log |ae^{-x} + b| + c \\
 &= -\frac{1}{a} \log \left| \frac{a + be^x}{e^x} \right| + c \\
 &= \frac{1}{a} \log \left| \frac{e^x}{a + be^x} \right| + c
 \end{aligned}$$

$$\begin{aligned}
 97. \quad \text{Put } e^x + e^{-x} &= t \\
 \Rightarrow (e^{-x} + e^x) dx &= dt \\
 \Rightarrow e^x (e^{-x} + e^{-x}) dx &= dt \\
 \Rightarrow (e^{-x} + e^{-x}) dx &= \frac{dt}{e^x} \\
 \therefore \int \frac{e^{-x} + e^{-x}}{e^x + e^{-x}} dx &= \frac{1}{e^x} \int \frac{dt}{t} = \frac{1}{e^x} \log |t| + c \\
 &= \frac{1}{e^x} \log |e^x + e^{-x}| + c
 \end{aligned}$$

$$\begin{aligned}
 98. \quad \text{Let } I &= \int (x^2 + 1) \sqrt{x^2 + 1} dx \\
 \text{Put } \sqrt{x^2 + 1} &= t \Rightarrow x^2 + 1 = t^2 \\
 \Rightarrow x^2 &= t^2 - 1 \Rightarrow dx = 2t dt \\
 \therefore I &= \int \left((t^2 - 1)^2 + 1 \right) (t) (2t dt) \\
 &= \int (t^4 - 2t^2 + 2) (2t^2) dt \\
 &= 2 \left[\int t^6 dt - 2 \int t^4 dt + 2 \int t^2 dt \right]
 \end{aligned}$$

$$\begin{aligned}
 &= 2 \left[\frac{t^7}{7} - \frac{2t^5}{5} + \frac{2t^3}{3} \right] + c \\
 &= 2 \left[\frac{(x^2 + 1)^{7/2}}{7} - \frac{2(x^2 + 1)^{5/2}}{5} + \frac{2(x^2 + 1)^{3/2}}{3} \right] + c
 \end{aligned}$$

$$\begin{aligned}
 99. \quad \text{Let } I &= \int \frac{x^3 dx}{(x^2 + 1)^3} \\
 \text{Put } x^2 + 1 &= t \Rightarrow 2x dx = dt \\
 \therefore I &= \frac{1}{2} \int \frac{(t-1) dt}{t^3} = \frac{1}{2} \int (t^{-2} - t^{-3}) dt \\
 &= \frac{1}{2} \left[\frac{t^{-1}}{-1} - \frac{t^{-2}}{-2} \right] + K \\
 &= \frac{1}{2} \left[\frac{1}{2(x^2 + 1)^2} - \frac{1}{x^2 + 1} \right] + K \\
 &= \frac{1}{2} \left[\frac{1 - 2(x^2 + 1)}{2(x^2 + 1)^2} \right] + K \\
 &= \frac{1}{2} \left[\frac{-(1 + 2x^2)}{2(x^2 + 1)^2} \right] + K \\
 &= \frac{-1}{4} \left[\frac{2x^2 + 1}{(x^2 + 1)^2} \right] + K
 \end{aligned}$$

$$\begin{aligned}
 100. \quad \text{Let } I &= \int \frac{x^2 - 1}{4 + 3x^2 + 1} dx \\
 &= \int \frac{1 - \frac{1}{x^2}}{x^2 + \frac{1}{x^2} + 3} dx = \int \frac{1 - \frac{1}{x^2}}{\left(x + \frac{1}{x}\right)^2 + 1} dx
 \end{aligned}$$

$$\text{Put } x + \frac{1}{x} = t \Rightarrow \left(1 - \frac{1}{x^2}\right) dx = dt$$

$$\begin{aligned}
 \therefore I &= \int \frac{dt}{1 + t^2} \\
 &= \tan^{-1} t + c \\
 &= \tan^{-1} \left(x + \frac{1}{x} \right) + c
 \end{aligned}$$

$$\begin{aligned}
 101. \quad \text{Let } I &= \int \frac{2 - \sin x}{2 + \cos x} dx \\
 &= \int \frac{2}{2 + \cos x} dx - \int \frac{\sin x}{2 + \cos x} dx \\
 &= I_1 - I_2 \\
 I_1 &= \int \frac{2}{2 + \cos x} dx
 \end{aligned}$$



$$\text{Put } \tan\left(\frac{\theta}{2}\right) = t$$

$$\therefore d = \frac{2dt}{1+t^2} \text{ and } \cos \theta = \frac{1-t^2}{1+t^2}$$

$$\begin{aligned} \therefore I_1 &= 2 \int \frac{1}{2 + \frac{1-t^2}{1+t^2}} \cdot \frac{2dt}{1+t^2} \\ &= \int \frac{2 \cdot 2dt}{t^2+3} = 4 \int \frac{dt}{t^2+(\sqrt{3})^2} \end{aligned}$$

$$\begin{aligned} &= \frac{4}{\sqrt{3}} \tan^{-1}\left(\frac{t}{\sqrt{3}}\right) + c_1 \\ &= \frac{4}{\sqrt{3}} \tan^{-1}\left(\frac{\tan(\theta/2)}{\sqrt{3}}\right) + c_1 \end{aligned}$$

$$\text{and } I_2 = \int \frac{\sin \theta}{2 + \cos \theta} d\theta = -\log(2 + \cos \theta) + c_2$$

$$\begin{aligned} \therefore I &= I_1 - I_2 \\ &= \frac{4}{\sqrt{3}} \tan^{-1}\left(\frac{\tan(\theta/2)}{\sqrt{3}}\right) + \log(2 + \cos \theta) + c \end{aligned}$$

$$102. \text{ Let } I = \int \frac{2d}{\sqrt{1-d^2}}$$

$$\text{Put } 1-d^2 = t^2 \Rightarrow d = -2tdt$$

$$\begin{aligned} \therefore I &= -2 \int \frac{(1-t^2)^2 t dt}{t} = -2 \int (1-t^2)^2 dt \\ &= -2 \int (1+t^4-2t^2) dt \\ &= -2 \left[t + \frac{t^5}{5} - \frac{2t^3}{3} \right] \\ &= -2t \left[\frac{15+3t^4-10t^2}{15} \right] \\ &= \frac{-2}{15} \sqrt{1-d^2} [15+3(1-d^2)^2-10(1-d^2)] \\ &= \frac{-2}{15} \sqrt{1-d^2} (3-d^2+4-d^2+8) \\ \therefore P &= \frac{-2}{15} \end{aligned}$$

$$103. \text{ Put } \theta = \tan^{-1} t \Rightarrow d\theta = \frac{1}{1+t^2} dt$$

$$\begin{aligned} \therefore f(\theta) &= \int \frac{\tan^2 \theta \sec^2 \theta d\theta}{\sec^2 \theta (1 + \sec \theta)} \\ &= \int \frac{\tan^2 \theta d\theta}{1 + \sec \theta} = \int \frac{\sin^2 \theta d\theta}{\cos \theta (1 + \cos \theta)} \\ &= \int \frac{1 - \cos^2 \theta d\theta}{\cos \theta (1 + \cos \theta)} \end{aligned}$$

$$= \int \frac{(1 - \cos \theta) d\theta}{\cos \theta} = \int \sec \theta d\theta - \int d\theta$$

$$= \log(\sec \theta + \tan \theta) - \theta + c$$

$$\therefore f(\theta) = \log\left(\frac{1+\sqrt{1+t^2}}{t}\right) - \tan^{-1} t + c$$

$$\begin{aligned} \therefore f(0) &= \log(0 + \sqrt{1+0}) - \tan^{-1}(0) + c \\ \Rightarrow 0 &= \log 1 - 0 + c \Rightarrow c = 0 \end{aligned}$$

$$\therefore f(\theta) = \log\left(\frac{1+\sqrt{1+t^2}}{t}\right) - \tan^{-1} t$$

$$\begin{aligned} \therefore f(1) &= \log\left(\frac{1+\sqrt{1+1^2}}{1}\right) - \tan^{-1}(1) \\ &= \log(1+\sqrt{2}) - \frac{\pi}{4} \end{aligned}$$

$$\begin{aligned} 104. \text{ } J - I &= \int \left(\frac{e^3}{e^4 + e^2 + 1} - \frac{e}{e^4 + e^2 + 1} \right) d \\ &= \int \frac{(e^2 - 1)e}{e^4 + e^2 + 1} d \end{aligned}$$

$$\text{Put } e = t \Rightarrow e d = dt$$

$$\therefore J - I = \int \frac{t^2 - 1}{t^4 + t^2 + 1} dt = \int \frac{1 - \frac{1}{t^2}}{\left(t + \frac{1}{t}\right)^2 - 1} dt$$

$$\text{Put } t + \frac{1}{t} = u$$

$$\Rightarrow \left(1 - \frac{1}{t^2}\right) dt = du$$

$$\therefore J - I = \int \frac{du}{u^2 - 1} = \frac{1}{2} \log \left| \frac{u-1}{u+1} \right| + C$$

$$= \frac{1}{2} \log \left| \frac{t + \frac{1}{t} - 1}{t + \frac{1}{t} + 1} \right| + C$$

$$= \frac{1}{2} \log \left| \frac{t^2 - t + 1}{t^2 + t + 1} \right| + C$$

$$= \frac{1}{2} \log \left| \frac{e^2 - e + 1}{e^2 + e + 1} \right| + C$$

$$105. \text{ Let } I = \int \frac{\sec^2 \theta}{(\sec \theta + \tan \theta)^2} d\theta$$

$$\text{Put } \sec \theta + \tan \theta = t \quad \dots(i)$$

$$\Rightarrow \sec \theta (\sec \theta + \tan \theta) d\theta = dt$$

$$\Rightarrow \sec \theta d\theta = \frac{1}{t} dt$$



$$\text{Also, } \sec \theta - \tan \theta = \frac{1}{t} \quad \dots \text{(ii)}$$

Adding (i) and (ii), we get

$$\sec \theta = \frac{1}{2} \left(t + \frac{1}{t} \right)$$

$$\begin{aligned} \therefore I &= \frac{1}{2} \int \frac{\left(t + \frac{1}{t} \right)}{t^2} \cdot \frac{1}{t} dt = \frac{1}{2} \int \left[\frac{1}{t^2} + \frac{1}{t^3} \right] dt \\ &= -\frac{1}{7t^{\frac{7}{2}}} - \frac{1}{11t^{\frac{11}{2}}} + K = -\frac{1}{t^{\frac{11}{2}}} \left(\frac{t^2}{7} + \frac{1}{11} \right) + K \\ &= \frac{-1}{(\sec \theta + \tan \theta)^{\frac{11}{2}}} \left[\frac{1}{11} + \frac{1}{7} (\sec \theta + \tan \theta)^2 \right] + K \end{aligned}$$

$$\begin{aligned} 106. \int f(\theta) \cdot g(\theta) d\theta &= \int \sin \theta d\theta \\ &= -\cos \theta + \int \cos \theta d\theta \\ &= -\cos \theta + \sin \theta + c \end{aligned}$$

$$\begin{aligned} 107. \int \sin \theta d\theta &= -\cos \theta + \int \cos \theta d\theta \\ &= -\cos \theta + \sin \theta + \text{constant} \end{aligned}$$

$$\therefore A = \sin \theta + \text{constant}$$

$$\begin{aligned} 108. \int \log_{10} \theta d\theta &= \int \frac{\log \theta}{\log 10} d\theta \\ &= \frac{1}{\log 10} (\log \theta - \theta) + c \\ &= (\log_{10} \theta - \log_{10} e) + c \end{aligned}$$

$$\begin{aligned} 109. \int \sin^2 \theta d\theta &= \int \frac{1 - \cos 2\theta}{2} d\theta \\ &= \frac{\theta}{2} - \frac{\sin 2\theta}{4} + c \\ &= \frac{\theta^2}{2} - \frac{\sin 2\theta}{4} + \frac{\cos 2\theta}{4} + c \end{aligned}$$

$$\begin{aligned} 110. \int \sin^2 \theta d\theta &= \int \left(\frac{1 - \cos 2\theta}{2} \right) d\theta \\ &= \frac{1}{2} \left[\int d\theta - \int \cos 2\theta d\theta \right] \\ &= \frac{1}{2} \left[\frac{\theta^2}{2} - \frac{\sin 2\theta}{2} + \int \frac{\sin 2\theta}{2} d\theta \right] \\ &= \frac{\theta^2}{4} - \frac{\sin 2\theta}{4} - \frac{\cos 2\theta}{8} + c \end{aligned}$$

$$\begin{aligned} 111. \int \cos^2 \theta d\theta &= \int \left(\frac{1 + \cos 2\theta}{2} \right) d\theta \\ &= \frac{1}{2} \left[\int d\theta + \int \cos 2\theta d\theta \right] \\ &= \frac{1}{2} \left[\frac{\theta^2}{2} + \frac{\sin 2\theta}{2} - \int \frac{\sin 2\theta}{2} d\theta \right] \\ &= \frac{\theta^2}{4} + \frac{\sin 2\theta}{4} + \frac{\cos 2\theta}{8} + c \end{aligned}$$

$$\begin{aligned} 112. \int \cos^{-1} \theta d\theta &= \cos^{-1} \theta \cdot \theta + \int \frac{1}{\sqrt{1-\theta^2}} \cdot \theta d\theta \\ &= \cos^{-1} \theta - \frac{1}{2} \int \frac{-2\theta}{\sqrt{1-\theta^2}} d\theta \\ &= \cos^{-1} \theta - \sqrt{1-\theta^2} + c \end{aligned}$$

$$\begin{aligned} 113. \int \sin(\log \theta) d\theta + \int \cos(\log \theta) d\theta \\ &= \sin(\log \theta) - \int \frac{\cos(\log \theta)}{\theta} d\theta + \int \cos(\log \theta) d\theta + c \\ &= \sin(\log \theta) + c \end{aligned}$$

$$\begin{aligned} 114. \int \frac{(\sin \theta + \cos \theta)(2 - \sin 2\theta)}{\sin^2 2\theta} d\theta \\ &= -\int \frac{(\sin \theta + \cos \theta)(\sin 2\theta - 2)}{\sin^2 2\theta} d\theta \\ &= -\int \frac{(\sin \theta + \cos \theta)(2\sin \theta \cos \theta - 2)}{4\sin^2 \theta \cos^2 \theta} d\theta \\ &= -\int \frac{2\sin^2 \theta \cos \theta - 2\sin \theta + 2\sin \theta \cos^2 \theta - 2\cos \theta}{4\sin^2 \theta \cos^2 \theta} d\theta \\ &= -\int \left[\frac{1}{2\cos \theta} + \frac{1}{2\sin \theta} - \frac{1}{2\sin^2 \theta \cos \theta} - \frac{1}{2\cos^2 \theta \sin \theta} \right] d\theta \\ &= \frac{-1}{2} \left[\int \sec \theta d\theta + \int \operatorname{cosec} \theta d\theta - \int \frac{\operatorname{cosec}^2 \theta}{\cos \theta} d\theta - \int \frac{\sec^2 \theta}{\sin \theta} d\theta \right] \\ &= \frac{-1}{2} \left[\int \sec \theta d\theta + \int \operatorname{cosec} \theta d\theta - \int \frac{1 + \cot^2 \theta}{\cos \theta} d\theta - \int \frac{1 + \tan^2 \theta}{\sin \theta} d\theta \right] \end{aligned}$$



$$\begin{aligned}
 &= \frac{-1}{2} \left[\int \sec d + \int \operatorname{cosec} d - \int \sec d \right. \\
 &\quad \left. - \int \operatorname{cosec} \cot d - \int \operatorname{cosec} d - \int \sec \tan d \right] \\
 &= \frac{-1}{2} \left[-\int \operatorname{cosec} \cot d - \int \sec \tan d \right] \\
 &= \frac{\sec - \operatorname{cosec}}{2} + c \\
 &= \frac{\sin - \cos}{2 \sin \cos} + c = \frac{\sin - \cos}{\sin 2} + c
 \end{aligned}$$

115. Let $I = \int e^{\sin} (\cos - \sec \tan) d$

$$\begin{aligned}
 &= \int e^{\sin} \cos d - \int e^{\sin} \sec \tan d \\
 I_1 &= \int e^{\sin} \cos d \\
 &= \int e^{\sin} \cos d - \int e^{\sin} d + c_1 \\
 &= e^{\sin} - \int e^{\sin} d + c_1 \\
 I_2 &= \int e^{\sin} \sec \tan d \\
 &= e^{\sin} \int \sec \tan d - \int \sec \cos e^{\sin} d + c_2 \\
 &= \sec e^{\sin} - \int e^{\sin} d + c_2 \\
 I &= I_1 - I_2 \\
 &= e^{\sin} - \int e^{\sin} d - \sec e^{\sin} + \int e^{\sin} d + c \\
 &= e^{\sin} (-\sec) + c
 \end{aligned}$$

116. Let $I = \int e^{\sin} \left(\frac{\cos^3 - \sin}{\cos^2} \right) d$

$$\begin{aligned}
 &= \int e^{\sin} (\cos - \sec \tan) d \\
 &= e^{\sin} (-\sec) + c
 \end{aligned}$$

117. Let $I = \int \sin(10 +) \cdot \sin^9 d$

$$\begin{aligned}
 &= \int \sin(10 +) \cdot \sin^9 d \\
 &= \int \sin 10 \cos \sin^9 d \\
 &\quad + \int \cos 10 \sin \sin^9 d \\
 &= \sin 10 \int \cos \sin^9 d \\
 &\quad - \int 10 \cos 10 \int \cos \sin^9 d \\
 &\quad + \int \cos 10 \sin^{10} d \\
 &= \frac{\sin 10 \cdot \sin^{10}}{10} - \frac{10}{10} \int \cos 10 \sin^{10} d \\
 &\quad + \int \cos 10 \sin^{10} d \\
 &= \frac{\sin 10 \cdot \sin^{10}}{10} + c
 \end{aligned}$$

118. Let $I = \int [e^2 f() + e^2 f'()] d$

$$\begin{aligned}
 &= \int e^2 f() d + \int e^2 f'() d \\
 &= f() \int e^2 d - \int \left[\int e^2 d \right] f'() d \\
 &\quad + g() + c \\
 &= \frac{e^2 f()}{2} - \int \frac{e^2}{2} f'() d + g() + c \\
 &= \frac{e^2 f()}{2} - \frac{g()}{2} + g() + c \\
 &= \frac{e^2 f()}{2} + \frac{g()}{2} + c \\
 &= \frac{1}{2} [e^2 f() + g()] + c
 \end{aligned}$$

119. Let $I = \int e^{\sqrt{}} d$

Put $\sqrt{} = t$

$$\begin{aligned}
 &\Rightarrow \frac{1}{2\sqrt{}} d = dt \\
 &\Rightarrow d = 2t dt \\
 \therefore I &= \int e^t \cdot 2t dt = 2(t \cdot e^t - e^t) + A \\
 &= 2(\sqrt{} \cdot e^{\sqrt{}} - e^{\sqrt{}}) + A \\
 &= 2(\sqrt{} - 1) e^{\sqrt{}} + A
 \end{aligned}$$

120. Let $I = \int \cos(\log) d$

Put $\log = t \Rightarrow = e^t \Rightarrow d = e^t dt$

$$\begin{aligned}
 \therefore I &= \int e^t \cos t dt \\
 &= e^t \cos t - \int e^t (-\sin t) dt + c_1 \\
 &= e^t \cos t + \int e^t \sin t dt + c_1 \\
 &= e^t \cos t + e^t \sin t - \int e^t \cos t dt + c_2 \\
 &= e^t \cos t + e^t \sin t - I + c_2 \\
 \Rightarrow 2I &= e^t (\cos t + \sin t) + c_2 \\
 \Rightarrow I &= \frac{e^t}{2} (\cos t + \sin t) + c \\
 &= \frac{1}{2} [\cos(\log) + \sin(\log)] + c
 \end{aligned}$$

121. Put $\sin^{-1} = t$

$$\begin{aligned}
 &\Rightarrow \frac{1}{\sqrt{1-t^2}} d = dt \\
 \therefore \int \frac{\sin^{-1}}{\sqrt{1-t^2}} d &= \int t \sin t dt = -t \cos t + \sin t + c \\
 &= -\sin^{-1} \cos(\sin^{-1}) + \sin(\sin^{-1}) + c \\
 &= -\sin^{-1} \cdot \sqrt{1-t^2} + c
 \end{aligned}$$



122. Let $I = \int \cos(\log_e x) dx$
 Put $\log_e x = t \Rightarrow x = e^t \Rightarrow dx = e^t dt$

$$\begin{aligned} \therefore I &= \int \cos t \cdot e^t dt \\ &= \cos t \cdot e^t - \int (-\sin t) \cdot e^t dt \\ &= \cos t \cdot e^t + \left[\sin t \cdot e^t - \int \cos t \cdot e^t dt \right] \end{aligned}$$

$$\begin{aligned} \therefore I &= \cos t \cdot e^t + \sin t \cdot e^t - I + c_1 \\ \Rightarrow 2I &= \cos t \cdot e^t + \sin t \cdot e^t + c_1 \\ \Rightarrow I &= \frac{1}{2} [\cos(\log_e x) + \sin(\log_e x)] + c, \end{aligned}$$

where $c = \frac{c_1}{2}$

123. $\int 32 x^3 (\log x)^2 dx$

$$\begin{aligned} &= 32 \int x^3 (\log x)^2 dx \\ &= 32 \left[(\log x)^2 \cdot \frac{x^4}{4} - \int 2 \log x \cdot \frac{1}{4} \cdot x^4 dx \right] \\ &= 32 \left[(\log x)^2 \cdot \frac{x^4}{4} - \frac{1}{2} \int x^3 \log x dx \right] \\ &= 32 \left[\frac{(\log x)^2 \cdot x^4}{4} - \frac{1}{2} \left(\frac{\log x \cdot x^4}{4} - \int \frac{1}{4} \cdot x^4 dx \right) \right] \\ &= 32 \left[\frac{(\log x)^2 \cdot x^4}{4} - \frac{1}{2} \left(\frac{x^4 \log x}{4} - \frac{1}{4} \cdot \frac{x^4}{4} \right) \right] + c \\ &= 8 \left[(\log x)^2 \cdot x^4 - \frac{1}{2} \left(x^4 \log x - \frac{x^4}{4} \right) \right] + c \\ &= 8 x^4 \left[(\log x)^2 - \frac{\log x}{2} + \frac{1}{8} \right] + c \\ &= x^4 [8(\log x)^2 - 4 \log x + 1] + c \end{aligned}$$

124. Let $I = \int x^4 e^x dx$

$$\begin{aligned} &= \frac{x^4 e^x}{2} - \int 4 x^3 \cdot \frac{e^x}{2} dx \\ &= \frac{x^4 e^x}{2} - 2 \int x^3 e^x dx \\ &= \frac{x^4 e^x}{2} - 2 \left[\int \frac{x^3 e^x}{2} dx - \int 3 x^2 \cdot \frac{e^x}{2} dx \right] \\ &= \frac{x^4 e^x}{2} - x^3 e^x + 3 \int x^2 e^x dx \\ &= \frac{x^4 e^x}{2} - x^3 e^x + 3 \left[\frac{x^2 e^x}{2} - \int 2 x \cdot \frac{e^x}{2} dx \right] \end{aligned}$$

$$\begin{aligned} &= \frac{x^4 e^x}{2} - x^3 e^x + \frac{3 x^2 e^x}{2} - \frac{3 x e^x}{2} + 3 \int \frac{e^x}{2} dx \\ &= \frac{x^4 e^x}{2} - x^3 e^x + \frac{3 x^2 e^x}{2} - \frac{3 x e^x}{2} + \frac{3 e^x}{4} + c \\ &= \frac{e^x}{4} [2 x^4 - 4 x^3 + 6 x^2 - 6 x + 3] + c \end{aligned}$$

125. $\int x^3 e^x dx = x^3 \cdot \frac{e^x}{5} - \int 3 x^2 \cdot \frac{e^x}{5} dx$

$$\begin{aligned} &= \frac{x^3 e^x}{5} - \frac{3}{5} \int x^2 e^x dx + \frac{3}{5} \int 2 x \cdot \frac{e^x}{5} dx \\ &= \frac{x^3 e^x}{5} - \frac{3}{25} \int x^2 e^x dx + \frac{6}{25} \int x e^x dx - \frac{6}{25} \int \frac{e^x}{25} dx + c \end{aligned}$$

$$\therefore \int x^3 e^x dx = \frac{e^x}{5^4} (5^3 x^3 - 75 x^2 + 30 x - 6) + c$$

$$\therefore f(x) = 5^3 x^3 - 75 x^2 + 30 x - 6$$

126. $\int \log(a^2 + x^2) dx$

$$\begin{aligned} &= \log(a^2 + x^2) \cdot x - \int \frac{1}{a^2 + x^2} \cdot 2x \cdot dx \\ &= x \log(a^2 + x^2) - 2 \int \frac{x^2}{a^2 + x^2} dx \\ &= x \log(a^2 + x^2) - 2 \int \left(1 - \frac{a^2}{a^2 + x^2} \right) dx \\ &= x \log(a^2 + x^2) - 2x + 2a^2 \cdot \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + c \\ &= x \log(a^2 + x^2) - 2x + 2a \tan^{-1} \left(\frac{x}{a} \right) + c \end{aligned}$$

127. Put $\log x = t$
 $\Rightarrow x = e^t$
 $\Rightarrow dx = e^t dt$

$$\begin{aligned} \therefore \int (\log x)^5 dx &= \int t^5 e^t dt \\ &= e^t (t^5 - 5t^4 + 20t^3 - 60t^2 + 120t - 120) \\ &\quad + \text{constant} \\ &= [(\log x)^5 - 5(\log x)^4 + 20(\log x)^3 - 60(\log x)^2 \\ &\quad + 120 \log x - 120] + \text{constant} \end{aligned}$$

$$\therefore A = 1, B = -5, C = 20, D = -60, E = 120 \text{ and } F = -120$$

$$\therefore A + B + C + D + E + F = -44$$

128. Put $x = \sec \theta \Rightarrow dx = \sec \theta \tan \theta d\theta$

$$\begin{aligned} \therefore \int \cos^{-1} \left(\frac{1}{x} \right) dx &= \int \cos^{-1} \left(\frac{1}{\sec \theta} \right) \cdot \sec \theta \tan \theta d\theta \\ &= \int \cos^{-1} (\cos \theta) \cdot \sec \theta \tan \theta d\theta \end{aligned}$$



$$\begin{aligned}
 &= \int \theta \cdot (\sec\theta \tan\theta) d\theta \\
 &= \theta \sec\theta - \int 1 \cdot \sec\theta d\theta \\
 &= \theta \sec\theta - \log |\tan\theta + \sec\theta| + c \\
 &= \theta \sec\theta - \log |\sqrt{(\sec^2\theta - 1)} + \sec\theta| + c \\
 &= \sec^{-1} - \log |\sqrt{(x^2 - 1)} + x| + c
 \end{aligned}$$

$$\begin{aligned}
 129. \int e^{\left[\frac{1+\log x}{x}\right]} dx &= \int e^{\left(\log x + \frac{1}{x}\right)} dx \\
 &= e^{\log x + c} \\
 \dots \left[\because \int e^{[f(x)+f'(x)]} dx &= e^{f(x)+c} \right]
 \end{aligned}$$

$$\begin{aligned}
 130. \int e^{\sin x (\sin x + 2\cos x)} dx \\
 &= \int e^{(\sin^2 x + 2\sin x \cos x)} dx \\
 &= e^{\sin^2 x} + c
 \end{aligned}$$

$$\begin{aligned}
 131. \int e^{\left[\frac{2+\sin 2x}{1+\cos 2x}\right]} dx &= \int e^{\left[\frac{2(1+\sin x \cos x)}{2\cos^2 x}\right]} dx \\
 &= \int e^{(\sec^2 x + \tan x)} dx \\
 &= e^{\tan x} + c \\
 \dots \left[\because \int e^{[f(x)+f'(x)]} dx &= e^{f(x)+c} \right]
 \end{aligned}$$

$$\begin{aligned}
 132. \int \frac{(x+3)e^x}{(x+4)^2} dx &= \int \frac{(x+4-1)e^x}{(x+4)^2} dx \\
 &= \int e^x \left(\frac{1}{x+4} - \frac{1}{(x+4)^2} \right) dx \\
 &= \frac{e^x}{x+4} + c
 \end{aligned}$$

$$\begin{aligned}
 133. \int \left(\frac{x+2}{x+4} \right)^2 e^x dx \\
 &= \int e^x \left[\frac{x^2+4x+4}{(x+4)^2} \right] dx \\
 &= \int e^x \left[\frac{(x+4)}{(x+4)^2} + \frac{4}{(x+4)^2} \right] dx \\
 &= \int e^x \left[\frac{1}{x+4} + \frac{4}{(x+4)^2} \right] dx = e^x \left(\frac{1}{x+4} \right) + c
 \end{aligned}$$

$$\begin{aligned}
 134. \int \frac{e^x}{(x+1)^2} dx &= \int e^x \left[\frac{x+1-1}{(x+1)^2} \right] dx \\
 &= \int e^x \left[\frac{1}{x+1} - \frac{1}{(x+1)^2} \right] dx
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{e^x}{x+1} + c \\
 \dots \left[\because \int e^{[f(x)+f'(x)]} dx &= e^{f(x)+c} \right]
 \end{aligned}$$

$$\begin{aligned}
 135. \int \frac{e^{(1+\sin x)}}{1+\cos x} dx &= \int \frac{e^{\left[1+2\sin\left(\frac{x}{2}\right)\cos\left(\frac{x}{2}\right)\right]}}{2\cos^2\left(\frac{x}{2}\right)} dx \\
 &= \int e^{\left(\frac{1}{2}\sec^2\frac{x}{2} + \tan\frac{x}{2}\right)} dx \\
 &= e^{\tan\frac{x}{2}} + c \\
 \dots \left[\because \int e^{[f(x)+f'(x)]} dx &= e^{f(x)+c} \right]
 \end{aligned}$$

$$\begin{aligned}
 136. \int \frac{e^{(x^2 \tan^{-1} x + \tan^{-1} x + 1)}}{x^2 + 1} dx \\
 &= \int e^{[(x^2+1)\tan^{-1} x + 1]} dx \\
 &= \int e^{\left(\tan^{-1} x + \frac{1}{1+x^2}\right)} dx \\
 &= e^{\tan^{-1} x} + c
 \end{aligned}$$

$$\begin{aligned}
 137. \text{ Let } I &= \int e^{\tan x} (\sec^2 x + \sec^3 x \cdot \sin x) dx \\
 &= \int e^{\tan x} (1 + \tan x) \sec^2 x dx \\
 \text{Put } \tan x &= t \Rightarrow \sec^2 x dx = dt \\
 \therefore I &= \int e^t (1+t) dt = e^t \cdot t + c = \tan x \cdot e^{\tan x} + c
 \end{aligned}$$

$$\begin{aligned}
 138. \text{ Let } I &= \int e^{\sin x} \left(\frac{\sin x + 1}{\sec x} \right) dx \\
 &= \int e^{\sin x} (\sin x + 1) \cos x dx \\
 \text{Put } \sin x &= t \Rightarrow \cos x dx = dt \\
 \therefore I &= \int e^t (1+t) dt \\
 &= te^t + c \dots \left[\because \int e^{[f(x)+f'(x)]} dx = e^{f(x)} \right] \\
 &= \sin x \cdot e^{\sin x} + c
 \end{aligned}$$

$$\begin{aligned}
 139. \text{ Put } \tan^{-1} x &= t \Rightarrow \frac{dx}{1+x^2} = dt \\
 \therefore \int e^{\tan^{-1} x} \left(\frac{1+x^2+x^2}{1+x^2} \right) dx &= \int e^t (\tan t + \sec^2 t) dt \\
 &= e^t \tan t + c \\
 &= e^{\tan^{-1} x} + c
 \end{aligned}$$



$$140. \text{ Put } \cot^{-1} t = t \Rightarrow \frac{-d}{1+t^2} = dt$$

$$\begin{aligned} \therefore \int e^{\cot^{-1} t} \left(\frac{2-t}{1+t^2} + 1 \right) dt \\ &= -\int e^t (\cot^2 t - \cot t + 1) dt \\ &= -\int e^t (\operatorname{cosec}^2 t - \cot t) dt \\ &= \int e^t (\cot t - \operatorname{cosec}^2 t) dt \\ &= e^t \cot t + c \\ &= e^{\cot^{-1} t} + c \end{aligned}$$

$$141. \int e^2 \left(\frac{1}{2} - \frac{1}{2^2} \right) d = \frac{e^2}{2} + c$$

$$\dots \left[\because e^m \left[f(x) + \frac{f'(x)}{m} \right] dx = \frac{e^m f(x)}{m} + c \right]$$

$$142. \int (1 + e^{-x}) e^{+x} dx$$

$$= \int \left[e^{+x} \left(1 - \frac{1}{2} \right) + e^{+x} \right] dx$$

$$= e^{+x} + c$$

$$\dots \left[\because \int [f'(x) + f(x)] dx = f(x) + c \right]$$

$$143. \int \frac{2x+3}{x^2-5x+6} dx = \int \frac{2x+3}{(x-3)(x-2)} dx$$

$$= \int \left(\frac{9}{x-3} - \frac{7}{x-2} \right) dx$$

$$= 9 \log(x-3) - 7 \log(x-2) + c$$

$$\therefore A = \text{constant}$$

$$144. \int \left(\frac{1}{x-3} - \frac{1}{x^2-3} \right) dx$$

$$= \int \left[\frac{1}{x-3} - \frac{1}{x(x-3)} \right] dx$$

$$= \int \left[\frac{1}{x-3} + \frac{1}{3} - \frac{1}{3(x-3)} \right] dx$$

$$= \int \left[\frac{2}{3(x-3)} + \frac{1}{3} \right] dx$$

$$= \frac{2}{3} \log(x-3) + \frac{1}{3} \log x + c$$

$$= \frac{2}{3} \log(x-3) + \frac{2}{3} \log \sqrt{x} + c$$

$$= \frac{2}{3} \log \left[\sqrt{x} (x-3) \right] + c$$

$$145. \text{ Let } I = \int \frac{1}{(x^2+4)(x^2+9)} dx$$

$$\therefore I = \int \frac{1}{5} \left[\frac{1}{x^2+4} - \frac{1}{x^2+9} \right] dx$$

$$\dots \left\{ \frac{1}{\alpha\beta} = \frac{1}{\beta-\alpha} \left[\frac{1}{\alpha} - \frac{1}{\beta} \right] \right\}$$

$$= \frac{1}{5} \left[\frac{1}{2} \tan^{-1} \left(\frac{x}{2} \right) - \frac{1}{3} \tan^{-1} \left(\frac{x}{3} \right) \right] + c$$

$$= \frac{1}{10} \tan^{-1} \left(\frac{x}{2} \right) - \frac{1}{15} \tan^{-1} \left(\frac{x}{3} \right) + c$$

$$\text{Comparing with } A \tan^{-1} \left(\frac{x}{2} \right) + B \tan^{-1} \left(\frac{x}{3} \right) + c,$$

we get

$$A = \frac{1}{10}, B = \frac{-1}{15}$$

$$A - B = \frac{1}{10} + \frac{1}{15} = \frac{1}{6}$$

$$146. \int \frac{1}{(x^2-a^2)(x^2-b^2)} dx$$

$$= \frac{1}{a^2-b^2} \left[\int \frac{1}{x^2-a^2} dx - \int \frac{1}{x^2-b^2} dx \right]$$

$$= \frac{1}{2(a^2-b^2)} \left[\log(x^2-a^2) - \log(x^2-b^2) \right] + c$$

$$= \frac{1}{2(a^2-b^2)} \log \left(\frac{x^2-a^2}{x^2-b^2} \right) + c$$

$$147. \text{ Let } \frac{2x^2+1}{(x^2-4)(x^2-1)} = \frac{A}{x^2-4} + \frac{B}{x^2-1}$$

$$\therefore 2x^2+1 = A(x^2-1) + B(x^2-4)$$

Comparing the coefficient of x^2 and constant term on both sides, we get

$$A + B = 2 \text{ and } -A - 4B = 1$$

Solving these two equations, we get

$$A = 3 \text{ and } B = -1$$

$$\therefore \int \frac{2x^2+1}{(x^2-4)(x^2-1)} dx$$

$$= \int \left[\frac{3}{x^2-4} - \frac{1}{x^2-1} \right] dx$$

$$= \frac{3}{2 \times 2} \log \left| \frac{x-2}{x+2} \right| - \frac{1}{2} \log \left| \frac{x-1}{x+1} \right| + c$$

$$= \log \left| \frac{-2}{x+2} \right|^{\frac{3}{4}} + \log \left| \frac{x+1}{-1} \right|^{\frac{1}{2}} + c$$



$$= \log \left[\left(\frac{-1}{-1} \right)^{\frac{1}{2}} \left(\frac{-2}{+2} \right)^{\frac{3}{4}} \right] + c$$

$$\therefore a = \frac{1}{2} \text{ and } b = \frac{3}{4}$$

$$\begin{aligned} 148. \int \frac{2^2+3}{(x^2-1)(x^2-4)} dx &= \int \frac{-\frac{5}{3}}{x^2-1} dx + \int \frac{\frac{11}{3}}{x^2-4} dx \\ &= \frac{-5}{3} \cdot \frac{1}{2 \times 1} \log \left| \frac{-1}{+1} \right| + \frac{11}{3} \cdot \frac{1}{2 \times 2} \log \left| \frac{-2}{+2} \right| + c \\ &= \log \left| \frac{-1}{-1} \right|^{\frac{5}{6}} + \log \left| \frac{-2}{+2} \right|^{\frac{11}{12}} + c \\ &= \log \left[\left(\frac{-1}{-1} \right)^{\frac{5}{6}} \left(\frac{-2}{+2} \right)^{\frac{11}{12}} \right] + c \end{aligned}$$

$$\therefore a = \frac{11}{12} \text{ and } b = \frac{5}{6}$$

$$\begin{aligned} 149. \text{ Let } I &= \int \frac{5x^2+3}{x^2(x^2-2)} dx \\ &= \int \frac{5}{x^2-2} dx + \int \frac{3}{x^2(x^2-2)} dx \\ &= \frac{5}{2\sqrt{2}} \log \left| \frac{-\sqrt{2}}{+\sqrt{2}} \right| \\ &\quad + \frac{3}{2} \int \left(\frac{1}{x^2-2} - \frac{1}{x^2} \right) dx \\ &= \frac{5}{2\sqrt{2}} \log \left| \frac{-\sqrt{2}}{+\sqrt{2}} \right| \\ &\quad + \frac{3}{2} \left[\frac{1}{2\sqrt{2}} \log \left| \frac{-\sqrt{2}}{+\sqrt{2}} \right| + \frac{1}{x} \right] + c \\ &= \frac{13}{4\sqrt{2}} \log \left| \frac{-\sqrt{2}}{+\sqrt{2}} \right| + \frac{3}{2} + c \end{aligned}$$

$$\begin{aligned} 150. \int \frac{dx}{x^6+x^4} &= \int \frac{(x^2+1)dx}{x^4(x^2+1)} - \int \frac{x^2 dx}{x^4(x^2+1)} \\ &= \int \frac{1}{x^4} dx - \int \frac{dx}{x^2(x^2+1)} \\ &= -\frac{1}{3x^3} - \int \frac{dx}{x^2} + \int \frac{dx}{x^2+1} \\ &= \frac{-1}{3x^3} + \frac{1}{x} + \tan^{-1} x + c \end{aligned}$$

$$\begin{aligned} 151. \int \frac{dx}{e^2-3e} &= \int \frac{dx}{e(e-3)} \\ &= -\frac{1}{3} \left[\int \frac{dx}{e} - \int \frac{dx}{e-3} \right] \\ &= \frac{1}{3} \int \frac{dx}{e-3} - \frac{1}{3} \int \frac{dx}{e} \\ &= \frac{1}{3} \int \frac{e^{-x}}{1-3e^{-x}} dx - \frac{1}{3} \int e^{-x} dx \\ &= \frac{1}{9} \log(1-3e^{-x}) + \frac{1}{3e} + c \\ &= \frac{1}{9} \log \left(\frac{e-3}{e} \right) + \frac{1}{3e} + c \\ &= \frac{1}{3e} + \frac{1}{9} \log(e-3) - \frac{1}{9} \log e + c \\ &= \frac{1}{3e} + \frac{1}{9} \log(e-3) - \frac{1}{9} + c \end{aligned}$$

$$\begin{aligned} 152. \int \frac{dx}{(x^2+1)(x-1)} dx \\ &= \frac{1}{2} \int \frac{1}{x-1} dx - \frac{1}{2} \int \frac{-1}{x^2+1} dx \\ &= \frac{1}{2} \int \frac{1}{x-1} dx - \frac{1}{2} \int \frac{1}{x^2+1} dx + \frac{1}{2} \int \frac{1}{x^2+1} dx \\ &= \frac{1}{2} \log|x-1| - \frac{1}{4} \int \frac{2}{x^2+1} dx + \frac{1}{2} \tan^{-1} x + D_1 \\ &= \frac{1}{2} \log|x-1| - \frac{1}{4} \log|x^2+1| + \frac{1}{2} \tan^{-1} x + D \end{aligned}$$

Comparing with

$A \log|x^2+1| + B \tan^{-1} x + C \log|x-1| + D$, we get

$$A = \frac{-1}{4}, B = \frac{1}{2}, C = \frac{1}{2}$$

$$\therefore A + B + C = \frac{-1}{4} + \frac{1}{2} + \frac{1}{2} = \frac{3}{4}$$

$$\begin{aligned} 153. \int \frac{2x+3}{(x-1)(x^2+1)} dx \\ &= \int \frac{5dx}{2(x-1)} + \int \frac{-\left(\frac{5}{2} + \frac{1}{2}\right)}{x^2+1} dx \\ &= \frac{5}{2} \log(x-1) - \frac{5}{2} \int \frac{dx}{1+x^2} - \frac{1}{2} \int \frac{dx}{1+x^2} \\ &= \frac{5}{2} \log(x-1) - \frac{5}{4} \log(1+x^2) - \frac{1}{2} \tan^{-1} x + A \\ &= \log(x-1)^{\frac{5}{2}} (1+x^2)^{-\frac{5}{4}} - \frac{1}{2} \tan^{-1} x + A \end{aligned}$$

$$\therefore a = -\frac{5}{4}$$



$$\begin{aligned}
 154. \text{ Let } I &= \int \frac{1}{(x^4+1)} dx \\
 &= \int \frac{x^3}{x^4(x^4+1)} dx \\
 \text{Put } x^4 &= t \Rightarrow 4x^3 dx = dt \\
 \therefore I &= \frac{1}{4} \int \frac{dt}{t(1+t)} \\
 &= \frac{1}{4} \int \left(\frac{1}{t} - \frac{1}{1+t} \right) dt \\
 &= \frac{1}{4} [\log |t| - \log |1+t|] + c \\
 &= \frac{1}{4} \log \left| \frac{t}{1+t} \right| + c \\
 &= \frac{1}{4} \log \left| \frac{x^4}{1+x^4} \right| + c
 \end{aligned}$$

$$\begin{aligned}
 155. \int \frac{dx}{x^3+3x^2+2} &= \int \frac{1}{(x^2+3x+2)} dx \\
 &= \int \frac{1}{(x+2)(x+1)} dx \\
 &= \int \frac{1}{2} dx + \int \frac{1}{2(x+2)} dx - \int \frac{1}{x+1} dx \\
 &= \frac{1}{2} \log |x| + \frac{1}{2} \log |x+2| - \log |x+1| + c \\
 &= \frac{1}{2} \log |(x+2)| - \frac{1}{2} \log (x+1)^2 + c \\
 &= \frac{1}{2} \log \left[\frac{|x^2+2x|}{(x+1)^2} \right] + c
 \end{aligned}$$

$$\begin{aligned}
 156. \text{ Let } I &= \int \frac{dx}{\sin x + \sin 2x} \\
 &= \int \frac{dx}{\sin x(1+2\cos x)} \\
 &= \int \frac{\sin x dx}{\sin^2 x(1+2\cos x)} \\
 &= \int \frac{\sin x dx}{(1-\cos x)(1+\cos x)(1+2\cos x)} \\
 \text{Put } \cos x &= t \\
 \Rightarrow -\sin x dx &= dt \\
 \therefore I &= - \int \frac{dt}{(1-t)(1+t)(1+2t)}
 \end{aligned}$$

$$\begin{aligned}
 \therefore I &= - \int \left[\frac{1}{6(1-t)} - \frac{1}{2(1+t)} + \frac{4}{3(1+2t)} \right] dt \\
 &= \frac{1}{6} \log(1-t) + \frac{1}{2} \log(1+t) - \frac{2}{3} \log(1+2t) + c \\
 &= \frac{1}{6} \log(1-\cos x) + \frac{1}{2} \log(1+\cos x) \\
 &\quad - \frac{2}{3} \log(1+2\cos x) + c
 \end{aligned}$$

$$\begin{aligned}
 157. \int \frac{f(x)}{\log \cos x} dx &= -\log(\log \cos x) + c \\
 \text{Differentiating on both sides, we get} \\
 \frac{f(x)}{\log \cos x} &= \frac{-1}{\log \cos x} \times \frac{1}{\cos x} \times (-\sin x) \\
 \Rightarrow \frac{f(x)}{\log \cos x} &= \frac{\tan x}{\log \cos x} \Rightarrow f(x) = \tan x
 \end{aligned}$$

$$\begin{aligned}
 158. \int \frac{f(x)}{\log(\sin x)} dx &= \log(\log \sin x) + c \\
 \text{Differentiating on both sides, we get} \\
 \frac{f(x)}{\log(\sin x)} &= \frac{1}{\log(\sin x)} \times \frac{1}{\sin x} \times \cos x \\
 \Rightarrow \frac{f(x)}{\log(\sin x)} &= \frac{\cot x}{\log(\sin x)} \Rightarrow f(x) = \cot x
 \end{aligned}$$

$$\begin{aligned}
 159. \int f(x) \cos x dx &= \frac{1}{2} [f(x)]^2 + c \\
 \text{Differentiating both sides w.r.t. } x, \text{ we get} \\
 f(x) \cos x &= f(x) \cdot f'(x) \\
 \Rightarrow f'(x) &= \cos x \\
 \Rightarrow \int f'(x) dx &= \int \cos x dx \\
 \Rightarrow f(x) &= \sin x + c \\
 \therefore f\left(\frac{\pi}{2}\right) &= 1 + c
 \end{aligned}$$

$$\begin{aligned}
 160. \int f(x) \cdot \cos x dx &= \frac{1}{2} [f(x)]^2 + c \\
 \text{Differentiating w.r.t. } x, \text{ we get} \\
 f(x) \cdot \cos x &= \frac{1}{2} \times 2 f(x) \cdot f'(x) \\
 \Rightarrow \cos x &= f'(x) \\
 \Rightarrow \cos 0 &= f'(0) \\
 \Rightarrow f'(0) &= 1
 \end{aligned}$$



$$161. f'(x) = 2 - \frac{5}{4}$$

$$\therefore f(x) = \int \left(2 - \frac{5}{4}\right) dx = 2x + \frac{5}{3 \cdot 3} + c$$

$$f(1) = 2(1) + \frac{5}{3(1)^3} + c$$

$$\Rightarrow \frac{14}{3} = 2 + \frac{5}{3} + c \Rightarrow c = 1$$

$$\therefore f(x) = 2x + \frac{5}{3x^3} + 1$$

$$f(-1) = 2(-1) + \frac{5}{3(-1)^3} + 1$$

$$= -2 - \frac{5}{3} + 1 = \frac{-8}{3}$$

$$162. \int f(x) \sin \cos x dx = \frac{1}{2(b^2 - a^2)} \log f(x) + c$$

Differentiating both sides w.r.t. x , we get

$$f(x) \sin \cos x = \frac{1}{2(b^2 - a^2)} \cdot \left[\frac{f'(x)}{f(x)} \right]$$

$$\Rightarrow [f(x)]^2 \sin 2x = \frac{1}{(b^2 - a^2)} f'(x)$$

$$\Rightarrow \sin 2x = \frac{1}{b^2 - a^2} \left(\frac{d}{dx} \right)$$

...[Putting $f(x) = \dots$]

$$\Rightarrow \frac{d}{dx} = (b^2 - a^2) \sin 2x dx$$

$$\Rightarrow \int \frac{d}{dx} = (b^2 - a^2) \int \sin 2x dx = \sin 2x dx$$

$$\Rightarrow \frac{-1}{2} = \frac{-(b^2 - a^2) \cos 2x}{2}$$

$$\Rightarrow \frac{1}{(b^2 - a^2) \cos 2x} = f(x)$$

$$163. \frac{d}{dx}[f(x)] = \cos x + \sin x$$

$$\Rightarrow f(x) = \int (\cos x + \sin x) dx = \sin x + c$$

Since, $f(0) = 2 \Rightarrow c = 2$

$$\therefore f(x) = \sin x + 2$$

$$164. \int \log(x^2 + 1) dx = \int \log[(x + 1)] dx$$

$$= \int \log x dx + \int \log(x + 1) dx$$

$$= \log x - \int \frac{1}{x} dx + \log(x + 1) - \int \frac{1}{x + 1} dx$$

$$= \log x - \int \frac{1}{x} dx + \log(x + 1) - \int \left(\frac{x + 1 - 1}{x + 1} \right) dx$$

$$= \log x - \int \frac{1}{x} dx + \log(x + 1) - \int \left(1 - \frac{1}{x + 1} \right) dx$$

$$= \log x - \int \frac{1}{x} dx + \log(x + 1) - x + \log|x + 1| + c$$

$$= [\log x + \log(x + 1)] - 2x + \log|x + 1| + c$$

$$= \log(x^2 + x) - 2x + \log|x + 1| + c$$

$$\therefore A = -2x + \log|x + 1| + c$$

$$165. I_1 = \int \sin^{-1} x dx$$

Put $\sin^{-1} x = \theta \Rightarrow x = \sin \theta \Rightarrow dx = \cos \theta d\theta$

$$\therefore I_1 = \int \theta \cos \theta d\theta = \theta \sin \theta - \int 1 \cdot \sin \theta d\theta$$

$$= \theta \sin \theta + \cos \theta$$

$$= \sin^{-1} x + \sqrt{1 - x^2}$$

Now, $I_2 = \int \sin^{-1} \sqrt{1 - x^2} dx = \int \cos^{-1} x dx$

Put $\cos^{-1} x = \phi \Rightarrow x = \cos \phi \Rightarrow dx = -\sin \phi d\phi$

$$\therefore I_2 = -\int \phi \sin \phi d\phi = \phi \cos \phi + \int 1 \cdot (-\cos \phi) d\phi$$

$$= \phi \cos \phi - \sin \phi = \cos^{-1} x - \sqrt{1 - x^2}$$

$$\therefore I_1 + I_2 = (\sin^{-1} x + \cos^{-1} x) = \frac{\pi}{2}$$

$$166. \frac{d}{dt} = f''(t) \cos t + f'(t) \sin t$$

$$\therefore \frac{d}{dt} = -f''(t) \sin t + f'''(t) \cos t + f''(t) \sin t$$

$$+ f'(t) \cos t$$

$$= f'''(t) \cos t + f'(t) \cos t$$

$$= \cos t [f'''(t) + f'(t)]$$

$$= -f''(t) \sin t + f'(t) \cos t$$

$$\therefore \frac{d}{dt} = -f'''(t) \sin t - f''(t) \cos t + f''(t) \cos t$$

$$- f'(t) \sin t$$

$$= -\sin t [f'''(t) + f'(t)]$$

$$\therefore \left(\frac{d}{dt} \right)^2 + \left(\frac{d}{dt} \right)^2 = [\sin^2 t + \cos^2 t] [f'''(t) + f'(t)]^2$$

$$= [f'''(t) + f'(t)]^2$$

$$\text{Let } I = \int \left[\left(\frac{d}{dt} \right)^2 + \left(\frac{d}{dt} \right)^2 \right]^{\frac{1}{2}} dt$$

$$= \int \{ [f'''(t) + f'(t)]^2 \}^{\frac{1}{2}} dt$$

$$= \int [f'''(t) + f'(t)] dt$$

$$= f''(t) + f(t) + c$$



Evaluation Test

$$\begin{aligned}
 1. \quad \text{Let } I &= \int \sqrt{\frac{5+t^{10}}{16}} dt \\
 &= \int \sqrt{\frac{5+t^{10}}{10}} \cdot \frac{1}{11} dt \\
 &= \int \sqrt{\frac{5}{10} + 1} \cdot \frac{1}{11} dt
 \end{aligned}$$

$$\text{Put } \frac{5}{10} + 1 = t$$

$$\therefore 5(-10)^{-11} dt = dt$$

$$\therefore \frac{1}{11} dt = -\frac{1}{50} dt$$

$$\therefore I = \int t^{\frac{1}{2}} \left(-\frac{1}{50} \right) dt$$

$$= -\frac{1}{50} \cdot \frac{t^{3/2}}{3/2} + c$$

$$= -\frac{1}{75} \left(1 + \frac{5}{10} \right)^{3/2} + c$$

2. Multiplying N^r and D^r by $\sin 3$, we get

$$\int \frac{\cos 5 + \cos 4}{1 - 2\cos 3} dt$$

$$= \int \frac{\sin 3 \cos 5 + \sin 3 \cos 4}{\sin 3 - 2\sin 3 \cos 3} dt$$

$$= \int \frac{\sin 3 (\cos 5 + \cos 4)}{\sin 3 - \sin 6} dt$$

$$= \int \frac{\left(2\sin \frac{3}{2} \cos \frac{3}{2} \right) \left(2\cos \frac{9}{2} \cos \frac{3}{2} \right)}{-2\cos \frac{9}{2} \sin \frac{3}{2}} dt$$

$$= - \int 2\cos \frac{3}{2} \cos \frac{3}{2} dt$$

$$= - \int (\cos 3 + \cos 6) dt$$

$$= - \left[\frac{1}{2} \sin 3 + \sin 6 \right] + c$$

$$3. \quad \text{Let } I = \int \sin^{-1} \sqrt{\frac{a+t}{a+t^2}} dt$$

$$\text{Put } t = a \tan^2 t$$

$$\therefore dt = 2a \tan t \sec^2 t dt$$

$$\begin{aligned}
 \therefore I &= \int \sin^{-1} \sqrt{\frac{a \tan^2 t}{a + a \tan^2 t}} \times 2a \tan t \sec^2 t dt \\
 &= \int \sin^{-1}(\sin t) \times 2a \tan t \sec^2 t dt \\
 &= 2a \int t \tan t \sec^2 t dt \\
 &= 2a \left[t \int \tan t \sec^2 t dt - \int \left\{ \frac{d}{dt}(t) \int \tan t \sec^2 t dt \right\} dt \right]
 \end{aligned}$$

$$= 2a \left[t \cdot \frac{\tan^2 t}{2} - \int 1 \cdot \frac{\tan^2 t}{2} dt \right]$$

$$= a \left[t \tan^2 t - \int (\sec^2 t - 1) dt \right]$$

$$= a \left[t \tan^2 t - \tan t + t \right] + c, \text{ where } t = \tan^{-1} \sqrt{\frac{a}{a+t}}$$

$$= a \left[-\tan^{-1} \sqrt{\frac{a}{a+t}} - \sqrt{\frac{a}{a+t}} + \tan^{-1} \sqrt{\frac{a}{a+t}} \right] + c$$

$$4. \quad \text{Let } I = \int \sqrt{\operatorname{cosec}^{-1} - 1} dt$$

$$= \int \sqrt{\frac{1}{\sin} - 1} dt$$

$$= \int \sqrt{\frac{1 - \sin}{\sin}} dt$$

$$= \int \sqrt{\frac{1 - \sin}{\sin}} \times \frac{1 + \sin}{1 + \sin} dt$$

$$= \int \frac{\cos}{\sqrt{\sin^2 + \sin}} dt$$

$$\text{Put } \sin t = t$$

$$\therefore \cos t dt = dt$$

$$\therefore I = \int \frac{1}{\sqrt{t^2 + t}} dt$$

$$= \int \frac{1}{\sqrt{t^2 + t + \frac{1}{4} - \frac{1}{4}}} dt$$

$$= \int \frac{1}{\sqrt{\left(t + \frac{1}{2}\right)^2 - \left(\frac{1}{2}\right)^2}} dt$$

$$= \log \left| t + \frac{1}{2} + \sqrt{t^2 + t} \right| + c, \text{ where } t = \sin$$

$$= \log \left| \sin t + \frac{1}{2} + \sqrt{\sin^2 t + \sin t} \right| + c$$



5. Let $I = \int \sqrt{\tan} \, d$
 Put $\tan = t^2$
 $\therefore \sec^2 d = 2t dt$
 $\therefore d = \frac{2t}{1+t^4} dt$
 $\therefore I = \int \sqrt{t^2} \cdot \frac{2t}{1+t^4} dt = 2 \int \frac{t^2}{1+t^4} dt$
 $= \int \frac{t^2+1+t^2-1}{t^4+1} dt = \int \frac{t^2+1}{t^4+1} dt + \int \frac{t^2-1}{t^4+1} dt$
 $= I_1 + I_2 \text{ (say)} \quad \dots \text{(i)}$

$I_1 = \int \frac{t^2+1}{t^4+1} dt$
 $= \int \frac{1+\frac{1}{t^2}}{t^2+\frac{1}{t^2}} dt$
 $= \int \frac{1}{\left(t-\frac{1}{t}\right)^2+2} \left(1+\frac{1}{t^2}\right) dt$
 $= \int \frac{1}{t^2+2} d, \text{ where } t - \frac{1}{t} =$
 $= \frac{1}{\sqrt{2}} \tan^{-1}\left(\frac{t-\frac{1}{t}}{\sqrt{2}}\right) = \frac{1}{\sqrt{2}} \tan^{-1}\left(\frac{t^2-1}{\sqrt{2}t}\right)$
 $= \frac{1}{\sqrt{2}} \tan^{-1}\left(\frac{t^2-1}{\sqrt{2}t}\right)$

$I_2 = \int \frac{t^2-1}{t^4+1} dt = \int \frac{1-\frac{1}{t^2}}{t^2+\frac{1}{t^2}} dt$
 $= \int \frac{1}{\left(t+\frac{1}{t}\right)^2-2} \left(1-\frac{1}{t^2}\right) dt$
 $= \int \frac{1}{m^2-2} dm, \text{ where } t + \frac{1}{t} = m$
 $= \frac{1}{2\sqrt{2}} \log \left| \frac{m-\sqrt{2}}{m+\sqrt{2}} \right| = \frac{1}{2\sqrt{2}} \log \left| \frac{t+\frac{1}{t}-\sqrt{2}}{t+\frac{1}{t}+\sqrt{2}} \right|$
 $= \frac{1}{2\sqrt{2}} \log \left| \frac{t^2-\sqrt{2}t+1}{t^2+\sqrt{2}t+1} \right|$

\therefore From (i),
 $I = \frac{1}{\sqrt{2}} \tan^{-1}\left(\frac{t^2-1}{\sqrt{2}t}\right) + \frac{1}{2\sqrt{2}} \log \left| \frac{t^2-\sqrt{2}t+1}{t^2+\sqrt{2}t+1} \right| + c$
 $= \frac{1}{\sqrt{2}} \tan^{-1}\left(\frac{\tan-1}{\sqrt{2}\tan}\right)$
 $+ \frac{1}{2\sqrt{2}} \log \left| \frac{\tan-\sqrt{2}\tan+1}{\tan+\sqrt{2}\tan+1} \right| + c$

6. Let $I = \int (1+t^{5/2})^{1/2} d$
 $= \int (1+t^{5/2})^{1/2} \cdot \frac{5}{2} t^{3/2} dt$
 Put $1+t^{5/2} = t$
 $\therefore \frac{5}{2} t^{3/2} dt = dt, \quad \therefore t^{3/2} dt = \frac{2}{5} dt$
 $\therefore I = \int (t-1)^2 \cdot t^{1/2} \cdot \frac{2}{5} dt$
 $= \frac{2}{5} \int (t^{5/2} - 2t^{3/2} + t^{1/2}) dt$
 $= \frac{2}{5} \left[\frac{2}{7} t^{7/2} - \frac{4}{5} t^{5/2} + \frac{2}{3} t^{3/2} \right] + c, \text{ where } t = 1+t^{5/2}$
 $= \frac{2}{5} \left[\frac{2}{7} (1+t^{5/2})^{7/2} - \frac{4}{5} (1+t^{5/2})^{5/2} \right. \\ \left. + \frac{2}{3} (1+t^{5/2})^{3/2} \right] + c$

7. Let $I = \int \frac{\tan}{1+\tan+\tan^2} d$
 $= \int \frac{\tan}{\sec^2+\tan} d$
 $= \int \frac{\frac{\sin}{\cos}}{\frac{1}{\cos^2}+\frac{\sin}{\cos}} d = \int \frac{\sin \cos}{1+\sin \cos} d$
 $= \int \frac{\frac{1}{2} \sin 2}{1+\frac{1}{2} \sin 2} d = \int \frac{\sin 2}{2+\sin 2} d$
 $= \int \frac{2+\sin 2-2}{2+\sin 2} d = \int \left(1 - \frac{2}{2+\sin 2}\right) d$
 $= -I_1 \text{ (say)} \quad \dots \text{(i)}$
 $I_1 = \int \frac{2}{2+\sin 2} d$



$$\text{Put } \tan^{-1} t = \theta \Rightarrow \sec^2 \theta \, d\theta = dt \Rightarrow d\theta = \frac{1}{1+t^2} dt$$

$$\sin 2\theta = \frac{2 \tan \theta}{1 + \tan^2 \theta} = \frac{2t}{1+t^2}$$

$$\therefore I_1 = \int \frac{2}{2 + \frac{2t}{1+t^2}} \times \frac{1}{1+t^2} dt$$

$$= \int \frac{1}{t^2 + t + 1} dt$$

$$= \int \frac{1}{t^2 + t + \frac{1}{4} + \frac{3}{4}} dt$$

$$= \int \frac{1}{\left(t + \frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2} dt$$

$$= \frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{t + \frac{1}{2}}{\frac{\sqrt{3}}{2}} \right) + c_1$$

$$= \frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{2 \tan^{-1} t + 1}{\sqrt{3}} \right) + c_1$$

\(\therefore\) From (i),

$$I = -\frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{2 \tan^{-1} t + 1}{\sqrt{3}} \right) + c$$

$$\therefore \sqrt{A} = \sqrt{3}$$

$$\therefore A = 3$$

$$8. \text{ Let } I = \int \log \left(\frac{-1}{+1} \right) \cdot \frac{1}{x^2 - 1} dx$$

$$\text{Put } \log \left(\frac{-1}{+1} \right) = t$$

$$\therefore \left(\frac{1}{-1} - \frac{1}{+1} \right) dx = dt$$

$$\therefore \frac{1}{x^2 - 1} dx = \frac{1}{2} dt$$

$$\therefore I = \int t \cdot \frac{1}{2} dt = \frac{1}{4} t^2 + c$$

$$= \frac{1}{4} \left[\log \left(\frac{-1}{+1} \right) \right]^2 + c$$

$$\therefore A = \frac{1}{4}$$

$$9. \text{ Let } I = \int \frac{1}{(x^2 + 2x + 2)^2} dx$$

$$= \int \frac{1}{[(x+1)^2 + 1]^2} dx$$

$$\text{Put } x+1 = \tan \theta$$

$$\therefore dx = \sec^2 \theta \, d\theta$$

$$\therefore I = \int \frac{\sec^2 \theta}{(\tan^2 \theta + 1)^2} d\theta$$

$$= \int \frac{\sec^2 \theta}{\sec^4 \theta} d\theta$$

$$= \int \cos^2 \theta \, d\theta$$


$$= \frac{1}{2} \int (1 + \cos 2\theta) d\theta$$

$$= \frac{1}{2} \left[\theta + \frac{\sin 2\theta}{2} \right] + c$$

$$= \frac{1}{2} (\theta + \sin \theta \cos \theta) + c$$

$$= \frac{1}{2} \left[\tan^{-1}(x+1) + \frac{x+1}{\sqrt{x^2+2x+2}} \cdot \frac{1}{\sqrt{x^2+2x+2}} \right] + c$$

$$= \frac{1}{2} \left[\tan^{-1}(x+1) + \frac{x+1}{x^2+2x+2} \right] + c$$



$$10. \text{ Let } I = \int \frac{1}{\cos^6 x + \sin^6 x} dx$$

$$\text{Since, } a^3 + b^3 = (a+b)^3 - 3ab(a+b)$$

$$\therefore \cos^6 x + \sin^6 x = 1 - 3 \sin^2 x \cos^2 x$$

$$\dots [\because a+b = \cos^2 x + \sin^2 x = 1]$$

$$\therefore I = \int \frac{1}{1 - 3 \sin^2 x \cos^2 x} dx$$

$$= \int \frac{1}{1 - \frac{3}{4} \sin^2 2x} dx$$

$$= \int \frac{4}{4 - 3 \sin^2 2x} dx$$

$$= \int \frac{4 \operatorname{cosec}^2 2x}{4 \operatorname{cosec}^2 2x - 3} dx$$

$$\int \frac{4 \operatorname{cosec}^2 2x}{4(1 + \cot^2 2x) - 3} dx$$

$$= - \int \frac{1}{4 \cot^2 2x + 1} (-4 \operatorname{cosec}^2 2x) dx$$

$$\text{Put } 2 \cot 2x = t$$

$$\therefore -4 \operatorname{cosec}^2 2x \, dx = dt$$



$$\begin{aligned} \therefore I &= -\int \frac{1}{t^2+1} dt \\ &= -\tan^{-1}(t) + c \\ &= -\tan^{-1}(2 \cot 2) + c \\ &= -\tan^{-1}\left(\frac{2 \cos^2 - 2 \sin^2}{2 \sin \cos}\right) + c \\ &= -\tan^{-1}(\cot - \tan) + c \\ &= \tan^{-1}(\tan - \cot) + c \end{aligned}$$

$$11. \int f(x) \sin \cos x dx = \frac{1}{2(b^2 - a^2)} \log[f(x)] + c$$

$$\therefore \frac{d}{dx} \left[\frac{1}{2(b^2 - a^2)} \log[f(x)] + c \right] = f(x) \sin \cos x$$

$$\therefore \frac{1}{2(b^2 - a^2)} \cdot \frac{1}{f(x)} \cdot f'(x) = f(x) \sin \cos x$$

$$\therefore \frac{f'(x)}{[f(x)]^2} = 2(b^2 - a^2) \sin \cos x$$

Integrating on both sides, we get

$$\int \frac{f'(x)}{[f(x)]^2} dx = (b^2 - a^2) \int 2 \sin \cos x dx$$

$$\begin{aligned} \therefore -\frac{1}{f(x)} &= (b^2 - a^2) \int 2 \sin \cos x dx \\ &= b^2 \int 2 \sin \cos x dx - a^2 \int 2 \sin \cos x dx \\ &= b^2 (-\cos^2 x) - a^2 (\sin^2 x) \end{aligned}$$

$$\therefore -\frac{1}{f(x)} = -b^2 \cos^2 x - a^2 \sin^2 x$$

$$\therefore f(x) = \frac{1}{a^2 \sin^2 x + b^2 \cos^2 x}$$

$$12. \text{ Let } I = \int e^{\sin \theta} [\log(\sin \theta) + \operatorname{cosec}^2 \theta] \cos \theta d\theta$$

Put $\sin \theta = t$

$$\therefore \cos \theta d\theta = dt$$

$$\therefore I = \int e^t \left[\log t + \frac{1}{t^2} \right] dt$$

$$= \int e^t \left[\log t - \frac{1}{t} + \frac{1}{t} + \frac{1}{t^2} \right] dt$$

$$= e^t \left(\log t - \frac{1}{t} \right) + c$$

$$\dots \left[\because \frac{d}{dt} \left(\log t - \frac{1}{t} \right) = \frac{1}{t} + \frac{1}{t^2} \right]$$

$$= e^{\sin \theta} \left[\log(\sin \theta) - \frac{1}{\sin \theta} \right] + c$$

$$= e^{\sin \theta} [\log(\sin \theta) - \operatorname{cosec} \theta] + c$$

$$\begin{aligned} 13. \log \left(\frac{\cos \theta + \sin \theta}{\cos \theta - \sin \theta} \right) &= \log \left(\frac{1 + \tan \theta}{1 - \tan \theta} \right) \\ &= \log \tan \left(\frac{\pi}{4} + \theta \right) \end{aligned}$$

$$\text{Since, } \int \sec 2\theta d\theta = \frac{1}{2} \log \tan \left(\frac{\pi}{4} + \theta \right)$$

$$\therefore \frac{d}{d\theta} \log \tan \left(\frac{\pi}{4} + \theta \right) = 2 \sec 2\theta \quad \dots (i)$$

Integrating the given expression by parts, we get

$$I = \log \tan \left(\frac{\pi}{4} + \theta \right) \cdot \frac{1}{2} \sin 2\theta - \int \frac{\sin 2\theta}{2} \cdot 2 \sec 2\theta d\theta$$

....[From (i)]

$$= \frac{1}{2} \sin 2\theta \log \tan \left(\frac{\pi}{4} + \theta \right) - \int \tan 2\theta d\theta$$

$$= \frac{1}{2} \sin 2\theta \log \tan \left(\frac{\pi}{4} + \theta \right) - \frac{1}{2} \log(\sec 2\theta) + c$$

$$14. \text{ Let } \frac{3 - 4}{3 + 4} = t$$

$$\Rightarrow \frac{(3 - 4) + (3 + 4)}{(3 - 4) - (3 + 4)} = \frac{t + 1}{t - 1}$$

$$\Rightarrow \frac{6}{-8} = \frac{t + 1}{t - 1} \Rightarrow = -\frac{4}{3} \left(\frac{t + 1}{t - 1} \right)$$

$$\Rightarrow + 2 = -\frac{4t + 4}{3t - 3} + 2$$

$$= \frac{-4t - 4 + 6t - 6}{3t - 3} = \frac{2t - 10}{3t - 3}$$

$$\text{Given, } f \left(\frac{3 - 4}{3 + 4} \right) = + 2$$

$$\therefore f(t) = \frac{2t - 10}{3t - 3} = \frac{2}{3} \left(\frac{t - 5}{t - 1} \right)$$

$$= \frac{2}{3} \left(\frac{t - 1 - 4}{t - 1} \right) = \frac{2}{3} \left(1 - \frac{4}{t - 1} \right) = \frac{2}{3} - \frac{8}{3(t - 1)}$$

$$\therefore f(x) = \frac{2}{3} - \frac{8}{3(x - 1)}$$

$$\therefore \int f(x) dx = \int \left\{ \frac{2}{3} - \frac{8}{3(x - 1)} \right\} dx$$

$$= \frac{2}{3} x - \frac{8}{3} \log |x - 1| + c$$



$$\begin{aligned}
 15. \quad \text{Let } I &= \int \log(1-\sqrt{x}) dx \\
 &= \int \log(1-\sqrt{x})(1) dx \\
 &= \log(1-\sqrt{x}) \int 1 dx - \int \left\{ \frac{d}{dx} \log(1-\sqrt{x}) \int 1 dx \right\} dx \\
 &= \log(1-\sqrt{x}) \cdot x - \int \frac{1}{1-\sqrt{x}} \left(-\frac{1}{2\sqrt{x}} \right) dx \\
 &= \log(1-\sqrt{x}) + \frac{1}{2} \int \frac{\sqrt{x}}{1-\sqrt{x}} dx \\
 &= \log(1-\sqrt{x}) + \frac{1}{2} I_1 \quad \dots(i)
 \end{aligned}$$

$$\text{Now, } I_1 = \int \frac{\sqrt{x}}{1-\sqrt{x}} dx$$

$$\text{Put } x = t^2,$$

$$\therefore \frac{dx}{dt} = 2t \quad dt$$

$$\begin{aligned}
 \therefore I_1 &= \int \frac{\sqrt{t^2}}{1-\sqrt{t^2}} \cdot 2t dt = 2 \int \frac{t^2}{1-t} dt \\
 &= -2 \int \frac{1-t^2-1}{1-t} dt \\
 &= -2 \int \left(1+t - \frac{1}{1-t} \right) dt \\
 &= 2 \int \left(\frac{1}{1-t} - 1 - t \right) dt \\
 &= 2 \left[\frac{1}{-1} \log(1-t) - t - \frac{t^2}{2} \right] + c_1 \\
 &= -2 \left[\log(1-\sqrt{x}) + \sqrt{x} + \frac{1}{2} \right] + c_1
 \end{aligned}$$

\therefore From (i),

$$\begin{aligned}
 I &= \log(1-\sqrt{x}) \\
 &\quad - \frac{1}{2} \cdot 2 \left[\log(1-\sqrt{x}) + \sqrt{x} + \frac{1}{2} \right] + c \\
 &= (-1) \log(1-\sqrt{x}) - \sqrt{x} - \frac{1}{2} + c
 \end{aligned}$$

$$16. \quad P(x) = \int \frac{x^3}{3x-2} dx, \quad Q(x) = \int \frac{1}{3x-2} dx$$

$$\begin{aligned}
 \therefore P(x) + Q(x) &= \int \frac{x^3+1}{3x-2} dx \\
 &= \int \frac{x^3-2^2+2^2+1}{3x-2} dx \\
 &= \int \left(1 + \frac{2^2+1}{3x-2} \right) dx \\
 &= x + I \quad \dots(i)
 \end{aligned}$$

$$I = \int \frac{x^2+1}{2(3x-2)} dx$$

$$\text{Put } \frac{x^2+1}{2(3x-2)} = \frac{A}{3x-2} + \frac{B}{x} + \frac{C}{2}$$

$$\therefore x^2+1 = A(3x-2) + B(3x-2)x + C(3x-2) \quad \dots(ii)$$

$$\text{Putting } x=0 \text{ in (ii), } C = -1$$

$$\text{Putting } x=1 \text{ in (ii), } A = 2$$

$$\text{Putting } x=-1 \text{ in (ii), } B = -1$$

$$\begin{aligned}
 \therefore I &= \int \left(\frac{2}{-1} - \frac{1}{x} - \frac{1}{2} \right) dx \\
 &= 2 \log| -1 | - \log|x| + \frac{1}{2}
 \end{aligned}$$

\therefore From (i),

$$P(x) + Q(x) = x + 2 \log| -1 | - \log|x| + \frac{1}{2} + c$$

$$\therefore (P+Q)(2) = P(2) + Q(2)$$

$$= 2 + 2 \log 1 - \log 2 + \frac{1}{2} + c$$

$$\therefore \frac{5}{2} = \frac{5}{2} - \log 2 + c \quad \dots \left[\because (P+Q)(2) = \frac{5}{2} \right]$$

$$\therefore c = \log 2$$

$$\therefore P(x) + Q(x) = x + 2 \log| -1 | - \log|x| + \frac{1}{2} + \log 2$$

$$\therefore P(3) + Q(3) = 3 + 2 \log 2 - \log 3 + \frac{1}{2} + \log 2$$

$$= \frac{10}{3} + \log \frac{8}{3}$$

$$17. \quad \text{Let } I = \int \frac{2a \sin x + b \sin 2x}{(b+a \cos x)^3} dx$$

$$= 2 \int \frac{(a+b \cos x)}{(b+a \cos x)^3} \cdot \sin x dx$$

$$\text{Put } b+a \cos x = t$$

$$\therefore -a \sin x dx = dt$$

$$\therefore \sin x dx = -\frac{1}{a} dt$$

$$\therefore I = 2 \int \frac{a+b \left(\frac{t-b}{a} \right)}{t^3} \left(-\frac{1}{a} \right) dt$$

$$= -\frac{2}{a} \int \frac{a^2+bt-b^2}{at^3} dt$$

$$= -\frac{2}{a^2} \int [(a^2-b^2)t^{-3} + bt^{-2}] dt$$



$$= -\frac{2}{a^2} \left[\frac{a^2 - b^2}{-2t^2} + \frac{b}{-t} \right] + c$$

$$= \frac{1}{a^2} \cdot \frac{a^2 - b^2}{t^2} + \frac{2b}{a^2 t} + c$$

18. Let $I = \int \frac{1}{(-1)\sqrt{t^2 + 4}} dt$

Put $-1 = \frac{1}{t}$, $\therefore d = -\frac{1}{t^2} dt$

$$\therefore I = \int \frac{1}{\frac{1}{t} \sqrt{\left(\frac{1}{t} + 1\right)^2 + 4}} \left(-\frac{1}{t^2}\right) dt$$

$$= - \int \frac{1}{\sqrt{\frac{1}{t^2} + \frac{2}{t} + 1 + 4}} \left(\frac{1}{t}\right) dt$$

$$= - \int \frac{1}{\sqrt{1 + 2t + 5t^2}} dt$$

$$= -\frac{1}{\sqrt{5}} \int \frac{1}{\sqrt{t^2 + \frac{2}{5}t + \frac{1}{5}}} dt$$

$$= -\frac{1}{\sqrt{5}} \int \frac{1}{\sqrt{t^2 + \frac{2}{5}t + \frac{1}{25} + \frac{4}{25}}} dt$$

$$= -\frac{1}{\sqrt{5}} \int \frac{1}{\sqrt{\left(t + \frac{1}{5}\right)^2 + \left(\frac{2}{5}\right)^2}} dt$$

$$= -\frac{1}{\sqrt{5}} \log \left| t + \frac{1}{5} + \sqrt{t^2 + \frac{2t}{5} + \frac{1}{5}} \right| + c$$

$$= -\frac{1}{\sqrt{5}} \log \left| \frac{1}{-1} + \frac{1}{5} + \sqrt{\frac{1}{(-1)^2} + \frac{2}{5(-1)} + \frac{1}{5}} \right| + c$$

$$= -\frac{1}{\sqrt{5}} \log \left| \frac{1}{-1} + \frac{1}{5} + \sqrt{\frac{2+4}{5(-1)^2}} \right| + c$$

19. Let $I = \int \frac{1 + \cos}{\left\{1 - \left(e^{\sin}\right)^2\right\}} dt$

$$= \int \frac{e^{\sin} (1 + \cos)}{e^{\sin} \left\{1 - \left(e^{\sin}\right)^2\right\}} dt$$

Put $e^{\sin} = t$

$$\therefore [e^{\sin} \cos + e^{\sin} (1)] dt = dt$$

$$\therefore e^{\sin} (1 + \cos) dt = dt$$

$$\therefore I = \int \frac{1}{t(1-t^2)} dt = \int \frac{1}{t(1-t)(1+t)} dt$$

Put $\frac{1}{t(1-t)(1+t)} = \frac{A}{t} + \frac{B}{1-t} + \frac{C}{1+t}$

$$\therefore 1 = A(1-t)(1+t) + Bt(1+t) + Ct(1-t) \quad \dots(i)$$

Putting $t = 0$ in (i), we get

$$A = 1$$

Putting $t = 1$ in (i), we get

$$B = \frac{1}{2}$$

Putting $t = -1$ in (i), we get

$$C = -\frac{1}{2}$$

$$\therefore I = \int \left(\frac{1}{t} + \frac{1/2}{1-t} - \frac{1/2}{1+t} \right) dt$$

$$= \log|t| - \frac{1}{2} \log|1-t| - \frac{1}{2} \log|1+t| + c$$

$$= \frac{1}{2} \log \left| \frac{t^2}{1-t^2} \right| + c = \frac{1}{2} \log \left| \frac{e^{2\sin}}{1 - e^{2\sin}} \right| + c$$

20. Let $I = \int \tan(\sin^{-1}) dt$

$$= \int \tan \left[\tan^{-1} \left(\frac{t}{\sqrt{1-t^2}} \right) \right] dt$$

$$= \int \frac{t}{\sqrt{1-t^2}} dt$$

Put $1 - t^2 = t$,

$$\therefore -2 dt = dt,$$

$$\therefore dt = -\frac{1}{2} dt$$

$$\therefore I = \int \frac{1}{\sqrt{t}} \left(-\frac{1}{2}\right) dt$$

$$= - \int \frac{1}{2\sqrt{t}} dt = -\sqrt{t} + c = -\sqrt{1-t^2} + c$$

21. Let $I = \int \sec^{25/13} \operatorname{cosec}^{27/13} dt$

$$= \int \cos^{-25/13} \sin^{-27/13} dt$$

Now $-\frac{25}{13} - \frac{27}{13} = -\frac{52}{13} = -4$

Multiplying and dividing by \cos^4 , we get

$$I = \int \cos^4 \cos^{-25/13} \sin^{-27/13} \sec^4 dt$$

$$= \int \tan^{-27/13} (1 + \tan^2) \sec^2 dt$$



$$\text{Put } \tan^{-1} t = t, \quad \therefore \sec^2 t \, dt = dt$$

$$\begin{aligned} \therefore I &= \int t^{-27/13} (1+t^2) dt \\ &= \int (t^{-27/13} + t^{-1/13}) dt \\ &= -\frac{13}{14} t^{-14/13} + \frac{13}{12} t^{12/13} + c \\ &= -\frac{13}{14} (\tan^{-1} t)^{-14/13} + \frac{13}{12} (\tan^{-1} t)^{12/13} + c \end{aligned}$$

$$22. \text{ Let } I = \int \frac{1}{\sqrt{t^2 - 2t + 1}} dt$$

$$\text{Put } \sqrt{t^2 - 2t + 1} = t$$

$$\therefore \sqrt{t^2 - 2t + 1} = t - 1$$

$$\therefore t^2 - 2t + 1 = t^2 - 2t + 1$$

$$\therefore = \frac{t^2 - 1}{2t - 1}$$

$$\therefore \frac{d}{dt} = \frac{(2t-1) \cdot 2t - (t^2-1) \cdot 2}{(2t-1)^2}$$

$$\therefore dt = \frac{2t^2 - 2t + 2}{(2t-1)^2} dt$$

$$\therefore I = \int \frac{1}{t} \times \frac{2t^2 - 2t + 2}{(2t-1)^2} dt$$

$$= 2 \int \frac{t^2 - t + 1}{t(2t-1)^2} dt$$

$$\text{Put } \frac{t^2 - t + 1}{t(2t-1)^2} = \frac{A}{t} + \frac{B}{2t-1} + \frac{C}{(2t-1)^2}$$

$$\therefore t^2 - t + 1 = A(2t-1)^2 + Bt(2t-1) + Ct$$

....(i)

Putting $t = 0$ in (i), we get

$$A = 1$$

Putting $t = \frac{1}{2}$ in (i), we get

$$C = \frac{3}{2}$$

Putting $t = 1$ in (i), we get

$$1 = A + B + C$$

$$\Rightarrow B = 1 - 1 - \frac{3}{2} \Rightarrow B = -\frac{3}{2}$$

$$\therefore I = 2 \int \left(\frac{1}{t} - \frac{3}{2(2t-1)} + \frac{3}{2} \cdot \frac{1}{(2t-1)^2} \right) dt$$

$$= 2 \log t - \frac{3}{2} \log(2t-1) - \frac{3}{2} \cdot \frac{1}{2t-1} + c$$

$$= 2 \log t - \frac{3}{2} \log(2t-1) - \frac{1}{2} \left(\frac{3}{2t-1} \right) + c,$$

$$\text{where } t = \sqrt{t^2 - 2t + 1} + 1$$

$$\text{and } 2t - 1 = 2 - 1 + 2\sqrt{t^2 - 2t + 1}$$

$$\therefore P = 2, Q = -\frac{3}{2}, R = -\frac{1}{2}$$

$$23. \text{ Let } I = \int \frac{1}{(1+t^2)\sqrt{1-t^2}} dt$$

$$\text{Put } t = \frac{1}{t}$$

$$\therefore dt = -\frac{1}{t^2} dt$$

$$\therefore I = \int \frac{1}{\left(1 + \frac{1}{t^2}\right)\sqrt{1 - \frac{1}{t^2}}} \left(-\frac{1}{t^2}\right) dt$$

$$= -\int \frac{t \, dt}{(t^2+1)\sqrt{t^2-1}}$$

$$\text{Put } t^2 - 1 = m^2$$

$$\therefore 2t \, dt = 2m \, dm,$$

$$\therefore t \, dt = m \, dm$$

$$\therefore I = -\int \frac{m \, dm}{(m^2+2)\sqrt{m^2}}$$

$$= -\int \frac{1}{m^2 + (\sqrt{2})^2} dm$$

$$= -\frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{m}{\sqrt{2}} \right) + c$$

$$= -\frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{t^2-1}}{\sqrt{2}} \right) + c$$

$$= -\frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{\frac{1}{t^2}-1}}{\sqrt{2}} \right) + c$$

$$= -\frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{1-t^2}}{\sqrt{2}} \right) + c$$

$$= -\frac{1}{\sqrt{2}} \left[\frac{\pi}{2} - \cot^{-1} \left(\frac{\sqrt{1-t^2}}{\sqrt{2}} \right) \right] + c$$

$$= -\frac{1}{\sqrt{2}} \left[\frac{\pi}{2} - \tan^{-1} \left(\frac{\sqrt{2}}{\sqrt{1-t^2}} \right) \right] + c$$

$$= \frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{2}}{\sqrt{1-t^2}} \right) - \frac{\pi}{2\sqrt{2}} + c$$



24. Let $I = \int \frac{2^x - 1}{\sqrt[3]{2^{4x} - 2^{2x} + 1}} dx$
 Dividing N^r and D^r by 2^{5x} , we get

$$I = \int \frac{\left(\frac{1}{3} - \frac{1}{5}\right)}{\sqrt{2 - \frac{2}{2} + \frac{1}{4}}} dx$$

 Put $2 - \frac{2}{2} + \frac{1}{4} = t \Rightarrow \left(\frac{4}{3} - \frac{4}{5}\right) dx = dt$

$$\therefore I = \frac{1}{4} \int \frac{dt}{\sqrt{t}} = \frac{1}{2} \sqrt{t} + c$$

$$= \frac{1}{2} \sqrt{2 - \frac{2}{2} + \frac{1}{4}} + c$$
25. $\int \frac{\log}{(x+1)^2} dx = \int \log(x+1)^{-2} dx$

$$= \log(x+1) \cdot \frac{(x+1)^{-1}}{-1} - \int \frac{1 \cdot (x+1)^{-1}}{-1} dx$$

$$= -\frac{\log}{x+1} + \int \left(\frac{1}{x+1}\right) dx$$

$$= -\frac{\log}{x+1} + \int \left(\frac{1}{x} - \frac{1}{x+1}\right) dx$$

$$= -\frac{\log}{x+1} + \log|x| - \log|x+1| + c$$
26. $I_n = \int \sin^n x dx$

$$= \int \sin^{n-1} x \cdot \sin x dx$$

$$= \sin^{n-1} x \int \sin x dx - \int \left\{ \frac{d}{dx} (\sin^{n-1} x) \int \sin x dx \right\} dx$$

$$= \sin^{n-1} x (-\cos x) - \int (n-1) \sin^{n-2} x \cos x (-\cos x) dx$$

$$= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x \cos^2 x dx$$

$$= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x (1 - \sin^2 x) dx$$

$$= -\sin^{n-1} x \cos x + (n-1) \int (\sin^{n-2} x - \sin^n x) dx$$

$$= -\sin^{n-1} x \cos x + (n-1) \int \sin^{n-2} x dx - (n-1) \int \sin^n x dx$$

$$\therefore I_n = -\sin^{n-1} x \cos x + (n-1) I_{n-2} - (n-1) I_n$$

$$\therefore I_n + (n-1) I_n - (n-1) I_{n-2} = -\sin^{n-1} x \cos x$$

$$\therefore n I_n - (n-1) I_{n-2} = -\sin^{n-1} x \cos x$$

27. $u = -f''(\theta) \sin \theta + f'(\theta) \cos \theta$

$$\therefore \frac{du}{d\theta} = -f'''(\theta) \sin \theta - f''(\theta) \cos \theta$$

$$+ f''(\theta) \cos \theta - f'(\theta) \sin \theta$$

$$= -f'''(\theta) \sin \theta - f'(\theta) \sin \theta$$

$$v = f''(\theta) \cos \theta + f'(\theta) \sin \theta$$

$$\therefore \frac{dv}{d\theta} = -f''(\theta) \sin \theta + f'''(\theta) \cos \theta$$

$$+ f'(\theta) \cos \theta + f''(\theta) \sin \theta$$

$$= f'''(\theta) \cos \theta + f'(\theta) \cos \theta$$

$$\therefore \left(\frac{du}{d\theta}\right)^2 + \left(\frac{dv}{d\theta}\right)^2 = [-f'''(\theta) \sin \theta - f'(\theta) \sin \theta]^2$$

$$+ [f'''(\theta) \cos \theta + f'(\theta) \cos \theta]^2$$

$$= [f'''(\theta)]^2 \sin^2 \theta + 2f'''(\theta) f'(\theta) \sin^2 \theta$$

$$+ [f'(\theta)]^2 \sin^2 \theta + [f'''(\theta)]^2 \cos^2 \theta$$

$$+ 2f'''(\theta) f'(\theta) \cos^2 \theta + [f'(\theta)]^2 \cos^2 \theta$$

$$= [f'''(\theta)]^2 + 2f'''(\theta) f'(\theta) + [f'(\theta)]^2$$

$$\dots [\because \sin^2 \theta + \cos^2 \theta = 1]$$

$$= [f'''(\theta) + f'(\theta)]^2$$

$$\therefore \int \left[\left(\frac{du}{d\theta}\right)^2 + \left(\frac{dv}{d\theta}\right)^2 \right]^{1/2} d\theta = \int [f'''(\theta) + f'(\theta)] d\theta$$

$$= f''(\theta) + f(\theta) + c$$
28. Let $I = \int \frac{-1}{(x+1)\sqrt{3x^2+2x}} dx$

$$= \int \frac{2^x - 1}{(x+1)^2 \sqrt{3x^2+2x}} dx$$

$$= \int \frac{2^x - 1}{(x^2+2x+1)\sqrt{3x^2+2x}} dx$$

$$= \int \frac{1 - \frac{1}{2}}{\left(x + \frac{1}{2}\right) \sqrt{\frac{1}{4} + \frac{1}{4} + 1}} dx$$

 Put $x + \frac{1}{2} + 1 = t^2$

$$\therefore \left(1 - \frac{1}{2}\right) dx = 2t dt$$

$$\therefore I = \int \frac{2t}{(t^2+1)\sqrt{t^2}} dt = 2 \int \frac{1}{t^2+1} dt = 2 \tan^{-1} t + c$$

$$= 2 \tan^{-1} \left(\sqrt{\frac{1}{4} + \frac{1}{4} + 1} \right) + c$$



$$\begin{aligned} 29. \quad \text{Let } I &= \int (\sqrt{\tan x} + \sqrt{\cot x}) dx \\ &= \int \left(\sqrt{\frac{\sin x}{\cos x}} + \sqrt{\frac{\cos x}{\sin x}} \right) dx \\ &= \int \frac{\sin x + \cos x}{\sqrt{\sin x \cos x}} dx \\ &= \int \frac{\sqrt{2}(\sin x + \cos x)}{\sqrt{2\sin x \cos x}} dx \\ &= \sqrt{2} \int \frac{\sin x + \cos x}{\sqrt{1 - 1 + 2\sin x \cos x}} dx \\ &= \sqrt{2} \int \frac{\sin x + \cos x}{\sqrt{1 - (1 - 2\sin x \cos x)}} dx \\ &= \sqrt{2} \int \frac{\sin x + \cos x}{\sqrt{1 - (\sin^2 x - 2\sin x \cos x + \cos^2 x)}} dx \\ &= \sqrt{2} \int \frac{1}{\sqrt{1 - (\sin x - \cos x)^2}} (\sin x + \cos x) dx \end{aligned}$$

Put $\sin x - \cos x = t$

$$\therefore (\cos x + \sin x) dx = dt$$

$$\therefore I = \sqrt{2} \int \frac{1}{\sqrt{1 - t^2}} dt$$

$$= \sqrt{2} \sin^{-1}(t) + c$$

$$= \sqrt{2} \sin^{-1}(\sin x - \cos x) + c$$

05 Definite Integrals



Hints



Classical Thinking

- $$\int_1^e \frac{1}{x} dx \quad [\log x]_1^e = \log_e e - \log 1 = 1$$
- $$\int_1^3 (-1)(-2)(-3) dx$$

$$= \int_1^3 (-x^2 + 11x - 6) dx$$

$$= \left[-\frac{x^3}{3} + \frac{11x^2}{2} - 6x \right]_1^3 = 0$$
- $$\int_0^1 (1-x)^9 dx = \left[-\frac{(1-x)^{10}}{10} \right]_0^1 = \frac{1}{10}$$
- $$\int_0^1 e^{2 \log x} dx = \int_0^1 e^{\log x^2} dx$$

$$= \int_0^1 x^2 dx = \left[\frac{x^3}{3} \right]_0^1 = \frac{1}{3}$$
- $$\int_0^{\pi/3} \cos 3x dx = \left[\frac{\sin 3x}{3} \right]_0^{\pi/3} = 0$$
- $$\int_{\pi/4}^{\pi/2} \operatorname{cosec}^2 x dx = [-\cot x]_{\pi/4}^{\pi/2} = -\left[\cot \frac{\pi}{2} - \cot \frac{\pi}{4} \right] = 1$$
- $$\int_0^{2\pi} (\sin x + \cos x) dx = [-\cos x + \sin x]_0^{2\pi} = 0$$
- $$\frac{1}{2} \int_0^{\pi/8} \sec^2 2x dx = \frac{1}{4} [\tan 2x]_0^{\pi/8} = \frac{1}{4} (1) = \frac{1}{4}$$
- $$\int_{\pi/6}^{\pi/4} \operatorname{cosec} 2x dx = \frac{1}{2} [\log \tan x]_{\pi/6}^{\pi/4}$$

$$= \frac{1}{2} \left[\log \tan \frac{\pi}{4} - \log \tan \frac{\pi}{6} \right] = \frac{1}{2} \log \sqrt{3}$$
- Put $1 + \log x = t \Rightarrow \frac{1}{x} dx = dt$

When $x = 1, t = 1$ and when $x = e, t = 2$

$$\therefore \int_1^e \frac{1 + \log x}{x} dx = \int_1^2 t dt = \left[\frac{t^2}{2} \right]_1^2 = \frac{3}{2}$$

- Put $t = \frac{1}{x} \Rightarrow dt = -\frac{1}{x^2} dx$

When $x = 1, t = 1$ and when $x = 2, t = \frac{1}{2}$

$$\therefore \int_1^2 \frac{1}{x^2} e^{-\frac{1}{x}} dx = \int_{\frac{1}{2}}^1 e^t dt = [e^t]_{\frac{1}{2}}^1$$

$$= e^{\frac{1}{2}} - e^{-1} = \frac{\sqrt{e} - 1}{e}$$
- Put $\log x = t \Rightarrow \frac{1}{x} dx = dt$

When $x = 1, t = 0$ and when $x = 2, t = \log 2$

$$\therefore \int_1^2 \frac{\cos(\log x)}{x} dx = \int_0^{\log 2} \cos t dt$$

$$= [\sin t]_0^{\log 2} = \sin(\log 2)$$
- $$\int_1^{\sqrt{3}} \frac{1}{1+x^2} dx = [\tan^{-1} x]_1^{\sqrt{3}} = \frac{\pi}{3} - \frac{\pi}{4} = \frac{\pi}{12}$$
- Put $\tan^{-1} x = t \Rightarrow \frac{1}{1+x^2} dx = dt$

When $x = 0, t = 0$ and when $x = 1, t = \frac{\pi}{4}$

$$\therefore \int_0^1 \frac{\tan^{-1} x}{1+x^2} dx = \int_0^{\pi/4} t dt = \left[\frac{t^2}{2} \right]_0^{\pi/4} = \frac{\pi^2}{32}$$
- $$\int_0^1 \frac{dx}{x^2 - 2x + 2} = \int_0^1 \frac{dx}{(x-1)^2 + 1}$$

$$= [\tan^{-1}(x-1)]_0^1$$

$$= 0 - \left(-\frac{\pi}{4} \right) = \frac{\pi}{4}$$
- $$\int_1^2 \log x dx = [\log x - x]_1^2$$

$$= 2 \log 2 - 2 + 1$$

$$= \log 4 - 1 = \log 4 - \log e = \log \left(\frac{4}{e} \right)$$
- $$\int_1^2 e \left(\frac{1}{x} - \frac{1}{2} \right) dx = \left[\frac{1}{x} e \right]_1^2 = \frac{e^2}{2} - e$$



$$\begin{aligned}
 18. \quad \int_2^3 \frac{d}{2-d} &= \int_2^3 \frac{d}{(-1)} = \int_2^3 \left[\frac{1}{-1} - \frac{1}{-1} \right] d \\
 &= \int_2^3 \frac{1}{(-1)} d - \int_2^3 \frac{1}{-1} d \\
 &= [\log(-1)]_2^3 - [\log(-1)]_2^3 \\
 &= (\log 2 - \log 1) - (\log 3 - \log 2) = 2\log 2 - \log 3 \\
 &= \log \left(\frac{4}{3} \right)
 \end{aligned}$$

19. Put $x = a - t \Rightarrow dx = -dt$
 When $x = 0$, $t = a$ and when $x = a$, $t = 0$

$$\begin{aligned}
 \therefore \int_0^a f(x) dx &= - \int_a^0 f(a-t) dt \\
 &= \int_0^a f(a-t) dt \quad \dots \left[\int_a^b f(x) dx = - \int_b^a f(x) dx \right] \\
 &= \int_0^a f(a-x) dx \quad \dots \left[\int_a^b f(x) dx = \int_a^b f(t) dt \right]
 \end{aligned}$$

$$\begin{aligned}
 20. \quad \text{Let } I &= \int_0^{\frac{\pi}{2}} \frac{\sin x}{\sin x + \cos x} dx \quad \dots(i) \\
 &= \int_0^{\frac{\pi}{2}} \frac{\sin\left(\frac{\pi}{2}-x\right)}{\sin\left(\frac{\pi}{2}-x\right) + \cos\left(\frac{\pi}{2}-x\right)} dx \\
 &\quad \dots \left[\int_0^a f(x) dx = \int_0^a f(a-x) dx \right] \\
 &= \int_0^{\frac{\pi}{2}} \frac{\cos x}{\cos x + \sin x} dx \quad \dots(ii)
 \end{aligned}$$

Adding (i) and (ii), we get

$$\begin{aligned}
 2I &= \int_0^{\frac{\pi}{2}} dx = [x]_0^{\frac{\pi}{2}} \\
 \therefore 2I &= \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4}
 \end{aligned}$$

$$21. \quad \text{Let } I = \int_0^{\pi/2} \frac{\sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \quad \dots(i)$$

$$\begin{aligned}
 \therefore I &= \int_0^{\pi/2} \frac{\sqrt{\cos\left(\frac{\pi}{2}-x\right)}}{\sqrt{\sin\left(\frac{\pi}{2}-x\right)} + \sqrt{\cos\left(\frac{\pi}{2}-x\right)}} dx \\
 &\quad \dots \left[\int_0^a f(x) dx = \int_0^a f(a-x) dx \right]
 \end{aligned}$$

$$\therefore I = \int_0^{\pi/2} \frac{\sqrt{\sin x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_0^{\pi/2} dx = \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4}$$

$$22. \quad \text{Let } I = \int_0^{\pi} \sin x dx \quad \dots(i)$$

$$\begin{aligned}
 \therefore I &= \int_0^{\pi} (\pi-x) \sin(\pi-x) dx \\
 &\quad \dots \left[\int_0^a f(x) dx = \int_0^a f(a-x) dx \right]
 \end{aligned}$$

$$\therefore I = \int_0^{\pi} (\pi-x) \sin x dx \quad \dots(ii)$$

Adding (i) and (ii), we get

$$\begin{aligned}
 2I &= \int_0^{\pi} \sin x dx = \pi [-\cos x]_0^{\pi} = 2\pi \\
 \Rightarrow I &= \pi
 \end{aligned}$$

$$23. \quad \text{Let } I = \int_2^3 \frac{\sqrt{x}}{\sqrt{5-x} + \sqrt{x}} dx \quad \dots(i)$$

$$\begin{aligned}
 \therefore I &= \int_2^3 \frac{\sqrt{5-x}}{\sqrt{x} + \sqrt{5-x}} dx \quad \dots(ii) \\
 &\quad \dots \left[\int_a^b f(x) dx = \int_a^b f(a+b-x) dx \right]
 \end{aligned}$$

Adding (i) and (ii), we get

$$\begin{aligned}
 2I &= \int_2^3 dx = [x]_2^3 = 3 - 2 = 1 \\
 \Rightarrow I &= \frac{1}{2}
 \end{aligned}$$

$$\begin{aligned}
 24. \quad \text{Let } f(x) &= x^{17} \cos^4 x \\
 \therefore f(-x) &= (-x)^{17} \{\cos(-x)\}^4 = -f(x) \\
 \therefore f(x) &\text{ is an odd function.}
 \end{aligned}$$

$$\therefore \int_{-1}^1 x^{17} \cos^4 x dx = 0$$

$$25. \quad \text{Since, } \sin^{11} x \text{ is an odd function.}$$

$$\therefore \int_{-1}^1 \sin^{11} x dx = 0$$

$$26. \quad \text{Since, } 3 \sin x + \sin^3 x \text{ is an odd function.}$$

$$\therefore \int_{-\pi/2}^{\pi/2} (3 \sin x + \sin^3 x) dx = 0$$



Critical Thinking

1. $\int_{-2}^2 (a^3 + b + c)d = \left[\frac{a^4}{4} + \frac{b^2}{2} + c \right]_{-2}^2 = 4c$

Hence, the value depends on c.

2.
$$\int_0^1 \frac{d}{\sqrt{1+d} - \sqrt{d}}$$

$$= \int_0^1 \frac{(\sqrt{1+d} + \sqrt{d})}{(\sqrt{1+d} - \sqrt{d})(\sqrt{1+d} + \sqrt{d})} d$$

$$= \int_0^1 \frac{(\sqrt{1+d} + \sqrt{d})}{1+d-d} d = \int_0^1 \sqrt{1+d} d + \int_0^1 \sqrt{d} d$$

$$= \left[\frac{2}{3}(1+d)^{\frac{3}{2}} \right]_0^1 + \left[\frac{2}{3}(d)^{\frac{3}{2}} \right]_0^1$$

$$= \frac{4\sqrt{2}}{3} - \frac{2}{3} + \frac{2}{3} - 0 = \frac{4\sqrt{2}}{3}$$

3. $I + J = \int_0^{\pi/4} (\sin^2 + \cos^2) d = \int_0^{\pi/4} d = \frac{\pi}{4}$

$\therefore I = \frac{\pi}{4} - J$

4.
$$\int_0^{\pi/4} \tan^2 d = \int_0^{\pi/4} (\sec^2 - 1) d$$

$$= [\tan d]_0^{\pi/4} - [d]_0^{\pi/4}$$

$$= 1 - \frac{\pi}{4}$$

5.
$$\int_0^{\pi} \frac{d}{1 + \sin d} = \int_0^{\pi} \frac{1 - \sin d}{\cos^2 d} d$$

$$= \int_0^{\pi} (\sec^2 d - \sec d \tan d) d$$

$$= [\tan d - \sec d]_0^{\pi} = \tan \pi - \sec \pi + 1$$

$$= 0 + 1 + 1 = 2$$

6.
$$\int_{\pi/4}^{3\pi/4} \frac{d}{1 + \cos d}$$

$$= \int_{\pi/4}^{3\pi/4} \frac{1 - \cos d}{1 - \cos^2 d} d = \int_{\pi/4}^{3\pi/4} \frac{1 - \cos d}{\sin^2 d} d$$

$$= \int_{\pi/4}^{3\pi/4} (\operatorname{cosec}^2 d - \cot d \operatorname{cosec} d) d$$

$$= [-\cot d + \operatorname{cosec} d]_{\pi/4}^{3\pi/4} = 2$$

7.
$$I = \int_0^{\pi/2} \frac{(\sin d + \cos d)^2}{\sqrt{1 + \sin 2d}} d$$

$$= \int_0^{\pi/2} \frac{(\sin d + \cos d)^2}{\sqrt{(\sin d + \cos d)^2}} d$$

$$= \int_0^{\pi/2} (\sin d + \cos d) d = [-\cos d + \sin d]_0^{\pi/2}$$

$$= 2$$

8.
$$\int_0^{2\pi} \sqrt{1 + \sin \frac{d}{2}} d = \int_0^{2\pi} \sqrt{\left(\sin \frac{d}{4} + \cos \frac{d}{4}\right)^2} d$$

$$= \int_0^{2\pi} \left(\sin \frac{d}{4} + \cos \frac{d}{4}\right) d$$

.... $\left[\because d \in (0, 2\pi), \therefore \left(\sin \frac{d}{4} + \cos \frac{d}{4}\right) > 0 \right]$

$$= 4 \left[-\cos \frac{d}{4} + \sin \frac{d}{4} \right]_0^{2\pi}$$

$$= 4(0 + 1 + 1 - 0)$$

$$= 8$$

9.
$$\int_{-1}^3 \left\{ \tan^{-1} \left(\frac{d}{d^2 + 1} \right) + \tan^{-1} \left(\frac{d^2 + 1}{d} \right) \right\} d$$

$$= \int_{-1}^3 \left\{ \tan^{-1} \left(\frac{d}{d^2 + 1} \right) + \cot^{-1} \left(\frac{d}{d^2 + 1} \right) \right\} d$$

$$= \int_{-1}^3 \frac{\pi}{2} d \quad \text{....} \left[\because \tan^{-1} x + \cot^{-1} x = \frac{\pi}{2} \right]$$

$$= \frac{\pi}{2} [d^2]_{-1}^3 = 2\pi$$

10. Put $\tan d = t \Rightarrow \sec^2 d \cdot dd = dt$
 When $d = 0, t = 0$ and when $d = \frac{\pi}{4}, t = 1$
 $\therefore \int_0^{\pi/4} \tan^6 d \sec^2 d = \int_0^1 t^6 dt = \frac{1}{7} [t^7]_0^1 = \frac{1}{7}$

11. Let $I = \int_{\pi/4}^{\pi/2} \cos \theta \frac{1}{\sin^2 \theta} d\theta$
 Put $\sin \theta = t \Rightarrow \cos \theta d\theta = dt$
 When $\theta = \frac{\pi}{4}, t = \frac{1}{\sqrt{2}}$ and when $\theta = \frac{\pi}{2}, t = 1$
 $\therefore I = \int_{1/\sqrt{2}}^1 \frac{1}{t^2} dt = \left[\frac{-1}{t} \right]_{1/\sqrt{2}}^1$

$$= \sqrt{2} - 1$$



$$12. \text{ Let } I = \int_0^{\pi/6} \frac{\sin}{\cos^3} d = \int_0^{\pi/6} \tan \sec^2 d$$

$$\text{Put } \tan = t \Rightarrow \sec^2 d = dt$$

$$\text{When } = 0, t = 0 \text{ and when } = \frac{\pi}{6}, t = \frac{1}{\sqrt{3}}$$

$$\therefore I = \int_0^{\frac{1}{\sqrt{3}}} t dt = \left[\frac{t^2}{2} \right]_0^{\frac{1}{\sqrt{3}}} = \frac{1}{6}$$

$$13. \text{ Let } I = \int_0^{\pi/4} \sec^7 \theta \cdot \sin^3 \theta d\theta = \int_0^{\pi/4} \tan^3 \theta \sec^4 \theta d\theta$$

$$\text{Put } \tan \theta = t \Rightarrow \sec^2 \theta d\theta = dt$$

$$\text{When } \theta = 0, t = 0 \text{ and when } \theta = \frac{\pi}{4}, t = 1$$

$$\therefore I = \int_0^1 t^3 (1+t^2) dt = \left[\frac{t^4}{4} + \frac{t^6}{6} \right]_0^1 = \frac{5}{12}$$

$$14. \text{ Put } t^3 = t \Rightarrow t^2 d = \frac{dt}{3}$$

$$\text{When } = 0, t = 0 \text{ and when } = a, t = a^3$$

$$\therefore \int_0^a t^2 \sin^{-3} d = \frac{1}{3} \int_0^{a^3} \sin t dt = -\frac{1}{3} [\cos t]_0^{a^3}$$

$$= -\frac{1}{3} (\cos a^3 - 1) = \frac{1}{3} (1 - \cos a^3)$$

$$15. \text{ Put } \sqrt{t} = t \Rightarrow \frac{1}{\sqrt{t}} d = 2 dt$$

$$\text{When } = 0, t = 0 \text{ and when } = 2, t = \sqrt{2}$$

$$\therefore \int_0^2 \frac{3^{\sqrt{t}}}{\sqrt{t}} d = 2 \int_0^{\sqrt{2}} 3^t dt = 2 \left[\frac{3^t}{\log 3} \right]_0^{\sqrt{2}} = \frac{2}{\log 3} (3^{\sqrt{2}} - 1)$$

$$16. \text{ Let } I = \int_0^{\pi/2} \sin \sin 2 d = 2 \int_0^{\pi/2} \sin^2 \cos d$$

$$\text{Put } \sin = t \Rightarrow \cos d = dt$$

$$\text{When } = 0, t = 0 \text{ and when } = \frac{\pi}{2}, t = 1$$

$$\therefore I = 2 \int_0^1 t^2 dt = \frac{2}{3} [t^3]_0^1 = \frac{2}{3}$$

$$17. \text{ Let } I = \int_1^{e^2} \frac{d}{(1 + \log)^2}$$

$$\text{Put } (1 + \log) = t \Rightarrow \frac{1}{t} d = dt$$

$$\text{When } = 1, t = 1 \text{ and when } = e^2, t = 3$$

$$\therefore I = \int_1^3 \frac{dt}{t^2} = \left[\frac{-1}{t} \right]_1^3 = -\left(\frac{1}{3} - 1 \right) = \frac{2}{3}$$

$$18. \text{ Let } I = \int_0^{\pi/4} \sec \log(\sec + \tan) d$$

$$\text{Put } \log(\sec + \tan) = t \Rightarrow \sec d = dt$$

$$\therefore I = \int_0^{\log(\sqrt{2}+1)} t dt = \left[\frac{t^2}{2} \right]_0^{\log(\sqrt{2}+1)} = \frac{[\log(\sqrt{2}+1)]^2}{2}$$

$$19. \int_0^{-\pi/4} \frac{1 + \tan}{1 - \tan} d = \int_0^{-\pi/4} \tan\left(\frac{\pi}{4} + \right) d$$

$$= \left[\log \left\{ \sec\left(\frac{\pi}{4} + \right) \right\} \right]_0^{-\pi/4}$$

$$= -\frac{1}{2} \log 2$$

$$20. \text{ Since, } \sin \theta \text{ is positive in interval } (0, \pi).$$

$$\therefore \int_0^{\pi} |\sin^3 \theta| d\theta = \int_0^{\pi} \sin^3 \theta d\theta = \int_0^{\pi} \sin \theta (1 - \cos^2 \theta) d\theta$$

$$= \int_0^{\pi} \sin \theta d\theta + \int_0^{\pi} (-\sin \theta) \cos^2 \theta d\theta$$

$$= [-\cos \theta]_0^{\pi} + \left[\frac{\cos^3 \theta}{3} \right]_0^{\pi} = \frac{4}{3}$$

$$21. \text{ Let } I = \int_0^{\pi/8} \cos^3 4\theta d\theta = \int_0^{\pi/8} \cos^2 4\theta \cdot \cos 4\theta d\theta$$

$$= \int_0^{\pi/8} (1 - \sin^2 4\theta) \cos 4\theta d\theta$$

$$\text{Put } \sin 4\theta = t \Rightarrow \cos 4\theta d\theta = \frac{dt}{4}$$

$$\text{When } \theta = 0, t = 0 \text{ and when } \theta = \frac{\pi}{8}, t = 1$$

$$\therefore I = \frac{1}{4} \int_0^1 (1 - t^2) dt = \frac{1}{4} \left[t - \frac{t^3}{3} \right]_0^1 = \frac{1}{6}$$

$$22. \text{ Let } I = \int_{\pi/3}^{\pi/2} \frac{\sqrt{1 + \cos}}{(1 - \cos)^{5/2}} d$$

$$= \int_{\pi/3}^{\pi/2} \frac{\sqrt{1 + \cos}}{(1 - \cos)^{5/2}} \times \frac{\sqrt{1 - \cos}}{\sqrt{1 - \cos}} d$$

$$= \int_{\pi/3}^{\pi/2} \frac{\sin}{(1 - \cos)^3} d$$

$$\text{Put } 1 - \cos = t$$

$$\Rightarrow \sin d = dt$$

$$\therefore I = \int_{1/2}^1 \frac{dt}{t^3} = \left[\frac{t^{-2}}{-2} \right]_{1/2}^1 = \frac{3}{2}$$



23. Put $\sin^2 = t \Rightarrow 2 \sin \cos d = dt$
 When $= 0, t = 0$ and when $= \frac{\pi}{2}, t = 1$

$$\therefore \int_0^{\pi/2} \frac{\sin \cos}{1 + \sin^4} d = \frac{1}{2} \int_0^1 \frac{1}{1+t^2} dt$$

$$= \frac{1}{2} [\tan^{-1}t]_0^1 = \frac{\pi}{8}$$

24. Let $I = \int_0^{\pi/4} \frac{4\sin 2\theta d\theta}{\sin^4\theta + \cos^4\theta} = 4 \int_0^{\pi/4} \frac{2\sin\theta\cos\theta d\theta}{\sin^4\theta + \cos^4\theta}$

$$= 4 \int_0^{\pi/4} \frac{2\tan\theta\sec^2\theta d\theta}{\tan^4\theta + 1}$$

Put $\tan^2\theta = t \Rightarrow 2 \tan\theta \sec^2\theta d\theta = dt$

$$\therefore I = 4 \int_0^1 \frac{dt}{t^2+1} = 4 [\tan^{-1}t]_0^1 = 4 \left[\frac{\pi}{4} - 0 \right] = \pi$$

25. $k \int_0^1 f(3^x) dx = \int_0^3 tf(t) dt \quad \dots(i)$

Put $3^x = t \Rightarrow dx = \frac{dt}{3}$

$$\therefore k \int_0^1 f(3^x) dx = k \int_{\frac{1}{3}}^3 \frac{t}{3} \cdot f(t) \cdot \frac{dt}{3} = \frac{k}{9} \int_0^3 tf(t) dt$$

From (i),

$$\frac{k}{9} \int_0^3 tf(t) dt = \int_0^3 tf(t) dt$$

$$\Rightarrow \frac{k}{9} = 1 \Rightarrow k = 9$$

26. $\int_0^{2/3} \frac{d}{4+9d^2} = \frac{1}{9} \int_0^{2/3} \frac{d}{\left(\frac{2}{3}\right)^2 + d^2}$

$$= \frac{1}{9} \times \frac{1}{\frac{2}{3}} \left[\tan^{-1} \frac{d}{\frac{2}{3}} \right]_0^{2/3}$$

$$= \frac{1}{6} \times \frac{\pi}{4} = \frac{\pi}{24}$$

27. $\int_0^k \frac{d}{2+8d^2} = \frac{\pi}{16}$

$$\Rightarrow \frac{1}{2} \int_0^k \frac{1}{1^2 + (2d)^2} d = \frac{\pi}{16}$$

$$\Rightarrow \frac{1}{2} \left[\frac{\tan^{-1}(2d)}{2} \right]_0^k = \frac{\pi}{16}$$

$$\Rightarrow \frac{1}{4} (\tan^{-1} 2k) = \frac{\pi}{16}$$

$$\Rightarrow \tan^{-1} 2k = \frac{\pi}{4} \Rightarrow k = \frac{1}{2}$$

28. $\int_0^1 \frac{d}{d^2 - d + 1} = \int_0^1 \frac{d}{\left(-\frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2}$

$$= \frac{2}{\sqrt{3}} \left[\tan^{-1} \left(\frac{2d-1}{\sqrt{3}} \right) \right]_0^1$$

$$= \frac{2}{\sqrt{3}} \left[\tan^{-1} \left(\frac{1}{\sqrt{3}} \right) - \tan^{-1} \left(-\frac{1}{\sqrt{3}} \right) \right]$$

$$= \frac{2}{\sqrt{3}} \left[\frac{\pi}{6} - \left(-\frac{\pi}{6} \right) \right] = \frac{2}{\sqrt{3}} \cdot \frac{\pi}{3} = \frac{2\pi}{3\sqrt{3}}$$

29. Let $I = \int_0^1 \frac{d}{e^d + e^{-d}} = \int_0^1 \frac{e^d}{1+e^{2d}} d$

Put $e^d = t \Rightarrow e^d d = dt$

$$\therefore I = \int_1^e \frac{dt}{1+t^2} = [\tan^{-1}t]_1^e = \tan^{-1} e - \tan^{-1} 1$$

$$= \tan^{-1} \left(\frac{e-1}{e+1} \right)$$

$$\dots \left[\because \tan^{-1} x - \tan^{-1} y = \tan^{-1} \left(\frac{x-y}{1+xy} \right) \right]$$

30. $\int_{1/4}^{1/2} \frac{d}{\sqrt{d^2 - 2}} = \int_{1/4}^{1/2} \frac{d}{\sqrt{\left(\frac{1}{2}\right)^2 - \left(-\frac{1}{2}\right)^2}}$

$$= \left[\sin^{-1} \left(\frac{-\frac{1}{2}}{\frac{1}{2}} \right) \right]_{1/4}^{1/2}$$

$$= [\sin^{-1}(2d-1)]_{1/4}^{1/2} = \frac{\pi}{6}$$

31. $\int_3^5 \frac{d^2}{d^2-4} d = \int_3^5 \left(1 + \frac{4}{d^2-4} \right) d$

$$= \left[d + \frac{4}{2(2)} \log \left| \frac{d-2}{d+2} \right| \right]_3^5$$

$$= 2 + \log_e \left(\frac{15}{7} \right)$$



$$32. \text{ Let } I = \int_0^1 \frac{1}{[a + b(1 - t)]^2} dt$$

$$= \int_0^1 \frac{1}{[(a-b) + b]^2} dt$$

Put $(a-b) + b = t \Rightarrow (a-b) dt = dt$
 When $t = 0, t = b$ and when $t = 1, t = a$

$$\therefore I = \frac{1}{a-b} \int_b^a \frac{1}{t^2} dt$$

$$= \frac{1}{(a-b)} \left[-\frac{1}{t} \right]_b^a$$

$$= \frac{1}{(a-b)} \left(\frac{a-b}{ab} \right)$$

$$\therefore I = \frac{1}{ab}$$

$$33. \text{ Put } a^2 + t^2 = t \Rightarrow 2t dt = dt$$

When $t = 0, t = a^2$ and when $t = a, t = 2a^2$

$$\therefore \int_0^a \frac{dt}{\sqrt{a^2 + t^2}} = \frac{1}{2} \int_{a^2}^{2a^2} \frac{1}{\sqrt{t}} dt$$

$$= \left[\sqrt{t} \right]_{a^2}^{2a^2} = (2a^2)^{\frac{1}{2}} - (a^2)^{\frac{1}{2}} = a(\sqrt{2} - 1)$$

$$34. \text{ Put } 1 + e^{-t} = t \Rightarrow -e^{-t} dt = dt$$

When $t = 0, t = 2$ and when $t = 1, t = 1 + \frac{1}{e}$

$$\therefore \int_0^1 \frac{e^{-t}}{1 + e^{-t}} dt = \int_2^{1+\frac{1}{e}} \frac{(t-1)(-dt)}{t} = \int_2^{1+\frac{1}{e}} \left(\frac{1}{t} - 1 \right) dt$$

$$= \left[\log t - t \right]_2^{1+\frac{1}{e}}$$

$$= \log \left(1 + \frac{1}{e} \right) - \left(1 + \frac{1}{e} \right) - \log 2 + 2$$

$$= \log \left(\frac{e+1}{2e} \right) - \frac{1}{e} + 1$$

$$35. \text{ Let } I = \int_0^{\pi/2} \frac{d}{a^2 \cos^2 t + b^2 \sin^2 t}$$

Dividing N^r and D^r by $\cos^2 t$, we get

$$I = \int_0^{\pi/2} \frac{\sec^2 t}{a^2 + b^2 \tan^2 t} dt$$

Put $b \tan t = t \Rightarrow b \sec^2 t dt = dt$

When $t = 0, t = 0$ and when $t = \frac{\pi}{2}, t = \infty$

$$\therefore I = \int_0^{\infty} \frac{dt}{a^2 + t^2} = \frac{1}{b} \left[\frac{1}{a} \tan^{-1} \left(\frac{t}{a} \right) \right]_0^{\infty}$$

$$= \frac{1}{ab} (\tan^{-1} \infty - \tan^{-1} 0)$$

$$= \frac{1}{ab} \left(\frac{\pi}{2} - 0 \right) = \frac{\pi}{2ab}$$

$$36. \text{ Put } e^{-1} - 1 = t^2 \Rightarrow e^{-t} dt = 2t dt$$

When $t = 0, t = 0$ and when $t = \log 5, t = 2$

$$\therefore \int_0^{\log 5} \frac{e^{-t} \sqrt{e^{-t} - 1}}{e^{-t} + 3} dt = \int_0^2 \frac{2t^2}{t^2 + 4} dt$$

$$= 2 \int_0^2 \left(1 - \frac{4}{t^2 + 4} \right) dt$$

$$= 2 \left[t - 4 \cdot \frac{1}{2} \tan^{-1} \frac{t}{2} \right]_0^2$$

$$= 2 \left(2 - 2 \cdot \frac{\pi}{4} \right) = 4 - \pi$$

$$37. \text{ Put } t = 2 \cos \theta \Rightarrow dt = -2 \sin \theta d\theta$$

$$\therefore \int_0^2 \sqrt{\frac{2+t}{2-t}} dt = -2 \int_{\pi/2}^0 \sqrt{\frac{1+\cos \theta}{1-\cos \theta}} \sin \theta d\theta$$

$$= -4 \int_{\pi/2}^0 \frac{\cos \left(\frac{\theta}{2} \right)}{\sin \left(\frac{\theta}{2} \right)} \sin \frac{\theta}{2} \cos \frac{\theta}{2} d\theta$$

$$= -2 \int_{\pi/2}^0 (1 + \cos \theta) d\theta$$

$$= -2 \left[\theta + \sin \theta \right]_{\pi/2}^0$$

$$= 2 \left(\frac{\pi}{2} + 1 \right) = \pi + 2$$

$$38. \text{ Since, } \int_a^b \sqrt{\frac{-a}{b-t}} dt = \frac{\pi}{2} (b-a)$$

$$\therefore \int_3^4 \sqrt{\frac{-3}{4-t}} dt = \frac{\pi}{2} (4-3) = \frac{\pi}{2}$$

$$39. \text{ Since, } \int_a^b \sqrt{(b-a)(b-t)} dt = \frac{\pi}{8} (b-a)^2$$

$$\therefore \int_3^7 \sqrt{(7-3)(7-t)} dt = \frac{\pi}{8} (7-3)^2$$

$$= \frac{\pi}{8} \times 16 = 2\pi$$



40. Put $x^2 + 1 = t \Rightarrow 2x dx = dt$
 When $x = 0, t = 1$ and when $x = 2, t = 5$

$$\begin{aligned} \therefore \int_0^2 \frac{x^3}{(x^2+1)^{\frac{3}{2}}} dx &= \frac{1}{2} \int_1^5 \frac{(t-1)}{t^{\frac{3}{2}}} dt \\ &= \frac{1}{2} \int_1^5 \left(t^{-\frac{1}{2}} - t^{-\frac{3}{2}} \right) dt \\ &= \frac{1}{2} \left[2\sqrt{t} + 2\frac{1}{\sqrt{t}} \right]_1^5 \\ &= \frac{1}{2} \left[2\sqrt{5} + \frac{2}{\sqrt{5}} - 2 - 2 \right] \\ &= \sqrt{5} + \frac{1}{\sqrt{5}} - 2 = \frac{6-2\sqrt{5}}{\sqrt{5}} \end{aligned}$$

41. Put $x = a \tan \theta \Rightarrow dx = a \sec^2 \theta d\theta$

$$\begin{aligned} \therefore \int_0^a \frac{x^4}{(a^2+x^2)^4} dx &= \int_0^{\frac{\pi}{4}} \frac{a^4 \tan^4 \theta \cdot a \sec^2 \theta}{a^8 \sec^8 \theta} d\theta \\ &= \frac{1}{a^3} \int_0^{\frac{\pi}{4}} \sin^4 \theta \cos^2 \theta d\theta \\ &= \frac{1}{a^3} \int_0^{\frac{\pi}{4}} (\sin^4 \theta - \sin^6 \theta) d\theta \\ &= \frac{1}{a^3} \int_0^{\frac{\pi}{4}} \left[\frac{(1-\cos 2\theta)^2}{4} - \frac{(1-\cos 2\theta)^3}{8} \right] d\theta \\ &= \frac{1}{8a^3} \int_0^{\frac{\pi}{4}} (1+\cos 2\theta)(1-\cos 2\theta)^2 d\theta \\ &= \frac{1}{8a^3} \int_0^{\frac{\pi}{4}} (1-\cos 2\theta - \cos^2 2\theta + \cos^3 2\theta) d\theta \\ &= \frac{1}{8a^3} \int_0^{\frac{\pi}{4}} \frac{1}{4} [2 - \cos 2\theta - 2\cos 4\theta + \cos 6\theta] d\theta \\ &\quad \dots \left[\begin{aligned} \because \cos^2 A &= \frac{1+\cos 2A}{2} \\ \text{and } \cos^3 A &= \frac{\cos 3A + 3\cos A}{4} \end{aligned} \right] \\ &= \frac{1}{32a^3} \left[2\theta - \frac{\sin 2\theta}{2} - \frac{\sin 4\theta}{2} + \frac{\sin 6\theta}{6} \right]_0^{\frac{\pi}{4}} \\ &= \frac{1}{16a^3} \left(\frac{\pi}{4} - \frac{1}{3} \right) \end{aligned}$$

$$\begin{aligned} 42. \int_0^{\pi} \frac{dx}{1-2a \cos x + a^2} &= \int_0^{\pi} \frac{dx}{(1+a^2) \left(\cos^2 \frac{x}{2} + \sin^2 \frac{x}{2} \right) - 2a \left(\cos^2 \frac{x}{2} - \sin^2 \frac{x}{2} \right)} \\ &= \int_0^{\pi} \frac{dx}{(1-a)^2 \cos^2 \frac{x}{2} + (1+a)^2 \sin^2 \frac{x}{2}} \\ &= \int_0^{\pi} \frac{\sec^2 \frac{x}{2}}{(1-a)^2 + (1+a)^2 \tan^2 \frac{x}{2}} dx \\ &= \frac{2}{(1+a)^2} \int_0^{\infty} \frac{dt}{\left\{ \frac{(1-a)}{(1+a)} \right\}^2 + t^2} \\ &\quad \dots \left[\text{Put } t = \tan \frac{x}{2} \Rightarrow dt = \frac{1}{2} \sec^2 \frac{x}{2} dx \right] \\ &= \frac{2}{(1+a)^2} \cdot \frac{(1+a)}{(1-a)} \left[\tan^{-1} \left(\frac{1+a}{1-a} \cdot t \right) \right]_0^{\infty} \\ &= \frac{2}{(1-a)^2} (\tan^{-1} \infty - \tan^{-1} 0) = \frac{\pi}{1-a^2} \end{aligned}$$

$$\begin{aligned} 43. \int_0^1 x^2 e^x dx &= [x^2 \cdot e^x]_0^1 - \int_0^1 2x e^x dx \\ &= e - 2 [x e^x - e^x]_0^1 \\ &= e - 2 [e - e - (0 - 1)] = e - 2 \end{aligned}$$

$$\begin{aligned} 44. \text{ Let } I &= \int_{-\frac{\pi}{4}}^{\frac{\pi}{2}} e^{-x} \sin x dx \\ \therefore I &= [-e^{-x} \sin x - e^{-x} \cos x]_{-\frac{\pi}{4}}^{\frac{\pi}{2}} - \int_{-\frac{\pi}{4}}^{\frac{\pi}{2}} e^{-x} \sin x dx \\ \Rightarrow 2I &= [e^{-x} (-\sin x - \cos x)]_{-\frac{\pi}{4}}^{\frac{\pi}{2}} \\ \Rightarrow I &= \frac{1}{2} \left[e^{-\frac{\pi}{2}} (-1-0) - \left\{ e^{\frac{\pi}{4}} \left(\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} \right) \right\} \right] \\ \Rightarrow I &= -\frac{1}{2} e^{-\frac{\pi}{2}} \end{aligned}$$

$$\begin{aligned} 45. \int_0^1 \tan^{-1} x dx &= [(\tan^{-1} x)]_0^1 - \int_0^1 \frac{1}{1+x^2} \cdot dx \\ &= \left[\tan^{-1} x - \frac{1}{2} \log |1+x^2| \right]_0^1 \\ &= \frac{\pi}{4} - \frac{1}{2} \log 2 \end{aligned}$$



$$46. \int_0^1 \cos^{-1} x \, dx = \left[\cos^{-1} x - \sqrt{1-x^2} \right]_0^1 = 1$$

$$47. \text{ Put } x = t^2 \Rightarrow dx = 2t \, dt$$

$$\text{When } x = 0, t = 0 \text{ and when } x = \frac{\pi^2}{4}, t = \frac{\pi}{2}$$

$$\begin{aligned} \therefore \int_0^{\frac{\pi^2}{4}} \sin \sqrt{x} \, dx &= 2 \int_0^{\frac{\pi}{2}} t \sin t \, dt \\ &= 2[-t \cos t + \sin t]_0^{\pi/2} = 2 \end{aligned}$$

$$48. \text{ Put } x = \tan \theta \Rightarrow dx = \sec^2 \theta \, d\theta$$

$$\text{When } x = 0, \theta = 0 \text{ and when } x = 1, \theta = \frac{\pi}{4}$$

$$\begin{aligned} \therefore \int_0^1 \sin^{-1} \left(\frac{2}{1+x^2} \right) dx &= \int_0^{\pi/4} \sin^{-1} \left(\frac{2 \tan \theta}{1 + \tan^2 \theta} \right) \sec^2 \theta \, d\theta \\ &= 2 \int_0^{\pi/4} \theta \sec^2 \theta \, d\theta \\ &= 2[\theta \tan \theta]_0^{\pi/4} - 2 \int_0^{\pi/4} \tan \theta \, d\theta \\ &= \frac{\pi}{2} + 2[\log \cos \theta]_0^{\pi/4} \\ &= \frac{\pi}{2} - 2 \log \sqrt{2} \end{aligned}$$

$$49. \int_0^{\frac{\pi}{2}} \frac{1 + \sin x}{1 + \cos x} dx = \int_0^{\frac{\pi}{2}} \left[\frac{1 + 2 \sin \frac{x}{2} \cdot \cos \frac{x}{2}}{2 \cos^2 \frac{x}{2}} \right] dx$$

$$= \frac{1}{2} \int_0^{\frac{\pi}{2}} \left[\sec^2 \frac{x}{2} + 2 \tan \frac{x}{2} \right] dx$$

$$= \frac{1}{2} \left[2 \tan \frac{x}{2} \right]_0^{\pi/2} - \int_0^{\frac{\pi}{2}} \tan \frac{x}{2} \, dx + \int_0^{\frac{\pi}{2}} \tan \frac{x}{2} \, dx$$

$$= \left[\tan \frac{x}{2} \right]_0^{\pi/2} = \frac{\pi}{2} \tan \frac{\pi}{4} = \frac{\pi}{2}$$

$$50. \int_0^{\frac{\pi}{6}} (2+3x^2) \cos 3x \, dx$$

$$= \left[(2+3x^2) \cdot \frac{\sin 3x}{3} \right]_0^{\pi/6} - \int_0^{\pi/6} 6x \cdot \frac{\sin 3x}{3} dx$$

$$= \frac{2}{3} + \frac{\pi^2}{36} + \left[\frac{2 \cos 3x}{3} \right]_0^{\pi/6} - 2 \int_0^{\pi/6} \cos 3x \, dx$$

$$= \frac{2}{3} + \frac{\pi^2}{36} + 0 - \frac{2}{9} [\sin 3x]_0^{\pi/6}$$

$$= \frac{2}{3} + \frac{\pi^2}{36} - \frac{2}{9} = \frac{1}{36} (\pi^2 + 16)$$

$$51. \int_2^e \left[\frac{1}{\log x} - \frac{1}{(\log x)^2} \right] dx$$

$$= \int_2^e \frac{1}{\log x} dx - \int_2^e \frac{1}{(\log x)^2} dx$$

$$= \left[\frac{x}{\log x} \right]_2^e - \int_2^e \left\{ -\frac{1}{(\log x)^2} \right\} dx - \int_2^e \frac{1}{(\log x)^2} dx$$

$$= \left[\frac{x}{\log x} \right]_2^e = e - \frac{2}{\log 2}$$

$$\therefore \alpha = e, \beta = -2$$

$$52. \int_1^e e^{-(1+\log x)} dx = \int_1^e e^{-1} e^{-\log x} dx$$

$$= [e^{-1} \cdot \log x]_1^e = e^{-1}$$

$$53. \int_{\pi/4}^{\pi/2} e^{(\log \sin x + \cot x)} dx$$

$$= [e^{\log \sin x}]_{\pi/4}^{\pi/2}$$

$$= e^{\frac{\pi}{2} \log \sin \frac{\pi}{2}} - e^{\frac{\pi}{4} \log \sin \frac{\pi}{4}} = \frac{1}{2} e^{\frac{\pi}{4} \log 2}$$

$$54. \int_0^1 \frac{e^{-(x-1)}}{(x+1)^3} dx = \int_0^1 \frac{e^{-(x+1-2)}}{(x+1)^3} dx$$

$$= \int_0^1 e^{-x} \left[\frac{1}{(x+1)^2} + \frac{-2}{(x+1)^3} \right] dx$$

$$= \left[\frac{e^{-x}}{(x+1)^2} \right]_0^1 = \frac{e^{-1}}{4} - 1$$

$$55. \phi(x) = \frac{1}{(x^4+1)} = \frac{1}{x^4} - \frac{3}{x^4+1}$$

$$\therefore \int_1^2 \phi(x) dx = \int_1^2 \left(\frac{1}{x^4} - \frac{3}{x^4+1} \right) dx$$

$$= [\log x]_1^2 - \left[\frac{1}{4} \log(x^4+1) \right]_1^2$$

$$= \frac{1}{4} \log \frac{32}{17}$$



56. Put $\sin \theta = t \Rightarrow \cos \theta d\theta = dt$

When $\theta = 0, t = 0$ and when $\theta = \frac{\pi}{2}, t = 1$

$$\begin{aligned} \therefore \int_0^{\pi/2} \frac{\cos \theta}{(1 + \sin \theta)(2 + \sin \theta)} d\theta &= \int_0^1 \frac{dt}{(1+t)(2+t)} \\ &= \int_0^1 \left(\frac{1}{1+t} - \frac{1}{2+t} \right) dt \\ &= [\log(1+t) - \log(2+t)]_0^1 \\ &= \log\left(\frac{2}{3}\right) - \log\left(\frac{1}{2}\right) = \log\left(\frac{4}{3}\right) \end{aligned}$$

57. Put $1 + \tan \theta = t \Rightarrow \sec^2 \theta d\theta = dt$

When $\theta = 0, t = 1$ and when $\theta = \frac{\pi}{4}, t = 2$

$$\begin{aligned} \therefore \int_0^{\pi/4} \frac{\sec^2 \theta}{(1 + \tan \theta)(2 + \tan \theta)} d\theta &= \int_1^2 \frac{dt}{t(1+t)} = \int_1^2 \frac{dt}{t} - \int_1^2 \frac{dt}{1+t} \\ &= [\log t - \log(1+t)]_1^2 \\ &= \log_e 2 - \log_e 3 + \log_e 2 = \log_e \left(\frac{4}{3}\right) \end{aligned}$$

58. Let $I = \int_0^{\pi/4} \frac{\sec \theta}{1 + 2\sin^2 \theta} d\theta$

$$\begin{aligned} &= \int_0^{\pi/4} \frac{\cos \theta}{\cos^2 \theta (1 + 2\sin^2 \theta)} d\theta \\ &= \int_0^{\pi/4} \frac{\cos \theta}{(1 - \sin^2 \theta)(1 + 2\sin^2 \theta)} d\theta \end{aligned}$$

Put $\sin \theta = t \Rightarrow \cos \theta d\theta = dt$

$$\begin{aligned} \therefore I &= \int_0^{1/\sqrt{2}} \frac{1}{(1-t^2)(1+2t^2)} dt \\ &= \frac{1}{3} \int_0^{1/\sqrt{2}} \left(\frac{1}{1-t^2} + \frac{2}{1+2t^2} \right) dt \\ &= \frac{1}{3} \left[\frac{1}{2.1} \log\left(\frac{1+t}{1-t}\right) + \frac{2}{\sqrt{2}} \tan^{-1}(\sqrt{2}t) \right]_0^{1/\sqrt{2}} \\ &= \frac{1}{3} \left[\frac{1}{2} \log\left(\frac{\sqrt{2}+1}{\sqrt{2}-1}\right) + \sqrt{2} \tan^{-1} 1 \right] \\ &= \frac{1}{3} \left[\frac{1}{2} \log(\sqrt{2}+1)^2 + \sqrt{2} \cdot \frac{\pi}{4} \right] \\ &= \frac{1}{3} \left[\log(\sqrt{2}+1) + \frac{\pi}{2\sqrt{2}} \right] \end{aligned}$$

59. $\int_{1/e}^{\tan^{-1} 1} \frac{t}{1+t^2} dt + \int_{1/e}^{\cot^{-1} 1} \frac{dt}{t(1+t^2)}$

$$\begin{aligned} &= \frac{1}{2} \int_{1/e}^{\tan^{-1} 1} \frac{2t}{1+t^2} dt + \int_{1/e}^{\cot^{-1} 1} \left(\frac{1}{t} - \frac{t}{1+t^2} \right) dt \\ &= \frac{1}{2} \left[\log(1+t^2) \right]_{1/e}^{\tan^{-1} 1} + \left[\log t - \frac{1}{2} \log(1+t^2) \right]_{1/e}^{\cot^{-1} 1} \\ &= \frac{1}{2} \left[\log(\sec^2 \theta) - \log\left(1 + \frac{1}{e^2}\right) \right] + \log(\cot \theta) \\ &\quad - \log\left(\frac{1}{e}\right) - \frac{1}{2} \left[\log(\operatorname{cosec}^2 \theta) - \log\left(1 + \frac{1}{e^2}\right) \right] \\ &= -\log\left(\frac{1}{e}\right) = \log e = 1 \end{aligned}$$

60. $\int_1^4 f(x) dx = \int_1^2 (4x+3) dx + \int_2^4 (3x+5) dx$

$$\begin{aligned} &= \left[2x^2 + 3x \right]_1^2 + \left[\frac{3}{2}x^2 + 5x \right]_2^4 = 37 \end{aligned}$$

61. $\int_{-1}^2 |x| dx = \int_{-1}^0 (-x) dx + \int_0^2 x dx$

$$\begin{aligned} &= -\left[\frac{x^2}{2} \right]_{-1}^0 + \left[\frac{x^2}{2} \right]_0^2 \\ &= -\left(0 - \frac{1}{2} \right) + 2 \\ &= 2 + \frac{1}{2} = \frac{5}{2} \end{aligned}$$

62. $\int_0^3 |2-x| dx = \int_0^2 (2-x) dx + \int_2^3 -(2-x) dx$

$$\begin{aligned} &= \left[2x - \frac{x^2}{2} \right]_0^2 - \left[2x - \frac{x^2}{2} \right]_2^3 \\ &= (4-2) - \left[6 - \frac{9}{2} - (4-2) \right] \\ &= 2 - \left(4 - \frac{9}{2} \right) = \frac{5}{2} \end{aligned}$$

63. $\int_{-4}^4 |x+2| dx = -\int_{-4}^{-2} (x+2) dx + \int_{-2}^4 (x+2) dx$

$$\begin{aligned} &= \left[\frac{-x^2}{2} - 2x \right]_{-4}^{-2} + \left[\frac{x^2}{2} + 2x \right]_{-2}^4 = 20 \end{aligned}$$



64. Since, \sin is positive in the interval $(0, \pi)$ and negative in the interval $(\pi, 2\pi)$.

$$\begin{aligned} \therefore \int_0^{2\pi} |\sin x| dx &= \int_0^{\pi} \sin x dx + \int_{\pi}^{2\pi} (-\sin x) dx \\ &= [-\cos x]_0^{\pi} + [\cos x]_{\pi}^{2\pi} \\ &= 1 + 1 + 1 + 1 = 4 \end{aligned}$$

$$\begin{aligned} 65. \int_0^{2\pi} (\sin x + |\sin x|) dx &= \int_0^{\pi} 2\sin x dx + \int_{\pi}^{2\pi} 0 dx \\ &= 2[-\cos x]_0^{\pi} + 0 \\ &= -2(\cos \pi - \cos 0) \\ &= -2(-1 - 1) = 4 \end{aligned}$$

$$\begin{aligned} 66. \int_0^{\pi} \sqrt{\frac{1 + \cos 2x}{2}} dx &= \int_0^{\pi} |\cos x| dx \\ &= \int_0^{\pi/2} \cos x dx - \int_{\pi/2}^{\pi} \cos x dx \\ &= [\sin x]_0^{\pi/2} - [\sin x]_{\pi/2}^{\pi} \\ &= \left[\sin \frac{\pi}{2} - \sin 0 \right] - \left[\sin \pi - \sin \frac{\pi}{2} \right] = 1 + 1 = 2 \end{aligned}$$

$$\begin{aligned} 67. \int_0^2 x^2 dx &= \int_0^1 x^2 dx + \int_1^2 x^2 dx \\ &= \int_0^1 x^2(0) dx + \int_1^2 x^2(1) dx \\ &= 0 + \left[\frac{x^3}{3} \right]_1^2 = \frac{7}{3} \end{aligned}$$

$$68. \text{ Let } I = \int_0^{\pi/2} \frac{\cos x - \sin x}{1 + \sin x \cos x} dx \quad \dots(i)$$

$$\begin{aligned} \therefore I &= \int_0^{\pi/2} \frac{\cos\left(\frac{\pi}{2} - x\right) - \sin\left(\frac{\pi}{2} - x\right)}{1 + \sin\left(\frac{\pi}{2} - x\right)\cos\left(\frac{\pi}{2} - x\right)} dx \\ &\quad \dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right] \end{aligned}$$

$$\therefore I = \int_0^{\pi/2} \frac{\sin x - \cos x}{1 + \cos x \sin x} dx \quad \dots(ii)$$

Adding (i) and (ii), we get
 $2I = 0 \Rightarrow I = 0$

$$\begin{aligned} 69. \int_0^{\pi/2} \log \tan x dx &= \int_0^{\pi/2} \log \left(\frac{\sin x}{\cos x} \right) dx \\ &= \int_0^{\pi/2} \log \sin x dx - \int_0^{\pi/2} \log \cos x dx \end{aligned}$$

$$\begin{aligned} &= \int_0^{\pi/2} \log \sin x dx - \int_0^{\pi/2} \log \sin x dx \\ &\quad \dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right] \\ &= 0 \end{aligned}$$

$$70. \text{ Let } I = \int_0^{2a} \frac{f(x)}{f(x) + f(2a-x)} dx \quad \dots(i)$$

$$\therefore I = \int_0^{2a} \frac{f(2a-x)}{f(2a-x) + f(x)} dx \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_0^{2a} dx = 2a$$

$$\Rightarrow I = a$$

$$71. \text{ Let } I = \int_0^{\pi/2} \frac{1000^{\sin x}}{1000^{\sin x} + 1000^{\cos x}} dx \quad \dots(i)$$

$$\begin{aligned} \therefore I &= \int_0^{\pi/2} \frac{1000^{\sin\left(\frac{\pi}{2} - x\right)}}{1000^{\sin\left(\frac{\pi}{2} - x\right)} + 1000^{\cos\left(\frac{\pi}{2} - x\right)}} dx \\ &\quad \dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right] \end{aligned}$$

$$\therefore I = \int_0^{\pi/2} \frac{1000^{\cos x}}{1000^{\cos x} + 1000^{\sin x}} dx \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_0^{\pi/2} 1 dx = \left[x \right]_0^{\pi/2} = \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4}$$

$$72. \text{ Let } I = \int_0^{\pi/2} \frac{e^{2x}}{e^{2x} + e^{\left(\frac{\pi}{2} - x\right)^2}} dx \quad \dots(i)$$

$$\therefore I = \int_0^{\pi/2} \frac{e^{\left(\frac{\pi}{2} - x\right)^2}}{e^{\left(\frac{\pi}{2} - x\right)^2} + e^{2x}} dx \quad \dots(ii)$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \int_0^{\pi/2} dx = \left[x \right]_0^{\pi/2} = \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4}$$



$$73. \text{ Let } I = \int_0^{\pi/2} \frac{\sin^{\frac{3}{2}} d}{\cos^{\frac{3}{2}} + \sin^{\frac{3}{2}}} \dots(i)$$

$$= \int_0^{\pi/2} \frac{\sin^{\frac{3}{2}} \left(\frac{\pi}{2} - \right)}{\cos^{\frac{3}{2}} \left(\frac{\pi}{2} - \right) + \sin^{\frac{3}{2}} \left(\frac{\pi}{2} - \right)} d$$

$$= \int_0^{\pi/2} \frac{\cos^{\frac{3}{2}} d}{\sin^{\frac{3}{2}} + \cos^{\frac{3}{2}}} \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_0^{\pi/2} d = []_0^{\pi/2} = \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4}$$

Alternate Method:

$$\int_0^{\pi/2} \frac{\sin^n}{\sin^n + \cos^n} d = \frac{\pi}{4}$$

$$74. \text{ Let } I = \int_0^{\frac{\pi}{2}} \frac{d}{1 + \tan^3}$$

$$= \int_0^{\frac{\pi}{2}} \frac{\cos^3}{\sin^3 + \cos^3} d \dots(i)$$

$$\therefore I = \int_0^{\frac{\pi}{2}} \frac{\sin^3}{\cos^3 + \sin^3} d \dots(ii)$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \int_0^{\frac{\pi}{2}} d = []_0^{\pi/2}$$

$$\therefore 2I = \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4}$$

$$75. \text{ Let } I = \int_0^{\pi} e^{\cos^2} \cos^5 3 d$$

$$= \int_0^{\pi} e^{\cos^2(\pi-x)} \cos^5 3(\pi-x) dx$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$\therefore I = -\int_0^{\pi} e^{\cos^2} \cos^5 3 d = -I$$

$$\Rightarrow 2I = 0 \Rightarrow I = 0$$

$$76. \text{ Let } I = \int_0^{\pi/2} \log \left(\frac{4+3 \sin}{4+3 \cos} \right) d$$

$$= \int_0^{\pi/2} \log \left(\frac{4+3 \cos}{4+3 \sin} \right) d$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$\therefore I = -\int_0^{\pi/2} \log \left(\frac{4+3 \sin}{4+3 \cos} \right) d = -I$$

$$\Rightarrow 2I = 0 \Rightarrow I = 0$$

$$77. \text{ Let } I = \int_0^{\frac{\pi}{2}} \frac{d\theta}{1 + \tan \theta} \dots(i)$$

$$= \int_0^{\frac{\pi}{2}} \frac{d\theta}{1 + \tan \left(\frac{\pi}{2} - \theta \right)}$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$\therefore I = \int_0^{\frac{\pi}{2}} \frac{d\theta}{1 + \cot \theta} \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_0^{\frac{\pi}{2}} \left(\frac{1}{1 + \tan \theta} + \frac{1}{1 + \cot \theta} \right) d\theta$$

$$= \int_0^{\frac{\pi}{2}} \left(\frac{1}{1 + \tan \theta} + \frac{\tan \theta}{\tan \theta + 1} \right) d\theta$$

$$= \int_0^{\frac{\pi}{2}} d\theta = [\theta]_0^{\pi/2}$$

$$\therefore 2I = \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4}$$

$$78. \text{ Let } I = \int_0^{\pi} \sin^3 d \dots(i)$$

$$= \int_0^{\pi} (\pi-x) \sin^3 d \dots(ii)$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

Adding (i) and (ii), we get



$$\begin{aligned}
 2I &= \pi \int_0^{\pi} \sin^3 x \, dx = \frac{\pi}{4} \int_0^{\pi} (3 \sin x - \sin 3x) \, dx \\
 &= \frac{\pi}{4} \left[-3 \cos x + \frac{\cos 3x}{3} \right]_0^{\pi} \\
 &= \frac{\pi}{4} \left[3 - \frac{1}{3} + 3 - \frac{1}{3} \right] \\
 &= \frac{4\pi}{3}
 \end{aligned}$$

$$\therefore I = \frac{2\pi}{3}$$

$$\begin{aligned}
 79. \quad \text{Let } I &= \int_0^{\frac{\pi}{2}} \log \sin x \, dx \\
 &= \int_0^{\frac{\pi}{4}} (\log \sin x + \log \cos x) \, dx \\
 &\dots \left[\because \int_0^{2a} f(x) \, dx = \int_0^a [f(x) + f(2a-x)] \, dx \right] \\
 &= \int_0^{\frac{\pi}{4}} \log \sin x \cos x \, dx \\
 &= \int_0^{\frac{\pi}{4}} \log \left(\frac{\sin 2x}{2} \right) \, dx \\
 &= \int_0^{\frac{\pi}{4}} \log \sin 2x \, dx - \int_0^{\frac{\pi}{4}} \log 2 \, dx
 \end{aligned}$$

In 1st integral, put $2x = t \Rightarrow 2 \, dx = dt$

$$\begin{aligned}
 \therefore I &= \frac{1}{2} \int_0^{\frac{\pi}{2}} \log \sin t \, dt - \frac{\pi}{4} \log 2 \\
 &= \frac{1}{2} \int_0^{\frac{\pi}{2}} \log \sin x \, dx - \frac{\pi}{4} \log 2 \\
 &\dots \left[\because \int_a^b f(x) \, dx = \int_a^b f(t) \, dt \right]
 \end{aligned}$$

$$\therefore I = \frac{1}{2} I - \frac{\pi}{4} \log 2$$

$$\therefore I = -\frac{\pi}{2} \log 2$$

$$80. \quad \text{Let } I = \int_0^{\pi} \log \sin x \, dx \quad \dots \text{(i)}$$

$$\therefore I = \int_0^{\pi} (\pi - x) \log \sin x \, dx \quad \dots \text{(ii)}$$

$$\dots \left[\because \int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx \right]$$

Adding (i) and (ii), we get

$$2I = \pi \int_0^{\pi} \log \sin x \, dx = 2\pi \int_0^{\frac{\pi}{2}} \log \sin x \, dx$$

$$\therefore 2I = 2\pi \left(-\frac{\pi}{2} \log 2 \right)$$

$$\dots \left[\because \int_0^{\frac{\pi}{2}} \log \sin x \, dx = -\frac{\pi}{2} \log 2 \right]$$

$$\Rightarrow I = \pi \left(\frac{\pi}{2} \log \frac{1}{2} \right) = \frac{\pi^2}{2} \log \frac{1}{2}$$

$$81. \quad \text{Let } I = \int_0^{\frac{\pi}{4}} \log(1 + \tan \theta) \, d\theta$$

$$= \int_0^{\frac{\pi}{4}} \log \left[1 + \tan \left(\frac{\pi}{4} - \theta \right) \right] \, d\theta$$

$$\dots \left[\because \int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx \right]$$

$$= \int_0^{\frac{\pi}{4}} \log \left(1 + \frac{1 - \tan \theta}{1 + \tan \theta} \right) \, d\theta$$

$$= \int_0^{\frac{\pi}{4}} \log 2 \, d\theta - \int_0^{\frac{\pi}{4}} \log(1 + \tan \theta) \, d\theta$$

$$\therefore 2I = \int_0^{\frac{\pi}{4}} \log 2 \, d\theta \Rightarrow I = \frac{\log 2}{2} [\theta]_0^{\pi/4} = \frac{\pi}{8} \log 2$$

$$82. \quad \int_0^1 \tan^{-1} \left(\frac{2x-1}{1-x^2} \right) \, dx$$

$$= \int_0^1 \tan^{-1} \left(\frac{+(x-1)}{1-(x-1)} \right) \, dx$$

$$= \int_0^1 (\tan^{-1} x + \tan^{-1} (x-1)) \, dx$$



$$\begin{aligned}
 &= \int_0^1 \tan^{-1} x \, dx + \int_0^1 \tan^{-1}(1-x) \, dx \\
 &= \int_0^1 \tan^{-1} x \, dx + \int_0^1 \tan^{-1}(1-x) \, dx \\
 &\quad \dots \left[\because \int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx \right] \\
 &= \int_0^1 \tan^{-1} x \, dx - \int_0^1 \tan^{-1} x \, dx = 0
 \end{aligned}$$

83. Let $I = \int_0^{\pi} \frac{\tan x}{\sec x + \tan x} dx \quad \dots(i)$

$\therefore I = \int_0^{\pi} \frac{(\pi-x)\tan x}{\sec x + \tan x} dx \quad \dots(ii)$

$\dots \left[\because \int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx \right]$

Adding (i) and (ii), we get

$$\begin{aligned}
 2I &= \pi \int_0^{\pi} \frac{\tan x}{\sec x + \tan x} dx \\
 \therefore I &= \frac{\pi}{2} \int_0^{\pi} \frac{\sin x}{1 + \sin x} dx \\
 &= \frac{\pi}{2} \left[\int_0^{\pi} 1 dx - \int_0^{\pi} \frac{dx}{1 + \sin x} \right]
 \end{aligned}$$

On solving, we get

$$I = \frac{\pi}{2}(\pi - 2) = \pi \left(\frac{\pi}{2} - 1 \right)$$

84. Let $I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{dx}{1 + \sqrt{\cot x}}$

$$= \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \quad \dots(i)$$

$\therefore I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx \quad \dots(ii)$

$\dots \left[\because \int_a^b f(x) \, dx = \int_a^b f(a+b-x) \, dx \right]$

Adding (i) and (ii), we get

$$\begin{aligned}
 2I &= \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} dx \\
 \therefore I &= \frac{1}{2} \left(\frac{\pi}{3} - \frac{\pi}{6} \right) = \frac{\pi}{12}
 \end{aligned}$$

85. Let $I = \int_a^b f(x) \, dx$

$$= \int_a^b f(a+b-x) f(a+b-x) \, dx$$

$\therefore I = \int_a^b (a+b)f(x) \, dx - \int_a^b f(x) \, dx$

$\dots [\because f(a+b-x) = f(x) \text{ (given)}]$

$$\Rightarrow 2I = (a+b) \int_a^b f(x) \, dx \Rightarrow I = \frac{a+b}{2} \int_a^b f(x) \, dx$$

86. Let $I = \int_{0.5}^{1.5} f(x) \, dx = \int_{0.5}^{1.5} (2-x)f(2-x) \, dx$

$\dots \left[\because \int_a^b f(x) \, dx = \int_a^b f(a+b-x) \, dx \right]$

$$= \int_{0.5}^{1.5} (2-x)f(x) \, dx$$

$\dots [\because f(x) = f(2-x) \text{ (given)}]$

$\therefore I = 2 \int_{0.5}^{1.5} f(x) \, dx - I$

$$\Rightarrow 2I = 2 \int_{0.5}^{1.5} f(x) \, dx \Rightarrow I = \int_{0.5}^{1.5} f(x) \, dx$$

87. $\int_0^{2a} f(x) \, dx = \int_0^a f(x) \, dx + \int_a^{2a} f(x) \, dx$

Let $I_1 = \int_a^{2a} f(x) \, dx$

Put $x = 2a - t \Rightarrow dx = -dt$

$\therefore I_1 = - \int_a^0 f(2a-t) dt$

$$= \int_0^a f(2a-t) dt = \int_0^a f(2a-x) dx$$

$\therefore \int_0^{2a} f(x) \, dx = \int_0^a f(x) \, dx + \int_0^a f(2a-x) dx$

$$= 2 \int_0^a f(x) \, dx, \text{ if } f(2a-x) = f(x)$$



$$88. \text{ Let } I = \int_0^{\pi} \sin^2 x \, dx = 2 \int_0^{\pi/2} \sin^2 x \, dx$$

$$\dots \left[\because \int_0^{2a} f(x) \, dx = 2 \int_0^a f(x) \, dx, \text{ if } f(2a-x) = f(x) \right]$$

$$\therefore I = 2 \times \frac{1}{2} \times \frac{\pi}{2}$$

$$= \frac{\pi}{2}$$

$$89. \int_0^{\pi} |\cos x| \, dx = 2 \int_0^{\pi/2} |\cos x| \, dx$$

$$\dots \left[\because \int_0^{2a} f(x) \, dx = 2 \int_0^a f(x) \, dx, \right.$$

$$\left. \text{if } f(2a-x) = f(x) \right]$$

$$= 2 \left[\sin x \right]_0^{\pi/2} = 2$$

$$90. \int_0^{2\pi} \cos^{99} x \, dx = 2 \int_0^{\pi} \cos^{99} x \, dx$$

$$\dots \left[\because \int_0^{2a} f(x) \, dx = 2 \int_0^a f(x) \, dx, \text{ if } f(2a-x) = f(x) \right]$$

$$\text{Let } I_1 = \int_0^{\pi} \cos^{99} x \, dx$$

$$\Rightarrow I_1 = - \int_0^{\pi} \cos^{99} x \, dx$$

$$\dots \left[\because \int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx \right]$$

$$\Rightarrow I_1 = -I_1 \Rightarrow 2I_1 = 0 \Rightarrow I_1 = 0$$

$$\therefore \int_0^{2\pi} \cos^{99} x \, dx = 2(0) = 0$$

$$91. \int_0^{\pi} \log \sin^2 x \, dx = \int_0^{\pi} 2 \log \sin x \, dx = 2 \int_0^{\pi/2} \log \sin x \, dx$$

$$= 2 \int_0^{\pi/2} [\log \sin x + \log \sin(\pi-x)] \, dx$$

$$\dots \left[\because \int_0^{2a} f(x) \, dx = \int_0^a [f(x) + f(2a-x)] \, dx \right]$$

$$= 4 \int_0^{\pi/2} \log \sin x \, dx$$

$$= 4 \times \left(-\frac{\pi}{2} \log 2 \right) = -2\pi \log_e 2 = 2\pi \log_e \left(\frac{1}{2} \right)$$

$$92. \text{ Let } I = \int_0^{\pi} \frac{dx}{a^2 \cos^2 x + b^2 \sin^2 x} \dots (i)$$

$$\therefore I = \int_0^{\pi} \frac{\pi-x}{a^2 \cos^2 x + b^2 \sin^2 x} \, dx \dots (ii)$$

$$\dots \left[\because \int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx \right]$$

Adding (i) and (ii), we get

$$2I = \pi \int_0^{\pi} \frac{dx}{a^2 \cos^2 x + b^2 \sin^2 x}$$

$$\therefore I = \frac{\pi}{2} \int_0^{\pi} \frac{dx}{a^2 \cos^2 x + b^2 \sin^2 x}$$

$$= 2 \cdot \frac{\pi}{2} \int_0^{\pi/2} \frac{dx}{a^2 \cos^2 x + b^2 \sin^2 x}$$

$$\dots \left[\because \int_0^{2a} f(x) \, dx = 2 \int_0^a f(x) \, dx, \right.$$

$$\left. \text{if } f(2a-x) = f(x) \right]$$

$$= \pi \int_0^{\pi/2} \frac{\sec^2 x}{a^2 + b^2 \tan^2 x} \, dx$$

Put $b \tan x = t \Rightarrow b \sec^2 x \, dx = dt$

$$\therefore I = \frac{\pi}{b} \int_0^{\infty} \frac{dt}{a^2 + t^2} = \frac{\pi}{b} \cdot \frac{1}{a} \left[\tan^{-1} \frac{t}{a} \right]_0^{\infty}$$

$$= \frac{\pi}{ab} \left(\frac{\pi}{2} - 0 \right) = \frac{\pi^2}{2ab}$$

$$93. \int_{-1}^1 f(x) \, dx = \int_{-1}^0 f(x) \, dx + \int_0^1 f(x) \, dx$$

In 1st integral, put $x = -t \Rightarrow dx = -dt$

$$\therefore \int_{-1}^0 f(x) \, dx = - \int_1^0 f(-t) \, dt$$

$$= \int_0^1 f(-t) \, dt$$

$$= \int_0^1 f(-x) \, dx$$

$$\therefore \int_{-1}^1 f(x) \, dx = \int_0^1 f(-x) \, dx + \int_0^1 f(x) \, dx$$

$$= 0, \text{ if } f(-x) = -f(x)$$



94. Since, $\int_{-a}^a f(x) dx = 0$, if $f(-x) = -f(x)$

$$\therefore \int_{-1}^1 f(x) dx = 0$$

$$\Rightarrow \int_{-1}^0 f(x) dx + \int_0^1 f(x) dx = 0$$

$$\Rightarrow \int_{-1}^0 f(x) dx = -5$$

$$\Rightarrow \int_{-1}^0 f(t) dt = -5$$

95. Let $f(x) = |x|$

$$\therefore f(-x) = |-x| = -|x| = -f(x)$$

$\therefore f(x)$ is an odd function.

$$\therefore \int_{-1}^1 |x| dx = 0$$

96. Since, $|\sin x|$ is an even function.

$$\begin{aligned} \therefore I &= 2 \int_0^{\frac{\pi}{2}} |\sin x| dx = 2 \int_0^{\frac{\pi}{2}} \sin x dx = 2[-\cos x]_0^{\frac{\pi}{2}} \\ &= 2(-0 + 1) = 2 \end{aligned}$$

97. Since, $\frac{1}{x^3}$ is an odd function.

$$\therefore \int_{-a}^a \frac{1}{x^3} dx = 0$$

98. Let $f(x) = \sin^{-1}(\cos x)$

$$\therefore f(-x) = \sin^{-1}(\cos(-x)) = \sin^{-1}(\cos x) = -f(x)$$

$\therefore f(x)$ is an odd function.

$$\therefore \int_{-a}^a f(x) dx = 0$$

99. Let $f(x) = \frac{\sin x}{1 + \cos^2 x} e^{-\cos^2 x}$

$$\therefore f(-x) = \frac{\sin(-x)}{1 + \cos^2(-x)} e^{-\cos^2(-x)} = -f(x)$$

$\therefore f(x)$ is an odd function.

$$\therefore \int_{-\pi/2}^{\pi/2} f(x) dx = 0$$

100. Let $f(x) = \sqrt{1+x^2} - \sqrt{1-x^2}$

$$\therefore f(-x) = \sqrt{1+(-x)^2} - \sqrt{1-(-x)^2} = -f(x)$$

$\therefore f(x)$ is an odd function.

$$\therefore \int_{-1}^1 f(x) dx = 0$$

101. Let $f(x) = (e^x + e^{-x})(e^x - e^{-x})$

$$\begin{aligned} \therefore f(-x) &= (e^{-x} + e^x)(e^{-x} - e^x) \\ &= -(e^x + e^{-x})(e^x - e^{-x}) = -f(x) \end{aligned}$$

$\therefore f(x)$ is an odd function.

$$\therefore \int_{-1}^1 (e^x + e^{-x})(e^x - e^{-x}) dx = 0$$

102. Let $f(x) = \log\left(\frac{1+x}{1-x}\right)$

$$\therefore f(-x) = \log\left(\frac{1+(-x)}{1-(-x)}\right) = -\log\left(\frac{1+x}{1-x}\right) = -f(x)$$

$\therefore f(x)$ is an odd function.

$$\therefore \int_{-1}^1 \log\left(\frac{1+x}{1-x}\right) dx = 0$$

103. Let $f(x) = \cos^{-1}\left(\log\left(\frac{1-x}{1+x}\right)\right)$

$$\begin{aligned} \therefore f(-x) &= \cos^{-1}\left(\log\left(\frac{1-(-x)}{1+(-x)}\right)\right) \\ &= \cos^{-1}\left(\log\left(\frac{1-x}{1+x}\right)\right) = -f(x) \end{aligned}$$

$\therefore f(x)$ is an odd function.

$$\therefore \int_{-\frac{1}{2}}^{\frac{1}{2}} \cos^{-1}\left(\log\left(\frac{1-x}{1+x}\right)\right) dx = 0$$

104. Let $f(\theta) = \log(\sec \theta - \tan \theta)$

$$\begin{aligned} \therefore f(-\theta) &= \log(\sec(-\theta) + \tan(-\theta)) \\ &= \log\left(\frac{1}{\sec \theta - \tan \theta}\right) \\ &= -\log(\sec \theta - \tan \theta) = -f(\theta) \end{aligned}$$

$\therefore f(\theta)$ is an odd function.

$$\therefore \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \log(\sec \theta - \tan \theta) d\theta = 0$$

105. Let $f(x) = \log(\sqrt{1+x^2} + x)$

$$\begin{aligned} \therefore f(-x) &= \log(\sqrt{1+(-x)^2} - x) \\ &= \log(\sqrt{1+x^2} - x) \cdot \frac{(\sqrt{1+x^2} + x)}{(\sqrt{1+x^2} + x)} \end{aligned}$$



$$\begin{aligned}
 &= \log \left(\frac{1 + \sqrt{1+x^2}}{\sqrt{1+x^2} + 1} \right) \\
 &= \log 1 - \log(\sqrt{1+x^2} + 1) \\
 &= -\log(\sqrt{1+x^2} + 1) = -f(x)
 \end{aligned}$$

$\therefore f(x)$ is an odd function.

$$\therefore \int_{-1}^1 \log(\sqrt{1+x^2} + 1) dx = 0$$

$$\begin{aligned}
 106. \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^2 x dx &= 2 \int_0^{\frac{\pi}{2}} \sin^2 x dx \\
 &\dots [\because \sin^2 x \text{ is an even function}]
 \end{aligned}$$

$$\text{Since, } \int_0^{\frac{\pi}{2}} \sin^n x dx = \frac{(n-1)(n-3)\dots 1}{n(n-2)\dots 2} \cdot \frac{\pi}{2},$$

if n is even

$$\therefore \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^2 x dx = 2 \left(\frac{2-1}{2} \cdot \frac{\pi}{2} \right) = \frac{\pi}{2}$$

$$\begin{aligned}
 107. \text{ Let } I &= \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \sin^{-4} x dx = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \operatorname{cosec}^4 x dx \\
 &= \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \operatorname{cosec}^2 x (1 + \cot^2 x) dx
 \end{aligned}$$

$$\text{Put } \cot x = t \Rightarrow \operatorname{cosec}^2 x dx = -dt$$

$$\begin{aligned}
 \therefore I &= - \int_{-1}^1 (1+t^2) dt \\
 &= -2 \int_0^1 (1+t^2) dt = -2 \left[t + \frac{t^3}{3} \right]_0^1 \\
 &= -2 \left(1 + \frac{1}{3} \right) = -\frac{8}{3}
 \end{aligned}$$

$$\begin{aligned}
 108. \int_{-1}^1 \frac{1+x^3}{9-x^2} dx &= \int_{-1}^1 \frac{1}{9-x^2} dx + \int_{-1}^1 \frac{x^3}{9-x^2} dx \\
 &= 2 \int_0^1 \frac{1}{9-x^2} dx + 0 \\
 &\dots \left[\begin{array}{l} \because \frac{1}{9-x^2} \text{ is an even function and} \\ \frac{x^3}{9-x^2} \text{ is an odd function.} \end{array} \right]
 \end{aligned}$$

$$= 2 \left[\frac{1}{2 \times 3} \log \left| \frac{3+x}{3-x} \right| \right]_0^1 = \frac{1}{3} (\log 2 - \log 1) = \frac{1}{3} \log 2$$

$$\begin{aligned}
 109. \text{ Let } I &= \int_{-\pi}^{\pi} \frac{2(1+\sin x)}{1+\cos^2 x} dx \\
 &= \int_{-\pi}^{\pi} \frac{2}{1+\cos^2 x} dx + \int_{-\pi}^{\pi} \frac{2 \sin x}{1+\cos^2 x} dx
 \end{aligned}$$

Since, $\frac{2}{1+\cos^2 x}$ is an odd function

and $\frac{2 \sin x}{1+\cos^2 x}$ is an even function.

$$\begin{aligned}
 \therefore I &= 0 + 2 \int_0^{\pi} \frac{2 \sin x}{1+\cos^2 x} dx \\
 \Rightarrow I &= 4 \int_0^{\pi} \frac{\sin x}{1+\cos^2 x} dx \quad \dots (i)
 \end{aligned}$$

$$\Rightarrow I = 4 \int_0^{\pi} \frac{(\pi-x) \sin x}{1+\cos^2 x} dx \quad \dots (ii)$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = 4 \int_0^{\pi} \frac{\pi \sin x}{1+\cos^2 x} dx \Rightarrow I = 2\pi \int_0^{\pi} \frac{\sin x}{1+\cos^2 x} dx$$

$$\text{Put } \cos x = t \Rightarrow -\sin x dx = dt$$

$$\begin{aligned}
 \therefore I &= 2\pi \int_1^{-1} \frac{-dt}{1+t^2} \\
 \Rightarrow I &= -2\pi \left[\tan^{-1} t \right]_1^{-1} = -2\pi \left(\frac{-\pi}{4} - \frac{\pi}{4} \right) = \pi^2
 \end{aligned}$$

$$\begin{aligned}
 110. \text{ Let } I &= \int_0^{\pi/2} \frac{\sin x}{1+\cos^2 x} dx \\
 \text{Put } \cos x &= t \Rightarrow -\sin x dx = dt \\
 \therefore I &= \int_1^0 \frac{-dt}{1+t^2} = \int_0^1 \frac{dt}{1+t^2} = \left[\tan^{-1} t \right]_0^1 = \frac{\pi}{4}
 \end{aligned}$$

$$\begin{aligned}
 111. \text{ Let } I &= \int_0^{\pi/2} \sqrt{\cos \theta} \sin^3 \theta d\theta \\
 \text{Put } t &= \cos \theta \Rightarrow dt = -\sin \theta d\theta \\
 \therefore I &= - \int_1^0 t^{1/2} (1-t^2) dt = \int_0^1 (t^{1/2} - t^{5/2}) dt \\
 &= \left[\frac{2}{3} t^{3/2} - \frac{2}{7} t^{7/2} \right]_0^1 = \frac{8}{21}
 \end{aligned}$$



$$112. \int_0^1 \frac{d}{+\sqrt{1-d^2}} = \int_0^{\pi/2} \frac{\cos \theta d\theta}{\sin \theta + \cos \theta}$$

....[Put $d = \sin \theta \Rightarrow dd = \cos \theta d\theta$]

$$= \frac{\pi}{4}$$

113. Since, $\sqrt{1+d^2} > d$, for all $d \in (1,2)$

$$\Rightarrow \frac{1}{\sqrt{1+d^2}} < \frac{1}{d}, \text{ for all } d \in (1,2)$$

$$\Rightarrow \int_1^2 \frac{d}{\sqrt{1+d^2}} < \int_1^2 \frac{d}{d}$$

$$\Rightarrow I_1 < I_2$$

114. Let $I = \int_0^{\pi/2} \frac{\sin \cos}{\cos^2 + 3 \cos + 2} d$

Put $\cos = t \Rightarrow -\sin d = dt$

When $d = 0, t = 1$ and when $d = \frac{\pi}{2}, t = 0$

$$\therefore I = -\int_1^0 \frac{t}{t^2 + 3t + 2} dt$$

$$= \int_0^1 \frac{t}{(t+2)(t+1)} dt \quad \dots \left[\because \int_a^b f(x) dx = -\int_b^a f(x) dx \right]$$

$$= \int_0^1 \left(\frac{2}{t+2} - \frac{1}{t+1} \right) dt$$

$$= [2 \log(t+2) - \log(t+1)]_0^1$$

$$= 2 \log 3 - \log 2 - 2 \log 2$$

$$= 2 \log 3 - 3 \log 2 = \log 9 - \log 8 = \log \left(\frac{9}{8} \right)$$

115. Put $d+1 = t^2 \Rightarrow dd = 2t dt$

When $d = 3, t = 2$ and when $d = 8, t = 3$

$$\therefore \int_3^8 \frac{2-3}{\sqrt{1+d}} d = 2 \int_2^3 \frac{2-3(t^2-1)}{t^2-1} dt$$

$$= 2 \int_2^3 \left(\frac{2}{t^2-1} - 3 \right) dt$$

$$= 2 \left[2 \cdot \frac{1}{2 \times 1} \log \left(\frac{t-1}{t+1} \right) - 3t \right]_2^3$$

$$= 2 \left(\log \frac{1}{2} - \log \frac{1}{3} - 3 \right)$$

$$= 2 \left(\log \frac{3}{2} - 3 \log e \right) = 2 \left(\log \frac{3}{2} - \log e^3 \right)$$

$$= 2 \log \left(\frac{3}{2e^3} \right)$$



Competitive Thinking

1. $\int_0^1 \sqrt{d} d = \left[\frac{2}{3} d^{3/2} \right]_0^1 = \frac{2}{3}$

2. $\int_0^{\pi/2} \frac{\sin + \cos}{\sqrt{1+\sin 2}} d = \int_0^{\pi/2} \frac{\sin + \cos}{\sqrt{(\sin + \cos)^2}} d$

$$= \int_0^{\pi/2} d = \frac{\pi}{2}$$

3. $\int_0^{\pi/2} \log \sec d = \int_0^{\pi/2} \log \frac{1}{\cos} d$

$$= -\int_0^{\pi/2} \log \cos d$$

$$= -\frac{\pi}{2} \log \frac{1}{2}$$

$$= \frac{\pi}{2} \log 2$$

4. $\int_0^1 \tan^{-1} \left(\frac{1-d}{1+d} \right) d = \int_0^1 \tan^{-1} d - \int_0^1 \tan^{-1} \frac{1}{d} d$

$$= (\tan^{-1} 1) []_0^1 - p$$

$$= \frac{\pi}{4} - p$$

5. $\int_{-\pi/4}^{\pi/4} \frac{d}{1+\cos 2}$

$$= \int_{-\pi/4}^{\pi/4} \frac{d}{2 \cos^2} = \frac{1}{2} \int_{-\pi/4}^{\pi/4} \sec^2 d$$

$$= \frac{1}{2} [\tan]_{-\pi/4}^{\pi/4} = \frac{1}{2} [1 - (-1)]$$

$$= 1$$

6. $\int_0^{\pi/2} \frac{\cos 2}{\cos + \sin} d$

$$= \int_0^{\pi/2} \frac{\cos^2 - \sin^2}{\cos + \sin} d$$

$$= \int_0^{\pi/2} (\cos - \sin) d$$

$$= [\sin + \cos]_0^{\pi/2} = 0$$



$$7. \int_0^{\frac{\pi}{2}} \cos^n x \, dx = \frac{(n-1)(n-3)\dots 1}{n(n-2)\dots 2} \cdot \frac{\pi}{2}, \text{ if } n \text{ is even.}$$

$$\therefore \int_0^{\frac{\pi}{2}} \cos^6 x \, dx = \frac{(6-1)(6-3)(6-5)}{6(6-2)(6-4)} \cdot \frac{\pi}{2} \\ = \frac{5\pi}{32}$$

$$8. \text{ Let } I = \int_{-1}^0 \frac{dx}{x^2 + 2x + 2} \\ = \int_{-1}^0 \frac{dx}{x^2 + 2x + 1 + 1} \\ = \int_{-1}^0 \frac{dx}{(x+1)^2 + 1} \\ = \left[\tan^{-1}(x+1) \right]_{-1}^0 \\ = \frac{\pi}{4}$$

$$9. \int_0^{\sqrt{2}} \sqrt{2-x^2} \, dx = \left[-\frac{1}{2}\sqrt{2-x^2} + \frac{1}{2}\sin^{-1}\left(\frac{x}{\sqrt{2}}\right) \right]_0^{\sqrt{2}} \\ = \sin^{-1} 1 \\ = \frac{\pi}{2}$$

$$10. \text{ Let } I = \int_0^1 \frac{dx}{e^x + e^{-x}} = \int_0^1 \frac{e^x dx}{(e^x)^2 + 1}$$

$$\text{Put } e^x = t \Rightarrow e^x dx = dt$$

$$\therefore I = \int_1^e \frac{dt}{1+t^2} \\ = \left[\tan^{-1} t \right]_1^e \\ = \tan^{-1} e - \frac{\pi}{4}$$

$$11. 3a \int_0^1 \left(\frac{a-x}{a-1} \right)^2 dx = \frac{3a}{(a-1)^2} \int_0^1 (a-x)^2 dx \\ = \frac{3a \left[(a-x)^3 \right]_0^1}{3a(a-1)^2} \\ = \frac{1}{(a-1)^2} \left[(a-1)^3 + 1 \right] \\ = (a-1) + (a-1)^{-2}$$

$$12. \int_0^{\frac{\pi}{4}} \sin(x) dx = \left[-\cos(x) \right]_0^{\frac{\pi}{4}}$$

$$= \int_0^{\frac{\pi}{4}} \sin(x-0) dx = \left[-\cos(x) \right]_0^{\frac{\pi}{4}}$$

$$= \int_0^{\frac{\pi}{4}} \sin x \, dx$$

$$= \left[-\cos x \right]_0^{\pi/4} = -\cos \frac{\pi}{4} + \cos 0 = 1 - \frac{1}{\sqrt{2}}$$

$$13. L(x) = \int_1^x \frac{1}{t} dt = \left[\log t \right]_1^x = \log x - \log 1 = \log x$$

$$\therefore L(x) = \log(x) = \log(x) + \log(1) = L(x) + L(1)$$

$$14. \text{ Given, } \int_a^b \{f(x) - 3\} dx = a^2 - b^2$$

$$\Rightarrow \int_a^b f(x) dx - 3 \int_a^b dx = a^2 - b^2$$

$$\Rightarrow \int_a^b f(x) dx - \frac{3}{2}(b^2 - a^2) = a^2 - b^2$$

$$\Rightarrow \int_a^b f(x) dx = \frac{1}{2}(b^2 - a^2)$$

$$\therefore f(x) = \dots \left[\because \int_a^b dx = \frac{1}{2}(b^2 - a^2) \right]$$

$$\therefore f\left(\frac{\pi}{6}\right) = \frac{\pi}{6}$$

$$15. I = \int_0^1 (1-x)^n dx$$

$$\therefore -I = \int_0^1 -(1-x)^n dx$$

$$= \int_0^1 (1-x-1)(1-x)^n dx$$

$$= \int_0^1 (1-x)^{n+1} dx - \int_0^1 (1-x)^n dx$$

$$= \left[\frac{(1-x)^{n+2}}{-(n+2)} \right]_0^1 - \left[\frac{(1-x)^{n+1}}{-(n+1)} \right]_0^1$$

$$\therefore -I = \frac{1}{n+2} - \frac{1}{n+1}$$

$$\therefore I = \frac{1}{n+1} - \frac{1}{n+2}$$



$$\begin{aligned}
 16. \quad & \int_0^{\frac{\pi}{4}} (\cos x - \sin x) dx + \int_{\frac{\pi}{4}}^{\frac{5\pi}{4}} (\sin x - \cos x) dx \\
 & + \int_{2\pi}^{\frac{\pi}{4}} (\cos x - \sin x) dx \\
 & = [\sin x + \cos x]_0^{\pi/4} - [\cos x + \sin x]_{\pi/4}^{5\pi/4} \\
 & \quad + [\sin x + \cos x]_{2\pi}^{\pi/4} \\
 & = \left(\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} - 1 \right) - \left[-\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} - \left(\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} \right) \right] \\
 & \quad + \left(\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} - 1 \right) \\
 & = (\sqrt{2} - 1) - (-\sqrt{2} - \sqrt{2}) + (\sqrt{2} - 1) \\
 & = 4\sqrt{2} - 2
 \end{aligned}$$

$$\begin{aligned}
 17. \quad & \frac{d}{dx} f(x) = \frac{e^{\sin x}}{x} \\
 \Rightarrow & f(x) = \int \frac{e^{\sin x}}{x} dx \\
 \text{Let } I &= \int_1^4 \frac{3e^{\sin^3 x}}{x} dx \\
 & = \int_1^4 \frac{3^2 e^{\sin^3 x}}{3} dx \\
 \text{Put } t &= \sin^3 x \Rightarrow 3^2 dx = dt \\
 \therefore I &= \int_1^{64} \frac{e^{\sin t}}{t} dt \\
 & = [f(t)]_1^{64} \\
 & = f(64) - f(1) \\
 \Rightarrow k &= 64
 \end{aligned}$$

$$\begin{aligned}
 18. \quad & \text{Let } I = \int_1^3 \frac{\sin 2x}{x} dx \\
 \text{Put } 2x &= t \Rightarrow 2dx = dt \Rightarrow dx = \frac{dt}{2} \\
 \therefore I &= \int_2^6 \frac{\sin t}{\frac{t}{2}} \cdot \frac{dt}{2} \\
 & = \int_2^6 \frac{\sin t}{t} dt = [F(t)]_2^6 \\
 & = F(6) - F(2)
 \end{aligned}$$

$$\begin{aligned}
 19. \quad & \text{Put } \frac{1}{x} = t \Rightarrow -\frac{1}{x^2} dx = dt \\
 \text{When } x &= \frac{1}{\pi}, t = \pi \text{ and when } x = \frac{2}{\pi}, t = \frac{\pi}{2} \\
 \therefore \int_{1/\pi}^{2/\pi} \frac{\sin\left(\frac{1}{x}\right)}{x^2} dx &= - \int_{\pi}^{\pi/2} \sin t dt = [\cos t]_{\pi}^{\pi/2} \\
 &= 0 - (-1) = 1
 \end{aligned}$$

$$\begin{aligned}
 20. \quad & \text{Let } I = \int_0^{\frac{\pi}{4}} \frac{\sqrt{\tan x}}{\sin x \cos x} dx \\
 & = \int_0^{\frac{\pi}{4}} \frac{\tan x}{\sin x \cos \sqrt{\tan x}} dx \\
 & = \int_0^{\frac{\pi}{4}} \frac{\sec^2 x}{\sqrt{\tan x}} dx \\
 \text{Put } \tan x &= t \Rightarrow \sec^2 x dx = dt \\
 \text{When } x &= 0, t = 0 \text{ and when } x = \frac{\pi}{4}, t = 1
 \end{aligned}$$

$$\therefore I = \int_0^1 \frac{dt}{\sqrt{t}} = [2\sqrt{t}]_0^1 = 2$$

$$\begin{aligned}
 21. \quad & I_8 + I_6 = \int_0^{\frac{\pi}{4}} (\tan^8 \theta + \tan^6 \theta) d\theta \\
 & = \int_0^{\frac{\pi}{4}} \tan^6 \theta \sec^2 \theta d\theta \\
 \text{Put } \tan \theta &= t \Rightarrow \sec^2 \theta d\theta = dt \\
 \text{When } \theta &= 0, t = 0 \text{ and when } \theta = \frac{\pi}{4}, t = 1
 \end{aligned}$$

$$\therefore I_8 + I_6 = \int_0^1 t^6 dt = \left[\frac{t^7}{7} \right]_0^1 = \frac{1}{7}$$

$$\begin{aligned}
 22. \quad & \text{Let } I = \int_1^2 [f(g(x))]^{-1} f'[g(x)] g'(x) dx \\
 \text{Put } f[g(x)] &= z \Rightarrow f'[g(x)] g'(x) dx = dz \\
 \text{When } x &= 1, z = f[g(1)] \\
 \text{and when } x &= 2, z = f[g(2)] \\
 \therefore I &= \int_{f[g(1)]}^{f[g(2)]} \frac{1}{z} dz = [\log z]_{f[g(1)]}^{f[g(2)]} \\
 & = \log f[g(2)] - \log f[g(1)] \\
 & = 0 \quad \dots [\because g(1) = g(2) \text{ (given)}]
 \end{aligned}$$



$$23. \int_0^k \frac{d}{2+18^{-2}} = \frac{1}{18} \int_0^k \frac{d}{2 + \frac{1}{9}}$$

$$\Rightarrow \frac{\pi}{24} = \frac{1}{18} \int_0^k \frac{d}{2 + \left(\frac{1}{3}\right)^2}$$

$$= \frac{1}{18} \cdot \frac{1}{\left(\frac{1}{3}\right)} \left[\tan^{-1} \frac{1}{\left(\frac{1}{3}\right)} \right]_0^k$$

$$= \frac{1}{6} \left[\tan^{-1} 3 \right]_0^k$$

$$\Rightarrow \frac{\pi}{24} = \frac{1}{6} (\tan^{-1} 3k - 0)$$

$$\Rightarrow \frac{\pi}{4} = \tan^{-1} 3k$$

$$\Rightarrow \tan \frac{\pi}{4} = 3k$$

$$\Rightarrow 3k = 1$$

$$\Rightarrow k = \frac{1}{3}$$

$$24. \int_{-1}^0 \frac{d}{x^2+2x+2} = \int_{-1}^0 \frac{d}{(x+1)^2+1}$$

$$= \left[\tan^{-1}(x+1) \right]_{-1}^0 = \tan^{-1} 1 - \tan^{-1} 0 = \frac{\pi}{4}$$

$$25. \int_0^1 \frac{x^2}{1+x^2} dx = \int_0^1 \left(1 - \frac{1}{1+x^2} \right) dx$$

$$= \left[x - \tan^{-1} x \right]_0^1$$

$$= 1 - \frac{\pi}{4}$$

$$26. \int_0^1 \frac{x^4+1}{x^2+1} dx = \int_0^1 \frac{x^4-1}{x^2+1} dx + 2 \int_0^1 \frac{dx}{x^2+1}$$

$$= \int_0^1 (x^2-1) dx + 2 \int_0^1 \frac{dx}{x^2+1}$$

$$= \left[\frac{x^3}{3} - x \right]_0^1 + \left[2 \tan^{-1} x \right]_0^1$$

$$= -\frac{2}{3} + \frac{\pi}{2}$$

$$= \frac{3\pi-4}{6}$$

$$27. \int_0^3 \frac{3x+1}{x^2+9} dx = \frac{3}{2} \int_0^3 \frac{2x}{x^2+9} dx + \int_0^3 \frac{dx}{x^2+9}$$

$$= \left[\frac{3}{2} \log(x^2+9) + \frac{1}{3} \tan^{-1} \left(\frac{x}{3} \right) \right]_0^3$$

$$= \frac{3}{2} (\log 18 - \log 9) + \frac{1}{3} \left(\frac{\pi}{4} \right)$$

$$= \frac{3}{2} \log 2 + \frac{\pi}{12} = \log(2\sqrt{2}) + \frac{\pi}{12}$$

$$28. \int_0^1 \frac{x^4(1-x)^4}{1+x^2} dx$$

$$= \int_0^1 \left(x^6 - 4x^5 + 5x^4 - 4x^3 + 4x - \frac{4}{1+x^2} \right) dx$$

$$= \left[\frac{x^7}{7} - \frac{2x^6}{3} + \frac{5x^5}{5} - \frac{4x^4}{4} + 4x - 4 \tan^{-1} x \right]_0^1$$

$$= \frac{1}{7} - \frac{2}{3} + 1 - \frac{4}{3} + 4 - 4 \left(\frac{\pi}{4} \right)$$

$$= \frac{22}{7} - \pi$$

$$29. \int_0^1 \sqrt{\frac{1-x}{1+x}} dx$$

$$= \int_0^1 \sqrt{\frac{1-x}{1+x}} \cdot \frac{1-x}{1-x} dx = \int_0^1 \frac{1-x}{\sqrt{1-x^2}} dx$$

$$= \int_0^1 \frac{dx}{\sqrt{1-x^2}} - \int_0^1 \frac{x}{\sqrt{1-x^2}} dx$$

$$= \left[\sin^{-1} x \right]_0^1 + \left[\sqrt{1-x^2} \right]_0^1$$

$$= \frac{\pi}{2} - 1$$

$$30. \int_{-1}^1 \sqrt{\frac{1-x}{1+x}} dx = \int_{-1}^1 \sqrt{\frac{1-x}{1+x}} \cdot \frac{1-x}{1-x} dx$$

$$= \int_{-1}^1 \frac{1-x}{\sqrt{1-x^2}} dx$$

$$= \int_{-1}^1 \frac{dx}{\sqrt{1-x^2}} - \int_{-1}^1 \frac{x}{\sqrt{1-x^2}} dx$$

$$= \left[\sin^{-1} x \right]_{-1}^1 + \left[\sqrt{1-x^2} \right]_{-1}^1$$

$$= \sin^{-1} 1 - \sin^{-1}(-1) + 0$$

$$= 2 \cdot \frac{\pi}{2} = \pi$$



$$\begin{aligned}
 31. \quad \text{Let } I &= \int_0^{\frac{\pi}{2}} \frac{1}{a^2 \cdot \sin^2 + b^2 \cdot \cos^2} dx \\
 &= \int_0^{\frac{\pi}{2}} \frac{1}{\cos^2 (a^2 \tan^2 + b^2)} dx \\
 &= \int_0^{\frac{\pi}{2}} \frac{\sec^2}{b^2 + a^2 \tan^2} dx
 \end{aligned}$$

$$\text{Put } a \tan = t \Rightarrow a \sec^2 dx = dt$$

$$\begin{aligned}
 \therefore I &= \frac{1}{a} \int_0^{\infty} \frac{dt}{b^2 + t^2} \\
 &= \frac{1}{ab} \left[\tan^{-1} \left(\frac{t}{b} \right) \right]_0^{\infty} = \frac{\pi}{2ab}
 \end{aligned}$$

$$32. \quad \text{Let } I = \int_0^{1/2} \frac{dx}{(1+x^2)\sqrt{1-x^2}}$$

$$\text{Put } x = \sin \theta \Rightarrow dx = \cos \theta d\theta$$

$$\begin{aligned}
 \therefore I &= \int_0^{\pi/6} \frac{\cos \theta d\theta}{(1 + \sin^2 \theta) \cos \theta} \\
 &= \int_0^{\pi/6} \frac{\sec^2 \theta}{\sec^2 \theta + \tan^2 \theta} d\theta \\
 &= \int_0^{\pi/6} \frac{\sec^2 \theta}{1 + 2 \tan^2 \theta} d\theta \\
 &= \int_0^{\pi/6} \frac{\sec^2 \theta}{1 + (\sqrt{2} \tan \theta)^2} d\theta \\
 &= \frac{1}{\sqrt{2}} \int_0^{\pi/6} \frac{\sqrt{2} \sec^2 \theta}{1 + (\sqrt{2} \tan \theta)^2} d\theta \\
 &= \frac{1}{\sqrt{2}} \left[\tan^{-1} (\sqrt{2} \tan \theta) \right]_0^{\pi/6} \\
 &= \frac{1}{\sqrt{2}} \tan^{-1} \sqrt{\frac{2}{3}}
 \end{aligned}$$

$$33. \quad \text{Put } \tan \frac{\alpha}{2} = t$$

$$\therefore dx = \frac{2dt}{1+t^2} \text{ and } \cos \alpha = \frac{1-t^2}{1+t^2}$$

$$\begin{aligned}
 \therefore \int_0^{\frac{\pi}{2}} \frac{dx}{2 + \cos \alpha} &= \int_0^1 \frac{2dt}{3+t^2} = \left[\frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{t}{\sqrt{3}} \right) \right]_0^1 \\
 &= \frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{1}{\sqrt{3}} \right)
 \end{aligned}$$

$$34. \quad \text{Put } \tan \frac{\alpha}{2} = t$$

$$\therefore dx = \frac{2dt}{1+t^2} \text{ and } \cos \alpha = \frac{1-t^2}{1+t^2}$$

$$\begin{aligned}
 \therefore \int_0^{\frac{\pi}{2}} \frac{dx}{5+4\cos \alpha} &= \int_0^{\infty} \frac{2dt}{9+t^2} \\
 &= \left[\frac{2}{3} \tan^{-1} \left(\frac{t}{3} \right) \right]_0^{\infty} \\
 &= \frac{2}{3} (\tan^{-1} \infty - 0) \\
 &= \frac{2}{3} \cdot \frac{\pi}{2} = \frac{\pi}{3}
 \end{aligned}$$

$$35. \quad \int_0^{\frac{\pi}{2}} \frac{\cos \alpha}{1 + \cos \alpha + \sin \alpha} dx$$

$$= \int_0^{\frac{\pi}{2}} \frac{\cos^2(\alpha/2) - \sin^2(\alpha/2)}{2\cos^2(\alpha/2) + 2\sin(\alpha/2)\cos(\alpha/2)} dx$$

$$= \frac{1}{2} \int_0^{\frac{\pi}{2}} \frac{1 - \tan^2(\alpha/2)}{1 + \tan(\alpha/2)} dx$$

$$= \frac{1}{2} \int_0^{\frac{\pi}{2}} \left[1 - \tan \left(\frac{\alpha}{2} \right) \right] dx$$

$$= \frac{1}{2} \left[x + 2 \log \left| \cos \left(\frac{\alpha}{2} \right) \right| \right]_0^{\frac{\pi}{2}}$$

$$= \frac{\pi}{4} + \log \frac{1}{\sqrt{2}} = \frac{\pi}{4} - \frac{1}{2} \log 2$$

$$36. \quad \int_0^1 \frac{dx}{x^2 + 2 \cos \alpha + 1} = \int_0^1 \frac{dx}{(x + \cos \alpha)^2 + 1 - \cos^2 \alpha}$$

$$= \int_0^1 \frac{dx}{(x + \cos \alpha)^2 + \sin^2 \alpha}$$

$$= \left[\frac{1}{\sin \alpha} \tan^{-1} \left(\frac{x + \cos \alpha}{\sin \alpha} \right) \right]_0^1$$

$$= \frac{1}{\sin \alpha} \left[\tan^{-1} \left(\cot \frac{\alpha}{2} \right) - \tan^{-1} (\cot \alpha) \right]$$

$$= \frac{1}{\sin \alpha} \left[\tan^{-1} \left(\tan \left(\frac{\pi}{2} - \frac{\alpha}{2} \right) \right) - \tan^{-1} \left(\tan \left(\frac{\pi}{2} - \alpha \right) \right) \right]$$

$$= \frac{\alpha}{2} (\sin \alpha)^{-1}$$



$$\begin{aligned}
 37. \quad \text{Let } I &= \int_0^{\pi/4} (\sqrt{\tan x} + \sqrt{\cot x}) dx \\
 &= \int_0^{\pi/4} \frac{\sin x + \cos x}{\sqrt{\sin x \cos x}} dx \\
 &= \sqrt{2} \int_0^{\pi/4} \frac{\sin x + \cos x}{\sqrt{1 - (\sin x - \cos x)^2}} dx
 \end{aligned}$$

$$\text{Put } \sin x - \cos x = t \Rightarrow (\cos x + \sin x) dx = dt$$

$$\begin{aligned}
 \therefore I &= \sqrt{2} \int_{-1}^0 \frac{dt}{\sqrt{1-t^2}} \\
 &= \sqrt{2} [\sin^{-1} t]_{-1}^0 \\
 &= \sqrt{2} \left[0 - \left(-\frac{\pi}{2} \right) \right] = \frac{\pi}{\sqrt{2}}
 \end{aligned}$$

$$38. \quad \int_{\log 2}^a \frac{e^x}{\sqrt{e^x - 1}} dx = 2$$

$$\text{Put } e^x - 1 = t \Rightarrow e^x dx = dt$$

$$\begin{aligned}
 \therefore \int_1^{e^a - 1} \frac{dt}{\sqrt{t}} &= 2 \\
 \Rightarrow [2\sqrt{t}]_1^{e^a - 1} &= 2 \\
 \Rightarrow \sqrt{e^a - 1} - 1 &= 1 \\
 \Rightarrow \sqrt{e^a - 1} &= 2 \\
 \Rightarrow e^a - 1 &= 4 \\
 \Rightarrow e^a &= 5 \\
 \Rightarrow a &= \log 5
 \end{aligned}$$

$$39. \quad \int_{\log 2}^u \frac{du}{(e^u - 1)^{1/2}} = \frac{\pi}{6}$$

$$\Rightarrow \int_{\log 2}^u \frac{e^u}{e^u (e^u - 1)^{1/2}} du = \frac{\pi}{6}$$

$$\text{Put } e^u - 1 = t^2 \Rightarrow e^u du = 2t dt$$

$$\text{When } u = \log 2, t = 1$$

$$\text{and when } u = u, t = \sqrt{e^u - 1}$$

$$\begin{aligned}
 \therefore \int_1^{\sqrt{e^u - 1}} \frac{2}{1+t^2} dt &= \frac{\pi}{6} \\
 \Rightarrow 2[\tan^{-1} t]_1^{\sqrt{e^u - 1}} &= \frac{\pi}{6} \\
 \Rightarrow \tan^{-1}(\sqrt{e^u - 1}) - \frac{\pi}{4} &= \frac{\pi}{12} \\
 \Rightarrow \sqrt{e^u - 1} = \tan \frac{\pi}{3} &\Rightarrow \sqrt{e^u - 1} = \sqrt{3} \Rightarrow e^u = 4
 \end{aligned}$$

$$\begin{aligned}
 40. \quad \frac{\pi^2}{\log 3} \int_{\frac{7}{6}}^{\frac{5}{6}} \sec(\pi x) dx \\
 = \frac{\pi^2}{\log 3} \times \frac{1}{\pi} [\log |\sec \pi x + \tan \pi x|]_{7/6}^{5/6} \\
 = \frac{\pi}{\log 3} \left[\log \left| \sec \frac{5\pi}{6} + \tan \frac{5\pi}{6} \right| \right. \\
 \left. - \log \left| \sec \frac{7\pi}{6} + \tan \frac{7\pi}{6} \right| \right]
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{\pi}{\log 3} \left[\log \left(\frac{2}{\sqrt{3}} + \frac{1}{\sqrt{3}} \right) - \log \left(\frac{2}{\sqrt{3}} - \frac{1}{\sqrt{3}} \right) \right] \\
 &= \frac{\pi}{\log 3} \left[\log \sqrt{3} - \log \left(\frac{1}{\sqrt{3}} \right) \right] = \frac{\pi}{\log 3} (\log 3) = \pi
 \end{aligned}$$

$$41. \quad \text{Put } x = \cos \theta \Rightarrow dx = -\sin \theta d\theta$$

$$\begin{aligned}
 \therefore \int_0^1 \sin \left(2 \tan^{-1} \sqrt{\frac{1+x}{1-x}} \right) dx \\
 = -\int_{\frac{\pi}{2}}^0 \sin \left(2 \tan^{-1} \sqrt{\frac{1+\cos \theta}{1-\cos \theta}} \right) \cdot \sin \theta d\theta \\
 = -\int_{\frac{\pi}{2}}^0 \sin \left[2 \tan^{-1} \left(\cot \frac{\theta}{2} \right) \right] \cdot \sin \theta d\theta \\
 = -\int_{\frac{\pi}{2}}^0 \sin \left[2 \tan^{-1} \left\{ \tan \left(\frac{\pi}{2} - \frac{\theta}{2} \right) \right\} \right] \cdot \sin \theta d\theta \\
 = -\int_{\frac{\pi}{2}}^0 \sin \left[2 \left(\frac{\pi}{2} - \frac{\theta}{2} \right) \right] \cdot \sin \theta d\theta \\
 = -\int_{\frac{\pi}{2}}^0 \sin(\pi - \theta) \cdot \sin \theta d\theta \\
 = -\int_{\frac{\pi}{2}}^0 \sin \theta \cdot \sin \theta d\theta \\
 = -\int_{\frac{\pi}{2}}^0 \left(\frac{1 - \cos 2\theta}{2} \right) d\theta \\
 = -\frac{1}{2} \left[\theta - \frac{\sin 2\theta}{2} \right]_{\frac{\pi}{2}}^0 \\
 = \frac{\pi}{4}
 \end{aligned}$$



$$\begin{aligned}
 42. \int_0^{\frac{\pi}{4}} \sec^2 x \, dx &= \left[\tan x \right]_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} \tan x \, dx \\
 &= \left[\frac{\pi}{4} - 0 \right] - \left[\log |\sec x| \right]_0^{\frac{\pi}{4}} \\
 &= \frac{\pi}{4} - \log \left| \sec \frac{\pi}{4} \right| + \log |\sec 0| \\
 &= \frac{\pi}{4} - \log \sqrt{2} + \log 1 \\
 &= \frac{\pi}{4} - \log \sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 43. \text{ Let } I &= \int_0^{\frac{\pi}{2}} e^{\sin x} \sin x \, dx \\
 &= \left[\sin x e^{\sin x} \right]_0^{\frac{\pi}{2}} - \int_0^{\frac{\pi}{2}} \cos x \cdot e^{\sin x} \, dx \\
 \therefore I &= \left[e^{\sin x} \sin x \right]_0^{\frac{\pi}{2}} - \left[\cos x e^{\sin x} \right]_0^{\frac{\pi}{2}} - \int_0^{\frac{\pi}{2}} \sin x \cdot e^{\sin x} \, dx \\
 \Rightarrow 2I &= \left[e^{\sin x} (\sin x - \cos x) \right]_0^{\frac{\pi}{2}} \\
 \Rightarrow 2I &= e^{\pi/2} + 1 \\
 \Rightarrow I &= \frac{e^{\pi/2} + 1}{2}
 \end{aligned}$$

$$\begin{aligned}
 44. \text{ Let } I &= \int_a^b \frac{1}{x} \log x \, dx \\
 \Rightarrow I &= \left[\log x \log x \right]_a^b - \int_a^b \frac{1}{x} \log x \, dx \\
 \Rightarrow 2I &= \left[(\log x)^2 \right]_a^b \\
 \Rightarrow I &= \frac{1}{2} \left[(\log b)^2 - (\log a)^2 \right] \\
 &= \frac{1}{2} \left[(\log b + \log a)(\log b - \log a) \right] \\
 &= \frac{1}{2} \log(ab) \log \left(\frac{b}{a} \right)
 \end{aligned}$$

$$\begin{aligned}
 45. \text{ Let } I &= \int_0^1 \tan^{-1} x \, dx \\
 &= \tan^{-1} x \int_0^1 dx - \int_0^1 \frac{1}{1+x^2} \cdot \frac{1}{2} dx \\
 &= \left[\frac{1}{2} \tan^{-1} x \right]_0^1 - \frac{1}{2} \int_0^1 \frac{1+x^2-1}{1+x^2} dx
 \end{aligned}$$

$$\begin{aligned}
 &= \left(\frac{\pi}{4} \cdot \frac{1}{2} - 0 \right) - \frac{1}{2} \left[-\tan^{-1} x \right]_0^1 \\
 &= \frac{\pi}{8} - \frac{1}{2} \left[(1-0) - \left(\frac{\pi}{4} - 0 \right) \right] \\
 &= \frac{\pi}{8} - \frac{1}{2} + \frac{\pi}{8} = \frac{\pi}{4} - \frac{1}{2}
 \end{aligned}$$

$$\begin{aligned}
 46. \text{ Let } I &= \int_0^{\frac{\pi}{2}} \frac{\sin^{-1} x}{(1-x^2)^{\frac{3}{2}}} dx \\
 \text{Put } \sin^{-1} x &= t \Rightarrow \frac{1}{\sqrt{1-x^2}} dx = dt \\
 \text{When } x &= 0, t = 0 \text{ and when } x = \frac{1}{\sqrt{2}}, t = \frac{\pi}{4}
 \end{aligned}$$

$$\therefore I = \int_0^{\pi/4} t \cdot \sec^2 t \, dt = \frac{\pi}{4} - \frac{1}{2} \log 2$$

$$47. \text{ Put } t = \sin^{-1} x \Rightarrow dt = \frac{1}{\sqrt{1-x^2}} dx$$

$$\begin{aligned}
 \therefore \int_0^{1/2} \frac{\sin^{-1} x}{\sqrt{1-x^2}} dx &= \int_0^{\pi/6} t \sin t \, dt \\
 &= \left[-t \cos t + \sin t \right]_0^{\pi/6} \\
 &= -\frac{\pi}{6} \cdot \frac{\sqrt{3}}{2} + \frac{1}{2} = \frac{1}{2} - \frac{\sqrt{3}\pi}{12}
 \end{aligned}$$

$$\begin{aligned}
 48. \text{ Let } I &= \int_{\pi/4}^{3\pi/4} \frac{1}{1+\sin x} dx \\
 &= \int_{\pi/4}^{3\pi/4} \frac{\sec x}{\sec x + \tan x} dx
 \end{aligned}$$

$$\text{Let } I_1 = \int \frac{\sec x}{\sec x + \tan x} dx$$

$$\text{Put } \frac{1}{\sec x + \tan x} = t$$

$$\Rightarrow - \frac{(\sec x \tan x + \sec^2 x)}{(\sec x + \tan x)^2} dx = dt$$

$$\begin{aligned}
 \therefore I_1 &= - \int \frac{-\sec x (\sec x + \tan x)}{(\sec x + \tan x)^2} dx \\
 &= - \int dt \\
 &= -t + c \\
 &= \frac{-1}{\sec x + \tan x} + c
 \end{aligned}$$



$$\begin{aligned} \therefore I &= \left[\frac{-1}{\sec \theta + \tan \theta} \right]_{\pi/4}^{3\pi/4} - \int_{\pi/4}^{3\pi/4} \frac{-1}{\sec \theta + \tan \theta} d\theta \\ &= \left[\int_a^b (uv) d\theta = \left[u \int v d\theta \right]_a^b - \int_a^b \left[\frac{du}{d\theta} \int v d\theta \right] d\theta \right] \\ &= \left(\frac{-3\pi}{4} \right) - \left(\frac{-\pi}{4} \right) + \int_{\pi/4}^{3\pi/4} \frac{\cos \theta}{1 + \sin \theta} d\theta \\ &= \frac{\pi}{1 + \sqrt{2}} + \left[\log |1 + \sin \theta| \right]_{\pi/4}^{3\pi/4} \\ &= \frac{\pi}{1 + \sqrt{2}} + \log \left| 1 + \frac{1}{\sqrt{2}} \right| - \log \left| 1 + \frac{1}{\sqrt{2}} \right| \\ &= \frac{\pi}{1 + \sqrt{2}} \\ &= \pi(\sqrt{2} - 1) \end{aligned}$$

$$\begin{aligned} 49. \quad F(t) &= \int_0^t f(t-x)g(x) dx \\ &= \int_0^t e^{-t+x} dx = e^{-t} \int_0^t e^x dx \\ &= -e^{-t} [e^{-x} + e^{-x}]_0^t \\ &= -e^{-t}(te^{-t} + e^{-t} - 0 - 1) \\ &= e^{-t} - (1+t) \end{aligned}$$

$$\begin{aligned} 50. \quad \text{Let } I &= \int_0^{2\pi} e^{\frac{x}{2}} \cdot \sin\left(\frac{x}{2} + \frac{\pi}{4}\right) dx \\ \text{Put } \frac{x}{2} &= t \\ \Rightarrow dx &= 2dt \\ \therefore I &= 2 \int_0^{\pi} e^t \sin\left(t + \frac{\pi}{4}\right) dt \\ &= 2 \left[\frac{e^t}{\sqrt{1+1}} \sin\left(t + \frac{\pi}{4} - \tan^{-1} \frac{1}{1}\right) \right]_0^{\pi} \\ &= \left[\int e^a \sin b dx = \frac{e^a}{\sqrt{a^2 + b^2}} \sin\left(b - \tan^{-1} \frac{b}{a}\right) + c \right] \\ &= \frac{2}{\sqrt{2}} [e^t \sin t]_0^{\pi} \\ &= \frac{2}{\sqrt{2}} [0] = 0 \end{aligned}$$

$$\begin{aligned} 51. \quad I_{10} &= \int_0^{\frac{\pi}{2}} 10 \sin^9 \theta d\theta \\ &= \left[-10 \cos \theta \right]_0^{\pi/2} - 10 \int_0^{\pi/2} 9 \cos^8 \theta (-\cos \theta) d\theta \\ &= \left[-\left(\frac{\pi}{2}\right)^{10} \cos \frac{\pi}{2} + 0 \right] \\ &\quad + 10 \left[\int_0^{\pi/2} 9 \sin^8 \theta d\theta - \int_0^{\pi/2} 9^8 (\sin^8 \theta) d\theta \right] \\ &= 10 \left[9 \sin^7 \theta \right]_0^{\pi/2} - 90 \int_0^{\pi/2} 8 \sin^6 \theta d\theta \end{aligned}$$

$$\begin{aligned} \therefore I_{10} &= 10 \left(\frac{\pi}{2}\right)^9 \sin \frac{\pi}{2} - 90 I_8 \\ \Rightarrow I_{10} + 90 I_8 &= 10 \left(\frac{\pi}{2}\right)^9 \end{aligned}$$

$$\begin{aligned} 52. \quad \int_0^1 \log\left(1 + \frac{x}{2}\right) dx &= \left[\log\left(1 + \frac{x}{2}\right) \cdot \frac{x}{2} - \int_0^1 \frac{1}{1 + \frac{x}{2}} \cdot \frac{1}{2} \cdot \frac{x}{2} dx \right] \\ &= \frac{1}{2} \log \frac{3}{2} - \frac{1}{2} \int_0^1 \frac{x}{x+2} dx \\ &= \frac{1}{2} \log \frac{3}{2} - \frac{1}{2} \int_0^1 \left[-\frac{2}{x+2} \right] dx \\ &= \frac{1}{2} \log \frac{3}{2} - \frac{1}{2} \left[-\frac{2}{x+2} \right]_0^1 + \int_0^1 \frac{1}{x+2} dx \\ &= \frac{1}{2} \log \frac{3}{2} - \frac{1}{4} + \int_0^1 \left[1 - \frac{2}{x+2} \right] dx \\ &= \frac{1}{2} \log \frac{3}{2} - \frac{1}{4} + \left[-2 \log(x+2) \right]_0^1 \\ &= \frac{1}{2} \log \frac{3}{2} - \frac{1}{4} + 1 - 2 \log 3 + 2 \log 2 \\ &= \frac{3}{4} + \frac{3}{2} \log \frac{2}{3} \\ \therefore a &= \frac{3}{4}, b = \frac{3}{2} \end{aligned}$$



$$53. \quad I(m, n) = \int_0^1 t^m (1-t)^n dt$$

$$\therefore I(m+1, n-1) = \int_0^1 t^{m+1} (1-t)^{n-1} dt$$

$$\Rightarrow I(m+1, n-1) = \left[-\frac{t^{m+1} (1-t)^n}{n} \right]_0^1 + \frac{m+1}{n} \int_0^1 t^m (1-t)^n dt$$

$$\Rightarrow I(m+1, n-1) = 0 + \frac{m+1}{n} I(m, n)$$

$$\Rightarrow I(m, n) = \frac{n}{m+1} I(m+1, n-1)$$

$$54. \quad \text{Let } I_1 = \int_0^1 (1-50)^{100} dx \quad \text{and } I_2 = \int_0^1 (1-50)^{101} dx$$

Now, $I_2 = \int_0^1 (1-50)^{101} \cdot 1 dx$

$$= \left[(1-50)^{101} \cdot x \right]_0^1 + 5050 \int_0^1 (1-50)^{100} \cdot 49 dx$$

$$= -5050 \int_0^1 (1-50)^{100} \{ (1-50) - 1 \} dx$$

$$= -5050 \int_0^1 (1-50)^{101} dx + 5050 \int_0^1 (1-50)^{100} dx$$

$$\therefore I_2 = -5050 I_2 + 5050 I_1$$

$$\Rightarrow \frac{5050 I_1}{I_2} = 5051$$

$$55. \quad \int_5^{10} \frac{1}{(x-1)(x-2)} dx = \int_5^{10} \left(\frac{1}{x-2} - \frac{1}{x-1} \right) dx$$

$$= [\log(x-2) - \log(x-1)]_5^{10}$$

$$= \log 8 - \log 9 - (\log 3 - \log 4)$$

$$= \log \frac{8}{9} - \log \frac{3}{4} = \log \left(\frac{8}{9} \times \frac{4}{3} \right)$$

$$= \log \left(\frac{32}{27} \right)$$

$$56. \quad \int_1^3 \frac{dx}{(1+x^2)} = \int_1^3 \left(\frac{1}{1+x^2} \right) dx$$

$$= \int_1^3 \frac{1}{1+x^2} dx - \frac{1}{2} \int_1^3 \frac{2}{1+x^2} dx$$

$$= [\log |x|]_1^3 - \frac{1}{2} [\log(1+x^2)]_1^3$$

$$= \log 3 - \log 1 - \frac{1}{2} (\log 10 - \log 2)$$

$$= \log 3 - \frac{1}{2} \log 5$$

$$= \frac{1}{2} \log 3^2 - \frac{1}{2} \log 5$$

$$= \frac{1}{2} (\log 9 - \log 5) = \frac{1}{2} \log \left(\frac{9}{5} \right)$$

$$57. \quad \int_2^3 \frac{x+1}{x^2(-1)} dx = \int_2^3 \left(-\frac{1}{x} - \frac{2}{x} + \frac{2}{-1} \right) dx$$

$$= \left[-\frac{1}{x} \right]_2^3 - 2 [\log x]_2^3 + 2 [\log(-1)]_2^3$$

$$= \frac{1}{3} - \frac{1}{2} - 2 \log \frac{3}{2} + 2 \log 2$$

$$= \log \frac{16}{9} - \frac{1}{6}$$

$$58. \quad \text{Let } I = \int_0^{\frac{\pi}{4}} \frac{\sin x + \cos x}{9 + 16 \sin^2 x} dx$$

$$= \int_0^{\frac{\pi}{4}} \frac{\sin x + \cos x}{9 + 16 [1 - (\sin^2 x - \cos^2 x)]} dx$$

Put $\sin^2 x - \cos^2 x = t$
 $\Rightarrow (\cos^2 x + \sin^2 x) dx = dt$

When $x = 0$, $t = -1$ and when $x = \frac{\pi}{4}$, $t = 0$

$$\therefore I = \int_{-1}^0 \frac{dt}{9 + 16(1-t^2)}$$

$$= \int_{-1}^0 \frac{1}{25 - 16t^2} dt$$

$$= \int_{-1}^0 \frac{1}{(5-4t)(5+4t)} dt$$

$$= \int_{-1}^0 \left[\frac{\frac{1}{10}}{5-4t} + \frac{\frac{1}{10}}{5+4t} \right] dt$$

$$= \frac{1}{10} \left[-\frac{1}{4} \log(5-4t) + \frac{1}{4} \log(5+4t) \right]_{-1}^0$$

$$= \frac{1}{40} (\log 9 - \log 1)$$

$$= \frac{1}{20} \log 3$$



$$\begin{aligned}
 59. \quad & \int_0^2 \frac{\log(x^2+2)}{(x+2)^2} dx \\
 &= -\left[\frac{\log(x^2+2)}{x+2} \right]_0^2 + \int_0^2 \frac{2}{(x^2+2)(x+2)} dx \\
 &= -\frac{1}{4} \log 6 + \frac{1}{2} \log 2 + \int_0^2 \left\{ \frac{-2}{3(x+2)} + \frac{\frac{2}{3} + \frac{2}{3}}{x^2+2} \right\} dx \\
 &= -\frac{1}{4} \log 3 - \frac{1}{4} \log 2 + \frac{1}{2} \log 2 \\
 &+ \left[-\frac{2}{3} \log(x+2) + \frac{1}{3} \log(x^2+2) + \frac{\sqrt{2}}{3} \tan^{-1} \left(\frac{x}{\sqrt{2}} \right) \right]_0^2 \\
 &= \frac{1}{4} \log 2 - \frac{1}{4} \log 3 \\
 &\quad + \left(-\frac{2}{3} \log 2 + \frac{1}{3} \log 3 + \frac{\sqrt{2}}{3} \tan^{-1} \sqrt{2} \right) \\
 &= \frac{\sqrt{2}}{3} \tan^{-1} \sqrt{2} - \frac{5}{12} \log 2 + \frac{1}{12} \log 3
 \end{aligned}$$

$$60. \quad \text{Put } \sin^{-1} \left(\frac{x}{2} \right) = t \Rightarrow x = 2 \sin t \Rightarrow dx = 2 \cos t dt$$

$$\begin{aligned}
 \therefore \int_0^1 \frac{\sin^{-1} \left(\frac{x}{2} \right)}{x} dx &= \int_0^{\frac{\pi}{6}} \frac{t}{2 \sin t} (2 \cos t dt) \\
 &= \int_0^{\frac{\pi}{6}} \frac{t}{\tan t} dt = \int_0^{\frac{\pi}{6}} \frac{t}{\tan t} dt \\
 &\quad \dots \left[\because \int_a^b f(x) dx = \int_a^b f(t) dt \right]
 \end{aligned}$$

$$61. \quad I_1 = \int_e^{e^2} \frac{dx}{\log x}$$

$$\text{Put } \log x = t$$

$$\Rightarrow dx = e^t dt = e^t dt$$

$$\text{When } x = e, t = 1 \text{ and when } x = e^2, t = 2$$

$$\begin{aligned}
 \therefore I_1 &= \int_1^2 \frac{e^t}{t} dt \\
 &= \int_1^2 \frac{e}{t} dt \quad \dots \left[\because \int_a^b f(x) dx = \int_a^b f(t) dt \right]
 \end{aligned}$$

$$\therefore I_1 = I_2$$

$$\begin{aligned}
 62. \quad & \int_0^3 (3a^2 + 2b^2 + c) dx = \int_1^3 (3a^2 + 2b^2 + c) dx \\
 & \Rightarrow \int_0^1 (3a^2 + 2b^2 + c) dx + \int_1^3 (3a^2 + 2b^2 + c) dx \\
 & \qquad \qquad \qquad = \int_1^3 (3a^2 + 2b^2 + c) dx
 \end{aligned}$$

$$\Rightarrow \int_0^1 (3a^2 + 2b^2 + c) dx = 0$$

$$\Rightarrow \left[\frac{3a^3}{3} + \frac{2b^2}{2} + cx \right]_0^1 = 0 \Rightarrow a + b + c = 0$$

$$63. \quad \int_2^4 (3 - f(x)) dx = 7$$

$$\Rightarrow \int_2^4 3 dx - \int_2^4 f(x) dx = 7$$

$$\Rightarrow 3 \left[x \right]_2^4 - 7 = \int_2^4 f(x) dx$$

$$\Rightarrow \int_2^4 f(x) dx = 3(4-2) - 7$$

$$= -1$$

$$\int_{-1}^4 f(x) dx = \int_{-1}^2 f(x) dx + \int_2^4 f(x) dx$$

$$\dots \left[\because \int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx \right. \\ \left. \dots \left[\text{if } a < c < b \right] \right]$$

$$\Rightarrow 4 = \int_{-1}^2 f(x) dx - 1$$

$$\Rightarrow \int_{-1}^2 f(x) dx = 5$$

$$64. \quad g(x + \pi) = \int_0^{x+\pi} \cos^4 t dt$$

$$= \int_0^{\pi} \cos^4 t dt + \int_{\pi}^{x+\pi} \cos^4 t dt$$

$$\text{In 2nd integral, put } t = u + \pi \Rightarrow dt = du$$

$$\therefore \int_{\pi}^{x+\pi} \cos^4 t dt = \int_0^x \cos^4(\pi + u) du$$

$$= \int_0^x \cos^4 u du = g(x)$$

$$\therefore g(x + \pi) = g(x) + g(\pi)$$



65. Since, $|-1| = -(-1)$, if $-1 < 0$ i.e., $x < 1$
 $= -1$, if $-1 \geq 0$ i.e., $x \geq 1$

$$\begin{aligned} \therefore \int_0^2 f(x) dx &= \int_0^1 |-1| dx + \int_1^2 (-1) dx \\ &= \int_0^1 (1-x) dx + \int_1^2 (-x) dx \\ &= \left[\frac{x^2}{2} - x \right]_0^1 + \left[-\frac{x^2}{2} \right]_1^2 \\ &= \left(\frac{1}{2} - 1 \right) + \left(-2 + \frac{1}{2} \right) = -1 \end{aligned}$$

$$\begin{aligned} 66. \int_{-5}^5 |x+2| dx &= -\int_{-5}^{-2} (x+2) dx + \int_{-2}^5 (x+2) dx \\ &= \left[\frac{x^2}{2} + 2x \right]_{-5}^{-2} + \left[\frac{x^2}{2} + 2x \right]_{-2}^5 \\ &= 29 \end{aligned}$$

$$\begin{aligned} 67. f(x) &= \int_{-1}^x |t| dt \\ &= \int_{-1}^0 (-t) dt + \int_0^x t dt \\ &= \left[-\frac{t^2}{2} \right]_{-1}^0 + \left[\frac{t^2}{2} \right]_0^x = \frac{1}{2} + \frac{x^2}{2} = \frac{1}{2}(1+x^2) \end{aligned}$$

$$\begin{aligned} 68. \int_0^1 |3x^2-1| dx &= \int_0^{\frac{1}{\sqrt{3}}} (1-3x^2) dx + \int_{\frac{1}{\sqrt{3}}}^1 (3x^2-1) dx \\ &= \left[x - x^3 \right]_0^{\frac{1}{\sqrt{3}}} + \left[x^3 - x \right]_{\frac{1}{\sqrt{3}}}^1 \\ &= \frac{1}{\sqrt{3}} - \frac{1}{3\sqrt{3}} - \frac{1}{3\sqrt{3}} + \frac{1}{\sqrt{3}} = \frac{4}{3\sqrt{3}} \end{aligned}$$

$$\begin{aligned} 69. \int_{-2}^2 |1-x^2| dx &= \int_{-2}^{-1} |1-x^2| dx + \int_{-1}^1 |1-x^2| dx + \int_1^2 |1-x^2| dx \\ &= -\int_{-2}^{-1} (1-x^2) dx + \int_{-1}^1 (1-x^2) dx - \int_1^2 (1-x^2) dx \\ &= -\left[x - \frac{x^3}{3} \right]_{-2}^{-1} + \left[x - \frac{x^3}{3} \right]_{-1}^1 - \left[x - \frac{x^3}{3} \right]_1^2 \\ &= \frac{4}{3} + \frac{4}{3} + \frac{4}{3} = 4 \end{aligned}$$

70. Since, $\left| -\frac{1}{2} \right| = -\left(-\frac{1}{2} \right)$, if $x < \frac{1}{2}$
 $= -\frac{1}{2}$, if $x \geq \frac{1}{2}$

$$\begin{aligned} \therefore \int_0^1 \left| -\frac{1}{2} \right| dx &= -\int_0^{\frac{1}{2}} \left(-\frac{1}{2} \right) dx + \int_{\frac{1}{2}}^1 \left(-\frac{1}{2} \right) dx \\ &= \frac{1}{2} \int_0^{\frac{1}{2}} \left(\frac{1}{2} - x^2 \right) dx + \int_{\frac{1}{2}}^1 \left(x^2 - \frac{1}{2} \right) dx \\ &= \left[\frac{x^2}{4} - \frac{x^3}{3} \right]_0^{\frac{1}{2}} + \left[\frac{x^3}{3} - \frac{x}{2} \right]_{\frac{1}{2}}^1 \\ &= \left(\frac{1}{16} - \frac{1}{24} \right) + \left(\frac{1}{3} - \frac{1}{4} + \frac{1}{16} - \frac{1}{24} \right) = \frac{1}{8} \end{aligned}$$

$$\begin{aligned} 71. \text{ Let } I &= \int_0^{100\pi} |\cos x| dx \\ &= 200 \int_0^{\frac{\pi}{2}} |\cos x| dx \\ \dots \because \int_0^{2a} f(x) dx &= 2 \int_0^a f(x) dx, \text{ if } f(2a-x) = f(x) \end{aligned}$$

Since $\cos x$ is positive in the interval $\left(0, \frac{\pi}{2} \right)$

$$\begin{aligned} \therefore I &= 200 \int_0^{\frac{\pi}{2}} \cos x dx \\ &= 200 \left[\sin x \right]_0^{\frac{\pi}{2}} \\ &= 200 \end{aligned}$$

$$\begin{aligned} 72. \int_0^{\pi/2} |\sin x - \cos x| dx &= -\int_0^{\pi/4} (\sin x - \cos x) dx + \int_{\pi/4}^{\pi/2} (\sin x - \cos x) dx \\ &= -\left[-\cos x - \sin x \right]_0^{\pi/4} + \left[-\cos x - \sin x \right]_{\pi/4}^{\pi/2} \\ &= 2(\sqrt{2}-1) \end{aligned}$$



$$\begin{aligned}
73. \quad & \int_0^{\pi} \sqrt{1 + 4\sin^2 \frac{x}{2} - 4\sin \frac{x}{2}} dx \\
&= \int_0^{\pi} \left| 2\sin \frac{x}{2} - 1 \right| dx \\
&= \int_0^{\frac{\pi}{3}} \left| 2\sin \frac{x}{2} - 1 \right| dx + \int_{\frac{\pi}{3}}^{\pi} \left| 2\sin \frac{x}{2} - 1 \right| dx \\
&= \int_0^{\frac{\pi}{3}} \left(1 - 2\sin \frac{x}{2} \right) dx + \int_{\frac{\pi}{3}}^{\pi} \left(2\sin \frac{x}{2} - 1 \right) dx \\
&= \left[x + 4\cos \frac{x}{2} \right]_0^{\frac{\pi}{3}} + \left[-4\cos \frac{x}{2} - x \right]_{\frac{\pi}{3}}^{\pi} \\
&= \frac{\pi}{3} + 4 \left(\frac{\sqrt{3}}{2} - 1 \right) + \left[-4 \left(0 - \frac{\sqrt{3}}{2} \right) - \left(\pi - \frac{\pi}{3} \right) \right] \\
&= 4\sqrt{3} - 4 - \frac{\pi}{3}
\end{aligned}$$

74. Since, $|\log x| = -\log x$, if $\frac{1}{e} < x < 1$
 $= \log x$, if $1 < x < e$

$$\begin{aligned}
\therefore I &= \int_{1/e}^1 \frac{(-\log x)}{2} dx + \int_1^e \frac{\log x}{2} dx \\
&= - \left[-\frac{\log x}{2} - \frac{1}{2} \right]_{1/e}^1 + \left[\frac{\log x}{2} - \frac{1}{2} \right]_1^e \\
&= \left[0 + 1 - \left(\frac{\log \frac{1}{e}}{2} + \frac{1}{2} \right) \right] - \left[\frac{\log e}{2} + \frac{1}{2} - (0 + 1) \right] \\
&= 1 - (-e + e) - \left(\frac{2}{e} - 1 \right) \\
&= 2 - \frac{2}{e} = 2 \left(1 - \frac{1}{e} \right)
\end{aligned}$$

75. Let $I = \int_0^3 [x] dx$

$$\begin{aligned}
&= \int_0^1 0 dx + \int_1^2 1 dx + \int_2^3 2 dx \\
&= [x]_0^1 + 2[x]_1^2 \\
&= (2 - 1) + 2(3 - 2) \\
&= 3
\end{aligned}$$

$$\begin{aligned}
76. \quad & \int_{-1}^1 (x - [x]) dx = \int_{-1}^0 (x - [x]) dx + \int_0^1 (x - [x]) dx \\
&= \int_{-1}^0 (x + 1) dx + \int_0^1 (x - 0) dx \\
&= \left[\frac{(x+1)^2}{2} \right]_{-1}^0 + \left[\frac{x^2}{2} \right]_0^1 \\
&= \frac{1}{2} + \frac{1}{2} = 1
\end{aligned}$$

$$\begin{aligned}
77. \quad & \int_0^{1.5} [x^2] dx = \int_0^1 [x^2] dx + \int_1^{\sqrt{2}} [x^2] dx + \int_{\sqrt{2}}^{1.5} [x^2] dx \\
&= \int_0^1 0 dx + \int_1^{\sqrt{2}} 1 dx + \int_{\sqrt{2}}^{1.5} 2 dx \\
&= \sqrt{2} - 1 + 3 - 2\sqrt{2} = 2 - \sqrt{2}
\end{aligned}$$

$$\begin{aligned}
78. \quad & \int_0^9 [\sqrt{x} + 2] dx \\
&= \int_0^1 [\sqrt{x} + 2] dx + \int_1^4 [\sqrt{x} + 2] dx + \int_4^9 [\sqrt{x} + 2] dx \\
&= \int_0^1 2 dx + \int_1^4 3 dx + \int_4^9 4 dx \\
&= 2 + (12 - 3) + (36 - 16) = 2 + 9 + 20 = 31
\end{aligned}$$

$$\begin{aligned}
79. \quad & \int_{\frac{\pi}{2}}^{\frac{3\pi}{2}} [2\sin x] dx = \int_{\frac{\pi}{2}}^{\frac{5\pi}{6}} [2\sin x] dx + \int_{\frac{5\pi}{6}}^{\pi} [2\sin x] dx \\
&\quad + \int_{\pi}^{\frac{7\pi}{6}} [2\sin x] dx + \int_{\frac{7\pi}{6}}^{\frac{3\pi}{2}} [2\sin x] dx \\
&= \int_{\frac{\pi}{2}}^{\frac{5\pi}{6}} 1 dx + \int_{\frac{5\pi}{6}}^{\pi} 0 dx + \int_{\pi}^{\frac{7\pi}{6}} (-1) dx + \int_{\frac{7\pi}{6}}^{\frac{3\pi}{2}} (-2) dx \\
&= \left(\frac{5\pi}{6} - \frac{\pi}{2} \right) - \left(\frac{7\pi}{6} - \pi \right) - 2 \left(\frac{3\pi}{2} - \frac{7\pi}{6} \right) = -\frac{\pi}{2}
\end{aligned}$$

80. Let $I = \int_0^{\frac{\pi}{2}} (x - [\cos x]) dx$

$$\begin{aligned}
&= \int_0^{\frac{\pi}{2}} x dx - \int_0^{\frac{\pi}{2}} [\cos x] dx \\
&= \left[\frac{x^2}{2} \right]_0^{\pi/2} - 0 = \frac{\pi^2}{8}
\end{aligned}$$



$$\begin{aligned}
 81. \quad & \int_1^a [f'(x)] dx \\
 &= \int_1^2 f'(x) dx + \int_2^3 f'(x) dx + \dots + \int_{[a]}^a f'(x) dx \\
 &= [f(2) - f(1)] + 2[f(3) - f(2)] \\
 &\quad + \dots + [a][f(a) - f([a])] \\
 &= [a] f(a) - \{f(1) + f(2) + \dots + f([a])\}
 \end{aligned}$$

$$\begin{aligned}
 82. \quad & \int_0^2 (|x-2| + [x]) dx = \int_0^1 |x-2| dx + \int_1^2 [x] dx \\
 &= -\int_0^2 (x-2) dx + \int_0^1 [x] dx + \int_1^2 [x] dx \\
 &= \int_0^2 (2-x) dx + \int_0^1 0 \cdot dx + \int_1^2 1 \cdot dx \\
 &= \left[2x - \frac{x^2}{2} \right]_0^2 + [x]_1^2 \\
 &= (4-2) + (2-1) = 3
 \end{aligned}$$

$$\begin{aligned}
 83. \quad & \int_{-2}^2 |x| dx \\
 &= \int_{-2}^{-1} |x| dx + \int_{-1}^0 |x| dx + \int_0^1 |x| dx + \int_1^2 |x| dx \\
 &= \int_{-2}^{-1} -x dx + \int_{-1}^0 -x dx + \int_0^1 x dx + \int_1^2 x dx \\
 &= 2 \int_{-2}^{-1} -x dx + \int_{-1}^0 -x dx + \int_1^2 x dx \\
 &= 2 \left[-\frac{x^2}{2} \right]_{-2}^{-1} + \left[-\frac{x^2}{2} \right]_{-1}^0 + \left[\frac{x^2}{2} \right]_1^2 \\
 &= 2(-1+2) + (0+1) + (2-1) \\
 &= 2+1+1=4
 \end{aligned}$$

$$\begin{aligned}
 84. \quad & \int_1^5 [x-3] dx \\
 &= \int_1^3 [x-3] dx + \int_3^5 [x-3] dx \\
 &= \int_1^2 [x-3] dx + \int_2^3 [x-3] dx \\
 &\quad + \int_3^4 [x-3] dx + \int_4^5 [x-3] dx \\
 &= \int_1^2 1 \cdot dx + \int_2^3 0 \cdot dx + \int_3^4 0 \cdot dx + \int_4^5 1 \cdot dx \\
 &= [x]_1^2 + [x]_4^5 = (2-1) + (5-4) = 2
 \end{aligned}$$

$$\begin{aligned}
 85. \quad & \text{Let } I = \int_0^{11} \frac{(11-x)^2}{x^2 + (11-x)^2} dx \quad \dots(i) \\
 &= \int_0^{11} \frac{x^2}{(11-x)^2 + x^2} dx \quad \dots(ii) \\
 &\quad \dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]
 \end{aligned}$$

Adding (i) and (ii), we get

$$\begin{aligned}
 2I &= \int_0^{11} 1 dx \\
 \Rightarrow 2I &= [x]_0^{11} \\
 \Rightarrow I &= \frac{11}{2}
 \end{aligned}$$

$$\begin{aligned}
 86. \quad & \text{Let } I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \quad \dots(i) \\
 &= \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin\left(\frac{\pi}{2}-x\right)}}{\sqrt{\sin\left(\frac{\pi}{2}-x\right)} + \sqrt{\cos\left(\frac{\pi}{2}-x\right)}} dx \\
 &\quad \dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]
 \end{aligned}$$

$$\therefore I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx \quad \dots(ii)$$

Adding (i) and (ii), we get

$$\begin{aligned}
 2I &= \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x} + \sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \\
 \Rightarrow 2I &= \left[x \right]_0^{\frac{\pi}{2}} \\
 &= \frac{\pi}{2} \\
 \Rightarrow I &= \frac{\pi}{4}
 \end{aligned}$$

$$\begin{aligned}
 87. \quad & \text{Since, } \int_0^{\frac{\pi}{2}} \frac{\sin^n x}{\sin^n x + \cos^n x} dx = \frac{\pi}{4} \\
 \therefore & \int_0^{\frac{\pi}{2}} \frac{\sin^{1000} x}{\sin^{1000} x + \cos^{1000} x} dx = \frac{\pi}{4}
 \end{aligned}$$



$$88. \text{ Let } I = \int_0^{\frac{\pi}{2}} \frac{\tan^7}{\cot^7 + \tan^7} d \quad \dots(i)$$

$$= \int_0^{\frac{\pi}{2}} \frac{\tan^7\left(\frac{\pi}{2} - \right)}{\cot^7\left(\frac{\pi}{2} - \right) + \tan^7\left(\frac{\pi}{2} - \right)} d$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$\therefore I = \int_0^{\frac{\pi}{2}} \frac{\cot^7}{\tan^7 + \cot^7} d \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_0^{\frac{\pi}{2}} d$$

$$\Rightarrow 2I = \left[x \right]_0^{\frac{\pi}{2}} \Rightarrow I = \frac{\pi}{4}$$

$$89. \text{ Let } I = \int_0^{\pi/2} \frac{\sqrt{\cot}}{\sqrt{\cot} + \sqrt{\tan}} d \quad \dots(i)$$

$$= \int_0^{\pi/2} \frac{\sqrt{\cot\left(\frac{\pi}{2} - \right)}}{\sqrt{\cot\left(\frac{\pi}{2} - \right)} + \sqrt{\tan\left(\frac{\pi}{2} - \right)}} d$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$= \int_0^{\pi/2} \frac{\sqrt{\tan}}{\sqrt{\tan} + \sqrt{\cot}} d \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_0^{\pi/2} d = \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4}$$

$$90. \text{ Let } I = \int_0^{\frac{\pi}{2}} \left(\frac{\sqrt[n]{\sec}}{\sqrt[n]{\sec} + \sqrt[n]{\csc}} \right) d \quad \dots(i)$$

$$\therefore I = \int_0^{\frac{\pi}{2}} \left(\frac{\sqrt[n]{\csc}}{\sqrt[n]{\csc} + \sqrt[n]{\sec}} \right) d \quad \dots(ii)$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \int_0^{\frac{\pi}{2}} d = \left[x \right]_0^{\frac{\pi}{2}} = \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4}$$

$$91. \text{ Let } I = \int_0^{\frac{\pi}{2}} \frac{d}{1 + \sqrt{\tan}} \quad \dots(i)$$

$$= \int_0^{\frac{\pi}{2}} \frac{d}{1 + \sqrt{\cot}} \quad \dots(ii)$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \int_0^{\frac{\pi}{2}} \left(\frac{1}{1 + \sqrt{\tan}} + \frac{1}{1 + \sqrt{\cot}} \right) d$$

$$\Rightarrow 2I = \int_0^{\frac{\pi}{2}} d = \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4}$$

$$92. \text{ Let } I = \int_0^{\frac{\pi}{2}} \frac{d}{1 + (\tan)^{\sqrt{2018}}} \quad \dots(i)$$

$$\therefore I = \int_0^{\frac{\pi}{2}} \frac{d}{1 + (\cot)^{\sqrt{2018}}} \quad \dots(ii)$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \int_0^{\frac{\pi}{2}} \left[\frac{1}{1 + (\tan)^{\sqrt{2018}}} + \frac{1}{1 + (\cot)^{\sqrt{2018}}} \right] d$$

$$= \int_0^{\frac{\pi}{2}} \frac{1}{1 + (\tan)^{\sqrt{2018}}} + \frac{1}{1 + \left(\frac{1}{\tan} \right)^{\sqrt{2018}}} d$$

$$= \int_0^{\frac{\pi}{2}} d$$

$$\Rightarrow I = \frac{\pi}{4}$$

$$93. \text{ Let } I = \int_0^{\frac{\pi}{2}} \frac{2^{\sin}}{2^{\sin} + 2^{\cos}} d \quad \dots(i)$$

$$\therefore I = \int_0^{\frac{\pi}{2}} \frac{2^{\cos}}{2^{\sin} + 2^{\cos}} d \quad \dots(ii)$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$



Adding (i) and (ii), we get

$$2I = \int_0^{\frac{\pi}{2}} \frac{2^{\sin} + 2^{\cos}}{2^{\sin} + 2^{\cos}} dx = \left[x \right]_0^{\frac{\pi}{2}} = \frac{\pi}{2}$$

$$\therefore I = \frac{\pi}{4}$$

94. Let $I = \int_0^{\frac{\pi}{2}} \frac{2008^{\sin}}{2008^{\sin} + 2008^{\cos}} dx \quad \dots(i)$

$$\therefore I = \int_0^{\frac{\pi}{2}} \frac{2008^{\cos}}{2008^{\cos} + 2008^{\sin}} dx \quad \dots(ii)$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \int_0^{\frac{\pi}{2}} dx \quad \left[\right]_0^{\pi/2} = \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4}$$

95. Let $I = \int_0^{\frac{\pi}{2}} \frac{\cos^3}{\sin + \cos} dx \quad \dots(i)$

$$I = \int_0^{\frac{\pi}{2}} \frac{\sin^3}{\cos + \sin} dx \quad \dots(ii)$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \int_0^{\frac{\pi}{2}} \frac{\sin^3 + \cos^3}{\sin + \cos} dx$$

$$= \int_0^{\frac{\pi}{2}} (\sin^2 - \sin \cos + \cos^2) dx$$

$$= \int_0^{\frac{\pi}{2}} (1 - \sin \cos) dx$$

$$= \int_0^{\frac{\pi}{2}} 1 dx - \int_0^{\frac{\pi}{2}} \sin \cos dx$$

$$= \left[x \right]_0^{\pi/2} - \left[\frac{\sin^2}{2} \right]_0^{\pi/2} = \frac{\pi}{2} - \frac{1}{2}$$

$$\therefore 2I = \frac{\pi-1}{2} \Rightarrow I = \frac{\pi-1}{4}$$

96. Let $I = \int_0^{\frac{\pi}{2}} \log(\cot x) dx \quad \dots(i)$

$$= \int_0^{\frac{\pi}{2}} \log(\tan x) dx \quad \dots(ii)$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \int_0^{\frac{\pi}{2}} \log(\cot \tan x) dx$$

$$\therefore 2I = \int_0^{\frac{\pi}{2}} \log 1 dx$$

$$\Rightarrow 2I = 0 \Rightarrow I = 0$$

97. $\int_0^{\frac{\pi}{2}} \log(\operatorname{cosec} x) dx = \int_0^{\frac{\pi}{2}} \log(\sec x) dx$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$= \int_0^{\frac{\pi}{2}} \log\left(\frac{1}{\cos}\right) dx$$

$$= \int_0^{\frac{\pi}{2}} [\log 1 - \log(\cos x)] dx$$

$$= - \int_0^{\frac{\pi}{2}} \log(\cos x) dx$$

$$= \frac{\pi}{2} \log 2$$

98. Let $I = \int_0^{\pi/2} \sin 2x \log \tan x dx$

$$= \int_0^{\pi/2} \sin 2\left(\frac{\pi}{2} - x\right) \log \tan\left(\frac{\pi}{2} - x\right) dx$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$= \int_0^{\pi/2} \sin 2x \log \cot x dx$$

$$= - \int_0^{\pi/2} \sin 2x \log \tan x dx$$

$$\therefore I = -I \Rightarrow 2I = 0 \Rightarrow I = 0$$



$$99. \text{ Let } I = \int_0^{\frac{\alpha}{3}} \frac{f(x)}{f(x) + f\left(\frac{\alpha-x}{3}\right)} dx$$

$$\Rightarrow I = \int_0^{\frac{\alpha}{3}} \frac{f(x)}{f(x) + f\left(\frac{\alpha-x}{3}\right)} dx \quad \dots(i)$$

$$\Rightarrow I = \int_0^{\frac{\alpha}{3}} \frac{f\left(\frac{\alpha-x}{3}\right)}{f\left(\frac{\alpha-x}{3}\right) + f(x)} dx \quad \dots(ii)$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \int_0^{\frac{\alpha}{3}} dx = \left[x \right]_0^{\alpha/3} = \frac{\alpha}{3}$$

$$\Rightarrow I = \frac{\alpha}{6}$$

$$100. \text{ } I = \int_0^a \frac{dx}{1+f(x)} \quad \dots(i)$$

$$= \int_0^a \frac{dx}{1+f(a-x)} \quad \dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$= \int_0^a \frac{dx}{1+\frac{1}{f(x)}} \quad \dots \left[\because f(x)f(a-x) = 1 \right]$$

$$\therefore I = \int_0^a \frac{f(x)}{1+f(x)} dx \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_0^a dx = \left[x \right]_0^a$$

$$\Rightarrow I = \frac{a}{2}$$

$$101. \text{ Let } I = \int_0^{\pi} \frac{dx}{1+\sin x} \quad \dots(i)$$

$$\therefore I = \int_0^{\pi} \frac{\pi-x}{1+\sin(\pi-x)} dx \quad \dots(ii)$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \int_0^{\pi} \frac{\pi}{1+\sin x} dx$$

$$= \pi \int_0^{\pi} \frac{1-\sin x}{(1+\sin x)(1-\sin x)} dx$$

$$= \pi \int_0^{\pi} \frac{1-\sin x}{\cos^2 x} dx = \pi \int_0^{\pi} (\sec^2 x - \sec x \tan x) dx$$

$$= \pi [\tan x - \sec x]_0^{\pi}$$

$$\therefore 2I = \pi[0 - (-1) - (0 - 1)] = 2\pi$$

$$\Rightarrow I = \pi$$

$$102. \text{ Let } I = \int_0^{\pi} \frac{dx}{4\cos^2 x + 9\sin^2 x} \quad \dots(i)$$

$$\therefore I = \int_0^{\pi} \frac{(\pi-x) dx}{4\cos^2(\pi-x) + 9\sin^2(\pi-x)}$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$\therefore I = \int_0^{\pi} \frac{(\pi-x) dx}{4\cos^2 x + 9\sin^2 x} \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_0^{\pi} \frac{\pi dx}{4\cos^2 x + 9\sin^2 x}$$

$$\therefore I = \frac{\pi}{2} \int_0^{\pi} \frac{dx}{4\cos^2 x + 9\sin^2 x}$$

$$= 2 \left(\frac{\pi}{2} \right) \int_0^{\frac{\pi}{2}} \frac{dx}{4\cos^2 x + 9\sin^2 x}$$

$$\dots \left[\because \int_0^a f(x) dx = 2 \int_0^{\frac{a}{2}} f(x) dx \right]$$

$$= \pi \int_0^{\frac{\pi}{2}} \frac{\sec^2 x}{4 + 9\tan^2 x} dx$$

$$= \frac{\pi}{9} \int_0^{\frac{\pi}{2}} \frac{\sec^2 x}{\frac{4}{9} + \tan^2 x} dx$$

Put $\tan x = t \Rightarrow \sec^2 x dx = dt$

$$\therefore I = \frac{\pi}{9} \int_0^{\infty} \frac{dt}{\frac{4}{9} + t^2}$$

$$= \frac{\pi}{9} \times \frac{3}{2} \left[\tan^{-1} \frac{3t}{2} \right]_0^{\infty} = \frac{\pi}{6} \left[\frac{\pi}{2} - 0 \right] = \frac{\pi^2}{12}$$



$$103. \text{ Let } I = \int_0^{\pi} \frac{\tan}{\sec + \cos} d \quad \dots(i)$$

$$\therefore I = \int_0^{\pi} \frac{(\pi -) \tan}{\sec + \cos} d \quad \dots(ii)$$

$$\dots \left[\because \int_0^a f() d = \int_0^a f(a -) d \right]$$

Adding (i) and (ii), we get

$$2I = \pi \int_0^{\pi} \frac{\tan}{\sec + \cos} d$$

$$\Rightarrow I = \frac{\pi}{2} \int_0^{\pi} \frac{\sin}{1 + \cos^2} d$$

Put $\cos = t \Rightarrow \sin d = -dt$

$$\therefore I = -\frac{\pi}{2} \int_1^{-1} \frac{dt}{1+t^2} = -\frac{\pi}{2} [\tan^{-1} t]_1^{-1}$$

$$= \left(-\frac{\pi}{2}\right) \left(-\frac{\pi}{2}\right) = \frac{\pi^2}{4}$$

$$104. \text{ Let } I = \int_0^{\pi/2} \frac{\sin \cos}{\cos^4 + \sin^4} d \quad \dots(i)$$

$$\therefore I = \int_0^{\pi/2} \frac{\left(\frac{\pi}{2} - \right) \cos \sin}{\sin^4 + \cos^4} d \quad \dots(ii)$$

$$\dots \left[\because \int_0^a f() d = \int_0^a f(a -) d \right]$$

Adding (i) and (ii), we get

$$2I = \frac{\pi}{2} \int_0^{\pi/2} \frac{\cos \sin}{\cos^4 + \sin^4} d$$

$$\Rightarrow I = \frac{\pi}{4} \int_0^{\pi/2} \frac{\tan \sec^2}{1 + \tan^4} d$$

Put $\tan^2 = t \Rightarrow \tan \sec^2 d = \frac{dt}{2}$

$$\therefore I = \frac{\pi}{8} \int_0^{\infty} \frac{dt}{1+t^2}$$

$$= \frac{\pi}{8} [\tan^{-1} t]_0^{\infty} = \frac{\pi^2}{16}$$

$$105. \text{ Let } I = \int_0^1 \frac{\log(1 +)}{1 + ^2} d$$

Put $= \tan \theta \Rightarrow d = \sec^2 \theta d\theta$

$$\therefore I = \int_0^{\pi/4} \frac{\log(1 + \tan \theta)}{1 + \tan^2 \theta} \cdot \sec^2 \theta d\theta$$

$$= \int_0^{\pi/4} \frac{\log(1 + \tan \theta)}{\sec^2 \theta} \cdot \sec^2 \theta d\theta$$

$$\therefore I = \int_0^{\pi/4} \log(1 + \tan \theta) d\theta \quad \dots(i)$$

$$= \int_0^{\pi/4} \log \left[1 + \tan \left(\frac{\pi}{4} - \theta \right) \right] d\theta$$

$$= \int_0^{\pi/4} \log \left[1 + \frac{1 - \tan \theta}{1 + \tan \theta} \right] d\theta$$

$$= \int_0^{\pi/4} \log \left(\frac{2}{1 + \tan \theta} \right) d\theta$$

$$\therefore I = \int_0^{\pi/4} \log 2 d\theta - \int_0^{\pi/4} \log(1 + \tan \theta) d\theta$$

$$\Rightarrow I = \int_0^{\pi/4} \log 2 d\theta - I \quad \dots[\text{From (i)}]$$

$$\Rightarrow 2I = \int_0^{\pi/4} \log 2 d\theta$$

$$= \frac{\pi}{4} \log 2$$

$$\Rightarrow I = \frac{\pi}{8} \log 2$$

106. Put $= \tan \theta \Rightarrow d = \sec^2 \theta d\theta$

$$\therefore \int_0^1 \frac{8 \log(1 +)}{1 + ^2} d = \int_0^{\pi/4} \frac{8 \log(1 + \tan \theta)}{1 + \tan^2 \theta} (\sec^2 \theta d\theta)$$

$$= 8 \int_0^{\pi/4} \log(1 + \tan \theta) d\theta = 8 \left(\frac{\pi}{8} \log 2 \right) = \pi \log 2$$

$$107. \text{ Let } I = \int_0^{\pi} [\cot] d \quad \dots(i)$$

$$\Rightarrow I = \int_0^{\pi} [\cot(\pi -)] d$$

$$\dots \left[\because \int_0^a f() d = \int_0^a f(a -) d \right]$$

$$\Rightarrow I = \int_0^{\pi} [-\cot] d \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_0^{\pi} \{ [\cot] + [-\cot] \} d$$

$$\Rightarrow 2I = \int_0^{\pi} -1 d \quad \dots[\because [] + [-] = -1, \text{ if } \notin \mathbb{Z}]$$

$$\Rightarrow 2I = -\pi \Rightarrow I = -\frac{\pi}{2}$$



$$108. \text{ Let } I = \int_2^8 \frac{\sqrt{10-x}}{\sqrt{x} + \sqrt{10-x}} dx \quad \dots(i)$$

$$\therefore I = \int_2^8 \frac{\sqrt{x}}{\sqrt{10-x} + \sqrt{x}} dx \quad \dots(ii)$$

$$\dots \left[\because \int_a^b f(x) dx = \int_a^b f(a+b-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \int_2^8 dx = [x]_2^8 = 8 - 2 = 6 \Rightarrow I = \frac{6}{2} = 3$$

$$109. \text{ Let } I = \int_{2016}^{2017} \frac{\sqrt{x}}{\sqrt{x} + \sqrt{4033-x}} dx \quad \dots(i)$$

$$= \int_{2016}^{2017} \frac{\sqrt{4033-x}}{\sqrt{4033-x} + \sqrt{x}} dx \quad \dots(ii)$$

$$\dots \left[\because \int_a^b f(x) dx = \int_a^b f(a+b-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \int_{2016}^{2017} dx = [x]_{2016}^{2017} = 1$$

$$\therefore I = \frac{1}{2}$$

$$110. \text{ Let } I = \int_{\frac{\pi}{4}}^{\frac{3\pi}{4}} \frac{dx}{1 + \cos x} \quad \dots(i)$$

$$\therefore I = \int_{\frac{\pi}{4}}^{\frac{3\pi}{4}} \frac{dx}{1 + \cos(\pi - x)}$$

$$\dots \left[\because \int_a^b f(x) dx = \int_a^b f(b+a-x) dx \right]$$

$$\therefore I = \int_{\frac{\pi}{4}}^{\frac{3\pi}{4}} \frac{dx}{1 - \cos x} \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_{\frac{\pi}{4}}^{\frac{3\pi}{4}} \frac{2 dx}{1 - \cos^2 x}$$

$$\therefore I = \int_{\frac{\pi}{4}}^{\frac{3\pi}{4}} \operatorname{cosec}^2 x dx$$

$$= -[\cot x]_{\pi/4}^{3\pi/4} = 2$$

$$111. \text{ Let } I = \int_2^4 \frac{\log^2 x}{\log^2 x + \log(36 - 12x + x^2)} dx \quad \dots(i)$$

$$\therefore I = \int_2^4 \frac{\log(6-x)^2}{\log(6-x)^2 + \log^2 x} dx$$

$$\dots(ii) \left[\because \int_a^b f(x) dx = \int_a^b f(a+b-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \int_2^4 dx = [x]_2^4 = 4 - 2 = 2$$

$$\therefore I = 1$$

$$112. \text{ Let } I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{dx}{e^{\sin x} + 1} = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{dx}{e^{\sin(\pi - x)} + 1}$$

$$\therefore I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{e^{-\sin x}}{1 + e^{-\sin x}} dx \quad \dots(i)$$

$$\text{Also, } I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{1}{e^{\sin x} + 1} dx$$

$$= \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{1}{e^{\sin(\pi - x)} + 1} dx$$

$$\dots \left[\because \int_a^b f(x) dx = \int_a^b f(b+a-x) dx \right]$$

$$\therefore I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{1}{e^{-\sin x} + 1} dx \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} dx$$

$$\Rightarrow 2I = [x]_{-\pi/2}^{\pi/2} \Rightarrow I = \frac{\pi}{2}$$

$$113. \text{ Let } I = \int_{\sqrt{\log 2}}^{\sqrt{\log 3}} \frac{\sin^2 x}{\sqrt{\log 2} \sin^2 x + \sin(\log 6 - x^2)} dx$$

$$\text{Put } x^2 = t \Rightarrow 2x dx = dt$$

$$\therefore I = \frac{1}{2} \int_{\log 2}^{\log 3} \frac{\sin t}{\sin t + \sin(\log 6 - t)} dt \quad \dots(i)$$



$$\therefore I = \frac{1}{2} \int_{\log 2}^{\log 3} \frac{\sin(\log 6 - t)}{\sin(\log 6 - t) + \sin t} dt \quad \dots(ii)$$

$$\dots \left[\because \int_a^b f(x) dx = \int_a^b f(a+b-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \frac{1}{2} \int_{\log 2}^{\log 3} dt = \frac{1}{2} (\log 3 - \log 2) = \frac{1}{2} \log \left(\frac{3}{2} \right)$$

$$\Rightarrow I = \frac{1}{4} \log \left(\frac{3}{2} \right)$$

114. $I = \int_{1/2014}^{2014} \frac{\tan^{-1} t}{t} dt \quad \dots(i)$

Put $t = \frac{1}{t} \Rightarrow dt = \frac{-1}{t^2} dt$

$$\therefore I = \int_{2014}^{1/2014} \frac{\tan^{-1} \left(\frac{1}{t} \right)}{\frac{1}{t}} \left(\frac{-1}{t^2} \right) dt$$

$$= \int_{2014}^{1/2014} \frac{-\cot^{-1} t}{t} dt = \int_{1/2014}^{2014} \frac{\cot^{-1} t}{t} dt$$

$\therefore I = \int_{1/2014}^{2014} \frac{\cot^{-1} t}{t} dt \quad \dots(ii)$

Adding (i) and (ii), we get

$$2I = \int_{1/2014}^{2014} \frac{\tan^{-1} t + \cot^{-1} t}{t} dt$$

$$= \frac{\pi}{2} \int_{1/2014}^{2014} \frac{dt}{t} = \frac{\pi}{2} [\log t]_{1/2014}^{2014}$$

$$= \frac{\pi}{2} \left[\log 2014 - \log \frac{1}{2014} \right]$$

$$= \frac{\pi}{2} \times 2 \log 2014$$

$$\Rightarrow I = \frac{\pi}{2} \log 2014$$

115. $I = \int_{\pi/2}^{5\pi/2} \frac{e^{\tan^{-1}(\sin x)}}{e^{\tan^{-1}(\sin x)} + e^{\tan^{-1}(\cos x)}} dx$

$$= \int_{\pi/2}^0 \frac{e^{\tan^{-1}(\sin x)}}{e^{\tan^{-1}(\sin x)} + e^{\tan^{-1}(\cos x)}} dx$$

$$+ \int_0^{5\pi/2} \frac{e^{\tan^{-1}(\sin x)}}{e^{\tan^{-1}(\sin x)} + e^{\tan^{-1}(\cos x)}} dx$$

$$\therefore I = \int_0^{5\pi/2} \frac{e^{\tan^{-1}(\sin x)}}{e^{\tan^{-1}(\sin x)} + e^{\tan^{-1}(\cos x)}} dx$$

$$- \int_0^{\pi/2} \frac{e^{\tan^{-1}(\sin x)}}{e^{\tan^{-1}(\sin x)} + e^{\tan^{-1}(\cos x)}} dx \quad \dots(i)$$

$$= \int_0^{5\pi/2} \frac{e^{\tan^{-1} \sin \left(\frac{5\pi}{2} - x \right)}}{e^{\tan^{-1} \sin \left(\frac{5\pi}{2} - x \right)} + e^{\tan^{-1} \cos \left(\frac{5\pi}{2} - x \right)}} dx$$

$$- \int_0^{\pi/2} \frac{e^{\tan^{-1} \sin \left(\frac{\pi}{2} - x \right)}}{e^{\tan^{-1} \sin \left(\frac{\pi}{2} - x \right)} + e^{\tan^{-1} \cos \left(\frac{\pi}{2} - x \right)}} dx$$

$\therefore I = \int_0^{5\pi/2} \frac{e^{\tan^{-1}(\cos x)}}{e^{\tan^{-1}(\cos x)} + e^{\tan^{-1}(\sin x)}} dx$

$$- \int_0^{\pi/2} \frac{e^{\tan^{-1}(\cos x)}}{e^{\tan^{-1}(\cos x)} + e^{\tan^{-1}(\sin x)}} dx \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_0^{5\pi/2} \frac{e^{\tan^{-1}(\sin x)} + e^{\tan^{-1}(\cos x)}}{e^{\tan^{-1}(\sin x)} + e^{\tan^{-1}(\cos x)}} dx$$

$$- \int_0^{\pi/2} \frac{e^{\tan^{-1}(\sin x)} + e^{\tan^{-1}(\cos x)}}{e^{\tan^{-1}(\sin x)} + e^{\tan^{-1}(\cos x)}} dx$$

$$\Rightarrow 2I = \frac{5\pi}{2} - \frac{\pi}{2} = 2\pi$$

$$\Rightarrow I = \pi$$

116. Let $I = \int_e^\pi f(x) dx$

$$= \int_e^\pi (e + \pi - x) f(x) dx$$

$$\dots \left[\because \int_a^b f(x) dx = \int_a^b f(a+b-x) dx \right]$$

$$= \int_e^\pi (e + \pi - x) f(x) dx$$

$\dots [\because f(x) = f(\pi + e - x) \text{ (given)}]$

$\therefore I = \int_e^\pi (e + \pi) f(x) dx - I$

$$\Rightarrow 2I = (e + \pi) \int_e^\pi f(x) dx \Rightarrow 2I = (e + \pi) \cdot \frac{2}{e + \pi}$$

$$\Rightarrow I = 1$$



$$117. \text{ Let } I = \int_{\frac{\pi}{4}}^{\frac{3\pi}{4}} \frac{\phi}{1 + \sin \phi} d\phi \quad \dots(i)$$

$$= \int_{\frac{\pi}{4}}^{\frac{3\pi}{4}} \frac{\pi - \phi}{1 + \sin(\pi - \phi)} d\phi$$

$$\dots \left[\because \int_a^b f(x) dx = \int_a^b f(a+b-x) dx \right]$$

$$\therefore I = \int_{\frac{\pi}{4}}^{\frac{3\pi}{4}} \frac{\pi - \phi}{1 + \sin \phi} d\phi \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_{\frac{\pi}{4}}^{\frac{3\pi}{4}} \frac{\pi}{1 + \sin \phi} d\phi$$

On solving, we get

$$2I = 2\pi(\sqrt{2} - 1)$$

$$\therefore I = \pi(\sqrt{2} - 1) = \pi \tan \frac{\pi}{8}$$

$$118. I_1 = \int_a^{\pi-a} f(\sin x) dx$$

$$= \int_a^{\pi-a} (\pi - x) f(\sin(\pi - x)) dx$$

$$\dots \left[\because \int_a^b f(x) dx = \int_a^b f(a+b-x) dx \right]$$

$$= \int_a^{\pi-a} (\pi - x) f(\sin x) dx$$

$$\therefore I_1 = \int_a^{\pi-a} \pi f(\sin x) dx - I_1$$

$$\Rightarrow 2I_1 = \pi I_2 \Rightarrow I_2 = \frac{2}{\pi} I_1$$

$$119. f(x) = \frac{e^x}{1 + e^x}$$

$$\therefore f(a) + f(-a) = \frac{e^a}{1 + e^a} + \frac{e^{-a}}{1 + e^{-a}}$$

$$\Rightarrow f(a) + f(-a) = \frac{e^a}{1 + e^a} + \frac{1}{1 + e^a} = 1$$

$$\text{Using } \int_a^b f(x) dx = \int_a^b f(a+b-x) dx, \text{ we have}$$

$$I_1 = \int_{f(-a)}^{f(a)} (1 - x) g((1 - x)) dx$$

$$\dots [\because f(a) + f(-a) = 1]$$

$$\Rightarrow I_1 = \int_{f(-a)}^{f(a)} g((1 - x)) dx - \int_{f(-a)}^{f(a)} g((1 - x)) dx$$

$$\Rightarrow I_1 = I_2 - I_1 \Rightarrow 2I_1 = I_2 \Rightarrow \frac{I_2}{I_1} = 2$$

$$120. \text{ Let } f(x) = e^{\cos^2 x} \cos^3(2n+1)x$$

$$\therefore f(\pi - x) = e^{\cos^2(\pi - x)} \cos^3[(2n+1)(\pi - x)]$$

$$= e^{\cos^2 x} \cos^3[(2n+1)\pi - (2n+1)x]$$

$$= -e^{\cos^2 x} \cos^3(2n+1)x = -f(x)$$

$$\text{Since, } \int_0^{2a} f(x) dx = 0, \text{ if } f(2a - x) = -f(x)$$

$$\therefore \int_0^{\pi} e^{\cos^2 x} \cos^3(2n+1)x dx = 0$$

$$121. \text{ Let } I = \int_0^{\pi} f(\sin x) dx \quad \dots(i)$$

$$\therefore I = \int_0^{\pi} (\pi - x) f(\sin x) dx \quad \dots(ii)$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \pi \int_0^{\pi} f(\sin x) dx$$

$$\Rightarrow 2I = 2\pi \int_0^{\frac{\pi}{2}} f(\sin x) dx \quad \dots \left[\because \int_0^a f(x) dx = 2 \int_0^{\frac{a}{2}} f(x) dx, \text{ if } f(2a-x) = f(x) \right]$$

$$\Rightarrow I = \pi \int_0^{\frac{\pi}{2}} f(\sin x) dx$$

$$\Rightarrow I = \pi \int_0^{\frac{\pi}{2}} f\left(\sin\left(\frac{\pi}{2} - x\right)\right) dx$$

$$\dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$= \pi \int_0^{\frac{\pi}{2}} f(\cos x) dx$$



$$\begin{aligned}
 122. \quad I_1 &= \int_0^{\frac{\pi}{2}} f(\sin 2x) \sin x \, dx \\
 &= \int_0^{\frac{\pi}{4}} \left[f(\sin 2x) \sin x + f\left\{\sin 2\left(\frac{\pi}{2}-x\right)\right\} \sin\left(\frac{\pi}{2}-x\right) \right] dx \\
 &\quad \dots \left[\because \int_0^a f(x) \, dx = \int_0^a [f(x) + f(2a-x)] \, dx \right] \\
 &= \int_0^{\frac{\pi}{4}} \left[f(\sin 2x) \sin x + f\{\sin(\pi-2x)\} \cos x \right] dx \\
 \therefore \quad I_1 &= \int_0^{\frac{\pi}{4}} [f(\sin 2x) \sin x + f(\sin 2x) \cos x] \, dx \quad \dots(i) \\
 I_2 &= \int_0^{\frac{\pi}{4}} f(\cos 2x) \cos x \, dx \\
 &= \int_0^{\frac{\pi}{4}} \left[\cos 2\left(\frac{\pi}{4}-x\right) \right] \cdot \cos\left(\frac{\pi}{4}-x\right) \, dx \\
 &\quad \dots \left[\because \int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx \right] \\
 &= \int_0^{\frac{\pi}{4}} \left[\cos\left(\frac{\pi}{2}-2x\right) \right] \cdot \left(\cos \frac{\pi}{4} \cos x + \sin \frac{\pi}{4} \sin x \right) \, dx \\
 &= \int_0^{\frac{\pi}{4}} f(\sin 2x) \left(\frac{1}{\sqrt{2}} \cos x + \frac{1}{\sqrt{2}} \sin x \right) \, dx \\
 &= \frac{1}{\sqrt{2}} \int_0^{\frac{\pi}{4}} [f(\sin 2x) \cos x + f(\sin 2x) \sin x] \, dx \\
 \therefore \quad I_2 &= \frac{1}{\sqrt{2}} I_1 \quad \dots[\text{From (i)}] \\
 \therefore \quad \frac{I_1}{I_2} &= \sqrt{2} \\
 123. \quad I &= \int_0^{100\pi} \sqrt{1-\cos 2x} \, dx = \int_0^{100\pi} \sqrt{2 \sin^2 x} \, dx \\
 &= \sqrt{2} \int_0^{100\pi} \sin x \, dx \\
 &= 100\sqrt{2} \int_0^{\pi} \sin x \, dx \\
 &\quad \dots \left[\because \int_0^{2a} f(x) \, dx = 2 \int_0^a f(x) \, dx, \text{ if } f(2a-x) = f(x) \right] \\
 &= 100\sqrt{2} [-\cos x]_0^{\pi} = 200\sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 124. \quad \text{Let } f(x) &= |x| \\
 f(-x) &= -|x| = -f(x) \\
 \therefore \quad f(x) &\text{ is an odd function} \\
 \therefore \quad \int_{-1}^1 |x| \, dx &= 0 \\
 \text{Let } I &= \int_0^{\frac{\pi}{2}} \left[1 + \log \left(\frac{4+3 \sin x}{4+3 \cos x} \right) \right] \, dx \quad \dots(i) \\
 &= \int_0^{\frac{\pi}{2}} \left[1 + \log \left(\frac{4+3 \sin\left(\frac{\pi}{2}-x\right)}{4+3 \cos\left(\frac{\pi}{2}-x\right)} \right) \right] \, dx \\
 &\quad \dots \left[\because \int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx \right] \\
 \therefore \quad I &= \int_0^{\frac{\pi}{2}} \left[1 + \log \left(\frac{4+3 \cos x}{4+3 \sin x} \right) \right] \, dx \quad \dots(ii) \\
 \text{Adding (i) and (ii), we get} \\
 2I &= \int_0^{\frac{\pi}{2}} \left[1 + \log \left(\frac{4+3 \sin x}{4+3 \cos x} \right) + 1 \right. \\
 &\quad \left. + \log \left(\frac{4+3 \cos x}{4+3 \sin x} \right) \right] \, dx \\
 &= \int_0^{\frac{\pi}{2}} \left[2 + \log \left(\frac{4+3 \sin x}{4+3 \cos x} \right) - \log \left(\frac{4+3 \sin x}{4+3 \cos x} \right) \right] \, dx \\
 &= \int_0^{\frac{\pi}{2}} 2 \, dx = 2 \left[x \right]_0^{\frac{\pi}{2}} \\
 \therefore \quad 2I &= 2 \left(\frac{\pi}{2} \right) \\
 \therefore \quad I &= \frac{\pi}{2} \\
 125. \quad \text{Since, } \sin^3 x \cos^2 x &\text{ is an odd function.} \\
 \therefore \quad \int_{-1}^1 \sin^3 x \cos^2 x \, dx &= 0 \\
 126. \quad \text{Since, } \sin^{103} x \cos^{101} x &\text{ is an odd function.} \\
 \therefore \quad \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \sin^{103} x \cos^{101} x \, dx &= 0 \\
 127. \quad \text{Since, } \cos x + \sin x &\text{ is an odd function.} \\
 \therefore \quad \int_{-2}^2 (\cos x + \sin x) \, dx &= \int_{-2}^2 dx = 0 + [x^2]_{-2}^2 = 4
 \end{aligned}$$



128. Let $f(x) = \log\left(\frac{9-x}{9+x}\right)$

$$\begin{aligned}\therefore f(-x) &= \log\left(\frac{9-(-x)}{9+(-x)}\right)^{-1} \\ &= -\log\left(\frac{9-x}{9+x}\right) = -f(x)\end{aligned}$$

$\therefore f(x)$ is an odd function.

$$\therefore \int_{-4}^4 \log\left(\frac{9-x}{9+x}\right) dx = 0$$

129. Let $f(x) = \log\left(\frac{2-\sin x}{2+\sin x}\right)$

$$\begin{aligned}\therefore f(-x) &= \log\left(\frac{2-\sin(-x)}{2+\sin(-x)}\right) \\ &= -\log\left(\frac{2-\sin x}{2+\sin x}\right) = -f(x)\end{aligned}$$

$\therefore f(x)$ is an odd function.

$$\therefore \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \log\left(\frac{2-\sin x}{2+\sin x}\right) dx = 0$$

130. Let $I = \int_{-2}^2 (p^2 + q + s)x dx$

$$\begin{aligned}&= \int_{-2}^2 (p^2 + s)x dx + q \int_{-2}^2 dx \\ &= 2 \int_0^2 (p^2 + s)x dx + 0 = \frac{4}{3}(4p + 3s)\end{aligned}$$

Thus, to find the numerical value of I , it is necessary to know the values of p and s .

131. $\int_{-2}^3 f(x) dx = \int_{-2}^2 f(x) dx + \int_2^3 f(x) dx$

$$= \int_{-2}^2 e^{\cos x} \sin x dx + \int_2^3 2 dx$$

Since, $e^{\cos x} \sin x$ is an odd function.

$$\therefore \int_{-2}^3 f(x) dx = 0 + 2(3-2) = 2$$

132. Let $I = \int_{-2}^0 [x^3 + 3x^2 + 3x + (x+1)\cos(x+1)] dx$

$$= \int_{-2}^0 [(x+1)^3 + 2 + (x+1)\cos(x+1)] dx$$

Put $x+1 = t \Rightarrow dx = dt$

$$\therefore I = \int_{-1}^1 (t^3 + 2 + t \cos t) dt$$

Since, t^3 and $t \cos t$ are odd functions.

$$\therefore I = \int_{-1}^1 2 dt = [2t]_{-1}^1 = 4$$

133. $g(x) = \frac{f(-x) - f(x)}{x^2 + 3}$

$$= \frac{4x^2 + 3 - 1 - 4x^2 + 3 - 1}{x^2 + 3} = \frac{6}{x^2 + 3}$$

$$\therefore g(-x) = -\frac{6}{x^2 + 3} = -g(x)$$

$\therefore g(x)$ is an odd function.

$$\therefore \int_{-2}^2 g(x) dx = 0$$

134. Let $I = \int_{-2}^2 |x| dx$

$$= 2 \int_0^2 x dx \quad \left[\because \int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx \right. \\ \left. \text{if } f(x) \text{ is an even function} \right]$$

$$= 2 \left[\frac{x^2}{2} \right]_0^2 = 2 \left(\frac{4}{2} \right) = 4$$

135. Let $I = \int_{-2}^2 |\cos \pi x| dx$

$$= 2 \int_0^2 |\cos \pi x| dx$$

$$\left[\because \int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx \right. \\ \left. \text{if } f(x) \text{ is an even function} \right]$$

$$= 2 \left[\int_0^{1/2} \cos \pi x dx - \int_{1/2}^{3/2} \cos \pi x dx \right. \\ \left. + \int_{3/2}^2 \cos \pi x dx \right]$$

Since $\int \cos \pi x = \frac{\sin \pi x}{\pi} + \frac{\cos \pi x}{\pi^2}$

$$\therefore I = 2 \left[\left(\frac{1}{2\pi} - \frac{1}{\pi^2} \right) - \left(\frac{-3}{2\pi} - \frac{1}{2\pi} \right) + \left(\frac{1}{\pi^2} + \frac{3}{2\pi} \right) \right]$$

$$= 2 \left[\frac{1}{2\pi} - \frac{1}{\pi^2} + \frac{3}{2\pi} + \frac{1}{2\pi} + \frac{1}{\pi^2} + \frac{3}{2\pi} \right]$$

$$= 2 \left(\frac{8}{2\pi} \right) = \frac{8}{\pi}$$



136. Since, \tan^{-1} is an even function.

$$\begin{aligned} \therefore \int_{-1}^1 \tan^{-1} d &= 2 \int_0^1 \tan^{-1} d \\ &= \left[2 \tan^{-1} \cdot \frac{2}{2} \right]_0^1 - 2 \int_0^1 \frac{1}{1 + \frac{2}{2}} \cdot \frac{2}{2} d \\ &= \left[2 \tan^{-1} \right]_0^1 - \int_0^1 \frac{2 + 1 - 1}{1 + \frac{2}{2}} d \\ &= \left[2 \tan^{-1} \right]_0^1 - \left[\frac{2}{2} \right]_0^1 + \left[\tan^{-1} \right]_0^1 \\ &= \frac{\pi}{4} - 1 + \frac{\pi}{4} = \frac{\pi}{2} - 1 \end{aligned}$$

137. Let $I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\sin^4}{\sin^4 + \cos^4} d$

Since, $\frac{\sin^4}{\sin^4 + \cos^4}$ is an even function.

$$\therefore I = 2 \int_0^{\frac{\pi}{2}} \frac{\sin^4}{\sin^4 + \cos^4} d = 2 \left(\frac{\pi}{4} \right) = \frac{\pi}{2}$$

138. Let $I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\sin^2}{1 + 2} d \dots(i)$

$$I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\sin^2}{1 + 2} d \dots(ii)$$

$$\dots \left[\because \int_a^b f(x) dx = \int_a^b f(a+b-x) dx \right]$$

Adding (i) and (ii), we get

$$2I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^2 d$$

$$\Rightarrow 2I = 2 \int_0^{\frac{\pi}{2}} \sin^2 d$$

$$\dots \left[\because \int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx \right. \\ \left. \text{if } f(x) \text{ is an even function} \right]$$

$$\Rightarrow I = \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos 2}{2} \right) d$$

$$= \frac{1}{2} \left[-\frac{\sin 2}{2} \right]_0^{\frac{\pi}{2}} = \frac{1}{2} \left(\frac{\pi}{2} - 0 \right) = \frac{\pi}{4}$$

139. Let $I = \int_{-1}^1 \frac{\sin^{-2}}{3 - |x|} d$

$$= \int_{-1}^1 \frac{\sin}{3 - |x|} d - \int_{-1}^1 \frac{2}{3 - |x|} d$$

Since, $\frac{\sin}{3 - |x|}$ is an odd function and $\frac{2}{3 - |x|}$ is an even function.

$$\therefore I = 0 - 2 \int_0^1 \frac{2}{3 - |x|} d = 2 \int_0^1 \frac{-2}{3 - |x|} d$$

140. $\int_{-\pi}^{\pi} (\cos a - \sin b)^2 d$

$$= \int_{-\pi}^{\pi} (\cos^2 a + \sin^2 b - 2 \cos a \sin b) d$$

$$= \int_{-\pi}^{\pi} (\cos^2 a + \sin^2 b) d - 2 \int_{-\pi}^{\pi} \cos a \sin b d$$

$$= 2 \int_0^{\pi} (\cos^2 a + \sin^2 b) d - 0$$

$\left[\because \cos a \sin b \text{ is an odd function and } (\cos^2 a + \sin^2 b) \text{ is an even function} \right]$

$$= 2 \int_0^{\pi} \left(\frac{1 + \cos 2a}{2} + \frac{1 - \cos 2b}{2} \right) d$$

$$= \int_0^{\pi} (2 + \cos 2a - \cos 2b) d = 2\pi$$

141. Let $I = \int_{-\frac{3\pi}{2}}^{-\frac{\pi}{2}} \left[(\pi + t)^3 + \cos^2(\pi + 3\pi) \right] dt$

Put $\pi + t = u \Rightarrow dt = du$

$$\therefore I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \left[u^3 + \cos^2(2\pi + u) \right] du$$

$$= \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} u^3 du + \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^2 t dt$$

Since, t^3 is an odd function and $\cos^2 t$ is an even function.

$$\therefore I = 0 + 2 \int_0^{\frac{\pi}{2}} \cos^2 t dt = 2 \times \frac{\pi}{4} = \frac{\pi}{2}$$



$$\begin{aligned}
 142. \text{ Let } I &= \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \left\{ x^2 + \log\left(\frac{\pi+x}{\pi-x}\right) \right\} \cos x \, dx \\
 &= \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} x^2 \cos x \, dx + \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \log\left(\frac{\pi+x}{\pi-x}\right) \cos x \, dx \\
 &= 2 \int_0^{\frac{\pi}{2}} x^2 \cos x \, dx + 0
 \end{aligned}$$

$$\begin{aligned}
 143. \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{x + \frac{\pi}{4}}{2 - \cos 2x} \, dx \\
 &= \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{x}{2 - \cos 2x} \, dx + \frac{\pi}{4} \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{1}{2 - \cos 2x} \, dx \\
 &= 0 + 2 \cdot \frac{\pi}{4} \int_0^{\frac{\pi}{4}} \frac{1}{2 - \cos 2x} \, dx \\
 &\quad \left[\begin{array}{l} \because \frac{x}{2 - \cos 2x} \text{ is an odd function} \\ \text{and } \frac{1}{2 - \cos 2x} \text{ is an even function} \end{array} \right] \\
 &= \frac{\pi}{2} \int_0^{\frac{\pi}{4}} \frac{1}{2 - \cos 2x} \, dx \\
 &= \frac{\pi}{2} \int_0^{\frac{\pi}{4}} \frac{1}{2 - \frac{1-t^2}{1+t^2}} \cdot \frac{dt}{1+t^2} \quad \dots [\text{Put } \tan x = t] \\
 &= \frac{\pi}{2} \int_0^{\frac{1}{\sqrt{3}}} \frac{dt}{1+3t^2} = \frac{\pi}{2} \cdot \frac{1}{\sqrt{3}} \left[\tan^{-1}(\sqrt{3}t) \right]_0^{\frac{1}{\sqrt{3}}} \\
 &= \frac{\pi}{2\sqrt{3}} \left[\tan^{-1}(\sqrt{3}) - 0 \right] \\
 &= \frac{\pi}{2\sqrt{3}} \cdot \frac{\pi}{3} = \frac{\pi^2}{6\sqrt{3}} \\
 &\quad \left[\begin{array}{l} \because \log\left(\frac{\pi+x}{\pi-x}\right) \cos x \text{ is an odd function} \\ \text{and } x^2 \cos x \text{ is an even function.} \end{array} \right] \\
 &= 2 \left[x^2 \sin x + 2x \cos x - 2 \sin x \right]_0^{\pi/2} \\
 &= 2 \left(\frac{\pi^2}{4} - 2 \right) = \frac{\pi^2}{2} - 4
 \end{aligned}$$

$$\begin{aligned}
 144. \quad p'(x) &= p'(1-x) \\
 \text{Integrating on both sides, we get} \\
 p(x) &= -p(1-x) + c \quad \dots (i) \\
 \therefore p(0) &= -p(1-0) + c \\
 \Rightarrow 1 &= -41 + c \Rightarrow c = 42 \\
 \therefore p(x) + p(1-x) &= 42 \quad \dots (ii) [\text{From (i)}] \\
 \text{Let } I &= \int_0^1 p(x) \, dx \\
 &= \int_0^1 p(1-x) \, dx \quad \dots \left[\because \int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx \right] \\
 &= \int_0^1 [42 - p(x)] \, dx \quad \dots [\text{From (ii)}] \\
 \therefore I &= 42 \int_0^1 dx - I \\
 \Rightarrow 2I &= 42 \Rightarrow I = 21
 \end{aligned}$$

$$\begin{aligned}
 145. \quad A(x) &= \begin{vmatrix} 1 & 2 & 3 \\ x+1 & 2x+1 & 3x+1 \\ x^2+1 & 2x^2+1 & 3x^2+1 \end{vmatrix} \\
 \text{Applying } C_2 &\rightarrow C_2 - 2C_1, C_3 \rightarrow C_3 - 3C_1, \text{ we get} \\
 A(x) &= \begin{vmatrix} 1 & 0 & 0 \\ x+1 & -1 & -2 \\ x^2+1 & -1 & -2 \end{vmatrix} = 1(2-2) - 0 + 0 = 0 \\
 \therefore \int_0^1 A(x) \, dx &= 0
 \end{aligned}$$

$$\begin{aligned}
 146. \text{ Let } I &= \int_0^{\sin^2} \sin^{-1} \sqrt{t} \, dt + \int_0^{\cos^2} \cos^{-1} \sqrt{t} \, dt \\
 \text{Putting } t &= \sin^2 u \text{ in the first integral and} \\
 t &= \cos^2 v \text{ in the second integral, we get} \\
 I &= \int_0^{\frac{\pi}{2}} u \sin 2u \, du - \int_{\frac{\pi}{2}}^0 v \sin 2v \, dv \\
 &= \int_0^{\frac{\pi}{2}} u \sin 2u \, du + \int_{\frac{\pi}{2}}^0 u \sin 2u \, du - \int_{\frac{\pi}{2}}^0 v \sin 2v \, dv \\
 &= \int_0^{\frac{\pi}{2}} u \sin 2u \, du \quad \dots \left[\because \int_a^b f(x) \, dx = \int_a^b f(t) \, dt \right] \\
 &= \left[\frac{-u \cos 2u}{2} \right]_0^{\pi/2} + \frac{1}{2} \int_0^{\pi/2} \cos 2u \, du \\
 &= \left[\frac{-u \cos 2u}{2} \right]_0^{\pi/2} + \frac{1}{4} [\sin 2u]_0^{\pi/2} = \frac{\pi}{4}
 \end{aligned}$$



$$147. N = \int_0^{\frac{\pi}{4}} \frac{\sin \cos}{(+ 1)^2} d = \frac{1}{2} \int_0^{\frac{\pi}{4}} \frac{\sin 2}{(+ 1)^2} d$$

$$= \frac{1}{2} \left[\left[\sin 2 \left(-\frac{1}{+1} \right) \right]_0^{\frac{\pi}{4}} + \int_0^{\frac{\pi}{4}} \frac{2 \cos 2}{(+ 1)} d \right]$$

$$= -\frac{2}{\pi + 4} + \int_0^{\frac{\pi}{4}} \frac{\cos 2}{+1} d = \frac{-2}{\pi + 4} + I_2$$

In I_2 , put $2 = t \Rightarrow d = \frac{dt}{2}$

$$\therefore I_2 = \int_0^{\frac{\pi}{2}} \frac{\cos t}{t + 2} dt = \int_0^{\frac{\pi}{2}} \frac{\cos}{+2} d = M$$

$$\therefore N = -\frac{2}{\pi + 4} + M \quad \therefore M - N = \frac{2}{\pi + 4}$$

$$148. af() + bf \left(\frac{1}{-} \right) = \frac{1}{-} - 5 \quad \dots(i)$$

Replacing by $\frac{1}{-}$ in (i), we get

$$af \left(\frac{1}{-} \right) + bf() = -5 \quad \dots(ii)$$

Eliminating $f \left(\frac{1}{-} \right)$ from (i) and (ii), we get

$$(a^2 - b^2)f() = \frac{a}{-} - b - 5a + 5b$$

$$\therefore (a^2 - b^2) \int_1^2 f() d$$

$$= \left[a \log \left| -\frac{b}{2} \right|^2 - 5(a - b) \right]_1^2$$

$$= a \log 2 - 2b - 10(a - b) - a \log 1 + \frac{b}{2} + 5(a - b)$$

$$= a \log 2 - 5a + \frac{7}{2} b$$

$$\therefore \int_1^2 f() d = \frac{1}{(a^2 - b^2)} \left[a \log 2 - 5a + \frac{7}{2} b \right]$$

$$149. \text{ Let } I = \frac{2}{\pi} \int_{-\pi}^{\pi} f() d$$

$$= \frac{2}{\pi} \int_{-\pi}^{\pi} \frac{\sin \left(\frac{9}{2} \right)}{\sin \left(\frac{2}{2} \right)} d = \frac{4}{\pi} \int_0^{\pi} \frac{\sin \left(\frac{9}{2} \right)}{\sin \left(\frac{2}{2} \right)} d$$

....[$\because f()$ is an even function]

Put $\frac{2}{2} = \theta \Rightarrow d = 2d\theta$

$$\therefore I = \frac{8}{\pi} \int_0^{\frac{\pi}{2}} \frac{\sin 9\theta}{\sin \theta} d\theta$$

$$= \frac{8}{\pi} \int_0^{\frac{\pi}{2}} \frac{(\sin 9\theta - \sin 7\theta) + (\sin 7\theta - \sin 5\theta) + (\sin 5\theta - \sin 3\theta) + (\sin 3\theta - \sin \theta) + \sin \theta}{\sin \theta} d\theta$$

$$= \frac{8}{\pi} \int_0^{\frac{\pi}{2}} (2 \cos 8\theta + 2 \cos 6\theta + 2 \cos 4\theta + 2 \cos 2\theta + 1) d\theta$$

$$= \frac{8}{\pi} \times \frac{\pi}{2} = 4$$



Evaluation Test

$$1. \text{ Let } I = \int_0^1 \frac{\log(1 +)}{1 + ^2} d$$

Put $= \tan t \Rightarrow d = \sec^2 t dt$

When $= 0$, $t = 0$ and when $= 1$, $t = \frac{\pi}{4}$

$$\therefore I = \int_0^{\frac{\pi}{4}} \frac{\log(1 + \tan t)}{1 + \tan^2 t} \cdot \sec^2 t dt = \int_0^{\frac{\pi}{4}} \log(1 + \tan t) dt$$

$$= \int_0^{\frac{\pi}{4}} \log \left[1 + \tan \left(\frac{\pi}{4} - t \right) \right] dt$$

$$= \int_0^{\frac{\pi}{4}} \log \left(1 + \frac{1 - \tan t}{1 + \tan t} \right) dt = \int_0^{\frac{\pi}{4}} \log \left(\frac{2}{1 + \tan t} \right) dt$$

$$= \int_0^{\frac{\pi}{4}} [\log 2 - \log(1 + \tan t)] dt$$

$$\therefore I = \int_0^{\frac{\pi}{4}} (\log 2) dt - I$$

$$\therefore 2I = \log 2 [t]_0^{\pi/4} = \frac{\pi}{4} \log 2$$

$$\therefore I = \frac{\pi}{8} \log 2$$



$$\begin{aligned}
 2. \quad \int_0^5 f(x) dx &= \int_0^1 f(x) dx + \int_1^2 f(x) dx + \int_2^3 f(x) dx \\
 &\quad + \int_3^4 f(x) dx + \int_4^5 f(x) dx \\
 &= 0 + \int_1^2 1^2 dx + \int_2^3 2^2 dx + \int_3^4 3^2 dx + \int_4^5 4^2 dx \\
 &= 1(2-1) + 4(3-2) + 9(4-3) + 16(5-4) \\
 &= 1 + 4 + 9 + 16 = 30
 \end{aligned}$$

$$\begin{aligned}
 3. \quad \text{Let } I &= \int_0^3 \frac{1}{1+2^{f(x)}} dx \quad \dots (i) \\
 &= \int_0^3 \frac{1}{1+2^{f(3-x)}} dx \\
 &\quad \dots \left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right] \\
 &= \int_0^3 \frac{1}{1+2^{-f(x)}} dx \\
 &\quad \dots [\because f(x) + f(3-x) = 0 \text{ (given)}]
 \end{aligned}$$

$$\therefore I = \int_0^3 \frac{2^{f(x)}}{2^{f(x)} + 1} dx \quad \dots (ii)$$

Adding (i) and (ii), we get

$$\begin{aligned}
 2I &= \int_0^3 \frac{1}{1+2^{f(x)}} dx + \int_0^3 \frac{2^{f(x)}}{2^{f(x)} + 1} dx \\
 &= \int_0^3 \frac{1+2^{f(x)}}{1+2^{f(x)}} dx = \int_0^3 1 dx = 3
 \end{aligned}$$

$$\therefore I = \frac{3}{2}$$

$$\begin{aligned}
 4. \quad \text{Let } I &= \int_0^{\infty} \frac{\log x}{(1+x^2)^2} dx \\
 &= \int_0^1 \frac{\log x}{(1+x^2)^2} dx + \int_1^{\infty} \frac{\log x}{(1+x^2)^2} dx \\
 &= I_1 + I_2 \text{ (Say)} \quad \dots (i)
 \end{aligned}$$

$$I_2 = \int_0^{\infty} \frac{\log x}{(1+x^2)^2} dx$$

$$\text{Put } x = \frac{1}{t}$$

$$\therefore dx = -\frac{1}{t^2} dt$$

When $x = 1$, $t = 1$ and when $x \rightarrow \infty$, $t \rightarrow 0$

$$\begin{aligned}
 \therefore I_2 &= \int_1^0 \frac{\frac{1}{t} \log\left(\frac{1}{t}\right)}{\left(1+\frac{1}{t^2}\right)^2} \left(-\frac{1}{t^2}\right) dt \\
 &= \int_1^0 \frac{\log t}{(1+t^2)^2} dt \quad \dots \left[\because \log\left(\frac{1}{t}\right) = -\log t \right] \\
 &= -\int_0^1 \frac{\log t}{(1+t^2)^2} dt \\
 &= -\int_0^1 \frac{\log t}{(1+t^2)^2} dt
 \end{aligned}$$

$$\begin{aligned}
 \therefore I_2 &= -I_1 \\
 \therefore \text{From (i), } I &= I_1 + I_2 = 0
 \end{aligned}$$

$$\begin{aligned}
 5. \quad \int_0^n [x] dx &= \int_0^1 [x] dx + \int_1^2 [x] dx + \int_2^3 [x] dx + \dots + \int_{n-1}^n [x] dx \\
 &= \int_0^1 0 dx + \int_1^2 1 dx + \int_2^3 2 dx + \dots + \int_{n-1}^n (n-1) dx \\
 &= 0 + [x]_1^2 + 2[x]_2^3 + \dots + (n-1)[x]_{n-1}^n \\
 &= (2-1) + 2(3-2) + \dots + (n-1)(n-n+1) \\
 &= 1 + 2(1) + \dots + (n-1)(1) \\
 &= 1 + 2 + \dots + (n-1) \\
 &= \frac{(n-1)n}{2} = \frac{n(n-1)}{2}
 \end{aligned}$$

$$\begin{aligned}
 6. \quad I_1 &= \int_0^{\pi/2} \frac{\sin^2 x}{\sin^2 x} dx = \frac{\pi}{2} \\
 I_2 &= \int_0^{\pi/2} \frac{\sin^2 2x}{\sin^2 x} dx = \int_0^{\pi/2} \frac{(2 \sin x \cos x)^2}{\sin^2 x} dx \\
 &= \int_0^{\pi/2} 4 \cos^2 x dx = 4 \times \frac{\pi}{4} = \pi
 \end{aligned}$$

$$\begin{aligned}
 I_3 &= \int_0^{\pi/2} \frac{\sin^2 3x}{\sin^2 x} dx \\
 &= \int_0^{\pi/2} \frac{(3 \sin x - 4 \sin^3 x)^2}{\sin^2 x} dx \\
 &= \int_0^{\pi/2} (9 - 24 \sin^2 x + 16 \sin^4 x) dx \\
 &= \frac{9\pi}{2} - 24 \cdot \frac{\pi}{4} + 16 \cdot \frac{3.1}{4.2} \cdot \frac{\pi}{2} = \frac{3\pi}{2}
 \end{aligned}$$

$$\therefore I_1 + I_3 = \frac{\pi}{2} + \frac{3\pi}{2} = 2\pi = 2I_2$$

$\therefore I_1, I_2, I_3$ are in A.P.



$$7. \text{ Let } I = \int_{\alpha}^{\beta} \frac{1}{\sqrt{(-\alpha)(\beta-x)}} dx$$

Put $x = \alpha \sin^2 t + \beta \cos^2 t$

$$\therefore dx = (\alpha \cdot 2 \sin t \cos t + \beta \cdot 2 \cos t (-\sin t)) dt$$

$$= 2(\alpha - \beta) \sin t \cos t dt$$

When $x = \alpha$, $\alpha = \alpha \sin^2 t + \beta(1 - \sin^2 t)$

$$\alpha = \beta + (\alpha - \beta) \sin^2 t$$

$$\therefore \sin t = 1, \quad \therefore t = \frac{\pi}{2}$$

When $x = \beta$,

$$\beta = \alpha(1 - \cos^2 t) + \beta \cos^2 t$$

$$= \alpha + (\beta - \alpha) \cos^2 t$$

$$\therefore \cos t = 1, t = 0$$

$$(-\alpha)(\beta - x) = (\alpha \sin^2 t + \beta \cos^2 t - \alpha)$$

$$= (\beta - \alpha \sin^2 t - \beta \cos^2 t)$$

$$= [\beta \cos^2 t - \alpha(1 - \sin^2 t)]$$

$$= [\beta(1 - \cos^2 t) - \alpha \sin^2 t]$$

$$= (\beta - \alpha) \cos^2 t (\beta - \alpha) \sin^2 t$$

Since, $\beta > \alpha$

$$\therefore \sqrt{(-\alpha)(\beta-x)} = (\beta - \alpha) \sin t \cos t$$

$$\therefore I = 2 \int_{\frac{\pi}{2}}^0 \frac{(\alpha - \beta) \sin t \cos t}{(\beta - \alpha) \sin t \cos t} dt$$

$$= 2 \int_{\frac{\pi}{2}}^0 (-1) dt$$

$$= 2 \int_0^{\frac{\pi}{2}} 1 dt$$

$$= 2 [t]_0^{\pi/2}$$

$$= 2 \left(\frac{\pi}{2} - 0 \right)$$

$$= \pi$$

$$8. \text{ Let } h(x) = {}^3f(x) = {}^3\left(\frac{e+x}{e-x}\right)$$

$$\therefore h(-x) = (-x) {}^3\left(\frac{e^{-x}+1}{e^{-x}-1}\right) = - {}^3\left(\frac{1+e}{1-e}\right)$$

$$= {}^3\left(\frac{e+1}{e-1}\right)$$

$$\therefore h(-x) = h(x)$$

$\therefore h(x)$ is an even function.

$$\therefore \int_{-1}^1 t^3 f(t) dt = \int_{-1}^1 h(t) dt = 2 \int_0^1 h(t) dt$$

$$= 2 \int_0^1 t^3 f(t) dt$$

$$= 2 \int_0^1 {}^3f(x) dx$$

$$= 2\alpha \quad \dots \left[\because \int_0^1 {}^3f(x) dx = \alpha \right]$$

$$9. f(m, n) = \int_0^1 (\log x)^m x^{n-1} dx$$

$$= \left[(\log x)^m \int_0^1 x^{n-1} dx \right]_0^1 - \int_0^1 \left\{ \frac{d}{dx} (\log x)^m \int_0^1 x^{n-1} dx \right\} dx$$

$$= \left[(\log x)^m \cdot \frac{x^n}{n} \right]_0^1 - \int_0^1 m (\log x)^{m-1} \cdot \frac{1}{n} x^{n-1} dx$$

$$= 0 - 0 - \frac{m}{n} \int_0^1 (\log x)^{m-1} \cdot x^{n-1} dx \quad \dots [\because \log 1 = 0]$$

$$= -\frac{m}{n} f(m-1, n)$$

$$10. \phi(x) = \int_{\frac{7\pi}{6}}^x (4 \sin t + 3 \cos t) dt$$

$$\therefore \phi'(x) = 4 \sin x + 3 \cos x$$

$$\text{If } x \in \left[\frac{7\pi}{6}, \frac{4\pi}{3} \right],$$

then x is in the third quadrant.

$\therefore \sin x$ and $\cos x$ are both negative.

$$\therefore \phi'(x) = 4 \sin x + 3 \cos x < 0$$

$$\therefore \phi(x) \text{ is decreasing on the interval } \left[\frac{7\pi}{6}, \frac{4\pi}{3} \right]$$

$$\therefore \text{Minimum (least) value of } \phi(x) \text{ on } \left[\frac{7\pi}{6}, \frac{4\pi}{3} \right]$$

$$\text{is } \phi\left(\frac{4\pi}{3}\right) = \int_{7\pi/6}^{4\pi/3} (4 \sin t + 3 \cos t) dt$$

$$= [-4 \cos t + 3 \sin t]_{7\pi/6}^{4\pi/3}$$

$$= -4 \left(\cos \frac{4\pi}{3} - \cos \frac{7\pi}{6} \right) + 3 \left(\sin \frac{4\pi}{3} - \sin \frac{7\pi}{6} \right)$$

$$= -4 \left(-\frac{1}{2} + \frac{\sqrt{3}}{2} \right) + 3 \left(-\frac{\sqrt{3}}{2} + \frac{1}{2} \right)$$

$$= \frac{7(1-\sqrt{3})}{2}$$



11. Let $I = \int_a^b f(x) dx$
 Put $x = (b-a)t + a$
 $\therefore dx = (b-a)dt$
 When $x = a$, $t = 0$ and when $x = b$, $t = 1$
 $\therefore I = \int_0^1 f[(b-a)t + a](b-a)dt$
 $= (b-a) \int_0^1 f[(b-a)t + a] dt$
 $= (b-a) \int_0^1 f[(b-a)t + a] d$
 $\therefore \lambda = b - a$
12. If $0 \leq x < 1$, then $0 \leq x^2 < 1$, $\therefore [x^2] = 0$
 If $1 \leq x < \sqrt{2}$, then $1 \leq x^2 < 2$, $\therefore [x^2] = 1$
 If $\sqrt{2} \leq x \leq 1.5$, then $2 \leq x^2 \leq 2.25$, $\therefore [x^2] = 2$
 $\therefore \int_0^{1.5} [x^2] dx = \int_0^1 [x^2] dx + \int_1^{\sqrt{2}} [x^2] dx + \int_{\sqrt{2}}^{1.5} [x^2] dx$
 $= \int_0^1 0 dx + \int_1^{\sqrt{2}} 1 dx + \int_{\sqrt{2}}^{1.5} 2 dx$
 $= 0 + [x]_1^{\sqrt{2}} + [2x]_{\sqrt{2}}^{1.5}$
 $= \sqrt{2} - 1 + 2(1.5 - \sqrt{2})$
 $= \sqrt{2} - 1 + 3 - 2\sqrt{2}$
 $= 2 - \sqrt{2}$
13. $f\left(\frac{1}{x}\right) + x^2 f(x) = 0 \dots$ (given)
 $\therefore f(x) = -\frac{1}{x^2} f\left(\frac{1}{x}\right)$
 Let $I = \int_{\cos \theta}^{\sec \theta} f(x) dx = \int_{\cos \theta}^{\sec \theta} -\frac{1}{x^2} f\left(\frac{1}{x}\right) dx$
 Put $\frac{1}{x} = t$, $\therefore -\frac{1}{x^2} dx = dt$
 When $x = \cos \theta$, $t = \sec \theta$
 and when $x = \sec \theta$, $t = \cos \theta$
 $\therefore I = \int_{\sec \theta}^{\cos \theta} f(t) dt = -\int_{\cos \theta}^{\sec \theta} f(t) dt = -\int_{\cos \theta}^{\sec \theta} f(x) dx = -I$
 $\therefore I + I = 0$
 $\therefore 2I = 0$
 $\therefore I = 0$

14. $\lim_{n \rightarrow \infty} \frac{1}{n} \left[1 + \sqrt{\frac{n}{n+1}} + \sqrt{\frac{n}{n+2}} + \sqrt{\frac{n}{n+3}} + \dots + \sqrt{\frac{n}{n+3(n-1)}} \right]$
 $= \lim_{n \rightarrow \infty} \frac{1}{n} \left[\frac{1}{\sqrt{1+\frac{0}{n}}} + \frac{1}{\sqrt{1+\frac{1}{n}}} + \frac{1}{\sqrt{1+\frac{2}{n}}} + \dots + \frac{1}{\sqrt{1+\frac{3(n-1)}{n}}} \right]$
 $= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{r=0}^{3(n-1)} \frac{1}{\sqrt{1+\frac{r}{n}}}$
 $= \int_0^3 \frac{1}{\sqrt{1+x}} dx$
 $= [2\sqrt{1+x}]_0^3$
 $= 2(\sqrt{1+3} - \sqrt{1+0})$
 $= 2(2 - 1) = 2(1) = 2$
15. $\lim_{n \rightarrow \infty} \sum_{r=1}^{4n} \frac{\sqrt{n}}{\sqrt{r}(3\sqrt{r} + 4\sqrt{n})^2}$
 $= \lim_{n \rightarrow \infty} \sum_{r=1}^{4n} \frac{1}{n} \cdot \frac{1}{\sqrt{\frac{r}{n}}} \cdot \frac{1}{(3\sqrt{r} + 4\sqrt{n})^2}$
 $= \lim_{n \rightarrow \infty} \sum_{r=1}^{4n} \frac{1}{n} \cdot \frac{1}{\sqrt{\frac{r}{n}}} \cdot \frac{1}{\left(3\sqrt{\frac{r}{n}} + 4\right)^2}$
 $= \int_0^4 \frac{1}{\sqrt{x}(3\sqrt{x} + 4)^2} dx$
 Put $3\sqrt{x} + 4 = t$
 $\therefore 3 \cdot \frac{1}{2\sqrt{x}} dx = dt$
 $\therefore \frac{1}{\sqrt{x}} dx = \frac{2}{3} dt$
 When $x = 0$, $t = 4$ and when $x = 4$, $t = 10$
 $\therefore I = \int_4^{10} \frac{1}{t^2} \cdot \frac{2}{3} dt = -\frac{2}{3} \left[\frac{1}{t} \right]_4^{10}$
 $= -\frac{2}{3} \left(\frac{1}{10} - \frac{1}{4} \right) = -\frac{2}{3} \left(\frac{2-5}{20} \right)$
 $= -\frac{2}{3} \left(-\frac{3}{20} \right) = \frac{1}{10}$



16. $f'(x) = f(x) \Rightarrow \frac{f'(x)}{f(x)} = 1$

Integrating on both sides, we get

$\log f(x) = x + \log c \Rightarrow f(x) = ce^x$

$\therefore f(0) = c \Rightarrow c = 1$

$\therefore f(x) = e^x$

Now, $f(x) + g(x) = e^{2x}$

$\Rightarrow g(x) = e^{2x} - e^x$

$\therefore \int_0^1 f(x)g(x) dx = \int_0^1 e^x (e^{2x} - e^x) dx$
 $= \int_0^1 e^{3x} dx - \int_0^1 e^{2x} dx$
 $= \left[\frac{e^{3x}}{3} - \frac{e^{2x}}{2} \right]_0^1$
 $= e - \frac{1}{2}e^2 - \frac{3}{2}$

17. Let $I = \int_0^{100\pi} (|\sin^3 x| + |\cos^3 x|) dx$

$= \int_0^{200 \times \frac{\pi}{2}} (|\sin^3 x| + |\cos^3 x|) dx$

$= 200 \int_0^{\frac{\pi}{2}} (|\sin^3 x| + |\cos^3 x|) dx$

..... $\left[\begin{array}{l} \because |\sin^3 x| + |\cos^3 x| \text{ is a periodic} \\ \text{function with period } \frac{\pi}{2} \end{array} \right]$

$\therefore I = 200 \int_0^{\frac{\pi}{2}} (\sin^3 x + \cos^3 x) dx$

$= 200 \left[\int_0^{\frac{\pi}{2}} \sin^3 x dx + \int_0^{\frac{\pi}{2}} \cos^3 x dx \right]$

$= 200[I_1 + I_2]$ (Say)(i)

Where $I_1 = \int_0^{\frac{\pi}{2}} \sin^3 x dx$

$= -\int_0^{\frac{\pi}{2}} (1 - \cos^2 x)(-\sin x) dx$

Put $\cos x = t, \therefore -\sin x dx = dt$

When $x = 0, t = \cos 0 = 1$ and when $x = \frac{\pi}{2},$

$t = \cos \frac{\pi}{2} = 0$

$\therefore I_1 = -\int_1^0 (1 - t^2) dt$

$= \int_0^1 (1 - t^2) dt$

$= \left[t - \frac{t^3}{3} \right]_0^1 = 1 - \frac{1}{3} = \frac{2}{3}$

$I_2 = \int_0^{\frac{\pi}{2}} \cos^3 x dx = \int_0^{\frac{\pi}{2}} \cos^2 x \left(\frac{\pi}{2} - x \right) dx$

$= \int_0^{\frac{\pi}{2}} \sin^3 x dx = I_1 = \frac{2}{3}$

\therefore From (i),

$I = 200 \left(\frac{2}{3} + \frac{2}{3} \right) = \frac{800}{3}$

18. $I_1 = \int_0^{\frac{\pi}{2}} \frac{\sin x - \cos x}{1 - \sin x \cos x} dx$

$= \int_0^{\frac{\pi}{2}} \frac{\cos x - \sin x}{1 - \sin x \cos x} dx$

..... $\left[\begin{array}{l} \because \int_0^{\frac{\pi}{2}} f(x) dx = \int_0^{\frac{\pi}{2}} f\left(\frac{\pi}{2} - x\right) dx \end{array} \right]$

$= -\int_0^{\frac{\pi}{2}} \frac{\sin x - \cos x}{1 - \sin x \cos x} dx = -I_1$

$\therefore 2I_1 = 0 \Rightarrow I_1 = 0$

$\therefore I_2 = \int_0^{2\pi} \cos^6 x dx$

$= 2 \int_0^{\pi} \cos^6 x dx$

..... $\left[\because \cos^6 x \text{ is a periodic } f^n \text{ with period } \pi \right]$

$= 2 \int_0^{\frac{\pi}{2}} (\cos^6 x + \cos^6(\pi - x)) dx$

..... $\left[\because \int_0^{2a} f(x) dx = \int_0^a [f(x) + f(2a - x)] dx \right]$

$= 2.2 \int_0^{\frac{\pi}{2}} \cos^6 x dx$

$= 4 \cdot \frac{(6-1)(6-3)(6-5)}{6(6-2)(6-4)} \cdot \frac{\pi}{2} = \frac{5\pi}{8}$



$$\therefore I_2 \neq 0$$

$$I_3 = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^3 x \, dx = 0$$

....[$\because \sin^3$ is an odd function]

$$I_4 = \int_0^1 \log\left(\frac{1-x}{1+x}\right) dx$$

$$= \int_0^1 \log\left(\frac{1-x}{1+x}\right) dx$$

....[$\because \int_0^a f(x) dx = \int_0^a f(a-x) dx$]

$$= \int_0^1 \log\left(\frac{1-x}{1+x}\right) dx$$

$$= -\int_0^1 \log\left(\frac{1-x}{1+x}\right) dx$$

$$\therefore I_4 = -I_4$$

$$\therefore 2I_4 = 0$$

$$\therefore I_4 = 0$$

$$\therefore I_1 = I_3 = I_4 = 0, \text{ but } I_2 \neq 0$$

19. Let $I = \int_0^{2\pi} \frac{\sin^{2n} x}{\sin^{2n} x + \cos^{2n} x} dx$

$$= \int_0^{2\pi} \frac{(2\pi-x)\sin^{2n}(2\pi-x)}{\sin^{2n}(2\pi-x) + \cos^{2n}(2\pi-x)} dx$$

$$= 2\pi \int_0^{2\pi} \frac{\sin^{2n} x}{\sin^{2n} x + \cos^{2n} x} dx - I$$

$$\therefore I = \pi \times 2 \int_0^{\pi} \frac{\sin^{2n} x}{\sin^{2n} x + \cos^{2n} x} dx$$

($\because \frac{\sin^{2n} x}{\sin^{2n} x + \cos^{2n} x}$ is a periodic f^n with period π)

$$= 4\pi \int_0^{\frac{\pi}{2}} \frac{\sin^{2n} x}{\sin^{2n} x + \cos^{2n} x} dx$$

....[$\because \int_0^{2a} f(x) dx = 2 \int_0^a f(x) dx$
if $f(2a-x) = f(x)$]

$$= 4\pi \times \frac{\pi}{4}$$

$$= \pi^2$$

20. $\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \left\{ a |\sin x| + \frac{b \sin x}{1 + \cos x} + c \right\} dx = 0$

i.e., $I_1 + I_2 + I_3 = 0$

$$I_1 = a \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} |\sin x| dx$$

$$= a \left[\int_{-\frac{\pi}{4}}^0 |\sin x| dx + \int_0^{\frac{\pi}{4}} |\sin x| dx \right]$$

$$= a \left[\int_{-\frac{\pi}{4}}^0 (-\sin x) dx + \int_0^{\frac{\pi}{4}} \sin x dx \right]$$

$$= a \left[[\cos x]_{-\pi/4}^0 - [\cos x]_0^{\pi/4} \right]$$

$$= a \left(1 - \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} + 1 \right) = a(2 - \sqrt{2})$$

$$I_2 = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{b \sin x}{1 + \cos x} dx$$

$$= [-b \log |1 + \cos x|]_{-\pi/4}^{\pi/4}$$

$$= 0 \quad \dots \left[\because \cos\left(\frac{-\pi}{4}\right) = \cos\frac{\pi}{4} \right]$$

$$I_3 = c \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} 1 dx = c \left[\frac{\pi}{4} + \frac{\pi}{4} \right] = \frac{c\pi}{2}$$

$\therefore I_1 + I_2 + I_3 = 0$ becomes

$$a(2 - \sqrt{2}) + \frac{c\pi}{2} = 0 \quad \dots [\because I_2 = 0]$$

\therefore The given equation is a relation between a and c .

21. Let $I = \int_0^{\sqrt{\log\left(\frac{\pi}{2}\right)}} \cos(e^{-t}) \cdot 2e^{-2t} dt$

Put $e^{-t} = t \Rightarrow 2e^{-2t} dt = dt$

When $t = 0, t = e^0 = 1$

When $t = \sqrt{\log\frac{\pi}{2}}, t = e^{\log\pi/2} = \frac{\pi}{2}$

$\therefore I = \int_1^{\frac{\pi}{2}} \cos t dt = [\sin t]_1^{\pi/2}$

$$= \sin \frac{\pi}{2} - \sin 1 = 1 - \sin 1$$



$$\begin{aligned}
 22. \quad f(\pi) &= \int_0^{\pi} \sin^6 t \, dt \\
 \therefore f(\pi + \pi) &= \int_0^{+\pi} \sin^6 t \, dt = \int_0^{\pi} \sin^6 t \, dt + \int_{\pi}^{+\pi} \sin^6 t \, dt \\
 &= f(\pi) + \int_0^{\pi} \sin^6(u + \pi) \, du, \text{ where } t = u + \pi \\
 &= f(\pi) + \int_0^{\pi} \sin^6 u \, du \\
 &= f(\pi) + \int_0^{\pi} \sin^6 t \, dt = f(\pi) + f(\pi) \\
 \therefore f(\pi + \pi) &= f(\pi) + f(\pi)
 \end{aligned}$$

$$\begin{aligned}
 23. \quad \text{Let } f(x) &= ax^2 + bx + c \\
 \therefore f'(x) &= 2ax + b \\
 f''(x) &= 2a \\
 f(0) &= c = 3 \\
 f'(0) &= b = -7 \\
 f''(0) &= 2a = 8 \\
 \therefore a &= 4 \\
 \therefore f(x) &= 4x^2 - 7x + 3 \\
 \int_1^2 f(x) \, dx &= \int_1^2 (4x^2 - 7x + 3) \, dx \\
 &= \left[\frac{4x^3}{3} - \frac{7x^2}{2} + 3x \right]_1^2 \\
 &= \frac{32}{3} - 14 + 6 - \left(\frac{4}{3} - \frac{7}{2} + 3 \right) \\
 &= \frac{32 - 42 + 18}{3} - \left(\frac{8 - 21 + 18}{6} \right) \\
 &= \frac{8}{3} - \frac{5}{6} = \frac{16 - 5}{6} = \frac{11}{6}
 \end{aligned}$$

$$\begin{aligned}
 24. \quad \int_0^1 f(x) \, dx &= \frac{2A}{\pi} \\
 \Rightarrow \int_0^1 \left[A \sin\left(\frac{\pi}{2}x\right) + B \right] \, dx &= \frac{2A}{\pi} \\
 \Rightarrow \left[-\frac{2A}{\pi} \cos\left(\frac{\pi}{2}x\right) + Bx \right]_0^1 &= \frac{2A}{\pi} \\
 \Rightarrow \frac{2A}{\pi} + B = \frac{2A}{\pi} &\Rightarrow B = 0 \\
 \text{Now, } f(x) &= A \sin\left(\frac{\pi}{2}x\right) + B \\
 \Rightarrow f'(x) &= A \cos\left(\frac{\pi}{2}x\right) \cdot \frac{\pi}{2}
 \end{aligned}$$

$$\begin{aligned}
 \Rightarrow f'\left(\frac{1}{2}\right) &= \frac{\pi A}{2} \cdot \frac{1}{\sqrt{2}} \\
 \Rightarrow \sqrt{2} &= \frac{\pi A}{2\sqrt{2}} \Rightarrow A = \frac{4}{\pi}
 \end{aligned}$$

$$\begin{aligned}
 25. \quad f(x) &= \begin{vmatrix} \sin x + \sin 2x + \sin 3x & \sin 2x & \sin 3x \\ 3 + 4\sin x & 3 & 4\sin x \\ 1 + \sin x & \sin x & 1 \end{vmatrix} \\
 &= \begin{vmatrix} \sin x & \sin 2x & \sin 3x \\ 0 & 3 & 4\sin x \\ 0 & \sin x & 1 \end{vmatrix} \quad (C_1 \rightarrow C_1 - C_2 - C_3) \\
 &= \sin x (3 - 4\sin^2 x) = 3\sin x - 4\sin^3 x = \sin 3x \\
 \therefore \int_0^{\frac{\pi}{2}} f(x) \, dx &= \int_0^{\frac{\pi}{2}} \sin 3x \, dx \\
 &= -\frac{1}{3} [\cos 3x]_0^{\frac{\pi}{2}} \\
 &= -\frac{1}{3} \left[\cos \frac{3\pi}{2} - \cos 0 \right] \\
 &= -\frac{1}{3} (0 - 1) = \frac{1}{3}
 \end{aligned}$$

$$\begin{aligned}
 26. \quad \int_0^{\frac{\pi}{2}} \left\{ a^2 \left(\frac{\cos 3t}{4} + \frac{3}{4} \cos t \right) + a \sin t - 20 \cos t \right\} \, dt \\
 &= \left[a^2 \left(\frac{1}{12} \sin 3t + \frac{3}{4} \sin t \right) - a \cos t - 20 \sin t \right]_0^{\frac{\pi}{2}} \\
 &= a^2 \left(\frac{1}{12} (-1) + \frac{3}{4} (1) - 0 \right) - a(0 - 1) - 20(1 - 0) \\
 &= a^2 \left(\frac{3}{4} - \frac{1}{12} \right) + a - 20 \\
 &= \frac{2a^2}{3} + a - 20 \\
 \therefore \text{from the given condition,} \\
 \frac{2a^2}{3} + a - 20 &\leq \frac{-a^2}{3} \\
 \therefore \frac{3a^2}{3} + a - 20 &\leq 0 \\
 \therefore a^2 + a - 20 &\leq 0 \\
 \therefore (a + 5)(a - 4) &\leq 0 \\
 \therefore -5 \leq a &\leq 4 \\
 \text{The positive integer values of } a \text{ satisfying the} \\
 \text{above inequality are } 1, 2, 3, 4. \\
 \therefore \text{There are 4 such values.}
 \end{aligned}$$



27. Since, $-1 \leq \sin \leq 1 \Rightarrow -2 \leq 2\sin \leq 2$

$$\begin{aligned} \int_{\frac{\pi}{2}}^{\frac{3\pi}{2}} [2\sin] d &= \int_{\frac{\pi}{2}}^{\frac{5\pi}{6}} [2\sin] d + \int_{\frac{5\pi}{6}}^{\pi} [2\sin] d \\ &+ \int_{\pi}^{\frac{7\pi}{6}} [2\sin] d + \int_{\frac{7\pi}{6}}^{\frac{3\pi}{2}} [2\sin] d \\ &= \int_{\frac{\pi}{2}}^{\frac{5\pi}{6}} (1) d + \int_{\frac{5\pi}{6}}^{\pi} (0) d + \int_{\pi}^{\frac{7\pi}{6}} (-1) d + \int_{\frac{7\pi}{6}}^{\frac{3\pi}{2}} (-2) d \\ &= \left(\frac{5\pi}{6} - \frac{\pi}{2} \right) + 0 - \left(\frac{7\pi}{6} - \pi \right) - 2 \left(\frac{3\pi}{2} - \frac{7\pi}{6} \right) \\ &= \frac{2\pi}{6} - \frac{\pi}{6} - \frac{4\pi}{6} = -\frac{\pi}{2} \end{aligned}$$

28. Applying $R_1 \rightarrow R_1 - \sec R_3$, we get

$$\begin{aligned} f(x) &= \begin{vmatrix} 0 & 0 & \sec^2 + \cot \operatorname{cosec} - \cos \\ \cos^2 & \cos^2 & \operatorname{cosec}^2 \\ 1 & \cos^2 & \cos^2 \end{vmatrix} \\ &= (\sec^2 + \cot \operatorname{cosec} - \cos)(\cos^4 - \cos^2) \\ &= (\sec^2 + \cot \operatorname{cosec} - \cos)(-\cos^2 \sin^2) \\ &= -\sin^2 - \cos^3 + \cos^3 \sin^2 \\ &= -\sin^2 - \cos^3 (1 - \sin^2) \\ &= -\sin^2 - \cos^5 \end{aligned}$$

$$\begin{aligned} \therefore \int_0^{\frac{\pi}{2}} f(x) dx &= -\int_0^{\frac{\pi}{2}} (\sin^2 + \cos^5) dx \\ &= -\left(\frac{1}{2} \times \frac{\pi}{2} + \frac{4.2}{5.3.1} \right) = -\frac{\pi}{4} - \frac{8}{15} \end{aligned}$$

29. $\frac{1}{\sqrt{a}} \int_1^a \left(\frac{3}{2} \sqrt{x} + 1 - \frac{1}{\sqrt{x}} \right) dx < 4$

$$\therefore \frac{1}{\sqrt{a}} \left[\frac{3}{2} \cdot \frac{3}{2} x^{3/2} + x - 2\sqrt{x} \right]_1^a < 4$$

$$\therefore \frac{1}{\sqrt{a}} [a\sqrt{a} - 1 + a - 1 - 2\sqrt{a} + 2] < 4$$

$$\therefore a + \sqrt{a} - 2 < 4$$

$$\therefore a + \sqrt{a} - 6 < 0$$

$$\therefore (\sqrt{a} + 3)(\sqrt{a} - 2) < 0$$

$$\therefore -3 < \sqrt{a} < 2$$

But \sqrt{a} cannot be negative and according to the problem, $a \neq 0$

$$\therefore 0 < \sqrt{a} < 2$$

$$\therefore 0 < a < 4$$

30. Let $I = \int_1^4 \frac{3e^{\sin^3}}{3} dx = \int_1^4 \frac{3^2 e^{\sin^3}}{3} dx$

Put $\sin^3 = t \Rightarrow 3^2 dx = dt$

$$\begin{aligned} \therefore I &= \int_1^{64} \frac{e^{\sin t}}{t} dt \\ &= \int_1^{64} \frac{e^{\sin}}{1} dx \end{aligned}$$

$$= [f(x)]_1^{64} \quad \dots \left[\because \frac{d}{dx} [f(x)] = \frac{e^{\sin}}{1} \right]$$

$$= f(64) - f(1)$$

$$\therefore k = 64$$



Hints



Classical Thinking

1. Required area = $\int_1^4 x^3 dx = \left[\frac{x^4}{4} \right]_1^4 = \frac{255}{4}$ sq. units


2. Required area = $\int_1^4 \frac{1}{x} dx = c \int_1^4 \frac{1}{x} dx$
= $2c \log 2$ sq. units.

3. Required area = $\int_0^4 \sqrt{3x+4} dx$
= $\left[\frac{(3x+4)^{3/2}}{3 \cdot \frac{3}{2}} \right]_0^4$
= $\frac{2}{9} \times 56 = \frac{112}{9}$ sq. units

4. Required area = $\int_2^4 \left(1 + \frac{8}{x}\right) dx = \left[x - \frac{8}{x} \right]_2^4$
= $(4-2) - (2-4)$
= $2+2=4$

5. Required area = $\int_1^2 x dx = \int_1^2 \log x dx$
= $\left[x \log x - x \right]_1^2$
= $2 \log 2 - 1$
= $(\log 4 - 1)$ sq. units

6.



Required area = $\int_0^{\pi/2} \sin x dx$
= $[-\cos x]_0^{\pi/2}$
= $-\left(\cos \frac{\pi}{2} - \cos 0\right)$
= 1 sq. unit

7. Required area = $\int_0^2 (4-x^2) dx$
= $\left[2x^2 - \frac{x^3}{3} \right]_0^2$
= $8 - \frac{8}{3}$
= $\frac{16}{3}$

8. Required area = $\int_0^{\pi/2} (2 + \sin x) dx$
= $\left[2x - \cos x \right]_0^{\pi/2}$
= $\left(\frac{\pi^2}{4} - \cos \frac{\pi}{2} \right) - (0 - \cos 0)$
= $\frac{\pi^2}{4} - 0 - (0 - 1)$
= $\frac{\pi^2}{4} + 1$

9. Required area = $\int_0^a x dx = \int_0^a e^{2x} dx$

Put $2x = t \Rightarrow dx = \frac{dt}{2}$

\therefore required area = $\frac{1}{2} \int_0^{a^2} e^t dt$
= $\frac{1}{2} [e^t]_0^{a^2}$
= $\frac{e^{a^2} - 1}{2}$ sq. units




10. Required area = $\int_0^{\pi/2} \sin^2 x \, dx$
 $= \int_0^{\pi/2} \left(\frac{1 - \cos 2x}{2} \right) dx$
 $= \frac{1}{2} \left[x \right]_0^{\pi/2} - \frac{1}{4} \left[\sin 2x \right]_0^{\pi/2} = \frac{\pi}{4}$


11. Required area = $\int_0^2 \frac{x^2}{4} dx$
 $= \frac{1}{12} \left[x^3 \right]_0^2$
 $= \frac{8}{12}$
 $= \frac{2}{3}$



Critical Thinking



Required area = $\int_1^4 x \, dx = \int_1^4 \frac{\sqrt{x}}{2} dx$
 $= \frac{1}{2} \cdot \frac{2}{3} \left[x^{3/2} \right]_1^4$
 $= \frac{7}{3}$ sq. units



$x^2 = -2 \Rightarrow x = \sqrt{-2}$

\therefore required area = $\int_4^6 \sqrt{-2} dx = \left[\frac{(-2)^{3/2}}{\frac{3}{2}} \right]_4^6$
 $= \frac{2}{3} \left[4^{3/2} - 2^{3/2} \right]$
 $= \frac{2}{3} (8 - 2\sqrt{2})$

3. Required area = $\int_0^4 2\sqrt{x} \, dx$

$= 2 \left[\frac{3}{2} x^{3/2} \right]_0^4 = \frac{4}{3} \left[4^{3/2} - 0 \right]$
 $= \frac{4}{3} (8) = \frac{32}{3}$

4. $A_1 = \int_0^{\pi/3} \cos x \, dx = \frac{\sqrt{3}}{2}$

$A_2 = \int_0^{\pi/3} \cos 2x \, dx = \frac{\sqrt{3}}{4}$

$\therefore A_1 : A_2 = 2 : 1$

5. $-3x - 2x - 10 = 0$

$\Rightarrow (-x) = 3x + 10$

$\Rightarrow x = \frac{3x + 10}{-2}$

\therefore Required area = $\int_3^4 x \, dx = \int_3^4 \frac{3x + 10}{-2} dx$
 $= \left[\frac{3}{2}x^2 + 5x \right]_3^4$
 $= 3 + 16 \log 2$ sq. units

6. According to the given condition,

$\int_1^3 (3x^2 - 4x + k) dx = 20$

$\Rightarrow \left[x^3 - 2x^2 + kx \right]_1^3 = 20$

$\Rightarrow (27 - 18 + 3k) - (1 - 2 + k) = 20$

$\Rightarrow 9 + 3k + 1 - k = 20 \Rightarrow 2k = 10$

$\Rightarrow k = 5$

7. For X-axis, $y = 0$

$\therefore 4 + 3x - x^2 = 0$

$\Rightarrow (x + 1)(x - 4) = 0 \Rightarrow x = -1$ or $x = 4$


\therefore Required area = $\int_{-1}^4 (4 + 3x - x^2) dx = \frac{125}{6}$





8. For Y-axis, $y = 0$
 $\therefore x^2 - 1 = 0$
 $\Rightarrow (x - 1)(x + 1) = 0$
 $\Rightarrow x = 0$ or $x = 1$
 \therefore required area $= \int_0^1 (x^2 - 1) dx = \left[\frac{x^3}{3} - x \right]_0^1$
 $= \frac{1}{3} - 1 = \left| \frac{-2}{3} \right| = \frac{2}{3}$ sq. units


9. For X-axis, $y = 0$
 $\therefore 4 - x^2 = 0$
 $\Rightarrow (2 - x)(2 + x) = 0 \Rightarrow x = 0, 2$
 Required area $= \int_0^2 (4 - x^2) dx$
 $= \left[4x - \frac{x^3}{3} \right]_0^2$
 $= 8 - \frac{8}{3} = \frac{16}{3}$ sq. units

10. According to the given condition,
 $\int_1^b f(x) dx = (b - 1) \sin(3b + 4)$
 Differentiating w.r.t.b, we get
 $f(b) \cdot 1 = 3(b - 1) \cos(3b + 4) + \sin(3b + 4)$
 $\therefore f(x) = 3(x - 1) \cos(3x + 4) + \sin(3x + 4)$

11. 
 Required area $= \int_0^{2a} \sqrt{2a - x^3} dx$
 Put $x = 2a \sin^2 \theta$
 $\Rightarrow dx = 4a \sin \theta \cos \theta d\theta$
 \therefore required area $= \int_0^{\frac{\pi}{2}} \frac{\sqrt{8a^3 \sin^6 \theta} \cdot 4a \sin \theta \cos \theta}{\sqrt{2a - 2a \sin^2 \theta}} d\theta$
 $= \int_0^{\frac{\pi}{2}} \frac{2a \cdot \sin^3 \theta \cdot 4a \sin \theta \cos \theta}{\cos \theta} d\theta$
 $= 8a^2 \int_0^{\frac{\pi}{2}} \sin^4 \theta d\theta = 8a^2 \cdot \frac{3.1}{4.2} \cdot \frac{\pi}{2} = \frac{3\pi a^2}{2}$


12. 
 Since, the curve is symmetrical about X-axis.
 \therefore Required area $= 2 \int_0^4 \sqrt{x} dx$
 $= 2 \int_0^4 x^{\frac{1}{2}} dx = \frac{28\sqrt{2}}{3}$ sq. units

13. 
 Since, the curve is symmetrical about X-axis.
 \therefore Required area $= 2 \int_0^2 \sqrt{8 - x^2} dx$
 $= 2 \int_0^2 \sqrt{8} dx = 4\sqrt{2} \int_0^2 \sqrt{1 - \frac{x^2}{4}} dx$
 $= 4\sqrt{2} \left[\frac{3/2}{2} \right]_0^2 = \frac{8\sqrt{2}}{3} (2\sqrt{2})$
 $= \frac{32}{3}$ sq. units

14. 
 Required area $= 2 \int_0^a \sqrt{4a - x^2} dx$
 $= 2 \times 2\sqrt{a} \times \frac{2}{3} \left[\frac{3/2}{2} \right]_0^a$
 $= \frac{8}{3} a^2$ sq. units




15.



$$\begin{aligned} \text{Required area} &= 2 \int_0^4 \sqrt{4-x} \, dx = 2 \left[-\frac{2}{3}(4-x)^{3/2} \right]_0^4 \\ &= 2 \left[0 + \frac{2}{3}(4)^{3/2} \right] = \frac{4 \times 8}{3} \\ &= \frac{32}{3} \text{ sq. units} \end{aligned}$$


16.



$$\begin{aligned} \text{Required area} &= 4(\text{area of the region OABO}) \\ &= 4 \int_0^1 (1-x^2) \, dx \\ &= 4 \left[x - \frac{x^3}{3} \right]_0^1 \\ &= \frac{8}{3} \end{aligned}$$


17. Required area = $2 \int_0^{\pi/3} \tan x \, dx$
 $= 2 [\log \sec x]_0^{\pi/3} = 2 \log(2)$

18.




$$\begin{aligned} \text{Required area} &= A_1 + A_2 + A_3 + A_4 \\ &= 4 A_1 \\ &= 4 \int_0^{\pi/2} \cos x \, dx \\ &= 4 [\sin x]_0^{\pi/2} = 4 \left(\sin \frac{\pi}{2} - \sin 0 \right) \\ &= 4(1 - 0) \\ &= 4 \end{aligned}$$

19.



$$\begin{aligned} \text{Required area} &= 4 \int_0^{\pi/2} \sin x \, dx \\ &= 4 [-\cos x]_0^{\pi/2} \\ &= -4 \left(\cos \frac{\pi}{2} - \cos 0 \right) \\ &= 4 \text{ sq. units} \end{aligned}$$


20.



$$\begin{aligned} \text{Area of smaller part} &= 2 \int_1^2 \sqrt{4-x^2} \, dx \\ &= 2 \left[\frac{x}{2} \sqrt{4-x^2} + 2 \sin^{-1} \frac{x}{2} \right]_1^2 \\ &= 2 \left[2 \cdot \frac{\pi}{2} - \left(\frac{\sqrt{3}}{2} + 2 \cdot \frac{\pi}{6} \right) \right] \\ &= \frac{4\pi}{3} - \sqrt{3} \end{aligned}$$



21.



Since, the curve is symmetrical about Y-axis as well as X-axis.

∴ the area of the given ellipse

$$= 4(\text{area of OABO}) = 4 \int_0^a \frac{b}{a} \sqrt{a^2 - x^2} dx$$

$$= 4 \int_0^{\frac{\pi}{2}} \frac{b}{a} \cdot a \cos \theta \cdot a \sin \theta d\theta \quad \dots [\text{Put } x = a \sin \theta]$$

$$= 4ab \int_0^{\frac{\pi}{2}} \left(\frac{1 + \cos 2\theta}{2} \right) d\theta$$

$$= 2ab \left[\theta \right]_0^{\pi/2} + \left[\frac{\sin 2\theta}{2} \right]_0^{\pi/2} = \pi ab \text{ sq. units}$$

22. $16x^2 + 9y^2 = 144 \Rightarrow \frac{x^2}{9} + \frac{y^2}{16} = 1$

Area of ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is πab sq. units.

Here, $a = 3, b = 4$


∴ Required area = $\pi ab = \pi (3)(4) = 12\pi$

23. Required area = $\int_{-1}^1 |x| dx$

$$= \left| \int_{-1}^0 -x dx \right| + \int_0^1 x dx$$

$$= \left| \frac{-x^2}{2} \right|_{-1}^0 + \frac{x^2}{2} \Big|_0^1 = \frac{1}{2} + \frac{1}{2} = 1$$

24.




Required area = $\left| \int_{-1}^{-\frac{2}{3}} (3x+2) dx \right| + \int_{-\frac{2}{3}}^1 (3x+2) dx$

$$= \left| \left[\frac{3x^2}{2} + 2x \right]_{-1}^{-\frac{2}{3}} \right| + \left[\frac{3x^2}{2} + 2x \right]_{-\frac{2}{3}}^1$$

$$= \frac{1}{6} + \frac{25}{6} = \frac{13}{3} \text{ sq. units.}$$

25.




Required area = $\int_0^{\sqrt{3}} \sqrt{4-x^2} dx + \int_{\sqrt{3}}^2 \sqrt{4-x^2} dx$

$$= \frac{1}{\sqrt{3}} \left[\frac{x^2}{2} \right]_0^{\sqrt{3}} + \left[-\frac{1}{2} \sqrt{4-x^2} + \frac{4}{2} \sin^{-1} \frac{x}{2} \right]_{\sqrt{3}}^2$$

$$= \frac{\sqrt{3}}{2} + \left[\pi - \frac{\sqrt{3}}{2} - \frac{2\pi}{3} \right] = \frac{\pi}{3}$$

26.




Required area = $\int_0^2 [(2x+2) - (2-x)] dx$

$$= \left[\frac{x^2}{2} + 2x \right]_0^2 = 4 \text{ sq. units}$$



27.




$$\begin{aligned} \text{Required area} &= \int_0^3 [(7x + 1) - (5x + 1)] dx \\ &= \int_0^3 2 dx \\ &= 2 \left[\frac{x^2}{2} \right]_0^3 = 9 \text{ sq. units} \end{aligned}$$

28. The curves $y = \cos x$ and $y = 1 + \sin x$ intersect at $(0, 0)$ and (π, π) .

\therefore required area


$$\begin{aligned} &= \int_0^\pi (\cos x + \sin x) dx - \int_0^\pi 1 dx = \int_0^\pi \sin x dx \\ &= [-\cos x]_0^\pi = -\cos \pi + \cos 0 \\ &= -(-1) + 1 = 2 \end{aligned}$$

29. The two curves intersect at $(0, 0)$ and $(1, 1)$.




$$\begin{aligned} \text{Required area} &= \int_0^1 (\sqrt{x} - x^2) dx \\ &= \left[\frac{2}{3} x^{3/2} - \frac{x^3}{3} \right]_0^1 = \frac{1}{3} \end{aligned}$$

30. The two parabolas intersect at $(0, 0)$ and $(4, 4)$.



$$\begin{aligned} \text{Required area} &= \int_0^4 \left(2\sqrt{x} - \frac{x^2}{4} \right) dx \\ &= 2 \left[\frac{2}{3} x^{3/2} \right]_0^4 - \frac{1}{4} \left[\frac{x^3}{3} \right]_0^4 \\ &= \frac{32}{3} - \frac{16}{3} = \frac{16}{3} \text{ sq. units} \end{aligned}$$

31. The two curves intersect at $O(0, 0)$ and $P\left(\frac{1}{a}, \frac{1}{a}\right)$.



According to the given condition,

$$\begin{aligned} \int_0^{1/a} (\sqrt{x} - x^2) dx &= 1 \\ \Rightarrow \left[\frac{2}{3\sqrt{a}} x^{3/2} - \frac{x^3}{3} \right]_0^{1/a} &= 1 \\ \Rightarrow \frac{2}{3\sqrt{a}} \times \frac{1}{a^{3/2}} - \frac{1}{3} \times \frac{1}{a^3} &= 1 \\ \Rightarrow \frac{2}{3a^2} - \frac{1}{3a^2} &= 1 \Rightarrow \frac{1}{3a^2} = 1 \\ \Rightarrow a &= \frac{1}{\sqrt{3}} \quad \dots [\because a > 0] \end{aligned}$$




32. The area of the region bounded by $y^2 = 4a$ and $y^2 = 4b$ is $\frac{16ab}{3}$ sq. units.

Given parabolas are $y^2 = \frac{9}{4}$ and $y^2 = \frac{16}{3}$

Here, $a = \frac{9}{16}$, $b = \frac{4}{3}$


\therefore Required area = $\frac{16}{3} \times \frac{9}{16} \times \frac{4}{3}$
= 4 sq. units

33.



$$\begin{aligned} \text{Required area} &= \int_{-2}^2 (x^2 + 4 - 2x^2) dx \\ &= \int_{-2}^2 (4 - x^2) dx = \left[4x - \frac{x^3}{3} \right]_{-2}^2 \\ &= \left(8 - \frac{8}{3} \right) - \left(-8 + \frac{8}{3} \right) = \frac{32}{3} \end{aligned}$$

34. The two curves intersect at (2, 1) and (-2, 1).



$$\begin{aligned} \text{Required area} &= 2 \int_0^2 \left(\frac{6 - x^2}{2} - \frac{x^2}{4} \right) dx \\ &= 2 \int_0^2 \left(3 - \frac{3x^2}{4} \right) dx = 6 \left[x - \frac{x^3}{12} \right]_0^2 \\ &= 6 \left(2 - \frac{8}{12} \right) = 6 \times \frac{16}{12} \\ &= 8 \text{ sq. units} \end{aligned}$$

35. The area of the region bounded by the parabola $y^2 = 4a$ and the line $y = m$ is $\frac{8a^2}{3m^3}$ sq. units.

36. The area bounded by $y^2 = 4a$ and the line $y = m$ is $\frac{8a^2 m^3}{3}$.

Given, $y^2 = 2 \Rightarrow y^2 = 4 \left(\frac{1}{2} \right)$ and $m = 3$

Here, $a = \frac{1}{2}$ and $m = 3$

\therefore Required area = $\frac{8}{3} \times \frac{1}{4} \times 3 \times 3 \times 3 = 18$ sq. units

37. Given curves are $y = x^2$ and $y = x$.
On solving, we get $x = 0$, $x = 1$

\therefore Required area = $\int_0^1 (x - x^2) dx$
= $\left[\frac{x^2}{2} - \frac{x^3}{3} \right]_0^1$
= $\frac{1}{2} - \frac{1}{3} = \frac{1}{6}$

38. $y^2 = x$ and $2y = x$

$\therefore \left(\frac{x}{2} \right)^2 = x \Rightarrow x^2 = 4x \Rightarrow x = 0, 4$


\therefore Required area = $\int_0^4 \left(\sqrt{x} - \frac{x}{2} \right) dx$
= $\left[\frac{2x^{3/2}}{3} - \frac{x^2}{4} \right]_0^4$
= $\frac{4}{3}$ sq. units

39. Given curves are $y = x^3$ and $y = \sqrt{x}$.
On solving, we get $x = 0$, $x = 1$

\therefore Required area = $\int_0^1 (\sqrt{x} - x^3) dx$
= $\left[\frac{2x^{3/2}}{3} - \frac{x^4}{4} \right]_0^1$
= $\frac{2}{3} - \frac{1}{4} = \frac{5}{12}$



40.




Required area = 4(area of the region OABO)

$$= 4 \int_0^1 (1 - x^2) dx$$

$$= 4 \left[x - \frac{x^3}{3} \right]_0^1$$

$$= \frac{8}{3}$$

41.




Required area = $\int_0^1 [(2 - x^2) - x^2] dx$

$$= \int_0^1 (2 - 2x^2) dx$$

$$= \left[2x - \frac{2x^3}{3} \right]_0^1 = \frac{1}{3}$$

42. The points of intersection of $x^2 = 4$ and $y = 4 - 2x^2$ are $(2, 1)$ and $(-1, \frac{1}{4})$.



Required area

$$= \int_{-1}^2 \frac{1}{4} (x^2 + 2) dx - \int_{-1}^2 \frac{1}{4} x^2 dx$$

$$= \frac{1}{4} \left[\frac{x^3}{3} + 2x \right]_{-1}^2 - \frac{1}{4} \left[\frac{x^3}{3} \right]_{-1}^2 = \frac{9}{8} \text{ sq. units}$$


43. The two curves $x^2 = 4a$ and $y = m$ intersect at $(0, 0)$ and $(\frac{4a}{m^2}, \frac{4a}{m})$.

According to the given condition,

$$\int_0^{\frac{4a}{m^2}} (\sqrt{4a} - m) dx = \frac{a^2}{3}$$

$$\Rightarrow \frac{8}{3} \cdot \frac{a^2}{m^3} = \frac{a^2}{3} \Rightarrow m^3 = 8 \Rightarrow m = 2$$


44.



Required area = $\int_0^1 (e^x - e^{-x}) dx$

$$= [e^x + e^{-x}]_0^1 = e + \frac{1}{e} - 2$$

45.



Required area = $\int_0^2 [2 - (2 - x^2)] dx$


$$= \left[\frac{2}{\log 2} x^2 + \frac{x^3}{3} \right]_0^2$$

$$= \frac{4}{\log 2} - 4 + \frac{8}{3} - \frac{1}{\log 2}$$

$$= \frac{3}{\log 2} - \frac{4}{3}$$




46.



$$\begin{aligned} \text{Required area} &= \int_1^e [\log_e x - (\log_e x)^2] dx \\ &= \int_1^e \log_e x \, dx - \int_1^e (\log_e x)^2 \, dx \\ &= [x \log_e x - x]_1^e - \left[\frac{(\log_e x)^3}{3} - 2 \log_e x + 2x \right]_1^e \\ &= [e - e - (-1)] - \left[\frac{e}{3} - 2e + 2e - \left(\frac{1}{3} - 2 + 2 \right) \right] \\ &= 1 - (e - 2) = 3 - e \end{aligned}$$

47.



$$\begin{aligned} \text{Required area} &= \int_0^2 (1 - |x - 1|) dx \\ &= \int_0^1 (1 - (1 - x)) dx + \int_1^2 (1 - (x - 1)) dx \\ &= \int_0^1 x dx + \int_1^2 (2 - x) dx \\ &= \left[\frac{x^2}{2} \right]_0^1 + \left[2x - \frac{x^2}{2} \right]_1^2 \\ &= \frac{1}{2} + \left[4 - 2 - \left(2 - \frac{1}{2} \right) \right] \\ &= \frac{1}{2} + \left(\frac{1}{2} \right) = 1 \end{aligned}$$

48. The area between $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and the straight

line $\frac{x}{a} + \frac{y}{b} = 1$ is $\frac{1}{4} \pi ab - \frac{1}{2} ab$ sq. units.

Here, $a = 3, b = 2$

$$\begin{aligned} \therefore \text{Required area} &= \frac{1}{4} \pi(3)(2) - \frac{1}{2} (3)(2) \\ &= \frac{3}{2} (\pi - 2) \text{ sq. units} \end{aligned}$$



Competitive Thinking

1. Required area = $\int_3^5 (3 - 5) dx$

$$= \frac{3}{2} [x^2]_3^5 - 5[x]_3^5 = \frac{3}{2} (25 - 9) - 5(5 - 3)$$


$$= 24 - 10 = 14 \text{ sq. units}$$

2. According to the given condition,

$$\int_1^2 m x^2 dx = 6 \Rightarrow m \left[\frac{x^3}{3} \right]_1^2 = 6$$

$$\Rightarrow m \left(\frac{8}{3} - \frac{1}{3} \right) = 6 \Rightarrow m = 4$$

3.




$$\begin{aligned} \text{Required area} &= 2 \int_0^1 (1 - x) dx = 2 \left[x - \frac{x^2}{2} \right]_0^1 \\ &= 2 \left[1 - \frac{1}{2} \right] = 1 \text{ sq. unit.} \end{aligned}$$

Alternate method:

$$\begin{aligned} \text{Required area} &= \frac{1}{2} \times \text{base} \times \text{height} \\ &= \frac{1}{2} \times 2 \times 1 = 1 \text{ sq. unit.} \end{aligned}$$


4.



$$\begin{aligned} \text{Area bounded by the lines} &= \int_0^4 (3x + 1 - 2x - 1) dx \\ &= \int_0^4 x dx = \left[\frac{x^2}{2} \right]_0^4 \\ &= 8 \text{ sq. units} \end{aligned}$$



5.




$$\begin{aligned} \text{Required area} &= \left| \int_{-1}^0 d \right| + \int_0^2 d \\ &= \left| \int_{-1}^0 d \right| + \int_0^2 d = \left| \frac{-1}{2} \right| + 2 \\ &= 2 + \frac{1}{2} = \frac{5}{2} \text{ sq. units} \end{aligned}$$

Alternate method:

$$\begin{aligned} \text{Required area} &= \frac{1}{2} \times 1 \times 1 + \frac{1}{2} \times 2 \times 2 \\ &= \frac{1}{2} + 2 \\ &= \frac{5}{2} \text{ sq. units.} \end{aligned}$$

6.




$$\begin{aligned} \text{Required area} &= \int_0^8 d = \int_0^8 x^{1/3} dx \\ &= \frac{3}{4} \left[x^{4/3} \right]_0^8 = \frac{3}{4} (8^{4/3} - 0) \\ &= 12 \end{aligned}$$

7.

$$\begin{aligned} \text{Required area} &= \int_1^3 |x - 2| dx \\ &= \int_1^2 (2 - x) dx + \int_2^3 (x - 2) dx \\ &= \left[2x - \frac{x^2}{2} \right]_1^2 + \left[\frac{x^2}{2} - 2x \right]_2^3 \\ &= \frac{1}{2} + \frac{1}{2} = 1 \end{aligned}$$

8.



$$\begin{aligned} \text{Required area} &= \int_{1-e}^0 \log_e(x+e) dx \\ &= \int_1^e \log t dt \quad \dots [\text{Put } x+e=t] \\ &= [t \log t - t]_1^e = 1 \text{ sq. unit} \end{aligned}$$

9. For X-axis, $y = 0$

$$\therefore 2 - x^2 = 0$$

$$\Rightarrow (2 - x) = 0 \Rightarrow x = 0, 2$$

$$\begin{aligned} \text{Required area} &= \int_0^2 (2 - x^2) dx \\ &= \left[2x - \frac{x^3}{3} \right]_0^2 \\ &= 4 - \frac{8}{3} \\ &= \frac{4}{3} \text{ sq. units} \end{aligned}$$

10. For X-axis, $y = 0$

$$\therefore 1 - x - 6x^2 = 0$$


$$\Rightarrow (2x + 1)(3x - 1) = 0$$

$$\Rightarrow x = \frac{-1}{2} \text{ or } x = \frac{1}{3}$$

$$\begin{aligned} \therefore \text{Required area} &= \int_{-1/2}^{1/3} (1 - x - 6x^2) dx \\ &= \left[x - \frac{x^2}{2} - 2x^3 \right]_{-1/2}^{1/3} \\ &= \left[\frac{1}{3} - \frac{1}{18} - \frac{2}{27} \right] - \left[\frac{-1}{2} - \frac{1}{8} + \frac{1}{4} \right] \\ &= \frac{11}{54} + \frac{3}{8} \\ &= \frac{125}{216} \text{ sq. units.} \end{aligned}$$



11.



$$\begin{aligned} \text{Area of the required region} &= 2 \int_0^{\sqrt{3}} (-x^2 + 3) dx \\ &= 2 \left[-\frac{x^3}{3} + 3x \right]_0^{\sqrt{3}} \\ &= 4\sqrt{3} \text{ sq. Units} \end{aligned}$$

12. In both cases area will be same.

$$\therefore A : B = 1 : 1$$

13. According to the given condition,

$$\begin{aligned} \int_1^b f(x) dx &= \sqrt{b^2 + 1} - \sqrt{2} \\ &= \sqrt{b^2 + 1} - \sqrt{1 + 1} = [\sqrt{x^2 + 1}]_1^b \end{aligned}$$

$$\therefore f(x) = \frac{d}{dx} (\sqrt{x^2 + 1}) = \frac{2x}{2\sqrt{x^2 + 1}} = \frac{x}{\sqrt{x^2 + 1}}$$


14. According to the given condition,

$$\int_{\frac{\pi}{4}}^{\beta} f(x) dx = \beta \sin \beta + \frac{\pi}{4} \cos \beta + \sqrt{2} \beta$$

Differentiating w.r.t. β , we get

$$f(\beta) = \sin \beta + \beta \cos \beta - \frac{\pi}{4} \sin \beta + \sqrt{2}$$

$$\begin{aligned} \therefore f\left(\frac{\pi}{2}\right) &= \sin \frac{\pi}{2} + \frac{\pi}{2} \cos \frac{\pi}{2} - \frac{\pi}{4} \sin \frac{\pi}{2} + \sqrt{2} \\ &= 1 - \frac{\pi}{4} + \sqrt{2} \end{aligned}$$



$$\begin{aligned} &= \left[x - \frac{8}{x} \right]_2^a \\ &= a - \frac{8}{a} + 2 \end{aligned}$$

$$A_2 = \int_a^4 \left(1 + \frac{8}{x}\right) dx = \left[x - \frac{8}{x} \right]_a^4 = 2 - a + \frac{8}{a}$$

According to the given condition,

$$\begin{aligned} a - \frac{8}{a} + 2 &= 2 - a + \frac{8}{a} \\ \Rightarrow 2a &= \frac{16}{a} \Rightarrow a^2 = 8 \\ \Rightarrow a &= 2\sqrt{2} \quad \dots[\because a > 0] \end{aligned}$$

16. The given curve passes through (1, 2).

$$\therefore 2 = a + b \quad \dots(i)$$

According to the given condition,

$$\begin{aligned} \int_0^4 (a\sqrt{x} + b) dx &= 8 \\ \Rightarrow \frac{2a}{3} [x^{3/2}]_0^4 + \frac{b}{2} [x^2]_0^4 &= 8 \Rightarrow \frac{2a}{3} \cdot 8 + 8b = 8 \\ \Rightarrow 2a + 3b &= 3 \quad \dots(ii) \end{aligned}$$

From (i) and (ii), we get

$$a = 3, b = -1$$

17. $R_1 = \int_{-1}^2 f(x) dx$

$$\begin{aligned} &= \int_{-1}^2 (1-x) f(1-x) dx \\ &\dots \left[\because \int_a^b f(x) dx = \int_a^b f(a+b-x) dx \right] \\ &= \int_{-1}^2 (1-x) f(x) dx \\ &\dots[\because f(x) = f(1-x) \text{ (given)}] \end{aligned}$$


$$\therefore R_1 = \int_{-1}^2 f(x) dx - R_1 \Rightarrow 2R_1 = \int_{-1}^2 f(x) dx$$

According to the given condition,

$$R_2 = \int_{-1}^2 f(x) dx$$

$$\therefore R_2 = 2R_1$$


18.



$$\begin{aligned} \text{Required area} &= 2 \int_0^{\pi/2} \cos x dx = 2 [\sin x]_0^{\pi/2} \\ &= 2 \text{ sq. units} \end{aligned}$$




19.



$$\begin{aligned} \text{Required area} &= \int_{\frac{\pi}{2}}^{\pi} \sin^2 d = \int_{\frac{\pi}{2}}^{\pi} \frac{1 - \cos 2}{2} d \\ &= \left[\frac{1}{2}d - \frac{\sin 2}{4} \right]_{\frac{\pi}{2}}^{\pi} \\ &= \frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4} \text{ sq. units} \end{aligned}$$

20.




Area of the region AOB

$$= 2 \int_0^a x^2 dx = 2 \int_0^a \sqrt{x} dx = 2 \left[\frac{3}{2} x^{3/2} \right]_0^a = \frac{4}{3} a^3$$

$$\begin{aligned} \text{Now, area of } \triangle AOB &= \frac{1}{2} \times AB \times OC \\ &= \frac{1}{2} \times 2a \times a^2 = a^3 \end{aligned}$$

$$\therefore \frac{\text{Area of } \triangle AOB}{\text{Area of the region AOB}} = \frac{a^3}{\frac{4}{3}a^3} = \frac{3}{4}$$

21.



Since the curve is symmetrical about X-axis.


$$\therefore \text{Required area } A = 2 \int_0^a \sqrt{a-x} dx$$

$$\text{Put } x = a \sin^2 \theta$$

$$\Rightarrow dx = 2a \sin \theta \cos \theta d\theta$$


$$\begin{aligned} \therefore A &= 2 \int_0^{\pi/2} a \sqrt{\frac{a \cos^2 \theta}{a \sin^2 \theta}} \cdot 2a \sin \theta \cos \theta d\theta \\ &= 4a^2 \int_0^{\pi/2} \frac{\cos \theta}{\sin \theta} \sin \theta \cos \theta d\theta \\ &= 4a^2 \int_0^{\pi/2} \cos^2 \theta d\theta = 4a^2 \cdot \frac{1}{2} \cdot \frac{\pi}{2} = \pi a^2 \end{aligned}$$

22.



$$\begin{aligned} \text{Required area} &= \int_1^4 x^2 dx \\ &= \int_1^4 2\sqrt{x} dx \\ &= 2 \left(\frac{2}{3} \right) \left[x^{3/2} \right]_1^4 \\ &= \frac{28}{3} \text{ sq. units.} \end{aligned}$$


23.



$$\begin{aligned} \text{Required area} &= \int_{-2}^{-1} (x + 2) dx + \int_{-1}^0 x^2 dx \\ &= \left[\frac{x^2}{2} + 2x \right]_{-2}^{-1} + \left[\frac{x^3}{3} \right]_{-1}^0 \\ &= \left[\left(\frac{1}{2} - 2 \right) + 2(-1 + 2) \right] + \left(0 + \frac{1}{3} \right) \\ &= \frac{5}{6} \end{aligned}$$




24.



$$\begin{aligned} \text{Required area} &= \int_0^1 \frac{1}{e} dx + \int_1^e \frac{1}{x} dx \\ &= \left[\frac{x}{e} \right]_0^1 + \left[\log x \right]_1^e \\ &= \frac{1}{e} + 1 \\ &= \frac{3}{2} \text{ sq. units} \end{aligned}$$

25.




Since the curve is symmetrical about X-axis and Y-axis,

Area of region between the two latus-rectum = 4 (Area of the shaded region)


$$\begin{aligned} &= 4 \int_0^{ae} dx \\ &= 4 \int_0^{ae} \frac{b}{a} \sqrt{a^2 - x^2} dx \\ &= \frac{4b}{a} \left[\frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \left(\frac{x}{a} \right) \right]_0^{ae} \\ &= \frac{4b}{a} \left[\frac{ae}{2} \sqrt{a^2(1-e^2)} + \frac{a^2}{2} \sin^{-1} e \right] \\ &= \frac{4b}{a} \left[\frac{abe}{2} + \frac{a^2}{2} \sin^{-1} e \right] \dots \left[\because b = a\sqrt{1-e^2} \right] \\ &= 2b (be + a \sin^{-1} e) \end{aligned}$$

26.



$$\begin{aligned} \text{Required area} &= \int_0^{\pi/4} (\cos x - \sin x) dx \\ &= [\sin x + \cos x]_0^{\pi/4} \\ &= \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} - 1 = \frac{2}{\sqrt{2}} - 1 = \sqrt{2} - 1 \end{aligned}$$

27. The two curves intersect at (0, 0) and (4a, 4a).




$$\begin{aligned} \text{Required area} &= \int_0^{4a} 2x dx - \int_0^{4a} x^2 dx \\ &= \frac{32}{3} a^2 - \frac{16}{3} a^2 = \frac{16}{3} a^2 \text{ sq. units} \end{aligned}$$

28. The two parabolas intersect at (0, 0) and (1, 1).

$$\therefore \text{required area} = \int_0^1 (\sqrt{x} - x^2) dx$$

$$= \left[\frac{2}{3} x^{3/2} - \frac{x^3}{3} \right]_0^1 = \frac{2}{3} - \frac{1}{3} = \frac{1}{3}$$

29.



Area bounded by the parabolas


$$= 2 \int_0^1 (1 - 3x^2 + 2x^2) dx$$



$$= 2 \int_0^1 (1 - x^2) dx = 2 \left[x - \frac{x^3}{3} \right]_0^1$$

$$= 2 \left(1 - \frac{1}{3} \right) = \frac{4}{3} \text{ sq. units}$$

30.



Area bounded by $y = x^2$ and line $y = 16$ is

$$2 \int_0^4 (x^2 - 16) dx$$

$$= 2 \left[\frac{x^3}{3} - 16x \right]_0^4 = \frac{-256}{3}$$

But area cannot be negative

∴ Required area = $\frac{256}{3}$ sq. units

31.

The points of intersection of $y^2 = 4a$ and $y = 2a$ are given by


$$(2a)^2 = 4a$$

$$\Rightarrow 4a(a - 1) = 0$$

$$\Rightarrow a = 0 \text{ or } a = \frac{1}{a}$$

When $a = 0$, $y = 0$ and when $a = \frac{1}{a}$, $y = 2$

∴ the points of intersection are $(0, 0)$ and $(\frac{1}{a}, 2)$.



$$\text{Required area} = \int_0^{1/a} [\sqrt{4a} - 2a] dx$$


$$= \frac{1}{3a} \text{ sq. units}$$

32. The area of the region bounded by the parabola $y^2 = 4ax$ and the line $y = m$ is $\frac{8a^2}{3m^3}$ sq. units.

Here, $a = \frac{1}{2}$ and $m = 1$

∴ Required area = $\frac{8(\frac{1}{2})^2}{3(1)^3} = \frac{2}{3}$ sq. units.

33.



The points of intersection of $y = x^2 - 2$ and $y = x$ are $(-1, -1)$ and $(2, 2)$

∴ Required area = $\int_{-1}^2 (x^2 - 2 - x) dx$

$$= \left[\frac{x^3}{3} - 2x - \frac{x^2}{2} \right]_{-1}^2$$


$$= \left[\frac{8}{3} - 4 - \frac{4}{2} \right] - \left[-\frac{1}{3} + 2 - \frac{1}{2} \right]$$

$$= \frac{-9}{2}$$

But area cannot be negative.

∴ Required area = $\frac{9}{2}$ sq. Units

34.




The two curves intersect at $(0, 0)$ and $(\frac{1}{a}, \frac{1}{a})$.



$$\begin{aligned} \text{Required area} &= \int_0^{\frac{1}{a}} \sqrt{\frac{1}{a}} d - \int_0^{\frac{1}{a}} a^{-2} d \\ \Rightarrow 1 &= \frac{2}{3\sqrt{a}} \left[\frac{3}{2} \right]_0^{\frac{1}{a}} - \frac{a}{3} \left[\frac{1}{a} \right]_0^{\frac{1}{a}} \\ &= \frac{2}{3\sqrt{a}} \left(\frac{1}{a} \right)^{\frac{3}{2}} - \frac{a}{3} \left(\frac{1}{a} \right)^3 = \frac{2}{3a^2} - \frac{1}{3a^2} \\ \Rightarrow \frac{1}{3a^2} &= 1 \Rightarrow a = \frac{1}{\sqrt{3}} \end{aligned}$$

35.




The given curves intersect at $O(0, 0)$ and $P(12, 12)$.

$$\begin{aligned} A_1 &= \int_0^{12} \sqrt{12} d - \int_0^{12} \frac{y^2}{12} d \\ &= \sqrt{12} \left[\frac{2}{3} \right]_0^{12} - \left[\frac{y^3}{36} \right]_0^{12} = \frac{45}{4} \end{aligned}$$

$$\begin{aligned} A_2 &= \int_3^{12} \sqrt{12} d - \int_3^{12} \frac{y^2}{12} d \\ &= \sqrt{12} \left[\frac{2}{3} \right]_3^{12} - \left[\frac{y^3}{36} \right]_3^{12} = \frac{147}{4} \end{aligned}$$

$$\therefore A_1 : A_2 = 45 : 147 = 15 : 49$$

36.



The two parabolas $y^2 = 4x$ and $x^2 = 4y$ intersect at $O(0, 0)$ and $P(4, 4)$.

$$\therefore S_3 = \int_0^4 \frac{y^2}{4} d - \frac{1}{4} \left[\frac{y^3}{3} \right]_0^4 = \frac{16}{3}$$

$$\text{and } S_2 + S_3 = \int_0^4 \sqrt{4x} d = 2 \int_0^4 \sqrt{x} d = \frac{4}{3} \left[\frac{3}{2} x^{3/2} \right]_0^4$$

$$\therefore S_2 + S_3 = \frac{32}{3} \quad \dots(i)$$

$$\therefore S_2 = \frac{32}{3} - \frac{16}{3} = \frac{16}{3}$$

Since, $S_1 + S_2 + S_3 = \text{Area of square OAPB} = 4 \times 4$


$$\therefore S_1 + S_2 + S_3 = 16$$

$$\Rightarrow S_1 + \frac{32}{3} = 16 \quad \dots[\text{From (i)}]$$

$$\Rightarrow S_1 = \frac{16}{3}$$


$$\therefore S_1 : S_2 : S_3 = 1 : 1 : 1$$

37.



$$\begin{aligned} \text{Required area} &= \int_0^9 \sqrt{x} d - \int_0^9 \left(\frac{-3}{2} \right) d \\ &= \left[\frac{2}{3} x^{3/2} \right]_0^9 - \frac{1}{2} \left[\frac{x^2}{2} - 3x \right]_0^9 \\ &= \frac{2}{3} (27 - 0) - \frac{1}{2} (36 - 18) \\ &= 9 \text{ sq. units} \end{aligned}$$

38.



$$\text{Here, } R_1 = \int_0^b (1-x)^2 d \quad \text{and } R_2 = \int_b^1 (1-x)^2 d$$



$$\therefore R_1 = \left[\frac{(-1)^3}{3} \right]_0^b \text{ and } R_2 = \left[\frac{(-1)^3}{3} \right]_b^1$$

$$\Rightarrow R_1 = \frac{(b-1)^3}{3} + \frac{1}{3} \text{ and } R_2 = -\frac{(b-1)^3}{3}$$

Since, $R_1 - R_2 = \frac{1}{4}$


$$\therefore \frac{(b-1)^3}{3} + \frac{1}{3} + \frac{(b-1)^3}{3} = \frac{1}{4}$$

$$\Rightarrow \frac{2}{3} (b-1)^3 = -\frac{1}{12} \Rightarrow (b-1)^3 = -\frac{1}{8}$$

$$\Rightarrow b-1 = -\frac{1}{2} \Rightarrow b = \frac{1}{2}$$

39. $\frac{d}{d} = 3^2 \Rightarrow \left(\frac{d}{d} \right)_{(1,1)} = 3$


\therefore equation of the tangent at (1, 1) is $-1 = 3(-1) \Rightarrow = 3 - 2$



Required area = $\int_0^1 3x^2 dx - \int_{\frac{2}{3}}^1 (3x - 2) dx$

$$= \left[\frac{3x^3}{3} \right]_0^1 - \left[\frac{3x^2}{2} - 2x \right]_{\frac{2}{3}}^1 = \frac{1}{2}$$

40.




Required area = $2 \int_0^1 (x - x^2) dx$

$$= 2 \left[\frac{x^2}{2} - \frac{x^3}{3} \right]_0^1$$

$$= 2 \left(\frac{1}{2} - \frac{1}{3} \right) = \frac{1}{3} \text{ sq. units}$$

41.




Required area = $2 \int_0^1 (-x + 2 - x) dx$

$$= 2 \int_0^1 (-2x + 2) dx$$

$$= 4 \left[-\frac{x^2}{2} + x \right]_0^1 = 2 \text{ sq. units}$$

42.




$3 - |x| = |x - 1| \Rightarrow |x| + |x - 1| = 3 \Rightarrow x = -1, 2$

Required area = $\int_{-1}^2 (3 - |x| - |x - 1|) dx$

$$= \left[3x - \frac{|x|^2}{2} - \frac{(|x - 1|)^2}{2} \right]_{-1}^2$$

$$= \left(6 - 2 - \frac{1}{2} \right) - \left(-3 + \frac{1}{2} + 2 \right) = 4 \text{ sq. units}$$

43.



The given curves intersect at (-1, 1) and (1, 1).
 \therefore Required area




$$= 2 \left[\int_0^1 \frac{2}{(1+x^2)} dx - \int_0^1 x^2 dx \right]$$

$$= 2 \left\{ 2 \left[\tan^{-1} x \right]_0^1 - \left[\frac{x^3}{3} \right]_0^1 \right\}$$

$$= 2 \left[2 \left(\frac{\pi}{4} - 0 \right) - \left(\frac{1}{3} - 0 \right) \right] = \pi - \frac{2}{3}$$

44.



The given curves intersect at $(-2a, a)$ and $(2a, a)$.

$$\therefore \text{Required area} = 2 \left[\int_0^{2a} \frac{8a^3}{x^2 + 4a^2} dx - \int_0^{2a} \frac{x^2}{4a} dx \right]$$

$$= 2 \left\{ \frac{8a^3}{2a} \left[\tan^{-1} \left(\frac{x}{2a} \right) \right]_0^{2a} - \frac{1}{4a} \left[\frac{x^3}{3} \right]_0^{2a} \right\}$$

$$= 2 \left[4a^2 (\tan^{-1} 1 - 0) - \frac{1}{4a} \left(\frac{8a^3}{3} - 0 \right) \right]$$

$$= 2 \left(4a^2 \cdot \frac{\pi}{4} - \frac{2a^2}{3} \right) = a^2 \left(2\pi - \frac{4}{3} \right)$$


45. The points of intersection of $x^2 + y^2 = 1$ and $x + y = 1$ are given by

$$x^2 + (1-x)^2 = 1$$

$$\Rightarrow 2x(x-1) = 0 \Rightarrow x = 0 \text{ or } x = 1$$

When $x = 0$, $y = 1$ and when $x = 1$, $y = 0$

\therefore the points of intersection are $(0, 1)$ and $(1, 0)$.




Required area

$$= \int_0^1 \left[\sqrt{1-x^2} - (1-x) \right] dx$$

$$= \left[\frac{\sqrt{1-x^2}}{2} + \frac{1}{2} \sin^{-1} x - x + \frac{x^2}{2} \right]_0^1$$

$$= \frac{1}{2} \cdot \frac{\pi}{2} - 1 + \frac{1}{2} = \frac{\pi}{4} - \frac{1}{2}$$

46.




Required area

$$= \frac{\pi(2)^2}{4} - \int_0^2 \sqrt{2x} dx$$

$$= \pi - \frac{2\sqrt{2}}{3} \left[\frac{3}{2} x^{3/2} \right]_0^2 = \pi - \frac{8}{3}$$

47.




Required area

$$= \frac{\pi a^2}{4} - \int_0^a \sqrt{ax} dx$$

$$= \frac{\pi a^2}{4} - \frac{2\sqrt{a}}{3} \left[\frac{3}{2} x^{3/2} \right]_0^a$$

$$= \frac{\pi a^2}{4} - \frac{2a^2}{3} = a^2 \left(\frac{\pi}{4} - \frac{2}{3} \right) \text{ sq. units.}$$

48.




Required area

$$= \frac{1}{2} \times \pi + 2 \int_0^1 \sqrt{1-x} dx$$

$$= \frac{\pi}{2} + 2 \left[\frac{(1-x)^{3/2}}{-3/2} \right]_0^1 = \frac{\pi}{2} - \frac{4}{3} (0-1) = \frac{\pi}{2} + \frac{4}{3}$$



49.



Putting $x = \frac{t}{2}$ in $y = 4 - x^2$, we get

$$y = 4 - \left(\frac{t}{2}\right)^2 \Rightarrow 2t^2 - t - 1 = 0$$

$$\Rightarrow (2t - 1)(t + 1) = 0 \Rightarrow t = 1, -\frac{1}{2}$$

$$\therefore \text{required area} = \int_{-1/2}^1 \left(\frac{4-t^2}{4}\right) dt - \int_{-1/2}^1 \frac{t^2}{2} dt$$

$$= \frac{1}{4} \left[\frac{t^2}{2} + \frac{t^3}{3} \right]_{-1/2}^1 - \frac{1}{2} \left[\frac{t^3}{3} \right]_{-1/2}^1$$

$$= \frac{1}{4} \left[\left(\frac{1}{2} + \frac{1}{8}\right) + \left(1 + \frac{1}{2}\right) \right] - \frac{1}{2} \left(\frac{1}{3} + \frac{1}{24}\right)$$


$$= \frac{1}{4} \left(\frac{15}{8}\right) - \frac{1}{2} \left(\frac{9}{24}\right) = \frac{15}{32} - \frac{3}{16} = \frac{9}{32}$$

50.

$$y = 4 - x^2$$

$$\therefore \frac{dy}{dx} = 4 - 2x$$

for minimum, $\frac{dy}{dx} = 0$



$$\begin{aligned} \text{Required area} &= 2 \int_0^{1/\sqrt{2}} (4 - x^2) dx \\ &= 2 \left[\frac{4x}{1} - \frac{x^3}{3} \right]_0^{1/\sqrt{2}} \\ &= 2 \left[\frac{4}{5(\sqrt{2})^5} - \frac{1}{3(\sqrt{2})^3} \right] \\ &= \frac{2}{(\sqrt{2})^3} \left[\frac{1}{5(\sqrt{2})^2} - \frac{1}{3} \right] \\ &= \frac{1}{\sqrt{2}} \left[\frac{1}{10} - \frac{1}{3} \right] = \frac{-7}{30\sqrt{2}} \end{aligned}$$


But, area cannot be negative.

$$\therefore \text{Required area} = \frac{7}{30\sqrt{2}}$$



Evaluation Test

1.




$$\text{Required area} = \int_{-\pi}^{\pi} |\sin x| dx$$

$$\begin{aligned} &= \int_{-\pi}^0 |\sin x| dx + \int_0^{\pi} |\sin x| dx \\ &= \int_{-\pi}^0 (-\sin x) dx + \int_0^{\pi} \sin x dx \\ &= [\cos x]_{-\pi}^0 - [\cos x]_0^{\pi} \\ &= \cos 0 - \cos(-\pi) - (\cos \pi - \cos 0) \\ &= 1 - (-1) - (-1 - 1) \\ &= 1 + 1 - (-2) \\ &= 4 \text{ sq. units} \end{aligned}$$



2.




$$= \sin^{-1}$$

$$\therefore = \sin$$

$$\begin{aligned} \text{Required area} &= 2 \int_0^{\frac{\pi}{2}} d = 2 \int_0^{\frac{\pi}{2}} \sin d \\ &= -2 [\cos]_0^{\frac{\pi}{2}} \\ &= -2(0 - 1) = 2 \text{ sq. units} \end{aligned}$$


3.



$$\begin{aligned} \text{Required area} &= \int_{-1}^2 \sqrt{5-x^2} dx - \int_{-1}^2 (x-1) dx \\ &= \left[\frac{x}{2} \sqrt{5-x^2} + \frac{5}{2} \sin^{-1} \frac{x}{\sqrt{5}} \right]_{-1}^2 - \left[\frac{x^2}{2} - x \right]_{-1}^2 \\ &= 1 + \frac{5}{2} \sin^{-1} \left(\frac{2}{\sqrt{5}} \right) - \left(-\frac{1}{2} \times 2 + \frac{5}{2} \sin^{-1} \left(-\frac{1}{\sqrt{5}} \right) \right) \\ &\quad - \left[1 - \frac{1}{2} - \left(-1 - \frac{1}{2} \right) \right] - \left(2 - 2 - \frac{1}{2} + 1 \right) \\ &= 1 + \frac{5}{2} \sin^{-1} \left(\frac{2}{\sqrt{5}} \right) + 1 + \frac{5}{2} \sin^{-1} \left(\frac{1}{\sqrt{5}} \right) - 2 - \frac{1}{2} \\ &= \frac{5}{2} \left[\sin^{-1} \left(\frac{2}{\sqrt{5}} \right) + \sin^{-1} \left(\frac{1}{\sqrt{5}} \right) \right] - \frac{1}{2} \end{aligned}$$

$$\begin{aligned} &= \frac{5}{2} \left[\sin^{-1} \left(\frac{2}{\sqrt{5}} \right) + \cos^{-1} \left(\frac{2}{\sqrt{5}} \right) \right] - \frac{1}{2} \\ &\quad \dots \left[\because \sin^{-1} = \cos^{-1} \sqrt{1-x^2} \right] \\ &= \frac{5}{2} \cdot \frac{\pi}{2} - \frac{1}{2} = \left(\frac{5\pi}{4} - \frac{1}{2} \right) \text{ sq. unit} \end{aligned}$$


4.



$$\begin{aligned} \text{Required area} &= 4 (\text{area of } \triangle OAB) \\ &= 4 \int_0^1 dx \\ &= 4 \int_0^1 (-x+1) dx \\ &= 4 \left[-\frac{x^2}{2} + x \right]_0^1 = 2 \text{ sq. units} \end{aligned}$$

5.

The given equation can be written as $x^2 - 6x + 9 + y^2 - 4y + 4 = 9 + 4 - 12$
 $\therefore (x-3)^2 + (y-2)^2 = 1$
 This is a circle with centre at (3, 2) and radius 1.




\therefore

$$\begin{aligned} \text{Required area} &= \int_2^{\frac{5}{2}} dx - \int_2^{\frac{5}{2}} \left[2 - \sqrt{1 - (x-3)^2} \right] dx \end{aligned}$$



$$\begin{aligned}
 &= \left[\frac{x^2}{2} \right]_2^{5/2} - \left[2x \right]_2^{5/2} \\
 &\quad + \left[\left(\frac{-3}{2} \right) \sqrt{1 - (-3)^2} + \frac{1}{2} \sin^{-1} \left(\frac{-3}{1} \right) \right]_2^{5/2} \\
 &= \frac{1}{2} \left(\frac{25}{4} - 4 \right) - 2 \left(\frac{5}{2} - 2 \right) + \left(\frac{-1}{2} \right) \sqrt{1 - \frac{1}{4}} \\
 &\quad + \frac{1}{2} \sin^{-1} \left(-\frac{1}{2} \right) - \left[0 + \frac{1}{2} \sin^{-1}(-1) \right] \\
 &= \frac{9}{8} - 1 - \frac{1}{4} \cdot \frac{\sqrt{3}}{2} + \frac{1}{2} \left(-\frac{\pi}{6} \right) - \frac{1}{2} \left(-\frac{\pi}{2} \right) \\
 &= \frac{1}{8} - \frac{\sqrt{3}}{8} + \frac{\pi}{4} - \frac{\pi}{12} = \left(\frac{\pi}{6} - \frac{\sqrt{3}-1}{8} \right) \text{sq. unit}
 \end{aligned}$$


6. The point of intersection of the curve $y = 2 - x^2$ and the line $y = -x$ are $(0, 0)$ and $(3, -3)$.



∴ Required area

$$\begin{aligned}
 &= \int_0^3 [(2 - x^2) - (-x)] dx \\
 &= \int_0^3 (3 - x^2) dx = \left[\frac{3x}{1} - \frac{x^3}{3} \right]_0^3 = \frac{9}{2} \text{ sq. unit}
 \end{aligned}$$


- 7.



$$|\cos x| = 1 \Rightarrow \cos x = 1 \text{ or } \cos x = -1$$

$$\begin{aligned}
 \text{Required Area} &= \int_{-1}^1 \cos x \, dx \\
 &= 2 \int_0^1 \cos x \, dx \\
 &\quad \dots [\because \cos x \text{ is an even function}] \\
 &= 2 [\sin x]_0^1 \\
 &= 2(\sin 1 - 0) \\
 &= 2 \sin 1
 \end{aligned}$$

- 8.




Required area

$$\begin{aligned}
 &= \int_0^{\frac{3\pi}{2}} |\cos x - \sin x| dx \\
 &= \int_0^{\frac{\pi}{4}} (\cos x - \sin x) dx + \int_{\frac{\pi}{4}}^{\frac{5\pi}{4}} (\sin x - \cos x) dx \\
 &\quad + \int_{\frac{5\pi}{4}}^{\frac{3\pi}{2}} (\cos x - \sin x) dx \\
 &= [\sin x + \cos x]_0^{\pi/4} - [\cos x + \sin x]_{\pi/4}^{5\pi/4} \\
 &\quad + [\sin x + \cos x]_{5\pi/4}^{3\pi/2} \\
 &= \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} - (0 + 1) - \left[-\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} - \left(\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} \right) \right] \\
 &\quad + (-1) + 0 - \left[-\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} \right] \\
 &= \sqrt{2} - 1 + \sqrt{2} + \sqrt{2} - 1 + \sqrt{2} \\
 &= (4\sqrt{2} - 2) \text{ sq. unit}
 \end{aligned}$$




9.



$$\begin{aligned} \text{Required area} &= \int_0^3 (x^2 + 2 - 3) dx \\ &= \left[\frac{x^3}{3} + 2x - 3x \right]_0^3 \\ &= 9 + 6 - \frac{9}{2} - 0 = \frac{21}{2} \text{ sq. unit} \end{aligned}$$


10.



Putting $x = 0$ in the given equation, we get $y = 1$ or $y = -2$


$$\begin{aligned} \therefore \text{Required Area} &= \int_{-2}^1 x dy = \int_{-2}^1 (2 - y^2) dy \\ &= \left[2y - \frac{y^3}{3} \right]_{-2}^1 \\ &= 2 - \frac{1}{3} - \left[-4 - \frac{8}{3} \right] \\ &= 2 - \frac{1}{3} + 4 + \frac{8}{3} \\ &= 8 - 3 - \frac{1}{3} = \frac{9}{2} \text{ sq. units} \end{aligned}$$

11.



$$\begin{aligned} \text{Required area} &= \int_0^{4a} \left(\frac{x}{4a^2} + 1 - 1 \right) dy \\ &= \int_0^{4a} \frac{x}{4a^2} dy = \frac{1}{4a^2} \left[\frac{y^3}{3} \right]_0^{4a} \\ &= \frac{1}{4a^2} \times \frac{1}{3} \times (64a^3 - 0) \\ &= \frac{16a}{3} \text{ sq. units} \end{aligned}$$


12.



$$\begin{aligned} \text{Required area} &= 2 \int_{-1}^0 \sqrt{y^2 + 1} dy + 2 \int_0^1 \sqrt{-y^2 + 1} dy \\ &= 2 \cdot \frac{2}{3} \left[(y^2 + 1)^{3/2} \right]_{-1}^0 + 2 \cdot \frac{2}{3} \left[\frac{(-y^2 + 1)^{3/2}}{-1} \right]_0^1 \\ &= \frac{4}{3} + \frac{4}{3} \\ &= \frac{8}{3} \text{ sq. units} \end{aligned}$$



13.



$$x^2 + 1 = -3x \Rightarrow \frac{d}{dx} = 2x + 1$$

$$\therefore \left(\frac{d}{dx}\right)_{(1,3)} = 2(1) + 1 = 2 + 1 = 3$$

\therefore The equation of the tangent at the point (1, 3) is $y - 3 = 3(x - 1)$ i.e., $y = 3x$.

\therefore It passes through origin.

\therefore Required area

= area of the region OABCO + area of the region OCDO

$$= \int_{-1}^0 x^2 + 1 dx + \int_0^1 (-3x - x^2) dx$$

$$= \int_{-1}^0 (x^2 + 1) dx + \int_0^1 (-x^2 - 3x) dx$$


$$= \int_{-1}^0 (x^2 + 1) dx + \int_0^1 (-x^2 - 2x + 1) dx$$

$$= \left[\frac{x^3}{3} + \frac{2x}{2} + \dots \right]_{-1}^0 + \left[-\frac{x^3}{3} - x^2 + \dots \right]_0^1$$

$$= 0 - \left(-\frac{1}{3} + \frac{1}{2} - 1 \right) + \frac{1}{3} - 1 + 1 - 0$$

$$= \frac{4}{3} - \frac{1}{2} + \frac{1}{3} = \frac{8-3+2}{6} = \frac{7}{6} \text{ sq. unit}$$

14.



The equation $x^2 + 4y^2 = 4$ is of ellipse with centre at origin and the equation $y^2 = 3x$ is of a parabola with vertex at origin.

Solving the equations, we get $x^2 + 3y^2 - 4 = 0$

$$\therefore (x+4)(x-1) = 0$$

But $x = -4$ is not possible, since both points of intersection lie on the right hand side of Y-axis.

$$\therefore x = 1 \text{ and } y = \pm \frac{\sqrt{3}}{2}$$

\therefore The points of intersection are $A\left(1, \frac{\sqrt{3}}{2}\right)$ and

$$B\left(1, -\frac{\sqrt{3}}{2}\right).$$

\therefore Required area

$$= \int_{-\frac{\sqrt{3}}{2}}^{\frac{\sqrt{3}}{2}} (x - y^2) dy$$

$$= \int_{-\frac{\sqrt{3}}{2}}^{\frac{\sqrt{3}}{2}} \left[\sqrt{4-4y^2} - \frac{4y^2}{3} \right] dy$$

$$= 2 \int_0^{\frac{\sqrt{3}}{2}} \left[\sqrt{4-4y^2} - \frac{4y^2}{3} \right] dy$$

....[\because the function is even]

$$= 4 \int_0^{\frac{\sqrt{3}}{2}} \sqrt{1-y^2} dy - \frac{8}{3} \int_0^{\frac{\sqrt{3}}{2}} y^2 dy$$

$$= 4 \left[\frac{y}{2} \sqrt{1-y^2} + \frac{1}{2} \sin^{-1}(y) \right]_0^{\frac{\sqrt{3}}{2}} - \frac{8}{3} \left[\frac{y^3}{3} \right]_0^{\frac{\sqrt{3}}{2}}$$

$$= 2 \left[\frac{\sqrt{3}}{2} \sqrt{\frac{1}{4}} + \sin^{-1}\left(\frac{\sqrt{3}}{2}\right) - 0 \right] - \frac{8}{3 \times 3} \times \left(\frac{3\sqrt{3}}{8} - 0 \right)$$

$$= 2 \left[\frac{\sqrt{3}}{4} + \frac{\pi}{3} \right] - \frac{8}{9} \times \frac{3\sqrt{3}}{8}$$


$$= \frac{2\pi}{3} + \frac{2\sqrt{3}}{4} - \frac{\sqrt{3}}{3}$$

$$= \frac{2\pi}{3} + \frac{2\sqrt{3}}{12}$$

$$= \left(\frac{2\pi}{3} + \frac{1}{2\sqrt{3}} \right) \text{ sq. unit}$$



15.



Slope of tangent = $\frac{dy}{dx} = 2x + 1$

$\therefore \int (2x + 1) dx = x^2 + x + c$

The curve passes through the point (1, 2).

$\therefore 2 = 1^2 + 1 + c \quad \therefore c = 0$

\therefore The equation of the curve is $y = x^2$, which is a parabola as shown in the figure.

\therefore Required area = $\int_0^1 (x^2) dx = \left[\frac{x^3}{3} + \frac{x^2}{2} \right]_0^1$
 $= \frac{1}{3} + \frac{1}{2} - 0 = \frac{5}{6}$ sq. unit

16. Draw $AP \perp$ to X-axis.


$A_1 = A(\Delta OAP) = \frac{1}{2} \times a \times a^2 = \frac{a^3}{2}$

$A_2 =$ Area bounded the curve OA and the lines OP and AP

$= \int_0^a x dx = \int_0^a x^2 dx = \left[\frac{x^3}{3} \right]_0^a = \frac{a^3}{3}$

\therefore Required area = $A_1 - A_2 = \frac{a^3}{2} - \frac{a^3}{3} = \frac{a^3}{6}$ sq. unit


17.



Required area = area of the rectangle OABC - area of the region OBCO

$= \frac{\pi}{4} \times \frac{1}{\sqrt{2}} - \int_0^{\frac{\pi}{4}} \sin x dx = \frac{\pi}{4\sqrt{2}} - [-\cos x]_0^{\pi/4}$
 $= \left[\frac{\pi}{4\sqrt{2}} + \left(\frac{1}{\sqrt{2}} - 1 \right) \right]$ sq. units

18.



The equation of the parabola is $(x - 2)^2 = -4(y - 3)$

$(x - 2)^2 = -4(y - 3)$

Diff. w.r.t. x , we get

$2(x - 2) \frac{dx}{dx} = -4$

$\therefore \frac{dx}{dx} = \frac{-4}{2(x - 2)}$

$\therefore \left(\frac{dx}{dx} \right)_{(2,3)} = \frac{1}{2(3-2)} = \frac{1}{2}$

\therefore Equation of tangent is $y - 3 = \frac{1}{2}(x - 2)$

$\therefore 2y - 6 = x - 2$

$\therefore x - 2 + 4 = 0$

It cuts the X-axis at the point Q(-4, 0) and the parabola cuts the X-axis at the point R(5, 0).

\therefore required area = $\int_0^3 ((x - 2)^2 - 4) dx$

$= \int_0^3 [(x - 2)^2 + 1 - (x - 4)] dx$

$= \int_0^3 (x^2 - 6x + 9) dx$

$= \left[\frac{x^3}{3} - 3x^2 + 9x \right]_0^3$

$= 9 - 27 + 27 - 0$

$= 9$ sq. units



$$\begin{aligned} 19. \quad \text{Required area} &= \int_0^{\frac{\pi}{4}} \left(\sqrt{\frac{1+\sin}{\cos}} - \sqrt{\frac{1-\sin}{\cos}} \right) d \\ &\quad \dots \left[\because \frac{1+\sin}{\cos} > \frac{1-\sin}{\cos} > 0 \right] \\ &= \int_0^{\frac{\pi}{4}} \left(\sqrt{\frac{\left(\cos \frac{\theta}{2} + \sin \frac{\theta}{2}\right)^2}{\cos^2 \frac{\theta}{2} - \sin^2 \frac{\theta}{2}}} - \sqrt{\frac{\left(\cos \frac{\theta}{2} - \sin \frac{\theta}{2}\right)^2}{\cos^2 \frac{\theta}{2} - \sin^2 \frac{\theta}{2}}} \right) d \\ &= \int_0^{\frac{\pi}{4}} \left(\sqrt{\frac{\cos \frac{\theta}{2} + \sin \frac{\theta}{2}}{\cos \frac{\theta}{2} - \sin \frac{\theta}{2}}} - \sqrt{\frac{\cos \frac{\theta}{2} - \sin \frac{\theta}{2}}{\cos \frac{\theta}{2} + \sin \frac{\theta}{2}}} \right) d \\ &= \int_0^{\frac{\pi}{4}} \left(\sqrt{\frac{1 + \tan \frac{\theta}{2}}{1 - \tan \frac{\theta}{2}}} - \sqrt{\frac{1 - \tan \frac{\theta}{2}}{1 + \tan \frac{\theta}{2}}} \right) d \\ &= \int_0^{\frac{\pi}{4}} \frac{1 + \tan \frac{\theta}{2} - 1 + \tan \frac{\theta}{2}}{\sqrt{1 - \tan^2 \frac{\theta}{2}}} d = \int_0^{\frac{\pi}{4}} \frac{2 \tan \frac{\theta}{2}}{\sqrt{1 - \tan^2 \frac{\theta}{2}}} d \\ \text{Put } \tan \frac{\theta}{2} = t &\Rightarrow \frac{1}{2} \sec^2 \frac{\theta}{2} d = dt \\ \therefore \text{required area} &= \int_0^{\tan \frac{\pi}{8}} \frac{4t}{(1+t^2)\sqrt{1-t^2}} dt \\ &= \int_0^{\sqrt{2}-1} \frac{4t}{(1+t^2)\sqrt{1-t^2}} dt \\ &\quad \dots \left[\because \tan \frac{\pi}{8} = \sqrt{2} - 1 \right] \end{aligned}$$

07 Differential Equations



Hints



Classical Thinking

- Here, the highest order derivative is $\frac{d^2s}{dt^2}$ with power 2.
 \therefore order = 2 and degree = 2
- Here, the highest order derivative is $\frac{d^2}{d^2}$ with power 3.
 \therefore order = 2 and degree = 3
- Here, the highest order derivative is $\frac{d^2}{d^2}$ with power 3.
 \therefore order = 2 and degree = 3
- Here, the highest order derivative is $\frac{d^3}{d^3}$ with power 1.
 \therefore order = 3 and degree = 1
- In option (B), " " is the highest order derivative, of order 2.
 \therefore option (B) is the correct answer.
- Here, the highest order derivative is $\frac{d^4}{d^4}$ with power 1.
 \therefore order = 4 and degree = 1
- $y = 4 \sin 3x$ (i)
 $\Rightarrow \frac{dy}{dx} = 12 \cos 3x$
 $\Rightarrow \frac{d^2y}{dx^2} = -36 \sin 3x = -9 \times 4 \sin 3x = -9y$ [From (i)]
 $\Rightarrow \frac{d^2y}{dx^2} + 9y = 0$
- $y = A \sin x + B \cos x$ (i)
 $\Rightarrow \frac{dy}{dx} = A \cos x - B \sin x$
 $\Rightarrow \frac{d^2y}{dx^2} = -A \sin x - B \cos x = -(A \sin x + B \cos x) = -y$ [From (i)]
 $\therefore \frac{d^2y}{dx^2} + y = 0$

- $y = a \cos(x + b)$ (i)
 $\Rightarrow \frac{dy}{dx} = -a \sin(x + b)$
 $\Rightarrow \frac{d^2y}{dx^2} = -a \cos(x + b) = -y$ [From (i)]
 $\Rightarrow \frac{d^2y}{dx^2} + y = 0$
- $y = ce^{\sin^{-1}x}$ (i)
 $\Rightarrow \frac{dy}{dx} = ce^{\sin^{-1}x} \cdot \frac{1}{\sqrt{1-x^2}}$
 $\Rightarrow \frac{dy}{dx} = \frac{y}{\sqrt{1-x^2}}$ [From (i)]
- $y = (x + k)e^{-x}$ (i)
 $\Rightarrow \frac{dy}{dx} = -(x + k)e^{-x} + e^{-x}$
 $\Rightarrow \frac{dy}{dx} = -y + e^{-x}$ [From (i)]
 $\Rightarrow \frac{dy}{dx} + y = e^{-x}$
- $y^2 = a$
Differentiating w.r.t x , we get
 $2y \frac{dy}{dx} + 2x = 0$
 $\Rightarrow \frac{dy}{dx} + \frac{2x}{y} = 0$
- $x^2 + y^2 = a^2$
Differentiating w.r.t x , we get
 $2x + 2y \frac{dy}{dx} = 0 \Rightarrow x + y \frac{dy}{dx} = 0$
- $\frac{d^2y}{dx^2} = 2$
Integrating on both sides, we get
 $\int d^2y = \int \frac{2}{2} dx + c$
 $\Rightarrow y = -\frac{2}{2}x + c$



$$15. \frac{d}{d} = 2 + \sin 3$$

Integrating on both sides, we get

$$\int d = \int (2 + \sin 3) d + c$$

$$\Rightarrow = \frac{2}{3} - \frac{\cos 3}{3} + c$$

$$16. \frac{d}{d} = (ae^b + c \cos m)$$

Integrating on both sides, we get

$$\int dy = \int (ae^{bx} + c \cos m) d + k$$

$$\Rightarrow = \frac{ae^b}{b} + \frac{c \sin(m)}{m} + k$$

$$17. \frac{d}{d} = \sec (\sec + \tan)$$

Integrating on both sides, we get

$$\int d = \int (\sec^2 + \sec \tan) d + c$$

$$\Rightarrow = \tan + \sec + c$$

$$18. \frac{d}{d} = e (\sin + \cos)$$

Integrating on both sides, we get

$$\int d = \int e (\sin + \cos) d + c$$

$$\Rightarrow = e \sin + c$$

$$19. \frac{d}{d} = e + \cos + \tan$$

Integrating on both sides, we get

$$\int d = \int (e + \cos + \tan) d + c$$

$$\Rightarrow = e + \sin + \frac{2}{2} + \log \sec + c$$

$$20. (1 + 2) \frac{d}{d} = 1$$

Integrating on both sides, we get

$$\int d = \int \frac{1}{1+2} d + c$$

$$\Rightarrow = \tan^{-1} + c$$

$$21. \frac{d}{d} + \frac{1}{\sqrt{1-2}} = 0$$

Integrating on both sides, we get

$$\int d + \int \frac{1}{\sqrt{1-2}} d = c$$

$$\Rightarrow + \sin^{-1} = c$$

$$22. \frac{d}{d} + \sin^2 = 0$$

$$\Rightarrow \frac{d}{d} = -\sin^2$$

$$\Rightarrow \frac{d}{d} = -\frac{1}{\operatorname{cosec}^2}$$

Integrating on both sides, we get

$$\int d = -\int \operatorname{cosec}^2 d + c$$

$$\Rightarrow = \cot + c$$

$$23. \frac{d}{d} + \frac{1+2}{d} = 0$$

Integrating on both sides, we get

$$\int d + \int \left(\frac{1}{d} + \frac{2}{d} \right) d = c$$

$$\Rightarrow + \log + \frac{2}{2} = c$$

$$24. (1 + 2) \frac{d}{d} = \Rightarrow d = \frac{d}{1+2}$$

Integrating on both sides, we get

$$\int d = \int \frac{1}{1+2} d + c$$

$$\Rightarrow = \frac{1}{2} \log_e(1 + 2) + c$$

$$25. \frac{d}{d} = \left(- \right)^{1/3}$$

$$\Rightarrow \frac{d}{1/3} = \frac{d}{1/3}$$

Integrating on both sides, we get

$$\int \frac{d}{1/3} - \int \frac{d}{1/3} = c_1$$

$$\Rightarrow \frac{3}{2} 2^{2/3} - \frac{3}{2} 2^{2/3} = c_1$$

$$\Rightarrow 2^{2/3} - 2^{2/3} = c, \text{ where } c = \frac{2c_1}{3}$$

$$26. \frac{d}{d} = (1 + 2)(1 + 2)$$

Integrating on both sides, we get

$$\int \frac{d}{1+2} = \int (1 + 2) d + c$$

$$\Rightarrow \tan^{-1} = \frac{2}{2} + c$$

$$\Rightarrow = \tan \left(\frac{2}{2} + c \right)$$



$$27. \frac{d}{d} = \log \Rightarrow d = \log d$$

Integrating on both sides, we get

$$\int d = \int \log d + c$$

$$\Rightarrow = \frac{2}{2} \log - \frac{2}{4} + c$$

$$28. \frac{d}{d} = (+)^2 \quad \dots(i)$$

$$\text{Put } + = v \quad \dots(ii)$$

$$\Rightarrow \frac{d}{d} = \frac{dv}{d} - 1 \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$\frac{dv}{d} - 1 = v^2$$

$$\Rightarrow \frac{dv}{d} = v^2 + 1$$

$$\Rightarrow \frac{dv}{v^2 + 1} = d$$

Integrating on both sides, we get

$$\tan^{-1} v = + c \Rightarrow v = \tan(+ c)$$

$$\Rightarrow + = \tan(+ c)$$

$$29. \frac{d}{d} + - = ^3 - 3$$

This is the linear differential equation of the form

$$\frac{d}{d} + P. = Q, \text{ where } P = \frac{1}{d}$$

$$\therefore \text{I.F.} = e^{\int P d} = e^{\int \frac{1}{d} d} = e^{\log} =$$

30. The given equation is of the form

$$\frac{d}{d} + P = Q.$$

$$\text{Here, } P = \frac{1}{3} \text{ and } Q = 1$$

$$\therefore \text{I.F.} = e^{\int \frac{1}{3} d} = e^{\frac{d}{3}}$$

\therefore solution of the given equation is

$$(\text{I.F.}) = \int Q(\text{I.F.})d + c$$

$$\Rightarrow . e^{\frac{d}{3}} = \int 1.e^{\frac{d}{3}} d + c$$

$$\Rightarrow . e^{\frac{d}{3}} = 3e^{\frac{d}{3}} + c$$

$$\Rightarrow = 3 + c.e^{-\frac{d}{3}}$$

$$31. \log\left(\frac{d}{d}\right) = +$$

$$\Rightarrow \frac{d}{d} = e^+$$

$$\Rightarrow \frac{d}{d} = e \cdot e$$

Integrating on both sides, we get

$$\int e d - \int e^{-} d = c$$

$$\Rightarrow e + e^{-} = c$$

$$32. \text{ Here, } P = \frac{1}{d} \text{ and } Q = ^2$$

$$\therefore \text{I.F.} = e^{\int \frac{1}{d} d} = e^{\log} =$$

\therefore solution of the given equation is

$$= \int ^2 \cdot d + c_1$$

$$\Rightarrow = \frac{4}{4} + c_1 \Rightarrow 4 = ^4 + c, \text{ where } c = 4 c_1$$

$$33. \frac{d}{d} + 3 = \Rightarrow \frac{d}{d} + \frac{3}{d} = 1$$

$$\therefore \text{I.F.} = e^{\int \frac{3}{d} d} = e^{3 \log} = ^3$$

\therefore solution of the given equation is

$$^3 = \int ^3 \cdot 1d + c \Rightarrow ^3 = \frac{4}{4} + c$$

$$34. \frac{d}{d} + - = \sin$$

$$\therefore \text{I.F.} = e^{\int - d} = e^{\log} =$$

\therefore solution of the given equation is

$$. = \int \sin d + c$$

$$\Rightarrow = - \cos + \sin + c$$

$$\Rightarrow (+ \cos) = \sin + c$$

$$35. \frac{d}{d} + = \cos$$

Here, $P = 1$ and $Q = \cos$

$$\therefore \text{I.F.} = e^{\int 1 d} = e$$

\therefore solution of the given equation is

$$.e = \int \cos .e d + c$$

$$\Rightarrow .e = \frac{e (\cos + \sin)}{2} + c$$

$$\Rightarrow = \frac{1}{2} (\cos + \sin) + c.e^{-}$$



$$36. \cos \frac{d}{d} + \sin = 1$$

$$\Rightarrow \frac{d}{d} + \tan = \sec$$

$$\therefore \text{I.F.} = e^{\int \tan d} = e^{\log \sec} = \sec$$

\therefore solution of the given equation is

$$\sec = \int \sec^2 + c = \tan + c$$

$$37. \text{I.F.} = e^{\int Pd} = e^{\int \cot d} = e^{\log \sin} = \sin$$

\therefore solution of the given equation is

$$\sin = \int 2 \cos \sin d + c_1$$

$$\Rightarrow \sin = \int \sin 2d + c_1$$

$$\Rightarrow \sin = -\frac{1}{2} \cos 2 + c_1$$

$$\Rightarrow 2 \sin + \cos 2 = c, \text{ where } c = 2c_1$$



Critical Thinking

$$1. \left(\frac{d^2}{d^2}\right)^3 = \left(1 + \frac{d}{d}\right)^{1/2} \Rightarrow \left(\frac{d^2}{d^2}\right)^6 = 1 + \frac{d}{d}$$

Here, the highest order derivative is $\frac{d^2}{d^2}$ with power 6.

\therefore degree = 6

$$2. \frac{d^2}{d^2} + \sqrt{1 + \left(\frac{d}{d}\right)^3} = 0$$

$$\Rightarrow \left(\frac{d^2}{d^2}\right)^2 = \left[-\sqrt{1 + \left(\frac{d}{d}\right)^3}\right]^2$$

$$\Rightarrow \left(\frac{d^2}{d^2}\right)^2 = 1 + \left(\frac{d}{d}\right)^3$$

Here, the highest order derivative is $\frac{d^2}{d^2}$ with power 2.

\therefore degree = 2

$$3. 3 \frac{d^2}{d^2} = \left\{1 + \left(\frac{d}{d}\right)^2\right\}^{\frac{3}{2}}$$

$$\Rightarrow 9 \left(\frac{d^2}{d^2}\right)^2 = \left\{1 + \left(\frac{d}{d}\right)^2\right\}^3$$

Here, the highest order derivative is $\frac{d^2}{d^2}$ with power 2.

\therefore degree = 2

$$4. \frac{d^2}{d^2} + \left(\frac{d}{d}\right)^{\frac{1}{3}} + \frac{1}{4} = 0$$

$$\Rightarrow \left(\frac{d^2}{d^2} + \frac{1}{4}\right)^3 = \left[-\left(\frac{d}{d}\right)^{\frac{1}{3}}\right]^3$$

$$\Rightarrow \left(\frac{d^2}{d^2} + \frac{1}{4}\right)^3 = -\frac{d}{d}$$

Here, the highest order derivative is $\frac{d^2}{d^2}$ with power 3.

\therefore order = 2 and degree = 3

$$5. \frac{d^2}{d^2} = \left\{ + \left(\frac{d}{d}\right)^2 \right\}^{1/4}$$

$$\Rightarrow \left(\frac{d^2}{d^2}\right)^4 = \left\{ \left[+ \left(\frac{d}{d}\right)^2 \right]^{1/4} \right\}^4$$

$$\Rightarrow \left(\frac{d^2}{d^2}\right)^4 = + \left(\frac{d}{d}\right)^2$$

Here, the highest order derivative is $\frac{d^2}{d^2}$ with power 4.

\therefore order = 2 and degree = 4

6. Since, the given differential equation cannot be expressed as a polynomial in differential coefficients, the degree is not defined.

$$7. \sqrt{\frac{d}{d}} - 4 \frac{d}{d} - 7 = 0$$

$$\Rightarrow \left(\sqrt{\frac{d}{d}}\right)^2 = \left(4 \frac{d}{d} + 7\right)^2$$

$$\Rightarrow \frac{d}{d} = 16 \left(\frac{d}{d}\right)^2 + 56 \frac{d}{d} + 49$$

This is a differential equation of order 1 and degree 2.

$$8. \frac{d^2}{d^2} - \sqrt{\frac{d}{d}} - 3 =$$

$$\Rightarrow \left(\frac{d^2}{d^2} - \right)^2 = \left(\sqrt{\frac{d}{d}} - 3\right)^2$$

$$\Rightarrow \left(\frac{d^2}{d^2}\right)^2 - 2 \cdot \frac{d^2}{d^2} + \frac{d}{d} - 3$$

Here, the highest order derivative is $\frac{d^2}{d^2}$ with power 2.

\therefore degree = 2



$$9. \left[1 + \left(\frac{d}{d} \right)^2 \right]^{3/4} = \left(\frac{d^2}{d^2} \right)^{1/3}$$

$$\Rightarrow \left\{ \left[1 + \left(\frac{d}{d} \right)^2 \right]^{3/4} \right\}^4 = \left(\frac{d^2}{d^2} \right)^{4/3}$$

$$\Rightarrow \left\{ \left[1 + \left(\frac{d}{d} \right)^2 \right]^3 \right\} = \left\{ \left[\frac{d^2}{d^2} \right]^{4/3} \right\}^3$$

$$\Rightarrow \left[1 + \left(\frac{d}{d} \right)^2 \right]^9 = \left(\frac{d^2}{d^2} \right)^4$$

Here, the highest order derivative is $\frac{d^2}{d^2}$ with power 4.

∴ degree = 4

$$10. = \frac{d}{d} + \sqrt{a^2 \left(\frac{d}{d} \right)^2 + b^2}$$

$$\Rightarrow - \frac{d}{d} = \sqrt{a^2 \left(\frac{d}{d} \right)^2 + b^2}$$

Squaring on both sides, we get

$$^2 - 2 \frac{d}{d} + ^2 \left(\frac{d}{d} \right)^2 = a^2 \left(\frac{d}{d} \right)^2 + b^2$$

This is a differential equation of order 1 and degree 2.

$$11. \left[1 + \left(\frac{d^2}{d^2} \right)^3 \right]^{4/5} = \left(\frac{m}{m+1} \right) \frac{d^3}{d^3}$$

$$\Rightarrow \left\{ \left[1 + \left(\frac{d^2}{d^2} \right)^3 \right]^{4/5} \right\}^5 = \left(\frac{m}{m+1} \right)^5 \left(\frac{d^3}{d^3} \right)^5$$

$$\Rightarrow \left[1 + \left(\frac{d^2}{d^2} \right)^3 \right]^4 = \left(\frac{m}{m+1} \right)^5 \left(\frac{d^3}{d^3} \right)^5$$

Here, the highest order derivative is $\frac{d^3}{d^3}$ with power 5.

∴ order = 3 and degree = 5

$$12. \left(\frac{d^2}{d^2} \right)^5 + 4 \frac{\left(\frac{d^2}{d^2} \right)^3}{\left(\frac{d^3}{d^3} \right)} + \frac{d^3}{d^3} = ^2 - 1$$

$$\therefore \left(\frac{d^2}{d^2} \right)^5 \cdot \left(\frac{d^3}{d^3} \right) + 4 \left(\frac{d^2}{d^2} \right)^3 + \left(\frac{d^3}{d^3} \right)^2 = (^2 - 1) \cdot \frac{d^3}{d^3}$$

Here, the highest order derivative is $\frac{d^3}{d^3}$ with power 2.

∴ order = 3 and degree = 2

∴ m = 3 and n = 2

13. Option (A) has order = 4, degree = 1

Option (B) has order = 3, degree = 4

Consider option (C),

$$\left[1 + \left(\frac{d}{d} \right)^3 \right]^{2/3} = 4 \frac{d^3}{d^3}$$

Cubing on both sides, we get

$$\left[1 + \left(\frac{d}{d} \right)^3 \right]^2 = 4^3 \left(\frac{d^3}{d^3} \right)^3$$

Here, order = 3 and degree = 3

∴ option (C) is the correct answer.

14. Since, the given equation has 3 arbitrary constants i.e., g, f and c, therefore order of the given differential equation is 3.

15. Since, the given equation has 3 arbitrary constants i.e., a, b and c, therefore order of the given differential equation is 3.

16. The equation of a family of circles of radius r passing through the origin and having centre on Y-axis is $(x - 0)^2 + (y - r)^2 = r^2$ or $x^2 + y^2 - 2ry = 0$.

Since this equation has one arbitrary constant, its order is 1.

17. The equation of the family of circles which touch both the axes is $(x - a)^2 + (y - a)^2 = a^2$, where a is a parameter.

Since this equation has one arbitrary constant, its order is 1.

$$18. = ae^m + be^{-m} \quad \dots(i)$$

$$\Rightarrow \frac{d}{d} = mae^m - mbe^{-m}$$

$$\Rightarrow \frac{d^2}{d^2} = m^2ae^m + m^2be^{-m}$$

$$\Rightarrow \frac{d^2}{d^2} = m^2(ae^m + be^{-m}) = m^2 \quad \dots[\text{From (i)}]$$

$$\Rightarrow \frac{d^2}{d^2} - m^2 = 0$$



$$19. \quad = c + c - c^3 \quad \dots(i)$$

$$\Rightarrow \frac{d}{d} = c \quad \dots(ii)$$

Substituting (ii) in (i), we get

$$= \frac{d}{d} + \frac{d}{d} - \left(\frac{d}{d}\right)^3$$

$$20. \quad = A \cos \omega t + B \sin \omega t \quad \dots(i)$$

$$\Rightarrow ' = -A \omega \sin \omega t + B \omega \cos \omega t$$

$$\Rightarrow '' = -A \omega^2 \cos \omega t - B \omega^2 \sin \omega t$$

$$\Rightarrow '' = -\omega^2 (A \cos \omega t + B \sin \omega t)$$

$$\Rightarrow '' = -\omega^2 \quad \dots[\text{From (i)}]$$

$$21. \quad = a^{n+1} + b^{-n} \quad \dots(i)$$

$$\Rightarrow \frac{d}{d} = a(n+1)n^{-n} - bn^{-n-1}$$

$$\Rightarrow \frac{d^2}{d^2} = a(n+1)n^{-n-1} + n(n+1)b^{-n-2}$$

$$\Rightarrow \frac{d^2}{d^2} = a(n+1)n^{-n-1} + bn(n+1) \cdot^{-n}$$

$$= n(n+1)(a^{n+1} + b^{-n})$$

$$\therefore \frac{d^2}{d^2} = n(n+1) \quad \dots[\text{From (i)}]$$

$$22. \quad = c_1 \cos a + c_2 \sin a \quad \dots(i)$$

$$\Rightarrow \frac{d}{d} = -c_1 a \sin a + c_2 a \cos a$$

$$\Rightarrow \frac{d^2}{d^2} = -c_1 a^2 \cos a - c_2 a^2 \sin a$$

$$\Rightarrow \frac{d^2}{d^2} = -a^2 (c_1 \cos a + c_2 \sin a)$$

$$\Rightarrow \frac{d^2}{d^2} = -a^2 \quad \dots[\text{From (i)}]$$

$$\Rightarrow \frac{d^2}{d^2} + a^2 = 0$$

$$23. \quad \sin^{-1} + \sin^{-1} = c$$

Differentiating w.r.t. , we get

$$\frac{1}{\sqrt{1-x^2}} + \frac{1}{\sqrt{1-x^2}} \cdot \frac{d}{d} = 0$$

$$\Rightarrow \frac{d}{d} = -\frac{\sqrt{1-x^2}}{\sqrt{1-x^2}}$$

$$\Rightarrow \sqrt{1-x^2} \frac{d}{d} + \sqrt{1-x^2} \frac{d}{d} = 0$$

$$24. \quad = (\sin^{-1})^2 + A \cos^{-1} + B$$

$$\Rightarrow = (\sin^{-1})^2 + A \left(\frac{\pi}{2} - \sin^{-1} \right) + B$$

$$\dots \left[\because \sin^{-1} + \cos^{-1} = \frac{\pi}{2} \right]$$

$$\Rightarrow = (\sin^{-1})^2 - A \sin^{-1} + \frac{\pi A}{2} + B \quad \dots(i)$$

Differentiating w.r.t. , we get

$$\frac{d}{d} = \frac{2 \sin^{-1}}{\sqrt{1-x^2}} - \frac{A}{\sqrt{1-x^2}}$$

$$\Rightarrow (1-x^2) \left(\frac{d}{d} \right)^2 = (2 \sin^{-1} - A)^2$$

$$= 4(\sin^{-1})^2 - 4A \sin^{-1} + A^2$$

$$= 4[(\sin^{-1})^2 - A \sin^{-1}] + A^2$$

$$= 4 \left(-\frac{\pi A}{2} - B \right) + A^2$$

....[From (i)]

$$\therefore (1-x^2) \left(\frac{d}{d} \right)^2 = 4 - 2\pi A - 4B + A^2$$

Differentiating w.r.t. , we get

$$(1-x^2) \cdot 2 \frac{d}{d} \cdot \frac{d^2}{d^2} - 2 \left(\frac{d}{d} \right)^2 = 4 \frac{d}{d}$$

$$\Rightarrow (1-x^2) \frac{d^2}{d^2} - \frac{d}{d} = 2$$

25. The equation of all the straight lines passing through the origin is

$$= m \quad \dots(i)$$

$$\therefore \frac{d}{d} = m$$

$$\Rightarrow \frac{d}{d} = - \quad \dots[\text{From (i)}]$$

$$26. \quad v = \frac{A}{r} + B$$

Differentiating w.r.t. r, we get

$$\frac{dv}{dr} = -\frac{A}{r^2} \quad \dots(i)$$

$$\therefore \frac{d^2 v}{dr^2} = 2A \cdot r^{-3}$$

$$= -2 r^2 \frac{dv}{dr} \cdot r^{-3} \quad \dots[\text{From (i)}]$$

$$\therefore \frac{d^2 v}{dr^2} + \frac{2}{r} \cdot \frac{dv}{dr} = 0$$



27. The equation of the family of lines passing through $(1, -1)$ is

$$y + 1 = m(x - 1)$$

$$\Rightarrow y = m(x - 1) - 1 \quad \dots(i)$$

$$\Rightarrow \frac{dy}{dx} = m$$

Substituting the value of m in (i), we get

$$= \frac{dy}{dx} (x - 1) - 1$$

28. $y = e^x$

Taking logarithm on both sides, we get

$$\log y = \log e^x + c \quad \dots(i)$$

$$\Rightarrow c = \frac{1}{x} \log y - \dots(ii)$$

Differentiating (i) w.r.t. x , we get

$$\frac{1}{y} \cdot \frac{dy}{dx} = \frac{1}{x} + c$$

$$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = \frac{1}{x} + \frac{1}{x} \log y \quad \dots[\text{From (ii)}]$$

$$\Rightarrow \frac{dy}{dx} = - \left[1 + \log y \right]$$

29. The system of circles which passes through origin and whose centre lies on $Y = a$ is

$$x^2 + y^2 - 2ay = 0 \quad \dots(i)$$

Differentiating w.r.t. x , we get

$$2x + 2y \frac{dy}{dx} - 2a \frac{dy}{dx} = 0$$

$$\Rightarrow 2a = 2x + 2y \frac{dy}{dx} \quad \dots(ii)$$

Substituting (ii) in (i), we get

$$x^2 + y^2 - 2x^2 - 2y \frac{dy}{dx} = 0$$

$$\Rightarrow (x^2 - y^2) \frac{dy}{dx} - 2xy = 0$$

30. $\frac{dy}{dx} = e^{-x} + 2e^{-2x}$

$$\Rightarrow \frac{dy}{dx} = e^{-x} (e^x + 2)$$

Integrating on both sides, we get

$$\int e^x dy = \int (e^x + 2) dx + c$$

$$\Rightarrow e^x y = e^{\frac{3x}{3}} + c$$

31. $\log\left(\frac{dy}{dx}\right) = x + c$

$$\Rightarrow \frac{dy}{dx} = e^{x+c} \Rightarrow \frac{dy}{dx} = e^x \cdot e^c$$

Integrating on both sides, we get

$$\int e^x dy - \int e^{-x} dx = c$$

$$\Rightarrow e^x + e^{-x} = c$$

32. $\cos x = (e^{\log x} + e^{-\log x})^x$

$$\Rightarrow \cos x = e^{\left(\log x + \frac{1}{x}\right) x}$$

Integrating on both sides, we get

$$\sin x = e^{\log x + c}$$

33. $\frac{dy}{dx} = 2^{-x}$

$$\Rightarrow \frac{dy}{dx} = \frac{2}{2^x}$$

Integrating on both sides, we get

$$\int 2^{-x} dy - \int 2^{-x} dx = c_1$$

$$\Rightarrow \frac{-2}{\log 2} + \frac{2}{\log 2} = c_1$$

$$\Rightarrow \frac{1}{2} - \frac{1}{2} = c_1 \log 2$$

$$\Rightarrow \frac{1}{2} - \frac{1}{2} = c, \text{ where } c = c_1 \log 2$$

34. $\frac{dy}{dx} + 2y = 1 \Rightarrow \frac{dy}{dx} = (1 - 2y)$

Integrating on both sides, we get

$$\int \frac{dy}{1-2y} = \int (1-2y) dx + c_1$$

$$\Rightarrow \log y = -2x + c_1$$

$$\Rightarrow y = e^{-2x} \cdot e^{c_1}$$

$$\Rightarrow y = c \cdot e^{-2x}, \text{ where } c = e^{c_1}$$

35. $\frac{dy}{dx} = \frac{1 - \cos x}{1 + \cos x} = \tan^2 \frac{x}{2}$

$$\Rightarrow dy = \left(\sec^2 \frac{x}{2} - 1 \right) dx$$

Integrating on both sides, we get

$$= 2 \tan \frac{x}{2} - x + c$$



$$36. \frac{d}{d} + \frac{1 + \cos 2}{1 - \cos 2} = 0$$

$$\Rightarrow \frac{d}{d} + \frac{2 \cos^2}{2 \sin^2} = 0$$

$$\Rightarrow \frac{d}{\cos^2} + \frac{d}{\sin^2} = 0$$

Integrating on both sides, we get

$$\int \sec^2 d + \int \operatorname{cosec}^2 d = c$$

$$\Rightarrow \tan - \cot = c$$

$$37. (e^2 - 1)d + (e^{-2} - 1)e d = 0$$

$$\Rightarrow (e^2 - 1)d = (1 - e^{-2})e d$$

Integrating on both sides, we get

$$\int \frac{e^2 - 1}{e} d - \int \frac{1 - e^{-2}}{e} d = c$$

$$\Rightarrow \int e d - \int e^{-1} d = \int \frac{1}{e} d - \int \frac{1}{e} d + c$$

$$\Rightarrow e + e^{-1} = \log - \frac{1}{e} + c$$

$$38. \frac{d^2}{d} = -2 \frac{d}{d}$$

$$\Rightarrow \frac{d}{d} = -\frac{2}{d} d$$

Integrating on both sides, we get

$$\int \frac{2}{d} d + \int \frac{d}{d} = \log c$$

$$\Rightarrow \log d^2 + \log d = \log c$$

$$\Rightarrow \log d^3 = \log c \Rightarrow d^3 = c$$

$$39. \cot d = \frac{d}{d}$$

Integrating on both sides, we get

$$\int \frac{d}{d} = \int \tan d + \log c$$

$$\Rightarrow \log d = \log (\sec d) + \log c$$

$$\Rightarrow \log d = \log (c \cdot \sec d) \Rightarrow d = c \sec d$$

$$40. \frac{d}{d} = \cot d - \cot d$$

$$\Rightarrow \cot d - \tan d = 0$$

Integrating on both sides, we get

$$\log (\sin d) - \log (\sec d) = \log c$$

$$\Rightarrow \log \left(\frac{\sin d}{\sec d} \right) = \log c \Rightarrow \sin d = c \sec d$$

$$41. \sec \frac{d}{d} = 1 \Rightarrow \sec d = \frac{d}{d}$$

Integrating on both sides, we get

$$\log (\sec d + \tan d) = \log d + \log c$$

$$\Rightarrow \log (\sec d + \tan d) = \log (c d)$$

$$\Rightarrow \sec d + \tan d = c d$$

$$42. (e + 1) \cos d + e \sin d = 0$$

$$\Rightarrow \frac{e}{e + 1} d + \frac{\cos}{\sin} d = 0$$

Integrating on both sides, we get

$$\int \frac{e}{e + 1} d + \int \frac{\cos}{\sin} d = \log c$$

$$\Rightarrow \log (e + 1) + \log (\sin d) = \log c$$

$$\Rightarrow (e + 1) \sin d = c$$

$$43. \frac{d}{d} + (1 + d^2) \tan^{-1} d = 0$$

Integrating on both sides, we get

$$\int \frac{d}{(1 + d^2) \tan^{-1} d} + \int \frac{d}{d} = \log c$$

$$\Rightarrow \log (\tan^{-1} d) + \log d = \log c$$

$$\Rightarrow \log (\tan^{-1} d \cdot d) = \log c$$

$$\Rightarrow \tan^{-1} d = \frac{c}{d}$$

$$44. 3e \tan d + (1 - e) \sec^2 d = 0$$

$$\Rightarrow \frac{\sec^2}{\tan} d = -3 \frac{e}{1 - e} d$$

Integrating on both sides, we get

$$\int \frac{\sec^2}{\tan} d = -3 \int \frac{e}{1 - e} d + \log c$$

$$\Rightarrow \log (\tan d) - 3 \log (1 - e) + \log c$$

$$\Rightarrow \log (\tan d) = \log [(1 - e)^3 c]$$

$$\Rightarrow \tan d = c(1 - e)^3$$

$$45. (\sin d + \cos d) d + (\cos d - \sin d) d = 0$$

$$\Rightarrow d = -\left(\frac{\cos d - \sin d}{\sin d + \cos d} \right) d$$

Integrating on both sides, we get

$$= -\log (\sin d + \cos d) + \log c$$

$$\Rightarrow = \log \left(\frac{c}{\sin d + \cos d} \right)$$

$$\Rightarrow e (\sin d + \cos d) = c$$

$$46. \frac{d}{d} = \frac{1 + d}{1 + d}$$

$$\Rightarrow \left(\frac{1 + d}{1 + d} \right) d = \left(\frac{1 + d}{1 + d} \right) d$$

Integrating on both sides, we get

$$\log d + \log (1 + d) = \log d + \log (1 + d) + \log A$$

$$\Rightarrow \log \left(\frac{d(1 + d)}{A} \right) = \log d + \log (1 + d) \Rightarrow \frac{d(1 + d)}{A} = d(1 + d)$$



$$47. \quad \frac{d}{d} + = 2 \Rightarrow \frac{d}{d} = 2 -$$

$$\Rightarrow \frac{d}{2 -} = \frac{d}{d}$$

$$\Rightarrow \left[\frac{1}{-1} - \frac{1}{-} \right] d = \frac{d}{d}$$

Integrating on both sides, we get

$$\log(-1) - \log = \log + \log c$$

$$\Rightarrow \log \left(\frac{-1}{-} \right) = \log(c)$$

$$\Rightarrow \frac{-1}{-} = c \Rightarrow = 1 + c$$

$$48. \quad (2 - 1)d - (2 + 3)d = 0$$

Integrating on both sides, we get

$$\int \frac{d}{2 + 3} - \int \frac{d}{2 - 1} = \log c_1$$

$$\Rightarrow \frac{1}{2} \log(2 + 3) - \frac{1}{2} \log(2 - 1) = \log c_1$$

$$\Rightarrow \log(2 + 3) - \log(2 - 1) = 2 \log c_1$$

$$\Rightarrow \log \left(\frac{2 + 3}{2 - 1} \right) = \log c_1^2$$

$$\Rightarrow \frac{2 + 3}{2 - 1} = c, \text{ where } c = c_1^2$$

$$49. \quad (-^2)d = (-^2)d$$

$$\Rightarrow (1 - ^2)d = (1 - ^2)d$$

Integrating on both sides, we get

$$\int \frac{d}{1 - ^2} - \int \frac{d}{1 - ^2} = \log c$$

$$\Rightarrow -\frac{1}{2} \log(1 - ^2) + \frac{1}{2} \log(1 - ^2) = \log c$$

$$\Rightarrow \log(1 - ^2) - \log(1 - ^2) = 2 \log c$$

$$\Rightarrow \frac{1 - ^2}{1 - ^2} = c^2$$

$$\Rightarrow 1 - ^2 = c^2(1 - ^2)$$

$$50. \quad (1 - ^2)d + d = ^2d$$

$$\Rightarrow (1 - ^2)d = (^2 -)d$$

Integrating on both sides, we get

$$\int \frac{d}{(-1)} = \int \frac{d}{1 - ^2} + \log c$$

$$\Rightarrow \int \left(\frac{1}{-1} - \frac{1}{-} \right) d = \frac{-1}{2} \int \frac{-2}{1 - ^2} d + \log c$$

$$\Rightarrow \log(-1) - \log = \frac{-1}{2} \log(1 - ^2) + \log c$$

$$\Rightarrow 2 \log(-1) + \log(1 - ^2) = 2 \log c + 2 \log$$

$$\Rightarrow \log[(-1)^2(1 - ^2)] = \log c^2$$

$$\Rightarrow (-1)^2(1 - ^2) = c^2$$

$$51. \quad (^2 - ^2) \frac{d}{d} + ^2 + ^2 = 0$$

$$\Rightarrow ^2(1 -) \frac{d}{d} + ^2(1 +) = 0$$

$$\Rightarrow \frac{(1 -)}{2} d + \frac{(1 +)}{2} d = 0$$

Integrating on both sides, we get

$$\int \left(\frac{1}{2} - \frac{1}{-} \right) d + \int \left(\frac{1}{2} + \frac{1}{-} \right) d = c$$

$$\Rightarrow -\frac{1}{-} - \log - \frac{1}{-} + \log = c$$

$$\Rightarrow \log \left(- \right) = \frac{1}{-} + \frac{1}{-} + c$$

$$52. \quad \frac{d}{d} \tan = \sin(+) + \sin(-)$$

$$\Rightarrow \frac{d}{d} \cdot \frac{\sin}{\cos} = 2 \sin \cos$$

$$\Rightarrow \frac{\sin}{\cos^2} d = 2 \sin d$$

Integrating on both sides, we get

$$\int \frac{\sin}{\cos^2} d - 2 \int \sin d = c$$

$$\Rightarrow \frac{1}{\cos} + 2 \cos = c$$

$$\Rightarrow \sec + 2 \cos = c$$

$$53. \quad \frac{d}{d} = \frac{1 + ^2}{1 + ^2}$$

$$\Rightarrow \frac{d}{1 + ^2} - \frac{d}{1 + ^2} = 0$$

Integrating on both sides, we get

$$\int \frac{d}{1 + ^2} - \int \frac{d}{1 + ^2} = \tan^{-1} c$$

$$\Rightarrow \tan^{-1} - \tan^{-1} = \tan^{-1} c$$

$$\Rightarrow \tan^{-1} \left(\frac{-}{1 +} \right) = \tan^{-1} c$$

$$\Rightarrow - = c(1 +)$$



$$54. \quad \frac{d}{d} = \frac{(1+x^2)(1+x^{-2})}{(1+x^2)}$$

Integrating on both sides, we get

$$\begin{aligned} \int \frac{d}{1+x^2} d &= \int \frac{1+x^2+x^{-2}}{(1+x^2)} d + c \\ \Rightarrow \frac{1}{2} \int \frac{2}{1+x^2} d &= \int \frac{1}{1+x^2} d + \int \frac{d}{1+x^2} + c \\ \Rightarrow \frac{1}{2} \log(1+x^2) &= \log x + \tan^{-1} x + c \end{aligned}$$

$$55. \quad (\operatorname{cosec} \log x) d + (x^2) d = 0 \\ \Rightarrow \frac{1}{x} \log x + x^2 \sin x = 0$$

Integrating on both sides, we get

$$\begin{aligned} \frac{(\log x)^2}{2} + [x^2(-\cos x) + \int 2x \cos x dx] &= c \\ \Rightarrow \frac{(\log x)^2}{2} - x^2 \cos x + 2(\sin x + \cos x) &= c \\ \Rightarrow \frac{(\log x)^2}{2} + (2-x^2) \cos x + 2 \sin x &= c \end{aligned}$$

$$56. \quad \frac{d}{d} = \frac{\log^2 x + \sin x + \cos x}{\sin x + \cos x}$$

Integrating on both sides, we get

$$\begin{aligned} \int (\sin x + \cos x) d &= \int (\log^2 x + \sin x + \cos x) d + c \\ \Rightarrow -\cos x + \sin x + \cos x &= \frac{2}{2} \log^2 x - \int \frac{2}{2} \cdot \frac{1}{2} \cdot 2x dx + \int dx + c \\ \Rightarrow \sin x &= \frac{2}{2} \times 2 \log x - \int dx + \int dx + c \\ \Rightarrow \sin x &= 2 \log x + c \end{aligned}$$

$$57. \quad \cos \log(\sec x + \tan x) d \\ = \cos \log(\sec x + \tan x) d$$

Integrating on both sides, we get

$$\begin{aligned} \int \sec \log(\sec x + \tan x) d \\ = \int \sec \log(\sec x + \tan x) d + c \end{aligned}$$

$$\text{Put } \log(\sec x + \tan x) = t \Rightarrow \sec x dx = dt$$

$$\text{and } \log(\sec x + \tan x) = z \Rightarrow \sec x dx = dz$$

$$\therefore \int t dt = \int z dz + c$$

$$\Rightarrow \frac{t^2}{2} = \frac{z^2}{2} + c$$

$$\Rightarrow \frac{[\log(\sec x + \tan x)]^2}{2} = \frac{[\log(\sec x + \tan x)]^2}{2} + c$$

$$58. \quad \sqrt{a+x} \frac{d}{d} + \frac{d}{d} = 0$$

Integrating on both sides, we get

$$\begin{aligned} \int d + \int \frac{d}{\sqrt{a+x}} d &= c \\ \Rightarrow \int \frac{+a-a}{\sqrt{a+x}} d &= c \\ \Rightarrow \int \left(\sqrt{a+x} - \frac{a}{\sqrt{a+x}} \right) d &= c \\ \Rightarrow \frac{2}{3} (a+x)^{\frac{3}{2}} - 2a\sqrt{a+x} &= c \end{aligned}$$

$$\Rightarrow 3 + 2(a+x)^{\frac{3}{2}} - 6a\sqrt{a+x} = 3c$$

$$\Rightarrow 3 + 2\sqrt{a+x} (a+x - 3a) = 3c$$

$$\Rightarrow 3 + 2\sqrt{a+x} (x - 2a) = 3c$$

$$59. \quad d + d + x^2 d - x^2 d = 0$$

$$\Rightarrow \frac{d}{2} + \frac{d}{2} + \frac{d}{2} - \frac{d}{2} = 0$$

$$\Rightarrow \frac{d}{2} + \frac{d}{2} - \frac{d}{2} = 0$$

Integrating on both sides, we get

$$-\frac{1}{x} + \log x - \log x = k$$

$$\Rightarrow \log x = \frac{1}{x} + k$$

$$60. \quad e^{-x} d - (e^{-x} + x^3) d = 0$$

$$\Rightarrow e^{-x} (d - d) = x^3 d$$

$$\Rightarrow e^{-x} \frac{(d - d)}{2} = d$$

$$\Rightarrow e^{-x} d \left(- \right) = d$$

Integrating on both sides, we get

$$-e^{-x} = \frac{x^2}{2} + c \Rightarrow \frac{x^2}{2} + e^{-x} = k,$$

where $k = -c$

$$61. \quad x' = 1 + x^2 + x^{-2}$$

$$\Rightarrow \frac{d}{d} = (1+x^2)(1+x^{-2})$$

Integrating on both sides, we get

$$\int \frac{d}{1+x^2} = \int (1+x^2) d + c$$

$$\Rightarrow \tan^{-1} x = x + \frac{2}{x} + c \quad \dots(i)$$



Since, $(0) = 0$ i.e., $= 0$, when $= 0$

$$\therefore \tan^{-1}(0) = 0 + c \Rightarrow c = 0$$

$$\therefore \tan^{-1} = + \frac{^2}{2} \quad \dots[\text{From (i)}]$$

$$\Rightarrow = \tan\left(+ \frac{^2}{2}\right)$$

62. $' - = 1$

$$\Rightarrow \frac{d}{d} - = 1$$

$$\Rightarrow \frac{d}{d} = 1 +$$

Integrating on both sides, we get

$$\int \frac{d}{1+} = \int d + c$$

$$\Rightarrow \log(1 +) = + c$$

Since, $(0) = 1$ i.e., $= 1$, when $= 0$

$$\therefore \log(1 + 1) = 0 + c \Rightarrow c = \log 2$$

$$\therefore \log(1 +) = + \log 2$$

$$\Rightarrow \log\left(\frac{1+}{2}\right) =$$

$$\Rightarrow \frac{1+}{2} = e$$

$$\Rightarrow = 2e^{-1}$$

$$\Rightarrow () = 2\exp() - 1$$

63. $e^{d/d} = (+ 1)$

$$\Rightarrow \frac{d}{d} = \log(+ 1)$$

Integrating on both sides, we get

$$\int d = \int \log(+ 1) d + c$$

$$\Rightarrow = \log(+ 1) - \int \frac{d}{+1} + c$$

$$= \log(+ 1) - \int \frac{+1-1}{+1} d + c$$

$$= \log(+ 1) - \int \left(1 - \frac{1}{+1}\right) d + c$$

$$\therefore = \log(+ 1) - + \log(+ 1) + c \quad \dots(i)$$

Since, $(0) = 3$ i.e., $= 3$, when $= 0$

$$\therefore 3 = 0 + c \Rightarrow c = 3$$

$$\therefore = \log(+ 1) + \log(+ 1) - + 3$$

$\dots[\text{From (i)}]$

$$\therefore = (+ 1) \log(+ 1) - + 3$$

64. $\frac{d}{d} = e^{+} - 1 \quad \dots(i)$

Put $+ = v \quad \dots(ii)$

$$\Rightarrow 1 + \frac{d}{d} = \frac{dv}{d}$$

$$\Rightarrow \frac{d}{d} = \frac{dv}{d} - 1 \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$\frac{dv}{d} = e^v$$

Integrating on both sides, we get

$$\int e^{-v} dv = \int d + c$$

$$\Rightarrow -e^{-v} = + c$$

$$\Rightarrow + e^{-v} + c = 0$$

$$\Rightarrow + e^{-(+)} + c = 0$$

65. $\frac{d}{d} = \sin(+) \quad \dots(i)$

Put $+ = v \quad \dots(ii)$

$$\Rightarrow \frac{d}{d} = \frac{dv}{d} - 1 \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$\frac{dv}{d} - 1 = \sin v$$

$$\Rightarrow \frac{dv}{1 + \sin v} = d \Rightarrow \frac{1 - \sin v}{\cos^2 v} dv = d$$

Integrating on both sides, we get

$$\int \sec^2 v dv - \int \sec v \tan v dv = \int d + c$$

$$\Rightarrow \tan v - \sec v = + c$$

$$\Rightarrow \tan(+) - \sec(+) = + c$$

66. $\frac{d}{d} = \sin(+) \tan(+) - 1 \quad \dots(i)$

Put $+ = v \quad \dots(ii)$

$$\Rightarrow 1 + \frac{d}{d} = \frac{dv}{d}$$

$$\Rightarrow \frac{d}{d} = \frac{dv}{d} - 1 \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$\frac{dv}{d} = \sin v \tan v \Rightarrow \frac{dv}{d} = \frac{\sin^2 v}{\cos v}$$

Integrating on both sides, we get

$$\int d - \int \frac{\cos v}{\sin^2 v} dv = c \Rightarrow - \left(-\frac{1}{\sin v} \right) = c$$

$\dots[\text{Put } \sin v = t \Rightarrow \cos v dv = dt]$

$$\Rightarrow + \operatorname{cosec} v = c$$

$$\Rightarrow + \operatorname{cosec}(+) = c$$



$$67. \quad \frac{d}{d} = \cos(\quad) + \sin(\quad) \quad \dots(i)$$

$$\text{Put } \quad = v \quad \dots(ii)$$

$$\Rightarrow 1 + \frac{d}{d} = \frac{dv}{d}$$

$$\Rightarrow \frac{d}{d} = \frac{dv}{d} - 1 \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$\frac{dv}{d} - 1 = \cos v + \sin v$$

$$\Rightarrow \frac{dv}{d} = 1 + \cos v + \sin v$$

Integrating on both sides, we get

$$\int \frac{dv}{1 + \cos v + \sin v} = \int d + c$$

$$\Rightarrow \int \frac{dv}{1 + \frac{1 - \tan^2 v/2}{1 + \tan^2 v/2} + \frac{2 \tan v/2}{1 + \tan^2 v/2}} = + c$$

$$\Rightarrow \int \frac{\sec^2(v/2)}{2(1 + \tan v/2)} dv = + c$$

$$\Rightarrow \log|1 + \tan v/2| = + c$$

$$\Rightarrow \log \left| 1 + \tan \left(\frac{\quad}{2} \right) \right| = + c$$

$$68. \quad \frac{d}{d} = \frac{\quad^2 + \quad^2}{2} \quad \dots(i)$$

$$\text{Put } \quad = v \quad \dots(ii)$$

$$\therefore \frac{d}{d} = v + \frac{dv}{d} \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = \frac{\quad^2 + v^2 + v^2}{2} = 1 + v + v^2$$

$$\Rightarrow \frac{dv}{d} = 1 + v^2 \Rightarrow \frac{dv}{1 + v^2} = \frac{d}{d}$$

Integrating on both sides, we get

$$\tan^{-1} v = \log + c$$

$$\Rightarrow \tan^{-1} \left(\frac{\quad}{\quad} \right) = \log + c$$

$$69. \quad \frac{d}{d} = - \left(\log - + 1 \right) \quad \dots(i)$$

$$\text{Put } \quad = v \quad \dots(ii)$$

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = v(\log v + 1) \Rightarrow \frac{dv}{d} = v \log v$$

Integrating on both sides, we get

$$\int \frac{dv}{v \log v} = \int \frac{d}{\quad} + \log c$$

$$\Rightarrow \log(\log v) = \log + \log c$$

$$\dots \left[\text{Put } \log v = t \Rightarrow \frac{1}{v} dv = dt \right]$$

$$\Rightarrow \log v = c \Rightarrow \log \left(\frac{\quad}{\quad} \right) = c$$

$$70. \quad (\quad) d + \quad = 0$$

$$\Rightarrow \frac{d}{d} = - \left(\frac{\quad}{\quad} \right) \quad \dots(i)$$

$$\text{Put } \quad = v \quad \dots(ii)$$

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = \frac{-v}{\quad} = -1 - v$$

$$\Rightarrow \frac{dv}{d} = -1 - 2v$$

Integrating on both sides, we get

$$\int \frac{dv}{1 + 2v} = - \int \frac{d}{\quad} + \log c_1$$

$$\Rightarrow \frac{1}{2} \log(1 + 2v) = - \log + \log c_1$$

$$\Rightarrow \log \left(1 + 2 \frac{\quad}{\quad} \right) = 2 \log \frac{c_1}{\quad}$$

$$\Rightarrow \frac{\quad + 2}{\quad} = \left(\frac{c_1}{\quad} \right)^2$$

$$\Rightarrow \quad + 2 = c_1^2$$

$$\Rightarrow \quad + 2 = c, \text{ where } c = c_1^2$$

$$71. \quad + \frac{d}{d} = 2 \Rightarrow - + \frac{d}{d} = 2 \quad \dots(i)$$

$$\text{Put } \quad = v \quad \dots(ii)$$

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$\frac{1}{v} + v + \frac{dv}{d} = 2$$

$$\Rightarrow v + \frac{dv}{d} = \frac{2v - 1}{v} \Rightarrow \frac{v}{(v - 1)^2} dv = - \frac{d}{\quad}$$

$$\Rightarrow \frac{v - 1 + 1}{(v - 1)^2} dv = - \frac{d}{\quad}$$

$$\Rightarrow \left[\frac{1}{(v - 1)} + \frac{1}{(v - 1)^2} \right] dv = - \frac{d}{\quad}$$



Integrating on both sides, we get

$$\int \frac{dv}{v-1} + \int \frac{dv}{(v-1)^2} = - \int \frac{d}{\dots} + c$$

$$\Rightarrow \log(v-1) - \frac{1}{v-1} = - \log \dots + c$$

$$\Rightarrow \log(\dots) = \dots + c$$

72. $2d + (2 - \dots + 2)d = 0$

$$\Rightarrow \frac{d}{d} = \frac{\dots}{2 - \dots + 2} \dots(i)$$

Put $\dots = v \dots(ii)$

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = \frac{-v^2 + 2}{2 - \dots + v^2 + 2}$$

$$\Rightarrow \frac{dv}{d} = \frac{-v^2}{1 - v + v^2} - v$$

$$\Rightarrow \frac{dv}{d} = \frac{-v - v^3}{1 - v + v^2}$$

Integrating on both sides, we get

$$\int \frac{v^2 - v + 1}{v(v^2 + 1)} dv = - \int \frac{d}{\dots} + c$$

$$\Rightarrow \int \left(\frac{1}{v} - \frac{1}{v^2 + 1} \right) dv = - \int \frac{d}{\dots} + c$$

$$\Rightarrow \log v - \tan^{-1} v = - \log \dots + c$$

$$\Rightarrow \log \left(\dots \right) - \tan^{-1} \left(\dots \right) = - \log \dots + c$$

$$\Rightarrow \log \dots = \tan^{-1} \left(\dots \right) + c$$

73. $2 \frac{d}{d} = 2 + 3 \dots^2$

$$\Rightarrow \frac{d}{d} = \frac{2 + 3 \dots^2}{2} \dots(i)$$

Put $\dots = v \dots(ii)$

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = \frac{2 + 3v^2}{2v^2}$$

$$\Rightarrow \frac{dv}{d} = \frac{2(1 + 3v^2)}{2v^2} - v$$

$$\Rightarrow \frac{dv}{d} = \frac{1 + 3v^2}{2v} - v$$

$$\Rightarrow \frac{dv}{d} = \frac{1 + v^2}{2v}$$

Integrating on both sides, we get

$$\int \frac{2v}{1 + v^2} dv = \int \frac{d}{\dots} + \log p$$

$$\Rightarrow \log(1 + v^2) = \log \dots + \log p$$

$$\Rightarrow \log \left(\frac{1 + v^2}{\dots} \right) = \log p$$

$$\Rightarrow \frac{1 + v^2}{\dots} = p \Rightarrow \frac{2 + \dots^2}{3} = p$$

$$\Rightarrow 2 + \dots^2 = p \cdot 3$$

74. $\frac{d}{d} = \frac{\dots}{2 - \dots} \dots(i)$

Put $\dots = v \dots(ii)$

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = \frac{1}{2v - \dots} = \frac{1}{2v - 1}$$

$$\Rightarrow \frac{dv}{d} = \frac{1}{2v - 1} - v = \frac{1 - 2v^2 + v}{2v - 1}$$

$$\Rightarrow \frac{dv}{d} = - \frac{(v-1)(2v+1)}{2v-1}$$

$$\Rightarrow \frac{(2v-1)}{(2v+1)(v-1)} dv = \frac{-d}{\dots}$$

$$\Rightarrow \frac{1}{3(v-1)} + \frac{4}{3(2v+1)} = \frac{-d}{\dots}$$

Integrating on both sides, we get

$$\frac{1}{3} \log(v-1) + \frac{4}{3} \cdot \frac{1}{2} \log(2v+1)$$

$$= - \log \dots + \log c_1$$

$$\Rightarrow \log(v-1)^{1/3} + \log(2v+1)^{2/3} = \log \frac{c_1}{\dots}$$

$$\Rightarrow (v-1)^{1/3} (2v+1)^{2/3} = \frac{c_1}{\dots}$$

$$\Rightarrow \left(\dots \right) \left(\frac{2 + \dots}{\dots} \right)^2 = \frac{c_1^3}{3}$$

$$\Rightarrow (\dots)(\dots + 2)^2 = c, \text{ where } c = -c_1^3$$



$$75. \quad \frac{d}{d} = \frac{-}{+} \quad \dots(i)$$

$$\text{Put } = v \quad \dots(ii)$$

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = \frac{-v}{+v}$$

$$\Rightarrow \frac{dv}{d} = \frac{1-v}{1+v} - v$$

$$\Rightarrow \frac{dv}{d} = \frac{1-2v-v^2}{1+v}$$

Integrating on both sides, we get

$$\int \frac{1+v}{2-(1+v)^2} dv = \int \frac{d}{d} + \log c_1$$

$$\Rightarrow -\frac{1}{2} \log[2-(1+v)^2] = \log + \log c_1$$

$$\Rightarrow -\frac{1}{2} \log(1-2v-v^2) = \log(c_1)$$

$$\Rightarrow \log\left(\frac{1}{\sqrt{1-2v-v^2}}\right) = \log(c_1)$$

$$\Rightarrow \frac{1}{\sqrt{1-2v-v^2}} = c_1$$

$$\Rightarrow \frac{1}{1-2v-v^2} = {}^2c_1^2$$

$$\Rightarrow {}^2c_1^2(1-2v-v^2) = 1$$

$$\Rightarrow {}^2c_1^2\left(1-\frac{2}{-}-\frac{2}{-}\right) = 1$$

$$\Rightarrow c_1^2({}^2-2-{}^2) = 1$$

$$\Rightarrow {}^2+2-{}^2 = c, \text{ where } c = \frac{-1}{c_1^2}$$

76. A differential equation in which the dependent variable (y) and its differential coefficient occur only in the first degree and are not multiplied together is called a linear differential equation.

Hence, $\frac{d}{d} + 4 = 0$ is a non-linear differential equation.

77. $\frac{d}{d} + = e$ can be written as

$$\frac{d}{d} + \frac{e}{2} = \frac{e}{2}, \text{ which is a linear equation.}$$

$$78. \quad \frac{d}{d} + \frac{2}{1+{}^2} = \frac{{}^2-1}{{}^2+1}$$

$$\therefore \text{I.F.} = e^{\int \frac{2}{1+{}^2} d} = e^{\log(1+{}^2)} = 1+{}^2$$

$$79. \quad (\log) \frac{d}{d} + = 2 \log$$

$$\Rightarrow \frac{d}{d} + \frac{1}{\log} = \frac{2}{\log}$$

$$\therefore \text{I.F.} = e^{\int \frac{1}{\log} d} = e^{\log(\log)} = \log$$

$$80. \quad (1-{}^2) \frac{d}{d} - = 1$$

$$\Rightarrow \frac{d}{d} - \frac{1}{1-{}^2} = \frac{1}{1-{}^2}$$

$$\therefore \text{I.F.} = e^{\int \frac{-}{1-{}^2} d} = e^{\frac{1}{2} \log(1-{}^2)} = e^{\log(\sqrt{1-{}^2})} = \sqrt{1-{}^2}$$

$$81. \quad \frac{d}{d} + \log = .e^{-\frac{1}{2} \log}$$

$$\Rightarrow \frac{d}{d} + \frac{\log}{d} = e^{-\frac{1}{2} \log}$$

$$\therefore \text{I.F.} = e^{\int \frac{\log}{d} d} = e^{\frac{1}{2}(\log)^2}$$

$$= \left(e^{\frac{1}{2} \log} \right)^{\log} \quad \dots[\because (a^m)^n = a^{mn}]$$

$$= (\sqrt{})^{\log}$$

$$82. \quad (1+{}^2) \frac{d}{d} - (\tan^{-1} -) \frac{d}{d} = 0$$

$$\Rightarrow (1+{}^2) \frac{d}{d} = (\tan^{-1} -) \frac{d}{d}$$

$$\Rightarrow \frac{d}{d} = \frac{\tan^{-1} -}{1+{}^2}$$

$$\Rightarrow \frac{d}{d} + \frac{1}{1+{}^2} = \frac{\tan^{-1}}{1+{}^2}$$

This is the linear differential equation of the

$$\text{form } \frac{d}{d} + P = Q, \text{ where } P = \frac{1}{1+{}^2}$$

$$\therefore \text{I.F.} = e^{\int P d} = e^{\int \frac{1}{1+{}^2} d} = e^{\tan^{-1}}$$

$$83. \quad \frac{d}{d} + 2 \cot = 3 {}^2 \text{cosec}^2$$

$$\therefore \text{I.F.} = e^{\int 2 \cot d} = e^{2 \log \sin} = \sin^2$$

\therefore solution of the given equation is

$$\sin^2 = \int 3 {}^2 \text{cosec}^2 \cdot \sin^2 d + c$$

$$\Rightarrow \sin^2 = \int 3 {}^2 d + c \Rightarrow \sin^2 = {}^3 + c$$



$$84. \quad \frac{d}{d} + 2 \tan = \sin$$

Here, $P = 2 \tan$ and $Q = \sin$

$$\therefore \text{I.F.} = e^{\int 2 \tan d} \\ = e^{2 \log(\sec)} = e^{\log \sec^2} = \sec^2$$

\therefore solution of the given equation is

$$(\sec^2) = \int \sin \sec^2 d + c \\ \Rightarrow \sec^2 = \int \sec \tan d + c \\ \Rightarrow \sec^2 = \sec + c$$

$$85. \quad \frac{d}{d} = \frac{1}{+ + 1} \Rightarrow \frac{d}{d} = + + 1 \\ \Rightarrow \frac{d}{d} - = + 1$$

$$\therefore \text{I.F.} = e^{\int -1 d} = e^{-}$$

\therefore solution of the given equation is

$$.e^{-} = \int (+1)e^{-} d + c \\ \Rightarrow e^{-} = e^{-} (- - 2) + c \\ \Rightarrow = ce - - 2$$

$$86. \quad \frac{d}{d} + = 2 + 3 + 2 \Rightarrow \frac{d}{d} + -- = + 3 + \frac{2}{-}$$

Here, $P = \frac{1}{-}$, $Q = + 3 + \frac{2}{-}$

$$\therefore \text{I.F.} = e^{\int \frac{1}{-} d} = e^{\log} =$$

\therefore solution of the given equation is

$$. = \int \left(+ 3 + \frac{2}{-} \right) d + c \\ \Rightarrow = \int ^2 d + \int 3 d + \int 2d + c \\ \Rightarrow = \frac{3}{3} + \frac{3^2}{2} + 2 + c$$

$$87. \quad \log \frac{d}{d} + = 2 \log$$

$$\Rightarrow \frac{d}{d} + \frac{1}{\log} = \frac{2}{-}$$

$$\therefore \text{I.F.} = e^{\int \frac{1}{\log} d} = e^{\log(\log)} = \log$$

\therefore solution of the given equation is

$$\log = \int \frac{2}{-} \cdot \log d + c \\ \Rightarrow \log = (\log)^2 + c$$

$$88. \quad \frac{d}{d} + \frac{3^2}{1+^3} = \frac{\sin^2}{1+^3}$$

Here, $P = \frac{3^2}{1+^3}$ and $Q = \frac{\sin^2}{1+^3}$

$$\therefore \text{I.F.} = e^{\int \frac{3^2}{1+^3} d} = e^{\log(1+^3)} = 1+^3$$

\therefore solution of the given equation is

$$.(1+^3) = \int \frac{\sin^2}{1+^3} (1+^3) d \\ \Rightarrow (1+^3) = \int \frac{1 - \cos 2}{2} d \\ \Rightarrow (1+^3) = \frac{1}{2} - \frac{\sin 2}{4} + c$$

$$89. \quad \frac{d}{d} + \sec^2 \tan \sec^2$$

Here, $P = \sec^2$, $Q = \tan \sec^2$

$$\therefore \text{I.F.} = e^{\int \sec^2 d} = e^{\tan}$$

\therefore solution of the given equation is

$$.e^{\tan} = \int \tan \cdot \sec^2 e^{\tan} d + c$$

Put $\tan = t \Rightarrow \sec^2 d = dt$

$$\therefore e^{\tan} = \int t e^t dt + c$$

$$\Rightarrow e^{\tan} = t e^t - e^t + c$$

$$\Rightarrow e^{\tan} = e^{\tan} (\tan - 1) + c$$

$$\Rightarrow = \tan - 1 + c \cdot e^{-\tan}$$

$$90. \quad (+ 2^3) \frac{d}{d} - = 0$$

$$\Rightarrow \frac{d}{d} = \frac{-}{+ 2^3}$$

$$\Rightarrow \frac{d}{d} = \frac{- + 2^3}{-} \Rightarrow \frac{d}{d} - - = 2^2$$

$$\therefore \text{I.F.} = e^{\int \frac{-1}{-} d} = e^{-\log} = \frac{1}{-}$$

\therefore solution of the given equation is

$$(\text{I.F.}) = \int Q(\text{I.F.}) d + c$$

$$\Rightarrow \frac{1}{-} = \int 2^2 \cdot \frac{1}{-} d + c$$

$$\Rightarrow - = ^2 + c$$

$$\Rightarrow = ^3 + c.$$

$$\Rightarrow ^3 - = -c$$

$$\Rightarrow ^3 - = A, \text{ where } A = -c$$



$$91. \quad d + d + \log d = 0$$

$$\Rightarrow d + d = -\log d$$

$$\Rightarrow \frac{d}{d} + \frac{d}{d} = -\log d \Rightarrow \frac{d}{d} + \frac{d}{d} = -\frac{\log d}{d}$$

\therefore I.F. = $e^{\int -1/d} = e^{-\log d} = \frac{1}{d}$
 \therefore solution of the given equation is

$$\frac{1}{d} \cdot d = -\int \frac{\log d}{d} + c$$

$$\Rightarrow \frac{d}{d} = -(\log d - 1) + c$$

$$\Rightarrow \frac{d}{d} + (\log d - 1) = c$$

$$92. \quad \frac{d}{d} = \tan^{-2} \sec$$

$$\Rightarrow \frac{1}{2} \cdot \frac{d}{d} - \frac{1}{2} \tan^{-2} = -\sec \quad \dots(i)$$

Put $v = \tan^{-1}$

$$\Rightarrow \frac{dv}{d} = -\frac{1}{2} \cdot \frac{d}{d}$$

$$\therefore -\frac{dv}{d} - v \tan^{-2} = -\sec \quad \dots[\text{From (i)}]$$

$$\Rightarrow \frac{dv}{d} + v \tan^{-2} = \sec$$

This is the standard form of the linear differential equation.

$$\therefore \text{I.F.} = e^{\int \tan^{-2} d} = e^{\log \sec} = \sec$$

$$93. \quad \frac{d}{d} = 2 + 3e \Rightarrow \frac{d}{d} - \frac{2}{d} = 2e$$

$$\therefore \text{I.F.} = e^{\int -2/d} = e^{-2 \log d} = e^{\log d^{-2}} = \frac{1}{d^2}$$

\therefore solution of the given equation is

$$\frac{1}{d^2} \cdot d = \int 2e \cdot \frac{1}{d^2} d + c$$

$$\Rightarrow \frac{d}{d^2} = e + c \quad \dots(i)$$

Since, $d = 0$, when $t = 1$

$$\therefore 0 = e^1 + c \Rightarrow c = -e$$

$$\therefore \frac{d}{d^2} = e - e \quad \dots[\text{From (i)}]$$

$$\Rightarrow \frac{d}{d^2} = 2(e - e)$$

$$94. \quad d = (d + d)$$

$$\Rightarrow d = (-2)d \Rightarrow \frac{d}{d} + \left(-\frac{1}{d}\right) = -$$

$$\therefore \text{I.F.} = e^{\int -1/d} = e^{-\log d} = \frac{1}{d}$$

\therefore solution of the given equation is

$$\frac{1}{d} = \int -\frac{1}{d} d + c$$

$$\Rightarrow \frac{1}{d} = -\frac{1}{d} + c \quad \dots(i)$$

Since, $(1) = 1$ i.e., $d = 1$, when $t = 1$

$$\therefore 1 = -1 + c \Rightarrow c = 2$$

$$\therefore \frac{1}{d} = -\frac{1}{d} + 2 \quad \dots[\text{From (i)}]$$

Putting $d = -3$, we get

$$-\frac{3}{d} = -\frac{1}{d} + 2$$

$$\Rightarrow \frac{3}{d} - \frac{1}{d} - 2 = 0$$

$$\Rightarrow (-3 - 1) \left(\frac{1}{d}\right) = 2$$

Since $(d) > 0$, $d = -3$

$$95. \quad \frac{d}{d} = \frac{1}{d} + 2$$

$$\therefore \frac{d}{d} - \frac{1}{d} = 2$$

$$\therefore \text{I.F.} = e^{\int -1/d} = e^{-\log d} = \frac{1}{d}$$

\therefore solution of the given equation is

$$\frac{1}{d} \cdot d = \int 2 \cdot \frac{1}{d} d + c$$

$$\frac{d}{d} = 2 \log d + c$$

$$\therefore \frac{d}{d} = 2(\log d + 1) + c$$

$$\therefore \frac{d}{d} = 2(\log d + 1) + c \quad \dots(i)$$

Since, the curve passes through origin $(0, 0)$.

$$\therefore 0 = 2(\log 0 + 1) + c$$

$$\Rightarrow c = -2$$

$$\therefore \frac{d}{d} = 2(\log d + 1) - 2$$

$$\therefore \frac{d}{d} + 2(\log d + 1) = 2e \quad \dots[\text{From (i)}]$$

96. Let P be the population at time t years.

$$\text{Then, } \frac{dP}{dt} = kP$$

$$\Rightarrow \frac{dP}{P} = k dt$$

Integrating on both sides, we get

$$\log P = kt + c$$

When $t = 0$, $P = 40000$

$$\therefore \log 40000 = 0 + c \Rightarrow c = \log 40000$$

$$\therefore \log P = kt + \log 40000$$

$$\Rightarrow \log \left(\frac{P}{40000}\right) = kt \quad \dots(i)$$

When $t = 40$ yrs, $P = 60000$

$$\therefore \log \left(\frac{60000}{40000}\right) = 40k \Rightarrow k = \frac{1}{40} \log \left(\frac{3}{2}\right)$$



$$\therefore \log \left(\frac{P}{40000} \right) = \frac{t}{40} \log \left(\frac{3}{2} \right) \quad \dots[\text{From (i)}]$$

When $t = 60$ yrs, we have

$$\log \left(\frac{P}{40000} \right) = \frac{60}{40} \log \left(\frac{3}{2} \right)$$

$$\Rightarrow \frac{P}{40000} = \left(\frac{3}{2} \right)^{\frac{3}{2}} \Rightarrow \frac{P}{40000} = \frac{3}{2} \left(\frac{3}{2} \right)^{\frac{1}{2}}$$

$$\Rightarrow P = 40000 \times \frac{3}{2} \times 1.2247 = 73482$$

97. Let P_0 be the initial population and let the population after t years be P . Then,

$$\frac{dP}{dt} = kP \Rightarrow \frac{dP}{P} = kdt$$

Integrating on both sides, we get

$$\log P = kt + c$$

When $t = 0$, $P = P_0$

$$\therefore \log P_0 = 0 + c \Rightarrow c = \log P_0$$

$$\therefore \log P = kt + \log P_0$$

$$\Rightarrow \log \frac{P}{P_0} = kt \quad \dots(\text{i})$$

When $t = 5$ hrs, $P = 2P_0$

$$\therefore \log \frac{2P_0}{P_0} = 5k$$

$$\Rightarrow k = \frac{\log 2}{5}$$

$$\therefore \log \frac{P}{P_0} = \frac{\log 2}{5} t \quad \dots[\text{From (i)}]$$

When $t = 25$ hrs, we have

$$\log \frac{P}{P_0} = \frac{\log 2}{5} \times 25 = 5 \log 2 = \log 32$$

$$\therefore P = 32P_0$$

98. Let ' θ ' be the temperature of the body at any time ' t '.

$$\therefore \frac{d\theta}{dt} \propto (\theta - 20)$$

$$\therefore \frac{d\theta}{dt} = k(\theta - 20)$$

Integrating on both sides, we get

$$\log (\theta - 20) = kt + c$$

When $t = 0$, $\theta = 100^\circ \text{C}$

$$\therefore \log (100 - 20) = k(0) + c \Rightarrow c = \log 80$$

$$\therefore \log (\theta - 20) = kt + \log 80 \quad \dots(\text{i})$$

When $t = 20$, $\theta = 60^\circ \text{C}$

$$\therefore \log (60 - 20) = k(20) + \log 80$$

$$\Rightarrow k = \frac{1}{20} \log \left(\frac{1}{2} \right)$$

$$\therefore \log (\theta - 20) = \frac{t}{20} \log \left(\frac{1}{2} \right) + \log 80$$

$\dots[\text{From (i)}]$

When $\theta = 30^\circ \text{C}$, we have

$$\log (30 - 20) = t \left(\frac{1}{20} \right) \log \left(\frac{1}{2} \right) + \log 80$$

$$\Rightarrow \log 10 - \log 80 = \frac{t}{20} \log \left(\frac{1}{2} \right)$$

$$\Rightarrow \log \left(\frac{1}{8} \right) = \frac{t}{20} \log \left(\frac{1}{2} \right)$$

$$\Rightarrow 3 \log \left(\frac{1}{2} \right) = \frac{t}{20} \log \left(\frac{1}{2} \right)$$

$$\Rightarrow \frac{t}{20} = 3$$

$$\Rightarrow t = 60 \text{ minutes}$$

99. Let ' d ' be the number of bacteria present at time ' t '.

$$\therefore \frac{d}{dt} \propto$$

$$\therefore \frac{d}{dt} = k$$

Integrating on both sides, we get

$$\log d = kt + c$$

When $t = 0$, $d = 1000$

$$\therefore \log (1000) = k(0) + c$$

$$\Rightarrow c = \log (1000)$$

$$\therefore \log d = kt + \log (1000) \quad \dots(\text{i})$$

When $t = 1$, $d = 2000$

$$\therefore \log (2000) = k(1) + \log (1000)$$

$$\Rightarrow k = \log \left(\frac{2000}{1000} \right) = \log 2$$

$$\therefore \log d = t \log 2 + \log (1000) \quad \dots[\text{From (i)}]$$

When $t = 2 \frac{1}{2} = \frac{5}{2}$, we have

$$\log d = \left(\frac{5}{2} \right) \log 2 + \log (1000)$$

$$= \log \left(2^{\frac{5}{2}} \right) + \log (1000)$$

$$= \log (4\sqrt{2}) + \log (1000)$$

$$= \log (4000\sqrt{2})$$

$$= \log (4000 \times 1.414)$$

$$\therefore \log d = \log (5656)$$

$$\Rightarrow d = 5656$$

**Competitive Thinking**

1. Here, the highest order derivative is $\frac{d^3}{d^3}$.

\therefore order = 3

2. Here, the order of the differential equation is 1.

3. Here, the highest order derivative is $\frac{d^3}{d^3}$.

\therefore order = 3

6. Here, the highest order derivative is $\frac{d^3}{d^3}$ with power 2.

\therefore order = 3, degree = 2

$$7. = \frac{d}{d} + \frac{2}{\frac{d}{d}}$$

$$\Rightarrow \frac{d}{d} = \left(\frac{d}{d}\right)^2 + 2$$

\therefore order = 1, degree = 2

$$8. = \left(\frac{d}{d}\right)^2 + \left(\frac{d}{d}\right)^2$$

$$\Rightarrow \left(\frac{d}{d}\right)^2 = \left(\frac{d}{d}\right)^4 + 1$$

\therefore order = 1, degree = 4

$$9. \left(1 + 3\frac{d}{d}\right)^{\frac{2}{3}} = 4\frac{d^3}{d^3}$$

$$\Rightarrow \left(1 + 3\frac{d}{d}\right)^2 = 4^3 \left(\frac{d^3}{d^3}\right)^3$$

Here, the highest order derivative is $\frac{d^3}{d^3}$ with power 3.

\therefore order = 3 and degree = 3

$$10. \frac{d^2}{d^2} = \sqrt[3]{1 + \left(\frac{d}{d}\right)^2}$$

$$\Rightarrow \left(\frac{d^2}{d^2}\right)^3 = 1 + \left(\frac{d}{d}\right)^2$$

Here, the highest order derivatives is $\frac{d^2}{d^2}$ with power 3

\therefore Order = 2 and degree = 3

$$11. \left[1 + \left(\frac{d}{d}\right)^3\right]^{\frac{7}{3}} = 7\left(\frac{d^2}{d^2}\right)$$

$$\Rightarrow \left[1 + \left(\frac{d}{d}\right)^3\right]^7 = 7^3 \left(\frac{d^2}{d^2}\right)^3$$

Here, the highest order derivative is $\frac{d^2}{d^2}$ with power 3.

\therefore order = 2 and degree = 3

$$12. \frac{d^3}{d^3} = \sqrt[5]{1 - \left(\frac{d}{d}\right)^7}$$

$$\Rightarrow \left(\frac{d^3}{d^3}\right)^5 = 1 - \left(\frac{d}{d}\right)^7$$

\therefore order = 3, degree = 5

$$13. p. \frac{d^2}{d^2} = \left[1 + \left(\frac{d}{d}\right)^2\right]^{\frac{3}{2}}$$

$$\Rightarrow \left(p. \frac{d^2}{d^2}\right)^2 = \left[1 + \left(\frac{d}{d}\right)^2\right]^3$$

\therefore order = 2 and degree = 2

$$14. \left(\frac{d^2}{d^2}\right)^{1/3} + \left(+\frac{d}{d}\right)^{1/2} = 0$$

$$\Rightarrow \left(\frac{d^2}{d^2}\right)^{1/3} = -\left(+\frac{d}{d}\right)^{1/2}$$

$$\Rightarrow \left(\frac{d^2}{d^2}\right)^2 = \left(+\frac{d}{d}\right)^3$$

\therefore order = 2 and degree = 2

$$15. (1 + \frac{d^2}{d^2})^{2/3} = 2$$

$$\Rightarrow (1 + \frac{d^2}{d^2})^2 = (2)^3$$

\therefore order(n) = 2, degree(m) = 3

$$\therefore \frac{m+n}{m-n} = \frac{3+2}{3-2} = 5$$

$$16. = p + \sqrt[3]{a^2 p^2 + b^2}$$

$$\Rightarrow (-p)^3 = a^2 p^2 + b^2$$

$$\Rightarrow -p^3 - 3^2 p^2 + 3p^2 - p^3 = a^2 p^2 + b^2$$

Here, $p = \frac{d}{d}$

\therefore order = 1, degree = 3



$$17. \left[1 + \left(\frac{d}{d} \right)^2 + \sin \left(\frac{d}{d} \right) \right]^4 = \frac{d^2}{d^2}$$

$$\Rightarrow \left[1 + \left(\frac{d}{d} \right)^2 + \sin \left(\frac{d}{d} \right) \right]^3 = \left(\frac{d^2}{d^2} \right)^4$$

Here, the highest order derivative is $\frac{d^2}{d^2}$.

∴ order = 2
 Since, the given differential equation cannot be expressed as polynomial in differential coefficients, the degree is not defined.

$$18. \frac{3}{2} - \frac{1}{1} - 4 = 0$$

$$\Rightarrow \frac{3}{2} = \frac{1}{1} + 4$$

Squaring on both sides, we get

$$\frac{3}{2} = \left(\frac{1}{1} + 4 \right)^2 = \frac{1}{1} + 16 + 8 \frac{1}{1}$$

$$\Rightarrow \frac{3}{2} - \frac{1}{1} - 16 = 8 \frac{1}{1}$$

Squaring on both sides, we get

$$\left(\frac{3}{2} - \frac{1}{1} - 16 \right)^2 = 64 \frac{1}{1}$$

Here, the highest order derivative is $\frac{1}{2}$ with power 6.

∴ degree = 6

$$19. \sqrt{\sin} (d + d) = \sqrt{\cos} (d - d)$$

$$\Rightarrow \frac{\sqrt{\sin}}{\sqrt{\cos}} (d + d) = d - d$$

$$\Rightarrow \sqrt{\tan} \left(1 + \frac{d}{d} \right) = 1 - \frac{d}{d}$$

$$\Rightarrow \frac{d}{d} = \frac{1 - \sqrt{\tan}}{1 + \sqrt{\tan}}$$

This is a differential equation of order 1 and degree 1.

$$20. () = 1 + \frac{d}{d} + \frac{1}{1.2} \left(\frac{d}{d} \right)^2 + \frac{1}{1.2.3} \left(\frac{d}{d} \right)^3 + \dots$$

$$\Rightarrow () = e^{\frac{d}{d}} \dots \left[\because e = 1 + \frac{2}{2!} + \frac{3}{3!} + \dots \right]$$

$$\Rightarrow \frac{d}{d} = \log$$

This is a differential equation of degree 1.

$$21. = C_1 e^{+C_2} + C_3 e + C_4 \sin(+ C_5)$$

$$= C_1 \cdot e^{C_2} e^2 + C_3 e + C_4 (\sin \cos C_5 + \cos \sin C_5)$$

$$= A e^2 + C_3 e + B \sin + D \cos ,$$

where $A = C_1 e^{C_2}$, $B = C_4 \cos C_5$, $D = C_4 \sin C_5$

Since, this equation consists of four arbitrary constants.

∴ order of differential equation = 4

$$22. \text{ Consider option (C),}$$

$$= 2 - 4$$

$$\therefore \frac{d}{d} = 2$$

$$\therefore \left(\frac{d}{d} \right)^2 - \frac{d}{d} + 2^2 - 2 + 2 - 4 = 0$$

$$23. = a + \frac{b}{d}$$

$$\Rightarrow \frac{d}{d} = -\frac{b}{2} \dots(i)$$

$$\Rightarrow \frac{d^2}{d^2} = \frac{2b}{3} \Rightarrow \frac{d^2}{d^2} = \frac{2b}{2}$$

$$\Rightarrow \frac{d^2}{d^2} - \frac{2b}{2} = 0$$

$$\Rightarrow \frac{d^2}{d^2} + \frac{2d}{d} = 0 \dots[\text{From (i)}]$$

$$24. = e^{-} \cos 2$$

$$\Rightarrow \frac{d}{d} = -2e^{-} \sin 2 - e^{-} \cos 2$$

$$\Rightarrow \frac{d^2}{d^2} = 4e^{-} \sin 2 - 3e^{-} \cos 2$$

$$\therefore \frac{d^2}{d^2} + \frac{2d}{d} + 5 = 0$$

$$25. = m + \frac{4}{m} \dots(i)$$

$$\Rightarrow \frac{d}{d} = m$$

Putting $m = \frac{d}{d}$ in (i), we get

$$\left(\frac{d}{d} \right) = \left(\frac{d}{d} \right)^2 + 4$$

$$26. = a e^b \dots(i)$$

$$\Rightarrow \frac{d}{d} = a b e^b$$

$$\Rightarrow \frac{d}{d} = b \dots(ii) [\text{From (i)}]$$

$$\Rightarrow \frac{d^2}{d^2} = b \frac{d}{d} \Rightarrow \frac{d^2}{d^2} = b \frac{d}{d}$$

$$\Rightarrow \frac{d^2}{d^2} - \left(\frac{d}{d} \right)^2 = 0 \dots[\text{From (ii)}]$$



27. $y = \frac{A}{x} + Bx^2$
 $\Rightarrow y = A + Bx^3$
 Differentiating w.r.t. x , we get
 $\frac{dy}{dx} + 3Bx^2 = 3Bx^2 \quad \dots(i)$
 Again, differentiating w.r.t. x , we get
 $\frac{d^2y}{dx^2} + 2 \frac{dy}{dx} = 6B$
 $\Rightarrow 2 \frac{d^2y}{dx^2} + 2 \frac{dy}{dx} = 6B$
 $\Rightarrow 2 \frac{d^2y}{dx^2} + 2(3Bx^2 - \frac{A}{x^2}) = 6B \quad \dots[\text{From (i)}]$
 $\Rightarrow 2 \frac{d^2y}{dx^2} = 2$

28. $y = e^m$
 $\Rightarrow \log y = m \quad \dots(i)$
 Differentiating w.r.t. x , we get
 $\frac{1}{y} \cdot \frac{dy}{dx} = m$
 $\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = \frac{\log y}{x} \quad \dots[\text{From (i)}]$
 $\Rightarrow \frac{dy}{dx} = \left(-\frac{1}{y}\right) \log y$

29. $y = e(A \cos x + B \sin x)$
 $\Rightarrow y' = e(A \cos x + B \sin x) + e(B \cos x - A \sin x)$
 $\Rightarrow y' = e(A \cos x + B \sin x) + e(B \cos x - A \sin x) \quad \dots(i)$
 $\therefore y'' = y' + e(B \cos x - A \sin x) - e(A \cos x + B \sin x)$
 $\Rightarrow y'' = y' + (B \cos x - A \sin x) - (A \cos x + B \sin x) \quad \dots[\text{From (i)}]$
 $\Rightarrow y'' - 2y' + 2y = 0$

30. $y = a \sin(\log x) + b \cos(\log x) \quad \dots(i)$
 $\Rightarrow \frac{dy}{dx} = \frac{a \cos(\log x) - b \sin(\log x)}{x}$
 $\Rightarrow \frac{dy}{dx} = a \cos(\log x) - b \sin(\log x)$
 Differentiating w.r.t. x , we get
 $\frac{d^2y}{dx^2} + \frac{dy}{dx} = \frac{-a \sin(\log x) - b \cos(\log x)}{x^2}$
 $\Rightarrow \frac{d^2y}{dx^2} + \frac{dy}{dx} = -\frac{1}{x^2} [a \sin(\log x) + b \cos(\log x)]$
 $\Rightarrow 2 \frac{d^2y}{dx^2} + \frac{dy}{dx} = -\frac{2}{x^2} \quad \dots[\text{From (i)}]$
 $\Rightarrow 2 \frac{d^2y}{dx^2} + \frac{dy}{dx} + \frac{2}{x^2} = 0$

31. Differentiating the given equation, we get
 $\frac{dy}{dx} = A$
 $\therefore y = \left(\frac{dy}{dx}\right)^3$, which is of degree 3

32. $y = A \cos(nt + \alpha)$
 Differentiating w.r.t. t , we get
 $\frac{dy}{dt} = -A \sin(nt + \alpha) \cdot n$
 $\Rightarrow \frac{dy}{dt} = -An \sin(nt + \alpha)$
 Differentiating w.r.t. t , we get
 $\frac{d^2y}{dt^2} = -An^2 \cos(nt + \alpha)$
 $\Rightarrow \frac{d^2y}{dt^2} = -n^2 y$
 $\Rightarrow \frac{d^2y}{dt^2} + n^2 y = 0$

33. $y = e^{a \sin x} \quad \dots(i)$
 $\Rightarrow \log y = a \sin x \quad \dots(ii)$
 Differentiating w.r.t. x , we get
 $\frac{1}{y} \cdot \frac{dy}{dx} = a \cos x$
 $\Rightarrow a = \frac{1}{\cos x} \cdot \frac{dy}{dx}$
 Putting the value of a in (i), we get
 $\log y = \tan x \cdot \frac{dy}{dx}$

34. $y^2 = 2d \left(x + \sqrt{d} \right) \quad \dots(i)$
 Differentiating w.r.t. x , we get
 $2 \frac{dy}{dx} = 2d \quad \dots(ii)$
 Substituting (ii) in (i), we get
 $y^2 = 2 \frac{dy}{dx} \left(x + \sqrt{\frac{dy}{dx}} \right)$
 $\Rightarrow y^2 = 2 \frac{dy}{dx} x + 2 \frac{dy}{dx} \cdot \sqrt{\frac{dy}{dx}}$
 $\Rightarrow \left(-2 \frac{dy}{dx} \right)^2 = 4 \left(\frac{dy}{dx} \right)^3$
 This is a differential equation of order 1 and degree 3.

35. Required equation of parabola is
 $(y - k)^2 = 4a(x - h)$
 Since, this equation has two arbitrary constants, its order is 2.



36. Equation of family of parabolas whose axis is X-axis is $y^2 = 4a(x - h)$

Differentiating w.r.t. x , we get

$$2 \frac{dy}{dx} = 4a$$

$$\Rightarrow \frac{dy}{dx} = 2a$$

Again, differentiating w.r.t. x , we get

$$\frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^2 = 0$$

\therefore order = 2 and degree = 1

37. $y = a \cos\left(\frac{1}{x+b}\right) \dots(i)$

$$\Rightarrow \frac{dy}{dx} = -a \sin\left(\frac{1}{x+b}\right) \cdot \left(-\frac{1}{x^2}\right) + a \cos\left(\frac{1}{x+b}\right)$$

$$\Rightarrow \frac{dy}{dx} = \frac{a}{x^2} \sin\left(\frac{1}{x+b}\right) + a \cos\left(\frac{1}{x+b}\right)$$

$$\begin{aligned} \Rightarrow \frac{d^2y}{dx^2} &= \frac{a}{x^2} \cos\left(\frac{1}{x+b}\right) \cdot \left(-\frac{1}{x^3}\right) \\ &\quad - \frac{a}{x^2} \sin\left(\frac{1}{x+b}\right) - a \sin\left(\frac{1}{x+b}\right) \cdot \left(-\frac{1}{x^2}\right) \\ &= -\frac{a}{x^3} \cos\left(\frac{1}{x+b}\right) - \frac{a}{x^4} \cos\left(\frac{1}{x+b}\right) \end{aligned}$$

$\therefore \frac{d^2y}{dx^2} = -\frac{a}{x^4} \dots[\text{From (i)}]$

$$\Rightarrow \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^2 = 0 \Rightarrow \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^2 = 0$$

38. The differential equation representing the family of parabolas having vertex at origin is $y^2 = 4ax \dots(i)$

Differentiating w.r.t. x , we get

$$2 \frac{dy}{dx} = 4a$$

$$\Rightarrow 2 \frac{dy}{dx} = \frac{2y}{x} \dots[\text{From (i)}]$$

$$\Rightarrow 2 \frac{dy}{dx} = \frac{2y}{x}$$

$$\Rightarrow x^2 - 2x \frac{dy}{dx} = 0$$

39. Equation of family of parabolas with focus at $(0, 0)$ and X-axis is $y^2 = 4a(x + a) \dots(i)$

Differentiating (i) w.r.t. x , we get

$$2 \frac{dy}{dx} = 4a \dots(ii)$$

Substituting (ii) in (i), we get

$$y^2 = 2 \frac{dy}{dx} \left(x + \frac{dy}{dx} \right)$$

$$\Rightarrow y^2 = 2 \frac{dy}{dx} x + \left(\frac{dy}{dx}\right)^2$$

$$\Rightarrow - \left(\frac{dy}{dx}\right)^2 = 2 \frac{dy}{dx} x - y^2$$

40. Equation of family of parabolas with focus at $(0, 0)$ and X-axis is $y^2 = 4a(x + a) \dots(i)$

Differentiating (i) w.r.t. x , we get

$$2 \frac{dy}{dx} = 4a \dots(ii)$$

Substituting (ii) in (i), we get

$$y^2 = 2 \frac{dy}{dx} \left(x + \frac{dy}{dx} \right)$$

$$\Rightarrow y^2 = 2 \frac{dy}{dx} x + \left(\frac{dy}{dx}\right)^2$$

\therefore order = $m = 1$, degree = $n = 2$

$$\text{Now, } mn - m + n = 1(2) - 1 + 2 = 3$$

41. Axis of parabola = X-axis

vertex = $(m, 0)$

Equation of all parabolas is

$$(y - 0)^2 = 4a(x - m)$$

$$\Rightarrow y^2 = 4ax - 4am$$

$\therefore 2 \frac{dy}{dx} = 4a$

$$\Rightarrow \frac{dy}{dx} = 2a$$

$\therefore \frac{d^2y}{dx^2} + \frac{dy}{dx} \left(\frac{dy}{dx}\right) = 0$

$$\Rightarrow \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^2 = 0$$

42. Axis of parabola = Y-axis

Vertex = $(0, m)$

\therefore Equation of parabola is

$$(y - 0)^2 = 4a(x - m)$$

$$\Rightarrow y^2 = 4ax - 4am$$

Differentiating w.r.t. x , we get

$$2y = 4a \frac{dy}{dx}$$

$$\Rightarrow \frac{1}{y} \frac{dy}{dx} = \frac{1}{2a} \Rightarrow \frac{1}{y} \frac{d^2y}{dx^2} - \frac{dy}{dx} \frac{1}{y^2} = 0$$

$$\Rightarrow \frac{d^2y}{dx^2} - \frac{dy}{dx} = 0$$



43. The differential equation of the family of circles touching Y-axis at the origin is
 $x^2 + y^2 - 2ax = 0$ (i)

Differentiating w.r.t. x , we get

$$2x + 2y \frac{dy}{dx} - 2a = 0$$

$$\Rightarrow 2a = 2x + 2y \frac{dy}{dx} \quad \dots\text{(ii)}$$

Substituting (ii) in (i), we get

$$x^2 + y^2 - 2x^2 - 2y \frac{dy}{dx} = 0$$

$$\Rightarrow x^2 - y^2 + 2y \frac{dy}{dx} = 0$$

44. The system of circles which passes through origin and whose centre lies on X-axis is
 $x^2 + y^2 - 2bx = 0$ (i)

Differentiating w.r.t. x , we get

$$2x + 2y \frac{dy}{dx} = 2b \quad \dots\text{(ii)}$$

Substituting (ii) in (i), we get

$$x^2 + y^2 - 2x^2 - 2y \frac{dy}{dx} = 0$$

$$x^2 - y^2 - 2y \frac{dy}{dx} = 0$$

45. The equation of the family of lines which are at a unit distance from the origin is
 $\cos \alpha + \sin \alpha = 1$ (i)

Differentiating w.r.t. x , we get

$$\cos \alpha + \sin \alpha \frac{d\alpha}{dx} = 0 \quad \dots\text{(ii)}$$

By (i) \times (ii), we get

$$\sin \alpha \left(-\frac{d\alpha}{dx} \right) = 1$$

$$\Rightarrow -\frac{d\alpha}{dx} = \operatorname{cosec} \alpha \quad \dots\text{(iii)}$$

$$\text{From (ii), } \left(\frac{d\alpha}{dx} \right)^2 = \cot^2 \alpha = \operatorname{cosec}^2 \alpha - 1$$

$$\therefore \left(\frac{d\alpha}{dx} \right)^2 = \left(-\frac{d\alpha}{dx} \right)^2 - 1 \quad \dots\text{[From (iii)]}$$

$$\therefore 1 + \left(\frac{d\alpha}{dx} \right)^2 = \left(-\frac{d\alpha}{dx} \right)^2$$

46. $\sec \alpha - \operatorname{cosec} \alpha = 0$

$$\Rightarrow \cos \alpha - \sin \alpha = 0$$

Integrating on both sides, we get

$$\sin \alpha + \cos \alpha = c$$

$$47. \frac{d}{dx} = 1 + \dots + \dots$$

$$\Rightarrow \frac{d}{dx} = (1 + \dots)(1 + \dots) \Rightarrow \frac{d}{1 + \dots} = (1 + \dots) dx$$

Integrating on both sides, we get

$$\log(1 + \dots) = \frac{\dots^2}{2} + c$$

$$48. 9 \frac{d}{dx} + 4 = 0$$

Integrating on both sides, we get

$$9 \int \frac{d}{dx} + 4 \int dx = c_1$$

$$\Rightarrow 9 \cdot \frac{\dots^2}{2} + 4 \cdot \frac{\dots^2}{2} = c_1$$

$$\Rightarrow \frac{\dots^2}{4} + \frac{\dots^2}{9} = \frac{c_1}{18}$$

$$\Rightarrow \frac{\dots^2}{4} + \frac{\dots^2}{9} = c, \text{ where } c = \frac{c_1}{18}$$

$$49. (1 + \dots) \tan^{-1} \dots + (1 + \dots)^2 \frac{d}{dx} = 0$$

$$\Rightarrow \frac{\tan^{-1} \dots}{1 + \dots} + \frac{2}{1 + \dots} \frac{d}{dx} = 0$$

Integrating on both sides, we get

$$\frac{(\tan^{-1} \dots)^2}{2} + \log |1 + \dots| = c_1$$

$$\Rightarrow (\tan^{-1} \dots)^2 + 2 \log |1 + \dots| = c, \text{ where } c = 2c_1$$

$$50. \left(\frac{d}{dx} \right)^2 + 2\sqrt{\dots} \frac{d}{dx} + \dots = 0$$

$$\Rightarrow \left(\sqrt{\dots} \frac{d}{dx} + \sqrt{\dots} \right)^2 = 0$$

$$\Rightarrow \sqrt{\dots} \frac{d}{dx} + \sqrt{\dots} = 0$$

Integrating on both sides, we get

$$\int \frac{d}{\sqrt{\dots}} + \int \frac{d}{\sqrt{\dots}} = c$$

$$\Rightarrow 2\sqrt{\dots} + 2\sqrt{\dots} = c$$

$$\Rightarrow \sqrt{\dots} + \sqrt{\dots} = \sqrt{a}, \text{ where } \sqrt{a} = \frac{c}{2}$$

$$51. \frac{d}{dx} + \frac{d}{dx} = 0$$

Integrating on both sides, we get

$$\log \dots + \log \dots = \log c$$

$$\Rightarrow \log(\dots) = \log c \Rightarrow \dots = c$$



$$52. \quad \frac{d}{d} - = 3$$

$$\Rightarrow \frac{d}{d} = 3 +$$

$$\Rightarrow \int \frac{1}{3+} d = \int 1 d$$

$$\Rightarrow \log| + 3| = \log| | + \log c$$

$$\Rightarrow + 3 = c$$

$$\Rightarrow = c - 3$$

This is the equation of family of straight line.

$$53. \quad \frac{d}{d} = \frac{(1+)}{(-1)}$$

$$\Rightarrow \frac{-1}{d} = \frac{(1+)}{d}$$

$$\Rightarrow \left(1 - \frac{1}{d}\right) d = \left(1 + \frac{1}{d}\right) d$$

Integrating on both sides, we get

$$+ \log = - \log + c$$

$$\Rightarrow - + \log = c$$

$$54. \quad (1 + \log) \frac{d}{d} - \log = 0$$

$$\Rightarrow (1 + \log) \frac{d}{d} = \log$$

$$\Rightarrow \left(\frac{1 + \log}{\log}\right) d = \frac{d}{\log}$$

Integrating on both sides, we get

$$\log(\log) + \log = \log + \log c$$

$$\Rightarrow \log(\log) = \log(c)$$

$$\Rightarrow \log = c$$

$$55. \quad - \frac{d}{d} = a \left(+ \frac{d}{d} \right)$$

$$\Rightarrow - a = a \frac{d}{d} + \frac{d}{d}$$

$$\Rightarrow (1 - a) = (a +) \frac{d}{d}$$

Integrating on both sides, we get

$$\int \frac{d}{a+} = \int \frac{d}{(1-a)} + \log c$$

$$\Rightarrow \int \frac{d}{a+} = \int \left[\frac{1}{1-a} + \frac{a}{1-a} \right] d + \log c$$

$$\Rightarrow \log(a +) = \log - \log(1 - a) + \log c$$

$$\Rightarrow \log[(a +)(1 - a)] = \log c$$

$$\Rightarrow (+ a)(1 - a) = c$$

$$56. \quad \frac{d}{d} + \sqrt{\frac{1-^2}{1-^2}} = 0$$

Integrating on both sides, we get

$$\int \frac{d}{\sqrt{1-^2}} + \int \frac{d}{\sqrt{1-^2}} = \sin^{-1} c$$

$$\Rightarrow \sin^{-1} + \sin^{-1} = \sin^{-1} c$$

$$\Rightarrow \sin^{-1} \left(\sqrt{1-^2} + \sqrt{1-^2} \right) = \sin^{-1} c$$

$$\Rightarrow \sqrt{1-^2} + \sqrt{1-^2} = c$$

$$57. \quad \frac{d}{d} + \sin\left(\frac{+}{2}\right) = \sin\left(\frac{-}{2}\right)$$

$$\Rightarrow \frac{d}{d} = \sin\left(\frac{-}{2}\right) - \sin\left(\frac{+}{2}\right)$$

$$\Rightarrow \frac{d}{d} = -2 \sin\left(\frac{-}{2}\right) \cdot \cos\left(\frac{-}{2}\right)$$

Integrating on both sides, we get

$$\int \operatorname{cosec}\left(\frac{-}{2}\right) d = - \int 2 \cos\left(\frac{-}{2}\right) d + c_1$$

$$\Rightarrow \frac{\log \tan\left(\frac{-}{4}\right)}{\frac{1}{2}} = - \frac{2 \sin\left(\frac{-}{2}\right)}{\frac{1}{2}} + c_1$$

$$\Rightarrow \log \tan\left(\frac{-}{4}\right) = c - 2 \sin\left(\frac{-}{2}\right), \text{ where } c = \frac{1}{2} c_1$$

$$58. \quad \frac{d}{d} = 1 + + ^2 + + + ^2$$

$$\Rightarrow \frac{d}{d} = (1 + + ^2)(+ 1)$$

Integrating on both sides, we get

$$\int \frac{d}{1+ + ^2} = \int (+ 1) d + c_1$$

$$\Rightarrow \int \frac{d}{\left(+\frac{1}{2}\right)^2 + \frac{3}{4}} = \frac{^2}{2} + + c_1$$

$$\Rightarrow \frac{1}{\sqrt{3}/2} \cdot \tan^{-1} \left(\frac{+\frac{1}{2}}{\sqrt{3}/2} \right) = \frac{^2}{2} + + c_1$$

$$\Rightarrow 4 \tan^{-1} \left(\frac{2+1}{\sqrt{3}} \right) = \sqrt{3} (^2 + 2) + c,$$

$$\text{where } c = 2\sqrt{3} c_1$$



$$59. \quad \frac{d}{d} = 3^+ \\ \Rightarrow 3^d - 3^{-d} = 0 \\ \text{Integrating on both sides, we get} \\ 3^d + 3^{-d} = c \\ \text{When } d = 0 = , \\ 3^0 + 3^0 = c \Rightarrow c = 2 \\ \therefore 3^d + 3^{-d} = 2 \\ \Rightarrow 3^d + 3^{-d} - 2 = 0$$

$$60. \quad d + 2 \frac{d}{d} = 0 \\ \Rightarrow \frac{d}{d} + 2 \frac{d}{d} = 0 \\ \Rightarrow \int \frac{d}{d} + 2 \int \frac{d}{d} = 0 \\ \Rightarrow \log d + 2 \log d = \log c \\ \Rightarrow \log d + \log d^2 = \log c \\ \Rightarrow \log d^3 = \log c \\ \Rightarrow d^3 = c \\ \text{Given that } d = 2 \text{ and } = 1 \\ \therefore (2)^3 = c \\ \Rightarrow c = 8 \\ \therefore d^3 = 8 \text{ is the particular solution.}$$

$$61. \quad 2 \frac{d}{d} - = 0 \\ \Rightarrow 2 \frac{d}{d} = d \\ \Rightarrow 2 \int \frac{1}{d} = \int \frac{1}{d} \\ \Rightarrow 2 \log d = \log d + \log c \\ \Rightarrow d^2 = c \\ \text{Since, } (1) = 2 \text{ i.e., } = 2 \text{ when } = 1 \\ \therefore 2^2 = 1 \times c \Rightarrow c = 4 \\ \therefore d^2 = 4 \\ \text{This represents the equation of parabola.}$$

$$62. \quad \frac{d}{e^d} = \\ \Rightarrow \frac{d}{d} = \log \\ \Rightarrow \int d = \int \log d \\ \Rightarrow = \log () - \int d + c \\ \Rightarrow = \log - + c \\ \text{Since, } (1) = 3 \text{ i.e., } = 3 \text{ when } = 1 \\ \therefore 3 = \log 1 - 1 + c \\ \Rightarrow c = 4 \\ \therefore = \log - + 4$$

$$63. \quad \log \frac{d}{d} = \\ \Rightarrow \frac{d}{d} = e \Rightarrow d = e^d \\ \text{Integrating on both sides, we get} \\ = e + c \\ \text{At } = 0 \text{ and } = 1, \\ 1 = e^0 + c \\ \Rightarrow c = 0 \\ \therefore = e$$

$$64. \quad \log \left(\frac{d}{d} \right) = 3 + 4 \\ \Rightarrow \frac{d}{d} = e^{3+4} \\ \Rightarrow e^{-4} d = e^3 d \\ \text{Integrating on both sides, we get} \\ \frac{e^{-4}}{-4} = \frac{e^3}{3} + c \\ \text{When } = 0 = , \\ -\frac{1}{4} = \frac{1}{3} + c \Rightarrow c = -\frac{7}{12} \\ \therefore \frac{e^{-4}}{-4} = \frac{e^3}{3} - \frac{7}{12} \\ \Rightarrow 4e^3 + 3e^{-4} - 7 = 0$$

$$65. \quad \frac{d}{d} = + 3 \\ \Rightarrow \frac{d}{+3} = d \\ \text{Integrating on both sides, we get} \\ \int \frac{d}{+3} = \int d + c \\ \Rightarrow \log(+3) = + c \quad \dots(i) \\ \text{Since, } (0) = 2 \text{ i.e., } = 2, \text{ when } = 0 \\ \therefore \log(2+3) = 0 + c \Rightarrow c = \log 5 \\ \therefore \log(+3) = + \log 5 \quad \dots[\text{From (i)}] \\ \Rightarrow + 3 = 5e \\ \Rightarrow = 5e - 3 \\ \therefore (\log 2) = 5e^{\log 2} - 3 = 10 - 3 \\ = 7$$

$$66. \quad \frac{d}{d} = 1 - \\ \int \frac{d}{1-} = \int d + c \\ \Rightarrow -\log(1 -) = + \log c \\ -\log(1 -) - \log c = - \\ \log(1 -)c = -$$



$$(1 -)c = e^{-} \quad \dots(i)$$

Since, $(0) = 3$ i.e., $= 3$, when $= 0$

$$\therefore -2c = e^0 \Rightarrow c = \frac{-1}{2}$$

$$\therefore (1 -)\left(\frac{-1}{2}\right) = e^{-} \quad \dots[\text{From (i)}]$$

$$\Rightarrow -1 = 2e^{-}$$

$$\Rightarrow = 2e^{-} + 1$$

$$\Rightarrow = 2e^{-\log e^8} + 1$$

$$\Rightarrow = 2 \times \frac{1}{8} + 1 = \frac{5}{4}$$

67. $(1 + \log) \frac{d}{d} - \log = 0$

$$\Rightarrow \frac{1 + \log}{\log} d = \frac{d}{d}$$

Integrating on both sides, we get

$$\log (\log) = \log + \log c$$

$$\Rightarrow \log (\log) = \log (c)$$

$$\Rightarrow \log \quad c \quad \dots(i)$$

When $= e$, $= e^2$

$$\therefore e \log e = e^2 c \Rightarrow c = \frac{1}{e}$$

$$\therefore \log = \frac{1}{e} \quad \dots[\text{From (i)}]$$

$$\Rightarrow = e \log$$

68. $\sin\left(\frac{d}{d}\right) = a$

$$\Rightarrow \frac{d}{d} = \sin^{-1} a \Rightarrow d = \sin^{-1} a \, d$$

Integrating on both sides, we get

$$= (\sin^{-1} a) + c \quad \dots(i)$$

Since, $(0) = 1$ i.e., $= 1$, when $= 0$

$$\therefore 1 = 0 + c \Rightarrow c = 1$$

$$\therefore = \sin^{-1} a + 1 \quad \dots[\text{From (i)}]$$

$$\Rightarrow \frac{-1}{d} = \sin^{-1} a$$

$$\Rightarrow \sin\left(\frac{-1}{d}\right) = a$$

69. $\left(\frac{2 + \sin}{1 +}\right) \frac{d}{d} = -\cos$

$$\Rightarrow \frac{d}{1 +} = \left(\frac{-\cos}{2 + \sin}\right) d$$

Integrating on both sides, we get

$$\int \frac{d}{1 +} + \int \frac{\cos}{2 + \sin} d = \log c$$

$$\Rightarrow \log(1 +) + \log(2 + \sin) = \log c$$

$$\Rightarrow (+ 1)(2 + \sin) = c \quad \dots(i)$$

Since, $(0) = 1$ i.e., $= 1$, when $= 0$

$$\therefore (1 + 1)(2 + \sin 0) = c \Rightarrow c = 4$$

$$\therefore (+ 1)(2 + \sin) = 4 \quad \dots[\text{From (i)}]$$

$$\Rightarrow = \frac{4}{2 + \sin} - 1$$

$$\therefore \left(\frac{\pi}{2}\right) = \frac{4}{2 + \sin \frac{\pi}{2}} - 1 = \frac{4}{3} - 1 = \frac{1}{3}$$

70. $(1 +) d = d$

$$\Rightarrow \frac{d - d}{2} = - d$$

$$\Rightarrow d \left(-\right) = - d$$

Integrating on both sides, we get

$$- = \frac{-^2}{2} + c \quad \dots(i)$$

Since, the curve passes through $(1, -1)$.

$$\therefore -1 = \frac{-1}{2} + c \Rightarrow c = \frac{-1}{2}$$

$$\therefore - = \frac{-^2}{2} - \frac{1}{2} \quad \dots[\text{From (i)}]$$

$$\Rightarrow = \frac{-2}{^2 + 1}$$

i.e., $f() = \frac{-2}{^2 + 1}$

$$\therefore f\left(-\frac{1}{2}\right) = \frac{4}{5}$$

71. $f() = \frac{-2}{1 + ^2}$

$$\therefore f\left(\frac{1}{2}\right) = \frac{-4}{5}$$

72. $\frac{d}{d} = +$

$$\Rightarrow \frac{1}{1 +} d = d$$

Integrating on both sides, we get

$$\int \frac{1}{1 +} d = \int d + c$$

$$\log(1 +) = \frac{^2}{2} + c \quad \dots(i)$$

Since, the required curve passes through $(0, 1)$.

$$\therefore c = \log 2$$



$$\therefore \log(1 + \frac{x}{2}) = \frac{x^2}{2} + \log 2 \quad \dots[\text{From (i)}]$$

$$\Rightarrow \log\left(\frac{1+x}{2}\right) = \frac{x^2}{2}$$

$$\Rightarrow \frac{1+x}{2} = 2e^{\frac{x^2}{2}} - 1$$

$$73. \frac{d}{dx} = \frac{x+1}{x-1} \quad \dots(\text{i})$$

$$\text{Put } x+1 = v \quad \dots(\text{ii})$$

$$\Rightarrow \frac{dv}{dx} = \frac{dv}{dx} - 1 \quad \dots(\text{iii})$$

Substituting (ii) and (iii) in (i), we get

$$\frac{dv}{dx} - 1 = \frac{v+1}{v-1}$$

$$\Rightarrow \frac{dv}{dx} = \frac{2v}{v-1} \Rightarrow \frac{v-1}{2v} dv = dx$$

Integrating on both sides, we get

$$\frac{v}{2} - \frac{1}{2} \log v = x + c_1$$

$$\Rightarrow v - \log v = 2x + 2c_1$$

$$\Rightarrow x - \log(x+1) = 2x + 2c_1$$

$$\Rightarrow x = -\log(x+1) + c, \text{ where } c = 2c_1$$

$$74. (x+1)^2 \frac{d}{dx} = a^2$$

$$\text{Put } x+1 = v \Rightarrow 1 + \frac{dv}{dx} = \frac{dv}{dx}$$

$$\therefore v^2 \left(\frac{dv}{dx} - 1\right) = a^2$$

$$\Rightarrow v^2 \frac{dv}{dx} - v^2 = a^2$$

$$\Rightarrow \frac{v^2}{v^2+a^2} dv = dx$$

$$\Rightarrow \int \frac{v^2}{v^2+a^2} dv = \int dx$$

$$\Rightarrow \int \frac{v^2+a^2-a^2}{v^2+a^2} dv = \int dx$$

$$\Rightarrow v - a \tan^{-1} \frac{v}{a} + c =$$

$$\Rightarrow x - a \tan^{-1} \left(\frac{x+1}{a}\right) + c =$$

$$\Rightarrow \tan^{-1} \left(\frac{x+1}{a}\right) = \frac{x+c}{a}$$

$$\Rightarrow \frac{x+1}{a} = \tan \left(\frac{x+c}{a}\right)$$

$$75. (x+1)dx + (2x^2-3)dx = 0$$

$$\Rightarrow \frac{d}{dx} = -\left(\frac{x+1}{2x^2-3}\right) \quad \dots(\text{i})$$

$$\text{Put } x+1 = v \quad \dots(\text{ii})$$

$$\Rightarrow 1 + \frac{dv}{dx} = \frac{dv}{dx}$$

$$\Rightarrow \frac{dv}{dx} = \frac{dv}{dx} - 1 \quad \dots(\text{iii})$$

Substituting (ii) and (iii) in (i), we get

$$\frac{dv}{dx} - 1 = -\left(\frac{v-1}{2v-3}\right)$$

$$\Rightarrow \frac{dv}{dx} = \frac{1-v}{2v-3} + 1 \Rightarrow \frac{dv}{dx} = \frac{v-2}{2v-3}$$

Integrating on both sides, we get

$$\int \frac{2v-3}{v-2} dv = \int dx + c$$

$$\Rightarrow \int \frac{2(v-2)+1}{v-2} dv = x + c$$

$$\Rightarrow 2v + \log(v-2) = x + c$$

$$\Rightarrow 2(x+1) + \log(x+1-2) = x + c$$

$$\Rightarrow 2x + 1 + \log(x-1) = c$$

$$76. \frac{d}{dx} = \frac{-2x+1}{2(x-2)} \quad \dots(\text{i})$$

$$\text{Put } -2x+1 = v \quad \dots(\text{ii})$$

$$\Rightarrow 1 - 2\frac{dv}{dx} = \frac{dv}{dx}$$

$$\Rightarrow \frac{dv}{dx} = \frac{1}{2} \left(1 - \frac{dv}{dx}\right) \quad \dots(\text{iii})$$

Substituting (ii) and (iii) in (i), we get

$$\frac{1}{2} \left(1 - \frac{dv}{dx}\right) = \frac{v+1}{2v}$$

$$\Rightarrow \frac{dv}{dx} = -\frac{1}{v}$$

Integrating on both sides, we get

$$\int v dv = -\int dx + c_1$$

$$\Rightarrow \frac{v^2}{2} = -x + c_1$$

$$\Rightarrow (-2x+1)^2 = -2x + 2c_1$$

$$\Rightarrow (-2x+1)^2 + 2x = c, \text{ where } c = 2c_1$$



$$77. \quad \frac{d}{d} = -\frac{-2 + 1}{2(-2) + 3} \quad \dots(i)$$

$$\text{Put } -2 = v \quad \dots(ii)$$

$$\Rightarrow 1 - \frac{2d}{d} = \frac{dv}{d}$$

$$\Rightarrow \frac{d}{d} = \frac{1}{2} \left(1 - \frac{dv}{d} \right) \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$\frac{1}{2} \left(1 - \frac{dv}{d} \right) = -\frac{v+1}{2v+3}$$

$$\Rightarrow \frac{2v+3}{4v+5} dv = d$$

Integrating on both sides, we get

$$\int \left[\frac{\frac{1}{2}(4v+5) + \frac{1}{2}}{4v+5} \right] dv = \int d + c_1$$

$$\Rightarrow \frac{1}{2}v + \frac{1}{2} \cdot \frac{1}{4} \log(4v+5) = + c_1$$

$$\Rightarrow \frac{1}{2}(-2) + \frac{1}{8} \log[4(-2)+5] = + c_1$$

$$\Rightarrow \log[4(-2)+5] = 8 - 4(-2) + 8c_1$$

$$\Rightarrow \log[4(-2)+5] = 4(+2) + c,$$

where $c = 8c_1$

$$78. \quad \frac{d}{d} = \frac{1}{\cos(+)} \quad \dots(i)$$

$$\text{Put } + = v \quad \dots(ii)$$

$$\Rightarrow 1 + \frac{d}{d} = \frac{dv}{d}$$

$$\Rightarrow \frac{d}{d} = \frac{dv}{d} - 1 \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$\frac{dv}{d} - 1 = \frac{1}{\cos v}$$

$$\Rightarrow \frac{dv}{d} = \frac{1}{\cos v} + 1 \Rightarrow \frac{\cos v}{1 + \cos v} dv = d$$

$$\Rightarrow \left[\frac{2\cos^2(v/2) - 1}{2\cos^2(v/2)} \right] dv = d$$

$$\Rightarrow \left[1 - \frac{1}{2}\sec^2(v/2) \right] dv = d$$

Integrating on both sides, we get

$$\int dv - \frac{1}{2} \int \sec^2(v/2) dv + c = \int d$$

$$\Rightarrow v - \tan\left(\frac{v}{2}\right) + c =$$

$$\Rightarrow + - \tan\left(\frac{+}{2}\right) + c =$$

$$\Rightarrow + c = \tan\left(\frac{+}{2}\right)$$

$$79. \quad \frac{d}{d} = \tan(-) + -$$

$$\text{Put } - = v \Rightarrow = v$$

$$\therefore \frac{dy}{d} = v + \frac{dv}{d}$$

$$\therefore v + \frac{dv}{d} = \tan v + v$$

$$\Rightarrow \frac{dv}{d} = \tan v \Rightarrow \int \frac{dv}{\tan v} = \int \frac{d}{d}$$

$$\Rightarrow \log |\sin v| = \log + \log c$$

$$\Rightarrow \log \sin(-) = \log c$$

$$\Rightarrow \sin(-) = c$$

$$80. \quad \frac{d}{d} = \frac{+}{-} \quad \dots(i)$$

$$\text{Put } = v \quad \dots(ii)$$

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = \frac{1+v}{1-v}$$

$$\Rightarrow \frac{dv}{d} = \frac{1+v^2}{1-v} \Rightarrow \frac{1-v}{1+v^2} dv = \frac{d}{d}$$

Integrating both sides, we get

$$\int \frac{1-v}{1+v^2} dv = \int \frac{d}{d}$$

$$\Rightarrow \int \frac{1}{1+v^2} dv - \frac{1}{2} \int \frac{2v}{1+v^2} = \log + c$$

$$\Rightarrow \tan^{-1} v - \frac{1}{2} \log(1+v^2) = \log + c$$

$$\Rightarrow \tan^{-1}(-) = \frac{1}{2} \log \frac{2+^2}{2} + \log + c$$

$$\Rightarrow \tan^{-1}(-) = \log \sqrt{\frac{2+^2}{2}} + \log + c$$

$$\Rightarrow \tan^{-1}(-) = \log \sqrt{2+^2} + c$$



$$81. \quad \frac{d}{d} = \frac{2}{-2} \quad \dots(i)$$

$$\text{Put } = v \quad \dots(ii)$$

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = \frac{v^2 - 2}{v - 2} = \frac{v^2}{v-1}$$

$$\Rightarrow \frac{dv}{d} = \frac{v^2}{v-1} - v \Rightarrow \frac{dv}{d} = \frac{v}{v-1}$$

Integrating on both sides, we get

$$\int \left(\frac{v-1}{v} \right) dv = \int \frac{d}{v-1} + \log k$$

$$\Rightarrow v - \log v = \log v + \log k$$

$$\Rightarrow v = \log(v.k)$$

$$\Rightarrow e^v = vk \Rightarrow e^{v/k} = k$$

$$82. \quad (x^2 + y^2)' = 2x$$

$$\therefore \frac{d}{d} = \frac{2x}{x^2 + y^2} \quad \dots(i)$$

$$\text{Put } = v \quad \dots(ii)$$

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = \frac{v^2 - 2}{x^2 + v^2}$$

$$\Rightarrow v + \frac{dv}{d} = \frac{v^2}{1+v}$$

$$\Rightarrow \frac{dv}{d} = \frac{v^2}{1+v} - v$$

$$\Rightarrow \frac{dv}{d} = \frac{-v}{1+v}$$

$$\Rightarrow \int -\frac{1+v}{v} dv = \int \frac{1}{d} - d$$

$$\Rightarrow -(\log v + v) = \log d + \log c$$

$$\Rightarrow -v = \log vc$$

$$\Rightarrow -v = \log c \Rightarrow e^{-v} = c$$

$$83. \quad (x^2 + y^2)d = 2xy$$

$$\Rightarrow \frac{d}{d} = \frac{1 + \left(\frac{y}{x}\right)^2}{2\left(\frac{y}{x}\right)} \quad \dots(i)$$

$$\text{Put } = v \quad \dots(ii)$$

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = \frac{1+v^2}{2v}$$

$$\Rightarrow \frac{dv}{d} = \frac{1+v^2}{2v} - v$$

$$\Rightarrow \frac{dv}{d} = \frac{1-v^2}{2v}$$

Integrating on both sides, we get

$$\int \frac{d}{d} - \int \frac{2v}{1-v^2} dv = c_1$$

$$\Rightarrow \log d + \log(1-v^2) = c_1$$

$$\Rightarrow (1-v^2) \cdot d = e^{c_1}$$

$$\Rightarrow \left(1 - \frac{y^2}{x^2}\right) \cdot d = e^{c_1}$$

$$\Rightarrow x^2 - y^2 = e^{c_1}$$

$$\Rightarrow c \cdot (x^2 - y^2) = e^{c_1}, \text{ where } c = \frac{1}{e^{c_1}}$$

$$84. \quad \frac{d}{d} = \frac{-y}{x} \quad \dots(i)$$

$$\text{Put } = v \quad \dots(ii)$$

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = \frac{v - \frac{1}{v}}{v + \frac{1}{v}} = \frac{v-1}{v+1}$$

$$\Rightarrow \frac{dv}{d} = \frac{v-1}{v+1} - v \Rightarrow \frac{dv}{d} = \frac{-(v^2+1)}{v+1}$$

Integrating on both sides, we get

$$\int \frac{v+1}{v^2+1} dv = -\int \frac{d}{v+1} + c_1$$

$$\Rightarrow \frac{1}{2} \int \frac{2v}{v^2+1} dv + \int \frac{1}{v^2+1} dv = -\int \frac{d}{v+1} + c_1$$

$$\Rightarrow \frac{1}{2} \log(v^2+1) + \tan^{-1} v = -\log d + c_1$$

$$\Rightarrow \log \left(\frac{x^2 + y^2}{2} \right) + 2 \tan^{-1} \left(\frac{y}{x} \right) = -2 \log d + 2c_1$$

$$\Rightarrow \log(x^2 + y^2) - 2 \log d + 2 \tan^{-1} \left(\frac{y}{x} \right) = -2 \log d + 2c_1$$

$$\Rightarrow \log(x^2 + y^2) + 2 \tan^{-1} \left(\frac{y}{x} \right) = c, \text{ where } c = 2c_1$$



$$85. \quad d - d = \sqrt{2 + 2} d$$

$$\Rightarrow d = \sqrt{2 + 2} d + d$$

$$\Rightarrow \frac{d}{d} = \frac{\sqrt{2 + 2} + 1}{d} \quad \dots(i)$$

Put $v = \frac{d}{d}$ (ii)

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = \frac{\sqrt{2 + v^2} + v}{d}$$

$$\Rightarrow v + \frac{dv}{d} = \sqrt{1 + v^2} + v$$

$$\Rightarrow \frac{dv}{d} = \sqrt{1 + v^2}$$

Integrating on both sides, we get

$$\int \frac{dv}{\sqrt{1 + v^2}} = \int \frac{d}{d} + \log c$$

$$\Rightarrow \log(v + \sqrt{1 + v^2}) = \log d + \log c$$

$$\Rightarrow v + \sqrt{1 + v^2} = c$$

$$\Rightarrow - + \sqrt{1 + \frac{2}{2}} = c$$

$$\Rightarrow + \sqrt{2 + 2} = 2c$$

86. Putting $v = \frac{d}{d}$ and $\frac{d}{d} = v + \frac{dv}{d}$ in the given equation, we get

$$v \left(v + \frac{dv}{d} \right) = \left[v^2 + \frac{\phi(v^2)}{\phi'(v^2)} \right]$$

Integrating on both sides, we get

$$\int \frac{v\phi'(v^2)}{\phi(v^2)} dv = \int \frac{d}{d} + \log c_1$$

$$\Rightarrow \frac{1}{2} \log [\phi(v^2)] = \log d + \log c_1$$

$$\Rightarrow \log \left[\phi \left(\frac{2}{2} \right) \right] = 2 \log d + 2 \log c_1$$

$$\Rightarrow \log \left[\phi \left(\frac{2}{2} \right) \right] = \log 2c_1^2$$

$$\Rightarrow \phi \left(\frac{2}{2} \right) = c^2, \text{ where } c = c_1^2$$

$$87. \quad \left(1 + e^{-x} \right) d + e^{-x} \left(1 - \frac{d}{d} \right) d = 0$$

$$\Rightarrow \frac{d}{d} = \frac{-e^{-x} \left(1 - \frac{d}{d} \right)}{1 + e^{-x}} \quad \dots(i)$$

Put $u = 1 + e^{-x}$ (ii)

$$\Rightarrow \frac{d}{d} = u + \frac{du}{d} \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$u + \frac{du}{d} = \frac{-e^{-x} (1 - u)}{1 + e^{-x}}$$

$$\therefore \frac{du}{d} = \frac{-e^{-x} - u}{1 + e^{-x}}$$

$$\therefore \frac{1 + e^{-x}}{u + e^{-x}} du = -\frac{d}{d}$$

Integrating on both sides, we get

$$\int \frac{1 + e^{-x}}{u + e^{-x}} du = -\int \frac{d}{d}$$

$$\therefore \log |u + e^{-x}| = -\log |d| + \log |c|$$

$$\therefore \log \left| 1 + e^{-x} \right| = \log \left| \frac{c}{d} \right|$$

$$\therefore 1 + e^{-x} = \frac{c}{d}$$

$$\therefore e^{-x} + 1 = c$$

$$88. \quad \cos \left(\frac{d}{d} - \frac{d}{d} \right) + \sin \left(\frac{d}{d} + \frac{d}{d} \right) = 0$$

$$\Rightarrow -\cos \left(\frac{d}{d} - \frac{d}{d} \right) + \sin \left(\frac{d}{d} + \frac{d}{d} \right) = 0$$

....(i)

Put $v = \frac{d}{d}$ (ii)

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \quad \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v \cos v \left(v + \frac{dv}{d} - v \right) + \sin v \left(v + \frac{dv}{d} + v \right) = 0$$

$$\Rightarrow (v \cos v) \frac{dv}{d} + \sin v \left(2v + \frac{dv}{d} \right) = 0$$

$$\Rightarrow (v \cos v + \sin v) \frac{dv}{d} + 2v \sin v = 0$$

Integrating on both sides, we get

$$\int \frac{v \cos v + \sin v}{v \sin v} dv = -2 \int \frac{d}{d} + \log c$$



$$\Rightarrow \log(v \sin v) = -2 \log \dots + \log c$$

$$\Rightarrow \log(v \sin v) = \log \frac{c}{2}$$

$$\Rightarrow v \sin v = \frac{c}{2}$$

$$\Rightarrow -\sin \dots = -\frac{c}{2} \Rightarrow \sin \dots = \frac{c}{2}$$

Since, $(1) = \frac{\pi}{2}$, i.e., $\dots = \frac{\pi}{2}$, when $\dots = 1$

$$\therefore \frac{\pi}{2} \sin \frac{\pi}{2} = c \Rightarrow c = \frac{\pi}{2}$$

$$\therefore \sin \dots = \frac{\pi}{2}$$

$$89. \frac{d}{d} = \frac{2 + 2}{2} \dots(i)$$

$$\text{Put } \dots = v \dots(ii)$$

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \dots(iii)$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = \frac{2 + v^2 + 2}{2v} = \frac{1 + v^2}{2v}$$

$$\Rightarrow \frac{dv}{d} = \frac{1 + v^2}{2v} - v \Rightarrow \frac{dv}{d} = \frac{1 - v^2}{2v}$$

Integrating on both sides, we get

$$\int \frac{d}{d} - \int \frac{2v}{1 - v^2} dv = \log c$$

$$\Rightarrow \log \dots + \log(1 - v^2) = \log c$$

$$\Rightarrow \log \dots + \log\left(1 - \frac{2}{2}\right) = \log c$$

$$\Rightarrow \left(1 - \frac{2}{2}\right) = c$$

$$\Rightarrow 2 - 2 = c \dots(iv)$$

Since, the required curve passes through (2, 1).

$$\therefore 4 - 1 = 2c \Rightarrow c = \frac{3}{2}$$

$$\therefore 2 - 2 = \frac{3}{2} \dots[\text{From (iv)}]$$

$$\Rightarrow 2(2 - 2) = 3$$

$$90. \frac{d}{d} \dots = 3 - 3$$

$$\therefore \text{I. F.} = e^{\int -1 d} = e^{-\log} = e^{\log^{-1}} = \frac{1}{\dots}$$

$$91. \frac{d}{d} + 2 = 2 \Rightarrow \frac{d}{d} + \frac{2}{\dots} =$$

$$\therefore \text{I.F.} = e^{\int 2 d} = e^{2 \log} = e^{\log^2} = 2$$

$$92. \frac{d}{d} + \frac{\dots}{1 + \dots^2} = \frac{e^{\tan^{-1}}}{1 + \dots^2}$$

$$\therefore \text{I.F.} = e^{\int \frac{1}{1 + \dots^2} d} = e^{\tan^{-1}}$$

$$93. \cos \frac{d}{d} + \sin \dots = 1$$

$$\Rightarrow \frac{d}{d} + \tan \dots = \sec$$

$$\therefore \text{I.F.} = e^{\int \tan d} = e^{\log \sec} = \sec$$

$$94. \frac{d}{d} + \dots = \frac{1 + \dots}{\dots}$$

$$\Rightarrow \frac{d}{d} + \dots = \frac{1}{\dots} + \dots$$

$$\Rightarrow \frac{d}{d} + \left(1 - \frac{1}{\dots}\right) = \frac{1}{\dots}$$

$$\therefore \text{I.F.} = e^{\int (1 - \frac{1}{\dots}) d} = e^{-\log} = \frac{e}{\dots}$$

$$95. \frac{d}{d} = \frac{1}{\dots + \dots + 2}$$

$$\Rightarrow \frac{d}{d} = \dots + 2 \Rightarrow \frac{d}{d} - \dots = \dots + 2$$

$$\therefore \text{I.F.} = e^{\int -d} = e^{-}$$

$$96. \text{I. F.} = \sin$$

$$\therefore e^{\int Pd} = \sin$$

$$\Rightarrow \int Pd = \log(\sin \dots) \Rightarrow P = \frac{d}{d} [\log(\sin \dots)]$$

$$\Rightarrow P = \frac{1}{\sin} \times \cos = \cot$$

$$97. \frac{d}{d} + P(\dots) = 0$$

Here, $Q = 0$

$$\therefore \text{I.F.} = e^{\int Pd}$$

\therefore solution of the given equation is

$$(\text{I.F.}) = \int Q(\text{I.F.}) d + c$$

$$\Rightarrow e^{\int Pd} = 0 + c \Rightarrow \dots = c e^{-\int Pd}$$



98. I.F. = $e^{\int ad} = e^a$
 \therefore solution of the given equation is

$$.e^a = \int e^m .e^a d + c = \frac{e^{(a+m)}}{a+m} + c$$

$$\Rightarrow = \frac{e^m}{a+m} + ce^{-a}$$

$$\Rightarrow (a+m) = e^m + c(a+m) e^{-a}$$
99. $\frac{d}{d} + = 1$
 I.F. = $e^{\int d} = e$
 \therefore Solution of the differential equation is

$$e = \int e d + c_1$$

$$\Rightarrow e = e + c_1 \Rightarrow e(1 -) = -c_1$$

$$\Rightarrow \log e + \log(1 -) = -\log c_1$$

$$\Rightarrow + c = -\log(1 -)$$

$$\Rightarrow \log \left| \frac{1}{1 - } \right| = + c$$
100. $\frac{d}{d} = - +$

$$\Rightarrow \frac{d}{d} - - =$$

 \therefore I.F. = $e^{\int -1 d} = e^{-\log} = \frac{1}{}$
 \therefore solution of the given equation is

$$- = \int . \frac{1}{d} + a$$

$$\Rightarrow - = + a \Rightarrow = ^2 + a$$
101. $\frac{d}{d} + 2 = ^2$

$$\Rightarrow \frac{d}{d} + \left(\frac{2}{} \right) =$$

 \therefore I.F. = $e^{2 \int \frac{1}{d}} = e^{2 \log} = ^2$
 \therefore solution of the given equation is

$$^2 = \int ^3 d + c_1$$

$$\Rightarrow ^2 = \frac{^4}{4} + c_1 = \frac{^4 + C}{4}, \text{ where } C = 4 c_1$$

$$\Rightarrow = \frac{^4 + C}{4^2}$$

102. $+ ^2 = \frac{d}{d} \Rightarrow \frac{d}{d} - = ^2$
 \therefore I.F. = $e^{\int -d} = e^{-}$
 \therefore solution of the given equation is

$$e^{-} = \int e^{-} . ^2 d + c$$

$$\Rightarrow .e^{-} = - ^2 .e^{-} - 2 e^{-} - 2e^{-} + c$$

$$\Rightarrow + ^2 + 2 + 2 = c.e$$
103. $(1 + ^2) + \left(-e^{\tan^{-1}} \right) \frac{d}{d} = 0$

$$\Rightarrow \frac{d}{d} + \frac{^2}{1 + ^2} = \frac{e^{\tan^{-1}}}{1 + ^2}$$

 \therefore I.F. = $e^{\int \frac{1}{1 + ^2} d} = e^{\tan^{-1}}$
 \therefore solution of the given equation is

$$.e^{\tan^{-1}} = \int \frac{e^{\tan^{-1}}}{1 + ^2} .e^{\tan^{-1}} d + c_1$$

$$\Rightarrow e^{\tan^{-1}} = \int \frac{e^{2 \tan^{-1}}}{1 + ^2} d + c_1$$

$$\Rightarrow e^{\tan^{-1}} = \frac{e^{2 \tan^{-1}}}{2} + c_1$$

$$\Rightarrow 2 e^{\tan^{-1}} = e^{2 \tan^{-1}} + c, \text{ where } c = 2c_1$$
104. $(- 3 ^2) d + d = 0$

$$\Rightarrow (- 3 ^2) = - \frac{d}{d} \Rightarrow \frac{d}{d} = - + 3$$

$$\Rightarrow \frac{d}{d} + \frac{1}{()} = 3$$

 \therefore I.F. = $e^{\int \frac{1}{d}} = e^{\log} =$
 \therefore Solution of the given differential equation is

$$= \int 3 ^2 + c$$

$$\Rightarrow = \frac{3^3}{3} + c$$

$$\Rightarrow = ^2 + \frac{c}{}$$
105. $(- 4 ^3) \frac{d}{d} - = 0$

$$\Rightarrow \frac{d}{d} = \frac{-}{-4^3}$$

$$\Rightarrow \frac{d}{d} = \frac{-4^3}{-} \Rightarrow \frac{d}{d} - - = -4^2$$

 \therefore I.F. = $e^{\int -\frac{1}{d}} = e^{-\log} = \frac{1}{}$



∴ Solution of the given equation is

$$\frac{1}{x} = \int -4x^{-2} \cdot \frac{1}{x} dx + c$$

$$\Rightarrow -\frac{1}{x} = -2x^{-2} + c \Rightarrow \frac{1}{x^2} = c - \frac{1}{x^2}$$

106. $\frac{d}{dx} + \left(\frac{1}{x^2}\right) \frac{d}{dx} = 0$

$$\Rightarrow \frac{dv}{dx} + \frac{1}{x^2} v = 0$$

$$\Rightarrow -\frac{1}{2} \cdot \frac{dv}{dx} + \left(\frac{-1}{x^2}\right) \cdot \frac{1}{x} = 1 \quad \dots(i)$$

Put $v = \frac{1}{x}$

$$\Rightarrow \frac{dv}{dx} = -\frac{1}{x^2} \cdot \frac{d}{dx}$$

∴ $\frac{dv}{dx} + \left(\frac{-1}{x^2}\right) \cdot v = 1 \quad \dots[\text{From (i)}]$

∴ I.F. = $e^{\int -\frac{1}{x^2} dx} = e^{-\log x} = \frac{1}{x}$

∴ solution of the given equation is

$$v \cdot \frac{1}{x} = \int \frac{1}{x^2} dx + c_1$$

$$\Rightarrow \frac{1}{x^2} = \log x + c_1$$

$$\Rightarrow -\frac{1}{x} + \log x = -c_1$$

$$\Rightarrow -\frac{1}{x} + \log x = c, \text{ where } c = -c_1$$

107. $\frac{d}{dx} + \frac{1}{x} = x^{-2}$

$$\Rightarrow \frac{1}{2} \cdot \frac{dv}{dx} + \frac{1}{x} \left(\frac{1}{x}\right) = 1 \quad \dots(i)$$

Put $v = \frac{1}{x}$

$$\Rightarrow \frac{dv}{dx} = -\frac{1}{x^2} \cdot \frac{d}{dx}$$

∴ $-\frac{dv}{dx} + \frac{1}{x} \cdot v = 1 \quad \dots[\text{From (i)}]$

$$\Rightarrow \frac{dv}{dx} + \left(\frac{-1}{x}\right) \cdot v = -1$$

∴ I.F. = $e^{\int -\frac{1}{x} dx} = e^{-\log x} = \frac{1}{x}$

∴ solution of the given equation is

$$v \cdot \left(\frac{1}{x}\right) = \int (-1) \cdot \frac{1}{x} dx + c$$

$$\Rightarrow \frac{1}{x^2} = -\log x + c \Rightarrow \frac{1}{x^2} = c - \log x$$

108. $\cos x \frac{d}{dx} = (\sin x - \cos x) \frac{d}{dx}$

$$\Rightarrow \frac{dv}{dx} = \tan x - \sec x$$

$$\Rightarrow \frac{1}{2} \cdot \frac{dv}{dx} + \tan x \left(\frac{-1}{2}\right) = -\sec x \quad \dots(i)$$

Put $v = \frac{1}{2}$ $\Rightarrow \frac{dv}{dx} = \frac{1}{2} \cdot \frac{d}{dx}$

∴ $\frac{dv}{dx} + (\tan x) v = -\sec x \quad \dots[\text{From (i)}]$

∴ I.F. = $e^{\int \tan x dx} = e^{\log(\sec x)} = \sec x$

∴ solution of the given equation is

$$v \cdot \sec x = \int -\sec x \cdot \sec x dx + c_1$$

$$\Rightarrow v \sec x = -\tan x + c_1$$

$$\Rightarrow \frac{1}{2} \sec x = -\tan x + c_1$$

$$\Rightarrow \sec x = (\tan x + c), \text{ where } c = -2c_1$$

109. $\left(\frac{d}{dx} + \frac{1}{x^4}\right) \frac{d}{dx} - \frac{d}{dx} = 0$

$$\Rightarrow \frac{dv}{dx} = \frac{1}{x^4} + \frac{d}{dx}$$

$$\Rightarrow \frac{dv}{dx} - \frac{d}{dx} = \frac{1}{x^4}$$

$$\Rightarrow \frac{1}{3} \cdot \frac{dv}{dx} - \frac{d}{dx} = 1 \quad \dots(i)$$

Put $v = \frac{1}{3}$

$$\Rightarrow \frac{dv}{dx} = -3 \cdot \frac{d}{dx}$$

∴ $-\frac{1}{3} \cdot \frac{dv}{dx} - \frac{v}{3} = 1 \quad \dots[\text{From (i)}]$

$$\Rightarrow \frac{dv}{dx} + \frac{3}{3} \cdot v = -3$$

∴ I.F. = $e^{\int \frac{3}{3} dx} = e^{3 \log x} = x^3$

∴ solution of the given equation is

$$v \cdot x^3 = \int -3 \cdot x^3 dx + c_1$$

$$\Rightarrow v \cdot x^3 = \frac{-3x^4}{4} + c_1 \Rightarrow \frac{x^3}{3} = -\frac{3}{4} x^4 + c_1$$

$$\Rightarrow 4x^3 + 3x^4 = 4c_1$$

$$\Rightarrow 4x^3 + 3x^4 = c, \text{ where } c = 4c_1$$



110. $\frac{d}{d} + \tan = \sec$

I.F. = $e^{\int \tan d} = e^{\log \sec} = \sec$

∴ Solution of the given differential equation is

$\sec = \int \sec^2 + c$

$\Rightarrow \sec = \tan + c$

$(0) = 0 \Rightarrow c = 0$

∴ $\sec = \tan$

111. I.F. = $e^{\int \frac{d}{\log_e} = e^{\log(\log_e)} = \log_e$

∴ solution of the given equation is

$\log_e = \int \frac{\log_e}{\log_e} d + c$

$\Rightarrow \log_e = \frac{(\log_e)^2}{2} + c$

When $= e, = 1$

∴ $1 = \frac{1}{2} + c \Rightarrow c = \frac{1}{2}$

∴ $\log_e = \frac{(\log_e)^2}{2} + \frac{1}{2}$

$\Rightarrow 2 = \log_e + \frac{1}{\log_e}$

112. Given, $(\log) \frac{d}{d} + = 2 \log$

When $= 1, = 0$

$(\log) \frac{d}{d} + = 2 \log$

∴ $\frac{d}{d} + \frac{1}{\log} = 2$

∴ I.F. = $e^{\int \frac{1}{\log} d} = e^{\log(\log)} = \log$

∴ solution of the given equation is

$\log = \int 2 \log d + c$

∴ $\log = 2(\log -) + c$

When $= 1, = 0$

$0 = -2 + c \Rightarrow c = 2$

∴ $\log = 2(\log -) + 2$

∴ $(e) = 2(e - e) + 2 = 2$

113. $^2d + \left(-\frac{1}{2}\right)d = 0$

$\Rightarrow \frac{d}{d} + \frac{1}{2} = \frac{1}{3}$

∴ I.F. = $e^{\int \frac{1}{2} d} = e^{\frac{1}{2}d}$

∴ solution of the given equation is

$\cdot e^{-\frac{1}{3}} = \int \frac{1}{3} \cdot e^{-\frac{1}{3}} d + c$

$\Rightarrow e^{-\frac{1}{3}} = e^{-\frac{1}{3}} \left(\frac{1}{3} + 1\right) + c \dots(i)$

Since, $(1) = 1$ i.e., $= 1$, when $= 1$

∴ $1 \cdot e^{-1} = e^{-1}(1 + 1) + c \Rightarrow c = -e^{-1}$

∴ $e^{-\frac{1}{3}} = e^{-\frac{1}{3}} \left(\frac{1}{3} + 1\right) - e^{-1} \dots[\text{From (i)}]$

$\Rightarrow = 1 + \frac{1}{3} - \frac{e^{1/3}}{e}$

114. $(1 + ^2) \frac{d}{d} + 2 - 4^2 = 0$

$\Rightarrow \frac{d}{d} + \left(\frac{2}{1 + ^2}\right) = \frac{4^2}{1 + ^2}$

∴ I.F. = $e^{\int \frac{2}{1 + ^2} d} = e^{\log(1 + ^2)} = 1 + ^2$

$\cdot (\text{I.F.}) = \int Q (\text{I.F.}) d + c$

$\Rightarrow (1 + ^2) = \int \frac{4^2}{1 + ^2} \times (1 + ^2) d + c$

$\Rightarrow (1 + ^2) = \frac{4}{3} ^3 + c$

Since, $(0) = -1$, i.e., when $= 0, = -1$

∴ $c = -1$

$\Rightarrow (1 + ^2) = \frac{4}{3} ^3 - 1$

$\Rightarrow = \frac{4^3}{3(1 + ^2)} - \frac{1}{1 + ^2}$

∴ $(1) = \frac{4}{6} - \frac{1}{2} = \frac{1}{6}$

115. $(1 + t) \frac{d}{dt} - t = 1 \Rightarrow \frac{d}{dt} - \frac{t}{1 + t} = \frac{1}{1 + t}$

∴ I.F. = $e^{\int \frac{-t}{1 + t} dt} = e^{-\int \frac{1 + t - 1}{1 + t} dt}$

$= e^{-\int \left(1 - \frac{1}{1 + t}\right) dt} = e^{-t + \log(1 + t)} = (1 + t) \cdot e^{-t}$

∴ solution of the given equation is

$\cdot (1 + t) \cdot e^{-t} = \int (1 + t) \cdot e^{-t} \cdot \frac{1}{1 + t} dt + c$

$= \int e^{-t} dt + c$

∴ $(1 + t) e^{-t} = -e^{-t} + c$

$\Rightarrow (1 + t) = -1 + c \cdot e^t \dots(i)$

Since, $(0) = -1$ i.e., $= -1$, when $t = 0$

∴ $-1(1 + 0) = -1 + c \cdot e^0 \Rightarrow c = 0$



$$\therefore (1+t) = -1 \quad \dots[\text{From (i)}]$$

$$\Rightarrow = \frac{-1}{1+t}$$

$$\therefore (1) = \frac{-1}{1+1} = \frac{-1}{2}$$

$$116. \frac{d}{dx} = \cos(2 - \operatorname{cosec} x)$$

$$\Rightarrow \frac{d}{dx} = 2 \cos x - \cot x$$

$$\Rightarrow \frac{d}{dx} + \cot x = 2 \cos x$$

$$\therefore \text{I.F.} = e^{\int \cot x dx} = e^{\log \sin x} = \sin x$$

\(\therefore\) solution of the given equation is

$$\sin x \cdot \frac{d}{dx} = \int 2 \cos x \cdot \sin x dx + c_1$$

$$\begin{aligned} \therefore \sin x \cdot \frac{d}{dx} &= \int \sin 2x dx + c_1 = -\frac{\cos 2x}{2} + c_1 \\ &= \frac{2 \sin^2 x - 1}{2} + c_1 = \sin^2 x - \frac{1}{2} + c_1 \end{aligned}$$

$$\therefore \sin x \cdot \frac{d}{dx} = \sin^2 x + c, \text{ where } c = c_1 - \frac{1}{2}$$

$$\text{When } x = \frac{\pi}{2}, \frac{d}{dx} = 2$$

$$\therefore 2 \sin \frac{\pi}{2} = \sin^2 \frac{\pi}{2} + c$$

$$\Rightarrow 2 = 1 + c \Rightarrow c = 1$$

$$\therefore \sin x \cdot \frac{d}{dx} = \sin^2 x + 1 \Rightarrow \frac{d}{dx} = \sin x + \operatorname{cosec} x$$

$$117. \frac{d}{dx} (g(x)) + g'(x) (g(x)) = g(x) g'(x)$$

$$\therefore \text{I.F.} = e^{\int g'(x) dx} = e^{g(x)}$$

\(\therefore\) solution of the given equation is

$$(g(x)) \cdot e^{g(x)} = \int g(x) g'(x) \cdot e^{g(x)} dx + c$$

$$\Rightarrow (g(x)) \cdot e^{g(x)} = e^{g(x)} [g(x) - 1] + c \quad \dots(\text{i})$$

Putting $x = 0$ in (i), we get

$$0 = e^0(0 - 1) + c \quad \dots[\because (0) = 0, g(0) = 0 \text{ (given)}]$$

$$\Rightarrow c = 1$$

$$\therefore (g(x)) \cdot e^{g(x)} = e^{g(x)} [g(x) - 1] + 1 \quad \dots[\text{From (i)}]$$

Putting $x = 2$, we get

$$(2) e^0 = e^0(0 - 1) + 1 \quad \dots[\because g(2) = 0 \text{ (given)}]$$

$$\Rightarrow (2) = 0$$

$$118. \sin \frac{d}{dx} + \cos x = 4$$

$$\Rightarrow \frac{d}{dx} + \cot x = \frac{4}{\sin x}$$

$$\text{I.F.} = e^{\int \cot x dx} = e^{\log \sin x} = \sin x$$

General solution of the given equation is

$$\sin x \cdot \frac{d}{dx} = \int \sin x \cdot \frac{4}{\sin x} dx + c$$

$$\Rightarrow \sin x \cdot \frac{d}{dx} = 2x^2 + c \quad \dots(\text{i})$$

$$\left(\frac{\pi}{2}\right) = 0 \text{ i.e. } = 0 \text{ when } x = \frac{\pi}{2}$$

$$0 = 2 \cdot \frac{\pi^2}{4} + c \Rightarrow c = -\frac{\pi^2}{2}$$

$$\sin x \cdot \frac{d}{dx} = 2x^2 - \frac{\pi^2}{2} \quad \dots[\text{From (i)}]$$

$$\text{When } x = \frac{\pi}{6},$$

$$\frac{1}{2} = 2 \cdot \frac{\pi^2}{36} - \frac{\pi^2}{2}$$

$$\Rightarrow \left(\frac{\pi}{6}\right) = \frac{-8}{9} \pi^2$$

$$119. \frac{d}{dx} - \tan x = 2 \sec x$$

$$\Rightarrow \frac{d}{dx} + (-\tan x) = 2 \sec x$$

$$\therefore \text{I.F.} = e^{-\int \tan x dx} = e^{\log \cos x} = \cos x$$

\(\therefore\) solution of the given equation is

$$\cos x \cdot \frac{d}{dx} = \int 2 \sec x \cdot \cos x dx + c$$

$$\Rightarrow \cos x \cdot \frac{d}{dx} = 2x + c \quad \dots(\text{i})$$

Since, $(0) = 0$ i.e., $x = 0$, when $x = 0$

$$\therefore 0 = 0 + c \Rightarrow c = 0$$

$$\therefore \cos x \cdot \frac{d}{dx} = 2x \quad \dots[\text{From (i)}]$$

$$\Rightarrow \frac{d}{dx} = 2x \sec x \quad \dots(\text{ii})$$

$$\Rightarrow \frac{d}{dx} = 2x \sec x \tan x + 2 \sec x \quad \dots(\text{iii})$$

Putting $x = \frac{\pi}{4}$ in (ii) and (iii), we get

$$\left(\frac{\pi}{4}\right) = \frac{\pi^2}{8\sqrt{2}} \text{ and } \left(\frac{\pi}{4}\right) = \frac{\pi^2}{8\sqrt{2}} + \frac{\pi}{\sqrt{2}}$$

Putting $x = \frac{\pi}{3}$ in (ii) and (iii), we get

$$\left(\frac{\pi}{3}\right) = \frac{2\pi^2}{9} \text{ and } \left(\frac{\pi}{3}\right) = \frac{2\pi^2}{3\sqrt{3}} + \frac{4\pi}{3}$$



120. 'p' is the population at time 't'.

$$\therefore \frac{dp}{dt} = \frac{3}{100} p$$

$$\Rightarrow \frac{dp}{p} = \frac{3}{100} dt$$

Integrating on both sides, we get

$$\int \frac{dp}{p} = \frac{3}{100} \int dt$$

$$\Rightarrow \log p = \frac{3}{100} t + c_1$$

$$\Rightarrow p = e^{\frac{3}{100}t + c_1} \Rightarrow p = e^{\frac{3}{100}t} e^{c_1}$$

$$\Rightarrow p = c e^{\frac{3}{100}t} \quad \dots[\text{where } e^{c_1} = c]$$

121. $\frac{dV}{dt} = -k(T - t)$

$$\Rightarrow dV = -k(T - t)dt$$

Integrating on both sides, we get

$$\int dV = -k \int (T - t)dt + c$$

$$\Rightarrow V(t) = \frac{k(T - t)^2}{2} + c \quad \dots(i)$$

Initially i.e., when $t = 0$, $V(t) = I$

$$\therefore I = \frac{kT^2}{2} + c \Rightarrow c = I - \frac{kT^2}{2}$$

$$\therefore V(t) = \frac{k(T - t)^2}{2} + I - \frac{kT^2}{2} \quad \dots[\text{From (i)}]$$

When $t = T$,

$$V(T) = I - \frac{kT^2}{2}$$

122. $\frac{d(p(t))}{dt} = \frac{1}{2}p(t) - 200$

Integrating on both sides, we get

$$\int \frac{d(p(t))}{\frac{1}{2}p(t) - 200} = \int dt + c_1$$

$$\Rightarrow 2 \log \left(\frac{p(t)}{2} - 200 \right) = t + c_1$$

$$\Rightarrow \frac{p(t)}{2} - 200 = e^{\frac{t}{2}} \cdot c, \quad \left(\text{where } c = e^{\frac{c_1}{2}} \right) \dots(i)$$

Putting $t = 0$, we get

$$\frac{p(0)}{2} - 200 = e^0 \cdot c$$

$$\Rightarrow \frac{100}{2} - 200 = c \Rightarrow c = -150$$

$$\therefore \frac{p(t)}{2} - 200 = e^{\frac{t}{2}} (-150) \quad \dots[\text{From (i)}]$$

$$\Rightarrow p(t) = 400 - 300e^{\frac{t}{2}}$$



Evaluation Test

1. The given equation is

$$\sqrt{1 - a^4} + \sqrt{1 - a^4} = a(1 - a^2)$$

$$\text{Put } 1 - a^2 = \sin \alpha, \quad 1 - a^2 = \sin \beta$$

The equation becomes

$$\cos \alpha + \cos \beta = a(\sin \alpha - \sin \beta)$$

$$2 \cos \left(\frac{\alpha + \beta}{2} \right) \cos \left(\frac{\alpha - \beta}{2} \right) = 2a \sin \left(\frac{\alpha - \beta}{2} \right) \cos \left(\frac{\alpha + \beta}{2} \right)$$

$$\therefore \cot \left(\frac{\alpha - \beta}{2} \right) = a$$

$$\therefore \alpha - \beta = 2 \cot^{-1} a$$

$$\therefore \sin^{-1} 1 - \sin^{-1} 1 = 2 \cot^{-1} a$$

Differentiating w.r.t. a , we get

$$\frac{1}{\sqrt{1 - 1^2}} \cdot 2 - \frac{1}{\sqrt{1 - 1^2}} \cdot 2 \frac{d}{da} = 0$$

$$\therefore \frac{d}{da} = -\sqrt{\frac{1 - a^4}{1 - a^4}}$$

\(\therefore\) Degree and order are both 1.

2. Since, the given differential equation cannot be expressed as a polynomial in differential coefficients, so its degree is not defined.

3. The equation of tangent at any point $P(x, y)$ is

$$Y - y = \frac{d}{dx}(X - x)$$

This meets the X-axis at $A\left(-\frac{d}{d}, 0\right)$.

Similarly, it meets the Y-axis at

$$B\left(0, -\frac{d}{d}\right)$$

According to the given condition,

P is the mid-point of AB.



$$\therefore 2 = -\frac{d}{d} \text{ and } 2 = -\frac{d}{d}$$

$$\therefore +\frac{d}{d} = 0 \text{ and } +\frac{d}{d} = 0$$

Both of these equations reduce to

$$\frac{1}{-d} + \frac{1}{-d} = 0$$

Integrating both sides, we get

$$\log + \log = \log c$$

$$\therefore \log(\) = \log c$$

$\therefore = c$, which is the equation of rectangular hyperbola.

$$4. \sqrt{1+^2} + \sqrt{1+^2} = A \left(\sqrt{1+^2} - \sqrt{1+^2} \right)$$

Put $= \tan \alpha$, $= \tan \beta$

The equation becomes

$$\sec \alpha + \sec \beta = A (\tan \alpha \sec \beta - \tan \beta \sec \alpha)$$

$$\therefore \frac{1}{\cos \alpha} + \frac{1}{\cos \beta} = A \left(\frac{\sin \alpha}{\cos \alpha} \cdot \frac{1}{\cos \beta} - \frac{\sin \beta}{\cos \beta} \cdot \frac{1}{\cos \alpha} \right)$$

$$\therefore \frac{\cos \alpha + \cos \beta}{\cos \alpha \cos \beta} = A \left(\frac{\sin \alpha - \sin \beta}{\cos \alpha \cos \beta} \right)$$

$$\therefore \cos \alpha + \cos \beta = A (\sin \alpha - \sin \beta)$$

$$\begin{aligned} \therefore 2 \cos \left(\frac{\alpha + \beta}{2} \right) \cos \left(\frac{\alpha - \beta}{2} \right) \\ = A \cdot 2 \sin \left(\frac{\alpha - \beta}{2} \right) \cos \left(\frac{\alpha + \beta}{2} \right) \end{aligned}$$

$$\therefore \cot \left(\frac{\alpha - \beta}{2} \right) = A$$

$$\therefore \alpha - \beta = 2 \cot^{-1} A$$

$$\therefore \tan^{-1} - \tan^{-1} = 2 \cot^{-1} A$$

Differentiating w.r.t. , we get

$$\frac{1}{1+^2} - \frac{1}{1+^2} \cdot \frac{d}{d} = 0$$

$$\therefore \frac{d}{d} = \frac{1+^2}{1+^2}$$

\therefore Degree and order of the differential equation are both 1.

$$5. \frac{d}{d} = - + \frac{f(-)}{f'(-)}$$

Put $= v$

$$\therefore \frac{d}{d} = v + \frac{dv}{d}$$

\therefore The given equation becomes,

$$v + \frac{dv}{d} = v + \frac{f(v)}{f'(v)}$$

$$\therefore \frac{1}{-d} = \frac{f'(v)}{f(v)} dv$$

Integrating on both sides, we get

$$\log = \log f(v) + \log K$$

$$\Rightarrow = f(v)K$$

$$\Rightarrow = Kf\left(-\right)$$

$$\therefore f\left(-\right) = \frac{1}{K} \cdot = c, \text{ where } c = \frac{1}{K}$$

$$6. \text{ The given equation is } \frac{d}{d} + f'(\) = f(\)f'(\)$$

$$\therefore \text{I.F.} = e^{\int f'(\) d} = e^{f(\)}$$

\therefore the required solution is

$$.e^{f(\)} = \int e^{f(\)} \cdot f(\) f'(\) d$$

$$= \int e^t \cdot t dt, \text{ where } f(\) = t$$

$$= t \cdot e^t - \int e^t dt$$

$$= t e^t - e^t + c$$

$$\therefore .e^{f(\)} = f(\) e^{f(\)} - e^{f(\)} + c$$

$$\therefore = f(\) - 1 + c e^{-f(\)}$$

7. The given equation is

$$(\ + 1) f'(\) - 2(\ ^2 +) f(\) = \frac{e^2}{+1}$$

If $= f(\)$, the equation is

$$\frac{d}{d} - 2 \frac{e^2}{(\ + 1)^2}, \text{ which is a linear equation}$$

$$\therefore \text{I.F.} = e^{-\int 2 d} = e^{-2}$$

\therefore the required solution is

$$.e^{-2} = \int \frac{1}{(\ + 1)^2} d + c = -\frac{1}{+1} + c$$

When $= 0$, $= 5$

$$\therefore c = 6$$

$$\therefore .e^{-2} = -\frac{1}{+1} + 6$$

$$= \frac{-1+6 \ + 6}{+1} = \frac{6 \ + 5}{+1}$$

$$\therefore = f(\) = \left(\frac{6 \ + 5}{+1} \right) e^2$$



$$8. \quad \frac{d}{d} = \frac{x^2 + x^2 + 1}{2}$$

$$\therefore 2 \frac{d}{d} = (x^2 + 1) \frac{d}{d} + x^2 \frac{d}{d}$$

$$\therefore 2 \frac{d}{d} - x^2 \frac{d}{d} = (x^2 + 1) \frac{d}{d}$$

Dividing by x^2 , we get

$$\frac{2 \frac{d}{d} - x^2 \frac{d}{d}}{x^2} = \left(\frac{x^2 + 1}{x^2} \right) \frac{d}{d}$$

$$\therefore d \left(\frac{2}{x^2} \right) = \left(1 + \frac{1}{x^2} \right) \frac{d}{d}$$

Integrating both sides, we get

$$\frac{2}{-x} = \frac{1}{-x} + c$$

$$\therefore \frac{2}{x} = \frac{1}{x} - 1 + c$$

When $x = 1$, $\frac{2}{1} = \frac{1}{1} - 1 + c \therefore c = 0$

\therefore the required solution is $\frac{2}{x} = \frac{1}{x} - 1$
i.e., $\frac{2}{x} - \frac{1}{x} = -1$, which is the equation of a hyperbola.

$$9. \quad \frac{d}{d} + \frac{d}{d} + \frac{d - d}{x^2 + x^2} = 0$$

$$\therefore \frac{1}{2} (2 \frac{d}{d} + 2 \frac{d}{d}) + \frac{1}{2} \left(\frac{d - d}{x^2 + x^2} \right) = 0$$

$$\therefore \frac{1}{2} d \left(\frac{2}{x^2 + x^2} \right) + d \left(\tan^{-1} \frac{1}{x} \right) = 0$$

Integrating both sides, we get

$$\frac{1}{2} \left(\frac{2}{x^2 + x^2} \right) + \tan^{-1} \left(\frac{1}{x} \right) = \frac{c}{2}$$

$$\therefore \frac{2}{x^2 + x^2} + 2 \tan^{-1} \left(\frac{1}{x} \right) = c$$

$$\therefore 2 \tan^{-1} \left(\frac{1}{x} \right) = c - \frac{2}{x^2 + x^2}$$

$$\therefore \tan^{-1} \left(\frac{1}{x} \right) = \frac{c - \frac{2}{x^2 + x^2}}{2}$$

$$\therefore \frac{1}{x} = \tan \left(\frac{c - \frac{2}{x^2 + x^2}}{2} \right)$$

$$\therefore \frac{1}{x} = \tan \left(\frac{c - \frac{2}{x^2 + x^2}}{2} \right)$$

$$10. \quad 2x^2 \frac{d}{d} = \tan(x^2 - 2) - 2x^2$$

$$\therefore 2x^2 \frac{d}{d} + 2x^2 = \tan(x^2 - 2)$$

$$\therefore \frac{d}{d} (x^2 - 2) = \tan(x^2 - 2)$$

$$\therefore \frac{dz}{d} = \tan z, \text{ where } z = x^2 - 2$$

$$\therefore d = \cot z dz$$

Integrating both sides, we get

$$= \log(\sin z) + c \Rightarrow = \log(\sin(x^2 - 2)) + c$$

When $x = 1$, $z = \frac{\pi}{2}$, $\therefore z = \frac{\pi}{2}$

$$\therefore 1 = \log 1 + c, \therefore c = 0$$

the required solution is

$$= \log(\sin(x^2 - 2)) + 1$$

$$\therefore \log(\sin(x^2 - 2)) = -1$$

$$\therefore \sin(x^2 - 2) = e^{-1}$$

$$11. \quad 3 \frac{d}{d} + 4x^2 \tan = e \sec$$

Dividing by $x^3 \sec$, we get

$$\frac{1}{\sec} \cdot \frac{d}{d} + \frac{4 \tan}{\sec} = \frac{e}{x^3}$$

$$\therefore \cos \frac{d}{d} + (4 \sin) \cdot \frac{1}{x} = \frac{e}{x^3}$$

Put $\sin = t$, $\therefore \cos \frac{d}{d} = \frac{dt}{d}$

$$\therefore$$
 the equation becomes,

$$\frac{dt}{d} + \left(\frac{4}{t} \right) t = \frac{e}{x^3}$$
, which is a linear equation
$$\therefore \text{I.F.} = e^{\int \frac{4}{t} dt} = e^{4 \log t} = e^{\log t^4} = t^4$$

$$\therefore$$
 the required solution is

$$t^4 = \int e \cdot d$$

$$\therefore t^4 = e - e + c$$

$$\therefore \sin^4 = e - e + c$$

When $x = 1$, $\sin = 0$

$$\therefore c = 0$$

$$\therefore \sin^4 = e - e$$

$$\therefore \sin = (e - 1) e^{-\frac{1}{x^3}}$$

$$12. \quad \text{We have, } \frac{d}{d} = \tan(x - 2) \sec$$

$$\therefore \frac{1}{2} \cdot \frac{d}{d} = \frac{1}{2} \tan(x - 2) \sec$$

$$\therefore \frac{1}{2} \cdot \frac{d}{d} - \frac{1}{2} \tan(x - 2) = -\sec$$

Put $\frac{1}{2} = t$, $\therefore -\frac{1}{2} \cdot \frac{d}{d} = \frac{dt}{d}$

$$\therefore$$
 The equation becomes,

$$\frac{dt}{d} + (\tan(x - 2)) t = \sec$$
, which is linear equation.



$$\begin{aligned}\therefore \text{I.F.} &= e^{\int \tan x \, dx} \\ &= e^{\log(\sec x)} = \sec x\end{aligned}$$

$$\begin{aligned}\therefore \text{the required solution is} \\ t \sec x &= \int \sec^2 x \, dx + c\end{aligned}$$

$$\therefore \frac{1}{\sec x} = \tan x + c$$

$$\therefore \sec x = (c + \tan x)$$

$$13. \left(\frac{1}{x^2} - \frac{2}{x^3} \right) \frac{dx}{x} = \frac{1}{x^2} - \frac{2}{x^3}$$

$$\therefore \frac{1}{x^2} \frac{dx}{x} = \frac{1}{x^2} - \frac{2}{x^3}$$

Dividing by x^2 , we get

$$\frac{1}{x^2} \frac{dx}{x} = \frac{1}{x^2} - \frac{2}{x^3}$$

$$\therefore -\frac{1}{2} \frac{dx}{x} + \frac{1}{3} \frac{dx}{x^3} = \frac{1}{x^2}$$

$$\text{Put } \frac{1}{x} = t$$

$$\therefore -\frac{1}{2} \frac{dx}{x} = \frac{dt}{d}$$

The equation becomes

$$\frac{dt}{d} + \frac{1}{2} \cdot t = \frac{1}{2}, \text{ which is a linear equation}$$

$$\therefore \text{I.F.} = e^{\int \frac{1}{2} dx} = e^{\log x} = x$$

the required solution is

$$t = \int \frac{1}{x} dx + c$$

$$\therefore t = \log x + c$$

$$\therefore \frac{1}{x} = \log x + c$$

$$\therefore \frac{1}{x} = (\log x + c)$$

The curve passes through the point $(-1, 1)$.

$$\therefore 1 = -1(0 + c), \therefore c = -1$$

$$\therefore \text{the required solution is } \frac{1}{x} = (\log x - 1).$$

$$14. \frac{dy}{dx} = e^{-x} (1 - e^{-x})$$

$$\therefore e^{-x} \frac{dy}{dx} = e^{-x} (1 - e^{-x})$$

$$\therefore e^{-x} \frac{dy}{dx} = e^{-x} - e^{-2x}$$

$$\therefore e^{-x} \frac{dy}{dx} + e^{-x} y = e^{-x}$$

$$\text{Put } e^{-x} = t, \therefore e^{-x} \frac{dy}{dx} = \frac{dt}{dx}$$

The given equation becomes

$$\frac{dt}{dx} + e^{-x} \cdot t = e^{-x}, \text{ which is a linear equation.}$$

$$\therefore \text{I.F.} = e^{\int e^{-x} dx} = e^{-x}$$

the required solution is

$$\begin{aligned}t \cdot e^{-x} &= \int e^{-x} \cdot e^{-x} dx \\ &= \int e^{-2x} dx, \text{ where } e^{-x} = z \\ &= -\frac{1}{2} e^{-2x} + c\end{aligned}$$

$$\therefore t \cdot e^{-x} = -\frac{1}{2} e^{-2x} + c$$

$$\therefore e^{-x} \cdot e^{-x} = -\frac{1}{2} e^{-2x} + c$$

15. The equation of the tangent to the curve $y = f(x)$ at $P(x_1, y_1)$ is

$$Y - y_1 = \frac{dy}{dx} (X - x_1)$$

$$\text{This meets the X-axis at } \left(x_1 - \frac{y_1}{\frac{dy}{dx}}, 0 \right).$$

According to the given condition,

$$-\frac{1}{2} = \frac{y_1}{\frac{dy}{dx}}$$

$$\therefore -\frac{1}{2} = \frac{y_1}{\frac{dy}{dx}}$$

$$\therefore \frac{dy}{dx} = -2y_1, \text{ which is a homogeneous d.E.}$$

$$\text{Put } y = vx$$

$$\therefore \frac{dy}{dx} = v + x \frac{dv}{dx}$$

the equation becomes,

$$v + x \frac{dv}{dx} = \frac{v}{-v} = \frac{v}{1-v}$$

$$\begin{aligned}\therefore \frac{dv}{dx} &= \frac{v}{1-v} - v \\ &= \frac{v - v + v^2}{1-v}\end{aligned}$$

$$\therefore \frac{1-v}{v^2} dv = \frac{1}{x} dx$$

Integrating both sides, we get

$$\int \left(v^{-2} - \frac{1}{v} \right) dv = \int \frac{1}{x} dx + c$$



$$\begin{aligned} \therefore \frac{1}{v} \log v &= \log v + c \\ \therefore \log\left(\frac{1}{v}\right) &= \log v + c \\ \therefore -\log v + \log v &= \log v + c \\ \therefore -\log v &= c \end{aligned}$$

This curve passes through (1, 1).

$$\begin{aligned} \therefore c &= -1 \\ \therefore -\log v &= -1 \\ \therefore -\log v + 1 &= 0 \\ \therefore \log v &= 1 \\ \therefore v &= e \end{aligned}$$

16. $\frac{d}{d} = \sqrt{1-x^2}$

$$\therefore \int \frac{1}{\sqrt{1-x^2}} dx = \int 1 dx$$

$$\therefore -\sqrt{1-x^2} = x + c$$

$$\therefore (x+c)^2 = 1-x^2$$

$$\therefore (x+c)^2 + x^2 = 1$$

\therefore Radius is fixed, which is 1 and the centre is $(-c, 0)$ which is a variable centre on the X-axis.

17. (A) $f(x, y) = \frac{y-x}{x^2+y^2} = x^{-1} \left(\frac{y-x}{x^2+y^2} \right)$
 $= x^{-1} f\left(\frac{y-x}{x}, \frac{y-x}{x}\right)$

\therefore Homogeneous of degree -1 .

(B) $f(x, y) = (x)^{\frac{1}{3}} (y)^{-\frac{2}{3}} \tan^{-1}\left(\frac{y}{x}\right)$
 $= (x)^{\frac{1}{3}} (x)^{-\frac{1}{3}} (x)^{-\frac{2}{3}} \tan^{-1}\left(\frac{y}{x}\right) = x^{-\frac{1}{3}} f\left(\frac{y}{x}, \frac{y}{x}\right)$

\therefore Homogeneous of degree $-\frac{1}{3}$.

(C) $f(x, y) = y \left[\log \sqrt{y^2+x^2} - \log x \right]$
 $= y \left[\log \frac{\sqrt{y^2+x^2}}{x} \right] + y e^{-\frac{1}{x}}$

$$\begin{aligned} &= y \left[\log \frac{\sqrt{y^2+x^2}}{x} \right] + y e^{-\frac{1}{x}} \\ &= y \left[\left(\log \sqrt{y^2+x^2} - \log x \right) \right] + y e^{-\frac{1}{x}} \\ &= y f\left(\frac{y}{x}, \frac{y}{x}\right) \end{aligned}$$

\therefore Homogeneous of degree 1.

(D) $f(x, y) = x \left\{ \log \left[\frac{2x^2+y^2}{x} \right] - \log(x+y) \right\}$
 $+ x^2 \tan\left(\frac{y+2x}{3x-y}\right)$
 $= x \left[\log \frac{2x^2+y^2}{(x+y)} \right] + x^2 \tan\left(\frac{y+2x}{3x-y}\right)$

\therefore Non-Homogeneous.

18. $\frac{d}{d} = \frac{\sin 2}{1+\tan 2}$

$$\therefore \frac{d}{d} = \frac{1+\tan 2}{\sin 2}$$

$$\therefore \frac{d}{d} - \frac{1}{\sin 2} = \frac{\tan 2}{\sin 2},$$

which is a linear equation

$$\begin{aligned} \therefore \text{I.F.} &= e^{-\int \operatorname{cosec} 2 \, d} = e^{-\frac{1}{2} \log(\tan 2)} = e^{\log(\tan 2) \cdot \frac{-1}{2}} \\ &= e^{\log \sqrt{\cot 2}} \\ &= \sqrt{\cot 2} \end{aligned}$$

\therefore the required solution is

$$\begin{aligned} \sqrt{\cot 2} \cdot d &= \int \frac{\tan 2}{\sin 2} \sqrt{\cot 2} \, d + c \\ &= \int \frac{1}{\sqrt{\tan 2}} \cdot \frac{\sin 2}{\cos 2} \cdot \frac{1}{2 \sin 2 \cos 2} \, d + c \\ &= \int \frac{1}{2\sqrt{\tan 2}} \sec^2 2 \, d + c \end{aligned}$$

$$\therefore \sqrt{\cot 2} \cdot d = \sqrt{\tan 2} + c$$

The curve passes through $\left(1, \frac{\pi}{4}\right)$.

$$\therefore 1 = 1 + c, \quad \therefore c = 0$$

\therefore the equation of the curve is $d = \tan 2$



19. The equation of hyperbola is $x^2 - y^2 = 2$
 $\therefore \frac{dy}{dx} = \frac{2}{x}$
 $\therefore m_1 = \frac{dy}{dx} = -\frac{2}{x}$ (slope of tangent to the hyperbola)
 $m_2 = \frac{dy}{dx} =$ slope of tangent to the required family of curves.
 The curves are intersecting orthogonally,
 $m_1 m_2 = -1$
 $\therefore \frac{dy}{dx} \times \left(-\frac{2}{x}\right) = -1$
 $\therefore \frac{dy}{dx} = \frac{x}{2}$
 Integrating both sides, we get $y = \frac{x^2}{6} + c$,
 which is the equation of required family of curves.

20. The given equation is
 $\frac{dy}{dx} = -\frac{(x+3)}{1+x^2} = \frac{-(x+3)}{1+(x^2)}$
 $\therefore \frac{dy}{dx} = -\frac{1+(x^2)}{(1+x^2)}$
 $= -\frac{1}{(1+x^2)}$
 $\therefore \frac{dy}{dx} + \frac{1}{x} = -\frac{1}{(1+x^2)}$, which is a linear equation

- \therefore I.F. $= e^{\int \frac{1}{x} dx} = e^{\log x} = x$
 \therefore the required solution is
 $y = -\int \frac{1}{1+x^2} dx + c$
 $\therefore y = -\tan^{-1} x + c$
 The curve passes through $(0, 1) \therefore c = \frac{\pi}{4}$
 \therefore the required equation of the curve is
 $y + \tan^{-1} x = \frac{\pi}{4}$

21. Slope of tangent $= \frac{dy}{dx}$
 \therefore slope of normal $= -\frac{1}{\frac{dy}{dx}}$

The equation of the normal is

$$Y - y = -\frac{1}{\frac{dy}{dx}}(X - x)$$

$$\therefore (Y - y) \frac{dy}{dx} + (X - x) = 0$$

The normal passes through the point $(3, 0)$.

$$\therefore (0 - y) \frac{dy}{dx} + (3 - x) = 0$$

$$\therefore \frac{dy}{dx} = 3 - x$$

$$\therefore dx = (3 - x) dy$$

Integrating both sides, we get $\frac{x^2}{2} = 3y - \frac{y^2}{2} + c$

The curve passes through $(3, 4)$, $\therefore c = \frac{7}{2}$

$$\therefore \text{the equation of the curve is } \frac{x^2}{2} = 3y - \frac{y^2}{2} + \frac{7}{2}$$

$$\therefore x^2 + y^2 - 6y - 7 = 0$$

22. The given equation can be written as

$$x^2 - dx = x^3(1 + \log x) dx$$

$$\therefore -\left(\frac{dx - d^2}{2}\right) = (1 + \log x) dx$$

$$\therefore -d\left(\frac{x}{2}\right) = (1 + \log x) dx$$

$$\therefore -\frac{dx}{2} = (1 + \log x) dx$$

Integrating both sides, we get

$$-\frac{(x)^2}{2} = (1 + \log x) \int x^2 dx - \int \left\{ \frac{dx}{x} (1 + \log x) \int x^2 dx \right\} dx + \frac{c}{2}$$

$$\therefore -\frac{x^2}{2} = (1 + \log x) \cdot \frac{x^3}{3} - \int \frac{1}{x} \cdot \frac{x^3}{3} dx + \frac{c}{2}$$

$$\therefore -\frac{x^2}{2} = (1 + \log x) \cdot \frac{x^3}{3} - \frac{x^3}{9} + \frac{c}{2}$$

$$\therefore -\frac{x^2}{2} = \frac{2x^3}{3} (1 + \log x) - \frac{2x^3}{9} + c$$

$$= \frac{2x^3}{3} \left(1 + \log x - \frac{1}{3}\right) + c$$

$$\therefore -\frac{x^2}{2} = \frac{2x^3}{3} \left(\frac{2}{3} + \log x\right) + c$$



23. The equation of the tangent at P(,) is

$$Y - = \frac{d}{d}(X -)$$

This meets the Y-axis at $\left(0, - \frac{d}{d}\right)$.

According to the given condition,

$$- \frac{d}{d} = ^3$$

$$\Rightarrow \frac{d}{d} + \left(-\frac{1}{d}\right) = - ^2$$

$$\therefore \text{I.F.} = e^{\int -\frac{1}{d}d} = e^{-\log} = \frac{1}{d}$$

\therefore solution of the given equation is

$$\frac{1}{d} = \int - ^2 \cdot \frac{1}{d} + c$$

$$\Rightarrow - = -\frac{^2}{2} + c \Rightarrow = -\frac{^3}{2} + c$$

$$\therefore f() = -\frac{^3}{2} + c \quad \dots\text{(i)}$$

$$\therefore f(1) = -\frac{1}{2} + c$$

$$\Rightarrow 1 = -\frac{1}{2} + c \Rightarrow c = \frac{3}{2}$$

$$\therefore f() = -\frac{^3}{2} + \frac{3}{2} \quad \dots\text{[From (i)]}$$

$$\therefore f(-3) = \frac{27}{2} - \frac{9}{2} = 9$$

24. $\frac{dV}{dt} = -k(T - t)$

$$\Rightarrow dV = -k(T - t)dt$$

Integrating on both sides, we get

$$\int dV = -k \int (T - t)dt + c$$

$$\Rightarrow V(t) = \frac{k(T - t)^2}{2} + c \quad \dots\text{(i)}$$

Initially i.e., when $t = 0$, $V(t) = I$

$$\therefore I = \frac{kT^2}{2} + c \Rightarrow c = I - \frac{kT^2}{2}$$

$$\therefore V(t) = \frac{k(T - t)^2}{2} + I - \frac{kT^2}{2} \quad \dots\text{[From (i)]}$$

When $t = T$,

$$V(T) = I - \frac{kT^2}{2}$$

25. $\frac{d}{d} = - - \cos^2 - \quad \dots\text{(i)}$

Put $= v \quad \dots\text{(ii)}$

$$\Rightarrow \frac{d}{d} = v + \frac{dv}{d} \quad \dots\text{(iii)}$$

Substituting (ii) and (iii) in (i), we get

$$v + \frac{dv}{d} = v - \cos^2 v \Rightarrow \frac{dv}{d} = - \cos^2 v$$

Integrating on both sides, we get

$$\int \sec^2 v dv = - \int \frac{d}{d} + c$$

$$\Rightarrow \tan v = - \log + c$$

$$\Rightarrow \tan - = - \log + c \quad \dots\text{(iv)}$$

Since, the required curve passes through $\left(1, \frac{\pi}{4}\right)$.

$$\therefore \tan \frac{\pi}{4} = - \log 1 + c \Rightarrow c = 1$$

$$\therefore \tan - = - \log + 1 \quad \dots\text{[From (iv)]}$$

$$\Rightarrow \tan - = - \log + \log e$$

$$\Rightarrow = \tan^{-1} \left[\log \left(\frac{e}{-} \right) \right]$$

08 Probability Distribution



Hints



Classical Thinking

1. The sum of all the probabilities in a probability distribution is always unity.

$$\begin{aligned} \therefore 0.1 + k + 0.2 + 2k + 0.3 + k &= 1 \\ \Rightarrow 4k + 0.6 &= 1 \Rightarrow 4k = 0.4 \Rightarrow k = 0.1 \end{aligned}$$

2. Since, $\sum_{i=1}^4 P(X = x_i) = 1$

$$\begin{aligned} \therefore \frac{1}{8} + \frac{1}{2} + \frac{1}{4} + k &= 1 \\ \Rightarrow k + \frac{1+4+2}{8} &= 1 \\ \Rightarrow k &= 1 - \frac{7}{8} = \frac{1}{8} \end{aligned}$$

3. Since, $\sum_{i=0}^4 P(X = x_i) = 1$

$$\begin{aligned} \therefore k + 2k + 3k + 2k + k &= 1 \\ \Rightarrow 9k &= 1 \Rightarrow k = \frac{1}{9} \end{aligned}$$

4. The probability distribution of X is

X	0	1	2
P(X)	k	2k	3k

Since, $\sum_{i=0}^2 P(X = x_i) = 1$

$$\begin{aligned} \therefore k + 2k + 3k &= 1 \\ \Rightarrow 6k &= 1 \Rightarrow k = \frac{1}{6} \end{aligned}$$

5. $P(X \leq 2) = P(X = 0) + P(X = 1) + P(X = 2)$

$$\begin{aligned} &= \frac{{}^5C_0}{2^5} + \frac{{}^5C_1}{2^5} + \frac{{}^5C_2}{2^5} \\ &= \frac{1+5+10}{2^5} = \frac{16}{32} \end{aligned}$$

6. $F(-1) = 0.2 + 0.3 = 0.5$

7. $F(x_1) = p_1 = 0.05$

$$F(x_2) = p_1 + p_2 = 0.05 + 0.2 = 0.25$$

$$F(x_3) = p_1 + p_2 + p_3 = 0.25 + 0.15 = 0.4$$

$$F(x_4) = p_1 + p_2 + p_3 + p_4 = 0.4 + 0.25 = 0.65$$

$$F(x_5) = p_1 + p_2 + p_3 + p_4 + p_5 = 0.65 + 0.35 = 1$$

$$\begin{aligned} 8. E(X) &= \sum x_i \cdot P(x_i) \\ &= 0 \left(\frac{1}{8} \right) + 1 \left(\frac{3}{8} \right) + 2 \left(\frac{3}{8} \right) + 3 \left(\frac{1}{8} \right) \\ &= \frac{3}{8} + \frac{6}{8} + \frac{3}{8} = \frac{3}{2} \end{aligned}$$

$$\begin{aligned} 9. E(X) &= \sum x_i \cdot P(x_i) \\ &= 0(0.2) + 1(0.5) + 3(0.2) + 5(0.1) \\ &= 0 + 0.5 + 0.6 + 0.5 \\ &= 1.6 \end{aligned}$$

$$\begin{aligned} \text{Variance} &= \sum x_i^2 \cdot P(x_i) - [E(X)]^2 \\ &= (0)^2(0.2) + (1)^2(0.5) + (3)^2(0.2) \\ &\quad + (5)^2(0.1) - (1.6)^2 \\ &= 4.8 - 2.56 = 2.24 \end{aligned}$$

$$\begin{aligned} 10. \text{Since, } \text{Var}(X) &= E(X^2) - [E(X)]^2 \\ \therefore 4 &= 13 - [E(X)]^2 \\ \therefore [E(X)]^2 &= 13 - 4 = 9 \\ \therefore E(X) &= 3 \end{aligned}$$

$$\begin{aligned} 11. \text{Since, } \text{Var}(X) &= E(X^2) - [E(X)]^2 \\ \therefore 6 &= E(X^2) - (5)^2 \\ \therefore E(X^2) &= 25 + 6 = 31 \end{aligned}$$

$$\begin{aligned} 12. \text{Mean} = E(X) &= \sum x_i \cdot P(x_i) \\ &= \frac{1}{6}(1) + \frac{1}{3}(2) + \frac{1}{3}(3) + \frac{1}{6}(4) \\ &= \frac{1}{6} + \frac{2}{3} + 1 + \frac{4}{6} = \frac{1+4+6+4}{6} \\ &= \frac{15}{6} = \frac{5}{2} \end{aligned}$$

$$\begin{aligned} \text{Variance} &= \sum x_i^2 \cdot P(x_i) - [E(X)]^2 \\ &= \frac{1}{6}(1)^2 + \frac{1}{3}(2)^2 + \frac{1}{3}(3)^2 + \frac{1}{6}(4)^2 - \left(\frac{5}{2} \right)^2 \\ &= \frac{1}{6} + \frac{4}{3} + \frac{9}{3} + \frac{16}{6} - \frac{25}{4} \\ &= \frac{2+16+36+32-75}{12} \\ &= \frac{86-75}{12} = \frac{11}{12} \end{aligned}$$



$$13. E(X) = 1\left(\frac{1}{7}\right) + 2\left(\frac{2}{7}\right) + 3\left(\frac{3}{7}\right) + 4\left(\frac{1}{7}\right) = \frac{18}{7}$$

$$E(X^2) = (1^2)\left(\frac{1}{7}\right) + (2^2)\left(\frac{2}{7}\right) + (3^2)\left(\frac{3}{7}\right) + (4^2)\left(\frac{1}{7}\right)$$

$$= \frac{1}{7} + \frac{8}{7} + \frac{27}{7} + \frac{16}{7} = \frac{52}{7}$$

$$\therefore \text{Var}(X) = E(X^2) - [E(X)]^2$$

$$= \frac{52}{7} - \left(\frac{18}{7}\right)^2 = \frac{40}{49}$$

$$14. P(1 < X < 3) = \int_1^3 f(x) dx$$

$$= \int_1^3 \frac{1}{5} dx = \frac{1}{5} [x]_1^3 = \frac{2}{5}$$

$$15. P\left(\frac{1}{3} < X < \frac{1}{2}\right) = \int_{1/3}^{1/2} f(x) dx$$

$$= \int_{1/3}^{1/2} 2x dx = [x^2]_{1/3}^{1/2} = \frac{5}{36}$$

$$16. P(0.5 \leq X \leq 1.5) = \int_{0.5}^{1.5} f(x) dx = \int_{0.5}^{1.5} 0.5x dx$$

$$= 0.5 \left[\frac{x^2}{2} \right]_{0.5}^{1.5} = \frac{1}{2}$$

$$17. P(X > 3) = \int_3^4 f(x) dx$$

$$= \int_3^4 \frac{1}{8} dx = \frac{1}{8} \left[\frac{x^2}{2} \right]_3^4 = \frac{7}{16}$$

18. Since, $f(x)$ is the p.d.f. of X .

$$\therefore \int_{-\infty}^{\infty} f(x) dx = 1$$

$$\Rightarrow \int_0^2 kx^2 dx = 1 \Rightarrow k \left[\frac{x^3}{3} \right]_0^2 = 1$$

$$\Rightarrow k \left[\frac{8}{3} \right] = 1 \Rightarrow k = \frac{3}{8}$$

19. The c.d.f. of X is

$$F(x) = \int_{-1}^x \frac{2}{3} dx = \frac{1}{3} \left[\frac{x^3}{3} \right]_{-1}^x$$

$$= \frac{1}{3} \left(\frac{x^3}{3} + \frac{1}{3} \right) = \frac{x^3}{9} + \frac{1}{9}$$



Critical Thinking

1. The sum of all the probabilities in a probability distribution is always unity.
In option (A), we have $0.3 + 0.2 + 0.4 + 0.1 = 1$

2. Since, $\sum_{i=1}^4 P(X = x_i) = 1$

$$\therefore k + 2k + 3k + 4k = 1$$

$$\Rightarrow 10k = 1 \Rightarrow k = \frac{1}{10}$$

$$\text{Now, } P(X < 3) = P(X = 1) + P(X = 2)$$

$$= k + 2k$$

$$= 3k = 3 \left(\frac{1}{10} \right) = 0.3$$

3. Since, $\sum_{i=0}^4 P(X = x_i) = 1$

$$\therefore k + 3k + 5k + 2k + k = 1$$

$$\Rightarrow 12k = 1 \Rightarrow k = \frac{1}{12}$$

$$\text{Now, } P(X \geq 2) = P(X = 2) + P(X = 3) + P(X = 4)$$

$$= 5k + 2k + k$$

$$= 8k = 8 \left(\frac{1}{12} \right) = \frac{2}{3}$$

4. Since, $\sum_{i=0}^6 P(X = x_i) = 1$

$$\therefore k + 3k + 5k + 7k + 9k + 11k + 13k = 1$$

$$\Rightarrow 49k = 1 \Rightarrow k = \frac{1}{49}$$

$$\therefore P(0 < X < 4)$$

$$= P(X = 1) + P(X = 2) + P(X = 3)$$

$$= 3k + 5k + 7k = 15k = \frac{15}{49}$$

5. $P(X \text{ is odd})$
 $= P(X = -3) + P(X = -1) + P(X = 1) + P(X = 3)$
 $= 0.05 + 0.15 + 0.25 + 0.10 = 0.55$

6.

$X =$	0	1	2	3	4
$P(X = x_i)$	k	4k	6k	4k	k

Since, $\sum_{i=0}^4 P(X = x_i) = 1$

$$\therefore k + 4k + 6k + 4k + k = 1$$

$$\Rightarrow 16k = 1$$

$$\Rightarrow k = \frac{1}{16}$$



7. Since, $\sum_{=0}^7 P(X =) = 1$

$$\therefore 0 + P + 2P + 2P + 3P + P^2 + 2P^2 + 7P^2 + P = 1$$

$$\therefore 10P^2 + 9P - 1 = 0$$

$$\therefore (P + 1)(10P - 1) = 0$$

$$\therefore P = \frac{1}{10} \quad \dots[\because P \geq 0, \therefore P + 1 \neq 0]$$

8. Since, $\sum_{=0}^2 P(X =) = 1$

$$\therefore 3k^3 + 4k - 10k^2 + 5k - 1 = 1$$

$$\Rightarrow 3k^3 - 10k^2 + 9k - 2 = 0$$

$$\Rightarrow (k - 1)(k - 2)(3k - 1) = 0$$

$$\Rightarrow k = 1 \text{ or } k = 2 \text{ or } k = \frac{1}{3}$$

For $k = 1$ or $k = 2$, $P(X = 1) < 0$, which is not possible.

$$\therefore k = \frac{1}{3}$$

9. Since, $\sum_{=1}^5 P(X =) = 1$

$$\therefore \frac{1}{20} + \frac{3}{20} + a + b + \frac{1}{20} = 1$$

$$\Rightarrow a + b = 1 - \frac{5}{20}$$

$$\Rightarrow a + 2a = 1 - \frac{1}{4} \quad \dots[\because b = 2a \text{ (given)}]$$

$$\Rightarrow 3a = \frac{3}{4}$$

$$\Rightarrow a = \frac{1}{4} \text{ and } b = 2\left(\frac{1}{4}\right) = \frac{1}{2}$$

10. The probability distribution of X is

X	0	1	2	3	4
P(X)	0.1	k	2k	2k	k

$$\text{Since, } \sum_{=0}^4 P(X =) = 1$$

$$\Rightarrow 0.1 + k + 2k + 2k + k = 1$$

$$\Rightarrow 6k = 0.9 \quad \Rightarrow k = 0.15$$

11. Let $P(X = 3) = k$. Then $P(X = 1) = \frac{k}{2}$,

$$P(X = 2) = \frac{k}{3}, P(X = 4) = \frac{k}{5}$$

$$\text{Since, } P(X = 1) + P(X = 2) + P(X = 3) + P(X = 4) = 1$$

$$\therefore \frac{k}{2} + \frac{k}{3} + k + \frac{k}{5} = 1 \quad \Rightarrow k = \frac{30}{61}$$

\therefore option (A) is the correct answer.

12. Let X denotes the number of heads. Thus, the possible values of X are 0, 1, 2 and 3.

$$P(X = 0) = P(\text{getting no head})$$

$$= P(\text{TTT}) = \frac{1}{8}$$

$$P(X = 1) = P(\text{getting one head})$$

$$= P(\text{HTT, THT, TTH}) = \frac{3}{8}$$

$$P(X = 2) = P(\text{getting two heads})$$

$$= P(\text{HHT, THH, HTH}) = \frac{3}{8}$$

$$P(X = 3) = P(\text{getting three heads})$$

$$= P(\text{HHH}) = \frac{1}{8}$$

\therefore Option (D) is the correct answer.

13. Let X denote the number of red balls drawn from the bag. There are 4 red balls and X can take values 0, 1, 2 and 3.

$$P(X = 0) = \text{Probability of getting no red ball}$$

$$= \frac{{}^6C_3}{{}^{10}C_3} = \frac{1}{6}$$

$$P(X = 1) = \text{Probability of getting one red ball}$$

$$= \frac{{}^4C_1 \times {}^6C_2}{{}^{10}C_3} = \frac{1}{2}$$

$$P(X = 2) = \text{Probability of getting two red balls}$$

$$= \frac{{}^4C_2 \times {}^6C_1}{{}^{10}C_3} = \frac{3}{10}$$

$$P(X = 3) = \text{Probability of getting three red balls}$$

$$= \frac{{}^4C_3}{{}^{10}C_3} = \frac{1}{30}$$

14. Let X denote the number of defective mangoes from the bag. X can take values 0, 1, 2, 3 and 4.

$$P(X = 0) = \text{Probability of getting no defective mango}$$

$$= \frac{{}^{15}C_4}{{}^{20}C_4} = \frac{91}{323}$$

$$P(X = 1) = \text{Probability of getting one defective mango}$$

$$= \frac{{}^5C_1 \times {}^{15}C_3}{{}^{20}C_4} = \frac{455}{969}$$



$$P(X = 2) = \text{Probability of getting two defective mangoes} = \frac{{}^5C_2 \times {}^{15}C_2}{{}^{20}C_4} = \frac{70}{323}$$

$$P(X = 3) = \text{Probability of getting three defective mangoes} = \frac{{}^5C_3 \times {}^{15}C_1}{{}^{20}C_4} = \frac{10}{323}$$

$$P(X = 4) = \text{Probability of getting four defective mangoes} = \frac{{}^5C_4}{{}^{20}C_4} = \frac{1}{969}$$

$$\begin{aligned} 15. \quad P(X = 2) &= F(2) - F(1) = 0.43 - 0.18 = 0.25 \\ P(X = 3) &= F(3) - F(2) = 0.54 - 0.43 = 0.11 \\ P(X = 4) &= F(4) - F(3) = 0.68 - 0.54 = 0.14 \\ \therefore P(1 < X < 5) &= P(X = 2) + P(X = 3) + P(X = 4) \\ &= 0.25 + 0.11 + 0.14 = 0.50 \end{aligned}$$

$$\begin{aligned} 16. \quad P(X = 1) &= F(1) - F(0) = 0.65 - 0.5 = 0.15 \\ P(X = 3) &= F(3) - F(2) = 0.75 - 0.65 = 0.10 \\ P(X = 5) &= 0.85 - 0.75 = 0.10 \\ P(X = 7) &= 0.90 - 0.85 = 0.05 \\ P(X = 9) &= 1 - 0.90 = 0.10 \end{aligned}$$

$$\begin{aligned} \therefore P(X \leq 3 | X > 0) &= \frac{P(X=1) + P(X=3)}{P(X=1) + P(X=3) + P(X=5) + P(X=7) + P(X=9)} \\ &= \frac{0.15 + 0.1}{0.15 + 0.1 + 0.1 + 0.05 + 0.1} \\ &= \frac{0.25}{0.50} = \frac{1}{2} \end{aligned}$$

17. The sum of all the probabilities in a probability distribution is always unity.

$$\begin{aligned} \therefore 0.1 + k + 0.2 + 2k + 0.3 + k &= 1 \\ \Rightarrow 0.6 + 4k &= 1 \\ \Rightarrow 4k &= 0.4 \\ \Rightarrow k &= 0.1 \end{aligned}$$

$$\begin{aligned} \therefore E(X) &= (-2)(0.1) + (-1)(0.1) + 0(0.2) \\ &\quad + 1(2 \times 0.1) + 2(0.3) + 3(0.1) = 0.8 \end{aligned}$$

18. The sum of all the probabilities in a probability distribution is always unity.

$$\begin{aligned} \therefore 0.2 + 0.1 + 0.3 + k &= 1 \\ \therefore k &= 1 - 0.6 = 0.4 \end{aligned}$$

$$\begin{aligned} E(X) &= \sum x_i \cdot P(x_i) \\ &= 1(0.2) + 2(0.1) + 3(0.3) + 4(0.4) \\ &= 0.2 + 0.2 + 0.9 + 1.6 = 2.9 \\ \text{Var}(X) &= E(X^2) - [E(X)]^2 \\ &= (1)^2(0.2) + (2)^2(0.1) + (3)^2(0.3) \\ &\quad + (4)^2(0.4) - (2.9)^2 \\ &= 0.2 + 0.4 + 2.7 + 6.4 - 8.41 \\ &= 9.7 - 8.41 = 1.29 \end{aligned}$$

19. The sum of all the probabilities in a probability distribution is always unity.

$$\begin{aligned} \therefore k + 3k + 3k + k &= 1 \\ \Rightarrow 8k &= 1 \\ \Rightarrow k &= \frac{1}{8} \end{aligned}$$

$$\begin{aligned} E(X) &= 0\left(\frac{1}{8}\right) + 1\left(\frac{3}{8}\right) + 2\left(\frac{3}{8}\right) + 3\left(\frac{1}{8}\right) \\ &= \frac{3}{2} \end{aligned}$$

$$\begin{aligned} \text{Var}(X) &= E(X^2) - [E(X)]^2 \\ &= 0^2\left(\frac{1}{8}\right) + 1^2\left(\frac{3}{8}\right) + 2^2\left(\frac{3}{8}\right) + 3^2\left(\frac{1}{8}\right) - \left(\frac{3}{2}\right)^2 \\ &= \frac{3}{4} \end{aligned}$$

$$\begin{aligned} 20. \quad \text{Mean} &= E(X) = \sum x_i \cdot P(x_i) \\ &= 1(0.1) + 2(0.2) + 3(0.3) + 4(0.4) = 3 \\ \text{Var}(X) &= \sum x_i^2 \cdot P(x_i) - [E(X)]^2 \\ &= 1^2(0.1) + 2^2(0.2) + 3^2(0.3) + 4^2(0.4) - (3)^2 \\ &= 0.1 + 0.8 + 2.7 + 6.4 - 9 = 10 - 9 = 1 \\ \therefore \text{S.D.} &= 1 \end{aligned}$$

21. The p.m.f. of the r.v. X is as follows:

X =	-1	0	1	2
P(X =)	$\frac{2}{5}$	$\frac{3}{10}$	$\frac{1}{5}$	$\frac{1}{10}$

$$\therefore E(X) = -1\left(\frac{2}{5}\right) + 0 + 1\left(\frac{1}{5}\right) + 2\left(\frac{1}{10}\right) = 0$$

22.

X =	1	2	3	4
P(X =)	k	4k	9k	16k

Since, $P(1) + P(2) + P(3) + P(4) = 1$

$$\begin{aligned} \therefore k + 4k + 9k + 16k &= 1 \\ \Rightarrow 30k &= 1 \\ \Rightarrow k &= \frac{1}{30} \end{aligned}$$

$$\begin{aligned} \therefore E(X) &= 1 \cdot \frac{1}{30} + 2 \cdot \frac{4}{30} + 3 \cdot \frac{9}{30} + 4 \cdot \frac{16}{30} \\ &= \frac{100}{30} \\ &= \frac{10}{3} \end{aligned}$$



$$23. \quad P(1) = \frac{C}{1^3}, \quad P(2) = \frac{C}{2^3}, \quad P(3) = \frac{C}{3^3}$$

$$\text{Since, } P(1) + P(2) + P(3) = 1$$

$$\therefore \frac{C}{1^3} + \frac{C}{2^3} + \frac{C}{3^3} = 1$$

$$\Rightarrow C \left(\frac{1}{1} + \frac{1}{8} + \frac{1}{27} \right) = 1$$

$$\Rightarrow C \left(\frac{216 + 27 + 8}{216} \right) = 1$$

$$\Rightarrow C = \frac{216}{251}$$

$$\begin{aligned} \therefore E(X) &= (1) \frac{C}{1^3} + (2) \frac{C}{2^3} + (3) \frac{C}{3^3} \\ &= C \left(1 + \frac{1}{4} + \frac{1}{9} \right) = C \left(\frac{36 + 9 + 4}{36} \right) \\ &= \frac{216}{251} \times \frac{49}{36} = \frac{294}{251} \end{aligned}$$

$$24. \quad E(X) = \sum_i x_i \cdot P(x_i) = 1.6$$

$$\begin{aligned} \text{Var}(X) &= \sum_i x_i^2 \cdot P(x_i) - [E(X)]^2 \\ &= 4.8 - 2.56 \\ &= 2.24 \end{aligned}$$

$$\text{Now, } 4 E(X^2) - \text{Var}(X)$$

$$\begin{aligned} &= 4 \sum_i x_i^2 \cdot P(x_i) - \text{Var}(X) = 4(4.8) - 2.24 \\ &= 19.2 - 2.24 \\ &= 16.96 \end{aligned}$$

$$\begin{aligned} 25. \quad E(X) &= \sum_i x_i \cdot P(x_i) \\ &= 0(q^2) + 1(2pq) + 2(p^2) \\ &= 2pq + 2p^2 \\ &= 2p(q + p) \\ &= 2p \quad \dots[\because p + q = 1] \end{aligned}$$

$$\begin{aligned} \text{Var}(X) &= E(X^2) - [E(X)]^2 \\ &= 0(q^2) + 1^2(2pq) + 2^2(p^2) - (2p)^2 \\ &= 2pq + 4p^2 - 4p^2 \\ &= 2pq \end{aligned}$$

$$\begin{aligned} 26. \quad E(X) &= \sum_i x_i \cdot P(x_i) \\ &= 0(q^3) + 1(3q^2p) + 2(3qp^2) + 3(p^3) \\ &= 3pq(q + 2p) + 3p^3 \\ &= 3pq[(p + q) + p] + 3p^3 \\ &= 3pq(1 + p) + 3p^3 \quad \dots[\because p + q = 1] \\ &= 3pq + 3p^2q + 3p^3 \\ &= 3pq + 3p^2(q + p) \\ &= 3p(q + p) \quad \dots[\because p + q = 1] \\ &= 3p(1) \\ &= 3p \end{aligned}$$

27. X can take values 0, 1, 2 and 3.

$$\begin{aligned} P(X = 0) &= \text{Probability of getting no head} \\ &= \frac{1}{8} \end{aligned}$$

$$\begin{aligned} P(X = 1) &= \text{Probability of getting one head} \\ &= \frac{3}{8} \end{aligned}$$

$$\begin{aligned} P(X = 2) &= \text{Probability of getting two heads} \\ &= \frac{3}{8} \end{aligned}$$

$$\begin{aligned} P(X = 3) &= \text{Probability of getting three heads} \\ &= \frac{1}{8} \end{aligned}$$

$$\begin{aligned} \therefore E(X) &= (0) \left(\frac{1}{8} \right) + (1) \left(\frac{3}{8} \right) + (2) \left(\frac{3}{8} \right) + (3) \left(\frac{1}{8} \right) \\ &= 0 + \frac{3}{8} + \frac{3}{4} + \frac{3}{8} = \frac{12}{8} = \frac{3}{2} \end{aligned}$$

28. X can take values 0, 1 and 2.

$$\begin{aligned} P(X = 0) &= \text{Probability of getting no tail} \\ &= \frac{1}{4} \end{aligned}$$

$$\begin{aligned} P(X = 1) &= \text{Probability of getting one tail} \\ &= \frac{1}{2} \end{aligned}$$

$$\begin{aligned} P(X = 2) &= \text{Probability of getting two tails} \\ &= \frac{1}{4} \end{aligned}$$

$$\begin{aligned} \therefore E(X) &= 0 \left(\frac{1}{4} \right) + (1) \left(\frac{1}{2} \right) + 2 \left(\frac{1}{4} \right) \\ &= 0 + \frac{1}{2} + \frac{1}{2} = 1 \end{aligned}$$

$$\begin{aligned} \text{Var}(X) &= E(X^2) - [E(X)]^2 \\ &= 0^2 \left(\frac{1}{4} \right) + 1^2 \left(\frac{1}{2} \right) + 2^2 \left(\frac{1}{4} \right) - (1)^2 \\ &= \frac{1}{2} \end{aligned}$$

29. X can take values 0, 1 and 2.

$$P(X = 0) = \text{Probability of not getting six} = \frac{25}{36}$$

$$P(X = 1) = \text{Probability of getting one six} = \frac{10}{36}$$

$$P(X = 2) = \text{Probability of getting two sixes} = \frac{1}{36}$$

\therefore the probability distribution of X is given by



X	0	1	2
P(X)	$\frac{25}{36}$	$\frac{10}{36}$	$\frac{1}{36}$

$$\begin{aligned} \therefore E(X) &= \sum_i P(i) = 0 \times \frac{25}{36} + 1 \times \frac{10}{36} + 2 \times \frac{1}{36} \\ &= \frac{10}{36} + \frac{2}{36} = \frac{1}{3} \end{aligned}$$

30. In a single throw of a pair of dice, the sum of the numbers on them can be 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12. So X can take values 2, 3, 4, ..., 12. The probability distribution of X is

X: 2 3 4 5 6 7 8 9 10 11 12
 P(X): $\frac{1}{36}$ $\frac{2}{36}$ $\frac{3}{36}$ $\frac{4}{36}$ $\frac{5}{36}$ $\frac{6}{36}$ $\frac{5}{36}$ $\frac{4}{36}$ $\frac{3}{36}$ $\frac{2}{36}$ $\frac{1}{36}$

$$\begin{aligned} \therefore E(X) &= \frac{1}{36} \times 2 + \frac{2}{36} \times 3 + \frac{3}{36} \times 4 + \frac{4}{36} \times 5 \\ &\quad + \frac{5}{36} \times 6 + \frac{6}{36} \times 7 + \frac{5}{36} \times 8 + \frac{4}{36} \times 9 \\ &\quad + \frac{3}{36} \times 10 + \frac{2}{36} \times 11 + \frac{1}{36} \times 12 \end{aligned}$$

$$\Rightarrow E(X) = \frac{1}{36} (2 + 6 + 12 + 20 + 30 + 42 + 40 + 36 + 30 + 22 + 12)$$

$$\Rightarrow E(X) = \frac{252}{36} = 7$$

31. $E(X) = \sum_i P(i)$
 $= 1\left(\frac{1}{15}\right) + 2\left(\frac{1}{15}\right) + \dots + 14\left(\frac{1}{15}\right) + 15\left(\frac{1}{15}\right)$
 $= \frac{1}{15} (1 + 2 + 3 + \dots + 14 + 15)$
 $= \frac{1}{15} \left(\frac{15 \times 16}{2}\right) \dots \left[\because \sum_{r=1}^n r = \frac{n(n+1)}{2}\right]$
 $= 8$

32.

X	1	2	3	...	n
P(X)	$\frac{2}{n(n+1)}$	$\frac{4}{n(n+1)}$	$\frac{6}{n(n+1)}$...	$\frac{2n}{n(n+1)}$

$$\begin{aligned} E(X) &= \sum_i P(i) \\ &= 1 \cdot \frac{2}{n(n+1)} + 2 \cdot \frac{4}{n(n+1)} + 3 \cdot \frac{6}{n(n+1)} \\ &\quad + \dots + n \cdot \frac{2n}{n(n+1)} \end{aligned}$$

$$\begin{aligned} &= \frac{2}{n(n+1)} (1 + 4 + 9 + \dots + n^2) \\ &= \frac{2}{n(n+1)} (1^2 + 2^2 + 3^2 + \dots + n^2) \\ &= \frac{2}{n(n+1)} \cdot \frac{n(n+1)(2n+1)}{6} \\ &= \frac{2n+1}{3} \end{aligned}$$

33. $E(X) = \sum_i P(i)$
 $= 1\left(\frac{1}{n}\right) + 2\left(\frac{1}{n}\right) + 3\left(\frac{1}{n}\right) + \dots + n\left(\frac{1}{n}\right)$
 $= \frac{1+2+3+\dots+n}{n}$
 $= \frac{1}{n} \times \frac{n(n+1)}{2}$
 $= \frac{n+1}{2}$

$$\begin{aligned} E(X^2) &= \sum_i P(i) \\ &= 1^2\left(\frac{1}{n}\right) + 2^2\left(\frac{1}{n}\right) + 3^2\left(\frac{1}{n}\right) + \dots + n^2\left(\frac{1}{n}\right) \\ &= \frac{1^2 + 2^2 + 3^2 + \dots + n^2}{n} \\ &= \frac{1}{n} \times \frac{n(n+1)(2n+1)}{6} \\ &= \frac{(n+1)(2n+1)}{6} \end{aligned}$$

$$\begin{aligned} \therefore \text{Var}(X) &= E(X^2) - [E(X)]^2 \\ &= \frac{(n+1)(2n+1)}{6} - \frac{(n+1)^2}{4} \\ &= \frac{n+1}{2} \left(\frac{2n+1}{3} - \frac{n+1}{2}\right) \\ &= \frac{n^2-1}{12} \end{aligned}$$

\therefore Standard deviation of X
 $= \sqrt{\text{Var}(X)} = \sqrt{\frac{n^2-1}{12}}$

34. Let X = demand for each type of cake (according to the profit)

$$P(X = 3) = 10\% = \frac{10}{100} = 0.1$$

$$P(X = 2.5) = 5\% = \frac{5}{100} = 0.05$$



$$P(X = 2) = 20\% = \frac{20}{100} = 0.2$$

$$P(X = 1.5) = 50\% = \frac{50}{100} = 0.5$$

$$P(X = 1) = 15\% = \frac{15}{100} = 0.15$$

∴ The probability distribution table is as follows:

X	3	2.5	2	1.5	1
P(X)	0.1	0.05	0.2	0.5	0.15

$$\begin{aligned} E(X) &= \sum x_i \cdot P(x_i) \\ &= 3(0.1) + 2.5(0.05) + 2(0.2) + 1.5(0.5) \\ &\quad + 1(0.15) \\ &= 0.3 + 0.125 + 0.4 + 0.75 + 0.15 = 1.725 \end{aligned}$$

35. Since, $f(x)$ is the p.d.f. of X

$$\therefore \int_{-\infty}^{\infty} f(x) dx = 1$$

$$\Rightarrow \int_0^3 C(9 - x^2) dx = 1$$

$$\Rightarrow C \left[9x - \frac{x^3}{3} \right]_0^3 = 1$$

$$\Rightarrow C(27 - 9) = 1 \Rightarrow C = \frac{1}{18}$$

36. Since, $f(x)$ is the p.d.f. of X .

$$\therefore \int_{-\infty}^{\infty} f(x) dx = 1$$

$$\Rightarrow \int_{-\infty}^0 f(x) dx + \int_0^1 f(x) dx + \int_1^{\infty} f(x) dx = 1$$

$$\Rightarrow 0 + \int_0^1 k^2(1-x) dx + 0 = 1$$

$$\Rightarrow k \left[\frac{x^3}{3} - \frac{x^4}{4} \right]_0^1 = 1 \Rightarrow k = 12$$

37. Since, $f(x)$ is the p.d.f. of X .

$$\therefore \int_{-\infty}^{\infty} f(x) dx = 1$$

$$\Rightarrow \int_0^3 \left(\frac{x}{6} + k \right) dx = 1$$

$$\Rightarrow \left[\frac{x^2}{12} + kx \right]_0^3 = 1 \Rightarrow \frac{3}{4} + 3k = 1$$

$$\Rightarrow 3k = \frac{1}{4} \Rightarrow k = \frac{1}{12}$$

$$\begin{aligned} 38. \quad P\left(\frac{1}{4} < X < \frac{1}{3}\right) &= \int_{\frac{1}{4}}^{\frac{1}{3}} f(x) dx = \int_{\frac{1}{4}}^{\frac{1}{3}} 3(1 - 2^{-2x}) dx \\ &= [3x - 2^{-2x}]_{1/4}^{1/3} \\ &= \left(1 - \frac{2}{27}\right) - \left(\frac{3}{4} - \frac{1}{32}\right) \\ &= \frac{1}{4} + \frac{1}{32} - \frac{2}{27} = \frac{179}{864} \end{aligned}$$

$$39. \quad \int_{-\infty}^{\infty} f(x) dx = 1$$

$$\therefore \int_0^4 \frac{K}{\sqrt{x}} dx = 1$$

$$\Rightarrow K [2\sqrt{x}]_0^4 = 1$$

$$\Rightarrow 2K [\sqrt{4} - \sqrt{0}] = 1$$

$$\Rightarrow 4K = 1$$

$$\Rightarrow K = \frac{1}{4}$$

$$\therefore P(X \geq 1) = P(1 \leq X < 4)$$

$$= \int_1^4 f(x) dx = 2K [\sqrt{x}]_1^4$$

$$= 2 \times \frac{1}{4} (2 - 1) = \frac{1}{2} = 0.5$$

$$40. \quad P(|X| < 1) = P(-1 < X < 1)$$

$$= \int_{-1}^1 \left(\frac{x+2}{18} \right) dx$$

$$= \frac{1}{18} \int_{-1}^1 (x+2) dx$$

$$= \frac{1}{18} \left[\frac{x^2}{2} + 2x \right]_{-1}^1$$

$$= \frac{1}{18} \left(\frac{5}{2} + 3 \right) = \frac{4}{18} = \frac{2}{9}$$

$$41. \quad P(0.2 \leq X \leq 0.5) = \int_{0.2}^{0.5} \frac{2}{8} dx = \left[\frac{x^3}{24} \right]_{0.2}^{0.5}$$

$$= \frac{1}{24} [(0.5)^3 - (0.2)^3]$$

$$= \frac{0.125 - 0.008}{24} = \frac{0.117}{24}$$



42. Since, $f(\cdot)$ is the p.d.f. of X .

$$\therefore \int_{-\infty}^{\infty} f(\cdot) d = 1$$

$$\therefore \int_0^{\infty} K.e^{-\theta} d = 1$$

$$\therefore K \left[\frac{e^{-\theta}}{-\theta} \right]_0^{\infty} = 1$$

$$\Rightarrow -\frac{K}{\theta} \left[\frac{1}{e^{\theta}} \right]_0^{\infty} = 1$$

$$\Rightarrow -\frac{K}{\theta} \left[\frac{1}{e^{\infty}} - \frac{1}{e^0} \right] = 1$$

$$\Rightarrow -\frac{K}{\theta} \left[\frac{1}{\infty} - \frac{1}{1} \right] = 1$$

$$\Rightarrow \frac{K}{\theta} = 1 \Rightarrow K = \theta$$

43. $P(0 < X < K) = 0.5$

$$\Rightarrow \int_0^K f(\cdot) d = \frac{1}{2}$$

$$\Rightarrow \int_0^K ae^{-a} d = \frac{1}{2}$$

$$\Rightarrow a \left[\frac{e^{-a}}{-a} \right]_0^K = \frac{1}{2}$$

$$\Rightarrow - \left[e^{-a} \right]_0^K = \frac{1}{2}$$

$$\Rightarrow -(e^{-aK} - e^0) = \frac{1}{2}$$

$$\Rightarrow -e^{-aK} + 1 = \frac{1}{2}$$

$$\Rightarrow e^{-aK} = \frac{1}{2}$$

$$\Rightarrow -aK = \log \left(\frac{1}{2} \right)$$

$$\Rightarrow aK = \log 2$$

$$\Rightarrow K = \frac{1}{a} \log 2$$

44. Since, $f_X(\cdot)$ is the p.d.f. of X .

$$\therefore \int_0^4 \frac{k}{\sqrt{d}} d = 1$$

$$\Rightarrow k \left[2\sqrt{d} \right]_0^4 = 1$$

$$\Rightarrow k = \frac{1}{4} \quad \dots(i)$$

$$\begin{aligned} F(\cdot) &= \int_0^{\cdot} \frac{k}{\sqrt{d}} d \\ &= k \left[2\sqrt{d} \right]_0^{\cdot} \\ &= \frac{\sqrt{d}}{2} \quad \dots[\text{From (i)}] \end{aligned}$$

$$\begin{aligned} 45. \quad P(C_1 \cup C_2) &= P(C_1) + P(C_2) \\ &= \int_1^2 f(\cdot) d + \int_4^5 f(\cdot) d \\ &= \int_1^2 \frac{1}{2} d + \int_4^5 \frac{1}{2} d \\ &= \left[\frac{-1}{2} \right]_1^2 + \left[\frac{-1}{2} \right]_4^5 \\ &= -\frac{1}{2} + 1 - \frac{1}{5} + \frac{1}{4} \\ &= \frac{11}{20} \end{aligned}$$

46. Since, $f(\cdot)$ is the p.d.f. of X .

$$\therefore \int_0^2 f(\cdot) d = 1$$

$$\Rightarrow \int_0^2 (k^2) d = 1$$

$$\Rightarrow k \left[\frac{d^3}{3} \right]_0^2 = 1$$

$$\Rightarrow k = \frac{3}{8}$$

$$\begin{aligned} \therefore \text{Required probability} &= P(X \leq 1) = \int_0^1 f(\cdot) d \\ &= \int_0^1 (k^2) d \\ &= \frac{3}{8} \int_0^1 d^2 d \\ &= \frac{3}{8} \left[\frac{d^3}{3} \right]_0^1 \\ &= \frac{3}{8} \left(\frac{1}{3} - 0 \right) \\ &= \frac{1}{8} \end{aligned}$$



Competitive Thinking

1. Since, $\sum_{i=1}^3 P(X=i) = 1$

$$\begin{aligned} \therefore 0.3 + k + 2k + 2k &= 1 \\ \Rightarrow 5k &= 0.7 \\ \Rightarrow k &= 0.14 \end{aligned}$$

2. Since, $\sum_{i=1}^6 P(X=i) = 1$

$$\begin{aligned} \therefore 0.1 + 2k + k + 0.2 + 3k + 0.1 &= 1 \\ \therefore 6k &= 1 - 0.4 = 0.6 \\ \therefore k &= \frac{0.6}{6} = 0.1 \end{aligned}$$

3. When we get 1, positive divisors = 1
 When we get 2, positive divisors = 2
 When we get 3, positive divisors = 2
 When we get 4, positive divisors = 3
 When we get 5, positive divisors = 2
 When we get 6, positive divisors = 4
 \therefore range of random variable $X = \{1, 2, 3, 4\}$

4. When a coin is tossed 3 times possibilities are

	HHH	TTT	HHT	HTH
Absolute difference between Heads and Tails($X=i$)	$3 - 0 = 3$	$3 - 0 = 3$	$2 - 1 = 1$	$2 - 1 = 1$

	THH	HTT	TTH	THT
Absolute difference between Heads and Tails($X=i$)	$2 - 1 = 1$	$2 - 1 = 1$	$2 - 1 = 1$	$2 - 1 = 1$

$$\therefore P(X=1) = \frac{6}{8} = \frac{3}{4}$$

5.

$X = k$	0	1	2	3	4	5
$P(X = k)$	a	a	$\frac{3a}{4}$	$\frac{4a}{8}$	$\frac{5a}{16}$	$\frac{6a}{32}$

$$\text{Since, } \sum_{k=0}^5 P(X=k) = 1$$

$$\begin{aligned} \therefore a + a + \frac{3a}{4} + \frac{4a}{8} + \frac{5a}{16} + \frac{6a}{32} &= 1 \\ \Rightarrow \frac{15}{4}a &= 1 \Rightarrow a = \frac{4}{15} \end{aligned}$$

$$\begin{aligned} \text{Now, } P(X = \text{prime value}) &= P(X=2) + P(X=3) + P(X=5) \\ &= \frac{3a}{4} + \frac{4a}{8} + \frac{6a}{32} \\ &= \frac{23a}{16} \\ &= \frac{23}{16} \times \frac{4}{15} \\ &= \frac{23}{60} \end{aligned}$$

6. Mean = $(1)\left(\frac{1}{4}\right) + (2)\left(\frac{1}{8}\right) + (3)\left(\frac{5}{8}\right)$
 $= \frac{19}{8}$

7. Mean = $1\left(\frac{1}{6}\right) + 2\left(\frac{1}{3}\right) + 3\left(\frac{1}{3}\right) + 4\left(\frac{1}{6}\right)$
 $= \frac{1}{6} + \frac{2}{3} + \frac{3}{3} + \frac{2}{3}$
 $= \frac{1}{6} + \frac{7}{3}$
 $= \frac{15}{6}$
 $= \frac{5}{2}$

8. Since $\sum_{i=1}^6 P(X=i) = 1$

$$\begin{aligned} \therefore a + a + a + b + b + 0.3 &= 1 \\ \Rightarrow 3a + 2b &= 0.7 \quad \dots(i) \\ \text{Mean} &= a + 2a + 3a + 4b + 5b + 6(0.3) \\ \Rightarrow 4.2 &= 6a + 9b + 1.8 \\ \Rightarrow 6a + 9b &= 2.4 \quad \dots(ii) \end{aligned}$$

On solving (i) and (ii), we get
 $a = 0.1, b = 0.2$

9. $E(X) = 3 \times \frac{1}{3} + 4 \times \frac{1}{4} + 12 \times \frac{5}{12}$
 $= 7$

10. $= 2$

	0	1	2	3
	0	2	4	6
$P(i)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

$$\begin{aligned} \therefore \text{Expected gain} &= \sum_i P(i) \\ &= 0\left(\frac{1}{8}\right) + 2\left(\frac{3}{8}\right) + 4\left(\frac{3}{8}\right) + 6\left(\frac{1}{8}\right) = 3 \end{aligned}$$



11. $E(X) = 2(0.3) + 3(0.4) + 4(0.3)$
 $= 0.6 + 1.2 + 1.2 = 3$
 $\therefore \text{Var}(X) = E(X^2) - [E(X)]^2$
 $= 4(0.3) + 9(0.4) + 16(0.3) - (3)^2$
 $= 9.6 - 9 = 0.6$
12. $E(X) = \sum_i x_i \cdot P(x_i)$
 $= 0(0.1) + 1(0.4) + 2(0.3) + 3(0.2) + 4(0)$
 $= 0 + 0.4 + 0.6 + 0.6 + 0$
 $= 1.6$
 $\text{Variance} = \sum_i x_i^2 \cdot P(x_i) - [E(X)]^2$
 $= 0^2(0.1) + 1^2(0.4) + 2^2(0.3)$
 $+ 3^2(0.2) + 4^2(0) - 1.6^2$
 $= 0 + 0.4 + 1.2 + 1.8 - 2.56$
 $= 0.84$
13. $E(X) = \sum_i x_i P(x_i)$
 $= 0\left(\frac{25}{16}\right) + 1\left(\frac{5}{18}\right) + 2\left(\frac{1}{36}\right) = \frac{1}{3}$
 $V(X) = \sum_i x_i^2 P(x_i) - [E(X)]^2$
 $= (0)^2\left(\frac{25}{36}\right) + (1)^2\left(\frac{5}{18}\right) + (2)^2\left(\frac{1}{36}\right)$
 $- \left(\frac{1}{3}\right)^2$
 $= \frac{7}{18} - \frac{1}{9} = \frac{5}{18}$
 $\text{S.D.} = \sqrt{\text{var}(X)} = \sqrt{\frac{5}{18}} = \frac{1}{3}\sqrt{\frac{5}{2}}$
14. $E(X) = \sum_i x_i \cdot P(x_i) = -\frac{1}{3} + 0 + \frac{1}{6} + \frac{2}{3} = \frac{1}{2}$
 $\text{Var}(X) = E(X^2) - [E(X)]^2$
 $= \frac{(-1)^2}{3} + 0 + \frac{1^2}{6} + \frac{2^2}{3} - \left(\frac{1}{2}\right)^2$
 $= \frac{1}{3} + \frac{1}{6} + \frac{4}{3} - \frac{1}{4}$
 $= \frac{11}{6} - \frac{1}{4} = \frac{19}{12}$
 $\therefore 6E(X^2) - \text{Var}(X)$
 $= 6\left(\frac{1}{3} + 0 + \frac{1}{6} + \frac{4}{3}\right) - \frac{19}{12}$
 $= 11 - \frac{19}{12}$
 $= \frac{113}{12}$

15. Let $P(X = 3) = a$, then
 $P(X = 1) = \frac{a}{2}$, $P(X = 2) = \frac{a}{3}$ and $P(X = 4) = \frac{a}{5}$
 Since, $P(X = 1) + P(X = 2) + P(X = 3)$
 $+ P(X = 4) = 1$

$$\therefore \frac{a}{2} + \frac{a}{3} + a + \frac{a}{5} = 1$$

$$\Rightarrow a = \frac{30}{61}$$

Now,

X =	1	2	3	4
P(X =)	$\frac{a}{2}$	$\frac{a}{3}$	a	$\frac{a}{5}$

$$\text{Now, } \mu = \text{mean} = \frac{1}{2}a + \frac{2}{3}a + 3a + \frac{4}{5}a$$

$$= \frac{149}{30}a$$

$\sigma^2 = \text{variance}$

$$= \frac{1}{2}a + \frac{4}{3}a + 9a + \frac{16}{5}a - \left(\frac{149}{30}a\right)^2$$

$$= \frac{421}{30}a - \left(\frac{149}{30}a\right)^2$$

$$\text{Now, } \sigma^2 + \mu^2 = \frac{421}{30}a - \left(\frac{149}{30}a\right)^2 + \left(\frac{149}{30}a\right)^2$$

$$= \frac{421}{30} \times \frac{30}{61} = \frac{421}{61}$$

16. $\text{Var}(X) = \sigma^2 = 5^2 = 25$
 $\text{Var}(X) = E(X^2) - [E(X)]^2$
 $\Rightarrow 25 = E(X^2) - 10^2$
 $\Rightarrow E(X^2) = 125$
 $E\left(\frac{X-15}{5}\right)^2 = E\left(\frac{X^2 - 30X + 225}{25}\right)$
 $= \frac{1}{25} [E(X^2) - 30E(X) + 225]$
 $= \frac{1}{25} (125 - 300 + 225)$
 $= 2$

17. Let denote number of defective pens.
 can take the values 0, 1, 2.

$$P(X = 0) = \frac{{}^4C_2}{{}^6C_2} = \frac{2}{5}$$



$$P(X=1) = \frac{{}^2C_1 \times {}^4C_1}{{}^6C_2} = \frac{8}{15}$$

$$P(X=2) = \frac{{}^2C_2}{{}^6C_2} = \frac{1}{15}$$

X =	0	1	2
P()	$\frac{2}{5}$	$\frac{8}{15}$	$\frac{1}{15}$

$$\begin{aligned} E(X) &= \sum_i P(x_i) \\ &= 0\left(\frac{2}{5}\right) + 1\left(\frac{8}{15}\right) + 2\left(\frac{1}{15}\right) \\ &= \frac{10}{15} \\ &= \frac{2}{3} \end{aligned}$$

$$\begin{aligned} E(X^2) &= \sum_i x_i^2 P(x_i) \\ &= 0\left(\frac{2}{5}\right) + 1\left(\frac{8}{15}\right) + 4\left(\frac{1}{15}\right) \\ &= \frac{12}{15} = \frac{4}{5} \end{aligned}$$

$$\begin{aligned} \text{Standard deviation } (\sigma) &= \sqrt{E(X^2) - [E(X)]^2} \\ &= \sqrt{\frac{4}{5} - \frac{4}{9}} \\ &= \frac{4}{3\sqrt{5}} \end{aligned}$$

$$\begin{aligned} 19. \text{ Required probability} &= \int_0^4 f(x) dx \\ &= \int_0^4 \frac{1}{5} dx \\ &= \frac{1}{5} [x]_0^4 = \frac{4}{5} = 0.8 \end{aligned}$$

$$\begin{aligned} 20. \quad P(X=4) &= F(4) - F(3) = 0.62 - 0.48 = 0.14 \\ P(X=5) &= F(5) - F(4) = 0.85 - 0.62 = 0.23 \\ P(3 < X \leq 5) &= P(X=4) + P(X=5) \\ &= 0.14 + 0.23 = 0.37 \end{aligned}$$



Evaluation Test

1. Given,
 $P(X=3) = 2P(X=1)$ and $P(X=2) = 0.3$ (i)
 Now, mean = 1.3
 $\therefore 0 \times P(X=0) + 1 \times P(X=1) + 2 \times P(X=2) + 3 \times P(X=3) = 1.3$
 $\Rightarrow 7P(X=1) = 0.7$ [From (i)]
 $\Rightarrow P(X=1) = 0.1$
 Also, $P(X=0) + P(X=1) + P(X=2) + P(X=3) = 1$
 $\Rightarrow P(X=0) + 3P(X=1) = 0.7$ [From (i)]
 $\Rightarrow P(X=0) + 0.3 = 0.7$
 $\Rightarrow P(X=0) = 0.4$

2. $\sum_{x=0}^8 P(X=x) = 1$
 $\Rightarrow a + 3a + 5a + 7a + 9a + 11a + 13a + 15a + 17a = 1$
 $\Rightarrow 81a = 1$
 $\Rightarrow a = \frac{1}{81}$

3. $P(E) = P(X=2 \text{ or } X=3 \text{ or } X=5 \text{ or } X=7)$
 $= P(X=2) + P(X=3) + P(X=5) + P(X=7)$
 $= 0.23 + 0.12 + 0.20 + 0.07 = 0.62$
 $P(F) = P(X < 4)$
 $= P(X=1) + P(X=2) + P(X=3)$
 $= 0.15 + 0.23 + 0.12 = 0.50$
 $P(E \cap F) = P(X \text{ is a prime number less than } 4)$
 $= P(X=2) + P(X=3)$
 $= 0.23 + 0.12 = 0.35$
 $\therefore P(E \cup F) = P(E) + P(F) - P(E \cap F)$
 $= 0.62 + 0.50 - 0.35 = 0.77$

4. Here, $\frac{1+3p}{4}$, $\frac{1-p}{4}$, $\frac{1+2p}{4}$ and $\frac{1-4p}{4}$ are probabilities when X takes values -1, 0, 1 and 2 respectively. Therefore, each is greater than or equal to 0 and less than or equal to 1.
 i.e., $0 \leq \frac{1+3p}{4} \leq 1$, $0 \leq \frac{1-p}{4} \leq 1$,
 $0 \leq \frac{1+2p}{4} \leq 1$ and $0 \leq \frac{1-4p}{4} \leq 1$



$$\Rightarrow -\frac{1}{3} \leq p \leq \frac{1}{4}$$

$$\begin{aligned} \text{Mean}(X) &= -1 \times \frac{1+3p}{4} + 0 \times \frac{1-p}{4} + 1 \times \frac{1+2p}{4} \\ &\quad + 2 \times \frac{1-4p}{4} \\ &= \frac{2-9p}{4} \end{aligned}$$

$$\text{Now, } -\frac{1}{3} \leq p \leq \frac{1}{4}$$

$$\Rightarrow 3 \geq -9p \geq -\frac{9}{4}$$

$$\Rightarrow -\frac{1}{4} \leq 2-9p \leq 5$$

$$\Rightarrow -\frac{1}{16} \leq \frac{2-9p}{4} \leq \frac{5}{4}$$

$$5. \quad P(X > 1.5) = \int_{1.5}^2 \frac{1}{2} dx = \left[\frac{x}{2} \right]_{1.5}^2 = 0.4375$$

$$\text{and } P(X > 1) = \int_1^2 \frac{1}{2} dx = \left[\frac{x}{2} \right]_1^2 = 0.75$$

$$\therefore P\left(\frac{X > 1.5}{X > 1}\right) = \frac{P(X > 1.5)}{P(X > 1)} = \frac{0.4375}{0.75} = \frac{7}{12}$$

$$6. \quad P(X = i) = ki, \text{ where } 1 \leq i \leq 10$$

$$\therefore \sum P(X = i) = 1$$

$$\Rightarrow (1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 + 9 + 10)k = 1$$

$$\Rightarrow k = \frac{1}{55}$$

$$7. \quad \text{We have, } \sum_{n=0}^{\infty} P(X = n) = 1$$

$$\Rightarrow k \sum_{n=0}^{\infty} (n+1) \left(\frac{1}{5}\right)^n = 1$$

$$\Rightarrow k \left[1 + 2 \left(\frac{1}{5}\right) + 3 \left(\frac{1}{5}\right)^2 + 4 \left(\frac{1}{5}\right)^3 + \dots \right] = 1$$

$$\Rightarrow k \left[\frac{1}{1 - \frac{1}{5}} + \frac{1 \times \frac{1}{5}}{\left(1 - \frac{1}{5}\right)^2} \right] = 1$$

$$\dots \left[\begin{aligned} &\because a + (a+d)r + (a+2d)r^2 + \dots \\ &= \frac{a}{1-r} + \frac{dr}{(1-r)^2} \end{aligned} \right]$$

$$\Rightarrow k \left(\frac{5}{4} + \frac{5}{16} \right) = 1$$

$$\Rightarrow \frac{25k}{16} = 1$$

$$\Rightarrow k = \frac{16}{25}$$



Hints



Classical Thinking

2. $P(X = 1) = {}^{10}C_1 (0.2) (0.8)^9 = 0.2684$
3. Probability of getting head is $p = \frac{1}{2}$
 $\therefore q = 1 - \frac{1}{2} = \frac{1}{2}$
 Also, $n = 4$
 \therefore Required probability = $P(X = 3)$
 $= {}^4C_3 \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right) = \frac{1}{4}$
4. Here $p = \frac{1}{2}$, $q = \frac{1}{2}$, $n = 10$
 \therefore Required probability = $P(X = 5)$
 $= {}^{10}C_5 \left(\frac{1}{2}\right)^5 \cdot \left(\frac{1}{2}\right)^5 = \frac{63}{256}$
5. Probability of obtaining 5 is $p = \frac{1}{6}$
 $\therefore q = 1 - \frac{1}{6} = \frac{5}{6}$
 Also, $n = 7$
 \therefore Required probability = $P(X = 4)$
 $= {}^7C_4 \left(\frac{1}{6}\right)^4 \left(\frac{5}{6}\right)^3$
6. Probability of getting on even number is
 $p = \frac{3}{6} = \frac{1}{2}$
 $\therefore q = 1 - \frac{1}{2} = \frac{1}{2}$ and $n = 5$, $r = 3$
 \therefore Required probability
 $= {}^5C_3 \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^2 = \frac{5}{16}$
7. Probability of getting an odd number, $p = \frac{3}{6} = \frac{1}{2}$
 $\therefore q = 1 - \frac{1}{2} = \frac{1}{2}$

- Also, $n = 2$
 \therefore Required probability = $P(X = 2)$
 $= {}^2C_2 \left(\frac{1}{2}\right)^2 \left(\frac{1}{2}\right)^0 = \frac{1}{4}$
8. Here, $p = \frac{1}{2}$, $q = \frac{1}{2}$, $n = 3$
 \therefore Required probability = $P(X \geq 2)$
 $= {}^3C_2 \left(\frac{1}{2}\right)^2 \left(\frac{1}{2}\right) + {}^3C_3 \left(\frac{1}{2}\right)^3$
 $= \frac{4}{8} = \frac{1}{2}$
 9. Probability of getting an odd number,
 $p = \frac{3}{6} = \frac{1}{2}$
 $\therefore q = 1 - \frac{1}{2} = \frac{1}{2}$ Also, $n = 5$
 \therefore Variance = $npq = 5 \cdot \left(\frac{1}{2}\right) \cdot \left(\frac{1}{2}\right) = \frac{5}{4}$



Critical Thinking

1. Here, $q = \frac{1}{5}$
 $\Rightarrow p = 1 - \frac{1}{5} = \frac{4}{5}$
 Also, $n = 5$
 \therefore Required probability = ${}^5C_1 \left(\frac{4}{5}\right) \left(\frac{1}{5}\right)^4$
 {Here exactly one student is swimmer}
2. Probability of success is $p = \frac{3}{5}$
 $\Rightarrow q = 1 - p = \frac{2}{5}$
 Also, $n = 5$
 \therefore Required probability = $P(X = 2)$
 $= {}^5C_2 \left(\frac{3}{5}\right)^2 \left(\frac{2}{5}\right)^3 = \frac{144}{625}$



3. Probability that bulb will fuse, $p = 0.05$

$$= \frac{1}{20}$$

\therefore Probability that bulb will not fuse,

$$q = 1 - p = 1 - \frac{1}{20} = \frac{19}{20}$$

Also, $n = 5$

\therefore Probability that out of 5 bulbs none will fuse

$$= {}^5C_0 \left(\frac{1}{20}\right)^0 \left(\frac{19}{20}\right)^5 = \left(\frac{19}{20}\right)^5$$

4. Probability of correct prediction,

$$p = \frac{1}{3} \Rightarrow q = 1 - \frac{1}{3} = \frac{2}{3}$$

Also, $n = 7$

\therefore Required probability = $P(X = 4)$

$$= {}^7C_4 \left(\frac{1}{3}\right)^4 \left(\frac{2}{3}\right)^3 = \frac{280}{3^7}$$

5. Here, $p = \frac{1}{2}$, $q = \frac{1}{2}$, $n = 3$

\therefore Required probability

= Probability of getting exactly one head +
 probability of getting exactly two heads

$$= {}^3C_1 \left(\frac{1}{2}\right)^1 \cdot \left(\frac{1}{2}\right)^2 + {}^3C_2 \left(\frac{1}{2}\right)^2 \cdot \left(\frac{1}{2}\right)^1$$

$$= \frac{3}{8} + \frac{3}{8} = \frac{6}{8} = \frac{3}{4}$$

6. Here, $p = \frac{1}{2}$, $q = \frac{1}{2}$, $n = 10$

\therefore Required probability = $P(X = 4)$

$$= {}^{10}C_4 \left(\frac{1}{2}\right)^4 \left(\frac{1}{2}\right)^6 = {}^{10}C_4 \left(\frac{1}{2}\right)^{10}$$

$$= {}^{10}C_6 \left(\frac{1}{2}\right)^{10} \quad \dots [\because {}^nC_r = {}^nC_{n-r}]$$

7. $9 P(X = 4) = P(X = 2)$

$$\therefore 9 \cdot {}^6C_4 p^4 q^2 = {}^6C_2 p^2 q^4 \Rightarrow 9 p^2 = q^2$$

Putting $q = 1 - p$, we get

$$p = \frac{1}{4}$$

8. Required probability

= $P(\text{exactly two success})$
 + $P(\text{exactly three success})$

$$= {}^3C_2 \cdot \left(\frac{2}{6}\right)^2 \left(\frac{4}{6}\right) + {}^3C_3 \left(\frac{2}{6}\right)^3$$

$$= \frac{2}{9} + \frac{1}{27} = \frac{7}{27}$$

9. We have, $p = \frac{3}{4}$

$$\Rightarrow q = \frac{1}{4} \text{ and } n = 5$$

\therefore Required probability = $P(X \geq 3)$

$$= {}^5C_3 \left(\frac{3}{4}\right)^3 \left(\frac{1}{4}\right)^2 + {}^5C_4 \left(\frac{3}{4}\right)^4 \left(\frac{1}{4}\right) + {}^5C_5 \left(\frac{3}{4}\right)^5$$

$$= \frac{(10)(27)}{4^5} + \frac{(5)(81)}{4^5} + \frac{243}{4^5}$$

$$= \frac{270 + 405 + 243}{1024}$$

$$= \frac{459}{512}$$

10. Required probability = $P(X \geq 1)$

$$= {}^3C_1 \left(\frac{1}{6}\right) \left(\frac{5}{6}\right)^2 + {}^3C_2 \left(\frac{1}{6}\right)^2 \left(\frac{5}{6}\right) + {}^3C_3 \left(\frac{1}{6}\right)^3 \left(\frac{5}{6}\right)^0$$

$$= \frac{91}{216}$$

11. Required probability = $P(X \geq 6)$

$$= {}^8C_6 \left(\frac{1}{2}\right)^6 \left(\frac{1}{2}\right)^2 + {}^8C_7 \left(\frac{1}{2}\right)^7 \left(\frac{1}{2}\right) + {}^8C_8 \left(\frac{1}{2}\right)^8$$

$$= \frac{37}{256}$$

12. Let the coin be tossed n times.

$$\text{Then, } P(7 \text{ heads}) = {}^nC_7 \left(\frac{1}{2}\right)^7 \left(\frac{1}{2}\right)^{n-7} = {}^nC_7 \left(\frac{1}{2}\right)^n$$

$$\text{and } P(9 \text{ heads}) = {}^nC_9 \left(\frac{1}{2}\right)^9 \left(\frac{1}{2}\right)^{n-9} = {}^nC_9 \left(\frac{1}{2}\right)^n$$

Now, $P(7 \text{ heads}) = P(9 \text{ heads})$

$$\Rightarrow {}^nC_7 = {}^nC_9$$

$$\Rightarrow n = 16$$

$$\therefore P(3 \text{ heads}) = {}^{16}C_3 \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^{16-3}$$

$$= {}^{16}C_3 \left(\frac{1}{2}\right)^{16} = \frac{35}{2^{12}}$$

14. Here, $n = 3$, $p = \frac{1}{6}$, $q = \frac{5}{6}$

$$\text{Mean} = np = 3 \times \frac{1}{6} = \frac{1}{2}$$

$$\text{Variance} = npq = 3 \times \frac{1}{6} \times \frac{5}{6} = \frac{5}{12}$$



$$15. \text{ Here, } p = \frac{2}{6} = \frac{1}{3} \Rightarrow q = 1 - \frac{1}{3} = \frac{2}{3}$$

$$\text{Also, } n = 2$$

$$\therefore \text{ Variance} = npq = 2 \times \frac{1}{3} \times \frac{2}{3} = \frac{4}{9}$$

$$16. \text{ We have, mean} = np = 2$$

$$\text{and variance} = npq = 1$$

$$\Rightarrow q = \frac{1}{2}, p = \frac{1}{2} \text{ and } n = 4$$

$$\therefore P(X \geq 1) = 1 - P(X = 0)$$

$$= 1 - {}^4C_0 \left(\frac{1}{2}\right)^4$$

$$= \frac{15}{16}$$

$$17. \text{ Here, } np = 4 \text{ and } npq = 3$$

$$\Rightarrow p = \frac{1}{4}, q = \frac{3}{4}$$

$$\text{Also, } n = 16$$

$$\therefore P(X = 6) = {}^{16}C_6 \left(\frac{1}{4}\right)^6 \left(\frac{3}{4}\right)^{10}$$

$$18. \text{ Probability of getting a red card is}$$

$$p = \frac{26}{52} = \frac{1}{2}$$

$$\therefore q = 1 - \frac{1}{2} = \frac{1}{2} \text{ Also, } n = 4$$

$$\text{Mean} = np = 4 \left(\frac{1}{2}\right) = 2$$

$$\text{Variance} = npq = 4 \left(\frac{1}{2}\right) \left(\frac{1}{2}\right) = 1$$

$$19. \frac{P(X = k)}{P(X = k - 1)} = \frac{{}^n C_k (p)^k (q)^{n-k}}{{}^n C_{k-1} (p)^{k-1} (q)^{n-k+1}}$$

$$= \frac{{}^n C_k}{{}^n C_{k-1}} \cdot \frac{p}{q}$$

$$\therefore \frac{P(X = k)}{P(X = k - 1)} = \frac{n - k + 1}{k} \cdot \frac{p}{q}$$

$$20. \text{ Let } X \text{ denote the number of aces obtained in two draws. Then, } X \text{ follows binomial}$$

$$\text{distribution with } n = 2, p = \frac{4}{52} = \frac{1}{13} \text{ and}$$

$$q = \frac{12}{13}$$

$$\therefore \text{ Mean of number of aces} = np = \frac{2}{13}$$

**Competitive Thinking**

$$1. \text{ Here, } n = 5, p = \frac{1}{3}$$

$$\text{and } q = 1 - \frac{1}{3} = \frac{2}{3}$$

$$\therefore P(2 < X < 4) = P(X = 3)$$

$$= {}^5C_3 \left(\frac{1}{3}\right)^3 \left(\frac{2}{3}\right)^2 = \frac{40}{243}$$

$$2. \text{ Probability of getting head, } p = \frac{1}{2}$$

$$\therefore q = 1 - p = 1 - \frac{1}{2} = \frac{1}{2}$$

$$\text{Also, } n = 10$$

$$\therefore \text{ Required probability} = P(X = 6)$$

$$= {}^{10}C_6 \left(\frac{1}{2}\right)^6 \left(\frac{1}{2}\right)^4$$

$$= \frac{10!}{6!4!} \cdot \frac{1}{2^{10}} = \frac{105}{512}$$

$$3. \text{ Probability of occurrence of '4' is } p = \frac{1}{6}$$

$$\therefore q = 1 - \frac{1}{6} = \frac{5}{6}$$

$$\text{Also, } n = 2,$$

$$\therefore \text{ Required probability} = P(X \geq 1)$$

$$= {}^2C_1 \left(\frac{1}{6}\right) \left(\frac{5}{6}\right) + {}^2C_2 \left(\frac{1}{6}\right)^2 \left(\frac{5}{6}\right)^0$$

$$= \frac{11}{36}$$

$$4. \text{ Probability that person will develop immunity (p) = 0.8}$$

$$q = 1 - p = 0.2$$

$$\therefore \text{ Required probability} = {}^8C_0 (0.8)^8 (0.2)^0 = (0.8)^8$$

$$5. \text{ Probability of getting rotten egg is}$$

$$p = \frac{10}{100} = \frac{1}{10}$$

$$\therefore q = 1 - \frac{1}{10} = \frac{9}{10}$$

$$\text{Also, } n = 5$$

$$\therefore \text{ The probability that no egg is rotten}$$

$$= {}^5C_0 \cdot \left(\frac{1}{10}\right)^0 \left(\frac{9}{10}\right)^5 = \left(\frac{9}{10}\right)^5$$



6. Probability of disease to be fatal = $p = 10\%$

$$p = \frac{10}{100} = \frac{1}{10}, q = \frac{9}{10}$$

Number of patients, $n = 6$

$$\begin{aligned} \therefore \text{Required probability} &= {}^6C_3 \left(\frac{1}{10}\right)^3 \left(\frac{9}{10}\right)^3 \\ &= 1458 \times 10^{-5} \end{aligned}$$

7. Probability of getting a 'six' in one throw is

$$p = \frac{1}{6}$$

$$\therefore q = 1 - \frac{1}{6} = \frac{5}{6}$$

Also, $n = 4$

\therefore Required probability

$$\begin{aligned} = P(X = 4) &= {}^4C_4 \left(\frac{1}{6}\right)^4 \cdot \left(\frac{5}{6}\right)^0 \\ &= \frac{1}{1296} \end{aligned}$$

8. $P(\text{without defect}) = \frac{8}{10} = \frac{4}{5} = p$

$$P(\text{defected}) = \frac{2}{10} = \frac{1}{5} = q \text{ and } n = 2, r = 2$$

$$\therefore \text{Required probability} = {}^2C_2 \left(\frac{4}{5}\right)^2 \cdot \left(\frac{1}{5}\right)^0 = \frac{16}{25}$$

9. $2P(2) = 3P(3)$

$$\Rightarrow 2 \cdot {}^6C_2 p^2 q^4 = 3 \cdot {}^6C_3 p^3 q^3$$

Putting $q = 1 - p$, we get

$$p = \frac{1}{3}$$

10. $4P(X = 4) = P(X = 2)$

$$\Rightarrow 4 \cdot {}^6C_4 p^4 q^2 = {}^6C_2 p^2 q^4$$

$$\Rightarrow 4p^2 = q^2$$

$$\Rightarrow 4p^2 = (1 - p)^2$$

$$\Rightarrow 3p^2 + 2p - 1 = 0$$

$$\Rightarrow p = \frac{1}{3}$$

11. Here, $p = q = \frac{1}{2}$

Probability that head occurs 6 times

$$= {}^nC_6 \left(\frac{1}{2}\right)^6 \left(\frac{1}{2}\right)^{n-6} \text{ and probability that head}$$

$$\text{occurs 8 times} = {}^nC_8 \left(\frac{1}{2}\right)^8 \left(\frac{1}{2}\right)^{n-8}$$

$$\therefore {}^nC_6 \left(\frac{1}{2}\right)^6 \left(\frac{1}{2}\right)^{n-6} = {}^nC_8 \left(\frac{1}{2}\right)^8 \left(\frac{1}{2}\right)^{n-8}$$

$$\Rightarrow {}^nC_6 = {}^nC_8 \Rightarrow (n-6)(n-7) = 56 \Rightarrow n = 14$$

12. We have, ${}^{100}C_{50} p^{50} (1-p)^{50} = {}^{100}C_{51} p^{51} (1-p)^{49}$

$$\Rightarrow \frac{1-p}{p} = \frac{100!}{51! \cdot 49!} \times \frac{50! \cdot 50!}{100!}$$

$$= \frac{50}{51}$$

$$\Rightarrow 51 - 51p = 50p \Rightarrow p = \frac{51}{101}$$

13. The required probability

= 1 - Probability of equal number of heads and tails

$$= 1 - {}^{2n}C_n \left(\frac{1}{2}\right)^n \left(\frac{1}{2}\right)^{2n-n}$$

$$= 1 - \frac{(2n)!}{n!n!} \left(\frac{1}{4}\right)^n$$

$$= 1 - \frac{(2n)!}{(n!)^2} \times \frac{1}{4^n}$$

14. Probability of failure, $q = \frac{1}{3}$

Probability for getting success, $p = 1 - \frac{1}{3} = \frac{2}{3}$

Also, $n = 4$

\therefore Required probability = $P(X \geq 3)$

$$= {}^4C_4 \left(\frac{2}{3}\right)^4 \left(\frac{1}{3}\right)^0 + {}^4C_3 \left(\frac{2}{3}\right)^3 \left(\frac{1}{3}\right)$$

$$= \left(\frac{2}{3}\right)^4 + 4 \left(\frac{2}{3}\right)^3 \left(\frac{1}{3}\right)$$

$$= \frac{16}{27}$$

15. Probability for white ball, $p = \frac{2}{6} = \frac{1}{3}$

Probability for black ball, $q = \frac{4}{6} = \frac{2}{3}$

Also, $n = 5$

\therefore Required probability = $P(X \geq 4)$

$$= {}^5C_5 \left(\frac{1}{3}\right)^5 \left(\frac{2}{3}\right)^0 + {}^5C_4 \left(\frac{1}{3}\right)^4 \left(\frac{2}{3}\right)$$

$$= \left(\frac{1}{3}\right)^4 \left[\frac{1}{3} + (5) \frac{2}{3} \right]$$

$$= \frac{11}{3^5} = \frac{11}{243}$$



$$16. \text{ Required probability} = P(X < 2)$$

$$= {}^8C_1 \left(\frac{1}{20}\right)^1 \left(\frac{19}{20}\right)^7 + {}^8C_0 \left(\frac{1}{20}\right)^0 \left(\frac{19}{20}\right)^8$$

$$= \frac{27}{20} \left(\frac{19}{20}\right)^7$$

$$17. P(\text{minimum face value not less than 2 and maximum face value is not greater than 5})$$

$$= P(2 \text{ or } 3 \text{ or } 4 \text{ or } 5)$$

$$= \frac{4}{6} = \frac{2}{3}$$

$$\therefore \text{ required probability} = {}^4C_4 \left(\frac{2}{3}\right)^4 \left(\frac{1}{3}\right)^0 = \frac{16}{81}$$

$$18. \text{ Here, } p = \text{probability of getting perfect square in any throw} = \frac{2}{6} = \frac{1}{3}$$

$$\therefore q = \frac{2}{3} \text{ and } n = 4$$

Now,

$$P(\text{getting perfect square in at least one throw})$$

$$= 1 - P(\text{not getting perfect square in any throw})$$

$$\Rightarrow P(X \geq 1) = 1 - P(X = 0)$$

$$= 1 - {}^4C_0 \left(\frac{1}{3}\right)^0 \left(\frac{2}{3}\right)^4$$

$$= 1 - \left(\frac{2}{3}\right)^4 = \frac{65}{81}$$

$$19. P(\text{answer is correct}) = p = \frac{1}{2}$$

$$\therefore q = 1 - \frac{1}{2} = \frac{1}{2}$$

Also, $n = 10$

$$\therefore P(\text{at least 7 answers are correct}) = P(X \geq 7)$$

$$= P(X = 7) + P(X = 8) + P(X = 9) + P(X = 10)$$

$$= {}^{10}C_7 \left(\frac{1}{2}\right)^7 \left(\frac{1}{2}\right)^3 + {}^{10}C_8 \left(\frac{1}{2}\right)^8 \left(\frac{1}{2}\right)^2$$

$$+ {}^{10}C_9 \left(\frac{1}{2}\right)^9 \left(\frac{1}{2}\right) + {}^{10}C_{10} \left(\frac{1}{2}\right)^{10}$$

$$= \left({}^{10}C_7 + {}^{10}C_8 + {}^{10}C_9 + {}^{10}C_{10}\right) \frac{1}{2^{10}}$$

$$= (120 + 45 + 10 + 1) \frac{1}{1024}$$

$$= \frac{176}{1024}$$

$$= \frac{11}{64}$$

$$20. \text{ Probability of green ball } (p) = \frac{15}{25} = \frac{3}{5}$$

$$\text{Probability of yellow ball } (q) = \frac{10}{25} = \frac{2}{5}$$

Also, $n = 10$

$$\therefore \text{ Variance} = npq$$

$$= 10 \left(\frac{3}{5}\right) \left(\frac{2}{5}\right)$$

$$= \frac{12}{5}$$

$$21. \text{ Probability of occurrence of event } A \text{ is}$$

$$p = 0.3$$

$$\therefore q = 0.7$$

Also, $n = 6$

$$\therefore \text{ Variance} = npq$$

$$= 6 \times 0.3 \times 0.7 = 1.26$$

$$22. n = 10, p = 0.4$$

$$E(X) = np = 4$$

$$V(X) = npq = 10 \times 0.4 \times 0.6 = 2.4$$

$$V(X) = E(X^2) - [E(X)]^2$$

$$\therefore 2.4 = E(X^2) - [4]^2$$

$$\Rightarrow E(X^2) = 18.4$$

$$23. \text{ Given } np = 6, npq = 4$$

$$\therefore \frac{npq}{np} = \frac{4}{6}$$

$$\Rightarrow q = \frac{2}{3} \text{ and } p = \frac{1}{3}$$

$$\therefore np = 6$$

$$\Rightarrow n \times \frac{1}{3} = 6$$

$$\Rightarrow n = 18$$

$$24. \text{ Mean} = np = 18$$

$$\text{Variance} = npq = 12$$

$$\therefore \frac{npq}{np} = \frac{12}{18} \Rightarrow q = \frac{2}{3}$$

$$p = 1 - q = 1 - \frac{2}{3} = \frac{1}{3}$$

Now, $np = 18$

$$\Rightarrow n \left(\frac{1}{3}\right) = 18$$

$$\Rightarrow n = 54$$

\therefore Values of are 0, 1, 2, 3, ..., 54 = 55 values



$$25. \quad \left. \begin{array}{l} np = 4 \\ npq = 2 \end{array} \right\} \Rightarrow q = \frac{1}{2}, p = \frac{1}{2}, n = 8$$

$$\begin{aligned} \therefore P(X=1) &= {}^8C_1 \left(\frac{1}{2}\right) \left(\frac{1}{2}\right)^7 \\ &= 8 \cdot \frac{1}{2^8} = \frac{1}{2^5} = \frac{1}{32} \end{aligned}$$

$$26. \quad \left. \begin{array}{l} np = 8 \\ npq = 4 \end{array} \right\} \Rightarrow q = \frac{1}{2}, p = \frac{1}{2}, n = 16$$

$$\begin{aligned} \therefore P(X=1) &= {}^{16}C_1 \left(\frac{1}{2}\right) \left(\frac{1}{2}\right)^{15} \\ &= 16 \times \frac{1}{2} \times \frac{1}{2^{15}} \\ &= 2^4 \times \frac{1}{2} \times \frac{1}{2^{15}} = \frac{1}{2^{12}} \end{aligned}$$

$$27. \quad \left. \begin{array}{l} np = 4 \\ npq = 2 \end{array} \right\} \Rightarrow q = \frac{1}{2}, p = \frac{1}{2}, n = 8$$

$$\begin{aligned} \therefore P(X=2) &= {}^8C_2 \left(\frac{1}{2}\right)^2 \left(\frac{1}{2}\right)^6 \\ &= 28 \cdot \frac{1}{2^8} = \frac{28}{256} \end{aligned}$$

$$\begin{aligned} 28. \quad E(X) &= 5 \text{ and } \text{Var}(X) = 2.5 \\ \Rightarrow np &= 5 \text{ and } npq = 2.5 \\ \Rightarrow p &= \frac{1}{2}, q = \frac{1}{2} \text{ and } n = 10 \end{aligned}$$

$$\begin{aligned} \therefore P(X < 1) &= P(X = 0) \\ &= {}^{10}C_0 \left(\frac{1}{2}\right)^0 \left(\frac{1}{2}\right)^{10} = \left(\frac{1}{2}\right)^{10} \end{aligned}$$

$$\begin{aligned} 29. \quad E(X) &= 6 \text{ and } V(X) = 2 \\ \Rightarrow np &= 6 \text{ and } npq = 2 \\ \Rightarrow q &= \frac{1}{3}, p = \frac{2}{3} \text{ and } n = 9 \end{aligned}$$

$$\begin{aligned} P(5 \leq X \leq 7) &= P(X=5) + P(X=6) + P(X=7) \\ &= {}^9C_5 \left(\frac{2}{3}\right)^5 \left(\frac{1}{3}\right)^4 + {}^9C_6 \left(\frac{2}{3}\right)^6 \left(\frac{1}{3}\right)^3 \\ &\quad + {}^9C_7 \left(\frac{2}{3}\right)^7 \left(\frac{1}{3}\right)^2 \\ &= \frac{2^5}{3^9} [{}^9C_5 + {}^9C_6 \times 2 + {}^9C_7 \times 4] \\ &= \frac{2^5}{3^9} [126 + 168 + 144] \\ &= \frac{2^5 \times 438}{3^9} = \frac{2^5 \times 146}{3^9} = \frac{4672}{6561} \end{aligned}$$

$$30. \quad \text{Probability of getting a success, } p = \frac{1}{4}$$

$$\text{Probability of not getting success, } q = \frac{3}{4}$$

$$\text{Standard deviation} = \sqrt{\text{Variance}}$$

$$\Rightarrow \text{Variance} = 9$$

$$\Rightarrow npq = 9 \Rightarrow n \cdot \frac{1}{4} \cdot \frac{3}{4} = 9 \Rightarrow n = 48$$

$$\text{Mean} = np = \frac{1}{4} \times 48 = 12$$

$$31. \quad \text{Let } X = \text{Number of heads appear in } n \text{ tosses}$$

$$X \sim B\left(n, \frac{1}{2}\right)$$

$$\text{Now, } P(X \geq 1) = 1 - P(X = 0) = 1 - \frac{1}{2^n}$$

$$\text{Since, } P(X \geq 1) \geq 0.9$$

$$\therefore 1 - \frac{1}{2^n} \geq 0.9$$

$$\Rightarrow \frac{1}{2^n} \leq \frac{1}{10} \Rightarrow 2^n \geq 10 \Rightarrow n \geq 4$$

$$\therefore \text{minimum number of tosses} = 4$$



Evaluation Test

$$1. \quad \text{We have, } p = \frac{10}{100} = \frac{1}{10}$$

$$\therefore q = 1 - \frac{1}{10} = \frac{9}{10}$$

According to the given condition,

$$P(X \geq 1) \geq \frac{50}{100}$$

$$\Rightarrow 1 - P(X = 0) \geq \frac{1}{2}$$

$$\Rightarrow P(X = 0) \leq \frac{1}{2}$$

$$\Rightarrow \left(\frac{9}{10}\right)^n \leq \frac{1}{2},$$

which is possible if n is at least 7.

$$\therefore n = 7$$



2. $P(X = 1) = 8 \cdot P(X = 3)$, if $n = 5$

$$\Rightarrow {}^5C_1 q^4 p^1 = 8 \cdot {}^5C_3 q^2 p^3$$

$$\Rightarrow \frac{5q^2}{p^2} = 8(10) \Rightarrow \frac{q^2}{p^2} = 16$$

$$\Rightarrow q = 4p$$

$$\Rightarrow 1 - p = 4p$$

$$\Rightarrow 5p = 1 \Rightarrow p = \frac{1}{5}$$

3. $P(X = 0) = \frac{16}{81}$

$$\Rightarrow {}^4C_0 p^0 q^4 = \frac{16}{81} \Rightarrow q^4 = \left(\frac{2}{3}\right)^4$$

$$\Rightarrow q = \frac{2}{3} \Rightarrow p = \frac{1}{3}$$

$$\therefore P(X = 4) = {}^4C_4 p^4 q^0 = p^4 = \left(\frac{1}{3}\right)^4 = \frac{1}{81}$$

4. Here $p = \frac{3}{6} = \frac{1}{2}$

$$\text{and } q = 1 - p = 1 - \frac{1}{2} = \frac{1}{2}$$

$$n = 100$$

$$\therefore \text{variance} = npq = 100 \times \frac{1}{2} \times \frac{1}{2} = 25$$

5. Here $n = 8$,

$$p = \text{Probability of getting 1 or 3} = \frac{2}{6} = \frac{1}{3}$$

$$\therefore q = 1 - \frac{1}{3} = \frac{2}{3}$$

$$\therefore \text{S.D.} = \sqrt{npq} = \sqrt{8 \times \frac{1}{3} \times \frac{2}{3}} = \sqrt{\frac{16}{9}} = \frac{4}{3}$$

6. Let X denote the number of failures in 5 trials.

$$\text{Then, } P(X = r) = {}^5C_r (1 - p)^r p^{5-r}; r = 0, 1, 2, \dots, 5$$

$$\therefore P(X \geq 1) \geq \frac{31}{32}$$

$$\Rightarrow 1 - P(X = 0) \geq \frac{31}{32}$$

$$\Rightarrow 1 - p^5 \geq \frac{31}{32}$$

$$\Rightarrow \frac{1}{32} \geq p^5$$

$$\Rightarrow p \leq \frac{1}{2} \Rightarrow p \in \left[0, \frac{1}{2}\right]$$

7. Required probability

$$= {}^3C_0 \left(\frac{1}{2}\right)^3 \times {}^3C_0 \left(\frac{1}{2}\right)^3 + {}^3C_1 \left(\frac{1}{2}\right)^3 \times {}^3C_1 \left(\frac{1}{2}\right)^3$$

$$+ {}^3C_2 \left(\frac{1}{2}\right)^3 \times {}^3C_2 \left(\frac{1}{2}\right)^3 + {}^3C_3 \left(\frac{1}{2}\right)^3 \times {}^3C_3 \left(\frac{1}{2}\right)^3$$

$$= (1 + 9 + 9 + 1) \times \frac{1}{8} \times \frac{1}{8} = \frac{5}{16}$$

8. $P(\text{at least one success}) \geq \frac{9}{10}$

$$\Rightarrow P(X \geq 1) \geq \frac{9}{10}$$

$$\Rightarrow 1 - P(X = 0) \geq \frac{9}{10}$$

$$\Rightarrow P(X = 0) \leq \frac{1}{10}$$

$$\Rightarrow {}^n C_0 \left(\frac{1}{4}\right)^0 \left(\frac{3}{4}\right)^n \leq \frac{1}{10}$$

$$\Rightarrow \left(\frac{3}{4}\right)^n \leq \frac{1}{10}$$

$$\Rightarrow \left(\frac{4}{3}\right)^n \geq 10$$

$$\Rightarrow n \log_{10} \left(\frac{4}{3}\right) \geq \log_{10} 10$$

$$\Rightarrow n(\log_{10} 4 - \log_{10} 3) \geq 1$$

$$\Rightarrow n \geq \frac{1}{\log_{10} 4 - \log_{10} 3}$$

9. According to the given condition,

$$np + npq = 15 \text{ and } (np)^2 + (npq)^2 = 117$$

$$\therefore \frac{n^2 p^2 (1 + q^2)}{(np + npq)^2} = \frac{117}{15^2}$$

$$\Rightarrow \frac{1 + q^2}{(1 + q)^2} = \frac{117}{225}$$

$$\Rightarrow 6q^2 - 13q + 6 = 0 \Rightarrow q = \frac{2}{3}$$

$$\therefore p = 1 - \frac{2}{3} = \frac{1}{3}$$

$$\text{Since, } np + npq = 15$$

$$\Rightarrow n \times \frac{1}{3} + n \times \frac{2}{9} = 15$$

$$\Rightarrow n = 27$$

$$\therefore \text{mean} = np = 27 \times \frac{1}{3} = 9$$



$$10. \quad P(\text{getting head}) = p = \frac{1}{2}$$

$$\therefore q = 1 - \frac{1}{2} = \frac{1}{2}$$

$$\begin{aligned} \text{Here, } P(X=r) &= {}^n C_r p^r q^{n-r} = {}^n C_r \left(\frac{1}{2}\right)^r \left(\frac{1}{2}\right)^{n-r} \\ &= {}^n C_r \left(\frac{1}{2}\right)^n \end{aligned}$$

Since, $P(X=4)$, $P(X=5)$ and $P(X=6)$ are in A.P.

$$\Rightarrow 2P(X=5) = P(X=4) + P(X=6)$$

$$\Rightarrow 2 {}^n C_5 \left(\frac{1}{2}\right)^n = {}^n C_4 \left(\frac{1}{2}\right)^n + {}^n C_6 \left(\frac{1}{2}\right)^n$$

$$\Rightarrow 2 {}^n C_5 = {}^n C_4 + {}^n C_6$$

$$\Rightarrow 2 \frac{n!}{5!(n-5)!} = \frac{n!}{4!(n-4)!} + \frac{n!}{6!(n-6)!}$$

$$\Rightarrow \frac{2}{5(n-5)} = \frac{1}{(n-4)(n-5)} + \frac{1}{6 \times 5}$$

$$\Rightarrow n^2 - 21n + 98 = 0$$

$$\Rightarrow (n-7)(n-14) = 0$$

$$\Rightarrow n = 7 \text{ or } 14$$

11. Let the probability of success and failure be p and q respectively.

$$\therefore p = 2q$$

$$\text{Since, } p + q = 1$$

$$\therefore 3q = 1 \Rightarrow q = \frac{1}{3}$$

$$\therefore p = 1 - \frac{1}{3} = \frac{2}{3}$$

\therefore required probability

$$\begin{aligned} &= {}^6 C_4 \left(\frac{2}{3}\right)^4 \left(\frac{1}{3}\right)^2 + {}^6 C_5 \left(\frac{2}{3}\right)^5 \left(\frac{1}{3}\right) \\ &\quad + {}^6 C_6 \left(\frac{2}{3}\right)^6 \left(\frac{1}{3}\right)^0 \end{aligned}$$

$$= \frac{240}{729} + \frac{192}{729} + \frac{64}{729} = \frac{496}{729}$$

12. Mean = np and variance = npq

$$\therefore np = 20 \text{ and } npq = 16$$

$$\therefore 20q = 16 \Rightarrow q = \frac{4}{5}$$

$$\therefore p = 1 - \frac{4}{5} = \frac{1}{5}$$


$$\text{Since, } np = 20$$

$$\therefore n \times \frac{1}{5} = 20 \Rightarrow n = 100$$



MHT - CET

BASED ON STD. XI & XII SYLLABUS OF MHT-CET



AVAILABLE SUBJECTS:

- MHT - CET Triumph Physics
- MHT - CET Triumph Chemistry
- MHT - CET Triumph Mathematics
- MHT - CET Triumph Biology

BUY NOW

SALIENT FEATURES:

- Extensive coverage of Std. 12th syllabus and relevant chapters for Std. 11th
- Exhaustive subtopic wise coverage of MCQs in accordance with the latest paper pattern
- Inclusion of solved MCQs from previous year's question papers
- Evaluation test provided at the end of each chapter
- Two Model Question Papers with Answer Key to assess the level of preparation

Target Publications Pvt. Ltd.

88799 39712 / 13 / 14 / 15

mail@targetpublications.org

www.targetpublications.org