University of Cumbria

Holt, Claire, Nevin, Owen, Smith, Darrell and Convery, Ian (2018) Environmental niche overlap between snow leopard and four prey species in Kazakhstan. Ecological Informatics, 48 . pp. 97-103.

Downloaded from: http://insight.cumbria.ac.uk/id/eprint/4093/

Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available here) for educational and not-for-profit activities

provided that

• the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form

• a hyperlink/URL to the original Insight record of that item is included in any citations of the work

• the content is not changed in any way

• all files required for usage of the item are kept together with the main item file.

You may not

• sell any part of an item

• refer to any part of an item without citation

• amend any item or contextualise it in a way that will impugn the creator's reputation

• remove or alter the copyright statement on an item.

The full policy can be found here.
Alternatively contact the University of Cumbria Repository Editor by emailing insight@cumbria.ac.uk.

Article

Title: Environmental niche overlap between snow leopard and four prey species in Kazakhstan.

Running Head: Environmental niche overlaps.

Authors: Claire Denice Stevenson Holt[a], Owen Thomas Nevin[b], Darrell Smith[a] and Ian Convery[a]

Authors affiliations: [a]Department of Science, Natural Resources and Outdoor Studies, University of Cumbria, Ambleside, UK; [b]Associate Vice-Chanceller, Central Queensland University, Gladstone, Australia.

Authors Email: claire.holt@cumbria.ac.uk, o.nevin@cqu.edu.au, darrell.smith@cumbria.ac.uk, ian.convery@cumbria.ac.uk.

Corresponding author Email: claire.holt@cumbria.ac.uk

Abstract

The snow leopard *Panthera uncia* has declined due to habitat loss, habitat fragmentation and human persecution. Predator distribution is heavily dependent on prey species availability and distribution. With increasing pressures from farming practices encroaching into native species range and persecution of snow leopards in response to livestock depredation, it is vital to assess current predator and prey species distribution to highlight sensitive areas of overlap for protection. This study uses MaxEnt, a presence-only Species Distribution Model (SDM) to assess snow leopard and four prey species habitat suitability along the southern and eastern borders of Kazakhstan using environmental data. This area is considered an important corridor between snow leopard populations in the north and south of their range. Each of the five SDM's produced models of 'good' discriminating abilities. We then compared the potential niche overlap between snow leopard and four prey species using ENMTools to highlight areas of important niche overlap within the corridor. The results indicated a very high degree of overlap between snow leopard and Siberian ibex *Capra sibirica* and high degrees with red deer *Cervus elaphus*, argali *Ovis ammon* and urial *Ovis orientalis*. The snow leopard population in this region is also found to be using forested areas below 2500 m, much lower than recorded in other areas of their range. The results highlight areas needed for protection but also pose additional conservation questions regarding the importance of prey species to transitory individuals.

Keywords: SDM, MaxEnt, Management, Conservation, dispersal, distribution

Acknowledgements: The authors would like to say thank you to the WWF for suppling species record data and The Snow Leopard Trust for use of their current snow leopard distribution shapefile.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

68 **Main Text**

69 ## 1. INTRODUCTION

70 Land use change due to human modification is a global issue that is not restricted to the
71 local environment where the change occurs (Foley *et al.*, 2005). Changes to forests,
72 farmland, waterways and air are driven by human need for resources and are causing
73 considerable losses to biodiversity (Foley *et al.*, 2005; WWF, 2016). Human activities,
74 geographical barriers and ecological processes, competition and predation impact
75 animal populations and can force populations out of their fundamental niche (all
76 suitable habitat) into a much smaller area (Phillips *et al.*, 2006; Pulliam, 2000, WWF
77 2016); the realised niche (Hutchinson, 1957). Many studies have used species
78 distribution models (SDM also known as habitat suitability models HSM and climate
79 envelopes) to estimate the relationship between species records and the characteristics
80 of the landscape (Elith *et al.*, 2011; Ward, 2007; Su *et al*., 2015: Aryal *et al.*, 2016;
81 Lamsal *et al*., 2018a; Lamsal *et al*., 2018b). SDM models require a set of known species
82 locations and predictor variables such as land cover, elevation and climate data to train
83 the model and predict species distribution (Phillips & Dudik, 2008). By identifying
84 suitable habitat, SDM models can produce starting points for further discussion and
85 research in particular areas, for example, highlighting the fundamental niche for a
86 species and comparing it to the realised niche and assessing what impact human activity
87 is having upon distribution.

88 The snow leopard *Panthera uncia* is one species that has declined due to habitat
89 loss, human persecution and reduction in prey species distribution (Jackson et al.,
90 2014). Until recently the snow leopard was listed as an endangered species but has now
91 been down listed to vulnerable (Aryal, A. 2017). This has occurred despite much debate
92 by experts as to the current populations size, with experts believing that many animals

93 are poached and deaths are unreported (Aryal, A. 2017). The estimated population size
94 published in 2003 was between 4080-6590 individuals (Jackson et al. 2008). It is
95 suggested that snow leopards are found between 2500 to 5500 m in alpine and sub-
96 alpine areas (Aryal *et al.*, 2016) in habitats such as grassland, bare areas and agricultural
97 mosaic (Forrest *et al.*, 2012). The current population inhabit the mountain regions of the
98 Himalaya, thorough the Quighai-Tibet Plateau and central Asia to southern Siberia
99 (Jackson *et al.* 2008). Human wildlife conflict occurs when these animals depredate
100 domesticated species in farmed areas, unless there are conservation incentives in place
101 to dissuade hunting (Bagchi & Mishra, 2006). There is no one singly important prey
102 species for snow leopard survival, as prey varies in different areas of the snow leopards
103 range (Lyngdoh *et al.*, 2014). Wild goat and sheep species are commonly taken by snow
104 leopards with Siberian ibex *Capra sibirica*, Himalayan tahr *Hemitragus jemlahicus*,
105 blue sheep *Pseudois nayaur* and argali *Ovis ammon* being the four favoured species
106 (Lyngdoh *et al.*, 2014). However, domesticated goat and sheep species are also taken
107 (Aryal *et al.*, 2014a; Aryal *et al.*, 2014b; Aryal *et al*., 2014c; Lyngdoh et al., 2014).
108 These domesticated species are often farmed within the same areas of the landscape that
109 the wild goat and sheep species occur and in some areas the density of livestock is
110 higher than the native wild ungulates (Bagchi & Mishra, 2006; Aryal *et al.*, 2014a;
111 Aryal *et al.*, 2014b; Aryal *et al*., 2014c). In areas of the snow leopards range in
112 Pakistan, livestock out compete wild species for food and have caused dramatic declines
113 in wild species such as the urial *Ovis orientalis* which is now classed as vulnerable
114 (Siraj-ud-Din *et al.*, 2016, Valdez, 2008). The decline in some native prey species
115 forces snow leopard to prey on livestock, behaviour that can cause human wildlife
116 conflict with the local farming population (Bagchi & Mishra, 2006).

A recent camera trap study has identified individual snow leopards within a reserve in south Kazakhstan in lower elevations with tree cover (Convery et al. 2015). Snow leopard distribution in the mountainous areas of Kazakhstan is suggested to be between 750 m and 5500 m (Jackson *et al*., 2014). This is much lower than Aryla *et al*.'s (2016) suggestion of a lower limit of 2500 m, though in one of the earliest papers on snow leopard ecology, Hemmer (1973) reports that 'seasonal migration from higher to lower elevations may depend on climatic conditions and the movements of ungulate herds, and during winter, it may descend to the lower zones.' Riordan et al. (2015) least-cost connectivity study suggested that the Tian Shan Mountain range, which runs through Kyrgyzstan and borders south Kazakhstan and north China, is a potentially sensitive corridor between southern and northern snow leopard populations (Riordan *et al.*, 2015). The use of lower elevations seen in Convery *et al*. (2015) and Jackson *et al*. (2008) could potentially be in response to prey species distribution and the functional connectivity of habitat in this area acting as a movement corridor. The movement patterns suggested by Convery *et al*. (2015) suggest that snow leopards frequently crossed valley bottoms when moving between alpine mountain ridges or from ridges to forested areas. This will bring them closer to areas of human activity and habitation and make them susceptible to anthropogenic disturbance, potential poaching and increase the likelihood of livestock depredation.

Recent SDM studies have focused primarily on snow leopard distribution in the Himalayan portion of the species range and how habitat may shift in response to climate change altering population distribution (Aryal *et al.*, 2016; Aryal *et al.*, 2013; Forrest *et al.*, 2012). However, Aryal et al (2016) suggests that predator species distribution models should be compared to prey species distribution due to the influence prey availability has on predator distribution. There are still gaps in our current knowledge

142 of global, national and local snow leopard population sizes and fine scale species
143 distribution modelling is needed to aid conservation and help map current distribution
144 (Network, 2014). Forrest *et al's* (2012) SDM suggests that climate change will effect
145 snow leopard distribution through changes in habitat loss and fragmentation rather than
146 temperature and precipitation. We suggest that habitat type, elevation and movement of
147 prey species is important to snow leopard distribution. As elevation and temperature co-
148 vary, in this study, elevation was chosen as an environmental layer to represent changes
149 in mountainous areas. Previous studies have used Maxent, a SDM, to assess snow
150 leopard distribution (Li *et al.* 2016; Li *et al.* 2014) and to assess prey species in the
151 Himalayan area (Aryal et al (2016). However, this study aims to use MaxEnt (Elith *et*
152 *al.*, 2011) to assess snow leopard habitat suitability along the Kazakhstan south and
153 eastern border, which has been highlighted by Riordan et al. (2015) least-cost
154 connectivity study as part of a potentially sensitive corridor between southern and
155 northern snow leopard populations. Due to the importance of prey species presence on
156 snow leopard distribution (Aryal et al (2016), we will then compare the potential niche
157 overlap between snow leopard and four prey species to highlight areas of importance
158 within this dispersal corridor.

159 ## 2. MATERIALS AND METHODS

160 ### 2.1. Species distribution data

161 This study focuses on snow leopard distribution along a potential corridor on the
162 southern and eastern border of Kazakhstan. This area includes the Western Tian Shan
163 and Kyrgyz Alatau mountain ranges, which run along part of the southern border with
164 Kyrgyzstan, the Borohoro, Junggar Alatau, Saur, and Tarbagatai ranges, which are on
165 the border of Kazakhstan and China, and the Altai which is on the border with

Mongolia, China and Russian (Jackson *et al*., 2014); Figure 1). Often there is limited systematic survey data available on the presence/absence of elusive animals in the environment (Elith *et al.*, 2011). In many cases only presence data is available which has either been collected systematically during surveys, or in the majority of cases, is acquired from natural history museums and databases (Elith *et al.*, 2011). Ideally using data that has been systematically collected over the survey area would best, however often this data is not available and data from museums and databases are used instead. One fundamental limitation of this data is sample selection bias, where some areas of the study area are sampled more intensively that others, but at times this is the only data available (Elith *et al.*, 2011). Snow leopard sightings (N= 125) data was obtained from WWF. These data were collected by multiple specialists over 50 years for WWF using multiple survey techniques within different studies, due to the different techniques used it is acknowledged there may be biases present within the data such as sample selection bias. Four prey species were also selected from data available from WWF, two of these, the Siberian ibex (N= 194) and argali (N= 317), are favoured prey species (Lyngdoh *et al.*, 2014). The two other species were the threatened urial (N= 49) and the red deer *Cervus elaphus* (N= 129), both of which are known prey species (Jackson *et al*., 2014). Within the study area, no data was recorded for the favoured prey species the blue sheep or the Himalayan tahr therefore species distribution model cannot be constructed for these species in this area currently.



Figure 1. Current suggested range of the snow leopard highlighted study area along the southeast border of Kazakhstan. * Snow leopard range shapefile curtsey of The Snow Leopard Trust.

2.2. Environmental Layer

Land cover data was obtained from the European Space Agency GlobCover data set (Medsia-France 2008). These data are divided into 22 land cover categories and are in raster (gridded data) format at a resolution of 300 m as used in Forrest et al (2012). It is acknowledged that these data have biases in what classes they differentiate.

196 Global Multi-resolution Terrain Elevation data (GMTED2010) was obtained from
197 United States Geological Service (USGS) Earth Explorer at a resolution of 250 m.
198 Aspect and slope were calculated from the elevation data using ArcGIS (ESRI,
199 Redlands, CA) Spatial Analysis extension toolbox and the slope and aspect tools. The
200 elevation, slope and aspect raster data sets were resized using the Spatial Analysis
201 Extract by Mask tool using nearest neighbour to 300 m resolution (this was done to
202 match the land cover extend which is the largest resolution). MaxEnt requires all
203 environmental layers to have the same co-ordinate systems, map extent and raster cell
204 size, all of which can be altered using the Extract by Mask tool. All environmental
205 layers must also be converted to an Ascii file type using the Conversion Tools Raster to
206 Ascii tool for modelling purposes.

207 **2.3. MaxEnt Modelling**

208 The models for all species were run in MaxEnt Version 3.3.3k, using primarily default
209 settings (regularisation multiplier = 1; duplicate occurrences removed; maximum
210 number of background points = 10000, as used in Kramer-Schadt *et al.*, 2013). MaxEnt
211 can select a proportion of random points to be used as test data or this can be defined by
212 the user. In this study, the distribution data was split so that 25 % of the distribution
213 locations were used for testing the model and 75 % for model training. Five-fold cross
214 validation was used to calculate mean Area Under Receiver Operating Characteristic
215 (ROC) Curve or AUC and extrinsic omission rates (the average proportion of test points
216 that fall outside the area predicted to be suitable), following use of the occupancy
217 threshold rule that maximises the sum of test sensitivity and specificity (as
218 recommended by Liu *et al.*, 2013). AUC is used to assess model performance with
219 values of 0.5 and below indicating the model is no better than random and values closer

220 to 1.0 indicating better model performance. Hawlitschek et al. (2011) define AUC >0.9
221 as having 'very good' discriminating abilities, >0.8 are 'good' and >0.7 is 'useful'
222 (based on the definition of (Swets, 1988)). The 10 % minimum training logistic
223 threshold found in the MaxEnt results table was used to define suitable and unsuitable
224 habitat for each species (Aryal *et al* 2016; Warren *et al.*, 2010). The ENMTools
225 software (Warren *et al.*, 2010) was then used to compare the ecological niche of the
226 snow leopard and their prey species using the niche overlap tool. Schoener's D (1968)
227 and the I statistic (Warren *et al.*, 2008) were the statistics used to measure niche
228 overlap.

229 **2.4. Accounting for pseudoabsences**

230

231 Within MaxEnt background samples, known as pseudoabsences can be randomly
232 selected within the programme to create absence points (Elith *et al.*, 2011). The
233 background samples used can have significant effects on the model outputs (Elith *et al.*,
234 2011). In MaxEnt points are selected typically from a large rectangular area that may
235 contain suitable habitat where no species sightings have been recorded (Brown, 2014).
236 When models select background points from these areas it causes false positives
237 (Brown, 2014). To overcome this bias Brown (2014) and Edith et al. (2011) suggest
238 reducing the area where background points can be selected by using a minimum convex
239 polygon (MCP) based on the presence data. In this study the SDMtoolbox was added in
240 to ArcGIS and a background file was created using the Sample by Buffered MCP tool.
241 This creates a bias file which can then be used within the MaxEnt interface. MaxEnt
242 will then only select background sample points from within the designated 1 km
243 buffered MCP. Therefore in this study, the background point selection is limited to the

areas where sightings have occurred and are assumed to have been surveyed for snow leopards.

3. RESULTS

3.1. Predicted snow leopard habitat

The results from the MaxEnt five-fold cross-validation test showed that the model for snow leopard distribution has 'good' discriminating abilities (Table 1) with a mean AUC of 0.817. The predicted areas of suitable habitat for the snow leopard along the Kazakhstan southern border included areas that are currently designated as within the snow leopards range (Figure 2). However, the MaxEnt model also highlighted additional areas (shown in green in Figure 4) within the fundamental niche that are potentially suitable for snow leopards. Mainly these areas are on the border between Kazakhstan and Kyrgyzstan and Kazakhstan and China. The study area contains multiple areas of suitable habitat, which vary in size and are surrounded by less suitable areas of the landscape (less suitable areas of the landscape shown in blue in Figure 4). The current predicted range does also include some areas where the MaxEnt model has not highlighted as highly suitable. Based on jackknife estimates assessing the importance of each environmental layer added into the MaxEnt model (land cover, elevation, slope and aspect), elevation is seen as a significant factor in defining the predicted area. Elevation influences the suitable habitat by contributing 74.1 % to the model, slope is second but with a much lower 17.6 % contribution, land cover is third (6.7 %) and aspect forth (1.6 %) (Figure 2). The partial dependency plots also indicates that probability of snow leopard presence is highest at an elevation of 2500m (Figure 3).

Table 1. MaxEnt results for snow leopard and prey species.

Species	Train set	Test set	Train AUC	Test AUC	Test gain	Test omission
Snow leopard	79	20	0.858 ±0.005	0.817 ±0.010	0.763 ±0.763	0.010*
Siberian ibex	117	29	0.801 ±0.009	0.736 ±0.020	0.395 ±0.124	0.020*
Argali	124	31	0.850 ±0.009	0.808 ±0.020	0.619 ±0232	0.038*
Urial	30	7	0.826 ±0.018	0.740 ±0.084	0.338 ±0.514	0.082
Red deer	78	19	0.913 ±0.002	0.898 ±0.003	1.485 ±0.129	0.031*

Train set = the average number of training samples, Test set = average number of test samples, Test omission = Balance training omission, predicted area and threshold value test. Asterisk = $p<0.05$, ± = Standard deviation from mean.



Figure 2. The probability of presence based on the effect of each variable for each species.



276

277 Figure 3. Partial dependency plot from MaxEnt displaying the partial effect elevation has

278 on the probability of presence of the snow leopard.

279

280



Figure 4. Predicted habitat suitability for snow leopard compared to current range data available from The Snow Leopard Network.

3.2. Comparison of predicted habitat suitability for snow leopard and prey species

The results from the MaxEnt five-fold cross-validation test for the four prey species, Siberian ibex, argali, urial and red deer, indicated that the models varied in performance but all performed better than random and are classed as having 'good' discriminating abilities (Table 1, Figure 5). Red deer had the highest mean test AUC of all models with a value of 0.898, argali had a lower value of 0.808, and urial and Siberian ibex test AUC values were 0.740 and 0.736, respectively. The highest degree of niche overlap using

293 the mean Schoener's *D* index value and the *I*-statistic was between snow leopard and
294 Siberian ibex with a value of $D = 0.716$ and $I = 0.921$. Values of 0 represent little
295 overlap and values closer to 1 represent high degree of overlap and *I*-statistic values are
296 generally higher than *D* values (Hawlitschek *et al.*, 2011). A lower degree of niche
297 overlap was seen between the snow leopard and red deer ($D = 0.665$ and $I = 0.889$) and
298 argali ($D = 0.629$ and $I = 0.876$) and the lowest overlap is seen between snow leopard
299 and urial ($D = 0.452$ and $I = 0.751$).
300 At elevation of 2500 m – 5500 m there is highly suitable habitat for snow
301 leopard ($\chi 2$ 8.26, df = 2, p=0.01) and Siberian ibex ($\chi 2$ 52.91, df = 2, $p<0.05$) ,
302 whereas argali ($\chi 2$ 396.20, df = 2, $p<0.05$), urial (chi 61.69, df = 2, $p<0.05$) and red
303 deer ($\chi 2$ 36.40, df 2, $p<0.05$) highly suitable habitat is lower than 2500 m.



Figure 5. Predicted suitable habitat for snow leopard and four prey species along the Kazakhstan south-eastern border. Suitable habitat based on 10 percentile training presence logistic threshold in MaxEnt.

4. DISCUSSION AND CONCLUSIONS

MaxEnt SDM has been used to assess the habitat suitability for the snow leopard in south and east Kazakhstan. The results provided a model of 'good' discriminatory abilities that indicated that there is a substantial amount of highly suitable habitat for the

313 snow leopard along the Kazakhstan border. These habitats are connected to other highly
314 suitable habitats in Kyrgyzstan, China, Mongolia and Russia. The habitat along the
315 Kazakhstan border forms part of an important narrow corridor between snow leopard
316 populations in the north and south of their range (Riordan *et al.*, 2015) and could
317 potentially have both resident and dispersing individuals using the habitat in these areas.
318 Kazakhstan is thought to have a population of 100-110 snow leopards (2.5% of the
319 global population) (Jackson *et al*. 2008) and two stable populations are thought to
320 inhabit Almaty State Nature Reserve (area = 915 km^2, population of 30-35 individuals)
321 and Aksu Zhabagly State Reserve (area = 744 km^2) both situated near the Kazakhstan
322 and Kyrgyzstan border (Convery et al. 2015; Jackson *et al.* 2014; Saparbayev &
323 Woodward, 2008). Individuals with these reserves are protected, but can potentially
324 disperse into unprotected areas of highly suitable habitat shown in the SDM. The
325 unsuitable habitat indicated in the SDM are seen at lower elevations and are mainly
326 comprised of urban and agricultural land cover types. However, MaxEnt jackknife
327 analysis identified elevation is the key variable in determining areas that are highly
328 suitable for snow leopards (contributing 74.1 % to the SDM) not land cover type. In
329 this study, snow leopard and Siberian ibex are shown to have the highest degree of
330 niche overlap. The SDM indicates that the majority of the landscape is shared by these
331 two species, with a small amount of the landscape highlighted as prey only. This
332 overlap suggests that the environmental space for both predator and prey is similar and
333 they can potentially inhabit similar areas (Lyngdoh *et al.*, 2014). This is consistent with
334 other studies that have found that the main prey species are blue sheep and the Siberian
335 ibex, both of which are found in higher elevations (Aryal *et al.*, 2016; Lyngdoh *et al.*,
336 2014). The snow leopard sightings locations in this study were located between 1188 m
337 to 4789 m, 1312 m lower than Aryal et al. (Aryal *et al.*, 2016) suggestion of a lower

threshold of 2500 m for the Himalaya population, though within the range suggested by Jackson et al. (2008) for Kazakhstan, where snow leopard can be found in the mountain ranges between 750 m to 5500 m (Jackson *et al*., 2014; Saparbayev & Woodward, 2008). Although, there were significantly more distribution points above 2500 m indicating a preference, 46 individual points were seen below 2500 m.

In the Almaty Nature Reserve, south Kazakhstan, snow leopards use elevations lower than 2500 m in winter months as they are following their main prey species, the Siberian ibex, to sheltered forested areas (Saparbayev & Woodward, 2008, Altynbek, (2015) pers com)). This is supported by anecdotal evidence from the ranger team at Almaty State Nature Reserve (ASNR), and in particular the head ranger Janyspayer Altynbek, who has over 30 years experience of working within ASNR. Aryal et al. (2016) suggest that climate change will reduce the degree of overlap between blue sheep and snow leopard in the Himalaya regions with prey species shifting their current range. The snow leopard population in Kazakhstan are showing levels of adaptability in that they are currently seen to use lower elevations at certain times of the year (Saparbayev & Woodward, 2008). Once in these lower areas, snow leopards are sharing a niche with other potential prey species such as the argali, urial and red deer which are seen in this study to have a 'good' degree of niche overlap with the snow leopard. Pilot studies using Fuzzy Logic modelling to assess the impact of climate change on snow leopard distributions in ASNR (Convery *et al*. 2015) suggest that changes in the elevation at which seasonal snow pack accumulates will have a strong driving influence on elevational range occupied. Snow leopards are seen to prey upon different species in different regions depending upon what is available (Lyngdoh *et al.*, 2014). The potential snow leopard niche area in Kazakhstan as indicated in the SDM suggests that snow leopard have opportunities to prey on a range of species, particular at lower elevations.

However, this will bring the snow leopard in to areas of the landscape that are farmed and may have the potential to cause human-wildlife conflicts.

To conserve the snow leopard population in Kazakhstan there are areas of research that must be prioritised. First, it is essential that presence/absence of snow leopards is ascertained for all areas in the landscape that have been highlighted as highly suitable in the SDM. A key assumption of SDM is that sightings data are reliable and collected in a systematic way (Elith *et al*., 2011). Although the data used in this study was collected over the last 50 years by WWF, it is acknowledged that there may be biases and subjectivity within the sightings data. Investigating presence/absence and population data for snow leopards globally has been highlighted as a current research priority by the Snow Leopard Network (Jackson *et al*., 2014). Previous studies have used methods such as questionnaires and interviews with the general public and farming communities to establish presence/absence (Taubmann *et al.*, 2016), while others have relied on tracks and signs in the environment and more recently using camera traps to assess presence and population size (Convery *et al*., 2015). By gathering this information, it will aid conversation efforts by highlighted areas that are currently inhabited by snow leopard but not protected and areas that are highly suitable but where snow leopards are missing.

Secondly, it is imperative to understand the predator/prey relationship in Kazakhstan. The areas highlighted in green in Figure 4 identify where the snow leopard share a niche with the four prey species. These areas need to be studied to see whether prey distribution is correctly predicted within the SDM and to determine the viability and health of the current prey population. A decline in prey species has been listed as one of the main causes of snow leopard population declines (Jackson *et al*., 2014). Although species like the markor and urial are seen as unimportant to snow leopards

388 current diet (Lyngdoh *et al.*, 2014), this may change in the future with the effects of
389 increased pressure from agriculture and due to climate change. It is clear from the
390 impact farming practices has had on the urial population in Pakistan, that presence of
391 domesticated animals can severely impact wild species populations size and niche area
392 (Siraj-ud-Din *et al.*, 2016). Also if regularly using lower elevations the snow leopards
393 are potentially coming in to regular contact with the farming community which may
394 cause human/wildlife conflicts. As the SDM has highlighted a high degree of overlap
395 between the snow leopard and red deer, argali, urial and, particularly, Siberian ibex, it is
396 vital that studies are undertaken to understand population dynamics and seasonal
397 movements of each species. It would be interesting to investigate the movements of
398 resident and dispersing snow leopard individuals to see whether any individuals are
399 found at elevations <2500 m all year round and whether they are seen to switch prey
400 species more regularly as they share a niche area with multiple species.
401 The SDM has highlighted highly suitable areas of the landscape within
402 Kazakhstan for both predator and prey species and where these species share a niche.
403 These populations are important to snow leopard conservations as they form part of a
404 corridor between the north and south snow leopard world wide range. Mountain habitats
405 are vulnerable to environmental change and anthropogenic influences, and climate
406 change poses a range of serious threats, including melting glaciers, changing rainfall
407 patterns, unpredictable weather conditions, and increasing temperatures. For mountain
408 species like snow leopards, climate change has immediate impacts with temperature,
409 competition from other predators, precipitation changes and increasing human activity
410 fragmenting suitable habitat (Riordan et al., 2015). A widespread upward encroachment
411 of subalpine forests would displace regionally unique alpine tundra habitats and
412 possibly cause the loss of alpine species. Therefore, the warmer and wetter conditions

consistent with climate change predictions in this region may result in vegetation communities at higher altitudes, with forests ascending into alpine areas, the snow leopards' preferred habitat (Forrest et al., 2012). Similar to Forest et al. (2012), we assume that the impacts of climate change on snow leopards will be primarily through changes in habitat, rather than through direct physiological impacts of temperature and precipitation. With additional pressure from farming practices and the threat of species shifts in relation to habitat shifts related to climate change, there is still information about current population dynamics that need to be understood before mitigation strategies can be developed for the future. Within Kazakhstan there may be resident and transient snow leopard individuals. However, these individuals are seen to follow prey to lower forested elevations. This leads to questions about the degree of adaptability the snow leopard has regarding prey species and habitat usage which need further investigation.

References:

Aryal, A., Shrestha, U.B., Ji, W., Ale, S.B., Shrestha, S., Ingty, T., Maraseni, T., Cockfield, G. & Raubenheimer, D. 2016. Predicting the distributions of predator (*snow leopard*) and prey (*blue sheep*) under climate change in the Himalaya. *Ecology and Evolution* 6: 4065-4075.

Aryal, A., Brunton, D., McCarthy, T., Karmachharya, D., Ji, W., Bencini, R., Raubenheimer, D. 2014a. Multipronged strategy including genetic analysis for assessing conservation options for the snow leopard in the central Himalaya. *Journal of Mammalogy* 95(6): 871-881.

Aryal, A., Brunton, D., Ji, W., Barraclough, R.K., Raubenheimer, D. 2014b. Humman-carnivore conflict: Ecological and economical sustainability of predation on livestock

437 by snow leopard and other carnivores in the Himalaya. *Sustainability Science* 9(3):321-
438 329.

439 Aryal, A., Brunton, D., Raubenheimer, D. 2014c. Impacts of Climate change on human-
440 wildlife- ecosystem interactions in the Trans-Himalayan region of Nepal. *Theoretical*
441 *and Applied Climatology* 115:517-529.

442 Aryal, A., Raubenheimer, D., Brunton, D. 2013. Habitat assessment for the
443 translocation of blue sheep to maintains a viable snow leopard population in the Mt
444 Everest Region, Nepal. *Zoology and Ecology* 23(1):66-82.

445 Bagchi, S. & Mishra, C. 2006. Living with large carnivores: predation on livestock by
446 the snow leopard (*Uncia uncia*). *Journal of Zoology* 268: 217-224.

447 Brown, J.L. 2014. SDMtoolbox: a python-based GIS toolkit for landscape genetic,
448 biogeographic and species distribution model analyses. *Methods in Ecology and*
449 *Evolution* 5: 694-700.

450 Convery, I., Baibagysov, A., Baiturbayev, K., Deecke, V.B., Harpley, D., Holt, C.,
451 Janyspayev, A.D, Nevin, O.T., Nurtazin, S, Smith, D.J. & van der Velden, N.K. 2015.
452 *Fuzzy Logic Modelling of Snow Leopard Populations in Response to Threats from*
453 *Climate Change*. - Centre for Wildlife Conservation, University of Cumbria,
454 Ambleside, U.K

455 Elith, J., Phillips, S.J., Hastie, T., Dudík, M., Chee, Y.E. & Yates, C.J. 2011. A
456 statistical explanation of MaxEnt for ecologists. *Diversity and Distributions* 17: 43-57.

457 Foley, J.A., Defries, R., Asner, G.P., Barford, C., Bonan, G., Carpenter, S.R., Chapin,
458 F.S., Coe, M.T., Daily, G.C., Gibbs, H.K., Helkowski, J.H., Holloway, T., Howard,

459 E.A., Kucharik, C.J., Monfreda, C., Patz, J.A., Prentice, I.C., Ramankutty, N. & Snyder,
460 P.K. 2005. Global consequences of land use. *Science* 309: 570-574.

461 Forrest, J.L., Wikramanayake, E., Shrestha, R., Areendran, G., Gyeltshen, K.,
462 Maheshwari, A., Mazumdar, S., Naidoo, R., Thapa, G.J. & Thapa, K. 2012.
463 Conservation and climate change: Assessing the vulnerability of snow leopard habitat to
464 treeline shift in the Himalaya. *Biological Conservation* 150: 129-135.

465 Hawlitschek, O., Porch, N., Hendrich, L. & Balke, M. 2011. Ecological niche modelling
466 and nDNA sequencing support a new, morphologically cryptic beetle species unveiled
467 by DNA barcoding. *PLoS One* 6: e16662.

468 Jackson, R., Mallon, D., McCarthy, T., Chundaway, R.A. & Habib, B. 2008. *Panthera*
469 *uncia*. The IUCN Red List of Threatened Species 2008: e.T22732A9381126. Retrieved
470 from http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T22732A9381126.en.

471 Jackson, R., Mallon, D., Mishra, C. Noras, S., Sharma R., & Suryawanshi, K . 2014.
472 Snow leopard survival strategy. - Snow Leopard Network, *Seattle, Washington, USA,* 1-
473 145.

474 Kramer-Schadt, S., Niedballa, J., Pilgrim, J.D., Schröder, B., Lindenborn, J., Reinfelder,
475 V., Stillfried, M., Heckmann, I., Scharf, A.K. & Augeri, D.M. 2013. The importance of
476 correcting for sampling bias in MaxEnt species distribution models. *Diversity and*
477 *Distributions* 19: 1366-1379.

478 Lamsal, P., Kumar, L., Aryal, A., Atreya, K. 2018a. Invasive alien plant species
479 dynamics in the Himalayan region under climate change. *AMBIO A Journal of the*
480 *Human Environment*. DOI 10.1007/s13280-018-1017-z.

481 Lamsal, P., Kumar, L., Aryal, A., Atreya, K. 2018. Future climate and habitat
482 distribution of Himalayan Musk Deer (*Moschus chrysogaster*). *Ecological*
483 *Informatics*. https://doi.org/10.1016/j.ecoinf.2018.02.004

484 Li, J., McCarthy, T. M., Wang, H., Weckworth, B. V., Schaller, G. B., Mishra, C., Lu,
485 Z. and Beissinger, S. R. 2016. Climate refugia of snow leopards in High Asia.
486 *Biological Conservation* 203: 188-196.Li, J., Wang, D., Yin, H., Zhaxi, D., Jiagong, Z.,
487 Schaller, G. B., Mishra, C., McCarthy, T. M, Wang, H., Wu, L, Xiao, L., Basang, L.,
488 Zhang, Y., Zhou, Y., and Zhi, L. (2014). Role of Tibetan Buddhist monasteries in snow
489 leopard conservation. *Conservation biology* 28: 87-94.

490 Liu, C., White, M. & Newell, G. 2013. Selecting thresholds for the prediction of species
491 occurrence with presence-only data. *Journal of Biogeography* 40: 778-789.

492 Lyngdoh, S., Shrotriya, S., Goyal, S.P., Clements, H., Hayward, M.W. & Habib, B.
493 2014. Prey preferences of the snow leopard (*Panthera uncia*): regional diet specificity
494 holds global significance for conservation. *PloS One* 9: e88349.

495 Phillips, S.J., Anderson, R.P. & Schapire, R.E. 2006. Maximum entropy modeling of
496 species geographic distributions. *Ecological Modelling* 190: 231-259.

497 Phillips, S.J. & Dudik, M. 2008. Modeling of species distributions with Maxent: new
498 extensions and a comprehensive evaluation. *Ecography* 31: 161-175.

499 Pulliam, H.R. 2000. On the relationship between niche and distribution. *Ecology Letters*
500 3: 349-361.

501 Riordan, P., Cushman, S.A., Mallon, D., Shi, K. & Hughes, J. 2015. Predicting global
502 population connectivity and targeting conservation action for snow leopard across its
503 range. *Ecography* 39: 419-426.

504 Saparbayev, S.K. & Woodward, D.B. 2008. Snow Leopard (*Uncia uncia*) as an
505 Indicator Species and Increasing Recreation Loads in the Almaty Nature Reserve. -
506 Paper presented at the meeting of The Fourth International Conference on Monitoring
507 and Management of Visitor Flows in Recreational and Protected Areas. Tuscany, Italy.

508 Schoener, T.W. 1968. The Anolis lizards of Bimini: resource partitioning in a complex
509 fauna. *Ecology* 49: 704-726.

510 Siraj-ud-Din, M., Minhas, R.A., Khan, M., Ali, U., Bibi, S.S., Ahmed, B. & Awan,
511 M.S. 2016. Conservation Status of Ladakh Urial (*Ovis vignei vignei* Blyth, 1841) in
512 Gilgit Baltistan, Pakistan. *Pakistan Journal of Zoology* 48: 1353-1365.

513 Su, J., Aryal, A., Nan, Z., Ji, W. 2015. Climate change-induced range expansion of a
514 subterranean rodent: Implications for rangeland management in Qinghai-Tibetan
515 Plateau. *PloSONE* 10(9):e0138969.

516 Swets, J.A. 1988. Measuring the accuracy of diagnostic systems. *Science* 240: 1285-
517 1293.

518 Taubmann, J., Sharma, K., Uulu, K.Z., Hines, J.E. & Mishra, C. 2016. Status
519 assessment of the Endangered snow leopard *Panthera uncia* and other large mammals
520 in the Kyrgyz Alay, using community knowledge corrected for imperfect detection.
521 *Oryx* 50: 220-230.

522 Valdez, R. 2008. *Ovis orientalis*. The IUCN Red List of Threatened Species 2008:
523 e.T15739A5076068. Retrieved from
524 http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T15739A5076068.en.

525 Ward, D.F. 2007. Modelling the potential geographic distribution of invasive ant
526 species in New Zealand. *Biological Invasions* 9: 723-735.

527 Warren, D.L., Glor, R.E. & Turelli, M. 2010. ENMTools: a toolbox for comparative
528 studies of environmental niche models. *Ecography* 33: 607-611.

529 Warren, D.L., Glor, R.E. & Turelli, M. 2008. Environmental niche equivalency versus
530 conservatism: quantitative approaches to niche evolution. *Evolution* 62: 2868-2883.

531 WWF (2016). Living Planet Report 2016. Risk and resilience in a new era. WWF
532 International, Gland, Switzerland. Retrieved from
533 http://awsassets.panda.org/downloads/lpr_living_planet_report_2016.pdf.

534
535
536
537
538
539