



# Armada® 50WP

## FUNGICIDE

BACKED  
by BAYER™

3844076B 100506A

For the enhancement of greener and more dense sodfarm turfgrass and for the control of certain foliar, stem, and root diseases of turfgrass including institutional, commercial and residential lawns, sod farms, sports fields, parks, municipal grounds and cemeteries, and for control of foliar and stem diseases of ornamentals and fruit trees grown in interiorscapes (to include shopping malls, shopping centers, hotel lobbies), residential and commercial landscapes.

Intended For Professional Use Only.

**Active Ingredient:**

Trifloxystrobin (CAS No. 141517-21-7).....	8.33%
Triadimefon (CAS No. 43121-43-3).....	41.67%

**Other Ingredients:**.....50.00%

**Total:**.....100.00%

EPA Reg. No. 432-1412

EPA Est. indicated by 3rd and 4th digits of the batch number on this package. (01, 03, 09, 14 or 19)=264-MO-003, (98)=33967-NJ-1

## KEEP OUT OF REACH OF CHILDREN. CAUTION

### FIRST AID

<b>If in eyes</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment.

**Note to Physician:** If ingested, induce emesis or lavage stomach. Treat symptomatically.

# PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Cause moderate eye irritation. Harmful if swallowed, inhaled, or absorbed through the skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid breathing dust or spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

### Personal Protective Equipment (PPE)

**Mixers/loaders, applicators and other handlers must wear:**

Long-sleeved shirt and long pants.

- Chemical-resistant gloves made of waterproof material such as neoprene, butyl rubber, barrier laminate or nitrile rubber when mixing/loading, when using handheld equipment or handheld nozzles.
- Shoes plus socks
- Chemical-resistant apron, when mixing/loading, or cleaning spills or equipment.

See engineering controls for additional requirements.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate.

Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Control Statements

Water-soluble packaging when used correctly qualifies as a closed mixing/loading system under the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240(d)(4)].

Mixers, loaders, applicators, and other handlers using water soluble packets must:

- wear the personal protective equipment on this labeling for mixers/loaders, however, they do not need to wear a respirator, and
- be provided and have immediately available for use in an emergency, such as a broken package, spill, or equipment breakdown, a NIOSH-approved respirator with any N, R, P or HE filter.

Pilots must use an enclosed cockpit in a manner that is consistent with the WPS for Agricultural Pesticides [40 CFR 170.240(d)(6)].

Pilots must wear the PPE required on this labeling for applicators.

## User Safety Recommendations

### Users should:

- Wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

### Ground Water Advisory

The active ingredients in this product have properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination.

Do not make applications when weather conditions favor drift from target area.

## PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill, or store near heat or open flame.

## DIRECTIONS FOR USE

**It is a violation of federal law to use this product in a manner inconsistent with its labeling.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

**Do not apply by aerial application in the State of New York.**

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Do not store below freezing (32°F). Exposure to moisture or excessive handling of water soluble packets may cause breakage. Store packets in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Material that cannot be used as directed should be disposed of as directed below. In spill or leak incidents, keep unauthorized people away.

**Pesticide Disposal:** Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be used according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

**Container Disposal:**

**Foil Pouches**

Non-refillable container. Do not reuse or refill this container. Dispose of empty pouches in a sanitary landfill.

**Mini drum**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available. Then dispose of empty carton in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## GENERAL INFORMATION

### Mixing Procedures

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Agitation is necessary for proper dispersal of the product. Maintain agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

### Mixing Procedures (Water Soluble Packets)

The enclosed packets containing ARMADA 50 WP are water soluble. Do not allow packets to become wet prior to adding to the spray tank. Do not handle with wet hands. Reseal outer carton/container to protect remaining packets.

To prepare the spray mixture, drop the required number of unopened packets into the spray tank while filling the water to the desired level. Operate the agitator while mixing. Depending on the water temperature and the degree of agitation, the packets should be completely dissolved within a few minutes from the time they were added to the water.

**Attention: Do not use ARMADA50 WP water soluble packets in a tank-mix with products that contain boron or release free chlorine. The resultant reaction of PVA and boron or free chlorine is a plastic, which is not soluble in water or solvents such as diesel oils, kerosene, gasoline or alcohol. Use of chlorinated water is acceptable. Do not attempt to use the PVA packets directly in liquid fertilizer, diesel oils or summer spray type oils as in ULV or LV uses. PVA Packets are water soluble not fertilizer or oil soluble. If liquid fertilizer or oils are to be added to the mix water, add ARMADA 50 WP packets first and wait until they are dissolved before adding liquid fertilizer or oil.**

**ARMADA 50 WP Alone:** Add 1/2 of the required water to the mix tank. With the agitator running, add the required number of ARMADA 50 WP packets to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after ARMADA 50 WP packets have completely dissolved and dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

**ARMADA 50 WP + Tank Mixtures:** Add 1/2 of the required amount of water to the tank mix. Start the agitation running before adding ARMADA 50 WP Fungicide water soluble packets and any tank mix partner(s). In general, tank mix partners should be added in this order: (1) products packaged in water soluble packaging (such as ARMADA 50 WP), wettable powders, wettable granules (dry flowables); (2) liquid flowables, liquids; and (3) emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

**Note:** When using ARMADA 50 WP in tank mixes, all products in water soluble packaging (including ARMADA 50 WP) should be added to the tank before any other tank mix partner. Allow the water soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using ARMADA 50 WP in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations that appear on the tank mix product label. No label dosage rate should be exceeded, and the most restrictive label precautions and limitations should be followed. This product should not be mixed with any product that prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

ARMADA 50 WP is compatible with most insecticide, fungicide, and foliar nutrient products. However, the compatibility of ARMADA 50 WP with tank mix partners should be tested before use.

To determine biological compatibility with other products, mix the products in the desired proportions, spray on target plants and observe for phytotoxicity seven days after the application.

Observe all directions, precautions, and limitations on labeling of all products used in tank mixes. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

**Use with additives:** Use of spray additives are not required. Any spray additive should be evaluated prior to use. Do not use in conjunction with organosilicate-based products, or plant injury may occur. Label directions are based on data with no additives.

**Chemigation:** Do not apply this product through any type of irrigation system.

**Resistance Management:** ARMADA 50 WP contains the strobilurin class of chemistry, which exhibits no known cross-resistance to other chemical classes including sterol inhibitors, dicarboximides, benzimidazoles, anilinopyrimidines, or phenylamides. However, certain fungal pathogens are known to develop resistance to products used repeatedly. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies. Such strategies may include rotating and/or tank mixing with products having different modes of action; or limiting the total number of applications per season. Bayer encourages responsible product stewardship to ensure effective long-term control of the fungal diseases on this label. See specific recommendations in the turf section and the ornamentals section.

**Application:** ARMADA 50 WP is absorbed rapidly and works systemically from within the plant. Good coverage and wetting of the foliage are necessary. Rainfall or sprinkler irrigation, within 30 minutes after application does not decrease effectiveness. Control may be less effective on plants suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

In all cases apply when plants are fully established and actively growing. Applications must be made at prescribed intervals to maintain disease control.

This product cannot be mixed with any product containing a label prohibition against such mixing.

Do not use on crops grown for food or forage.

For Residential and commercial turf sites apply product with spray equipment such as back pack sprayer, hand pump sprayer, tank and hand-held spray gun, boom sprayer, and or ride-on sprayer.

For Residential and commercial Landscapes apply product with spray equipment such as back pack sprayer, hand pump sprayer, tank and hand-held spray gun or wand.

For Noncommercial Greenhouse and Interior Plantscapes apply product with spray equipment such as back pack sprayer, hand pump sprayer, tank and hand-held spray gun or wand.

## **TURFGRASS DISEASE CONTROL**

ARMADA 50 WP is a preventive and curative fungicide that may be applied to turf sites including institutional, commercial and residential lawns, sod farms, sports fields, parks, municipal grounds and cemeteries.

ARMADA 50 WP is a broad spectrum fungicide for the control of dollar spot (*Sclerotinia homoeocarpa*), brown patch (*Rhizoctonia solani*), anthracnose (*Colletotrichum graminicola*), red thread (*Corticium fuciforme*), pink patch (*Limonomyces roseipellis*), leaf spot (*Bipolaris* spp., *Drechslera* spp.), gray leaf spot (*Pyricularia grisea*), rust (*Puccinia* spp.), pink snowmold (*Microdochium nivale*), fusarium patch (*Fusarium nivale*), southern blight (*Sclerotium rofskii*), stripe smut (*Ustilago striiformis*), rapid blight, and summer patch (*Magnaporthe poae*). When conditions are favorable for Pythium blight (*Pythium* spp.), mix a Pythium control fungicide with ARMADA 50 WP.

1. FUNGICIDES SUCH AS ARMADA 50 WP ARE BEST USED IN A PREVENTIVE DISEASE CONTROL PROGRAM. CURATIVE APPLICATIONS CAN BE MADE FOR CERTAIN DISEASES.
2. Apply in 1-2 gal of water per 1,000 sq ft to ensure thorough coverage. For soil-borne diseases, such as summer patch and take-all patch, use sufficient water to reach the crown and upper root zone, or irrigate immediately after application with sufficient water to move the active ingredient down into the crown and root zone of the turf.
3. Apply after mowing OR allow sprayed area to completely dry before mowing.
4. For control of both foliar and soil-borne diseases, allow sprayed area to completely dry before irrigation.
5. Under conditions optimum for high disease pressure, use the higher rate and the shorter interval.
6. For optimum turf quality and disease control, use ARMADA 50 WP in conjunction with turf management practices that promote good health and optimum disease control.
7. Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.

## **Plant Health**

Apply Armada 50 WP at 1.2 to 1.5 oz/1,000 sq ft to sod production of warm season grasses such as St. Augustine's, centipede, and Bermuda grass. The following plant health enhancements may be seen 30 to 60 days after application of Armada 50 WP; visible leaf greenness and density. Sod laid under shade trees also benefits in turf quality when this application is made prior to transfer. Apply Armada 50 WP at 1.2-1.5 oz /1,000 sq ft 15-30 days pre-harvest of sodfarm turf for plant health benefits.

## **Resistance Management for Turfgrass**

Do not apply more than 2 sequential ARMADA 50 WP fungicide applications for gray leaf spot control. Do not apply more than 3 sequential applications of ARMADA 50 WP fungicide for all other diseases. Alternate with a fungicide having a different mode of action.

## **Tank Mixes**

For more broad-spectrum control ARMADA 50 WP can be tank mixed with other fungicides. Check compatibility before tank mixing. ARMADA 50 WP CAN BE APPLIED TO ALL MAJOR TURFGRASS SPECIES. RATES ARE DEPENDENT UPON DISEASE AND THE ORGANISM THAT IT IS INCITED BY.

## Turfgrass Diseases Controlled with ARMADA 50 WP

Disease	ARMADA 50 WP	OZ OF ARMADA 50 WP PER 1,000 SQ FT	Application Interval/Timing
Dollar Spot	Apply when conditions are favorable for disease development.	0.6-1.5	14-28 days
Brown Patch	Apply when conditions are favorable for disease development.	0.6-1.5	14-28 days
Leaf Spot	Apply when conditions are favorable for disease development.	0.6	14 days
	For curative applications.	1.2-1.5	21-28 days
Anthracnose	Apply when conditions are favorable for disease development.	0.6-1.5	14-28 days
Summer Patch	Make initial application when the soil temperatures reach 65°F. Subsequent applications at 21-28 day interval. Do not make more than 3 sequential applications.	1.2-1.5	21-28 days
Gray Leaf Spot and Rapid Blight	Apply when conditions are favorable for disease development. For best results on Rapid Blight apply ARMADA 50 WP at the highest rate.	0.6-1.5	14-28 days
Red Thread, Pink Patch	Apply when conditions are favorable for disease development.	0.6-1.5	14-28 days
Rust	Apply when conditions are favorable for disease development.	0.6-1.5	14-28 days
Southern Blight	Preventive applications only	0.6-1.5	Repeat at 14-day intervals.
Stripe Smut	Make the first application in the spring just before the turf breaks dormancy followed by a second application just prior to the summer heat stress period and a third application when the cool nighttime temperatures of the late summer or early fall return.	0.6	3 applications/season Spring, Early Summer, and Early Fall
Fusarium Patch	Apply when conditions are favorable for disease development.	1.2-1.5	Fall-Early Spring
Pink Snow Mold	Apply one application late fall before snow cover or early spring after snow melts. Do not apply on top of snow.	1.2-1.5	Late Fall
Take-All Patch	Apply 1 to 2 applications preventatively.	1.2-1.5	28 days
Necrotic Ring Spot	Apply 1 to 2 applications preventatively.	1.2-1.5	28 days
Large Patch	Apply 1 to 2 applications preventatively when conditions are favorable for disease development.	1.2-1.5	28 days
Fairy Ring	First application is prescribed in the spring when 5 day average daily soil temperatures taken at 2 inches are 60-65°F. The second application is prescribed 14 to 28 days later. Make applications in 2 gallons water/1,000 sq ft. Water in application with 1/8 to 1/4 inch of irrigation within 6-8 hours after application. Tank mixing with Revolution wetting agent will decrease the fungicidal efficacy of Armada 50 WP. Apply wetting agents 2 weeks after the Armada 50 WP application on routine schedule for wetting agents.	1.2-1.5	14-28 days

For preventive applications apply when conditions are favorable for disease. Use the higher rate/shorter interval for curative applications or under severe disease conditions.

Use Rate - ARMADA 50WP Turf	Area Covered (square feet)	
	oz/1,000 sq ft	3 oz PVA Packet
0.6	5,000 sq ft	15,000 sq ft
1.2	2,500 sq ft	7,500 sq ft

### Maximum Use Rates

For turfgrass, up to 207.1 oz. of ARMADA 50 WP can be applied per acre per year.

### ORNAMENTAL DISEASE CONTROL

ARMADA 50 WP is a broad-spectrum fungicide for the control of certain foliar and stem diseases of ornamentals grown in interiorscapes (to include shopping malls, shopping centers, hotel lobbies) and residential and commercial landscapes.

**Foliar Diseases:** ARMADA 50 WP will control foliar diseases of ornamental plants when applied as a foliar spray to the plant species listed on this label. Apply 3 to 9 oz ARMADA 50 WP per 100 gallons of spray solution. Apply as a full-coverage spray to the point of drip and repeat at 14 to 28 day intervals until the threat of disease is over. Begin applications when conditions are favorable for disease development and continue until the threat of disease is over. Use higher rates or shorter intervals under high disease pressure.

**Spray Additives:** Use of spray additives such as spreaders, stickers, extenders, trace elements or fertilizers should be evaluated prior to use. The label directions given here are based on data obtained with no additives; use of any product with Armada 50 WP may affect the result.

## Plants

Flowering and Foliage Plants		Ornamental Shrubs, Trees, and Fruit Trees		Shade Trees	
Ageratum	Hollyhock	Amelanchier	Leucothoe	Ash	Maple
Aster	Hydrangea	Apple (non-bearing)	Lilac	Aspen	Oak
Begonia	Iris	Azalea	Mock-Orange	Birch	Pine
Calendula	Kalanchoe	Barberry	Mountain Laurel	Buckeye	Poplar
Canna	Marigold	Buckthorn	Ninebark	Chestnut	Russian Olive
Carnation	Nephthytis	Camellia	Paulownia	Cottonwood	Olive
Chrysanthemum	Pansy	Cedar	Pear (non-bearing)	Elm	Sycamore
Cineraria	Petunia	Cherry(non-bearing)	Photinia	Fir	Walnut
Crassula	Phlox	Crabapple	Pittosporum	Locust	Willow
Dahlia	Poinsettia	Crape myrtle	Plum (non-bearing)		
Daisy	Rose	Cypress, Leyland	Potentilla		
Delphinium	Salvia	Dogwood	Privet		
Dianthus	Sedum	Euonymus	Pyracantha		
Four O'Clock	Snapdragon	Gardenia	Rhododendron		
Geranium	Sunflowers	Hawthorn	Spirea		
Gerbera	Zinnia	Hemlock	Viburnam		
Grape Leaf Ivy		Holly	Vitex		
		Juniper			

**NOTICE TO USER:** Plant tolerance to ARMADA 50 WP has been found to be acceptable on all ornamentals that it has been tested on with the exception of Petunia, Violets, and New Guinea impatiens. Due to the large number of species and varieties of ornamentals plants, it is impossible to test every one for tolerance to ARMADA 50 WP. Neither the Manufacturer nor the Seller has determined whether or not ARMADA 50 WP can be used safely on ornamental plants not specified on this label. The professional user should determine if ARMADA 50 WP can be used safely prior to commercial use. In a small area test the recommended rates on a small number of plants for phytotoxicity prior to widespread use. Before using ARMADA 50 WP in tank mixture with other products, test the mixture on a small number of plants for phytotoxicity prior to widespread use.

Before using ARMADA 50 WP on plants that are not listed on this label, test ARMADA 50 WP on a small scale first.

### Common and scientific names of diseases controlled by ARMADA 50 WP

Common Name	Scientific Name		
Anthraxnose	<i>Apiognomonina veneta</i> <i>Colletotrichum gloeosporioides</i>	<i>Discula quercina</i> <i>Gloeosporium aridum</i>	<i>Glomerella cingulata</i>
Black spot	<i>Diplocarpon rosae</i>		
Downy mildew	<i>Peronospora</i> spp.		
Leaf spot	<i>Cercospora</i> spp.	<i>Entomosporium</i> spp.	<i>Septoria</i> spp.
Powdery mildew	<i>Erysiphe</i> spp. <i>Microsphaera</i> spp.	<i>Oidium</i> spp. <i>Podosphaera</i> spp.	<i>Sphaerotheca</i> spp. <i>Uncinula</i> spp.
Rust	<i>Coleosporium</i> spp. <i>Gymnosporangium</i> spp.	<i>Melampsorium</i> spp. <i>Phragmidium</i> spp.	<i>Puccinia</i> spp. <i>Uromyces</i> spp.
Scab	<i>Cladosporium</i> spp.		
		<i>Venturia inaequalis</i>	

Use Rate ARMADA 50WP Ornamentals	Total Spray Solution (gallons)	
Oz/100 Gallons	3 oz PVA Packet	9 oz PVA Packet
3	100 gallons	300 gallons
6	50 gallons	150 gallons
9	33 gallons	100 gallons

### Maximum Use Rates in Ornamentals

For ornamental plants do not exceed 207.1 oz of ARMADA 50 WP per acre per year.

### Restrictions

1. For ground application, use a minimum of 50 gals of water/A.
2. For aerial application, use a minimum of 10 gals of water/A.
3. To avoid spray drift, do not apply when conditions favor drift beyond the target area. Avoid spray overlap. For information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural experiment station recommendations.
4. Use of spray additives are not required. Any spray additive should be evaluated prior to use. Do not use in conjunction with organosilicate-based products, or plant injury may occur. Label directions are based on data with no additives.

### Resistance Management for Ornamentals

ARMADA 50 WP contains a modern, site-specific fungicide belonging to the strobilurin class of chemistry. Fungal pathogens are known to develop resistance to fungicides with a specific mode of action. When site-specific fungicides are introduced without a clear resistance management strategy, resistance development may be rapid, particularly with greenhouse use.

ARMADA 50 WP exhibits cross-resistance to other strobilurins and fungicides within the **Strobilurin Type Action** and **Resistance group (STAR compounds)**, but there is no known cross-resistance to fungicides of other classes including sterol inhibitors, dicarboximides, benzimidazoles, anilino-pyrimidines, phenylpyrroles, or phenylamides.

Many fungi which attack ornamentals and flowering plants including Botrytis and powdery mildews have a history of fungicide resistance development. Because resistance development cannot be predicted, implementation of suitable strategies to manage the resistance risk to ARMADA 50 WP is needed. To minimize the risk of resistance development to ARMADA 50 WP, the following practices are recommended.

1. Use ARMADA 50 WP preventively.
  2. For Leaf Spots and diseases other than Powdery Mildew, Downy Mildew, and Botrytis:
    - Use no more than two (2) applications of ARMADA 50 WP before rotating to another effective product that is not in the strobilurin class of chemistry for two (2) applications before rotating back to ARMADA 50 WP.
- OR**
- Rotate to another fungicide of nonstrobilurin chemistry after each ARMADA 50 WP application.
  3. For Powdery Mildew, Downy Mildew, and Botrytis:
    - Between each ARMADA 50 WP application, make two (2) applications of a fungicide of nonstrobilurin chemistry before rotating back to ARMADA 50 WP.
- OR**
- Rotate to another fungicide of nonstrobilurin chemistry after each ARMADA 50 WP application.
4. Make no more than four (4) foliar applications of ARMADA 50 WP per season for each at risk pathogen.

### IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, OR APPLICATION OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer (reg'd), the Bayer Cross (reg'd), Armada™ and Backed by Bayer™ are trademarks of Bayer.



Produced for:

**Bayer Environmental Science**

A Division of Bayer CropScience LP

2 T. W. Alexander Drive

Research Triangle Park, NC 27709