

# Lecture 2: Linear Regression

Feng Li

Shandong University

*fli@sdu.edu.cn*

December 28, 2021


# Lecture 2: Linear Regression

- 1 Supervised Learning: Regression and Classification
- 2 Linear Regression
- 3 Gradient Descent Algorithm
- 4 Stochastic Gradient Descent
- 5 Revisiting Least Square
- 6 A Probabilistic Interpretation to Linear Regression

# Supervised Learning


- Regression: Predict a continuous value
- Classification: Predict a discrete value, the class

Living area (feet <sup>2</sup> )	Price (1000\$s)
2104	400
1600	330
2400	369
1416	232
3000	540
⋮	⋮




# Supervised Learning (Contd.)

- Features: input variables,  $x$ ;
- Target: output variable,  $y$ ;
- Training example:  $(x^{(i)}, y^{(i)})$ ,  $i = 1, 2, 3, \dots, m$
- Hypothesis:  $h : \mathcal{X} \rightarrow \mathcal{Y}$ .




# Linear Regression

- Linear hypothesis:  $h(x) = \theta_1 x + \theta_0$ .
- $\theta_i$  ( $i = 1, 2$  for 2D cases): Parameters to estimate.
- How to choose  $\theta_i$ 's?




# Linear Regression (Contd.)



- Input: Training set  $(x^{(i)}, y^{(i)}) \in \mathbb{R}^2$  ( $i = 1, \dots, m$ )
- Goal: Model the relationship between  $x$  and  $y$  such that we can predict the corresponding target according to a given new feature.

# Linear Regression (Contd.)



- The relationship between  $x$  and  $y$  is modeled as a linear function.
- The linear function in the 2D plane is a straight line.
- Hypothesis:  $h_{\theta}(x) = \theta_0 + \theta_1 x$  (where  $\theta_0$  and  $\theta_1$  are parameters)

# Linear Regression (Contd.)

- Given data  $x \in \mathbb{R}^n$ , we then have  $\theta \in \mathbb{R}^{n+1}$
- Thus  $h_{\theta}(x) = \sum_{i=0}^n \theta_i x_i = \theta^T x$ , where  $x_0 = 1$
- What is the best choice of  $\theta$  ?


$$\min_{\theta} J(\theta) = \frac{1}{2} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

where  $J(\theta)$  is so-called a cost function



# Linear Regression (Contd.)

$$\min_{\theta} J(\theta) = \frac{1}{2} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$



## Definition

**Directional Derivative:** The directional derivative of function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  in the direction  $u \in \mathbb{R}^n$  is

$$\nabla_u f(x) = \lim_{h \rightarrow 0} \frac{f(x + hu) - f(x)}{h}$$

- $\nabla_u f(x)$  represents the rate at which  $f$  is increased in direction  $u$
- When  $u$  is the  $i$ -th standard unit vector  $e_i$ ,

$$\nabla_u f(x) = f'_i(x)$$

where  $f'_i(x) = \frac{\partial f(x)}{\partial x_i}$  is the partial derivative of  $f(x)$  w.r.t.  $x_i$

## Theorem

*For any  $n$ -dimensional vector  $u$ , the directional derivative of  $f$  in the direction of  $u$  can be represented as*

$$\nabla_u f(x) = \sum_{i=1}^n f'_i(x) \cdot u_i$$

# Gradient (Contd.)

## Proof.

Letting  $g(h) = f(x + hu)$ , we have

$$g'(0) = \lim_{h \rightarrow 0} \frac{g(h) - g(0)}{h} = \lim_{h \rightarrow 0} \frac{f(x + hu) - f(x)}{h} = \nabla_u f(x) \quad (1)$$

On the other hand, by the chain rule,

$$g'(h) = \sum_{i=1}^n f'_i(x) \frac{d}{dh}(x_i + hu_i) = \sum_{i=1}^n f'_i(x) u_i \quad (2)$$

Let  $h = 0$ , then  $g'(0) = \sum_{i=1}^n f'_i(x) u_i$ , by substituting which into (1), we complete the proof. □

## Definition

**Gradient:** The gradient of  $f$  is a vector function  $\nabla f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  defined by

$$\nabla f(x) = \sum_{i=1}^n \frac{\partial f}{\partial x_i} e_i$$

where  $e_i$  is the  $i$ -th standard unit vector. In another simple form,

$$\nabla f(x) = \left[ \frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right]^T$$

## Gradient (Contd.)

- $\nabla_u f(x) = \nabla f(x) \cdot u = \|\nabla f(x)\| \|u\| \cos a$  where  $a$  is the angle between  $\nabla f(x)$  and  $u$
- Without loss of generality, assume  $u$  is a unit vector,

$$\nabla_u f(x) = \|\nabla f(x)\| \cos a$$

- When  $u = \nabla f(x)$  such that  $a = 0$  (and thus  $\cos a = 1$ , we have the maximum directional derivative of  $f$ , which implies that  $\nabla f(x)$  is **the direction of steepest ascent** of  $f$ .

# Gradient Descent (GD) Algorithm

- If the multi-variable function  $J(\theta)$  is differentiable in a neighborhood of a point  $\theta$ , then  $J(\theta)$  decreases fastest if one goes from  $\theta$  in the direction of the negative gradient of  $J$  at  $\theta$
- Find a local minimum of a differentiable function using gradient descent

---

## Algorithm 1 Gradient Descent

---

- 1: **Given** a starting point  $\theta \in \text{dom } J$
  - 2: **repeat**
  - 3:   Calculate gradient  $\nabla J(\theta)$ ;
  - 4:   Update  $\theta \leftarrow \theta - \alpha \nabla J(\theta)$
  - 5: **until** convergence criterion is satisfied
- 

- $\theta$  is usually initialized randomly
- $\alpha$  is so-called learning rate

- Stopping criterion (i.e., conditions to convergence)
  - the gradient has its magnitude less than or equal to a predefined threshold (say  $\varepsilon$ ), i.e.

$$\|\nabla f(\mathbf{x})\|_2 \leq \varepsilon$$

where  $\|\cdot\|_2$  is  $\ell_2$  norm, such that the values of the objective function differ very slightly in successive iterations

- Set a fixed value for the maximum number of iterations, such that the algorithm is terminated after the number of the iterations exceeds the threshold.



# GD Algorithm (Contd.)

- In more details, we update each component of  $\theta$  according to the following rule


$$\theta_j \leftarrow \theta_j - \alpha \frac{\partial J(\theta)}{\partial \theta_j}, \quad \forall j = 0, 1, \dots, n$$

- Calculating the gradient for linear regression

$$\begin{aligned} \frac{\partial J(\theta)}{\partial \theta_j} &= \frac{\partial}{\partial \theta_j} \frac{1}{2} \sum_{i=1}^m (\theta^T x^{(i)} - y^{(i)})^2 \\ &= \frac{\partial}{\partial \theta_j} \frac{1}{2} \sum_{i=1}^m \left( \sum_{j=0}^n \theta_j x_j^{(i)} - y^{(i)} \right)^2 \\ &= \sum_{i=1}^m (\theta^T x^{(i)} - y^{(i)}) x_j^{(i)} \end{aligned}$$

# GD Algorithm (Contd.)

- An illustration of gradient descent algorithm
- The objective function is decreased fastest along the gradient



# GD Algorithm (Contd.)


- Another commonly used form

$$\min_{\theta} J(\theta) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

- What's the difference?
  - $m$  is introduced to scale the objective function to deal with differently sized training set.
- Gradient ascent algorithm
  - Maximize the differentiable function  $J(\theta)$
  - The gradient represents the direction along which  $J$  increases fastest
  - Therefore, we have

$$\theta_j \leftarrow \theta_j + \alpha \frac{\partial J(\theta)}{\partial \theta_j}$$

# Convergence under Different Step Sizes



# Stochastic Gradient Descent (SGD)

- What if the training set is huge?
  - In the above batch gradient descent algorithm, we have to run through the entire training set in each iteration
  - A considerable computation cost is induced!
- Stochastic gradient descent (SGD), also known as incremental gradient descent, is a stochastic approximation of the gradient descent optimization method
  - In each iteration, the parameters are updated according to the gradient of the error with respect to one training sample only

# Stochastic Gradient Descent (Contd.)

---

**Algorithm 2** Stochastic Gradient Descent for Linear Regression

---

- 1: **Given** a starting point  $\theta \in \text{dom } J$
  - 2: **repeat**
  - 3:   Randomly shuffle the training data;
  - 4:   **for**  $i = 1, 2, \dots, m$  **do**
  - 5:      $\theta \leftarrow \theta - \alpha \nabla J(\theta; x^{(i)}, y^{(i)})$
  - 6:   **end for**
  - 7: **until** convergence criterion is satisfied
-

# More About SGD

- The objective does not always decrease for each iteration
- Usually, SGD has  $\theta$  approaching the minimum much faster than batch GD
- SGD may never converge to the minimum, and oscillating may happen
- A variants: Mini-batch, say pick up a small group of samples and do average, which may accelerate and smoothen the convergence

# Matrix Derivatives<sup>1</sup>

- A function  $f : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$
- The derivative of  $f$  with respect to  $A$  is defined as

$$\nabla f(A) = \begin{bmatrix} \frac{\partial f}{\partial A_{11}} & \cdots & \frac{\partial f}{\partial A_{1n}} \\ \vdots & \ddots & \vdots \\ \frac{\partial f}{\partial A_{m1}} & \cdots & \frac{\partial f}{\partial A_{mn}} \end{bmatrix}$$

- For an  $n \times n$  matrix, its trace is defined as  $\text{tr}A = \sum_{i=1}^n A_{ii}$ 
  - $\text{tr}ABCD = \text{tr}DABC = \text{tr}CDAB = \text{tr}BCDA$
  - $\text{tr}A = \text{tr}A^T$ ,  $\text{tr}(A+B) = \text{tr}A + \text{tr}B$ ,  $\text{tr}A = a \text{tr}A$
  - $\nabla_A \text{tr}AB = B^T$ ,  $\nabla_{A^T} f(A) = (\nabla_A f(A))^T$
  - $\nabla_A \text{tr}ABA^T C = CAB + C^T AB^T$ ,  $\nabla_A |A| = |A|(A^{-1})^T$
  - Funky trace derivative  $\nabla_{A^T} \text{tr}ABA^T C = B^T A^T C^T + BA^T C$

---

<sup>1</sup>Details can be found in “Properties of the Trace and Matrix Derivatives” by John Duchi



# Revisiting Least Square

- Assume

$$X = \begin{bmatrix} (x^{(1)})^T \\ \vdots \\ (x^{(m)})^T \end{bmatrix} \quad Y = \begin{bmatrix} y^{(1)} \\ \vdots \\ y^{(m)} \end{bmatrix}$$

- Therefore, we have

$$X\theta - Y = \begin{bmatrix} (x^{(1)})^T \theta \\ \vdots \\ (x^{(m)})^T \theta \end{bmatrix} - \begin{bmatrix} y^{(1)} \\ \vdots \\ y^{(m)} \end{bmatrix} = \begin{bmatrix} h_{\theta}(x^{(1)}) - y^{(1)} \\ \vdots \\ h_{\theta}(x^{(m)}) - y^{(m)} \end{bmatrix}$$

- $J(\theta) = \frac{1}{2} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2 = \frac{1}{2} (X\theta - Y)^T (X\theta - Y)$

## Revisiting Least Square (Contd.)

- Minimize  $J(\theta) = \frac{1}{2}(Y - X\theta)^T(Y - X\theta)$
- Calculate its derivatives with respect to  $\theta$

$$\begin{aligned}\nabla_{\theta} J(\theta) &= \nabla_{\theta} \frac{1}{2}(Y - X\theta)^T(Y - X\theta) \\&= \frac{1}{2} \nabla_{\theta} (Y^T - \theta^T X^T)(Y - X\theta) \\&= \frac{1}{2} \nabla_{\theta} \text{tr}(Y^T Y - Y^T X\theta - \theta^T X^T Y + \theta^T X^T X\theta) \\&= \frac{1}{2} \nabla_{\theta} \text{tr}(\theta^T X^T X\theta) - X^T Y \\&= \frac{1}{2} (X^T X\theta + X^T X\theta) - X^T Y \\&= X^T X\theta - X^T Y\end{aligned}$$

- Tip: Funky trace derivative  $\nabla_A \text{tr} ABA^T C = B^T A^T C^T + BA^T C$

# Revisiting Least Square (Contd.)

- **Theorem:**

The matrix  $A^T A$  is invertible if and only if the columns of  $A$  are linearly independent. In this case, there exists only one least-squares solution

$$\theta = (X^T X)^{-1} X^T Y$$

- Prove the above theorem in Problem Set 1.

# Probabilistic Interpretation

- The target variables and the inputs are related

$$y = \theta^T x + \epsilon$$

- $\epsilon$ 's denote the errors and are independently and identically distributed (i.i.d.) according to a Gaussian distribution  $\mathcal{N}(0, \sigma^2)$
- The density of  $\epsilon^{(i)}$  is given by

$$f(\epsilon) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{\epsilon^2}{2\sigma^2}\right)$$

- The conditional probability density function of  $y$

$$y \mid x; \theta \sim \mathcal{N}(\theta^T x, \sigma^2)$$

# Probabilistic Interpretation (Contd.)

- The training data  $\{x^{(i)}, y^{(i)}\}_{i=1, \dots, m}$  are sampled identically and independently

$$p(y = y^{(i)} \mid x = x^{(i)}; \theta) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y^{(i)} - \theta^T x^{(i)})^2}{2\sigma^2}\right)$$

- Likelihood function

$$\begin{aligned} L(\theta) &= \prod_i p(y^{(i)} \mid x^{(i)}; \theta) \\ &= \prod_i \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y^{(i)} - \theta^T x^{(i)})^2}{2\sigma^2}\right) \end{aligned}$$

# Probabilistic Interpretation (Contd.)

- Maximizing the likelihood  $L(\theta)$
- Since  $L(\theta)$  is complicated, we maximize an increasing function of  $L(\theta)$  instead

$$\begin{aligned}\ell(\theta) &= \log L(\theta) \\ &= \log \prod_i^m \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y^{(i)} - \theta^T x^{(i)})^2}{2\sigma^2}\right) \\ &= \sum_i^m \log \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(y^{(i)} - \theta^T x^{(i)})^2}{2\sigma^2}\right) \\ &= m \log \frac{1}{\sqrt{2\pi}\sigma} - \frac{1}{2\sigma^2} \sum_i (y^{(i)} - \theta^T x^{(i)})^2\end{aligned}$$

- Apparently, maximizing  $L(\theta)$  (thus  $\ell(\theta)$ ) is equivalent to minimizing

$$\frac{1}{2} \sum_i^m (y^{(i)} - \theta^T x^{(i)})^2$$

# Thanks!

Q & A