

Lumbar Spine Stenosis Algorithm

Definition: Clinical Syndrome of gradual onset of Intermittent buttock or lower extremity pain, with or without back pain^{1,8}. Symptoms are aggravated by standing or walking (neurogenic claudication^{3,6}) and relieved by sitting or leaning forward^{1,3,4,5,7,8}. Spinal stenosis is associated with decreased space available for neural elements (thecal sac and/or exiting nerve roots³) and epidural veins in the lumbar spine^{1,7,8}.

The typical patient has a stooped forward posture, restricted lumbar extension, thigh pain with 30 seconds of lumbar extension.

Must differentiate from other conditions such as:

Peripheral Vascular Disease (Measure ankle brachial index)^{1,2,3,4,5,6,8}

Diabetes^{1,4,6,8}

Peripheral Neuropathy^{1,4,5,6,8}

Hip/Knee Osteoarthritis^{1,5,6,8}

```
graph TD; A[Initial assessment of any back pain patient] --> B[Establish Standardized Score:  
Ask patient to complete the Zurich Claudication Questionnaire9 and document score  
Ask patient to complete the STarT Back Screening Tool (SBST) and document score]; B --> C[Assess your patient:  
Conduct a focused history and physical examination  
Review the standardized questionnaire scores  
Classify your patient into one of five broach categories];
```

Initial assessment of any back pain patient

Establish Standardized Score:

Ask patient to complete the Zurich Claudication Questionnaire⁹ and document score

Ask patient to complete the STarT Back Screening Tool (SBST) and document score

Assess your patient:

Conduct a focused history and physical examination

Review the standardized questionnaire scores

Classify your patient into one of five broach categories

Presence of Red Flags:

New Bowel/Bladder incontinence or retention; recent severe trauma; progressive paraparesis, quadraparesis, neurologic signs → Send to emergency in your RUIS (Réseau Universitaire Intégré de Santé)

Unexplained weight loss (>10 lbs over 6 months), fever, chills, saddle anaesthesia without new bowel/bladder incontinence or retention; acute pain not eased by recumbent position; incremental non-relenting pain → page spine resident in the Emergency room of your RUIS

Unsteadiness of gait, weakness, numb/clumsy fingers → go to myelopathy algorithm

Lumbar Spinal Stenosis
Typical patient > 50yr old^{1,3,4,5,8}

Non-Specific Low Back Pain (NSLBP) → Go to Acute/subacute non-specific low back pain algorithm

Unilateral leg pain below the knee (with/without numbness and weakness) → Go to radiculopathy algorithm

Educate:

Reassure patients → development of cauda equina syndrome or severe progressive neurologic deficit rare^{3,4,5,6,7}.

Advise patients to stay active

Promote self management

Discourage bed rest

Prescribe medication for pain relief (if needed):

Gabapentin^{4,10}

NSAIDs^{1,3,6}

Non-narcotic analgesics⁶

Narcotic analgesics (short course of maximum 2 weeks only if necessary)^{3,5}

Prescribe Physiotherapy:

Strengthening of core muscles^{1,3,5,11}

Stretching of lower extremity muscles (hamstrings, quadriceps, hip flexors)^{1,11}

Lumbar flexion exercises (e.g. cycling)^{3,4,5,11}

Avoid Lumbar extension exercises^{4,8}

Elastic Lumbar binder (Wear only for brief periods to avoid deconditioning of para-spinal muscles)⁸

Follow-up visit (4-6 weeks after initial visit):

Purpose: keep the diagnosis under review and re-assess patient's symptoms

Establish standardized scores again

Address any yellow flags or red flags

Improvement

Continue with same treatment
Re-assess regularly (every 6 months)

No improvement

Diagnostic Imaging:
MRI or CT myelograph (if MRI inconclusive or contraindicated)
CT scan (if MRI and CT myelograph are inconclusive, contraindicated or inappropriate)
Maintain intermittent communication with patient while awaiting imaging results

MRI reported as mild-moderate spinal canal or neuroforaminal stenosis

Fluoroscopically guided epidural injection (ESI)^{1,12,13,14}
by:
Physiatrist
Pain management specialist


Follow-up in 3-4 weeks

MRI reported as severe spinal canal or neuroforaminal stenosis

Send the McGill Consult referral Form to the McGill Spine Program

MRI showed no evidence of spinal canal or neuroforaminal stenosis

Not a neurocompressive disorder
Further work-up for neuropathy/other pathology
Refer patient to Neurologist for further assessment



Improvement

No-Improvement

Prescribe
Physiotherapy

Send the McGill
Consult referral
Form to the McGill
Spine Program


Follow-up in 3-4 weeks

Redevelopment of symptoms

Continued symptom resolution

2nd ESI (improvement with 1st) or
opioid analgesics

Continue with same treatment
Reassess regularly (every 6 months)



Triage

Pre-visit Triage: Triager reviews patient consult and MRI/CT

Spine program initial visit: Keep diagnosis under review, looking for pathoanatomical causes related to symptoms

Establish standardized score:
Ask patient to complete the spine program survey (Including the Zurich Claudication Questionnaire) and compare to latest score from referring physician's office

Assess patient and make a decision

Tertiary care not required

Patient reassurance provided

Patient needs further non-operative specialized, multidisciplinary care

Psychologist
Physiatrist
Physiotherapist
GP – Sports Medicine

Patient is a surgical candidate

Request further imaging for surgical planning:
Standing AP and lateral
Scoliosis series
Flexion/Extension X-rays of
the Lumbar Spine

Surgeon decides on different surgical options:
Decompressive Surgery
Lumbar fusion

References

1. Doorly TP, Lambing CL, Malanga GA, Maurer PM, Rashbaum RF. Algorithmic approach to the management of the patient with lumbar spinal stenosis. *J Fam Pract.* 2010 Aug;59(8 Suppl Algorithmic):S1-8.
2. Manchikanti L, Helm S, Singh V, Benyamin RM, Datta S, Hayek SM, Fellows B, Boswell MV; ASIPP. An algorithmic approach for clinical management of chronic spinal pain. *Pain Physician.* 2009 Jul-Aug;12(4):E225-64.
3. Issack PS, Cunningham ME, Pumberger M, Hughes AP, Cammisa FP Jr. Degenerative lumbar spinal stenosis: evaluation and management. *J Am Acad Orthop Surg.* 2012 Aug;20(8):527-35.
4. Markman JD, Frazer ME, Girgis PS, McCormick KR. Diagnosis and Management Approaches to Lumbar Spinal Stenosis. *J Current clinical care.* 2011 Jan
5. Haig AJ, Tomkins CC. Diagnosis and management of lumbar spinal stenosis. *JAMA.* 2010 Jan 6;303(1):71-2.
6. Alexander JT. Lumbar Spinal Stenosis: Diagnosis and Treatment Options. Available from: <http://www.dcmsonline.org/jax-medicine/1999journals/june1999/lumbar.htm>
7. Kreiner DS, Baisden J, Gilbert T, Shaffer WO, Summers J, Toton J, Hwang S, Mendel R, Reitman C. Diagnosis and Treatment of Degenerative Lumbar Spinal Stenosis; Clinical Guidelines for Multidisciplinary Spine Care. North American Spine Society 2011
8. Yuan PS, Albert TJ. Managing degenerative lumbar spinal stenosis. *The Journal of Musculoskeletal Medicine.* 2009 June. Vol. 26 No. 6
9. Stucki G, Daltroy L, Liang MH, Lipson SJ, Fossel AH, Katz JN. Measurement properties of a self-administered outcome measure in lumbar spinal stenosis. *Spine (Phila Pa 1976).* 1996 Apr 1;21(7):796-803.
10. Yaksi A, Ozgönel L, Ozgönel B. The efficiency of gabapentin therapy in patients with lumbar spinal stenosis. *Spine (Phila Pa 1976).* 2007 Apr 20;32(9):939-42.
11. Goren A, Yildiz N, Topuz O, Findikoglu G, Ardic F. Efficacy of exercise and ultrasound in patients with lumbar spinal stenosis: a prospective randomized controlled trial. *Clin Rehabil.* 2010 Jul;24(7):623-31. Epub 2010 Jun 8.
12. Fukusaki M, Kobayashi I, Hara T, Sumikawa K. Symptoms of spinal stenosis do not improve after epidural steroid injection. *Clin J Pain.* 1998 Jun;14(2):148-51
13. Koc Z, Ozcakar S, Sivrioglu K, Gurbet A, Kucukoglu S. Effectiveness of physical therapy and epidural steroid injections in lumbar spinal stenosis. *Spine (Phila Pa 1976).* 2009 May 1;34(10):985-9.
14. Cuckler JM, Bernini PA, Wiesel SW, Booth RE Jr, Rothman RH, Pickens GT. The use of epidural steroids in the treatment of lumbar radicular pain. A prospective, randomized, double-blind study. *J Bone Joint Surg Am.* 1985 Jan;67(1):63-6.