

Physics Reference Tables*

Contents:

Table A. Metric Prefixes.....	1	Table R. Absolute Indices of Refraction.....	10
Table B. Physical Constants.....	2	Table S. Fluid Mechanics Formulas and Equations.....	11
Table C. Quantities, Variables and Units.....	3	Table T. Planetary Data.....	11
Table D. Mechanics Formulas and Equations.....	4	Table U. Sun & Moon Data.....	11
Table E. Approximate Coefficients of Friction.....	4	Table V. Atomic & Particle Physics (Modern Physics).....	12
Table F. Angular/Rotational Mechanics Formulas and Equations.....	5	Figure W. Quantum Energy Levels.....	12
Table G. Moments of Inertia.....	5	Figure X. Particle Sizes.....	13
Table H. Heat and Thermal Physics Formulas and Equations ..	6	Figure Y. Classification of Matter.....	13
Table I. Thermal Properties of Selected Materials.....	6	Table Z. The Standard Model of Elementary Particles.....	13
Table J. Electricity Formulas & Equations.....	7	Figure AA. Periodic Table of the Elements.....	14
Table K. Electricity & Magnetism Formulas & Equations.....	8	Table BB. Symbols Used in Nuclear Physics.....	15
Table L. Resistor Color Code.....	8	Table CC. Selected Radioisotopes.....	15
Table M. Symbols Used in Electrical Circuit Diagrams.....	8	Table DD. Constants Used in Nuclear Physics.....	15
Table N. Resistivities at 20°C.....	8	Figure EE. Neutron/Proton Stability Band.....	15
Table O. Waves & Optics Formulas & Equations.....	9	Table FF. Geometry & Trigonometry Formulas.....	16
Figure P. The Electromagnetic Spectrum.....	9	Table GG. Values of Trigonometric Functions.....	17
Table Q. Properties of Water and Air.....	10	Table HH. Some Exact and Approximate Conversions.....	18
		Table II. Greek Alphabet.....	18

Table A. Metric Prefixes			
Factor		Prefix	Symbol
1 000 000 000 000 000 000 000 000	10^{24}	yotta	Y
1 000 000 000 000 000 000 000 000	10^{21}	zeta	Z
1 000 000 000 000 000 000 000	10^{18}	exa	E
1 000 000 000 000 000 000	10^{15}	peta	P
1 000 000 000 000	10^{12}	tera	T
1 000 000 000	10^9	giga	G
1 000 000	10^6	mega	M
1 000	10^3	kilo	k
100	10^2	hecto	h
10	10^1	deca	da
1	10^0	—	—
0.1	10^{-1}	deci	d
0.01	10^{-2}	centi	c
0.001	10^{-3}	milli	m
0.000 001	10^{-6}	micro	μ
0.000 000 001	10^{-9}	nano	n
0.000 000 000 001	10^{-12}	pico	p
0.000 000 000 000 001	10^{-15}	femto	f
0.000 000 000 000 000 001	10^{-18}	atto	a
0.000 000 000 000 000 000 001	10^{-21}	zepto	z
0.000 000 000 000 000 000 000 001	10^{-24}	yocto	y

↑ ↓
 ↑ ↓
 ↑ ↓
 ↑ ↓
 ↑ ↓
 ↑ ↓
 ↑ ↓
 ↑ ↓
 ↑ ↓
 ↑ ↓
 ↑ ↓
 ↑ ↓
 ↑ ↓
 ↑ ↓

Move Decimal Point to the Left
 Move Decimal Point to the Right

* Data from various sources, including: The University of the State of New York, The State Education Department. Albany, NY, *Reference Tables for Physical Setting/Physics, 2006 Edition.* <http://www.p12.nysed.gov/apda/reftable/physics-rt/physics06tbl.pdf>, SparkNotes: SAT Physics website. <http://www.sparknotes.com/testprep/books/sat2/physics/>, The Engineering Toolbox: <https://www.engineeringtoolbox.com>, and The College Board: *Equations and Constants for AP® Physics 1 and AP® Physics 2.*

Table B. Physical Constants			
Description	Symbol	Precise Value	Common Approximation
acceleration due to gravity / strength of gravity field on Earth's surface	g	$9.7639 \frac{m}{s^2}$ to $9.8337 \frac{m}{s^2}$ average value at sea level is $9.80665 \frac{m}{s^2}$	$9.8 \frac{m}{s^2} \equiv 9.8 \frac{N}{kg}$ or $10 \frac{m}{s^2} \equiv 10 \frac{N}{kg}$
universal gravitational constant	G	$6.67384(80) \times 10^{-11} \frac{Nm^2}{kg^2}$	$6.67 \times 10^{-11} \frac{Nm^2}{kg^2}$
speed of light in a vacuum	c	$299792458 \frac{m}{s}^*$	$3.00 \times 10^8 \frac{m}{s}$
elementary charge (proton or electron)	e	$\pm 1.602176634 \times 10^{-19} C^*$	$\pm 1.60 \times 10^{-19} C$
1 coulomb (C)		$6.241509074 \times 10^{18}$ elementary charges	6.24×10^{18} elementary charges
(electric) permittivity of a vacuum	ϵ_0	$8.85418782 \times 10^{-12} \frac{A^2 \cdot s^4}{kg \cdot m^3}$	$8.85 \times 10^{-12} \frac{A^2 \cdot s^4}{kg \cdot m^3}$
(magnetic) permeability of a vacuum	μ_0	$4\pi \times 10^{-7} = 1.25663706 \times 10^{-6} \frac{T \cdot m}{A}$	$1.26 \times 10^{-6} \frac{T \cdot m}{A}$
electrostatic constant	k	$\frac{1}{4\pi\epsilon_0} = 8.9875517873681764 \times 10^9 \frac{Nm^2}{C^2}^*$	$8.99 \times 10^9 \frac{Nm^2}{C^2}$
1 electron volt (eV)		$1.602176565(35) \times 10^{-19} J$	$1.60 \times 10^{-19} J$
Planck's constant	h	$6.62607015 \times 10^{-34} J \cdot s^*$	$6.63 \times 10^{-34} J \cdot s$
1 universal (atomic) mass unit (u)		$931.494061(21) MeV/c^2$ $1.660538921(73) \times 10^{-27} kg$	$931 MeV/c^2$ $1.66 \times 10^{-27} kg$
Avogadro's constant	N_A	$6.02214076 \times 10^{23} mol^{-1}^*$	$6.02 \times 10^{23} mol^{-1}$
Boltzmann constant	k_B	$1.380649 \times 10^{-23} \frac{J}{K}^*$	$1.38 \times 10^{-23} \frac{J}{K}$
universal gas constant	R	$8.3144621(75) \frac{J}{mol \cdot K}$	$8.31 \frac{J}{mol \cdot K}$
Rydberg constant	R_H	$\frac{m_e e^4}{8\epsilon_0^2 h^3 c} = 10973731.6 \frac{1}{m}$	$1.10 \times 10^7 m^{-1}$
Stefan-Boltzmann constant	σ	$\frac{2\pi^5 R^4}{15h^3 c^2} = 5.670374419 \times 10^{-8} \frac{J}{m^2 \cdot s \cdot K^4}$	$5.67 \times 10^{-8} \frac{J}{m^2 \cdot s \cdot K^4}$
standard atmospheric pressure at sea level		$101325 Pa \equiv 1.01325 bar^*$	$100000 Pa \equiv 1.0 bar$
rest mass of an electron	m_e	$9.10938215(45) \times 10^{-31} kg$	$9.11 \times 10^{-31} kg$
mass of a proton	m_p	$1.672621777(74) \times 10^{-27} kg$	$1.67 \times 10^{-27} kg$
mass of a neutron	m_n	$1.674927351(74) \times 10^{-27} kg$	$1.67 \times 10^{-27} kg$

*denotes an exact value (by definition)

Table C. Quantities, Variables and Units				
Quantity	Variable	MKS Unit Name	MKS Unit Symbol	S.I. Base Unit
position	\vec{x}	meter*	m	m
distance/displacement, (length, height)	$d, \vec{d}, (L, h)$	meter*	m	m
angle	θ	radian, degree	—, °	—
area	A	square meter	m ²	m ²
volume	V	cubic meter, liter	m ³	m ³
time	t	second*	s	s
velocity	\vec{v}	meter/second	$\frac{m}{s}$	$\frac{m}{s}$
speed of light	c		$\frac{m}{s}$	$\frac{m}{s}$
angular velocity	$\vec{\omega}$	radians/second	$\frac{1}{s^2}, s^{-1}$	$\frac{1}{s^2}, s^{-1}$
acceleration	\vec{a}	meter/second ²	$\frac{m}{s^2}$	$\frac{m}{s^2}$
acceleration due to gravity	\vec{g}		$\frac{m}{s^2}$	$\frac{m}{s^2}$
angular acceleration	$\vec{\alpha}$	radians/second ²	$\frac{1}{s^2}, s^{-2}$	$\frac{1}{s^2}, s^{-2}$
mass	m	kilogram*	kg	kg
force	\vec{F}	newton	N	$\frac{kg \cdot m}{s^2}$
gravitational field	\vec{g}	newton/kilogram	$\frac{N}{kg}$	$\frac{m}{s^2}$
pressure	P	pascal	Pa	$\frac{kg}{m \cdot s^2}$
energy (generic)	E			
potential energy	U			
kinetic energy	K, E_k	joule	J	$\frac{kg \cdot m^2}{s^2}$
heat	Q			
work	W	joule, newton-meter	J, N·m	$\frac{kg \cdot m^2}{s^2}$
torque	$\vec{\tau}$	newton-meter	N·m	$\frac{kg \cdot m^2}{s^2}$
power	P	watt	W	$\frac{kg \cdot m^2}{s^3}$
momentum	\vec{p}	newton-second	N·s	$\frac{kg \cdot m}{s}$
impulse	\vec{j}			$\frac{kg \cdot m}{s}$
moment of inertia	I	kilogram-meter ²	kg·m ²	kg·m ²
angular momentum	\vec{L}	newton-meter-second	N·m·s	$\frac{kg \cdot m^2}{s}$
frequency	f	hertz	Hz	s ⁻¹
wavelength	λ	meter	m	m
period	T	second	s	s
index of refraction	n	—	—	—
electric current	\vec{I}	ampere*	A	A
electric charge	q	coulomb	C	A·s
electric potential potential difference (voltage) electromotive force (emf)	V ΔV ϵ	volt	V	$\frac{kg \cdot m^2}{A \cdot s^3}$
electrical resistance	R	ohm	Ω	$\frac{kg \cdot m^2}{A^2 \cdot s^3}$
capacitance	C	farad	F	$\frac{A^2 \cdot s^4}{m^2 \cdot kg}$
electric field	\vec{E}	newton/coulomb volt/meter	$\frac{N}{C}, \frac{V}{m}$	$\frac{kg \cdot m}{A \cdot s^3}$
magnetic field	\vec{B}	tesla	T	$\frac{kg}{A \cdot s^2}$
temperature	T	kelvin*	K	K
amount of substance	n	mole*	mol	mol
luminous intensity	I_v	candela*	cd	cd

Variables representing vector quantities are typeset in **bold italics** with **arrows**. * = S.I. base unit

Table D. Mechanics Formulas and Equations	
<p>Kinematics (Distance, Velocity & Acceleration)</p>	$\vec{d} = \Delta\vec{x} = \vec{x} - \vec{x}_o$ $\frac{\vec{x} - \vec{x}_o}{t} = \frac{\vec{d}}{t} = \frac{\vec{v}_o + \vec{v}}{2} = \vec{v}_{ave.}$ $\Delta\vec{v} = \vec{v} - \vec{v}_o = \vec{a}t$ $\vec{x} - \vec{x}_o = \vec{d} = \vec{v}_o t + \frac{1}{2}\vec{a}t^2$ $\vec{v}^2 - \vec{v}_o^2 = 2\vec{a}\vec{d} = 2\vec{a}(\vec{x} - \vec{x}_o)$
<p>Forces & Dynamics</p>	$\sum \vec{F} = \vec{F}_{net} = m\vec{a}$ $F_f \leq \mu_s F_N \quad F_f = \mu_k F_N$ $\vec{F}_g = m\vec{g} = \frac{Gm_1 m_2}{r^2}$
<p>Circular/ Centripetal Motion & Force</p>	$a_c = \frac{v^2}{r}$ $F_c = ma_c$
<p>Simple Harmonic Motion</p>	$T = \frac{1}{f}$ $T_s = 2\pi\sqrt{\frac{m}{k}} \quad T_p = 2\pi\sqrt{\frac{L}{g}}$ $\vec{F}_s = -k\vec{x}$ $U_s = \frac{1}{2}kx^2$
<p>Momentum</p>	$\vec{p} = m\vec{v}$ $\sum m_i \vec{v}_i + \vec{J} = \sum m_f \vec{v}_f$ $\vec{J} = \Delta\vec{p} = \vec{F}_{net} \Delta t$
<p>Energy, Work & Power</p>	$U_g = mgh = \frac{Gm_1 m_2}{r}$ $K = \frac{1}{2}mv^2 = \frac{p^2}{2m}$ $W = \Delta E = \Delta(K + U)$ $W = F_{ }d = \vec{F} \bullet \vec{d} = Fd \cos \theta$ $E_{total} = U + K (+Q + \dots)$ $P = \frac{W}{t} = \vec{F} \bullet \vec{v} = Fv \cos \theta$
<p><i>var.</i> = name of quantity (unit)</p> <p>Δ = change in something (E.g., Δx means change in x)</p> <p>Σ = sum</p> <p>d = distance (m)</p> <p>\vec{d} = displacement (m)</p> <p>\vec{x} = position (m)</p> <p>t = time (s)</p> <p>\vec{v} = velocity ($\frac{m}{s}$)</p> <p>$\vec{v}_{ave.}$ = average velocity ($\frac{m}{s}$)</p> <p>\vec{a} = acceleration ($\frac{m}{s^2}$)</p> <p>f = frequency (Hz = $\frac{1}{s}$)</p> <p>\vec{F} = force (N)</p> <p>\vec{F}_{net} = net force (N)</p> <p>F_f = force due to friction (N)</p> <p>\vec{F}_g = force due to gravity (N)</p> <p>\vec{F}_N = normal force (N)</p> <p>m = mass (kg)</p> <p>\vec{g} = acceleration due to gravity = $10 \frac{m}{s^2}$ on Earth</p> <p>\vec{g} = strength of gravity field = $10 \frac{N}{kg}$ on Earth</p> <p>G = gravitational constant = $6.67 \times 10^{-11} \frac{N \cdot m^2}{kg^2}$</p> <p>$r$ = radius (m)</p> <p>μ = coefficient of friction* (dimensionless)</p> <p>θ = angle ($^\circ$, rad)</p> <p>k = spring constant ($\frac{N}{m}$)</p> <p>\vec{x} = displacement of spring (m)</p> <p>L = length of pendulum (m)</p> <p>E = energy (J)</p> <p>$K = E_k$ = kinetic energy (J)</p> <p>U = potential energy (J)</p> <p>h = height (m)</p> <p>Q = heat (J)</p> <p>P = power (W)</p> <p>W = work (J, N·m)</p> <p>T = (time) period (Hz)</p> <p>\vec{p} = momentum (N·s)</p> <p>\vec{J} = impulse (N·s)</p> <p>π = pi (mathematical constant) = 3.14159 26535 89793...</p> <p>*characteristic property of a substance (to be looked up)</p>	

Table E. Approximate Coefficients of Friction					
Substance	Static (μ_s)	Kinetic (μ_k)	Substance	Static (μ_s)	Kinetic (μ_k)
rubber on concrete (dry)	0.90	0.68	wood on wood (dry)	0.42	0.30
rubber on concrete (wet)		0.58	wood on wood (wet)	0.2	
rubber on asphalt (dry)	0.85	0.67	wood on metal	0.3	
rubber on asphalt (wet)		0.53	wood on brick	0.6	
rubber on ice		0.15	wood on concrete	0.62	
steel on ice	0.03	0.01	Teflon on Teflon	0.04	0.04
waxed ski on snow	0.14	0.05	Teflon on steel	0.04	0.04
aluminum on aluminum	1.2	1.4	graphite on steel	0.1	
cast iron on cast iron	1.1	0.15	leather on wood	0.3–0.4	
steel on steel	0.74	0.57	leather on metal (dry)	0.6	
copper on steel	0.53	0.36	leather on metal (wet)	0.4	
diamond on diamond	0.1		glass on glass	0.9–1.0	0.4
diamond on metal	0.1–0.15		metal on glass	0.5–0.7	

Table F. Angular/Rotational Mechanics Formulas and Equations		
Angular Kinematics (Distance, Velocity & Acceleration)	$\Delta\vec{\theta} = \vec{\theta} - \vec{\theta}_o$ $\frac{\vec{\theta} - \vec{\theta}_o}{t} = \frac{\Delta\vec{\theta}}{t} = \frac{\vec{\omega}_o + \vec{\omega}}{2} = \vec{\omega}_{ave.}$ $\Delta\vec{\omega} = \vec{\omega} - \vec{\omega}_o = \vec{\alpha}t$ $\Delta\vec{\theta} = \vec{\omega}_o t + \frac{1}{2}\vec{\alpha}t^2$ $\vec{\omega}^2 - \vec{\omega}_o^2 = 2\vec{\alpha}(\Delta\vec{\theta})$	var. = name of quantity (unit) Δ = change in something (E.g., Δx = change in x) Σ = sum s = arc length (m) t = time (s) a_c = centripetal acceleration ($\frac{m}{s^2}$)
Circular/Centripetal Motion	$s = r\Delta\theta \quad v_r = r\omega \quad a_r = r\alpha$ $a_c = \frac{v^2}{r} = \omega^2 r$	F_c = centripetal force (N) m = mass (kg) r = radius (m) \vec{r} = radius (vector) θ = angle ($^\circ$, rad) $\vec{\omega}$ = angular velocity ($\frac{rad}{s}$) $\vec{\alpha}$ = angular velocity ($\frac{rad}{s^2}$)
Rotational Dynamics	$x_{cm} = \frac{\sum m_i x_i}{\sum m_i}$ $I = \int_0^m r^2 dm$ $F_c = ma_c = mr\omega^2$ $\vec{\tau} = \vec{r} \times \vec{F} \quad \tau = rF \sin\theta = r_{\perp} F$ $\sum \vec{\tau} = \vec{\tau}_{net} = I\vec{\alpha}$	$\vec{\tau}$ = torque (N·m) x = position (m) f = frequency (Hz) A = amplitude (m) ϕ = phase offset ($^\circ$, rad) E = energy (J)
Simple Harmonic Motion	$T = \frac{1}{f} = \frac{2\pi}{\omega}$ $x = A \cos(2\pi ft) + \phi$	$K = E_k$ = kinetic energy (J) K_t = translational kinetic energy (J) K_r = rotational kinetic energy (J)
Angular Momentum	$\vec{L} = \vec{r} \times \vec{p} = I\vec{\omega} \quad L = rp \sin\theta = I\omega$ $\Delta\vec{L} = \vec{\tau}\Delta t$	P = power (W) W = work (J, N·m) \vec{p} = momentum (N·s) \vec{L} = angular momentum (N·m·s)
Angular/Rotational Energy, Work & Power	$K_r = \frac{1}{2}I\omega^2$ $K = K_t + K_r = \frac{1}{2}mv^2 + \frac{1}{2}I\omega^2$ $W_r = \tau\Delta\theta$ $P = \frac{W}{t} = \tau\omega$	









Table G. Moments of Inertia			
 Point Mass: $I = mr^2$	 Hollow Cylinder: $I = mr^2$	 Solid Cylinder: $I = \frac{1}{2}mr^2$	 Hoop About Diameter: $I = \frac{1}{2}mr^2$
 Hollow Sphere: $I = \frac{2}{3}mr^2$	 Solid Sphere: $I = \frac{2}{5}mr^2$	 Rod About the Middle: $I = \frac{1}{12}mL^2$	 Rod About the End: $I = \frac{1}{3}mL^2$

Table H. Heat and Thermal Physics Formulas and Equations		
Temperature	$T_F = 1.8(T_C) + 32$	<i>var. = name of quantity (unit)</i> Δ = change in something (E.g., Δx = change in x) T_K = Kelvin temperature (K) T_F = Fahrenheit temperature ($^{\circ}F$) T_C = Celsius temperature ($^{\circ}C$) T = temperature (K) Q = heat (J, kJ) m = mass (kg) C = specific heat capacity* ($\frac{kJ}{kg \cdot ^{\circ}C}, \frac{J}{g \cdot ^{\circ}C}$) t = time (s) L = length (m) k = coefficient of thermal conductivity* ($\frac{J}{m \cdot s \cdot ^{\circ}C}, \frac{W}{m \cdot ^{\circ}C}$) ϵ = emissivity* (dimensionless)
	$T_K = T_C + 273.15$	
Heat	$Q = mC \Delta T$ $Q_{melt} = m \Delta H_{fus}$ $Q_{boil} = m \Delta H_{vap}$ $C_p - C_v = R$ $\Delta L = \alpha L_i \Delta T \quad \Delta V = \beta V_i \Delta T$ $P = \frac{Q}{t} = kA \frac{\Delta T}{L} = \frac{1}{R_i} A \Delta T$ $P = \frac{Q}{t} = \epsilon \sigma A T^4$	H_{fus} = latent heat of fusion H_{vap} = heat of vaporization σ = Stefan-Boltzmann constant $= 5.67 \times 10^{-8} \frac{J}{m^2 \cdot s \cdot K^4}$ V = volume (m^3) α = linear coefficient of thermal expansion* ($^{\circ}C^{-1}$) β = volumetric coefficient of thermal expansion* ($^{\circ}C^{-1}$) P = pressure n = (number of) moles R = gas constant = $8.31 \frac{J}{mol \cdot K}$ k_B = Boltzmann constant = $1.38 \times 10^{-23} \frac{J}{K}$ U = internal energy (J) W = work (J)
	Thermodynamics	$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$ $PV = nRT \quad P \Delta V = nR \Delta T$ $PV = Nk_B T \quad P \Delta V = Nk_B \Delta T$ $\Delta U = Q + W$ $U = \frac{3}{2} nRT \quad \Delta U = \frac{3}{2} nR \Delta T$ $W = -\Delta(PV) = -P \Delta V$
		*characteristic property of a substance (to be looked up)

Table I. Thermal Properties of Selected Materials									
Substance	Melting Point ($^{\circ}C$)	Boiling Point ($^{\circ}C$)	Heat of Fusion ΔH_{fus} ($\frac{kJ}{kg}, \frac{J}{g}$)	Heat of Vaporization ΔH_{vap} ($\frac{kJ}{kg}, \frac{J}{g}$)	Specific Heat Capacity C_p ($\frac{kJ}{kg \cdot ^{\circ}C}, \frac{J}{g \cdot ^{\circ}C}$) at 25 $^{\circ}C$	Thermal Conductivity k ($\frac{J}{ms \cdot ^{\circ}C}$) at 25 $^{\circ}C$	Emissivity ϵ black body = 1	Coefficients of Expansion at 20 $^{\circ}C$	
								Linear α ($^{\circ}C^{-1}$)	Volumetric β ($^{\circ}C^{-1}$)
air (gas)	—	—	—	—	1.012	0.024	—	—	—
aluminum (solid)	659	2467	395	10460	0.897	250	0.09*	2.3×10^{-5}	6.9×10^{-5}
ammonia (gas)	-75	-33.3	339	1369	4.7	0.024	—	—	—
argon (gas)	-189	-186	29.5	161	0.520	0.016	—	—	—
carbon dioxide (gas)	—	-78	—	574	0.839	0.0146	—	—	—
copper (solid)	1086	1187	134	5063	0.385	401	0.03*	1.7×10^{-5}	5.1×10^{-5}
brass (solid)	—	—	—	—	0.380	120	0.03*	1.9×10^{-5}	5.6×10^{-5}
diamond (solid)	3550	4827	10 000	30 000	0.509	2200	—	1×10^{-6}	3×10^{-6}
ethanol (liquid)	-117	78	104	858	2.44	0.171	—	2.5×10^{-4}	7.5×10^{-4}
glass (solid)	—	—	—	—	0.84	0.96–1.05	0.92	8.5×10^{-6}	2.55×10^{-5}
gold (solid)	1063	2660	64.4	1577	0.129	310	0.025*	1.4×10^{-5}	4.2×10^{-5}
granite (solid)	1240	—	—	—	0.790	1.7–4.0	0.96	—	—
helium (gas)	—	-269	—	21	5.193	0.142	—	—	—
hydrogen (gas)	-259	-253	58.6	452	14.30	0.168	—	—	—
iron (solid)	1535	2750	289	6360	0.450	80	0.31	1.18×10^{-5}	3.33×10^{-5}
lead (solid)	327	1750	24.7	870	0.160	35	0.06	2.9×10^{-5}	8.7×10^{-5}
mercury (liquid)	-39	357	11.3	293	0.140	8	—	6.1×10^{-5}	1.82×10^{-4}
paraffin wax (solid)	46–68	~300	~210	—	2.5	0.25	—	—	—
silver (solid)	962	2212	111	2360	0.233	429	0.025*	1.8×10^{-5}	5.4×10^{-5}
steam (gas) @ 100 $^{\circ}C$	—	—	—	—	2.080	0.016	—	—	—
water (liq.) @ 25 $^{\circ}C$	0	100	—	2260	4.181	0.58	0.95	6.9×10^{-5}	2.07×10^{-4}
ice (solid) @ -10 $^{\circ}C$	—	—	334	—	2.11	2.18	0.97	—	—

*polished surface

Table J. Electricity Formulas & Equations		var. = name of quantity (unit)
Electrostatic Charges & Electric Fields	$\vec{F}_e = \frac{kq_1q_2}{r^2} = \frac{1}{4\pi\epsilon_0} \frac{q_1q_2}{r^2}$ $\vec{E} = \frac{\vec{F}_e}{q} = \frac{Q}{\epsilon_0 A} \quad \vec{E} = \frac{kq}{r^2} = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} = \frac{\Delta V}{\Delta r}$ $W = q\vec{E} \cdot \vec{d} = qEd_{\parallel} = qEd \cos\theta$ $\Delta V = \frac{W}{q} = \vec{E} \cdot \vec{d} = Ed_{\parallel} = \frac{1}{4\pi\epsilon_0} \frac{q}{r}$ $\Delta U_e = q\Delta V \quad U_e = \frac{kq_1q_2}{r}$	<p>Δ = change in something. (E.g., Δx = change in x)</p> <p>\vec{F}_e = force due to electric field (N)</p> <p>ϵ_0 = electric permittivity of a vacuum = $8.85 \times 10^{-12} \frac{A^2 \cdot s^4}{kg \cdot m^3}$</p> <p>$k$ = electrostatic constant = $\frac{1}{4\pi\epsilon_0} = 9.0 \times 10^9 \frac{N \cdot m^2}{C^2}$</p> <p>$q$ = point charge (C)</p> <p>Q = charge (C)</p> <p>\vec{E} = electric field ($\frac{N}{C}, \frac{V}{m}$)</p> <p>V = electric potential (V)</p> <p>ΔV = voltage = electric potential difference (V)</p> <p>\mathcal{E} = emf = electromotive force (V)</p> <p>W = work (J, N·m)</p> <p>$\kappa = \epsilon_r$ = relative permittivity* (<i>dimensionless</i>)</p> <p>d = distance (m)</p> <p>r = radius (m)</p> <p>\vec{I} = current (A)</p> <p>t = time (s)</p> <p>R = resistance (Ω)</p> <p>P = power (W)</p> <p>ρ = resistivity ($\Omega \cdot m$)</p> <p>L = length (m)</p> <p>A = cross-sectional area (m^2)</p> <p>C = capacitance (F)</p> <p>U = potential energy (J)</p> <p>π = pi (mathematical constant) = 3.14159 26535 89793...</p> <p>e = Euler's number (mathematical constant) = 2.78182 81812 84590...</p>
Circuits and Electrical Components	$\Delta V = IR \quad I = \frac{\Delta Q}{\Delta t} = \frac{\Delta V}{R}$ $\mathcal{E} = IR$ $P = I\Delta V = I^2R = \frac{(\Delta V)^2}{R}$ $W = Pt = I\Delta Vt$ $R = \frac{\rho L}{A}$ $C = \kappa\epsilon_0 \frac{A}{d}$ $Q = C\Delta V$ $U_{capacitor} = \frac{1}{2}Q\Delta V = \frac{1}{2}C(\Delta V)^2$ $P_{total} = P_1 + P_2 + P_3 + \dots = \sum P_i$ $U_{total} = U_1 + U_2 + U_3 + \dots = \sum U_i$	
Series Circuits (or Series Sections of Circuits)	$I_{total} = I_1 = I_2 = I_3 = \dots$ $\Delta V_{total} = \Delta V_1 + \Delta V_2 + \Delta V_3 + \dots = \sum \Delta V_i$ $R_{equiv.} = R_1 + R_2 + R_3 + \dots = \sum R_i$ $Q_{total} = Q_1 = Q_2 = Q_3 = \dots$ $\frac{1}{C_{total}} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3} + \dots = \sum \frac{1}{C_i}$	
Parallel Circuits (or Parallel Sections of Circuits)	$I_{total} = I_1 + I_2 + I_3 + \dots = \sum I_i$ $\Delta V_{total} = \Delta V_1 = \Delta V_2 = \Delta V_3 = \dots$ $\frac{1}{R_{equiv.}} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots = \sum \frac{1}{R_i}$ $Q_{total} = Q_1 + Q_2 + Q_3 + \dots = \sum Q_i$ $C_{total} = C_1 + C_2 + C_3 + \dots = \sum C_i$	
Resistor-Capacitor (RC) Circuits	<p>charging: $\frac{I}{I_o} = e^{-t/RC}$</p> <p>charging: $\frac{Q}{Q_{max}} = 1 - e^{-t/RC}$</p> <p>discharging: $\frac{I}{I_o} = \frac{V}{V_o} = \frac{Q}{Q_{max}} = e^{-t/RC}$</p>	

*characteristic property of a substance (to be looked up)

Table K. Electricity & Magnetism Formulas & Equations	
<p>Magnetism and Electromagnetism</p> $\vec{F}_M = q(\vec{v} \times \vec{B}) \quad F_M = qvB \sin \theta$ $\vec{F}_M = \ell(\vec{I} \times \vec{B}) \quad F_M = \ell IB \sin \theta$ $\Delta V = \ell(\vec{v} \times \vec{B}) \quad \Delta V = \ell vB \sin \theta$ $B = \frac{\mu_0 I}{2\pi r}$ $\Phi_B = \vec{B} \cdot \vec{A} = BA \cos \theta$ $\mathcal{E} = \frac{\Delta \Phi_B}{\Delta t} = BLv$	<p><i>var.</i> = name of quantity (unit)</p> <p>Δ = change in something. (E.g., Δx = change in x)</p> <p>\vec{F}_e = force due to electric field (N)</p> <p>\vec{v} = velocity (of moving charge or wire) ($\frac{m}{s}$)</p> <p>q = point charge (C)</p> <p>ΔV = voltage = electric potential difference (V)</p> <p>\mathcal{E} = emf = electromotive force (V)</p> <p>r = radius (m) = distance from wire</p> <p>\vec{I} = current (A)</p> <p>L = length (m)</p> <p>t = time (s)</p> <p>A = cross-sectional area (m^2)</p> <p>\vec{B} = magnetic field (T)</p> <p>μ_0 = magnetic permeability of a vacuum = $4\pi \times 10^{-7} \frac{T \cdot m}{A}$</p> <p>$\Phi_B$ = magnetic flux ($T \cdot m^2$)</p>
<p>Electromagnetic Induction</p> $\frac{\#turns_{in}}{\#turns_{out}} = \frac{V_{in}}{V_{out}} = \frac{I_{out}}{I_{in}}$ $P_{in} = P_{out}$	

Table L. Resistor Color Code		
Color	Digit	Multiplier
black	0	$\times 10^0$
brown	1	$\times 10^1$
red	2	$\times 10^2$
orange	3	$\times 10^3$
yellow	4	$\times 10^4$
green	5	$\times 10^5$
blue	6	$\times 10^6$
violet	7	$\times 10^7$
gray	8	$\times 10^8$
white	9	$\times 10^9$
gold		$\pm 5\%$
silver		$\pm 10\%$

Table M. Symbols Used in Electrical Circuit Diagrams			
Component	Symbol	Component	Symbol
wire	—	battery	
switch		ground	
fuse		resistor	
voltmeter		variable resistor (rheostat, potentiometer, dimmer)	
ammeter		lamp (light bulb)	
ohmmeter		capacitor	
		diode	

Table N. Resistivities at 20°C					
Conductors		Semiconductors		Insulators	
Substance	Resistivity ($\Omega \cdot m$)	Substance	Resistivity ($\Omega \cdot m$)	Substance	Resistivity ($\Omega \cdot m$)
silver	1.59×10^{-8}	germanium	0.001 to 0.5	deionized water	1.8×10^5
copper	1.72×10^{-8}	silicon	0.1 to 60	glass	1×10^9 to 1×10^{13}
gold	2.44×10^{-8}	sea water	0.2	rubber, hard	1×10^{13} to 1×10^{13}
aluminum	2.82×10^{-8}	drinking water	20 to 2000	paraffin (wax)	1×10^{13} to 1×10^{17}
tungsten	5.60×10^{-8}			air	1.3×10^{16} to 3.3×10^{16}
iron	9.71×10^{-8}			quartz, fused	7.5×10^{17}
nichrome	1.50×10^{-6}				
graphite	3×10^{-5} to 6×10^{-4}				

Table O. Waves & Optics Formulas & Equations		
<p>Waves</p>	$v = \lambda f$ $f = \frac{1}{T}$ $v_{\text{wave on a string}} = \sqrt{\frac{F_T}{\mu}}$ $f_{\text{doppler shifted}} = f \left(\frac{\vec{v}_{\text{wave}} + \vec{v}_{\text{detector}}}{\vec{v}_{\text{wave}} + \vec{v}_{\text{source}}} \right)$ $x = A \cos(2\pi ft + \phi)$	<p><i>var. = name of quantity (unit)</i></p> <p>Δ = change in something (E.g., Δx = change in x)</p> <p>v = velocity of wave ($\frac{m}{s}$)</p> <p>\vec{v} = velocity of source or detector ($\frac{m}{s}$)</p> <p>f = frequency (Hz)</p> <p>λ = wavelength (m)</p> <p>A = amplitude (m)</p> <p>x = position (m)</p> <p>T = period (of time) (s)</p> <p>F_T = tension (force) on string (N)</p> <p>μ = elastic modulus of string ($\frac{kg}{m}$)</p> <p>θ = angle ($^\circ$, rad)</p> <p>ϕ = phase offset ($^\circ$, rad)</p> <p>θ_i = angle of incidence ($^\circ$, rad)</p> <p>θ_r = angle of reflection ($^\circ$, rad)</p> <p>θ_c = critical angle ($^\circ$, rad)</p> <p>n = index of refraction* (<i>dimensionless</i>)</p> <p>c = speed of light in a vacuum = $3.00 \times 10^8 \frac{m}{s}$</p> <p>$f = s_f = d_f$ = distance to focus of mirror/lens (m)</p> <p>r_c = radius of curvature of spherical mirror (m)</p> <p>$s_i = d_i$ = distance from mirror/lens to image (m)</p> <p>$s_o = d_o$ = distance from mirror/lens to object (m)</p> <p>h_i = height of image (m)</p> <p>h_o = height of object (m)</p> <p>M = magnification (<i>dimensionless</i>)</p> <p>d = separation (m)</p> <p>L = distance from the opening (m)</p> <p>m = an integer</p>
<p>Reflection, Refraction & Diffraction</p>	$\theta_i = \theta_r$ $n = \frac{c}{v}$ $n_1 \sin \theta_1 = n_2 \sin \theta_2$ $\theta_c = \sin^{-1} \left(\frac{n_2}{n_1} \right)$ $\frac{n_2}{n_1} = \frac{v_1}{v_2} = \frac{\lambda_1}{\lambda_2}$ $\Delta L = m \lambda = d \sin \theta$	<p>*characteristic property of a substance (to be looked up)</p>
<p>Mirrors & Lenses</p>	$f = \frac{r_c}{2}$ $\frac{1}{s_i} + \frac{1}{s_o} = \frac{1}{f}$ $M = \frac{h_i}{h_o} = -\frac{s_i}{s_o}$	




Table Q. Properties of Water and Air					
Temp. (°C)	Water			Air	
	Density $\left(\frac{\text{kg}}{\text{m}^3}\right)$	Speed of Sound $\left(\frac{\text{m}}{\text{s}}\right)$	Vapor Pressure (Pa)	Density $\left(\frac{\text{kg}}{\text{m}^3}\right)$	Speed of Sound $\left(\frac{\text{m}}{\text{s}}\right)$
0	999.78	1 403	611.73	1.288	331.30
5	999.94	1 427	872.60	1.265	334.32
10	999.69	1 447	1 228.1	1.243	337.31
20	998.19	1 481	2 338.8	1.200	343.22
25	997.02	1 496	3 169.1	1.180	346.13
30	995.61	1 507	4 245.5	1.161	349.02
40	992.17	1 526	7 381.4	1.124	354.73
50	990.17	1 541	9 589.8	1.089	360.35
60	983.16	1 552	19 932	1.056	365.88
70	980.53	1 555	25 022	1.025	371.33
80	971.79	1 555	47 373	0.996	376.71
90	965.33	1 550	70 117	0.969	382.00
100	954.75	1 543	101 325	0.943	387.23

Table R. Absolute Indices of Refraction			
Measured using $f = 5.89 \times 10^{14}$ Hz (yellow light) at 20 °C unless otherwise specified			
Substance	Index of Refraction	Substance	Index of Refraction
air (0 °C and 1 atm)	1.000293	silica (quartz), fused	1.459
ice (0 °C)	1.309	Plexiglas	1.488
water	1.3330	Lucite	1.495
ethyl alcohol	1.36	glass, borosilicate (Pyrex)	1.474
human eye, cornea	1.38	glass, crown	1.50–1.54
human eye, lens	1.41	glass, flint	1.569–1.805
safflower oil	1.466	sodium chloride, solid	1.516
corn oil	1.47	PET (#1 plastic)	1.575
glycerol	1.473	zircon	1.777–1.987
honey	1.484–1.504	cubic zirconia	2.173–2.21
silicone oil	1.52	diamond	2.417
carbon disulfide	1.628	silicon	3.96

Table S. Fluid Mechanics Formulas and Equations	
<p>Density & Pressure</p>	$\rho = \frac{m}{V}$ $p = \frac{F}{A}$ $\frac{F_1}{A_1} = \frac{F_2}{A_2}$ $P_{\text{hydrostatic}} = P_H = \rho gh$ $P_{\text{dynamic}} = P_D = \frac{1}{2} \rho v^2$ $A_1 v_1 = A_2 v_2$ $P_{\text{total}} = P_{\text{ext.}} + P_H + P_D$ $P_1 + P_{H,1} + P_{D,1} = P_2 + P_{H,2} + P_{D,2}$ $P_1 + \rho gh_1 + \frac{1}{2} \rho v_1^2 = P_2 + \rho gh_2 + \frac{1}{2} \rho v_2^2$
<p>Forces, Work & Energy</p>	$F_B = \rho V_d g$ $PV = nRT = Nk_B T \quad P\Delta V = nR\Delta T$ $\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$ $K_{(\text{molecular})} = \frac{3}{2} RT$ $U = \frac{3}{2} nRT = \frac{3}{2} Nk_B T \quad \Delta U = \frac{3}{2} nR\Delta T$ $v_{\text{rms}} = \sqrt{\frac{3RT}{M}} = \sqrt{\frac{3k_B T}{\mu}}$ $W = -P\Delta V$
<p><i>var.</i> = name of quantity (unit)</p> <p>Δ = change in something. (E.g., Δx = change in x)</p> <p>ρ = density ($\frac{\text{kg}}{\text{m}^3}$)</p> <p>$m$ = mass (kg)</p> <p>V = volume (m^3)</p> <p>P = pressure (Pa)</p> <p>g = gravitational field = $9.8 \frac{\text{N}}{\text{kg}} \approx 10 \frac{\text{N}}{\text{kg}}$</p> <p>$h$ = height or depth (m)</p> <p>A = area (m^2)</p> <p>v = velocity (of fluid) ($\frac{\text{m}}{\text{s}}$)</p> <p>$F$ = force (N)</p> <p>n = number of moles (mol)</p> <p>R = gas constant = $8.31 \frac{\text{J}}{\text{mol}\cdot\text{K}}$</p> <p>$N$ = number of molecules</p> <p>k_B = Boltzmann's constant = $1.38 \times 10^{-23} \frac{\text{J}}{\text{K}}$</p> <p>$T$ = temperature (K)</p> <p>v_{rms} = root mean square speed ($\frac{\text{m}}{\text{s}}$)</p> <p>$M$ = molar mass* ($\frac{\text{kg}}{\text{mol}}$)</p> <p>$\mu$ = molecular mass* (kg)</p> <p>K = kinetic energy (J)</p> <p>W = work (J, N·m)</p> <p>*characteristic property of a substance (to be looked up)</p>	

Table T. Planetary Data									
	Mercury	Venus	Earth	Mars	Jupiter	Saturn	Uranus	Neptune	Pluto
Distance from Sun (m)	5.79×10^{10}	1.08×10^{11}	1.50×10^{11}	2.28×10^{11}	7.79×10^{11}	1.43×10^{12}	2.87×10^{12}	4.50×10^{12}	5.91×10^{12}
Radius (m)	2.44×10^6	6.05×10^6	6.38×10^6	3.40×10^6	7.15×10^7	6.03×10^7	2.56×10^7	2.48×10^7	1.19×10^6
Mass (kg)	3.30×10^{23}	4.87×10^{24}	5.97×10^{24}	6.42×10^{23}	1.90×10^{27}	5.68×10^{26}	8.68×10^{25}	1.02×10^{26}	1.46×10^{22}
Density ($\frac{\text{kg}}{\text{m}^3}$)	5427	5243	5514	3340	3933	687	1271	1638	2095
Orbit (years)	0.24	0.61	1.00	1.88	11.8	29	84	164	248
Rotation Period (hours)	1408	-5833	23.9	24.6	9.9	10.7	-17.2	16.1	-153.3
Tilt of axis	0.034°	177.4°	23.4°	25.2°	3.1°	26.7°	97.8°	28.3°	122.5°
# of observed satellites	0	0	1	2	79	82	27	14	14
Mean temperature ($^\circ\text{C}$)	167	464	15	-20	-65	-110	-195	-200	-225
Global magnetic field?	Yes	No	Yes	No	Yes	Yes	Yes	Yes	Yes

Data from NASA Planetary Fact Sheet, <https://nssdc.gsfc.nasa.gov/planetary/factsheet/> accessed 03 February 2020.

Table U. Sun & Moon Data	
Radius of the sun (m)	6.96×10^8
Mass of the sun (kg)	1.99×10^{30}
Radius of the moon (m)	1.74×10^6
Mass of the moon (kg)	7.35×10^{22}
Distance of moon from Earth (m)	3.84×10^8

Table V. Atomic & Particle Physics (Modern Physics)		
Energy	$E_{\text{photon}} = hf = \frac{hc}{\lambda} = pc = \hbar\omega$ $E_{k,\text{max}} = hf - \phi$ $\lambda = \frac{h}{p}$ $E_{\text{photon}} = E_i - E_f$ $E^2 = (pc)^2 + (mc^2)^2$ $\frac{1}{\lambda} = R_H \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$	<p><i>var.</i> = name of quantity (unit)</p> <p>Δ = change in something. (E.g., Δx = change in x)</p> <p>E = energy (J)</p> <p>h = Planck's constant = 6.63×10^{-34} J·s</p> <p>\hbar = reduced Planck's constant = $\frac{h}{2\pi} = 1.05 \times 10^{-34}$ J·s</p> <p>f = frequency (Hz)</p> <p>v = velocity ($\frac{m}{s}$)</p> <p>c = speed of light = 3.00×10^8 $\frac{m}{s}$</p> <p>λ = wavelength (m)</p> <p>p = momentum (N·s)</p> <p>m = mass (kg)</p> <p>K = kinetic energy (J)</p> <p>ϕ = work function* (J)</p> <p>R_H = Rydberg constant = 1.10×10^7 m⁻¹</p>
	Special Relativity	$\gamma = \frac{1}{\sqrt{1 - v^2/c^2}}$ $\gamma = \frac{L_o}{L} = \frac{\Delta t'}{\Delta t} = \frac{m_{rel}}{m_o}$




Figure X. Particle Sizes




Figure Y. Classification of Matter




Table Z. The Standard Model of Elementary Particles

Standard Model of Elementary Particles


	three generations of matter (fermions)			interactions / force carriers (bosons)	
	I	II	III		
mass	$\approx 2.2 \text{ MeV}/c^2$	$\approx 1.28 \text{ GeV}/c^2$	$\approx 173.1 \text{ GeV}/c^2$	0	$\approx 124.97 \text{ GeV}/c^2$
charge	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	0	0
spin	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	0
QUARKS	u up	c charm	t top	g gluon	H higgs
	$\approx 4.7 \text{ MeV}/c^2$	$\approx 96 \text{ MeV}/c^2$	$\approx 4.18 \text{ GeV}/c^2$	0	
	$-\frac{1}{3}$	$-\frac{1}{3}$	$-\frac{1}{3}$	0	
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	
	d down	s strange	b bottom	γ photon	
LEPTONS	$\approx 0.511 \text{ MeV}/c^2$	$\approx 105.66 \text{ MeV}/c^2$	$\approx 1.7768 \text{ GeV}/c^2$	$\approx 91.19 \text{ GeV}/c^2$	
	-1	-1	-1	0	
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	
	e electron	μ muon	τ tau	Z Z boson	
	$<1.0 \text{ eV}/c^2$	$<0.17 \text{ MeV}/c^2$	$<18.2 \text{ MeV}/c^2$	$\approx 80.39 \text{ GeV}/c^2$	
	0	0	0	± 1	
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	W W boson	
					GAUGE BOSONS VECTOR BOSONS
					SCALAR BOSONS

Figure AA. Periodic Table of the Elements

Period	1 IA	18 VIII A																													
		1 H hydrogen 1.008																	2 He helium 4.003												
	2	2 II A																10													
		Li lithium 6.968	Be beryllium 9.012															Ne neon 20.18													
	3	3 III B																18													
		Na sodium 22.99	Mg magnesium 24.31															Ar argon 39.95													
	4	4 IV B		5 VB		6 VIB		7 VII B		8 VIII B		9 VIII B		10 VIII B		11 IB		12 IIB													
		K potassium 39.10	Ca calcium 40.08	Ti titanium 47.87	V vanadium 50.94	Cr chromium 52.00	Mn manganese 54.94	Fe iron 55.85	Co cobalt 58.93	Ni nickel 58.69	Cu copper 63.55	Zn zinc 65.38	Ga gallium 69.72	Ge germanium 72.63	As arsenic 74.92	Se selenium 78.97	Br bromine 79.90	Kr krypton 83.80													
	5	3 III B		4 IV B		5 VB		6 VIB		7 VII B		8 VIII B		9 VIII B		10 VIII B		11 IB		12 IIB											
		Rb rubidium 85.47	Sr strontium 87.62	Y yttrium 88.91	Zr zirconium 91.22	Nb niobium 92.91	Mo molybdenum 95.95	Tc technetium 98	Ru ruthenium 101.1	Rh rhodium 102.9	Pd palladium 106.4	Ag silver 107.9	Cd cadmium 112.4	In indium 114.8	Sn tin 118.7	Sb antimony 121.8	Te tellurium 127.6	I iodine 126.9	Xe xenon 131.3												
	6	4 IV B		5 VB		6 VIB		7 VII B		8 VIII B		9 VIII B		10 VIII B		11 IB		12 IIB		13 IIIA		14 IVA		15 VA		16 VIA		17 VIIA		18 VIIIA	
		Cs cesium 132.9	Ba barium 137.3	Lu lutetium 175.0	Hf hafnium 178.5	Ta tantalum 180.9	W tungsten 183.8	Re rhenium 186.2	Os osmium 190.2	Ir iridium 192.2	Pt platinum 195.1	Au gold 197.0	Hg mercury 200.6	Tl thallium 204.4	Pb lead 207.2	Bi bismuth 209.0	Po polonium 209	At astatine 210	Rn radon 222												
	7	5 VB		6 VIB		7 VII B		8 VIII B		9 VIII B		10 VIII B		11 IB		12 IIB		13 IIIA		14 IVA		15 VA		16 VIA		17 VIIA		18 VIIIA			
		Fr francium 223	Ra radium 226	Lr lawrencium 262	Rf rutherfordium 267	Db dubnium 268	Sg seaborgium 271	Bh bohrium 272	Hs hassium 270	Mt meitnerium 276	Ds darmstadtium 281	Rg roentgenium 280	Cn copernicium 285	Nh nihonium 284	Fl flerovium 289	Mc moscovium 288	Lv livermorium 293	Ts tennessine 292	Og oganeson 294												
	lanthanides (rare earth metals)		57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.2	61 Pm promethium 145	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1															
	actinides		89 Ac actinium 227	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium 237	94 Pu plutonium 244	95 Am americium 243	96 Cm curium 247	97 Bk berkelium 247	98 Cf californium 251	99 Es einsteinium 252	100 Fm fermium 257	101 Md mendelevium 258	102 No nobelium 259															

Name	Notation	Symbol
alpha particle	${}^4_2\text{He}$ or ${}^4_2\alpha$	α
beta particle (electron)	${}^0_{-1}e$ or ${}^0_{-1}\beta$	β^-
gamma radiation	${}^0_0\gamma$	γ
neutron	1_0n	n
proton	${}^1_1\text{H}$ or 1_1p	p
positron	${}^0_{+1}e$ or ${}^0_{+1}\beta$	β^+

Constant	Value
mass of an electron (m_e)	0.00055 amu
mass of a proton (m_p)	1.00728 amu
mass of a neutron (m_n)	1.00867 amu
Bequerel (Bq)	1 disintegration/second
Curie (Ci)	3.7×10^{10} Bq



Nuclide	Half-Life	Decay Mode
${}^3\text{H}$	12.26 y	β^-
${}^{14}\text{C}$	5730 y	β^-
${}^{16}\text{N}$	7.2 s	β^-
${}^{19}\text{Ne}$	17.2 s	β^+
${}^{24}\text{Na}$	15 h	β^-
${}^{27}\text{Mg}$	9.5 min	β^-
${}^{32}\text{P}$	14.3 d	β^-
${}^{36}\text{Cl}$	3.01×10^5 y	β^-
${}^{37}\text{K}$	1.23 s	β^+
${}^{40}\text{K}$	1.26×10^9 y	β^+
${}^{42}\text{K}$	12.4 h	β^-
${}^{37}\text{Ca}$	0.175 s	β^-
${}^{51}\text{Cr}$	27.7 d	α
${}^{53}\text{Fe}$	8.51 min	β^-
${}^{59}\text{Fe}$	46.3 d	β^-
${}^{60}\text{Co}$	5.26 y	β^-
${}^{85}\text{Kr}$	10.76 y	β^-
${}^{87}\text{Rb}$	4.8×10^{10} y	β^-
${}^{90}\text{Sr}$	28.1 y	β^-
${}^{99}\text{Tc}$	2.13×10^5 y	β^-
${}^{131}\text{I}$	8.07 d	β^-
${}^{137}\text{Cs}$	30.23 y	β^-
${}^{153}\text{Sm}$	1.93 d	β^-
${}^{198}\text{Au}$	2.69 d	β^-
${}^{222}\text{Rn}$	3.82 d	α
${}^{220}\text{Fr}$	27.5 s	α
${}^{226}\text{Ra}$	1600 y	α
${}^{232}\text{Th}$	1.4×10^{10} y	α
${}^{233}\text{U}$	1.62×10^5 y	α
${}^{235}\text{U}$	7.1×10^8 y	α
${}^{238}\text{U}$	4.51×10^9 y	α
${}^{239}\text{Pu}$	2.44×10^4 y	α
${}^{241}\text{Am}$	432 y	α



Table FF. Geometry & Trigonometry Formulas		
Triangles	$A = \frac{1}{2}bh$ $c^2 = a^2 + b^2 - 2ab\cos C$ $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$	
Right Triangles	$c^2 = a^2 + b^2$ $\sin \theta = \frac{a}{c} = \frac{\text{opposite}}{\text{hypotenuse}}$ $\cos \theta = \frac{b}{c} = \frac{\text{adjacent}}{\text{hypotenuse}}$ $\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{a}{b} = \frac{\text{opposite}}{\text{adjacent}}$ $b = c \cos \theta$ $a = c \sin \theta$	
Rectangles, Parallelograms and Trapezoids	$A = \bar{b}h$	<p>a, b, c = length of a side of a triangle θ = angle A = area C = circumference S = surface area V = volume b = base h = height L = length w = width r = radius</p>
Rectangular Solids	$V = Lwh$	
Circles	$C = 2\pi r$ $A = \pi r^2$	
Cylinders	$S = 2\pi rL + 2\pi r^2 = 2\pi r(L + r)$ $V = \pi r^2L$	
Spheres	$S = 4\pi r^2$ $V = \frac{4}{3}\pi r^3$	

Table GG. Values of Trigonometric Functions									
degree	radian	sine	cosine	tangent	degree	radian	sine	cosine	tangent
0°	0.000	0.000	1.000	0.000					
1°	0.017	0.017	1.000	0.017	46°	0.803	0.719	0.695	1.036
2°	0.035	0.035	0.999	0.035	47°	0.820	0.731	0.682	1.072
3°	0.052	0.052	0.999	0.052	48°	0.838	0.743	0.669	1.111
4°	0.070	0.070	0.998	0.070	49°	0.855	0.755	0.656	1.150
5°	0.087	0.087	0.996	0.087	50°	0.873	0.766	0.643	1.192
6°	0.105	0.105	0.995	0.105	51°	0.890	0.777	0.629	1.235
7°	0.122	0.122	0.993	0.123	52°	0.908	0.788	0.616	1.280
8°	0.140	0.139	0.990	0.141	53°	0.925	0.799	0.602	1.327
9°	0.157	0.156	0.988	0.158	54°	0.942	0.809	0.588	1.376
10°	0.175	0.174	0.985	0.176	55°	0.960	0.819	0.574	1.428
11°	0.192	0.191	0.982	0.194	56°	0.977	0.829	0.559	1.483
12°	0.209	0.208	0.978	0.213	57°	0.995	0.839	0.545	1.540
13°	0.227	0.225	0.974	0.231	58°	1.012	0.848	0.530	1.600
14°	0.244	0.242	0.970	0.249	59°	1.030	0.857	0.515	1.664
15°	0.262	0.259	0.966	0.268	60°	1.047	0.866	0.500	1.732
16°	0.279	0.276	0.961	0.287	61°	1.065	0.875	0.485	1.804
17°	0.297	0.292	0.956	0.306	62°	1.082	0.883	0.469	1.881
18°	0.314	0.309	0.951	0.325	63°	1.100	0.891	0.454	1.963
19°	0.332	0.326	0.946	0.344	64°	1.117	0.899	0.438	2.050
20°	0.349	0.342	0.940	0.364	65°	1.134	0.906	0.423	2.145
21°	0.367	0.358	0.934	0.384	66°	1.152	0.914	0.407	2.246
22°	0.384	0.375	0.927	0.404	67°	1.169	0.921	0.391	2.356
23°	0.401	0.391	0.921	0.424	68°	1.187	0.927	0.375	2.475
24°	0.419	0.407	0.914	0.445	69°	1.204	0.934	0.358	2.605
25°	0.436	0.423	0.906	0.466	70°	1.222	0.940	0.342	2.747
26°	0.454	0.438	0.899	0.488	71°	1.239	0.946	0.326	2.904
27°	0.471	0.454	0.891	0.510	72°	1.257	0.951	0.309	3.078
28°	0.489	0.469	0.883	0.532	73°	1.274	0.956	0.292	3.271
29°	0.506	0.485	0.875	0.554	74°	1.292	0.961	0.276	3.487
30°	0.524	0.500	0.866	0.577	75°	1.309	0.966	0.259	3.732
31°	0.541	0.515	0.857	0.601	76°	1.326	0.970	0.242	4.011
32°	0.559	0.530	0.848	0.625	77°	1.344	0.974	0.225	4.331
33°	0.576	0.545	0.839	0.649	78°	1.361	0.978	0.208	4.705
34°	0.593	0.559	0.829	0.675	79°	1.379	0.982	0.191	5.145
35°	0.611	0.574	0.819	0.700	80°	1.396	0.985	0.174	5.671
36°	0.628	0.588	0.809	0.727	81°	1.414	0.988	0.156	6.314
37°	0.646	0.602	0.799	0.754	82°	1.431	0.990	0.139	7.115
38°	0.663	0.616	0.788	0.781	83°	1.449	0.993	0.122	8.144
39°	0.681	0.629	0.777	0.810	84°	1.466	0.995	0.105	9.514
40°	0.698	0.643	0.766	0.839	85°	1.484	0.996	0.087	11.430
41°	0.716	0.656	0.755	0.869	86°	1.501	0.998	0.070	14.301
42°	0.733	0.669	0.743	0.900	87°	1.518	0.999	0.052	19.081
43°	0.750	0.682	0.731	0.933	88°	1.536	0.999	0.035	28.636
44°	0.768	0.695	0.719	0.966	89°	1.553	1.000	0.017	57.290
45°	0.785	0.707	0.707	1.000	90°	1.571	1.000	0.000	∞

Table HH. Some Exact and Approximate Conversions			
Length	1 cm	≈	width of a small paper clip
	1 inch (in.)	≡	2.54 cm
	length of a US dollar bill	=	6.14 in. = 15.6 cm
	12 in.	≡	1 foot (ft.) ≈ 30 cm
	3 ft.	≡	1 yard (yd.) ≈ 1 m
	1 m	≡	0.3048 ft. = 39.37 in.
	1 km	≈	0.6 mi.
	5,280 ft.	≡	1 mile (mi.) ≈ 1.6 km
Mass / Weight	1 small paper clip	≈	0.5 g
	US 1¢ coin (1983–present)	=	2.5 g
	US 5¢ coin	=	5 g
	1 oz.	≈	30 g
	one medium-sized apple	≈	1 N ≈ 3.6 oz.
	1 pound (lb.)	≡	16 oz. ≈ 454 g
	1 pound (lb.)	≈	4.45 N
	1 ton	≡	2000 lb. ≈ 0.9 tonne
1 tonne	≡	1000 kg ≈ 1.1 ton	
Volume	1 pinch	≈	$\frac{1}{16}$ teaspoon (tsp.)
	1 dash	≈	$\frac{1}{8}$ teaspoon (tsp.)
	1 mL	≈	10 drops
	1 tsp.	≈	5 mL ≈ 60 drops
	3 tsp.	≡	1 tablespoon (Tbsp.) ≈ 15 mL
	2 Tbsp.	≡	1 fluid ounce (fl. oz.) ≈ 30 mL
	8 fl. oz.	≡	1 cup (C) ≈ 250 mL
	16 fl. oz.	≡	1 U.S. pint (pt.) ≈ 500 mL
	20 fl. oz.	≡	1 Imperial pint (UK) ≈ 600 mL
	2 pt. (U.S.)	≡	1 U.S. quart (qt.) ≈ 1 L
4 qt. (U.S.)	≡	1 U.S. gallon (gal.) ≈ 3.8 L	
4 qt. (UK) ≡ 5 qt. (U.S.)	≡	1 Imperial gal. (UK) ≈ 4.7 L	
Speed / Velocity	1 m/s	=	3.6 km/h ≈ 2.24 mi./h
	60 mi./h	≈	100 km/h ≈ 27 m/s
Energy	1 cal	≈	4.18 J
	1 Calorie (food)	≡	1 kcal ≈ 4.18 kJ
	1 BTU	≈	1.06 kJ
Power	1 hp	≈	746 W
	1 kW	≈	1.34 hp
Temperature	0 K	≡	-273.15°C = absolute zero
	0°R	≡	-459.67°F = absolute zero
	0°F	≈	-18°C ≡ 459.67°R
	32°F	=	0°C ≡ 273.15 K = water freezes
	70°F	≈	21°C ≈ room temperature
	212°F	=	100°C = water boils
Speed of light	300 000 000 m/s	≈	186 000 mi./s ≈ 1 ft./ns

Table II. Greek Alphabet		
A	α	alpha
B	β	beta
Γ	γ	gamma
Δ	δ	delta
E	ε	epsilon
Z	ζ	zeta
H	η	eta
Θ	θ	theta
I	ι	iota
K	κ	kappa
Λ	λ	lambda
M	μ	mu
N	ν	nu
Ξ	ξ	xi
O	ο	omicron
Π	π	pi
P	ρ	rho
Σ	σ	sigma
T	τ	tau
Υ	υ	upsilon
Φ	φ	phi
X	χ	chi
Ψ	ψ	psi
Ω	ω	omega