RD Sharma Solutions For Class 8 Chapter 7 Factorization

Find the greatest common factor (GCF/HCF) of the following polynomials: (1-14)

1. 2x² and 12x²

Solution:

We know that the numerical coefficients of given numerical are 2 and 12

The greatest common factor of 2 and 12 is 2

The common literals appearing in given monomial is x

The smallest power of x in two monomials is 2

The monomial of common literals with smallest power is x²

: The greatest common factor = $2x^2$

2. 6x³y and 18x²y³

Solution:

We know that the numerical coefficients of given numerical are 6 and 18

The greatest common factor of 6 and 18 is 6

Common literals appearing in given numerical are x and y

Smallest power of x in three monomial is 2

Smallest power of y in three monomial is 1

Monomial of common literals with smallest power is x²y

: The greatest common factor = $6x^2y$

3. 7x, 21x² and 14xy²

Solution:

We know that the numerical coefficients of given numerical are 7, 21 and 14

Greatest common factor of 7, 21 and 14 is 7

Common literals appearing in given numerical are x and y

Smallest power of x in three monomials is 1

Smallest power of y in three monomials is 0

Monomials of common literals with smallest power is x

 \therefore The greatest common factor = 7x

4. 42x²yz and 63x³y²z³

Solution:

We know that the numerical coefficients of given numerical are 42 and 63.

Greatest common factor of 42, 63 is 21. Common literals appearing in given numerical are x, y and z Smallest power of x in two monomials is 2 Smallest power of y in two monomials is 1 Smallest power of z in two monomials is 1 Monomials of common literals with smallest power is x^2yz \therefore The greatest common factor = $21x^2yz$

5. 12ax², 6a²x³ and 2a³x⁵

Solution:

We know that the numerical coefficients of given numerical are 12, 6 and 2

Greatest common factor of 12, 6 and 2 is 2.

Common literals appearing in given numerical are a and x

Smallest power of x in three monomials is 2

Smallest power of a in three monomials is 1

Monomials of common literals with smallest power is ax²

: The greatest common factor = 2ax²

6. 9x², 15x²y³, 6xy² and 21x²y²

Solution:

We know that the numerical coefficients of given numerical are 9, 15, 16 and 21

Greatest common factor of 9, 15, 16 and 21 is 3.

Common literals appearing in given numerical are x and y

Smallest power of x in four monomials is 1

Smallest power of y in four monomials is 0

Monomials of common literals with smallest power is x

∴ The greatest common factor = 3x

7. 4a²b³, -12a³b, 18a⁴b³

Solution:

We know that the numerical coefficients of given numerical are 4, -12 and 18.

Greatest common factor of 4, -12 and 18 is 2.

Common literals appearing in given numerical are a and b

Smallest power of a in three monomials is 2

Smallest power of b in three monomials is 1

Monomials of common literals with smallest power is a²b

: The greatest common factor = 2a²b

8. 6x²y², 9xy³, 3x³y²

Solution:

We know that the numerical coefficients of given numerical are 6, 9 and 3

Greatest common factor of 6, 9 and 3 is 3.

Common literals appearing in given numerical are x and y

Smallest power of x in three monomials is 1

Smallest power of y in three monomials is 2

Monomials of common literals with smallest power is xy²

: The greatest common factor = 3xy²

9. a²b³, a³b²

Solution:

We know that the numerical coefficients of given numerical are 0

Common literals appearing in given numerical are a and b

Smallest power of a in two monomials = 2

Smallest power of b in two monomials = 2

Monomials of common literals with smallest power is a²b²

: The greatest common factor = a²b²

10. 36a²b²c⁴, 54a⁵c², 90a⁴b²c²

Solution:

We know that the numerical coefficients of given numerical are 36, 54 and 90

Greatest common factor of 36, 54 and 90 is 18.

Common literals appearing in given numerical are a, b and c

Smallest power of a in three monomials is 2

Smallest power of b in three monomials is 0

Smallest power of c in three monomials is 2

Monomials of common literals with smallest power is a²c²

: The greatest common factor = 18a²c²

11. x³, -yx²

Solution:

We know that the numerical coefficients of given numerical are 0 Common literals appearing in given numerical are x and y Smallest power of x in two monomials is 2 Smallest power of y in two monomials is 0

Monomials of common literals with smallest power is x²

: The greatest common factor = x²

12. 15a3, -45a2, -150a

Solution:

We know that the numerical coefficients of given numerical are 15, -45 and 150

Greatest common factor of 15, -45 and 150 is 15.

Common literals appearing in given numerical is a

Smallest power of a in three monomials is 1

Monomials of common literals with smallest power is a

: The greatest common factor = 15a

13. 2x³y², 10x²y³, 14xy

Solution:

We know that the numerical coefficients of given numerical are 2, 10 and 14.

Greatest common factor of 2, 10 and 14 is 2.

Common literals appearing in given numerical are x and y

Smallest power of x in three monomials is 1

Smallest power of y in three monomials is 1

Monomials of common literals with smallest power is xy

∴ The greatest common factor = 2xy

14. 14x³y⁵, 10x⁵y³, 2x²y²

Solution:

We know that the numerical coefficients of given numerical are 14, 10 and 2.

Greatest common factor of 14, 10 and 2 is 2.

Common literals appearing in given numerical are x and y

Smallest power of x in three monomials is 2

Smallest power of y in three monomials is 2

Monomials of common literals with smallest power is x²y²

: The greatest common factor = $2x^2y^2$

Find the greatest common factor of the terms in each of the following expressions:

15. 5a⁴ + 10a³ – 15a²

Solution:

The greatest common factor of the three terms is 5a²

16. 2xyz + 3x²y + 4y²

Solution:

The greatest common factor of the three terms is y

17. $3a^2b^2 + 4b^2c^2 + 12a^2b^2c^2$

Solution:

The greatest common factor of the three terms is b².

EXERCISE 7.2 PAGE NO: 7.5

Factorize the following:

1. 3x – 9

Solution:

The greatest common factor in the given two terms is 3

3x – 9

3 (x – 3)

2. 5x - 15x²

Solution:

The greatest common factor in the given two terms is 5x

 $5x - 15x^2$

5x (1 – 3x)

3. 20a¹²b² – 15a⁸b⁴

Solution:

Greatest common factor in the given two terms is 5a8b2

20a12b2 - 15a8b4

5a⁸b² (4a⁴ - 3b²)

4. 72x⁶y⁷ – 96x⁷y⁶ Solution:

Greatest common factor in the given two terms is 24x⁶y⁶

 $72x^6y^7 - 96x^7y^6$

 $24x^{6}y^{6}(3y-4x)$

5. 20x³ - 40x² + 80x

Solution:

Greatest common factor in the given three terms is 20x

 $20x^3 - 40x^2 + 80x$

 $20x(x^2-2x+4)$

6. 2x³y² - 4x²y³ + 8xy⁴

Solution:

Greatest common factor in the given three terms is 2xy²

 $2x^{3}y^{2} - 4x^{2}y^{3} + 8xy^{4}$

 $2xy^{2}(x^{2}-2xy+4y^{2})$

7. 10m³n² + 15m⁴n – 20m²n³

Solution:

Greatest common factor in the given three terms is 5mn²

 $10m^{3}n^{2} + 15m^{4}n - 20m^{2}n^{3}$

 $5m^2n(2mn + 3m^2 - 4n^2)$

8. 2a^₄b^₄ – 3a^₃b^₅ + 4a^₂b^₅

Solution:

Greatest common factor in the given three terms is a2b4

2a4b4 - 3a3b5 + 4a2b5

a²b⁴ (2a² - 3ab + 4b)

9. 28a² + 14a²b² - 21a⁴

Solution:

Greatest common factor in the given three terms is 7a²

28a² + 14a²b² - 21a⁴

7a²(4a + 2b² – 3a²)

10. a⁴b – 3a²b² – 6ab³

Solution:

Greatest common factor in the given three terms is ab

 $a^4b - 3a^2b^2 - 6ab^3$

 $ab(a^{3}-3ab-6b^{2})$

11. 2l²mn – 3lm²n + 4lmn²

Solution:

Greatest common factor in the given three terms is Imn

 $2l^2mn - 3lm^2n + 4lmn^2$

lmn (2l - 3m + 4n)

12. $x^4y^2 - x^2y^4 - x^4y^4$

Greatest common factor in the given three terms is x²y²

 $X^4 y^2 - X^2 y^4 - X^4 y^4$

 $X^{2}Y^{2}(X^{2} - Y^{2} - X^{2}Y^{2})$

13. 9x²y + 3axy

Solution:

Greatest common factor in the given three terms is 3xy

 $9x^2y + 3axy$

3xy(3x + a)

14. 16m – 4m²

Solution:

Greatest common factor in the given two terms is 4m

16m – 4m²

4m (4 – m)

15. -4a² + 4ab – 4ca

Solution:

Greatest common factor in the given three terms is - 4a

-4a² + 4ab – 4ca

-4a (a – b + c)

```
16. x<sup>2</sup>yz + xy<sup>2</sup>z + xyz<sup>2</sup>
```

Solution:

Greatest common factor in the given three terms is xyz

 $x^2yz + xy^2z + xyz^2$

xyz (x + y + z)

17. ax²y + bxy² + cxyz

Solution:

Greatest common factor in the given three terms is xy

 $ax^2y + bxy^2 + cxyz$

xy (ax + by + cz)

EXERCISE 7.3 PAGE NO: 7.7

Factorize each of the following algebraic expressions:

```
1. 6x (2x - y) + 7y (2x - y)
```

We have,

6x (2x−y) + 7y (2x−y)

By taking (2x - y) as common we get,

(6x + 7y) (2x – y)

2. 2r(y - x) + s(x - y)

Solution:

We have,

2r(y-x) + s(x-y)

By taking (-1) as common we get,

-2r(x - y) + s(x - y)

By taking (x - y) as common we get,

(x – y) (-2r + s)

(x - y) (s - 2r)

3. 7a (2x – 3) + 3b (2x – 3)

Solution:

We have,

7a (2x - 3) + 3b (2x - 3)

By taking (2x - 3) as common we get,

(7a + 3b) (2x - 3)

4. 9a (6a - 5b) - 12a² (6a - 5b)

Solution:

We have,

9a (6a – 5b) – 12a² (6a – 5b)

By taking (6a - 5b) as common we get,

(9a – 12a²) (6a – 5b)

3a(3 – 4a) (6a – 5b)

5. 5 $(x - 2y)^2$ + 3 (x - 2y)

Solution:

We have,

 $5(x-2y)^2 + 3(x-2y)$

By taking (x - 2y) as common we get,

(x - 2y) [5 (x - 2y) + 3](x - 2y) (5x - 10y + 3)6. 16 $(2I - 3m)^2 - 12 (3m - 2I)$ Solution: We have, $16 (2I - 3m)^2 - 12 (3m - 2I)$ By taking (-1) as common we get, $16 (2l - 3m)^2 + 12 (2l - 3m)$ By taking 4(2l - 3m) as common we get, 4(2I - 3m) [4 (2I - 3m) + 3]4(2I - 3m)(8I - 12m + 3)7. 3a (x - 2y) - b (x - 2y)Solution: We have, 3a(x-2y) - b(x-2y)By taking (x - 2y) as common we get, (3a - b) (x - 2y)8. $a^{2}(x + y) + b^{2}(x + y) + c^{2}(x + y)$ Solution: We have, $a^{2}(x + y) + b^{2}(x + y) + c^{2}(x + y)$ By taking (x + y) as common we get, $(a^2 + b^2 + c^2) (x + y)$ 9. $(x - y)^2 + (x - y)$ Solution: We have, $(x - y)^2 + (x - y)$ By taking (x - y) as common we get, (x - y) (x - y + 1)10. 6 $(a + 2b) - 4 (a + 2b)^2$ Solution: We have, 6 (a + 2b) - 4 (a + 2b)²

By taking (a + 2b) as common we get,

[6 - 4 (a + 2b)] (a + 2b)(6 - 4a - 8b) (a + 2b)

2(3 – 2a – 4b) (a + 2b)

11. a (x - y) + 2b (y - x) + c (x - y)²

Solution:

We have,

a $(x - y) + 2b (y - x) + c (x - y)^2$

By taking (-1) as common we get,

a $(x - y) - 2b (x - y) + c (x - y)^2$

By taking (x - y) as common we get,

[a-2b + c(x - y)] (x - y)(x - y) (a - 2b + cx - cy)

12. $-4(x-2y)^2 + 8(x-2y)$

Solution:

We have,

 $-4 (x - 2y)^2 + 8 (x - 2y)$

By taking 4(x - 2y) as common we get,

 $\frac{[-(x-2y)+2]}{4(x-2y)} + \frac{4(x-2y)}{(-x+2y+2)}$

13. x³ (a – 2b) + x² (a – 2b)

Solution:

We have,

 $x^{3}(a-2b) + x^{2}(a-2b)$

By taking x^2 (a – 2b) as common we get,

 $(x + 1) [x^2 (a - 2b)]$

 $x^{2}(a-2b)(x+1)$

14. (2x - 3y) (a + b) + (3x - 2y) (a + b)

Solution:

We have,

(2x - 3y) (a + b) + (3x - 2y) (a + b)By taking (a + b) as common we get, (a + b) [(2x - 3y) + (3x - 2y)](a + b) [2x - 3y + 3x - 2y] (a + b) [5x - 5y] (a + b) 5(x - y) **15.** 4(x + y) (3a - b) + 6(x + y) (2b - 3a) **Solution:** We have, 4(x + y) (3a - b) + 6(x + y) (2b - 3a) By taking (x + y) as common we get, (x + y) [4(3a - b) + 6(2b - 3a)] (x + y) [12a - 4b + 12b - 18a] (x + y) [-6a + 8b] (x + y) 2(-3a + 4b) (x + y) 2(4b - 3a)

EXERCISE 7.4 PAGE NO: 7.12

Factorize each of the following expressions:

1. qr – pr + qs – ps Solution: We have, qr - pr + qs - ps By grouping similar terms we get, qr + qs – pr – ps q(r + s) - p(r + s)(q - p) (r + s)2. p²q – pr² – pq + r² Solution: We have, $p^2q - pr^2 - pq + r^2$ By grouping similar terms we get, $p^2q - pq - pr^2 + r^2$ $pq(p-1) - r^2(p-1)$ $(p-1)(pq-r^2)$ 3. $1 + x + xy + x^2y$

We have,

 $1 + x + xy + x^2y$ 1(1 + x) + xy(1 + x)(1 + x) (1 + xy)4. ax + ay – bx – by Solution: We have, ax + ay - bx - bya(x + y) - b(x + y)(a - b) (x + y)5. xa² + xb² - ya² - yb² Solution: We have, $xa^{2} + xb^{2} - ya^{2} - yb^{2}$ $x(a^2 + b^2) - y(a^2 + b^2)$ $(x - y) (a^2 + b^2)$ 6. x² + xy + xz + yz Solution: We have, $x^2 + xy + xz + yz$

x (x + y) + z (x + y)

(x + y) (x + z)

7. 2ax + bx + 2ay + by

Solution:

We have,

2ax + bx + 2ay + by

By grouping similar terms we get,

2ax + 2ay + bx + by

2a (x + y) + b (x + y)

(2a + b) (x + y)

8. ab – by – ay + y²

Solution:

We have,

```
ab - by - ay + y^2
```

By grouping similar terms we get,

 $Ab - ay - by + y^2$

 $a\ (b-y)-y\ (b-y)$

$$(a - y) (b - y)$$

9. axy + bcxy – az – bcz

Solution:

We have,

axy + bcxy – az – bcz

By grouping similar terms we get,

axy - az + bcxy - bcz

$$a(xy-z) + bc(xy-z)$$

(a + bc) (xy – z)

10. lm² – mn² – lm + n²

Solution:

We have,

 $Im^2 - mn^2 - Im + n^2$

By grouping similar terms we get,

```
Im^2 - Im - mn^2 + n^2
```

 $lm (m - 1) - n^2 (m - 1)$

 $(Im - n^2) (m - 1)$

11. $x^3 - y^2 + x - x^2y^2$

Solution:

We have,

 $x^3 - y^2 + x - x^2y^2$

By grouping similar terms we get,

 $x + x^3 - y^2 - x^2y^2$

 $x(1 + x^2) - y^2(1 + x^2)$

 $(x - y^2)(1 + x^2)$

12. 6xy + 6 – 9y – 4x

Solution:

We have,

6xy + 6 - 9y - 4xBy grouping similar terms we get, 6xy - 4x - 9y + 62x(3y-2) - 3(3y-2)(2x - 3)(3y - 2)13. $x^2 - 2ax - 2ab + bx$ Solution: We have, $x^2 - 2ax - 2ab + bx$ By grouping similar terms we get, x^2 + bx – 2ax – 2ab x(x + b) - 2a(x + b)(x - 2a) (x + b)14. $x^3 - 2x^2y + 3xy^2 - 6y^3$ Solution: We have, $x^3 - 2x^2y + 3xy^2 - 6y^3$ By grouping similar terms we get, $x^3 + 3xy^2 - 2x^2y - 6y^3$ $x(x^2 + 3y^2) - 2y(x^2 + 3y^2)$ $(x - 2y) (x^2 + 3y^2)$ 15. abx² + (ay – b) x – y Solution: We have, $abx^{2} + (ay - b)x - y$ $abx^{2} + ayx - bx - y$ By grouping similar terms we get, $abx^2 - bx + ayx - y$ bx(ax - 1) + y(ax - 1)(bx + y) (ax - 1)16. $(ax + by)^2 + (bx - ay)^2$ Solution: We have,

```
(ax + by)^2 + (bx - ay)^2

a^2x^2 + b^2y^2 + 2axby + b^2x^2 + a^2y^2 - 2axby

a^2x^2 + b^2y^2 + b^2x^2 + a^2y^2

By grouping similar terms we get,

a^2x^2 + a^2y^2 + b^2y^2 + b^2x^2

a^2 (x^2 + y^2) + b^2 (x^2 + y^2)

(a^2 + b^2) (x^2 + y^2)

17. 16 (a - b)^3 - 24 (a - b)^2

Solution:
```

We have,

 $16(a - b)^{3} - 24(a - b)^{2}$ $8 (a - b)^{2} [2 (a - b) - 3]$ $8 (a - b)^{2} (2a - 2b - 3)$ $18. ab (x^{2} + 1) + x (a^{2} + b^{2})$

Solution:

We have,

```
ab(x^{2} + 1) + x(a^{2} + b^{2})
abx^2 + ab + xa^2 + xb^2
By grouping similar terms we get,
abx^{2} + xa^{2} + xb^{2} + ab
ax(bx + a) + b(bx + a)
(ax + b)(bx + a)
19. a<sup>2</sup>x<sup>2</sup> + (ax<sup>2</sup> + 1)x + a
Solution:
We have,
a^{2}x^{2} + (ax^{2} + 1)x + a
a^{2}x^{2} + ax^{3} + x + a
ax^{2}(a + x) + 1(x + a)
(x + a) (ax^{2} + 1)
20. a(a - 2b - c) + 2bc
Solution:
We have,
```

a(a - 2b - c) + 2bc

 $a^2 - 2ab - ac + 2bc$ a(a-2b) - c(a-2b)(a – 2b) (a – c) 21. a (a + b – c) – bc Solution: We have, a(a+b-c)-bc $a^2 + ab - ac - bc$ a(a + b) - c(a + b)(a + b) (a - c)22. $x^2 - 11xy - x + 11y$ Solution: We have, $x^2 - 11xy - x + 11y$ By grouping similar terms we get, $x^2 - x - 11xy + 11y$ x(x-1) - 11y(x-1)(x - 11y)(x - 1)23. ab – a – b + 1 Solution: We have, ab – a – b + 1 a (b - 1) - 1 (b - 1) (a – 1) (b – 1) 24. $x^2 + y - xy - x$ Solution: We have, $x^2 + y - xy - x$ By grouping similar terms we get, $x^{2} - x + y - xy$ x(x-1) - y(x-1)(x - y) (x - 1)

EXERCISE 7.5 PAGE NO: 7.17

Factorize each of the following expressions: 1. 16x² – 25y² Solution: We have, $16x^2 - 25y^2$ $(4x)^2 - (5y)^2$ By using the formula $(a^2 - b^2) = (a + b) (a - b)$ we get, (4x + 5y) (4x - 5y)2. 27x² – 12y² Solution: We have, $27x^2 - 12y^2$ By taking 3 as common we get, $3[(3x)^2 - (2y)^2]$ By using the formula $(a^2 - b^2) = (a-b)(a+b)$ 3(3x + 2y)(3x - 2y)3. 144a² - 289b² Solution: We have, 144a² - 289b² (12a)² – (17b)² By using the formula $(a^2 - b^2) = (a-b) (a+b)$ (12a + 17b) (12a – 17b) 4. 12m² – 27 Solution: We have, 12m² – 27 By taking 3 as common we get, $3(4m^2 - 9)$ $3[(2m)^2 - 3^2]$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ 3 (2m + 3) (2m - 3)

5. 125x² – 45y² Solution: We have, $125x^2 - 45y^2$ By taking 5 as common we get, $5(25x^2 - 9y^2)$ $5[(5x)^2 - (3y)^2]$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ 5(5x + 3y)(5x - 3y)6. 144a² – 169b² Solution: We have, 144a² - 169b² $(12a)^2 - (13b)^2$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ (12a + 13b) (12a - 13b) 7. (2a – b)² – 16c² Solution: We have, $(2a - b)^2 - 16c^2$ $(2a - b)^2 - (4c)^2$ By using the formula $(a^2 - b^2) = (a-b)(a+b)$ (2a - b + 4c) (2a - b - 4c)8. $(x + 2y)^2 - 4 (2x - y)^2$ Solution: We have, $(x + 2y)^2 - 4 (2x - y)^2$ $(x + 2y)^2 - [2(2x - y)]^2$ By using the formula $(a^2 - b^2) = (a + b) (a - b)$ we get, [(x + 2y) + 2(2x - y)] [x + 2y - 2(2x - y)](x + 4x + 2y - 2y) (x - 4x + 2y + 2y)(5x)(4y - 3x)9. 3a^₅ – 48a^₃

We have, 3a⁵ – 48a³ By taking 3 as common we get, $3a^3(a^2-16)$ $3a^3(a^2-4^2)$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $3a^{3}(a+4)(a-4)$ 10. a⁴ – 16b⁴ Solution: We have, a⁴ - 16b⁴ $(a^2)^2 - (4b^2)^2$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $(a^2 + 4b^2) (a^2 - 4b^2)$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $(a^2 + 4b^2) (a + 2b) (a - 2b)$ 11. x⁸ − 1 Solution: We have, $x^{\scriptscriptstyle 8}-1$ $(X^4)^2 - (1)^2$ By using the formula $(a^2 - b^2) = (a-b)(a+b)$ $(x^4 + 1) (x^4 - 1)$ By using the formula $(a^2 - b^2) = (a-b)(a+b)$ $(x^{4} + 1) (x^{2} + 1) (x - 1) (x + 1)$ 12. 64 – (a + 1)² Solution: We have, $64 - (a + 1)^2$ $8^2 - (a + 1)^2$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ [8 + (a + 1)] [8 - (a + 1)]

```
(a + 9) (7 - a)
13. 36l<sup>2</sup> – (m + n)<sup>2</sup>
Solution:
We have,
36l^2 - (m + n)^2
(6I)^2 - (m + n)^2
By using the formula (a^2 - b^2) = (a-b) (a+b)
(6l + m + n) (6l - m - n)
14. 25x⁴y⁴ – 1
Solution:
We have,
25x^{4}y^{4} - 1
(5x^2y^2)^2 - (1)^2
By using the formula (a^2 - b^2) = (a-b) (a+b)
(5x^2y^2 - 1)(5x^2y^2 + 1)
15. a⁴ – 1/b⁴
Solution:
We have,
a4 – 1/b4
(a^2)^2 - (1/b^2)^2
By using the formula (a^2 - b^2) = (a-b)(a+b)
(a^2 + 1/b^2) (a^2 - 1/b^2)
By using the formula (a^2 - b^2) = (a-b) (a+b)
(a^2 + 1/b^2) (a - 1/b) (a + 1/b)
16. x<sup>3</sup> – 144x
Solution:
We have,
x<sup>3</sup> - 144x
x [x^2 - (12)^2]
By using the formula (a^2 - b^2) = (a-b) (a+b)
x (x + 12) (x - 12)
17. (x - 4y)<sup>2</sup> - 625
Solution:
```

We have,

 $(x - 4y)^2 - 625$ $(x - 4y)^2 - (25)^2$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$

(x - 4y + 25) (x - 4y - 25)

18. 9 (a – b)² – 100 (x – y)²

Solution:

We have,

9 $(a - b)^2 - 100 (x - y)^2$ [3 (a - b)]² - [10 (x - y)]² By using the formula $(a^2 - b^2) = (a-b) (a+b)$

[3 (a - b) + 10 (x + y)] [3 (a - b) - 10 (x - y)] [3a - 3b + 10x - 10y] [3a - 3b - 10x + 10y]**19.** $(3 + 2a)^2 - 25a^2$

Solution:

We have,

(3 + 2a)² - 25a²

$$(3 + 2a)^2 - (5a)^2$$

By using the formula $(a^2 - b^2) = (a-b)(a+b)$

(3 + 2a + 5a)(3 + 2a - 5a)

(3 + 7a) (3 – 3a)

(3 + 7a) 3(1 - a)

20. $(x + y)^2 - (a - b)^2$

Solution:

We have,

 $(x + y)^2 - (a - b)^2$

By using the formula $(a^2 - b^2) = (a-b) (a+b)$

 $\begin{array}{l} [(x + y) + (a - b)] [(x + y) - (a - b)] \\ (x + y + a - b) (x + y - a + b) \end{array}$

21. 1/16x²y² - 4/49y²z²

Solution:

We have,

 $1/16x^2y^2 - 4/49y^2z^2$

 $(1/4xy)^2 - (2/7yz)^2$

By using the formula $(a^2 - b^2) = (a-b) (a+b)$

(xy/4 + 2yz/7) (xy/4 - 2yz/7) $y^{2}(x/4 + 2/7z)(x/4 - 2/7z)$ 22. 75a³b² – 108ab⁴ Solution: We have, 75a3b2 - 108ab4 3ab² (25a² - 36b²) $3ab^{2}[(5a)^{2}-(6b)^{2}]$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $3ab^{2}(5a + 6b)(5a - 6b)$ 23. x⁵ – 16x³ Solution: We have, x⁵ − 16x³ $x^{3}(x^{2}-16)$ $x^{3}(x^{2}-4^{2})$ By using the formula $(a^2 - b^2) = (a-b)(a+b)$ $x^{3}(x + 4)(x - 4)$ 24. 50/x² - 2x²/81 Solution: We have, $50/x^2 - 2x^2/81$ 2 (25/x² - x²/81) $2[(5/x)^2 - (x/9)^2]$ By using the formula $(a^2 - b^2) = (a-b)(a+b)$ 2(5/x+x/9)(5/x-x/9)25. 256x³ – 81x Solution: We have, 256x³ – 81x x (256x₄ – 81) $x [(16x^2)^2 - 9^2]$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$

We have, $p^2q^2 - p^4q^4$ $(pq)^2 - (p^2q^2)^2$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $(pq + p^2q^2) (pq - p^2q^2)$ $p^{2}q^{2}(1 + pq)(1 - pq)$ 29. 3x³y - 243xy³ Solution: We have, 3x³y - 243xy³ $3xy(x^2 - 81y^2)$ $3xy [x^2 - (9y)^2]$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ (3xy) (x + 9y) (x - 9y)30. a⁴b⁴ – 16c⁴

 $[(3x + 4y)^{2} + x^{2}] [(3x + 4y)^{2} - x^{2}] [(3x + 4y)^{2} + x^{2}] [3x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y] [2x + 4y] [2x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y] [2x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y] [2x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y + x] [3x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y + x] [3x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y + x] [3x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y + x] [3x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y + x] [3x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y + x] [3x + 4y + x] [3x + 4y - x] [(3x + 4y)^{2} + x^{2}] [4x + 4y + x] [3x + 4y$ $+ 4y [(3x + 4y)^{2} + x^{2}] 8[x + 2y] [x + y]$ **28.** $p^2q^2 - p^4q^4$

 $[(3x + 4y)^2]^2 - (x^2)^2$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$

By using the formula $(a^2 - b^2) = (a-b) (a+b)$

By using the formula $(a^2 - b^2) = (a-b) (a+b)$

 $[a^2 + (2b + c)^2] [a + 2b + c] [a - 2b - c]$

 $(3x + 4y)^4 - x^4$

Solution.

27. (3x + 4y)⁴ - x⁴

 $x (4x + 3) (4x - 3) (16x^{2} + 9)$

 $[a^2 + (2b + c)^2] [a^2 - (2b + c)^2]$

26. a^₄ – (2b + c)^₄

Solution: We have,

a⁴ - (2b + c)⁴

(a²)² - [(2b + c)²]²

vve nave,

Solution:

We have, a4b4 - 16c4 $(a^2b^2)^2 - (4c^2)^2$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $(a^{2}b^{2} + 4c^{2})(a^{2}b^{2} - 4c^{2})$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $(a^{2}b^{2} + 4c^{2})(ab + 2c)(ab - 2c)$ 31. x⁴ – 625 Solution: We have, x⁴ - 625 $(X^2)^2 - (25)^2$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $(x^2 + 25) (x^2 - 25)$ $(x^2 + 25) (x^2 - 5^2)$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $(x^{2} + 25) (x + 5) (x - 5)$ 32. x⁴ – 1 Solution: We have, x4 - 1 $(X^2)^2 - (1)^2$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $(x^2 + 1) (x^2 - 1)$

By using the formula $(a^2 - b^2) = (a-b) (a+b)$

 $(x^{2} + 1)(x + 1)(x - 1)$

33. 49(a - b)² - 25(a + b)²

Solution:

We have,

 $49(a - b)^2 - 25(a + b)^2$

 $[7 (a - b)]^2 - [5 (a + b)]^2$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ [7 (a - b) + 5 (a + b)] [7 (a - b) - 5 (a + b)](7a - 7b + 5a + 5b) (7a - 7b - 5a - 5b) (12a - 2b) (2a - 12b) 2 (6a - b) 2 (a - 6b) 4 (6a - b) (a - 6b) **34. x - y - x² + y²**

Solution:

We have,

 $x - y - x^2 + y^2$

 $x - y - (x^2 - y^2)$

By using the formula $(a^2 - b^2) = (a-b) (a+b)$

x - y - (x + y) (x - y)

(x - y) (1 - x - y)

35. 16(2x - 1)² - 25y²

Solution:

We have,

 $16(2x-1)^2 - 25y^2$

 $[4 (2x - 1)]^2 - (5y)^2$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$

(8x + 5y - 4) (8x - 5y - 4)

36. $4(xy + 1)^2 - 9(x - 1)^2$

Solution:

We have,

 $4(xy + 1)^2 - 9(x - 1)^2$

 $[2 (xy + 1)]^2 - [3 (x - 1)]^2$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$

(2xy + 2 + 3x - 3)(2xy + 2 - 3x + 3)

(2xy + 3x - 1)(2xy - 3x + 5)

37. (2x + 1)² - 9x⁴

Solution:

We have,

 $(2x + 1)^2 - 9x^4$

 $(2x + 1)^2 - (3x^2)^2$

By using the formula $(a^2 - b^2) = (a-b) (a+b)$

 $(2x + 1 + 3x^2)(2x + 1 - 3x^2)$ $(3x^2 + 2x + 1)(-3x^2 + 2x + 1)$ 38. x⁴ - (2y - 3z)² Solution: We have, $x^4 - (2y - 3z)^2$ $(x^2)^2 - (2y - 3z)^2$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $(x^2 + 2y - 3z) (x^2 - 2y + 3z)$ **39.** $a^2 - b^2 + a - b$ Solution: We have, $a^2 - b^2 + a - b$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ (a + b) (a - b) + (a - b)(a - b) (a + b + 1)40. 16a⁴ – b⁴ Solution: We have, 16a⁴ – b⁴ $(4a^2)^2 - (b^2)^2$ $(4a^2 + b^2) (4a^2 - b^2)$ By using the formula $(a^2 - b^2) = (a-b)(a+b)$ $(4a^2 + b^2) (2a + b) (2a - b)$ 41. a^₄ – 16(b – c)^₄ Solution: We have, $a^4 - 16(b - c)^4$ $(a^2)^2 - [4 (b - c)^2]$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $[a^2 + 4 (b - c)^2] [a^2 - 4 (b - c)^2]$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $[a^2 + 4 (b - c)^2] [(a + 2b - 2c) (a - 2b + 2c)]$ 42. 2a^₅ – 32a

We have, 2a⁵ – 32a 2a (a⁴ – 16) $2a[(a^2)^2 - (4)^2]$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $2a(a^2 + 4)(a^2 - 4)$ $2a(a^2 + 4)(a^2 - 2^2)$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $2a(a^{2}+4)(a+2)(a-2)$ 43. a⁴b⁴ – 81c⁴ Solution: We have, a4b4 - 81c4 $(a^2b^2)^2 - (9c^2)^2$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $(a^2b^2 + 9c^2)(a^2b^2 - 9c^2)$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $(a^{2}b^{2} + 9c^{2})(ab + 3c)(ab - 3c)$ 44. xy⁹ – yx⁹ Solution: We have, $xy^9 - yx^9$ $-xy(x^{8}-y^{8})$ -xy $[(x^4)^2 - (y^4)^2]$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $-xy(x^{4} + y^{4})(x^{4} - y^{4})$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $-xy(x^{4} + y^{4})(x^{2} + y^{2})(x^{2} - y^{2})$ By using the formula $(a^2 - b^2) = (a-b) (a+b)$ $-xy(x^{4} + y^{4})(x^{2} + y^{2})(x + y)(x - y)$ 45. x³ − x Solution:

We have,

 $x^{3} - x$ $x (x^{2} - 1)$ By using the formula $(a^{2} - b^{2}) = (a-b) (a+b)$ x (x + 1) (x - 1) **46.** 18a²x² - 32 **Solution:** We have, $18a^{2}x^{2} - 32$ $2 [(3ax)^{2} - (4)^{2}]$ By using the formula $(a^{2} - b^{2}) = (a-b) (a+b)$ 2 (3ax + 4) (3ax - 4)

EXERCISE 7.6 PAGE NO: 7.22

```
Factorize each of the following algebraic expressions:
1. 4x<sup>2</sup> + 12xy + 9y<sup>2</sup>
Solution:
We have,
4x^2 + 12xy + 9y^2
By using the formula (x + y)^2 = x^2 + y^2 + 2xy
(2x)^2 + (3y)^2 + 2(2x)(3y)
(2x + 3y)^2
(2x + 3y) (2x + 3y)
2. 9a<sup>2</sup> - 24ab + 16b<sup>2</sup>
Solution:
We have,
9a<sup>2</sup> - 24ab + 16b<sup>2</sup>
By using the formula (x - y)^2 = x^2 + y^2 - 2xy
Here x = 3a, y = 4b So,
(3a)^2 + (4b)^2 - 2 (3a) (4b)
(3a – 4b)<sup>2</sup>
(3a - 4b) (3a - 4b)
```

3. p²q² – 6pqr + 9r²

Solution:

We have,

 $p^2q^2 - 6pqr + 9r^2$

By using the formula $(a - b)^2 = a^2 + b^2 - 2ab$

 $(pq)^{2} + (3r)^{2} - 2 (pq) (3r)$

(pq - 3r)²

(pq - 3r) (pq - 3r)

4. 36a² + 36a + 9

Solution:

We have,

```
36a<sup>2</sup> + 36a + 9
```

```
(6a)^2 + 2 \times 6a \times 3 + 3^2
```

(6a + 3)²

5. a² + 2ab + b² - 16

Solution:

We have,

a² + 2ab + b² - 16

By using the formula $(a - b)^2 = a^2 + b^2 - 2ab$

 $(a + b)^2 - 4^2$

By using the formula $(a^2 - b^2) = (a+b) (a-b)$

(a + b + 4) (a + b - 4)

6. $9z^2 - x^2 + 4xy - 4y^2$

Solution:

We have,

 $9z^2 - x^2 + 4xy - 4y^2$

 $(3z)^2 - [x^2 - 2(x)(2y) + (2y)^2]$

By using the formula $(a - b)^2 = a^2 + b^2 - 2ab$

 $(3z)^2 - (x - 2y)^2$

By using the formula $(a^2 - b^2) = (a+b) (a-b)$

[(x - 2y) + 3z] [-x + 2y + 3z]7. 9a⁴ - 24a²b² + 16b⁴ - 256

Solution:

We have, $9a^4 - 24a^2b^2 + 16b^4 - 256$ $(3a^2)^2 - 2(4a^2)(3b^2) + (4b^2)^2 - (16)^2$ By using the formula $(a - b)^2 = a^2 + b^2 - 2ab$ $(3a^2 - 4b^2)^2 - (16)^2$ By using the formula $(a^2 - b^2) = (a+b)(a-b)$ $(3a^2 - 4b^2 + 16)(3a^2 - 4b^2 - 16)$ 8. 16 - a⁶ + 4a³b³ - 4b⁶

Solution:

We have,

 $16 - a^6 + 4a^3b^3 - 4b^6$ $4^2 - [(a^3)^2 - 2 (a^3) (2b^3) + (2b^3)^2]$

By using the formula $(a - b)^2 = a^2 + b^2 - 2ab$

 $4^2 - (a^3 - 2b^3)^2$

By using the formula $(a^2 - b^2) = (a+b) (a-b)$

 $\begin{array}{l} [4 + (a^{_3} - 2b^{_3})] \ [4 - (a^{_3} - 2b^{_3})] \\ \textbf{9. } a^{_2} \textbf{-2ab} + b^{_2} \textbf{-c}^{_2} \end{array}$

Solution:

We have, $a^2 - 2ab + b^2 - c^2$ By using the formula $(a - b)^2 = a^2 + b^2 - 2ab$ $(a - b)^2 - c^2$ By using the formula $(a^2 - b^2) = (a+b)$ (a-b) (a - b + c) (a - b - c)10. $x^2 + 2x + 1 - 9y^2$ Solution: We have, $x^2 + 2x + 1 - 9y^2$ By using the formula $(a - b)^2 = a^2 + b^2 - 2ab$ $(x + 1)^2 - (3y)^2$ By using the formula $(a^2 - b^2) = (a+b)$ (a-b) (x + 3y + 1) (x - 3y + 1)11. $a^2 + 4ab + 3b^2$

We have,

a² + 4ab + 3b²

By using factors for 3 i.e., 3 and 1

```
a^{2} + ab + 3ab + 3b^{2}
```

By grouping we get,

a(a + b) + 3b(a + b)

(a + 3b) (a + b)

12. 96 – 4x – x²

Solution:

We have,

 $96 - 4x - x^2$

 $-x^{2} - 4x + 96$

By using factors for 96 i.e., 12 and 8

 $-x^2 - 12x + 8x + 96$

By grouping we get,

```
-x (x + 12) + 8 (x + 12)
```

(x + 12) (-x + 8)

13. a⁴ + 3a² + 4

Solution:

We have,

```
a^{4} + 3a^{2} + 4
(a^{2})^{2} + (a^{2})^{2} + 2 (2a^{2}) + 4 - a^{2}
(a^{2} + 2)^{2} + (-a^{2})
By using the formula (a^{2} - b^{2}) = (a+b) (a-b)
(a^{2} + 2 + a) (a^{2} + 2 - a)
(a^{2} + a + 2) (a^{2} - a + 2)
14. 4x<sup>4</sup> + 1
Solution:
We have,
4x^{4} + 1
(2x^{2})^{2} + 1 + 4x^{2} - 4x^{2}
(2x^{2} + 1)^{2} - 4x^{2}
```

By using the formula $(a^2 - b^2) = (a+b) (a-b)$ $(2x^2 + 1 + 2x)(2x^2 + 1 - 2x)$ $(2x^2 + 2x + 1)(2x^2 - 2x + 1)$ 15. 4x⁴ + y⁴ Solution: We have, 4x⁴ + y⁴ $(2x^2)^2 + (y^2)^2 + 4x^2y^2 - 4x^2y^2$ $(2x^2 + y^2)^2 - 4x^2y^2$ By using the formula $(a^2 - b^2) = (a+b) (a-b)$ $(2x^2 + y^2 + 2xy) (2x^2 + y^2 - 2xy)$ 16. $(x + 2)^4 - 6(x + 2) + 9$ Solution: We have, $(x + 2)^4 - 6(x + 2) + 9$ $(x^2 + 2^2)^2 - 6x - 12 + 9$ $(x^2 + 2^2 + 2(2)(x)) - 6x - 12 + 9$ $x^{2} + 4 + 4x - 6x - 12 + 9$ $x^2 - 2x + 1$ By using the formula $(a - b)^2 = a^2 + b^2 - 2ab$ $(x - 1)^2$ 17. 25 – p² – q² – 2pq Solution: We have, $25 - p^2 - q^2 - 2pq$ $25 - (p^2 + q^2 + 2pq)$ $(5)^2 - (p + q)^2$ By using the formula $(a^2 - b^2) = (a+b) (a-b)$ (5 + p + q) (5 - p - q)-(p+q+5)(p+q-5)18. $x^2 + 9y^2 - 6xy - 25a^2$ Solution: We have,

 $x^{2} + 9y^{2} - 6xy - 25a^{2}$ $(x - 3y)^2 - (5a)^2$ By using the formula $(a^2 - b^2) = (a+b) (a-b)$ (x - 3y + 5a) (x - 3y - 5a)19. 49 - a² + 8ab - 16b² Solution: We have, 49 - a² + 8ab - 16b² $49 - (a^2 - 8ab + 16b^2)$ $49 - (a - 4b)^2$ By using the formula $(a^2 - b^2) = (a + b) (a - b)$ (7 + a - 4b) (7 - a + 4b)-(a - 4b + 7) (a - 4b - 7) 20. a² - 8ab + 16b² - 25c² Solution: We have, a² - 8ab + 16b² - 25c² $(a - 4b)^2 - (5c)^2$ By using the formula $(a^2 - b^2) = (a+b) (a-b)$ (a - 4b + 5c) (a - 4b - 5c)21. $x^2 - y^2 + 6y - 9$ Solution: We have, $x^2 - y^2 + 6y - 9$ $x^2 + 6y - (y^2 - 6y + 9)$ $x^2 - (y - 3)^2$ By using the formula $(a^2 - b^2) = (a+b) (a-b)$ (x + y - 3)(x - y + 3)22. 25x² - 10x + 1 - 36y² Solution: We have, $25x^2 - 10x + 1 - 36y^2$ $(5x)^2 - 2(5x) + 1 - (6y)^2$

 $(5x - 1)^2 - (6y)^2$ By using the formula $(a^2 - b^2) = (a+b) (a-b)$ (5x - 6y - 1)(5x + 6y - 1)23. $a^2 - b^2 + 2bc - c^2$ Solution: We have, $a^2 - b^2 + 2bc - c^2$ $a^2 - (b^2 - 2bc + c^2)$ $a^{2} - (b - c)^{2}$ By using the formula $(a^2 - b^2) = (a+b) (a-b)$ (a + b - c) (a - b + c)24. a^2 + 2ab + b^2 - c^2 Solution: We have, a² + 2ab + b² - c² $(a + b)^2 - c^2$ By using the formula $(a^2 - b^2) = (a+b) (a-b)$ (a + b + c) (a + b - c)25. 49 - x² - y² + 2xy Solution: We have, $49 - x^2 - y^2 + 2xy$ $49 - (x^2 + y^2 - 2xy)$ $7^2 - (x - y)^2$ By using the formula $(a^2 - b^2) = (a+b) (a-b)$ $\begin{array}{l} \left[7 + (x - y)\right] \left[7 - x + y\right] \\ (x - y + 7) \left(y - x + 7\right) \end{array}$ 26. a² + 4b² - 4ab - 4c² Solution: We have, $a^{2} + 4b^{2} - 4ab - 4c^{2}$ $a^{2} - 2(a)(2b) + (2b)^{2} - (2c)^{2}$ $(a - 2b)^2 - (2c)^2$

By using the formula $(a^2 - b^2) = (a+b) (a-b)$ (a - 2b + 2c) (a - 2b - 2c) **27.** $x^2 - y^2 - 4xz + 4z^2$ **Solution:** We have, $x^2 - y^2 - 4xz + 4z^2$ $x^2 - 2 (x) (2z) + (2z)^2 - y^2$ As $(a-b)^2 = a^2 + b^2 - 2ab$ $(x - 2z)^2 - y^2$ By using the formula $(a^2 - b^2) = (a+b) (a-b)$ (x + y - 2z) (x - y - 2z)

EXERCISE 7.7 PAGE NO: 7.27

Factorize each of the following algebraic expressions:

1. x² + 12x - 45

Solution:

We have,

x² + 12x - 45

To factorize the given expression we have to find two numbers p and q such that p+q = 12 and pq = -45

So we can replace 12x by 15x - 3x

-45 by 15 × 3

 $x^2 + 12x - 45 = x^2 + 15x - 3x - 45$

= x (x + 15) - 3 (x + 15)

= (x - 3) (x + 15)

2. 40 +
$$3x - x^2$$

Solution:

We have,

 $40 + 3x - x^2$

 $-(x^2 - 3x - 40)$

By considering, p+q = -3 and pq = -40

So we can replace -3x by 5x - 8x

```
-40 by 5 × -8
-(x^2 - 3x - 40) = x^2 + 5x - 8x - 40
= -x (x + 5) - 8 (x + 5)
= -(x - 8) (x + 5)
= (-x + 8) (x + 5)
3. a<sup>2</sup> + 3a - 88
Solution:
We have,
a<sup>2</sup> + 3a – 88
By considering, p+q = 3 and pq = -88
So we can replace 3a by 11a - 8a
-40 by -11 × 8
a^{2} + 3a - 88 = a^{2} + 11a - 8a - 88
= a (a + 11) - 8 (a + 11)
= (a - 8) (a + 11)
4. a<sup>2</sup> – 14a – 51
Solution:
We have,
a<sup>2</sup> - 14a - 51
By considering, p+q = -14 and pq = -51
So we can replace -14a by 3a - 17a
-51 by -17 × 3
a^2 - 14a - 51 = a^2 + 3a - 17a - 51
= a (a + 3) - 17 (a + 3)
= (a - 17) (a + 3)
5. x<sup>2</sup> + 14x + 45
Solution:
We have,
x^2 + 14x + 45
By considering, p+q = 14 and pq = 45
So we can replace 14x by 5x + 9x
45 by 5 × 9
x^{2} + 14x + 45 = x^{2} + 5x + 9x + 45
```

= x (x + 5) - 9 (x + 5)= (x + 9) (x + 5)6. $x^2 - 22x + 120$ Solution: We have, $x^2 - 22x + 120$ By considering, p+q = -22 and pq = 120So we can replace -22x by -12x -10x 120 by -12 × -10 $x^2 - 22x + 120 = x^2 - 12x - 10x + 120$ = x (x - 12) - 10 (x - 12)= (x - 10) (x - 12)7. $x^2 - 11x - 42$ Solution: We have, $x^2 - 11x - 42$ By considering, p+q = -11 and pq = -42So we can replace -11x by 3x -14x -42 by 3 × -14 $x^2 - 11x - 42 = x^2 + 3x - 14x - 42$ = x (x + 3) - 14 (x + 3)= (x - 14) (x + 3)8. a² + 2a – 3 Solution: We have, a² + 2a – 3 By considering, p+q = 2 and pq = -3So we can replace 2a by 3a -a -3 by 3 × -1 $a^{2} + 2a - 3 = a^{2} + 3a - a - 3$ = a (a + 3) - 1 (a + 3)= (a - 1) (a + 3)9. a² + 14a + 48

Solution:

We have, a² + 14a + 48 By considering, p+q = 14 and pq = 48So we can replace 14a by 8a + 6a 48 by 8 × 6 a^{2} + 14a + 48 = a^{2} + 8a + 6a + 48 = a (a + 8) + 6 (a + 8)= (a + 6) (a + 8)10. $x^2 - 4x - 21$ Solution: We have, $x^2 - 4x - 21$ By considering, p+q = -4 and pq = -21So we can replace -4x by 3x - 7x-21 by 3 × -7 $x^{2} + 4x - 21 = x^{2} + 3x - 7x - 21$ = x (x + 3) - 7 (x + 3)= (x - 7) (x + 3)11. y² + 5y - 36 Solution: We have, $y^2 + 5y - 36$ By considering, p+q = 5 and pq = -36So we can replace 5y by 9y – 4y -36 by 9 × -4 $y^2 + 5y - 36 = y^2 + 9y - 4y - 36$ = y (y + 9) - 4 (y + 9)= (y - 4) (y + 9)12. (a² - 5a)² - 36 Solution: We have, $(a^2 - 5a)^2 - 36$

 $(a^2 - 5a)^2 - 6^2$ By using the formula $(a^2 - b^2) = (a+b) (a-b)$ $(a^2 - 5a)^2 - 6^2 = (a^2 - 5a + 6)(a^2 - 5a - 6)$ So now we shall factorize the expression $(a^2 - 5a + 6)$ By considering, p+q = -5 and pq = 6So we can replace -5a by a -6a 6 by 1 × -6 $a^2 - 5a - 6 = a^2 + a - 6a - 6$ = a (a + 1) - 6(a + 1)= (a - 6) (a + 1)So now we shall factorize the expression $(a^2 - 5a + 6)$ By considering, p+q = -5 and pq = -6So we can replace -5a by -2a -3a $6 \text{ by } -2 \times -3$ $a^2 - 5a + 6 = a^2 - 2a - 3a + 6$ = a (a - 2) - 3 (a - 2)= (a – 3) (a – 2) $\therefore (a^2 - 5a)^2 - 36 = (a^2 - 5a + 6) (a^2 - 5a - 6)$ = (a + 1) (a - 6) (a - 2) (a - 3)13. (a + 7) (a – 10) + 16 Solution: We have, (a + 7) (a - 10) + 16 $a^2 - 10a + 7a - 70 + 16$ a² – 3a – 54 By considering, p+q = -3 and pq = -54So we can replace -3a by 6a - 9a -54 by 6 × -9 $a^2 - 3a - 54 = a^2 + 6a - 9a - 54$ = a (a + 6) - 9 (a + 6)= (a - 9) (a + 6)

EXERCISE 7.8 PAGE NO: 7.30

Resolve each of the following quadratic trinomials into factors:

1. 2x² + 5x + 3

Solution:

We have,

 $2x^2 + 5x + 3$

The coefficient of x² is 2

The coefficient of x is 5

Constant term is 3

We shall split up the center term i.e., 5 into two parts such that their sum p+q is 5 and product $pq = 2 \times 3$ is 6

So, we express the middle term 5x as 2x + 3x

$$2x^2 + 5x + 3 = 2x^2 + 2x + 3x + 3$$

$$= 2x(x + 1) + 3(x + 1)$$

= (2x + 3) (x + 1)

2. $2x^2 - 3x - 2$

Solution:

We have,

 $2x^2 - 3x - 2$

The coefficient of x² is 2

The coefficient of x is -3

Constant term is -2

So, we express the middle term -3x as -4x + x

$$2x^2 - 3x - 2 = 2x^2 - 4x + x - 2$$

$$= 2x (x - 2) + 1 (x - 2)$$

= (x - 2) (2x + 1)

Solution:

We have,

 $3x^2 + 10x + 3$

The coefficient of x² is 3

The coefficient of x is 10

Constant term is 3

So, we express the middle term 10x as 9x + x $3x^2 + 10x + 3 = 3x^2 + 9x + x + 3$ = 3x(x + 3) + 1(x + 3)= (3x + 1)(x + 3)4. $7x - 6 - 2x^2$ Solution: We have, $7x - 6 - 2x^2$ $-2x^{2}+7x-6$ $2x^2 - 7x + 6$ The coefficient of x² is 2 The coefficient of x is -7 Constant term is 6 So, we express the middle term -7x as -4x - 3x $2x^2 - 7x + 6 = 2x^2 - 4x - 3x + 6$ = 2x(x-2) - 3(x-2)= (x - 2) (2x - 3)5. $7x^2 - 19x - 6$ Solution: We have, $7x^2 - 19x - 6$ The coefficient of x^2 is 7 The coefficient of x is -19 Constant term is -6 So, we express the middle term -19x as 2x - 21x $7x^2 - 19x - 6 = 7x^2 + 2x - 21x - 6$ = x (7x + 2) - 3 (7x + 2)= (7x + 2) (x - 3)6. $28 - 31x - 5x^2$ Solution: We have, $28 - 31x - 5x^2$ $-5x^{2}-31x+28$

 $5x^2 + 31x - 28$ The coefficient of x^2 is 5 The coefficient of x is 31 Constant term is -28 So, we express the middle term 31x as -4x + 35x $5x^2 + 31x - 28 = 5x^2 - 4x + 35x - 28$ = x (5x - 4) + 7 (5x - 4)= (x + 7) (5x - 4)7.3 + 23y - $8y^2$ Solution: We have, $3 + 23y - 8y^2$ $-8y^{2}+23y+3$ $8y^2 - 23y - 3$ The coefficient of y² is 8 The coefficient of y is -23 Constant term is -3 So, we express the middle term -23y as -24y + y $8y^2 - 23y - 3 = 8y^2 - 24y + y - 3$ = 8y(y-3) + 1(y-3)= (8y + 1)(y - 3)8. $11x^2 - 54x + 63$ Solution: We have, $11x^2 - 54x + 63$ The coefficient of x² is 11 The coefficient of x is -54 Constant term is 63 So, we express the middle term -54x as -33x - 21x $11x^2 - 54x + 63 = 11x^2 - 33x - 21x + 63$ = 11x (x - 3) - 21 (x - 3)= (11x - 21)(x - 3)9. $7x - 6x^2 + 20$

Solution:

We have,

 $7x - 6x^2 + 20$

 $-6x^{2} + 7x + 20$

 $6x^2 - 7x - 20$

The coefficient of x² is 6

The coefficient of x is -7

Constant term is -20

So, we express the middle term -7x as -15x + 8x

 $6x^2 - 7x - 20 = 6x^2 - 15x + 8x - 20$

= 3x (2x - 5) + 4 (2x - 5)

= (3x + 4) (2x - 5)

10. 3x² + 22x + 35

Solution:

We have,

 $3x^2 + 22x + 35$

The coefficient of x² is 3

The coefficient of x is 22

Constant term is 35

So, we express the middle term 22x as 15x + 7x

 $3x^2 + 22x + 35 = 3x^2 + 15x + 7x + 35$

= 3x (x + 5) + 7 (x + 5)

= (3x + 7) (x + 5)

11. 12x² – 17xy + 6y²

Solution:

We have,

 $12x^2 - 17xy + 6y^2$

The coefficient of x² is 12

The coefficient of x is -17y

Constant term is 6y²

So, we express the middle term -17xy as -9xy - 8xy

 $12x^2 - 17xy + 6y^2 = 12x^2 - 9xy - 8xy + 6y^2$

= 3x (4x - 3y) - 2y (4x - 3y)

= (3x - 2y) (4x - 3y)12. 6x² - 5xy - 6y² Solution: We have, $6x^2 - 5xy - 6y^2$ The coefficient of x² is 6 The coefficient of x is -5y Constant term is -6y² So, we express the middle term -5xy as 4xy – 9xy $6x^2 - 5xy - 6y^2 = 6x^2 + 4xy - 9xy - 6y^2$ = 2x (3x + 2y) - 3y (3x + 2y)= (2x - 3y) (3x + 2y)13. $6x^2 - 13xy + 2y^2$ Solution: We have, $6x^2 - 13xy + 2y^2$ The coefficient of x² is 6 The coefficient of x is -13y Constant term is 2y² So, we express the middle term -13xy as -12xy - xy $6x^2 - 13xy + 2y^2 = 6x^2 - 12xy - xy + 2y^2$ = 6x (x - 2y) - y (x - 2y)= (6x - y) (x - 2y)14. $14x^2 + 11xy - 15y^2$ Solution: We have, $14x^2 + 11xy - 15y^2$ The coefficient of x² is 14 The coefficient of x is 11y Constant term is -15y² So, we express the middle term 11xy as 21xy - 10xy $14x^2 + 11xy - 15y^2 = 14x^2 + 21xy - 10xy - 15y^2$ = 2x (7x - 5y) + 3y (7x - 5y)

= (2x + 3y) (7x - 5y)15. 6a² + 17ab - 3b² Solution: We have, 6a² + 17ab - 3b² The coefficient of a² is 6 The coefficient of a is 17b Constant term is -3b² So, we express the middle term 17ab as 18ab – ab 6a² +17ab- 3b² = 6a² + 18ab - ab - 3b² = 6a (a + 3b) - b (a + 3b)= (6a - b) (a + 3b)16. 36a² + 12abc - 15b²c² Solution: We have, 36a² + 12abc - 15b²c² The coefficient of a² is 36 The coefficient of a is 12bc Constant term is -15b²c² So, we express the middle term 12abc as 30abc - 18abc 36a² –12abc– 15b²c² = 36a² + 30abc – 18abc – 15b²c² = 6a (6a + 5bc) - 3bc (6a + 5bc)= (6a + 5bc) (6a - 3bc)= (6a + 5bc) 3(2a - bc)17. 15x² – 16xyz – 15y²z² Solution: We have, $15x^2 - 16xyz - 15y^2z^2$ The coefficient of x² is 15 The coefficient of x is -16yz Constant term is -15y²z² So, we express the middle term -16xyz as -25xyz + 9xyz $15x^2 - 16xyz - 15y^2z^2 = 15x^2 - 25yz + 9yz - 15y^2z^2$

= 5x (3x - 5yz) + 3yz (3x - 5yz)= (5x + 3yz) (3x - 5yz)18. $(x - 2y)^2 - 5(x - 2y) + 6$ Solution: We have. $(x - 2y)^2 - 5(x - 2y) + 6$ The coefficient of (x-2y)² is 1 The coefficient of (x-2y) is -5 Constant term is 6 So, we express the middle term -5(x - 2y) as -2(x - 2y) - 3(x - 2y) $(x-2y)^2 - 5(x-2y) + 6 = (x-2y)^2 - 2(x-2y) - 3(x-2y) + 6$ = (x - 2y - 2) (x - 2y - 3)19. $(2a - b)^2 + 2(2a - b) - 8$ Solution: We have, $(2a - b)^2 + 2(2a - b) - 8$ The coefficient of (2a-b)² is 1 The coefficient of (2a-b) is 2 Constant term is -8 So, we express the middle term 2(2a - b) as 4(2a - b) - 2(2a - b) $(2a - b)^{2} + 2(2a - b) - 8 = (2a - b)^{2} + 4(2a - b) - 2(2a - b) - 8$ = (2a - b) (2a - b + 4) - 2 (2a - b + 4)= (2a - b + 4) (2a - b - 2)

EXERCISE 7.9 PAGE NO: 7.32

Factorize each of the following quadratic polynomials by using the method of completing the square:

1. p² + 6p + 8

Solution:

We have,

p² + 6p + 8

Coefficient of p² is unity. So, we add and subtract square of half of coefficient of p.

 $p^{2} + 6p + 8 = p^{2} + 6p + 3^{2} - 3^{2} + 8 \text{ (Adding and subtracting 3^{2})}$ $= (p + 3)^{2} - 1^{2} \text{ (By completing the square)}$ By using the formula $(a^{2} - b^{2}) = (a+b) (a-b)$ = (p + 3 - 1) (p + 3 + 1) = (p + 2) (p + 4)2. $q^{2} - 10q + 21$ Solution: We have, $q^{2} - 10q + 21$ Coefficient of q^{2} is unity. So, we add and subtract square of half of coefficient of q. $q^{2} - 10q + 21 = q^{2} - 10q + 5^{2} - 5^{2} + 21 \text{ (Adding and subtracting 5^{2})}$ $= (q - 5)^{2} - 2^{2} \text{ (By completing the square)}$ By using the formula $(a^{2} - b^{2}) = (a+b) (a-b)$ = (q - 5 - 2) (q - 5 + 2) = (q - 3) (q - 7)

3. 4y² + 12y + 5

Solution:

We have,

4y² + 12y + 5

 $4(y^2 + 3y + 5/4)$

Coefficient of y² is unity. So, we add and subtract square of half of coefficient of y.

 $4(y^2 + 3y + 5/4) = 4[y^2 + 3y + (3/2)^2 - (3/2)^2 + 5/4]$ (Adding and subtracting $(3/2)^2$)

```
= 4 [(y + 3/2)^2 - 1^2] (Completing the square)
```

```
By using the formula (a^2 - b^2) = (a+b) (a-b)
```

= 4 (y + 3/2 + 1) (y + 3/2 - 1)

= 4 (y + 1/2) (y + 5/2) (by taking LCM)

$$= 4 [(2y + 1)/2] [(2y + 5)/2]$$

$$= (2y + 1) (2y + 5)$$

4. p² + 6p - 16

Solution:

We have,

p² + 6p - 16

Coefficient of p² is unity. So, we add and subtract square of half of coefficient of p.

 $p^{2} + 6p - 16 = p^{2} + 6p + 3^{2} - 3^{2} - 16$ (Adding and subtracting 3^{2}) = $(p + 3)^{2} - 5^{2}$ (Completing the square) By using the formula $(a^{2} - b^{2}) = (a+b)$ (a-b) = (p + 3 + 5) (p + 3 - 5)= (p + 8) (p - 2)5. $x^{2} + 12x + 20$

Solution:

We have,

x² + 12x + 20

Coefficient of x² is unity. So, we add and subtract square of half of coefficient of x.

 x^{2} + 12x + 20 = x^{2} + 12x + 6^{2} - 6^{2} + 20 (Adding and subtracting 6^{2})

= $(x + 6)^2 - 4^2$ (Completing the square)

By using the formula $(a^2 - b^2) = (a+b) (a-b)$

= (x + 6 + 4) (x + 6 - 4)

= (x + 2) (x + 10)

6. a² – 14a – 51

Solution:

We have,

a² - 14a - 51

Coefficient of a² is unity. So, we add and subtract square of half of coefficient of a.

 $a^{2} - 14a - 51 = a^{2} - 14a + 7^{2} - 7^{2} - 51$ (Adding and subtracting 7²)

= $(a - 7)^2 - 10^2$ (Completing the square)

By using the formula $(a^2 - b^2) = (a+b) (a-b)$

= (a – 7 + 10) (9 – 7 – 10)

= (a – 17) (a + 3)

7. a² + 2a – 3

Solution:

We have,

a² + 2a – 3

Coefficient of a² is unity. So, we add and subtract square of half of coefficient of a.

 $a^{2} + 2a - 3 = a^{2} + 2a + 1^{2} - 1^{2} - 3$ (Adding and subtracting 1²)

= $(a + 1)^2 - 2^2$ (Completing the square)

By using the formula $(a^2 - b^2) = (a+b) (a-b)$

= (a + 1 + 2) (a + 1 - 2)= (a + 3) (a – 1) 8. $4x^2 - 12x + 5$ Solution: We have. $4x^2 - 12x + 5$ $4(x^2 - 3x + 5/4)$ Coefficient of x² is unity. So, we add and subtract square of half of coefficient of x. $4(x^2 - 3x + 5/4) = 4[x^2 - 3x + (3/2)^2 - (3/2)^2 + 5/4]$ (Adding and subtracting $(3/2)^2$) = 4 $[(x - 3/2)^2 - 1^2]$ (Completing the square) By using the formula $(a^2 - b^2) = (a+b)(a-b)$ = 4 (x - 3/2 + 1) (x - 3/2 - 1)= 4 (x - 1/2) (x - 5/2) (by taking LCM) = 4 [(2x-1)/2] [(2x-5)/2]= (2x - 5)(2x - 1)9. y² - 7y + 12 Solution: We have, $v^2 - 7v + 12$ Coefficient of y² is unity. So, we add and subtract square of half of coefficient of y. $y^2 - 7y + 12 = y^2 - 7y + (7/2)^2 - (7/2)^2 + 12$ [Adding and subtracting $(7/2)^2$] = $(y - 7/2)^2 - (7/2)^2$ (Completing the square) By using the formula $(a^2 - b^2) = (a+b) (a-b)$ = (y - (7/2 - 1/2)) (y - (7/2 + 1/2))= (y - 3) (y - 4)10. z² – 4z – 12 Solution: We have, $z^2 - 4z - 12$ Coefficient of z² is unity. So, we add and subtract square of half of coefficient of z. $z^2 - 4z - 12 = z^2 - 4z + 2^2 - 2^2 - 12$ [Adding and subtracting 2²] = $(z - 2)^2 - 4^2$ (Completing the square)

By using the formula $(a^2 - b^2) = (a+b) (a-b)$

= (z - 2 + 4) (z - 2 - 4)= (z - 6) (z + 2)