

Research Article

Improvement of D-Ribose Production from Corn Starch Hydrolysate by a Transketolase-Deficient Strain *Bacillus subtilis* UJS0717

Zhuan Wei,¹ Jue Zhou,² WenJing Sun,² FengJie Cui,²
QinHua Xu,³ and ChangFeng Liu³

¹Hebei Chemical and Pharmaceutical College, Shijiazhuang 050026, China

²School of Food and Biological Engineering, Jiangsu University, Zhenjiang 212013, China

³Shandong Depu Chemical Technology Co., Ltd., Tai'an 271200, China

Correspondence should be addressed to WenJing Sun; sunwenjing1919@163.com and FengJie Cui; fengjiecui@163.com

Received 16 September 2015; Accepted 15 November 2015

Academic Editor: Denise Freire

Copyright © 2015 Zhuan Wei et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

D-Ribose is a five-carbon sugar and generally used as an energy source to improve athletic performance and the ability. The culture conditions for maximum D-ribose production performance from cheap raw material corn starch hydrolysate were improved by using one-factor-at-a-time experiments and a three-level Box-Behnken factorial design. The optimal fermentation parameters were obtained as 36°C culture temperature, 10% inoculum volume, and 7.0 initial pH. The mathematical model was then developed to show the effect of each medium composition and their interactions on the production of D-ribose and estimated that the optimized D-ribose production performance with the concentration of 62.13 g/L, yield of 0.40 g/g, and volumetric productivity of 0.86 g/L·h could be obtained when the medium compositions were set as 157 g/L glucose, 21 g/L corn steep liquor, 3.2 g/L (NH₄)₂SO₄, 1 g/L yeast extract, 0.05 g/L MnSO₄·H₂O, and 20 g/L CaCO₃. These findings indicated the D-ribose production performance was significantly improved compared to that under original conditions.

1. Introduction

D-Ribose (C₅H₁₀O₅) is a functional five-carbon sugar and plays the important role in life as a ribosyl residue for ATP, RNA, NAD, NADP, FAD, and coenzyme A. D-Ribose has been used to improve athletic performance/ability as an energy source [1] and produce riboflavin (vitamin B₂), animal feed additives, cosmetics and foods [2], and antiviral and anticancer drugs [3].

Two methods including yeast RNA hydrolysis and chemical synthesis from gluconic acids, glucose, arabinose, and xylose are previously used for preparing D-ribose [4, 5]. For example, D-ribose with a yield of 60–94% was produced by epimerizing D-arabinose in the presence of molybdenic and boric acids [4]. However, chemical synthesis processes for D-ribose production suffered from significant disadvantages such as low yield, complex scheme, and recovering/purifying

burdens. Currently, almost all the 2000–3000 tonne of D-ribose produced annually worldwide are obtained by microbial fermentation due to high selectivity, high rate, and high yield of conversion [6]. The microorganisms from genera *Bacillus* are the main D-ribose producers including *B. subtilis* and *B. pumilus*. Most of those strains, however, restricted their usefulness for commercial production due to certain disadvantages such as long fermentation time and lower ribose concentration and productivity. For example, about 40 g/L of D-ribose was produced from 200 g/L of glucose by *Bacillus subtilis* ATCC 21951 after 7-d fermentation [7]. Generally, only *Bacillus* strains deficient in the transketolase and/or D-ribulose-5-phosphate-3-epimerase have the ability to accumulate ribose in the fermented broth since these enzymes will further catalyze the produced ribose to the aromatic amino acids [6]. Our group has focused on the D-ribose

fermentation since 2000 and screened several transketolase-deficient *Bacillus* strains with industrial application potency [8–10]. One of the strains, *Bacillus subtilis* B941, was screened to produce 41.8 g/L of D-ribose from 180 g/L of glucose. After mutation with UV irradiation, the mutant strain *Bacillus subtilis* Buvp-24 produced the maximum D-ribose concentration of 55 g/L [9].

Starchy biomass is a promising feedstock for chemical bioproduction due to its abundant availability. Commercial scale for converting starchy biomass to chemicals such as ethanol, lactic acid, and 2-keto-gluconic acid has been realized [11]. Corn starch is an abundant inexpensive renewable resource and larger output in China and the United States [12, 13] while to date the fermentative production of D-ribose from corn starch has not been yet reported.

Statistical techniques have been used for optimizing the culture media to produce microbial metabolites [14–16]. Response surface methodology (RSM), as a collection of statistical techniques for experiment designing, model developing, factors evaluating, and optimum conditions searching, has been extensively applied in optimization of medium composition, conditions of enzymatic hydrolysis, fermentation, and food manufacturing processes [17, 18]. However, very few references with application of statistical techniques to maximize D-ribose production are available. Hence, the objective of the present study was to optimize the process parameters and medium compositions to increase the D-ribose production performance by a transketolase-deficient strain *Bacillus subtilis* UJS0717 using one-factor-at-a-time experiments followed by a three-level Box-Behnken factorial design combining with response surface methodology (RSM).

2. Materials and Methods

2.1. Bacterial Strain and Media. The D-ribose producer *B. subtilis* UJS0717 was a mutant from Industrial Microbiology Laboratory in Shanxi Institute of Biology and maintained at 4°C in ampoule tube and kept in our laboratory. Stock medium contained 5 g/L of D-sorbitol, 10 g/L of peptone, 2 g/L of NaCl, 2 g of yeast extract, and 20 g/L of agar. A loopful of the stock culture was diluted with sterilized water and inoculated into 20 mL of seed medium containing glucose 20 g/L, yeast extract 3 g/L, K₂HPO₄ 3 g/L, and KH₂PO₄ 1 g/L, in a 250 mL Erlenmeyer flask, incubated at 36°C, 240 rpm, for 20 h, and used as a seed culture.

Corn starch hydrolysate (CSH) was obtained from Shandong Depu Chemical Technology Co., Ltd. (Taian, Shandong, China), produced by liquefaction and saccharification processes with amylases and glucoamylase. The corn starch hydrolysate contains approximately 30% (w/v) of glucose and 0.6% of protein (total nitrogen × 6.38). CSH was diluted with deionized water to obtain various concentrations of glucose. Other analytical chemicals were obtained from Sigma Aldrich (St. Louis, MO, USA). The basic fermentation medium contained glucose 120 g/L, corn steep liquor 15 g/L, (NH₄)₂SO₄ 7.5 g/L, yeast extract 1 g/L, and MnSO₄·H₂O

0.05 g/L. Glucose and other nutrients were sterilized separately at 121°C for 20 min. 20.0 g/L of CaCO₃ was added to the media for balancing the broth pH.

2.2. One-Factor-at-a-Time Experiments. For one-factor-at-a-time experiments investigating effect of glucose, corn steep liquor, and (NH₄)₂SO₄ concentration on D-ribose production, the glucose, corn steep liquor, and (NH₄)₂SO₄ concentrations in the basic fermentation medium were adjusted ranging from 100 g/L to 210 g/L, 5 g/L to 25 g/L, and 1.5 g/L to 9 g/L, respectively. Four fermentation temperatures from 30°C to 39°C, four inoculum volumes from 5% to 20% (v/v), and five initial pH values from 6.0 to 8.0 were used for selecting the optimal temperature, inoculum volume, and initial pH using one-factor-at-a-time experimental design. The fermentation time was 72 hours if not further specified.

2.3. Optimization of Fermentation Medium Compositions. To find the interactions and give the precise levels of media compositions including glucose, corn steep liquor, and (NH₄)₂SO₄ significantly influencing D-ribose production, Box-Behnken design for 3 variables at three levels (+1, 0, and -1) was further employed to optimize their concentrations to maximize the D-ribose production by *B. subtilis* UJS0717 based on the one-factor-at-a-time experiments [18]. The statistical matrix included 15 runs of experiments for fitting a second-order response surface. Table 5 gives the variables, their values, and the experimental design, respectively.

A mathematical model, describing the relationships between the process indices (D-ribose concentration) and the medium component contents in second-order equation, was developed with the least squares method as follows:

$$Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \beta_{11} X_1^2 + \beta_{22} X_2^2 + \beta_{33} X_3^2 + \beta_{12} X_1 X_2 + \beta_{13} X_1 X_3 + \beta_{23} X_2 X_3, \quad (1)$$

where Y is the measured response; β_0 model constant; X_1 , X_2 , and X_3 are independent variables; β_1 , β_2 , and β_3 are linear coefficients; β_{12} , β_{13} , and β_{23} are cross product coefficients; and β_{11} , β_{22} , and β_{33} are the quadratic coefficients [19]. The accuracy and general ability of the above polynomial model could be evaluated by the coefficient of determination R^2 .

2.4. Analytical Methods. The cell growth was represented with dry cell weight (DCW, g/L) by neutralizing the residual CaCO₃ with 1 M HCl solution, centrifuging at 10000 ×g for 5 min to obtain *B. subtilis* cells, and drying the cells at 80°C to the constant weight.

Glucose and D-ribose were measured using high performance liquid chromatography (Agilent 1100 series, MN, USA) equipped with a SUGAR SH1011 column (Shodex, 8.0 mm ID × 300 mm) and a differential refractometer (Agilent 1100 series). The mobile phase was 0.005 M H₂SO₄ at a flow rate of 0.6 mL/min. The column temperature was maintained at 50°C. About 3 mL samples were taken and filtered with Whatman 0.45 μm syringe filter to obtain about 1 mL permeate for HPLC analysis.

The performance of D-ribose production was evaluated based on D-ribose concentration, D-ribose productivity,

glucose conversion ratio, and D-ribose yield. D-Ribose productivity was defined as the amount of D-ribose produced per hour liter. D-Ribose yield was calculated by dividing the amount of D-ribose produced by the amount of glucose consumed. All fermentation tests were run in duplicate. Data analysis including analysis of variance was conducted using the SAS System (SAS Institute, Cary, NC, USA).

3. Results and Discussion

3.1. Effect of Temperature. In order to understand the influence of temperature on the production of D-ribose, fermentation with an initial concentration of 10% (v/v), pH 7.0, and basic fermentation medium was conducted at four temperatures ranging from 30°C to 39°C (Figure 1). D-Ribose productivity was significantly affected by the temperature ($P < 0.05$). The maximum D-ribose productivity of 0.50 g/L-h, glucose utilization ratio of 97.05%, and highest cell concentration of 8.70 g/L were obtained at 36°C after 72 h of fermentation. D-Ribose productivity of 0.44 g/L-h and 0.48 g/L-h and glucose utilization ratio of 90.49% and 93.75% were obtained at 33°C and 39°C, respectively. At 30°C, the ribose productivity and glucose utilization ratio decreased to 0.42 g/L-h and 89.04%, respectively. Kishimoto et al. (1990) found that temperature about 37°C was suitable for D-ribose production by *B. pumilus* NO.716 [23]. Therefore, 36°C appeared to be the optimal temperature for D-ribose production from glucose by *B. subtilis* UJS0717.

3.2. Effect of Inoculum Volume. The influence of inoculum volume, ranging from 5 to 20% (v/v), on D-ribose production with conditions of 36°C, pH 7.0, and basic fermentation medium was investigated. As shown in Table 1, D-ribose productivity increased substantially from 0.43 g/L-h to 0.50 g/L-h when inoculum volume increased from 5% to 10% (v/v) ($P < 0.05$). Further increases in inoculum volume (beyond 10%) had no significant effect on the D-ribose production ($P > 0.05$). Similar results were also observed by Ren et al. that high inoculum volume resulted in a negative impact on D-ribose production by *B. subtilis* ptn15-1 [29]. For the present study, an inoculum volume of 10% (v/v) was selected.

3.3. Effect of Initial pH. The effect of initial pH, ranging from 6.0 to 8.0, on the performance of D-ribose production by *B. subtilis* UJS0717 was investigated at 36°C, inoculum volume of 10% (v/v) with basic fermentation medium. As shown in Table 2, D-ribose production and cell growth were influenced by the initial pH. D-Ribose productivity increased substantially from 0.43 to 0.50 and glucose utilization ratio increased visibly from 90.78% to 96.31% when initial pH adjusted from 6.0 to 7.0 ($P < 0.05$). Further increases in initial pH (beyond 7.0) had a negative impact on D-ribose production. Park and Seo also found that initial pH about 7.0 was suitable for D-ribose production by *B. subtilis* JY1 [30]. Therefore, pH 7.0 appeared to be optimal for D-ribose production from glucose by *B. subtilis* UJS0717.

3.4. Effect of Glucose Concentration. The effect of glucose concentration ranging from 100 g/L to 210 g/L on the performance of D-ribose production by *B. subtilis* UJS0717 was investigated at 36°C, pH 7.0, and inoculum volume of 10% (v/v). As shown in Figures 2(a) and 2(b), D-ribose concentration increased from 25.81 g/L to 48.01 g/L and glucose concentration decreased gradually from 50.01 g/L to 6.65 g/L with the increase of fermentation time from 48 h to 72 h at glucose concentration of 150 g/L. Within the 48 h fermentation, 100 g/L of glucose was consumed completely. The maximum D-ribose concentration of 48.15 g/L appeared at the initial glucose concentration of 150 g/L. Too high glucose concentrations seemed to possess the inhibition on the glucose consumption and D-ribose production. D-Ribose concentration kept the lower level below 30 g/L and approximately 72% of glucose was consumed during 96 h fermentation when the initial glucose concentration was set as 210 g/L.

Similar trends of cell growth could be observed (Figure 2(c)). *B. subtilis* cells were in the exponential phase with the lower D-ribose production in the 48 h of fermentation and then entered the stationary phase at the 48–96 h with the higher D-ribose production. Glucose concentration of 150 g/L benefited the cell growth and reached maximum concentration of 10.59 g/L.

Figures 2(e) and 2(f) showed the D-ribose yield and productivity of strain *B. subtilis* UJS0717 during the overall fermentation process. *B. subtilis* UJS0717 gave the total D-ribose yield of 0.32 g/g with the medium composed of 150 g/L glucose, followed by 0.30 g/g with 120 g/L and 0.24 g/g with 180 g/L of glucose. High concentrations of glucose (over 180 g/L) had the lower D-ribose yield of 0.17 g/g. The highest volumetric productivity (0.67 g/L-h) was reached at 72 h with the 150 g/L of glucose. With 100, 120, 180, and 210 g/L of glucose, their productivity reached 0.37 g/L-h, 0.49 g/L-h, 0.57 g/L-h, and 0.35 g/L-h at 48 h, 48 h, 72 h, and 72 h, respectively. Therefore, glucose concentration of about 150 g/L appeared to be optimal for D-ribose production by *B. subtilis* UJS0717.

3.5. Effect of Corn Steep Liquor Concentration. Corn steep liquor is a major byproduct of the corn wet milling industry and is a low-cost nutrient source available on a large scale [31]. It is the cost effective medium composition due to its high content of nitrogen, water soluble vitamins, amino acids, minerals, and other growth factors [32]. Corn steep liquor as the essential microbial nutrient has been used for production of organic acids, solvents, and enzymes. Previous report also proved that corn steep liquor was an efficient nitrogen source and growth factor for industrial D-ribose fermentation [9, 33]. Herein, to select the optimal corn steep liquor concentration for D-ribose production, five concentrations ranging from 5 g/L to 25 g/L were used. Other culture conditions were 36°C, pH 7.0, inoculum volume of 10% (v/v), and glucose concentration of about 150 g/L. The D-ribose production performances were concluded in Table 3. After 72 h fermentation, cell concentrations increased from 5.71 g/L to 10.71 g/L with the increase of corn steep liquor concentration from 5 g/L to 20 g/L. Lower corn steep liquor




FIGURE 1: Effect of temperature on D-ribose production performance of *B. subtilis* UJS0717 (fermentation time: 72 h; initial pH 7.0; inoculum volume: 10%, v/v; * $P < 0.05$ compared to 30 °C group).

TABLE 1: Effect of inoculum volume on D-ribose production performance by *B. subtilis* UJS0717.

Inoculum volume (v/v)	Initial glucose (g/L)	Residual glucose (g/L)	Glucose consumption rate (%)	Glucose consumption rate (g/L-h)	Cell concentration (g/L)	D-Ribose (g/L)	D-Ribose yield (g/g)	D-Ribose productivity (g/L-h)
5%	122.0	12.75 ± 0.14	89.55 ± 0.11	1.52 ± 0.00	7.79 ± 0.01	31.23 ± 0.35	0.29 ± 0.00	0.43 ± 0.00
10%	122.0	4.50 ± 0.35*	96.31 ± 0.29*	1.63 ± 0.00*	8.70 ± 0.32*	36.00 ± 0.75*	0.31 ± 0.00*	0.50 ± 0.00*
15%	122.0	6.53 ± 0.58*	94.65 ± 0.47*	1.60 ± 0.00*	9.30 ± 0.20*	34.73 ± 0.66*	0.30 ± 0.00*	0.48 ± 0.00*
20%	122.0	6.51 ± 0.67*	94.66 ± 0.55*	1.60 ± 0.00*	9.37 ± 0.10*	34.91 ± 0.58*	0.30 ± 0.00*	0.49 ± 0.00*

* P < 0.05 compared to 5% inoculum volume (v/v) group.

TABLE 2: Effect of the initial pH on D-ribose production performance by *B. subtilis* UJS0717.

Initial pH	Initial glucose (g/L)	Residual glucose (g/L)	Glucose consumption rate (%)	Glucose consumption rate (g/L.h)	Cell concentration (g/L)	D-Ribose (g/L)	D-Ribose yield (g/g)	D-Ribose productivity (g/L.h)
6.0	122.0	11.25 ± 0.12	90.78 ± 0.10	1.54 ± 0.00	7.95 ± 0.15	30.92 ± 0.47	0.28 ± 0.00	0.43 ± 0.00
6.5	122.0	8.95 ± 0.05*	92.66 ± 0.04*	1.57 ± 0.00*	8.60 ± 0.25*	31.35 ± 0.35	0.28 ± 0.00	0.43 ± 0.00
7.0	122.0	4.50 ± 0.35*	96.31 ± 0.29*	1.63 ± 0.00*	8.70 ± 0.32*	36.00 ± 0.75*	0.31 ± 0.00*	0.50 ± 0.00*
7.5	122.0	6.85 ± 0.18*	94.39 ± 0.15*	1.60 ± 0.00*	8.35 ± 0.35*	34.57 ± 0.55*	0.30 ± 0.00*	0.48 ± 0.01*
8.0	122.0	8.55 ± 0.14*	92.99 ± 0.11*	1.57 ± 0.00*	8.15 ± 0.25*	31.76 ± 0.68	0.28 ± 0.00	0.44 ± 0.00*

* $P < 0.05$ compared to pH 6.0 group.




FIGURE 2: Effect of glucose concentration on D-ribose production performance of *B. subtilis* UJS0717 (corn steep liquor: 15 g/L; $(\text{NH}_4)_2\text{SO}_4$: 7.5 g/L).

TABLE 3: Effect of corn steep liquor concentration on D-ribose production performance by *B. subtilis* UJS0717.

Corn steep liquor concentration (g/L)	Initial glucose (g/L)	Residual glucose (g/L)	Glucose consumption rate (%)	Glucose consumption rate (g/L-h)	Cell concentration (g/L)	D-Ribose (g/L)	D-Ribose yield (g/g)	D-Ribose productivity (g/L-h)
5	152.0	28.00 ± 1.14	81.58 ± 0.93	1.72 ± 0.02	5.71 ± 0.11	28.89 ± 0.94	0.23 ± 0.01	0.40 ± 0.00
10	152.0	18.00 ± 0.14*	88.16 ± 0.09*	1.86 ± 0.00*	7.41 ± 0.10*	34.15 ± 1.54*	0.25 ± 0.01*	0.47 ± 0.00*
15	152.0	6.40 ± 0.06*	95.79 ± 0.04*	2.02 ± 0.00*	10.16 ± 0.16*	48.25 ± 0.24*	0.33 ± 0.00*	0.67 ± 0.00*
20	152.0	0.13 ± 0.04*	99.91 ± 0.03*	2.11 ± 0.00*	10.71 ± 0.21*	54.02 ± 0.89*	0.36 ± 0.00*	0.75 ± 0.00*
25	152.0	0.18 ± 0.04*	99.88 ± 0.03*	2.11 ± 0.00*	10.21 ± 0.09*	48.84 ± 0.26*	0.32 ± 0.00*	0.68 ± 0.00*

* P < 0.05 compared to corn steep liquor concentration of 5g/L group.

concentrations (<15 g/L) resulted in the high residual glucose with the concentration of over 18 g/L and lower D-ribose production of about 28 g/L. With the increase of corn steep liquor concentration to 20 g/L, D-ribose production reached the maximum level of 54.02 g/L with the highest productivity of 0.75 g/L·h and yield of 0.36 g/g. Too high levels of corn steep liquor concentration (25 g/L) seemed to be negative for the D-ribose production (48.84 g/L) and cell growth (10.21 g/L). Therefore, concentration of corn steep liquor of 20 g/L appeared to be optimal for D-ribose production.

3.6. Effect of $(\text{NH}_4)_2\text{SO}_4$ Concentration. Nitrogen substrates such as $(\text{NH}_4)_2\text{SO}_4$ have been shown to be useful for large-scale D-ribose production [25]. In order to select the optimal $(\text{NH}_4)_2\text{SO}_4$ concentration on the production of D-ribose, fermentation with 36°C, pH 7.0, inoculum volume of 10% (v/v), glucose concentration of about 150 g/L, and corn steep liquor of 20 g/L was conducted at six $(\text{NH}_4)_2\text{SO}_4$ concentrations ranging from 1.5 g/L to 9.0 g/L (Table 4).

D-Ribose productivity was significantly affected by the $(\text{NH}_4)_2\text{SO}_4$ concentrations ($P < 0.05$). The maximum D-ribose productivity and yield of 0.84 g/L·h and 0.40 g/g were obtained at $(\text{NH}_4)_2\text{SO}_4$ concentration of 3.0 g/L. After 72 h fermentation, cell concentrations increased from 9.40 g/L to 10.77 g/L with the increase of $(\text{NH}_4)_2\text{SO}_4$ concentration from 1.5 g/L to 3.0 g/L. Higher $(\text{NH}_4)_2\text{SO}_4$ concentrations (>4.5 g/L) resulted in the residual glucose. With $(\text{NH}_4)_2\text{SO}_4$ concentration of 6.0 g/L, D-ribose production reached the minimum level of 42.89 g/L with the lowest productivity of 0.60 g/L·h and yield of 0.29 g/g. Too high levels of $(\text{NH}_4)_2\text{SO}_4$ concentration (>3.0 g/L) seemed not to benefit the D-ribose production and cell growth, while Srivastava and Wangikar found that 5.0 g/L of $(\text{NH}_4)_2\text{SO}_4$ was optimum for D-ribose yield and too high concentrations of $(\text{NH}_4)_2\text{SO}_4$ resulted in lower quantities of D-ribose and large quantities of acetic acid and acetoin of 20 g/L and 30 g/L, respectively [34]. Therefore, in our study, concentration of $(\text{NH}_4)_2\text{SO}_4$ of 3.0 g/L appeared to be optimal for D-ribose production from glucose by *B. subtilis* UJS0717.

Based on one-factor-at-one-time experimental results, it could be concluded that the optimal culture conditions for D-ribose production from glucose by *B. subtilis* UJS0717 were 36°C, inoculum volume of 10% (v/v), pH 7.0, glucose concentration of 150 g/L, corn steep liquor concentration of 20 g/L, and $(\text{NH}_4)_2\text{SO}_4$ concentration of 3.0 g/L. However, one-factor-at-a-time experiments are incapable of reaching the true optimum especially due to interactions among various factors while RSM statistically designs and builds models, evaluates the effects of factors, and searches optimum conditions of factors for the desirable responses. Hence, following response surface methodology combined Box-Behnken design was applied to find the precise levels and interactions among significant factors.

3.7. Optimization of Fermentation Media Using Box-Behnken Design (BBD). In this work, the actual levels of the variables for each of the experiments in the design matrix were calculated and experimental results obtained as given in

Table 5. Table 6 shows the results of the statistical analysis. F value of 83.91 and low P value ($P < 0.01$) indicated that the model was highly significant and yielded good predictions of the experimental results. The value of the coefficient of determination ($R^2 = 0.9934$) also reflected the good fit of the response model [35], which showed that 99.34% of the sample variation in the experiments was explained by the independent variables. The coefficient of variation ($\text{CV} = 2.02\% < 10\%$) indicated the experiment was accurate and reliable. The value of 0.0556 for lack of fit implies that it is not significant comparing to the pure error and that the model equation was adequate for predicting D-ribose concentration. All the above analytical results demonstrated that the model for D-ribose concentration was appropriate in terms of the Box-Behnken design.

The following second-order polynomial equation based on the multiple regression analysis explained the relationship between variables and D-ribose concentration:

$$Y = 60.29 + 5.09X_1 + 0.93X_2 + 1.42X_3 - 11.52X_1^2 - 3.42X_2^2 - 4.00X_3^2 - 0.24X_1X_2 - 0.19X_1X_3 - 1.15X_2X_3, \quad (2)$$

where Y stands for the response variable (D-ribose concentration) and X_1 , X_2 , and X_3 are the actual values of D-glucose, corn steep liquor, and $(\text{NH}_4)_2\text{SO}_4$ concentrations, respectively. In this equation, the signs of the linear coefficients of X_1 , X_2 , and X_3 were positive. This result indicated that glucose, corn steep liquor, and $(\text{NH}_4)_2\text{SO}_4$ had a synergistic effect on the production of D-ribose. The signs of the coefficients of X_1X_2 , X_1X_3 , X_2X_3 , X_1^2 , X_2^2 , and X_3^2 were negative, indicating that they had an inverse effect on D-ribose concentration [35]. The first-order and the quadratic main effects of glucose concentration were highly significant ($P < 0.01$) according to the P value of the model. This result indicated that the factor had significant effect on the production of D-ribose. Therefore, the different ratio of the factor in fermentation would affect the production of D-ribose, and small variations in the factor would produce large changes in the results.

The three-dimensional (3D) response surface plots drawn by Design-Expert 8.0.6.1 software based on the model equations were used to explain the interactions among the variables and to determine the optimal ratios of each component for the production of D-ribose. The response surface shapes reflected the nature and range of different components, and the peaks suggested that the optimum points were within the design limits. In this experiment, each plot was generated for the interactive effects of two variables on the production of D-ribose while holding the other factor at "zero" levels. Figure 3 was the 3D-surface plot and 2D-projection.

The maximum D-ribose concentration of 61.01 g/L was obtained by solving the model regression equation with the medium composed of 157 g/L glucose, 21 g/L corn steep liquor, and 3.2 g/L $(\text{NH}_4)_2\text{SO}_4$. To confirm the predicted results and verify the model, the above-calculated critical levels of the three variables were used to produce D-ribose, and the mean value of D-ribose production was 62.13 ±

TABLE 4: Effect of $(\text{NH}_4)_2\text{SO}_4$ concentration on D-ribose production performance by *B. subtilis* UJS0717.

$(\text{NH}_4)_2\text{SO}_4$ concentration (g/L)	Initial glucose (g/L)	Residual glucose (g/L)	Glucose consumption rate (%)	Glucose consumption rate (g/L.h)	Cell concentration (g/L)	D-Ribose (g/L)	D-Ribose yield (g/g)	D-Ribose productivity (g/L.h)
1.5	151.0	3.85 ± 0.14	97.45 ± 0.09	2.04 ± 0.00	9.40 ± 0.30	45.81 ± 1.23	0.31 ± 0.01	0.64 ± 0.01
3.0	151.0	0.12 ± 0.01*	99.92 ± 0.00*	2.09 ± 0.00*	10.77 ± 0.17*	60.11 ± 0.12*	0.40 ± 0.00*	0.84 ± 0.01*
4.5	151.0	0.29 ± 0.01*	99.81 ± 0.01*	2.09 ± 0.00*	10.65 ± 0.15*	56.25 ± 0.67*	0.37 ± 0.00*	0.78 ± 0.01*
6.0	151.0	5.59 ± 0.21*	96.30 ± 0.14*	2.02 ± 0.00*	9.15 ± 0.15	42.89 ± 0.17*	0.29 ± 0.00*	0.60 ± 0.00*
7.5	151.0	0.80 ± 0.14*	99.47 ± 0.09*	2.09 ± 0.00*	10.55 ± 0.35*	54.56 ± 0.30*	0.36 ± 0.00*	0.76 ± 0.00*
9.0	151.0	5.43 ± 0.04*	96.40 ± 0.28*	2.02 ± 0.00*	9.25 ± 0.25	43.04 ± 0.64*	0.30 ± 0.00*	0.60 ± 0.00*

* $P < 0.05$ compared to $(\text{NH}_4)_2\text{SO}_4$ concentration of 1.5 g/L group.

TABLE 5: Box-Behnken design matrix along with the experimental and predicted values.

Run	Actual and coded level of variables			D-Ribose production (g/L)	
	X_1 (g/L)	X_2 (g/L)	X_3 (g/L)	Experimental	Predicted
1	150 (0)	20 (0)	3.0 (0)	60.64 ± 1.34	60.29
2	150 (0)	25 (+1)	4.5 (+1)	54.32 ± 2.12	54.25
3	180 (+1)	25 (+1)	3.0 (0)	52.09 ± 1.22	51.31
4	150 (0)	15 (-1)	1.5 (-1)	49.49 ± 2.01	49.56
5	150 (0)	20 (0)	3.0 (0)	60.20 ± 1.67	60.29
6	180 (+1)	20 (0)	4.5 (+1)	50.23 ± 1.34	51.08
7	120 (-1)	25 (+1)	3.0 (0)	40.82 ± 3.10	41.61
8	120 (-1)	20 (0)	4.5 (+1)	42.00 ± 3.12	41.28
9	180 (+1)	20 (0)	1.5 (-1)	47.92 ± 1.45	48.64
10	180 (+1)	15 (-1)	3.0 (0)	50.72 ± 1.33	49.93
11	150 (0)	20 (0)	3.0 (0)	60.04 ± 2.03	60.29
12	150 (0)	25 (+1)	1.5 (-1)	53.64 ± 1.34	53.71
13	120 (-1)	20 (0)	1.5 (-1)	38.92 ± 1.43	38.07
14	120 (-1)	15 (-1)	3.0 (0)	38.48 ± 1.98	39.26
15	150 (0)	15 (-1)	4.5 (+1)	54.75 ± 2.50	54.68

TABLE 6: Analysis of variance for the response surface quadratic model of D-ribose concentration of Box-Behnken design.

Source	Sum of squares	df	Mean square	F value	P value Prob. > F
Model	779.43	9	86.60	83.91	<0.0001**
X_1	207.47	1	207.47	201.01	<0.0001**
X_2	6.90	1	6.90	6.69	0.0491*
X_3	16.05	1	16.05	15.55	0.0109*
AB	0.24	1	0.24	0.23	0.6532
AC	0.15	1	0.15	0.14	0.7203
BC	5.24	1	5.24	5.08	0.0739
A^2	490.36	1	490.36	475.09	<0.0001**
B^2	38.80	1	38.80	37.59	0.0017**
C^2	59.13	1	59.13	57.29	0.0006**
Residual	5.16	5	1.03		
Lack of fit	4.97	3	1.66	17.15	0.0556
Pure error	0.19	2	0.097		
Cor. total	784.59	14			

Note: CV% = 2.02; $R^2 = 0.9934$; Adj. $R^2 = 0.9816$; Pred. $R^2 = 0.8981$. Note: X_1 = glucose (g/L); X_2 = corn steep liquor (g/L); X_3 = $(\text{NH}_4)_2\text{SO}_4$ (g/L).

* $P < 0.05$.

** $P < 0.01$.

1.16 g/L, which was in agreement with the predicted value (61.01 g/L).

Figure 4 compared the residual glucose concentration, pH, cell concentration, and D-ribose concentration before and after optimization. Cell concentration reached 9.05 g/L during 72 h cultivation, and relative lower D-ribose concentration of 35.95 g/L, yield of 0.31 g/g, and productivity of 0.50 g/L-h were finally obtained under the original fermentation conditions (Figure 4(a)). After optimization, the substrate glucose was utilized completely and cell concentration

was achieved at 10.99 g/L after 72 h cultivation. D-Ribose concentration, yield, and productivity reached 62.13 g/L, 0.40 g/g, and 0.86 g/L-h by *B. subtilis* UJS0717, which were higher than those under the original conditions (Figure 4(b)).

Genus *Bacillus* is the main D-ribose producer which utilizes glucose, gluconic acid, or xylose as substrate. Table 7 presented the D-ribose producing strains and their fermentation performance from this work and from literature reports. Most D-ribose producing strains in Table 7 had the capacity to produce D-ribose over 60 g/L. The strain *B. subtilis* UJS0717 used in this work is a comparable D-ribose producing bacterium that is able to produce D-ribose with the concentration of 62.13 g/L from 157 g/L glucose and high efficiency of converting glucose to D-ribose.

4. Conclusion

The transketolase-deficient strain *B. subtilis* UJS0717 produced D-ribose at level of 62.13 g/L with yield of 0.40 g/g and volumetric productivity of 0.86 g/L-h, which was therefore potentially useful as an industrial D-ribose producer from cheap raw material corn starch hydrolysate. Semicontinuous/continuous fermentation and scale-up experiments for D-ribose production are ongoing in our lab for further improving the D-ribose production performance and evaluating the technical feasibility.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Authors' Contribution

Zhuan Wei and Jue Zhou are the co-first authors and have the equal contributions.




FIGURE 3: (a) The 3D-plot and 2D-projection showing the interaction between glucose and corn steep liquor at 3.0 g/L $(NH_4)_2SO_4$ on D-ribose concentration (Y). (b) The 3D-plot and 2D-projection showing the interaction between glucose and $(NH_4)_2SO_4$ at 20 g/L corn steep liquor on D-ribose concentration (Y). (c) The 3D-plot and 2D-projection showing the interaction between corn steep liquor and $(NH_4)_2SO_4$ at 150 g/L glucose on D-ribose concentration (Y).




FIGURE 4: Comparison of D-ribose production performance of *B. subtilis* UJS0717 before and after optimization ((a) before optimization; (b) after optimization).

TABLE 7: Summarized results of D-ribose biosynthesis previously described in literature.

Strain	Culture mode	Glucose (g/L)	D-Ribose (g/L)	Fermentation time (h)	Reference
<i>B. subtilis</i> UJS0717	Batch	157	62.13	72	In the present work
<i>B. pumilus</i> ATCC 21357	Batch	125	31	55	[20]
<i>B. subtilis</i> ATCC 31092	Batch	150	67	60	[21]
<i>Bacillus</i> sp. EMP-58	Batch	140	64	55	[22]
<i>B. subtilis</i> ATCC 21951	Batch	200	95	72	[23]
<i>B. subtilis</i> IFO 1538	Batch	160	62	72	[24]
<i>B. subtilis</i> ATCC 21951	Batch	100/100 ^a	60	110	[25]
<i>B. subtilis</i> ATCC 21951	Batch	100/50 ^b	45	84	[8]
<i>B. subtilis</i> C1-B941	Batch	180	60.9	68	[9]
<i>B. subtilis</i> SPK1	Fed-batch	20/20 + 200/50 ^c	46.6	63	[2]
<i>B. subtilis</i> EC2	Batch	200	83.4	42	[26]
<i>B. subtilis</i> NJT-1507	Shake-flask	172.75	88.57	72	[27]
<i>B. subtilis</i> NJT-1507	Batch	172.75	95.27	72	[27]
<i>B. subtilis</i> XB02	Single-stage, continuous	200 ^d	68.7	160	[28]

^a100 g/L glucose plus 100 g/L D-gluconic acid.

^b100 g/L glucose plus 50 g/L D-gluconic acid.

^cAfter initial sugars of 20 g/L xylose and 20 g/L glucose were consumed completely, a sugar mixture of 200 g/L xylose and 50 g/L glucose was fed stepwise into a bioreactor.

^dInitial glucose 200 g/L, starting time 24 h, dilution rates 0.006/h, and influent glucose concentration 200 g/L.

Acknowledgments

This work was supported by funds from the Technology Research and Development Program of Shandong Province (2012YD21008), Science Foundation for Youths of Hebei Department of Education (2010289), Technology Research and Development Program of Tai'an (201340629), and 2012 Excellent Key Young Teachers Project of Jiangsu University and Research Foundation of Hebei Chemical and Pharmaceutical College (YZ201311).

References

- [1] J. E. Teitelbaum, C. Johnson, and J. St. Cyr, "The use of D-ribose in chronic fatigue syndrome and fibromyalgia: a pilot study," *Journal of Alternative and Complementary Medicine*, vol. 12, no. 9, pp. 857–862, 2006.
- [2] Y.-C. Park, S.-G. Kim, K. Park, K. H. Lee, and J.-H. Seo, "Fed-batch production of D-ribose from sugar mixtures by transketolase-deficient *Bacillus subtilis* SPK1," *Applied Microbiology and Biotechnology*, vol. 66, no. 3, pp. 297–302, 2004.
- [3] A. J. Cooper and R. G. Salomon, "Total synthesis of halichondrins: enantioselective construction of a homochiral pentacyclic C1-C15 intermediate from D-ribose," *Tetrahedron Letters*, vol. 31, no. 27, pp. 3813–3816, 1990.
- [4] N. Hiroshi, T. Takeshi, and H. Masahiko, "Method of producing solution containing D-ribose," US patent 4,602,086, 1986.
- [5] B. Lacourt-Gadras, M. Grignon-Dubois, and B. Rezzonico, "Nouvelle voie d'accès au D-ribose et au D-lyxose," *Carbohydrate Research*, vol. 235, pp. 281–288, 1992.
- [6] P. De Wulf and E. J. Vandamme, "Production of D-ribose by fermentation," *Applied Microbiology and Biotechnology*, vol. 48, no. 2, pp. 141–148, 1997.
- [7] P. De Wulf, W. Soetaert, D. Schwengers, and E. J. Vandamme, "Optimization of D-ribose production with a transketolase-affected *Bacillus subtilis* mutant strain in glucose and gluconic acid-based media," *Journal of Applied Microbiology*, vol. 83, no. 1, pp. 25–30, 1997.
- [8] H. Xie, W. J. Sun, Q. W. Yang, R. X. Gao, F. M. Zhao, and M. Z. Jiang, "Study on D-ribose fermentation," *Food and Fermentation Industries*, vol. 26, no. 2, pp. 7–10, 2000 (Chinese).
- [9] M. H. Wang, W. J. Sun, J. Q. Guo, F. M. Zhao, L. Qin, and Q. W. Yang, "Breeding of D-ribose producing mutants by protoplast mutagenesis of ultraviolet irradiation," *Industrial Microbiology*, vol. 35, no. 1, pp. 24–27, 2005 (Chinese).
- [10] M. H. Wang, W. J. Sun, F. M. Zhao, Q. W. Yang, J. Q. Guo, and L. E. Yang, "UV mutation of D-ribose high-yielding strain by sugar-resisting breeding," *Journal of Microbiology*, vol. 27, no. 1, pp. 95–98, 2007 (Chinese).
- [11] W.-J. Sun, Q.-Q. Yun, Y.-Z. Zhou et al., "Continuous 2-keto-gluconic acid (2KGA) production from corn starch hydrolysate by *Pseudomonas fluorescens* AR4," *Biochemical Engineering Journal*, vol. 77, pp. 97–102, 2013.
- [12] Y. L. Huang, Z. Wu, L. Zhang, C. M. Cheung, and S.-T. Yang, "Production of carboxylic acids from hydrolyzed corn meal by immobilized cell fermentation in a fibrous-bed bioreactor," *Bioresource Technology*, vol. 82, no. 1, pp. 51–59, 2002.
- [13] M.-L. Yuan, Z.-H. Lu, Y.-Q. Cheng, and L.-T. Li, "Effect of spontaneous fermentation on the physical properties of corn starch and rheological characteristics of corn starch noodle," *Journal of Food Engineering*, vol. 85, no. 1, pp. 12–17, 2008.
- [14] S. Djekrif-Dakhmouche, Z. Gheribi-Aoulmi, Z. Meraihi, and L. Bennamoun, "Application of a statistical design to the optimization of culture medium for α -amylase production by *Aspergillus niger* ATCC 16404 grown on orange waste powder," *Journal of Food Engineering*, vol. 73, no. 2, pp. 190–197, 2006.
- [15] J. Ren, W.-T. Lin, Y.-J. Shen, J.-F. Wang, X.-C. Luo, and M.-Q. Xie, "Optimization of fermentation media for nitrite oxidizing bacteria using sequential statistical design," *Bioresource Technology*, vol. 99, no. 17, pp. 7923–7927, 2008 (Chinese).

- [16] Y.-L. Feng, W.-Q. Li, X.-Q. Wu, J.-W. Cheng, and S.-Y. Ma, "Statistical optimization of media for mycelial growth and exopolysaccharide production by *Lentinus edodes* and a kinetic model study of two growth morphologies," *Biochemical Engineering Journal*, vol. 49, no. 1, pp. 104–112, 2010.
- [17] S. K. Psomas, M. Liakopoulou-Kyriakides, and D. A. Kyriakidis, "Optimization study of xanthan gum production using response surface methodology," *Biochemical Engineering Journal*, vol. 35, no. 3, pp. 273–280, 2007.
- [18] T. Khajvand, M. J. Chaichi, O. Nazari, and H. Golchoubian, "Application of Box–Behnken design in the optimization of catalytic behavior of a new mixed chelate of copper (II) complex in chemiluminescence reaction of luminol," *Journal of Luminescence*, vol. 131, no. 5, pp. 838–842, 2011.
- [19] C. D. Montgomery, *Design and Analysis of Experiments*, John Wiley & Sons, Singapore, 2002.
- [20] M. Yoneda and K. Sasajima, "Biotechnisches Verfahren zur Herstellung von D-Ribose," German patent DE 1904265, 1969.
- [21] K. Sasajima and M. Doi, "Verfahung zur Herstellung von D-Ribose," German patent DE 2454931 C2, 1975.
- [22] T. Asai, M. Doi, T. Kono, and H. Fukada, "Kinetic study on the production of D-ribose by *Bacillus* sp.," *Journal of Fermentation Technology*, vol. 56, pp. 91–95, 1978.
- [23] K. Kishimoto, K. Kintaka, and N. Ochiyama, "Production of D-ribose," US Patent 4 904 587, 1990.
- [24] K. Miyagawa, J. Miyazaki, and N. Kanzaki, "Method of producing D-ribose," European patent 0 501 765 A1, 1992.
- [25] K.-I. Sasajima and M. Yoneda, "Carbohydrate metabolism-mutants of a *Bacillus* species. Part II. D-ribose accumulation by pentose phosphate pathway mutants," *Agricultural and Biological Chemistry*, vol. 35, no. 4, pp. 509–517, 1971.
- [26] L. Wu, Z. M. Li, and Q. Ye, "Enhanced D-ribose biosynthesis in batch culture of a transketolase-deficient *Bacillus subtilis* strain by citrate," *Journal of Industrial Microbiology and Biotechnology*, vol. 36, no. 10, pp. 1289–1296, 2009.
- [27] T. Fang, X. Chen, N. Li et al., "Optimization of medium components for d-ribose production by transketolase-deficient *Bacillus subtilis* NJT-1507," *Korean Journal of Chemical Engineering*, vol. 27, no. 6, pp. 1725–1729, 2010.
- [28] S. K. Zhang, S. L. Yan, H. Y. Huang et al., "Study on production of D-ribose single-stage continuous fermentation," *Food and Fermentation Industries*, vol. 38, no. 1, pp. 92–95, 2012 (Chinese).
- [29] P. F. Ren, Q. A. Peng, T. Y. Yang, and C. Xu, "Study on the conditions of D-ribose fermentation," *Journal of Biotechnology*, vol. 18, no. 5, pp. 75–77, 2008 (Chinese).
- [30] Y.-C. Park and J.-H. Seo, "Optimization of culture conditions for D-ribose production by transketolase-deficient *Bacillus subtilis* JY1," *Journal of Microbiology and Biotechnology*, vol. 14, no. 4, pp. 665–672, 2004.
- [31] T. Niwa, U. Doi, Y. Kato, and T. Osawa, "Antioxidative properties of phenolic antioxidants isolated from corn steep liquor," *Journal of Agricultural and Food Chemistry*, vol. 49, no. 1, pp. 177–182, 2001.
- [32] J. Saxena and R. S. Tanner, "Optimization of a corn steep medium for production of ethanol from synthesis gas fermentation by *Clostridium ragsdalei*," *World Journal of Microbiology and Biotechnology*, vol. 28, no. 4, pp. 1553–1561, 2012.
- [33] P. De Wulf, W. Soetaert, D. Schwengers, and E. J. Vandamme, "D-Glucose does not catabolite repress a transketolase-deficient D-ribose producing *Bacillus subtilis* mutant strain," *Journal of Industrial Microbiology*, vol. 17, no. 2, pp. 104–109, 1996.
- [34] R. K. Srivastava and P. P. Wangikar, "Combined effects of carbon, nitrogen and phosphorus substrates on D-ribose production via transketolase deficient strain of *Bacillus pumilus*," *Journal of Chemical Technology and Biotechnology*, vol. 83, no. 8, pp. 1110–1119, 2008.
- [35] F. G. Zhang, Z. Zhu, B. Wang et al., "Optimization of *Trichoderma harzianum* T-E5 biomass and determining the degradation sequence of biopolymers by FTIR in solid-state fermentation," *Industrial Crops and Products*, vol. 49, pp. 619–627, 2013.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

