

Response of Nutrient Solution and Photosynthetic Photon Flux Density for Growth and Accumulation of Antioxidant in *Agastache rugosa* under Hydroponic Culture Systems

Sung Jin Kim, Kwon Jung Bok, Vu Phong Lam, and Jong Seok Park*

Department of Horticultural Science, Chungnam National University, Daejeon 34134, Korea

Abstract. *Agastache rugosa*, is a perennial medicinal plant commonly used in Chinese herbalism, and may have anti-atherogenic and antibacterial properties. Here in this study, we investigated the growth and variations in antioxidant contents of *A. rugosa* in response to nutrient solution and photosynthetic photon flux density (PPFD) with artificial lighting for a hydroponics culture. Fluorescent light at 150, and 200 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ PPFD with a 16/8 (light/dark) photoperiod, combined with four different nutrient solutions [developed by Horticulture experiment station in Japan (HES), University of Seoul (UOS), Europe vegetable research center (EVR), Otsuka-house 1A (OTS)], were used in a hydroponics culture system for 6 weeks. The shoot and root dry weights of *A. rugosa* grown with the OTS were significantly higher than those of other nutrient solutions. The amount of tilianin was the highest grown with the OTS, followed by EVR, HES, and UOS. Total acacetin content was the highest in *A. rugosa* grown under EVR which was statistically similar with OTS. The *A. rugosa* grown under 200 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ PPFD produced higher fresh weight and both acacetin and tilianin contents than that grown under 150 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ PPFD. The present results suggested that OTS along with 200 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ PPFD could be an optimum growing condition for better growth and higher accumulation of tilianin and acacetin contents in *A. rugosa* with hydroponic culture systems in a plant factory.

Additional key words : acacetin, hydroponics, medicinal plant, PPFD, tilianin

Introduction

Agastache rugosa, is a perennial medicinal plant belonging to the family Labiatae, usually grown throughout East Asian countries especially Korea and China. In Korea, this plant is a popular herb and has been used as an herbal drug in traditional therapies as well as a wild vegetable (Ahn and Yang, 1991). *A. rugosa* contains several types of essential oils, sesquiterpenes, diterpenes, triterpenes, flavonoids, and carotenoids (Han, 1987; Lee et al., 1995; Choi and Lee, 1999). Among these, tilianin, which is considered to be the main flavonoid, and rosmarinic acid have been proven to contribute to the medicinal potential of *A. rugosa* (Kim et al., 1999; Oh et al., 2006). Tilianin and acacetin are found in various plants as a glucose-glycoside compound of a flavonoid. Natural product chemists and clinicians have been interested in tilianin because of its important biological activities, for example, its anti-inflammatory, antiatherogenic, antihypertensive, and vasorelax-

ant effects (Hong et al., 2001; Nam et al., 2006; Hernandez et al., 2009). Studies have indicated that phytochemicals from *A. rugosa*, tilianin, acacetin, and rosmarinic acid, has pharmacological activities such as anti-HIV integration activities and antifungal effects (Kim et al., 1999; Shin, 2004). Extracts of *A. rugosa* are also believed to be valuable in the treatment of inflammatory and oxidative stress-induced disorders (Hong et al., 2001; Lee et al., 2002). Daily consumption of herbal or medicinal plants containing bioactive compounds has been increased recently in a public health (Wu and Kubota, 2008). Developing fresh produce containing greater concentrations of phytochemicals to have increasing effect on some health aspects can be an alternative approach to increase the exposure to the bioactive compounds (Kubota et al., 2006).

By controlling environmental conditions such as temperature, nutrient solution, light intensity or quality etc., manipulation of target compounds in plants such as phytochemicals and antioxidants has been studied as a research area attracting scientific interest (Afreen et al., 2006; Kubota et al., 2006; Zobayed et al., 2007). Also, increasing demand of herbal or medicinal plants for industrial area such as health beverages, functional cosmetic, and medica-

*Corresponding author: jongseok@cnu.ac.kr

Received September 05, 2017; Revised September 20, 2017;

Accepted September 22, 2017

tions has led to their mass cultivation under a closed or semi closed-type plant production system in a large scale manner. Closed systems with a hydroponic culture and artificial lights for plant production have several potential benefits, such as higher quality of transplants (Kozai et al., 2013), optimization of plant production systems (Seginer and Ioslovich, 1999), shorter production period, and manipulation of phytochemicals by environmental control (Park, et al., 2014), and year-round production, compared with open-type production systems (Kozai et al., 2006). To produce some wild plant in a hydroponic culture system, strength or combination of each nutrient solution and light intensity of artificial light are crucial for plant growth and can affect the concentration of phytochemical content during cultivation (Wu and Kubota, 2008). Especially, light intensity irradiated from fluorescent lamps in a plant production system affects greatly initial cost for building of the production systems. A commercial plant factory system is designed with an artificial light source which has to be irradiated for the proper crops growth and quality with the economic point of view. To produce leafy vegetables, five plant factories in Korea I have visited have supplied photosynthetic photon flux density (PPFD) from 110 to $165\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ on the seedlings canopy level. Although this values can be considered quite low level as compared to outdoor production environment, burden of the electricity charge from the lighting source is the big part of the plant factory operation cost, especially, in Japan, and the increase of the light intensity, PPFD, greatly affects the initial installation cost, which increases the depreciation cost of the plant factory. Considering a realistic situation, It is necessary to understand how the difference in light intensity from 150 to $200\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ PPFD affects the growth and phytochemicals content in *A. rugosa* grown in the plant factory systems.

Up to now, many studies related to nutrient solution and PPFD have been reported for growth of leaf vegetable, but no information about *A. rugosa* is available related to nutrient solution and suitable PPFD which might affect the growth and phytochemical contents in hydroponic culture systems. To produce *A. rugosa* efficiently for industrial application, selection of optimum nutrient solution and proper light intensity with fluorescent lamps have to be investigated under the conditions in closed plant production systems. Therefore, the present study was undertaken getting a proper nutrient solution and an economic PPFD

condition for maximizing growth and higher accumulation of secondary metabolites in *A. rugosa* grown in a plant factory system.

Materials and methods

1. Plant Materials & Culture Systems

Seeds of *A. rugosa* were purchased from Danong Seed Co. Ltd, Seoul, Korea. Seeds were sown in a crushed rock-wool media tray (240 holes, UR rockwool, Suwon, Korea) in a plant culture room and then placed in a dark condition for germination at 26°C for first 2 days. The culture room was maintained at $22\pm 1^\circ\text{C}$ (day) and $18\pm 1^\circ\text{C}$ (night) using an air conditioner (AF30FSAM1EEN, Samsung Co. Ltd., Suwon, Korea). Fluorescent lamps (TL5 14W/865 Philips, Korea) were installed in it providing a PPFD of $120\pm 10\mu\text{mol m}^{-2} \text{s}^{-1}$ on the surface of the culture media. Immediately after appearance of hypocotyl, the seedlings were placed in a half-strength Yamazaki Nutrient Solution (6 me L⁻¹ NO₃-N, 0.5 me L⁻¹ NH₄-N, 2 me L⁻¹ Ca, 1.5 me L⁻¹ PO₄-P, 1 me L⁻¹ Mg, and 4 me L⁻¹ K with micro nutrient solutions) during seedling growth stage. Fifty two seedlings at 42 days aged (at four-leaf stage) were selected for four nutrient solution treatments. Thirteen seedlings for one treatment were planted maintaining 120 mm planting distance and cultured at $22\pm 1^\circ\text{C}$ (day) and $18\pm 1^\circ\text{C}$ (night) in a culture container (460 mm long × 330 mm wide × 110 mm high) for 4 weeks. Fluorescent lamps provided with PPFDs of 153 ± 10 , 151 ± 13 , 152 ± 11 , or $153\pm 11\mu\text{mol m}^{-2} \text{s}^{-1}$ (mean±SD) was averaged at 9 points at the canopy level for four conditions of nutrient solutions with a 16/8h photoperiod (Ex-1). The PPFDs with fluorescent lamps was controlled to 152 ± 11 and $201\pm 15\mu\text{mol m}^{-2} \text{s}^{-1}$ (mean±SD) to find out the optimum light condition for increasing plant growth and phytochemicals in *A. rugosa* with a Otsuka-house 1A solution for 6 weeks under the same environmental conditions as above (Ex-2).

2. Nutrient Solution Treatment

Four nutrient solutions, developed by Horticultural Experiment Station in Japan (HES), The University of Seoul (UOS), European Vegetable R&D Center (EVR), and Otsuka-house 1A (OTS) were used as experimental treatment in this study (Table 1). HES and OTS have higher nitrogen content, potassium, and calcium content than UOS and EVR. Electrical conductivity of OTS and

Table 1. The composition of UOS, HES, EVR, and OTS nutrient solution for *A. rugosa* in hydroponics.

Nutrient solution ^z	Nutrient concentration (me L ⁻¹)						EC (dS m ⁻¹)
	NO ₃ -N	NH ₄ -N	P	K	Ca	Mg	
UOS	11.6	1.2	3.6	5.8	5.8	3.0	1.0
HES	16.0	1.3	4.0	8.0	8.0	4.0	1.2
EVR	10.1	0.5	5.4	5.4	4.8	2.0	1.0
OTS	16.8	1.8	5.1	8.6	8.2	3.0	1.3

^zUOS: The nutrient solution developed by University of Seoul.

HES: The nutrient solution of Horticultural Experiment Station in Japan.

EVR: The nutrient solution of European Vegetable R&D Center.

OTS: The nutrient solution, Otsuka House1-A developed by Otsuka Co. Ltd., in Japan.

HES was 1.3 and 1.2 dS m⁻¹, respectively, and UOS and EVR were 1.0 dS m⁻¹. Each nutrient solution was put in a tank (460mm long × 330mm wide × 110mm high) installed under the culture container and was supplied by a pump (HJ-950-20W, Amazon, China). The nutrient solution was maintained with a constant level of 60mm height in the culture container and supplied for 24 hours, continuously.

3. Measurement

Leaf number was counted which developed over 10 mm after detaching from the basal ends of the stems. Length and width of the each detached leaf and stem length were measured with a digital caliper (Digipa, Mitutoyo Co., Ltd., Tokyo, Japan). Shoot and root fresh weight was measured with a digital balance (ARG224 OHAUS, Sigma-Aldrich Co. LLC, Seoul, Korea) after absorbing surface water of roots by compressing the roots with a paper (Kimwiper S200, Japan Paper Crexia Co., Ltd., Tokyo, Japan). Then, the shoot and roots were placed in a dry oven (HB-502M, Hanback Sci, Suwon, Korea) at 66°C for 7 d, and the shoot and root dry weight were also measured with the balance.

4. Extraction and HPLC Analysis for Tilianin and Acacetin

Samples (0.1 g) were frozen in liquid nitrogen, ground to a fine powder using a mortar and pestle, and then extracted with 10mL of 70% ethanol for 1 h at 60°C. After centrifugation, the supernatant was filtered through a 0.45µm poly filter (Acrodisc Syringe Filters, Pall, Port Washington, NY) and analyzed by HPLC. The analysis was performed with a C18 column (250mm × 4.6mm, 5µm; RS tech, Daejon, Korea) at 30°C. The mobile phase was a gradient mixture of acetonitrile, methanol, and 0.2% acetic acid. The flow rate was maintained at 1.0mL·min⁻¹, the injection volume was 20µL, and the detection wavelength was 275 nm. The

concentration of tilianin and acacetin in the samples was calculated by using a standard curve. Mean values were obtained from three independent replicates.

5. Statistical Analysis

Each experiment was conducted twice with completely randomized designs. All data were subjected to ANOVA followed by Duncan's multiple rate test at $P \leq 0.05$, 0.01 (JMP, SAS Institute Inc., Cary, NC, USA).

Results and Discussion

1. Growth as Influenced by Different Nutrient Solution (Ex-1)

A. rugosa grown under different nutrient solution significantly influenced the growth of root, shoot and stem length (Fig. 1). Among the nutrient solution, the OTS performed the best for giving the highest stem length, shoot fresh and dry weights, and root fresh and dry weights of *A. rugosa*. The variation of root dry weight was much higher than any other growth parameters. Nutrient solution OTS dominated over other nutrient solution for the production of high amount of root dry matter compared to HES, UOS, or EVR treatment and its stem length was significantly longer compared to EVR culture medium (Fig. 1). However, the growth pattern for leaf number was slightly different compared to other growth parameters. Though OTS nutrient solution influenced markedly for achieving higher values for most of the growth parameters except leaf number, *A. rugosa* grown under nutrient solution of EVR produced the highest leaf number than that of other nutrient solution. The ratio of each cation and anion affects the uptake of the nutrient solution by the plants, and their growth changes. OTS nutrient solution contains higher potassium and cal-




Fig. 1. Growth parameters of *A. rugosa* grown under the different nutrient solutions in hydroponics. HES: The nutrient solution of Horticultural Experiment Station in Japan, UOS: The nutrient solution developed by The University of Seoul, EVR: The nutrient developed by The University of Seoul, EVR: The nutrient solution of European Vegetable R&D Center, OTS: The nutrient solution, Otsuka House1-A developed by Otsuka Co. Ltd., in Japan. Data represent means and standard deviation (n=13). Means with different letters are significantly different ($P < 0.05$ by DMRT).

cium ions compared to the other nutrient solutions, also EC was the highest among treatments. Fruit dry matter of tomato (*Solanum lycopersicum* ‘Lunarossa’) were increased by higher proportion of potassium in the nutrient solution (Fanasca et al., 2006). In an another study, Mossi et al, (2011) reported that growth of *Cunila galioides* in a hydroponics culture showed significant differences due to variation of concentration of nutrient solution. Here in this study

OTS nutrient solution which has higher potassium and calcium, or EC compared to other treatments performed the best for achieving higher values for several growth parameters in the hydroponic culture system for *A. rugosa*.

2. Tilianin and Acacetin Contents as Influenced by Different Nutrient Solution

At 4 weeks after transplanting, seedlings of *A. rugosa*




Fig. 2. Tilianin and total acacetin contents of *A. rugosa* grown under the different nutrient solutions in hydroponics for 4 weeks. HES: The nutrient solution of Horticultural Experiment Station in Japan, UOS: The nutrient solution developed by The University of Seoul, EVR: The nutrient solution of European Vegetable R&D Center, OTS: The nutrient solution, Otsuka House1-A developed by Otsuka Co. Ltd., in Japan. Data represent means and standard deviation ($n=3$). Means with different letters are significantly different ($P \leq 0.05$ by DMRT).

grown under different nutrient solution, were analyzed for detection of tilianin and acacetin in the hydroponics culture system. Both tilianin and acacetin content varied significantly in response to different nutrient solution (Fig. 2). The level of tilianin was the highest in *A. rugosa* grown under OTS condition achieving 51% higher tilianin than that in the UOS (Fig. 2). The content of tilianin was almost similar when *A. rugosa* grown under EVR and HES conditions. There was a significant difference in acacetin content of *A. rugosa* grown under different nutrient solution. For acacetin accumulation, EVR showed the best performance which was statistically similar grown under OTS whereas the lowest acacetin was accumulated in *A. rugosa* grown under UOS. Sorgoleone content in the root of sor-

ghum greatly influenced by the different concentration of Hoagland solution (Uddin et al., 2010). In the case of the application of full strength Hoagland solution, sorgoleone content increased with delayed application. Half-strength solution produced maximum root growth and sorgoleone content. The total phenolic content and antioxidant capacity per shoot of *Crepidiastrum denticulatum* significantly increased with increasing in nutrient solution concentration from 0.5 to 2.5 dS m^{-1} concentration in a plant factory (Park et al., 2016). The mean antioxidant activity of tomato (*Solanum lycopersicum*) grown under the NH_4^+ nutrient solution with high Cl^- and SO_4^{2-} levels ($\text{NO}_3^- : \text{NH}_4^+$ ratio =1:4) reduced 14% lower compared with NO_3^- nutrient solution with low Cl^- and intermediate SO_4^{2-} levels ($\text{NO}_3^- : \text{NH}_4^+$ ratio =4:1) (Toor et al., 2006). Total soluble solid content and lycopene content of tomato (*Solanum lycopersicum* ‘Lunarossa’) were increased by higher proportion of potassium in the nutrient solution (Fanasca et al., 2006). Also a higher proportion of potassium of 7 and 14 mmol L^{-1} increased total phenolic content of red and green peppers (*Capsicum annuum*) compared with lower potassium concentration of 0.2 and 2 mmol L^{-1} in nutrient solution (Marín et al., 2009). Higher EC of nutrient solution or composition of nutrient solution may have affected accumulation of secondary metabolites, which play roll in plant protection. However, Total phenolic compound levels in the two kinds of basil (*Ocimum basilicum* L.) ‘dark opal’ and ‘genovese’ increased with decreasing nitrogen concentration at the treatments of 0.1, 0.5, 1.0, and 5.0 mM nitrogen concentrations. Similarly, basil grown under the lowest nitrogen fertilization level generally contained significantly higher rosmarinic and caffeic acid concentrations ($p \leq 0.001$) than basil treated at higher nitrogen levels. However, the anthocyanin content of ‘dark opal’ was not affected by applied nitrogen level, but anthocyanin concentrations were significantly impacted by growing season ($p \leq 0.001$) (Nguyen and Niemeyer 2008). Application of different nutrient solution and phytohormones acted positively for accumulation of different secondary metabolites (Uddin et al., 2010; Kim et al., 2013; Uddin et al., 2013). Therefore, the composition of EVR which contains lower NH_4^+ nutrient and OTS nutrient solutions which was higher potassium and EC than other treatments may increase tilianin and acacetin contents of *A. rugosa* when growing in hydroponics culture systems.




Fig. 3. Time course of fresh weight of *A. rugosa* grown under the photosynthetic photon flux density (PPFD) of 150 and 200 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for 6 weeks. Data represent means and standard deviation ($n=4$). Means with asterisk are significantly different ($P \leq 0.05$ by DMRT).

3. Growth in Response to Different PPFDs Condition

A time course experiment was considered to investigate the growth and tilianin and acacetin contents in *A. rugosa* under two PPFDs conditions. Results of time course experiment revealed that the growth of *A. rugosa* was subjected to different PPFDs conditions which has crucial view point for plant growth in a plant factory. Fresh weight of *A. rugosa* was increased in 200 $\mu\text{mol m}^{-2} \text{s}^{-1}$ irradiated than that of 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$. There was a significantly difference in the fresh weights between two PPFDs treatments from the third week after transplanting. At the growth of 5 week, the difference of fresh weight was higher, after that these differences were reduced in 6 weeks (Fig. 3). The fresh and dry weights of shoot and leafy area were affected more by nutrient solution EC than by PPFD in green perilla (*Perilla frutescens*) grown in a plant factory system. Leaf photosynthetic rates were increased as PPFD increased from 100 to 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ in green and red perilla varieties (Lu et al., 2017). To appropriately grow *A. rugosa* in a plant factory system, optimum PPFD conditions may be better around 200 $\mu\text{mol m}^{-2} \text{s}^{-1}$ than 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$ at the canopy level.

4. Tilianin and Acacetin Contents in Response to Different PPFDs Condition

The total acacetin and tilianin contents of *A. rugosa* were shown to be significantly higher under the PPFD of




Fig. 4. Tilianin and total acacetin contents of *A. rugosa* grown under the different photosynthetic photon flux density (PPFD; 150 and 200 $\mu\text{mol m}^{-2} \text{s}^{-1}$) in hydroponics for 6 weeks. The values of PPFDs were measured on the planting board at the start of the experiment. Means with different letters are significantly different ($P \leq 0.01$ by DMRT).

200 $\mu\text{mol m}^{-2} \text{s}^{-1}$ condition at all dates of sampling (Fig. 4). In particular, by five weeks after transplanting, the total acacetin content under the PPFD of the 200 $\mu\text{mol m}^{-2} \text{s}^{-1}$ condition increased by approximately 49% compared to the PPFD 150 condition and the total tilianin content under the same condition increased by approximately 40% by six weeks after transplanting (Fig. 4). On reviewing the light intensity/photosynthesis curve of *Angelica gigas* on the basis of individual leaves, the light intensity at which photosynthesis increased maximally is observed under the condition of approximately 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PPFD except for slightly above the light compensation point. The light intensity that shows the maximum slope value of light intensity/photosynthesis curves can be said to be the most economic light condition, in many actual studies of leaf vegetables, increases in

growth under the PPFD of $200\mu\text{mol m}^{-2} \text{s}^{-1}$ condition were reported (Park et al., 2014). Given the growth and phytochemical content of *A. rugosa*, the PPFD of $200\mu\text{mol m}^{-2} \text{s}^{-1}$ conditions is considered to be more economic light intensity than the 150 PPFD condition. Light conditions have showed significant influence on accumulation of secondary metabolites (Li et al., 2013; Al-Dhabi et al., 2015). Perillaldehyde concentrations, standard functional materials in green perilla (*Perilla frutescens*), was significantly different between combinations of the highest PPFD (among 100, 200, and $300\mu\text{mol m}^{-2} \text{s}^{-1}$) with the highest EC (among 1.0, 2.0, and 3.0 dS m^{-1}), and the lowest PPFD with the lowest EC. However, rosmarinic acid concentration (mg g^{-1}) of green perilla was increased in a combination of low EC and high PPFD conditions (Lu et al., 2017). It varies depending on the environmental conditions, such as nutrient solution, light intensity, temperature, etc. for the plant growth and the kinds of secondary metabolites. It is believed that optimum light conditions in plant factories is crucial to induce optimum crop production, and that bioactive compounds of *A. rugosa* were also increased in the light intensity of $200\mu\text{mol m}^{-2} \text{s}^{-1}$ which is applied for plant growth. Additional experiments with increased PPFD for plant growth and accumulation of bioactive compounds of *A. rugosa* are required.

Conclusion

The optimum growth and accumulation of tlianin and acacetin of *A. rugosa* in response to different nutrient solution and PPFD considering economic may extend our understanding of the mechanisms related to the biosynthesis of tlianin and acacetin in plants. Moreover, our study also indicates increasing the production of tlianin and acacetin as well plant growth of *A. rugosa* in a plant factory system.

Acknowledgement

This work was supported by Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry and Fisheries (IPET) through Agri-Bio industry Technology Development Program, funded by Ministry of Agriculture, Food and Rural Affairs (MAFRA) (116115-03-1-SB010).

Literature Cited

- Ahn, B., and C. B. Yang. 1991. Volatile flavor components of Bangah herb. Korean J. Food Sci. Technol. 23:582-586.
- Al-Dhabi, M., V. Arasu, S.J. Kim, M.R. Uddin, W.T. Park, S.Y. Lee, and S.U. Park. 2015. Methyl jasmonate- and light-induced glucosinolate and anthocyanin biosynthesis in radish seedlings. Natural Products Communications 10:1211-1214.
- Afreen, F., S.M.A. Zobayed, and T. Kozai. 2006. Melatonin in *Glycyrrhiza uralensis*: response of plant roots to spectral quality of light and UV-B radiation. J. Pineal Res. 41:108-115.
- Choi, K.S., and H.Y. Lee. 1999. Characteristics of useful components in the leaves of baechohyang (*Agastache rugosa*, O. Kuntze). J. Korean Soc. Food Sci. Nutr. 28:326-332 In Korean.
- Fanasca, S, G. Colla, G. Maiani, E. Venneria, Y. Rouphael, E. Azzini and F. Saccardo. 2006. Changes in antioxidant content of tomato fruits in response to cultivar and nutrient solution composition. J. Agric. Food Chem. 54:4319-4325.
- Han, D. S. 1987. Triterpenes from the root of *Agastache rugosa*. Korean J. Pharmacogn. 18:50-53 In Korean.
- Hernández-Abreu, O., P. Castillo-España, I. León-Rivera, M. Ibarra-Barajas, R. Villalobos-Molina, J. González-Christen, J. Vergara-Galicia, and S. Estrada-Soto. 2009. Antihypertensive and vasorelaxant effects of tlianin isolated from *Agastache mexicana* are mediated by NO/cGMP pathway and potassium channel opening. Biochem. Pharmacol. 78:54-61.
- Hong, J.J., J.H. Choi, S.R. Oh, H.K. Lee, J.H. Park, K.Y. Lee, J.J. Kim, T.S. Jeong, and G.T. Oh. 2001. Inhibition of cytokine-induced vascular cell adhesion molecule-1 expression; possible mechanism for anti-atherogenic effect of *Agastache rugosa*. FEBS Lett. 495:142-147.
- Kim Y.B., J.K. Kim, M.R. Uddin, H.H. Xu, W.T. Park, P.A. Tuan, X. Li, E. Chung, J.H. Lee, and S.U. Park. 2013. Metabolomics analysis and biosynthesis of rosmarinic acid in *Agastache rugosa* Kuntze treated with methyl jasmonate. PLoS ONE 8(5):e64199.
- Kim, H.K., H.K. Lee, C.G. Shin, and H. Huh. 1999. HIV integrase inhibitory activity of *Agastache rugosa*. Arch. Pharm. Res. 22:520-523.
- Kozai, T. 2013. Resource use efficiency of closed plant production system with artificial light: Concept, estimation and application to plant factory. Proc. Jpn. Acad. Ser. B Phys. Biol. Sci. 89(10):447-461.
- Kozai, T., K. Ohyama, and C. Chun. 2006. Commercialized closed systems with artificial lighting for plant production. Acta Hort. 711:61-70.
- Kubota, C., C.A. Thomson, M. Wu, and J. Javanmardi. 2006. Controlled environments for production of value-added food crops with high phytochemical concentrations: lycopene in

- tomato as an example. *HortScience* 41:522-525.
- Lee, C., H. Kim, and Y. Kho. 2002. Agastinol and agastenol, novel lignans from *Agastache rugosa* and their evaluation in an apoptosis inhibition assay. *J. Nat. Prod.* 65:414-416.
- Lee, H.K., S.R. Oh, J.I. Kim, J.W. Kim, and C.O. Lee. 1995. Agastaquinone, a new cytotoxic diterpenoid quinone from *Agastache rugosa*. *J. Nat. Prod.* 58:1718-1721.
- Li, X., Y.B. Kim, M.R. Uddin, S. Lee, S.J. Kim, and S.U. Park. 2013. Influence of light on the free amino acid content and γ -Aminobutyric acid synthesis in *Brassica juncea* seedlings. *J. Agric. Food Chem.* 61:8624-8631.
- Lu, N., E.L. Bernardo, C. Tippayadarapanich, M. Takagaki, N. Kagawa, and W. Yamori. 2017. Growth and accumulation of secondary metabolites in perilla as affected by photosynthetic photon flux density and electrical conductivity of the nutrient solution. *Front. Plant Sci.* 708:1-12.
- Marín A., J.S. Rubio, V. Martinez, and M. I. Gil. 2009. Antioxidant compounds in green and red peppers as affected by irrigation frequency, salinity and nutrient solution composition. *J. Sci. Food Agric.* 89:1352-1359.
- Mossi, A.J., G.F. Pauletti, L. Rota, S. Echeverrigaray, I.B.I. Barros, J.V. Oliveira, N. Paroul, and R.L. Cansian. 2011. Effect of aluminum concentration on growth and secondary metabolites production in three chemotypes of *Cunila galioides* Benth. *Medicinal plant. Braz. J. Biol.* 71:1003-1009.
- Nam, K.H., J.H. Choi, Y.J. Seo, Y.M. Lee, Y.S. Won, M.R. Lee, M.N. Lee, J.G. Park, Y.M. Kim, H.C. Kim, C.H. Lee, H.K. Lee, S.R. Oh, and G.T. Oh. 2006. Inhibitory effects of tilianin on the expression of inducible nitric oxide synthase in low density lipoprotein receptor deficiency mice. *Exp. Mol. Med.* 38:445-452.
- Nguyen, P.M. and E.D. Niemeyer. 2008. Effects of nitrogen fertilization on the phenolic composition and antioxidant properties of basil (*Ocimum basilicum* L.),” *Brown Working Papers in the Arts and Sciences, Southwestern University, Vol. VIII.* <http://www.southwestern.edu/academic/bwp/vol8/niemeyer-vol8.pdf>.
- Oh, H.M., Y.J. Kang, Y.S. Lee, M.K. Park, S.H. Kim, H.J. Kim, H.G. Seo, J.H. Lee, and K.C. Chang. 2006. Protein kinase G-dependent heme oxygenase-1 induction by *Agastache rugosa* leaf extract protects RAW264.7 cells from hydrogen peroxide-induced injury. *J. Ethnopharmacol.* 103:229-235.
- Park, J.S., S.J. Kim, H.J. Kim, J.M. Choi, and G.I. Lee. 2014. Photosynthetic characteristics and growth analysis of *Angelica gigas* according to difference hydroponics methods. *CNU J. of Agri. Sci.* 41:321-326.
- Park, S.Y., S.B. Oh, S.M. Kim, Y.Y. Cho, and M.M. Oh. 2016. Evaluating the effects of a newly developed nutrient solution on growth, antioxidants, and chicoric acid contents in *Crepidiastrum denticulatum*. *Hortic. Environ. Biotechnol.* 57:478-486.
- Seginer, I., and I. Ioslovich. 1999. Optimal spacing and cultivation intensity for an industrialized crop production system. *Agric. Syst.* 62:143-157.
- Shin, S. 2004. Essential oil compounds from *Agastache rugosa* as antifungal agents against Trichophyton species. *Arch. Pharm. Res.* 27:295-299.
- Toor, R.k., G.P. Savage, and a. Heeb. 2006. Influence of different types of fertilisers on the major antioxidant components of tomatoes. *J. Food Composition and Analysis* 19:20-27.
- Uddin, M.R., A.A. Thwe, Y.B. Kim, W.T. Park, S.C. Chae, and S.U. Park. 2013. Effects of jasmonates on sorgoleone accumulation and expression of genes for sorgoleone biosynthesis in sorghum roots. *J. Chem. Ecol.* 39:712-722.
- Uddin. M.R., K.W. Park, Y.K. Kim, S.U. Park, and J.Y. Pyon. 2010. Enhancing sorgoleone levels in grain sorghum root exudates. *J. Chem. Ecol.* 36:914-922.
- Wu, M., and C. Kubota. 2008. Effects of high electrical conductivity of nutrient solution and its application timing on lycopene, chlorophyll and sugar concentrations of hydroponic tomatoes during ripening. *Sci. Hort.* 116:122-129.
- Zobayed, S.M.A., F. Afreen, and T. Kozai. 2007. Phytochemical and physiological changes in the leaves of St. John's wort plants under a water stress condition. *Environ. Exp. Bot.* 59:109-116.

식물공장에서 양액의 종류 및 PPF가 배초향의 성장 및 항산화 물질에 미치는 영향

김성진 · 복권정 · 푸동람 · 박종석*

충남대학교 원예학과

적 요. 배초향은 항동맥경화나 항박테리아의 특성을 가지는 한약재에 널리 사용되는 영년생 약용식물이다. 연구의 목적은 수경재배에서 배양액의 종류와 PPF값에 따른 배초향의 성장 및 항산화 물질의 변화를 조사하는 것이다. 배초향은 주야간 16:8 시간의 일장조건에서 150과 200 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ PPF 조건과 일본원시(HES), 서울시립대(UOS), 유럽채소연구소(EVR), 오오츠키 배양액(OTS)을 이용하여 6주간 재배하였다. OTS 배양액조건에서 자란 배초향의 지상부 및 지하부 건물중은 다른 배양액 처리구와 비교하여 유의적으로 높았다. 배초향의 티리아닌 함량은 OTS 처리에서 가장 높았으며 다음으로 EVR, HES, UOS 순서로 낮아졌다. 총 아카세틴의 함량은 EVR처리에서 가장 높았으나 OTS처리와는 유의적 차이를 보이지 않았다. 또한 200 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ PPF 조건에서 자란 배초향은 PPF 150처리구와 비교하여 유의적으로 생체중과 건물중이 증가하였으며 기능성 물질은 티리아닌과 아카세틴의 함량도 높았다. 본 연구는 수경재배 방식을 이용하여 식물공장에서 배초향을 재배할 경우 200 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ PPF 조건과 OTS 배양액 조건에서 경제적인 광원조건으로 최적 바이오매스 생산량과 티리아닌과 아카세틴의 함량을 증가시킬 수 있을 것으로 제안한다.

추가 주제어 : 아카세틴, 수경재배, 약용작물, 광합성광량자속밀도, 티리아닌