

Solar system

The **Solar System** or **solar system** consists of the Sun and the other celestial objects gravitationally bound to it: the eight planets, their 166 known moons,^[1] three dwarf planets (Ceres, Pluto, and Eris and their four known moons), and billions of small bodies. This last category includes asteroids, Kuiper belt objects, comets, meteoroids, and interplanetary dust.

In broad terms, the charted regions of the Solar System consist of the Sun, four terrestrial **inner planets**, an asteroid belt composed of small rocky bodies, four gas giant **outer planets**, and a second belt, called the Kuiper belt, composed of icy objects. Beyond the Kuiper belt is the scattered disc, the heliopause, and ultimately the hypothetical Oort cloud.


In order of their distances from the Sun, the planets are Mercury, Venus, Earth, Mars, Jupiter, Saturn, Uranus, and Neptune. Six of the eight planets are in turn orbited by natural satellites, usually termed "moons" after Earth's Moon, and each of the outer planets is encircled by planetary rings of dust and other particles. All the planets except Earth are named after gods and goddesses from Greco-Roman mythology. The three dwarf planets are Pluto, the largest known Kuiper belt object; Ceres, the largest object in the asteroid belt; and Eris, which lies in the scattered disc.

Terminology


See also: Definition of planet

Objects orbiting the Sun are divided into three classes: planets, dwarf planets, and small Solar System bodies.

A planet is any body in orbit around the Sun that a) has enough mass to form itself into a spherical shape and b) has cleared its immediate neighbourhood of all smaller objects. There are eight known planets: Mercury, Venus, Earth, Mars, Jupiter, Saturn, Uranus, and Neptune.



Major features of the Solar System; sizes and distances not to scale. From left to right): Pluto, Neptune, Uranus, Saturn, Jupiter, the asteroid belt, the Sun, Mercury, Venus, Earth and its Moon, and Mars. A comet is also seen on the left.



Planets and dwarf planets of the Solar System; while the sizes are to scale, the relative distances from the Sun are not.

On August 24 2006 the International Astronomical Union defined the term "planet" for the first time, excluding Pluto and reclassifying it under the new category of dwarf planet along with Eris and Ceres.^[2]

A dwarf planet is not required to clear its neighbourhood of other celestial bodies. Other objects that may become classified as dwarf planets are Sedna, Orcus, and Quaoar.

From the time of its discovery in 1930 until 2006, Pluto was considered the Solar System's ninth planet. But in the late 20th and early 21st centuries, many objects similar to Pluto were discovered in the outer Solar System, most notably Eris, which is slightly larger than Pluto.

The remainder of the objects in orbit around the Sun are small Solar System bodies (SSSBs).^[3]


Natural satellites, or moons, are those objects in orbit around planets, dwarf planets and SSSBs, rather than the Sun itself.

A planet's distance from the Sun varies in the course of its year. Its closest approach to the Sun is called its perihelion, while its farthest distance from the Sun is called its aphelion.

Astronomers usually measure distances within the Solar System in astronomical units (AU). One AU is the approximate distance between the Earth and the Sun, or roughly 149,598,000 km (93,000,000 mi). Pluto is roughly 38 AU from the Sun while Jupiter lies at roughly 5.2 AU. One light year, the best known unit of interstellar distance, is roughly 63,240 AU.

Informally, the Solar System is sometimes divided into separate zones. The **inner Solar System** includes the four terrestrial planets and the main asteroid belt. Some define the **outer Solar System** as comprising everything beyond the asteroids.^[4] Others define it as the region beyond Neptune, with the four gas giants considered a separate "middle zone".^[5]

Layout and structure



The ecliptic viewed in sunlight from behind the Moon in this Clementine image. From left to right: Mercury, Mars, Saturn.

The principal component of the Solar System is the Sun, a main sequence G2 star that contains 99.86% of the system's known mass and dominates it gravitationally.^[6] Jupiter and Saturn, the Sun's two largest orbiting bodies, account for more than 90% of the system's remaining mass.


Most large objects in orbit around the Sun lie near the plane of Earth's orbit, known as the ecliptic. The planets are very close to the ecliptic while comets and Kuiper belt objects are usually at significantly greater angles to it.

All of the planets and most other objects also orbit with the Sun's rotation in a counter-clockwise direction as viewed from a point above the Sun's north


pole. There are exceptions, such as Halley's Comet.

Objects travel around the Sun following Kepler's laws of planetary motion. Each object orbits along an approximate ellipse with the Sun at one focus of the ellipse. The closer an object is to the Sun, the faster it moves. The orbits of the planets are nearly circular, but many comets, asteroids and objects of the Kuiper belt follow highly-elliptical orbits.

To cope with the vast distances involved, many representations of the Solar System show orbits the same distance apart. In reality, with a few exceptions, the farther a planet or belt is from the Sun, the larger the distance between it and the previous orbit. For example, Venus is approximately 0.33 AU farther out than Mercury, while Saturn is 4.3 AU out from Jupiter, and Neptune lies 10.5 AU out from Uranus. Attempts have been made to determine a correlation between these orbital distances (see Titius-Bode law), but no such theory has been accepted.



Formation



The Solar System is believed to have formed according to the nebular hypothesis, first proposed in 1755 by Immanuel Kant and independently formulated by Pierre-Simon Laplace.^[7] This theory holds that 4.6 billion years ago the Solar System formed from the gravitational collapse of a giant molecular cloud. This initial cloud was likely several light-years across and probably birthed several stars.^[8] Studies of ancient meteorites reveal traces of elements only formed in the hearts of very large exploding stars, indicating that the Sun formed within a star cluster, and in range of a number of nearby supernovae explosions. The shock wave from these supernovae may have triggered the formation of the Sun by creating regions of overdensity in the surrounding nebula, allowing gravitational forces to overcome internal gas pressures and cause collapse.^[9]

The region that would become the Solar System, known as the pre-solar nebula,^[10] had a diameter of between 7000 and 20,000 AU^{[11] [12]} and a mass just over that of the Sun (by between 0.1 and 0.001 solar masses).^[13] As the nebula collapsed, conservation of angular momentum made it rotate faster. As the material within the nebula condensed, the atoms within it began to

collide with increasing frequency. The centre, where most of the mass collected, became increasingly hotter than the surrounding disc.^[14] As gravity, gas pressure, magnetic fields, and rotation acted on the contracting nebula, it began to flatten into a spinning protoplanetary disk with a diameter of roughly 200 AU^[15] and a hot, dense protostar at the center.^[16]
^[17]

Studies of T Tauri stars, young, pre-fusing solar mass stars believed to be similar to the Sun at this point in its evolution, show that they are often accompanied by discs of pre-planetary matter.^[18] These discs extend to several hundred AU and reach only a thousand kelvins at their hottest.^[19]

After 100 million years, the pressure and density of hydrogen in the centre of the collapsing nebula became great enough for the protosun to begin thermonuclear fusion. This increased until hydrostatic equilibrium was achieved, with the thermal energy countering the force of gravitational contraction. At this point the Sun became a full-fledged star.^[20]

From the remaining cloud of gas and dust (the "solar nebula"), the various planets formed. They are believed to have formed by accretion: the planets began as dust grains in orbit around the central protostar; then gathered by direct contact into clumps between one and ten metres in diameter; then collided to form larger bodies (planetesimals) of roughly 5 km in size; then gradually increased by further collisions at roughly 15 cm per year over the course of the next few million years.^[21]

The inner Solar System was too warm for volatile molecules like water and methane to condense, and so the planetesimals which formed there were relatively small (comprising only 0.6% the mass of the disc)^[22] and composed largely of compounds with high melting points, such as silicates and metals. These rocky bodies eventually became the terrestrial planets. Farther out, the gravitational effects of Jupiter made it impossible for the protoplanetary objects present to come together, leaving behind the asteroid belt.^[23]


Farther out still, beyond the frost line, where more volatile icy compounds could remain solid, Jupiter and Saturn became the gas giants. Uranus and Neptune captured much less material and are known as ice giants because their cores are believed to be made mostly of ices (hydrogen compounds).^[24]
^[25]

Once the young Sun began producing energy, the solar wind (see below) blew the gas and dust in the protoplanetary disk into interstellar space and ended the growth of the planets. T Tauri stars have far stronger stellar winds than more stable, older stars.^[26] ^[27]



Hubble image of protoplanetary disks in the Orion Nebula, a light-years-wide "stellar nursery" likely very similar to the primordial nebula from which our Sun formed.

Sun



The Sun as seen from Earth

The Sun is the Solar System's parent star, and far and away its chief component. Its large mass gives it an interior density high enough to sustain nuclear fusion, which releases enormous amounts of energy, mostly radiated into space as electromagnetic radiation such as visible light.


The Sun is classified as a moderately large yellow dwarf, but this name is misleading as, compared to stars in our galaxy, the Sun is rather large and bright. Stars are classified by the Hertzsprung-Russell diagram, a graph which plots the brightness of stars against their surface temperatures. Generally, hotter stars are brighter. Stars following this pattern are said to be on the main sequence; the Sun lies right in the middle of it. However, stars brighter and hotter than the Sun are rare, while stars dimmer and cooler are common.^[28]

It is believed that the Sun's position on the main sequence puts it in the "prime of life" for a star, in that it has not yet exhausted its store of hydrogen for nuclear fusion. The Sun is growing brighter; early in its history it was 75 percent as bright as it is today.^[29]

Calculations of the ratios of hydrogen and helium within the Sun suggest it is halfway through its life cycle. It will eventually move off the main sequence and become larger, brighter, cooler and redder, becoming a red giant in about five billion years.^[30] At that point its luminosity will be several thousand times its present value.


The Sun is a population I star; it was born in the later stages of the universe's evolution. It contains more elements heavier than hydrogen and helium

("metals" in astronomical parlance) than older population II stars.^[31] Elements heavier than hydrogen and helium were formed in the cores of ancient and exploding stars, so the first generation of stars had to die before the universe could be enriched with these atoms. The oldest stars contain few metals, while stars born later have more. This high metallicity is thought to have been crucial to the Sun's developing a planetary system, because planets form from accretion of metals.^[32]



The Hertzsprung-Russell diagram; the main sequence is from bottom right to top left.

Interplanetary medium




The heliospheric current sheet

Along with light, the Sun radiates a continuous stream of charged particles (a plasma) known as the solar wind. This stream of particles spreads outwards at roughly 1.5 million kilometres per hour,^[33] creating a tenuous atmosphere (the heliosphere) that permeates the Solar System out to at least 100 AU (see heliopause). This is known as the interplanetary medium. The Sun's 11-year sunspot cycle and frequent solar flares and coronal mass ejections disturb the

heliosphere, creating space weather.^[34] The Sun's rotating magnetic field acts on the interplanetary medium to create the heliospheric current sheet, the largest structure in the solar system.^[35]

Earth's magnetic field protects its atmosphere from interacting with the solar wind. Venus and Mars do not have magnetic fields, and the solar wind causes their atmospheres to gradually bleed away into space.^[36] The interaction of the solar wind with Earth's magnetic field creates the aurorae seen near the magnetic poles.

Cosmic rays originate outside the Solar System. The heliosphere partially shields the Solar System, and planetary magnetic fields (for planets which have them) also provide some protection. The density of cosmic rays in the interstellar medium and the strength of the Sun's magnetic field change on very long timescales, so the level of cosmic radiation in the Solar System varies, though by how much is unknown.^[37]



Aurora australis seen from orbit.

The interplanetary medium is home to at least two disc-like regions of cosmic dust. The first, the zodiacal dust cloud, lies in the inner Solar System and causes zodiacal light. It was likely formed by collisions within the asteroid belt brought on by interactions with the planets.^[38] The second extends from about 10 AU to about 40 AU, and was probably created by similar collisions within the Kuiper belt.^{[39] [40]}

The interplanetary medium is home to at least two disc-like regions of cosmic dust. The first, the zodiacal dust cloud, lies in the inner Solar System and causes zodiacal light. It was likely formed by collisions within the asteroid belt brought on by interactions with the planets.^[38] The second extends from about 10 AU to about 40 AU, and was probably created by similar collisions within the Kuiper belt.^{[39] [40]}

Inner Solar System

The inner Solar System is the traditional name for the region comprising the terrestrial planets and asteroids. Composed mainly of silicates and metals, the objects of the inner Solar System huddle very closely to the Sun; the radius of this entire region is shorter than the distance between Jupiter and Saturn. This region was, in old parlance, denoted inner space; the area outside the asteroid belt was denoted outer space.

Inner planets

Main article: Terrestrial planet

The four inner or terrestrial planets have dense, rocky compositions, few or no moons, and no ring systems. They are composed largely of minerals with high melting points, such as the silicates which form their solid crusts and semi-liquid mantles, and metals such as iron and nickel, which form their cores. Three of the four inner planets (Venus, Earth and Mars) have substantial atmospheres; all have impact craters and tectonic surface features such as rift valleys and volcanoes. The term *inner planet* should not be confused with *inferior planet*, which designates those planets which are closer to the Sun than Earth is (i.e. Mercury and Venus).



The inner planets. From left to right: Mercury, Venus, Earth, and Mars (sizes to scale)

Mercury (0.4 AU) is the closest planet to the Sun and the smallest planet (0.055 Earth masses). Mercury has no natural satellites, and its only known geological features besides impact craters are "wrinkle-ridges", probably produced by a period of contraction early in its history.^[41] Mercury's almost negligible atmosphere consists of atoms blasted off its surface by the solar wind.^[42] Its relatively large iron core and thin mantle have not yet been adequately explained. Hypotheses include that its outer layers were stripped off by a giant impact, and that it was prevented from fully accreting by the young Sun's energy.^[43]

Mercury

[44]

Venus

Venus (0.7 AU) is close in size to Earth (0.815 Earth masses) and, like Earth, has a thick silicate mantle around an iron core, a substantial atmosphere and evidence of internal geological activity. However, it is much drier than Earth and its atmosphere is ninety times as dense. Venus has no natural satellites. It is the hottest planet, with surface temperatures over 400 °C, most likely due to the amount of greenhouse gases in the atmosphere.^[45] No definitive evidence of current geological activity has been detected on Venus, but it has no magnetic field that would prevent depletion of its substantial atmosphere, which suggests that its atmosphere is regularly replenished by volcanic eruptions.^[46]

Earth

Earth (1 AU) is the largest and densest of the inner planets, the only one known to have current geological activity, and the only planet known to have life. Its liquid hydrosphere is unique among the terrestrial planets, and it is also the only planet where plate tectonics has been observed. Earth's atmosphere is radically different from those of the other planets, having been altered by the presence of life to contain 21% free

oxygen.^[47] It has one satellite, the Moon, the only large satellite of a terrestrial planet in the Solar System.

Mars

Mars (1.5 AU) is smaller than Earth and Venus (0.107 Earth masses). It possesses a tenuous atmosphere of mostly carbon dioxide. Its surface, peppered with vast volcanoes such as Olympus Mons and rift valleys such as Valles Marineris, shows geological activity that may have persisted until very recently.^[48] Mars has two tiny natural satellites (Deimos and Phobos) thought to be captured asteroids.^[49]

Asteroid belt

Main article: Asteroid belt

Asteroids are mostly small Solar System bodies composed mainly of rocky and metallic non-volatile minerals.

The main asteroid belt occupies the orbit between Mars and Jupiter, between 2.3 and 3.3 AU from the Sun. It is thought to be remnants from the Solar System's formation that failed to coalesce because of the gravitational interference of Jupiter.

Asteroids range in size from hundreds of kilometres across to microscopic. All asteroids save the largest, Ceres, are classified as small Solar System bodies, but some asteroids such as Vesta and Hygieia may be reclassified as dwarf planets if they are shown to have achieved hydrostatic equilibrium.

The asteroid belt contains tens of thousands, possibly millions, of objects over one kilometre in diameter.^[50] Despite this, the total mass of the main belt is unlikely to be more than a thousandth of that of the Earth.^[51] The main belt is very sparsely populated; spacecraft routinely pass through without incident. Asteroids with diameters between 10 and 10^{-4} m are called meteoroids.^[52]





Image of the main asteroid belt and the Trojan asteroids



Ceres

Ceres

Ceres (2.77 AU) is the largest body in the asteroid belt and its only dwarf planet. It has a diameter of slightly under 1000 km, large enough for its own gravity to pull it into a spherical shape. Ceres was considered a planet when it was discovered in the 19th century, but was reclassified as an asteroid in the 1850s as further observation revealed additional asteroids.^[53] It was again reclassified in

2006 as a dwarf planet.

Asteroid groups

Asteroids in the main belt are divided into asteroid groups and families based on their orbital characteristics. Asteroid moons are asteroids that orbit larger asteroids. They are not as clearly distinguished as planetary moons, sometimes being almost as large as their partners. The asteroid belt also contains main-belt comets^[54] which may have been the source of Earth's water.

Trojan asteroids are located in either of Jupiter's L_4 or L_5 points (gravitationally stable regions leading and trailing a planet in its orbit); the term "Trojan" is also used for small bodies in any other planetary or satellite Lagrange point. Hilda asteroids are in a 2:3 resonance with Jupiter; that is, they go around the Sun three times for every two Jupiter orbits.


The inner Solar System is also dusted with rogue asteroids, many of which cross the orbits of the inner planets.

Mid Solar System

The middle region of the Solar System is home to the gas giants and their planet-sized satellites. Many short period comets, including the centaurs, also lie in this region. It has no traditional name; it is occasionally referred to as the "outer Solar System", although recently that term has been more often applied to the region beyond Neptune. The solid objects in this region are composed of a higher proportion of "ices" (water, ammonia, methane) than the rocky denizens of the inner Solar System.

Outer planets

The four outer planets, or gas giants (sometimes called Jovian planets), collectively make up 99 percent of the mass known to orbit the Sun. Jupiter and Saturn's atmospheres are largely hydrogen and helium. Uranus and Neptune's atmospheres have a higher percentage of "ices", such as water, ammonia and methane. Some astronomers suggest they belong in their own category, "ice giants."^[55] All four gas giants have rings, although only Saturn's ring system is easily observed from Earth. The term *outer planet* should not be confused with *superior planet*, which designates planets outside Earth's orbit (the outer planets and Mars).



From top to bottom: Neptune, Uranus, Saturn, and Jupiter (not to scale)

Jupiter

Jupiter (5.2 AU), at 318 Earth masses, masses 2.5 times all the other planets put together. It is composed largely of hydrogen and helium. Jupiter's strong internal heat creates a number of semi-permanent features in its atmosphere, such as cloud bands and the Great Red Spot. Jupiter has sixty-three known satellites. The four largest, Ganymede, Callisto, Io, and Europa, show similarities to the terrestrial planets, such as volcanism and internal heating.^[56] Ganymede, the largest satellite in the Solar System, is larger than Mercury.

Saturn

Saturn (9.5 AU), famous for its extensive ring system, has similarities to Jupiter, such as its atmospheric composition. Saturn is far less massive, being only 95 Earth masses. Saturn has sixty known satellites (and 3 unconfirmed); two of which, Titan and Enceladus, show signs of geological activity, though they are largely made of ice.^[57] Titan is larger than Mercury and the only satellite in the Solar System with a substantial atmosphere.

Uranus

Uranus (19.6 AU), at 14 Earth masses, is the lightest of the outer planets. Uniquely among the planets, it orbits the Sun on its side; its axial tilt is over ninety degrees to the ecliptic. It has a much colder core than the other gas giants, and radiates very little heat into space.^[58] Uranus has twenty-seven known satellites, the largest ones being Titania, Oberon, Umbriel, Ariel and Miranda.

Neptune

Neptune (30 AU), though slightly smaller than Uranus, is more massive (equivalent to 17 Earths) and therefore denser. It radiates more internal heat, but not as much as Jupiter or Saturn.^[59] Neptune has thirteen known satellites. The largest, Triton, is geologically active, with geysers of liquid nitrogen.^[60] Triton is the only large satellite with a retrograde orbit. Neptune is accompanied in its orbit by a number of minor planets in a 1:1 resonance with it, termed Neptune Trojans.

Comets

Main article: Comet

Comets are small Solar System bodies, usually only a few kilometres across, composed largely of volatile ices. They have highly eccentric orbits, generally a perihelion within the orbits of the inner planets and an aphelion far beyond Pluto. When a comet enters the inner Solar System, its proximity to the Sun causes its icy surface to sublimate and ionise, creating a coma: a long tail of gas and dust often visible to the naked eye.

Short-period comets have orbits lasting less than two hundred years. Long-period comets have orbits lasting thousands of years. Short-period comets are believed to originate in the Kuiper belt, while long-period comets, such as Hale-Bopp, are believed to originate in the Oort cloud. Many comet groups, such as the Kreutz Sungrazers, formed from the breakup of a single parent.^[61] Some comets with hyperbolic orbits may originate outside the Solar System, but determining their precise orbits is difficult.^[62] Old comets that have had most of their volatiles driven out by solar warming are often categorised as asteroids.^[63]



Comet Hale-Bopp

Centaur

The centaurs, which extend from 9 to 30 AU, are icy comet-like bodies that orbit in the region between Jupiter and Neptune. The largest known centaur, 10199 Chariklo, has a diameter of between 200 and 250 km.^[64] The first centaur discovered, 2060 Chiron, has been called a comet since it develops a coma just as comets do when they approach the Sun.^[65] Some astronomers classify centaurs as inward-scattered Kuiper belt objects along with the outward-scattered residents of the scattered disc.^[66]



Trans-Neptunian region

The area beyond Neptune, often called the outer Solar System or the "trans-Neptunian region", is still largely unexplored. It appears to consist overwhelmingly of small worlds (the largest having a diameter only a fifth that of the Earth and a mass far smaller than that of the Moon) composed mainly of rock and ice.

Kuiper belt

Main article: Kuiper belt

The Kuiper belt, the region's first formation, is a great ring of debris similar to the asteroid belt, but composed mainly of ice. It extends between 30 and 50 AU from the Sun. This region is thought to be the source of short-period comets. It is composed mainly of small Solar System bodies, but many of the largest Kuiper belt objects, such as Quaoar, Varuna, , and Orcus, may be reclassified as dwarf planets. There are estimated to be over 100,000 Kuiper belt objects with a diameter greater than 50 km, but the total mass of the Kuiper belt is thought to be only a tenth or even a hundredth the mass of the Earth.^[67] Many Kuiper belt objects have multiple satellites, and most have orbits that take them outside the plane of the ecliptic.



The Kuiper belt can be roughly divided into the "resonant" belt and the "classical" belt. The resonant belt consists of objects with orbits linked to that of Neptune (e.g. orbiting twice for every three Neptune orbits, or once for every two). The resonant belt actually begins within the orbit of Neptune itself. The classical belt consists of objects having no resonance with Neptune, and extends from roughly 39.4 AU to 47.7 AU.^[68] Members of the classical Kuiper belt are classified as cubewanos, after the first of their kind to be discovered, .^[69]


Pluto and Charon

Pluto (39 AU average), a dwarf planet, is the largest known object in the Kuiper belt. When discovered in 1930 it was considered to be the ninth planet; this changed in 2006 with the adoption of a formal definition of

planet. Pluto has a relatively eccentric orbit inclined 17 degrees to the ecliptic plane and ranging from 29.7 AU from the Sun at perihelion (within the orbit of Neptune) to 49.5 AU at aphelion.

It is unclear whether Charon, Pluto's largest moon, will continue to be classified as such or as a dwarf planet itself. Both Pluto and Charon orbit a barycenter of gravity above their surfaces, making Pluto-Charon a binary system. Two much smaller moons, Nix and Hydra, orbit Pluto and Charon.

Pluto lies in the resonant belt, having a 3:2 resonance with Neptune (it orbits twice round the Sun for every three Neptunian orbits). Kuiper belt objects whose orbits share this resonance are called plutinos.^[70]




Pluto and its three known moons


Scattered disc

Main article: Scattered disc

The scattered disc overlaps the Kuiper belt but extends much further outwards. Scattered disc objects are believed to come from the Kuiper belt, having been ejected into erratic orbits by the gravitational influence of Neptune's early outward migration. Most scattered disc objects (SDOs) have perihelia within the Kuiper belt but aphelia as far as 150 AU from the Sun. SDOs' orbits are also highly inclined to the ecliptic plane, and are often almost perpendicular to it. Some astronomers consider the scattered disc to be merely another region of the Kuiper belt, and describe scattered disc objects as "scattered Kuiper belt objects."^[71]



Black: scattered; blue: classical; green: resonant



*Eris and its moon
Dysnomia*


Eris

Eris (68 AU average) is the largest known scattered disc object, and caused a debate about what constitutes a planet, since it is at least 5% larger than Pluto with an estimated diameter of 2400 km (1500 mi). It is the largest of the known dwarf planets.^[72] It has one moon, Dysnomia. Like Pluto, its orbit is highly eccentric, with a perihelion of 38.2 AU (roughly Pluto's distance from the Sun) and an aphelion of 97.6 AU, and steeply inclined to the ecliptic plane.

Farthest regions

The point at which the Solar System ends and interstellar space begins is not precisely defined, since its outer boundaries are shaped by two separate forces: the solar wind and the Sun's gravity. The solar wind is believed to surrender to the interstellar medium at roughly four times Pluto's distance. However, the Sun's Roche sphere, the effective range of its gravitational influence, is believed to extend up to a thousand times farther.

Heliopause



The Voyagers entering the heliosheath

The heliosphere is divided into two separate regions. The solar wind travels at its maximum velocity out to about 95 AU, or three times the orbit of Pluto. The edge of this region is the termination shock, the point at which the solar wind collides with the opposing winds of the interstellar medium. Here the wind slows, condenses and becomes more turbulent, forming a great oval structure known as the heliosheath that looks and behaves very


much like a comet's tail, extending outward for a further 40 AU at its stellar-windward side, but tailing many times that distance in the opposite direction. The outer boundary of the heliosphere, the heliopause, is the point at which the solar wind finally terminates, and is the beginning of interstellar space.^[73]

The shape and form of the outer edge of the heliosphere is likely affected by the fluid dynamics of interactions with the interstellar medium,^[74] as well as solar magnetic fields prevailing to the south, e.g. it is bluntly shaped with the northern hemisphere extending 9 AU (roughly 900 million miles) farther than the southern hemisphere. Beyond the heliopause, at around 230 AU, lies the bow shock, a plasma "wake" left by the Sun as it travels through the Milky Way.^[75]

No spacecraft have yet passed beyond the heliopause, so it is impossible to know for certain the conditions in local interstellar space. How well the heliosphere shields the Solar System from cosmic rays is poorly understood. A dedicated mission beyond the heliosphere has been suggested.^[76] ^[77]


Oort cloud

The hypothetical Oort cloud is a great mass of up to a trillion icy objects that is believed to be the source for all long-period comets and to surround the Solar System at around 50,000 AU, and possibly to as far as 100,000 AU. It is believed to be composed of comets which were ejected from the inner Solar System by gravitational interactions with the outer planets. Oort cloud objects move very slowly, and can be perturbed by infrequent events such as collisions, the gravitational effects of a passing star, or the galactic tide.^[78] ^[79]



Sedna and the inner Oort cloud


90377 Sedna is a large, reddish Pluto-like object with a gigantic, highly elliptical orbit that takes it from about 76 AU at perihelion to 928 AU at aphelion and takes 12,050 years to complete. Mike Brown, who discovered the object in 2003, asserts that it cannot be part of the scattered disc or the Kuiper Belt as its perihelion is too distant to have been affected by Neptune's migration. He and other astronomers consider it to be the first in an entirely new population, which also may include the object , which has a perihelion of 45 AU, an aphelion of 415 AU, and an orbital period of 3420 years.^[80] Brown terms this population the "Inner Oort cloud," as it may have formed through a similar process, although it is far closer to the Sun.^[81] Sedna is very likely a dwarf planet, though its shape has yet to be determined with certainty.



Boundaries

Much of our Solar System is still unknown. The Sun's gravitational field is estimated to dominate the gravitational forces of surrounding stars out to about two light years (125,000 AU). The outer extent of the Oort cloud, by contrast, may not extend farther than 50,000 AU.^[82] Despite discoveries such as Sedna, the region between the Kuiper belt and the Oort cloud, an area tens of thousands of AU in radius, is still virtually unmapped. There are also ongoing studies of the region between Mercury and the Sun.^[83] Objects may yet be discovered in the Solar System's uncharted regions.

Galactic context



The Solar System is located in the Milky Way galaxy, a barred spiral galaxy with a diameter of about 100,000 light years containing about 200 billion stars.^[84] Our Sun resides in one of the Milky Way's outer spiral arms, known as the Orion Arm or Local Spur.^[85] The Sun lies between 25,000 and 28,000 light years from the Galactic Centre, and its speed within the galaxy is about 220 kilometres per second, so that it completes one revolution every 225-250 million years. This revolution is known as the Solar System's galactic year.^[86]

The Solar System's location in the galaxy is very likely a factor in the evolution of life on Earth. Its orbit is close to being circular and is at roughly the same speed as that of the spiral arms, which means it passes through them only rarely. Since spiral arms are home to a far larger concentration of potentially dangerous supernovae, this has given Earth long periods of interstellar stability for life to evolve.^[87] The Solar System also lies well outside the star-crowded environs of the galactic centre. Near the centre, gravitational tugs from nearby stars could perturb bodies in the Oort Cloud and send many comets into the inner Solar System, producing collisions with potentially catastrophic implications for life on Earth. The intense radiation of the galactic centre could also interfere with the development of complex life.^[88] Even at the Solar System's current location, some scientists have hypothesised that recent supernovae may have adversely affected life in the last 35,000 years by flinging pieces of expelled stellar core towards the Sun in the form of radioactive dust grains and larger, comet-like bodies.^[89]


The solar apex, the direction of the Sun's path through interstellar space, is near the constellation of Hercules in the direction of the current location of the bright star Vega.^[90]

Neighbourhood

The immediate galactic neighbourhood of the Solar System is known as the Local Interstellar Cloud or Local Fluff, an area of dense cloud in an otherwise sparse region known as the Local Bubble, an hourglass-shaped cavity in the interstellar medium roughly 300 light years across. The bubble is suffused with high-temperature plasma that suggests it is the product of several recent supernovae.^[91]

There are relatively few stars within ten light years (95 trillion km) of the Sun.

The closest is the triple star system Alpha Centauri, which is about 4.4 light years away. Alpha Centauri A and B are a closely tied pair of Sun-like stars, while the small red dwarf Alpha Centauri C (also known as Proxima Centauri) orbits the pair at a distance of 0.2 light years. The stars next closest to the Sun are the red dwarfs Barnard's Star (at 6 light years), Wolf 359 (7.8 light years) and Lalande 21185 (8.3 light years). The largest star within ten light years is Sirius, a bright main sequence star roughly twice the Sun's mass and orbited by a white dwarf called Sirius B. It lies 8.6 light years away. The remaining systems within ten light years are the binary red dwarf system Luyten 726-8 (8.7 light years) and the solitary red dwarf Ross 154 (9.7 light years).^[92] Our closest solitary sunlike star is Tau Ceti, which lies 11.9 light years away. It has roughly 80 percent the Sun's mass, but only 60 percent its luminosity.^[93] The closest earth-like planet discovered, Gliese 581c, is 20.40 light years away.



Artist's conception of the Local Bubble

Discovery and exploration

Main articles: Geocentric model and Heliocentrism

For many thousands of years, people, with a few notable exceptions, did not believe the Solar System existed. The Earth was believed not only to be stationary at the centre of the universe, but to be categorically different from the divine or ethereal objects that moved through the sky. While Nicolaus Copernicus and his predecessors, such as the Indian mathematician-astronomer Aryabhata and the Greek philosopher Aristarchus of Samos, had speculated on a heliocentric reordering of the cosmos, it was the conceptual advances of the 17th century, led by Galileo Galilei, Johannes Kepler, and Isaac Newton, which led gradually to the acceptance of the idea not only that Earth moved round the Sun, but that the planets were governed by the same physical laws that governed the Earth, and therefore could be material worlds in their own right, with such earthly phenomena as craters, weather, geology, seasons and ice caps.

Telescopic observations

Main article: Timeline of solar system astronomy

The first exploration of the Solar System was conducted by telescope, when astronomers first began to map those objects too faint to be seen with the naked eye.

Galileo Galilei was the first to discover physical details about the individual bodies of the Solar System. He discovered that the Moon was cratered, that the Sun was marked with sunspots, and that Jupiter had four satellites in orbit around it.^[94] Christiaan Huygens followed on from Galileo's discoveries by discovering Saturn's moon

Titan and the shape of the rings of Saturn.^[95] Giovanni Domenico Cassini later discovered four more moons of Saturn, the Cassini division in Saturn's rings, and the Great Red Spot of Jupiter.^[96]

Edmond Halley realised in 1705 that repeated sightings of a comet were in fact recording the same object, returning regularly once every 75–76 years. This was the first evidence that anything other than the planets orbited the Sun.^[97]

In 1781, William Herschel was looking for binary stars in the constellation of Taurus when he observed what he thought was a new comet. In fact, its orbit revealed that it was a new planet, Uranus, the first ever discovered.^[98]

Giuseppe Piazzi discovered Ceres in 1801, a small world between Mars and Jupiter that was initially considered a new planet. However, subsequent discoveries of thousands of other small worlds in the same region led to their eventual reclassification as asteroids.^[99]

By 1846, discrepancies in the orbit of Uranus led many to suspect a large planet must be tugging at it from farther out. Urbain Le Verrier's calculations eventually led to the discovery of Neptune.^[100] The excess perihelion precession of Mercury's orbit led Le Verrier to postulate the intra-Mercurian planet Vulcan in 1859 - but that would turn out to be a red herring.

Further apparent discrepancies in the orbits of the outer planets led Percival Lowell to conclude that yet another planet, "Planet X," must still be out there. After his death, his Lowell Observatory conducted a search which ultimately led to Clyde Tombaugh's discovery of Pluto in 1930. Pluto was, however, found to be too small to have disrupted the orbits of the outer planets, and its discovery was therefore coincidental. Like Ceres, it was initially considered to be a planet, but after the discovery of many other similarly sized objects in its vicinity it was reclassified in 2006 as a dwarf planet by the IAU.^[101]

In 1992, astronomers David Jewitt of the University of Hawaii and Jane Luu of the Massachusetts Institute of Technology discovered . This object proved to be the first of a new population, which came to be known as the Kuiper



A replica of Isaac Newton's telescope

belt; an icy analogue to the asteroid belt of which such objects as Pluto and Charon were deemed a part.^[102] ^[103]


Mike Brown, Chad Trujillo and David Rabinowitz announced the discovery of Eris in 2005, a scattered disc object larger than Pluto and the largest object discovered in orbit round the Sun since Neptune.^[104]

Observations by spacecraft

Since the start of the Space Age, a great deal of exploration has been performed by robotic spacecraft missions that have been organized and executed by various space agencies.

All planets in the Solar System have now been visited to varying degrees by spacecraft launched from Earth. Through these unmanned missions, humans have been able to get close-up photographs of all of the planets and, in the case of landers, perform tests of the soils and atmospheres of some.


The first manmade object sent into space was the Soviet satellite *Sputnik 1*, launched in 1957, which successfully orbited the Earth for over a year. The American probe *Explorer 6*, launched in 1959, was the first satellite to image the Earth from space.



Artist's conception of Pioneer 10, which passed the orbit of Pluto in 1983. The last transmission was received in January 2003, sent from approximately 82 AU away. The 35-year-old space probe is now receding at over 43,400 km/h (27,000 mph) from the Sun.^[105]

Flybys

The first successful probe to fly by another Solar System body was *Luna 1*, which sped past the Moon in 1959. Originally meant to impact with the Moon, it instead missed its target and became the first manmade object to orbit the Sun. *Mariner 2* was the first probe to fly by another planet, Venus, in 1962. The first successful flyby of Mars was made by *Mariner 4* in 1964. Mercury was first encountered by *Mariner 10* in 1974.



A photo of Earth (circled) taken by Voyager 1, 6 billion km (4 billion miles) away. The streaks of light are diffraction spikes radiating from the Sun (off frame to the left).

The first probe to explore the outer planets was *Pioneer 10*, which flew by Jupiter in 1973. *Pioneer 11* was the first to visit Saturn, in 1979. The *Voyager* probes performed a grand tour of the outer planets following their launch in 1977, with both probes passing Jupiter in 1979 and Saturn in 1980 - 1981. *Voyager 2* then went on to make close approaches to Uranus in 1986 and Neptune in 1989. The *Voyager* probes are now far beyond Neptune's orbit, and are on course to find and study the termination shock, heliosheath, and heliopause. According to NASA, both *Voyager* probes have encountered the termination shock at a distance of approximately 93 AU from the Sun.^{[106] [107]}

The first flyby of a comet occurred in 1985, when the International Cometary Explorer (ICE) passed by the comet Giacobini-Zinner,^[108] while the first flybys of asteroids were conducted by the *Galileo* spacecraft, which imaged both 951 Gaspra (in 1991) and 243 Ida (in 1993) on its way to Jupiter.

No Kuiper belt object has yet been visited by a spacecraft. Launched on January 19 2006, the *New Horizons* probe is currently en route to becoming the first man-made spacecraft to explore this area. This unmanned mission is scheduled to fly by Pluto in July 2015. Should it prove feasible, the mission will then be extended to observe a number of other Kuiper belt objects.^[109]

Orbiters, landers and rovers

In 1966, the Moon became the first Solar System body beyond Earth to be orbited by an artificial satellite (*Luna 10*), followed by Mars in 1971 (*Mariner 9*), Venus in 1975 (*Venera 9*), Jupiter in 1995 (*Galileo*), the asteroid 433 Eros in 2000 (*NEAR Shoemaker*), and Saturn in 2004 (*Cassini-Huygens*). The MESSENGER probe is currently en route to commence the first orbit of Mercury in 2011, while the *Dawn* spacecraft is currently set to orbit the asteroid Vesta in 2011 and the dwarf planet Ceres in 2015.

The first probe to land on another Solar System body was the Soviet *Luna 2* probe, which impacted the Moon in 1959. Since then, increasingly distant planets have been reached, with probes landing on or impacting the surfaces of Venus in 1966 (*Venera 3*), Mars in 1971 (*Mars 3*, although a fully successful landing didn't occur until *Viking 1* in 1976), the asteroid 433 Eros in 2001 (*NEAR Shoemaker*), and Saturn's moon Titan (*Huygens*) and the comet Tempel 1 (*Deep Impact*) in 2005. The *Galileo* orbiter also dropped a

probe into Jupiter's atmosphere in 1995; since Jupiter has no physical surface, it was destroyed by increasing temperature and pressure as it descended.

To date, only two worlds in the Solar System, the Moon and Mars, have been visited by mobile rovers. The first rover to visit another celestial body was the Soviet *Lunokhod 1*, which landed on the Moon in 1970. The first to visit another planet was Sojourner, which travelled 500 metres across the surface of Mars in 1997. The only manned rover to visit another world was NASA's Lunar rover, which travelled with Apollos 15, 16 and 17 between 1971 and 1972.

Manned exploration

Manned exploration of the Solar System is currently confined to Earth's immediate environs. The first human being to reach space (defined as an altitude of over 100 km) and to orbit the Earth was Yuri Gagarin, a Soviet cosmonaut who was launched in *Vostok 1* on April 12, 1961. The first man to walk on the surface of another Solar System body was Neil Armstrong, who stepped onto the Moon on July 21, 1969 during the *Apollo 11* mission. The United States' Space Shuttle is the only reusable spacecraft to successfully make multiple orbital flights. The first orbital space station to host more than one crew was NASA's Skylab, which successfully held three crews from 1973 to 1974. The first true human settlement in space was the Soviet space station Mir, which was continuously occupied for close to ten years, from 1989 to 1999. It was decommissioned in 2001, and its successor, the International Space Station, has maintained a continuous human presence in space since then. In 2004, SpaceShipOne became the first privately funded vehicle to reach space on a suborbital flight. That same year, President George W. Bush announced the Vision for Space Exploration, which called for a replacement for the aging Shuttle, a return to the Moon and, ultimately, a manned mission to Mars.

See also

- List of Solar System objects:
 - By orbit
 - By mass
 - By radius
 - By name
 - By surface gravity
- Attributes of the largest solar system bodies
- Astronomical symbols
- Geological features of the solar system
- Numerical model of solar system
- Table of planetary attributes
- Timeline of discovery of Solar System planets and their natural satellites

- Solar system model
- Space colonization
- Solar System in fiction
- Celestia - Space-simulation on your computer (OpenGL)
- Family Portrait (Voyager)
- The Parable of the Solar System Model

Notes

1. Capitalization of the name varies. The IAU, the authoritative body regarding astronomical nomenclature, specifies capitalizing the names of all individual astronomical objects (**Solar System**). However, the name is commonly rendered in lower case (**solar system**) including in the *Oxford English Dictionary*, *Merriam-Webster's 11th Collegiate Dictionary*, and *Encyclopædia Britannica*.
2. The mass of the Solar System excluding the Sun, Jupiter and Saturn can be determined by adding together all the calculated masses for its largest objects and using rough calculations for the masses of the Oort cloud (estimated at roughly 3 Earth masses),^[110] the Kuiper Belt (estimated at roughly 0.1 Earth mass)^[111] and the asteroid belt (estimated to be 0.0005 Earth mass)^[112] for a total, rounded upwards, of ~37 Earth masses, or 8.9 percent the combined mass of Jupiter and Saturn.

References

- [1] Scott S. Sheppard. The Jupiter Satellite Page. *University of Hawaii*. Retrieved on 2006-07-23.
- [2] Akwagyiram, Alexis (2005-08-02). Farewell Pluto?. BBC News. Retrieved on 2006-03-05.
- [3] " The Final IAU Resolution on the definition of "planet" ready for voting", *IAU*, 2006-08-24. Retrieved on 2007-03-02.
- [4] nineplanets.org. An Overview of the Solar System. Retrieved on 2007-02-15.
- [5] Amir Alexander (2006). New Horizons Set to Launch on 9-Year Voyage to Pluto and the Kuiper Belt. *The Planetary Society*. Retrieved on 2006-11-08.
- [6] M Woolfson. The origin and evolution of the solar system (PDF). *University of York*. Retrieved on 2006-07-22.
- [7] See, T. J. J. (April 23 1909). " The Past History of the Earth as Inferred from the Mode of Formation of the Solar System". *Proceedings of the American Philosophical Society* **48** (191): 119-128. Retrieved on 2006-07-23.
- [8] Lecture 13: The Nebular Theory of the origin of the Solar System. *University of Arizona*. Retrieved on 2006-12-27.

- [9] Jeff Hester (2004). New Theory Proposed for Solar System Formation. *Arizona State University*. Retrieved on 2007-01-11.
- [10] Irvine, W. M.. The chemical composition of the pre-solar nebula. *Amherst College, Massachusetts*. Retrieved on 2007-02-15.
- [11] Yoshimi Kitamura, Munetake Momose, Sozo Yokogawa, Ryohei Kawabe, Shigeru Ida and Motohide Tamura (2002-12-10). " Investigation of the Physical Properties of Protoplanetary Disks around T Tauri Stars by a 1 Arcsecond Imaging Survey: Evolution and Diversity of the Disks in Their Accretion Stage". *The Astrophysical Journal* **581** (1): 357-380. DOI: 10.1086/344223. Retrieved on 2007-01-09.
- [12] Present Understanding of the Origin of Planetary Systems. National Academy of Sciences (April 5, 2000). Retrieved on 2007-01-19.
- [13] Manfred Küker, Thomas Henning and Günther Rüdiger (2003). Magnetic Star-Disk Coupling in Classical T Tauri Systems. *Science Magazine*. Retrieved on 2006-11-16.
- [14] Chrysostomou and Phil W Lucas The formation of stars. *Department of Physics Astronomy & Mathematics University of Hertfordshire*. Retrieved on 2007-05-02.
- [15] Peter Goldreich and William R. Ward (1973). The Formation of Planetesimals. *The American Astronomical Society*. Retrieved on 2006-11-16.
- [16] Jean-Marc Petit and Alessandro Morbidelli (2001). The Primordial Excitation and Clearing of the Asteroid Belt (PDF). *Centre National de la Recherche Scientifique, Observatoire de Nice*. Retrieved on 2006-11-19.
- [17] Mummma, M. J., M. A. DiSanti, N. Dello Russo, K. Magee-Sauer, E. Gibb, and R. Novak (June 2003). " Remote infrared observations of parent volatiles in comets: A window on the early solar system" (PDF). *Advances in Space Research* **31** (12): 2563-2575. DOI: 10.1016/S0273-1177(03)00578-7. Retrieved on 2006-11-16.
- [18] Edward W. Thommes, Martin J. Duncan and Harold F. Levison. The formation of Uranus and Neptune in the Jupiter-Saturn region of the Solar System. *Department of Physics, Queen's University, Kingston, Ontario; Space Studies Department, Southwest Research Institute, Boulder, Colorado*. Retrieved on 2007-04-02.
- [19] Elmegreen, B. G. (November 1979). " On the disruption of a protoplanetary disk nebula by a T Tauri like solar wind" (PDF). *Astronomy and Astrophysics* **80** (1): 77-78. Retrieved on 2007-02-11.
- [20] Heng Hao (November 1979). " Disc-Protoplanet interactions" (PDF). *Astronomy and Astrophysics* **80** (1): 77-78. Retrieved on 2006-11-19.
- [21] Smart, R. L.; Carollo, D.; Lattanzi, M. G.; McLean, B.; Spagna, A. (2001). The Second Guide Star Catalogue and Cool Stars. *Perkins Observatory*. Retrieved on 2006-12-26.
- [22] Kasting, J.F., Ackerman, T.P. (1986). "Climatic Consequences of Very High Carbon Dioxide Levels in the Earth's Early Atmosphere". *Science* **234**: 1383-1385.

- [23] Richard W. Pogge (1997). The Once and Future Sun. *Perkins Observatory*. Retrieved on 2006-06-23.
- [24] T. S. van Albada, Norman Baker (1973). "On the Two Oosterhoff Groups of Globular Clusters". *Astrophysical Journal* **185**: 477-498.
- [25] Charles H. Lineweaver (2000). An Estimate of the Age Distribution of Terrestrial Planets in the Universe: Quantifying Metallicity as a Selection Effect. *University of New South Wales*. Retrieved on 2006-07-23.
- [26] Solar Physics: The Solar Wind. *Marshall Space Flight Center* (2006). Retrieved on 2006-10-03.
- [27] Phillips, Tony (2001-02-15). The Sun Does a Flip. *Science@NASA*. Retrieved on 2007-02-04.
- [28] Artist's Conception of the Heliospheric Current Sheet. *Wilcox Solar Observatory*. Retrieved on 2006-06-22.
- [29] Langner, U. W., M.S. Potgieter (2005). " Effects of the position of the solar wind termination shock and the heliopause on the heliospheric modulation of cosmic rays". *Advances in Space Research* **35** (12): 2084-2090. DOI: 10.1016/j.asr.2004.12.005. Retrieved on 2007-02-11.
- [30] Long-term Evolution of the Zodiacal Cloud (1998). Retrieved on 2007-02-03.
- [31] ESA scientist discovers a way to shortlist stars that might have planets. *ESA Science and Technology* (2003). Retrieved on 2007-02-03.
- [32] Landgraf, M., Liou, J.-C.; Zook, H. A.; Grün, E. (May 2002). " Origins of Solar System Dust beyond Jupiter". *The Astronomical Journal* **123** (5): 2857-2861. DOI: 10.1086/339704. Retrieved on 2007-02-09.
- [33] Schenk P., Melosh H.J. (1994), *Lobate Thrust Scarps and the Thickness of Mercury's Lithosphere*, Abstracts of the 25th Lunar and Planetary Science Conference, 1994LPI....25.1203S
- [34] Bill Arnett (2006). Mercury. *The Nine Planets*. Retrieved on 2006-09-14.
- [35] Benz, W., Slattery, W. L., Cameron, A. G. W. (1988), *Collisional stripping of Mercury's mantle*, *Icarus*, v. 74, p. 516-528.
- [36] Cameron, A. G. W. (1985), *The partial volatilization of Mercury*, *Icarus*, v. 64, p. 285-294.
- [37] Mark Alan Bullock (1997). " *The Stability of Climate on Venus*" (PDF). Southwest Research Institute. Retrieved on 2006-12-26.
- [38] Paul Rincon (1999). Climate Change as a Regulator of Tectonics on Venus (PDF). *Johnson Space Center Houston, TX, Institute of Meteoritics, University of New Mexico, Albuquerque, NM*. Retrieved on 2006-11-19.
- [39] Anne E. Egger, M.A./M.S.. Earth's Atmosphere: Composition and Structure. *VisionLearning.com*. Retrieved on 2006-12-26.
- [40] David Noever (2004). Modern Martian Marvels: Volcanoes?. *NASA Astrobiology Magazine*. Retrieved on 2006-07-23.

- [41] Scott S. Sheppard, David Jewitt, and Jan Kleyna (2004). A Survey for Outer Satellites of Mars: Limits to Completeness. *The Astronomical Journal*. Retrieved on 2006-12-26.
- [42] New study reveals twice as many asteroids as previously believed. *ESA* (2002). Retrieved on 2006-06-23.
- [43] Krasinsky, G. A., Pitjeva, E. V.; Vasilyev, M. V.; Yagudina, E. I. (July 2002). "Hidden Mass in the Asteroid Belt". *Icarus* **158** (1): 98-105. DOI: 10.1006/icar.2002.6837.
- [44] Beech, M., Duncan I. Steel (September 1995). "On the Definition of the Term Meteoroid". *Quarterly Journal of the Royal Astronomical Society* **36** (3): 281-284. Retrieved on 2006-08-31.
- [45] History and Discovery of Asteroids (DOC). *NASA*. Retrieved on 2006-08-29.
- [46] Phil Berardelli (2006). Main-Belt Comets May Have Been Source Of Earths Water. *SpaceDaily*. Retrieved on 2006-06-23.
- [47] Jack J. Lissauer, David J. Stevenson (2006). Formation of Giant Planets (PDF). *NASA Ames Research Center; California Institute of Technology*. Retrieved on 2006-01-16.
- [48] Pappalardo, R T (1999). Geology of the Icy Galilean Satellites: A Framework for Compositional Studies. *Brown University*. Retrieved on 2006-01-16.
- [49] J. S. Kargel (1994). Cryovolcanism on the icy satellites. *U.S. Geological Survey*. Retrieved on 2006-01-16.
- [50] Hawksett, David; Longstaff, Alan; Cooper, Keith; Clark, Stuart (2005). 10 Mysteries of the Solar System. *Astronomy Now*. Retrieved on 2006-01-16.
- [51] Podolak, M.; Reynolds, R. T.; Young, R. (1990). Post Voyager comparisons of the interiors of Uranus and Neptune. *NASA, Ames Research Center*. Retrieved on 2006-01-16.
- [52] Duxbury, N.S., Brown, R.H. (1995). The Plausibility of Boiling Geysers on Triton. *Beacon eSpace*. Retrieved on 2006-01-16.
- [53] Sekanina, Zdenek (2001). "Kreutz sungrazers: the ultimate case of cometary fragmentation and disintegration?". *Publications of the Astronomical Institute of the Academy of Sciences of the Czech Republic* **89 p. 78-93**.
- [54] Królikowska, M. (2001). "A study of the original orbits of hyperbolic comets". *Astronomy & Astrophysics* **376** (1): 316-324. DOI: 10.1051/0004-6361:20010945. Retrieved on 2007-01-02.
- [55] Fred L. Whipple (04/1992). The activities of comets related to their aging and origin. Retrieved on 2006-12-26.
- [56] Stansberry (2005). TNO/Centaur diameters and albedos. Retrieved on 2006-11-08.
- [57] Patrick Vanouplines (1995). Chiron biography. *Vrije Universiteit Brussel*. Retrieved on 2006-06-23.

- [58] List Of Centaurs and Scattered-Disk Objects. *IAU: Minor Planet Center*. Retrieved on 2007-04-02.
- [59] Audrey Delsanti and David Jewitt (2006). The Solar System Beyond The Planets (PDF). *Institute for Astronomy, University of Hawaii*. Retrieved on 2007-01-03.
- [60] M. W. Buie, R. L. Millis, L. H. Wasserman, J. L. Elliot, S. D. Kern, K. B. Clancy, E. I. Chiang, A. B. Jordan, K. J. Meech, R. M. Wagner, D. E. Trilling (2005). Procedures, Resources and Selected Results of the Deep Ecliptic Survey. *Lowell Observatory, University of Pennsylvania, Large Binocular Telescope Observatory, Massachusetts Institute of Technology, University of Hawaii, University of California at Berkeley*. Retrieved on 2006-09-07.
- [61] E. Dotto¹, M.A. Barucci², and M. Fulchignoni (2006-08-24). Beyond Neptune, the new frontier of the Solar System (PDF). Retrieved on 2006-12-26.
- [63] David Jewitt (2005). The 1000 km Scale KBOs. *University of Hawaii*. Retrieved on 2006-07-16.
- [64] Mike Brown (2005). The discovery of 2003 UB313 Eris, the 10th planet largest known dwarf planet.. *CalTech*. Retrieved on 2006-09-15.
- [65] Voyager Enters Solar System's Final Frontier. *NASA*. Retrieved on 2007-04-02.
- [66] Fahr, H. J.; Kausch, T.; Scherer, H. (2000). A 5-fluid hydrodynamic approach to model the Solar System-interstellar medium interaction. *Institut für Astrophysik und Extraterrestrische Forschung der Universität Bonn*. Retrieved on 2006-06-23.
- [67] P. C. Frisch (2002). The Sun's Heliosphere & Heliopause. *University of Chicago*. Retrieved on 2006-06-23.
- [68] R. L. McNutt, Jr. et al. (2006). " Innovative Interstellar Explorer". *AIP Conference Proceedings* **858**: 341-347.
- [69] Interstellar space, and step on it!. *New Scientist* (2007-01-05). Retrieved on 2007-02-05.
- [70] Stern SA, Weissman PR. (2001). Rapid collisional evolution of comets during the formation of the Oort cloud.. *Space Studies Department, Southwest Research Institute, Boulder, Colorado*. Retrieved on 2006-11-19.
- [71] Bill Arnett (2006). The Kuiper Belt and the Oort Cloud. *nineplanets.org*. Retrieved on 2006-06-23.
- [72] David Jewitt (2004). Sedna - 2003 VB₁₂. *University of Hawaii*. Retrieved on 2006-06-23.
- [73] Mike Brown. Sedna. *CalTech*. Retrieved on 2007-05-02.
- [74] T. Encrenaz, JP. Bibring, M. Blanc, MA. Barucci, F. Roques, PH. Zarka; [] (2004). "[]" *The Solar System: Third edition* (),, (in), , : Springer.. URL accessed during . ""

- [75] Durda D.D.; Stern S.A.; Colwell W.B.; Parker J.W.; Levison H.F.; Hassler D.M. (2004). A New Observational Search for Vulcanoids in SOHO/LASCO Coronagraph Images. Retrieved on 2006-07-23.
- [76] A.D. Dolgov (2003). Magnetic fields in cosmology. Retrieved on 2006-07-23.
- [77] R. Drimmel, D. N. Spergel (2001). Three Dimensional Structure of the Milky Way Disk. Retrieved on 2006-07-23.
- [78] Leong, Stacy (2002). Period of the Sun's Orbit around the Galaxy (Cosmic Year). *The Physics Factbook*. Retrieved on 2007-04-02.
- [79] Leslie Mullen (2001). Galactic Habitable Zones. *Astrobiology Magazine*. Retrieved on 2006-06-23.
- [80] Supernova Explosion May Have Caused Mammoth Extinction. *Physorg.com* (2005). Retrieved on 2007-02-02.
- [81] C. Barbieri (2003). Elementi di Astronomia e Astrofisica per il Corso di Ingegneria Aerospaziale V settimana. *IdealStars.com*. Retrieved on 2007-02-12.
- [82] Near-Earth Supernovas. *NASA*. Retrieved on 2006-07-23.
- [83] Stars within 10 light years. *SolStation*. Retrieved on 2007-04-02.
- [84] Tau Ceti. *SolStation*. Retrieved on 2007-04-02.
- [85] Eric W. Weisstein (2006). Galileo Galilei (1564-1642). *Wolfram Research*. Retrieved on 2006-11-08.
- [86] Discoverer of Titan: Christiaan Huygens. *ESA Space Science* (2005). Retrieved on 2006-11-08.
- [87] Giovanni Domenico Cassini (June 8, 1625-September 14, 1712). *SEDS.org*. Retrieved on 2006-11-08.
- [88] Comet Halley. *University of Tennessee*. Retrieved on 2006-12-27.
- [89] Herschel, Sir William (1738-1822). *enotes.com*. Retrieved on 2006-11-08.
- [90] Discovery of Ceres: 2nd Centenary, 1 January 1801-1 January 2001. *astropa.unipa.it* (2000). Retrieved on 2006-11-08.
- [91] J. J. O'Connor and E. F. Robertson (1996). Mathematical discovery of planets. *St. Andrews University*. Retrieved on 2006-11-08.
- [92] Jane X. Luu -and David C. Jewitt - (2002). KUIPER BELT OBJECTS: Relics from the Accretion Disk of the Sun. *MIT, University of Hawaii*. Retrieved on 2006-11-09.
- [93] Minor Planet Center. List of Trans-Neptunian Objects. Retrieved on 2007-04-02.
- [94] Eris (2003 UB313). *Solstation.com* (2006). Retrieved on 2006-11-09.
- [95] Donald Savage; Michael Mewhinney (February 25, 2003). Farewell Pioneer 10. *NASA*. Retrieved on 2007-07-11.
- [96] Randy Culp (2002). Time Line of Space Exploration. Retrieved on 2006-07-01.
- [97] Comet Space Missions Accessed 2007-10-23

- [98] New Horizons NASA's Pluto-Kuiper Belt Mission (2006). Retrieved on 2006-07-01.
- [99] Alessandro Morbidelli (2006). ORIGIN AND DYNAMICAL EVOLUTION OF COMETS AND THEIR RESERVOIRS (PDF). *CNRS, Observatoire de la Côte d'Azur*. Retrieved on 2007-08-03.

External links

- Solar System Profile by NASA's Solar System Exploration
- NASA's Solar System Simulator
- NASA/JPL Solar System main page
- The Nine Planets - Comprehensive Solar System site by Bill Arnett
- SPACE.com: All About the Solar System
- Illustration of the distance between planets
- Illustration comparing the sizes of the planets with each other, the sun, and other stars

GNU Free Documentation License

Version 1.2, November 2002

Copyright (C) 2000,2001,2002 Free Software Foundation, Inc.
51 Franklin St, Fifth Floor, Boston, MA 02110-1301 USA
Everyone is permitted to copy and distribute verbatim copies
of this license document, but changing it is not allowed.

0. PREAMBLE

The purpose of this License is to make a manual, textbook, or other functional and useful document "free" in the sense of freedom: to assure everyone the effective freedom to copy and redistribute it, with or without modifying it, either commercially or noncommercially. Secondly, this License preserves for the author and publisher a way to get credit for their work, while not being considered responsible for modifications made by others.

This License is a kind of "copyleft", which means that derivative works of the document must themselves be free in the same sense. It complements the GNU General Public License, which is a copyleft license designed for free software.

We have designed this License in order to use it for manuals for free software, because free software needs free documentation: a free program should come with manuals providing the same freedoms that the software does. But this License is not limited to software manuals; it can be used for any textual work, regardless of subject matter or whether it is published as a printed book. We recommend this License principally for works whose purpose is instruction or reference.

1. APPLICABILITY AND DEFINITIONS

This License applies to any manual or other work, in any medium, that contains a notice placed by the copyright holder saying it can be distributed under the terms of this License. Such a notice grants a world-wide, royalty-free license, unlimited in duration, to use that work under the conditions stated herein. The "Document", below, refers to any such manual or work. Any member of the public is a licensee, and is addressed as "you". You accept the license if you copy, modify or distribute the work in a way requiring permission under copyright law.

A "Modified Version" of the Document means any work containing the Document or a portion of it, either copied verbatim, or with modifications and/or translated into another language.

A "Secondary Section" is a named appendix or a front-matter section of the Document that deals exclusively with the relationship of the publishers or authors of the Document to the Document's overall subject (or to related matters) and contains nothing that could fall directly within that overall subject. (Thus, if the Document is in part a textbook of mathematics, a Secondary Section may not explain any mathematics.) The relationship could be a matter of historical connection with the subject or with related matters, or of legal, commercial, philosophical, ethical or political position regarding them.

The "Invariant Sections" are certain Secondary Sections whose titles are designated, as being those of Invariant Sections, in the notice that says that the Document is released under this License. If a section does not fit the above definition of Secondary then it is not allowed to be designated as Invariant. The Document may contain zero Invariant Sections. If the Document does not identify any Invariant Sections then there are none.

The "Cover Texts" are certain short passages of text that are listed, as Front-Cover Texts or Back-Cover Texts, in the notice that says that the Document is released under this License. A Front-Cover Text may be at most 5 words, and a Back-Cover Text may be at most 25 words.

A "Transparent" copy of the Document means a machine-readable copy, represented in a format whose specification is available to the general public, that is suitable for revising the document straightforwardly with generic text editors or (for images composed of pixels) generic paint programs or (for drawings) some widely available drawing editor, and that is suitable for input to text formatters or for automatic translation to a variety of formats suitable for input to text formatters. A copy made in an otherwise Transparent file format whose markup, or absence of markup, has been arranged to thwart or discourage subsequent modification by readers is not Transparent. An image format is not Transparent if used for any substantial amount of text. A copy that is not "Transparent" is called "Opaque".

Examples of suitable formats for Transparent copies include plain ASCII without markup, Texinfo input format, LaTeX input format, SGML or XML using a publicly available DTD, and standard-conforming simple HTML, PostScript or PDF designed for human modification. Examples of transparent image formats include PNG, XCF and JPG. Opaque formats include proprietary formats that can be read and edited only by proprietary word processors, SGML or XML for which the DTD and/or processing tools are not generally available, and the machine-generated HTML, PostScript or PDF produced by some word processors for output purposes only.

The "Title Page" means, for a printed book, the title page itself, plus such following pages as are needed to hold, legibly, the material this License requires to appear in the title page. For works in formats which do not have any title page as such, "Title Page" means the text near the most prominent appearance of the work's title, preceding the beginning of the body of the text.

A section "Entitled XYZ" means a named subunit of the Document whose title either is precisely XYZ or contains XYZ in parentheses following text that translates XYZ in another language. (Here XYZ stands for a specific section name mentioned below, such as "Acknowledgements", "Dedications", "Endorsements", or "History".) To "Preserve the Title" of such a section when you modify the Document means that it remains a section "Entitled XYZ" according to this definition.

The Document may include Warranty Disclaimers next to the notice which states that this License applies to the Document. These Warranty Disclaimers are considered to be included by reference in this License, but only as regards disclaiming warranties: any other implication that these Warranty Disclaimers may have is void and has no effect on the meaning of this License.

2. VERBATIM COPYING

You may copy and distribute the Document in any medium, either commercially or noncommercially, provided that this License, the copyright notices, and the license notice saying this License applies to the Document are reproduced in all copies, and that you add no other conditions whatsoever to those of this License. You may not use technical measures to obstruct or control the reading or further copying of the copies you make or distribute. However, you may accept compensation in exchange for copies. If you distribute a large enough number of copies you must also follow the conditions in section 3.

You may also lend copies, under the same conditions stated above, and you may publicly display copies.

3. COPYING IN QUANTITY

If you publish printed copies (or copies in media that commonly have printed covers) of the Document, numbering more than 100, and the Document's license notice requires Cover Texts, you must enclose the copies in covers that carry, clearly and legibly, all these Cover Texts: Front-Cover Texts on the front cover, and Back-Cover Texts on the back cover. Both covers must also clearly and legibly identify you as the publisher of these copies. The front cover must present the full title with all words of the title equally prominent and visible. You may add other material on the covers in addition. Copying with changes limited to the covers, as long as they preserve the title of the Document and satisfy these conditions, can be treated as verbatim copying in other respects.

If the required texts for either cover are too voluminous to fit legibly, you should put the first ones listed (as many as fit reasonably) on the actual cover, and continue the rest onto adjacent pages.

If you publish or distribute Opaque copies of the Document numbering more than 100, you must either include a machine-readable Transparent copy along with each Opaque copy, or state in or with each Opaque copy a computer-network location from which the general network-using public has access to download using public-standard network protocols a complete Transparent copy of the Document, free of added material. If you use the latter option, you must take reasonably prudent steps, when you begin distribution of Opaque copies in quantity, to ensure that this Transparent copy will remain thus accessible at the stated location until at least one year after the last time you distribute an Opaque copy (directly or through your agents or retailers) of that edition to the public.

It is requested, but not required, that you contact the authors of the Document well before redistributing any large number of copies, to give them a chance to provide you with an updated version of the Document.

4. MODIFICATIONS

You may copy and distribute a Modified Version of the Document under the conditions of sections 2 and 3 above, provided that you release the Modified Version under precisely this License, with the Modified Version filling the role of the Document, thus licensing distribution and modification of the Modified Version to whoever possesses a copy of it. In addition, you must do these things in the Modified Version:

- A. Use in the Title Page (and on the covers, if any) a title distinct from that of the Document, and from those of previous versions (which should, if there were any, be listed in the History section of the Document). You may use the same title as a previous version if the original publisher of that version gives permission.
- B. List on the Title Page, as authors, one or more persons or entities responsible for authorship of the modifications in the Modified Version, together with at least five of the principal authors of the Document (all of its principal authors, if it has fewer than five), unless they release you from this requirement.

- C. State on the Title page the name of the publisher of the Modified Version, as the publisher.
- D. Preserve all the copyright notices of the Document.
- E. Add an appropriate copyright notice for your modifications adjacent to the other copyright notices.
- F. Include, immediately after the copyright notices, a license notice giving the public permission to use the Modified Version under the terms of this License, in the form shown in the Addendum below.
- G. Preserve in that license notice the full lists of Invariant Sections and required Cover Texts given in the Document's license notice.
- H. Include an unaltered copy of this License.
- I. Preserve the section Entitled "History", Preserve its Title, and add to it an item stating at least the title, year, new authors, and publisher of the Modified Version as given on the Title Page. If there is no section Entitled "History" in the Document, create one stating the title, year, authors, and publisher of the Document as given on its Title Page, then add an item describing the Modified Version as stated in the previous sentence.
- J. Preserve the network location, if any, given in the Document for public access to a Transparent copy of the Document, and likewise the network locations given in the Document for previous versions it was based on. These may be placed in the "History" section. You may omit a network location for a work that was published at least four years before the Document itself, or if the original publisher of the version it refers to gives permission.
- K. For any section Entitled "Acknowledgements" or "Dedications", Preserve the Title of the section, and preserve in the section all the substance and tone of each of the contributor acknowledgements and/or dedications given therein.
- L. Preserve all the Invariant Sections of the Document, unaltered in their text and in their titles. Section numbers or the equivalent are not considered part of the section titles.
- M. Delete any section Entitled "Endorsements". Such a section may not be included in the Modified Version.
- N. Do not retitle any existing section to be Entitled "Endorsements" or to conflict in title with any Invariant Section.
- O. Preserve any Warranty Disclaimers.

If the Modified Version includes new front-matter sections or appendices that qualify as Secondary Sections and contain no material copied from the Document, you may at your option designate some or all of these sections as invariant. To do this, add their titles to the list of Invariant Sections in the Modified Version's license notice. These titles must be distinct from any other section titles.

You may add a section Entitled "Endorsements", provided it contains nothing but endorsements of your Modified Version by various parties--for example, statements of peer review or that the text has been approved by an organization as the authoritative definition of a standard.

You may add a passage of up to five words as a Front-Cover Text, and a passage of up to 25 words as a Back-Cover Text, to the end of the list of Cover Texts in the Modified Version. Only one passage of Front-Cover Text and one of Back-Cover Text may be added by (or through arrangements made by) any one entity. If the Document already includes a cover text for the same cover, previously added by you or by arrangement made by the same entity you are acting on behalf of, you may not add another; but you may replace the old one, on explicit permission from the previous publisher that added the old one.

The author(s) and publisher(s) of the Document do not by this License give permission to use their names for publicity for or to assert or imply endorsement of any Modified Version.

5. COMBINING DOCUMENTS

You may combine the Document with other documents released under this License, under the terms defined in section 4 above for modified versions, provided that you include in the combination all of the Invariant Sections of all of the original documents, unmodified, and list them all as Invariant Sections of your combined work in its license notice, and that you preserve all their Warranty Disclaimers.

The combined work need only contain one copy of this License, and multiple identical Invariant Sections may be replaced with a single copy. If there are multiple Invariant Sections with the same name but different contents, make the title of each such section unique by adding at the end of it, in parentheses, the name of the original author or publisher of that section if known, or else a unique number. Make the same adjustment to the section titles in the list of Invariant Sections in the license notice of the combined work.

In the combination, you must combine any sections Entitled "History" in the various original documents, forming one section Entitled "History"; likewise combine any sections Entitled "Acknowledgements", and any sections Entitled "Dedications". You must delete all sections

Entitled "Endorsements."

6. COLLECTIONS OF DOCUMENTS

You may make a collection consisting of the Document and other documents released under this License, and replace the individual copies of this License in the various documents with a single copy that is included in the collection, provided that you follow the rules of this License for verbatim copying of each of the documents in all other respects.

You may extract a single document from such a collection, and distribute it individually under this License, provided you insert a copy of this License into the extracted document, and follow this License in all other respects regarding verbatim copying of that document.

7. AGGREGATION WITH INDEPENDENT WORKS

A compilation of the Document or its derivatives with other separate and independent documents or works, in or on a volume of a storage or distribution medium, is called an "aggregate" if the copyright resulting from the compilation is not used to limit the legal rights of the compilation's users beyond what the individual works permit. When the Document is included in an aggregate, this License does not apply to the other works in the aggregate which are not themselves derivative works of the Document.

If the Cover Text requirement of section 3 is applicable to these copies of the Document, then if the Document is less than one half of the entire aggregate, the Document's Cover Texts may be placed on covers that bracket the Document within the aggregate, or the electronic equivalent of covers if the Document is in electronic form. Otherwise they must appear on printed covers that bracket the whole aggregate.

8. TRANSLATION

Translation is considered a kind of modification, so you may distribute translations of the Document under the terms of section 4. Replacing Invariant Sections with translations requires special permission from their copyright holders, but you may include translations of some or all Invariant Sections in addition to the original versions of these Invariant Sections. You may include a translation of this License, and all the license notices in the Document, and any Warranty Disclaimers, provided that you also include the original English version of this License and the original versions of those notices and disclaimers. In case of a disagreement between the translation and the original version of this License or a notice or disclaimer, the original version will prevail.

If a section in the Document is Entitled "Acknowledgements", "Dedications", or "History", the requirement (section 4) to Preserve its Title (section 1) will typically require changing the actual title.

9. TERMINATION

You may not copy, modify, sublicense, or distribute the Document except as expressly provided for under this License. Any other attempt to copy, modify, sublicense or distribute the Document is void, and will automatically terminate your rights under this License. However, parties who have received copies, or rights, from you under this License will not have their licenses terminated so long as such parties remain in full compliance.

10. FUTURE REVISIONS OF THIS LICENSE

The Free Software Foundation may publish new, revised versions of the GNU Free Documentation License from time to time. Such new versions will be similar in spirit to the present version, but may differ in detail to address new problems or concerns. See <http://www.gnu.org/copyleft/>.

Each version of the License is given a distinguishing version number. If the Document specifies that a particular numbered version of this License "or any later version" applies to it, you have the option of following the terms and conditions either of that specified version or of any later version that has been published (not as a draft) by the Free Software Foundation. If the Document does not specify a version number of this License, you may choose any version ever published (not as a draft) by the Free Software Foundation.