



American College Health Association **National College Health Assessment**

SPRING 2018

Reference Group Executive Summary



Table of Contents

I. Introduction	2
II. Findings	
A. General Health of College Students	3
B. Disease and Injury Prevention	4
C. Academic Impacts	5
D. Violence, Abusive Relationships and Personal Safety	5
E. Tobacco, Alcohol, and Marijuana Use	6
F. Sexual Behavior	10
G. Nutrition and Exercise	12
H. Mental Health	13
I. Sleep	16
III. Demographics and Student Characteristics	17

ACHA, the nation's principal advocate and leadership organization for college and university health, represents a diverse membership that provides and supports the delivery of health care and prevention and wellness services for the nation's 20 million college students. For more information about the association's programs and services, visit www.acha.org, and www.acha-ncha.org.

Suggested citation for this document:

American College Health Association. American College Health Association-National College Health Assessment II: Reference Group Executive Summary Spring 2018. Silver Spring, MD: American College Health Association; 2018.

Introduction

The ACHA-National College Health Assessment II (ACHA-NCHA II) is a national research survey organized by the American College Health Association (ACHA) to assist college health service providers, health educators, counselors, and administrators in collecting data about their students' habits, behaviors, and perceptions on the most prevalent health topics.

ACHA initiated the original ACHA-NCHA in 2000 and the instrument was used nation wide through the spring 2008 data collection period. The ACHA-NCHA now provides the largest known comprehensive data set on the health of college students, providing the college health and higher education fields with a vast spectrum of information on student health. A revised survey, the ACHA-NCHA-II, has been in use since the fall 2008 data collection period.

Please note the ACHA-NCHA II is not appropriate for trend comparison with items from the original ACHA-NCHA survey. Directly comparing pre- and post-redesign estimates on similar data points, without taking into account the impact of the survey's redesign, can lead to an erroneous conclusion.

Notes about this report:

1. Missing values have been excluded from analysis and only valid percents are included in this document.
2. Students responding "not applicable" were excluded from several analyses, which are specifically noted throughout this document. This will often explain differences observed between this document and the full data report.
3. ***A note about the use of sex and gender in this report:*** Survey responses are reported by sex based on the responses to questions 47a, 47b, and 47c. For the purpose of the ACHA-NCHA report documents, respondents are reported as male or female only when their responses to these three questions are consistent with one another. If students' gender identity is consistent with their sex at birth AND the student selects "no" for transgender, then respondents are designated as either *male* or *female*. If respondents select "yes" for transgender OR their sex at birth is not consistent with their gender identity, then they are designated as *non-binary*. A respondent that skips any of the three questions is designated as *unknown*. Totals displayed in this report include *non-binary* and *unknown* students.

For additional information about the survey's development, design, and methodology, email Mary T Hoban, PhD, MCHES, (mhoban@acha.org), E. Victor Leino, PhD (vleino@acha.org), or visit www.acha-ncha.org.

This Executive Summary highlights results of the ACHA-NCHA II Spring 2018 survey for Reference Group consisting of 88,178 respondents.

The overall response proportion was 17%.

Findings

A. General Health of College Students

■ 47.5 % of college students surveyed (54.9 % male and 45.0 % female) described their health as *very good or excellent*.

■ 82.4 % of college students surveyed (85.2 % male and 81.9 % female) described their health as *good, very good or excellent*.

Proportion of college students who reported being diagnosed or treated by a professional for any of the following health problems within the last 12 months:

Allergies:	19.2 %	Hepatitis B or C:	0.3 %
Asthma:	9.5 %	High blood pressure:	3.3 %
Back pain:	13.2 %	High cholesterol:	3.1 %
Broken bone/Fracture/Sprain:	5.8 %	HIV infection:	0.2 %
Bronchitis:	5.6 %	Irritable Bowel Syndrome:	3.3 %
Chlamydia:	1.7 %	Migraine headache:	9.5 %
Diabetes:	1.3 %	Mononucleosis:	1.3 %
Ear infection:	6.8 %	Pelvic Inflammatory Disease:	0.3 %
Endometriosis:	1.1 %	Repetitive stress injury:	2.0 %
Genital herpes:	0.8 %	Sinus infection:	15.2 %
Genital warts/HPV:	0.9 %	Strep throat:	9.6 %
Gonorrhea:	0.4 %	Tuberculosis:	0.3 %
		Urinary tract infection:	10.2 %

■ 54.6 % of college students (43.1 % male, 59.8 % female) reported being diagnosed or treated by a professional with one or more of the above conditions within the last 12 months.

Proportion of college students who reported any of the following:

Attention Deficit and Hyperactivity Disorder (ADHD)	7.8 %
Chronic illness (e.g., cancer, diabetes, auto-immune disorders)	6.1 %
Deafness/Hearing loss	2.1 %
Learning disability	4.7 %
Mobility/Dexterity disability	1.2 %
Partial sightedness/Blindness	2.5 %
Psychiatric condition	9.2 %
Speech or language disorder	0.9 %
Other disability	2.8 %

B. Disease and Injury Prevention

College students reported receiving the following vaccinations (shots):

- 68.4 % reported receiving vaccination against hepatitis B.
- 56.1 % reported receiving vaccination against Human Papillomavirus/HPV (cervical cancer vaccine).
- 46.4 % reported receiving vaccination against influenza (flu) in the last 12 months (shot or nasal mist).
- 73.3 % reported receiving vaccination against measles, mumps, rubella.
- 65.1 % reported receiving vaccination against meningococcal meningitis.
- 65.6 % reported receiving vaccination against varicella (chicken pox).

Other disease prevention practices reported by college students:

- 72.0 % reported having a dental exam and cleaning in the last 12 months.
- 33.3 % of males reported performing a testicular self exam in the last 30 days.
- 35.6 % of females reported performing a breast self exam in the last 30 days.
- 40.5 % of females reported having a routine gynecological exam in the last 12 months.
- 52.4 % reported using sunscreen regularly with sun exposure.
- 28.8 % reported ever being tested for Human Immunodeficiency Virus (HIV) infection.

College students reported the following behaviors within the last 12 months:

<i>Percent (%)</i>	N/A, did not do this activity within the last 12 months		Never*	Rarely or sometimes*	Mostly or always*
Wear a seatbelt when you rode in a car	0.3		0.4	2.7	97.0
Wear a helmet when you rode a bicycle	49.6		36.1	24.9	39.0
Wear a helmet when you rode a motorcycle	85.6		9.1	6.6	84.3
Wear a helmet when you were inline skating	84.8		50.6	14.7	34.8

* Students responding "N/A, did not do this activity within the last 12 months" were excluded.

C. Academic Impacts

Within the last 12 months, students reported the following factors affecting their individual academic performance, defined as: received a lower grade on an exam, or an important project; received a lower grade in the course; received an incomplete or dropped the course; or experienced a significant disruption in thesis, dissertation, research, or practicum work; (listed alphabetically):

Alcohol use:	3.1 %	Gambling:	0.2 %
Allergies:	2.1 %	Homesickness:	4.3 %
Anxiety:	26.5 %	Injury:	2.3 %
Assault (physical):	0.8 %	Internet use/computer games:	9.9 %
Assault (sexual):	1.8 %	Learning disability:	3.5 %
Attention Deficit/Hyperactivity Disorder:	5.6 %	Participation in extracurricular activities:	8.9 %
Cold/Flu/Sore throat:	16.1 %	Pregnancy (yours or partner's):	0.7 %
Concern for a troubled friend or family member:	11.9 %	Relationship difficulties:	9.5 %
Chronic health problem or serious illness:	4.4 %	Roommate difficulties:	5.7 %
Chronic pain:	3.5 %	Sexually transmitted disease/infection (STD/I):	0.4 %
Death of a friend or family member:	6.5 %	Sinus infection/Ear infection/Bronchitis/Strep throat:	4.5 %
Depression:	18.7 %	Sleep difficulties:	21.8 %
Discrimination:	1.5 %	Stress:	33.2 %
Drug use:	1.7 %	Work:	15.3 %
Eating disorder/problem:	1.5 %	Other:	2.4 %
Finances:	7.6 %		

D. Violence, Abusive Relationships and Personal Safety

Within the last 12 months, college students reported experiencing:

	Percent (%)	Male	Female	Total
A physical fight		6.4	2.1	3.5
A physical assault (not sexual assault)		2.9	2.7	2.9
A verbal threat		21.2	15.8	17.7
Sexual touching without their consent		4.6	12.4	10.2
Sexual penetration attempt without their consent		1.0	5.0	3.9
Sexual penetration without their consent		0.7	3.2	2.5
Stalking		2.6	6.7	5.6
An emotionally abusive intimate relationship		6.5	10.7	9.6
A physically abusive intimate relationship		1.7	1.9	1.9
A sexually abusive intimate relationship		1.1	3.0	2.5

Findings continued

College students reported feeling *very safe* :

	<i>Percent (%)</i>	Male	Female	Total
On their campus (daytime)		86.0	79.1	80.8
On their campus (nighttime)		53.8	23.1	32.2
In the community surrounding their school (daytime)		59.0	47.1	50.4
In the community surrounding their school (nighttime)		32.1	12.8	18.6

E. Tobacco, Alcohol and Marijuana Use

Reported use versus perceived use - reported use for all students within the past 30 days compared with how often students perceived the typical student on campus used substances within the same time period. The last line of each table combines all categories of any use in the last 30 days.

Cigarette

	<i>Percent (%)</i>	Actual Use			Perceived Use		
		Male	Female	Total	Male	Female	Total
Never used		73.8	81.6	79.1	16.2	11.4	12.9
Used, but not in the last 30 days		16.1	12.2	13.5	16.6	13.0	14.1
Used 1-9 days		6.6	4.0	4.8	45.7	45.0	45.1
Used 10-29 days		1.4	0.8	1.0	12.6	16.2	15.1
Used all 30 days		2.1	1.4	1.6	8.9	14.4	12.8
<i>Any use within the last 30 days</i>		10.1	6.2	7.5	67.3	75.6	73.0

E-Cigarette

	<i>Percent (%)</i>	Actual Use			Perceived Use		
		Male	Female	Total	Male	Female	Total
Never used		75.2	84.1	81.4	14.4	11.6	12.5
Used, but not in the last 30 days		12.0	8.6	9.6	10.9	8.6	9.4
Used 1-9 days		7.0	4.6	5.3	40.9	39.6	39.9
Used 10-29 days		2.5	1.2	1.6	20.8	21.8	21.4
Used all 30 days		3.3	1.4	2.0	13.0	18.4	16.8
<i>Any use within the last 30 days</i>		12.8	7.3	9.0	74.7	79.8	78.1

Tobacco from a water pipe (hookah)

	<i>Percent (%)</i>	Actual Use			Perceived Use		
		Male	Female	Total	Male	Female	Total
Never used		79.6	84.0	82.6	24.2	17.0	19.3
Used, but not in the last 30 days		17.1	13.6	14.7	21.1	17.4	18.6
Used 1-9 days		2.7	2.1	2.3	43.5	48.2	46.6
Used 10-29 days		0.4	0.2	0.3	7.7	12.1	10.7
Used all 30 days		0.2	0.1	0.2	3.5	5.3	4.8
<i>Any use within the last 30 days</i>		3.3	2.4	2.7	54.7	65.6	62.1

Findings continued

Alcohol

	<i>Percent (%)</i>	Actual Use		
		Male	Female	Total
Never used		23.6	20.0	21.2
Used, but not in the last 30 days		15.9	17.4	17.1
Used 1-9 days		45.2	51.3	49.3
Used 10-29 days		13.9	10.7	11.6
Used all 30 days		1.5	0.5	0.8
<i>Any use within the last 30 days</i>		60.6	62.5	61.7

Perceived Use		
Male	Female	Total
5.6	3.7	4.4
3.0	2.2	2.5
43.0	39.0	40.2
37.1	41.3	39.9
11.4	13.8	13.0
91.4	94.1	93.2

Marijuana

	<i>Percent (%)</i>	Actual Use		
		Male	Female	Total
Never used		58.6	58.8	58.5
Used, but not in the last 30 days		19.6	21.1	20.6
Used 1-9 days		12.5	13.4	13.2
Used 10-29 days		5.5	4.1	4.5
Used all 30 days		3.8	2.7	3.1
<i>Any use within the last 30 days</i>		21.8	20.2	20.9

Perceived Use		
Male	Female	Total
9.9	6.7	7.8
6.9	4.4	5.1
45.2	39.9	41.4
26.5	32.4	30.6
11.5	16.6	15.1
83.2	88.9	87.1

Drinking and Driving

■ 1.4 % of college students reported driving after having **5 or more drinks** in the last 30 days.*

■ 19.8 % of college students reported driving after having **any alcohol** in the last 30 days.*

*Students responding "N/A, don't drive" and "N/A don't drink" were excluded from this analysis.

Estimated Blood Alcohol Concentration (or eBAC) of college students reporting 1 or more drinks the last time they "partied" or socialized. **Students reporting 0 drinks were excluded from the analysis.**

Due to the improbability of a student surviving a drinking episode resulting in an extremely high eBAC, all students with an eBAC of 0.50 or higher are also omitted from these eBAC figures. eBAC is an estimated figure based on the reported number of drinks consumed during the last time they "partied" or socialized, their approximate time of consumption, sex, weight, and the average rate of ethanol metabolism.

Estimated BAC	<i>Percent (%)</i>	Male	Female	Total
< .08		73.3	71.1	71.8
< .10		80.1	78.5	78.9

Mean	0.06	0.06	0.06
Median	0.03	0.04	0.04
Std Dev	0.07	0.07	0.07

Findings continued

Reported number of drinks consumed the last time students "partied" or socialized. Only students reporting one or more drinks were included.

Number of drinks*	Percent (%)	Male	Female	Total
4 or fewer		52.4	70.1	65.1
5		10.9	11.7	11.4
6		9.5	7.4	7.9
7 or more		27.2	10.9	15.5
Mean		5.23	3.74	4.16
Median		4.00	3.00	3.00
Std Dev		4.16	2.76	3.33

* Students reporting 0 drinks were excluded.

Reported number of times college students consumed five or more drinks in a sitting within the last two weeks:

	Percent (%)	Male	Female	Total
N/A don't drink		28.8	25.3	26.5
None		38.0	50.0	46.4
1-2 times		22.7	19.5	20.3
3-5 times		8.5	4.5	5.7
6 or more times		2.0	0.6	1.1

Percent of college students who reported using prescription drugs that were not prescribed to them within the last 12 months:

	Percent (%)	Male	Female	Total
Antidepressants		2.0	3.1	2.9
Erectile dysfunction drugs		0.9	0.6	0.7
Pain killers		4.0	4.2	4.2
Sedatives		3.0	2.9	3.0
Stimulants		6.6	5.4	5.8
<i>Used 1 or more of the above</i>		11.0	10.8	11.0

Findings continued

College students reported doing the following *most of the time* or *always* when they "partied" or socialized during the last 12 months:*

	Percent (%)	Male	Female	Total
Alternate non-alcoholic with alcoholic beverages		32.8	40.9	38.8
Avoid drinking games		34.8	42.1	40.3
Choose not to drink alcohol		21.0	26.3	24.9
Determine in advance not to exceed a set number of drinks		37.8	47.7	44.8
Eat before and/or during drinking		79.1	84.8	82.9
Have a friend let you know when you have had enough		32.9	47.4	43.2
Keep track of how many drinks being consumed		63.7	72.8	70.2
Pace drinks to one or fewer an hour		27.2	41.6	37.5
Stay with the same group of friends the entire time drinking		83.2	92.7	89.9
Stick with only one kind of alcohol when drinking		46.8	57.3	54.2
Use a designated driver		81.1	90.6	87.7
Reported one or more of the above		96.7	98.6	98.0

*Students responding "N/A, don't drink" were excluded from this analysis.

College students who drank alcohol reported experiencing the following in the last 12 months when drinking alcohol:*

	Percent (%)	Male	Female	Total
Did something you later regretted		31.8	32.0	31.9
Forgot where you were or what you did		28.3	26.8	27.1
Got in trouble with the police		2.7	1.4	1.8
Someone had sex with me without my consent		1.3	2.9	2.5
Had sex with someone without their consent		0.4	0.2	0.3
Had unprotected sex		22.0	21.4	21.5
Physically injured yourself		12.4	11.6	11.9
Physically injured another person		1.8	0.8	1.1
Seriously considered suicide		4.2	3.7	4.0
Reported one or more of the above		50.6	48.8	49.3

*Students responding "N/A, don't drink" were excluded from this analysis.

F. Sexual Behavior

College students reported having the following number of sexual partners (oral sex, vaginal or anal intercourse) within the last 12 months:

	Percent (%)	Male	Female	Total
None		34.7	31.4	32.5
1		40.5	46.1	44.2
2		8.3	9.0	8.8
3		5.4	5.0	5.1
4 or more		11.1	8.5	9.3

Number of partners among students reporting to have at least one sexual partner within the last 12 months:*

	Male	Female	Total
Mean	2.50	1.98	2.16
Median	1.00	1.00	1.00
Std Dev	4.09	2.71	3.39

*Students reporting 0 sexual partners within the last 12 months were excluded.

College students reported having oral, vaginal or anal sex in the last 30 days:

Oral sex within the past 30 days

	Percent (%)	Male	Female	Total
No, have never done this sexual activity		30.2	30.3	30.4
No, have done this sexual activity but not in the last 30 days		27.3	25.1	25.8
Yes		42.5	44.5	43.8

Vaginal sex within the past 30 days

	Percent (%)	Male	Female	Total
No, have never done this sexual activity		36.4	31.5	33.1
No, have done this sexual activity but not in the last 30 days		21.3	18.1	19.1
Yes		42.3	50.4	47.8

Anal sex within the past 30 days

	Percent (%)	Male	Female	Total
No, have never done this sexual activity		70.5	75.2	73.5
No, have done this sexual activity but not in the last 30 days		22.1	20.3	20.9
Yes		7.4	4.5	5.5

Findings continued

Using a condom or other protective barrier within the last 30 days (*mostly or always*):

	<i>Percent (%)</i>	Male	Female	Total
<i>Sexually active students reported*</i>				
Oral sex		5.7	4.8	5.1
Vaginal intercourse		50.2	42.4	44.4
Anal intercourse		36.2	19.7	25.6

*Students responding "Never did this sexual activity" or "Have not done this during the last thirty days" were excluded from the analysis.

Contraceptive use reported by students or their partner the last time they had vaginal intercourse:

	<i>Percent (%)</i>	Male	Female	Total
Yes, used a method of contraception		50.1	56.4	54.2
Not applicable/Didn't use a method/Don't know		49.9	43.6	45.8

If YES to contraceptive use the last time student had vaginal intercourse, reported means of birth control used among college students or their partner to prevent pregnancy:

	<i>Percent (%)</i>	Male	Female	Total
Birth control pills (monthly or extended cycle)		56.3	52.9	53.5
Birth control shots		4.2	3.3	3.6
Birth control implants		9.7	8.9	9.1
Birth control patch		1.6	0.8	1.0
Vaginal ring		3.0	2.7	2.8
Intrauterine device		13.4	14.2	14.1
Male condom		65.5	56.2	58.8
Female condom		0.9	0.6	0.7
Diaphragm or cervical cap		0.5	0.2	0.3
Contraceptive sponge		0.4	0.1	0.2
Spermicide (foam, jelly, cream)		4.6	2.2	2.9
Fertility awareness (calendar, mucous, basal body temperature)		6.3	8.0	7.6
Withdrawal		27.7	32.5	31.1
Sterilization (hysterectomy, tubes tied, vasectomy)		2.1	2.3	2.3
Other method		2.4	1.7	1.9
<i>Male condom use plus another method</i>		50.3	45.4	46.7
<i>Any two or more methods (excluding male condoms)</i>		32.8	33.9	33.6

■ 15.6 % of sexually active college students reported using (or reported their partner used) emergency contraception ("morning after pill") within the last 12 months.

(male: 13.1 %; female: 16.6 %).*

*Students responding "Not sexually active" were excluded from the analysis.

■ 1.2 % of college students who had vaginal intercourse within the last 12 months reported experiencing an unintentional pregnancy or got someone pregnant within the last 12 months.

(male: 1.1 %; female: 1.3 %).**

**Students responding "Have not had vaginal intercourse within the last 12 months" were excluded from the analysis.

G. Nutrition and Exercise

College students reported usually eating the following number of servings of fruits and vegetables per day:

	Percent (%)	Male	Female	Total
0 servings per day		10.2	7.5	8.4
1-2 per day		62.5	61.4	61.6
3-4 per day		23.1	26.1	25.2
5 or more per day		4.3	5.0	4.8

College students reported the following behaviors within the past 7 days:

Do moderate-intensity cardio or aerobic exercise for at least 30 minutes:

	Percent (%)	Male	Female	Total
0 days		21.4	21.5	21.7
1-4 days		55.9	59.8	58.6
5-7 days		22.7	18.6	19.8

Do vigorous-intensity cardio or aerobic exercise for at least 20 minutes:

	Percent (%)	Male	Female	Total
0 days		36.1	43.3	41.5
1-2 days		32.0	30.9	31.1
3-7 days		31.9	25.8	27.4

Physical Activity and Public Health: Updated Recommendations for Adults. From the American College of Sports Medicine and the American Heart Association (2007): Moderate-intensity cardio or aerobic exercise for at least 30 minutes on 5 or more days per week, or vigorous-intensity cardio or aerobic exercise for at least 20 minutes on 3 or more days per week.

Students meeting the Recommendations for moderate-intensity exercise, vigorous-intensity exercise, or a combination of the two (2 moderate-intensity exercise periods = 1 vigorous-intensity exercise period).

	Percent (%)	Male	Female	Total
Guidelines met		50.6	44.6	46.2

Estimated average Body Mass Index (BMI): This figure incorporates reported height, and weight to form a general indicator of physical health. Categories defined by The World Health Organization (WHO) 2000, reprinted 2004. Obesity: Preventing and Managing the Global Epidemic. WHO Tech Report Series: 894.

BMI	Percent (%)	Male	Female	Total
<18.5 Underweight		3.5	4.6	4.3
18.5-24.9 Healthy Weight		52.2	57.4	55.7
25-29.9 Overweight		29.0	21.7	23.8
30-34.9 Class I Obesity		9.9	9.4	9.6
35-39.9 Class II Obesity		3.5	4.2	4.0
≥40 Class III Obesity		1.9	2.8	2.6
Mean		25.37	25.09	25.20
Median		24.37	23.49	23.75
Std Dev		5.47	6.01	6.07

H. Mental Health

Students reported experiencing the following within the last 12 months:

Felt things were hopeless

	Percent (%)	Male	Female	Total
No, never		35.7	25.0	27.8
No, not last 12 months		19.2	18.9	18.8
Yes, last 2 weeks		17.2	20.9	20.2
Yes, last 30 days		8.0	11.2	10.3
Yes, in last 12 months		19.9	23.9	22.8
Any time within the last 12 months		45.1	56.1	53.4

Felt overwhelmed by all you had to do

	Percent (%)	Male	Female	Total
No, never		14.7	5.4	8.2
No, not last 12 months		7.5	3.2	4.4
Yes, last 2 weeks		40.6	57.0	52.4
Yes, last 30 days		15.3	16.6	16.1
Yes, in last 12 months		21.9	17.8	18.9
Any time within the last 12 months		77.8	91.4	87.4

Felt exhausted (not from physical activity)

	Percent (%)	Male	Female	Total
No, never		17.1	7.6	10.3
No, not last 12 months		7.8	4.5	5.4
Yes, last 2 weeks		41.8	55.7	51.9
Yes, last 30 days		15.2	16.1	15.7
Yes, in last 12 months		18.2	16.2	16.7
Any time within the last 12 months		75.2	87.9	84.3

Felt very lonely

	Percent (%)	Male	Female	Total
No, never		26.2	17.3	19.7
No, not last 12 months		18.7	17.2	17.5
Yes, last 2 weeks		23.0	28.3	27.1
Yes, last 30 days		10.7	14.2	13.2
Yes, in last 12 months		21.4	23.0	22.5
Any time within the last 12 months		55.1	65.5	62.8

Felt very sad

Percent (%)	Male	Female	Total
No, never	24.0	13.8	16.6
No, not last 12 months	17.9	13.6	14.7
Yes, last 2 weeks	23.2	32.0	29.9
Yes, last 30 days	11.3	15.3	14.1
Yes, in last 12 months	23.6	25.4	24.7
Any time within the last 12 months	58.1	72.6	68.7

Felt overwhelming anxiety

Percent (%)	Male	Female	Total
No, never	35.7	19.9	24.3
No, not last 12 months	15.4	11.1	12.3
Yes, last 2 weeks	18.2	30.9	27.6
Yes, last 30 days	10.5	14.8	13.6
Yes, in last 12 months	20.1	23.3	22.3
Any time within the last 12 months	48.8	68.9	63.4

Seriously considered suicide

Percent (%)	Male	Female	Total
No, never	75.4	70.7	71.2
No, not last 12 months	14.1	17.3	16.7
Yes, last 2 weeks	2.4	2.2	2.4
Yes, last 30 days	1.6	1.8	1.8
Yes, in last 12 months	6.5	7.9	7.8
Any time within the last 12 months	10.5	12.0	12.1

Intentionally cut, burned, bruised, or otherwise injured yourself

Percent (%)	Male	Female	Total
No, never	85.3	74.9	77.1
No, not last 12 months	9.8	16.7	15.1
Yes, last 2 weeks	1.3	1.9	1.9
Yes, last 30 days	0.8	1.3	1.2
Yes, in last 12 months	2.7	5.2	4.7
Any time within the last 12 months	4.8	8.4	7.8

Felt so depressed that it was difficult to function

Percent (%)	Male	Female	Total
No, never	45.0	34.9	37.3
No, not last 12 months	20.7	21.2	20.9
Yes, last 2 weeks	11.9	15.7	15.1
Yes, last 30 days	6.1	8.6	8.0
Yes, in last 12 months	16.3	19.7	18.8
Any time within the last 12 months	34.3	44.0	41.9

Felt overwhelming anger

Percent (%)	Male	Female	Total
No, never	40.2	33.7	35.3
No, not last 12 months	23.3	22.5	22.7
Yes, last 2 weeks	11.4	13.8	13.3
Yes, last 30 days	7.1	9.8	9.1
Yes, in last 12 months	17.9	20.2	19.6
Any time within the last 12 months	36.5	43.8	42.1

Attempted suicide

Percent (%)	Male	Female	Total
No, never	91.1	87.9	88.3
No, not last 12 months	7.3	10.5	10.0
Yes, last 2 weeks	0.4	0.2	0.3
Yes, last 30 days	0.2	0.1	0.2
Yes, in last 12 months	1.0	1.2	1.2
Any time within the last 12 months	1.6	1.6	1.7

Findings continued

Within the last 12 months, diagnosed or treated by a professional for the following:

	<i>Percent (%)</i>	Male	Female	Total
Anorexia		0.7	1.9	1.6
Anxiety		11.7	25.5	22.1
Attention Deficit and Hyperactivity Disorder		6.6	5.3	6.0
Bipolar Disorder		1.6	1.8	1.9
Bulimia		0.7	1.3	1.2
Depression		11.0	20.1	18.1
Insomnia		3.8	5.8	5.5
Other sleep disorder		2.7	2.4	2.6
Obsessive Compulsive Disorder		2.3	3.2	3.1
Panic attacks		4.5	13.2	11.0
Phobia		1.0	1.5	1.5
Schizophrenia		0.6	0.2	0.4
Substance abuse or addiction		1.7	1.0	1.2
Other addiction		1.1	0.4	0.6
Other mental health condition		2.8	3.8	3.8
<i>Students reporting none of the above</i>		79.1	66.9	69.8
<i>Students reporting only one of the above</i>		8.4	9.1	8.9
<i>Students reporting both Depression and Anxiety</i>		7.6	16.7	14.6
<i>Students reporting any two or more of the above excluding the combination of Depression and Anxiety</i>		5.8	10.1	9.3

Within the last 12 months, any of the following been traumatic or very difficult to handle:

	<i>Percent (%)</i>	Male	Female	Total
Academics		41.5	53.3	50.2
Career-related issue		25.2	30.8	29.3
Death of family member or friend		12.8	18.2	16.7
Family problems		21.3	34.9	31.3
Intimate relationships		27.8	32.6	31.4
Other social relationships		22.6	31.3	29.2
Finances		29.4	38.2	35.9
Health problem of family member or partner		15.8	24.1	21.8
Personal appearance		20.6	35.1	31.2
Personal health issue		17.3	27.1	24.6
Sleep difficulties		27.7	34.5	32.9
Other		9.0	10.9	10.8
<i>Students reporting none of the above</i>		31.0	19.2	22.4
<i>Students reporting only one of the above</i>		13.9	10.9	11.6
<i>Students reporting 2 of the above</i>		12.2	11.6	11.7
<i>Students reporting 3 or more of the above</i>		42.9	58.4	54.3

Findings continued

Within the last 12 months, how would you rate the overall level of stress experienced:

	<i>Percent (%)</i>	Male	Female	Total
No stress		3.5	0.8	1.6
Less than average stress		11.7	4.4	6.5
Average stress		37.9	33.3	34.3
More than average stress		37.7	47.9	44.9
Tremendous stress		9.2	13.7	12.7

I. Sleep

Past 7 days, getting enough sleep to feel rested in the morning:

	<i>Percent (%)</i>	Male	Female	Total
0 days		10.4	12.6	12.2
1-2 days		27.7	32.9	31.3
3-5 days		47.2	44.5	45.1
6+ days		14.6	10.0	11.3

Past 7 days, how often felt tired, dragged out, or sleepy during the day:

	<i>Percent (%)</i>	Male	Female	Total
0 days		12.8	6.4	8.3
1-2 days		34.1	27.6	29.2
3-5 days		40.1	46.0	44.3
6+ days		13.0	20.0	18.2

Past 7 days, how much of a problem with sleepiness during daytime activities:

	<i>Percent (%)</i>	Male	Female	Total
No problem		13.9	7.9	9.7
A little problem		49.1	46.4	46.9
More than a little problem		22.7	26.2	25.3
A big problem		10.2	13.7	12.8
A very big problem		4.1	5.8	5.4

Demographics and Student Characteristics

■ Age:

18 - 20 years:	45.1 %
21 - 24 years:	34.9 %
25 - 29 years:	10.9 %
30+ years:	9.1 %

■ Gender*

Female:	68.5 %
Male:	28.5 %
Non-binary	3.0 %

■ Student status:

1st year undergraduate:	20.8 %
2nd year undergraduate:	17.7 %
3rd year undergraduate:	21.9 %
4th year undergraduate:	17.8 %
5th year or more undergraduate:	6.7 %
Graduate or professional:	14.4 %
Not seeking a degree:	0.2 %
Other:	0.5 %

Full-time student:	91.9 %
Part-time student:	7.2 %
Other student:	0.9 %

■ Relationship status:

Not in a relationship:	49.8 %
In a relationship but not living together:	34.1 %
In a relationship and living together:	16.1 %

■ Marital status:

Single:	87.9 %
Married/Partnered:	9.5 %
Separated/Divorced/Other:	2.6 %

■ Primary Source of Health Insurance:

College/university sponsored plan:	11.0 %
Parents' plan:	65.4 %
Another plan:	18.5 %
Don't have health insurance:	3.4 %
Not sure if have plan:	1.7 %

■ Students describe themselves as:

White:	63.5 %
Black or African American:	4.8 %
Hispanic or Latino/a:	17.4 %
Asian or Pacific Islander:	14.2 %
American Indian, Alaskan Native or Native Hawaiian:	1.8 %
Biracial or Multiracial:	5.0 %
Other:	2.7 %

■ International Student:

International:	6.1 %
----------------	-------

■ Students describe themselves as:

Asexual:	1.0 %
Bisexual:	7.2 %
Gay:	1.9 %
Lesbian:	1.4 %
Pansexual:	1.9 %
Queer:	1.4 %
Questioning:	1.9 %
Same Gender Loving:	0.0 %
Straight/Heterosexual:	82.6 %
Another identity:	0.6 %

■ Housing:

Campus residence hall:	31.9 %
Fraternity or sorority house:	1.0 %
Other university housing:	4.2 %
Parent/guardian home:	19.4 %
Other off-campus housing:	38.2 %
Other:	5.3 %

■ Participated in organized college athletics:

Varsity:	6.5 %
Club sports:	8.0 %
Intramurals:	13.1 %

■ Member of a social fraternity or sorority:

Greek member:	8.7 %
---------------	-------

* See note on page 2 regarding gender categories

Demographics of Participating Institutions

One hundred seventy-three postsecondary institutions self-selected to participate in the Spring 2018 ACHA National College Health Assessment and 104,648 surveys were completed by students on these campuses. For the purpose of forming the Reference Group, only institutions located in the United States that surveyed all students or used a random sampling technique are included in the analysis, yielding a final data set consisting of 88,178 students at 140 schools. Demographic characteristics of the 140 campuses follow. *Note that schools surveying in the 30 days following their Spring Break are omitted from this report.*

Demographical Characteristics of the 140 US Postsecondary Institutions Included in the Spring 2018 ACHA-NCHA II Reference Group	
Campus Characteristic	<i>n</i>
Type of Institution	
Public	96
Private	44
2-year	3
4-year or above	137
Location of Campus	
Northeast (CT, ME, MA, NH, NJ, NY, PA, RI, VT)	23
Midwest (IL, IN, IA, KS, MI, MN, MO, NE, ND, OH, SD, WI)	33
South (AL, AR, DE, DC, FL, GA, KY, LA, MD, MS, NC, OK, SC, TN, TX, VA, WV)	33
West (AK, AZ, CA, CO, HI, ID, MT, NV, NM, OR, UT, WA, WY)	51
Campus Size	
< 2,500 students	35
2,500 – 4,999 students	16
5,000 – 9,999 students	26
10,000 – 19,999 students	25
20,000 students or more	38
Campus Setting	
Very large city (population over 500,000)	19
Large city (population 250,000-499,999)	15
Small city (population 50,000-249,999)	52
Large town (population 10,000 – 49,999)	34
Small town (population 2,500-9,999)	17
Rural community (population under 2,500)	3
Carnegie Classification	
Associates Colleges	3
Baccalaureate Colleges	31
Baccalaureate/Associates Colleges	1
Masters Colleges and Universities	54
Doctoral Universities	48
Special Focus Institutions	3
Miscellaneous/Not Classified	0

Demographical Characteristics of the 140 US Postsecondary Institutions Included in the Spring 2018 ACHA-NCHA II Reference Group	
Campus Characteristic	<i>n</i>
ACHA Membership Status	
Institutional Member	116
Nonmember	24
Religious Affiliation	
No	124
Yes	16
<u>If yes:</u>	
Catholic	4
Protestant or Other Christian	12
Postsecondary Minority Institution (US Department of Education)	
No	114
Yes	26
<u>*If yes:</u>	
Postsecondary Minority Institution	0
Historically Black College or University (HBCU)	1
High Hispanic Enrollment	4
Hispanic-serving Institution (HSI)	22
Tribal College or University	0
Predominately Black Institution	0
Asian American and Native American Pacific Islander-serving	3
Alaska Native-serving or Native Hawaiian-serving Institution	0
Native American-serving Nontribal Institution	2
*institutions may hold more than one type of minority status	

	All Spring 2018 Institutions	Paper Survey Administration	Web Survey Administration
Number of institutions	140	4	136
Number of students	88,178	2,358	85,820
Mean response proportion	17%	78%	15%
Median response proportion	14%	88%	14%