

**(Summary) A demonstration of the Titius–Bode law, the number max. 31 (including max. 9 main rings) of Saturn’s rings and the radius of Fine-Ring by using the Kerr-Newman solution of the relativity theory**

Fumitaka Inuyama. Senior Power Engineer (e-mail: inusanin@yahoo.co.jp)

Permanent address: 5-17-33, Torikai Jonan-ku, Fukuoka, 814-0103, Japan

Environment Dept., Kyudensangyo Inc. (Retired) (www.kyudensangyo.co.jp)

Sayuri Taniguchi. Amateur of starry sky

## 1. Introduction

The beautiful Titius-Bode law ( $\xi = 0.4 + 0.3 \times 2^n$ ) discovered 250 years ago (in 1766), is considered to be a mathematical coincidence rather than an "exact" law [1], because it has not yet been physically proved. However, considering the disturbance restoration and the stability of the asteroid belt orbit, there must be some underlying logical necessity.

Planetary orbits are often computed by Newtonian mechanics with the kinetic energy and the universal gravitation energy. However, applying the principle of energy-minimum to the Newtonian mechanics leads to the result that the stable orbital radius is only one value, which is totally incompatible with actual phenomena. This discrepancy must result from the shortage of elements which rule over the planetary orbits. Other elements to rule over the planetary orbits are the electric charge energy and the rotation energy, both of which are guided by the Kerr-Newman solution (discovered in 1965) of the general relativity theory (discovered in 1915). Here, I mathematically demonstrated the Titius-Bode law, and also calculated the number of Saturn’s rings, maximum 31 and the radius of Fine-Ring, by applying the principle of energy-minimum of Newtonian methods to the complicated energy equation which adopts mass, electric charge and rotation elements of the central core star and solving sole differential equation.

## 2. Methods

The outline of the solution method and the key equation numbers in this article are as follows. The detailed analysis process is provided in a separate sheet of paper [2].

- 1) The equation for energy in the space-time field is obtained from the Kerr-Newman solution, a strict solution of the Einstein equations of the general relativity theory.

$$f_1(\rho, \theta, d\rho/dt, d\theta/dt, d\varphi/dt, \varepsilon) = 0 \quad (\text{eq. 3})$$

- 2) This energy equation is partially differentiated by  $\theta$  to the minimum energy. The result is  $\theta=\pi/2$ , and the calculation below proceeds at  $\theta=\pi/2$ , i.e., on the equatorial plane.

$$f_2(\rho, \pi/2, d\rho/dt, 0, d\varphi/dt, \varepsilon) = 0$$

- 3) The angular momentum equivalent  $J$  is obtained by applying the variational principle to the Kerr-Newman solution to calculate  $d\varphi/dt$ .

$$\xi(\rho, d\varphi/dt, J) = 0 \quad (\text{eq. 6})$$

- 4) Because an additional radius is  $d\rho=0$  at the aphelion and perihelion distances  $R$ , the calculation below proceeds at distance  $R$ .

$$f_3(R, \pi/2, 0, 0, d\varphi/dt, \varepsilon) = 0 \quad (\text{eq. 7})$$

- 5) Substituting  $d\varphi/dt$  from  $\xi=0$  into  $f_3=0$  results in a relational expression of the radius, the angular momentum equivalent, and the energy.

$$f_4(R, \pi/2, 0, 0, J, \varepsilon) = 0 \quad (\text{eq. 9})$$

6) The orbital distance  $R$  is determined by the energy and the angular momentum equivalent, i.e.,  $R = R(\varepsilon, J)$ .  $R$  is partially differentiated by  $\varepsilon$ , that is,  $f_4$  is partially differentiated by  $\varepsilon$ .

$$g(R, J, \varepsilon, \partial R/\partial \varepsilon) = 0 \quad (\text{eq. 10})$$

7) Derive the angular momentum equivalent  $J$  from  $f_4(R, \pi/2, 0, 0, J, \varepsilon) = 0$  and substitute it into  $g(R, J, \varepsilon, \partial R/\partial \varepsilon) = 0$ . Make this into an important differential equation which is just composed of the radius and the energy to analyze unique characteristics of orbits.


$$h(R, \varepsilon, d\varepsilon/dR) = 0 \quad (\text{eq. 11})$$

8) Solving the differential equation  $h$  results in a complicated set of *arctan*, *log*, power functions and an integration constant  $K$ .

$$H(R, \varepsilon, K) = 0 \quad (\text{eq. 14}) \quad (\text{eq. 15})$$

9) By using that the minimum energy is  $d\varepsilon/dR=0$  in  $h(R, \varepsilon, d\varepsilon/dR) = 0$ , following simultaneous equations are obtained and solved.

$$h(r, \varepsilon_{min}, 0) = 0 \quad (1) \quad H(r, \varepsilon_{min}, K) = 0 \quad (2) \quad (\text{eq. 16})$$



10) Because the integration constant  $K$  is common to all orbits, the Titius–Bode law is demonstrated and also the number of Saturn’s rings and the radius of Fine-Ring are calculated.

$$I(r, K) = 0 \quad (\text{eq. 23}) \quad (\text{eq. 26}) \quad (\text{eq. 27})$$

## Chapter 1 The Energy Equation

### 1.1. Introduction to the Energy Equation

In the no-hair theorem, all black holes are thought to settle into the black holes of the Kerr-Newman solution. There are two preconditions for the following analysis besides it in the Kerr-Newman solution.

- 1) The analysis object must be sufficiently far from the center of mass.
- 2) The rotation speed of the center of mass must not be too fast. The characteristic Boyer-Lindquist coordinates in the Kerr solution are equal to general polar coordinates in the first-order term  $a/\rho$ [3].

The strict Boyer-Lindquist metric of the Kerr-Newman geometry [4] is as follows.

$$ds^2 = -\frac{R^2 \Delta}{\rho^2} (dt - a \sin^2 \theta d\phi)^2 + \frac{\rho^2}{R^2 \Delta} dr^2 + \rho^2 d\theta^2 + \frac{R^4 \sin^2 \theta}{\rho^2} \left( d\phi - \frac{a}{R^2} dt \right)^2$$

At the large radius  $r$ , the Boyer-Lindquist metric is as follows.

$$ds^2 \rightarrow -\left(1 - \frac{2M}{r}\right) dt^2 - \frac{4aM \sin^2 \theta}{r} dt d\phi + \left(1 + \frac{2M}{r}\right) dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2)$$

Symbols are changed from the Boyer-Lindquist metric to the general polar coordinate.

The Kerr-Newman solution of the general relativity theory is given by (eq.1). In this expression,  $m$ ,  $a$  and  $e$  are the mass, rotation and electric charge elements respectively.

$$(1) \quad ds^2 = \left(1 - \frac{2m\rho - e^2}{\rho^2 + a^2 \cos^2 \theta}\right) (c dt)^2 - \frac{\rho^2 + a^2 \cos^2 \theta}{\rho^2 + a^2 - 2m\rho + e^2} d\rho^2 - (\rho^2 + a^2 \cos^2 \theta) d\theta^2 \\ - \left[ (\rho^2 + a^2) + \frac{(2m\rho - e^2)a^2 \sin^2 \theta}{\rho^2 + a^2 \cos^2 \theta} \right] \sin^2 \theta d\phi^2 - \frac{2(2m\rho - e^2)a \sin^2 \theta}{\rho^2 + a^2 \cos^2 \theta} c dt \cdot d\phi$$

$\Gamma$  is as follows when  $ds$  is divided by the time elements ( $c dt$ ).  $\frac{1}{\Gamma^2} = \left(\frac{ds}{c dt}\right)^2$

The Lorentz transformation factor  $\gamma (= c dt/ds)$  in the Minkowski space-time of the special relativity theory is an important component of the energy  $E = M c^2 = M_0 \gamma c^2$ .  $\Gamma (= c dt/ds)$  of the Kerr-Newman solution of the general relativity theory is analogous to  $\gamma$ .

On this occasion, by following the principle of minimum energy, the sign of  $m$  is changed to  $-m$ ,  $a$  is changed to  $+a$ , and  $e$  is changed to  $+e$ . Therefore, the energy equation is  $E = \Gamma(\rho, \theta, \varphi, t, -m, a, e)$ .

$$(2) \quad \frac{1}{E^2} = \left(1 + \frac{2m\rho + e^2}{\rho^2 + a^2 \cos^2 \theta}\right) - \frac{\rho^2 + a^2 \cos^2 \theta}{\rho^2 + a^2 + 2m\rho + e^2} \left(\frac{d\rho}{cdt}\right)^2 - (\rho^2 + a^2 \cos^2 \theta) \left(\frac{d\theta}{cdt}\right)^2 \\ - \left[ (\rho^2 + a^2) - \frac{(2m\rho + e^2)a^2 \sin^2 \theta}{\rho^2 + a^2 \cos^2 \theta} \right] \sin^2 \theta \left(\frac{d\varphi}{cdt}\right)^2 + \frac{2(2m\rho + e^2)a \sin^2 \theta}{\rho^2 + a^2 \cos^2 \theta} \left(\frac{d\varphi}{cdt}\right)$$

Since  $E$  has a decisive massive energy  $M_0 c^2$ , it is converted into  $\varepsilon$   $1/E^2 = 1 - 2\varepsilon$  in (eq.3).

$$(3) \quad -2\varepsilon = \frac{2m\rho + e^2}{\rho^2 + a^2 \cos^2 \theta} - \frac{\rho^2 + a^2 \cos^2 \theta}{\rho^2 + a^2 + 2m\rho + e^2} \left(\frac{d\rho}{cdt}\right)^2 - (\rho^2 + a^2 \cos^2 \theta) \left(\frac{d\theta}{cdt}\right)^2 \\ - \left[ (\rho^2 + a^2) - \frac{(2m\rho + e^2)a^2 \sin^2 \theta}{\rho^2 + a^2 \cos^2 \theta} \right] \sin^2 \theta \left(\frac{d\varphi}{cdt}\right)^2 + \frac{2(2m\rho + e^2)a \sin^2 \theta}{\rho^2 + a^2 \cos^2 \theta} \left(\frac{d\varphi}{cdt}\right)$$

Partial differentiation is used to minimize the energy  $\varepsilon(\rho, \theta, \varphi, t)$  by using  $\partial\varepsilon/\partial\theta = 0$ .

$$\left\{ \begin{array}{l} \frac{(2m\rho + e^2)a^2}{(\rho^2 + a^2 \cos^2 \theta)^2} + \frac{a^2}{\rho^2 + a^2 + 2m\rho + e^2} \left(\frac{d\rho}{cdt}\right)^2 + a^2 \left(\frac{d\theta}{cdt}\right)^2 \\ - \left[ (\rho^2 + a^2) - \frac{(2m\rho + e^2)2a^2 \sin^2 \theta}{\rho^2 + a^2 \cos^2 \theta} - \frac{(2m\rho + e^2)a^4 \sin^4 \theta}{(\rho^2 + a^2 \cos^2 \theta)^2} \right] \left(\frac{d\varphi}{cdt}\right)^2 \\ + \left[ \frac{2(2m\rho + e^2)a}{\rho^2 + a^2 \cos^2 \theta} + \frac{2(2m\rho + e^2)a^3 \sin^2 \theta}{(\rho^2 + a^2 \cos^2 \theta)^2} \right] \left(\frac{d\varphi}{cdt}\right) \end{array} \right\} \cdot \sin 2\theta = 0$$

That is, the energy  $E$  and  $\varepsilon$  are minimized at  $\theta = \pi/2$  and the planets gather on the equatorial plane where the energy is stable.

## 1.2. Time component from the variational principle

When the rotation speed of the center of mass is not too fast, the Kerr-Newman solution expanded in the first order of  $a/\rho$  takes the form given in (eq.4):

$$(4) \quad \left(\frac{ds}{ds}\right)^2 = 1 = \left(1 - \frac{2m}{\rho} + \frac{e^2}{\rho^2}\right) \left(\frac{cdt}{ds}\right)^2 - \frac{1}{1 - \frac{2m}{\rho} + \frac{e^2}{\rho^2}} \left(\frac{d\rho}{ds}\right)^2 - \rho^2 \left(\frac{d\theta}{ds}\right)^2 - \rho^2 \sin^2 \theta \left(\frac{d\varphi}{ds}\right)^2 \\ - \frac{2a}{\rho} \left(2m - \frac{e^2}{\rho}\right) \sin^2 \theta \left(\frac{cdt}{ds}\right) \left(\frac{d\varphi}{ds}\right)$$

The Euler-Lagrange equation [5] is adopted by applying the variational principle to the Kerr-Newman solution.

$$\delta \int \left[ \left(1 - \frac{2m}{\rho} + \frac{e^2}{\rho^2}\right) \left(\frac{cdt}{ds}\right)^2 - \frac{1}{1 - \frac{2m}{\rho} + \frac{e^2}{\rho^2}} \left(\frac{d\rho}{ds}\right)^2 - \rho^2 \left\{ \left(\frac{d\theta}{ds}\right)^2 + \sin^2 \theta \left(\frac{d\varphi}{ds}\right)^2 \right\} \right. \\ \left. - \frac{2a}{\rho} \left(2m - \frac{e^2}{\rho}\right) \sin^2 \theta \left(\frac{cdt}{ds}\right) \left(\frac{d\varphi}{ds}\right) \right] ds = 0$$

Eventually, (eq.5) is obtained at the equatorial plane of the rotating center of mass where the energy is stable. Hereafter, I perform the calculation at the equatorial plane ( $\theta = \pi/2$ ) of the rotating center of mass.

$$(5) \quad \begin{cases} \frac{d}{ds} \left[ \left( 1 - \frac{2m}{\rho} + \frac{e^2}{\rho^2} \right) \left( \frac{cdt}{ds} \right) - \frac{a}{\rho} \left( 2m - \frac{e^2}{\rho} \right) \left( \frac{d\varphi}{ds} \right) \right] = 0 & \text{time component} \\ \frac{d}{ds} \left[ \rho^2 \left( \frac{d\varphi}{ds} \right) + \frac{a}{\rho} \left( 2m - \frac{e^2}{\rho} \right) \left( \frac{cdt}{ds} \right) \right] = 0 & \varphi \text{ component} \end{cases}$$

The two equations in (eq.5) are integrated over  $ds$ .  $d\varphi/dt$  (eq. 6) with an integration variable  $J$  is obtained by using the resulting pair of simultaneous equations,

$$(6) \quad \frac{d\varphi}{dt} = \frac{\left( \frac{d\varphi}{ds} \right)}{\left( \frac{dt}{ds} \right)} = \frac{J \left( \rho - 2m + \frac{e^2}{\rho} \right) + a \left( \frac{e^2}{\rho} - 2m \right)}{\rho^3 + Ja \left( 2m - \frac{e^2}{\rho} \right)} \cdot c \quad \begin{array}{l} J : \text{the angular momentum equivalent} \\ \text{(a kind of Carter constant in relativity theory)} \end{array}$$

The distance variables are defined as follows:

$\rho$  : An arbitrary orbital distance in two- or three-dimensional coordinates.

$R$  : The aphelion and perihelion distances at the equatorial plane of the rotating center of mass.

$r$  : The aphelion and perihelion distances, both of which are energetically stable at the equatorial plane.

### 1.3. Introduction of the angular momentum equivalent

Because additional  $\rho$  at the aphelion and perihelion distances is  $d\rho=0$ , the energy equation is given by (eq.7).

$$(7) \quad 0 = 2\varepsilon + \frac{2m}{R} + \frac{e^2}{R^2} - R^2 \left( \frac{d\varphi}{cdt} \right)^2 + \frac{4a}{R} \left( m + \frac{e^2}{2R} \right) \left( \frac{d\varphi}{cdt} \right)$$

$d\varphi/cdt$  (eq. 6) is composed of the angular momentum equivalent and is substituted into (eq.7).  $J$  is obtained as in (eq.8) by adopting the secondary order  $R$ .

$$(8) \quad J = \frac{4am + R\delta\sqrt{R(2\varepsilon R + 2m + C)}}{R^2(R - 2m + C) - a(2m - C)\delta\sqrt{R(2\varepsilon R + 2m + C)}} R^2$$

Here,  $\delta = \pm 1$  and  $C = e^2/R$ .  $\delta$  is related to the orbital rotation direction.

## Chapter 2 The Space Fantasy Differential Equation

### 2.1. Introduction of the Space Fantasy differential equation

It leads not to the numerical analysis but to the analytical unique characteristics. The relation of  $R$ ,  $\varepsilon$ , and  $J$  are given as (eq.9) at the aphelion and perihelion distances  $R$  by changing the angular momentum equivalent  $J$  (eq. 8).

(eq.9) is far more complicated than the Kepler-Newton equation  $2\varepsilon R^2 + 2mR - J^2 = 0$

$$(9) \quad 0 = 2\varepsilon + \frac{2m}{R} + \frac{e^2}{R^2} - R^2 \left[ \frac{J \left( R - 2m + \frac{e^2}{R} \right) + a \left( \frac{e^2}{R} - 2m \right)}{R^3 + Ja \left( 2m - \frac{e^2}{R} \right)} \right]^2 + \frac{4a}{R} \left( m + \frac{e^2}{2R} \right) \left[ \frac{J \left( R - 2m + \frac{e^2}{R} \right) + a \left( \frac{e^2}{R} - 2m \right)}{R^3 + Ja \left( 2m - \frac{e^2}{R} \right)} \right]$$

Since the orbital distance  $R$  is determined by the energy  $\varepsilon$  and the angular momentum equivalent  $J$ ,  $R = R(\varepsilon, J)$ . A new differential equation is given as (eq.10) by partially differentiating  $R$  by  $\varepsilon$ , then substituting  $J$  into this, and adopting the reciprocal of  $\partial R/\partial \varepsilon$ .

$$(10) \quad \frac{\partial \varepsilon}{\partial R} [R^3 + Ja(2m - C)]^2 = \frac{(m + C)[R^3 + Ja(2m - C)]^2}{R^2} + \frac{[J(R - 2m + C) - 2am + aC] [J(R - 2m + C) + 3aC] \cdot R}{1} + \frac{2R^2[J(R - 2m + C) - 4am] [J^2a(m - C) - JR^2(R - 3m + 2C) + aR^2(3m - 2C)]}{R^3 + Ja(2m - C)}$$

Here, by substituting  $J$  of (eq. 8) into (eq.10), the second order  $R$  is obtained.

Through all these extensive calculation processes, the relation between  $\varepsilon$  and  $R$  is summarized as (eq.11).

$$(11) \quad \frac{d\varepsilon}{dR} R^4 (R^2 - 4mR + 2CR + 4m^2) = mR^2(-R^2 + 8mR - 4CR - 12m^2) + \varepsilon \cdot 2R^3(-R^2 + 6mR - 4CR - 8m^2) + 2am(2R^2 + 2mR - CR - 12m^2)\delta\sqrt{R(2\varepsilon R + 2m + C)} + \varepsilon \cdot 4aR(3mR - 2CR - 6m^2 + 7Cm)\delta\sqrt{R(2\varepsilon R + 2m + C)} \quad C = e^2/R \quad (R \text{ 2ry order})$$

I call tentatively this second order equation (eq. 11) "the Space Fantasy (SF) differential equation".

The change of variables is performed to solve the SF differential equation for  $S$ . The result is (eq.12).

$$S = R\sqrt{R(2\varepsilon R + 2m + C)}$$

$$(12) \quad \frac{dS}{dR} = \frac{2e^2(e^2 + 2m^2)}{SR} + \frac{4a\delta m + S}{R} + \frac{6a\delta m S^2}{R^5} \quad (R \text{ 0 order})$$

The form of the differential equation in (eq.12) is more complicated than the Riccati's differential equation, which never has an exact general solution [6]. Since  $6a\delta m S^2/R^5$  is much smaller than  $S/R$  and  $4a\delta m/R$ , it can be treated as a constant  $\theta$ . Also, (eq.12) can be reduced to the problem of an approximate differential equation, and it is given as (eq.13).

$$\frac{dS}{dR} = \frac{1}{S} \left[ \frac{2E^4}{R} + \frac{4a\delta m S}{R} \left( 1 + \frac{6S^2}{4R^4} \right) + \frac{S^2}{R} \right] \quad E^4 = e^2(e^2 + 2m^2)$$

$$\cong \frac{1}{S} \left[ \frac{2E^4}{R} + \frac{4a\delta m S}{R} (1 + \theta) + \frac{S^2}{R} \right] \quad \theta = \frac{3S_0^2}{2R_0^4} \quad (S_0^2, R_0^4 \text{ are centroids } S^2/3, R^4/5)$$

$$(13) \quad \frac{SdS}{S^2 + 4a\delta m S(1 + \theta) + 2E^4} = \frac{dR}{R}$$

Solving (eq.13) by the quadrature formulae [7] leads to (eq.14), (eq.15) and (eq.16).

In the case that the discriminant is  $\Delta = E^4 - 2a^2m^2(1 + \theta)^2 > 0$ :

$$\frac{1}{2} \log[S^2 + 4a\delta m(1 + \theta)S + 2E^4] - \frac{4a\delta m(1 + \theta)}{2\sqrt{2E^4 - 4a^2m^2(1 + \theta)^2}} \arctan\left(\frac{2S + 4a\delta m(1 + \theta)}{2\sqrt{2E^4 - 4a^2m^2(1 + \theta)^2}}\right) = \log R + K$$

$$(14) \quad K = \frac{S^2 + 4a\delta m(1 + \theta)S + 2E^4}{R^2} \cdot \text{EXP} \left[ \frac{-4a\delta m(1 + \theta)}{\sqrt{2E^4 - 4a^2m^2(1 + \theta)^2}} \arctan\left(\frac{S + 2a\delta m(1 + \theta)}{\sqrt{2E^4 - 4a^2m^2(1 + \theta)^2}}\right) \right]$$

In the case that the discriminant is  $\Delta = E^4 - 2a^2m^2(1+\theta)^2 < 0$ :

$$(15) \quad K = \log \left[ \frac{\log[S^2 + 4a\delta mS(1+\theta) + 2E^4] - \frac{2a\delta m(1+\theta)}{\sqrt{4a^2m^2(1+\theta)^2 - 2E^4}} \cdot \log \left[ \frac{S + 2a\delta m(1+\theta) - \sqrt{4a^2m^2(1+\theta)^2 - 2E^4}}{S + 2a\delta m(1+\theta) + \sqrt{4a^2m^2(1+\theta)^2 - 2E^4}} \right]}{\frac{S^2 + 4a\delta mS(1+\theta) + 2E^4}{R^2} \left[ \frac{S + 2a\delta m(1+\theta) - \sqrt{4a^2m^2(1+\theta)^2 - 2E^4}}{S + 2a\delta m(1+\theta) + \sqrt{4a^2m^2(1+\theta)^2 - 2E^4}} \right]^{\frac{2a\delta m(1+\theta)}{\sqrt{4a^2m^2(1+\theta)^2 - 2E^4}}}} \right] = 2 \log R + K$$

In the case that the discriminant is  $\Delta = E^4 - 2a^2m^2(1+\theta)^2 = 0$ :

$$(16) \quad \frac{dS}{dR} = \frac{1}{S} \cdot \frac{(S + \sqrt{2}E^2)^2}{R} \quad \text{Solving this equation by the quadrature formulae leads}$$

$$K = \frac{S + \sqrt{2}E^2}{R} \text{EXP} \left[ \frac{\sqrt{2}E^2}{S + \sqrt{2}E^2} \right]$$

## 2.2. Conditions of the energy minimum orbit

Since the minimum energy is  $d\varepsilon/dR = 0$  in the SF differential equation (eq. 11), it is a cubic equation in  $\varepsilon$ .

$$0 = \varepsilon^3 \cdot 32a^2r^3(3mr - 2Cr - 6m^2 + 7Cm)^2 + \varepsilon^2 \cdot r^2 \left[ \begin{array}{l} 16a^2(3mr - 2Cr - 6m^2 + 7Cm)^2(2m + C) \\ + 32a^2m(2r^2 + 2mr - Cr - 12m^2)(3mr - 2Cr - 6m^2 + 7Cm) \\ - 4r^3(-r^2 + 6mr - 4Cr - 8m^2)^2 \end{array} \right] + \varepsilon \cdot 4mr \left[ \begin{array}{l} 2a^2m(2r^2 + 2mr - Cr - 12m^2)^2 \\ + 4a^2(2r^2 + 2mr - Cr - 12m^2)(3mr - 2Cr - 6m^2 + 7Cm)(2m + C) \\ - r^3(-r^2 + 8mr - 4Cr - 12m^2)(-r^2 + 6mr - 4Cr - 8m^2) \end{array} \right] + m^2[4a^2(2r^2 + 2mr - Cr - 12m^2)^2(2m + C) - r^3(-r^2 + 8mr - 4Cr - 12m^2)^2]$$

Solve this cubic equation by the method of  $\varepsilon^0$  small coefficient not by Cardano formula. A solution  $\varepsilon_{min}$  (eq. 17) very close to 0 is adopted in accordance with the principle of the energy minimum.

$$(17) \quad \varepsilon_{min} = \frac{-m}{4r} \cdot \frac{r^3(r^2 - 8mr + 4Cr + 12m^2)^2 - 4a^2(2m + C)(2r^2 + 2mr - Cr - 12m^2)^2}{\left[ \begin{array}{l} r^3(r^2 - 8mr + 4Cr + 12m^2)(r^2 - 6mr + 4Cr + 8m^2) \\ - 4a^2(2m + C)(2r^2 + 2mr - Cr - 12m^2)(3mr - 2Cr - 6m^2 + 7Cm) \\ - 2a^2m(2r^2 + 2mr - Cr - 12m^2)^2 \end{array} \right]}$$

$$\cong \frac{-m}{4r} \quad (r \text{ 0 order})$$

$\varepsilon_{min}$  (eq. 17) is substituted into the change of variables  $S = r\sqrt{r(2\varepsilon r + 2m + C)}$  (eq. 12).

$$S^2 = \frac{-mr^4}{2} \cdot \frac{r^4(r^2 - 8mr + 4e^2 + 12m^2)^2 - 4a^2(2mr + e^2)(2r^2 + 2mr - e^2 - 12m^2)^2}{\left[ \begin{array}{l} r^5(r^2 - 8mr + 4e^2 + 12m^2)(r^2 - 6mr + 4e^2 + 8m^2) \\ - 4a^2(2mr + e^2)(2r^2 + 2mr - e^2 - 12m^2)(3mr^2 - 2e^2r - 6m^2r + 7me^2) \\ - 2a^2mr^2(2r^2 + 2mr - e^2 - 12m^2)^2 \end{array} \right]} + r^2(2mr + e^2)$$

$$= \frac{r^4 \times [r^8 \text{ polynomial}] + r^2(2mr + e^2) \times [r^9 \text{ polynomial}]}{[r^9 \text{ polynomial}]} = \frac{r^2 \times P}{Q}$$

$$\cong \frac{3m}{2} r^3 \quad (r \text{ 0 order})$$

Here,  $P$  and  $Q$  are given by (eq.18) and (eq.19).

$$(18) \quad P = -mr^2/2 [ r^4(r^2 - 8mr + 4e^2 + 12m^2)^2 - 4a^2(2mr + e^2)(2r^2 + 2mr - e^2 - 12m^2)^2 ] \\ + (2mr + e^2) \times Q \quad [r^{10} \text{ polynomial}]$$

$$(19) \quad Q = r^5(r^2 - 8mr + 4e^2 + 12m^2)(r^2 - 6mr + 4e^2 + 8m^2) \\ - 4a^2(2mr + e^2)(2r^2 + 2mr - e^2 - 12m^2)(3mr^2 - 2e^2r - 6m^2r + 7me^2) \\ - 2a^2mr^2(2r^2 + 2mr - e^2 - 12m^2)^2 \quad [r^9 \text{ polynomial}]$$

And for  $\theta$ ,

$$\theta = \frac{3S_0^2}{2R_0^4} = \frac{5S^2}{2r^4} = \frac{5P}{2Qr^2} \cong \frac{15m}{4r} \quad (r \text{ at 0 order})$$

### Chapter 3 The Titius –Bode Law

#### 3.1. Demonstration of the Titius-Bode Law

In the case that the discriminant is  $\Delta = E^4 - 2a^2m^2(1 + \theta)^2 > 0$  of the SF differential equation, the function  $f(\theta)$  is given in (eq.14) and is subjected to a Maclaurin series expansion. Terms above  $\theta^2$  are neglected. The result is given in (eq.20).

$$f(\theta) = \frac{S^2 + 4a\delta m(1 + \theta)S + 2E^4}{R^2} \text{EXP} \left[ \frac{-4a\delta m(1 + \theta)}{\sqrt{2E^4 - 4a^2m^2(1 + \theta)^2}} \arctan \left( \frac{S + 2a\delta m(1 + \theta)}{\sqrt{2E^4 - 4a^2m^2(1 + \theta)^2}} \right) \right] \\ -K = 0$$

$$f(\theta) = f(0) + \frac{1}{1!} \cdot \frac{\partial f(0)}{\partial \theta} \theta + \frac{1}{2!} \cdot \frac{\partial^2 f(0)}{(\partial \theta)^2} \theta^2 + \dots = 0$$

$$(20) \quad f(\theta) = \frac{3mr}{2} \text{EXP} \left[ \frac{-4a\delta m}{\sqrt{2E^4 - 4a^2m^2}} \arctan \left( \frac{r\sqrt{3mr}}{2\sqrt{E^4 - 2a^2m^2}} \right) \right] \times \\ \times \left[ 1 - \frac{30a\delta m^2 E^4}{r[2E^4 - 4a^2m^2]^{\frac{3}{2}}} \times \arctan \left( \frac{r\sqrt{3mr}}{2\sqrt{E^4 - 2a^2m^2}} \right) \right] - K = 0$$

Since  $r$  is very large, it is given as  $\arctan \left( \frac{r\sqrt{3mr}}{2\sqrt{E^4 - 2a^2m^2}} \right) = \pi/2 + \pi N - \frac{2\sqrt{E^4 - 2a^2m^2}}{r\sqrt{3mr}}$ . This is substituted into (eq.20).

$$K = \frac{3mr}{2} \text{EXP} \left[ \frac{-2a\delta m\pi(1 + 2N)}{\sqrt{2E^4 - 4a^2m^2}} \right] \cdot \left[ 1 - \frac{30a\delta m^2 E^4}{r[2E^4 - 4a^2m^2]^{\frac{3}{2}}} \cdot \frac{\pi(1 + 2N)}{2} \right]$$

Since the integration constant  $K$  is common to all planets that orbit the center of mass, the base planet and the distance ratio to the base planet can be set as follows.  $r_1, N_1, N - N_1 = n - 1$ , and  $\xi = r/r_1$ . The result is given in (eq.21).

$$(21) \quad n - 1 = \frac{\sqrt{2E^4 - 4a^2m^2}}{4a\delta m\pi} \cdot \log \left[ \frac{\xi - \frac{15a\delta m^2 E^4 \pi (2N_1 + 2n - 1)}{r_1 [2E^4 - 4a^2m^2]^{\frac{3}{2}}}}{1 - \frac{15a\delta m^2 E^4 \pi (2N_1 + 1)}{r_1 [2E^4 - 4a^2m^2]^{\frac{3}{2}}}} \right]$$

On the other hand, the Titius-Bode law is changed into (eq.22).

$$\xi_{Earth} = 0.4 + 0.3 \times 2^n = 0.4 + 0.6 \times 2^{n-1} \quad (\xi_{Earth} : \text{the Earth basis } \xi)$$

$$(22) \quad n - 1 = \frac{1}{\log 2} \cdot \log \frac{\xi_{Earth} - 0.4}{1 - 0.4}$$

The Titius-Bode law (eq. 22) is remarkably similar to the solution (eq. 21) of the approximate SF differential equation. If the two coefficients are the same, the two equations are almost equal.

(The Earth is the base planet,  $n=1$ .)

$$\frac{1}{\log 2} = \frac{\sqrt{2E^4 - 4a^2m^2}}{4a\delta m\pi} \quad 0.4 = \frac{15a\delta m^2 E^4 \pi (2N_1 + 1)}{r_1 [2E^4 - 4a^2m^2]^{\frac{3}{2}}}$$

Since  $r_1 = 1.5 \times 10^8 km$  for the Earth, and,  $m = 1.476 km$  and  $a = 0.32 km$  [8] for the Sun, it is calculated that  $e = 2.1 km$ , and  $N_1 = 1.5 \times 10^7$ . The  $2n$  on the right side of (eq.21) is neglected because of the very large  $N_1$ . Thus,

$$(23) \quad \xi = \left[ 1 - \frac{30a\delta m^2 E^4 \pi N_1}{r_1 [2E^4 - 4a^2m^2]^{\frac{3}{2}}} \right] \cdot \text{EXP} \left[ \frac{4am\pi(n-1)}{\sqrt{2E^4 - 4a^2m^2}} \right] + \frac{30a\delta m^2 E^4 \pi N_1}{r_1 [2E^4 - 4a^2m^2]^{\frac{3}{2}}}$$

$\delta = \pm 1$  is related to the orbital rotation direction.  $\delta(n-1)$  is converted into  $(n-1)$ .

If the rotation direction of orbits is same that of the central core star,  $\delta = +1$ . Since the Titius-Bode law is using the major axis of planetary orbits, the aphelion and perihelion distances are shown by  $r_F$  and  $r_N$ .

$$\xi_{\delta=+1} = \frac{r_F + r_N}{r_{1F} + r_{1N}} = \left[ 1 - \frac{60am^2 E^4 \pi N_1}{(r_{1F} + r_{1N})(2E^4 - 4a^2m^2)^{\frac{3}{2}}} \right] \cdot \text{EXP} \left[ \frac{4am\pi(n-1)}{\sqrt{2E^4 - 4a^2m^2}} \right] + \frac{60am^2 E^4 \pi N_1}{(r_{1F} + r_{1N})[2E^4 - 4a^2m^2]^{\frac{3}{2}}}$$

The form of the upper equation is equal to the Titius-Bode law  $\xi_{Earth} = (1 - 0.4) \cdot 2^{n-1} + 0.4$ .

Therefore, the Titius-Bode law has been demonstrated.

### 3.2. The orbits with counter rotation direction against the fixed star

In the case that  $\delta = -1$  i.e., the rotation direction of orbits and that of the fixed star are opposite each other, the ratio of orbital radii in some exoplanets is reversed against the solar system planets.[9]

$$\xi_{\delta=-1} = \left[ 1 + \frac{60am^2 E^4 \pi N_1}{(r_{1F} + r_{1N})(2E^4 - 4a^2m^2)^{\frac{3}{2}}} \right] \cdot \text{EXP} \left[ \frac{4am\pi(n-1)}{\sqrt{2E^4 - 4a^2m^2}} \right] - \frac{60am^2 E^4 \pi N_1}{(r_{1F} + r_{1N})[2E^4 - 4a^2m^2]^{\frac{3}{2}}}$$

If  $\frac{1}{\log 2} = \frac{\sqrt{2E^4 - 4a^2m^2}}{4am\pi}$ ,  $0.4 = \frac{60am^2 E^4 \pi N_1}{(r_{1F} + r_{1N})[2E^4 - 4a^2m^2]^{\frac{3}{2}}}$  like as the value of solar system planets, then  $\xi_{\delta=+1}$ ,

$\xi_{\delta=-1}$  and  $r_n / r_{n-1}$  are shown in the table1.  $n$  is larger and larger,  $r_n / r_{n-1}$  nears 2.0 in both  $\xi_{\delta=+1}$  and  $\xi_{\delta=-1}$ , but  $r_n / r_{n-1}$  (in  $\xi_{\delta=+1}$ ) is  $1.4 < 2.0$  monotonic increase,  $r_n / r_{n-1}$  (in  $\xi_{\delta=-1}$ ) is  $3.3 > 2.0$  monotonic decrease at  $n=1$ . As a result, there are a few planetary systems with counter rotation direction orbits against the fixed star really.

$$\xi_{\delta=+1} = (1 - 0.4) \cdot 2^{n-1} + 0.4 \quad \xi_{\delta=-1} = (1 + 0.4) \cdot 2^{n-1} - 0.4$$

Table 1. the ratio of orbital radii  $r_n / r_{n-1}$  and  $n$

n	-3	-2	-1	0	1	2	3	4	5	6	7	8
$\xi_{\delta=+1} \quad (1 - 0.4) \cdot 2^{n-1} + 0.4$	0.4	0.5	0.6	0.7	1	1.6	2.8	5.2	10	20	39	77
$r_n / r_{n-1}$	1.0	1.1	1.2	1.3	1.4	1.6	1.8	1.9	1.9	2.0	2.0	2.0
$\xi_{\delta=-1} \quad (1 + 0.4) \cdot 2^{n-1} - 0.4$	-0.3	-0.2	-0.1	0.3	1	2.4	5.2	11	22	44	89	180
$r_n / r_{n-1}$	0.9	0.7	0.2	-6.0	3.3	2.4	2.2	2.1	2.0	2.0	2.0	2.0



4.1. Number of the Saturn's Rings

Since the autorotation of the Saturn is fast, the discriminant is  $\Delta = E^4 - 2a^2m^2(1 + \theta)^2 > 0$ . The solution (eq. 15) of the SF differential equation is as follows for considering the number of rings not for calculating orbits of rings.

$$K = \log \left[ \frac{\frac{S^2 + 4a\delta mS(1 + \theta) + 2E^4}{R^2}}{\left[ \frac{S + 2a\delta m(1 + \theta) - \sqrt{4a^2m^2(1 + \theta)^2 - 2E^4}}{S + 2a\delta m(1 + \theta) + \sqrt{4a^2m^2(1 + \theta)^2 - 2E^4}} \right]^{\frac{2a\delta m(1 + \theta)}{\sqrt{4a^2m^2(1 + \theta)^2 - 2E^4}}}} \right]$$

Since  $\left[ \frac{S + 2a\delta m(1 + \theta) - \sqrt{4a^2m^2(1 + \theta)^2 - 2E^4}}{S + 2a\delta m(1 + \theta) + \sqrt{4a^2m^2(1 + \theta)^2 - 2E^4}} \right]^{\frac{2a\delta m(1 + \theta)}{\sqrt{4a^2m^2(1 + \theta)^2 - 2E^4}}}$  is nearly 1, the denominator is expressed as  $(1 - \lambda)$ .  $\lambda$  is extremely small, but not zero. The solution of the SF differential equation is (eq.24).

$$(24) \quad K = \frac{S^2 + 4a\delta mS(1 + \theta) + 2E^4}{r^2} \cdot \frac{1}{(1 - \lambda)}$$

The integration constant  $K$  is common to all the rings that belong to the Saturn. For the base ring, the variables are  $r_1$  and  $F = K$  and the polynomial of  $S$  is (eq.25).

$$(25) \quad S^4 - 2S^2[F(1 - \lambda)r^2 - 2E^4 + 8a^2m^2(1 + \theta)^2] + [F(1 - \lambda)r^2 - 2E^4]^2 = 0$$

$P$  (eq. 18) and  $Q$  (eq. 19) are substituted into (eq.25) to give  $S$  and  $\theta$ . Finally, the polynomial of  $r$  is (eq.26).

$$(26) \quad Qr^2 (Pr^2 - Q[F(1 - \lambda)r^2 - 2E^4])^2 - 4a^2m^2P(2Qr^2 + 5P)^2 = 0$$

The degree of (eq.26) is the highest at the first term  $P^2Qr^6$ , and is  $r$  to the power of 35  $[10 \times 2 + 9 + 6]$ . That is, (eq.26) is a polynomial of  $r^{35}$  and has four micro roots because of large coefficient  $F$ . And it has maximum 9 main rings because of  $Q = 0$  by coefficient  $F^2$ . Thus, planets with rings such as the Saturn have a maximum of 31 rings (including maximum 9 main rings). The real number of rings decreases because of roots of complex number, minus roots, equal roots and the swelling of the central core star. It is expected to observe and determine the rotation element  $a$  and the electric charge element  $e$ . The coefficients  $\lambda$  and  $F$  contain  $\delta = \pm 1$  related to the orbital rotation direction.

4.2. The rings with counter direction against the central core star

$\delta = \pm 1$  is related to the orbital rotation direction and the coefficients  $\lambda$  and  $F$  contain  $\delta$ . In the case of  $\delta = -1$ , there are a few ring stars with the counter rotation direction rings against the central core star each other really. The planets with the counter rotation direction rings have a maximum of 31 rings. (eq.24) changes as follow.

$$1 - \lambda = \left[ \frac{S - 2am(1 + \theta) - \sqrt{4a^2m^2(1 + \theta)^2 - 2E^4}}{S - 2am(1 + \theta) + \sqrt{4a^2m^2(1 + \theta)^2 - 2E^4}} \right]^{\frac{-2am(1 + \theta)}{\sqrt{4a^2m^2(1 + \theta)^2 - 2E^4}}}$$

$$K = \frac{S^2 - 4amS(1 + \theta) + 2E^4}{r^2} \cdot \frac{1}{(1 - \lambda)}$$

## Chapter 5 Fine Ring star

In the case that the discriminant is  $\Delta = E^4 - 2a^2m^2(1 + \theta)^2 = 0$  because the balance of the mass, rotation and electric charge elements is exquisite.

$$\frac{dS}{dR} = \frac{1}{S} \left[ \frac{2E^4}{R} + \frac{4a\delta mS}{R} \left( 1 + \frac{3S^2}{2R^4} \right) + \frac{S^2}{R} \right] \quad E^4 = e^2(e^2 + 2m^2)$$

$$\text{the discriminant } \Delta = 2E^4 - 4a^2m^2 \left( 1 + \frac{3S^2}{2R^4} \right)^2 = 0 \quad \textcircled{1}$$

the differential equation  $dS/dR = 1/S \cdot (S + \sqrt{2}E^2)^2/R$  solving this equation

$$K = \frac{S + \sqrt{2}E^2}{R} \text{EXP} \left( \frac{\sqrt{2}E^2}{S + \sqrt{2}E^2} \right) \quad \textcircled{2}$$

$$\text{the energy stable equation } S = r\sqrt{3mr/2} \quad \textcircled{3}$$

$r, S$  and  $K$  are unknown, then solve simultaneous equations  $\textcircled{1}, \textcircled{2}, \textcircled{3}$

$$(27) \quad r = \frac{9a\delta m^2}{2\sqrt{2}(E^2 - \sqrt{2}a\delta m)} \quad (\text{in this case, } \delta = +1)$$

$m, e$  and  $a$  are small constant value, but if it is  $E^2 - \sqrt{2}a\delta m \simeq 0$  generally,  $r$  grows larger. That is, in the case of exquisite balances  $a^2 \simeq e^2(1 + e^2/2m^2)$ , Super Fine-Ring is formed. In the case that the discriminant is  $\Delta \neq 0$  slightly out of alignment, Fine-Ring is formed and has some components of the Saturn's rings or Solar System planets.

## Chapter 6 The Elliptical Sphere Planets System

In the case that the central core star of the elliptical sphere planets system has no rotation like an elliptical galaxy,  $a = 0$  in (eq.11) gives (eq.28)

$$(28) \quad \frac{d\varepsilon}{dR} + \frac{2(R^2 - 6mR + 4CR + 8m^2)}{R(R^2 - 4mR + 2CR + 4m^2)} \cdot \varepsilon = \frac{m(-R^2 + 8mR - 4CR - 12m^2)}{R^2(R^2 - 4mR + 2CR + 4m^2)}$$

Also, the condition of the minimum and maximum energy is  $\partial\varepsilon/\partial R = 0$ .

$$\varepsilon = \frac{m(-r^2 + 8mr - 4e^2 - 12m^2)}{2r(r^2 - 6mr + 4e^2 + 8m^2)}$$

Solve the differential equation (eq.28) and add the condition of the energy minimum, then the integration constant  $K$  is as follows.

$$K = \frac{mr^3(-r^2 + 8mr - 12m^2 - 4e^2)}{2(r^2 - 6mr + 8m^2 + 4e^2)(r^2 - 4mr + 4m^2 + 12e^2)} + \frac{(r^2 - 4mr + 14m^2 + e^2)mr - 6m^2(2m^2 + e^2)}{r^2 - 4mr + 4m^2 + 2e^2} - \frac{m(2m^2 + e^2)\pi(1 + 2N)}{2\sqrt{2}e}$$

The integration constant  $K$  is common to all orbits which belong to the elliptical sphere planets system. At the base orbit they are  $r_1, N_1, N - N_1 = n - 1$ . It is (eq.29).

$$(29) \quad \left[ \frac{r^3(-r^2 + 8mr - 12m^2 - 4e^2)(r^2 - 4mr + 4m^2 + 2e^2) + 2r(r^2 - 4mr + 14m^2 + e^2)(r^2 - 6mr + 8m^2 + 4e^2)(r^2 - 4mr + 4m^2 + 12e^2) - 12m(2m^2 + e^2)(r^2 - 6mr + 8m^2 + 4e^2)(r^2 - 4mr + 4m^2 + 12e^2)}{(r^2 - 6mr + 8m^2 + 4e^2)(r^2 - 4mr + 4m^2 + 12e^2)(r^2 - 4mr + 4m^2 + 2e^2)} \right]$$

$$= F - \frac{\sqrt{2}(2m^2 + e^2)\pi(n-1)}{e}$$

$F$  is equal to (eq.29) at  $r=r_1$ . (eq.29) is the polynomial of  $r^7$  with the minimum and maximum energy which has coefficients of natural variable number  $n$ . The elliptical sphere planets system is composed of maximum  $4 \times$  number  $n$  spherical surfaces with the minimum energy. There are planet orbits in the spherical surfaces. The real number of spherical surfaces decreases by roots of complex number, minus roots, equal roots and the swelling of the central core star.

## Chapter 7 Summary

The Titius-Bode law, discovered 250 years ago, is considered to be a mathematical coincidence rather than an "exact" law. However, I successfully proved the law and showed here that the Saturn can physically have a maximum of 31 rings. 250 years of astronomical mystery is now solved not by the computer analysis but by the theoretical analysis.

The Kerr-Newman solution of the Einstein's equation is considered as follows. The no-hair theorem postulates that all black hole solutions of the Einstein-Maxwell equations of gravitation and electromagnetism in general relativity can be completely characterized by only three externally observable classical parameters: mass, electric charge, and angular momentum. [10], [11]

In this manner, since this theory is based on the steady state of Kerr-Newman solution in the mature galaxies, it cannot be applied to the galaxies which are still young, unstable and transitional. Three important equations can be summarized as follows.

(eq.11) is a fundamental differential equation based on the steady state, and it can be applied to the Solar system, other planets and rings in the galaxies. There must be many solutions of (eq.11).

(eq.23) is one of the approximate solutions of (eq.11). Since this is energetically stable, it is applicable to the Solar system planets and many of the around 4000 extrasolar planets in the galaxies. A few planetary systems with counter direction against the celestial star exist. However, it is not applicable to still young, unstable and transitional planets like comets.

(eq.26) is one of the approximate solutions of (eq.11). This is also energetically stable and applicable to Saturn's rings and some other extrasolar planets' rings.

$m$ ,  $e$  and  $a$  are small constant value, but in the case of exquisite balances  $a^2 \approx e^2(1 + e^2/2m^2)$ , Super Fine Ring is formed (eq.27).

This theory is applied to planets, which belong to the center of mass in the universe, but it is not available to a bulge space near center of mass. It is important to know that (eq.11) is established by ignoring the energy momentum tensor of planets.

$$\begin{aligned}
 (11) \quad & \frac{d\varepsilon}{dR} R^4 (R^2 - 4mR + 2CR + 4m^2) \\
 & = mR^2(-R^2 + 8mR - 4CR - 12m^2) + \varepsilon \cdot 2R^3(-R^2 + 6mR - 4CR - 8m^2) \\
 & \quad + 2am(2R^2 + 2mR - CR - 12m^2)\delta\sqrt{R(2\varepsilon R + 2m + C)} \\
 & \quad + \varepsilon \cdot 4aR(3mR - 2CR - 6m^2 + 7Cm)\delta\sqrt{R(2\varepsilon R + 2m + C)} \quad C = e^2/R \quad (R \text{ 2ry order})
 \end{aligned}$$

$$(23) \quad \xi = \left[ 1 - \frac{30a\delta m^2 E^4 \pi N_1}{r_1 [2E^4 - 4a^2 m^2]^{\frac{3}{2}}} \right] \cdot EXP \left[ \frac{4am\pi(n-1)}{\sqrt{2E^4 - 4a^2 m^2}} \right] + \frac{30a\delta m^2 E^4 \pi N_1}{r_1 [2E^4 - 4a^2 m^2]^{\frac{3}{2}}}$$

$$(26) \quad Qr^2 (Pr^2 - Q [F(1-\lambda)r^2 - 2E^4])^2 - 4a^2 m^2 P (2Qr^2 + 5P)^2 = 0$$

$P$  is (eq.18) and  $Q$  is (eq.19).  $E^4 = e^2(e^2 + 2m^2)$

$$(27) \quad r = \frac{9a\delta m^2}{2\sqrt{2}(E^2 - \sqrt{2}a\delta m)}$$

By means of distance from the center of mass, spiral planets system and elliptical planets system are organized theoretically as follow, using mass element  $m (\neq 0)$ , electric charge element  $e$  and angular momentum element  $a$  of the center of mass. In considering attenuation of  $a$  for a long universal time, elliptical planets system is a late form of planets.

Spiral planets system

$$a \neq 0 \quad e^4 + 2e^2 m^2 - 2a^2 m^2 < 0 \quad r^{35} \text{ polynomial group solution (ex. Saturn's rings)}$$

$$a \neq 0 \quad e^4 + 2e^2 m^2 - 2a^2 m^2 > 0 \quad \text{periodic solution SF/Titius law (ex. Solar sytem planets)}$$

Elliptical planets system

$$a = 0 \quad e \neq 0 \quad \text{periodic solution * polynomial solution}$$

$$a = 0 \quad e = 0 \quad r^3 \text{ polynomial solution}$$

## Appendix

### Analysis solution and Numerical solution

Without use of Space Fantasy differential equation which leads the analysis solution, there is a method of numerical solution i.e. simultaneous equations consist of an integral constant equation and an energy minimum orbit equation, following.

However, a numerical solution composed of power algebraic expressions is difficult to understand the characteristics of orbits

$$(8) \quad J = \frac{4am + R\delta\sqrt{R(2\varepsilon R + 2m + C)}}{R^2(R - 2m + C) - a(2m - C)\delta\sqrt{R(2\varepsilon R + 2m + C)}} R^2 \quad \delta = \pm 1 \quad C = e^2/R$$

An integral constant equation  $K$

$$K = \frac{4am + R\delta\sqrt{R(2\varepsilon R + 2m + C)}}{R^2(R - 2m + C) - a(2m - C)\delta\sqrt{R(2\varepsilon R + 2m + C)}} R^2$$

An energy minimum orbit equation

$$0 = \lim \left( \frac{d\varepsilon}{dR} \rightarrow 0 \right) \left\{ \frac{d}{d\varepsilon} \frac{4am + R\delta\sqrt{R(2\varepsilon R + 2m + C)}}{R^2(R - 2m + C) - a(2m - C)\delta\sqrt{R(2\varepsilon R + 2m + C)}} R^2 \right\}$$

## References

- 1) Internet Titius-Bode law - Wikipedia  
[https://en.wikipedia.org/wiki/Titius%E2%80%93Bode\\_law](https://en.wikipedia.org/wiki/Titius%E2%80%93Bode_law), accessed in Jan 2018.

- 2) Internet    A demonstration of the Titius–Bode law and the number of Saturn’s rings by Newtonian methods using the Kerr-Newman solution of the general relativity theory  
<https://sayuri-fumitaka.icurus.jp>, accessed in Oct 2018.
- 3) Internet    Boyer–Lindquist coordinates - Wikipedia  
[https://en.wikipedia.org/wiki/Boyer%E2%80%93Lindquist\\_coordinates](https://en.wikipedia.org/wiki/Boyer%E2%80%93Lindquist_coordinates), accessed in Jan 2018.
- 4) Internet    General Relativity, Black Holes and Cosmology, Andrew J S. Hamilton  
[http://jila.colorado.edu/~ajsh/ast5770\\_14/grbook.pdf#search=%27general+relativity%2C+black+hole+and+cosmology%27](http://jila.colorado.edu/~ajsh/ast5770_14/grbook.pdf#search=%27general+relativity%2C+black+hole+and+cosmology%27), accessed in Jan 2018.
- 5) Internet    Euler-Lagrange Differential Equation  
<http://mathworld.wolfram.com/Euler-LagrangeDifferentialEquation.html>, accessed in Jan 2018.
- 6) Internet    Riccati equation - Wikipedia (similar to Japanese)  
[https://en.wikipedia.org/wiki/Riccati\\_equation](https://en.wikipedia.org/wiki/Riccati_equation), accessed in Jan 2018.
- 7) Formeln+Hilfen Höhere Mathematik, 2013 (translated into Japanese)  
 Gerhard Merziger, Günter Mühlbach, Detlef Wille, Thomas Wirth.
- 8) Exploring Black Holes: Introduction to General Relativity, 2000, Edwin F. Taylor, John Archibald Wheeler (p272, translated into Japanese by Nobuyoshi Makino)
- 9) Winn & Fabrycky (2015). “The Occurrence and Architecture of Exoplanetary Systems”. *Annual Review of Astronomy and Astrophysics* 53: p.409-
- 10) Internet    No-hair theorem - Wikipediaen.  
[https://en.wikipedia.org/wiki/No-hair\\_theorem](https://en.wikipedia.org/wiki/No-hair_theorem), accessed in Jan 2018.
- 11) Misner, Charles W.; Thorne, Kip S.; Wheeler, John Archibald (1973). *Gravitation*. San Francisco: W. H. Freeman. pp. 875–876. ISBN 0716703343. Retrieved 24 January 2013.
- 12) 2017 Autumn Annual Meeting Proceedings, Astronomical Society of Japan. (jp) ,    Fumitaka Inuyama  
 2018 Spring Annual Meeting Proceedings, Astronomical Society of Japan. (en) ,    Fumitaka Inuyama  
 2019 Autumn Annual Meeting Proceedings, The Japanese Society for Planetary Sciences (jp) ,  
Fumitaka Inuyama



spiral galaxy (reference page 3)



elliptical galaxy (reference page 9)



Solar System planets (reference page 7)



Saturn's rings (reference page 8)

courtesy of NASA



Fine Ring Nebula — captured here by the ESO Faint Object Spectrograph and Camera mounted on the New Technology Telescope at the La Silla Observatory in Chile. Credit ESO. (reference page 10)