

Surveillance for Infections in Long-term Care

Evelyn Cook, RN, CIC
Associate Director SPICE

**How confident are you that your facility has
a strong infection prevention program that
includes all the recommended elements?**


Completely confident

Somewhat confident


Not confident

Have no idea

Do you believe you have the skills and the qualifications to be the individual with oversight of the infection prevention program?



If you wanted to compare your infection surveillance data to another nursing home in your community that cared for a similar resident population, how confident are you that events will be tracked in the same way





What Standardized Definition Does Your Facility Use for Surveillance

National Healthcare
Safety Network (NHSN)

Revised McGeer
Definitions

Loeb criteria

When the physician
documents an infection

No standardized criteria

Start the presentation to see live content. Still no live content? Install the app or get help at PollEv.com/app

SPICE 



Objectives

- Discuss the importance of surveillance
- Describe standardized surveillance definitions
- Discuss ways to implement and apply surveillance definitions

- “Surveillance is a comprehensive method of measuring outcomes and related processes of care, analyzing the data, and providing information to members of the healthcare team to assist in improving those outcomes and processes”

Surveillance in LTCF

- The LTCF should have a system for ongoing collection of data on infections in the institution (Cat IC)





Types of Surveillance

- **Total (or Whole) House Surveillance**
 - All HAIs are monitored in the entire population
 - Calculate rates for specific population (not an overall facility wide rate)
 - Example: Monitoring all UTIs, report CAUTIs separately
- **Targeted Surveillance**
 - Care units
 - Infections related to medical devices
 - Organisms of epidemiological importance
- **Combination Surveillance Strategy**
 - Most use a combination and monitor targeted events that occur in defined populations while concurrently monitoring select HAIs and laboratory reports from house-wide locations

Selection of Processes and Outcomes

Processes

- Hand hygiene
- Urinary Catheter insertion/maintenance

Outcomes

- Acute respiratory infections
- Urinary tract infections
- Skin/Soft Tissue Infections
- Gastroenteritis



SPICE 

Consideration for choosing Outcomes

- Mandatory/required
- Frequency (incidence) of the infection
- Communicability
- System/resident cost (↑mortality, hospitalization)
- Early Detection

Outcomes selected for surveillance should be re-evaluated annually as a component of the IP risk assessment

SPICE 

Infections that should be included in routine surveillance

Points to Consider	Infections	Comments
Evidence of transmissibility in a healthcare setting	Viral respiratory tract infections, viral GE, and viral conjunctivitis	Associated with outbreaks among residents and HCP in LTCFs
Processes available to prevent acquisition of infection		
Clinically significant cause of morbidity or mortality	Pneumonia, UTI, GI tract infections, (including <i>C. difficile</i>) and SSTI	Associated with hospitalization and functional decline in LTCF residents
Specific pathogens causing serious outbreaks	Any invasive group A <i>Streptococcus</i> infection, acute viral hepatitis, norovirus, scabies, influenza-COVID-19	A single laboratory-confirmed case should prompt further investigation

Infections that could be included in routine surveillance

Points to Consider	Infections	Comments
Infections with limited transmissibility in a healthcare settings	Ear and sinus infections, fungal oral and skin infections and herpetic skin infections	Associated with underlying comorbid conditions and reactivation of endogenous infection
Infections with limited preventability		

Infections for which other accepted definitions should be applied in LTCF surveillance

Points to Consider	Infections	Comments
Infections with other accepted definitions (may apply to only specific at-risk residents)	Surgical site infections, central-line- associated bloodstream infections and ventilator-associated pneumonia	LTCF-specific definitions were not developed. Refer to the National Healthcare Safety Network's criteria



Sources of Data for Surveillance

- Clinical ward/unit rounds
- Medical Chart
- Lab reports
- Kardex/Patient Profile/Temperature logs
- Antibiotic Starts



IMPLEMENTING AND APPLYING SURVEILLANCE DEFINITIONS AND PRINCIPALS



Surveillance

- *The facility's surveillance system must include a data collection tool and the use of nationally-recognized surveillance criteria such as but not limited to CDC's National Healthcare Safety Network (NHSN) Long Term Care Criteria to define infections or updated McGeer criteria*



Guiding Principles for Using Standardized Definitions



Infection surveillance only

Identify trends in a population



Applied retrospectively as it relates to clinical diagnosis/treatment

Limited elements



Focus on transmissible/preventable infections

Not for clinical decision making

SPICE 

Minimum Criteria for Initiation of Antibiotics in Long-Term Care Residents

Suspected Urinary Tract Infection

Loeb et al. Development of Minimum Criteria for the Initiation of Antibiotics in Residents of Long-Term Care Facilities: Results of a Consensus Conference.

Inf Control Hosp Epi. 2001

SPICE 

The screenshot shows the CDC NHSN website. The browser address bar displays www.cdc.gov/nhsn/ltc/uti/index.html. The page header includes the CDC logo and the text "Centers for Disease Control and Prevention" with the tagline "CDC 24/7. Saving Lives. Protecting People™". A search bar is located in the top right corner.

The main navigation menu on the left lists various facility types: Ambulatory Surgery Centers, Acute Care Hospitals/Facilities, Long-term Acute Care Hospitals/Facilities, and Long-term Care Facilities. The "Surveillance for C. difficile and MRSA Infections" section is highlighted.

The main content area is titled "Surveillance for C. difficile, MRSA, and other Drug-resistant Infections". It includes social media icons for Facebook, Twitter, and YouTube. Below this, there is a section for "Resources for NHSN Users Already Enrolled" with expandable categories: Training, Protocol, Data Collection Forms, Supporting Material, and Analysis Resources. A "New Users - Start Here" sidebar provides a three-step enrollment process: Step 1: Enroll into NHSN, Step 2: Set up NHSN, and Step 3: Report, with a link to "Click here to enroll".

The Windows taskbar at the bottom shows the date and time as 5:32 PM on 9/13/2016, along with various application icons.

Purposes of NHSN



Provide facilities with risk-adjusted data that can be used for inter-facility comparisons and local quality improvement activities



Assist facilities in developing surveillance and analysis methods that permit timely recognition of patient and healthcare personnel safety problems and prompt intervention with appropriate measures




Conduct collaborative research studies with members



Data repository for reporting COVID-19

Long-term Care Facility Component NHSN



<https://www.cdc.gov/nhsn/training/ltc/index.html>

SPICE 

NHSN LTCF

COVID-19 Module for LTCF

Resident
Impact &
Facility
Capacity

Staff &
Personnel
Impact

Supplies &
Personal
Protective
Equipment

Ventilator
Capacity &
Supplies

SPICE 

Surveillance Definitions of Infections in Long-Term Care Facilities: Revisiting the McGeer Criteria

Nimalie D. Stone, MD;¹ Muhammad S. Ashraf, MD;² Jennifer Calder, PhD;³ Christopher J. Crnich, MD;⁴
Kent Crossley, MD;⁵ Paul J. Drinka, MD;⁶ Carolyn V. Gould, MD;¹ Manisha Juthani-Mehta, MD;⁷
Ebbing Lautenbach, MD;⁸ Mark Loeb, MD;⁹ Taranisia MacCannell, PhD;¹ Preeti N. Malani, MD;^{10,11} Lona Mody, MD;^{10,11}
Joseph M. Mylotte, MD;¹² Lindsay E. Nicolle, MD;¹³ Mary-Claire Roghmann, MD;¹⁴ Steven J. Schweon, MSN;¹⁵
Andrew E. Simor, MD;¹⁶ Philip W. Smith, MD;¹⁷ Kurt B. Stevenson, MD;¹⁸ Suzanne F. Bradley, MD^{10,11}
for the Society for Healthcare Epidemiology Long-Term Care Special Interest Group*



Attribution of infection to LTCF

- No evidence of an incubating infection at the time of admission to the facility
 - Basis of clinical documentation of appropriate signs and symptoms and not solely on screening microbiologic data
- Onset of clinical manifestation occurs > 2 calendar days after admission.



Attribution of infection to LTCF

- All symptoms must be new or acutely worse
- Non-infectious causes of signs and symptoms should always be considered prior to diagnosis
- Identification of an infection should not be based on a single piece of evidence
 - Clinical, microbiologic, radiologic
- Diagnosis by physician insufficient (based on definition)

Attribution of infection to LTCF

- A “new, nursing home onset” refers to COVID-19 cases that originated in the nursing home, and not cases where the nursing home admitted individuals from a hospital with a known COVID-19 positive status, or unknown COVID-19 status but became COVID-19 positive within 14 days after admission.

Constitutional Requirements

Fever:

- A single oral temperature $>37.8^{\circ}\text{C}$ [100°F], OR
- Repeated oral temperatures $>37.2^{\circ}\text{C}$ [99°F];
rectal temperature $>37.5^{\circ}$ (99.5°F) OR
- $>1.1^{\circ}\text{C}$ [2°F] over baseline from a temperature
taken at any site



SPICE 

Constitutional Requirements

Leukocytosis

- Neutrophilia > 14000 WBC/ mm^3
OR
- Left shift ($>6\%$ bands or ≥ 1500 bands/ mm^3)



SPICE 

Constitutional Requirements

Acute Change in Mental Status from Baseline

- Based on Confusion Assessment Method (CAM) criteria available in MDS

Change	Criteria
Acute Onset	Evidence of acute change in mental status from resident baseline
Fluctuating	Behavior fluctuating (e.g., coming and going or changing in severity during assessment)
Inattention	Resident has difficulty focusing attention (e.g., unable to keep track of discussion or easily distracted)
Disorganized Thinking	Resident's thinking is incoherent (e.g., rambling conversation, unclear flow of ideas)
Altered level of consciousness	Resident's level of consciousness is described as different from baseline (e.g., hyperalert, sleepy, drowsy, difficult arouse, nonresponsive)

SPICE 

Constitutional Requirements

Acute Functional Decline

- New 3 point increase in total ADL score (0-28) from baseline based on 7 ADLs {0 = independent; 4 = total dependence}
 1. Bed mobility
 2. Transfer
 3. Locomotion within LTCF
 4. Dressing
 5. Toilet use
 6. Personal hygiene
 7. Eating

SPICE 



Site Specific Definitions

Knowledge Checks



SPICE

Respiratory Tract Infections

Criteria	Comments
<p>A. <u>Common cold syndrome/pharyngitis</u></p> <p>At least two criteria present</p> <ol style="list-style-type: none">1. Runny nose or sneezing2. Stuffy nose (i.e., congestion)3. Sore throat or hoarseness or difficulty swallowing4. Dry cough5. Swollen or tender glands in neck	<p>Fever may or may not be present. Symptoms must be new, and not attributable to allergies</p>

SPICE

Respiratory Tract Infections

Criteria	Comments
<p>B. <u>Influenza-like Illness</u></p> <p>Both criteria 1 and 2 present</p> <ol style="list-style-type: none"> 1. Fever 2. At least three of the following symptom sub-criteria (a-f) present <ol style="list-style-type: none"> a. Chills b. New headache or eye pain c. Myalgias or body aches d. Malaise or loss of appetite e. Sore throat f. New or increased dry cough 	<p>If criteria for influenza-like illness and another upper or lower respiratory tract infection are met at the same time, only the diagnosis of influenza-like illness should be used</p> <p>Due to increasing uncertainty surrounding the timing of the start of influenza season, the peak of influenza activity and the length of the season, 'seasonality' is no longer part of the criteria to define influenza-like illness</p>




Respiratory Tract Infections

Criteria	Comments
<p>C. <u>Pneumonia</u></p> <p>All criteria 1-3 present</p> <ol style="list-style-type: none"> 1. Interpretation of chest radiograph as demonstrating pneumonia or the presence of <u>new</u> infiltrate 2. At least one of the following respiratory sub-criteria (a-f) present <ol style="list-style-type: none"> a. New or increased cough b. New or increased sputum production c. O₂ saturation <94% on room air or a reduction in O₂ saturation of more than 3% from baseline d. New or changed lung exam abnormalities e. Pleuritic chest pain f. Respiratory rate of ≥ 25/min 3. At least one constitutional criteria 	<p>For both pneumonia and lower respiratory tract infections, presence of underlying conditions which could mimic a respiratory tract infection presentation (congestive heart failure, interstitial lung disease), should be excluded by review of clinical records and an assessment of presenting symptoms and signs</p>



Respiratory Tract Infections

Criteria	Comments
D. <u>Lower respiratory tract (Bronchitis or Tracheo-bronchitis)</u>	
All criteria 1-3 present	For both pneumonia and lower respiratory tract infections, presence of underlying conditions which could mimic a respiratory tract infection presentation (congestive heart failure, interstitial lung disease), should be excluded by review of clinical records and an assessment of presenting symptoms and signs
1. Chest radiograph not performed <u>or</u> negative for pneumonia or new infiltrate.	
2. At least two of the following respiratory sub-criteria (a-f) present	
a. New or increased cough	
b. New or increased sputum production	
c. O ₂ saturation <94% on room air or a reduction in O ₂ saturation of more than 3% from baseline	
d. New or changed lung exam abnormalities	
e. Pleuritic chest pain	
f. Respiratory rate of ≥ 25/min	
3. At least one constitutional criteria	



Questions?

Urinary Specimens: What do the Guidelines Say?

- Specimens collected through the catheter present for more than a few days reflect biofilm microbiology.
- For residents with chronic indwelling catheters and symptomatic infection, changing the catheter immediately prior to instituting antimicrobial therapy allows collection of a bladder specimen, which is a more accurate reflection of infecting organisms.
- Urinary catheters coated with antimicrobial materials have the potential to decrease UTIs but have not been studied in the LTCF setting.

SHEA/APIC Guideline: Infection prevention and control in the long-term care facility Philip W. Smith, MD, Gail Bennett, RN, MSN, CICb Suzanne Bradley, MD, Paul Drinka, MD, Ebbing Lautenbach, MD, James Marx, RN, MS, CIC, Lona Mody, MD, Lindsay Nicolle, MD and Kurt Stevenson, MD July 2008




McGeer Urinary Tract Infections

Criteria		Comments
<p>A. <u>For Residents without an indwelling catheter</u></p> <p>Both criteria 1 and 2 present</p> <p>1. At least one of the following sign/symptom sub-criteria (a-c) present:</p> <p>a) Acute dysuria <u>or</u> acute pain, swelling, or tenderness of the testes, epididymis, or prostate</p> <p>b) Fever <u>or</u> leukocytosis and</p> <p>At least one of the following localizing urinary tract sub-criteria:</p> <p>i. Acute costovertebral angle pain or tenderness</p> <p>ii. Suprapubic pain</p> <p>iii. Gross hematuria</p> <p>iv. New or marked increase in incontinence</p> <p>v. New or marked increase in urgency</p> <p>vi. New or marked increase in frequency</p>	<p>c) In the absence of fever of leukocytosis, then at least two or more of the following localizing urinary symptoms</p> <p>i. Suprapubic pain</p> <p>ii. Gross hematuria</p> <p>iii. New or marked increase in incontinence</p> <p>iv. New or marked increase in urgency</p> <p>v. New or marked increase in frequency</p> <p>2. One of the following microbiologic sub-criteria</p> <p>a) $\geq 10^5$ cfu/ml of no more than 2 species of microorganisms in a voided urine</p> <p>b) $\geq 10^2$ cfu/ml of any number of organisms in a specimen collected by an in and out catheter</p>	<p>UTI should be diagnosed when there are localizing s/s <u>and</u> a positive urinary culture</p> <p>A diagnosis of UTI can be made without localizing symptoms if a blood culture isolate of the same organism isolated from the urine and there is no alternate sight of infection</p> <p>In the absence of a clear alternate source, fever or rigors with a positive urine culture in a non-catheterized resident will often be treated as a UTI. However evidence suggest most of the these episodes are not from a urinary source</p> <p>Pyuria does not differentiate symptomatic UTI from asymptomatic bacturia</p> <p>Absence of pyuria in diagnostic test excludes symptomatic UTI in residents of LTCF</p> <p>Urine specimens should be processed within 1-2 hours, or refrigerated and processed with in 24 hours.</p>

McGeer Urinary Tract Infections

Criteria	Comments
<p>B. <u>For the resident with an indwelling catheter</u></p> <p>Both criteria 1 and 2 present</p> <p>1. At least one of the following sign/symptom sub-criteria (a-d) present:</p> <ul style="list-style-type: none"> a) Fever, rigors, or new onset hypotension, with no alternate site of infection b) Either acute change in mental status <u>or</u> acute functional decline with no alternate diagnosis <u>and</u> Leukocytosis c) New onset suprapubic pain <u>or</u> costovertebral angle pain or tenderness d) Purulent discharge from around the catheter <u>or</u> acute pain, swelling, or tenderness of the testes, epididymis, or prostate <p>2. Urinary catheter culture with $\geq 10^5$ cfu/ml of any organism(s)</p>	<p>Recent catheter trauma, catheter obstruction or new onset hematuria are useful localizing signs consistent with UTI, but not necessary for diagnosis</p> <p>Urinary catheter specimens for culture should be collected following the replacement of the catheter (if current catheter has been in place for >14 days)</p>



NHSN
National Healthcare
Safety Network



NHSN Notes

- Indwelling urinary catheter should be in place for a minimum of 2 calendar days before infection onset (day 1 = day of insertion)
- Indwelling urinary catheter: a drainage tube that is inserted into the urinary bladder through the urethra, is left in place and is connected to a closed collection system, also called a foley catheter. Indwelling urinary catheters do not include straight in-and-out catheters or suprapubic catheters (these would be captures as SUTIs, not CA-SUTIs)
- *Indwelling catheters which have been in place for > 14 days should be changed prior to specimen collection but failure to change catheter does not exclude a UTI for surveillance purposes*



Must meet Criteria 1,2 OR 3		
<p>NHSN Urinary Tract Infections For Residents without an indwelling catheter</p>	<p>Criteria 1</p> <p>Either of the following:</p> <ol style="list-style-type: none"> 1. Acute dysuria 2. Acute pain, swelling or tenderness of the testes, epididymis or prostate <p style="text-align: center;">AND</p> <p>A positive urine culture with no more than 2 species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/ml</p>	<p>Criteria 2</p> <p>Either of the following:</p> <ol style="list-style-type: none"> 1. Fever: (Single temperature $>100^\circ$ F or $>99^\circ$ F on repeated occasions OR an increase of $>2^\circ$ F over baseline 2. Leukocytosis: $>14,000$ cells/mm³ or left shift (6% or 1, 500 bands/mm³) <p style="text-align: center;">AND</p> <p>One or more of the following (New or Marked increase):</p> <ol style="list-style-type: none"> 1. Costovertebral angle pain or tenderness 2. Suprapubic tenderness 3. Gross hematuria 4. Incontinence 5. Urgency 6. Frequency <p style="text-align: center;">AND</p> <p>A positive urine culture with no more than 2 species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/m</p>
	<p>Criteria 3</p> <p>Two or more of the following (New and/or marked increase):</p> <ol style="list-style-type: none"> 1. Costovertebral angle pain or tenderness 2. Incontinence 3. Urinary urgency 4. Urinary frequency 5. Suprapubic tenderness 6. Visible (gross) hematuria <p style="text-align: center;">AND</p> <p>A positive urine culture with no more than 2 species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/ml</p>	


Comments: Fever can be used to meet SUTI criteria even if the resident has another possible cause for the fever (for example, pneumonia)

NHSN Urinary Tract Infections

For the resident with an indwelling catheter

Criteria	Comments
<p><u>CA-SUTI</u></p> <p>One or more of the following (Signs and Symptoms and Laboratory and diagnostic Testing):</p> <ul style="list-style-type: none"> a) *Fever (Single temperature >100° F or >99° F on repeated occasions OR an increase of >2° F over baseline) b) Rigors c) New onset hypotension, with no alternate site of infection d) New onset confusion/functional decline AND Leukocytosis (>14,000 cells/mm³ or left shift (6% or 1, 500 bands/mm³)) e) New or marked increase in costovertebral angle pain or tenderness f) New or marked increase in suprapubic tenderness g) Acute pain, swelling, or tenderness of the testes, epididymis, or prostate h) Purulent discharge from around the catheter 	<p style="text-align: center;">AND</p> <p style="text-align: center;">↓</p> <p style="color: red;">A positive urine culture with no more than 2 species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/ml</p> <p><i>*Fever can be used to meet CA-SUTI criteria even if the resident has another possible cause for the fever (for example, pneumonia)</i></p>

Questions?



Skin, Soft Tissue and Mucosal Infections

Criteria	Comments
<p>A. <u>Cellulitis/soft tissue/wound infection</u></p> <p>At least one of the following criteria is present</p> <ol style="list-style-type: none"> 1. Pus present at a wound, skin, or soft tissue site 2. New or increasing presence of at least four of the following sign/symptom sub-criteria <ol style="list-style-type: none"> a) Heat at affected site b) Redness at affected site c) Swelling at affected site d) Tenderness or pain at affected site e) Serous drainage at affected site f) One constitutional criteria 	<p>More than one resident with streptococcal skin infection from the same serogroup (e.g., A, B, C, G) in a LTCF may suggest an outbreak</p> <p>For wound infections related to surgical procedures: LTCF should use the CDC's NHSN surgical site infection criteria and report these infections back to the institution performing the original surgery</p> <p>Presence of organisms cultured from the surface (e.g., superficial swab culture) of a wound is not enough evidence that the wound is infected</p>



Skin, Soft Tissue and Mucosal Infections

Criteria	Comments
<p>B. <u>Scabies</u></p> <p>Both criteria 1 and 2 present</p> <ol style="list-style-type: none"> 1. A maculopapular and/or itching rash 2. At least one of the following sub-criteria: <ol style="list-style-type: none"> a) Physician diagnosis b) Laboratory confirmation (scrapping or biopsy) c) Epidemiologic linkage to a case of scabies with laboratory confirmation 	<p>Care must be taken to rule out rashes due to skin irritation, allergic reactions, eczema, and other non-infectious skin conditions</p> <p>An epidemiologic linkage to a case can be considered if there is evidence of geographic proximity in the facility, temporal relationship to the onset of symptoms, or evidence of a common source of exposure (i.e., shared caregiver).</p>



Skin, Soft Tissue and Mucosal Infections

Criteria	Comments
<p>C. <u>Fungal oral/perioral and skin infections</u></p> <p><u>Oral candidiasis:</u></p> <p>Both criteria 1 and 2 present:</p> <ol style="list-style-type: none"> 1. Presence of raised white patches on inflamed mucosa, or plaques on oral mucosa 2. Medical or dental provider diagnosis <p><u>Fungal skin Infection:</u></p> <p>Both criteria 1 and 2 present:</p> <ol style="list-style-type: none"> 1. Characteristic rash or lesion 2. Either a medical provider diagnosis or laboratory confirmed fungal pathogen from scrapping or biopsy 	<p>Mucocutaneous candida infections are usually due to underlying clinical conditions such as poorly controlled diabetes or severe immunosuppression. Although not transmissible infections in the healthcare setting, they can be a marker for increased antibiotic exposure</p> <p>Dermatophytes have been known to cause occasional infections, and rare outbreaks, in the LTC setting.</p>



Skin, Soft Tissue and Mucosal Infections

Criteria	Comments
<p>D. <u>Herpes viral skin infections</u></p> <p><u>Herpes simplex infection</u></p> <p>Both criteria 1 and 2 present:</p> <ol style="list-style-type: none"> 1. A vesicular rash 2. Either physician diagnosis or laboratory confirmation <p><u>Herpes zoster infection</u></p> <p>Both criteria 1 and 2 present:</p> <ol style="list-style-type: none"> 1. A vesicular rash 2. Either physician diagnosis or laboratory confirmation 	<p>Reactivation of old herpes simplex (“cold sores”) or herpes zoster (“shingles”) is not considered a healthcare-associated infection</p> <p>Primary herpes viral skin infections are very uncommon in LTCF, except in pediatric populations where it should be considered healthcare-associated.</p>



Skin, Soft Tissue and Mucosal Infections

Criteria	Comments
<p>E. <u>Conjunctivitis</u></p> <p>At least one of the following criteria present:</p> <ol style="list-style-type: none">1. Pus appearing from one or both eyes, present for at least 24 hours2. New or increasing conjunctival erythema, with or without itching.3. New or increased conjunctival pain, present for at least 24 hours.	<p>Conjunctivitis symptoms (“pink eye”) should not be due to allergic reaction or trauma.</p>

Questions?



Gastrointestinal Tract Infections

Criteria	Comments
<p>A. <u>Gastroenteritis</u></p> <p>At least one of the following criteria present</p> <ol style="list-style-type: none"> 1. Diarrhea, three or more liquid or watery stools above what is normal for the resident within a 24-hour period 2. Vomiting, two or more episodes in a 24-hour period 3. Both of the following sign/symptom sub-criteria present: <ol style="list-style-type: none"> a) A stool specimen positive for a pathogen (such as Salmonella, Shigella, E. coli O157:H7, Campylobacter species, rotavirus) b) At least one of the following GI sub-criteria present <ol style="list-style-type: none"> i. Nausea ii. Vomiting iii. Abdominal pain iv. Diarrhea 	<p>Care must be taken to exclude non-infectious causes of symptoms. For instance, new medication may cause diarrhea, nausea/vomiting; initiation of new enteral feeding may be associated with diarrhea; nausea or vomiting may be associated with gallbladder disease.</p> <p>Presence of new GI symptoms in a single resident may prompt enhanced surveillance for additional cases.</p> <p>In the presence of an outbreak, stool from specimens should be sent to confirm the presence of norovirus, or other pathogens (such as rotavirus and E. coli O157:H7).</p>



Gastrointestinal Tract Infections

Criteria	Comments
<p>B. <u>Norovirus gastroenteritis</u></p> <p>Both criteria 1 and 2 present</p> <ol style="list-style-type: none"> 1. At least one of the following GI sub-criteria <ol style="list-style-type: none"> a) Diarrhea, three or more liquid or watery stools above what is normal for the resident within a 24-hour period b) Vomiting, two or more episodes in a 24-hour period 2. A stool specimen positive for detection of norovirus either by electron microscopy, enzyme immune assay, or by a molecular diagnostic test such as polymerase chain reaction (PCR). 	<p>In the absence of laboratory confirmation, an outbreak (2 or more cases occurring in a LTCF) of acute gastroenteritis due to norovirus infection in a LTCF may be assumed to be present if all of the following criteria are present ("Kaplan criteria")</p> <ol style="list-style-type: none"> a) Vomiting in more than half of affected persons b) A mean (or median) incubation period of 24-48 hours c) A mean (or median) duration of illness of 12-60 hours d) No bacterial pathogen is identified in stool culture.




Criteria	Comments
C. <u><i>Clostridium difficile</i></u> gastroenteritis	A "primary episode" of <i>C. difficile</i> infection (CDI) is defined as one that has occurred without any previous history of CDI., or that has occurred more than 8 weeks after the onset of a previous episode of CDI.
Both criteria 1 and 2 present	
1. One of the following GI sub-criteria	
a) Diarrhea, three or more liquid or watery stools above what is normal for the resident within a 24-hour period	A "recurrent episode" of CDI is defined as an episode of CDI that occurs 8 weeks or less after the onset of previous episode, provided the symptoms from the earlier (previous) episode resolved
b) The presence of toxic megacolon (abnormal dilation of the large bowel documented on radiology)	Individuals previously infected with <i>C. difficile</i> may continue to remain colonized even after symptoms resolve
2. One of the following diagnostic sub-criteria	
a) The stool sample yields a positive laboratory test result for <i>C. difficile</i> toxin A or B, or a toxin-producing <i>C. difficile</i> organism is identified in a stool culture or by a molecular diagnostic test such as PCR	In the setting of a GI outbreak, individuals could test positive for <i>C. difficile</i> toxin due to ongoing colonization and be co-infected with another pathogen. It is important that other surveillance criteria are used to differentiate infections in this situation.
b) Pseudomembranous colitis is identified during endoscopic examination or surgery, or in histopathologic examination of a biopsy specimen.	

Gastrointestinal Tract Infections



Why is *C. difficile* Surveillance Important?



CDI LabID Event

- *C. difficile* positive laboratory assay, tested on a loose-unformed stool specimen, and collected while a resident is receiving care from the LTCF, and the resident has no prior *C. difficile* positive laboratory assay collected in the previous two weeks (<14 days) while receiving care from the LTCF

SPICE 

Knowledge Check #1

Mr. Do Little has multiple co-morbidities including hypertension and acute respiratory failure. Vitals on admission WNL

On day seven after admission, the daughter tells the nurse “dad is not responding like he used to. He can not hold a conversation, tires easily and is not able to brush his teeth, eat or dress without assistance.”

SPICE 

Clinical Picture

Physical exam:

- Temp 100.7, pulse 107, RR 26 and O2 sat 93%
- Ronchi noted on auscultation of the chest the resident is confused

MD notified and orders urine and chest x-ray

Results:

- Culture + E. coli 10^2 cfu/ml and
- chest x-ray: no new findings

SPICE 

What surveillance criteria are met?

Resident has a cold

Resident has pneumonia

Resident has a lower respiratory tract infection

Resident is just “faking” to get daughter’s attention

Respiratory Tract Infections

Criteria	Comments
<p>D. <u>Lower respiratory tract (Bronchitis or Tracheo-bronchitis)</u></p> <p>All criteria 1-3 present</p> <ol style="list-style-type: none"> 1. Chest radiograph not performed <u>or negative for pneumonia or new infiltrate.</u> 2. At least two of the following respiratory sub-criteria (a-f) present <ol style="list-style-type: none"> a. New or increased cough b. New or increased sputum production c. O₂ saturation <94% on room air or a reduction in O₂ saturation of more than 3% from baseline d. New or changed lung exam abnormalities e. Pleuritic chest pain f. Respiratory rate of ≥ 25/min 3. At least one constitutional criteria 	<p>For both pneumonia and lower respiratory tract infections, presence of underlying conditions which could mimic a respiratory tract infection presentation (congestive heart failure, interstitial lung disease), should be excluded by review of clinical records and an assessment of presenting symptoms and signs</p>

SPICE

Knowledge Check # 2



1 Mar.

Mrs. Ross is a resident in your facility. An indwelling urinary catheter was inserted on March 1.



5 Mar.

On March 5, the nurse practitioner documented that Mrs. Ross complained of suprapubic pain.



6 Mar.

The following day, on March 6, a specimen collected from the Foley catheter was sent to the lab and subsequently tested positive for greater than 100,000 CFU/ml of E. coli and 100,000 CFU/ml of Candida auris.

SPICE

Does Mr. Ross have a CAUTI?

Yes

No

Start the presentation to see live content. Still no live content? Install the app or get help at PollEv.com/app

POLL

NHSN Urinary Tract Infections

For the resident with an indwelling catheter ✓

Criteria	Comments
<u>CA-SUTI</u>	
One or more of the following (Signs and Symptoms and Laboratory and diagnostic Testing):	
<ul style="list-style-type: none"> a) *Fever (Single temperature >100° F or >99° F on repeated occasions OR an increase of >2° F over baseline) b) Rigors c) New onset hypotension, with no alternate site of infection d) New onset confusion/functional decline AND Leukocytosis (>14,000 cells/mm³ or left shift (6% or 1, 500 bands/mm³)) e) New or marked increase in costovertebral angle pain or tenderness f) New or marked increase in suprapubic tenderness ✓ g) Acute pain, swelling, or tenderness of the testes, epididymis, or prostate h) Purulent discharge from around the catheter 	<p>AND</p> <p>↓</p> <p>A positive urine culture with no more than 2 species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/ml ✓</p> <p><i>*Fever can be used to meet CA-SUTI criteria even if the resident has another possible cause for the fever (for example, pneumonia)</i></p>

Knowledge Check # 3

Mr. U, a resident of LTC facility has a urinary catheter in place for 3 days for acute urinary retention. On day 3, he spikes a fever of 101°F and has a cough with shortness of breath.

The physician orders a urine culture, and it comes back positive with >100,000 CFU/ml of *Pseudomonas aeruginosa* and *Candida albicans*.

Upon further work, up Mr. U is determined not to have any other symptoms that meet the NHSN CA-SUTI criteria,

§ But, a chest X-ray does show infiltrates in the right upper lobe of the lung.

SPICE 

Does Mr. U Meet NHSN Definition?

YES, he meets NHSN criteria
for a CA-SUTI

NO, he does not meet NHSN
criteria for CA-SUTI because
the fever has another
alternative source
(respiratory infection)

NHSN Urinary Tract Infections

For the resident with an **indwelling catheter** ✓ > 2 days

Criteria	Comments
<u>CA-SUTI</u>	
One or more of the following (Signs and Symptoms and Laboratory and diagnostic Testing): ✓	
a) *Fever (Single temperature >100° F or >99° F on repeated occasions OR an increase of >2° F over baseline) ✓	AND ↓
b) Rigors	
c) New onset hypotension, with no alternate site of infection	A positive urine culture with no more than 2 species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/ml ✓
d) New onset confusion/functional decline AND Leukocytosis (>14,000 cells/mm ³ or left shift (6% or 1,500 bands/mm ³))	
e) New or marked increase in costovertebral angle pain or tenderness	<i>*Fever can be used to meet CA-SUTI criteria even if the resident has another possible cause for the fever (for example, pneumonia)</i>
f) New or marked increase in suprapubic tenderness	
g) Acute pain, swelling, or tenderness of the testes, epididymis, or prostate	
h) Purulent discharge from around the catheter	

Knowledge Check # 4

Day 1: Ms. R had an indwelling urinary catheter inserted in for a bladder outlet obstruction

Day 2: The indwelling urinary catheter remains in place

Day 3: The resident's indwelling urinary catheter remains in place. The resident had a single oral temp of 100.2°F. A urine culture was collected from the catheter

Ms. R
continued

Day 4: The indwelling urinary catheter remains in place. No symptoms documented

Day 5: The urine culture was positive for *Candida glabrata* 10^5 CFU/ml

SPICE 

Does Mr. R have a CA-SUTI?

Yes

No

Start the presentation to see live content. Still no live content? Install the app or get help at PollEv.com/app

SPICE 

What Standardized Definition is Met?

Meets NHSN and
McGeer definition

Meets NHSN definition
only

Meets McGeer definition
but not NHSN

Start the presentation to see live content. Still no live content? Install the app or get help at PollEv.com/app

SPICE

McGeer Urinary Tract Infections

Criteria	Comments
<p>B. <i>For the resident with an indwelling catheter</i> ✓</p> <p>Both criteria 1 and 2 present</p> <p>1. At least one of the following sign/symptom sub-criteria (a-d) present:</p> <p>a) Fever, rigors, or new onset hypotension, with no alternate site of infection ✓</p> <p>b) Either acute change in mental status <u>or</u> acute functional decline with no alternate diagnosis <u>and</u> Leukocytosis</p> <p>c) New onset suprapubic pain <u>or</u> costovertebral angle pain or tenderness</p> <p>d) Purulent discharge from around the catheter <u>or</u> acute pain, swelling, or tenderness of the testes, epididymis, or prostate</p> <p>2. Urinary catheter culture with $\geq 10^5$ cfu/ml of any organism(s) ✓</p>	<p>Recent catheter trauma, catheter obstruction or new onset hematuria are useful localizing signs consistent with UTI, but not necessary for diagnosis</p> <p>Urinary catheter specimens for culture should be collected following the replacement of the catheter (if current catheter has been in place for >14 days)</p>

SPICE

Knowledge Check # 5

- Mr. Bill was transferred to your facility from the local hospital on May 1.
- According to his admission record, he completed treatment for CDI prior to transfer.
- Two days after being transferred to your facility, the new NP ordered a C. diff test **“just to be on the safe side”**.
- On May 4, the stool specimen was positive for toxin A.



How Would You Classify This Event

C. difficile infection that is facility acquired

C. difficile infection that was acquired while in the hospital

Test not important because of history of C. difficile

This would meet the definition of CDI LabID event



Knowledge Check # 6

Mrs. Hammer is admitted to your facility for rehab after having hip replacement surgery at the local hospital. While in the hospital she received treatment for *C. difficile* infection

Two weeks later, resident complains that she has had multiple episodes of vomiting and diarrhea

Stool specimen is tested and is toxin negative for *C difficile* but PCR + *C. difficile*

SPICE 

Ms. Hammer

The nurse remembers that this is the 8th such case of diarrhea and vomiting and that the resident's roommate had similar symptoms 2 days ago.

When completing the line listing of infected cases the following data was noted:

6/8 residents had vomiting

5/8 residents had diarrhea

Most symptoms occurred within 48 hours of each other

Symptoms lasted on average of 36 hours (range 24-48 hrs)

SPICE 

What HAI Surveillance Criteria are Met?

Resident has a recurrent
C difficile infection **A**

Resident has a C.
difficile LabID event **B**

Resident has
gastroenteritis **C**

Resident has norovirus **D**

Start the presentation to see live content. For screen share software, share the entire screen. Get help at pollev.com/app

SPICE

Gastrointestinal Tract Infections

Criteria	Comments
<p>B. <i>Norovirus gastroenteritis</i></p> <p>Both criteria 1 and 2 present</p> <p>1. At least one of the following GI sub-criteria</p> <ul style="list-style-type: none"> a) Diarrhea, three or more liquid or watery stools above what is normal for the resident within a 24 hour period b) Vomiting, two or more episodes in a 24 hour period <p>2. A stool specimen positive for detection of norovirus either by electron microscopy, enzyme immune assay, or by a molecular diagnostic test such as polymerase chain reaction (PCR).</p>	<p>In the absence of laboratory confirmation, an outbreak (2 or more cases occurring in a LTCF) of acute gastroenteritis due to norovirus infection in a LTCF may be assumed to be present if all of the following criteria are present (“Kaplan criteria”)</p> <ul style="list-style-type: none"> a) Vomiting in more than half of affected persons b) A mean (or median) incubation period of 24-48 hours c) A mean (or median) duration of illness of 12-60 hours d) No bacterial pathogen is identified in stool culture.

SPICE

Gastrointestinal Tract Infections

Criteria	Comments
<p>A. <u>Gastroenteritis</u></p> <p>At least one of the following criteria present</p> <ol style="list-style-type: none">1. Diarrhea, three or more liquid or watery stools above what is normal for the resident within a 24 hour period2. Vomiting, two or more episodes in a 24 hour period3. Both of the following sign/symptom sub-criteria present:<ol style="list-style-type: none">a) A stool specimen positive for a pathogen (such as <i>Salmonella</i>, <i>Shigella</i>, <i>E. coli</i> 0157:H7, <i>Campylobacter</i> species, rotavirus)b) At least one of the following GI sub-criteria present<ol style="list-style-type: none">i. Nauseaii. Vomitingiii. Abdominal painiv. Diarrhea	<p>Care must be taken to exclude non-infectious causes of symptoms. For instance, new medication may cause diarrhea, nausea/vomiting; initiation of new enteral feeding may be associated with diarrhea; nausea or vomiting may be associated with gallbladder disease.</p> <p>Presence of new GI symptoms in a single resident may prompt enhanced surveillance for additional cases.</p> <p>In the presence of an outbreak, stool from specimens should be sent to confirm the presence of norovirus, or other pathogens (such as rotavirus and <i>E. coli</i> 0157:H7).</p>

Questions?

