

**The Multifactor Screener in the 2000
National Health Interview Survey Cancer
Control Supplement
(NHIS 2000)**

November 2019

THE MULTIFACTOR SCREENER: NHIS 2000

This document was originally published online on the Epidemiology and Genomic Research Program, National Cancer Institute's website as a reference for the Multifactor Screener in the 2000 National Health Interview Survey Cancer Control Supplement (NHIS 2000). The information contained in this document informs analyses of data from this screener. This information is archived and provided for reference purposes only.

This publication may be viewed and downloaded from
<https://epi.grants.cancer.gov/diet/screeners/files>.

Suggested citation for information contained in this report: The Multifactor Screener in the 2000 National Health Interview Survey Cancer Control Supplement. Epidemiology and Genomics Research Program. National Cancer Institute.
<https://epi.grants.cancer.gov/diet/screeners/files>. Updated November 20, 2019.

Acknowledgements

The Epidemiology and Genomics Research Program, Risk Factors Assessment branch, National Cancer Institute (EGRP, RFAB, NCI) acknowledge the work of Dr. Frances E. Thompson, Dr. Amy F. Subar, Douglas Midthune, Dr. Kevin Dodd, Dr. Victor Kipnis, and Lisa Kahle, whose work formed the basis of this document.

The EGRP, NCI would also like to acknowledge the important role of Dr. Edwina Wambogo and Jennifer Lerman who turned the webpages associated with this screener into this document.

If you have a comment or question about this document, please contact the EGRP, RFAB, NCI, at RFAB@mail.nih.gov.

THE MULTIFACTOR SCREENER: NHIS 2000

CONTENTS

| | |
|---|----|
| 1. Overview | 1 |
| 2. Definition of Acceptable Dietary Data Values | 2 |
| 3. Scoring Procedures | 3 |
| How Analytical Scoring Procedures Were Developed..... | 3 |
| Scoring Procedures | 4 |
| 4. Uses of Screener Estimates | 12 |
| Introduction | 12 |
| Variance-Adjustment Factor..... | 12 |
| What is the variance adjustment estimate and why do we need it? | 12 |
| How did we estimate the variance adjustment factors?..... | 13 |
| When do you use variance adjustment estimates? | 15 |
| Attenuation of Regression Parameters Using Screener Estimates | 15 |
| 5. Validation Results | 16 |
| 6. Computed Variables | 17 |
| 7. References | 18 |

THE MULTIFACTOR SCREENER: NHIS 2000

TABLES

Table 2- 1 Maximum daily average frequencies and acceptable values..... 2

Table 2- 2 Acceptable frequency reports for the weekly and monthly time periods..... 3

Table 3- 1 Converting frequency of intake into daily frequencies..... 4

Table 3- 2 Median Portion Size (Pk) in Grams per Mention by Age for Percentage Energy from Fat and Fiber Analyses: Men 5

Table 3- 3 Median Portion Size (Pk) in Grams per Mention by Age for Percentage Energy from Fat and Fiber Analyses: Women 7

Table 3- 4 Median Portion Size (Pk) in Pyramid Servings* per Mention by Gender and Age for Fruits and Vegetables Analyses 8

Table 3- 5 Estimated Regression Coefficients for Foods as Predictors of Percentage Energy from Fat and Grams of Fiber, by Gender..... 10

Table 3- 6 Estimated Regression Coefficients for Sum of Foods Predicting Pyramid Servings of Total Fruits and Vegetables and Fruits and Vegetables Excluding French Fries, by Gender 12

Table 4- 1 Variance Adjustment Factors for the NHIS Multifactor Screener, from the OPEN Study 14

Table 4- 2 Suggested procedures for various analytical objectives 15

Table 4- 3 Estimated attenuation factors for screener predicted intake in the OPEN study 16

THE MULTIFACTOR SCREENER: NHIS 2000

1. Overview

The Multifactor Screener may be useful to assess approximate intakes of fruits and vegetables, percentage energy from fat, and fiber. The screener asks respondents to report how frequently they consume foods in 16 categories. The screener also asks one question about the type of milk consumed. No portion size questions are asked. This screener does not attempt to assess total diet.

The foods selected to compose the screener were identified through an analysis of the U.S. Department of Agriculture's (USDA) [1994-96 Continuing Survey of Food Intakes of Individuals](#) (CSFII), a nationally representative survey of the food intakes of the U.S. population available from the USDA's Food Surveys Research Group. Staff in NCI's Risk Factor Assessment Branch (RFAB) used stepwise regression to identify the food groups that would best predict the three dietary exposures. Some of the foods in the screener predict all three exposures; some predict only one or two of the exposures.

The questions for the Multifactor Screener are provided below in two formats: self-report and in-person interviewer-administered.

- Self-report version of the Multifactor Screener, administered in the Observing Protein and Energy Nutrition (OPEN) Study.
- In-person interviewer-administered version of the Multifactor Screener on the NHIS CCM (questions NAC.010 - NAC.160).

You can view or print the Multifactor Screener in NHIS 2000 from the National Cancer Institute's (NCI) [Register of Validated Short Dietary Assessment Instruments](#).

In the 2000 National Health Interview Survey (NHIS), we applied rules for excluding extreme data responses, described in [Definition of Acceptable Dietary Data Values](#). The process of scoring the individual response data is described in [Scoring Procedures](#). A description and guidelines for the appropriate uses of the screener-estimated dietary intakes is found in [Uses of Screener Estimates](#). Validation data for the NHIS 2000 screener are presented in [Validation Results](#). Finally, the various dietary intake variables are found in [Computed Variables](#).

NOTE: The dietary variables on the NHIS 2000 dataset are in their natural units. For analyses, however, they must be transformed, first, to approximate normal distributions. For servings of fruits and vegetables, use the square root transformation; for fiber, use the cube-root transformation. No transformation is necessary for percentage energy from fat. After analyses, the result variables can be back-transformed for easier interpretation.

Different dietary screeners were used in the 2005, 2010, and 2015 NHIS, and these are also available in the [Register of Validated Short Dietary Assessment Instruments](#).

2. Definition of Acceptable Dietary Data Values

We used the U.S. Department of Agriculture's (USDA) [1994-96 Continuing Survey of Food Intakes of Individuals](#) (CSFII) data on reported intakes over two days of 24-hour recall to make judgments about reasonable frequencies of consumption that were reported on a per day basis. Maximum daily average frequencies (averaged for each individual across his two days of report) in the CSFII ranged from 1.5 times to 12 times per day for the relevant food groups. We accepted frequency values reported in the [National Health Interview Survey's Cancer Control Supplement](#) that were reported on a per day basis up to the maximum average values (rounded to the next whole number).

Table 2- 1 Maximum daily average frequencies and acceptable values

| Food Group | Maximum Daily Acceptable Value | Food Group | Maximum Daily Acceptable Value |
|-------------------|--------------------------------|----------------------|--------------------------------|
| Cold cereals | 10 | Fruit | 12 |
| Whole milk | 5 | Salad dressing | 3 |
| 2% milk | 6 | Salad | 5 |
| 1% milk | 6 | Fried potatoes | 3 |
| Skim milk | 5 | Other white potatoes | 3 |
| Other milk | 5 | Dried beans | 3 |
| Bacon or Sausage | 3 | Other vegetables | 9 |
| Hot dogs | 2 | Pasta | 3 |
| Whole grain bread | 5 | Nuts | 3 |
| 100% fruit juice | 4 | Chips | 3 |

In addition, we applied judgment to determine the acceptability of frequency reports for the weekly and monthly time period (see below). For example, a report of 25 times may be most logically associated with a month or year time period, but not so logically associated with a week time period. We applied this judgment to all foods.

THE MULTIFACTOR SCREENER: NHIS 2000

Table 2- 2 Acceptable frequency reports for the weekly and monthly time periods

| Time Period | Acceptable Frequency |
|--------------------|--|
| Week | 14 / week: acceptable > 14: assign a missing value |
| Month | 60 / month: acceptable > 60: assign a missing value |
| Year | any reported value is acceptable |

Depending on the intent of the analysis, the researcher could exclude a person with a missing value for any food, or only for foods of interest in the analysis. In our analyses of NHIS data, we excluded individuals only for the exposures in which they had missing data.

3. Scoring Procedures

How Analytical Scoring Procedures Were Developed

Scoring procedures were developed to convert a respondent's screener responses to estimates of individual dietary intake for percentage energy from fat, grams of fiber, and servings of fruits and vegetables, using the U.S. Department of Agriculture's (USDA) [1994-96 Continuing Survey of Food Intakes of Individuals \(CSFII 94-96\)](#) dietary recall data. The following equations were estimated in the CSFII 94-96:

For **percentage energy from fat and fiber**:

$$E(\text{Dietary Factor}) = b_0 + b_1 N_{FG1} P_1 + b_2 N_{FG2} P_2 + \dots + b_{20} N_{FG20} P_{20}$$

$E(\text{Dietary Factor})$ indicates the expected values for percentage energy from fat and for fiber, and assumes a normal distribution. In the CSFII 94-96 dataset percentage energy from fat was normally distributed. However, fiber was positively skewed and required a cube-root transformation to approximate normality. N_{FGk} is the usual number of times per day an individual consumed food group k ; P_k is the median portion size of group k ; and k indexes the 20 food groups. These 20 food groups were formed to reflect the same food groups on the screener. We calculated weighted least-squares estimates of the regression coefficients b_k , $k = 0, \dots, 20$ on CSFII 94-96 adults aged 18 and above, stratifying by gender and excluding extreme exposure values. We first included all 20 food groups in the regression model. After examining the results, we dropped food groups that failed to attain statistical significance at the $\alpha = 0.25$ level to form more parsimonious final models. In both the percentage energy from fat model and the fiber model, the lettuce food group was dropped. Because of the complex survey design, the analysis was performed using SUDAAN.

THE MULTIFACTOR SCREENER: NHIS 2000

For Pyramid servings of **fruits and vegetables** (defined by USDA in the 1992 Dietary Guidelines Food Guide Pyramid):

$$E(\text{Fruits and Veg}^{1/2}) = b_0 + b_1 (N_{FG1}P_1 + N_{FG2}P_2 + \dots + N_{FG7}P_7)^{1/2}$$

Pyramid servings of fruits and vegetables was square-root-transformed to approximate normality; N_{FGk} is the usual number of times per day an individual consumed food group k ; P_k is the median portion size of group k ; and k indexes the 7 fruit and vegetable food groups. We calculated weighted least-squares estimates of the regression coefficients b_0 and b_1 on the adults in the CSFII 94-96 sample, stratifying by gender and excluding extreme exposure values.

Scoring Procedures

After exclusion of extreme and missing values, we performed the following steps with the Cancer Control Supplement dietary data to estimate the individual's intake of servings of fruits and vegetables, percent energy from fat, and fiber.

1. **Estimation of N_{FGk} :** All reported frequencies were standardized to a common unit of time by converting them to daily frequencies.

Table 3- 1 Converting frequency of intake into daily frequencies.

| Time Period Reported | N_{FGk} : Daily Frequency |
|----------------------|-----------------------------------|
| Day | As reported |
| Week | Reported frequency divided by 7 |
| Month | Reported frequency divided by 30 |
| Year | Reported frequency divided by 365 |

2. **Estimation of P_k :** The median age- and gender-specific portion sizes for each food were estimated from CSFII 94-96. For percent energy from fat and fiber variables, the units were in grams ([Tables 3-2](#) and [Table 3-3](#)); for fruit and vegetable servings variables, the units were in Pyramid servings ([Table 3-4](#)).

A Pyramid serving is defined by the U.S. Department of Agriculture in the Dietary Guidelines Food Guide Pyramid (used from 1992-2004) as:

- vegetables: 1 cup raw leafy, 1/2 cup of other vegetables, or 3/4 cup vegetable juice; and
- fruit: 1 whole fruit, 1/2 cup of cut-up fruit, or 3/4 cup fruit juice.

Note: current dietary guidance uses cups rather than servings.

THE MULTIFACTOR SCREENER: NHIS 2000

Table 3- 2 Median Portion Size (Pk) in Grams per Mention by Age for Percentage Energy from Fat and Fiber Analyses: Men

| Food Group | Age Group | | | | | | |
|--|------------|------------|------------|------------|------------|------------|------------|
| | 18-27 | 28-37 | 38-47 | 48-57 | 58-67 | 68-77 | 78-99 |
| Men | | | | | | | |
| Cold cereals (P₁) | 74.666667 | 61.500000 | 57.500000 | 56.000000 | 46.000000 | 39.000000 | 33.000000 |
| Whole milk (P₂) | 305.000000 | 259.250000 | 306.710000 | 244.000000 | 244.000000 | 244.000000 | 203.333333 |
| 2% milk (P₃) | 259.250000 | 305.000000 | 244.000000 | 244.000000 | 244.000000 | 183.000000 | 183.000000 |
| 1% milk (P₄) | 341.600000 | 245.000000 | 245.000000 | 244.000000 | 213.500000 | 223.666667 | 183.000000 |
| Skim milk (P₅) | 366.666667 | 250.000000 | 250.000000 | 245.000000 | 214.375000 | 198.937500 | 160.725000 |
| Other milk (P₆) | 163.340000 | 163.340000 | 163.340000 | 163.340000 | 163.340000 | 163.340000 | 163.340000 |
| Bacon or sausage (P₇) | 25.000000 | 40.250000 | 32.000000 | 32.000000 | 27.000000 | 26.000000 | 24.000000 |
| Hotdogs (P₈) | 114.000000 | 85.500000 | 88.000000 | 114.000000 | 57.000000 | 57.000000 | 57.000000 |
| Whole grain bread (P₉) | 56.000000 | 54.000000 | 52.000000 | 52.000000 | 51.000000 | 48.250000 | 48.000000 |
| 100% fruit juice (P₁₀) | 372.000000 | 311.250000 | 249.000000 | 249.000000 | 248.000000 | 186.750000 | 186.750000 |
| Fruit (P₁₁) | 131.750000 | 128.000000 | 123.200000 | 127.500000 | 122.000000 | 118.000000 | 114.250000 |
| Salad dressing (P₁₂) | 23.543333 | 23.626667 | 22.030000 | 27.500000 | 24.585000 | 19.285000 | 15.600000 |
| Fried potatoes (P₁₃) | 112.500000 | 114.000000 | 100.000000 | 100.000000 | 85.500000 | 85.500000 | 97.000000 |

THE MULTIFACTOR SCREENER: NHIS 2000

| Food Group | Age Group | | | | | | |
|--|------------|------------|------------|------------|------------|------------|------------|
| | 18-27 | 28-37 | 38-47 | 48-57 | 58-67 | 68-77 | 78-99 |
| Other white potatoes (P₁₄) | 210.000000 | 196.000000 | 184.000000 | 161.000000 | 145.000000 | 127.000000 | 107.000000 |
| Dried beans (P₁₅) | 180.000000 | 130.000000 | 172.000000 | 172.000000 | 158.125000 | 175.000000 | 170.100000 |
| Other vegetables (P₁₆) | 60.013333 | 73.000000 | 74.063333 | 79.833333 | 76.500000 | 73.000000 | 67.520909 |
| Pasta (P₁₇) | 330.000000 | 280.000000 | 280.000000 | 247.500000 | 280.000000 | 210.000000 | 210.000000 |
| Nuts (P₁₈) | 31.625000 | 58.000000 | 35.500000 | 54.665000 | 39.250000 | 17.130000 | 35.916667 |
| Chips (P₁₉) | 40.000000 | 40.000000 | 31.895000 | 30.000000 | 26.000000 | 21.000000 | 17.500000 |

THE MULTIFACTOR SCREENER: NHIS 2000

Table 3- 3 Median Portion Size (Pk) in Grams per Mention by Age for Percentage Energy from Fat and Fiber Analyses: Women

| Food Group | Age Group | | | | | | |
|--|------------|------------|------------|------------|------------|------------|------------|
| | 18-27 | 28-37 | 38-47 | 48-57 | 58-67 | 68-77 | 78-99 |
| Women | | | | | | | |
| Cold cereals (P₁) | 50.000000 | 49.500000 | 44.000000 | 43.500000 | 33.000000 | 33.000000 | 33.500000 |
| Whole milk (P₂) | 244.000000 | 244.000000 | 244.000000 | 244.000000 | 198.250000 | 198.250000 | 196.400000 |
| 2% milk (P₃) | 244.000000 | 244.000000 | 244.000000 | 244.000000 | 183.000000 | 183.000000 | 152.500000 |
| 1% milk (P₄) | 244.000000 | 244.000000 | 183.000000 | 152.500000 | 183.000000 | 183.000000 | 218.583333 |
| Skim milk (P₅) | 245.000000 | 245.000000 | 244.800000 | 229.690000 | 196.000000 | 183.750000 | 183.750000 |
| Other milk (P₆) | 61.000000 | 61.000000 | 61.000000 | 61.000000 | 61.000000 | 61.000000 | 61.000000 |
| Bacon or sausage (P₇) | 26.000000 | 25.000000 | 24.000000 | 24.000000 | 18.000000 | 19.500000 | 16.000000 |
| Hotdogs (P₈) | 57.000000 | 57.000000 | 57.000000 | 114.000000 | 57.000000 | 57.000000 | 57.000000 |
| Whole grain bread (P₉) | 50.000000 | 48.000000 | 47.500000 | 45.000000 | 45.000000 | 42.400000 | 34.000000 |
| 100% fruit juice (P₁₀) | 280.125000 | 249.000000 | 248.800000 | 233.250000 | 189.755000 | 186.600000 | 186.700000 |
| Fruit (P₁₁) | 118.000000 | 118.000000 | 118.000000 | 118.000000 | 118.000000 | 112.427143 | 109.000000 |
| Salad dressing (P₁₂) | 17.140000 | 20.626667 | 23.020000 | 21.873333 | 22.035000 | 18.335000 | 10.210000 |
| Fried potatoes (P₁₃) | 79.500000 | 70.000000 | 70.000000 | 70.000000 | 66.000000 | 70.000000 | 64.000000 |

THE MULTIFACTOR SCREENER: NHIS 2000

| Food Group | Age Group | | | | | | |
|--|------------|------------|------------|------------|------------|------------|------------|
| | 18-27 | 28-37 | 38-47 | 48-57 | 58-67 | 68-77 | 78-99 |
| Other white potatoes (P₁₄) | 122.000000 | 127.000000 | 116.000000 | 122.000000 | 105.000000 | 105.000000 | 105.000000 |
| Dried beans (P₁₅) | 126.500000 | 89.000000 | 126.500000 | 126.500000 | 126.500000 | 126.500000 | 173.000000 |
| Other vegetables (P₁₆) | 53.750000 | 61.625000 | 61.500000 | 61.532500 | 63.165000 | 67.142857 | 71.333333 |
| Pasta (P₁₇) | 217.500000 | 217.500000 | 182.525000 | 185.000000 | 165.000000 | 160.000000 | 175.000000 |
| Nuts (P₁₈) | 18.000000 | 32.000000 | 20.655000 | 21.265000 | 18.250000 | 11.250000 | 25.500000 |
| Chips (P₁₉) | 28.000000 | 24.333333 | 27.000000 | 26.000000 | 20.000000 | 18.000000 | 14.000000 |

Table 3- 4 Median Portion Size (Pk) in Pyramid Servings* per Mention by Gender and Age for Fruits and Vegetables Analyses

| Food Group | Age Group | | | | | | |
|---|-----------|----------|----------|----------|----------|----------|----------|
| | 18-27 | 28-37 | 38-47 | 48-57 | 58-67 | 68-77 | 78-99 |
| Men | | | | | | | |
| 100% fruit juice (P₁) | 2.000000 | 1.667500 | 1.335000 | 1.335000 | 1.334000 | 1.001000 | 1.001000 |
| Fruit (P₂) | 1.301000 | 1.301000 | 1.229571 | 1.227333 | 1.168000 | 1.168000 | 1.052333 |
| Salad (P₃) | 0.545000 | 0.708000 | 0.754500 | 0.750000 | 0.833500 | 0.750000 | 0.822500 |
| Fried potatoes (P₄) | 2.000000 | 2.000000 | 1.773000 | 1.710000 | 1.400000 | 1.250000 | 1.250000 |

THE MULTIFACTOR SCREENER: NHIS 2000

| Food Group | Age Group | | | | | | |
|---|------------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | 18-27 | 28-37 | 38-47 | 48-57 | 58-67 | 68-77 | 78-99 |
| Other white potatoes (P₅) | 2.000000 | 2.000000 | 1.999000 | 1.999000 | 1.914000 | 1.544000 | 1.508000 |
| Dried beans (P₆) | 1.374000 | 1.047000 | 1.065000 | 1.227000 | 1.000000 | 1.000000 | 1.114000 |
| Other vegetables (P₇) | 0.750000 | 0.906000 | 0.974500 | 1.000000 | 1.000000 | 0.880000 | 0.833333 |
| Women | | | | | | | |
| 100% fruit juice (P₁) | 1.500500 | 1.334000 | 1.334000 | 1.251250 | 1.019500 | 1.000500 | 1.000500 |
| Fruit (P₂) | 1.168000 | 1.168000 | 1.168000 | 1.168000 | 1.150500 | 1.083833 | 1.000000 |
| Salad (P₃) | 0.613500 | 0.572500 | 0.833333 | 1.000000 | 0.795500 | 0.625000 | 0.750000 |
| Fried potatoes (P₄) | 1.481000 | 1.365500 | 1.272000 | 1.400000 | 1.000000 | 1.026000 | 1.000000 |
| Other white potatoes (P₅) | 1.544000 | 1.544000 | 1.528000 | 1.544000 | 1.499000 | 1.516000 | 1.272000 |
| Dried beans (P₆) | 0.964000 | 0.684000 | 0.800000 | 0.687000 | 0.822000 | 0.807000 | 1.000000 |
| Other vegetables (P₇) | 0.702200 | 0.779333 | 0.792500 | 0.788500 | 0.774000 | 0.833000 | 0.856750 |

* Defined by [Dietary Guidelines Food Guide Pyramid \(1992-2004\)](#).

THE MULTIFACTOR SCREENER: NHIS 2000

3. For >percentage energy from fat and fiber (grams) estimation of b_k , $k = 0, \dots, 19$: the values for each parameter, for each gender, are in the following table:

Table 3- 5 Estimated Regression Coefficients for Foods as Predictors of Percentage Energy from Fat and Grams of Fiber, by Gender

| Parameter | Men | | Women | |
|---|----------------------------|-------------------|----------------------------|-------------------|
| | Percentage Energy from Fat | Fiber (cube root) | Percentage Energy from Fat | Fiber (cube root) |
| Intercept (b_0) | 31.952688 | 2.083624 | 31.371597 | 1.898219 |
| Cold cereals (b_1) | -0.025863 | 0.002045 | -0.057206 | 0.003843 |
| Whole milk (b_2) | 0.006512 | 0.000129 | 0.008364 | 0.000095 |
| 2% milk (b_3) | 0.002045 | 0.000137 | 0.002634 | 0.000115 |
| 1% milk (b_4) | -0.001500 | 0.000225 | -0.002082 | 0.000245 |
| Skim milk (b_5) | -0.008536 | 0.000285 | -0.008775 | 0.000344 |
| Other milk (b_6) | -0.026527 | 0.001474 | -0.047247 | 0.003496 |
| Bacon or sausage (b_7) | 0.138079 | -0.001377 | 0.231718 | -0.002043 |
| Hotdogs (b_8) | 0.041135 | 0 | 0.101657 | -0.001416 |
| Whole grain bread (b_9) | 0 | 0.002827 | 0 | 0.003373 |
| 100% fruit juice (b_{10}) | -0.005354 | 0.000187 | -0.010099 | 0.000246 |
| Fruit (b_{11}) | -0.009290 | 0.001031 | -0.011894 | 0.001046 |
| Salad dressing (b_{12}) | 0.149646 | 0 | 0.238928 | 0 |
| Fried potatoes (b_{13}) | 0.027202 | 0.001603 | 0.042656 | 0.001563 |

THE MULTIFACTOR SCREENER: NHIS 2000

| Parameter | Men | | Women | |
|--|----------------------------|-------------------|----------------------------|-------------------|
| | Percentage Energy from Fat | Fiber (cube root) | Percentage Energy from Fat | Fiber (cube root) |
| Other white potatoes (b₁₄) | 0.005688 | 0.000716 | 0.006148 | 0.000658 |
| Dried beans (b₁₅) | -0.005231 | 0.002748 | -0.005934 | 0.003787 |
| Other vegetables (b₁₆) | 0 | 0.000839 | 0 | 0.000932 |
| Pasta (b₁₇) | -0.005029 | 0.000753 | -0.005416 | 0.000826 |
| Nuts (b₁₈) | 0.125658 | 0.005401 | 0.260003 | 0.006053 |
| Chips (b₁₉) | 0.053905 | 0.005276 | 0.130953 | 0.004592 |

THE MULTIFACTOR SCREENER: NHIS 2000

4. For Pyramid servings of **fruits and vegetables**, *estimation of b_0 and b_1* :

$$\text{The model is: } E(\text{Dietary Factor}^{1/2}) = b_0 + b_1 (N_{FG1}P_1 + N_{FG2}P_2 + \dots + N_{FG7}P_7)^{1/2}$$

For Pyramid servings of fruits and vegetables, including and excluding French fries, for each gender, the estimates of the parameters are:

Table 3- 6 Estimated Regression Coefficients for Sum of Foods Predicting Pyramid Servings of Total Fruits and Vegetables and Fruits and Vegetables Excluding French Fries, by Gender

| Parameter | Men | Women |
|--|----------|----------|
| Summary variable with French fries | | |
| Intercept (b_0) | 0.906793 | 0.819559 |
| b_1 | 0.758560 | 0.730865 |
| Summary variable excluding French fries | | |
| Intercept (b_0) | 0.940772 | 0.816265 |
| b_1 | 0.739056 | 0.730219 |

4. Uses of Screener Estimates

Introduction

Dietary intake estimates derived from the Multifactor Screener are rough estimates of usual intake of fruits and vegetables, fiber, calcium, servings of dairy, and added sugar. These estimates are not as accurate as those from more detailed methods (e.g., 24-hour recalls). However, [Validation Results](#) suggests that the estimates may be useful to characterize a population's median intakes, to discriminate among individuals or populations with regard to higher vs. lower intakes, to track dietary changes in individuals or populations over time, and to allow examination of interrelationships between diet and other variables. In addition, diet estimates from the Cancer Control Supplement (CCS) could be used as benchmark national data for smaller surveys, for example, in a particular state.

Variance-Adjustment Factor

What is the variance adjustment estimate and why do we need it?

Data from the Multifactor Screener are individuals' reports about their intake and, like all self-reports, contain some error. The algorithms we use to estimate servings of fruits and vegetables, grams of fiber, mg of calcium, servings of dairy, and teaspoons of added sugar calibrate the data to 24-hour recalls. The screener estimate of intake represents what we

THE MULTIFACTOR SCREENER: NHIS 2000

expect the person would have reported on his 24-hour recall, given what he reported on the individual items in the screener. As a result, the mean of the screener estimate of intake should equal the mean of the 24-hour recall estimate of intake in the population. (It would also equal the mean of true intake in the population if the 24-hour recalls were unbiased. However, there are many studies suggesting that recalls underestimate individuals' true intakes).

When describing a population's distribution of dietary intakes, the parameters needed are an estimate of central tendency (i.e. mean or median) and an estimate of spread (variance). The variance of the screener, however, is expected to be smaller than the variance of true intake, since the screener prediction formula estimates the conditional expectation of true intake given the screener responses, and in general the variance of a conditional expectation of a variable is smaller than the variance of itself. As a result, the screener estimates of intake cannot be used to estimate quantiles (other than median) or prevalence estimates of true intake without an adjustment. Procedures have been developed to estimate the variance of true intake using data from 24-hour recalls, by taking into consideration within person variability [1, 2].

We extended these procedures to allow estimation of the variance of true intake using data from the screener. The resulting variance adjustment factor adjusts the screener variance to approximate the variance of true intake in the population.

How did we estimate the variance adjustment factors?

We have estimated the adjustment factors in the various external validation datasets available to us. The results indicate that the adjustment factors differ by gender and dietary variable. Under the assumption that the variance adjustment factors appropriate to National Health Interview Study (NHIS) are similar to those in Observing Protein and Energy Nutrition Study (OPEN) [3], the variance-adjusted screener estimate of intake should have variance closer to the estimated variance of true intake that would have been obtained from repeat 24-hour recalls. For Pyramid servings of fruits and vegetables, the variance adjustment factors in OPEN and Eating at America's Table Study (EATS) [4] are quite similar, which gives us some indication that these factors might be relatively stable from population to population.

THE MULTIFACTOR SCREENER: NHIS 2000

Table 4- 1 Variance Adjustment Factors for the NHIS Multifactor Screener, from the OPEN Study

| Nutrient | Gender | Variance Adjustment Factor |
|--|--------|----------------------------|
| Total Fruit & Vegetable Intake (Pyramid Servings) | Male | 1.3 |
| | Female | 1.1 |
| Fruit & Vegetable Intake (excluding fried potatoes) (Pyramid Servings) | Male | 1.3 |
| | Female | 1.2 |
| Percentage Calories from Fat | Male | 1.5 |
| | Female | 1.3 |
| Fiber Intake (grams) | Male | 1.2 |
| | Female | 1.2 |

How do you use the variance adjustment estimates?

To estimate quantile values or prevalence estimates for an exposure, you should first adjust the screener so that it has approximately the same variance as true intake.

Adjust the screener estimate of intake by:

- multiplying intake by an adjustment factor (an estimate of the ratio of the standard deviation of true intake to the standard deviation of screener intake); and
- adding a constant so that the overall mean is unchanged.

The formula for the variance-adjusted screener is:

$$\text{variance-adjusted screener} = (\text{variance adjustment factor}) * (\text{unadjusted screener} - \text{mean}_{\text{unadj. scr.}}) + \text{mean}_{\text{unadj. scr.}}$$

This procedure is performed on the normally distributed version of the variable (i.e., Pyramid servings of fruits and vegetables is square-rooted, percentage energy from fat is untransformed, and fiber is cube rooted). For fruits and vegetables and fiber, the results can then be squared or cubed, respectively, to obtain estimates in the original units.

The variance adjustment procedure is used to estimate prevalence of obtaining recommended intakes for the 2000 NHIS in Thompson et al., [5].

THE MULTIFACTOR SCREENER: NHIS 2000

When do you use variance adjustment estimates?

The appropriate use of the screener information depends on the analytical objective. Following is a characterization of suggested procedures for various analytical objectives.

Table 4- 2 Suggested procedures for various analytical objectives

| Analytical Objective | Procedure |
|--|---|
| Estimate mean or median intake in the population or within subpopulations. | Use the unadjusted screener estimate of intake. |
| Estimate quantiles (other than median) of the distribution of intake in the population; estimate prevalence of attaining certain levels of dietary intake. | Use the variance-adjusted screener estimate. |
| Classify individuals into exposure categories (e.g., meeting recommended intake vs. not meeting recommended intake) for later use in a regression model. | Use the variance-adjusted screener estimates to determine appropriate classification into categories. |
| Use the screener estimate as a continuous covariate in a multivariate regression model. | Use the unadjusted screener estimate. |
| Use the screener estimate as a response (dependent) variable. | Use the variance-adjusted screener estimate. |

Attenuation of Regression Parameters Using Screener Estimates

When the screener estimate of dietary intake is used as a continuous covariate in a multivariate regression, the estimated regression coefficient will typically be attenuated (biased toward zero) due to measurement error in the screener. The "attenuation factor" [6] can be estimated in a calibration study and used to deattenuate the estimated regression coefficient (by dividing the estimated regression coefficient by the attenuation factor).

We estimated attenuation factors in the OPEN study (see below). If you use these factors to deattenuate estimated regression coefficients, note that the data come from a relatively small study that consists of a fairly homogeneous population (primarily white, well-educated individuals).

THE MULTIFACTOR SCREENER: NHIS 2000

Table 4- 3 Estimated attenuation factors for screener predicted intake in the OPEN study

| Gender | Square-Root Fruit & Veg (Pyramid Servings) | Square-Root Fruit & Veg (excluding French Fries) (Pyramid Servings) | Percentage Energy From Fat | Cube-Root Fiber (grams) |
|---------------|---|--|-----------------------------------|--------------------------------|
| Men | 0.75 | 0.79 | 0.96 | 0.70 |
| Women | 0.81 | 0.87 | 0.88 | 0.69 |

If you categorize the screener values into quantiles and use the resulting categorical variable in a linear or logistic regression, the bias (due to misclassification) is more complicated because the categorization can lead to differential misclassification in the screener [7]. Although methods may be available to correct for this [8, 9], it is not simple, nor are we comfortable suggesting how to do it at this time.

Even though the estimated regression coefficients are biased (due to measurement error in the screener or misclassification in the categorized screener), tests of whether the regression coefficient is different from zero are still valid. For example, if one used the SUDAAN REGRESS procedure with fruit and vegetable intake (estimated by the screener) as a covariate in the model, one could use the Wald F statistic provided by SUDAAN to test whether the regression coefficients were statistically significantly different from zero. This assumes that there is only one covariate in the model measured with error; when there are multiple covariates measured with error, the Wald F test that a single regression coefficient is zero may not be valid, although the test that the regression coefficients for all covariates measured with error are zero is still valid.

5. Validation Results

The NCI'S Risk Factor Assessment Branch (RFAB) staff have assessed the validity of the Multifactor Screener in several studies: NCI's Observing Protein and Energy (OPEN) Study [3], the Eating at America's Table Study (EATS) [4], and the joint [NIH-AARP Diet and Health Study](#). In all studies, multiple 24-hour recalls in conjunction with a measurement error model were used to assess validity. In general, the validation results reflect the Multifactor Screener's hierarchical design -- fruit and vegetable intake was estimated best by the screener, followed by percentage energy from fat, and lastly grams of fiber.

THE MULTIFACTOR SCREENER: NHIS 2000

In the OPEN Study, estimates of median intake were:

- Pyramid Servings of Fruits and Vegetables:
 - Men: recalls - 6.3; screener - 5.3;
 - Women: recalls - 5.4; screener - 4.7.
- Percentage Energy from Fat:
 - Men: recalls - 31.8; screener - 32.0;
 - Women: recalls - 32.0; screener - 30.5.
- Fiber:
 - Men: recalls - 21.2; screener - 18.3;
 - Women: recalls - 16.5; screener - 14.1.

These validation results suggest that dietary exposure estimates computed for the Cancer Control Supplement may be useful to compare subgroup means, especially for populations consuming mainstream diets. The estimates may be less useful for populations with more ethnic diets, including Asian and possibly Latino populations.

At the individual level, correlations between the screener and estimated true intake ranged from 0.54 (fiber for men) to 0.76 (Pyramid servings of fruits and vegetables for women); about 25 to 50 percent of the variability in the true intake will be captured by the screener questions. Thus, although significant error may be associated with these estimates of diet, we believe the exposure estimates still substantially reflect what individuals are actually consuming.

Validation results are reported in detail in Thompson et al., [10].

6. Computed Variables

The computed diet variables for the 2000 National Health Interview Survey Multifactor Screener are available for download in two formats -- SAS transport and comma-separated values (CSV). The files include the following variables:

- Pyramid servings of fruits and vegetables
- Adjusted Pyramid servings of fruits and vegetables
- Pyramid servings of fruits and vegetables excluding French fries
- Adjusted Pyramid servings of fruits and vegetables excluding French fries
- Fiber (gm)
- Adjusted fiber (gm)

THE MULTIFACTOR SCREENER: NHIS 2000

- Percent energy from fat
- Adjusted percent energy from fat
- Household serial number (HHX)
- Family serial number (FMX)
- Person number (PX)

These datasets, data dictionary, and SAS program for this screener are available for download on [NCI's Short Dietary Assessment Instruments](#) website.

The datasets are sorted in ascending order by the ID variables HHX, FMX, and PX. All numeric variables have been rounded to the nearest 0.000001.

- Comma-separated Values File (dietvars.2004_03_30.csv.zip) - This zip file contains the comma-separated values file, which includes 11 variables, 32,374 records, and an additional record for the variable names.
- SAS Transport File (dietvars.2004_03_30.v8x.zip) - The SAS transport file includes 32,374 records and 11 variables. To access the SAS dataset, unzip the file, then use proc cimport.

For example:

```
proc cimport file='dietvars.2004_03_30.v8x' data=dietvars'
```

Other Documents

The following documents may also be of use:

- Content tables of the SAS dataset: contents.dietvars.2004_03_30.pdf
- SAS program that created the dataset: create.dietvars.2004_03_30.sas

7. References

1. National Research Council. *Nutrient Adequacy: Assessment Using Food Consumption Surveys*. Washington, DC: *National Academy Press*, 1986. Available for download from: <https://www.nap.edu/catalog/618/nutrient-adequacy-assessment-using-food-consumption-surveys>.
2. Institute of Medicine. *Dietary Reference Intakes: Applications in Dietary Assessment*. Washington, DC: *National Academy Press*, 2000. Available for download from:

THE MULTIFACTOR SCREENER: NHIS 2000

- <https://www.nap.edu/catalog/9956/dietary-reference-intakes-applications-in-dietary-assessment>.
3. Subar AF, et al. [Using intake biomarkers to evaluate the extent of dietary misreporting in a large sample of adults: the OPEN study](#). *Am J Epidemiol*, 2003. 158(1):1-13.
 4. Subar AF, et al. [Comparative validation of the Block, Willett, and National Cancer Institute food frequency questionnaires: the Eating at America's Table Study](#). *Am J Epidemiol*, 2001. 154(12): 1089-99.
 5. Thompson FE, et al. [Dietary intake estimates in the National Health Interview Survey, 2000: methodology, results, and interpretation](#). *J Am Diet Assoc*, 2005. 105(3): 352-63.
 6. Rosner B, Willett WC, Spiegelman D. [Correction of logistic regression relative risk estimates and confidence intervals for systematic within-person measurement error](#). *Stat Med*, 1989. 8(9): 1051-69; discussion 1071-3.
 7. Flegal KM, Keyl PM, Nieto FJ. [Differential misclassification arising from nondifferential errors in exposure measurement](#). *Am J Epidemiol*, 1991. 134(10): 1233-44.
 8. Flegal KM, Brownie C, Haas JD. [The effects of exposure misclassification on estimates of relative risk](#). *Am J Epidemiol*, 1986. 123(4): 736-51.
 9. Morrissey MJ, Spiegelman D. [Matrix methods for estimating odds ratios with misclassified exposure data: extensions and comparisons](#). *Biometrics*, 1999. 55(2): 338-44.
 10. Thompson FE, et al. [Performance of a short tool to assess dietary intakes of fruits and vegetables, percentage energy from fat and fibre](#). *Public Health Nutr*, 2004. 7(8): 1097-105.