

# FELINE NUTRITIONAL MANAGEMENT DECISION TREE



**URINARY S/O**

**URINARY S/O**  
MODERATE CALORIE

URINARY + SATIETY

URINARY + CALM

URINARY + HYPOALLERGENIC



**URINARY S/O**

**URINARY S/O**  
SMALL DOGS **S**

**URINARY U/C**

**URINARY S/O**  
MODERATE CALORIE

**URINARY S/O** AGEING 7+

URINARY + SATIETY

URINARY + HYPOALLERGENIC

This resource has been produced with the greatest care and incorporates the most up-to-date scientific information and research findings. We recommend that you refer to our packaging as the primary reference for product information as this can change with time. Bearing in mind the diversity and complexity of clinical cases in dogs and cats, it is important to note that the suggestions in this resource for further tests and therapeutic options cannot be exhaustive. Under no circumstances should the treatments and dietary solutions suggested here replace examination by a veterinary surgeon.

The publishers and the authors will accept no responsibility for the failure of any of the treatments or nutritional solutions put forward in this resource. Not for use in Canada and in USA.



# URINARY S/O



Available in  
1.5, 3.5,  
and 7kg bags

## RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
2	39	4/8	33	3/8	26	2/8	
2.5	46	4/8	38	4/8	31	3/8	
3	52	5/8	44	4/8	35	3/8	
3.5	59	5/8	49	5/8	39	4/8	
4	64	6/8	54	5/8	43	4/8	
4.5	70	6/8	58	5/8	47	4/8	
5	75	7/8	63	6/8	50	5/8	
5.5	81	1	67	7/8	54	5/8	
6	86	1	72	7/8	57	5/8	
6.5	91	1	76	7/8	61	6/8	
7	96	1 + 1/8	80	7/8	64	6/8	
7.5	101	1 + 1/8	84	1	67	6/8	
8	105	1 + 2/8	88	1	70	6/8	
8.5	110	1 + 2/8	92	1 + 1/8	73	7/8	
9	115	1 + 3/8	96	1 + 1/8	76	7/8	
9.5	119	1 + 3/8	99	1 + 1/8	79	7/8	
10	123	1 + 3/8	103	1 + 2/8	82	1	

Urinary S/O is a complete dietetic feed for cats, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties and its low level of magnesium.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The cat should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** - the wet version of Urinary S/O Feline is recommended for recurrent idiopathic cystitis, in older cats renal function should be checked before recommending a new diet.

**[S/O] INDEX** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



### LOW MAGNESIUM

Reduced level of magnesium, a natural component of struvite crystals.

KEY VALUES	Per 1000 kcal	Per 100g as fed
Protein	89.1g	34.5g
Fat content	38.7g	15g
Carbohydrate	85.7g	33.2g
Crude fibre	7.5g	2.9g
Dietary fibre	17.6g	6.8g
Calcium	2.3g	0.9g
Phosphorus	2.3g	0.9g
Potassium	2.6g	1g
Sodium	3.4g	1.30g
Chloride	5.8g	2.26g
Magnesium	0.1g	0.05g
Copper	3.9mg	1.5mg
Zinc	43.9mg	17mg
Taurine	0.6g	0.23g
Vitamin A	6198.3IU	2400IU
Vitamin E	131.7mg	51mg
Omega 6	8.7g	3.38g
Omega 3	2.3g	0.88g
EPA + DHA	1g	0.37g

Metabolisable energy\* 387kcal/100g as fed  
\*Calculated according to NRC2006.

### COMPOSITION

Rice, wheat gluten\*\*, dehydrated poultry protein, maize flour, animal fats, hydrolysed animal proteins, maize gluten, minerals, vegetable fibres, fish oil, soya oil, fructo-oligo-saccharides, marigold extract (source of lutein)

\*\*L.I.P.: protein selected for its very high digestibility

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 21500 IU, Vitamin D3: 800IU, E1 (Iron): 37 mg, E2 (Iodine): 3.7mg, E4 (Copper): 11mg, E5 (Manganese): 48mg, E6 (Zinc): 136mg, E8 (Selenium): 0.06mg, Urine acidifying substances: calcium sulphate [1.25%], Preservatives - Antioxidants.

\*\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



# URINARY S/O



MORSELS IN GRAVY

Available in box  
of 12 pouches  
of 85g

## RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch	grams	pouch
2	170	2	140	1 + 1/2	110	1 + 1/2	
2.5	195	2 + 1/2	165	2	130	1 + 1/2	
3	225	2 + 1/2	185	2	150	2	
3.5	250	3	210	2 + 1/2	165	2	
4	275	3	230	2 + 1/2	185	2	
4.5	300	3 + 1/2	250	3	200	2 + 1/2	
5	320	4	270	3	215	2 + 1/2	
5.5	345	4	285	3 + 1/2	230	2 + 1/2	
6	365	4 + 1/2	305	3 + 1/2	245	3	
6.5	390	4 + 1/2	325	4	260	3	
7	410	5	340	4	275	3	
7.5	430	5	360	4	285	3 + 1/2	
8	450	5 + 1/2	375	4 + 1/2	300	3 + 1/2	
8.5	470	5 + 1/2	390	4 + 1/2	315	3 + 1/2	
9	490	6	410	5	325	4	
9.5	510	6	425	5	340	4	
10	530	6	440	5	350	4	

Urinary S/O is a complete dietetic feed for cats, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The cat should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** - in older cats renal function should be checked before recommending a new diet.

**[S/O] INDEX** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Feline idiopathic cystitis (FIC)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### IDIOPATHIC CYSTITIS

High moisture content increases urine dilution



### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.

KEY VALUES	Per 1000 kcal	Per 100g as fed
Protein	119.76g	10g
Fat content	41.92g	3.5g
Carbohydrate	44.31g	3.7g
Crude fibre	11.98g	1g
Dietary fibre	14.37g	1.2g
Calcium	2.63g	0.22g
Phosphorus	2.28g	0.19g
Potassium	2.16g	0.18g
Sodium	3.59g	0.3g
Chloride	5.03g	0.42g
Magnesium	0.14g	0.01g
Copper	4.55mg	0.38mg
Zinc	25.15mg	2.10mg
Taurine	1.2g	0.1g
Vitamin A	35928.14IU	3000IU
Vitamin E	147.31mg	12.30mg
Omega 6	8.38g	0.7g
Omega 3	1.8g	0.15g
EPA + DHA	1.1g	0.09g

Metabolisable energy\* 84kcal/100g as fed  
\*Calculated according to NRC2006.

### COMPOSITION

Poultry by-products, pork by-products, pork blood products, wheat flour, vital wheat gluten, powder cellulose, starch mixture, minerals, fish oil, sunflower oil refined, caramelised sugars, marigold extract (source of lutein).

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 225IU, E1 (Iron): 3mg, E2 (Iodine): 0.25mg, E4 (Copper): 2mg, E5 (Manganese): 1mg, E6 (Zinc): 10mg, Urine acidifying substances: calcium sulfate: 0.25%. Technological additives: acidity regulator) sodium bisulfate: 0.5g.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## URINARY S/O



LOAF

Available in box  
of 12 pouches  
of 85g

## RECOMMENDED DAILY INTAKE &amp; FEEDING DURATION

	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch	grams	pouch
ADULT CAT	2	170	2	140	1 + 1/2	110	1 + 1/2
	2.5	195	2 + 1/2	165	2	130	1 + 1/2
	3	225	2 + 1/2	185	2	150	2
	3.5	250	3	210	2 + 1/2	165	2
	4	275	3	230	2 + 1/2	185	2
	4.5	300	3 + 1/2	250	3	200	2 + 1/2
	5	320	4	270	3	215	2 + 1/2
	5.5	345	4	285	3 + 1/2	230	2 + 1/2
	6	365	4 + 1/2	305	3 + 1/2	245	3
	6.5	390	4 + 1/2	325	4	260	3
	7	410	5	340	4	275	3
7.5	430	5	360	4	285	3 + 1/2	
8	450	5 + 1/2	375	4 + 1/2	300	3 + 1/2	
8.5	470	5 + 1/2	390	4 + 1/2	315	3 + 1/2	
9	490	6	410	5	325	4	
9.5	510	6	425	5	340	4	
10	530	6	440	5	350	4	

Urinary S/O is a complete dietetic feed for cats, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The cat should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** - in older cats renal function should be checked before recommending a new diet.

**S/O® INDEX** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

## RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Feline idiopathic cystitis (FIC)

## NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



## IDIOPATHIC CYSTITIS

High moisture content increases urine dilution



## STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



## LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



## URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.

KEY VALUES	Per 1000 kcal	Per 100g as fed
Protein	130.18g	11g
Fat content	41.42g	3.5g
Carbohydrate	33.14g	2.8g
Crude fibre	11.83g	1g
Dietary fibre	17.75g	1.5g
Calcium	2.96g	0.25g
Phosphorus	2.96g	0.25g
Potassium	2.6g	0.22g
Sodium	3.55g	0.3g
Chloride	4.73g	0.4g
Magnesium	0.19g	0.02g
Copper	4.62mg	0.39g
Zinc	30.77mg	2.6mg
Taurine	1.18g	0.1g
Vitamin A	26035.5IU	2200IU
Vitamin E	118.34mg	10mg
Omega 6	8.88g	0.75g
Omega 3	1.89g	0.16g
EPA + DHA	1.07g	0.09g
Metabolisable energy*	85kcal/100g as fed	

\*Calculated according to NRC2006.

## COMPOSITION

Poultry by-products, pork by-products, powder cellulose, minerals, pork blood products, vital wheat gluten, starch mixture, fish oil, caramelised sugars, marigold extract (source of lutein).

## ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 225IU, E1 (Iron): 3mg, E2 (Iodine): 0.25mg, E4 (Copper): 2mg, E5 (Manganese): 1mg, E6 (Zinc): 10mg, Urine acidifying substances: calcium sulfate: 0.25%. Technological additives: (acidity regulator) sodium bisulfate: 0.5g.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## URINARY S/O

## MODERATE CALORIE



Available in  
1.5, 3.5  
and 7kg bags

## RECOMMENDED DAILY INTAKE &amp; FEEDING DURATION

	CAT'S WEIGHT (kg)	BODY CONDITION			
		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups
ADULT CAT	2	36	3/8	29	3/8
	2.5	42	4/8	34	3/8
	3	48	5/8	39	4/8
	3.5	54	5/8	43	4/8
	4	59	6/8	47	4/8
	4.5	64	6/8	51	5/8
	5	69	7/8	55	5/8
	5.5	74	7/8	59	6/8
	6	79	1	63	6/8
	6.5	84	1	67	6/8
	7	88	1	70	7/8
7.5	92	1 + 1/8	74	7/8	
8	97	1 + 1/8	77	7/8	
8.5	101	1 + 2/8	81	1	
9	105	1 + 2/8	84	1	
9.5	109	1 + 2/8	87	1	
10	113	1 + 3/8	91	1 + 1/8	

Urinary S/O Moderate Calorie is a complete dietetic feed for cats, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties and its low level of magnesium.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Moderate Calorie for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The cat should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** the wet version of Urinary S/O Moderate Calorie is recommended for recurrent idiopathic cystitis, in older cats renal function should be checked before recommending a new diet.

**S/O® INDEX** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

## RECOMMENDED FOR CATS WITH TENDENCY TO BE OVERWEIGHT IN CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence

## NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



## STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



## MODERATE CALORIE

Moderate calorie content to help maintain ideal weight.



## URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



## LOW RSS

Helps lower the concentration of ions contributing to crystal formation.

KEY VALUES	Per 1000 kcal	Per 100g as fed
Protein	96.73g	34g
Fat content	31.29g	11g
Carbohydrate	95.87g	33.7g
Crude fibre	19.91g	7g
Dietary fibre	36.98g	13g
Calcium	2.84g	1g
Phosphorus	2.56g	0.9g
Potassium	2.84g	1g
Sodium	3.41g	1.2g
Chloride	6.09g	2.14g
Magnesium	0.17g	0.06g
Copper	4.27mg	1.5mg
Zinc	48.65mg	17.1mg
Taurine	0.63g	0.22g
Vitamin A	6543.39IU	2300IU
Vitamin E	119.49mg	42mg
Omega 6	7.2g	2.56g
Omega 3	1.99g	0.7g
EPA + DHA	0.91g	0.32g
Metabolisable energy*	352kcal/100g as fed	

\*Calculated according to NRC2006.

## COMPOSITION

Rice, dehydrated poultry protein, wheat gluten\*\*, maize flour, vegetable fibres, maize gluten, hydrolysed animal proteins, minerals, animal fats, fish oil, soya oil, fructo-oligo-saccharides, marigold extract (source of lutein).

\*\*L.I.P.: protein selected for its very high digestibility

## ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 21500IU, Vitamin D3: 800IU, E1 (Iron): 35mg, E2 (Iodine): 3.5mg, E4 (Copper): 11mg, E5 (Manganese): 46mg, E6 (Zinc): 134mg, E8 (Selenium): 0.06mg, urine acidifying substances: calcium sulphate (1.3%), Preservatives - Antioxidants.

\*\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



# URINARY S/O

## MODERATE CALORIE



MORSELS IN GRAVY

Available in box of 12 pouches of 85g

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION			
		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups
	2	160	2	130	1 + 1/2
	2.5	185	2	150	2
	3	215	2 + 1/2	170	2
	3.5	240	3	190	2
	4	260	3	210	2 + 1/2
	4.5	285	3 + 1/2	225	2 + 1/2
	5	305	3 + 1/2	245	3
	5.5	330	4	260	3
	6	350	4	280	3 + 1/2
	6.5	370	4 + 1/2	295	3 + 1/2
	7	390	4 + 1/2	310	3 + 1/2
	7.5	410	5	325	4
	8	430	5	340	4
	8.5	445	5	355	4
	9	465	5 + 1/2	370	4 + 1/2
	9.5	485	5 + 1/2	385	4 + 1/2
	10	500	6	400	4 + 1/2

Urinary S/O Moderate Calorie is a complete dietetic feed for cats, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties and its low level of magnesium.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Moderate Calorie for 5 to 12 weeks for the dissolution of struvite stones and up to 6 struvite stone recurrence. The cat should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** - in older cats renal function should be checked before recommending a new diet.

**[S/O® INDEX]** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Feline idiopathic cystitis (FIC)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



### MODERATE CALORIE

Moderate calorie content to help maintain ideal weight.

KEY VALUES	Per 1000 kcal	Per 100g as fed
Protein	129.96g	9.5g
Fat content	36.94g	2.7g
Carbohydrate	45.14g	3.3g
Crude fibre	17.78g	1.3g
Dietary fibre	20.52g	1.5g
Calcium	3.01g	0.22g
Phosphorus	2.6g	0.19g
Potassium	2.19g	0.16g
Sodium	3.56g	0.26g
Chloride	4.51g	0.33g
Magnesium	0.14g	0.01g
Copper	4.92mg	0.36mg
Zinc	24.62mg	1.80mg
Taurine	1.16g	0.09g
Vitamin A	68399.45IU	5000IU
Vitamin E	136mg	10mg
Omega 6	5.06g	0.37g
Omega 3	2.33g	0.17g
EPA + DHA	1.09g	0.08g

Metabolisable energy\* 73kcal/100g as fed

\*Calculated according to NRC2006.

### COMPOSITION

Pork by-products, poultry by-products, salmon, pork blood products, vital wheat gluten, powdercellulose, wheat flour, starch mixture, minerals, fish oil, caramelised sugars, marigold extract [source of lutein].

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 170IU, E1 [Iron]: 2mg, E2 [Iodine]: 0.25mg, E4 [Copper]: 2mg, E5 [Manganese]: 0.5mg, E6 [Zinc]: 5mg, Urine acidifying substances: calcium sulfate: 0.25%.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## URINARY + SATIETY



Available in 1.5 and 3.5kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	TARGET BODY WEIGHT OF CAT	START		AFTER 4 WEEKS WITH NO WEIGHT LOSS	
		grams	cups	grams	cups
		2	28	3/8	23
2.5	33	3/8	26	3/8	
3	38	4/8	30	3/8	
3.5	42	4/8	34	3/8	
4	47	5/8	37	4/8	
4.5	51	5/8	40	4/8	
5	55	5/8	43	4/8	
5.5	58	6/8	46	5/8	
6	62	6/8	49	5/8	
6.5	66	6/8	52	5/8	
7	69	7/8	55	5/8	
7.5	73	7/8	58	6/8	
8	76	7/8	60	6/8	
8.5	80	1	63	6/8	
9	83	1	66	6/8	
9.5	86	1	68	7/8	
10	89	1 1/8	71	7/8	

For nutritional management of cats with struvite urolithiasis and excessive bodyweight: feed the diet for 5-12 weeks and re-evaluate based on clinical assessment of the response. To avoid recurrence of struvite uroliths, the diet should be fed for a further 6 months minimum followed by urinalysis. For detailed guidance on weight reduction please see product information for Satiety Feline. S/O-Kcal Control Feline can be maintained for the cat's lifetime. It is recommended to divide the daily intake into 2 meals.

**RECOMMENDATIONS:** Target Weight should be determined according to Body Condition Score of the individual. The optimal weight loss rate is 0.5-2% bodyweight weekly. Maintenance Ration: It is recommended to continue feeding the weight-loss diet once target weight is achieved. Energy intakes should be increased by 10% and Body Condition Score and weight monitored closely throughout the stabilisation period. Should the cat regain more than 5% of body weight, the daily ration should be decreased to weight loss levels. It is recommended to divide the daily intake into 2 meals.

**[S/O® INDEX]** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

**Feline lower urinary tract disease:** • Dissolution of struvite uroliths • Management of recurrent struvite and calcium oxalate urolithiasis. In older cats, renal function should be checked before S/O-Kcal Control Feline is recommended.

**+ Weight Management:** Obesity • Diabetes mellitus associated with obesity

### NOT RECOMMENDED IN CASE OF

Pregnancy, lactation, growth • Chronic diseases requiring an increased energy intake • Chronic Kidney Disease • Heart disease (where sodium restriction is required) • Concurrent use of urine-acidifying drugs • Conditions where high fibre content is not recommended



### SATIETOGENIC EFFECT

Contains a special blend of fibres which have a satiating effect on cats.



### HIGH PROTEIN

A high protein content (111 g/1000 kcal ME) helps maintain muscle mass throughout the weight loss period.



### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



### LOW RSS

Helps lower the concentration of ions contributing to struvite and calcium oxalate crystal formation.

KEY VALUES	Per 100g as fed
Protein	34g
Fat content	9g
Carbohydrate	18.4g
NFE	27.9g
Dietary fibre	23.5g
Crude fibre	14g
Omega 6	1.89mg
Omega 3	0.37mg
Epa+DHA+DHA	0.14g
Calcium	1.26g
Phosphorus	1.15g
Sodium	1.1g
L-Carnitine	20mg
Chondroitin sulphate + Glucosamine	50mg

Metabolisable energy\* 304.6 Kcal

\*Calculated according to NRC2006.

### COMPOSITION

Dehydrated poultry protein, vegetable fibres, tapioca, wheat gluten\*\*, maize gluten, wheat flour, hydrolysed animal proteins, minerals, animal fats, chicory pulp, fish oil, psyllium husks and seeds, hydrolysed crustaceans (source of glucosamine), hydrolysed cartilage (source of chondroitin).

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 23800IU, Vitamin D3: 900IU, E1 [Iron]: 29mg, E2 [Iodine]: 2.9mg, E4 [Copper]: 9mg, E5 [Manganese]: 37mg, E6 [Zinc]: 112mg, E8 [Selenium]: 0.04mg Preservatives - Antioxidants.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





## URINARY + CALM

Available in  
2 and 4kg bags

## RECOMMENDED DAILY INTAKE &amp; FEEDING DURATION

ADULT CAT	TARGET BODY WEIGHT OF CAT		NORMAL		OVERWEIGHT	
	grams	cups	grams	cups	grams	cups
2	36	3/8				
3	48	4/8				
4	58	5/8	47	4/8		
5	68	5/8	55	4/8		
6	78	6/8	62	5/8		
7	87	7/8	69	5/8		
8	95	7/8	76	6/8		
9	104	1	83	7/8		
10	112	1 1/8	89	7/8		

For nutritional management of cats with struvite urolithiasis: feed the diet for 5-12 weeks and re-evaluate based on clinical assessment of the response. To avoid recurrence of struvite uroliths, the diet should be fed for a further 6 months minimum followed by urinalysis. It is recommended to divide the daily intake into 2 meals.

**[S/O] INDEX** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

## RECOMMENDED FOR CASES OF

**Feline lower urinary tract disease** • Management of risk factors: tendency to be overweight • Insufficient water intake and exposure to increased environmental challenges • Dissolution of struvite uroliths • Management of recurrent struvite and calcium oxalate urolithiasis. **In older cats**, renal function should be checked before S/O-Biopeptide Feline is recommended.  
**+ Weight Management:** Obesity • Diabetes mellitus associated with obesity • Hyperlipidaemia associated with obesity

## NOT RECOMMENDED IN CASE OF

Pregnancy, lactation, growth • Chronic Kidney Disease  
 • Heart disease (where sodium restriction is required)  
 • Concurrent use of urine-acidifying drugs



## URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



## CALMING SUPPORT

Contains hydrolysed casein (milk protein) and L-tryptophan to help manage exposure to increased environmental challenges.



## STRUVITE DISSOLUTION

Helps dissolve struvite stones.



## MODERATE CALORIE

Moderate calorie content to help maintain ideal weight. Overweight is a risk factor for FLUTD.

KEY VALUES	Per 100g as fed
Protein	34g
Fat content	11g
Carbohydrate	27.4g
NFE	34.8g
Dietary fibre	13g
Crude fibre	5.6g
Omega 6	2.25mg
Omega 3	1.18mg
EPA+DHA	0.67g
Calcium	1.08g
Phosphorus	0.98g
Sodium	1.15g
L-Tryptophan	0.4mg
Hydrolysed milk protein	94mg

Metabolisable energy\* 356.5Kcal

\*Calculated according to NRC2006.

## COMPOSITION

Dehydrated poultry protein, rice, wheat gluten\*, maize, wheat, vegetable fibres, hydrolysed animal proteins, maize gluten, minerals, chicory pulp, fish oil, animal fats, fructo-oligo-saccharides, psyllium husks and seeds, soya oil, hydrolysed milk protein\*\*, marigold extract [source of lutein].

## ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 28100IU, Vitamin D3: 800IU, E1 (Iron): 34mg, E2 (Iodine): 3.4mg, E4 (Copper): 10mg, E5 (Manganese): 44mg, E6 (Zinc): 131mg, E8 (Selenium): 0.07mg  
 Preservatives - Antioxidants.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## URINARY + HYPOALLERGENIC

Available in  
2kg bags

## RECOMMENDED DAILY INTAKE &amp; FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
	grams	cups	grams	cups	grams	cups	
2	40	3/8	34	3/8			
3	54	4/8	45	4/8			
4	66	5/8	55	4/8	44	4/8	
5	78	6/8	65	5/8	52	4/8	
6	88	7/8	74	6/8	59	5/8	
7	99	1	82	7/8	66	5/8	
8	108	1 + 1/8	90	7/8	72	6/8	
9	118	1 + 2/8	98	1	79	6/8	
10	127	1 + 2/8	106	1 + 1/8	85	7/8	

For nutritional management of cats with struvite urolithiasis and Adverse Food Reaction: feed the diet for 5-12 weeks and re-evaluate based on clinical assessment of the response. To avoid recurrence of struvite uroliths, the diet should be fed for a further 6 months minimum followed by urinalysis. S/O-AFR Feline can be maintained for the cat's lifetime. It is recommended to divide the daily intake into 2 meals.

**[S/O] INDEX** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

## RECOMMENDED FOR CASES OF

**Feline lower urinary tract disease:** • Dissolution of struvite uroliths • Management of recurrent struvite and calcium oxalate urolithiasis. **In older cats**, renal function should be checked before S/O-AFR Feline is recommended.

**+ Dermatological/GI conditions:** Food elimination trial  
 • Food allergies • Food intolerance • Inflammatory Bowel Disease (IBD) • Chronic diarrhoea • Exocrine Pancreatic Insufficiency (EPI) • Small Intestinal Bacterial overgrowth (SIBO)

## NOT RECOMMENDED IN CASE OF

Pregnancy, lactation, growth • Chronic Kidney Disease  
 • Heart disease (where sodium restriction is required)  
 • Concurrent use of urine-acidifying drugs



## URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



## LOW RSS

Helps lower the concentration of ions contributing to struvite and calcium oxalate crystal formation.



## HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to help ensure very low allergenic potential.



## SKIN BARRIER

A complex to support the barrier effect of the skin.

KEY VALUES	Per 100g as fed
Protein	27g
Fat content	16g
Carbohydrate	32.8g
NFE	37.3g
Dietary fibre	9.8g
Crude fibre	5.3g
Omega 6	4.39mg
Omega 3	0.39mg
EPA+DHA	0.32g
Calcium	0.71g
Phosphorus	0.7g
Sodium	1.3g
Potassium	0.8mg
Vitamin D3	70IU

Metabolisable energy\* 376.6Kcal

\*Calculated according to NRC2006.

## COMPOSITION

Rice, hydrolysed soya protein isolate, animal fats, minerals, chicory pulp, vegetable fibre, soya oil, fish oil, fructo-oligo-saccharides, fatty acid salt, borage oil.


## ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 28500IU, Vitamin D3: 800IU, E1 (Iron): 43mg, E2 (Iodine): 3.9mg, E4 (Copper): 15mg, E5 (Manganese): 57mg, E6 (Zinc): 205mg, E8 (Selenium): 0.25mg  
 Technological additives: Clinoptilolite of sedimentary origin: 10g. Preservatives - Antioxidants.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



# CANINE NUTRITIONAL MANAGEMENT DECISION TREE



## URINARY S/O



Available in 2, 7.5 and 13kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
2	54	5/8	48	4/8	41	4/8	
4	91	1	80	7/8	69	6/8	
6	123	1 + 3/8	109	1 + 2/8	94	1 + 1/8	
8	153	1 + 6/8	135	1 + 5/8	116	1 + 3/8	
10	181	2 + 1/8	159	1 + 7/8	138	1 + 5/8	
15	245	2 + 7/8	216	2 + 4/8	187	2 + 1/8	
20	305	3 + 4/8	268	3 + 1/8	232	2 + 5/8	
25	360	4 + 1/8	317	3 + 5/8	274	3 + 1/8	
30	413	4 + 6/8	363	4 + 2/8	314	3 + 5/8	
35	463	5 + 3/8	408	4 + 6/8	352	4 + 1/8	
40	512	5 + 7/8	451	5 + 2/8	389	4 + 4/8	
45	560	6 + 4/8	492	5 + 6/8	425	4 + 7/8	
50	606	7	533	6 + 1/8	460	5 + 3/8	
55	650	7 + 4/8	572	6 + 5/8	494	5 + 6/8	
60	694	8	611	7 + 1/8	528	6 + 1/8	
70	779	9	686	8	592	6 + 7/8	
80	862	10	758	8 + 6/8	655	7 + 5/8	

Urinary S/O is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The dog should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** - renal function should be checked before recommending a new diet. - in older dogs, the use of Urinary S/O Ageing\*\* is recommended.

**[S/O] INDEX** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

- RECOMMENDED FOR CASES OF**
- Struvite uroliths: dissolution and management of recurrence
  - Calcium oxalate uroliths: management of recurrence
- NOT RECOMMENDED IN CASE OF**
- Chronic Kidney Disease (CKD)
  - Heart disease (when sodium restriction is sought)
  - Concurrent use of urine-acidifying medication
  - Growth, gestation/lactation

- STRUVITE DISSOLUTION**  
Helps dissolve all types of struvite stones.
- URINE DILUTION**  
Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.
- LOW RSS**  
Helps lower the concentration of ions contributing to crystal formation.
- LOW MAGNESIUM**  
Reduced level of magnesium, a natural component of struvite crystals.

KEY VALUES	Per 1000 kcal	Per 100g as fed
Protein	46.38g	18g
Fat content	43.8g	17g
Carbohydrate	120.85g	46.9g
Crude fibre	5.67g	2.2g
Dietary fibre	14.69g	5.7g
Calcium	1.29g	0.5g
Phosphorus	1.29g	0.5g
Potassium	2.06g	0.8g
Sodium	2.83g	1.1g
Chloride	5.51g	2.14g
Magnesium	0.1g	0.04g
Copper	3.86mg	1.5mg
Zinc	43.8mg	17mg
Taurine	0.54g	0.21g
Vitamin A	4380.31IU	1700IU
Vitamin E	157.18mg	61mg
Omega 6	9.25 g	3.59g
Omega 3	1.76g	0.63g
EPA + DHA	0.54g	0.21g

Metabolisable energy\* 388kcal/100g as fed  
\*Calculated according to NRC2006.

**COMPOSITION**  
Maize flour, rice, animal fats, dehydrated poultry protein, maize gluten, minerals, hydrolysed animal proteins, vegetable fibres, soya oil, fish oil, fructo-oligo-saccharides, mono - and diglycerides of palmitic and stearic acids esterified with citric acid, marigold extract (source of lutein). Protein sources: dehydrated poultry protein, hydrolysed animal proteins.

**ADDITIVES (KG)\*\***  
Nutritional additives: Vitamin A: 15500IU, Vitamin D3: 1000IU, E1 (Iron): 44mg, E2 (Iodine): 4.4mg, E4 (Copper): 13mg, E5 (Manganese): 57mg, E6 (Zinc): 147mg, E8 (Selenium): 0.06mg. Urine acidifying substances: calcium sulfate (0.88%). Preservatives - Antioxidants.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





## URINARY S/O



LOAF

Available in  
410g can

## RECOMMENDED DAILY INTAKE &amp; FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION						
		THIN		NORMAL		OVERWEIGHT		
		grams	can	grams	can	grams	can	
2	208	1/2	183	1/2	158	1/2	158	1/2
4	350	3/4	308	3/4	266	3/4	266	3/4
6	475	1 + 1/4	418	1	361	1	361	1
8	589	1 + 1/2	519	1 + 1/4	448	1	448	1
10	697	1 + 3/4	613	1 + 1/2	529	1 + 1/4	529	1 + 1/4
15	944	2 + 1/4	831	2	718	1 + 3/4	718	1 + 3/4
20	1172	2 + 3/4	1031	2 + 1/2	890	2 + 1/4	890	2 + 1/4
25	1385	3 + 1/2	1219	3	1053	2 + 1/2	1053	2 + 1/2
30	1588	3 + 3/4	1397	3 + 1/2	1207	3	1207	3
35	1783	4 + 1/4	1569	3 + 3/4	1355	3 + 1/4	1355	3 + 1/4
40	1970	4 + 3/4	1734	4 + 1/4	1498	3 + 3/4	1498	3 + 3/4
45	2158	5 + 1/4	1894	4 + 1/2	1636	4	1636	4
50	2329	5 + 3/4	2050	5	1770	4 + 1/4	1770	4 + 1/4
55	2502	6	2202	5 + 1/4	1902	4 + 3/4	1902	4 + 3/4
60	2671	3 + 1/2	2350	5 + 3/4	2030	5	2030	5
70	2998	7 + 1/4	26.38	6 + 1/2	2279	5 + 1/2	2279	5 + 1/2
80	3314	8	2916	7	2519	6 + 3/4	2519	6 + 3/4

Urinary S/O is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The dog should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** - renal function should be checked before recommending a new diet. - in older dogs, the use of Urinary S/O Ageing<sup>™</sup> is recommended.

**[S/O®]  
INDEX]** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

## RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

## NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



## STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



## URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



## LOW RSS

Helps lower the concentration of ions contributing to crystal formation.

KEY VALUES	Per 1000 kcal	Per 100g as fed
Protein	83.68g	8g
Fat content	62.76g	6g
Carbohydrate	48.12g	4.6g
Crude fibre	14.64g	1.4g
Dietary fibre	25.1g	2.4g
Calcium	2.2g	0.21g
Phosphorus	2.09g	0.2g
Potassium	2.09g	0.2g
Sodium	3.03g	0.29g
Chloride	4.18g	0.4g
Magnesium	0.17g	0.02g
Copper	3.14mg	0.3mg
Zinc	23.01mg	2.2mg
Taurine	1.26g	0.12g
Vitamin A	47071.13IU	4500IU
Vitamin E	198.74mg	19mg
Omega 6	25.1 g	2.4g
Omega 3	1.36g	0.13g
EPA + DHA	0.63g	0.06g

Metabolisable energy\* 96kcal/100g as fed

\*Calculated according to NRC2006.

## COMPOSITION

Pork by-products, poultry by-products, chicken meat, sunflower oil refined, powdercellulose, maize flour, starch mixture, minerals, pork blood products, caramelised sugars, fish oil, marigold extract (source of lutein).

## ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 190IU, E1 (Iron): 3.4mg, E2 (Iodine): 0.24 mg, E4 (Copper): 2mg, E5 (Manganese): 1mg, E6 (Zinc): 10mg.  
Urine acidifying substances: calcium sulfate: 0.25%.  
Technological additives: (acidity regulator) Sodium bisulfate: 0.56g.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## URINARY S/O



THIN SLICES IN GRAVY

Available in box  
of 12 pouches  
of 100g

## RECOMMENDED DAILY INTAKE &amp; FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch	grams	pouch
2	207	2	182	2	157	1 + 1/2	157
4	348	3 + 1/2	306	3	264	2 + 1/2	264
6	471	4 + 1/2	415	4	358	3 + 1/2	358
8	585	6	515	5	444	4 + 1/2	444
10	691	7	608	6	525	5 + 1/2	525
15	973	9 + 1/2	824	8	712	7	712
20	1162	11 + 1/2	1023	10	883	9	883
25	1374	13 + 1/2	1209	12	1044	10 + 1/2	1044
30	1576	16	1386	14	1197	12	1197
35	1769	17 + 1/2	1556	15 + 1/2	1344	13 + 1/2	1344
40	1955	19 + 1/2	1720	17	1486	15	1486

Urinary S/O is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The dog should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** - renal function should be checked before recommending a new diet. - in older dogs, the use of Urinary S/O Ageing<sup>™</sup> is recommended.

**[S/O®]  
INDEX]** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

## RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

## NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



## STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



## URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



## LOW RSS

Helps lower the concentration of ions contributing to crystal formation.

KEY VALUES	Per 1000 kcal	Per 100g as fed
Protein	86.11g	8g
Fat content	59.2g	5.5g
Carbohydrate	51.67g	4.8g
Crude fibre	12.92g	1.2g
Dietary fibre	15.07g	1.4g
Calcium	1.61g	0.15g
Phosphorus	1.51g	0.14g
Potassium	1.61g	0.15g
Sodium	2.91g	0.27g
Chloride	3.98g	0.37g
Magnesium	0.13g	0.01g
Copper	3.77mg	0.35mg
Zinc	23.14mg	2.15mg
Taurine	1.24g	0.12g
Vitamin A	46286.33IU	4300IU
Vitamin E	226.05mg	21mg
Omega 6	19.38 g	1.8g
Omega 3	1.35g	0.13g
EPA + DHA	0.65g	0.06g

Metabolisable energy\* 93kcal/100g as fed

\*Calculated according to NRC2006.

## COMPOSITION

Pork by-products, chicken meat, poultry by-products, wheat flour, sunflower oil refined, pork blood products, vital wheat gluten, powdercellulose, starch mixture, minerals, fish oil, caramelised sugars, marigold extract (source of lutein).

## ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 210IU, E1 (Iron): 3.5mg, E2 (Iodine): 0.25mg, E4 (Copper): 2mg, E5 (Manganese): 1.1mg, E6 (Zinc): 11mg.  
Urine acidifying substances: calcium sulfate: 0.25%.  
Technological additives: (acidity regulator) sodium bisulfate: 1.107g.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.









# URINARY S/O

## MODERATE CALORIE



Available in  
6.5 and  
12kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION			
		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups
2	49	5/8	41	4/8	
4	82	1	69	7/8	
6	111	1 + 3/8	93	1 + 1/8	
8	137	1 + 6/8	116	1 + 4/8	
10	163	2	137	1 + 6/8	
15	220	2 + 6/8	186	2 + 3/8	
20	273	3 + 4/8	230	2 + 7/8	
25	323	4 + 1/8	272	3 + 3/8	
30	370	4 + 5/8	312	4	
35	416	5 + 2/8	350	4 + 3/8	
40	460	5 + 6/8	387	4 + 7/8	
45	502	6 + 3/8	423	5 + 3/8	
50	543	6 + 7/8	458	5 + 6/8	
55	584	7 + 3/8	492	6 + 2/8	
60	623	7 + 7/8	525	6 + 5/8	
65	662	8 + 3/8	557	7	
70	699	8 + 7/8	589	7 + 3/8	

Urinary S/O Moderate Calorie is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Moderate Calorie for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The dog should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** - in older dogs renal function should be checked before recommending a new diet. It is recommended to divide the daily intake into 2 meals.

**S/O® INDEX** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease [when sodium restriction is sought]
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



### MODERATE CALORIE

Moderate calorie content to help maintain ideal weight.



### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.

KEY VALUES	Per 1000 kcal	Per 100g as fed
Protein	60.85g	20g
Fat content	33.47g	11g
Carbohydrate	139.34g	45.8g
Crude fibre	19.77g	6.5g
Dietary fibre	37.12g	12.2g
Calcium	1.73g	0.57g
Phosphorus	1.73g	0.57g
Potassium	2.43g	0.8g
Sodium	3.65g	1.2g
Chloride	6.54g	2.15g
Magnesium	0.18g	0.06g
Copper	4.56mg	1.5mg
Zinc	51.72mg	17mg
Taurine	0.52g	0.17g
Vitamin A	6997.26IU	2300IU
Vitamin E	136.9mg	45mg
Omega 6	7.18 g	2.36g
Omega 3	1.52g	0.5g
EPA + DHA	0.61g	0.2g

Metabolisable energy\* 329kcal/100g as fed

\*Calculated according to NRC2006.

### COMPOSITION

Maize flour, rice, dehydrated poultry protein, vegetable fibres, maize gluten, animal fats, minerals, hydrolysed animal proteins, chicory pulp, wheat gluten\*\*, soya oil, fish oil, fructo-oligo-saccharides, mono - and diglycerides of palmitic and stearic acids esterified with citric acid, marigold extract (source of lutein). Protein sources: dehydrated poultry protein, maize gluten, hydrolysed animal proteins, wheat gluten.

\*\*L.I.P.: protein selected for its very high digestibility

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 21500IU, Vitamin D3: 1400IU, E1 (Iron): 44mg, E2 (Iodine): 4.4mg, E4 (Copper): 14mg, E5 (Manganese): 58mg, E6 (Zinc): 146mg, E8 (Selenium): 0.11mg, Urine acidifying substances: calcium sulphate (1%). Preservatives - Antioxidants.

\*\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



# URINARY S/O

## MODERATE CALORIE



THIN SLICES IN GRAVY

Available in box  
of 12 pouches  
of 100g

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION			
		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch
2	181	2	152	1 + 1/2	
4	304	3	256	2 + 1/2	
6	412	4	347	3 + 1/2	
8	512	5	431	4 + 1/2	
10	605	6	509	5	
15	820	8	691	7	
20	1018	10	857	8 + 1/2	
25	1203	12	1013	10	
30	1379	14	1161	11 + 1/2	
35	1548	15 + 1/2	1304	13	
40	1711	17	1441	14 + 1/2	

Urinary S/O Moderate Calorie is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Moderate Calorie for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The dog should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** - in older dogs renal function should be checked before recommending a new diet. It is recommended to divide the daily intake into 2 meals.

**S/O® INDEX** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease [when sodium restriction is sought]
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



### MODERATE CALORIE

Moderate calorie content to help maintain ideal weight.



### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.

KEY VALUES	Per 1000 kcal	Per 100g as fed
Protein	97.32g	8g
Fat content	43.8g	3.6g
Carbohydrate	71.78g	5.9g
Crude fibre	12.17g	1g
Dietary fibre	17.64g	1.5g
Calcium	1.7g	0.14g
Phosphorus	1.58g	0.13g
Potassium	1.82g	0.15g
Sodium	2.92g	0.24g
Chloride	4.26g	0.35g
Magnesium	0.12g	0.01
Copper	3.65mg	0.30mg
Zinc	21.29mg	1.75mg
Taurine	1.22g	0.1g
Vitamin A	48661.8IU	4000IU
Vitamin E	182.48mg	15mg
Omega 6	973g	0.80g
Omega 3	1.34g	0.11g
EPA + DHA	0.61g	0.05g

Metabolisable energy\* 82kcal/100g as fed

\*Calculated according to NRC2006.

### COMPOSITION

Pork by-products, chicken meat, wheat flour, poultry by-products, pork blood products, vital wheat gluten, starch mixture, powdercellulose, minerals, sunflower oil refined, fish oil, caramelised sugars, marigold extract (source of lutein).

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 180IU, E1 (Iron): 2mg, E2 (Iodine): 0.2mg, E4 (Copper): 1.8mg, E5 (Manganese): 0.8mg, E6 (Zinc): 8mg, Sulphur: 0.2%, Urine acidifying substances: calcium sulfate: 0.3%. Technological additives: [acidity regulator] sodium bisulfate: 1.1g.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





# URINARY S/O AGEING 7+



Available in 1.5, 3.5 and 8kg bags

## RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
2	55	5/8	48	4/8	42	4/8	
4	92	1	81	7/8	70	6/8	
6	125	1 + 3/8	110	1 + 2/8	95	1	
8	155	1 + 6/8	136	1 + 4/8	117	1 + 2/8	
10	183	2	161	1 + 6/8	139	1 + 4/8	
15	248	2 + 6/8	218	2 + 3/8	188	2	
20	307	3 + 3/8	270	3	233	2 + 4/8	
25	363	4	320	3 + 4/8	276	3	
30	416	4 + 4/8	366	4	316	3 + 4/8	
35	467	5 + 1/8	411	4 + 4/8	355	3 + 7/8	
40	517	5 + 5/8	455	5	393	4 + 2/8	
45	564	6 + 1/8	497	5 + 4/8	429	4 + 6/8	
50	611	6 + 6/8	538	5 + 7/8	464	5 + 1/8	
55	656	7 + 2/8	577	6 + 3/8	499	5 + 4/8	
60	700	7 + 5/8	616	6 + 6/8	532	5 + 7/8	
70	786	8 + 5/8	692	7 + 5/8	597	6 + 4/8	
80	869	9 + 4/8	765	8 + 6/8	660	7 + 2/8	

Urinary S/O Ageing 7+ is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Ageing 7+ for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

**[S/O® INDEX]** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR AGEING DOGS (> 7 YEARS OLD) IN CASE OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Hyperkalemia
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### URINARY TRACT HEALTH

Effective struvite stone dissolution & specifically formulated for up to 80% of all urinary stones.



### AGEING SUPPORT

Support dogs' vitality and cognition with an exclusive blend of nutrients & antioxidants. Also support kidneys health.

KEY VALUES	Per 1000 kcal	Per 100g as fed
Protein	24.69g	20g
Fat content	44.18g	17g
Carbohydrate	114.35g	44g
Crude fibre	6.24g	2.4g
Dietary fibre	14.81g	5.7g
Calcium	1.74g	0.67g
Phosphorus	1.46g	0.56g
Potassium	4.55g	1.75g
Sodium	1.04g	0.4g
Chloride	3.82g	1.47g
Magnesium	0.23g	0.09g
Copper	3.9mg	1.5mg
Zinc	44.18mg	17mg
Taurine	0.49g	0.19g
Vitamin A	4417.88IU	1700IU
Vitamin E	124.74mg	48mg
Omega 6	8.97g	3.45g
Omega 3	2.39g	0.92g
EPA + DHA	1.04g	0.4g

Metabolisable energy\* 385kcal/100g as fed  
\*Calculated according to NRC2006.

### COMPOSITION

Rice, maize flour, dehydrated poultry protein, animal fats, wheat flour, minerals, hydrolysed animal proteins, maize gluten, cellulose powder, wheat gluten\*\*, egg powder, fish oil, soya oil, dried tomato pulp (source of lycopene), fructo-oligo-saccharides, hydrolysed crustaceans (source of glucosamine), marigold extract (source of lutein), hydrolysed cartilage (source of chondroitin). Protein sources: dehydrated poultry protein, hydrolysed animal proteins, maize gluten, wheat gluten, egg powder.

\*\*L.I.P.: protein selected for its very high digestibility

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 15500IU, Vitamin D3: 1000IU, E1 (Iron): 60mg, E2 (Iodine): 5.3mg, E4 (Copper): 14mg, E5 (Manganese): 67mg, E6 (Zinc): 145 mg, E8 (Selenium): 0.11mg. Urine acidifying substances: calcium sulphate (1.4%). Technological additive: sodium bisulphate (urine acidifying substance): 5g - Preservatives - Antioxidants.

\*\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



# URINARY S/O AGEING 7+



LOAF

Available in box of 12 pouches of 85g

## RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
2	203	2 + 1/2	178	2	154	2	
4	341	4	300	3 + 1/2	259	3	
6	462	5 + 1/2	406	5	351	4	
8	573	6 + 1/2	504	6	435	5	
10	677	8	596	7	515	6	
15	918	11	808	9 + 1/2	698	8	
20	1132	13 + 1/2	1002	12	866	10	
25	1346	16	1185	14	1024	12	
30	1544	18	1358	16	1173	14	
35	1733	20 + 1/2	1525	18	1317	15 + 1/2	
40	1915	22 + 1/2	1686	20	1456	17	

Urinary S/O Ageing 7+ is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Ageing 7+ for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

**[S/O® INDEX]** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR AGEING DOGS (> 7 YEARS OLD) IN CASE OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Hyperkalemia
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### URINARY TRACT HEALTH

Effective struvite stone dissolution & specifically formulated for up to 80% of all urinary stones.



### AGEING SUPPORT

Support dogs' vitality and cognition with an exclusive blend of nutrients & antioxidants. Also support kidneys health.

KEY VALUES	Per 1000 kcal	Per 100g as fed
Protein	91.68g	8.7g
Fat content	62.17g	5.9g
Carbohydrate	44.26g	4.2g
Crude fibre	17.91g	1.7g
Dietary fibre	23.18g	2.2g
Calcium	2g	0.19g
Phosphorus	1.48g	0.14g
Potassium	4.43g	0.42g
Sodium	1.05g	0.1g
Chloride	4.21g	0.4g
Magnesium	0.14g	0.01g
Copper	3.16mg	0.3mg
Zinc	27.4mg	2.6mg
Taurine	1.29g	0.12g
Vitamin A	50579.56IU	4800IU
Vitamin E	221.29mg	21mg
Omega 6	20.02g	1.9g
Omega 3	1.9g	0.18g
EPA + DHA	1.05g	0.1g

Metabolisable energy\* 95kcal/100g as fed  
\*Calculated according to NRC2006.

### COMPOSITION

Pork by-products, poultry by-products, chicken meat, corn starch mixture, sunflower oil refined, powdercellulose, minerals, pork blood products, crude lecithins, caramelised sugars, fish oil, dried tomato pulp, marigold extract (source of lutein).

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin D3: 210IU, E1 (Iron): 4mg, E2 (Iodine): 0.2mg, E4 (Copper): 1.5mg, E5 (Manganese): 1.3mg, E6 (Zinc): 13mg. Urine acidifying substances: calcium sulfate: 0.28%.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





## URINARY + SATIETY

Available in  
3.5 and 9kg bags

## RECOMMENDED DAILY INTAKE &amp; FEEDING DURATION

ADULT DOG	TARGET BODY WEIGHT OF DOG	START		AFTER 4 WEEKS WITH NO WEIGHT LOSS	
		grams	cups	grams	cups
	2	44	4/8	37	4/8
	4	73	7/8	63	6/8
	6	99	1 1/8	85	1
	8	123	1 4/8	106	1 2/8
	10	146	1 6/8	125	1 4/8
	12	167	2	143	1 6/8
	14	188	2 2/8	161	1 7/8
	16	207	2 4/8	178	2 1/8
	18	227	2 6/8	194	2 2/8
	20	245	2 7/8	210	2 4/8
	25	290	3 4/8	249	3
	30	332	4	285	3 3/8
	35	373	4 4/8	320	3 6/8
	40	413	4 7/8	354	4 2/8
	45	451	5 3/8	386	4 5/8
	50	488	5 6/8	418	5
	60	559	6 5/8	479	5 6/8
	70	628	7 4/8	538	6 3/8
	80	694	8 2/8	595	7 1/8
	90	758	9	650	7 6/8

For nutritional management of dogs at risk of struvite stone recurrence and reduction of excess bodyweight: feed the diet up to 6 months and re-evaluate based on clinical assessment of the response. If a urinary tract infection is present S/O-Kcal Control Canine, combined with an antibacterial agent, should be fed for at least one month after a negative urine culture result has been obtained. For detailed guidance on weight reduction please see product information for Satiety Canine.

**RECOMMENDATIONS:** Target Weight should be determined according to Body Condition Score of the individual. The optimal weight loss rate is 1-3% bodyweight weekly. It is recommended to continue feeding the weight-loss diet once target weight is achieved. Energy intakes should be increased by 10% and Body Condition Score and weight monitored closely throughout the stabilisation period. Should the dog regain more than 5% of body weight, the daily ration should be decreased to weight loss levels. It is recommended to divide the daily intake into 2 meals.

**[S/O® INDEX]** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

## RECOMMENDED FOR CASES OF

**Urinary disorders:** • Recurrent struvite and calcium oxalate urolithiasis • Bacterial cystitis • Dissolution of struvite uroliths. **In older dogs**, renal function should be checked before S/O-Kcal Control Canine is recommended.

+ **Weight Management:** Obesity • Diabetes mellitus associated with obesity • Hyperlipidaemia associated with obesity

## NOT RECOMMENDED IN CASE OF

Pregnancy, lactation, growth • Chronic Kidney Disease • Heart disease (where sodium restriction is required) • Concurrent use of urine-acidifying drugs • Conditions where high fibre content is not recommended



## SATIETOGENIC EFFECT

Contains a special blend of fibres which have a satiating effect on dogs.



## HIGH PROTEIN

A high protein content (111 g/1000 kcal ME) helps maintain muscle mass throughout the weight loss period.



## URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



## LOW RSS

Helps lower the concentration of ions contributing to struvite and calcium oxalate crystal formation.

KEY VALUES	Per 100g as fed
Protein	30g
Fat content	9.5g
Carbohydrate	16.9g
NFE	26.9g
Dietary fibre	26.7g
Crude fibre	16.7g
Omega 6	1.96mg
Omega 3	0.61mg
EPA+DHA	0.29g
Calcium	0.85g
Phosphorus	0.7g
Sodium	0.95g
L-Carnitine	30mg
Chondroitin sulphate + Glucosamine	100mg

Metabolisable energy\* 269.9Kcal

\*Calculated according to NRC2006.

## COMPOSITION

Dehydrated poultry protein, cellulose powder, wheat gluten\*\*, vegetable fibres, tapioca, wheat, hydrolysed animal proteins, chicory pulp, maize, animal fats, minerals, maize gluten, fish oil, psyllium husks and seeds, fatty acid salt, fructo-oligo-saccharides, soya oil, hydrolysed crustaceans (source of glucosamine), hydrolysed cartilage (source of chondroitin).

## ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 19800IU, Vitamin D3: 1000IU, E1 (Iron): 40mg, E2 (Iodine): 4mg, E4 (Copper): 12mg, E5 (Manganese): 52mg, E6 (Zinc): 154mg, E8 (Selenium): 0.07mg  
Preservatives - Antioxidants.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## URINARY + HYPOALLERGENIC

Available in  
3.5 and 10kg bags

## RECOMMENDED DAILY INTAKE &amp; FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
	2	50	5/8	43	4/8	41	4/8
	4	84	1	72	6/8	69	7/8
	6	113	1 + 2/8	98	1 + 1/8	94	1 + 1/8
	8	141	1 + 5/8	122	1 + 3/8	116	1 + 3/8
	10	166	1 + 7/8	144	1 + 5/8	138	1 + 5/8
	15	226	2 + 4/8	195	2 + 2/8	186	2 + 2/8
	20	280	3 + 1/8	242	2 + 6/8	231	2 + 6/8
	25	331	3 + 6/8	286	3 + 2/8	273	3 + 2/8
	30	379	4 + 2/8	328	3 + 6/8	313	3 + 6/8
	35	426	4 + 6/8	368	4 + 1/8	352	4 + 2/8
	40	471	5 + 2/8	407	4 + 5/8	389	4 + 5/8
	50	557	6 + 2/8	481	5 + 3/8	460	5 + 4/8
	60	638	7 + 1/8	551	6 + 2/8	527	6 + 2/8
	70	716	8 + 1/8	619	7	592	7
	80	792	8 + 7/8	684	7 + 6/8	654	7 + 6/8

For nutritional management of dogs with struvite urolithiasis and Adverse Food Reaction: feed the diet for 5-12 weeks and re-evaluate based on clinical assessment of the response. If a urinary tract infection is present S/O-AFR Canine, combined with an antibacterial agent, should be fed for at least one month after a negative urine culture result has been obtained. It is recommended to divide the daily intake into 2 meals.

**[S/O® INDEX]** This product creates a urinary environment unfavourable to the formation of struvite and calcium oxalate stones.

## LOWER URINARY TRACT DISEASE

## RECOMMENDED FOR CASES OF

**Urinary disorders:** • Dissolution of struvite uroliths • Recurrent struvite and calcium oxalate urolithiasis • Bacterial cystitis. **In older dogs**, renal function should be checked before S/O-AFR Canine is recommended.  
+ **Dermatological/GI conditions:** Food elimination trial • Food allergies • Food intolerance • Food responsive Inflammatory Bowel Disease (IBD) • Chronic diarrhoea • Small Intestinal Bacterial Overgrowth (SIBO) • Exocrine Pancreatic Insufficiency (EPI)

## NOT RECOMMENDED IN CASE OF

Pregnancy, lactation, growth • Chronic Kidney Disease • Heart disease (where sodium restriction is required) • Concurrent use of urine-acidifying drugs



## URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



## LOW RSS

Helps lower the concentration of ions contributing to struvite and calcium oxalate crystal formation.



## HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to help ensure very low allergenic potential.



## SKIN BARRIER

A complex to support the barrier effect of the skin.

KEY VALUES	Per 100g as fed
Protein	20g
Fat content	15g
Carbohydrate	40.4g
NFE	44.9g
Dietary fibre	6.9g
Crude fibre	2.4g
Omega 6	3.48mg
Omega 3	0.87mg
EPA+DHA	0.36g
Calcium	0.51g
Phosphorus	0.5g
Sodium	1.3g
Potassium	0.8mg
Vitamin D3	90IU

Metabolisable energy\* 371.6Kcal

\*Calculated according to NRC2006.

## COMPOSITION

Rice, hydrolysed soya protein isolate, animal fats, minerals, chicory pulp, vegetable fibre, soya oil, fish oil, fructo-oligo-saccharides, fatty acid salt, borage oil.

## ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 28500IU, Vitamin D3: 800IU, E1 (Iron): 43mg, E2 (Iodine): 3.9mg, E4 (Copper): 15mg, E5 (Manganese): 57mg, E6 (Zinc): 205mg, E8 (Selenium): 0.25mg  
Technological additives: Clinoptilolite of sedimentary origin: 10g. Preservatives - Antioxidants.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





[WWW.ROYALCANIN.CO.UK](http://WWW.ROYALCANIN.CO.UK)

**NUTRITIONAL HELPLINE**

**+44(0)800 717800**