

## What is Pet Poison Helpline?

- 24/7 animal poison control
- Veterinary & human expertise
  - - 20 DVMs, 50 CVTs
      - DABVT, DABT
         DACVECC
         DACVIM
    - 7 PharmDs
- Case fee of \$49 includes
- Unlimited consultation
- Fax or email of case report
- · Educational center
  - Free webinars (archived)
  - Tox tools

    - Wheel of Vomit
       Pot of Poisons (toxic plants)
  - Textbook
  - iPhone app
  - Newsletters for vet professionals
  - Free resources for clinics
    - VideosElectronic materialClings









1
Τ

# Best. Pet Insurance. Ever. whole pet with wellness

With more coverage than any other plan, Whole Pet with Wellness® works like pet insurance is supposed to!

90% on virtually everything

- -> from Bordetella to bilateral ACL surgeries
- -> from dental cleanings to dysplasias

It's the best pet insurance plan ever made And it's *only* from Nationwide!



Best. Pet Insurance. Ever.



Recommend Whole Pet with Wellness® for all puppies & kittens

- One annual \$250 deductible
- Exam fees (wellness and emergency)
- Preventive care: Spay/neuter, vaccinations and parasite preventives
- Internal and external parasite treatments (e.g. heartworm)
- Teeth cleaning and dental diseases
- · Hereditary and congenital conditions such as hip dysplasia
- Bilateral knee conditions such as ACL with no waiting period
- Prescribed nutritional supplements and therapeutic diets
- Elective procedures (e.g. gastropexy)

Best. Pet Insurance. Ever.



So much coverage, it's easier to tell you what's not covered.

- Pre-existing conditions
   Non-veterinary fees (e.g. tax & waste)
- 3. Boarding and grooming



Remember, if it doesn't include preventive care, it doesn't cover the whole pet!

#### Best. Pet Insurance. Ever.



#### Our goal is to keep pets healthy and happy!

\*Be sure that the pet insurance you recommend covers the whole pet and that includes preventive care and your examinations!



#### POISON HELPLINE 800.213.6680

# **Speaker Introduction**



#### Charlotte Flint, DVM

Senior Consulting Veterinarian, Clinical Toxicology

Pet Poison Helpline Minneapolis, Minnesota



#### POISON HELPLINE 800.213.6680

#### **Valentine Toxins**

- Valentine's Day flowers
- Valentine's Day treats
  - Chocolate
  - What else is in that chocolate?
  - Alcohol



- Valentine's Day medications
  - OTCheadache/hangover
  - medications

     Erectile dysfunction
  - drugs
     Hormonal birth
    control





#### **Common Flowers in Bouquets**

- Lilies
- Calla lilies
- Peruvian lilies
- Tulips
- Daisies
- Sunflowers

- Chrysanthemums
- Baby's breath
- Carnations







# What are these flowers??

- Florist
- Master gardener
- University horticulture or botanydepartment
- Extension service
- Garden center







#### Lilies – which are which?

- Lilium species
  - Easter lilies, tiger lilies, stargazer lilies
- Hemerocallis species
  - Day lilies
- Calla lilies
- Peruvian lilies (Alstromeria)
- Lily of the valley
- Peace lilies





#### Lilium and Hemerocallis species

- Renal toxin in cats
- All parts of the plant are toxic
- All exposures should be considered serious, including pollen and vase water exposures









#### Clinical Signs and Course of Toxicity

- <u>0-12 hours post</u> <u>ingestion</u>
  - Salivation, vomiting, anorexia, depression
- 12-24 hours post ingestion
  - Polyuria, dehydration
  - Azotemia with creatinine often disproportionately higher than BUN
- 24-72 hours post ingestion
  - Renal failure
  - Continued anorexia, vomiting, depression, weakness, anuria, death if untreated





#### PET POISON HELPLINE 800.213.6680

# Treatment of Lily Poisoning

- Emesis if recent
- Bathe if pollen exposure
- Activated charcoal x 1 dose if recent
- Baseline lab work and recheck q 24 hours for at least 48 hours or until normalized
- IV crystalloids x 48 hours minimum
- GI support with antiemetics, GI protectants as needed



## Calla Lilies - Zantedeschia sp

- Insoluble calcium oxalate crystals
- Can cause self-limiting drooling, oral irritation, vomiting, vocalizing, pawing at the mouth
- Treatment is symptomatic:
  - Rinse mouth, dilute milk or yogurt
  - Antiemetics, fluids, GI protectants prn



# Peruvian Lilies /Alstromeria

- Mild and self-limiting GI upset
- Treatment is symptomaticand supportive
  - Anti-emetics, fluids if needed







#### **Educational campaign**

- www.noliliesforkitties.com
- Download free graphics/lily info
- Free video
- List of safer flower alternatives

NO LILIES F	OR KITTIES!
	o (a
lapanes	e Show Lily
Lillium species ("true lilies") such as Easte	Easter Lily les dont mix!  r, tiger, Assatic, Oriental, etc., and dayfiles as or people). Indesting petals, leaves, pollen.
or even water in the vase can result in kidney lily, call Pet Poison Helpline, your local emer	y failure and death. If your cat eats any part of a gency clinic or your veterinarian immediately. It is imperative!
SEE BELOW FOR SAFER FLOWER Of Other lifty varieties shown here such as peace (Spath)s/flvm/n, Peruvian (Alstromensa) and calla (Zantedeschu) are much less dangerous and cause only minor symptoms when eaten.	HOICES FOR CATS
SAFER* cut flower choices include: Roses, sunflower carnations, mums, baby's breath, hyacinth and daffodili	
*These flowers may cause vomiting and clarrhea but are not considered life threatening. www.notillesforkittles.com	Pet Poison HELPLINE 800.213.6680 Wesperposonispine con

PET
800.213.668
800.213.008

## Roses (Rosa species)

- GI irritant with possibility for mechanical trauma
- Symptomatic and supportive treatment, if needed
  - Anti-emetics
  - Fluids
  - Check for trauma from thorns





# Lots of GI Irritant Flowers!

- Baby's breath
- Daisies
- Chrysanthemums
- Tulips
- Carnations
- Irises
- Orchids
- Dahlias
- Freesia
- Sunflowers

- Signs
  - Vomiting, diarrhea, anorexia
- Symptomatic and supportive treatment
  - None if small ingestion and signs are mild
  - Decontaminate if a large
  - ingestionAntiemetics
  - Fluids





#### Chocolate





#### POISON HELPLINE 800.213.6680

#### **Chocolate Questions**

- What kind of chocolate?
  - Unsweetened > semi-sweet > dark > milk > white
- What brand and product?
- What is the weight or amount?
- What other ingredients?
  - Xylitol, nuts, raisins, coffee, alcohol, more chocolate, marijuana
  - Recipes for baked goods?
- Foreign body potential?





## **Chocolate Signs**

- Mild signs >20 mg/kg theobromine
  - Vomiting, diarrhea, excitability, increased thirst
- Moderate signs >40-50 mg/kg theobromine
  - Tachycardia
- Severe signs >60 mg/kg theobromine
  - Tremors
  - Seizures







#### **Chocolate Treatment**

- Induce emesis up to 6 hours after ingestion
- Activated charcoal possibly multiple doses
- Antiemetics
- Fluids
- Frequent walks
- Monitoring HR, BP, CNS status
- Sedation (acepromazine 0.02-0.04 mg/kg, butorphanol 0.2-0.4mg/kg IV)





#### **Chocolate Treatment**

- Beta blockers or anti-arrhythmic if persistent tachycardia HR > 180
- Methocarbamol 55-220mg/kg IV to effect for tremors
- Benzodiazepine or other anticonvulsants for seizures
- Monitor for pancreatitis







#### Coffee and Caffeine

- Rapid onset of signs usually within 1-2 hours
- Signs
  - GI: vomiting, diarrhea
  - CNS: agitation, tremors, seizures
  - Cardiac stimulation
  - Hypokalemia
  - Hyperthermia







#### **Caffeine Treatment**

- Emesis?
- AC maybe multiple doses
- IV fluids
- Frequent walks or urinary catheter



- Sedation
- CNS and cardiac monitoring
- · Beta blockers
- Anticonvulsants
- Potassium supplementation





#### Marijuana

- Toxic dose of THC: Minimum lethal oral dose 3 g/kg, signs occur at much lower doses
- Clinical signs
  - Ataxia, hyperesthesia, urinary incontinence, sedation or agitation, hypothermia, vomiting
  - Severe cases- coma, respiratory depression, bradycardia, hypotension
- Treatment
  - Symptomatic and supportive decontamination, fluids, anti-emetics, monitoring of vitals
  - Possibly IV lipid therapy for severe cases





#### **Xylitol**



- Sources
  - Sugar free, low calorie, and/or low carb foods
  - Gums, mints
  - Dental products toothpaste, mouthwash and dental floss
  - Personal lubricants (Astroglide), lotions, cosmetics
- Other sugar alcohols
  - Sorbitol, mannitol, erythritol, maltitol, isomalt
  - GI signs only





## **Xylitol Signs**

- Hypoglycemia > 0.1 g/kg
  - Rapid and dose dependent rise in seruminsulin causes hypoglycemia
- Liver toxicity >0.5 g/kg
  - MOA unknown
- Rapid onset of signs
  - Vomiting
  - Hypoglycemia: lethargy, ataxia, recumbency, tremors, seizures
  - Hepatic necrosis: elevated ALT andtbili, vomiting, lethargy, anorexia, icterus, coagulopathy





#### **Xylitol Treatment**

- Check BG before inducing emesis
- No activated charcoal
- Antiemetics
- Small frequent meals
- Monitor BG q 2-6 hours depending on patient status and dose
- Monitor electrolytes (hypokalemia)
- Supplement dextrose +/- KCl
  - Supplement in fluids even with normal BG if a liver toxic





#### **Xylitol Treatment**

- Monitor liver enzymes q 24 hours for 48-72 hours or until normalized
  - Start liver protectants SAM-e
  - Consider N-acetylcysteine if liver enzymes are elevating or if a very high dose ingested
- Treatment for coagulopathy if indicated
  - Vitamin K1
  - Transfusion





#### **Alcohol**

- Beer, wine, hard alcohol Rapid onset of signs drinks
- Bread dough, pizza dough
- · Chocolate fillings, alcohol-soaked desserts



- within 30-90 minutes
  - Depression, weakness, ataxia
  - Hypoglycemia
  - Hypothermia
  - · Hypotension, tachycardia
  - GI signs
  - Metabolic acidosis



л	- 4	



#### **Alcohol Treatment**

- Decontamination
  - Emesis unlikely to be helpful with liquid alcohols but could be helpful if ingested as dough or with chocolate or other food
  - Activated charcoal does not bind alcohol
- IV crystalloids with dextrose supplementationas needed
- Anti-emetics
- Monitor BG, monitor vitals
- Nursing care thermoregulation, turn, lube eyes





#### **Valentine Medications**

- OTC headache/hangover medications
- Erectile dysfunction drugs
- Hormonal birth control





#### PET POISON HELPLINE 800.213.6680

#### **OTC Headache Medications**

- OTC NSAIDs
  - Ibuprofen
  - Naproxen
  - Aspirin
- Acetaminophen
- Combos
  - Acetaminophen, aspirin, and caffeine
- Cats are more sensitive to these drugs due to their limited glucuronylconjugating capacity







#### **OTC NSAID Clinical Signs**

- GI irritation and ulceration
  - Vomiting +/- blood
  - Melena
  - Anemia
- · Renal toxicity
- CNS signs
  - Sedation, seizures with high doses
- Aspirin OD can also cause tachypnea, hyperthermia, coagulopathy





#### **OTC NSAIDs Treatment**

- · Emesis if recent
- Multidose AC q 8 h up to 24 hours
- Baseline lab work and recheck q 24 h x 2-3 days
- Antiemetics
- GI protectants for 7-10 days
- IVF for 2-3 days if a renal dose
- Liver protectants
- Anticonvulsants, possibly naloxone prn





## Overdose of Acetaminophen

- Dogs
  - Liver toxicity, methemoglobinemia (high doses), KCS, rare renal toxicity
  - Depression, anorexia, vomiting, abdominal pain, with progression to liverfailure
- Cats
  - Methemoglobinemia then hepatic toxicity
  - Anorexia, salivation, vomiting, facial and paw edema with lacrimation and pruritus, depression, Heinz body anemia, rare renal toxicity





## Treatment of Acetaminophen OD


- Emesis if recent
- Activated charcoal
- Monitoring of lab work
- Antidote: Nacetylcystine (diluted) 140 – 280 mg/kg orally or IV once, followed by 70 mg/kg q 6 hours X 7-8 treatments or up to 17 doses if signs
- IV fluids
- Hepatoprotectants SAM-e, silymarin
- Antiemetics
- Antioxidants to prevent RBC damage Vit C
- Transfusion
- Oxygen
- GI protectants not needed!





# **Erectile Dysfunction Drugs**

- Phosphodiesterasetype5 (PDE5) inhibitors
  - Sildenafil (Viagra)
  - Tadalafil (Cialis)
  - Vardenafil (Levitra, Staxyn)
  - Avanafil (Stendra)



 Relax smooth muscle in the pulmonary vascular bed in the lungs and the corpus cavernosum of the penis





#### **Erectile Drugs Signs**

- Vasodilation
- Hypotension
- Tachycardia (reflex tachycardia)
- Weakness
- Vomiting
- Headaches (human)
- Priapism





#### PET POISON HELPLIN 800.213.668

#### **Erectile Drugs Treatment**

- · Rapidly absorbed
  - Consider emesis and AC if recent ingestion
- Monitor HR and BP closely
- Bloodwork, especially if prolonged hypotension develops
- IVF for BP support
  - Vasopressors if hypotension does not respondto
- Antiemetics





#### **Hormonal Birth Control**

- Oral contraceptives
- Patch
- Ortho Evra
  - Xulane
- Ring
  - Nuvaring



- Estrogens
  - Ethinyl estradiol
  - Estradiol valerate
- Progesterones
  - Norethindrone
  - Levonorgestrel
  - Etonogestrel
  - Norgestimate





#### **Hormonal Birth Control**

- May have different doses of hormone based on the week – ask brand name, pill colors, and number of each pill color!
- Placebo pills
  - Might contain iron and if so, typically have FE in the brand name







#### **Hormonal Birth Control**

- Usually not very toxic as hormone doses are low
- Estrogen >1 mg/kg can cause bone marrow suppression (anemia, thrombocytopenia, leukopenia)
- Progesterone can cause GI signs with ataxia and sedation at higher doses
- Consider foreign body potential with ingestion of patch, ring, or packaging





#### Hormonal Birth Control Example

- Lo Loestrin Fe
  - 24 active blue pills with
     1 mg norethidrone and
     10 mcg ethinyl estradiol
  - 2 active white tablets with 10 mcg ethinyl estradiol
  - 2 brown inactive tablets with 75 mg ferrous fumarate (25 mg elemental iron)
- A 10 kg dog eats all 28 of the Lo Loestrin Fe pills
- Total estrogen
  - $-0.26\,{\rm mg}$  or  $0.026\,{\rm mg/kg}$
- Total progesterone
  - 24 mg or 2.4 mg/kg
- Total iron
  - 50 mg or 5 mg/kg elemental iron





#### **Hormonal Birth Control Treatment**

- Consider emesis if recent and/or if foreign body concern
- Activated charcoal if >1 mg/kg estrogen
- Baseline CBC with recheck in 7 and 14 days or until normal
- Nuvaring is not radiopaque
- GI support antiemetics, fluids prn
- Transfusion if anemic
- Antibiotics if leukopenic
- Neupogen or Epogen to stimulate WBC and RBC production from bone marrow, respectively













