

# Wheat growth guide

**AHDB**  
CEREALS & OILSEEDS



# Managing wheat growth

## Crop managers need to respond to unexpected weather or growth throughout the season.

The steps in any management cycle are to:

1. Set targets
2. Assess progress
3. Adjust inputs
4. Monitor success

Measurement is vital for effective management at every stage of a crop's progress.

In addition to assessments on weeds, pests and diseases, managers must assess the crop itself. Crop assessments should be objective, targeted and, where possible, measured.

This guide presents measurements by which grain production targets can be set and progress monitored. It also explains how measurements interrelate.

## Contents

Managing wheat growth	2
Wheat growth stages and benchmarks	4
Crop life cycle	6
Establishment	8
Leaf emergence and tillering	10
Root growth and distribution	12
Nitrogen uptake	14
Canopy expansion and senescence	16
Dry matter growth	18
Stem elongation	20
Stem carbohydrate storage	21
Ear formation	22
Grain filling and ripening	23
Yield, quality and protein deposition	24
Measurements	26
Glossary	27



## Benchmarks

This symbol identifies a benchmark, a quantitative reference point against which a crop's performance can be compared. While benchmarks are compatible with good yields, they should not necessarily be regarded as management targets.

Page 4 gives the important growth stages and page 5 gives the benchmark values for key processes. All benchmarks are then explained in subsequent sections.

Each benchmark is based on observations of a high-yielding feed wheat with a slow rate of development at UK sites through several seasons. Unless otherwise stated, crops were sown in early October at 375 seeds/m<sup>2</sup> and grown with ample nutrition, complete crop protection and without lodging.

Note that some graphs used to illustrate growth processes are based on example crops and so may give data that differ from benchmarks. Some benchmark data for modern varieties can be found within the AHDB Recommended List datasets ([cereals.ahdb.org.uk/varieties](https://cereals.ahdb.org.uk/varieties)).


By assessing crops against benchmark values, growers can determine how best to manipulate husbandry. Some targets and husbandry responses are suggested but this guide is not an agronomy manual.



## Green Area Index

Canopy size can be expressed as Green Area Index (GAI) – the ratio of total green area (one side only) to the ground area occupied. These photographs illustrate typical GAIs. For more information, see page 26.

**Illustration of GAI = 2**  
(two areas of green leaf and stem to one area of ground)






## For more information

Publications and details of projects funded by AHDB Cereals & Oilseeds are all available at [cereals.ahdb.org.uk/publications](http://cereals.ahdb.org.uk/publications)


### AHDB Recommended Lists for cereals and oilseeds (annual)

- G67** Barley growth guide (2015)
- G65** Oilseed rape guide (2015)
- G63** Wheat disease management guide (2015)
- G61** Managing weeds in arable rotations – a guide (2014)
- G49** Cereal growth stages – a guide for crop treatments (2009)
- G48** Nitrogen for winter wheat – management guidelines (2009)



# Wheat growth stages and benchmarks

## Growth stages



Growth Stage	GS13	GS30	GS31	GS39	GS59	GS61	GS71	GS87	Harvest
--------------	------	------	------	------	------	------	------	------	---------

Growth Stage	Description of stage
<b>Seedling growth</b>	
GS10	First leaf through coleoptile
GS11	First leaf unfolded (ligule visible)
GS13	3 leaves unfolded
GS15	5 leaves unfolded
GS19	9 or more leaves unfolded
<b>Tillering</b>	
GS20	Main shoot only
GS21	Main shoot and 1 tiller
GS23	Main shoot and 3 tillers
GS25	Main shoot and 5 tillers
GS29	Main shoot and 9 or more tillers
<b>Stem elongation</b>	
GS30	Ear at 1 cm (pseudostem erect)
GS31	First node detectable
GS32	Second node detectable
GS33	Third node detectable
GS37	Flag leaf just visible
GS39	Flag leaf blade all visible
<b>Booting</b>	
GS41	Flag leaf sheath extending
GS43	Flag leaf sheath just visibly swollen
GS45	Flag leaf sheath swollen
GS47	Flag leaf sheath opening

Growth Stage	Description of stage
<b>Ear emergence</b>	
GS51	First spikelet of ear just visible above flag leaf ligule
GS55	Half of ear emerged above flag leaf ligule
GS59	Ear completely emerged above flag leaf ligule
<b>Flowering</b>	
GS61	Start of flowering
GS65	Flowering half-way
GS69	Flowering complete
<b>Milk development</b>	
GS71	Grain watery ripe
GS73	Early milk
GS75	Medium milk
GS77	Late milk
<b>Dough development</b>	
GS83	Early dough
GS85	Soft dough
GS87	Hard dough (thumbnail impression held)
<b>Ripening</b>	
GS91	Grain hard (difficult to divide)
GS92	Grain hard (not dented by thumbnail)
GS93	Grain loosening in daytime


## Wheat growth stages


The Decimal Code system for measuring wheat growth used throughout this guide is based on work published in Tottman, DR & Broad, H (1987) The decimal code for the growth stages of cereals, with illustrations. *Annals of Applied Biology* **110**, 441–454.





## Benchmarks


Benchmarks are reference values, compatible with high yields but they are not management targets.


	GS30	31 March	Ear at 1cm
	Plants	260/m <sup>2</sup>	70% of seeds sown.
	Shoots	941/m <sup>2</sup>	Tillering ceases when GAI > 1.
	Roots	0.4 t/ha; 12 km/m <sup>2</sup>	Little of the soil is yet fully rooted.
	GAI	1.6	Only enough to intercept 45% of light.


	GS31	10 April	First node detectable
	Shoots	902/m <sup>2</sup>	Shoot numbers usually start to decrease from GS31.
	Three leaves yet to emerge	28 April 9 May 19 May	One leaf emerges every 122 day degrees.
	Roots	0.5 t/ha; 15 km/m <sup>2</sup>	Roots now reach to about 1 m depth.
	N uptake	81 kg/ha	About 30% of final uptake.
	GAI	2.0	Only enough to intercept half the light.
	Total dry weight	1.9 t/ha	Only 10% of final dry weight.
	Growth rate	0.16 t/ha/day	During the Construction Phase (GS31–61).
	Height to top ligule	9 cm	Stem extension just starting.


	GS39	19 May	Flag leaf blade all visible
	Fertile shoots	655/m <sup>2</sup>	Some young shoots are still dying.
	Total leaf number on main shoot	14	No further leaves emerge.
	N uptake	189 kg/ha	Increasing by 2.5 kg/ha/day.
	GAI	6.2	Enough to intercept 95% of light.
	Total dry weight	6.9 t/ha	About 40% of maximum growth.
	Height to top ligule	34 cm	Late PGR reduces subsequent extension.

	GS59	6 June	Ear completely emerged
	Fertile	495/m <sup>2</sup>	Little further shoot death occurs.
	N uptake	233 kg/ha	36 kg N/ha green area.
	GAI	6.3	GAI reaches its maximum about now.
	Total dry weight	11.4 t/ha	Growth may slow if flowering is delayed.
	Height to collar	53 cm	Five internodes extend.

	GS61	11 June	Start of flowering
	Fertile shoots	460/m <sup>2</sup>	150 additional infertile shoots/m <sup>2</sup> remain until harvest.
	Roots	1.0 t/ha; 31 km/m <sup>2</sup>	Sufficient for full moisture capture to 70 cm depth. Deepest roots reach to ~1.5 m. Root growth now slows.
	N uptake	248 kg/ha	Only 30 kg/ha further uptake occurs.
	GAI	6.3	Canopy senescence is slow.
	Total dry weight	12.1 t/ha	About two-thirds of maximum dry weight.
	Growth rate	0.18 t/ha/day	During the Production Phase (GS61–GS87).
	Height to collar	69 cm	Little further extension occurs. Ear length adds about 10 cm.
	Stem dry weight	7.1 t/ha	33% is soluble, giving 2.3 t/ha for redistribution.
	Ears	1.9 t/ha	Ears have 48 grain sites after flowering. Each ear weighs 420 mg (dry) both now and as chaff at harvest.

	GS71	20 June	Grain watery ripe
	GAI	5.7	Rapid senescence now starts.
	Stem dry weight	7.6 t/ha	Now at its maximum. Rapid redistribution of soluble reserves begins at GS73.
	Total dry weight	13.7 t/ha	All further increase occurs in the grain.

	GS87	29 July	Grain at 'hard dough'
	GAI	1.3	All greenness will be lost in the next few days.
	Total dry weight	19.6 t/ha	About 0.8 t/ha is subsequently lost, mainly from the straw.
	Grain filling	Lasts 45 days	Grain filling stopped at about 45% moisture, about 3 days before GS87.

	Harvest ripe 9 August		
	Ears (fertile shoots)	460/m <sup>2</sup>	At least 400 shoots/m <sup>2</sup> required to avoid yield loss.
	N uptake	282 kg/ha	68% of final crop N is in grain, 32% in chaff, straw and stubble.
	Total dry weight	18.4 t/ha	51% grain, about 10% chaff, the rest as straw and stubble.
	Straw dry weight	7.3 t/ha	Includes stems and leaves; only 0.2 t/ha soluble sugars remains.
	Chaff dry weight	2.0 t/ha	Chaff is 430 mg/ear.
	Grain weight	50 mg at 15% moisture	Specific weight 78 kg/hl.
	Grain protein	11.5% (dry basis)	Calculated as 19.6% N x 5.7.
	Grain yield	11.0 t/ha at 15% moisture	Shedding losses are 0.03 t/ha.

# Crop life cycle

Throughout the growing season, the plant both changes in form (development) and accumulates dry matter (growth).

## Key facts:

- Development is governed by temperature and day length
- The rate at which wheat passes through its life cycle may only be managed through variety choice and sowing date

## The key phases

At successive growth stages, crop processes 'switch' on or off. Key stages are crop emergence (GS10), the start of stem extension (GS31), flowering (GS61) and the end of grain filling (GS87). These key stages separate the important phases.

Crop development can be divided into three phases: **Foundation, Construction and Production**


## Development phases

The duration of each phase is governed by:  
**Vernalisation:** A period of cool temperatures (0–12°C) advances floral development. Vernalisation reduces the duration of the Foundation Phase. Winter wheat varieties respond strongly to vernalisation; spring wheats may have a slight response.

**Temperature:** Affects the duration of all crop development phases. Warmth shortens phase length. More growth occurs in any phase during cool, rather than warm, temperatures as phase duration is prolonged.


**Day length:** Long days advance floral development in most varieties. Day length affects the duration of both Foundation and Construction Phases. Almost all commercial UK wheat varieties respond to day length.

Varieties vary in their response to vernalisation and day length. AHDB Recommended List tables give the relative speed of development of varieties to reach GS31 from early, medium and late sowings. They also provide conservative times for the latest safe sowing date for each variety to allow sufficient vernalisation to trigger flowering. The data are updated each year, so always use the current version.





Crops are at least risk if Construction and Production Phases start when frost risk is over and end before drought risk starts.



## Stress-sensitive stages

The apex is frost tolerant until reproductive development starts. Susceptibility to frost damage is highest when the ear is developing.

Frost risk falls significantly from April. Overall risk of damage is lowest in late May and early June; from July onwards, drought risk increases.

Radiation frosts can damage the ear, especially at flowering, which can significantly reduce yield. By comparison, damage to leaves results in less yield loss.

## What does it mean?

Development can only be managed by variety choice and sowing date.

Prolonging any development phase increases dry matter formation during that phase.

▷ Prolong development by sowing slow-developing varieties early.

For a given variety and sowing date, management after sowing influences growth, not development.

▷ For highest yields, feed and protect leaves that emerge during stem extension in preference to earlier, lower leaves.

Jun Jul Aug

### **B** Production Phase - two months

#### - Post-flowering

Grains fill and ripen. Light is bright but canopy survival and activity depend on good crop protection and ample uptake of nitrogen and water.

GS87

Grain filling

Flowering

Ripening

Root growth slows

GS61

GS92

100 2200 2300 2400 2500 2600 2700 2800 2900 3000 3100

# Establishment

**Establishment includes germination, emergence and overwinter survival.**

## Key facts:

- Poor establishment or low plant population density does not reduce yield, unless:
  - significant areas of the field have very few or no plants
  - conditions are unsuitable for compensatory tillering and root growth
- Establishment decreases from around 70% for sowings in September to less than 50% for sowings in November or later
- Soil type and cultivations can have large effects on establishment: average establishment for sandy soils is 90%, compared with 65% for loams and clays
- Plant density markedly affects crop structure but has little effect on grain yield, above a low threshold
- High seed rates, coupled with good establishment, increase lodging risk

## Sowing to emergence

- B** 150°C days
- 11 days in September
  - 15 days in October
  - 26 days in November


Germination and emergence require moisture and warmth. Initially, seeds imbibe water; roots start to grow; coleoptiles emerge and extend to the soil surface. Then first leaves emerge and seedlings establish.

A proportion of viable seeds fail to emerge due to pests, diseases and soil conditions. Establishment declines if sowing is delayed after mid-October. Establishment will be around 50% of seeds sown in mid-November.

While sowing to emergence takes longer in cold than in warm weather, the thermal time in each case is very similar. Delayed drilling, deep drilling, cool and dry conditions extend the interval between sowing and emergence.

Sowing too deep or too shallow can delay or decrease establishment. Optimum sowing depth is about 4 cm.

### Establishment declines as sowing is delayed



## Overwinter survival

- B** 70% of seeds sown
- B** 260 plants/m<sup>2</sup>

By the end of February, the benchmark for establishment is 70%. Plant damage, or loss over winter, may occur due to:

- frost damage, especially after early drilling of fast-developing varieties
- frost heave
- pest or disease damage
- poor or impeded drainage leading to poor rooting and waterlogging

Tillering tends to compensate for uneven establishment. It may be possible to counteract poor establishment by using fertiliser N to encourage tillering and tiller survival. Very few plants die after winter.

## Calculating seed rate

To achieve a target plant population in spring, seed rate should be set in seeds/m<sup>2</sup> and adjusted for expected establishment after overwinter losses.

$$\text{Seed rate (kg/ha)} = \frac{\text{Target plant population (plants/m}^2\text{)} \times \text{Thousand grain weight (g)}}{\text{Expected establishment (\%)}}$$






## Effects of plant population

Plant density depends on seeds sown and establishment. Eventual crop structure is markedly affected by surviving plants/m<sup>2</sup>. Higher plant densities cause:

- fewer crown roots on each plant
- fewer tillers on each plant
- more fertile shoots to survive, hence more ears/m<sup>2</sup>
- smaller culm leaves, but greater canopy size (GAI)
- fewer grains on each ear
- little change in grain yield, above a low ‘threshold’




## Lodging

High plant population density weakens plant anchorage. It also increases shoot height and decreases stem diameter and stem wall width. This weakens stem base strength.

The benefit of small plant populations is greatest for varieties with poor lodging resistance.

Shallow drilling can increase lodging risk.



## What does it mean?

**Soil type and cultivations** affect establishment. For example, average establishment on sandy soils is 90%, compared with 65% on loams and clays. Cultivations affect seedbed quality and establishment, depending on soil stability and type.

▷ Consider deep cultivation on unstable silt soils.

▷ Consider reduced tillage on stable clay soils.

**Other factors** that may affect emergence and establishment are:

- germination capacity (seed quality and vigour, which are affected by variety or seed crop ripening)
- some seed treatments may delay/reduce emergence, especially of deep-sown seeds
- high seed rates can lead to reductions in percentage establishment
- sowing too deep or too shallow
- disease or pest damage

**Seed rate:** Delayed sowing reduces the tillering period; for each month drilling is delayed, an extra 50 plants/m<sup>2</sup> are needed to compensate for reduced tillering.

▷ Increase seed rates with later sowings.

# Leaf emergence and tillering

**Leaf emergence indicates plant development and sets tillering potential.**

## Key facts:

- Rate of leaf emergence is controlled mainly by temperature
- Each main shoot produces 9 to 14 leaves, of which 5 to 7 are on the extended stem
- At least 400 fertile shoots/m<sup>2</sup> (400 ears/m<sup>2</sup>) are required for maximum yield
- All varieties can produce many tillers
- Early sowing and fertile soils increase tillering
- Tillering is prolonged at low plant populations
- Many tillers die between GS31 and flowering
- High N supplies encourage tiller survival

## Potential tiller production

**B 35 shoots/plant**

Tillering is the emergence of side shoots at leaf-stem junctions. It can continue until after the start of stem extension.

The first tiller emerges in the junction of the first leaf (or coleoptile) as the second, third or fourth leaves emerge. The next tiller develops in the second leaf junction one phyllochron later, and so on. Secondary tillers develop in leaf junctions of primary tillers. Well-spaced plants can produce fertile tillers until stem extension starts; tillers produced later are rarely fertile.

## Leaf emergence

**B 122°C days/leaf (14 main shoot leaves)**


Leaf emergence slows or stops in winter and progressively speeds up as temperatures rise in spring.

Other environmental factors have little effect on leaf emergence.

The phyllochron is the time between emergence of two successive leaves, measured in thermal time.


The phyllochron varies with variety and sowing date. Late sowing decreases both phyllochron and total leaves emerged, eg early September drilling may give 15 leaves; November drilling may give 9 leaves.

T1 fungicides generally target the third from last leaf, which normally coincides with GS32 depending on sowing date and variety.



**The first tiller emerges in the junction of the first leaf (or coleoptile) as the second, third or fourth leaves emerge.**

## Potential fertile tillers depend on leaves present at stem extension



The benchmark crop, sown in early October, produces 8–9 leaves by stem extension, so can produce a maximum of 35 shoots on each plant. November-sown crops may have only 5 leaves by stem extension with a potential for 11 or fewer shoots. This relationship is useful in assessing minimum plant numbers needed at establishment to give at least 400 ears/m<sup>2</sup>.



## Tillering patterns

**B** 1,020 shoots/m<sup>2</sup> (8 April)

The timing of tillering depends on sowing date, plant numbers established and temperature.

**Early sowing (early September):** Most tillering occurs in autumn. If more than 250 plants/m<sup>2</sup> are established, then tillering usually ends before winter. If fewer than 100 plants/m<sup>2</sup> are established, tillering can continue into spring.

**Late sowing (November):** Tillering is usually delayed until spring unless unusually warm temperatures follow drilling.

Tillering normally ends when Green Area Index (GAI) reaches about 1 (see page 3). In dense crops, this usually occurs just before GS31.

Limited resources, especially N, may limit tiller production.

The benchmark maximum for a late September/early October-sown crop with 260 plants/m<sup>2</sup> is 1,020 shoots/m<sup>2</sup>. The benchmark date of maximum shoot number is in early April. Early sown crops or crops with many plants/m<sup>2</sup> tend to have greater maximum shoot numbers. Varieties with large leaves or that reach stem extension early, tend to have lower maximum shoot numbers.

## Shoot survival

**B** 45% (460/m<sup>2</sup>)

Some tillers die between the start of stem extension and flowering, with the last formed dying first. Few die after flowering. Tiller losses are higher in crops with many shoots.

Shoot survival is as important as shoot production in determining final shoot number. Shoot survival varies significantly between varieties – under 40% to over 70%. However, all varieties can produce sufficient tillers.

Increasing N reduces tiller loss. Dry matter losses of up to 3 t/ha can occur as shoots die.

### What does it mean?


Leaf emergence, together with disease risk, determines spray timings.

▷ Generally, target T1 fungicides at the third from last leaf, normally GS32.

Tillering is the most important process governing canopy formation. Some tillers die between stem extension and flowering. Those that survive until flowering result in extra ears.

▷ Manage the crop to achieve at least 400 ears/m<sup>2</sup> for maximum potential yield.

### Tiller production and survival



Source: AHDB Cereals & Oilseeds-funded research

# Root growth and distribution

**Soil structure, management and drainage have major effects on root growth and distribution.**

## Key facts:

- A mature root system has 20+ main roots per plant, with many branches
- Root growth is slow in the Foundation Phase, more rapid in the Construction Phase, then slow during the Production Phase when dry matter is redistributed and roots senesce
- Good rooting, especially deep rooting, will enhance crop growth when water or nitrogen is short

## Foundation Phase


**B** 15 km roots/m<sup>2</sup> at GS31 (0.5 t/ha)

Roots begin to grow at germination, with three to six seminal roots emerging before the second leaf appears. These can grow deep and persist throughout the crop's life.

The number of additional crown roots, which develop from the stem base, relates to leaf and tiller numbers. Once the main shoot has three to four leaves, crown roots appear with thickened upper regions to anchor the plant. The mature root system has 20 or more roots on each plant, plus numerous branches.

In well-drained and well-structured soil, the rate of root extension depends on temperature. In autumn, if soils are warm, seminal roots can grow quickly (12 mm/day). Extension and branching slow during winter, then increase in spring. By GS31, maximum rooting depth can exceed 1 m and root dry weight is about 27% of that of the shoot.

### Root weight and length are proportional



## Construction Phase

**B** 31 km roots/m<sup>2</sup> at GS61 (1.05 t/ha)

During stem extension, roots grow rapidly. Root extension and branching increase as soil temperatures rise (main root extension of 18 mm/day).

This is the main period of crown root production. However, dry matter may be lost as some roots die and as assimilate is exuded or respired. At GS61, root dry weight is 1.05 t/ha (9% of the shoot) but twice as much assimilate may have been used in root growth.

With typical root distribution, total root length reaches 31 km/m<sup>2</sup> by anthesis and maximum rooting depth reaches 1.5–2.0 m.

## Production Phase

**B** 27 km roots/m<sup>2</sup> at GS87 (0.9 t/ha)

After anthesis, root growth slows – only 10% of total assimilate produced during grain filling is used by the root system.

As roots in the topsoil begin to die, those in the subsoil may continue growing.

Protecting leaves with fungicides can prolong root growth and N uptake after GS61.



**Sufficient roots are required for fully effective capture of water and nutrients at depth.**




## Root distribution: Water and nutrient uptake

The relative distribution of roots down the soil profile changes little between GS31 and anthesis. Over 70% of root length is found in the top 30 cm.

High rates of uptake of less mobile nutrients, eg P, only occur when root length densities exceed 5 cm/cm<sup>3</sup> of soil. Lower root length densities are adequate for K, and about 1 cm/cm<sup>3</sup> is needed for uptake of water and N.

Maximising root growth in the subsoil significantly improves soil water supply to the crop.




**Root distribution through the soil profile determines water and nutrient uptake.**

### Typical main root extension rates in deep well-structured soil

Autumn	12 mm/day
Winter	6 mm/day
Spring	18 mm/day

### Roots fully utilise water and nitrogen to about 40cm depth at GS31 and to 70 cm depth by GS61



## What does it mean?

Root systems can only be managed indirectly, through improved soil structure and drainage. Encouraging deep rooting will improve water or nitrogen supplies for crop growth. Varietal differences in crown root spread and low plant number can help to maximise root anchorage strength.

▷ Sow early to increase overall root system size at flowering.

Soil structure has a major impact on root growth and distribution. In some clay-rich soils, moisture extraction by roots promotes cracking, which improves soil structure and root access in following seasons. Hence, deep rooting can be self-sustaining, unless wheelings or cultivations destroy soil structure. With minimal tillage, enhanced earthworm activity creates long continuous pores in the subsoil to aid root penetration.

▷ Consider field drainage and soil management to ensure adequate pore spaces for aeration and root penetration into the subsoil.

Take-all reduces rooting at all depths.

▷ Consider take-all control measures, such as seed treatments or foliar fungicide applications.


# Nitrogen uptake

**Most N is taken up during the four to six week Construction Phase around May.**

## Key facts:

- N uptake can occur throughout the crop's life
- Soils in arable rotations supply sufficient N for wheat to produce roughly half its unlimited yield
- The other half of unlimited yield can be realised with applied N:
  - 40% from the first half of N applied
  - just 10% from the second half of N applied
- Fertiliser N controls canopy size, primarily through shoot number
- N uptake is closely linked to canopy expansion
- During grain filling, a large proportion of N in leaves and stems is redistributed to grain

Variation in soil N in UK arable fields



## Sources of N

**B** 75 kg/ha available soil N

Soil N release and crop recovery are both very variable.

Seed contributes about 5 kg N/ha while some 40 kg/ha comes from the atmosphere, in rain and N-containing gases. However, most is acquired from soil. The benchmark is 75 kg/ha available N (nitrate and ammonium) at soil N supply (SNS) Index 1.

Soil N availability is increased by unrecovered fertiliser from previous crops or organic residues, eg legume roots or animal manures.

Soil N release is stimulated in warm moist soils and after cultivations that thoroughly disturb the soil. Thus, much of the N from a previous crop is released in autumn, before sowing.

Crop residues with low N such as straw, can cause temporary unavailability through 'locking up' N.

Uptake of soil N continues throughout growth. Early sowing and unimpeded rooting improve soil N uptake.

## Canopy nitrogen requirement

**B** 36 kg N/ha of green canopy

N uptake has a major influence on a crop's green canopy. The way that N controls canopy expansion depends upon the stage of crop development:

Stage of development	N uptake affects canopy size by promoting:
Before stem extension	Tillering
During stem extension	Shoot survival, with some increase in final leaf size
After stem extension	Prolonged survival of yield-forming leaves

Soil N is particularly prone to leaching when uptake is low due to slow canopy expansion over winter. As temperatures rise, canopy expansion accelerates and demand for N increases.

Throughout development, the area of green tissues relates to the amount of N they contain; there are about 36 kg N per hectare of green tissue. Thus, it is possible to control canopy size by controlling N availability.



## Pattern of N uptake

**B** 81 kg/ha from sowing to first node (GS31)

**B** 167 kg/ha from first node to flowering

By harvest, a typical crop takes up 279 kg/ha:


- 30% before first node emergence
- 40% between first node and flag leaf (only 5 weeks); most is used to produce 'yield-forming leaves', ie the top four leaves within a crop's canopy
- 20% between flag leaf and flowering
- the remainder slowly after flowering

## N redistribution to grain

**B** 158 kg/ha transferred to grain

**B** 90 kg/ha left in chaff, straw and stubble

### Crop nitrogen uptake



This graph shows a pattern of N uptake compatible with other benchmarks. These values, and those for canopy N requirement and N redistribution, are greater than for most UK crops.

During grain filling, there is a massive redistribution of N within the crop as proteins in the leaves are degraded and N is transferred to form grain protein. This, not root uptake, is the main source of grain N. At harvest, chaff, straw and stubble contain 90 kg N/ha, 158 kg/ha less than at flowering.

Occasionally, crop N decreases slightly before harvest, probably due to loss of leaves.

## What does it mean?

Consider a 'canopy management' approach to N nutrition, using fertiliser N to achieve an optimum canopy size and adequate canopy survival through grain filling.

**Soil mineral N analysis** is a better predictor of available soil N than the SNS index, particularly where soil N residues may be large. It should include an estimate of crop N content at the time of soil sampling.

▷ Obtain a good estimate of eventual soil N supply from soil analysis.

**Autumn N** may occasionally be justified where soil N uptake over winter may be limiting.

▷ Consider autumn N only where N availability may be inadequate over winter, eg on light soils with large amounts of surface straw, and after minimal cultivation.

**Early spring N** is important for some crops.

▷ Apply early N to encourage tillering after poor establishment, or to overcome root restrictions where there is soil compaction or a risk of take-all.

**Late spring N, after tillering** is needed by most crops. It encourages rapid canopy expansion mainly through better tiller survival.

▷ Use late spring N before the canopy turns pale, unless canopy size is excessive.

**Early summer N** helps optimise canopy expansion and survival during grain filling.

▷ Use early summer N, particularly for crops with pale, small canopies or with high yield potential.

**Late summer N** ensures canopy survival through grain filling as well as adequate grain protein concentration for breadmaking.

▷ Consider late N (summer-applied), particularly for crops intended for breadmaking, where yield potential is high and where field history indicates a need.

# Canopy expansion and senescence


**Managing canopy expansion and senescence is key to optimising crop output.**

## Key facts:

- Canopies go through three distinct phases:
  - slow expansion
  - rapid expansion
  - senescence and death
- Canopy size determines the proportion of sunlight intercepted and subsequent dry matter increase

Canopy expansion starts at crop emergence and stops shortly after ear emergence. The canopy dies before harvest. Canopy size can be expressed as Green Area Index – the ratio of total green area (one side only) to the ground area occupied (page 3).

## Change in GAI over the growing season



## Foundation Phase – slow canopy expansion

**B** GAI = 2.0 by GS31

From emergence to early April, ground cover increases. Cover increases as leaves and tillers emerge during early autumn and winter, but GAI rarely exceeds 1 before March.

During this phase, photosynthesis and growth are slow because ground cover is incomplete; both light levels and temperatures are low.

## Construction Phase – rapid canopy expansion

**B** 1 GAI unit every 9 days

Light interception is sufficient for rapid growth when GAI = 3.


Canopy expansion accelerates in late April as temperatures rise and the largest leaves emerge. As stems and leaf sheaths extend, they contribute to GAI. The benchmark rate of canopy expansion is 0.1 GAI/day. At this rate, crops expand by 3 GAI units during May.

Nitrogen availability controls canopy expansion quite closely because crop nitrogen for each unit of green area remains constant at 36 kg/ha – the Canopy N Requirement (CNR). Nitrogen shortage curtails rapid canopy expansion and advances senescence.

At flag leaf emergence, leaf blades comprise about 85% of total GAI. The benchmark date for maximum canopy size, which occurs between flag leaf emergence and ear emergence, is 26 May. The benchmark maximum GAI is 6.9. Maximum canopy size occurs earlier in N-starved crops, as lower leaves begin to die.

Third and fourth leaves from the ear significantly increase GAI but contribute little to grain filling.

## Distribution of green area at flowering







## Production Phase – canopy senescence and death

**B** GAI falls to less than 2 by GS87

The canopy senesces from June onwards. Lowest leaves die first, unless disease intervenes. Leaf sheaths usually die last. GAI drops below 2 at the end of July, causing the end of rapid crop and grain growth. The canopy continues to respire, losing greenness and weight.

## Canopy size and light interception

**B** GAI at GS61 = 6.3

The crop canopy comprises all green surfaces – mainly leaf blades. At flowering, fertile shoots have a GAI of 6.0 and infertile shoots 0.3. The benchmark maximum green area for each fertile shoot is 130 cm<sup>2</sup>.


Husbandry has little effect on leaf number or size, so canopy management focuses on shoot number. As shoot number and GAI increase, the extra light intercepted decreases, eg an increase from GAI = 2 to 3 captures 15% more light, whereas only 2% extra is captured as GAI rises from 6 to 7. Upper leaves become more important for light interception as GAI increases (see diagram).

There is an optimum canopy size at flowering for grain production of about 6.

- **Small canopies** (under 4), which waste sunlight, can result from inadequate plant or shoot survival, as well as N deficiency
- **Large canopies** (over 7), which can result from high seed rates and high N supplies, cost more than is necessary to intercept all available sunlight. They are at high risk from foliar disease and lodging.

Optimum canopy size for varieties with erect leaves or low CNR only differ slightly from varieties with lax leaves. Differences in leaf greenness have little effect on photosynthesis.

Full light capture is hastened by early sowing, warm winters and springs, and adequate moisture and N.



**What does it mean?**

Achieving optimum canopy size is important for good yields.

Eventual canopy size tends to be increased by pre-sowing management:

- early sowing
- high seed rate
- plentiful soil N
- adequate P and K, and correct pH.

During the growing season canopy can be managed by:

- amount and timing of fertiliser N applied
- disease control measures.

Often canopy growth needs to be kept in check to avoid exceeding target GAI.

# Dry matter growth

**Dry matter growth represents the net effect of photosynthesis after losses from respiration (and any shedding).**

## Key facts:

- 90% of final crop dry matter is formed after GS31
- No grain dry matter accumulates during the Foundation Phase; the Construction Phase contributes 20–50% with 50–80% from the Production Phase
- The crop grows by 0.18 t/ha/day (1.3 t/ha/week) from May to July
- Crop growth on dull days is less than half that on bright days

## Foundation Phase

**1.9 t/ha growth by GS31 (9 April)**

Over winter, growth is slow as canopy cover is incomplete and sunlight limited. Just 10% of total growth occurs by GS31 (6 months after sowing). Dry matter formed in this period produces leaves which are all lost before flowering with only their nitrogen being redistributed in the plant.

Root growth of about 0.5 t/ha is additional to growth above ground during the Foundation Phase.

## Construction Phase

**10.2 t/ha increase in growth by GS61**


Rapid growth starts in late April as internodes start to extend, light interception approaches completeness and sunlight intensity increases. Over half of total growth occurs in this phase. Dry matter produced in this period supports at least 0.6 t/ha extra root growth and the formation of all the organs vital to grain production.

Stem reserves act as a buffer. Reserves accumulate when photosynthetic rate is more than adequate and fall when photosynthesis is inadequate, as in dull light.

Each fertile stem has a finite storage capacity for grain dry matter, determined by fertile floret numbers. The amount stored depends on how much dry matter is partitioned to the ear during booting and ear emergence. Partitioning differs between varieties.

Growth may slow towards the end of crop construction if storage capacity is already full.

## Change in crop dry weight over growing season



## Production Phase

**7.5 t/ha increase in growth by GS87, then 0.8 t/ha loss**

Rapid growth continues, although it slows slightly as leaves age and larger organs respire more in warmer weather. Soil water may also become limiting.


Only grains gain dry matter after flowering. Other plant parts (eg stems and leaves) lose weight, although chaff remains constant. As most soluble stem reserves produced pre-flowering are redistributed, grain growth always exceeds total crop growth during this phase.

Canopy senescence occurs as leaf and stem N moves to the grain. Thus, as grain proteins form, photosynthesis progressively slows. Senescence can be delayed if N and water uptake continue.

Rapid grain filling starts at GS71 and ends at about GS87, even if green tissues remain. Early canopy senescence, often due to drought or disease, brings grain filling to a premature end.

Crop dry weight often decreases from its maximum, mainly due to ongoing respiration but also through leaf loss. Dry matter is rarely lost from grain.

## Dry matter distribution at flowering





## Growth

**B** 0.18 t/ha/day from May to July

Slow growth results from incomplete light interception or dull conditions. Rapid growth arises from complete interception of intense sunlight.

Respiration reduces dry matter, particularly when tissues are senescing.

Growth is driven by solar energy. The uptake of carbon dioxide also requires water loss by transpiration. Photosynthetic rate is controlled by:

- light intensity
- green canopy size
- availability of water
- adequate storage for assimilates.

## Resources for growth

Assuming a full canopy, growth may be limited by the availability of solar energy, carbon dioxide or water. Low winter temperatures can also limit growth.

**Solar energy:** Dry matter growth in the UK usually relates directly to solar energy intercepted by the green canopy. About half of this radiation is photosynthetic.

Factors affecting interception of solar radiation are:

- region
- canopy size
- leaf posture
- foliar disease


Light levels reach a maximum in May–July. On sunny summer days, growth can be 0.25 t/ha. Clouds reduce light energy by about two-thirds, so on dull days growth is just 0.1 t/ha. Theoretically, based on solar energy and rainfall, the South West has the greatest potential for crop growth. However, cooler weather in the duller North prolongs wheat development and so increases grain yield.

**Potential growth (light limited) – total dry matter\***


Southern regions have brighter sunlight. This is not fully counteracted by longer summer days in the north. Coastal areas have less cloud.

\*assumes interception of 60% average annual solar radiation

### Solar energy received by a crop is mainly governed by season and cloud cover



**Carbon dioxide:** Atmospheric carbon dioxide is about 370 parts per million (ppm), and is increasing at about 25 ppm a decade. In this range, crop growth relates almost directly to carbon dioxide concentration, so atmospheric change is increasing growth by about 5% a decade. Variation in carbon dioxide concentrations is not significant on a regional or seasonal scale.



**Water:** To absorb carbon dioxide, leaves must lose water to the air in transpiration. On an average summer day, transpiration uses about 3 mm water. For each tonne per hectare of dry matter formed, the crop transpires about 20 mm water. Drought restricts growth.

Factors affecting water availability for transpiration are:

- region: rainfall amounts and distribution
- soil type: moisture retention through summer, when transpiration generally exceeds rainfall
- soil depth and rooting
- take-all and other diseases: reducing root function.



## What does it mean?

Growth can mainly be managed through the size of the crop's green canopy, taking light conditions and water availability into account.

# Stem elongation


**Crop height, determined by extension of the last five or six internodes, is a reflection of variety and growing conditions.**

## Key facts:

- Crop height is measured to the base of the last fully emerged leaf blade, or to the base of the ear – the ear adds a further 10 cm
- Crop height depends on extension of the five or six internodes forming the stem
- Crop height is affected mainly by variety, sowing date and plant growth regulator (PGR) use
- Crop height is only one of several determinants of lodging risk in wheat

## The extended stem

**B** 5 internodes measuring 69 cm



Height before stem extension is related to leaf sheath length and reaches only 9 cm by GS31. Final crop height results primarily from internode extension.


Each internode starts to extend when the previous one has reached half its final length. The benchmark number of nodes in a wheat stem is 4, giving 5 internodes. Crops sown earlier can have an extra internode and may be taller.

The benchmark final height is 69 cm, with a full PGR programme. Without PGR applications, crops can be as much as 20 cm taller.

By flag leaf emergence, stem height is only 34 cm (50% of final height). Stems extend to 53 cm by full ear emergence (GS59) and reach their final height around the time of flowering (GS61).

Stem height does not reflect stem reserves or drought tolerance as taller stems have a greater proportion of structural materials that cannot be remobilised.

## Increase in crop height



## Plant breeding and control of stem extension

Over recent decades plant breeders have introduced improved, shorter varieties containing reduced height ('Rht') and other dwarfing genes. In the AHDB Recommended Lists, height is measured to the top of the ear.

## Agronomy and stem extension

The benchmark heights are for crops receiving applications of PGR during both early and late stem extension. Early sowing, high N residues, or lack of PGR contribute to tallness. The main effect of fertiliser N is to extend the penultimate internode and peduncle.

## Height and lodging

In tall crops, the aerial parts of the plant impose a large force on the stem base and root system, but variation in crop height is only a minor contributor to lodging risk; other factors are weight distribution along the shoot, root anchorage and stem strength. All three components of lodging risk can be altered by choice of variety and husbandry – see **Avoiding lodging in winter wheat – practical guidelines (2005)**.

The earlier that lodging occurs during grain filling, the greater the yield loss. Lodging can also adversely affect quality characteristics such as Hagberg falling number and specific weight.

# Stem carbohydrate storage

**Stem reserves contribute 20–50% of grain yield.**

## Key facts:


- Carbohydrate reserves are measured by weighing and analysing the sugar content of stems – they are the major part of the dry matter redistributed from stems and leaves during grain filling; protein is also redistributed
- Soluble carbohydrate stem reserves reach a maximum of 2.7 t/ha between late booting and early grain filling
- Variety and growing conditions both influence stem reserves by up to 2 t/ha
- Grain filling normally depends on stem reserves and ongoing photosynthesis. By harvest, almost all stem reserves have been relocated to the grain or lost through respiration

## Accumulation of stem reserves – GS31 until early grain filling

More assimilate is produced during the Construction Phase than is needed for structural tissues. The surplus is stored in stems as fructan sugar, mainly in the pith of upper internodes. About 25% is in the peduncle, 30% in the penultimate internode and 45% in lower internodes.

Stem storage capacity is set by stem number and structure. Maximum capacity can be reached by late booting (GS47). Reserves may be utilised during temporary shortages caused by factors such as dull weather before and after flowering. Reserves fluctuate with growing conditions from booting to early grain filling; they then decrease.

### Accumulation of stem reserves from GS31 to early grain filling



## Maximum stem storage

**B 2.7 t/ha soluble carbohydrate**

The benchmark amount of stem reserves at flowering is 2.3 t/ha. An additional 0.4 t/ha accumulates by early grain filling. Sometimes, the maximum is reached before flowering.

Both variety and growing conditions can affect stem reserves by about 2 t/ha. Some crops can store more than 4 t/ha. Varietal differences may be due to both total stem weight and per cent soluble material. Recent UK varieties have 25–35% of stem biomass as soluble reserves. Stem height does not necessarily indicate larger stem reserves. Environmental differences in stem reserves commonly relate to differences in stem number. Soluble stem carbohydrate is not affected by PGR applications.


## Dry matter redistribution during grain filling

**B 3.1 t/ha soluble carbohydrate**

Redistribution of soluble reserves lasts from 26 days after flowering to the end of grain filling. The loss in straw dry matter between flowering and harvest is 3.1 t/ha comprising 1.9 t/ha soluble stem reserves, and protein from leaves and stems.

Reserves contribute significantly to yield under all post-flowering conditions. Varieties with highest yield potential tend to be those that accumulate greatest amounts of stem soluble carbohydrate. Stem reserves contribute similar amounts of assimilate in stressed or unstressed crops. However, because yields are reduced in stressed crops, reserves contribute a higher proportion.

### Soluble stem dry matter data for a range of varieties



# Ear formation

**Capacity for grain filling is set by grain number per unit area and the storage capacity of each grain.**

## Key facts:

- Ear weight increases rapidly during booting. Ear weight at flowering is the same as the weight of chaff at harvest, and relates closely to final grain number
- Grain number per ear is largely controlled by survival of flower initials (florets) while the last leaves and ear are emerging
- Grain number per ear is an important yield component
- Ear weight at flowering can indicate grain numbers and hence storage capacity of the ear


## Ear development

**B 20 days from flag leaf to ear emergence**

Ears are initiated during the Foundation Phase and spikelet initiation is completed as stem extension starts (GS31). Floret initiation and development then proceed until flowering. The number of potentially fertile florets depends on assimilate supplies to the ear, particularly during booting, and is affected by shoot numbers. At shoot numbers of over 400/m<sup>2</sup>, mutual shading results in fewer grains in each ear.

Weather conditions affect ear development, especially during booting and ear emergence. Cooler bright conditions in the one or two weeks before flowering can prolong or enhance the ear formation period and increase grain number per ear. However, inclement weather at flowering, such as heavy rain, heat or drought, occasionally impairs pollination and reduces the number of fertilised florets.

### Structure of a spikelet



## What does it mean?

In wheat, yield variation due to region, soil type and early crop management is strongly related to grain number, rather than weight of each grain. Severe disease or drought can significantly reduce grain size.

## Grain number


**B 48 grains/ear – 22,000 grains/m<sup>2</sup>**

The benchmark for grain number per ear is 48. With 460 surviving fertile shoots/m<sup>2</sup>, this gives 22,000 grains/m<sup>2</sup>.

Varieties with smaller culm leaves tend to have more shoots/m<sup>2</sup> and fewer grains per ear. Modern breadmaking varieties tend to have fewer grains/ear than other varieties.

Some insects, eg orange blossom midge, can reduce grain numbers and feed on developing grains.

### Grain numbers compensate above 400 shoots/m<sup>2</sup>



Source: AHDB Cereals & Oilseeds seed rate trials

## Ear weight at flowering

**B 420 mg/ear**

By flowering the benchmark ear weight is 420 mg/ear, comprising glumes, florets and rachis. These components remain of similar weight throughout grain filling and become the chaff at harvest, so ear weight at flowering multiplied by fertile ear number per unit area can indicate yield potential.

Photosynthesis in the ear contributes significantly to grain growth. The benchmark final ear weight at harvest is 2.6 g, of which 17% is chaff.

# Grain filling and ripening

**Grain filling depends on ear and leaf photosynthesis, as well as redistribution of stem reserves.**

## Key facts:

- Grain filling starts when flowering is complete and continues until grain reaches about 45% moisture
- After flowering, grains swell (largely by water uptake); rapid dry weight growth continues with starch and protein deposition in expanded grain cells – these are supplied by both current photosynthesis and redistribution of reserves
- Where green canopy persists to the end of grain growth, stem reserves are less important
- Ripening and moisture loss continue after grain filling until the grain is dry enough to harvest

## Grain filling

**B** 43 mg dry matter per grain in 45 days until 26 July

Grain filling determines final dry grain weight. This final stage in yield formation influences grain appearance and specific weight.

Grains accumulate more water than dry matter for about four weeks after flowering, when water content is at its maximum. Water enables cells first to divide, then expand. Dry matter accumulation accelerates as water uptake stops.

Suboptimal photosynthesis during the first two or three weeks of grain growth will reduce cell number and potential weight of each grain.

Grain filling depends on the capacity of both 'sink' (ie all grains in ear) and the 'source' (ie materials from photosynthesis and reserves).

Where 'source' does not satisfy 'sink', eg due to late drought or disease, grains will be inadequately filled and, after ripening, may appear shrivelled.

Where 'sink' is too small to store all of the assimilate from the 'source', grains will be plump, and the adverse effects of drought or disease on yield are reduced.

The benchmark period from flowering until maximum dry weight (the grain filling period) is 45 days but it varies considerably; it can be just 28 days in severe drought conditions. The benchmark weight per grain is 43 mg dry matter, which equates to a 'thousand grain weight' of 50 g at 15% moisture. Varietal differences in average grain weight are shown in the AHDB Recommended List.

## Canopy survival during grain filling

**B** GAI <2 a day after maximum grain weight


Benchmark canopies lose most greenness just after grain weight reaches its maximum. High N uptake, fungicide use or cool, moist weather all tend to delay senescence. Crops at northerly latitudes generally have high grain weight because cool temperatures prolong grain filling. Hot weather reduces grain weight by shortening the period of grain growth, even in bright and moist conditions. Grain weight is also reduced by leaf or root disease, pest infestation or early lodging.

## Ripening

**B** 45% to 20% moisture content in two weeks

After filling, moisture content provides the best indication of ripening until grains are dry enough to harvest. On average, grain takes about two weeks to dry from 45% to 20% moisture. Frequent rain re-wets grain and slows moisture loss, especially at low moisture contents. Lodged crops dry slowly.

### Grain filling continues until moisture content reaches 45%



## What does it mean?

No further grain filling occurs once grain moisture is below 45%, so yield cannot be affected by any treatment applied at or after this stage.

▷ Consider using a desiccant (approved for use at less than 30% moisture, GS91) if harvest timing is threatened by inclement weather.

# Yield, quality and protein deposition

**Marketable yield depends on number of ears, grains in each ear and weight of each grain.**

## Key facts:

- Grain dry weight usually constitutes about half of final crop dry weight
- Grain yield is measured as the weight of grain recovered from an area of crop, corrected to 15% moisture content
- Specific weight partly depends upon grain size, also on grain density and grain packing characteristics
- Hagberg falling number reflects the gelling properties of flour made from whole grain
- Grain protein is related directly to grain N: it is increased by protein deposition and diluted by other grain growth
- Grain protein levels acceptable for breadmaking are more consistently achieved in second, than in first, wheat crops

## Grain yield

**B 11.0 t/ha at 15% moisture**

Grain is the principal product of crop growth, especially in June and July. Yields therefore depend on the state of the crop leading into this period and then on growing conditions during this period.

In experiments, yields may be determined by combine harvester or by hand-harvesting crop from a quadrat of specified land area. The benchmark crop yielded 11.0 t/ha by both methods, when corrected to 15% moisture. Yields vary with variety (see AHDB Recommended List) and growing conditions, as affected by soil, weather and husbandry.

Estimating grain yield:

$$\begin{aligned} \text{Estimated yield} &= \text{Ears/m}^2 \\ &\times \text{Grains per ear} \\ &\times \text{Individual grain weight (g)} \end{aligned}$$

Individual grain weight can be derived from the thousand grain weight.

## Yield components

At 15% moisture:

**B 460 ears/m<sup>2</sup>**   **B 48 grains/ear**   **B 50 mg/grain**

Ear number/m<sup>2</sup>, grains per ear and grain weight are key yield-determining attributes at harvest. Their values are related to the success of different growth phases:

- ear number reflects growth from the start of tillering to flag leaf appearance (GS39)
- grain number per ear reflects growth from GS39 to flowering (GS61)
- individual weight per grain reflects growth after flowering


Each phase partially compensates for the outcome of earlier phases. A crop with a sparse shoot density tends to produce more grains per ear and heavier grains than a thick crop.

## Final distribution of dry matter

**B 18.4 t/ha dry matter with 51% Harvest Index**

Harvest Index is the ratio between grain yield on a dry basis and the total crop dry weight at harvest. This ratio is variable.

### Final dry matter distribution



The benchmark dry matter of the harvest-ripe crop is 18.4 t/ha, of which 9.4 t/ha is grain (equivalent to 11.0 t/ha at 15% moisture).

The remaining 9t/ha of dry matter includes straw, chaff and stubble but only about half of this can be baled as straw, even when the height of the combine cut is low. Much variability in crop dry weight comes through variable production of non-harvestable material. Plant breeders have been most successful in increasing harvest index; growers have been most successful in increasing total crop dry matter.





## Specific weight

**B** 78 kg/hl

Crops with a large grain weight tend to have high specific weight. This is also influenced by:

- the density of individual grains
- the range of grain sizes
- characteristics of grain surfaces that affect packing

Specific weight indicates the weight of grain that can be loaded onto lorries or ships. A typical requirement for milling or export is 76 kg/hl; samples below this standard are likely to incur price penalties or even exclusion from the intended market. Late harvesting can reduce specific weight through weathering.

## Hagberg falling number

**B** 270 seconds

Hagberg falling number is a measure of the of hot paste viscosity, expressed as the number of seconds (s) taken for a plunger to fall through a wholemeal water suspension. Lower viscosity results from starch breakdown by alpha-amylase. This may form during or after ripening.

The minimum value possible is 60 s. The minimum value required for breadmaking is 250 s and for soft wheat for export is 220 s.

Enzyme activity is mainly associated with initiation of germination and, hence, with sprouting. Alpha-amylase may also form in cool, wet weather during ripening, even in the absence of visible sprouting. Green grains on late tillers, or grains damaged by orange blossom midge also have high alpha-amylase.

Effects of husbandry on Hagberg falling number tend to be small and inconsistent compared to the effects of variety and weather. Varietal differences in specific weight, dormancy and Hagberg falling number are indicated in the AHDB Recommended List. Group 1 wheats are generally classed as 'very high'.

## What does it mean?

By GS91, management can only protect against losses of yield or quality.

▷ Depending on intended market and likely risks, consider:

- combining as soon as the crop is ripe
- assessing Hagberg falling number pre-harvest on hand-threshed, air-dried grain
- harvesting at high moisture to optimise chances of high Hagberg falling numbers

## Nitrogen redistribution

**B** 158 kg/ha N

Most grain protein is formed from redistributed nitrogen, 158 kg/ha N coming mainly from stems, leaves and roots as they die. Only an additional 31 kg/ha of grain N comes from uptake after flowering.

The straw and chaff contains 90 kg/ha N at harvest – 32% of total crop N – making the N harvest index 68%.

## Protein deposition

**B** 189 kg/ha N or 1.1 t/ha protein

The weight of grain protein relates directly to the weight of N, with a ratio of 5.7 to 1. The benchmark amount of grain N is 189 kg/ha, equating to about 1.1 t/ha protein. Soil type, season and husbandry all influence grain protein deposition.

Generally, the later that N fertiliser is applied, the more grain N is increased. Applying urea as a spray when grain is milky ripe (GS75) has the largest effect on grain N, but normally has little effect on yield.

## Grain protein concentration

**B** 11.6% on a dry basis

High concentrations of grain protein can arise either from large N uptake or poor starch formation during grain filling. First wheats tend to have grain percentages below those of second wheats if both are fertilised optimally. Protein contents of first wheats are usually diluted by their greater yields.

Conversely, factors that reduce yield without affecting nitrogen transport to grain, for example, drought, early lodging or disease (eg take-all) may raise protein. Powdery mildew is an exception; it lowers protein percentage by interfering with nitrogen transport.

# Measurements

**To support husbandry decisions, commercial crops are best monitored using careful in-field observations and measurements.**

For quantitative assessments, at least four samples should be taken (one from each quarter of the field). In a variable crop more samples are required. Each sample point should be selected to represent the crop, but away from headlands, gateways and atypical patches.

## Development or 'Growth Stages'

The decimal (or 'Zadoks') growth stage code should be used (see page 4). Assessments are restricted to main shoots until flag leaves emerge. To determine the growth stage of the crop as a whole, quote the middle (median) stage from an odd number of plants arranged in order.


### For example:

If five plants were taken and their growth stages were:



then the crop would be considered to be at GS37.

For more information see: G49 Cereal growth stages – a guide for crop treatments (2009)



## Plant populations

Take at least four samples (more if variable) across the field. Throw down a quadrat (a square frame) and count the number of plants inside it. Divide this number by the area (m<sup>2</sup>) of the quadrat to get the number of plants/m<sup>2</sup>.

Alternatively, use a ruler and count the number of plants growing along a measured length of row. Divide this by the row width (m) and length (m) to determine the number of plants per square metre, as follows:

$$\frac{\text{No. of plants counted in the row}}{\text{Row length (m) X Row width (m)}} = \text{Plants/m}^2$$

### For example:

$$\frac{15 \text{ plants counted in the row} \times 10,000}{50 \text{ (cm)} \times 15 \text{ (cm)}} = 200 \text{ plants/m}^2$$

The same method can be used to work out shoot numbers/m<sup>2</sup>, counting shoots instead of plants.

## Crop canopy

Crop canopies are measured by their Green Area Index (GAI). This is the surface area of green material (one side only) divided by the area of ground it occupies. For example if, when all the green parts (leaves, shoots and ears) from 1 m<sup>2</sup> of a field were separated and laid out adjacent and flat, they covered 2 m<sup>2</sup>, the crop would have a GAI of 2 (see page 3). If they covered 4 m<sup>2</sup>, the crop would have a GAI of 4, and so on.

To avoid such a long and tedious process, it is better to 'get your eye in' and judge the GAI from the crop's appearance (see page 3).

Instruments and free apps are available which can measure the reflectance or light intercepted by a crop and translate it into GAI. A combination of experience and calibration using such machines can give the assessor a sufficiently accurate picture of canopy size to support crop management decisions.



## Dry weight

Dry weight is less easy to assess in the field than canopy size. However, measurements of ear dry weight can help in anticipating yield after the grain has reached 45% moisture. These can be made by placing a known number of ears in a domestic microwave at high power for 10–15 minutes, turning 2–3 times during drying. The resulting weight can be measured on a domestic digital balance in grammes, divided by the ear number, multiplied by 0.82 to correct for chaff, and multiplied by fertile shoot number per square metre to give the estimated grain yield (g/m<sup>2</sup>) NB t/ha = g/m<sup>2</sup> ÷ 100.

# Glossary

**Anthesis:** In wheat, anthesis is normally recognised by appearance of pollen sacs (anthers) from florets within the ear. This signifies pollination. Anthesis is also known as flowering.

**Assimilate:** The product of the crop's synthetic processes, mainly photosynthesis. Measured as dry matter.

**Benchmark:** A defined measure of crop progress consistent with good final performance.

**Carbohydrates:** Synthesised entirely from carbon dioxide and water, these are mainly starch and cellulose which are not 'soluble' or mobile, and sugars, eg fructan, which dissolve in water, are mobile in the plant and are classed as 'soluble'.

**Coleoptile:** The first leaf structure to emerge from the seed at germination. It protects the first true leaves during emergence of the seedling. It has little chlorophyll but may give rise to tillers.

**Day degrees:** See thermal time.

**Dormancy:** A condition in which grains do not germinate in the presence of adequate moisture, temperature and air.

**Dry matter:** Crop constituents other than water, left after tissue has been dried. Often, 'total dry matter' refers to just the above ground parts of the crop.

Dry matter is measured by weighing crop material after drying in a forced-draught oven at 80°C until it reaches constant weight (for about a day).

**Floret:** The primary sub-component of a spikelet. Each floret can only bear one grain; while they retain this potential they are termed fertile florets.

**Frost heave:** Lifting of the soil surface, caused by freezing of moisture in the topsoil and expansion, often leading to stretching and breaking of roots and other sub-surface structures.

**Fructan:** A form of sugar, a polymer of fructose (the main component of soluble carbohydrate), used by wheat and other grass species as storage assimilate in stem tissues.

**GAI:** Green Area Index. The ratio between the total area of all green tissues, one side only, and the area of ground from which they came.

**Hagberg falling number:** A measure of the hot paste viscosity of a wholemeal suspension in water. In the laboratory a suspension of flour is heated in water for a fixed period. The time in seconds taken for a plunger to fall through the resultant gel is recorded as the 'Hagberg falling number'.

**Harvest Index:** The ratio between grain yield on a dry basis and the total crop dry weight at harvest.

**Internode:** The section of stem between two adjacent nodes.

**Leaf blade:** The upper portion of a leaf, from the tip to the ligule (junction with the sheath).

**Leaf sheath:** The basal portion of a leaf which encloses the stem and sheaths of younger leaves.

**Ligule:** A small structure at the junction of leaf sheath and leaf blade.

**Lodging:** Permanent displacement of a stem or stems from a vertical posture. Lodging can be considered as an event occurring within one day, although lodged stems may initially lean rather than lie horizontally.

**Main shoot:** The primary axis of the plant, on which the primary tillers are borne.

**Mean:** The average. The sum of all the values divided by the number of values.

**Median:** The middle value when all values are ranked by size. Medians may provide more robust summaries than *means* because they are not influenced by exceptional values.

**N:** Nitrogen.

**Node:** The point at which a leaf sheath is attached to a stem.

**Partitioning:** The division of dry matter between organs.

**Peduncle:** The topmost internode, between the flag leaf node and the base of the ear (the collar).

**PGR:** Plant growth regulator. The 'full PGR programme' used to grow the benchmark crops included chlormequat at the end of tillering and GS31, and Terpal at GS37–39.

**Photosynthesis:** Formation of carbohydrates by green tissues from absorbed carbon dioxide and water, driven by energy from sunlight.

**Phyllochron:** The interval in thermal time from emergence of one leaf tip on a shoot to emergence of the next. Phyllochron is the reciprocal of leaf emergence rate.

**Pollination:** Reception of pollen produced in the anthers and bearing the male genetic complement, by the stigma, leading to fertilisation of the ovum, bearing the female genetic complement. Fertilisation of wheat normally occurs within one floret, rather than between florets. (Flower parts are shown in a diagram on page 22).

**Rachis:** The portion of the stem within the ear (above the collar), bearing the spikelets.

**Respiration:** Degradation of sugars and the associated absorption of oxygen and emission of carbon dioxide (and water) to yield energy for crop metabolism.

**Ripening:** The changes that occur in the grain between completion of growth and maturity. These include drying, and development and loss of dormancy. Grain is considered 'ripe' when it is ready for harvest – at less than 20% moisture.

**Senescence:** Loss of greenness in photosynthetic tissues, normally brought about by ageing, but also by diseases or drought.

**Shoots:** All the axes of a plant with the potential to bear an ear. The main shoot and all tillers are included. Shoots retaining the potential to form grain are termed 'fertile shoots'.

**Soil stability:** The tendency for soil aggregates to retain their integrity when wetted and disturbed. It is measured by assessing how easily aggregates break up into fine particles.

**Specific weight:** Or bulk density, is the weight of grain (corrected for variation in moisture content) when packed into a standard container. It is expressed in kilograms per hectolitre (100 litres).

**Spikelet:** The primary sub-component of the ear. About 20 spikelets are borne on alternate sides of the ear stem or 'rachis', and there is one 'terminal spikelet'. Each spikelet is contained within two glumes, and consists of several fertile florets.

**Thermal time:** The sum of all daily temperatures (mean of maximum and minimum) above a base temperature below which the process in question stops. In the case of leaf development this is 0°C. Results are expressed in 'day degrees' (°C days).

**Tiller:** A side shoot. Thus 'tillers' do not include the main or primary shoot.

**Transpiration:** Loss of water vapour from a crop's green surfaces, mainly through leaf pores (stomata).

**Vernalisation:** A change in the physiological state of a plant from vegetative to reproductive brought about by a period of cold – can be applied to seeds or (in the case of wheat) to the young plant.

**Vigour:** Term used to describe the capacity of a seed, plant or organ to grow.

**Waterlogging:** Filling of soil pores with water to the extent that there is insufficient oxygen for normal root function.

## Acknowledgements

The previous editions of this guide were written by Professor Roger Sylvester-Bradley, Dr Pete Berry, Jonathan Blake, Dr Daniel Kindred and John Spink, ADAS; Dr Ian Bingham, SRUC; Dr Jim McVittie, AHDB; and Dr John Foulkes, University of Nottingham. It was edited by Dr Clive Edwards, AHDB and Geoff Dodgson, Chamberlain.

This edition was edited by Dr Emily Boys and Fiona Geary, AHDB.

AHDB is also grateful to many people who commented on draft versions of previous editions, including: Alan Bide and Steve Cook, Hampshire Arable Systems; Peter Kettlewell, Harper Adams University; Jemilah Bailey, Teri Gibbons, Dr Simon Hook, Professor Graham Jellis, Dr Clare Kelly and Dr Roger Williams, AHDB; Neil Watson, TAG; Keith Norman, Velcourt; and Andrew Cragg, farmer.

While the Agriculture and Horticulture Development Board seeks to ensure that the information contained within this document is accurate at the time of printing, no warranty is given in respect thereof and, to the maximum extent permitted by law, the Agriculture and Horticulture Development Board accepts no liability for loss, damage or injury howsoever caused (including that caused by negligence) or suffered directly or indirectly in relation to information and opinions contained in or omitted from this document.

Reference herein to trade names and proprietary products without stating that they are protected does not imply that they may be regarded as unprotected and thus free for general use. No endorsement of named products is intended, nor is any criticism implied of other alternative, but unnamed products.